

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All causes | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 816.5 | 685.2 | 29.9 | 16.8 | 80.1 | 102.1 | 193.5 | 427.0 | 910.3 | 2,164.6 | 5,275.1 | 13,823.5 |
| 2003..... | 841.9 | 700.0 | 31.5 | 17.0 | 81.5 | 103.6 | 201.6 | 433.2 | 940.9 | 2,255.0 | 5,463.1 | 14,593.3 |
| 2002..... | 847.3 | 695.0 | 31.2 | 17.4 | 81.4 | 103.6 | 202.9 | 430.1 | 952.4 | 2,314.7 | 5,556.9 | 14,828.3 |
| 2001..... | 848.5 | 683.4 | 33.3 | 17.3 | 80.7 | 105.2 | 203.6 | 428.9 | 964.6 | 2,353.3 | 5,582.4 | 15,112.8 |
| 2000..... | 854.0 | 736.7 | 32.4 | 18.0 | 79.9 | 101.4 | 198.9 | 425.6 | 992.2 | 2,399.1 | 5,666.5 | 15,524.4 |
| 1999..... | 857.0 | 736.0 | 34.2 | 18.6 | 79.3 | 102.2 | 198.0 | 418.2 | 1,005.0 | 2,457.3 | 5,714.5 | 15,554.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 817.6 | 753.7 | 32.4 | 19.2 | 114.7 | 139.5 | 243.6 | 543.5 | 1,128.8 | 2,644.8 | 6,394.3 | 15,031.1 |
| 2003..... | 840.3 | 777.4 | 35.1 | 19.8 | 116.5 | 141.4 | 255.0 | 552.2 | 1,165.5 | 2,771.7 | 6,641.8 | 15,794.0 |
| 2002..... | 846.6 | 761.5 | 35.2 | 20.0 | 117.3 | 142.2 | 257.5 | 547.5 | 1,184.0 | 2,855.3 | 6,760.5 | 16,254.5 |
| 2001..... | 846.4 | 749.8 | 37.0 | 19.8 | 117.0 | 143.7 | 259.6 | 545.1 | 1,192.7 | 2,911.5 | 6,833.0 | 16,744.8 |
| 2000..... | 853.0 | 806.5 | 35.9 | 20.9 | 114.9 | 138.6 | 255.2 | 542.8 | 1,230.7 | 2,979.6 | 6,972.6 | 17,501.4 |
| 1999..... | 859.2 | 805.0 | 37.9 | 21.5 | 113.1 | 139.7 | 254.9 | 533.1 | 1,252.0 | 3,073.7 | 7,083.3 | 17,597.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 815.4 | 613.4 | 27.3 | 14.3 | 43.6 | 63.5 | 143.5 | 314.3 | 707.4 | 1,761.4 | 4,521.8 | 13,280.3 |
| 2003..... | 843.4 | 619.1 | 27.8 | 14.0 | 44.4 | 64.6 | 148.5 | 318.4 | 732.7 | 1,823.0 | 4,675.5 | 14,062.5 |
| 2002..... | 848.0 | 625.3 | 27.0 | 14.7 | 43.7 | 64.0 | 148.8 | 316.9 | 738.0 | 1,864.7 | 4,757.9 | 14,209.6 |
| 2001..... | 850.4 | 613.9 | 29.5 | 14.6 | 42.6 | 66.0 | 148.2 | 316.8 | 754.0 | 1,890.8 | 4,760.5 | 14,429.9 |
| 2000..... | 855.0 | 663.4 | 28.7 | 15.0 | 43.1 | 63.5 | 143.2 | 312.5 | 772.2 | 1,921.2 | 4,814.7 | 14,719.2 |
| 1999..... | 854.9 | 663.6 | 30.3 | 15.6 | 43.7 | 64.1 | 141.8 | 307.6 | 777.6 | 1,952.3 | 4,825.4 | 14,731.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 296.2 | 586.9 | 27.3 | 14.4 | 76.5 | 78.2 | 138.7 | 313.5 | 684.0 | 1,607.5 | 3,926.7 | 9,483.8 |
| 2003..... | 305.8 | 610.1 | 30.2 | 15.7 | 77.1 | 83.9 | 144.9 | 327.8 | 714.3 | 1,685.6 | 4,093.5 | 10,431.5 |
| 2002..... | 302.2 | 592.7 | 29.8 | 15.5 | 77.2 | 79.3 | 145.8 | 321.3 | 719.8 | 1,734.1 | 4,143.1 | 10,679.0 |
| 2001..... | 306.8 | 572.7 | 30.6 | 14.7 | 75.6 | 84.5 | 154.8 | 334.4 | 722.2 | 1,776.0 | 4,262.6 | 11,821.8 |
| 2000..... | 303.8 | 596.3 | 29.6 | 15.7 | 72.8 | 84.6 | 157.6 | 330.2 | 744.4 | 1,803.8 | 4,335.9 | 11,822.9 |
| 1999..... | 305.7 | 583.8 | 30.9 | 15.6 | 71.3 | 85.5 | 158.2 | 333.2 | 752.5 | 1,853.0 | 4,356.3 | 12,177.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 321.1 | 636.5 | 30.2 | 16.7 | 115.3 | 109.2 | 184.2 | 417.8 | 874.0 | 1,994.3 | 4,791.6 | 9,932.8 |
| 2003..... | 330.7 | 665.5 | 33.8 | 18.4 | 112.9 | 118.1 | 189.8 | 436.4 | 920.8 | 2,115.7 | 4,948.5 | 11,029.2 |
| 2002..... | 328.7 | 644.0 | 34.2 | 17.4 | 114.4 | 112.5 | 192.5 | 423.4 | 937.4 | 2,193.4 | 5,043.5 | 11,674.1 |
| 2001..... | 332.9 | 624.4 | 33.8 | 16.6 | 111.5 | 118.0 | 208.5 | 443.9 | 923.9 | 2,242.6 | 5,258.0 | 12,888.3 |
| 2000..... | 331.3 | 637.1 | 31.5 | 17.9 | 107.7 | 120.2 | 211.0 | 439.0 | 965.7 | 2,287.9 | 5,395.3 | 13,086.2 |
| 1999..... | 332.6 | 623.4 | 32.9 | 17.8 | 104.1 | 120.6 | 215.1 | 444.4 | 974.8 | 2,368.9 | 5,379.2 | 13,485.9 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 269.7 | 535.1 | 24.3 | 12.0 | 31.8 | 40.9 | 88.2 | 208.3 | 511.3 | 1,297.2 | 3,329.6 | 9,253.0 |
| 2003..... | 279.3 | 552.4 | 26.3 | 12.8 | 35.5 | 43.0 | 95.5 | 219.1 | 527.9 | 1,342.9 | 3,506.4 | 10,128.1 |
| 2002..... | 274.0 | 539.1 | 25.3 | 13.5 | 34.1 | 40.0 | 94.9 | 219.8 | 524.3 | 1,368.7 | 3,526.4 | 10,186.0 |
| 2001..... | 279.0 | 518.9 | 27.2 | 12.7 | 33.7 | 45.2 | 97.0 | 226.7 | 543.0 | 1,408.0 | 3,589.8 | 11,300.5 |
| 2000..... | 274.6 | 553.6 | 27.5 | 13.4 | 31.7 | 43.4 | 100.5 | 223.8 | 548.4 | 1,423.2 | 3,624.5 | 11,202.8 |
| 1999..... | 277.2 | 542.3 | 28.7 | 13.3 | 32.9 | 44.9 | 97.4 | 224.9 | 555.8 | 1,448.8 | 3,675.7 | 11,547.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All causes | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 899.4 | 705.6 | 30.5 | 17.2 | 80.6 | 107.3 | 201.7 | 437.4 | 926.1 | 2,200.1 | 5,340.3 | 13,997.7 |
| 2003..... | 924.4 | 716.2 | 31.7 | 17.2 | 82.0 | 107.6 | 209.3 | 442.1 | 955.6 | 2,289.1 | 5,522.6 | 14,740.6 |
| 2002..... | 928.8 | 713.0 | 31.4 | 17.8 | 81.9 | 108.4 | 210.3 | 438.8 | 967.2 | 2,347.0 | 5,610.8 | 14,958.7 |
| 2001..... | 926.2 | 703.5 | 33.8 | 17.7 | 81.4 | 108.9 | 209.2 | 435.6 | 979.5 | 2,383.9 | 5,628.5 | 15,199.0 |
| 2000..... | 929.6 | 760.4 | 32.9 | 18.4 | 80.9 | 104.1 | 202.9 | 431.8 | 1,006.5 | 2,427.9 | 5,707.6 | 15,619.8 |
| 1999..... | 929.9 | 760.7 | 34.8 | 19.1 | 80.5 | 104.7 | 201.6 | 423.3 | 1,019.2 | 2,485.0 | 5,753.4 | 15,631.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 900.9 | 778.7 | 32.9 | 19.6 | 114.2 | 146.6 | 252.8 | 554.7 | 1,145.4 | 2,683.5 | 6,472.4 | 15,260.4 |
| 2003..... | 922.9 | 798.1 | 35.2 | 20.0 | 116.7 | 146.5 | 264.2 | 561.6 | 1,179.8 | 2,808.1 | 6,715.9 | 15,984.2 |
| 2002..... | 928.0 | 782.6 | 35.4 | 20.4 | 117.3 | 148.6 | 266.0 | 557.0 | 1,198.1 | 2,889.1 | 6,825.9 | 16,418.0 |
| 2001..... | 923.6 | 773.2 | 37.6 | 20.4 | 117.5 | 148.5 | 265.2 | 551.6 | 1,207.0 | 2,944.1 | 6,888.0 | 16,866.3 |
| 2000..... | 928.1 | 836.8 | 36.8 | 21.5 | 115.9 | 141.6 | 259.2 | 548.7 | 1,244.2 | 3,009.8 | 7,020.7 | 17,639.8 |
| 1999..... | 932.2 | 834.9 | 38.8 | 22.1 | 114.6 | 142.7 | 258.1 | 537.3 | 1,265.6 | 3,102.8 | 7,133.0 | 17,710.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 898.0 | 628.9 | 28.1 | 14.7 | 45.7 | 68.1 | 151.5 | 324.5 | 722.2 | 1,792.8 | 4,579.4 | 13,433.1 |
| 2003..... | 925.9 | 630.7 | 28.0 | 14.3 | 46.0 | 68.9 | 155.6 | 327.3 | 747.2 | 1,853.7 | 4,726.4 | 14,194.2 |
| 2002..... | 929.5 | 640.1 | 27.2 | 15.0 | 45.3 | 68.5 | 155.7 | 325.2 | 753.0 | 1,894.4 | 4,805.4 | 14,329.1 |
| 2001..... | 928.6 | 630.4 | 29.9 | 14.9 | 44.1 | 69.6 | 154.3 | 324.0 | 768.7 | 1,918.4 | 4,801.8 | 14,505.6 |
| 2000..... | 931.0 | 680.1 | 28.8 | 15.2 | 45.0 | 66.8 | 147.7 | 319.2 | 786.8 | 1,947.5 | 4,852.4 | 14,802.9 |
| 1999..... | 927.8 | 682.9 | 30.6 | 16.0 | 45.4 | 67.1 | 146.3 | 313.7 | 791.7 | 1,977.5 | 4,858.2 | 14,798.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 967.8 | 564.4 | 26.8 | 15.5 | 75.2 | 95.9 | 185.9 | 400.8 | 875.1 | 2,152.5 | 5,337.1 | 14,191.1 |
| 2003..... | 993.6 | 576.5 | 27.6 | 15.6 | 75.8 | 95.8 | 190.9 | 402.3 | 902.4 | 2,237.2 | 5,515.4 | 14,944.8 |
| 2002..... | 997.5 | 575.9 | 27.1 | 16.0 | 75.8 | 95.9 | 191.6 | 400.1 | 912.8 | 2,296.1 | 5,596.0 | 15,149.1 |
| 2001..... | 991.1 | 555.4 | 30.1 | 16.3 | 74.1 | 96.4 | 188.3 | 396.0 | 924.0 | 2,325.5 | 5,604.4 | 15,371.2 |
| 2000..... | 993.2 | 596.5 | 28.5 | 17.1 | 73.6 | 90.1 | 181.0 | 390.5 | 950.2 | 2,366.2 | 5,675.4 | 15,799.7 |
| 1999..... | 990.7 | 593.1 | 30.1 | 17.3 | 72.4 | 91.3 | 177.8 | 381.5 | 957.8 | 2,420.0 | 5,716.0 | 15,795.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 957.4 | 625.2 | 29.0 | 17.9 | 104.8 | 130.9 | 235.7 | 510.5 | 1,076.4 | 2,617.9 | 6,461.5 | 15,489.2 |
| 2003..... | 979.1 | 647.2 | 30.2 | 18.1 | 105.9 | 129.9 | 243.8 | 513.2 | 1,110.5 | 2,738.5 | 6,692.2 | 16,234.4 |
| 2002..... | 983.9 | 643.5 | 30.3 | 18.3 | 106.7 | 130.9 | 244.9 | 509.9 | 1,126.5 | 2,824.1 | 6,801.7 | 16,641.9 |
| 2001..... | 975.6 | 611.6 | 33.7 | 18.6 | 105.1 | 131.5 | 241.6 | 502.6 | 1,136.3 | 2,869.4 | 6,851.5 | 17,055.3 |
| 2000..... | 978.5 | 658.7 | 32.4 | 20.0 | 103.5 | 123.0 | 233.9 | 497.7 | 1,170.9 | 2,930.5 | 6,977.8 | 17,853.2 |
| 1999..... | 979.6 | 658.1 | 33.4 | 19.9 | 100.8 | 124.5 | 230.0 | 483.7 | 1,187.4 | 3,023.2 | 7,088.0 | 17,871.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 977.7 | 500.7 | 24.4 | 13.0 | 44.3 | 60.3 | 136.0 | 292.9 | 683.8 | 1,752.0 | 4,571.1 | 13,609.6 |
| 2003..... | 1,007.6 | 502.5 | 24.9 | 12.9 | 44.3 | 61.2 | 138.0 | 293.1 | 705.0 | 1,807.9 | 4,720.5 | 14,377.2 |
| 2002..... | 1,010.6 | 504.8 | 23.8 | 13.6 | 43.8 | 60.3 | 138.3 | 292.1 | 710.5 | 1,846.0 | 4,787.9 | 14,504.3 |
| 2001..... | 1,006.1 | 496.4 | 26.3 | 13.9 | 41.9 | 60.9 | 134.9 | 291.0 | 723.5 | 1,864.1 | 4,777.3 | 14,670.6 |
| 2000..... | 1,007.3 | 530.9 | 24.4 | 13.9 | 42.6 | 56.8 | 128.1 | 285.0 | 742.1 | 1,891.0 | 4,819.3 | 14,971.7 |
| 1999..... | 1,001.3 | 524.6 | 26.7 | 14.7 | 42.9 | 57.8 | 125.6 | 281.0 | 741.6 | 1,915.1 | 4,817.7 | 14,967.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All causes | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 768.8 | 1297.2 | 45.5 | 24.4 | 112.8 | 186.5 | 332.3 | 760.8 | 1,527.9 | 3,032.3 | 6,251.7 | 13,492.2 |
| 2003..... | 788.8 | 1290.0 | 48.0 | 23.8 | 116.6 | 188.9 | 354.2 | 788.3 | 1,578.4 | 3,158.7 | 6,433.0 | 14,107.7 |
| 2002..... | 792.8 | 1280.3 | 48.4 | 25.3 | 116.9 | 193.9 | 357.8 | 782.0 | 1,592.9 | 3,204.8 | 6,584.8 | 14,386.7 |
| 2001..... | 798.1 | 1354.4 | 49.5 | 24.0 | 121.4 | 191.5 | 365.8 | 786.9 | 1,602.5 | 3,300.7 | 6,645.3 | 14,660.8 |
| 2000..... | 805.5 | 1477.8 | 51.7 | 25.0 | 123.2 | 194.6 | 365.1 | 802.2 | 1,633.9 | 3,378.8 | 6,764.3 | 14,837.5 |
| 1999..... | 812.1 | 1510.2 | 54.6 | 27.5 | 125.5 | 191.1 | 376.9 | 796.7 | 1,681.7 | 3,454.5 | 6,805.4 | 14,839.2 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 818.7 | 1427.6 | 49.5 | 26.9 | 169.4 | 262.1 | 409.3 | 974.9 | 1,998.6 | 3,880.2 | 7,813.2 | 14,647.3 |
| 2003..... | 840.6 | 1427.7 | 55.1 | 27.9 | 176.5 | 266.6 | 440.0 | 1012.8 | 2,047.7 | 4,041.1 | 8,165.7 | 15,082.2 |
| 2002..... | 842.3 | 1369.3 | 55.8 | 29.8 | 177.4 | 274.3 | 445.7 | 1002.9 | 2,074.5 | 4,078.9 | 8,246.2 | 15,776.9 |
| 2001..... | 849.7 | 1484.2 | 55.1 | 28.6 | 186.2 | 267.9 | 456.3 | 1014.6 | 2,055.4 | 4,218.9 | 8,426.8 | 16,576.2 |
| 2000..... | 859.5 | 1623.2 | 56.1 | 29.1 | 187.2 | 270.2 | 462.7 | 1036.2 | 2,111.4 | 4,303.9 | 8,551.0 | 16,944.8 |
| 1999..... | 872.8 | 1650.8 | 61.5 | 33.1 | 191.4 | 267.9 | 479.5 | 1048.1 | 2,176.7 | 4,397.8 | 8,615.3 | 17,385.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 723.4 | 1161.6 | 41.3 | 21.8 | 55.3 | 117.2 | 264.1 | 577.0 | 1,149.9 | 2,427.9 | 5,370.3 | 13,035.6 |
| 2003..... | 741.6 | 1146.9 | 40.7 | 19.6 | 56.0 | 118.0 | 278.4 | 595.7 | 1,201.6 | 2,531.2 | 5,454.6 | 13,730.3 |
| 2002..... | 748.0 | 1187.5 | 40.8 | 20.7 | 56.1 | 121.3 | 280.1 | 592.7 | 1,206.3 | 2,584.8 | 5,644.9 | 13,855.4 |
| 2001..... | 751.2 | 1217.8 | 43.6 | 19.3 | 56.8 | 122.4 | 285.7 | 591.5 | 1,238.2 | 2,653.0 | 5,645.1 | 13,951.4 |
| 2000..... | 756.7 | 1327.7 | 47.1 | 20.7 | 60.1 | 126.5 | 278.8 | 601.2 | 1,249.4 | 2,728.0 | 5,758.6 | 14,054.9 |
| 1999..... | 757.3 | 1365.4 | 47.4 | 21.8 | 60.5 | 122.1 | 286.4 | 581.7 | 1,285.1 | 2,789.2 | 5,792.0 | 13,887.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| I. Certain infectious and parasitic diseases (A00-B99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 22.0 | 11.8 | 1.2 | 0.4 | 1.1 | 5.1 | 15.5 | 24.8 | 28.0 | 49.5 | 113.5 | 249.5 |
| 2003..... | 22.2 | 12.3 | 1.2 | 0.5 | 1.2 | 5.4 | 16.7 | 25.1 | 27.6 | 49.3 | 114.4 | 255.2 |
| 2002..... | 22.2 | 14.4 | 1.1 | 0.6 | 1.2 | 6.2 | 17.5 | 24.9 | 26.1 | 50.7 | 113.4 | 252.8 |
| 2001..... | 21.2 | 14.0 | 1.5 | 0.5 | 1.3 | 6.7 | 17.6 | 22.9 | 25.4 | 48.0 | 104.9 | 246.3 |
| 2000..... | 21.0 | 13.4 | 1.4 | 0.5 | 1.2 | 7.5 | 17.7 | 22.9 | 24.4 | 45.8 | 103.7 | 256.0 |
| 1999..... | 21.5 | 14.8 | 1.5 | 0.6 | 1.2 | 8.4 | 18.7 | 22.1 | 24.1 | 48.0 | 105.1 | 267.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 23.7 | 13.8 | 1.1 | 0.4 | 1.3 | 6.0 | 21.0 | 34.9 | 35.8 | 56.6 | 128.1 | 273.1 |
| 2003..... | 24.0 | 12.6 | 1.4 | 0.4 | 1.2 | 6.7 | 23.2 | 35.8 | 34.9 | 56.2 | 128.8 | 269.1 |
| 2002..... | 24.2 | 14.8 | 1.2 | 0.6 | 1.2 | 7.6 | 24.6 | 36.1 | 33.0 | 58.0 | 130.8 | 270.9 |
| 2001..... | 23.3 | 14.7 | 1.6 | 0.5 | 1.4 | 8.5 | 25.2 | 33.1 | 31.9 | 54.5 | 122.3 | 269.7 |
| 2000..... | 23.0 | 14.1 | 1.5 | 0.5 | 1.3 | 9.5 | 25.4 | 33.7 | 30.8 | 52.7 | 120.5 | 273.0 |
| 1999..... | 23.7 | 17.0 | 1.8 | 0.7 | 1.3 | 11.2 | 27.1 | 32.2 | 30.4 | 55.8 | 123.3 | 290.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 20.3 | 9.6 | 1.2 | 0.4 | 1.0 | 4.2 | 9.9 | 15.1 | 20.7 | 43.5 | 103.7 | 238.8 |
| 2003..... | 20.6 | 12.1 | 1.1 | 0.5 | 1.1 | 4.0 | 10.3 | 14.7 | 20.8 | 43.5 | 104.8 | 249.1 |
| 2002..... | 20.2 | 14.0 | 0.9 | 0.5 | 1.1 | 4.7 | 10.4 | 14.1 | 19.7 | 44.6 | 101.9 | 245.0 |
| 2001..... | 19.2 | 13.2 | 1.3 | 0.5 | 1.3 | 4.8 | 10.2 | 13.0 | 19.4 | 42.5 | 93.5 | 236.5 |
| 2000..... | 19.0 | 12.7 | 1.4 | 0.5 | 1.0 | 5.4 | 10.1 | 12.5 | 18.4 | 40.1 | 92.7 | 249.1 |
| 1999..... | 19.3 | 12.5 | 1.2 | 0.6 | 1.2 | 5.6 | 10.3 | 12.4 | 18.4 | 41.7 | 93.3 | 258.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 12.4 | 10.8 | 1.5 | 0.4 | 0.9 | 3.8 | 15.7 | 29.9 | 33.2 | 48.8 | 99.9 | 185.5 |
| 2003..... | 12.5 | 9.8 | 1.0 | 0.4 | 1.0 | 4.4 | 17.1 | 31.7 | 32.8 | 47.6 | 96.2 | 189.2 |
| 2002..... | 12.7 | 15.0 | 1.0 | 0.5 | 1.1 | 5.1 | 17.7 | 30.3 | 31.7 | 50.1 | 95.6 | 197.9 |
| 2001..... | 12.6 | 13.2 | 1.1 | 0.5 | 1.1 | 5.5 | 18.7 | 29.7 | 32.2 | 48.1 | 93.1 | 213.2 |
| 2000..... | 12.5 | 14.3 | 1.4 | 0.5 | 1.0 | 6.9 | 19.9 | 30.1 | 31.0 | 44.6 | 93.6 | 185.1 |
| 1999..... | 13.3 | 13.0 | 1.3 | 0.6 | 1.0 | 7.9 | 22.7 | 28.5 | 29.5 | 53.6 | 97.7 | 226.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 15.1 | 12.5 | 1.6 | * | 1.2 | 5.1 | 22.3 | 44.3 | 44.9 | 57.3 | 109.4 | 197.5 |
| 2003..... | 15.5 | 9.5 | * | * | 1.1 | 5.8 | 24.0 | 49.6 | 44.7 | 55.9 | 103.2 | 209.3 |
| 2002..... | 15.7 | 16.4 | * | 0.6 | 1.1 | 6.9 | 26.0 | 45.9 | 42.3 | 56.2 | 107.5 | 214.2 |
| 2001..... | 15.6 | 13.4 | * | 0.7 | 1.4 | 7.3 | 26.9 | 45.4 | 45.4 | 56.1 | 110.9 | 205.7 |
| 2000..... | 16.0 | 14.4 | 1.6 | * | 1.3 | 9.5 | 29.7 | 45.6 | 42.8 | 53.6 | 110.1 | 209.6 |
| 1999..... | 16.7 | 14.7 | 1.7 | * | 1.1 | 10.8 | 33.8 | 42.5 | 40.9 | 63.7 | 110.0 | 230.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 9.5 | 9.0 | 1.4 | * | 0.6 | 2.1 | 8.4 | 15.4 | 22.6 | 42.0 | 93.4 | 179.3 |
| 2003..... | 9.4 | 10.1 | * | * | 0.8 | 2.6 | 9.4 | 13.8 | 22.1 | 40.9 | 91.4 | 178.9 |
| 2002..... | 9.5 | 13.5 | * | * | 1.1 | 3.0 | 8.7 | 14.7 | 22.2 | 45.2 | 87.4 | 189.8 |
| 2001..... | 9.3 | 13.0 | * | * | 0.7 | 3.4 | 9.8 | 14.2 | 20.6 | 41.7 | 81.0 | 216.9 |
| 2000..... | 8.9 | 14.1 | * | * | 0.8 | 3.8 | 9.4 | 14.9 | 20.5 | 37.5 | 82.5 | 173.1 |
| 1999..... | 9.8 | 11.1 | * | * | 1.0 | 4.5 | 10.9 | 14.8 | 19.5 | 45.6 | 89.5 | 224.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| I. Certain infectious and parasitic diseases (A00-B99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 23.5 | 11.8 | 1.1 | 0.4 | 1.2 | 5.4 | 15.3 | 24.1 | 27.4 | 49.4 | 114.0 | 251.9 |
| 2003..... | 23.7 | 13.0 | 1.3 | 0.5 | 1.2 | 5.6 | 16.5 | 24.2 | 27.0 | 49.2 | 115.2 | 257.5 |
| 2002..... | 23.5 | 14.1 | 1.1 | 0.6 | 1.1 | 6.3 | 17.2 | 24.1 | 25.5 | 50.5 | 114.0 | 254.3 |
| 2001..... | 22.3 | 14.1 | 1.6 | 0.5 | 1.4 | 6.9 | 17.3 | 22.0 | 24.6 | 47.7 | 105.2 | 246.9 |
| 2000..... | 22.0 | 13.0 | 1.5 | 0.5 | 1.2 | 7.5 | 17.2 | 22.0 | 23.7 | 45.6 | 103.7 | 257.7 |
| 1999..... | 22.5 | 15.2 | 1.6 | 0.6 | 1.3 | 8.5 | 17.9 | 21.2 | 23.5 | 47.5 | 105.0 | 268.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 25.1 | 13.8 | 1.0 | 0.4 | 1.3 | 6.2 | 20.6 | 33.6 | 34.9 | 56.3 | 128.9 | 276.4 |
| 2003..... | 25.3 | 13.4 | 1.5 | 0.4 | 1.2 | 6.8 | 22.9 | 34.1 | 34.0 | 55.9 | 129.9 | 271.4 |
| 2002..... | 25.5 | 14.1 | 1.2 | 0.6 | 1.2 | 7.7 | 24.1 | 34.7 | 32.0 | 57.8 | 131.5 | 272.8 |
| 2001..... | 24.3 | 15.0 | 1.8 | 0.5 | 1.4 | 8.7 | 24.6 | 31.5 | 30.6 | 54.1 | 122.4 | 272.0 |
| 2000..... | 23.9 | 13.9 | 1.5 | 0.5 | 1.3 | 9.4 | 24.5 | 32.2 | 29.7 | 52.2 | 120.2 | 274.5 |
| 1999..... | 24.5 | 17.5 | 1.8 | 0.7 | 1.3 | 11.2 | 25.9 | 30.7 | 29.3 | 55.1 | 123.3 | 291.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 21.9 | 9.8 | 1.1 | 0.4 | 1.1 | 4.6 | 10.1 | 15.0 | 20.5 | 43.5 | 104.0 | 240.9 |
| 2003..... | 22.1 | 12.6 | 1.1 | 0.5 | 1.2 | 4.3 | 10.4 | 14.7 | 20.5 | 43.5 | 105.3 | 251.4 |
| 2002..... | 21.7 | 14.1 | 1.0 | 0.6 | 1.1 | 5.0 | 10.5 | 13.9 | 19.4 | 44.5 | 102.4 | 246.3 |
| 2001..... | 20.5 | 13.1 | 1.3 | 0.5 | 1.4 | 5.0 | 10.1 | 12.8 | 19.2 | 42.5 | 93.9 | 236.5 |
| 2000..... | 20.3 | 12.2 | 1.4 | 0.5 | 1.1 | 5.7 | 10.1 | 12.2 | 18.1 | 40.1 | 92.9 | 250.9 |
| 1999..... | 20.5 | 12.8 | 1.3 | 0.6 | 1.3 | 5.7 | 10.1 | 12.1 | 18.2 | 41.3 | 93.2 | 259.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 21.2 | 9.5 | 0.9 | 0.3 | 0.7 | 2.4 | 9.0 | 16.8 | 21.5 | 43.6 | 108.1 | 244.1 |
| 2003..... | 21.2 | 8.7 | 1.0 | 0.3 | 0.8 | 2.7 | 9.6 | 16.8 | 21.0 | 43.4 | 109.0 | 246.9 |
| 2002..... | 20.8 | 11.4 | 0.9 | 0.5 | 0.7 | 2.9 | 9.9 | 16.4 | 19.6 | 44.4 | 107.1 | 243.2 |
| 2001..... | 19.5 | 9.5 | 1.4 | 0.4 | 0.8 | 3.1 | 9.9 | 14.6 | 18.9 | 41.9 | 98.7 | 235.4 |
| 2000..... | 19.0 | 9.2 | 1.1 | 0.4 | 0.7 | 3.3 | 10.0 | 14.4 | 18.0 | 39.7 | 95.7 | 244.1 |
| 1999..... | 19.2 | 11.3 | 1.3 | 0.5 | 0.8 | 4.3 | 10.2 | 13.7 | 17.8 | 40.9 | 96.6 | 251.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 21.8 | 11.2 | 0.9 | 0.3 | 0.7 | 3.0 | 12.9 | 23.8 | 26.6 | 49.2 | 121.9 | 269.4 |
| 2003..... | 21.7 | 8.2 | 1.1 | 0.3 | 0.8 | 3.5 | 14.2 | 24.0 | 25.7 | 48.6 | 122.7 | 261.2 |
| 2002..... | 21.6 | 11.2 | 1.1 | 0.5 | 0.8 | 4.0 | 14.8 | 23.8 | 24.3 | 50.4 | 122.3 | 261.4 |
| 2001..... | 20.3 | 10.2 | 1.6 | 0.4 | 0.8 | 4.4 | 15.3 | 21.2 | 22.9 | 46.9 | 114.8 | 258.2 |
| 2000..... | 19.6 | 9.8 | 1.1 | 0.4 | 0.9 | 4.7 | 15.0 | 21.2 | 22.0 | 45.3 | 110.4 | 259.6 |
| 1999..... | 20.0 | 13.8 | 1.5 | 0.5 | 0.9 | 6.2 | 15.7 | 19.8 | 21.3 | 47.1 | 113.6 | 271.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 20.5 | 7.7 | 1.0 | 0.3 | 0.6 | 1.7 | 5.1 | 10.0 | 16.7 | 38.8 | 98.7 | 232.7 |
| 2003..... | 20.7 | 9.2 | 0.8 | 0.4 | 0.7 | 1.8 | 5.0 | 9.7 | 16.6 | 38.9 | 99.8 | 240.6 |
| 2002..... | 20.0 | 11.6 | 0.7 | 0.4 | 0.7 | 1.9 | 5.0 | 9.1 | 15.1 | 39.2 | 97.0 | 235.3 |
| 2001..... | 18.6 | 8.8 | 1.2 | 0.4 | 0.8 | 1.8 | 4.5 | 8.1 | 15.1 | 37.6 | 88.0 | 225.9 |
| 2000..... | 18.3 | 8.5 | 1.2 | 0.4 | 0.5 | 2.0 | 5.0 | 7.6 | 14.2 | 35.0 | 86.0 | 237.8 |
| 1999..... | 18.4 | 8.7 | 1.1 | 0.4 | 0.6 | 2.3 | 4.6 | 7.7 | 14.4 | 35.8 | 85.4 | 244.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| I. Certain infectious and parasitic diseases (A00-B99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 41.7 | 21.5 | 1.5 | 0.7 | 3.5 | 21.2 | 54.4 | 76.6 | 79.2 | 103.8 | 196.3 | 379.9 |
| 2003..... | 43.0 | 28.8 | 2.3 | 0.9 | 3.2 | 21.1 | 59.6 | 78.7 | 79.2 | 103.6 | 197.7 | 419.1 |
| 2002..... | 44.2 | 24.6 | 1.7 | 0.9 | 3.1 | 25.1 | 63.3 | 81.2 | 75.8 | 108.8 | 206.1 | 417.4 |
| 2001..... | 43.6 | 32.7 | 2.3 | 0.9 | 4.2 | 27.3 | 64.0 | 77.7 | 74.8 | 103.8 | 191.7 | 409.2 |
| 2000..... | 44.3 | 30.7 | 2.5 | 0.8 | 3.5 | 30.9 | 63.9 | 80.4 | 70.5 | 103.9 | 208.1 | 439.8 |
| 1999..... | 45.7 | 32.2 | 2.6 | 1.3 | 3.6 | 32.3 | 68.2 | 79.1 | 71.6 | 111.2 | 209.5 | 470.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 49.3 | 23.5 | * | * | 3.7 | 23.6 | 71.4 | 109.9 | 114.2 | 131.1 | 237.6 | 413.6 |
| 2003..... | 51.5 | 32.9 | 2.5 | 0.9 | 3.0 | 25.4 | 80.4 | 114.4 | 113.2 | 132.6 | 238.6 | 461.3 |
| 2002..... | 53.1 | 24.8 | 1.6 | 0.7 | 3.0 | 29.2 | 86.5 | 120.9 | 104.3 | 137.5 | 259.1 | 454.0 |
| 2001..... | 52.7 | 35.4 | 2.8 | 0.9 | 4.1 | 33.9 | 88.4 | 114.4 | 103.3 | 129.5 | 233.5 | 478.2 |
| 2000..... | 54.0 | 32.8 | 2.9 | 0.8 | 3.6 | 37.5 | 89.8 | 122.4 | 99.3 | 128.5 | 259.2 | 488.5 |
| 1999..... | 56.0 | 34.3 | 3.3 | 1.4 | 3.2 | 41.4 | 96.5 | 121.0 | 101.0 | 140.5 | 254.2 | 536.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 34.7 | 19.3 | * | 0.8 | 3.3 | 18.9 | 39.3 | 48.1 | 51.1 | 84.3 | 172.9 | 366.6 |
| 2003..... | 35.2 | 24.5 | 2.0 | 1.0 | 3.4 | 17.2 | 41.2 | 48.1 | 51.8 | 83.1 | 174.5 | 402.8 |
| 2002..... | 36.1 | 24.3 | 1.8 | 1.0 | 3.3 | 21.3 | 42.7 | 47.2 | 52.8 | 88.4 | 176.1 | 403.4 |
| 2001..... | 35.3 | 29.9 | 1.8 | 0.9 | 4.4 | 21.3 | 42.4 | 46.3 | 51.9 | 85.7 | 168.3 | 383.7 |
| 2000..... | 35.5 | 28.5 | 2.2 | 0.8 | 3.5 | 24.9 | 40.9 | 44.3 | 47.3 | 86.7 | 179.4 | 421.7 |
| 1999..... | 36.4 | 30.0 | 2.0 | 1.1 | 4.1 | 24.1 | 43.1 | 43.4 | 47.9 | 90.4 | 184.5 | 446.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intestinal infectious diseases (A00-A09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.4 | 13.2 | 35.1 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 9.6 | 25.4 |
| 2002..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 7.1 | 21.5 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.7 | 13.1 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.9 | 12.0 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 3.4 | 10.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 3.3 | 13.2 | 32.7 |
| 2003..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.6 | 2.3 | 9.6 | 21.9 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.6 | 2.4 | 7.3 | 21.1 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.8 | 12.1 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.2 | 11.6 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 1.0 | 3.7 | 9.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | * | 0.2 | 0.8 | 3.6 | 13.1 | 36.1 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.7 | 9.5 | 27.0 |
| 2002..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 7.1 | 21.7 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 4.6 | 13.5 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 1.2 | 3.7 | 12.2 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.0 | 3.2 | 11.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.6 | 13.9 | 37.5 |
| 2003..... | 1.4 | * | * | * | * | * | * | 0.2 | 0.7 | 2.7 | 10.1 | 26.8 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.5 | 2.2 | 7.6 | 22.6 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.9 | 13.8 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.2 | 12.5 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.1 | 3.5 | 11.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.9 | 3.4 | 13.9 | 34.7 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 10.2 | 23.2 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.5 | 2.4 | 7.7 | 22.2 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.4 | 4.9 | 12.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.4 | 4.4 | 12.1 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 1.1 | 3.8 | 10.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 0.2 | 0.9 | 3.8 | 13.9 | 38.7 |
| 2003..... | 1.8 | * | * | * | * | * | * | 0.1 | 0.7 | 2.9 | 10.0 | 28.4 |
| 2002..... | 1.4 | * | * | * | * | * | * | 0.2 | 0.5 | 2.0 | 7.5 | 22.7 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.3 | 1.3 | 4.8 | 14.2 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.0 | 12.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 1.1 | 3.3 | 11.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other intestinal infectious diseases (A01-A08) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.3 | 12.8 | 34.3 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 9.3 | 24.6 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 2.0 | 6.9 | 20.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.5 | 12.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 3.8 | 11.4 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.1 | 10.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.2 | 12.9 | 32.2 |
| 2003..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.6 | 2.3 | 9.3 | 21.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 2.2 | 7.0 | 20.3 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.6 | 11.8 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.0 | 10.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.9 | 3.5 | 9.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.8 | 3.5 | 12.7 | 35.2 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.1 | 0.7 | 2.6 | 9.3 | 26.1 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 6.8 | 20.8 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 4.4 | 13.0 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 1.1 | 3.6 | 11.7 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 0.9 | 2.8 | 10.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.5 | 13.5 | 36.6 |
| 2003..... | 1.4 | * | * | * | * | * | * | 0.2 | 0.7 | 2.6 | 9.8 | 25.8 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 | 7.4 | 21.6 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.7 | 13.3 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 4.0 | 11.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.9 | 3.2 | 10.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.9 | 3.3 | 13.6 | 34.2 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 9.8 | 22.4 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 2.3 | 7.5 | 21.3 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.4 | 4.7 | 12.3 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.2 | 11.1 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.9 | 3.7 | 9.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 0.2 | 0.9 | 3.7 | 13.5 | 37.7 |
| 2003..... | 1.7 | * | * | * | * | * | * | 0.1 | 0.7 | 2.8 | 9.7 | 27.4 |
| 2002..... | 1.3 | * | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.3 | 21.7 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.7 | 13.7 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | 0.2 | 1.2 | 3.9 | 12.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 1.0 | 2.9 | 10.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intestinal infections due to other specified organisms (A04,A07-A08) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 3.3 | 12.8 | 34.2 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.6 | 2.4 | 9.2 | 24.3 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.5 | 2.0 | 6.8 | 20.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.4 | 12.4 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.7 | 11.3 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.9 | 3.0 | 9.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.9 | 3.1 | 12.8 | 31.9 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.6 | 2.3 | 9.2 | 20.6 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.5 | 2.2 | 6.9 | 20.2 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.5 | 11.7 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.2 | 3.9 | 10.5 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.4 | 9.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.8 | 3.4 | 12.7 | 35.2 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.1 | 0.7 | 2.6 | 9.2 | 26.0 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 6.8 | 20.7 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 4.3 | 12.7 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 1.1 | 3.5 | 11.5 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 0.9 | 2.7 | 10.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.5 | 13.5 | 36.5 |
| 2003..... | 1.3 | * | * | * | * | * | * | 0.1 | 0.6 | 2.6 | 9.7 | 25.6 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.1 | 0.5 | 2.1 | 7.3 | 21.6 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.5 | 13.1 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 3.9 | 11.7 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.9 | 3.1 | 10.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 0.2 | 0.9 | 3.3 | 13.5 | 34.0 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.5 | 2.4 | 9.7 | 21.9 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.5 | 2.3 | 7.3 | 21.2 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.5 | 12.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.2 | 4.1 | 11.0 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 0.9 | 3.6 | 9.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 0.2 | 0.9 | 3.7 | 13.5 | 37.6 |
| 2003..... | 1.7 | * | * | * | * | * | * | 0.1 | 0.7 | 2.8 | 9.7 | 27.2 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | 0.5 | 1.9 | 7.3 | 21.7 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.6 | 13.4 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | 0.2 | 1.1 | 3.8 | 11.9 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 0.9 | 2.8 | 10.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diarrhea and gastroenteritis of infectious origin (A09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 0.8 |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 0.9 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 0.9 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 0.7 |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 0.9 |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 0.9 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 0.9 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 0.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 0.8 |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 0.9 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 1.0 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 0.7 |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 0.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.0 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.0 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 0.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Tuberculosis (A16-A19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.7 | 1.1 | 2.2 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.6 | 2.2 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.8 | 2.5 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.5 | 2.7 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.9 | 2.5 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.9 | 3.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.0 | 1.3 | 3.5 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.4 | 0.5 | 0.8 | 2.1 | 3.0 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.3 | 3.8 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 1.9 | 4.3 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.4 | 0.5 | 1.0 | 2.6 | 4.3 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.3 | 2.6 | 6.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.4 | 1.0 | 1.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 1.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 1.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 2.1 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.7 | 1.4 | 1.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 1.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 1.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 1.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.2 | 2.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.5 | 2.4 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.7 | 2.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.4 | 2.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | * | 0.6 | 1.5 | 3.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.8 | 4.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.1 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 1.4 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.1 | 1.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.1 | 1.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Respiratory tuberculosis (A16) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 0.8 | 2.0 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.3 | 1.7 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.2 | 2.4 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.6 | 1.6 | 2.2 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.0 | 3.3 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.3 | 0.4 | 0.6 | 1.8 | 2.4 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.8 | 3.3 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 0.8 | 1.6 | 3.8 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 2.2 | 3.8 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.1 | 2.3 | 6.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.3 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 1.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.2 | 1.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.2 | 1.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.2 | 2.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.0 | 1.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 1.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 1.1 | 1.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.5 | 0.7 | 2.2 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.2 | 1.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.4 | 1.1 | 2.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | * | 0.5 | 1.3 | 3.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.6 | 4.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 1.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 1.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 1.2 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Zoonotic and other bacterial diseases (A20-A49) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 12.0 | 7.5 | 0.7 | 0.2 | 0.4 | 0.9 | 2.1 | 5.7 | 13.7 | 33.9 | 85.6 | 194.8 |
| 2003..... | 12.3 | 7.9 | 0.7 | 0.2 | 0.5 | 0.9 | 2.2 | 5.6 | 13.9 | 34.2 | 88.7 | 209.5 |
| 2002..... | 12.3 | 8.6 | 0.6 | 0.3 | 0.4 | 0.9 | 2.1 | 5.5 | 13.3 | 36.2 | 90.2 | 210.5 |
| 2001..... | 11.9 | 8.9 | 0.9 | 0.3 | 0.5 | 0.8 | 2.0 | 5.4 | 13.0 | 34.3 | 85.9 | 213.7 |
| 2000..... | 11.6 | 8.6 | 0.8 | 0.2 | 0.4 | 0.8 | 2.1 | 5.3 | 12.6 | 32.3 | 83.8 | 222.5 |
| 1999..... | 11.7 | 9.1 | 0.8 | 0.3 | 0.5 | 0.8 | 2.0 | 5.0 | 12.1 | 33.2 | 83.9 | 231.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 11.0 | 8.9 | 0.7 | 0.2 | 0.4 | 0.9 | 2.1 | 6.3 | 14.7 | 37.7 | 97.8 | 214.7 |
| 2003..... | 11.1 | 8.4 | 0.8 | 0.2 | 0.5 | 0.9 | 2.2 | 6.2 | 15.1 | 38.2 | 99.4 | 223.2 |
| 2002..... | 11.1 | 8.6 | 0.7 | 0.3 | 0.4 | 0.9 | 2.2 | 6.1 | 14.4 | 39.7 | 103.7 | 222.5 |
| 2001..... | 10.8 | 9.5 | 0.9 | 0.3 | 0.5 | 0.8 | 2.2 | 5.9 | 14.1 | 38.4 | 99.9 | 233.5 |
| 2000..... | 10.4 | 8.9 | 0.9 | 0.2 | 0.5 | 0.9 | 2.2 | 6.1 | 13.7 | 36.1 | 96.0 | 233.1 |
| 1999..... | 10.5 | 10.4 | 0.9 | 0.3 | 0.5 | 0.9 | 2.3 | 5.6 | 13.3 | 37.6 | 97.5 | 246.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 12.9 | 6.1 | 0.7 | 0.2 | 0.4 | 0.9 | 2.0 | 5.2 | 12.7 | 30.7 | 77.5 | 185.8 |
| 2003..... | 13.5 | 7.5 | 0.5 | 0.3 | 0.5 | 0.8 | 2.2 | 5.1 | 12.8 | 30.8 | 81.5 | 203.5 |
| 2002..... | 13.5 | 8.5 | 0.6 | 0.3 | 0.4 | 0.8 | 2.0 | 4.9 | 12.3 | 33.3 | 81.3 | 205.3 |
| 2001..... | 12.9 | 8.3 | 0.8 | 0.2 | 0.5 | 0.8 | 1.8 | 4.9 | 12.0 | 30.8 | 76.6 | 205.4 |
| 2000..... | 12.9 | 8.3 | 0.7 | 0.2 | 0.4 | 0.7 | 2.0 | 4.5 | 11.5 | 29.2 | 75.9 | 218.2 |
| 1999..... | 12.9 | 7.7 | 0.7 | 0.3 | 0.4 | 0.8 | 1.8 | 4.5 | 11.1 | 29.7 | 75.0 | 225.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | 6.9 | 0.9 | * | * | 0.5 | 1.5 | 4.3 | 10.3 | 26.3 | 66.9 | 143.7 |
| 2003..... | 4.0 | 6.9 | * | * | 0.3 | 0.6 | 1.5 | 4.2 | 9.8 | 23.7 | 64.8 | 157.5 |
| 2002..... | 4.2 | 9.4 | * | * | 0.3 | 0.6 | 1.3 | 4.1 | 8.8 | 29.7 | 66.1 | 163.5 |
| 2001..... | 4.0 | 8.3 | 0.8 | 0.3 | * | 0.5 | 1.4 | 4.2 | 8.8 | 26.1 | 65.6 | 178.8 |
| 2000..... | 3.8 | 10.2 | 0.7 | * | 0.3 | 0.6 | 1.6 | 4.2 | 9.2 | 22.6 | 62.6 | 148.6 |
| 1999..... | 4.0 | 7.1 | * | * | * | 0.6 | 1.7 | 3.6 | 9.1 | 27.6 | 65.0 | 192.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.0 | 8.6 | * | * | * | 0.5 | 1.7 | 5.6 | 10.8 | 28.8 | 72.2 | 152.7 |
| 2003..... | 3.7 | 7.5 | * | * | * | 0.6 | 1.6 | 5.4 | 10.6 | 26.9 | 69.8 | 166.2 |
| 2002..... | 4.0 | 9.6 | * | * | * | 0.7 | 1.7 | 5.5 | 9.5 | 31.5 | 74.7 | 172.3 |
| 2001..... | 3.9 | 8.6 | * | * | * | 1.6 | 5.3 | 10.7 | 30.8 | 80.0 | 165.7 | |
| 2000..... | 3.6 | 9.9 | * | * | * | 0.7 | 1.8 | 5.2 | 10.5 | 27.0 | 72.8 | 159.2 |
| 1999..... | 3.8 | 8.3 | * | * | * | 0.6 | 2.1 | 4.4 | 11.5 | 30.8 | 76.6 | 194.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.5 | 5.2 | * | * | * | * | 1.3 | 2.9 | 9.9 | 24.3 | 63.2 | 139.0 |
| 2003..... | 4.3 | 6.4 | * | * | * | * | 1.4 | 3.1 | 9.0 | 21.0 | 61.4 | 153.1 |
| 2002..... | 4.4 | 9.1 | * | * | * | * | 0.8 | 2.7 | 8.2 | 28.3 | 60.1 | 159.2 |
| 2001..... | 4.2 | 8.0 | * | * | * | * | 1.2 | 3.2 | 7.1 | 22.5 | 55.9 | 185.2 |
| 2000..... | 3.9 | 10.6 | * | * | * | * | 1.4 | 3.2 | 8.0 | 19.1 | 55.8 | 143.4 |
| 1999..... | 4.2 | 6.0 | * | * | * | * | 1.2 | 2.8 | 7.0 | 25.1 | 57.2 | 191.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Zoonotic and other bacterial diseases (A20-A49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 13.2 | 7.5 | 0.6 | 0.2 | 0.5 | 1.0 | 2.2 | 5.9 | 13.9 | 34.4 | 86.5 | 196.8 |
| 2003..... | 13.6 | 8.2 | 0.8 | 0.3 | 0.5 | 0.9 | 2.3 | 5.8 | 14.2 | 34.8 | 89.7 | 211.3 |
| 2002..... | 13.5 | 8.2 | 0.6 | 0.3 | 0.4 | 0.9 | 2.2 | 5.7 | 13.6 | 36.5 | 91.1 | 211.7 |
| 2001..... | 13.0 | 9.0 | 0.9 | 0.2 | 0.5 | 0.9 | 2.1 | 5.5 | 13.2 | 34.7 | 86.6 | 214.5 |
| 2000..... | 12.7 | 8.1 | 0.8 | 0.2 | 0.4 | 0.8 | 2.1 | 5.3 | 12.8 | 32.8 | 84.4 | 224.4 |
| 1999..... | 12.7 | 9.5 | 0.8 | 0.3 | 0.5 | 0.9 | 2.1 | 5.1 | 12.3 | 33.5 | 84.3 | 231.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 12.2 | 8.8 | 0.6 | 0.2 | 0.5 | 1.0 | 2.2 | 6.3 | 14.9 | 38.3 | 99.1 | 217.4 |
| 2003..... | 12.3 | 8.6 | 0.9 | 0.2 | 0.5 | 0.9 | 2.3 | 6.2 | 15.3 | 38.8 | 100.8 | 225.4 |
| 2002..... | 12.3 | 8.1 | 0.6 | 0.4 | 0.4 | 1.0 | 2.3 | 6.2 | 14.7 | 40.0 | 104.7 | 224.2 |
| 2001..... | 11.9 | 9.7 | 1.0 | 0.2 | 0.6 | 0.9 | 2.2 | 5.9 | 14.2 | 38.7 | 100.5 | 236.2 |
| 2000..... | 11.3 | 8.6 | 0.9 | 0.2 | 0.5 | 0.9 | 2.2 | 6.1 | 13.8 | 36.5 | 96.5 | 235.2 |
| 1999..... | 11.4 | 10.9 | 0.9 | 0.3 | 0.5 | 0.9 | 2.3 | 5.6 | 13.3 | 37.9 | 97.9 | 247.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 14.2 | 6.3 | 0.6 | 0.3 | 0.4 | 1.0 | 2.1 | 5.4 | 12.9 | 31.1 | 78.1 | 187.6 |
| 2003..... | 14.8 | 7.8 | 0.6 | 0.3 | 0.5 | 0.9 | 2.3 | 5.3 | 13.1 | 31.5 | 82.4 | 205.0 |
| 2002..... | 14.8 | 8.3 | 0.6 | 0.3 | 0.4 | 0.9 | 2.1 | 5.2 | 12.6 | 33.5 | 82.1 | 206.4 |
| 2001..... | 14.1 | 8.4 | 0.7 | 0.2 | 0.5 | 0.9 | 1.9 | 5.1 | 12.3 | 31.3 | 77.4 | 205.6 |
| 2000..... | 14.0 | 7.6 | 0.8 | 0.2 | 0.4 | 0.8 | 2.1 | 4.6 | 11.7 | 29.8 | 76.5 | 220.0 |
| 1999..... | 14.0 | 8.1 | 0.8 | 0.3 | 0.5 | 0.8 | 1.9 | 4.6 | 11.4 | 29.8 | 75.6 | 225.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 13.2 | 5.8 | 0.5 | 0.2 | 0.4 | 0.8 | 1.7 | 4.7 | 11.8 | 30.8 | 81.1 | 187.6 |
| 2003..... | 13.5 | 5.1 | 0.6 | 0.2 | 0.5 | 0.7 | 1.8 | 4.5 | 12.0 | 30.9 | 84.6 | 199.6 |
| 2002..... | 13.5 | 6.6 | 0.6 | 0.3 | 0.4 | 0.7 | 1.8 | 4.6 | 11.3 | 32.5 | 85.0 | 200.1 |
| 2001..... | 12.9 | 6.1 | 0.8 | 0.2 | 0.4 | 0.6 | 1.6 | 4.3 | 11.0 | 30.9 | 81.0 | 203.3 |
| 2000..... | 12.5 | 5.9 | 0.7 | 0.2 | 0.4 | 0.6 | 1.7 | 4.1 | 10.6 | 28.9 | 77.5 | 211.0 |
| 1999..... | 12.4 | 7.1 | 0.8 | 0.3 | 0.4 | 0.7 | 1.6 | 4.0 | 10.1 | 29.2 | 77.1 | 216.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 12.1 | 6.8 | 0.5 | 0.2 | 0.4 | 0.8 | 1.8 | 5.1 | 12.6 | 34.2 | 92.8 | 209.7 |
| 2003..... | 12.1 | 4.9 | 0.8 | 0.2 | 0.5 | 0.7 | 1.9 | 4.9 | 12.9 | 34.3 | 95.0 | 214.3 |
| 2002..... | 12.1 | 6.4 | 0.7 | 0.4 | 0.4 | 0.8 | 1.8 | 4.9 | 12.5 | 35.7 | 96.8 | 212.8 |
| 2001..... | 11.6 | 6.8 | 1.0 | 0.2 | 0.5 | 0.6 | 1.8 | 4.6 | 11.8 | 34.2 | 94.1 | 223.0 |
| 2000..... | 11.0 | 6.1 | 0.8 | 0.2 | 0.5 | 0.6 | 1.8 | 4.7 | 11.6 | 32.4 | 88.8 | 221.2 |
| 1999..... | 11.0 | 8.4 | 0.8 | 0.2 | 0.5 | 0.7 | 1.7 | 4.3 | 10.6 | 32.8 | 89.6 | 230.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 14.3 | 4.8 | 0.5 | 0.2 | 0.3 | 0.8 | 1.7 | 4.3 | 11.1 | 27.8 | 73.2 | 177.7 |
| 2003..... | 14.9 | 5.4 | * | 0.2 | 0.4 | 0.7 | 1.8 | 4.2 | 11.3 | 28.0 | 77.6 | 193.1 |
| 2002..... | 14.8 | 6.8 | 0.5 | 0.2 | 0.4 | 0.7 | 1.8 | 4.2 | 10.2 | 29.8 | 77.1 | 194.6 |
| 2001..... | 14.2 | 5.3 | 0.6 | 0.2 | 0.4 | 0.7 | 1.4 | 4.0 | 10.3 | 28.1 | 72.4 | 195.1 |
| 2000..... | 13.9 | 5.6 | 0.7 | 0.2 | 0.3 | 0.6 | 1.6 | 3.6 | 9.6 | 26.0 | 70.0 | 206.9 |
| 1999..... | 13.9 | 5.8 | 0.7 | 0.3 | 0.4 | 0.6 | 1.4 | 3.6 | 9.7 | 26.1 | 68.8 | 210.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Zoonotic and other bacterial diseases (A20-A49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 16.8 | 14.7 | 1.0 | * | 0.9 | 2.2 | 5.0 | 14.4 | 33.6 | 73.7 | 164.5 | 333.3 |
| 2003..... | 17.5 | 19.8 | 1.3 | 0.4 | 0.8 | 2.1 | 5.5 | 14.9 | 34.4 | 76.1 | 164.3 | 381.7 |
| 2002..... | 17.5 | 14.5 | 0.9 | 0.5 | 0.6 | 2.1 | 5.1 | 13.9 | 35.1 | 78.5 | 177.5 | 380.2 |
| 2001..... | 17.1 | 21.3 | 1.2 | * | 1.0 | 2.1 | 5.0 | 14.4 | 34.3 | 75.3 | 166.8 | 377.1 |
| 2000..... | 17.4 | 18.7 | 1.3 | 0.3 | 0.8 | 2.2 | 5.2 | 14.7 | 32.9 | 74.9 | 179.7 | 405.4 |
| 1999..... | 17.6 | 20.0 | 0.9 | 0.3 | 0.8 | 2.0 | 5.5 | 14.0 | 32.2 | 78.9 | 178.9 | 430.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 15.9 | 16.1 | * | * | 0.9 | 2.4 | 5.1 | 16.0 | 38.9 | 87.1 | 200.5 | 357.7 |
| 2003..... | 16.6 | 22.7 | * | * | 0.7 | 2.4 | 5.6 | 17.0 | 40.8 | 90.6 | 196.5 | 420.8 |
| 2002..... | 16.2 | 14.8 | * | * | * | 2.0 | 5.5 | 15.9 | 38.2 | 90.6 | 220.4 | 408.0 |
| 2001..... | 16.1 | 22.0 | * | * | 1.1 | 2.1 | 5.2 | 16.0 | 39.4 | 90.2 | 202.2 | 442.4 |
| 2000..... | 16.1 | 20.5 | * | * | 0.9 | 2.6 | 5.3 | 18.0 | 37.0 | 84.7 | 214.4 | 445.0 |
| 1999..... | 16.8 | 21.5 | * | * | 0.8 | 2.2 | 6.5 | 16.5 | 39.7 | 96.2 | 216.4 | 470.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 17.6 | 13.2 | * | * | 0.8 | 2.0 | 4.8 | 13.1 | 29.3 | 64.1 | 144.1 | 323.7 |
| 2003..... | 18.3 | 16.8 | * | * | 0.9 | 1.8 | 5.3 | 13.0 | 29.3 | 65.8 | 146.2 | 366.5 |
| 2002..... | 18.6 | 14.2 | * | 0.7 | * | 2.2 | 4.7 | 12.1 | 32.6 | 69.9 | 153.2 | 369.6 |
| 2001..... | 17.9 | 20.5 | * | * | 0.9 | 2.1 | 4.9 | 13.0 | 30.3 | 64.8 | 146.9 | 352.9 |
| 2000..... | 18.5 | 16.9 | * | * | * | 1.9 | 5.2 | 11.9 | 29.5 | 67.9 | 160.1 | 390.7 |
| 1999..... | 18.3 | 18.6 | * | * | 0.8 | 1.9 | 4.6 | 12.0 | 26.2 | 66.7 | 157.9 | 414.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Septicemia (A40-A41) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 11.4 | 6.6 | 0.5 | 0.2 | 0.3 | 0.8 | 1.9 | 5.4 | 12.9 | 32.4 | 81.6 | 186.7 |
| 2003..... | 11.7 | 6.9 | 0.5 | 0.2 | 0.4 | 0.8 | 2.1 | 5.3 | 13.1 | 32.6 | 85.0 | 202.5 |
| 2002..... | 11.7 | 7.3 | 0.5 | 0.2 | 0.3 | 0.8 | 1.9 | 5.2 | 12.6 | 34.7 | 86.5 | 203.0 |
| 2001..... | 11.3 | 7.7 | 0.7 | 0.2 | 0.3 | 0.7 | 1.8 | 5.0 | 12.3 | 32.8 | 82.3 | 205.9 |
| 2000..... | 11.1 | 7.2 | 0.6 | 0.2 | 0.3 | 0.7 | 1.9 | 4.9 | 11.9 | 31.0 | 80.4 | 215.7 |
| 1999..... | 11.0 | 7.5 | 0.6 | 0.2 | 0.3 | 0.7 | 1.8 | 4.6 | 11.4 | 31.2 | 79.4 | 220.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 10.4 | 8.0 | 0.5 | 0.1 | 0.3 | 0.8 | 2.0 | 5.9 | 13.8 | 35.8 | 93.3 | 203.9 |
| 2003..... | 10.5 | 7.4 | 0.7 | 0.2 | 0.4 | 0.8 | 2.1 | 5.8 | 14.1 | 36.2 | 95.0 | 213.8 |
| 2002..... | 10.6 | 7.7 | 0.5 | 0.2 | 0.3 | 0.8 | 2.0 | 5.7 | 13.7 | 38.0 | 99.0 | 213.9 |
| 2001..... | 10.2 | 8.4 | 0.8 | 0.2 | 0.4 | 0.7 | 2.0 | 5.4 | 13.2 | 36.7 | 95.5 | 224.3 |
| 2000..... | 9.8 | 7.1 | 0.7 | 0.2 | 0.3 | 0.8 | 2.0 | 5.6 | 13.0 | 34.5 | 91.9 | 223.0 |
| 1999..... | 9.8 | 9.0 | 0.6 | 0.2 | 0.3 | 0.7 | 2.1 | 5.1 | 12.4 | 35.2 | 91.9 | 233.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 12.3 | 5.2 | 0.6 | 0.2 | 0.3 | 0.8 | 1.9 | 4.9 | 12.0 | 29.6 | 73.7 | 179.0 |
| 2003..... | 12.9 | 6.5 | 0.4 | 0.2 | 0.4 | 0.8 | 2.0 | 4.8 | 12.2 | 29.5 | 78.3 | 197.5 |
| 2002..... | 12.9 | 7.0 | 0.5 | 0.2 | 0.3 | 0.8 | 1.8 | 4.7 | 11.7 | 31.9 | 78.2 | 198.3 |
| 2001..... | 12.4 | 7.0 | 0.6 | 0.1 | 0.3 | 0.7 | 1.7 | 4.7 | 11.4 | 29.5 | 73.7 | 198.3 |
| 2000..... | 12.3 | 7.3 | 0.6 | 0.1 | 0.2 | 0.6 | 1.9 | 4.2 | 11.0 | 28.1 | 72.9 | 212.7 |
| 1999..... | 12.2 | 5.8 | 0.5 | 0.2 | 0.3 | 0.6 | 1.6 | 4.1 | 10.5 | 28.0 | 71.3 | 215.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | 6.1 | 0.7 | * | * | 0.5 | 1.4 | 4.0 | 9.7 | 25.1 | 63.8 | 138.0 |
| 2003..... | 3.8 | 5.8 | * | * | * | 0.5 | 1.5 | 3.9 | 9.1 | 22.1 | 62.0 | 151.8 |
| 2002..... | 3.9 | 7.8 | * | * | * | 0.5 | 1.2 | 3.7 | 8.4 | 28.9 | 63.7 | 156.3 |
| 2001..... | 3.7 | 6.1 | * | * | * | 0.3 | 1.3 | 3.9 | 8.2 | 25.4 | 62.4 | 171.3 |
| 2000..... | 3.5 | 8.0 | * | * | * | 0.5 | 1.5 | 3.9 | 8.5 | 21.8 | 59.3 | 144.7 |
| 1999..... | 3.6 | 5.7 | * | * | * | 0.4 | 1.5 | 3.1 | 8.0 | 25.6 | 61.7 | 180.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.8 | 7.8 | * | * | * | * | 1.6 | 5.2 | 10.3 | 27.2 | 69.7 | 147.1 |
| 2003..... | 3.5 | 6.6 | * | * | * | 0.5 | 1.5 | 4.9 | 9.6 | 25.3 | 67.1 | 160.1 |
| 2002..... | 3.7 | 8.0 | * | * | * | 0.6 | 1.6 | 5.2 | 9.0 | 30.8 | 72.2 | 164.0 |
| 2001..... | 3.6 | 7.0 | * | * | * | * | 1.5 | 4.7 | 9.9 | 29.8 | 76.0 | 160.0 |
| 2000..... | 3.4 | 8.4 | * | * | * | 0.6 | 1.7 | 5.0 | 10.0 | 26.6 | 70.3 | 157.2 |
| 1999..... | 3.5 | 7.0 | * | * | * | * | 2.0 | 3.8 | 10.2 | 29.4 | 75.1 | 181.2 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | * | 1.2 | 2.8 | 9.2 | 23.4 | 59.8 | 133.2 |
| 2003..... | 4.1 | 5.0 | * | * | * | * | 1.3 | 2.8 | 8.7 | 19.6 | 58.5 | 147.7 |
| 2002..... | 4.1 | 7.6 | * | * | * | * | 0.7 | 2.3 | 7.8 | 27.4 | 57.9 | 152.5 |
| 2001..... | 3.9 | 5.2 | * | * | * | * | 1.1 | 3.1 | 6.6 | 21.9 | 53.2 | 176.9 |
| 2000..... | 3.5 | 7.7 | * | * | * | * | 1.3 | 2.8 | 7.2 | 18.1 | 51.8 | 138.5 |
| 1999..... | 3.7 | * | * | * | * | * | 1.0 | 2.4 | 6.1 | 22.5 | 52.7 | 179.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Septicemia (A40-A41) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 12.5 | 6.7 | 0.5 | 0.2 | 0.4 | 0.9 | 2.0 | 5.5 | 13.1 | 32.9 | 82.5 | 188.7 |
| 2003..... | 12.9 | 7.2 | 0.6 | 0.2 | 0.4 | 0.8 | 2.1 | 5.4 | 13.3 | 33.2 | 86.0 | 204.1 |
| 2002..... | 12.9 | 7.1 | 0.5 | 0.2 | 0.3 | 0.8 | 2.0 | 5.3 | 12.9 | 34.9 | 87.3 | 204.2 |
| 2001..... | 12.4 | 8.1 | 0.7 | 0.1 | 0.4 | 0.8 | 1.9 | 5.1 | 12.5 | 33.1 | 83.0 | 206.8 |
| 2000..... | 12.1 | 6.9 | 0.7 | 0.2 | 0.3 | 0.7 | 2.0 | 5.0 | 12.1 | 31.4 | 81.0 | 217.5 |
| 1999..... | 12.0 | 7.9 | 0.6 | 0.2 | 0.3 | 0.7 | 1.8 | 4.7 | 11.6 | 31.4 | 79.8 | 221.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 11.5 | 7.9 | 0.4 | 0.1 | 0.4 | 0.9 | 2.0 | 5.9 | 14.0 | 36.3 | 94.6 | 206.3 |
| 2003..... | 11.6 | 7.6 | 0.8 | 0.2 | 0.4 | 0.9 | 2.1 | 5.8 | 14.3 | 36.8 | 96.2 | 215.9 |
| 2002..... | 11.6 | 7.3 | 0.5 | 0.3 | 0.3 | 0.9 | 2.0 | 5.7 | 14.0 | 38.2 | 99.8 | 215.6 |
| 2001..... | 11.2 | 8.8 | 0.9 | 0.1 | 0.4 | 0.8 | 2.1 | 5.5 | 13.4 | 37.0 | 96.1 | 226.8 |
| 2000..... | 10.7 | 6.7 | 0.7 | 0.2 | 0.3 | 0.8 | 2.0 | 5.6 | 13.1 | 34.8 | 92.3 | 224.8 |
| 1999..... | 10.7 | 9.4 | 0.6 | 0.2 | 0.4 | 0.8 | 2.1 | 5.2 | 12.5 | 35.4 | 92.2 | 234.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 13.5 | 5.4 | 0.5 | 0.2 | 0.3 | 0.9 | 2.0 | 5.1 | 12.2 | 30.0 | 74.3 | 180.8 |
| 2003..... | 14.2 | 6.8 | 0.4 | 0.2 | 0.4 | 0.8 | 2.1 | 5.0 | 12.4 | 30.1 | 79.1 | 199.0 |
| 2002..... | 14.1 | 6.8 | 0.5 | 0.2 | 0.3 | 0.8 | 2.0 | 4.9 | 11.9 | 32.1 | 79.0 | 199.3 |
| 2001..... | 13.5 | 7.4 | 0.6 | 0.2 | 0.3 | 0.8 | 1.7 | 4.8 | 11.8 | 29.9 | 74.4 | 198.5 |
| 2000..... | 13.5 | 7.0 | 0.6 | 0.1 | 0.2 | 0.7 | 1.9 | 4.3 | 11.2 | 28.7 | 73.6 | 214.5 |
| 1999..... | 13.2 | 6.2 | 0.5 | 0.2 | 0.3 | 0.7 | 1.6 | 4.2 | 10.8 | 28.2 | 71.8 | 215.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 12.5 | 4.8 | 0.4 | 0.2 | 0.3 | 0.7 | 1.6 | 4.4 | 11.1 | 29.3 | 77.2 | 179.6 |
| 2003..... | 12.9 | 4.2 | 0.5 | 0.2 | 0.3 | 0.6 | 1.7 | 4.3 | 11.3 | 29.3 | 80.8 | 192.4 |
| 2002..... | 12.9 | 5.5 | 0.5 | 0.2 | 0.3 | 0.7 | 1.6 | 4.3 | 10.7 | 31.0 | 81.3 | 192.6 |
| 2001..... | 12.3 | 5.1 | 0.7 | 0.1 | 0.3 | 0.6 | 1.4 | 4.0 | 10.4 | 29.4 | 77.6 | 195.6 |
| 2000..... | 11.9 | 4.8 | 0.5 | 0.2 | 0.2 | 0.5 | 1.5 | 3.8 | 10.0 | 27.7 | 74.2 | 204.3 |
| 1999..... | 11.7 | 5.3 | 0.5 | 0.2 | 0.3 | 0.6 | 1.4 | 3.6 | 9.5 | 27.3 | 73.0 | 206.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 11.4 | 5.7 | * | 0.2 | 0.3 | 0.7 | 1.6 | 4.8 | 11.7 | 32.4 | 88.5 | 198.5 |
| 2003..... | 11.4 | 3.9 | 0.6 | 0.2 | 0.4 | 0.6 | 1.7 | 4.6 | 12.0 | 32.5 | 90.5 | 204.8 |
| 2002..... | 11.5 | 5.6 | 0.5 | 0.3 | 0.3 | 0.7 | 1.6 | 4.6 | 11.8 | 34.0 | 92.1 | 203.9 |
| 2001..... | 11.0 | 5.8 | 0.9 | * | 0.3 | 0.5 | 1.7 | 4.2 | 11.1 | 32.5 | 89.7 | 213.6 |
| 2000..... | 10.4 | 4.7 | 0.5 | 0.2 | 0.3 | 0.6 | 1.6 | 4.3 | 10.9 | 30.9 | 84.7 | 211.2 |
| 1999..... | 10.2 | 6.7 | 0.5 | 0.2 | 0.3 | 0.7 | 1.5 | 4.0 | 9.9 | 30.5 | 84.6 | 218.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 13.6 | 3.9 | 0.5 | 0.2 | 0.2 | 0.7 | 1.5 | 4.1 | 10.4 | 26.7 | 69.5 | 171.1 |
| 2003..... | 14.3 | 4.5 | * | 0.2 | 0.3 | 0.7 | 1.7 | 3.9 | 10.6 | 26.6 | 74.3 | 187.0 |
| 2002..... | 14.2 | 5.4 | * | 0.2 | 0.3 | 0.6 | 1.7 | 3.9 | 9.7 | 28.5 | 74.0 | 187.7 |
| 2001..... | 13.6 | 4.4 | 0.5 | * | 0.3 | 0.6 | 1.2 | 3.8 | 9.9 | 26.8 | 69.6 | 188.1 |
| 2000..... | 13.4 | 4.8 | 0.6 | * | * | 0.5 | 1.5 | 3.3 | 9.2 | 24.9 | 67.2 | 201.6 |
| 1999..... | 13.1 | 3.8 | 0.5 | 0.2 | 0.2 | 0.5 | 1.2 | 3.3 | 9.1 | 24.5 | 65.4 | 201.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Septicemia (A40-A41) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 16.1 | 14.2 | * | * | 0.7 | 1.9 | 4.6 | 13.6 | 32.2 | 71.4 | 157.9 | 324.7 |
| 2003..... | 16.8 | 18.8 | 1.0 | * | 0.7 | 2.0 | 5.1 | 14.0 | 32.8 | 74.0 | 160.0 | 373.9 |
| 2002..... | 16.8 | 13.1 | * | 0.3 | 0.4 | 2.0 | 4.6 | 13.2 | 33.8 | 76.0 | 172.9 | 370.8 |
| 2001..... | 16.4 | 20.3 | 0.9 | * | 0.7 | 1.9 | 4.8 | 13.7 | 32.6 | 72.8 | 161.9 | 368.6 |
| 2000..... | 16.7 | 16.6 | 1.1 | * | 0.6 | 2.1 | 4.8 | 13.9 | 31.8 | 72.5 | 175.1 | 395.9 |
| 1999..... | 16.6 | 18.6 | * | * | 0.6 | 1.6 | 5.1 | 13.2 | 30.8 | 75.3 | 169.8 | 414.2 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 15.1 | 15.8 | * | * | 0.7 | 2.1 | 4.8 | 15.2 | 37.1 | 82.9 | 192.5 | 346.5 |
| 2003..... | 15.8 | 21.7 | * | * | * | 2.3 | 5.2 | 15.8 | 38.7 | 86.9 | 191.3 | 409.1 |
| 2002..... | 15.5 | 13.9 | * | * | * | 1.9 | 5.0 | 15.0 | 36.6 | 87.6 | 213.7 | 399.3 |
| 2001..... | 15.4 | 21.4 | * | * | 0.9 | 1.9 | 4.9 | 15.2 | 37.4 | 86.9 | 196.8 | 430.8 |
| 2000..... | 15.3 | 16.4 | * | * | * | 2.4 | 4.8 | 16.8 | 35.9 | 81.6 | 208.9 | 432.1 |
| 1999..... | 15.7 | 20.8 | * | * | * | 1.7 | 6.0 | 15.4 | 37.9 | 91.5 | 202.6 | 451.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 17.0 | 12.6 | * | * | 0.8 | 1.8 | 4.5 | 12.3 | 28.3 | 63.1 | 138.3 | 316.0 |
| 2003..... | 17.7 | 15.8 | * | * | 0.7 | 1.7 | 5.1 | 12.5 | 28.2 | 64.9 | 142.3 | 360.3 |
| 2002..... | 18.0 | 12.3 | * | * | * | 2.0 | 4.2 | 11.6 | 31.6 | 67.8 | 149.9 | 359.9 |
| 2001..... | 17.2 | 19.1 | * | * | * | 1.9 | 4.7 | 12.5 | 28.8 | 62.8 | 142.4 | 345.6 |
| 2000..... | 17.9 | 16.9 | * | * | * | 1.8 | 4.9 | 11.5 | 28.5 | 66.1 | 156.1 | 382.5 |
| 1999..... | 17.4 | 16.4 | * | * | * | 1.6 | 4.3 | 11.3 | 25.1 | 63.9 | 151.4 | 400.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 1.5 | 4.0 | 7.9 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 3.7 | 7.0 |
| 2002..... | 0.6 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.7 | 7.4 |
| 2001..... | 0.5 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 3.5 | 7.5 |
| 2000..... | 0.5 | 0.7 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.3 | 6.7 |
| 1999..... | 0.7 | 0.7 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.0 | 4.4 | 10.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 1.9 | 4.5 | 11.1 |
| 2003..... | 0.6 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.0 | 4.5 | 9.6 |
| 2002..... | 0.6 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.8 | 4.7 | 8.5 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.4 | 9.1 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.6 | 4.0 | 10.2 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.4 | 5.6 | 13.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.1 | 3.7 | 6.4 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.1 | 5.9 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.1 | 6.9 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.9 | 6.8 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.8 | 5.3 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.6 | 3.6 | 9.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 3.9 | 7.8 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.6 | 3.7 | 7.1 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.6 | 7.3 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 3.4 | 7.4 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 3.2 | 6.5 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.8 | 4.0 | 9.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.8 | 1.7 | 4.2 | 11.1 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 1.8 | 4.5 | 9.6 |
| 2002..... | 0.6 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.7 | 4.6 | 8.7 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.4 | 0.7 | 1.6 | 4.4 | 9.1 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.4 | 0.7 | 1.5 | 4.0 | 9.8 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.3 | 5.0 | 12.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 1.1 | 3.6 | 6.3 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.2 | 6.0 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 6.7 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.7 | 6.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.7 | 5.1 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.5 | 3.3 | 8.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.7 | 1.4 | 2.3 | 6.6 | 8.3 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.7 | 1.4 | 2.1 | 4.4 | 7.7 |
| 2002..... | 0.6 | * | * | * | * | * | 0.4 | 0.6 | 1.2 | 2.5 | 4.6 | 9.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.7 | 1.5 | 2.5 | 4.7 | 8.1 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.5 | 1.1 | 2.2 | 4.6 | 9.5 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.8 | 1.4 | 3.5 | 9.0 | 15.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | 1.8 | 4.0 | 8.0 | * |
| 2003..... | 0.7 | * | * | * | * | * | * | 1.1 | 1.9 | 3.8 | * | * |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | 6.7 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 1.8 | 3.3 | * | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 1999..... | 0.9 | * | * | * | * | * | * | * | * | 4.7 | 13.4 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | * | 5.8 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | 3.9 | * |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 2.1 | 3.3 | 9.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | * | 4.5 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | 4.0 | * |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 2.7 | 6.5 | 14.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral diseases (A80-B34) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.8 | 2.9 | 0.3 | 0.1 | 0.6 | 3.9 | 12.5 | 16.2 | 10.3 | 7.0 | 6.5 | 7.8 |
| 2003..... | 7.1 | 2.9 | 0.4 | 0.2 | 0.6 | 4.2 | 13.7 | 16.7 | 10.0 | 7.7 | 7.5 | 9.1 |
| 2002..... | 7.4 | 4.1 | 0.3 | 0.2 | 0.6 | 5.0 | 14.6 | 17.3 | 9.6 | 7.8 | 7.8 | 10.1 |
| 2001..... | 7.4 | 3.0 | 0.4 | 0.2 | 0.7 | 5.6 | 15.0 | 16.0 | 9.7 | 7.9 | 7.2 | 8.3 |
| 2000..... | 7.5 | 2.9 | 0.4 | 0.2 | 0.6 | 6.4 | 15.0 | 16.3 | 9.4 | 8.0 | 8.1 | 9.6 |
| 1999..... | 7.6 | 3.5 | 0.5 | 0.3 | 0.6 | 7.1 | 15.6 | 15.1 | 8.8 | 8.1 | 8.3 | 11.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 9.5 | 3.2 | 0.4 | 0.1 | 0.7 | 4.8 | 17.7 | 24.6 | 15.9 | 9.4 | 7.3 | 9.0 |
| 2003..... | 10.1 | 2.5 | 0.4 | 0.2 | 0.6 | 5.4 | 19.8 | 25.8 | 15.2 | 10.1 | 9.3 | 10.2 |
| 2002..... | 10.6 | 4.6 | 0.4 | 0.1 | 0.6 | 6.3 | 21.3 | 26.9 | 14.5 | 10.0 | 9.6 | 11.7 |
| 2001..... | 10.5 | 3.1 | 0.5 | 0.2 | 0.7 | 7.4 | 22.1 | 25.2 | 14.5 | 9.6 | 8.5 | 9.3 |
| 2000..... | 10.8 | 2.9 | 0.4 | 0.2 | 0.7 | 8.3 | 22.3 | 25.8 | 14.3 | 9.9 | 10.4 | 10.8 |
| 1999..... | 10.9 | 4.0 | 0.6 | 0.3 | 0.6 | 9.8 | 23.5 | 24.1 | 13.3 | 10.3 | 9.9 | 13.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.1 | 2.6 | 0.3 | 0.1 | 0.6 | 3.0 | 7.3 | 8.2 | 5.0 | 5.0 | 6.0 | 7.2 |
| 2003..... | 4.2 | 3.3 | 0.4 | 0.2 | 0.5 | 3.0 | 7.6 | 7.9 | 5.1 | 5.7 | 6.3 | 8.7 |
| 2002..... | 4.4 | 3.6 | * | 0.2 | 0.6 | 3.6 | 7.9 | 7.9 | 5.1 | 6.0 | 6.5 | 9.3 |
| 2001..... | 4.3 | 2.8 | 0.3 | 0.2 | 0.8 | 3.7 | 7.9 | 7.1 | 5.3 | 6.4 | 6.4 | 7.9 |
| 2000..... | 4.4 | 2.9 | 0.4 | 0.2 | 0.6 | 4.5 | 7.7 | 7.1 | 5.0 | 6.4 | 6.6 | 9.1 |
| 1999..... | 4.3 | 3.1 | 0.4 | 0.2 | 0.6 | 4.4 | 7.8 | 6.5 | 4.7 | 6.4 | 7.2 | 10.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 6.3 | 2.4 | * | * | 0.5 | 2.9 | 13.0 | 21.2 | 16.7 | 13.2 | 12.3 | * |
| 2003..... | 6.8 | * | * | * | 0.6 | 3.5 | 14.1 | 23.1 | 17.3 | 15.0 | 13.4 | * |
| 2002..... | 7.0 | 4.0 | * | * | 0.6 | 4.0 | 15.3 | 22.9 | 18.1 | 13.0 | 13.2 | * |
| 2001..... | 7.4 | 3.2 | * | * | 0.6 | 4.6 | 16.3 | 23.5 | 20.3 | 15.1 | 13.7 | 13.1 |
| 2000..... | 7.7 | * | * | * | 0.5 | 5.9 | 17.5 | 24.2 | 18.5 | 16.3 | 17.4 | * |
| 1999..... | 8.0 | 3.2 | * | * | 0.7 | 6.8 | 19.7 | 22.6 | 16.6 | 18.0 | 15.6 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 9.0 | * | * | * | 0.7 | 4.2 | 19.0 | 32.4 | 25.6 | 17.9 | 12.5 | * |
| 2003..... | 9.7 | * | * | * | 0.7 | 4.8 | 20.3 | 37.5 | 26.1 | 18.5 | 14.6 | * |
| 2002..... | 10.0 | 5.6 | * | * | 0.7 | 5.6 | 22.7 | 35.3 | 26.7 | 15.5 | 14.0 | * |
| 2001..... | 10.4 | * | * | * | 0.7 | 6.1 | 24.0 | 37.3 | 29.7 | 16.8 | 15.7 | * |
| 2000..... | 11.0 | * | * | * | 0.7 | 8.3 | 26.7 | 38.6 | 27.5 | 19.0 | 19.7 | * |
| 1999..... | 11.4 | * | * | * | 0.7 | 9.6 | 29.8 | 35.2 | 25.5 | 23.1 | 13.0 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | 1.4 | 6.3 | 9.8 | 8.5 | 9.5 | 12.1 | * |
| 2003..... | 3.7 | * | * | * | * | 1.9 | 7.3 | 8.7 | 9.3 | 12.2 | 12.6 | * |
| 2002..... | 3.9 | * | * | * | * | 2.1 | 7.3 | 10.5 | 10.4 | 10.9 | 12.6 | * |
| 2001..... | 4.2 | * | * | * | * | 2.8 | 8.0 | 9.8 | 11.9 | 13.7 | 12.4 | * |
| 2000..... | 4.1 | * | * | * | * | 3.1 | 7.6 | 10.2 | 10.6 | 14.3 | 15.8 | * |
| 1999..... | 4.4 | * | * | * | * | 3.5 | 8.9 | 10.3 | 8.6 | 14.1 | 17.3 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral diseases (A80-B34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.8 | 3.0 | 0.3 | 0.1 | 0.6 | 4.1 | 12.3 | 15.5 | 9.7 | 6.5 | 6.1 | 7.9 |
| 2003..... | 7.1 | 3.2 | 0.4 | 0.2 | 0.6 | 4.4 | 13.5 | 15.9 | 9.3 | 7.1 | 7.2 | 9.2 |
| 2002..... | 7.4 | 4.0 | 0.4 | 0.2 | 0.6 | 5.1 | 14.3 | 16.5 | 8.8 | 7.4 | 7.5 | 10.1 |
| 2001..... | 7.3 | 2.9 | 0.5 | 0.2 | 0.7 | 5.7 | 14.6 | 15.1 | 8.8 | 7.3 | 6.9 | 8.1 |
| 2000..... | 7.4 | 3.1 | 0.4 | 0.2 | 0.6 | 6.5 | 14.5 | 15.4 | 8.6 | 7.4 | 7.7 | 9.6 |
| 1999..... | 7.4 | 3.6 | 0.5 | 0.3 | 0.6 | 7.1 | 14.9 | 14.2 | 8.1 | 7.5 | 8.0 | 11.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 9.5 | 3.3 | * | 0.1 | 0.7 | 4.9 | 17.4 | 23.5 | 15.0 | 8.7 | 7.0 | 9.3 |
| 2003..... | 10.1 | 2.8 | 0.4 | 0.2 | 0.6 | 5.5 | 19.5 | 24.3 | 14.2 | 9.4 | 9.0 | 10.1 |
| 2002..... | 10.6 | 4.2 | 0.5 | 0.1 | 0.6 | 6.4 | 20.8 | 25.7 | 13.4 | 9.6 | 9.4 | 11.7 |
| 2001..... | 10.4 | 3.0 | 0.6 | 0.2 | 0.7 | 7.6 | 21.6 | 23.7 | 13.1 | 9.1 | 8.2 | 9.1 |
| 2000..... | 10.6 | 3.2 | 0.4 | 0.2 | 0.7 | 8.2 | 21.5 | 24.3 | 13.1 | 9.1 | 10.0 | 10.6 |
| 1999..... | 10.7 | 4.1 | 0.6 | 0.3 | 0.6 | 9.8 | 22.4 | 22.7 | 12.2 | 9.4 | 9.7 | 13.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.2 | 2.6 | * | 0.1 | 0.6 | 3.4 | 7.4 | 7.9 | 4.7 | 4.7 | 5.6 | 7.2 |
| 2003..... | 4.3 | 3.6 | 0.4 | 0.2 | 0.6 | 3.2 | 7.5 | 7.8 | 4.7 | 5.2 | 6.0 | 8.7 |
| 2002..... | 4.4 | 3.8 | * | 0.2 | 0.6 | 3.9 | 7.9 | 7.6 | 4.6 | 5.6 | 6.2 | 9.3 |
| 2001..... | 4.3 | 2.8 | 0.4 | 0.2 | 0.8 | 3.8 | 7.8 | 6.8 | 4.8 | 5.9 | 6.1 | 7.7 |
| 2000..... | 4.4 | 3.0 | 0.4 | 0.2 | 0.6 | 4.7 | 7.6 | 6.8 | 4.5 | 5.9 | 6.2 | 9.1 |
| 1999..... | 4.3 | 3.1 | 0.4 | 0.2 | 0.6 | 4.5 | 7.5 | 6.1 | 4.4 | 5.9 | 6.8 | 11.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | 2.6 | 0.2 | * | 0.2 | 1.4 | 6.5 | 9.5 | 6.2 | 4.8 | 5.6 | 7.9 |
| 2003..... | 4.5 | 2.5 | 0.3 | * | 0.3 | 1.8 | 7.2 | 9.9 | 5.8 | 5.4 | 6.6 | 9.4 |
| 2002..... | 4.7 | 3.5 | 0.3 | 0.1 | 0.2 | 2.0 | 7.5 | 10.1 | 5.7 | 5.6 | 6.8 | 10.3 |
| 2001..... | 4.5 | 1.9 | 0.4 | 0.1 | 0.3 | 2.3 | 7.8 | 9.0 | 5.6 | 5.7 | 6.3 | 8.2 |
| 2000..... | 4.6 | 2.1 | 0.2 | 0.2 | 0.2 | 2.6 | 7.9 | 9.2 | 5.5 | 5.8 | 7.2 | 9.6 |
| 1999..... | 4.6 | 2.7 | 0.4 | 0.1 | 0.2 | 3.3 | 8.0 | 8.3 | 5.1 | 5.8 | 7.5 | 11.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.2 | 2.9 | * | * | 0.2 | 2.1 | 10.2 | 15.1 | 9.7 | 6.3 | 6.4 | 9.1 |
| 2003..... | 6.6 | 2.1 | * | * | 0.3 | 2.7 | 11.5 | 15.8 | 9.0 | 6.9 | 8.1 | 10.3 |
| 2002..... | 6.9 | 3.4 | * | * | 0.3 | 3.1 | 12.2 | 16.4 | 8.6 | 7.1 | 8.5 | 11.7 |
| 2001..... | 6.7 | 1.9 | * | * | 0.3 | 3.6 | 12.9 | 15.0 | 8.4 | 7.1 | 7.5 | 9.4 |
| 2000..... | 6.8 | 2.1 | * | * | 0.3 | 3.9 | 12.7 | 15.1 | 8.2 | 6.8 | 9.3 | 10.7 |
| 1999..... | 6.8 | 3.4 | 0.5 | 0.2 | 0.3 | 5.2 | 13.2 | 13.6 | 7.6 | 7.2 | 9.3 | 12.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.3 | 2.4 | * | * | 0.2 | 0.8 | 2.9 | 4.0 | 2.9 | 3.6 | 5.0 | 7.3 |
| 2003..... | 2.5 | 2.9 | * | * | 0.2 | 0.9 | 2.8 | 4.1 | 2.9 | 4.0 | 5.6 | 8.9 |
| 2002..... | 2.5 | 3.6 | * | * | 0.2 | 1.0 | 2.8 | 4.0 | 3.0 | 4.4 | 5.7 | 9.7 |
| 2001..... | 2.3 | 2.0 | * | * | 0.3 | 1.0 | 2.8 | 3.2 | 3.0 | 4.5 | 5.5 | 7.7 |
| 2000..... | 2.5 | 2.0 | * | 0.2 | 0.2 | 1.3 | 3.1 | 3.3 | 3.0 | 4.9 | 5.8 | 9.1 |
| 1999..... | 2.5 | 2.0 | * | * | 0.2 | 1.4 | 2.8 | 3.1 | 2.8 | 4.6 | 6.3 | 11.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral diseases (A80-B34) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 22.1 | 4.3 | * | 0.4 | 2.5 | 18.3 | 47.7 | 58.8 | 39.2 | 19.3 | 10.5 | 7.5 |
| 2003..... | 22.9 | 5.2 | * | 0.5 | 2.1 | 18.3 | 52.6 | 59.4 | 38.6 | 20.4 | 12.8 | 8.0 |
| 2002..... | 24.2 | 5.9 | * | 0.4 | 2.3 | 22.1 | 56.4 | 63.4 | 34.0 | 21.7 | 13.7 | 7.6 |
| 2001..... | 24.3 | 6.9 | 0.9 | 0.5 | 3.1 | 24.4 | 57.5 | 60.2 | 35.0 | 19.1 | 11.8 | 6.2 |
| 2000..... | 24.6 | 7.5 | 1.0 | 0.4 | 2.6 | 27.9 | 56.7 | 62.3 | 32.9 | 20.0 | 12.7 | 8.6 |
| 1999..... | 24.8 | 7.2 | 1.2 | 0.8 | 2.5 | 28.8 | 59.6 | 59.8 | 32.2 | 20.9 | 11.2 | 11.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 30.2 | * | * | * | 2.6 | 20.4 | 63.9 | 88.7 | 65.6 | 31.4 | 12.5 | * |
| 2003..... | 32.0 | * | * | * | 2.1 | 22.1 | 72.9 | 91.2 | 64.0 | 32.5 | 17.6 | * |
| 2002..... | 33.9 | 6.7 | * | * | 2.3 | 26.1 | 78.6 | 99.4 | 56.2 | 34.8 | 20.5 | * |
| 2001..... | 34.0 | 8.0 | * | * | 2.7 | 31.2 | 81.0 | 93.6 | 57.0 | 27.1 | 15.9 | * |
| 2000..... | 35.1 | 7.5 | * | * | 2.5 | 34.0 | 82.1 | 99.3 | 55.3 | 31.9 | 19.1 | * |
| 1999..... | 35.7 | 7.6 | * | 0.9 | 2.1 | 37.6 | 86.8 | 97.9 | 52.9 | 30.5 | 13.1 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 14.7 | * | * | * | 2.4 | 16.4 | 33.3 | 33.1 | 18.1 | 10.7 | 9.3 | * |
| 2003..... | 14.6 | * | * | * | 2.2 | 14.9 | 34.7 | 32.0 | 18.1 | 11.8 | 10.1 | * |
| 2002..... | 15.3 | * | * | * | 2.3 | 18.5 | 36.9 | 32.7 | 16.2 | 12.4 | 9.8 | * |
| 2001..... | 15.5 | * | * | * | 3.4 | 18.2 | 36.6 | 31.6 | 17.3 | 13.5 | 9.4 | * |
| 2000..... | 15.2 | 7.4 | * | * | 2.7 | 22.4 | 34.3 | 30.6 | 14.8 | 11.7 | 9.0 | 8.7 |
| 1999..... | 15.0 | * | * | 0.7 | 2.9 | 20.8 | 35.7 | 27.2 | 15.7 | 14.1 | 10.2 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Arthropod-borne viral encephalitis (A83-A84,A85.2)

Non-Hispanic 4/, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk (*)

Non-Hispanic 4/, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk (*)

Non-Hispanic 4/, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk (*)

Non-Hispanic white, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk (*)

Non-Hispanic white, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk (*)

Non-Hispanic white, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk (*)

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Zoster (herpes zoster)(B02) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.7 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 2.2 |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.9 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.5 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.5 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.9 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.1 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.5 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 2.2 |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 2.0 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Smallpox (B03)

Non-Hispanic 4/, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Non-Hispanic 4/, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Non-Hispanic 4/, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Non-Hispanic white, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Non-Hispanic white, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Non-Hispanic white, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Measles (B05)

All origins 3/, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Measles (B05) All origins 3/, both sexes.

All origins 3/, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Measles (B05) All origins 3/, male.

All origins 3/, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Measles (B05) All origins 3/, female.

Hispanic, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Measles (B05) Hispanic, both sexes.

Hispanic, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Measles (B05) Hispanic, male.

Hispanic, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Measles (B05) Hispanic, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Rubella (German measles) (B06)

Non-Hispanic 4/, both sexes

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic 4/, male

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic 4/, female

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic white, both sexes

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic white, male

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic white, female

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), showing asterisks for all data points.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral hepatitis (B15-B19) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.2 | 1.4 | 5.3 | 4.3 | 3.4 | 3.6 | 2.7 |
| 2003..... | 1.9 | * | * | * | * | 0.2 | 1.5 | 5.5 | 3.9 | 3.8 | 4.2 | 2.4 |
| 2002..... | 2.0 | * | * | * | 0.1 | 0.2 | 1.6 | 5.8 | 4.0 | 4.2 | 4.6 | 3.9 |
| 2001..... | 2.0 | * | * | * | 0.1 | 0.2 | 1.8 | 5.3 | 4.0 | 4.6 | 4.9 | 3.8 |
| 2000..... | 1.9 | * | * | * | * | 0.2 | 1.7 | 5.0 | 3.7 | 4.6 | 5.3 | 4.0 |
| 1999..... | 1.7 | * | * | * | * | 0.2 | 1.6 | 4.1 | 3.3 | 4.8 | 5.3 | 4.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | 0.2 | 1.8 | 8.0 | 6.1 | 4.0 | 3.6 | 3.0 |
| 2003..... | 2.5 | * | * | * | * | 0.2 | 2.1 | 8.3 | 5.4 | 4.2 | 4.8 | 2.8 |
| 2002..... | 2.7 | * | * | * | * | 0.3 | 2.3 | 8.9 | 5.5 | 4.5 | 5.3 | 5.0 |
| 2001..... | 2.5 | * | * | * | * | 0.2 | 2.5 | 8.2 | 5.3 | 4.8 | 5.3 | 4.4 |
| 2000..... | 2.4 | * | * | * | * | 0.2 | 2.4 | 7.7 | 5.0 | 4.8 | 6.4 | 5.7 |
| 1999..... | 2.2 | * | * | * | * | 0.3 | 2.3 | 6.3 | 4.3 | 5.1 | 6.6 | 6.9 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.1 | 1.0 | 2.7 | 2.5 | 3.0 | 3.5 | 2.5 |
| 2003..... | 1.3 | * | * | * | * | 0.1 | 0.9 | 2.9 | 2.5 | 3.5 | 3.8 | 2.3 |
| 2002..... | 1.4 | * | * | * | * | 0.2 | 1.0 | 2.8 | 2.7 | 3.9 | 4.1 | 3.3 |
| 2001..... | 1.4 | * | * | * | * | 0.2 | 1.1 | 2.5 | 2.8 | 4.4 | 4.6 | 3.5 |
| 2000..... | 1.4 | * | * | * | * | 0.2 | 1.1 | 2.4 | 2.6 | 4.5 | 4.7 | 3.3 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.9 | 2.0 | 2.4 | 4.5 | 4.5 | 3.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 1.8 | 6.8 | 7.7 | 7.0 | 7.5 | * |
| 2003..... | 1.8 | * | * | * | * | * | 1.8 | 7.0 | 7.2 | 8.7 | 9.2 | * |
| 2002..... | 2.0 | * | * | * | * | * | 1.7 | 8.0 | 8.3 | 8.6 | 10.5 | * |
| 2001..... | 2.1 | * | * | * | * | * | 2.3 | 7.8 | 10.4 | 9.9 | 11.0 | * |
| 2000..... | 2.0 | * | * | * | * | * | 2.5 | 7.0 | 8.6 | 11.1 | 13.0 | * |
| 1999..... | 1.9 | * | * | * | * | * | 2.1 | 6.8 | 7.8 | 12.4 | 13.5 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | 2.6 | 10.2 | 10.4 | 8.0 | * | * |
| 2003..... | 2.3 | * | * | * | * | * | 2.6 | 10.8 | 10.4 | 7.8 | 7.7 | * |
| 2002..... | 2.5 | * | * | * | * | * | 2.5 | 12.0 | 11.6 | 7.6 | 10.3 | * |
| 2001..... | 2.6 | * | * | * | * | * | 3.1 | 12.2 | 14.2 | 7.0 | 10.3 | * |
| 2000..... | 2.5 | * | * | * | * | * | 3.8 | 10.6 | 11.7 | 11.0 | 12.3 | * |
| 1999..... | 2.3 | * | * | * | * | * | 3.3 | 10.5 | 9.8 | 13.0 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.9 | 3.4 | 5.2 | 6.2 | 9.1 | * |
| 2003..... | 1.3 | * | * | * | * | * | 1.0 | 3.1 | 4.4 | 9.4 | 10.3 | * |
| 2002..... | 1.4 | * | * | * | * | * | 0.9 | 4.0 | 5.4 | 9.4 | 10.7 | * |
| 2001..... | 1.7 | * | * | * | * | * | 1.4 | 3.5 | 7.1 | 12.3 | 11.5 | * |
| 2000..... | 1.5 | * | * | * | * | * | 1.0 | 3.3 | 5.8 | 11.3 | 13.5 | * |
| 1999..... | 1.5 | * | * | * | * | * | 0.8 | 3.3 | 6.0 | 12.0 | 16.3 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral hepatitis (B15-B19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.2 | 1.3 | 5.2 | 4.0 | 3.2 | 3.3 | 2.6 |
| 2003..... | 1.9 | * | * | * | * | 0.2 | 1.4 | 5.4 | 3.6 | 3.4 | 4.0 | 2.3 |
| 2002..... | 2.0 | * | * | * | * | 0.2 | 1.6 | 5.6 | 3.6 | 3.8 | 4.3 | 3.7 |
| 2001..... | 1.9 | * | * | * | * | 0.2 | 1.7 | 5.0 | 3.5 | 4.2 | 4.6 | 3.6 |
| 2000..... | 1.9 | * | * | * | * | 0.2 | 1.6 | 4.8 | 3.3 | 4.2 | 5.0 | 3.8 |
| 1999..... | 1.7 | * | * | * | * | 0.2 | 1.5 | 3.8 | 3.0 | 4.3 | 5.0 | 4.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | 0.2 | 1.7 | 7.8 | 5.8 | 3.7 | 3.5 | 3.0 |
| 2003..... | 2.5 | * | * | * | * | 0.2 | 2.0 | 8.0 | 5.0 | 3.9 | 4.7 | 2.7 |
| 2002..... | 2.7 | * | * | * | * | 0.3 | 2.3 | 8.6 | 4.9 | 4.3 | 5.1 | 4.8 |
| 2001..... | 2.5 | * | * | * | * | 0.2 | 2.4 | 7.7 | 4.6 | 4.6 | 5.0 | 4.3 |
| 2000..... | 2.4 | * | * | * | * | 0.2 | 2.2 | 7.4 | 4.4 | 4.4 | 6.1 | 5.4 |
| 1999..... | 2.2 | * | * | * | * | 0.2 | 2.2 | 5.9 | 3.9 | 4.7 | 6.4 | 6.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.2 | 1.0 | 2.7 | 2.3 | 2.7 | 3.2 | 2.4 |
| 2003..... | 1.3 | * | * | * | * | 0.1 | 0.8 | 2.8 | 2.3 | 3.0 | 3.5 | 2.2 |
| 2002..... | 1.4 | * | * | * | * | 0.2 | 1.0 | 2.7 | 2.4 | 3.5 | 3.8 | 3.2 |
| 2001..... | 1.4 | * | * | * | * | 0.2 | 1.0 | 2.3 | 2.5 | 3.8 | 4.3 | 3.3 |
| 2000..... | 1.4 | * | * | * | * | 0.2 | 1.0 | 2.4 | 2.3 | 4.0 | 4.3 | 3.2 |
| 1999..... | 1.2 | * | * | * | * | 0.2 | 0.9 | 1.8 | 2.1 | 4.0 | 4.0 | 3.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.1 | 1.3 | 4.9 | 3.4 | 2.7 | 3.0 | 2.4 |
| 2003..... | 1.8 | * | * | * | * | 0.2 | 1.4 | 5.2 | 2.9 | 2.9 | 3.6 | 2.2 |
| 2002..... | 1.9 | * | * | * | * | 0.2 | 1.6 | 5.3 | 3.0 | 3.3 | 4.1 | 3.5 |
| 2001..... | 1.8 | * | * | * | * | 0.2 | 1.7 | 4.6 | 2.9 | 3.7 | 4.3 | 3.3 |
| 2000..... | 1.8 | * | * | * | * | 0.1 | 1.6 | 4.3 | 2.7 | 3.7 | 4.7 | 3.7 |
| 1999..... | 1.6 | * | * | * | * | 0.2 | 1.4 | 3.4 | 2.5 | 3.7 | 4.7 | 4.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 1.7 | 7.4 | 4.8 | 3.2 | 3.2 | 2.7 |
| 2003..... | 2.4 | * | * | * | * | 0.2 | 2.0 | 7.7 | 4.2 | 3.4 | 4.2 | 2.7 |
| 2002..... | 2.6 | * | * | * | * | 0.3 | 2.3 | 8.2 | 4.1 | 3.9 | 4.9 | 4.6 |
| 2001..... | 2.5 | * | * | * | * | 0.2 | 2.3 | 7.2 | 4.0 | 4.3 | 5.1 | 4.4 |
| 2000..... | 2.3 | * | * | * | * | * | 2.1 | 6.7 | 3.6 | 3.8 | 5.9 | 5.4 |
| 1999..... | 2.1 | * | * | * | * | 0.2 | 2.1 | 5.2 | 3.4 | 4.3 | 6.4 | 6.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.9 | 2.5 | 2.0 | 2.3 | 2.8 | 2.3 |
| 2003..... | 1.2 | * | * | * | * | * | 0.8 | 2.6 | 1.8 | 2.5 | 3.1 | 2.0 |
| 2002..... | 1.3 | * | * | * | * | * | 0.9 | 2.5 | 2.0 | 2.8 | 3.5 | 3.1 |
| 2001..... | 1.2 | * | * | * | * | * | 1.0 | 2.0 | 2.0 | 3.1 | 3.8 | 2.9 |
| 2000..... | 1.3 | * | * | * | * | * | 1.0 | 2.0 | 1.8 | 3.6 | 4.0 | 3.1 |
| 1999..... | 1.1 | * | * | * | * | 0.2 | 0.8 | 1.7 | 1.6 | 3.3 | 3.6 | 3.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral hepatitis (B15-B19) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | 1.4 | 7.6 | 8.8 | 4.6 | 5.0 | * |
| 2003..... | 2.3 | * | * | * | * | * | 1.8 | 7.6 | 8.2 | 5.3 | 5.9 | * |
| 2002..... | 2.3 | * | * | * | * | 0.4 | 2.0 | 8.2 | 7.2 | 6.0 | 5.2 | * |
| 2001..... | 2.3 | * | * | * | * | * | 2.2 | 8.1 | 6.8 | 5.6 | 5.7 | * |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 2.1 | 8.0 | 5.8 | 5.8 | 6.3 | * |
| 1999..... | 2.1 | * | * | * | * | * | 2.2 | 7.0 | 5.1 | 6.8 | 5.9 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 1.5 | 11.9 | 14.1 | 6.7 | * | * |
| 2003..... | 3.1 | * | * | * | * | * | 2.4 | 11.7 | 12.1 | 5.9 | 6.1 | * |
| 2002..... | 3.1 | * | * | * | * | * | 2.6 | 12.6 | 10.7 | 7.0 | * | * |
| 2001..... | 2.8 | * | * | * | * | * | 3.0 | 12.0 | 9.0 | 5.0 | * | * |
| 2000..... | 2.9 | * | * | * | * | * | 3.0 | 12.3 | 8.1 | 7.2 | 7.7 | * |
| 1999..... | 2.6 | * | * | * | * | * | 3.1 | 11.8 | 6.3 | 5.7 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 1.4 | 3.9 | 4.5 | 3.1 | 5.3 | * |
| 2003..... | 1.6 | * | * | * | * | * | 1.2 | 4.1 | 5.1 | 4.9 | 5.9 | * |
| 2002..... | 1.7 | * | * | * | * | * | 1.5 | 4.4 | 4.4 | 5.2 | 5.3 | * |
| 2001..... | 1.8 | * | * | * | * | * | 1.4 | 4.7 | 5.0 | 6.0 | 6.7 | * |
| 2000..... | 1.6 | * | * | * | * | * | 1.4 | 4.3 | 3.9 | 4.7 | 5.6 | * |
| 1999..... | 1.6 | * | * | * | * | * | 1.4 | 2.9 | 4.1 | 7.5 | 6.7 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease (B20-B24) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | 0.1 | 0.5 | 3.7 | 10.9 | 10.6 | 5.4 | 2.4 | 0.8 | * |
| 2003..... | 4.7 | * | * | 0.1 | 0.4 | 4.0 | 12.0 | 10.9 | 5.4 | 2.4 | 0.7 | * |
| 2002..... | 4.9 | * | * | 0.1 | 0.4 | 4.6 | 12.7 | 11.2 | 5.1 | 2.2 | 0.8 | * |
| 2001..... | 5.0 | * | * | 0.1 | 0.6 | 5.3 | 13.0 | 10.5 | 5.2 | 2.1 | 0.7 | * |
| 2000..... | 5.1 | * | * | 0.1 | 0.5 | 6.1 | 13.1 | 11.0 | 5.1 | 2.2 | 0.7 | * |
| 1999..... | 5.3 | * | 0.2 | 0.2 | 0.5 | 6.8 | 13.8 | 10.7 | 4.8 | 2.2 | 0.6 | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.6 | * | * | * | 0.5 | 4.5 | 15.7 | 16.3 | 9.0 | 4.0 | 1.4 | * |
| 2003..... | 7.1 | * | * | * | 0.4 | 5.1 | 17.5 | 17.2 | 9.1 | 4.0 | 1.5 | * |
| 2002..... | 7.4 | * | * | * | 0.4 | 5.9 | 18.8 | 17.7 | 8.5 | 3.9 | 1.4 | * |
| 2001..... | 7.6 | * | * | 0.1 | 0.5 | 7.1 | 19.5 | 16.8 | 8.6 | 3.5 | 1.5 | * |
| 2000..... | 7.9 | * | * | 0.1 | 0.5 | 8.0 | 19.8 | 17.8 | 8.7 | 3.8 | 1.3 | * |
| 1999..... | 8.2 | * | * | 0.2 | 0.5 | 9.5 | 21.0 | 17.5 | 8.3 | 3.8 | 1.0 | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | 0.4 | 2.8 | 6.2 | 5.2 | 2.0 | 1.0 | 0.5 | * |
| 2003..... | 2.4 | * | * | * | 0.4 | 2.8 | 6.5 | 4.8 | 2.1 | 1.0 | 0.3 | * |
| 2002..... | 2.5 | * | * | * | 0.4 | 3.3 | 6.7 | 4.8 | 1.9 | 0.8 | 0.3 | * |
| 2001..... | 2.5 | * | * | * | 0.6 | 3.5 | 6.7 | 4.4 | 2.0 | 0.9 | * | * |
| 2000..... | 2.5 | * | * | 0.1 | 0.4 | 4.2 | 6.5 | 4.4 | 1.8 | 0.8 | 0.3 | * |
| 1999..... | 2.5 | * | * | 0.2 | 0.5 | 4.1 | 6.7 | 4.1 | 1.6 | 0.8 | 0.3 | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | 0.4 | 2.8 | 11.0 | 14.1 | 8.4 | 5.0 | * | * |
| 2003..... | 4.7 | * | * | * | 0.4 | 3.3 | 12.3 | 15.9 | 9.8 | 5.2 | * | * |
| 2002..... | 4.8 | * | * | * | 0.4 | 3.6 | 13.4 | 14.7 | 9.5 | 3.4 | * | * |
| 2001..... | 5.0 | * | * | * | 0.5 | 4.4 | 13.9 | 15.5 | 9.2 | 4.5 | * | * |
| 2000..... | 5.4 | * | * | * | 0.3 | 5.6 | 14.9 | 16.9 | 9.5 | 3.8 | * | * |
| 1999..... | 5.8 | * | * | * | 0.5 | 6.4 | 17.3 | 15.4 | 8.4 | 4.7 | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | * | * | 4.1 | 16.1 | 22.0 | 14.4 | 8.0 | * | * |
| 2003..... | 7.1 | * | * | * | 0.5 | 4.6 | 17.7 | 26.1 | 15.4 | 9.2 | * | * |
| 2002..... | 7.2 | * | * | * | * | 5.1 | 20.1 | 23.1 | 15.0 | 6.9 | * | * |
| 2001..... | 7.6 | * | * | * | 0.6 | 6.0 | 20.8 | 24.9 | 14.8 | 8.6 | * | * |
| 2000..... | 8.2 | * | * | * | * | 7.8 | 22.7 | 27.5 | 15.3 | 7.2 | * | * |
| 1999..... | 8.7 | * | * | * | 0.6 | 9.3 | 26.2 | 24.2 | 15.4 | 9.0 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 1.2 | 5.3 | 6.2 | 2.9 | * | * | * |
| 2003..... | 2.2 | * | * | * | * | 1.8 | 6.3 | 5.6 | 4.6 | * | * | * |
| 2002..... | 2.2 | * | * | * | * | 1.8 | 6.2 | 6.3 | 4.6 | * | * | * |
| 2001..... | 2.3 | * | * | * | * | 2.5 | 6.5 | 6.2 | 4.1 | * | * | * |
| 2000..... | 2.4 | * | * | * | * | 3.0 | 6.4 | 6.6 | 4.4 | * | * | * |
| 1999..... | 2.6 | * | * | * | * | 3.2 | 7.9 | 6.7 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease (B20-B24) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | 0.1 | 0.5 | 3.9 | 10.9 | 10.1 | 5.0 | 2.1 | 0.7 | * |
| 2003..... | 4.6 | * | * | 0.1 | 0.4 | 4.1 | 11.9 | 10.2 | 5.0 | 2.1 | 0.7 | * |
| 2002..... | 4.8 | * | * | 0.1 | 0.4 | 4.8 | 12.5 | 10.6 | 4.7 | 2.1 | 0.7 | * |
| 2001..... | 4.9 | * | * | 0.1 | 0.6 | 5.4 | 12.7 | 9.9 | 4.8 | 1.9 | 0.6 | * |
| 2000..... | 5.0 | * | * | 0.1 | 0.5 | 6.2 | 12.7 | 10.3 | 4.7 | 2.0 | 0.6 | * |
| 1999..... | 5.2 | * | * | 0.2 | 0.5 | 6.8 | 13.2 | 10.0 | 4.5 | 2.0 | 0.6 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | 0.5 | 4.6 | 15.5 | 15.5 | 8.5 | 3.6 | 1.2 | * |
| 2003..... | 7.0 | * | * | * | 0.4 | 5.2 | 17.3 | 16.0 | 8.4 | 3.6 | 1.4 | * |
| 2002..... | 7.3 | * | * | * | 0.4 | 6.0 | 18.4 | 16.9 | 7.9 | 3.7 | 1.3 | * |
| 2001..... | 7.5 | * | * | 0.1 | 0.5 | 7.3 | 19.0 | 15.8 | 8.0 | 3.1 | 1.3 | * |
| 2000..... | 7.7 | * | * | * | 0.5 | 7.9 | 19.1 | 16.7 | 8.0 | 3.4 | 1.2 | * |
| 1999..... | 8.0 | * | * | 0.2 | 0.5 | 9.4 | 20.1 | 16.5 | 7.6 | 3.4 | 1.0 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | 0.5 | 3.1 | 6.3 | 5.0 | 1.9 | 0.9 | 0.4 | * |
| 2003..... | 2.4 | * | * | * | 0.4 | 3.0 | 6.5 | 4.7 | 1.9 | 0.9 | * | * |
| 2002..... | 2.5 | * | * | * | 0.5 | 3.6 | 6.7 | 4.7 | 1.7 | 0.8 | 0.3 | * |
| 2001..... | 2.5 | * | * | * | 0.7 | 3.6 | 6.6 | 4.2 | 1.8 | 0.9 | * | * |
| 2000..... | 2.5 | * | * | 0.1 | 0.5 | 4.4 | 6.4 | 4.2 | 1.6 | 0.8 | * | * |
| 1999..... | 2.5 | * | * | 0.1 | 0.6 | 4.2 | 6.5 | 3.9 | 1.6 | 0.7 | 0.3 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | 0.1 | 1.2 | 5.1 | 4.3 | 2.2 | 0.9 | 0.4 | * |
| 2003..... | 2.1 | * | * | * | 0.1 | 1.5 | 5.6 | 4.5 | 2.2 | 0.8 | 0.3 | * |
| 2002..... | 2.1 | * | * | * | 0.1 | 1.7 | 5.8 | 4.6 | 2.2 | 0.8 | 0.3 | * |
| 2001..... | 2.2 | * | * | * | 0.1 | 2.1 | 6.0 | 4.2 | 2.1 | 0.7 | 0.3 | * |
| 2000..... | 2.3 | * | * | * | 0.1 | 2.4 | 6.2 | 4.6 | 2.2 | 0.8 | 0.3 | * |
| 1999..... | 2.4 | * | * | * | 0.1 | 3.0 | 6.4 | 4.6 | 2.0 | 0.8 | 0.3 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 1.8 | 8.4 | 7.4 | 4.0 | 1.7 | 0.7 | * |
| 2003..... | 3.6 | * | * | * | * | 2.4 | 9.4 | 7.8 | 4.0 | 1.5 | 0.8 | * |
| 2002..... | 3.7 | * | * | * | * | 2.7 | 9.8 | 8.0 | 4.0 | 1.5 | 0.5 | * |
| 2001..... | 3.8 | * | * | * | * | 3.2 | 10.4 | 7.5 | 3.8 | 1.4 | 0.6 | * |
| 2000..... | 4.0 | * | * | * | 0.2 | 3.7 | 10.4 | 8.2 | 4.0 | 1.5 | 0.6 | * |
| 1999..... | 4.2 | * | * | * | 0.2 | 4.9 | 10.9 | 8.2 | 3.6 | 1.5 | 0.6 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | 0.6 | 1.8 | 1.3 | 0.4 | * | * | * |
| 2003..... | 0.6 | * | * | * | * | 0.7 | 1.8 | 1.2 | 0.5 | 0.3 | * | * |
| 2002..... | 0.6 | * | * | * | * | 0.8 | 1.8 | 1.2 | 0.5 | * | * | * |
| 2001..... | 0.6 | * | * | * | * | 0.9 | 1.7 | 1.0 | 0.5 | * | * | * |
| 2000..... | 0.7 | * | * | * | * | 1.1 | 2.0 | 1.1 | 0.5 | * | * | * |
| 1999..... | 0.7 | * | * | * | * | 1.2 | 1.9 | 1.1 | 0.4 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease (B20-B24) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 19.4 | * | * | * | 2.3 | 17.9 | 46.0 | 50.9 | 29.9 | 13.6 | 4.6 | * |
| 2003..... | 20.1 | * | * | 0.5 | 1.9 | 18.1 | 50.5 | 51.3 | 29.9 | 14.4 | 4.7 | * |
| 2002..... | 21.3 | * | * | * | 2.1 | 21.5 | 54.0 | 54.8 | 26.3 | 14.6 | 5.7 | * |
| 2001..... | 21.6 | * | * | 0.4 | 2.8 | 24.0 | 55.0 | 51.9 | 28.0 | 12.8 | 5.3 | * |
| 2000..... | 22.0 | * | * | 0.4 | 2.2 | 27.2 | 54.5 | 53.9 | 26.3 | 13.2 | 4.6 | * |
| 1999..... | 22.3 | * | * | 0.6 | 2.3 | 28.3 | 57.2 | 52.2 | 26.4 | 13.2 | 3.7 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 26.7 | * | * | * | 2.4 | 20.0 | 62.1 | 76.3 | 50.8 | 23.4 | 7.7 | * |
| 2003..... | 28.5 | * | * | * | 1.9 | 21.7 | 70.0 | 79.0 | 51.4 | 25.3 | 8.9 | * |
| 2002..... | 30.3 | * | * | * | 2.0 | 25.5 | 75.6 | 86.2 | 44.8 | 26.6 | 12.3 | * |
| 2001..... | 30.8 | * | * | * | 2.4 | 30.8 | 77.8 | 81.4 | 47.7 | 21.4 | 11.4 | * |
| 2000..... | 31.7 | * | * | * | 2.2 | 33.4 | 79.0 | 86.6 | 46.4 | 23.8 | 8.9 | * |
| 1999..... | 32.6 | * | * | 0.7 | 1.9 | 37.1 | 83.5 | 85.5 | 45.6 | 24.0 | 6.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 12.8 | * | * | * | 2.2 | 15.9 | 31.6 | 29.0 | 13.1 | 6.6 | * | * |
| 2003..... | 12.6 | * | * | * | 2.0 | 14.8 | 33.2 | 27.6 | 12.6 | 6.6 | * | * |
| 2002..... | 13.2 | * | * | * | 2.2 | 17.9 | 34.9 | 28.0 | 11.5 | 6.0 | * | * |
| 2001..... | 13.3 | * | * | * | 3.2 | 17.8 | 34.8 | 26.6 | 12.1 | 6.7 | * | * |
| 2000..... | 13.1 | * | * | * | 2.3 | 21.7 | 32.8 | 25.8 | 10.1 | 5.8 | * | * |
| 1999..... | 13.0 | * | * | * | 2.8 | 20.3 | 34.1 | 23.8 | 11.0 | 5.6 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.2 | 1.5 | 4.4 | 4.2 | 1.9 | 0.9 | 0.3 | * |
| 2003..... | 1.8 | * | * | * | 0.2 | 1.6 | 4.5 | 4.3 | 2.0 | 0.8 | 0.2 | * |
| 2002..... | 1.9 | * | * | * | 0.2 | 1.9 | 5.1 | 4.5 | 1.9 | 0.7 | 0.2 | * |
| 2001..... | 1.9 | * | * | * | 0.2 | 2.1 | 5.1 | 4.2 | 1.9 | 0.7 | 0.2 | * |
| 2000..... | 2.0 | * | * | * | 0.2 | 2.4 | 5.3 | 4.4 | 1.8 | 0.8 | 0.3 | * |
| 1999..... | 2.1 | * | * | 0.1 | 0.2 | 2.8 | 5.6 | 4.4 | 1.6 | 0.7 | 0.3 | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 0.2 | 1.9 | 6.3 | 6.2 | 3.2 | 1.5 | 0.5 | * |
| 2003..... | 2.7 | * | * | * | 0.2 | 2.1 | 6.5 | 6.9 | 3.3 | 1.3 | 0.4 | * |
| 2002..... | 2.9 | * | * | * | 0.2 | 2.3 | 7.5 | 7.2 | 3.0 | 1.3 | 0.5 | * |
| 2001..... | 2.9 | * | * | * | 0.3 | 2.8 | 7.6 | 6.7 | 3.1 | 1.2 | 0.5 | * |
| 2000..... | 3.1 | * | * | * | 0.2 | 3.2 | 7.8 | 7.1 | 2.9 | 1.4 | 0.5 | * |
| 1999..... | 3.3 | * | * | * | 0.2 | 3.8 | 8.4 | 7.2 | 2.7 | 1.3 | 0.6 | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | 0.2 | 1.2 | 2.4 | 2.2 | 0.7 | 0.4 | * | * |
| 2003..... | 0.9 | * | * | * | 0.1 | 1.1 | 2.5 | 1.8 | 0.9 | 0.4 | * | * |
| 2002..... | 1.0 | * | * | * | 0.2 | 1.4 | 2.7 | 1.9 | 0.8 | 0.3 | * | * |
| 2001..... | 1.0 | * | * | * | 0.2 | 1.5 | 2.6 | 1.8 | 0.8 | 0.4 | * | * |
| 2000..... | 1.0 | * | * | * | 0.2 | 1.6 | 2.8 | 1.7 | 0.7 | 0.2 | * | * |
| 1999..... | 1.0 | * | * | * | 0.3 | 1.8 | 2.7 | 1.8 | 0.5 | 0.3 | * | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 1.3 | 5.2 | 5.9 | 3.8 | 2.3 | * | * |
| 2003..... | 2.1 | * | * | * | * | 1.5 | 5.3 | 7.4 | 4.4 | 2.0 | * | * |
| 2002..... | 2.2 | * | * | * | * | 1.7 | 6.1 | 6.4 | 4.6 | * | * | * |
| 2001..... | 2.3 | * | * | * | * | 2.0 | 6.5 | 7.0 | 4.2 | * | * | * |
| 2000..... | 2.5 | * | * | * | * | 2.7 | 7.0 | 7.6 | 3.8 | * | * | * |
| 1999..... | 2.6 | * | * | * | * | 3.1 | 7.6 | 7.1 | 3.0 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 1.9 | 7.8 | 8.8 | 6.5 | 3.5 | * | * |
| 2003..... | 3.2 | * | * | * | * | 2.1 | 7.8 | 11.9 | 6.5 | * | * | * |
| 2002..... | 3.2 | * | * | * | * | 2.2 | 8.9 | 10.1 | 7.3 | * | * | * |
| 2001..... | 3.4 | * | * | * | * | 2.9 | 9.5 | 11.2 | 6.1 | * | * | * |
| 2000..... | 3.7 | * | * | * | * | 3.8 | 10.6 | 12.3 | 5.5 | * | * | * |
| 1999..... | 3.8 | * | * | * | * | 4.2 | 11.1 | 11.0 | 5.6 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 2.3 | 2.9 | * | * | * | * |
| 2003..... | 1.0 | * | * | * | * | 0.8 | 2.6 | 2.8 | 2.6 | * | * | * |
| 2002..... | 1.0 | * | * | * | * | 1.0 | 2.9 | 2.8 | 2.1 | * | * | * |
| 2001..... | 1.1 | * | * | * | * | 1.0 | 3.2 | 2.9 | 2.5 | * | * | * |
| 2000..... | 1.2 | * | * | * | * | 1.4 | 3.2 | 3.0 | 2.3 | * | * | * |
| 1999..... | 1.3 | * | * | * | * | 1.8 | 3.8 | 3.3 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.2 | 1.6 | 4.2 | 3.9 | 1.7 | 0.8 | 0.3 | * |
| 2003..... | 1.7 | * | * | * | 0.2 | 1.6 | 4.3 | 3.9 | 1.8 | 0.7 | 0.2 | * |
| 2002..... | 1.9 | * | * | * | 0.2 | 1.9 | 4.9 | 4.2 | 1.7 | 0.7 | 0.2 | * |
| 2001..... | 1.9 | * | * | * | 0.2 | 2.1 | 4.8 | 3.9 | 1.7 | 0.7 | 0.2 | * |
| 2000..... | 1.9 | * | * | * | 0.2 | 2.4 | 5.0 | 4.0 | 1.6 | 0.7 | 0.2 | * |
| 1999..... | 2.0 | * | * | 0.1 | 0.2 | 2.7 | 5.2 | 4.1 | 1.4 | 0.7 | 0.2 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | 0.2 | 1.8 | 6.0 | 5.9 | 2.8 | 1.3 | 0.5 | * |
| 2003..... | 2.6 | * | * | * | 0.1 | 2.1 | 6.3 | 6.3 | 3.0 | 1.2 | * | * |
| 2002..... | 2.8 | * | * | * | 0.2 | 2.3 | 7.1 | 6.7 | 2.7 | 1.2 | 0.5 | * |
| 2001..... | 2.8 | * | * | * | 0.2 | 2.8 | 7.2 | 6.1 | 2.8 | 1.0 | 0.4 | * |
| 2000..... | 2.9 | * | * | * | 0.2 | 3.1 | 7.4 | 6.5 | 2.6 | 1.3 | 0.4 | * |
| 1999..... | 3.1 | * | * | * | 0.2 | 3.6 | 8.0 | 6.7 | 2.4 | 1.2 | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | 0.2 | 1.3 | 2.4 | 2.1 | 0.7 | 0.4 | * | * |
| 2003..... | 0.9 | * | * | * | 0.2 | 1.1 | 2.5 | 1.7 | 0.7 | 0.3 | * | * |
| 2002..... | 1.0 | * | * | * | 0.2 | 1.5 | 2.7 | 1.8 | 0.7 | 0.3 | * | * |
| 2001..... | 1.0 | * | * | * | 0.2 | 1.5 | 2.5 | 1.7 | 0.7 | 0.4 | * | * |
| 2000..... | 1.0 | * | * | * | 0.2 | 1.7 | 2.7 | 1.6 | 0.6 | 0.2 | * | * |
| 1999..... | 1.0 | * | * | * | 0.3 | 1.8 | 2.6 | 1.6 | 0.5 | 0.3 | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | 0.5 | 2.0 | 1.8 | 0.7 | 0.3 | * | * |
| 2003..... | 0.8 | * | * | * | * | 0.6 | 2.2 | 1.8 | 0.8 | 0.3 | * | * |
| 2002..... | 0.8 | * | * | * | * | 0.7 | 2.4 | 1.8 | 0.8 | 0.3 | * | * |
| 2001..... | 0.8 | * | * | * | * | 0.9 | 2.3 | 1.7 | 0.8 | 0.3 | * | * |
| 2000..... | 0.9 | * | * | * | 0.1 | 0.9 | 2.4 | 1.8 | 0.7 | 0.3 | * | * |
| 1999..... | 0.9 | * | * | * | * | 1.2 | 2.5 | 1.8 | 0.7 | 0.3 | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.7 | 3.3 | 2.9 | 1.3 | 0.6 | * | * |
| 2003..... | 1.3 | * | * | * | * | 1.0 | 3.6 | 3.0 | 1.3 | 0.5 | * | * |
| 2002..... | 1.4 | * | * | * | * | 1.0 | 3.8 | 3.1 | 1.4 | 0.5 | * | * |
| 2001..... | 1.4 | * | * | * | * | 1.3 | 3.9 | 3.0 | 1.4 | 0.4 | * | * |
| 2000..... | 1.5 | * | * | * | * | 1.4 | 3.9 | 3.2 | 1.3 | 0.6 | * | * |
| 1999..... | 1.6 | * | * | * | * | 1.8 | 4.2 | 3.2 | 1.2 | 0.5 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | 0.3 | 0.8 | 0.6 | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | 0.3 | 0.8 | 0.5 | 0.2 | * | * | * |
| 2002..... | 0.3 | * | * | * | * | 0.4 | 0.9 | 0.5 | 0.3 | * | * | * |
| 2001..... | 0.3 | * | * | * | * | 0.4 | 0.8 | 0.4 | 0.2 | * | * | * |
| 2000..... | 0.3 | * | * | * | * | 0.4 | 0.9 | 0.5 | 0.2 | * | * | * |
| 1999..... | 0.3 | * | * | * | * | 0.5 | 0.8 | 0.5 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 7.4 | * | * | * | 1.0 | 7.3 | 17.4 | 19.1 | 10.9 | 5.3 | * | * |
| 2003..... | 7.3 | * | * | * | 0.7 | 7.0 | 17.5 | 19.3 | 10.9 | 4.7 | * | * |
| 2002..... | 8.2 | * | * | * | 0.9 | 8.4 | 20.5 | 22.2 | 9.2 | 4.6 | * | * |
| 2001..... | 8.2 | * | * | * | 1.1 | 9.2 | 20.5 | 20.3 | 9.9 | 4.5 | * | * |
| 2000..... | 8.5 | * | * | * | 1.0 | 10.6 | 21.7 | 20.6 | 8.8 | 4.6 | * | * |
| 1999..... | 8.8 | * | * | * | 1.2 | 11.5 | 22.6 | 21.6 | 8.2 | 4.4 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 10.0 | * | * | * | 1.0 | 7.9 | 23.6 | 27.8 | 18.0 | 8.8 | * | * |
| 2003..... | 10.4 | * | * | * | 0.7 | 8.6 | 23.9 | 30.8 | 18.7 | 8.1 | * | * |
| 2002..... | 11.7 | * | * | * | 0.9 | 9.9 | 29.1 | 35.5 | 15.1 | 7.8 | * | * |
| 2001..... | 11.6 | * | * | * | 1.2 | 11.2 | 29.5 | 31.3 | 16.6 | 7.4 | * | * |
| 2000..... | 12.2 | * | * | * | 0.9 | 13.4 | 31.0 | 33.2 | 15.1 | 8.8 | * | * |
| 1999..... | 12.9 | * | * | * | 0.8 | 14.5 | 33.5 | 35.2 | 14.2 | 8.1 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | 1.1 | 6.7 | 11.9 | 11.6 | 5.2 | 2.8 | * | * |
| 2003..... | 4.4 | * | * | * | 0.7 | 5.5 | 11.8 | 9.4 | 4.7 | 2.3 | * | * |
| 2002..... | 5.0 | * | * | * | 0.8 | 7.0 | 13.0 | 10.7 | 4.4 | 2.3 | * | * |
| 2001..... | 5.1 | * | * | * | 1.1 | 7.4 | 12.5 | 10.9 | 4.5 | 2.4 | * | * |
| 2000..... | 5.1 | * | * | * | 1.1 | 8.1 | 13.4 | 9.7 | 3.8 | * | * | * |
| 1999..... | 5.2 | * | * | * | 1.5 | 8.7 | 13.0 | 10.0 | 3.4 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.7 | 2.3 | 2.9 | 2.0 | 1.2 | * | * |
| 2003..... | 1.0 | * | * | * | * | 0.8 | 2.3 | 2.6 | 2.2 | * | * | * |
| 2002..... | 1.1 | * | * | * | * | 0.8 | 2.7 | 3.1 | 1.4 | * | * | * |
| 2001..... | 1.1 | * | * | * | * | 1.2 | 2.7 | 2.5 | 2.5 | * | * | * |
| 2000..... | 1.2 | * | * | * | * | 1.1 | 3.1 | 2.9 | 1.6 | * | * | * |
| 1999..... | 1.2 | * | * | * | * | 1.5 | 3.0 | 2.8 | 1.9 | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.9 | 3.6 | 4.6 | 3.8 | * | * | * |
| 2003..... | 1.4 | * | * | * | * | 1.1 | 3.5 | 3.5 | 3.5 | * | * | * |
| 2002..... | 1.7 | * | * | * | * | 0.9 | 4.1 | 5.0 | 2.5 | * | * | * |
| 2001..... | 1.8 | * | * | * | * | 1.8 | 4.4 | 4.0 | 4.5 | * | * | * |
| 2000..... | 1.9 | * | * | * | * | 1.6 | 5.1 | 4.9 | 3.2 | * | * | * |
| 1999..... | 2.0 | * | * | * | * | 2.3 | 5.1 | 4.9 | 3.2 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 1.2 | 1.4 | * | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | 1.2 | 1.8 | * | * | * | * |
| 2002..... | 0.6 | * | * | * | * | * | 1.4 | 1.5 | * | * | * | * |
| 2001..... | 0.6 | * | * | * | * | * | 1.1 | 1.3 | * | * | * | * |
| 2000..... | 0.5 | * | * | * | * | 0.7 | 1.3 | 1.2 | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | 0.8 | 1.2 | 1.0 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.4 | 1.0 | 1.0 | 0.6 | 0.2 | * | * |
| 2003..... | 0.4 | * | * | * | * | 0.3 | 1.2 | 1.0 | 0.5 | 0.2 | * | * |
| 2002..... | 0.5 | * | * | * | * | 0.5 | 1.1 | 0.9 | 0.5 | 0.2 | * | * |
| 2001..... | 0.5 | * | * | * | * | 0.6 | 1.2 | 0.9 | 0.6 | 0.3 | * | * |
| 2000..... | 0.6 | * | * | * | * | 0.7 | 1.4 | 1.1 | 0.6 | 0.2 | * | * |
| 1999..... | 0.6 | * | * | * | * | 0.7 | 1.5 | 1.0 | 0.5 | 0.2 | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.4 | 1.5 | 1.6 | 0.9 | 0.3 | * | * |
| 2003..... | 0.7 | * | * | * | * | 0.5 | 1.8 | 1.6 | 0.9 | 0.4 | * | * |
| 2002..... | 0.7 | * | * | * | * | 0.7 | 1.8 | 1.5 | 1.0 | 0.4 | * | * |
| 2001..... | 0.8 | * | * | * | * | 0.8 | 2.0 | 1.5 | 0.9 | 0.4 | * | * |
| 2000..... | 0.9 | * | * | * | * | 1.0 | 2.3 | 1.9 | 1.0 | 0.3 | * | * |
| 1999..... | 0.9 | * | * | * | * | 1.1 | 2.4 | 1.7 | 0.9 | 0.4 | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.3 | 0.5 | 0.4 | 0.2 | * | * | * |
| 2003..... | 0.2 | * | * | * | * | 0.2 | 0.5 | 0.4 | * | * | * | * |
| 2002..... | 0.2 | * | * | * | * | 0.4 | 0.5 | 0.4 | * | * | * | * |
| 2001..... | 0.2 | * | * | * | * | 0.3 | 0.5 | 0.3 | 0.2 | * | * | * |
| 2000..... | 0.2 | * | * | * | * | 0.4 | 0.6 | 0.4 | * | * | * | * |
| 1999..... | 0.2 | * | * | * | * | 0.4 | 0.6 | 0.3 | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.1 | 0.5 | 0.6 | 0.3 | * | * | * |
| 2003..... | 0.2 | * | * | * | * | 0.2 | 0.6 | 0.5 | 0.2 | 0.1 | * | * |
| 2002..... | 0.2 | * | * | * | * | 0.2 | 0.6 | 0.5 | 0.3 | * | * | * |
| 2001..... | 0.2 | * | * | * | * | 0.3 | 0.6 | 0.5 | 0.3 | * | * | * |
| 2000..... | 0.3 | * | * | * | * | 0.3 | 0.9 | 0.6 | 0.3 | * | * | * |
| 1999..... | 0.3 | * | * | * | * | 0.4 | 0.9 | 0.6 | 0.2 | * | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.2 | 0.9 | 1.0 | 0.5 | * | * | * |
| 2003..... | 0.4 | * | * | * | * | 0.2 | 1.1 | 1.0 | 0.4 | * | * | * |
| 2002..... | 0.4 | * | * | * | * | 0.4 | 1.1 | 1.0 | 0.6 | * | * | * |
| 2001..... | 0.4 | * | * | * | * | 0.4 | 1.1 | 0.9 | 0.5 | * | * | * |
| 2000..... | 0.5 | * | * | * | * | 0.6 | 1.5 | 1.0 | 0.5 | * | * | * |
| 1999..... | 0.6 | * | * | * | * | 0.7 | 1.5 | 1.2 | 0.4 | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.2 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 1.5 | 4.0 | 4.5 | 3.0 | 1.2 | * | * |
| 2003..... | 1.7 | * | * | * | * | 1.4 | 4.5 | 4.3 | 3.1 | * | * | * |
| 2002..... | 1.7 | * | * | * | * | 2.1 | 4.4 | 3.8 | 2.7 | 1.5 | * | * |
| 2001..... | 2.0 | * | * | * | * | 2.3 | 5.0 | 4.0 | 3.0 | 1.8 | * | * |
| 2000..... | 2.2 | * | * | * | * | 2.9 | 5.3 | 5.3 | 3.0 | 1.4 | * | * |
| 1999..... | 2.2 | * | * | * | * | 2.8 | 6.0 | 4.1 | 3.1 | 1.4 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | 1.5 | 5.5 | 6.9 | 5.3 | * | * | * |
| 2003..... | 2.6 | * | * | * | * | 1.7 | 6.8 | 6.7 | 5.7 | * | * | * |
| 2002..... | 2.5 | * | * | * | * | 2.4 | 6.5 | 5.9 | 5.1 | 2.9 | * | * |
| 2001..... | 2.9 | * | * | * | * | 3.0 | 7.7 | 6.3 | 4.6 | * | * | * |
| 2000..... | 3.3 | * | * | * | * | 3.7 | 7.8 | 8.9 | 5.7 | * | * | * |
| 1999..... | 3.3 | * | * | * | * | 4.0 | 8.9 | 6.7 | 5.7 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 1.5 | 2.6 | 2.4 | * | * | * | * |
| 2003..... | 0.9 | * | * | * | * | 1.1 | 2.5 | 2.2 | * | * | * | * |
| 2002..... | 1.0 | * | * | * | * | 1.8 | 2.5 | 1.9 | * | * | * | * |
| 2001..... | 1.2 | * | * | * | * | 1.6 | 2.7 | 2.1 | 1.6 | * | * | * |
| 2000..... | 1.2 | * | * | * | * | 2.2 | 3.1 | 2.3 | * | * | * | * |
| 1999..... | 1.2 | * | * | * | * | 1.6 | 3.3 | 1.8 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in other conditions (B23) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.1 | 0.5 | 1.7 | 1.8 | 1.0 | 0.5 | 0.2 | * |
| 2003..... | 1.1 | * | * | * | 0.1 | 0.9 | 2.7 | 2.7 | 1.4 | 0.7 | 0.2 | * |
| 2002..... | 1.2 | * | * | * | 0.1 | 1.0 | 3.0 | 2.8 | 1.4 | 0.6 | 0.3 | * |
| 2001..... | 1.2 | * | * | * | 0.2 | 1.2 | 3.1 | 2.6 | 1.4 | 0.5 | 0.2 | * |
| 2000..... | 1.2 | * | * | * | 0.1 | 1.3 | 2.9 | 2.6 | 1.5 | 0.6 | 0.2 | * |
| 1999..... | 1.2 | * | * | 0.1 | 0.1 | 1.4 | 3.0 | 2.6 | 1.4 | 0.7 | * | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.7 | 2.4 | 2.7 | 1.7 | 0.8 | * | * |
| 2003..... | 1.6 | * | * | * | * | 1.0 | 3.8 | 4.1 | 2.2 | 1.1 | 0.4 | * |
| 2002..... | 1.7 | * | * | * | * | 1.3 | 4.2 | 4.2 | 2.3 | 1.0 | 0.4 | * |
| 2001..... | 1.8 | * | * | * | 0.1 | 1.5 | 4.4 | 4.1 | 2.3 | 0.9 | 0.4 | * |
| 2000..... | 1.8 | * | * | * | * | 1.6 | 4.2 | 4.0 | 2.4 | 0.9 | 0.5 | * |
| 1999..... | 1.8 | * | * | * | 0.1 | 1.9 | 4.3 | 4.1 | 2.3 | 1.2 | * | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.4 | 1.1 | 0.8 | 0.4 | * | * | * |
| 2003..... | 0.6 | * | * | * | 0.1 | 0.7 | 1.7 | 1.4 | 0.5 | 0.3 | * | * |
| 2002..... | 0.7 | * | * | * | * | 0.7 | 1.8 | 1.4 | 0.5 | 0.3 | * | * |
| 2001..... | 0.7 | * | * | * | 0.2 | 0.9 | 1.9 | 1.3 | 0.5 | * | * | * |
| 2000..... | 0.7 | * | * | * | 0.1 | 1.1 | 1.6 | 1.2 | 0.6 | 0.3 | * | * |
| 1999..... | 0.7 | * | * | * | 0.1 | 0.9 | 1.8 | 1.1 | 0.5 | 0.2 | * | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | 0.4 | 1.3 | 2.0 | 1.1 | * | * | * |
| 2003..... | 1.0 | * | * | * | * | 0.5 | 2.6 | 3.7 | 2.3 | * | * | * |
| 2002..... | 1.1 | * | * | * | * | 0.6 | 3.3 | 3.6 | 2.5 | * | * | * |
| 2001..... | 1.2 | * | * | * | * | 0.9 | 3.2 | 3.9 | 2.3 | * | * | * |
| 2000..... | 1.2 | * | * | * | * | 1.0 | 2.9 | 3.9 | 2.9 | * | * | * |
| 1999..... | 1.2 | * | * | * | * | 1.2 | 3.1 | 3.1 | 2.7 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | 0.6 | 1.9 | 3.0 | 1.9 | * | * | * |
| 2003..... | 1.5 | * | * | * | * | 0.7 | 3.4 | 5.8 | 3.6 | * | * | * |
| 2002..... | 1.6 | * | * | * | * | 0.7 | 4.8 | 5.6 | 3.6 | * | * | * |
| 2001..... | 1.7 | * | * | * | * | 1.0 | 4.6 | 5.8 | 3.9 | * | * | * |
| 2000..... | 1.7 | * | * | * | * | 1.3 | 4.3 | 6.3 | 4.5 | * | * | * |
| 1999..... | 1.7 | * | * | * | * | 1.8 | 4.3 | 4.4 | 4.9 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.7 | * | * | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | 1.8 | 1.5 | * | * | * | * |
| 2002..... | 0.6 | * | * | * | * | * | 1.7 | 1.6 | * | * | * | * |
| 2001..... | 0.7 | * | * | * | * | 0.7 | 1.8 | 1.9 | * | * | * | * |
| 2000..... | 0.6 | * | * | * | * | * | 1.5 | 1.5 | * | * | * | * |
| 1999..... | 0.6 | * | * | * | * | * | 1.8 | 1.9 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in other conditions (B23) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.1 | 0.6 | 1.8 | 1.7 | 1.0 | 0.4 | 0.2 | * |
| 2003..... | 1.1 | * | * | * | 0.1 | 0.9 | 2.7 | 2.6 | 1.3 | 0.6 | 0.2 | * |
| 2002..... | 1.2 | * | * | * | 0.1 | 1.1 | 2.9 | 2.6 | 1.3 | 0.6 | 0.2 | * |
| 2001..... | 1.2 | * | * | * | 0.2 | 1.3 | 3.1 | 2.5 | 1.2 | 0.4 | 0.2 | * |
| 2000..... | 1.2 | * | * | * | 0.1 | 1.4 | 2.9 | 2.4 | 1.3 | 0.5 | 0.2 | * |
| 1999..... | 1.2 | * | * | * | 0.1 | 1.4 | 3.0 | 2.4 | 1.3 | 0.6 | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.7 | 2.5 | 2.7 | 1.7 | 0.8 | * | * |
| 2003..... | 1.6 | * | * | * | * | 1.0 | 3.9 | 3.8 | 2.1 | 0.9 | 0.4 | * |
| 2002..... | 1.7 | * | * | * | * | 1.4 | 4.0 | 4.0 | 2.2 | 1.0 | * | * |
| 2001..... | 1.8 | * | * | * | * | 1.6 | 4.3 | 3.8 | 2.0 | 0.8 | 0.4 | * |
| 2000..... | 1.7 | * | * | * | * | 1.6 | 4.1 | 3.7 | 2.1 | 0.9 | * | * |
| 1999..... | 1.8 | * | * | * | * | 1.9 | 4.2 | 3.9 | 2.1 | 1.0 | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.4 | 1.1 | 0.8 | 0.4 | * | * | * |
| 2003..... | 0.7 | * | * | * | 0.1 | 0.8 | 1.7 | 1.4 | 0.5 | 0.3 | * | * |
| 2002..... | 0.7 | * | * | * | * | 0.8 | 1.8 | 1.4 | 0.5 | 0.3 | * | * |
| 2001..... | 0.7 | * | * | * | 0.2 | 0.9 | 1.9 | 1.2 | 0.5 | * | * | * |
| 2000..... | 0.7 | * | * | * | * | 1.2 | 1.6 | 1.2 | 0.6 | 0.3 | * | * |
| 1999..... | 0.7 | * | * | * | 0.2 | 1.0 | 1.8 | 1.0 | 0.5 | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | 0.1 | 0.7 | 0.6 | 0.4 | 0.2 | * | * |
| 2003..... | 0.4 | * | * | * | * | 0.2 | 1.0 | 0.9 | 0.5 | 0.2 | * | * |
| 2002..... | 0.4 | * | * | * | * | 0.3 | 1.1 | 0.9 | 0.4 | 0.2 | * | * |
| 2001..... | 0.4 | * | * | * | * | 0.4 | 1.2 | 0.8 | 0.5 | * | * | * |
| 2000..... | 0.5 | * | * | * | * | 0.4 | 1.2 | 0.9 | 0.5 | 0.2 | * | * |
| 1999..... | 0.5 | * | * | * | * | 0.5 | 1.2 | 0.8 | 0.4 | 0.2 | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | 0.2 | 1.1 | 1.0 | 0.7 | 0.4 | * | * |
| 2003..... | 0.7 | * | * | * | * | 0.3 | 1.7 | 1.5 | 0.9 | 0.3 | * | * |
| 2002..... | 0.7 | * | * | * | * | 0.5 | 1.8 | 1.4 | 0.9 | 0.3 | * | * |
| 2001..... | 0.7 | * | * | * | * | 0.6 | 1.9 | 1.4 | 0.9 | * | * | * |
| 2000..... | 0.7 | * | * | * | * | 0.6 | 2.0 | 1.4 | 0.9 | 0.3 | * | * |
| 1999..... | 0.8 | * | * | * | * | 0.7 | 2.0 | 1.5 | 0.8 | 0.3 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | 0.3 | 0.2 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.3 | 0.3 | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.4 | 0.3 | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | 0.2 | 0.5 | 0.2 | * | * | * | * |
| 2000..... | 0.2 | * | * | * | * | 0.3 | 0.5 | 0.3 | * | * | * | * |
| 1999..... | 0.2 | * | * | * | * | 0.2 | 0.5 | 0.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Human immunodeficiency virus (HIV) disease resulting in other conditions (B23) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | 0.3 | 3.0 | 8.5 | 9.8 | 6.5 | 2.6 | * | * |
| 2003..... | 5.5 | * | * | * | 0.6 | 4.6 | 13.2 | 14.4 | 8.1 | 4.5 | * | * |
| 2002..... | 5.9 | * | * | * | 0.5 | 5.3 | 14.1 | 15.5 | 8.6 | 4.7 | * | * |
| 2001..... | 5.9 | * | * | * | 0.7 | 6.1 | 14.6 | 14.6 | 7.8 | 3.4 | 2.2 | * |
| 2000..... | 5.6 | * | * | * | 0.5 | 6.7 | 13.3 | 14.2 | 8.3 | 3.9 | * | * |
| 1999..... | 5.9 | * | * | * | 0.6 | 6.9 | 14.6 | 14.6 | 8.3 | 4.4 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 5.2 | * | * | * | * | 3.7 | 11.3 | 15.5 | 10.6 | 4.8 | * | * |
| 2003..... | 7.4 | * | * | * | * | 5.1 | 17.9 | 21.4 | 13.6 | 7.4 | * | * |
| 2002..... | 8.2 | * | * | * | * | 6.7 | 19.0 | 23.7 | 14.5 | 8.6 | * | * |
| 2001..... | 8.2 | * | * | * | * | 7.7 | 19.7 | 22.7 | 13.4 | 6.2 | * | * |
| 2000..... | 7.9 | * | * | * | * | 7.8 | 18.7 | 22.3 | 13.7 | 6.9 | * | * |
| 1999..... | 8.5 | * | * | * | * | 8.9 | 20.1 | 23.6 | 13.9 | 8.4 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 2.4 | 5.9 | 4.9 | 3.2 | * | * | * |
| 2003..... | 3.7 | * | * | * | 0.7 | 4.2 | 9.0 | 8.4 | 3.6 | 2.5 | * | * |
| 2002..... | 3.8 | * | * | * | * | 4.1 | 9.7 | 8.5 | 3.9 | * | * | * |
| 2001..... | 3.7 | * | * | * | 1.0 | 4.6 | 10.1 | 7.7 | 3.4 | * | * | * |
| 2000..... | 3.6 | * | * | * | * | 5.7 | 8.5 | 7.3 | 3.9 | * | * | * |
| 1999..... | 3.6 | * | * | * | 0.7 | 5.1 | 9.7 | 6.9 | 3.8 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified human immunodeficiency virus (HIV) disease (B24) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.1 | 1.0 | 3.1 | 2.8 | 1.4 | 0.5 | 0.2 | * |
| 2003..... | 1.0 | * | * | * | 0.1 | 1.0 | 2.8 | 2.2 | 1.1 | 0.5 | 0.2 | * |
| 2002..... | 1.0 | * | * | * | 0.1 | 1.0 | 2.7 | 2.1 | 0.9 | 0.4 | * | * |
| 2001..... | 1.0 | * | * | * | 0.1 | 1.1 | 2.7 | 2.0 | 0.8 | 0.4 | * | * |
| 2000..... | 1.0 | * | * | * | 0.1 | 1.3 | 2.5 | 2.1 | 0.9 | 0.4 | * | * |
| 1999..... | 1.0 | * | * | * | 0.1 | 1.3 | 2.6 | 1.8 | 0.8 | 0.4 | * | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.1 | 1.2 | 4.3 | 4.2 | 2.3 | 0.9 | * | * |
| 2003..... | 1.5 | * | * | * | * | 1.3 | 4.0 | 3.4 | 1.8 | 0.8 | * | * |
| 2002..... | 1.5 | * | * | * | * | 1.2 | 3.9 | 3.3 | 1.5 | 0.7 | * | * |
| 2001..... | 1.5 | * | * | * | * | 1.5 | 4.0 | 3.4 | 1.4 | 0.7 | * | * |
| 2000..... | 1.5 | * | * | * | 0.1 | 1.6 | 3.8 | 3.3 | 1.6 | 0.7 | * | * |
| 1999..... | 1.5 | * | * | * | * | 1.8 | 4.0 | 3.0 | 1.5 | 0.7 | * | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.1 | 0.9 | 1.9 | 1.4 | 0.5 | 0.2 | * | * |
| 2003..... | 0.5 | * | * | * | 0.1 | 0.7 | 1.6 | 1.0 | 0.4 | 0.2 | * | * |
| 2002..... | 0.5 | * | * | * | 0.1 | 0.8 | 1.4 | 0.9 | 0.3 | * | * | * |
| 2001..... | 0.5 | * | * | * | 0.1 | 0.7 | 1.4 | 0.7 | 0.3 | * | * | * |
| 2000..... | 0.5 | * | * | * | * | 0.9 | 1.2 | 0.8 | 0.2 | * | * | * |
| 1999..... | 0.5 | * | * | * | * | 0.8 | 1.3 | 0.7 | 0.3 | * | * | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.6 | 2.5 | 3.6 | 2.4 | * | * | * |
| 2003..... | 0.8 | * | * | * | * | 0.7 | 2.1 | 2.2 | 1.5 | * | * | * |
| 2002..... | 0.7 | * | * | * | * | 0.6 | 1.8 | 2.4 | 1.3 | * | * | * |
| 2001..... | 0.6 | * | * | * | * | 0.5 | 1.8 | 2.3 | * | * | * | * |
| 2000..... | 0.8 | * | * | * | * | 0.9 | 2.3 | 2.5 | 1.4 | * | * | * |
| 1999..... | 1.0 | * | * | * | * | 1.0 | 3.0 | 2.5 | 1.3 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.8 | 3.6 | 5.7 | 4.0 | * | * | * |
| 2003..... | 1.1 | * | * | * | * | 1.0 | 2.8 | 3.6 | 2.7 | * | * | * |
| 2002..... | 1.0 | * | * | * | * | 0.9 | 2.7 | 3.6 | * | * | * | * |
| 2001..... | 1.0 | * | * | * | * | 0.6 | 2.7 | 3.9 | * | * | * | * |
| 2000..... | 1.2 | * | * | * | * | 1.1 | 3.6 | 3.9 | * | * | * | * |
| 1999..... | 1.5 | * | * | * | * | 1.4 | 4.6 | 4.3 | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 1.3 | 1.6 | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | 1.2 | * | * | * | * | * |
| 2002..... | 0.4 | * | * | * | * | * | 0.9 | 1.1 | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | 0.8 | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | * | 1.0 | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | 1.3 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified human immunodeficiency virus (HIV) disease (B24) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.1 | 1.1 | 3.2 | 2.7 | 1.3 | 0.5 | 0.2 | * |
| 2003..... | 1.0 | * | * | * | 0.1 | 1.0 | 2.8 | 2.1 | 1.0 | 0.4 | 0.2 | * |
| 2002..... | 1.0 | * | * | * | 0.1 | 1.1 | 2.8 | 2.1 | 0.8 | 0.4 | * | * |
| 2001..... | 1.0 | * | * | * | 0.1 | 1.2 | 2.8 | 1.9 | 0.8 | 0.4 | * | * |
| 2000..... | 1.0 | * | * | * | 0.1 | 1.3 | 2.5 | 2.0 | 0.8 | 0.4 | * | * |
| 1999..... | 1.0 | * | * | * | 0.1 | 1.4 | 2.5 | 1.7 | 0.8 | 0.3 | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.1 | 1.3 | 4.4 | 4.0 | 2.1 | 0.8 | * | * |
| 2003..... | 1.6 | * | * | * | * | 1.3 | 4.1 | 3.3 | 1.7 | 0.7 | * | * |
| 2002..... | 1.5 | * | * | * | * | 1.3 | 4.0 | 3.3 | 1.4 | 0.7 | * | * |
| 2001..... | 1.5 | * | * | * | * | 1.6 | 4.1 | 3.2 | 1.4 | 0.6 | * | * |
| 2000..... | 1.5 | * | * | * | 0.1 | 1.7 | 3.7 | 3.2 | 1.5 | 0.7 | * | * |
| 1999..... | 1.5 | * | * | * | * | 1.9 | 3.8 | 2.8 | 1.4 | 0.6 | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.1 | 1.0 | 2.0 | 1.4 | 0.5 | 0.2 | * | * |
| 2003..... | 0.6 | * | * | * | * | 0.7 | 1.6 | 1.0 | 0.3 | * | * | * |
| 2002..... | 0.5 | * | * | * | 0.1 | 0.9 | 1.5 | 0.9 | 0.3 | * | * | * |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.7 | 1.5 | 0.7 | 0.3 | * | * | * |
| 2000..... | 0.5 | * | * | * | * | 1.0 | 1.3 | 0.8 | 0.2 | * | * | * |
| 1999..... | 0.5 | * | * | * | * | 0.9 | 1.3 | 0.7 | 0.3 | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | 0.3 | 1.4 | 1.0 | 0.6 | 0.2 | * | * |
| 2003..... | 0.5 | * | * | * | * | 0.4 | 1.2 | 0.9 | 0.5 | * | * | * |
| 2002..... | 0.4 | * | * | * | * | 0.4 | 1.2 | 0.9 | 0.4 | 0.1 | * | * |
| 2001..... | 0.4 | * | * | * | * | 0.4 | 1.3 | 0.8 | 0.3 | 0.2 | * | * |
| 2000..... | 0.4 | * | * | * | * | 0.5 | 1.1 | 0.8 | 0.4 | 0.1 | * | * |
| 1999..... | 0.4 | * | * | * | * | 0.7 | 1.2 | 0.8 | 0.3 | 0.1 | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | 0.5 | 2.3 | 1.7 | 1.0 | 0.3 | * | * |
| 2003..... | 0.8 | * | * | * | * | 0.6 | 2.0 | 1.6 | 0.9 | * | * | * |
| 2002..... | 0.8 | * | * | * | * | 0.6 | 2.1 | 1.6 | 0.7 | * | * | * |
| 2001..... | 0.8 | * | * | * | * | 0.7 | 2.3 | 1.5 | 0.6 | 0.4 | * | * |
| 2000..... | 0.7 | * | * | * | * | 0.8 | 1.9 | 1.5 | 0.8 | 0.3 | * | * |
| 1999..... | 0.8 | * | * | * | * | 1.0 | 2.1 | 1.4 | 0.6 | 0.3 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.2 | 0.6 | 0.3 | 0.2 | * | * | * |
| 2003..... | 0.1 | * | * | * | * | 0.2 | 0.4 | 0.3 | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | 0.2 | 0.3 | 0.2 | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.3 | 0.2 | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | 0.2 | 0.4 | 0.2 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | 0.3 | 0.4 | 0.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified human immunodeficiency virus (HIV) disease (B24) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 0.6 | 5.4 | 13.8 | 14.6 | 7.5 | 3.3 | * | * |
| 2003..... | 4.6 | * | * | * | 0.4 | 4.4 | 13.0 | 10.7 | 5.7 | 3.1 | * | * |
| 2002..... | 4.4 | * | * | * | 0.5 | 4.9 | 12.3 | 10.2 | 4.4 | 2.9 | * | * |
| 2001..... | 4.5 | * | * | * | 0.6 | 5.2 | 12.2 | 10.4 | 4.8 | 2.2 | * | * |
| 2000..... | 4.5 | * | * | * | 0.5 | 5.9 | 11.2 | 10.9 | 4.5 | 2.5 | * | * |
| 1999..... | 4.2 | * | * | * | 0.4 | 5.7 | 11.1 | 9.2 | 4.9 | 2.0 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 7.5 | * | * | * | * | 5.9 | 18.1 | 21.5 | 13.2 | 5.6 | * | * |
| 2003..... | 6.5 | * | * | * | * | 5.2 | 17.9 | 16.6 | 9.9 | 5.9 | * | * |
| 2002..... | 6.2 | * | * | * | * | 5.5 | 16.9 | 16.1 | 7.7 | 5.3 | * | * |
| 2001..... | 6.4 | * | * | * | * | 7.0 | 16.5 | 17.1 | 8.5 | 3.7 | * | * |
| 2000..... | 6.5 | * | * | * | * | 6.9 | 16.5 | 17.4 | 8.8 | 4.5 | * | * |
| 1999..... | 6.0 | * | * | * | * | 7.4 | 15.9 | 15.1 | 8.5 | 3.3 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | 4.9 | 10.0 | 8.7 | 3.0 | * | * | * |
| 2003..... | 2.9 | * | * | * | * | 3.6 | 8.6 | 5.7 | 2.4 | * | * | * |
| 2002..... | 2.9 | * | * | * | * | 4.4 | 8.2 | 5.2 | 1.7 | * | * | * |
| 2001..... | 2.8 | * | * | * | 0.8 | 3.6 | 8.4 | 4.6 | 1.8 | * | * | * |
| 2000..... | 2.6 | * | * | * | * | 5.0 | 6.5 | 5.3 | * | * | * | * |
| 1999..... | 2.5 | * | * | * | * | 4.1 | 6.8 | 4.1 | 2.0 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.5 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 0.8 | 1.2 | 2.0 |
| 2003..... | 0.4 | 2.3 | 0.4 | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.5 | 1.0 | 1.3 | 2.8 |
| 2002..... | 0.4 | 3.2 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 1.0 | 1.4 | 3.0 |
| 2001..... | 0.3 | 1.9 | 0.4 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 1.0 | 2.2 |
| 2000..... | 0.4 | 1.9 | 0.3 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 1.4 | 3.4 |
| 1999..... | 0.4 | 2.7 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 1.7 | 4.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.8 | * | * | * | * | 0.1 | 0.2 | 0.6 | 0.9 | 1.5 | 2.4 |
| 2003..... | 0.4 | 1.9 | 0.3 | * | 0.1 | * | 0.2 | 0.2 | 0.6 | 1.1 | 1.4 | 2.9 |
| 2002..... | 0.4 | 3.3 | 0.4 | * | 0.1 | * | 0.2 | 0.3 | 0.4 | 1.0 | 1.6 | 2.9 |
| 2001..... | 0.3 | 1.8 | 0.4 | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 1.2 | 2.4 |
| 2000..... | 0.4 | 2.1 | * | * | 0.1 | * | 0.1 | 0.2 | 0.6 | 1.2 | 1.8 | 3.1 |
| 1999..... | 0.4 | 2.8 | 0.4 | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.6 | 3.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.1 | * | * | 0.1 | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.0 | 1.8 |
| 2003..... | 0.4 | 2.8 | 0.4 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.2 | 2.8 |
| 2002..... | 0.4 | 3.0 | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.2 | 3.0 |
| 2001..... | 0.3 | 2.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 0.9 | 2.1 |
| 2000..... | 0.4 | 1.8 | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.2 | 3.4 |
| 1999..... | 0.5 | 2.6 | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.8 | 1.8 | 4.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.2 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 0.8 | 1.3 | 2.1 |
| 2003..... | 0.4 | 1.8 | 0.3 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.6 | 1.0 | 1.4 | 3.0 |
| 2002..... | 0.4 | 2.9 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 1.0 | 1.4 | 3.2 |
| 2001..... | 0.3 | 1.2 | 0.3 | 0.1 | 0.1 | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.1 | 2.4 |
| 2000..... | 0.4 | 1.5 | 0.2 | 0.1 | 0.1 | * | 0.1 | 0.2 | 0.5 | 1.1 | 1.4 | 3.5 |
| 1999..... | 0.4 | 2.3 | 0.3 | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.0 | 1.8 | 4.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | 2.5 | * | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.7 | 2.6 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.2 | 1.5 | 3.1 |
| 2002..... | 0.4 | 2.6 | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 1.7 | 3.1 |
| 2001..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 1.3 | 2.6 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 1.8 | 3.4 |
| 1999..... | 0.4 | 2.6 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.7 | 4.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.0 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 1.8 |
| 2003..... | 0.4 | 2.2 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.3 | 2.9 |
| 2002..... | 0.4 | 3.2 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 1.2 | 3.3 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 0.9 | 2.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.2 | 3.6 |
| 1999..... | 0.5 | 2.0 | * | * | * | * | * | 0.3 | 0.7 | 0.9 | 1.9 | 4.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | 1.0 | * | * | 0.1 | 0.2 | 0.7 | 2.4 | 2.8 | 4.4 | 7.1 | 9.7 |
| 2003..... | 1.5 | 1.1 | * | * | 0.1 | 0.2 | 0.7 | 2.3 | 2.7 | 4.2 | 7.1 | 9.0 |
| 2002..... | 1.3 | 1.3 | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.7 | 2.2 | 3.7 | 6.4 | 8.3 |
| 2001..... | 1.1 | 1.6 | * | 0.0 | 0.1 | 0.2 | 0.5 | 1.2 | 1.9 | 3.6 | 5.4 | 8.2 |
| 2000..... | 1.0 | 1.3 | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 1.7 | 3.3 | 5.7 | 9.0 |
| 1999..... | 1.4 | 1.6 | 0.1 | 0.1 | 0.1 | 0.4 | 0.8 | 1.6 | 2.4 | 4.7 | 7.4 | 10.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.8 | 1.3 | * | * | 0.1 | 0.2 | 0.9 | 3.5 | 3.8 | 5.1 | 8.4 | 12.9 |
| 2003..... | 1.7 | 1.3 | * | * | * | 0.3 | 0.9 | 3.2 | 3.5 | 4.7 | 8.3 | 10.4 |
| 2002..... | 1.5 | 1.5 | * | * | * | 0.3 | 0.9 | 2.5 | 2.9 | 4.7 | 7.9 | 11.9 |
| 2001..... | 1.2 | 1.6 | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 2.3 | 3.8 | 7.1 | 10.1 |
| 2000..... | 1.1 | 1.5 | * | * | * | 0.2 | 0.7 | 1.3 | 2.0 | 4.3 | 6.9 | 12.6 |
| 1999..... | 1.6 | 1.9 | * | * | 0.1 | 0.4 | 1.0 | 1.9 | 2.7 | 5.4 | 9.4 | 13.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.0 | 3.8 | 6.2 | 8.3 |
| 2003..... | 1.3 | * | * | * | * | 0.2 | 0.5 | 1.4 | 1.9 | 3.9 | 6.3 | 8.3 |
| 2002..... | 1.0 | 1.2 | * | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 2.9 | 5.5 | 6.8 |
| 2001..... | 1.0 | 1.6 | * | * | * | 0.2 | 0.4 | 0.8 | 1.5 | 3.5 | 4.4 | 7.4 |
| 2000..... | 0.9 | 1.1 | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 2.5 | 4.9 | 7.6 |
| 1999..... | 1.3 | 1.2 | * | * | * | 0.4 | 0.6 | 1.3 | 2.1 | 4.0 | 6.2 | 8.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 1.1 | 3.9 | 4.9 | 5.4 | 9.3 | 11.9 |
| 2003..... | 1.2 | * | * | * | * | * | 1.2 | 3.8 | 4.6 | 6.1 | 9.8 | * |
| 2002..... | 1.0 | * | * | * | * | 0.5 | 0.9 | 2.9 | 3.6 | 4.5 | 7.8 | * |
| 2001..... | 0.7 | * | * | * | * | 0.3 | 0.7 | 1.6 | 1.8 | 4.4 | 6.3 | * |
| 2000..... | 0.7 | * | * | * | * | * | 0.8 | 1.3 | 2.0 | 3.6 | 6.7 | 13.3 |
| 1999..... | 0.9 | * | * | * | * | 0.4 | 1.1 | 1.5 | 2.3 | 4.8 | 9.8 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 1.4 | 5.6 | 7.0 | 5.7 | 12.2 | * |
| 2003..... | 1.5 | * | * | * | * | * | 1.7 | 5.9 | 6.4 | 6.8 | 9.6 | * |
| 2002..... | 1.3 | * | * | * | * | 0.5 | 1.3 | 4.6 | 4.9 | 5.7 | 9.0 | * |
| 2001..... | 0.9 | * | * | * | * | * | 1.0 | 2.1 | 2.9 | 5.2 | * | * |
| 2000..... | 0.9 | * | * | * | * | * | 1.2 | 1.4 | 3.2 | 4.6 | * | * |
| 1999..... | 0.9 | * | * | * | * | * | 1.5 | 1.6 | * | 5.5 | 12.0 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.7 | 2.2 | 2.9 | 5.3 | 7.4 | * |
| 2003..... | 0.9 | * | * | * | * | * | * | 1.8 | 3.0 | 5.6 | 10.0 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.2 | 2.4 | 3.5 | 7.0 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.8 | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.8 | * | * | * | * | * | * | 1.3 | 2.4 | 4.3 | 8.3 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | 1.0 | * | * | 0.1 | 0.2 | 0.7 | 2.2 | 2.7 | 4.3 | 6.9 | 9.6 |
| 2003..... | 1.5 | 1.2 | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 2.5 | 4.1 | 6.9 | 9.1 |
| 2002..... | 1.3 | 1.4 | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.6 | 2.1 | 3.7 | 6.3 | 8.3 |
| 2001..... | 1.1 | 1.6 | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 1.9 | 3.6 | 5.4 | 8.3 |
| 2000..... | 1.1 | 1.4 | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.7 | 3.3 | 5.6 | 8.9 |
| 1999..... | 1.5 | 1.5 | 0.2 | 0.1 | 0.1 | 0.4 | 0.8 | 1.6 | 2.4 | 4.6 | 7.3 | 9.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.8 | 1.4 | * | * | * | 0.2 | 0.8 | 3.2 | 3.5 | 5.0 | 8.1 | 13.0 |
| 2003..... | 1.7 | 1.5 | * | * | * | 0.3 | 0.8 | 3.0 | 3.2 | 4.5 | 8.2 | 10.5 |
| 2002..... | 1.5 | 1.6 | * | * | * | 0.2 | 0.8 | 2.3 | 2.8 | 4.6 | 7.8 | 11.7 |
| 2001..... | 1.2 | 1.8 | * | * | * | 0.2 | 0.7 | 1.5 | 2.2 | 3.7 | 7.0 | 10.3 |
| 2000..... | 1.2 | 1.5 | * | * | * | 0.2 | 0.6 | 1.2 | 1.9 | 4.2 | 6.7 | 12.3 |
| 1999..... | 1.6 | 1.7 | * | * | * | 0.4 | 1.0 | 1.9 | 2.7 | 5.4 | 9.3 | 13.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.2 | 0.5 | 1.3 | 1.9 | 3.7 | 6.1 | 8.1 |
| 2003..... | 1.3 | * | * | * | * | 0.2 | 0.5 | 1.4 | 1.8 | 3.7 | 6.1 | 8.5 |
| 2002..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.5 | 2.9 | 5.4 | 6.8 |
| 2001..... | 1.0 | 1.5 | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 3.4 | 4.3 | 7.5 |
| 2000..... | 1.0 | 1.4 | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 2.5 | 4.9 | 7.5 |
| 1999..... | 1.4 | * | * | * | * | 0.4 | 0.6 | 1.3 | 2.1 | 4.0 | 6.0 | 8.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.1 | 0.6 | 2.2 | 2.4 | 4.0 | 6.6 | 9.4 |
| 2003..... | 1.5 | * | * | * | * | 0.1 | 0.6 | 2.1 | 2.2 | 4.0 | 6.5 | 9.1 |
| 2002..... | 1.3 | 1.1 | * | * | * | 0.1 | 0.5 | 1.4 | 1.8 | 3.6 | 6.2 | 8.2 |
| 2001..... | 1.1 | 1.1 | * | * | * | 0.1 | 0.4 | 1.0 | 1.6 | 3.4 | 5.3 | 8.0 |
| 2000..... | 1.1 | 1.0 | * | * | * | 0.1 | 0.4 | 0.8 | 1.5 | 3.1 | 5.5 | 8.7 |
| 1999..... | 1.4 | 1.1 | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.0 | 4.3 | 7.0 | 9.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.7 | 3.2 | 3.1 | 4.8 | 7.8 | 13.3 |
| 2003..... | 1.7 | * | * | * | * | 0.2 | 0.8 | 2.9 | 2.9 | 4.3 | 7.7 | 10.6 |
| 2002..... | 1.5 | * | * | * | * | * | 0.7 | 2.1 | 2.4 | 4.5 | 7.6 | 11.7 |
| 2001..... | 1.2 | * | * | * | * | * | 0.5 | 1.3 | 2.0 | 3.6 | 6.9 | 10.1 |
| 2000..... | 1.1 | * | * | * | * | * | 0.5 | 1.1 | 1.7 | 4.0 | 6.3 | 11.7 |
| 1999..... | 1.5 | * | * | * | * | 0.3 | 0.8 | 1.6 | 2.3 | 5.1 | 8.8 | 12.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.5 | 1.3 | 1.8 | 3.4 | 5.8 | 7.7 |
| 2003..... | 1.3 | * | * | * | * | * | 0.4 | 1.3 | 1.6 | 3.7 | 5.6 | 8.4 |
| 2002..... | 1.1 | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.8 | 5.3 | 6.7 |
| 2001..... | 1.0 | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 3.2 | 4.2 | 7.1 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 4.9 | 7.4 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.3 | 0.9 | 1.6 | 3.7 | 5.8 | 8.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.5 | 1.2 | 2.4 | 4.4 | 5.5 | 9.4 | 13.5 |
| 2003..... | 1.6 | * | * | * | * | 0.5 | 1.1 | 3.2 | 4.2 | 4.1 | 9.8 | 9.2 |
| 2002..... | 1.5 | * | * | * | * | 0.7 | 1.3 | 3.0 | 4.2 | 3.8 | 6.5 | 9.4 |
| 2001..... | 1.3 | 3.9 | * | * | * | 0.6 | 1.1 | 2.1 | 3.4 | 5.0 | 5.8 | 11.5 |
| 2000..... | 1.3 | 3.5 | * | * | * | 0.6 | 1.3 | 2.0 | 3.0 | 4.6 | 7.0 | 11.1 |
| 1999..... | 2.3 | 3.9 | * | * | * | 1.3 | 2.5 | 4.2 | 5.1 | 7.6 | 10.2 | 14.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 1.5 | 3.8 | 6.7 | 5.6 | 12.0 | * |
| 2003..... | 1.8 | * | * | * | * | * | 1.4 | 4.2 | 5.7 | 5.3 | 11.3 | * |
| 2002..... | 1.8 | * | * | * | * | 0.9 | 1.7 | 4.3 | 6.5 | 5.0 | 7.0 | * |
| 2001..... | 1.5 | * | * | * | * | * | 1.6 | 3.0 | 4.2 | 5.8 | 6.9 | * |
| 2000..... | 1.5 | * | * | * | * | * | 1.7 | 2.7 | 3.9 | 6.4 | 9.9 | * |
| 1999..... | 2.3 | * | * | * | * | 1.5 | 2.5 | 4.7 | 5.1 | 8.7 | 13.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.9 | 1.1 | 2.5 | 5.4 | 7.9 | 13.3 |
| 2003..... | 1.4 | * | * | * | * | * | 1.0 | 2.4 | 3.1 | 3.2 | 9.0 | 9.1 |
| 2002..... | 1.2 | * | * | * | * | * | 0.9 | 1.9 | 2.4 | 3.0 | 6.1 | * |
| 2001..... | 1.2 | * | * | * | * | * | 0.7 | 1.3 | 2.8 | 4.5 | 5.2 | 12.0 |
| 2000..... | 1.1 | * | * | * | * | * | 0.9 | 1.5 | 2.4 | 3.4 | 5.4 | 9.2 |
| 1999..... | 2.3 | * | * | * | * | 1.2 | 2.5 | 3.8 | 5.1 | 6.9 | 8.2 | 11.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mycoses (B35-B49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | 0.9 | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 2.0 | 2.5 |
| 2003..... | 0.4 | 1.1 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.3 | 2.1 | 2.5 |
| 2002..... | 0.5 | 1.2 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 2.4 | 2.5 |
| 2001..... | 0.4 | 1.5 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 2.2 | 3.0 |
| 2000..... | 0.5 | 1.4 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.5 | 2.2 | 3.2 |
| 1999..... | 0.7 | 1.3 | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.2 | 2.3 | 3.2 | 3.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.3 | 2.7 | 3.8 |
| 2003..... | 0.4 | 1.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.3 | 2.8 | 3.5 |
| 2002..... | 0.6 | 1.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 2.0 | 3.5 | 4.1 |
| 2001..... | 0.5 | 1.6 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.4 | 3.0 | 3.9 |
| 2000..... | 0.5 | 1.5 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 2.9 | 4.9 |
| 1999..... | 0.8 | 1.5 | * | * | * | 0.2 | 0.4 | 0.7 | 1.4 | 2.8 | 4.5 | 6.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.5 | 1.9 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 1.6 | 2.0 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.7 | 1.8 |
| 2001..... | 0.4 | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 1.6 | 2.6 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.1 | 1.7 | 2.4 |
| 1999..... | 0.6 | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.0 | 1.8 | 2.3 | 2.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.8 | 2.3 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 1.9 | 2.3 |
| 2002..... | 0.4 | 0.9 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 2.3 | 2.4 |
| 2001..... | 0.4 | 1.0 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.0 | 2.6 |
| 2000..... | 0.4 | 1.0 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.1 | 2.9 |
| 1999..... | 0.6 | 0.9 | * | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 2.1 | 3.0 | 3.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.4 | 3.9 |
| 2003..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 2.7 | 3.6 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.9 | 3.4 | 4.1 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.1 | 2.8 | 3.5 |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.8 | 2.7 | 4.4 |
| 1999..... | 0.7 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.7 | 4.3 | 5.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.3 | 1.7 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 1.4 | 1.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 1.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 1.5 | 2.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 1.8 | 2.2 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.5 | 0.8 | 1.6 | 2.1 | 2.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mycoses (B35-B49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.4 | 0.6 | 1.5 | 1.7 | 4.1 | * |
| 2003..... | 0.6 | * | * | * | * | * | 0.4 | 0.7 | 1.6 | 1.6 | 3.5 | * |
| 2002..... | 0.7 | * | * | * | * | 0.4 | 0.5 | 1.0 | 1.6 | 2.2 | 3.0 | * |
| 2001..... | 0.7 | 3.8 | * | * | * | * | 0.4 | 0.7 | 1.6 | 2.8 | 3.6 | 7.2 |
| 2000..... | 0.7 | * | * | * | * | * | 0.6 | 0.8 | 1.5 | 2.6 | 3.0 | 6.7 |
| 1999..... | 1.0 | 3.5 | * | * | * | 0.6 | 0.8 | 1.4 | 2.2 | 4.2 | 5.2 | 9.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.9 | 2.2 | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 2.1 | * | * | * |
| 2002..... | 0.9 | * | * | * | * | * | * | 1.7 | 2.5 | 3.5 | * | * |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 2.0 | 3.9 | * | * |
| 2000..... | 0.7 | * | * | * | * | * | * | 1.1 | * | 3.3 | * | * |
| 1999..... | 1.1 | * | * | * | * | * | 1.0 | 1.8 | 2.2 | 4.8 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 3.5 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 2.1 | * | * |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.1 | 2.2 | 3.7 | 4.7 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Leishmaniasis (B55)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Leishmaniasis (B55) in Non-Hispanic 4/, both sexes.

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Leishmaniasis (B55) in Non-Hispanic 4/, male.

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Leishmaniasis (B55) in Non-Hispanic 4/, female.

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Leishmaniasis (B55) in Non-Hispanic white, both sexes.

Non-Hispanic white, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Leishmaniasis (B55) in Non-Hispanic white, male.

Non-Hispanic white, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Leishmaniasis (B55) in Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Sequelae of tuberculosis (B90) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.2 | 0.4 |
| 2003..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.2 | 0.4 |
| 2002..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.3 | 0.9 |
| 2001..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.3 | 0.7 |
| 2000..... | 0.0 | ... | * | * | * | * | * | * | * | 0.1 | 0.4 | 0.5 |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | 0.2 | 0.6 | 0.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.4 | * |
| 2000..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | * | 0.6 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.3 | * |
| 2002..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | 0.9 |
| 2001..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.3 | * |
| 2000..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.4 | * |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | * | 0.6 | 0.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.2 | 0.9 |
| 2001..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.3 | 0.6 |
| 2000..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.4 | * |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | * | 0.6 | 0.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2003..... | * | ... | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.5 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | 1.0 |
| 2001..... | 0.0 | ... | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | ... | * | * | * | * | * | * | * | * | 0.4 | * |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | * | 0.6 | 0.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Sequelae of poliomyelitis (B91) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | * | * | * | * | * | * | 0.1 | 0.5 | 0.9 | 1.4 |
| 2003..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.2 |
| 2002..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.3 |
| 2001..... | 0.1 | ... | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 0.9 | 1.3 |
| 2000..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.2 |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | * | * | * | * | * | * | * | 0.5 | 1.1 | 2.0 |
| 2003..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.5 |
| 2002..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.0 |
| 2001..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 1.8 |
| 2000..... | 0.1 | ... | * | * | * | * | * | * | * | 0.5 | 0.8 | * |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | 0.5 | 0.7 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | * | * | * | * | * | * | * | 0.4 | 0.8 | 1.2 |
| 2003..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 |
| 2002..... | 0.1 | ... | * | * | * | * | * | * | * | 0.5 | 0.9 | 1.1 |
| 2001..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.0 |
| 2000..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.1 |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | 0.4 | 0.6 | 1.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | * | * | * | * | * | * | 0.1 | 0.6 | 1.0 | 1.6 |
| 2003..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.3 |
| 2002..... | 0.2 | ... | * | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.5 |
| 2001..... | 0.2 | ... | * | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.4 |
| 2000..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.4 |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | * | * | * | * | * | * | * | 0.6 | 1.3 | 2.2 |
| 2003..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.6 |
| 2002..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.1 |
| 2001..... | 0.2 | ... | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.0 |
| 2000..... | 0.1 | ... | * | * | * | * | * | * | * | 0.6 | 0.9 | * |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | 0.5 | 0.8 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | * | * | * | * | * | * | * | 0.5 | 0.8 | 1.3 |
| 2003..... | 0.1 | ... | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.2 |
| 2002..... | 0.2 | ... | * | * | * | * | * | * | * | 0.5 | 1.0 | 1.2 |
| 2001..... | 0.2 | ... | * | * | * | * | * | * | 0.3 | 0.6 | 0.8 | 1.1 |
| 2000..... | 0.2 | ... | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.3 |
| 1999..... | 0.1 | ... | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.1 | 0.4 | 1.8 | 1.8 | 2.4 | 3.6 | 5.1 |
| 2003..... | 0.9 | * | * | * | * | 0.1 | 0.4 | 1.7 | 1.5 | 2.2 | 3.7 | 4.9 |
| 2002..... | 0.6 | * | * | * | * | * | 0.3 | 1.1 | 1.1 | 1.5 | 2.5 | 3.5 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.6 | 0.7 | 1.3 | 1.8 | 3.2 |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.9 | 3.7 |
| 1999..... | 0.5 | * | * | * | * | 0.1 | 0.4 | 0.7 | 0.8 | 1.5 | 2.6 | 3.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.5 | 2.7 | 2.4 | 2.8 | 3.8 | 6.3 |
| 2003..... | 1.1 | * | * | * | * | * | 0.5 | 2.5 | 2.0 | 2.4 | 4.2 | 4.9 |
| 2002..... | 0.7 | * | * | * | * | * | 0.4 | 1.6 | 1.4 | 1.7 | 2.8 | 4.7 |
| 2001..... | 0.5 | * | * | * | * | * | 0.3 | 0.9 | 0.9 | 1.3 | 2.2 | 3.4 |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.5 | 0.5 | 1.2 | 2.2 | 4.4 |
| 1999..... | 0.6 | * | * | * | * | * | 0.4 | 0.9 | 0.9 | 1.6 | 3.2 | 3.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.4 | 1.0 | 1.1 | 2.1 | 3.5 | 4.6 |
| 2003..... | 0.8 | * | * | * | * | * | 0.3 | 1.0 | 1.0 | 1.9 | 3.4 | 4.9 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.3 | 2.3 | 3.0 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 1.5 | 3.1 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.7 | 3.5 |
| 1999..... | 0.5 | * | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.4 | 2.2 | 3.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.4 | 1.9 | 1.7 | 2.2 | 3.5 | 4.9 |
| 2003..... | 0.9 | * | * | * | * | * | 0.4 | 1.7 | 1.4 | 2.1 | 3.4 | 4.9 |
| 2002..... | 0.6 | * | * | * | * | * | 0.3 | 1.0 | 0.9 | 1.4 | 2.5 | 3.4 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.6 | 0.6 | 1.3 | 1.8 | 3.2 |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.4 | 0.9 | 1.9 | 3.7 |
| 1999..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.4 | 2.5 | 3.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.5 | 2.8 | 2.3 | 2.7 | 3.7 | 6.3 |
| 2003..... | 1.1 | * | * | * | * | * | 0.5 | 2.5 | 1.9 | 2.3 | 3.8 | 4.8 |
| 2002..... | 0.7 | * | * | * | * | * | 0.4 | 1.6 | 1.2 | 1.7 | 2.8 | 4.6 |
| 2001..... | 0.5 | * | * | * | * | * | 0.3 | 0.8 | 0.8 | 1.3 | 2.1 | 3.5 |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.5 | 0.4 | 1.1 | 2.2 | 4.2 |
| 1999..... | 0.5 | * | * | * | * | * | 0.3 | 0.7 | 0.7 | 1.5 | 2.9 | 3.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.4 | 1.1 | 1.1 | 1.9 | 3.3 | 4.3 |
| 2003..... | 0.8 | * | * | * | * | * | 0.3 | 1.0 | 0.9 | 1.9 | 3.1 | 4.9 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.2 | 2.2 | 2.8 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.5 | 1.2 | 1.5 | 3.1 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.6 | 3.5 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 2.2 | 3.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.4 | 1.5 | 2.3 | 3.3 | 4.4 | 8.3 |
| 2003..... | 0.8 | * | * | * | * | * | 0.4 | 2.2 | 2.5 | 2.0 | 5.6 | * |
| 2002..... | 0.6 | * | * | * | * | * | 0.4 | 1.4 | 2.0 | * | 2.9 | * |
| 2001..... | 0.4 | * | * | * | * | * | 0.4 | 0.9 | 1.0 | 1.9 | * | * |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.6 | 1.2 | 1.2 | 2.9 | * |
| 1999..... | 1.0 | * | * | * | * | 0.4 | 1.1 | 2.1 | 2.5 | 2.9 | 4.4 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 2.5 | 3.6 | 3.4 | * | * |
| 2003..... | 1.0 | * | * | * | * | * | * | 3.0 | 3.3 | * | 6.6 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.9 | 3.2 | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | 1.3 | * | * | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 1.0 | * | * | * | * | * | 1.0 | 2.0 | 2.5 | 3.3 | 7.2 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 1.3 | 3.1 | 3.9 | 8.0 |
| 2003..... | 0.7 | * | * | * | * | * | * | 1.4 | 1.8 | * | 5.1 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.9 | * | * | * | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 2.3 | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 1.0 | * | * | * | * | * | 1.2 | 2.2 | 2.4 | 2.5 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| II. Neoplasms (C00-D48) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 193.2 | 3.6 | 2.8 | 2.7 | 4.4 | 9.5 | 34.1 | 120.7 | 337.9 | 767.8 | 1,316.2 | 1,731.0 |
| 2003..... | 196.2 | 3.4 | 2.8 | 2.8 | 4.2 | 9.8 | 35.7 | 123.8 | 347.4 | 783.1 | 1,339.0 | 1,778.8 |
| 2002..... | 197.9 | 3.5 | 3.0 | 2.8 | 4.5 | 10.1 | 36.6 | 125.6 | 355.6 | 805.0 | 1,348.0 | 1,802.6 |
| 2001..... | 199.2 | 2.9 | 3.1 | 2.7 | 4.5 | 10.5 | 37.6 | 128.3 | 361.4 | 816.9 | 1,352.8 | 1,848.3 |
| 2000..... | 201.3 | 4.0 | 3.1 | 2.7 | 4.6 | 10.2 | 37.4 | 129.3 | 371.7 | 830.2 | 1,373.2 | 1,904.3 |
| 1999..... | 201.8 | 3.3 | 3.1 | 2.7 | 4.7 | 10.4 | 37.9 | 129.4 | 379.3 | 841.6 | 1,368.0 | 1,888.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 203.1 | 3.5 | 2.9 | 3.0 | 5.1 | 9.1 | 29.9 | 126.2 | 382.1 | 923.3 | 1,707.7 | 2,448.4 |
| 2003..... | 206.0 | 3.2 | 3.2 | 3.0 | 4.9 | 9.4 | 31.7 | 129.3 | 391.9 | 947.7 | 1,741.5 | 2,515.6 |
| 2002..... | 208.5 | 3.5 | 3.1 | 3.1 | 5.1 | 9.7 | 32.4 | 129.9 | 405.1 | 980.6 | 1,757.2 | 2,592.2 |
| 2001..... | 210.1 | 2.8 | 3.3 | 2.8 | 5.3 | 9.7 | 33.4 | 132.4 | 410.7 | 1,002.0 | 1,775.3 | 2,721.3 |
| 2000..... | 212.0 | 4.2 | 3.4 | 2.9 | 5.4 | 9.6 | 33.6 | 133.0 | 421.5 | 1,018.9 | 1,808.1 | 2,819.6 |
| 1999..... | 213.7 | 3.4 | 3.3 | 2.9 | 5.4 | 10.2 | 34.2 | 133.7 | 434.1 | 1,037.8 | 1,813.1 | 2,831.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 183.7 | 3.7 | 2.7 | 2.4 | 3.6 | 10.0 | 38.3 | 115.3 | 296.9 | 637.2 | 1,052.8 | 1,408.2 |
| 2003..... | 186.6 | 3.6 | 2.4 | 2.6 | 3.6 | 10.3 | 39.7 | 118.6 | 306.2 | 645.5 | 1,070.0 | 1,453.0 |
| 2002..... | 187.6 | 3.6 | 2.8 | 2.6 | 3.8 | 10.6 | 40.8 | 121.5 | 309.7 | 658.8 | 1,076.3 | 1,460.0 |
| 2001..... | 188.7 | 3.0 | 2.9 | 2.6 | 3.7 | 11.3 | 41.8 | 124.4 | 315.9 | 663.5 | 1,075.1 | 1,483.0 |
| 2000..... | 191.1 | 3.8 | 2.8 | 2.5 | 3.8 | 10.8 | 41.2 | 125.8 | 325.8 | 674.8 | 1,089.5 | 1,531.6 |
| 1999..... | 190.3 | 3.2 | 2.9 | 2.6 | 4.0 | 10.6 | 41.5 | 125.3 | 328.9 | 681.0 | 1,079.0 | 1,508.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 60.8 | 4.1 | 2.2 | 2.9 | 4.5 | 7.3 | 22.4 | 76.8 | 213.4 | 480.7 | 895.5 | 1,179.7 |
| 2003..... | 61.8 | 3.2 | 3.2 | 3.1 | 4.4 | 7.5 | 23.3 | 81.8 | 217.8 | 493.9 | 919.0 | 1,283.1 |
| 2002..... | 61.2 | 4.2 | 2.5 | 3.2 | 4.6 | 7.1 | 23.8 | 79.7 | 218.1 | 504.9 | 919.6 | 1,357.5 |
| 2001..... | 62.0 | * | 3.2 | 2.8 | 4.4 | 7.5 | 26.3 | 84.3 | 227.1 | 512.8 | 937.0 | 1,421.2 |
| 2000..... | 61.4 | 3.0 | 2.8 | 3.1 | 4.3 | 7.7 | 26.0 | 83.5 | 224.1 | 525.3 | 977.8 | 1,453.8 |
| 1999..... | 61.1 | 3.2 | 2.9 | 3.0 | 4.3 | 7.8 | 25.0 | 86.2 | 228.8 | 531.1 | 949.9 | 1,451.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 62.4 | * | 1.8 | 3.1 | 5.7 | 6.8 | 19.2 | 79.1 | 243.0 | 595.5 | 1,202.1 | 1,566.3 |
| 2003..... | 63.0 | * | 3.7 | 3.4 | 5.3 | 7.3 | 18.7 | 82.3 | 250.4 | 627.6 | 1,197.6 | 1,721.0 |
| 2002..... | 62.6 | * | 2.4 | 3.3 | 5.4 | 6.5 | 18.9 | 79.9 | 258.0 | 632.6 | 1,222.4 | 1,925.9 |
| 2001..... | 63.6 | * | 3.3 | 2.8 | 5.7 | 6.5 | 22.6 | 83.2 | 257.5 | 652.8 | 1,288.2 | 2,037.9 |
| 2000..... | 62.6 | * | 3.0 | 3.7 | 4.8 | 7.3 | 20.7 | 80.6 | 256.2 | 660.9 | 1,333.0 | 2,136.4 |
| 1999..... | 62.5 | * | 2.6 | 3.0 | 4.9 | 7.0 | 20.6 | 86.4 | 272.0 | 679.3 | 1,277.4 | 1,995.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 59.2 | * | 2.6 | 2.6 | 3.1 | 7.9 | 25.9 | 74.5 | 186.6 | 388.5 | 683.8 | 981.0 |
| 2003..... | 60.6 | * | 2.6 | 2.7 | 3.4 | 7.7 | 28.4 | 81.3 | 188.2 | 387.4 | 727.6 | 1,060.9 |
| 2002..... | 59.6 | * | 2.6 | 3.2 | 3.7 | 7.7 | 29.0 | 79.5 | 182.2 | 403.3 | 712.2 | 1,076.0 |
| 2001..... | 60.2 | * | 3.1 | 2.8 | 3.0 | 8.8 | 30.3 | 85.4 | 200.0 | 402.3 | 699.6 | 1,119.8 |
| 2000..... | 60.1 | * | 2.6 | 2.4 | 3.8 | 8.1 | 31.7 | 86.3 | 195.6 | 418.6 | 739.2 | 1,118.8 |
| 1999..... | 59.6 | * | 3.3 | 2.9 | 3.6 | 8.7 | 29.7 | 86.0 | 190.6 | 414.9 | 732.0 | 1,189.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| II. Neoplasms (C00-D48) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 214.5 | 3.4 | 3.0 | 2.7 | 4.3 | 10.0 | 35.9 | 125.1 | 347.5 | 787.5 | 1,337.6 | 1,753.3 |
| 2003..... | 217.0 | 3.4 | 2.7 | 2.8 | 4.2 | 10.3 | 37.5 | 127.8 | 357.0 | 801.9 | 1,358.4 | 1,796.5 |
| 2002..... | 218.5 | 3.3 | 3.1 | 2.7 | 4.4 | 10.7 | 38.4 | 129.8 | 365.5 | 823.6 | 1,365.8 | 1,816.6 |
| 2001..... | 219.2 | 3.0 | 3.1 | 2.7 | 4.5 | 11.0 | 39.0 | 132.1 | 370.9 | 834.9 | 1,368.9 | 1,860.1 |
| 2000..... | 220.8 | 4.2 | 3.1 | 2.6 | 4.7 | 10.7 | 38.7 | 133.0 | 381.8 | 847.0 | 1,386.9 | 1,916.2 |
| 1999..... | 220.7 | 3.4 | 3.2 | 2.7 | 4.8 | 10.8 | 39.3 | 132.9 | 389.4 | 858.0 | 1,381.5 | 1,899.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 227.1 | 3.3 | 3.2 | 2.9 | 4.9 | 9.6 | 31.7 | 131.1 | 392.4 | 944.9 | 1,733.7 | 2,488.3 |
| 2003..... | 229.5 | 3.5 | 3.0 | 2.9 | 4.8 | 9.9 | 33.7 | 133.7 | 402.1 | 967.5 | 1,767.0 | 2,547.9 |
| 2002..... | 231.8 | 3.2 | 3.3 | 3.0 | 5.0 | 10.4 | 34.4 | 134.5 | 415.5 | 1,001.3 | 1,779.6 | 2,616.1 |
| 2001..... | 232.6 | 3.1 | 3.3 | 2.8 | 5.2 | 10.4 | 34.8 | 136.7 | 421.0 | 1,021.6 | 1,794.0 | 2,743.9 |
| 2000..... | 234.0 | 4.4 | 3.4 | 2.8 | 5.5 | 10.1 | 35.2 | 137.3 | 432.4 | 1,037.7 | 1,824.3 | 2,841.8 |
| 1999..... | 235.2 | 3.2 | 3.4 | 2.9 | 5.5 | 10.8 | 35.9 | 137.5 | 444.6 | 1,055.7 | 1,830.6 | 2,858.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 202.6 | 3.5 | 2.7 | 2.4 | 3.7 | 10.4 | 40.1 | 119.4 | 305.7 | 654.8 | 1,071.3 | 1,424.6 |
| 2003..... | 205.1 | 3.4 | 2.3 | 2.6 | 3.6 | 10.8 | 41.2 | 122.0 | 315.0 | 663.1 | 1,085.8 | 1,466.4 |
| 2002..... | 205.9 | 3.5 | 2.9 | 2.4 | 3.8 | 11.1 | 42.3 | 125.2 | 319.2 | 675.3 | 1,091.5 | 1,471.6 |
| 2001..... | 206.4 | 3.0 | 2.9 | 2.6 | 3.9 | 11.7 | 43.2 | 127.7 | 324.5 | 679.7 | 1,089.9 | 1,492.5 |
| 2000..... | 208.4 | 4.1 | 2.8 | 2.5 | 3.8 | 11.3 | 42.2 | 128.9 | 334.9 | 689.5 | 1,102.0 | 1,542.0 |
| 1999..... | 207.0 | 3.5 | 2.9 | 2.5 | 4.0 | 10.9 | 42.7 | 128.4 | 338.4 | 695.7 | 1,090.0 | 1,515.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 232.8 | 3.7 | 3.1 | 2.6 | 4.2 | 9.7 | 34.4 | 119.0 | 340.7 | 786.8 | 1,342.8 | 1,762.5 |
| 2003..... | 235.3 | 3.7 | 2.7 | 2.7 | 4.2 | 10.0 | 36.0 | 121.1 | 350.4 | 802.2 | 1,360.6 | 1,802.1 |
| 2002..... | 236.6 | 3.4 | 2.9 | 2.7 | 4.5 | 10.4 | 37.0 | 124.0 | 357.6 | 822.7 | 1,365.9 | 1,819.8 |
| 2001..... | 236.5 | 3.1 | 3.3 | 2.6 | 4.5 | 10.8 | 37.3 | 126.5 | 363.4 | 830.6 | 1,365.1 | 1,860.0 |
| 2000..... | 238.1 | 4.5 | 3.2 | 2.7 | 4.5 | 10.5 | 36.4 | 126.9 | 374.5 | 842.0 | 1,382.1 | 1,914.2 |
| 1999..... | 237.2 | 3.1 | 3.4 | 2.6 | 4.6 | 10.6 | 37.3 | 126.6 | 379.8 | 852.1 | 1,374.2 | 1,894.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 245.1 | 3.6 | 3.3 | 2.9 | 4.7 | 9.8 | 30.6 | 123.7 | 379.9 | 934.0 | 1,725.1 | 2,492.7 |
| 2003..... | 247.7 | 3.8 | 3.0 | 2.9 | 4.8 | 9.8 | 32.9 | 125.7 | 389.9 | 958.8 | 1,756.2 | 2,549.6 |
| 2002..... | 249.6 | 3.5 | 3.2 | 2.9 | 5.1 | 10.3 | 33.4 | 127.8 | 400.8 | 992.0 | 1,764.1 | 2,613.7 |
| 2001..... | 249.4 | 3.2 | 3.4 | 2.6 | 5.1 | 10.4 | 33.5 | 129.5 | 407.3 | 1,006.5 | 1,771.9 | 2,728.2 |
| 2000..... | 250.2 | 4.7 | 3.5 | 2.8 | 5.4 | 10.1 | 33.1 | 129.3 | 417.8 | 1,019.5 | 1,800.1 | 2,827.2 |
| 1999..... | 250.9 | 3.2 | 3.7 | 2.8 | 5.2 | 10.5 | 34.3 | 128.7 | 428.2 | 1,038.5 | 1,805.8 | 2,838.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 220.9 | 3.7 | 2.9 | 2.3 | 3.6 | 9.6 | 38.1 | 114.3 | 303.4 | 660.1 | 1,082.3 | 1,435.4 |
| 2003..... | 223.3 | 3.5 | 2.5 | 2.5 | 3.6 | 10.3 | 39.2 | 116.6 | 312.9 | 668.0 | 1,093.5 | 1,473.1 |
| 2002..... | 224.1 | 3.2 | 2.7 | 2.5 | 3.9 | 10.6 | 40.5 | 120.3 | 316.7 | 678.4 | 1,099.0 | 1,476.8 |
| 2001..... | 224.2 | 3.0 | 3.2 | 2.6 | 3.9 | 11.1 | 41.2 | 123.5 | 321.8 | 681.4 | 1,095.3 | 1,498.8 |
| 2000..... | 226.5 | 4.2 | 2.7 | 2.6 | 3.7 | 10.9 | 39.7 | 124.5 | 333.6 | 692.5 | 1,107.4 | 1,546.1 |
| 1999..... | 224.0 | 3.1 | 3.1 | 2.4 | 3.9 | 10.6 | 40.2 | 124.5 | 334.2 | 696.1 | 1,091.6 | 1,518.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| II. Neoplasms (C00-D48) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 170.7 | * | 2.8 | 2.9 | 5.3 | 13.2 | 49.5 | 184.1 | 473.5 | 944.9 | 1,497.9 | 1,852.1 |
| 2003..... | 173.0 | * | 2.6 | 3.0 | 4.3 | 13.2 | 51.7 | 192.2 | 486.2 | 954.4 | 1,537.8 | 1,926.4 |
| 2002..... | 174.7 | 3.2 | 3.5 | 2.9 | 4.3 | 13.4 | 52.7 | 188.5 | 500.2 | 990.0 | 1,569.1 | 1,968.7 |
| 2001..... | 176.0 | * | 2.2 | 3.1 | 4.8 | 13.8 | 53.9 | 190.7 | 500.5 | 1,020.1 | 1,591.4 | 2,020.2 |
| 2000..... | 178.1 | 3.5 | 3.2 | 2.5 | 5.6 | 13.0 | 56.4 | 199.4 | 510.8 | 1,034.4 | 1,614.1 | 2,079.4 |
| 1999..... | 180.2 | 4.4 | 2.6 | 3.0 | 6.1 | 13.2 | 56.5 | 199.4 | 536.9 | 1,049.5 | 1,626.0 | 2,086.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 185.5 | * | 3.1 | 2.9 | 5.9 | 10.8 | 41.0 | 204.3 | 588.5 | 1,247.2 | 2,156.8 | 2,843.1 |
| 2003..... | 187.8 | * | 3.1 | 3.2 | 4.9 | 11.3 | 44.1 | 214.0 | 604.0 | 1,258.6 | 2,212.5 | 2,927.8 |
| 2002..... | 190.7 | * | 3.4 | 3.4 | 4.9 | 12.0 | 45.2 | 204.5 | 629.1 | 1,307.9 | 2,286.9 | 3,089.3 |
| 2001..... | 193.9 | * | 2.7 | 3.4 | 5.6 | 11.4 | 46.9 | 211.5 | 620.7 | 1,372.3 | 2,352.8 | 3,326.3 |
| 2000..... | 198.0 | * | 3.3 | 2.9 | 6.7 | 11.0 | 50.8 | 221.8 | 644.4 | 1,402.2 | 2,408.7 | 3,366.4 |
| 1999..... | 200.5 | * | 2.9 | 3.2 | 7.1 | 12.7 | 49.8 | 228.2 | 678.2 | 1,406.2 | 2,416.2 | 3,425.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 157.2 | * | 2.5 | 2.9 | 4.7 | 15.3 | 56.9 | 166.7 | 381.2 | 729.4 | 1,125.9 | 1,460.5 |
| 2003..... | 159.6 | * | 2.2 | 2.7 | 3.8 | 14.9 | 58.4 | 173.5 | 391.7 | 738.1 | 1,156.8 | 1,538.5 |
| 2002..... | 160.1 | * | 3.7 | 2.3 | 3.8 | 14.7 | 59.2 | 174.8 | 396.8 | 764.6 | 1,163.1 | 1,540.5 |
| 2001..... | 159.8 | * | * | 2.7 | 4.0 | 15.9 | 60.1 | 172.8 | 403.7 | 771.6 | 1,163.9 | 1,536.5 |
| 2000..... | 160.2 | * | 3.1 | 2.2 | 4.4 | 14.8 | 61.4 | 180.1 | 403.3 | 775.7 | 1,166.9 | 1,601.3 |
| 1999..... | 161.8 | * | 2.2 | 2.9 | 5.1 | 13.7 | 62.5 | 174.8 | 423.6 | 797.8 | 1,183.5 | 1,586.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms (C00-C97) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 188.6 | 1.8 | 2.5 | 2.5 | 4.1 | 9.1 | 33.4 | 119.0 | 333.4 | 755.1 | 1,280.4 | 1,653.3 |
| 2003..... | 191.5 | 1.9 | 2.5 | 2.6 | 4.0 | 9.4 | 35.0 | 122.2 | 343.0 | 770.3 | 1,302.5 | 1,698.2 |
| 2002..... | 193.2 | 1.8 | 2.6 | 2.6 | 4.3 | 9.7 | 35.8 | 123.8 | 351.1 | 792.1 | 1,311.9 | 1,723.9 |
| 2001..... | 194.4 | 1.6 | 2.7 | 2.5 | 4.3 | 10.1 | 36.8 | 126.5 | 356.5 | 802.8 | 1,315.8 | 1,765.6 |
| 2000..... | 196.5 | 2.4 | 2.7 | 2.5 | 4.4 | 9.8 | 36.6 | 127.5 | 366.7 | 816.3 | 1,335.6 | 1,819.4 |
| 1999..... | 197.0 | 1.8 | 2.7 | 2.5 | 4.5 | 10.0 | 37.1 | 127.6 | 374.6 | 827.1 | 1,331.5 | 1,805.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 198.4 | 1.8 | 2.6 | 2.7 | 4.8 | 8.6 | 29.1 | 124.3 | 376.7 | 907.6 | 1,662.1 | 2,349.5 |
| 2003..... | 201.3 | 1.7 | 2.8 | 2.8 | 4.6 | 8.9 | 30.8 | 127.4 | 386.8 | 931.7 | 1,695.4 | 2,413.8 |
| 2002..... | 203.8 | 2.0 | 2.7 | 2.9 | 4.9 | 9.2 | 31.5 | 128.0 | 399.8 | 964.8 | 1,711.3 | 2,491.1 |
| 2001..... | 205.3 | 1.5 | 2.9 | 2.5 | 5.0 | 9.3 | 32.6 | 130.3 | 405.2 | 984.6 | 1,727.1 | 2,613.6 |
| 2000..... | 207.2 | 2.6 | 3.0 | 2.7 | 5.1 | 9.2 | 32.7 | 130.9 | 415.8 | 1,001.9 | 1,760.6 | 2,710.7 |
| 1999..... | 208.9 | 1.9 | 2.9 | 2.7 | 5.2 | 9.7 | 33.3 | 131.7 | 428.4 | 1,019.4 | 1,767.0 | 2,722.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 179.1 | 1.9 | 2.4 | 2.2 | 3.4 | 9.6 | 37.7 | 113.8 | 293.2 | 627.1 | 1,023.5 | 1,340.1 |
| 2003..... | 182.0 | 2.1 | 2.1 | 2.4 | 3.4 | 9.9 | 39.1 | 117.1 | 302.3 | 635.3 | 1,040.1 | 1,381.9 |
| 2002..... | 183.0 | 1.6 | 2.4 | 2.4 | 3.6 | 10.2 | 40.0 | 119.8 | 306.0 | 648.5 | 1,046.7 | 1,391.1 |
| 2001..... | 183.9 | 1.8 | 2.5 | 2.4 | 3.5 | 10.9 | 41.0 | 122.7 | 311.5 | 652.2 | 1,045.4 | 1,410.7 |
| 2000..... | 186.2 | 2.3 | 2.5 | 2.2 | 3.6 | 10.4 | 40.4 | 124.2 | 321.3 | 663.6 | 1,058.5 | 1,456.4 |
| 1999..... | 185.6 | 1.7 | 2.5 | 2.3 | 3.7 | 10.3 | 40.9 | 123.7 | 325.0 | 669.5 | 1,048.5 | 1,436.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 59.3 | * | 2.0 | 2.8 | 4.3 | 6.9 | 21.8 | 75.5 | 209.7 | 472.0 | 872.2 | 1,131.2 |
| 2003..... | 60.3 | * | 3.0 | 2.9 | 4.1 | 7.1 | 22.9 | 80.7 | 214.7 | 485.2 | 890.1 | 1,234.9 |
| 2002..... | 59.7 | * | 2.3 | 2.9 | 4.5 | 6.8 | 23.2 | 78.2 | 215.1 | 496.0 | 894.9 | 1,308.8 |
| 2001..... | 60.5 | * | 2.8 | 2.5 | 4.3 | 7.2 | 25.6 | 82.8 | 223.4 | 504.0 | 913.9 | 1,363.7 |
| 2000..... | 59.9 | * | 2.5 | 2.9 | 4.2 | 7.3 | 25.2 | 82.1 | 221.0 | 516.2 | 953.5 | 1,383.5 |
| 1999..... | 59.6 | * | 2.5 | 2.7 | 4.1 | 7.4 | 24.4 | 84.9 | 225.1 | 520.5 | 926.8 | 1,399.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 60.9 | * | 1.6 | 3.0 | 5.5 | 6.4 | 18.6 | 77.4 | 239.0 | 585.8 | 1,174.2 | 1,508.8 |
| 2003..... | 61.5 | * | 3.6 | 3.2 | 4.9 | 6.8 | 18.2 | 81.1 | 246.5 | 617.6 | 1,163.9 | 1,668.6 |
| 2002..... | 61.2 | * | 2.2 | 2.9 | 5.2 | 6.3 | 18.4 | 78.4 | 254.3 | 622.3 | 1,190.8 | 1,869.0 |
| 2001..... | 62.2 | * | 2.9 | 2.5 | 5.5 | 6.3 | 21.7 | 81.5 | 253.5 | 642.8 | 1,258.3 | 1,967.4 |
| 2000..... | 61.3 | * | 2.9 | 3.5 | 4.5 | 6.9 | 20.1 | 79.4 | 253.1 | 651.2 | 1,306.4 | 2,049.7 |
| 1999..... | 61.2 | * | 2.3 | 2.8 | 4.6 | 6.6 | 19.8 | 84.6 | 268.1 | 668.9 | 1,252.9 | 1,945.3 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 57.7 | * | 2.4 | 2.5 | 3.0 | 7.5 | 25.4 | 73.5 | 183.0 | 380.7 | 663.6 | 937.0 |
| 2003..... | 59.1 | * | 2.4 | 2.6 | 3.2 | 7.4 | 28.0 | 80.2 | 185.9 | 379.7 | 702.1 | 1,014.8 |
| 2002..... | 58.1 | * | 2.4 | 2.9 | 3.6 | 7.5 | 28.4 | 78.0 | 179.8 | 395.6 | 692.2 | 1,031.2 |
| 2001..... | 58.7 | * | 2.7 | 2.5 | 2.9 | 8.3 | 29.8 | 84.2 | 196.7 | 394.5 | 681.2 | 1,068.6 |
| 2000..... | 58.5 | * | 2.2 | 2.3 | 3.7 | 7.8 | 30.7 | 84.7 | 192.5 | 410.0 | 716.5 | 1,056.5 |
| 1999..... | 58.0 | * | 2.7 | 2.6 | 3.5 | 8.3 | 29.4 | 85.1 | 187.2 | 404.3 | 709.8 | 1,136.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms (C00-C97) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 209.4 | 1.7 | 2.6 | 2.4 | 4.0 | 9.6 | 35.2 | 123.4 | 342.9 | 774.5 | 1,301.1 | 1,674.3 |
| 2003..... | 211.9 | 2.0 | 2.3 | 2.6 | 4.0 | 9.9 | 36.7 | 126.0 | 352.4 | 788.9 | 1,321.6 | 1,714.7 |
| 2002..... | 213.4 | 1.7 | 2.6 | 2.5 | 4.2 | 10.3 | 37.6 | 128.0 | 360.9 | 810.5 | 1,329.2 | 1,736.8 |
| 2001..... | 213.9 | 1.6 | 2.7 | 2.4 | 4.2 | 10.6 | 38.2 | 130.3 | 365.9 | 820.5 | 1,331.4 | 1,776.6 |
| 2000..... | 215.6 | 2.5 | 2.7 | 2.4 | 4.4 | 10.3 | 37.9 | 131.2 | 376.6 | 832.8 | 1,348.9 | 1,831.1 |
| 1999..... | 215.5 | 1.7 | 2.7 | 2.4 | 4.5 | 10.4 | 38.6 | 131.0 | 384.6 | 843.2 | 1,344.5 | 1,816.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 221.8 | 1.8 | 2.8 | 2.7 | 4.6 | 9.1 | 30.8 | 129.2 | 387.0 | 928.8 | 1,687.2 | 2,387.3 |
| 2003..... | 224.3 | 1.9 | 2.6 | 2.7 | 4.5 | 9.4 | 32.8 | 131.9 | 396.9 | 951.1 | 1,720.2 | 2,444.0 |
| 2002..... | 226.7 | 1.9 | 2.8 | 2.8 | 4.8 | 9.9 | 33.5 | 132.6 | 410.0 | 985.1 | 1,733.2 | 2,513.2 |
| 2001..... | 227.2 | 1.6 | 2.9 | 2.5 | 4.9 | 9.9 | 34.0 | 134.6 | 415.3 | 1,003.8 | 1,745.1 | 2,634.8 |
| 2000..... | 228.7 | 2.6 | 2.9 | 2.5 | 5.2 | 9.7 | 34.3 | 135.1 | 426.6 | 1,020.2 | 1,776.0 | 2,732.4 |
| 1999..... | 229.9 | 1.6 | 3.0 | 2.6 | 5.3 | 10.3 | 35.0 | 135.4 | 438.8 | 1,036.8 | 1,783.7 | 2,747.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 197.5 | 1.6 | 2.4 | 2.2 | 3.4 | 10.0 | 39.5 | 117.8 | 301.9 | 644.5 | 1,041.6 | 1,355.6 |
| 2003..... | 200.0 | 2.1 | 2.1 | 2.4 | 3.4 | 10.4 | 40.6 | 120.5 | 311.1 | 652.7 | 1,055.6 | 1,394.3 |
| 2002..... | 200.8 | 1.5 | 2.4 | 2.2 | 3.6 | 10.6 | 41.6 | 123.5 | 315.4 | 664.7 | 1,061.5 | 1,401.8 |
| 2001..... | 201.2 | 1.6 | 2.5 | 2.3 | 3.6 | 11.3 | 42.4 | 126.0 | 320.0 | 668.1 | 1,059.8 | 1,419.6 |
| 2000..... | 203.1 | 2.4 | 2.6 | 2.2 | 3.6 | 10.9 | 41.4 | 127.3 | 330.3 | 678.1 | 1,070.6 | 1,466.6 |
| 1999..... | 201.9 | 1.8 | 2.5 | 2.2 | 3.7 | 10.5 | 42.1 | 126.8 | 334.5 | 684.2 | 1,059.4 | 1,443.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 227.0 | 1.9 | 2.7 | 2.4 | 3.9 | 9.3 | 33.7 | 117.3 | 336.2 | 773.6 | 1,305.4 | 1,680.9 |
| 2003..... | 229.5 | 2.2 | 2.4 | 2.6 | 4.0 | 9.6 | 35.2 | 119.4 | 345.9 | 788.8 | 1,322.6 | 1,717.8 |
| 2002..... | 230.9 | 1.6 | 2.6 | 2.5 | 4.3 | 10.0 | 36.2 | 122.3 | 353.0 | 809.2 | 1,328.4 | 1,737.9 |
| 2001..... | 230.6 | 1.8 | 2.9 | 2.4 | 4.3 | 10.3 | 36.5 | 124.6 | 358.3 | 815.9 | 1,326.5 | 1,775.2 |
| 2000..... | 232.2 | 2.6 | 2.8 | 2.4 | 4.3 | 10.1 | 35.6 | 125.1 | 369.3 | 827.7 | 1,342.9 | 1,826.6 |
| 1999..... | 231.5 | 1.6 | 3.0 | 2.3 | 4.3 | 10.2 | 36.6 | 124.8 | 375.0 | 837.1 | 1,336.4 | 1,808.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 239.2 | 2.0 | 2.8 | 2.7 | 4.4 | 9.3 | 29.9 | 121.9 | 374.6 | 917.5 | 1,677.3 | 2,387.1 |
| 2003..... | 241.8 | 2.1 | 2.6 | 2.7 | 4.6 | 9.4 | 31.9 | 123.8 | 384.8 | 942.0 | 1,707.8 | 2,441.7 |
| 2002..... | 243.8 | 2.0 | 2.8 | 2.7 | 4.9 | 9.8 | 32.5 | 125.9 | 395.5 | 975.3 | 1,716.5 | 2,507.7 |
| 2001..... | 243.4 | 1.7 | 3.1 | 2.4 | 4.9 | 10.0 | 32.6 | 127.3 | 401.7 | 988.2 | 1,721.8 | 2,616.8 |
| 2000..... | 244.4 | 2.9 | 3.1 | 2.6 | 5.0 | 9.7 | 32.3 | 127.2 | 412.0 | 1,002.1 | 1,750.2 | 2,714.1 |
| 1999..... | 245.1 | 1.7 | 3.3 | 2.6 | 5.0 | 10.1 | 33.5 | 126.7 | 422.3 | 1,019.3 | 1,757.7 | 2,721.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 215.3 | 1.8 | 2.5 | 2.1 | 3.3 | 9.3 | 37.5 | 112.9 | 299.8 | 649.8 | 1,052.0 | 1,364.5 |
| 2003..... | 217.6 | 2.3 | 2.2 | 2.4 | 3.4 | 9.8 | 38.6 | 115.1 | 308.9 | 657.6 | 1,062.4 | 1,399.1 |
| 2002..... | 218.5 | * | 2.3 | 2.3 | 3.6 | 10.3 | 39.9 | 118.7 | 312.8 | 667.7 | 1,068.3 | 1,405.4 |
| 2001..... | 218.4 | 1.8 | 2.8 | 2.4 | 3.6 | 10.7 | 40.4 | 121.9 | 317.3 | 669.7 | 1,064.4 | 1,425.1 |
| 2000..... | 220.6 | 2.4 | 2.4 | 2.3 | 3.5 | 10.5 | 38.9 | 123.0 | 328.9 | 681.0 | 1,075.3 | 1,468.7 |
| 1999..... | 218.4 | * | 2.7 | 2.1 | 3.6 | 10.3 | 39.7 | 123.0 | 330.5 | 684.6 | 1,060.5 | 1,444.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms (C00-C97) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 167.5 | * | 2.5 | 2.6 | 5.0 | 12.4 | 48.3 | 181.8 | 467.2 | 931.9 | 1,465.9 | 1,794.3 |
| 2003..... | 170.0 | * | 2.3 | 2.7 | 4.1 | 12.6 | 50.5 | 189.7 | 480.9 | 941.1 | 1,508.2 | 1,867.5 |
| 2002..... | 171.5 | * | 3.0 | 2.6 | 4.1 | 12.8 | 51.5 | 186.0 | 494.6 | 977.2 | 1,537.4 | 1,904.3 |
| 2001..... | 172.8 | * | 2.0 | 2.7 | 4.4 | 13.4 | 52.8 | 188.7 | 494.5 | 1,005.6 | 1,559.5 | 1,947.5 |
| 2000..... | 174.9 | * | 2.8 | 2.3 | 5.3 | 12.3 | 55.3 | 196.7 | 505.2 | 1,019.3 | 1,582.8 | 2,015.1 |
| 1999..... | 176.9 | * | 2.0 | 2.8 | 5.6 | 12.6 | 55.3 | 196.8 | 530.5 | 1,033.6 | 1,592.9 | 2,024.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 182.4 | * | 2.8 | 2.5 | 5.6 | 10.1 | 39.8 | 201.8 | 580.5 | 1,231.5 | 2,116.9 | 2,781.1 |
| 2003..... | 184.8 | * | 2.4 | 2.9 | 4.7 | 10.6 | 42.9 | 211.5 | 597.1 | 1,242.1 | 2,175.8 | 2,859.6 |
| 2002..... | 187.8 | * | 2.7 | 3.1 | 4.6 | 11.7 | 44.3 | 202.3 | 621.8 | 1,293.5 | 2,247.6 | 3,008.4 |
| 2001..... | 190.8 | * | 2.4 | 3.0 | 5.1 | 10.9 | 45.8 | 209.6 | 614.1 | 1,354.7 | 2,310.5 | 3,238.5 |
| 2000..... | 194.9 | * | 2.6 | 2.6 | 6.4 | 10.5 | 49.6 | 219.0 | 637.4 | 1,382.7 | 2,372.3 | 3,290.1 |
| 1999..... | 197.4 | * | 2.0 | 2.9 | 6.6 | 11.9 | 48.6 | 225.8 | 670.9 | 1,387.0 | 2,377.4 | 3,361.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 153.9 | * | 2.2 | 2.7 | 4.4 | 14.5 | 55.8 | 164.6 | 376.3 | 718.3 | 1,098.4 | 1,404.3 |
| 2003..... | 156.5 | * | 2.1 | 2.4 | 3.5 | 14.5 | 57.2 | 170.9 | 387.6 | 727.0 | 1,131.2 | 1,483.2 |
| 2002..... | 156.7 | * | 3.2 | 2.0 | 3.6 | 13.9 | 57.8 | 172.1 | 392.5 | 752.8 | 1,135.6 | 1,482.4 |
| 2001..... | 156.5 | * | * | 2.3 | 3.8 | 15.7 | 59.0 | 170.7 | 398.3 | 759.4 | 1,137.8 | 1,469.3 |
| 2000..... | 156.9 | * | 3.0 | 1.9 | 4.2 | 14.0 | 60.4 | 177.5 | 398.8 | 763.6 | 1,138.4 | 1,541.6 |
| 1999..... | 158.3 | * | 2.0 | 2.7 | 4.8 | 13.3 | 61.3 | 172.0 | 417.9 | 784.2 | 1,153.6 | 1,524.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | 0.1 | 0.1 | 0.6 | 3.0 | 6.3 | 10.4 | 13.9 | 18.2 |
| 2003..... | 3.0 | * | * | * | 0.1 | 0.2 | 0.7 | 2.9 | 6.8 | 10.5 | 13.3 | 18.7 |
| 2002..... | 3.0 | * | * | * | 0.1 | 0.2 | 0.6 | 2.9 | 7.1 | 10.7 | 13.7 | 18.7 |
| 2001..... | 3.0 | * | * | * | 0.1 | 0.2 | 0.7 | 2.8 | 7.1 | 10.6 | 14.3 | 18.4 |
| 2000..... | 2.9 | * | * | * | 0.1 | 0.1 | 0.7 | 3.0 | 6.9 | 10.8 | 13.3 | 19.5 |
| 1999..... | 2.9 | * | * | * | 0.1 | 0.1 | 0.7 | 3.0 | 7.0 | 10.5 | 14.3 | 19.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 0.2 | 0.9 | 4.8 | 10.3 | 15.8 | 20.7 | 24.9 |
| 2003..... | 4.1 | * | * | * | * | 0.2 | 1.0 | 4.8 | 10.9 | 16.0 | 19.5 | 27.1 |
| 2002..... | 4.1 | * | * | * | * | 0.2 | 0.8 | 4.6 | 11.3 | 16.2 | 19.8 | 26.1 |
| 2001..... | 4.1 | * | * | * | * | 0.2 | 1.0 | 4.6 | 11.6 | 16.2 | 20.6 | 25.8 |
| 2000..... | 3.9 | * | * | * | * | 0.1 | 1.0 | 4.6 | 11.1 | 16.1 | 19.4 | 26.5 |
| 1999..... | 4.0 | * | * | * | 0.1 | 0.2 | 1.0 | 4.8 | 11.0 | 15.8 | 21.0 | 27.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.4 | 1.3 | 2.6 | 5.8 | 9.4 | 15.1 |
| 2003..... | 1.9 | * | * | * | * | * | 0.4 | 1.1 | 2.9 | 5.9 | 9.2 | 15.0 |
| 2002..... | 1.9 | * | * | * | * | * | 0.4 | 1.2 | 3.2 | 6.1 | 9.7 | 15.3 |
| 2001..... | 1.9 | * | * | * | * | 0.1 | 0.4 | 1.2 | 2.9 | 6.0 | 10.2 | 15.3 |
| 2000..... | 2.0 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.0 | 6.5 | 9.3 | 16.6 |
| 1999..... | 2.0 | * | * | * | * | * | 0.4 | 1.4 | 3.2 | 6.0 | 9.9 | 16.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | 0.1 | 0.5 | 2.6 | 5.8 | 10.2 | 13.9 | 18.7 |
| 2003..... | 3.1 | * | * | * | * | 0.1 | 0.6 | 2.6 | 6.3 | 10.1 | 13.0 | 19.1 |
| 2002..... | 3.1 | * | * | * | * | 0.1 | 0.5 | 2.5 | 6.5 | 10.6 | 13.9 | 18.9 |
| 2001..... | 3.1 | * | * | * | * | 0.1 | 0.5 | 2.5 | 6.5 | 10.2 | 14.5 | 18.9 |
| 2000..... | 3.0 | * | * | * | * | 0.1 | 0.6 | 2.5 | 6.1 | 10.4 | 13.3 | 19.8 |
| 1999..... | 3.0 | * | * | * | * | 0.1 | 0.5 | 2.4 | 6.3 | 10.1 | 14.2 | 19.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 0.2 | 0.7 | 4.0 | 9.4 | 15.0 | 20.4 | 24.8 |
| 2003..... | 4.1 | * | * | * | * | 0.2 | 1.0 | 4.2 | 9.9 | 14.9 | 18.7 | 27.4 |
| 2002..... | 4.1 | * | * | * | * | 0.2 | 0.7 | 4.0 | 10.2 | 15.7 | 19.9 | 26.8 |
| 2001..... | 4.1 | * | * | * | * | 0.2 | 0.8 | 4.0 | 10.4 | 15.3 | 20.5 | 26.4 |
| 2000..... | 3.9 | * | * | * | * | * | 0.8 | 3.8 | 9.7 | 15.0 | 18.9 | 27.4 |
| 1999..... | 3.9 | * | * | * | * | * | 0.8 | 3.8 | 9.7 | 15.1 | 20.3 | 27.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | 0.3 | 1.1 | 2.4 | 6.0 | 9.6 | 16.0 |
| 2003..... | 2.0 | * | * | * | * | * | 0.2 | 0.9 | 2.9 | 5.9 | 9.2 | 15.4 |
| 2002..... | 2.1 | * | * | * | * | * | 0.3 | 1.0 | 2.9 | 6.2 | 9.8 | 15.5 |
| 2001..... | 2.1 | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 5.9 | 10.5 | 15.7 |
| 2000..... | 2.1 | * | * | * | * | * | 0.4 | 1.2 | 2.8 | 6.5 | 9.6 | 16.8 |
| 1999..... | 2.1 | * | * | * | * | * | 0.3 | 1.1 | 3.0 | 5.9 | 10.1 | 16.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 1.3 | 6.2 | 11.4 | 14.1 | 15.3 | 12.7 |
| 2003..... | 3.0 | * | * | * | * | * | 1.1 | 5.2 | 11.5 | 15.7 | 15.6 | 15.5 |
| 2002..... | 2.9 | * | * | * | * | * | 1.1 | 5.6 | 12.4 | 12.9 | 12.1 | 16.0 |
| 2001..... | 3.0 | * | * | * | * | * | 1.3 | 5.3 | 12.9 | 15.7 | 13.9 | 13.1 |
| 2000..... | 3.1 | * | * | * | * | * | 1.1 | 6.1 | 13.7 | 15.7 | 15.1 | 14.9 |
| 1999..... | 3.3 | * | * | * | * | * | 1.2 | 7.3 | 13.6 | 15.4 | 16.8 | 14.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | 1.9 | 10.7 | 20.3 | 27.1 | 26.2 | 27.4 |
| 2003..... | 4.8 | * | * | * | * | * | 1.2 | 9.2 | 21.8 | 29.1 | 28.6 | 25.6 |
| 2002..... | 4.4 | * | * | * | * | * | 1.6 | 9.8 | 22.5 | 23.0 | 18.8 | 26.3 |
| 2001..... | 4.6 | * | * | * | * | * | 1.7 | 8.9 | 23.9 | 27.3 | 23.4 | * |
| 2000..... | 4.9 | * | * | * | * | * | 1.8 | 10.7 | 24.7 | 28.6 | 29.0 | * |
| 1999..... | 5.0 | * | * | * | * | * | 1.8 | 12.3 | 23.5 | 25.4 | 31.9 | 25.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.9 | 2.3 | 4.3 | 4.8 | 9.2 | * |
| 2003..... | 1.4 | * | * | * | * | * | 1.1 | 1.8 | 3.3 | 6.2 | 8.3 | 11.6 |
| 2002..... | 1.5 | * | * | * | * | * | 0.7 | 2.1 | 4.3 | 5.8 | 8.3 | 12.1 |
| 2001..... | 1.5 | * | * | * | * | * | 0.9 | 2.1 | 4.0 | 7.5 | 8.6 | 10.7 |
| 2000..... | 1.5 | * | * | * | * | * | * | 2.1 | 4.9 | 6.6 | 7.3 | 14.4 |
| 1999..... | 1.7 | * | * | * | * | * | 0.7 | 3.0 | 5.7 | 8.4 | 8.4 | 10.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of tongue (C01-C02) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.8 | 1.5 | 2.5 | 2.9 | 4.3 |
| 2003..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.8 | 1.7 | 2.5 | 3.1 | 3.7 |
| 2002..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 | 2.5 | 3.3 | 4.0 |
| 2001..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 2.4 | 3.3 | 3.7 |
| 2000..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 2.4 | 3.0 | 4.0 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.8 | 1.6 | 2.4 | 3.1 | 4.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.3 | 1.2 | 2.6 | 3.7 | 4.0 | 5.0 |
| 2003..... | 1.0 | * | * | * | * | * | 0.3 | 1.2 | 2.6 | 3.7 | 4.3 | 5.0 |
| 2002..... | 0.9 | * | * | * | * | 0.3 | 1.1 | 2.7 | 3.7 | 4.5 | 5.0 | 5.0 |
| 2001..... | 0.9 | * | * | * | * | * | 0.3 | 1.2 | 2.9 | 3.7 | 4.3 | 4.1 |
| 2000..... | 0.9 | * | * | * | * | 0.3 | 1.1 | 2.8 | 3.4 | 3.7 | 4.3 | 4.8 |
| 1999..... | 0.9 | * | * | * | * | * | 0.3 | 1.2 | 2.4 | 3.8 | 4.5 | 5.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.5 | 2.2 | 3.9 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 2.3 | 3.1 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.9 | 1.5 | 2.5 | 3.6 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 2.5 | 3.5 |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.5 | 2.5 | 3.6 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.3 | 2.2 | 3.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.7 | 1.6 | 2.5 | 3.0 | 4.4 |
| 2003..... | 0.8 | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 2.5 | 3.1 | 3.8 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 2.5 | 3.5 | 4.0 |
| 2001..... | 0.8 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 2.3 | 3.3 | 3.9 |
| 2000..... | 0.7 | * | * | * | * | * | 0.2 | 0.7 | 1.6 | 2.4 | 3.1 | 4.2 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 2.4 | 3.2 | 4.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.2 | 1.1 | 2.6 | 3.6 | 4.2 | 5.0 |
| 2003..... | 1.0 | * | * | * | * | * | 0.3 | 1.2 | 2.6 | 3.6 | 4.2 | 5.1 |
| 2002..... | 1.0 | * | * | * | * | * | 0.3 | 1.0 | 2.6 | 3.5 | 4.6 | 5.0 |
| 2001..... | 1.0 | * | * | * | * | * | 0.2 | 1.1 | 2.7 | 3.6 | 4.4 | 4.4 |
| 2000..... | 0.9 | * | * | * | * | * | 0.3 | 1.0 | 2.7 | 3.3 | 3.6 | 5.1 |
| 1999..... | 0.9 | * | * | * | * | * | 0.3 | 1.0 | 2.2 | 3.7 | 4.6 | 5.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.6 | 2.3 | 4.1 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.8 | 1.5 | 2.3 | 3.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.9 | 1.6 | 2.7 | 3.6 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.6 | 3.7 |
| 2000..... | 0.6 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 2.7 | 3.8 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.2 | 2.3 | 3.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of pharynx (C10-C13,C14.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.3 | 1.3 | 2.9 | 4.6 | 5.7 | 5.0 |
| 2003..... | 1.2 | * | * | * | * | * | 0.3 | 1.2 | 3.0 | 4.8 | 5.0 | 6.0 |
| 2002..... | 1.2 | * | * | * | * | * | 0.2 | 1.3 | 3.1 | 4.6 | 5.1 | 5.4 |
| 2001..... | 1.2 | * | * | * | * | * | 0.3 | 1.3 | 3.2 | 4.6 | 5.6 | 5.9 |
| 2000..... | 1.2 | * | * | * | * | * | 0.3 | 1.2 | 3.2 | 4.8 | 5.3 | 5.5 |
| 1999..... | 1.2 | * | * | * | * | * | 0.3 | 1.3 | 3.2 | 4.6 | 5.6 | 5.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | 0.4 | 2.2 | 4.8 | 7.4 | 9.3 | 8.4 |
| 2003..... | 1.8 | * | * | * | * | * | 0.4 | 2.0 | 4.8 | 7.6 | 8.3 | 10.8 |
| 2002..... | 1.7 | * | * | * | * | * | 0.3 | 2.1 | 5.0 | 7.2 | 8.2 | 9.5 |
| 2001..... | 1.8 | * | * | * | * | * | 0.4 | 2.1 | 5.2 | 7.3 | 9.0 | 10.0 |
| 2000..... | 1.8 | * | * | * | * | * | 0.4 | 2.0 | 5.3 | 7.6 | 8.7 | 9.5 |
| 1999..... | 1.8 | * | * | * | * | * | 0.4 | 2.2 | 5.2 | 7.2 | 9.1 | 10.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 2.2 | 3.4 | 3.5 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 2.4 | 2.9 | 3.9 |
| 2002..... | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.4 | 3.1 | 3.6 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 2.3 | 3.4 | 4.2 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.5 | 1.3 | 2.6 | 3.1 | 3.8 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 2.6 | 3.3 | 3.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.2 | 1.0 | 2.4 | 4.3 | 5.5 | 5.0 |
| 2003..... | 1.2 | * | * | * | * | * | 0.2 | 0.9 | 2.6 | 4.4 | 4.7 | 5.8 |
| 2002..... | 1.2 | * | * | * | * | * | 0.2 | 1.0 | 2.6 | 4.4 | 4.9 | 5.2 |
| 2001..... | 1.2 | * | * | * | * | * | 0.2 | 0.9 | 2.7 | 4.2 | 5.4 | 5.9 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.9 | 2.6 | 4.4 | 5.1 | 5.2 |
| 1999..... | 1.1 | * | * | * | * | * | 0.2 | 0.9 | 2.7 | 4.3 | 5.2 | 5.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.3 | 1.6 | 4.0 | 6.7 | 8.8 | 7.9 |
| 2003..... | 1.7 | * | * | * | * | * | 0.3 | 1.5 | 4.1 | 6.7 | 7.5 | 10.6 |
| 2002..... | 1.7 | * | * | * | * | * | 0.2 | 1.6 | 4.2 | 6.8 | 7.8 | 9.2 |
| 2001..... | 1.7 | * | * | * | * | * | 0.3 | 1.6 | 4.4 | 6.5 | 8.6 | 9.9 |
| 2000..... | 1.6 | * | * | * | * | * | 0.2 | 1.4 | 4.2 | 6.8 | 8.2 | 9.2 |
| 1999..... | 1.6 | * | * | * | * | * | 0.3 | 1.5 | 4.4 | 6.5 | 8.3 | 10.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.4 | 1.0 | 2.1 | 3.3 | 3.7 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.3 | 1.2 | 2.4 | 2.8 | 3.7 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.4 | 1.1 | 2.4 | 2.9 | 3.5 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.3 | 1.1 | 2.3 | 3.3 | 4.3 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.4 | 1.1 | 2.4 | 3.1 | 3.6 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.4 | 1.2 | 2.4 | 3.2 | 3.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of pharynx (C10-C13,C14.0) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.6 | 3.3 | 7.1 | 8.1 | 8.2 | 5.8 |
| 2003..... | 1.6 | * | * | * | * | * | 0.5 | 2.7 | 6.2 | 8.7 | 8.0 | 7.1 |
| 2002..... | 1.5 | * | * | * | * | * | 0.6 | 2.9 | 6.5 | 6.1 | 6.4 | 7.3 |
| 2001..... | 1.5 | * | * | * | * | * | 0.7 | 2.9 | 6.6 | 7.7 | 7.4 | * |
| 2000..... | 1.7 | * | * | * | * | * | 0.6 | 2.9 | 7.9 | 8.7 | 8.0 | 7.3 |
| 1999..... | 1.7 | * | * | * | * | * | 0.6 | 3.7 | 7.0 | 8.2 | 9.8 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | 0.9 | 5.8 | 12.6 | 15.5 | 14.0 | * |
| 2003..... | 2.7 | * | * | * | * | * | * | 4.9 | 11.9 | 17.1 | 16.2 | * |
| 2002..... | 2.3 | * | * | * | * | * | 0.8 | 5.2 | 12.1 | 10.7 | 10.6 | * |
| 2001..... | 2.4 | * | * | * | * | * | 1.0 | 4.9 | 12.0 | 14.5 | 12.9 | * |
| 2000..... | 2.7 | * | * | * | * | * | 1.1 | 5.2 | 14.3 | 15.9 | 15.7 | * |
| 1999..... | 2.7 | * | * | * | * | * | 1.0 | 6.7 | 12.1 | 13.7 | 19.4 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.8 | 2.8 | 5.0 | * |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.8 | 1.6 | 2.8 | 3.4 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.9 | 2.1 | 2.9 | 4.0 | * |
| 2001..... | 0.7 | * | * | * | * | * | * | 1.2 | 2.3 | 2.8 | 4.4 | * |
| 2000..... | 0.7 | * | * | * | * | * | * | 1.0 | 2.8 | 3.7 | 3.6 | * |
| 1999..... | 0.8 | * | * | * | * | * | * | 1.2 | 2.9 | 4.3 | 4.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.2 | 0.9 | 1.8 | 3.2 | 5.0 | 8.4 |
| 2003..... | 1.0 | * | * | * | * | * | 0.2 | 0.9 | 2.1 | 3.0 | 5.0 | 8.8 |
| 2002..... | 1.0 | * | * | * | * | * | 0.2 | 0.9 | 2.2 | 3.5 | 5.1 | 8.9 |
| 2001..... | 1.0 | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 3.6 | 5.3 | 8.4 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 1.0 | 1.9 | 3.6 | 4.8 | 9.7 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.9 | 2.1 | 3.3 | 5.5 | 9.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.3 | 1.4 | 2.9 | 4.6 | 6.9 | 10.9 |
| 2003..... | 1.3 | * | * | * | * | * | 0.3 | 1.6 | 3.4 | 4.5 | 6.6 | 10.9 |
| 2002..... | 1.3 | * | * | * | * | * | 0.2 | 1.5 | 3.6 | 5.3 | 6.8 | 11.3 |
| 2001..... | 1.3 | * | * | * | * | * | 0.3 | 1.3 | 3.4 | 5.1 | 7.0 | 10.7 |
| 2000..... | 1.2 | * | * | * | * | * | 0.3 | 1.5 | 2.9 | 5.0 | 6.7 | 11.5 |
| 1999..... | 1.3 | * | * | * | * | * | 0.3 | 1.4 | 3.3 | 4.7 | 7.2 | 11.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.4 | 0.8 | 2.0 | 3.8 | 7.3 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 1.8 | 3.9 | 7.8 |
| 2002..... | 0.7 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.1 | 4.0 | 7.8 |
| 2001..... | 0.8 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 4.2 | 7.4 |
| 2000..... | 0.8 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 3.5 | 9.0 |
| 1999..... | 0.8 | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 2.2 | 4.3 | 9.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.8 | 1.8 | 3.3 | 5.2 | 8.8 |
| 2003..... | 1.1 | * | * | * | * | * | 0.2 | 0.9 | 2.0 | 3.1 | 5.0 | 9.1 |
| 2002..... | 1.1 | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 3.6 | 5.3 | 9.2 |
| 2001..... | 1.1 | * | * | * | * | * | 0.2 | 0.8 | 2.0 | 3.6 | 5.6 | 8.6 |
| 2000..... | 1.1 | * | * | * | * | * | 0.2 | 0.9 | 1.8 | 3.6 | 4.9 | 10.1 |
| 1999..... | 1.1 | * | * | * | * | * | 0.2 | 0.8 | 2.0 | 3.3 | 5.6 | 10.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.2 | 1.3 | 2.8 | 4.5 | 7.0 | 11.3 |
| 2003..... | 1.4 | * | * | * | * | * | 0.3 | 1.5 | 3.2 | 4.4 | 6.5 | 11.2 |
| 2002..... | 1.4 | * | * | * | * | * | 0.2 | 1.4 | 3.4 | 5.3 | 7.1 | 11.7 |
| 2001..... | 1.3 | * | * | * | * | * | 0.3 | 1.3 | 3.1 | 5.0 | 7.2 | 11.1 |
| 2000..... | 1.3 | * | * | * | * | * | 0.3 | 1.3 | 2.7 | 4.8 | 6.7 | 12.3 |
| 1999..... | 1.3 | * | * | * | * | * | 0.2 | 1.3 | 3.0 | 4.6 | 7.3 | 11.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.8 | 2.2 | 3.9 | 7.7 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.9 | 1.9 | 4.0 | 8.2 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 4.1 | 8.1 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.4 | 0.9 | 2.4 | 4.5 | 7.5 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.6 | 3.7 | 9.2 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.4 | 1.0 | 2.2 | 4.5 | 9.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.4 | 1.7 | 2.6 | 3.1 | 4.2 | * |
| 2003..... | 0.8 | * | * | * | * | * | 0.4 | 1.5 | 3.1 | 3.8 | 5.0 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.6 | 3.2 | 3.8 | 4.0 | 6.4 |
| 2001..... | 0.9 | * | * | * | * | * | * | 1.2 | 3.8 | 5.5 | 3.9 | * |
| 2000..... | 0.8 | * | * | * | * | * | * | 1.9 | 3.2 | 4.1 | 4.7 | * |
| 1999..... | 0.9 | * | * | * | * | * | 0.4 | 1.9 | 3.9 | 4.3 | 4.6 | 6.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 3.0 | 4.4 | 5.8 | 7.4 | * |
| 2003..... | 1.3 | * | * | * | * | * | * | 2.6 | 6.0 | 6.5 | 7.2 | * |
| 2002..... | 1.2 | * | * | * | * | * | * | 2.7 | 5.9 | 6.7 | * | * |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.9 | 7.0 | 8.5 | 6.6 | * |
| 2000..... | 1.3 | * | * | * | * | * | * | 3.2 | 5.8 | 7.6 | 8.3 | * |
| 1999..... | 1.3 | * | * | * | * | * | * | 2.9 | 6.7 | 6.9 | 7.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 3.7 | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 3.5 | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 3.3 | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.6 | * | * | * | * | * | * | 1.0 | 1.6 | 2.4 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of digestive organs (C15-C26) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 44.7 | * | 0.1 | * | 0.3 | 1.5 | 7.1 | 29.3 | 78.5 | 170.4 | 302.9 | 445.3 |
| 2003..... | 45.5 | * | 0.1 | 0.1 | 0.3 | 1.6 | 7.0 | 30.0 | 79.1 | 176.7 | 309.8 | 461.2 |
| 2002..... | 45.7 | * | 0.2 | 0.1 | 0.3 | 1.5 | 7.4 | 29.7 | 80.6 | 181.0 | 311.9 | 467.4 |
| 2001..... | 46.0 | * | * | * | 0.3 | 1.5 | 7.4 | 30.1 | 80.4 | 183.8 | 315.0 | 487.9 |
| 2000..... | 46.4 | * | 0.2 | 0.1 | 0.2 | 1.5 | 7.2 | 29.6 | 83.1 | 184.4 | 323.9 | 504.9 |
| 1999..... | 46.4 | * | 0.2 | * | 0.3 | 1.5 | 7.4 | 29.3 | 84.0 | 187.0 | 320.2 | 507.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 49.8 | * | * | * | 0.4 | 1.7 | 8.5 | 39.5 | 104.7 | 222.9 | 385.0 | 531.7 |
| 2003..... | 50.4 | * | * | * | 0.3 | 1.7 | 8.5 | 39.9 | 105.2 | 230.8 | 395.9 | 546.7 |
| 2002..... | 50.7 | * | 0.3 | * | 0.3 | 1.8 | 9.1 | 39.7 | 108.1 | 237.5 | 397.9 | 557.9 |
| 2001..... | 50.5 | * | * | * | 0.3 | 1.7 | 8.8 | 39.7 | 106.9 | 238.4 | 402.1 | 593.2 |
| 2000..... | 50.8 | * | * | * | 0.3 | 1.8 | 8.9 | 39.1 | 109.6 | 239.8 | 413.7 | 614.1 |
| 1999..... | 50.7 | * | 0.3 | * | 0.3 | 1.8 | 8.9 | 38.9 | 110.0 | 243.0 | 411.2 | 620.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 39.8 | * | * | * | 0.3 | 1.3 | 5.6 | 19.5 | 54.2 | 126.4 | 247.7 | 406.4 |
| 2003..... | 40.7 | * | * | * | 0.3 | 1.4 | 5.6 | 20.5 | 54.8 | 131.5 | 252.4 | 423.4 |
| 2002..... | 40.8 | * | * | * | 0.3 | 1.3 | 5.7 | 20.1 | 55.2 | 133.9 | 254.9 | 428.1 |
| 2001..... | 41.7 | * | * | * | 0.2 | 1.3 | 6.0 | 20.9 | 56.0 | 138.5 | 257.7 | 443.9 |
| 2000..... | 42.3 | * | * | * | 0.2 | 1.2 | 5.6 | 20.5 | 58.6 | 138.9 | 265.3 | 460.4 |
| 1999..... | 42.3 | * | * | * | 0.3 | 1.2 | 5.9 | 20.1 | 60.0 | 141.1 | 261.1 | 461.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 18.7 | * | * | * | * | 1.4 | 6.2 | 25.2 | 71.4 | 155.5 | 273.3 | 369.1 |
| 2003..... | 18.8 | * | * | * | * | 1.3 | 5.7 | 27.2 | 73.7 | 159.4 | 280.6 | 377.8 |
| 2002..... | 18.2 | * | * | * | * | 1.1 | 6.2 | 24.6 | 70.3 | 162.4 | 273.7 | 420.2 |
| 2001..... | 18.5 | * | * | * | * | 1.2 | 6.4 | 26.1 | 71.9 | 162.9 | 290.9 | 444.6 |
| 2000..... | 18.2 | * | * | * | * | 1.2 | 6.6 | 25.4 | 69.6 | 166.3 | 301.4 | 436.6 |
| 1999..... | 17.9 | * | * | * | * | 1.4 | 5.9 | 25.9 | 69.4 | 166.6 | 295.0 | 443.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 20.7 | * | * | * | * | 1.5 | 7.5 | 31.9 | 98.0 | 200.3 | 354.4 | 431.5 |
| 2003..... | 20.8 | * | * | * | * | 1.3 | 6.6 | 36.5 | 96.8 | 208.5 | 361.2 | 409.5 |
| 2002..... | 20.0 | * | * | * | * | 1.3 | 6.5 | 32.7 | 94.8 | 217.7 | 339.4 | 480.2 |
| 2001..... | 20.0 | * | * | * | * | 1.1 | 7.3 | 33.7 | 93.5 | 211.9 | 363.5 | 550.4 |
| 2000..... | 19.8 | * | * | * | * | 1.3 | 6.9 | 31.4 | 90.1 | 224.3 | 394.3 | 560.3 |
| 1999..... | 19.3 | * | * | * | * | 1.4 | 6.2 | 33.7 | 90.8 | 214.9 | 377.0 | 528.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 16.5 | * | * | * | * | 1.3 | 4.7 | 18.5 | 47.2 | 119.5 | 217.3 | 337.1 |
| 2003..... | 16.8 | * | * | * | * | 1.3 | 4.7 | 17.9 | 52.9 | 120.3 | 225.2 | 361.7 |
| 2002..... | 16.4 | * | * | * | * | 1.0 | 5.8 | 16.5 | 48.3 | 118.5 | 228.8 | 390.4 |
| 2001..... | 16.9 | * | * | * | * | 1.5 | 5.4 | 18.6 | 52.7 | 124.2 | 241.8 | 392.8 |
| 2000..... | 16.4 | * | * | * | * | 1.1 | 6.3 | 19.5 | 51.5 | 120.6 | 239.0 | 375.9 |
| 1999..... | 16.5 | * | * | * | * | 1.3 | 5.6 | 18.4 | 50.5 | 128.7 | 240.4 | 402.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of digestive organs (C15-C26) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 48.9 | * | 0.2 | * | 0.3 | 1.5 | 7.2 | 29.7 | 78.9 | 171.2 | 304.0 | 448.2 |
| 2003..... | 49.6 | * | * | * | 0.3 | 1.6 | 7.2 | 30.2 | 79.3 | 177.6 | 310.8 | 463.8 |
| 2002..... | 49.8 | * | 0.2 | * | 0.3 | 1.6 | 7.5 | 30.1 | 81.2 | 181.8 | 313.1 | 468.0 |
| 2001..... | 50.0 | * | * | * | 0.3 | 1.6 | 7.5 | 30.4 | 80.9 | 184.6 | 315.5 | 488.5 |
| 2000..... | 50.4 | * | 0.2 | * | 0.2 | 1.6 | 7.3 | 29.9 | 83.8 | 185.1 | 324.1 | 506.3 |
| 1999..... | 50.2 | * | 0.2 | * | 0.3 | 1.5 | 7.5 | 29.5 | 84.8 | 187.6 | 320.4 | 508.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 54.7 | * | * | * | 0.4 | 1.8 | 8.7 | 40.2 | 105.0 | 224.0 | 386.0 | 536.0 |
| 2003..... | 55.2 | * | * | * | 0.3 | 1.8 | 8.8 | 40.1 | 105.6 | 231.8 | 396.9 | 551.8 |
| 2002..... | 55.6 | * | * | * | 0.4 | 1.8 | 9.4 | 40.3 | 108.7 | 238.3 | 399.8 | 559.8 |
| 2001..... | 55.1 | * | * | * | 0.3 | 1.8 | 9.0 | 40.1 | 107.5 | 239.3 | 403.2 | 593.9 |
| 2000..... | 55.3 | * | * | * | 0.3 | 1.9 | 9.1 | 39.7 | 110.8 | 240.0 | 413.6 | 615.2 |
| 1999..... | 55.1 | * | * | * | 0.3 | 1.8 | 9.2 | 39.2 | 111.0 | 243.8 | 411.5 | 622.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 43.3 | * | * | * | 0.3 | 1.3 | 5.7 | 19.5 | 54.7 | 126.7 | 248.9 | 408.9 |
| 2003..... | 44.3 | * | * | * | 0.3 | 1.4 | 5.7 | 20.7 | 54.8 | 132.1 | 253.4 | 425.1 |
| 2002..... | 44.3 | * | * | * | 0.3 | 1.3 | 5.7 | 20.4 | 55.7 | 134.7 | 255.6 | 428.4 |
| 2001..... | 45.1 | * | * | * | 0.3 | 1.3 | 6.0 | 21.1 | 56.1 | 139.2 | 257.9 | 444.6 |
| 2000..... | 45.7 | * | * | * | 0.2 | 1.2 | 5.4 | 20.6 | 58.9 | 139.7 | 265.9 | 462.2 |
| 1999..... | 45.5 | * | * | * | 0.3 | 1.2 | 5.9 | 20.2 | 60.6 | 141.5 | 261.3 | 462.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 51.3 | * | * | * | 0.3 | 1.3 | 6.5 | 27.0 | 73.7 | 163.7 | 297.9 | 442.5 |
| 2003..... | 52.0 | * | * | * | 0.3 | 1.4 | 6.4 | 26.8 | 74.0 | 171.1 | 303.9 | 455.9 |
| 2002..... | 52.3 | * | 0.2 | * | 0.3 | 1.4 | 6.8 | 27.4 | 75.9 | 174.7 | 305.3 | 460.5 |
| 2001..... | 52.3 | * | * | * | 0.2 | 1.4 | 6.7 | 27.6 | 75.6 | 176.9 | 307.1 | 481.5 |
| 2000..... | 52.7 | * | * | * | 0.2 | 1.3 | 6.3 | 27.1 | 78.3 | 177.8 | 315.1 | 497.6 |
| 1999..... | 52.5 | * | 0.2 | * | 0.2 | 1.3 | 6.6 | 26.8 | 78.8 | 180.7 | 311.5 | 500.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 57.3 | * | * | * | 0.3 | 1.6 | 7.9 | 36.9 | 98.0 | 214.1 | 378.8 | 532.5 |
| 2003..... | 57.8 | * | * | * | 0.3 | 1.5 | 7.9 | 35.6 | 98.9 | 224.1 | 389.3 | 543.8 |
| 2002..... | 58.2 | * | * | * | 0.3 | 1.6 | 8.5 | 36.7 | 101.7 | 229.8 | 390.8 | 551.6 |
| 2001..... | 57.6 | * | * | * | 0.3 | 1.6 | 8.0 | 36.4 | 100.8 | 230.1 | 394.6 | 587.7 |
| 2000..... | 57.6 | * | * | * | 0.3 | 1.7 | 8.0 | 35.9 | 103.2 | 231.0 | 403.5 | 605.1 |
| 1999..... | 57.4 | * | * | * | 0.3 | 1.5 | 8.2 | 35.3 | 102.9 | 235.9 | 400.7 | 616.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 45.6 | * | * | * | 0.2 | 1.1 | 5.0 | 17.2 | 50.6 | 120.4 | 242.8 | 402.2 |
| 2003..... | 46.4 | * | * | * | 0.2 | 1.2 | 4.9 | 18.2 | 50.5 | 125.7 | 246.1 | 417.3 |
| 2002..... | 46.6 | * | * | * | 0.3 | 1.3 | 5.1 | 18.2 | 51.5 | 127.7 | 248.0 | 421.1 |
| 2001..... | 47.2 | * | * | * | 0.2 | 1.2 | 5.4 | 18.9 | 51.8 | 131.9 | 249.2 | 437.3 |
| 2000..... | 48.0 | * | * | * | 0.2 | 1.0 | 4.7 | 18.4 | 54.9 | 132.9 | 257.1 | 454.3 |
| 1999..... | 47.8 | * | * | * | 0.2 | 1.0 | 5.0 | 18.4 | 56.2 | 134.5 | 253.2 | 453.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of digestive organs (C15-C26) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 44.2 | * | * | * | 0.6 | 2.3 | 10.6 | 48.0 | 125.6 | 247.3 | 384.2 | 528.2 |
| 2003..... | 45.2 | * | * | * | 0.3 | 2.3 | 11.6 | 53.0 | 127.3 | 243.6 | 401.6 | 569.9 |
| 2002..... | 45.1 | * | * | * | 0.4 | 2.2 | 11.4 | 49.3 | 129.2 | 255.8 | 412.2 | 571.4 |
| 2001..... | 45.6 | * | * | * | * | 2.3 | 11.9 | 51.0 | 128.7 | 260.9 | 421.1 | 590.9 |
| 2000..... | 46.1 | * | * | * | * | 2.4 | 12.1 | 52.4 | 131.8 | 261.4 | 434.5 | 612.8 |
| 1999..... | 46.1 | * | * | * | 0.4 | 2.5 | 12.2 | 50.2 | 136.7 | 263.0 | 433.6 | 617.2 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 49.3 | * | * | * | 0.8 | 2.5 | 12.6 | 64.7 | 172.3 | 337.5 | 495.4 | 578.2 |
| 2003..... | 50.2 | * | * | * | * | 2.6 | 14.0 | 73.7 | 173.9 | 325.4 | 505.9 | 663.8 |
| 2002..... | 50.2 | * | * | * | * | 2.8 | 14.4 | 66.7 | 176.9 | 338.4 | 535.7 | 678.2 |
| 2001..... | 49.7 | * | * | * | * | 2.7 | 14.6 | 68.2 | 172.5 | 345.0 | 534.1 | 686.1 |
| 2000..... | 50.9 | * | * | * | * | 2.8 | 15.3 | 70.8 | 182.1 | 343.5 | 557.5 | 736.2 |
| 1999..... | 51.0 | * | * | * | * | 3.1 | 14.7 | 70.2 | 188.0 | 341.3 | 573.5 | 717.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 39.5 | * | * | * | * | 2.1 | 8.9 | 33.6 | 88.2 | 183.0 | 321.4 | 508.4 |
| 2003..... | 40.6 | * | * | * | * | 1.9 | 9.5 | 35.3 | 89.9 | 185.4 | 342.7 | 533.6 |
| 2002..... | 40.5 | * | * | * | * | 1.6 | 8.8 | 34.4 | 90.9 | 197.3 | 342.4 | 530.5 |
| 2001..... | 41.8 | * | * | * | * | 1.8 | 9.6 | 36.2 | 93.4 | 201.5 | 357.7 | 555.6 |
| 2000..... | 41.7 | * | * | * | * | 2.1 | 9.2 | 36.6 | 91.4 | 203.6 | 365.2 | 566.9 |
| 1999..... | 41.7 | * | * | * | * | 2.0 | 10.0 | 33.2 | 95.6 | 207.7 | 355.2 | 579.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of esophagus (C15) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | 0.1 | 0.7 | 3.4 | 10.3 | 19.2 | 26.7 | 25.8 |
| 2003..... | 4.4 | * | * | * | * | 0.1 | 0.6 | 3.4 | 10.5 | 19.7 | 26.4 | 25.0 |
| 2002..... | 4.4 | * | * | * | * | * | 0.7 | 3.5 | 10.4 | 20.2 | 26.6 | 24.6 |
| 2001..... | 4.4 | * | * | * | * | 0.1 | 0.7 | 3.6 | 10.1 | 20.6 | 26.3 | 25.8 |
| 2000..... | 4.3 | * | * | * | * | 0.1 | 0.7 | 3.6 | 10.8 | 20.0 | 25.6 | 25.9 |
| 1999..... | 4.3 | * | * | * | * | 0.1 | 0.7 | 3.5 | 10.6 | 20.1 | 24.4 | 26.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.0 | * | * | * | * | 0.1 | 1.1 | 5.9 | 17.9 | 33.4 | 48.8 | 48.5 |
| 2003..... | 6.9 | * | * | * | * | 0.1 | 1.1 | 6.0 | 18.5 | 34.1 | 47.8 | 45.3 |
| 2002..... | 6.9 | * | * | * | * | * | 1.1 | 6.0 | 18.2 | 35.6 | 48.3 | 43.4 |
| 2001..... | 6.9 | * | * | * | * | 0.1 | 1.1 | 6.3 | 17.6 | 36.0 | 47.4 | 48.1 |
| 2000..... | 6.7 | * | * | * | * | 0.2 | 1.2 | 6.3 | 18.6 | 34.1 | 45.8 | 47.8 |
| 1999..... | 6.6 | * | * | * | * | 0.1 | 1.1 | 6.1 | 18.3 | 35.3 | 44.4 | 48.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | 0.2 | 1.0 | 3.2 | 7.3 | 11.9 | 15.7 |
| 2003..... | 2.0 | * | * | * | * | * | 0.2 | 1.0 | 3.2 | 7.7 | 12.0 | 16.1 |
| 2002..... | 2.0 | * | * | * | * | * | 0.2 | 1.0 | 3.2 | 7.4 | 12.2 | 16.5 |
| 2001..... | 2.0 | * | * | * | * | * | 0.2 | 1.0 | 3.2 | 7.9 | 12.4 | 16.5 |
| 2000..... | 2.1 | * | * | * | * | * | 0.1 | 1.0 | 3.5 | 8.3 | 12.4 | 17.1 |
| 1999..... | 2.0 | * | * | * | * | * | 0.2 | 0.9 | 3.6 | 7.6 | 11.5 | 18.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 1.0 | 4.7 | 10.3 | 15.0 | 18.6 |
| 2003..... | 1.1 | * | * | * | * | * | * | 1.3 | 5.0 | 10.0 | 14.4 | 16.1 |
| 2002..... | 1.1 | * | * | * | * | * | 0.4 | 1.5 | 4.1 | 10.2 | 15.2 | 23.3 |
| 2001..... | 1.1 | * | * | * | * | * | * | 1.9 | 5.2 | 8.4 | 18.2 | 20.0 |
| 2000..... | 1.1 | * | * | * | * | * | * | 1.2 | 5.0 | 11.1 | 19.4 | 19.9 |
| 1999..... | 1.1 | * | * | * | * | * | * | 1.5 | 5.2 | 9.2 | 18.1 | 26.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 1.9 | 8.6 | 18.2 | 27.2 | 29.4 |
| 2003..... | 1.6 | * | * | * | * | * | * | 2.4 | 8.0 | 17.2 | 28.4 | * |
| 2002..... | 1.6 | * | * | * | * | * | * | 2.7 | 8.0 | 18.3 | 27.1 | 36.8 |
| 2001..... | 1.7 | * | * | * | * | * | * | 3.6 | 8.5 | 16.0 | 30.4 | 41.9 |
| 2000..... | 1.6 | * | * | * | * | * | * | 2.2 | 8.0 | 19.4 | 35.4 | 40.3 |
| 1999..... | 1.7 | * | * | * | * | * | * | 2.7 | 10.2 | 16.3 | 32.8 | 49.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | 4.0 | 6.7 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 2.4 | 4.3 | * | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | 3.6 | 7.0 | 16.6 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 2.2 | * | 10.0 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 2.4 | 4.6 | 8.6 | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 3.6 | 8.3 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of esophagus (C15) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | * | 0.1 | 0.7 | 3.7 | 10.7 | 19.9 | 27.3 | 26.2 |
| 2003..... | 4.9 | * | * | * | * | 0.1 | 0.7 | 3.7 | 10.9 | 20.4 | 27.0 | 25.4 |
| 2002..... | 4.9 | * | * | * | * | * | 0.7 | 3.7 | 10.8 | 20.8 | 27.2 | 24.5 |
| 2001..... | 4.9 | * | * | * | * | 0.1 | 0.7 | 3.7 | 10.4 | 21.3 | 26.6 | 26.0 |
| 2000..... | 4.8 | * | * | * | * | 0.1 | 0.7 | 3.8 | 11.2 | 20.4 | 25.8 | 26.1 |
| 1999..... | 4.7 | * | * | * | * | 0.1 | 0.7 | 3.6 | 11.0 | 20.6 | 24.6 | 26.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 7.9 | * | * | * | * | 0.2 | 1.2 | 6.3 | 18.6 | 34.5 | 49.9 | 49.4 |
| 2003..... | 7.8 | * | * | * | * | 0.1 | 1.2 | 6.4 | 19.3 | 35.3 | 48.8 | 46.3 |
| 2002..... | 7.8 | * | * | * | * | * | 1.2 | 6.3 | 18.9 | 36.6 | 49.3 | 43.5 |
| 2001..... | 7.7 | * | * | * | * | 0.1 | 1.3 | 6.5 | 18.2 | 37.1 | 48.1 | 48.4 |
| 2000..... | 7.5 | * | * | * | * | 0.2 | 1.3 | 6.6 | 19.4 | 34.8 | 46.1 | 47.8 |
| 1999..... | 7.4 | * | * | * | * | 0.2 | 1.2 | 6.4 | 18.8 | 36.2 | 44.7 | 48.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | 0.2 | 1.1 | 3.4 | 7.6 | 12.2 | 15.8 |
| 2003..... | 2.2 | * | * | * | * | * | 0.2 | 1.1 | 3.2 | 7.9 | 12.4 | 16.1 |
| 2002..... | 2.2 | * | * | * | * | * | 0.3 | 1.1 | 3.4 | 7.6 | 12.5 | 16.3 |
| 2001..... | 2.2 | * | * | * | * | * | 0.3 | 1.1 | 3.2 | 8.2 | 12.5 | 16.7 |
| 2000..... | 2.3 | * | * | * | * | * | 0.1 | 1.0 | 3.6 | 8.5 | 12.6 | 17.3 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.9 | 3.8 | 7.8 | 11.6 | 18.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | 0.1 | 0.8 | 3.6 | 10.5 | 19.9 | 27.8 | 26.5 |
| 2003..... | 5.3 | * | * | * | * | 0.1 | 0.7 | 3.4 | 10.6 | 20.3 | 27.3 | 26.1 |
| 2002..... | 5.3 | * | * | * | * | * | 0.7 | 3.6 | 10.4 | 20.5 | 27.3 | 25.1 |
| 2001..... | 5.1 | * | * | * | * | 0.1 | 0.8 | 3.5 | 9.7 | 20.5 | 26.8 | 26.5 |
| 2000..... | 5.1 | * | * | * | * | 0.1 | 0.7 | 3.5 | 10.4 | 20.0 | 25.5 | 26.7 |
| 1999..... | 4.9 | * | * | * | * | 0.1 | 0.7 | 3.3 | 10.2 | 20.0 | 24.4 | 27.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 8.7 | * | * | * | * | * | 1.3 | 6.3 | 18.4 | 34.6 | 51.0 | 50.1 |
| 2003..... | 8.5 | * | * | * | * | * | 1.2 | 6.0 | 18.8 | 35.1 | 49.9 | 47.9 |
| 2002..... | 8.4 | * | * | * | * | * | 1.3 | 6.3 | 18.2 | 36.3 | 49.9 | 44.7 |
| 2001..... | 8.1 | * | * | * | * | * | 1.3 | 6.1 | 17.0 | 36.0 | 48.6 | 49.0 |
| 2000..... | 8.0 | * | * | * | * | 0.2 | 1.4 | 6.3 | 18.3 | 34.3 | 45.9 | 49.4 |
| 1999..... | 7.7 | * | * | * | * | 0.2 | 1.2 | 6.0 | 17.5 | 35.4 | 44.4 | 48.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | 0.2 | 0.9 | 3.0 | 7.2 | 12.0 | 16.0 |
| 2003..... | 2.3 | * | * | * | * | * | 0.2 | 0.9 | 2.8 | 7.7 | 12.0 | 16.5 |
| 2002..... | 2.3 | * | * | * | * | * | 0.2 | 0.9 | 3.0 | 7.1 | 12.2 | 16.6 |
| 2001..... | 2.3 | * | * | * | * | * | 0.2 | 1.0 | 2.7 | 7.5 | 12.4 | 17.1 |
| 2000..... | 2.3 | * | * | * | * | * | * | 0.8 | 3.0 | 8.0 | 12.0 | 17.5 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.8 | 3.3 | 7.1 | 11.3 | 18.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of esophagus (C15) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | 0.7 | 5.3 | 15.8 | 25.2 | 27.4 | 21.9 |
| 2003..... | 4.3 | * | * | * | * | * | 0.9 | 6.1 | 17.3 | 26.1 | 28.4 | 21.4 |
| 2002..... | 4.3 | * | * | * | * | * | 1.0 | 5.2 | 17.7 | 28.9 | 29.9 | 20.6 |
| 2001..... | 4.7 | * | * | * | * | * | 1.0 | 6.1 | 18.9 | 32.8 | 30.0 | 24.0 |
| 2000..... | 4.7 | * | * | * | * | * | 0.8 | 6.8 | 20.2 | 28.6 | 34.1 | 22.6 |
| 1999..... | 4.7 | * | * | * | * | * | 1.0 | 6.6 | 20.3 | 30.2 | 31.4 | 23.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.0 | * | * | * | * | * | 1.0 | 8.5 | 26.0 | 42.2 | 47.6 | 39.6 |
| 2003..... | 6.5 | * | * | * | * | * | 1.4 | 10.4 | 29.4 | 45.2 | 46.7 | 38.4 |
| 2002..... | 6.5 | * | * | * | * | * | 1.5 | 8.4 | 30.7 | 49.8 | 52.2 | 38.3 |
| 2001..... | 7.2 | * | * | * | * | * | 1.6 | 10.5 | 33.0 | 56.2 | 55.5 | 47.4 |
| 2000..... | 7.0 | * | * | * | * | * | 1.2 | 11.5 | 34.8 | 47.6 | 58.9 | 38.7 |
| 1999..... | 7.2 | * | * | * | * | * | 1.7 | 11.5 | 35.3 | 51.1 | 57.2 | 49.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 2.5 | 7.6 | 13.1 | 15.9 | 14.9 |
| 2003..... | 2.4 | * | * | * | * | * | * | 2.5 | 7.5 | 12.5 | 18.1 | 14.9 |
| 2002..... | 2.4 | * | * | * | * | * | * | 2.4 | 7.2 | 14.0 | 17.2 | 13.8 |
| 2001..... | 2.4 | * | * | * | * | * | * | 2.3 | 7.5 | 16.2 | 15.7 | 15.4 |
| 2000..... | 2.6 | * | * | * | * | * | * | 2.7 | 8.5 | 15.2 | 20.1 | 16.6 |
| 1999..... | 2.4 | * | * | * | * | * | * | 2.4 | 8.3 | 15.5 | 17.0 | 13.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of stomach (C16) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | 0.3 | 1.1 | 2.7 | 6.5 | 14.9 | 26.3 | 42.2 |
| 2003..... | 4.2 | * | * | * | * | 0.3 | 1.1 | 3.0 | 6.7 | 15.2 | 27.6 | 43.6 |
| 2002..... | 4.2 | * | * | * | 0.1 | 0.3 | 1.0 | 2.8 | 6.9 | 15.4 | 28.7 | 47.6 |
| 2001..... | 4.3 | * | * | * | 0.1 | 0.3 | 1.0 | 2.9 | 7.0 | 16.5 | 30.2 | 45.4 |
| 2000..... | 4.5 | * | * | * | * | 0.3 | 1.1 | 2.9 | 7.4 | 16.6 | 31.0 | 52.7 |
| 1999..... | 4.6 | * | * | * | * | 0.3 | 1.1 | 3.0 | 7.8 | 17.1 | 31.6 | 50.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.8 | * | * | * | * | 0.3 | 1.2 | 3.5 | 9.1 | 20.6 | 37.2 | 62.9 |
| 2003..... | 4.9 | * | * | * | * | 0.3 | 1.2 | 3.8 | 9.5 | 21.9 | 38.4 | 60.0 |
| 2002..... | 5.0 | * | * | * | * | 0.4 | 1.1 | 3.7 | 9.5 | 22.5 | 40.6 | 67.2 |
| 2001..... | 5.1 | * | * | * | * | 0.4 | 1.1 | 3.8 | 10.2 | 23.2 | 43.1 | 62.6 |
| 2000..... | 5.3 | * | * | * | * | 0.3 | 1.3 | 3.8 | 10.5 | 23.7 | 43.9 | 75.6 |
| 1999..... | 5.5 | * | * | * | * | 0.4 | 1.3 | 4.2 | 11.0 | 24.2 | 46.5 | 71.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.3 | 0.9 | 2.0 | 4.1 | 10.1 | 19.0 | 32.9 |
| 2003..... | 3.4 | * | * | * | * | 0.3 | 1.0 | 2.1 | 4.1 | 9.7 | 20.3 | 36.4 |
| 2002..... | 3.5 | * | * | * | 0.1 | 0.3 | 0.9 | 1.9 | 4.5 | 9.5 | 20.8 | 39.1 |
| 2001..... | 3.5 | * | * | * | * | 0.3 | 0.9 | 2.0 | 4.1 | 10.9 | 21.7 | 38.3 |
| 2000..... | 3.7 | * | * | * | * | 0.3 | 0.9 | 2.1 | 4.6 | 10.8 | 22.6 | 43.4 |
| 1999..... | 3.7 | * | * | * | * | 0.3 | 1.0 | 1.9 | 4.9 | 11.3 | 21.9 | 41.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.5 | 2.0 | 4.6 | 11.7 | 24.4 | 44.0 | 62.8 |
| 2003..... | 3.1 | * | * | * | * | 0.5 | 1.6 | 5.1 | 11.4 | 23.8 | 43.6 | 62.7 |
| 2002..... | 3.1 | * | * | * | * | 0.5 | 1.7 | 4.2 | 11.0 | 21.9 | 47.2 | 72.1 |
| 2001..... | 3.2 | * | * | * | * | 0.6 | 1.8 | 5.4 | 11.3 | 26.4 | 45.6 | 63.8 |
| 2000..... | 3.1 | * | * | * | * | 0.4 | 2.0 | 4.8 | 9.8 | 28.4 | 50.2 | 65.7 |
| 1999..... | 3.1 | * | * | * | * | 0.6 | 1.7 | 4.1 | 11.7 | 26.2 | 49.8 | 73.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | 0.5 | 2.0 | 4.7 | 14.8 | 32.0 | 66.9 | 84.1 |
| 2003..... | 3.4 | * | * | * | * | * | 1.5 | 5.7 | 14.1 | 33.2 | 63.7 | 73.9 |
| 2002..... | 3.4 | * | * | * | * | 0.7 | 1.5 | 4.7 | 13.7 | 32.9 | 61.6 | 107.1 |
| 2001..... | 3.6 | * | * | * | * | * | 1.9 | 6.2 | 15.0 | 37.1 | 60.4 | 95.2 |
| 2000..... | 3.6 | * | * | * | * | * | 2.0 | 5.0 | 13.7 | 43.3 | 70.8 | 98.8 |
| 1999..... | 3.5 | * | * | * | * | 0.7 | 1.6 | 5.2 | 16.4 | 35.6 | 73.5 | 75.5 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | * | 2.1 | 4.5 | 8.8 | 18.3 | 28.2 | 51.9 |
| 2003..... | 2.8 | * | * | * | * | * | 1.6 | 4.5 | 9.0 | 16.3 | 29.8 | 57.0 |
| 2002..... | 2.7 | * | * | * | * | * | 1.9 | 3.8 | 8.7 | 13.2 | 37.4 | 54.7 |
| 2001..... | 2.8 | * | * | * | * | 0.7 | 1.7 | 4.6 | 8.1 | 18.0 | 35.7 | 48.4 |
| 2000..... | 2.7 | * | * | * | * | * | 2.1 | 4.6 | 6.3 | 16.8 | 36.3 | 49.5 |
| 1999..... | 2.7 | * | * | * | * | * | 1.8 | 3.0 | 7.6 | 18.8 | 34.0 | 72.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of stomach (C16) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | 0.3 | 0.9 | 2.5 | 6.0 | 14.2 | 25.3 | 41.2 |
| 2003..... | 4.3 | * | * | * | * | 0.3 | 1.0 | 2.7 | 6.3 | 14.6 | 26.7 | 42.7 |
| 2002..... | 4.4 | * | * | * | 0.1 | 0.3 | 0.9 | 2.6 | 6.5 | 14.9 | 27.7 | 46.3 |
| 2001..... | 4.5 | * | * | * | * | 0.2 | 0.9 | 2.6 | 6.6 | 15.8 | 29.4 | 44.7 |
| 2000..... | 4.7 | * | * | * | * | 0.3 | 0.9 | 2.7 | 7.2 | 15.8 | 30.1 | 52.0 |
| 1999..... | 4.7 | * | * | * | * | 0.3 | 1.0 | 2.9 | 7.5 | 16.5 | 30.8 | 49.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | * | 0.2 | 1.1 | 3.4 | 8.6 | 19.7 | 35.5 | 61.7 |
| 2003..... | 5.2 | * | * | * | * | 0.3 | 1.1 | 3.6 | 9.2 | 21.0 | 36.9 | 59.2 |
| 2002..... | 5.3 | * | * | * | * | 0.3 | 1.0 | 3.6 | 9.1 | 21.7 | 39.4 | 65.0 |
| 2001..... | 5.4 | * | * | * | * | 0.3 | 1.0 | 3.6 | 9.8 | 22.3 | 42.1 | 61.0 |
| 2000..... | 5.5 | * | * | * | * | 0.3 | 1.2 | 3.7 | 10.3 | 22.4 | 42.6 | 74.5 |
| 1999..... | 5.7 | * | * | * | * | 0.3 | 1.2 | 4.0 | 10.6 | 23.4 | 45.3 | 71.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.3 | 0.8 | 1.7 | 3.7 | 9.5 | 18.4 | 32.1 |
| 2003..... | 3.5 | * | * | * | * | 0.3 | 0.9 | 1.9 | 3.7 | 9.2 | 19.8 | 35.5 |
| 2002..... | 3.5 | * | * | * | * | 0.3 | 0.7 | 1.7 | 4.1 | 9.2 | 19.9 | 38.3 |
| 2001..... | 3.6 | * | * | * | * | 0.2 | 0.8 | 1.7 | 3.7 | 10.4 | 21.0 | 37.9 |
| 2000..... | 3.8 | * | * | * | * | 0.3 | 0.7 | 1.8 | 4.4 | 10.4 | 21.9 | 42.9 |
| 1999..... | 3.8 | * | * | * | * | 0.3 | 0.9 | 1.8 | 4.7 | 10.8 | 21.3 | 40.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.9 | 11.8 | 22.3 | 37.6 |
| 2003..... | 4.0 | * | * | * | * | 0.2 | 0.8 | 2.1 | 5.4 | 12.6 | 23.3 | 37.5 |
| 2002..... | 4.1 | * | * | * | * | 0.2 | 0.7 | 2.1 | 5.4 | 12.6 | 24.1 | 40.7 |
| 2001..... | 4.2 | * | * | * | * | 0.2 | 0.7 | 2.1 | 5.6 | 13.5 | 25.8 | 40.0 |
| 2000..... | 4.4 | * | * | * | * | 0.2 | 0.7 | 2.2 | 6.1 | 13.5 | 26.1 | 46.3 |
| 1999..... | 4.4 | * | * | * | * | 0.2 | 0.7 | 2.3 | 6.1 | 14.4 | 27.0 | 43.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | * | 0.2 | 0.9 | 2.7 | 7.1 | 16.3 | 31.1 | 56.9 |
| 2003..... | 4.8 | * | * | * | * | 0.2 | 0.9 | 2.8 | 7.7 | 18.0 | 32.4 | 52.3 |
| 2002..... | 5.0 | * | * | * | * | 0.2 | 0.8 | 2.9 | 7.8 | 18.6 | 35.0 | 58.0 |
| 2001..... | 5.0 | * | * | * | * | 0.2 | 0.8 | 2.8 | 8.2 | 18.8 | 37.4 | 55.7 |
| 2000..... | 5.2 | * | * | * | * | 0.2 | 0.8 | 3.0 | 8.9 | 19.3 | 37.4 | 67.3 |
| 1999..... | 5.4 | * | * | * | * | 0.2 | 0.9 | 3.3 | 8.7 | 20.6 | 39.9 | 62.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.9 | 8.0 | 16.3 | 28.9 |
| 2003..... | 3.3 | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.1 | 8.0 | 17.2 | 31.0 |
| 2002..... | 3.2 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.2 | 7.5 | 16.8 | 33.3 |
| 2001..... | 3.4 | * | * | * | * | * | 0.6 | 1.3 | 3.1 | 9.0 | 18.1 | 33.5 |
| 2000..... | 3.5 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.6 | 8.6 | 18.7 | 37.9 |
| 1999..... | 3.5 | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.7 | 9.1 | 18.6 | 35.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of stomach (C16) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | 0.5 | 1.7 | 5.4 | 12.9 | 30.8 | 50.9 | 71.4 |
| 2003..... | 5.5 | * | * | * | * | 0.4 | 1.8 | 5.8 | 12.9 | 27.5 | 55.0 | 86.0 |
| 2002..... | 5.8 | * | * | * | * | 0.6 | 1.6 | 5.5 | 13.2 | 31.5 | 57.7 | 100.1 |
| 2001..... | 5.8 | * | * | * | * | 0.5 | 2.2 | 5.5 | 13.1 | 31.0 | 59.1 | 89.3 |
| 2000..... | 6.1 | * | * | * | * | 0.7 | 2.1 | 5.8 | 14.1 | 32.3 | 64.3 | 102.7 |
| 1999..... | 6.3 | * | * | * | * | 0.7 | 2.3 | 6.0 | 16.1 | 31.5 | 65.1 | 103.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | * | 1.9 | 7.3 | 19.6 | 48.2 | 75.9 | 99.6 |
| 2003..... | 6.8 | * | * | * | * | * | 2.0 | 8.5 | 20.0 | 44.8 | 80.5 | 107.6 |
| 2002..... | 6.9 | * | * | * | * | 0.8 | 1.9 | 7.9 | 19.5 | 47.3 | 80.9 | 136.7 |
| 2001..... | 6.9 | * | * | * | * | * | 2.4 | 8.0 | 20.0 | 49.2 | 87.9 | 106.3 |
| 2000..... | 7.2 | * | * | * | * | 0.8 | 3.0 | 8.1 | 20.9 | 47.0 | 91.0 | 138.6 |
| 1999..... | 7.6 | * | * | * | * | * | 2.8 | 8.7 | 23.3 | 46.4 | 101.6 | 149.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | * | 1.5 | 3.7 | 7.6 | 18.4 | 36.9 | 60.2 |
| 2003..... | 4.4 | * | * | * | * | * | 1.6 | 3.5 | 7.3 | 15.2 | 40.6 | 77.6 |
| 2002..... | 4.9 | * | * | * | * | * | 1.4 | 3.4 | 8.1 | 20.3 | 44.6 | 86.1 |
| 2001..... | 4.7 | * | * | * | * | * | 2.0 | 3.3 | 7.6 | 18.1 | 43.0 | 83.0 |
| 2000..... | 5.2 | * | * | * | * | * | 1.4 | 3.9 | 8.6 | 22.0 | 49.3 | 89.4 |
| 1999..... | 5.1 | * | * | * | * | * | 1.9 | 3.7 | 10.2 | 21.0 | 44.6 | 85.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of small intestine (C17) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 2.4 | 3.1 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.3 | 2.5 | 3.0 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.4 | 2.8 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 2.5 | 3.2 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 2.5 | 3.0 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.4 | 2.4 | 3.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.0 | 3.1 | 4.4 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.9 | 1.7 | 2.7 | 3.5 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.6 | 2.9 | 3.8 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.9 | 3.0 | 4.1 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.8 | 3.0 | 4.2 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.5 | 0.9 | 1.7 | 3.3 | 4.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 1.9 | 2.5 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.4 | 2.7 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.1 | 2.4 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.1 | 2.8 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.7 | 1.2 | 2.1 | 2.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.4 | 1.2 | 1.9 | 2.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 2.4 | 2.9 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.4 | 2.8 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 2.3 | 2.7 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.4 | 3.0 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.4 | 3.1 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.4 | 2.3 | 3.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.8 | 1.9 | 3.0 | 4.2 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.9 | 1.6 | 2.6 | 3.2 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.7 | 1.6 | 2.8 | 3.6 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.8 | 2.9 | 3.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.9 | 1.8 | 2.8 | 4.3 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.5 | 0.8 | 1.7 | 3.1 | 4.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.9 | 2.4 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.3 | 2.7 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.0 | 2.3 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.1 | 2.7 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.1 | 2.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.4 | 1.2 | 1.7 | 2.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of colon, rectum and anus (C18-C21) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 18.3 | * | * | * | 0.1 | 0.7 | 3.0 | 10.3 | 28.2 | 64.7 | 128.9 | 225.6 |
| 2003..... | 19.2 | * | * | * | 0.1 | 0.7 | 3.0 | 10.9 | 29.8 | 70.5 | 134.7 | 239.7 |
| 2002..... | 19.7 | * | * | * | 0.1 | 0.7 | 3.2 | 11.4 | 31.7 | 73.3 | 137.4 | 241.4 |
| 2001..... | 20.0 | * | * | * | 0.1 | 0.8 | 3.2 | 11.6 | 31.5 | 75.3 | 139.4 | 255.7 |
| 2000..... | 20.4 | * | * | * | 0.1 | 0.7 | 3.0 | 11.2 | 32.7 | 76.7 | 146.0 | 270.4 |
| 1999..... | 20.5 | * | * | * | 0.1 | 0.7 | 3.0 | 11.1 | 33.5 | 77.4 | 146.7 | 271.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 18.6 | * | * | * | 0.2 | 0.7 | 3.2 | 11.9 | 34.2 | 80.6 | 156.9 | 254.7 |
| 2003..... | 19.6 | * | * | * | 0.1 | 0.8 | 3.2 | 12.5 | 36.0 | 88.0 | 164.4 | 273.3 |
| 2002..... | 20.1 | * | * | * | 0.2 | 0.8 | 3.6 | 13.0 | 39.4 | 91.3 | 167.4 | 275.1 |
| 2001..... | 20.2 | * | * | * | 0.1 | 0.8 | 3.5 | 13.1 | 38.3 | 93.7 | 169.5 | 298.3 |
| 2000..... | 20.6 | * | * | * | 0.1 | 0.8 | 3.4 | 12.8 | 39.6 | 95.4 | 179.5 | 315.6 |
| 1999..... | 20.6 | * | * | * | 0.1 | 0.8 | 3.0 | 12.5 | 40.5 | 95.1 | 181.9 | 324.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 18.0 | * | * | * | 0.1 | 0.6 | 2.9 | 8.7 | 22.7 | 51.4 | 110.0 | 212.5 |
| 2003..... | 18.9 | * | * | * | * | 0.7 | 2.7 | 9.3 | 24.0 | 55.9 | 114.8 | 224.8 |
| 2002..... | 19.2 | * | * | * | * | 0.7 | 2.8 | 9.8 | 24.5 | 58.3 | 117.5 | 226.8 |
| 2001..... | 19.8 | * | * | * | 0.1 | 0.7 | 2.9 | 10.2 | 25.3 | 60.1 | 119.6 | 237.8 |
| 2000..... | 20.2 | * | * | * | * | 0.6 | 2.7 | 9.8 | 26.4 | 61.2 | 124.2 | 251.9 |
| 1999..... | 20.3 | * | * | * | 0.1 | 0.6 | 2.9 | 9.7 | 27.0 | 62.9 | 123.8 | 250.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | * | 0.6 | 1.9 | 7.2 | 21.2 | 47.5 | 87.5 | 147.5 |
| 2003..... | 6.2 | * | * | * | * | 0.5 | 2.1 | 8.2 | 23.6 | 49.7 | 95.5 | 144.6 |
| 2002..... | 6.1 | * | * | * | * | 0.4 | 2.1 | 7.7 | 23.0 | 51.2 | 95.6 | 165.7 |
| 2001..... | 6.1 | * | * | * | * | 0.3 | 2.3 | 7.2 | 22.5 | 52.1 | 96.5 | 193.8 |
| 2000..... | 6.0 | * | * | * | * | 0.5 | 2.3 | 8.4 | 23.1 | 49.2 | 98.2 | 181.1 |
| 1999..... | 6.0 | * | * | * | * | 0.5 | 1.7 | 8.3 | 21.0 | 56.1 | 101.0 | 176.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | * | 0.5 | 2.2 | 8.5 | 26.6 | 59.4 | 114.1 | 172.3 |
| 2003..... | 6.6 | * | * | * | * | 0.5 | 2.3 | 10.0 | 29.8 | 64.8 | 120.4 | 172.4 |
| 2002..... | 6.4 | * | * | * | * | * | 2.0 | 9.3 | 31.1 | 67.1 | 121.1 | 174.0 |
| 2001..... | 6.2 | * | * | * | * | * | 2.7 | 7.4 | 28.2 | 66.5 | 116.7 | 257.1 |
| 2000..... | 6.2 | * | * | * | * | * | 2.0 | 9.0 | 28.2 | 63.5 | 131.3 | 235.8 |
| 1999..... | 6.1 | * | * | * | * | * | 1.5 | 9.1 | 24.1 | 72.7 | 134.0 | 211.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | * | 0.6 | 1.4 | 5.8 | 16.3 | 38.0 | 69.1 | 134.7 |
| 2003..... | 5.7 | * | * | * | * | * | 1.8 | 6.4 | 18.0 | 37.6 | 78.5 | 130.5 |
| 2002..... | 5.7 | * | * | * | * | * | 2.1 | 6.1 | 15.7 | 38.6 | 78.1 | 161.6 |
| 2001..... | 5.9 | * | * | * | * | * | 1.9 | 7.1 | 17.4 | 40.8 | 82.8 | 162.9 |
| 2000..... | 5.8 | * | * | * | * | * | 2.7 | 7.8 | 18.5 | 38.0 | 75.9 | 154.3 |
| 1999..... | 5.9 | * | * | * | * | * | 2.0 | 7.5 | 18.3 | 43.0 | 79.1 | 160.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of colon, rectum and anus (C18-C21) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.3 | * | * | * | 0.1 | 0.7 | 3.2 | 10.6 | 28.8 | 65.9 | 130.9 | 228.8 |
| 2003..... | 21.3 | * | * | * | 0.1 | 0.8 | 3.1 | 11.1 | 30.2 | 71.9 | 136.5 | 243.2 |
| 2002..... | 21.7 | * | * | * | 0.1 | 0.8 | 3.3 | 11.8 | 32.3 | 74.6 | 139.1 | 244.0 |
| 2001..... | 22.0 | * | * | * | 0.1 | 0.9 | 3.3 | 12.0 | 32.2 | 76.7 | 141.1 | 257.4 |
| 2000..... | 22.4 | * | * | * | 0.1 | 0.7 | 3.1 | 11.5 | 33.3 | 78.2 | 147.8 | 273.1 |
| 1999..... | 22.4 | * | * | * | 0.1 | 0.7 | 3.1 | 11.3 | 34.3 | 78.5 | 148.1 | 274.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 20.7 | * | * | * | 0.2 | 0.8 | 3.3 | 12.3 | 34.8 | 81.9 | 159.1 | 258.5 |
| 2003..... | 21.7 | * | * | * | * | 0.9 | 3.4 | 12.7 | 36.4 | 89.4 | 166.3 | 277.3 |
| 2002..... | 22.3 | * | * | * | 0.2 | 0.9 | 3.8 | 13.4 | 39.9 | 92.7 | 169.3 | 279.2 |
| 2001..... | 22.3 | * | * | * | 0.1 | 0.9 | 3.6 | 13.6 | 38.9 | 95.2 | 171.8 | 299.6 |
| 2000..... | 22.8 | * | * | * | 0.1 | 0.8 | 3.6 | 13.1 | 40.3 | 97.2 | 181.3 | 318.6 |
| 1999..... | 22.7 | * | * | * | 0.1 | 0.9 | 3.2 | 12.8 | 41.6 | 96.2 | 183.2 | 327.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 19.9 | * | * | * | * | 0.7 | 3.1 | 8.9 | 23.2 | 52.3 | 112.0 | 215.6 |
| 2003..... | 20.9 | * | * | * | * | 0.7 | 2.8 | 9.6 | 24.4 | 57.2 | 116.6 | 228.1 |
| 2002..... | 21.2 | * | * | * | * | 0.7 | 2.9 | 10.2 | 25.2 | 59.6 | 119.1 | 228.8 |
| 2001..... | 21.7 | * | * | * | * | 0.8 | 3.0 | 10.5 | 25.9 | 61.3 | 121.0 | 239.9 |
| 2000..... | 22.1 | * | * | * | * | 0.6 | 2.7 | 9.9 | 26.9 | 62.6 | 126.1 | 254.7 |
| 1999..... | 22.2 | * | * | * | * | 0.6 | 3.0 | 9.9 | 27.6 | 63.9 | 125.3 | 253.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 21.7 | * | * | * | 0.1 | 0.7 | 3.0 | 9.8 | 27.3 | 63.8 | 129.4 | 228.5 |
| 2003..... | 22.7 | * | * | * | * | 0.7 | 2.8 | 10.1 | 28.5 | 69.9 | 135.0 | 242.1 |
| 2002..... | 23.2 | * | * | * | 0.1 | 0.8 | 3.0 | 10.9 | 30.6 | 73.0 | 136.8 | 243.2 |
| 2001..... | 23.5 | * | * | * | 0.1 | 0.8 | 3.1 | 11.1 | 30.6 | 74.7 | 138.5 | 256.9 |
| 2000..... | 24.0 | * | * | * | 0.1 | 0.6 | 2.8 | 10.6 | 31.7 | 76.5 | 145.7 | 271.9 |
| 1999..... | 24.0 | * | * | * | 0.1 | 0.7 | 2.9 | 10.6 | 32.8 | 76.7 | 145.9 | 273.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 22.2 | * | * | * | * | 0.7 | 3.1 | 11.5 | 33.3 | 79.4 | 156.5 | 259.6 |
| 2003..... | 23.2 | * | * | * | * | 0.8 | 3.1 | 11.6 | 34.5 | 87.4 | 164.5 | 276.8 |
| 2002..... | 23.8 | * | * | * | 0.2 | 0.9 | 3.4 | 12.5 | 38.0 | 91.1 | 165.8 | 277.6 |
| 2001..... | 23.9 | * | * | * | * | 0.9 | 3.3 | 12.8 | 37.4 | 93.2 | 169.3 | 298.2 |
| 2000..... | 24.2 | * | * | * | * | 0.8 | 3.2 | 12.2 | 38.2 | 95.1 | 178.4 | 315.8 |
| 1999..... | 24.3 | * | * | * | * | 0.8 | 3.1 | 12.1 | 39.7 | 94.8 | 180.6 | 328.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 21.3 | * | * | * | * | 0.6 | 2.8 | 8.1 | 21.7 | 50.3 | 110.9 | 214.6 |
| 2003..... | 22.2 | * | * | * | * | 0.7 | 2.5 | 8.5 | 22.8 | 54.9 | 115.0 | 226.8 |
| 2002..... | 22.7 | * | * | * | * | 0.7 | 2.7 | 9.4 | 23.7 | 57.6 | 117.3 | 228.3 |
| 2001..... | 23.1 | * | * | * | * | 0.7 | 2.8 | 9.5 | 24.2 | 59.0 | 118.1 | 239.8 |
| 2000..... | 23.8 | * | * | * | * | 0.5 | 2.4 | 9.0 | 25.5 | 60.9 | 124.1 | 254.1 |
| 1999..... | 23.7 | * | * | * | * | 0.6 | 2.7 | 9.0 | 26.2 | 61.5 | 123.2 | 251.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of colon, rectum and anus (C18-C21) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 17.8 | * | * | * | * | 1.0 | 4.6 | 16.9 | 45.8 | 96.4 | 167.6 | 262.5 |
| 2003..... | 18.8 | * | * | * | * | 1.0 | 5.2 | 19.2 | 48.9 | 102.2 | 175.0 | 282.9 |
| 2002..... | 18.9 | * | * | * | * | 0.7 | 5.1 | 19.0 | 50.5 | 100.6 | 185.9 | 278.3 |
| 2001..... | 19.2 | * | * | * | * | 1.1 | 5.0 | 19.5 | 50.6 | 105.9 | 189.7 | 289.0 |
| 2000..... | 19.4 | * | * | * | * | 1.0 | 5.2 | 20.0 | 51.1 | 106.3 | 192.5 | 312.9 |
| 1999..... | 19.3 | * | * | * | * | 1.1 | 4.9 | 18.1 | 53.7 | 107.2 | 194.5 | 308.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 18.0 | * | * | * | * | 1.1 | 4.7 | 19.5 | 54.7 | 122.8 | 218.8 | 277.4 |
| 2003..... | 19.0 | * | * | * | * | 1.0 | 5.4 | 22.4 | 59.9 | 128.2 | 217.0 | 321.8 |
| 2002..... | 19.3 | * | * | * | * | 0.9 | 6.0 | 21.8 | 62.7 | 123.0 | 240.9 | 332.6 |
| 2001..... | 19.0 | * | * | * | * | 1.2 | 5.8 | 21.4 | 59.3 | 130.7 | 229.9 | 352.3 |
| 2000..... | 19.8 | * | * | * | * | 1.1 | 6.1 | 22.7 | 63.7 | 133.7 | 244.4 | 381.6 |
| 1999..... | 19.2 | * | * | * | * | 1.2 | 4.8 | 20.6 | 67.0 | 127.8 | 245.8 | 351.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 17.6 | * | * | * | * | 1.0 | 4.4 | 14.8 | 38.7 | 77.6 | 138.8 | 256.6 |
| 2003..... | 18.7 | * | * | * | * | 0.9 | 5.0 | 16.4 | 40.1 | 83.8 | 151.2 | 267.8 |
| 2002..... | 18.4 | * | * | * | * | * | 4.3 | 16.7 | 40.7 | 84.7 | 154.9 | 257.5 |
| 2001..... | 19.4 | * | * | * | * | 1.0 | 4.3 | 17.9 | 43.6 | 88.4 | 167.1 | 265.6 |
| 2000..... | 19.1 | * | * | * | * | 1.0 | 4.4 | 17.6 | 40.9 | 87.1 | 163.2 | 287.4 |
| 1999..... | 19.5 | * | * | * | * | 0.9 | 5.0 | 15.9 | 43.1 | 92.7 | 165.8 | 292.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Colon (C18) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.2 | * | * | * | 0.1 | 0.5 | 2.1 | 7.8 | 22.3 | 53.8 | 109.8 | 195.3 |
| 2003..... | 16.1 | * | * | * | 0.1 | 0.6 | 2.2 | 8.5 | 23.6 | 59.1 | 115.7 | 208.4 |
| 2002..... | 16.5 | * | * | * | 0.1 | 0.5 | 2.4 | 8.9 | 25.3 | 61.0 | 118.5 | 211.3 |
| 2001..... | 16.8 | * | * | * | 0.1 | 0.6 | 2.4 | 8.9 | 25.7 | 63.5 | 120.1 | 221.6 |
| 2000..... | 17.3 | * | * | * | 0.1 | 0.5 | 2.3 | 8.9 | 26.5 | 64.3 | 126.5 | 236.6 |
| 1999..... | 17.4 | * | * | * | 0.1 | 0.5 | 2.3 | 8.8 | 27.4 | 65.1 | 127.0 | 238.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 15.1 | * | * | * | 0.1 | 0.5 | 2.2 | 9.1 | 26.3 | 65.7 | 131.7 | 219.9 |
| 2003..... | 16.1 | * | * | * | 0.1 | 0.6 | 2.4 | 9.6 | 28.0 | 72.9 | 140.1 | 236.1 |
| 2002..... | 16.6 | * | * | * | 0.1 | 0.6 | 2.6 | 9.9 | 30.9 | 75.5 | 142.6 | 240.2 |
| 2001..... | 16.7 | * | * | * | * | 0.6 | 2.6 | 9.9 | 30.6 | 77.7 | 144.4 | 256.7 |
| 2000..... | 17.1 | * | * | * | * | 0.6 | 2.6 | 9.9 | 31.4 | 78.6 | 154.0 | 275.8 |
| 1999..... | 17.2 | * | * | * | 0.1 | 0.6 | 2.3 | 9.8 | 32.7 | 79.2 | 156.1 | 282.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 15.2 | * | * | * | * | 0.5 | 2.1 | 6.6 | 18.5 | 43.8 | 95.1 | 184.2 |
| 2003..... | 16.1 | * | * | * | * | 0.5 | 2.1 | 7.4 | 19.6 | 47.5 | 99.4 | 196.1 |
| 2002..... | 16.4 | * | * | * | * | 0.5 | 2.1 | 7.9 | 20.2 | 49.0 | 102.6 | 198.7 |
| 2001..... | 16.9 | * | * | * | 0.1 | 0.6 | 2.3 | 8.0 | 21.2 | 51.8 | 104.2 | 206.9 |
| 2000..... | 17.4 | * | * | * | * | 0.5 | 2.1 | 7.9 | 22.0 | 52.5 | 108.5 | 220.6 |
| 1999..... | 17.5 | * | * | * | * | 0.5 | 2.3 | 7.9 | 22.5 | 53.5 | 108.2 | 220.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | 0.4 | 1.3 | 5.9 | 16.5 | 40.0 | 77.4 | 129.9 |
| 2003..... | 5.1 | * | * | * | * | 0.3 | 1.4 | 6.3 | 18.4 | 41.8 | 82.4 | 122.8 |
| 2002..... | 5.1 | * | * | * | * | * | 1.7 | 6.0 | 18.1 | 43.1 | 84.4 | 145.8 |
| 2001..... | 5.0 | * | * | * | * | * | 1.7 | 5.4 | 18.4 | 43.4 | 81.7 | 172.6 |
| 2000..... | 5.0 | * | * | * | * | 0.4 | 1.8 | 6.7 | 18.1 | 42.0 | 83.6 | 160.6 |
| 1999..... | 5.0 | * | * | * | * | 0.4 | 1.4 | 6.9 | 16.6 | 46.8 | 88.3 | 154.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | * | * | 1.5 | 6.9 | 20.0 | 50.4 | 98.7 | 152.7 |
| 2003..... | 5.3 | * | * | * | * | * | 1.5 | 7.5 | 22.6 | 53.3 | 104.3 | 141.6 |
| 2002..... | 5.3 | * | * | * | * | * | 1.6 | 7.1 | 23.3 | 56.6 | 108.3 | 153.9 |
| 2001..... | 5.0 | * | * | * | * | * | 1.8 | 5.2 | 22.2 | 54.9 | 96.6 | 220.9 |
| 2000..... | 5.1 | * | * | * | * | * | 1.6 | 7.2 | 20.9 | 52.3 | 116.0 | 211.6 |
| 1999..... | 5.0 | * | * | * | * | * | 1.2 | 7.4 | 19.4 | 58.9 | 118.4 | 185.5 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | * | * | 1.2 | 4.9 | 13.3 | 31.6 | 62.7 | 118.1 |
| 2003..... | 4.8 | * | * | * | * | * | 1.3 | 5.1 | 14.7 | 32.6 | 67.4 | 113.3 |
| 2002..... | 4.9 | * | * | * | * | * | 1.8 | 4.9 | 13.5 | 32.4 | 68.0 | 141.7 |
| 2001..... | 5.1 | * | * | * | * | * | 1.6 | 5.7 | 14.9 | 34.3 | 71.6 | 148.9 |
| 2000..... | 4.8 | * | * | * | * | * | 2.1 | 6.2 | 15.5 | 33.8 | 61.7 | 135.5 |
| 1999..... | 5.0 | * | * | * | * | * | 1.7 | 6.5 | 14.2 | 37.3 | 68.3 | 139.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Colon (C18) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 16.8 | * | * | * | 0.1 | 0.5 | 2.2 | 8.0 | 22.7 | 54.7 | 111.5 | 198.0 |
| 2003..... | 17.8 | * | * | * | 0.1 | 0.6 | 2.4 | 8.7 | 24.0 | 60.2 | 117.2 | 211.5 |
| 2002..... | 18.2 | * | * | * | 0.1 | 0.6 | 2.5 | 9.2 | 25.8 | 62.1 | 119.9 | 213.5 |
| 2001..... | 18.5 | * | * | * | 0.1 | 0.7 | 2.5 | 9.2 | 26.2 | 64.7 | 121.7 | 222.9 |
| 2000..... | 19.0 | * | * | * | 0.1 | 0.6 | 2.4 | 9.1 | 27.1 | 65.6 | 128.1 | 238.9 |
| 1999..... | 19.0 | * | * | * | 0.1 | 0.6 | 2.4 | 9.0 | 28.1 | 66.0 | 128.2 | 240.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 16.9 | * | * | * | 0.1 | 0.5 | 2.3 | 9.3 | 26.8 | 66.7 | 133.4 | 223.0 |
| 2003..... | 17.9 | * | * | * | * | 0.7 | 2.5 | 9.8 | 28.4 | 74.1 | 141.7 | 239.9 |
| 2002..... | 18.4 | * | * | * | 0.2 | 0.6 | 2.8 | 10.2 | 31.4 | 76.6 | 144.0 | 243.7 |
| 2001..... | 18.5 | * | * | * | * | 0.7 | 2.7 | 10.3 | 31.1 | 78.9 | 146.5 | 257.8 |
| 2000..... | 18.9 | * | * | * | * | 0.6 | 2.7 | 10.1 | 32.1 | 80.1 | 155.3 | 278.2 |
| 1999..... | 19.0 | * | * | * | * | 0.6 | 2.5 | 10.0 | 33.5 | 80.2 | 157.1 | 285.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 16.8 | * | * | * | * | 0.5 | 2.2 | 6.8 | 18.9 | 44.6 | 96.8 | 186.8 |
| 2003..... | 17.8 | * | * | * | * | 0.6 | 2.2 | 7.6 | 19.9 | 48.6 | 100.9 | 199.0 |
| 2002..... | 18.1 | * | * | * | * | 0.5 | 2.2 | 8.2 | 20.7 | 50.1 | 104.0 | 200.5 |
| 2001..... | 18.6 | * | * | * | * | 0.6 | 2.4 | 8.2 | 21.6 | 52.9 | 105.4 | 208.4 |
| 2000..... | 19.1 | * | * | * | * | 0.5 | 2.1 | 8.0 | 22.3 | 53.6 | 110.4 | 223.0 |
| 1999..... | 19.0 | * | * | * | * | 0.5 | 2.3 | 8.0 | 23.0 | 54.3 | 109.5 | 222.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 17.9 | * | * | * | * | 0.5 | 2.1 | 7.3 | 21.3 | 52.5 | 109.6 | 197.7 |
| 2003..... | 19.0 | * | * | * | * | 0.6 | 2.1 | 7.7 | 22.5 | 58.3 | 115.8 | 210.2 |
| 2002..... | 19.4 | * | * | * | 0.1 | 0.6 | 2.3 | 8.4 | 24.3 | 60.4 | 117.7 | 212.6 |
| 2001..... | 19.7 | * | * | * | 0.1 | 0.6 | 2.3 | 8.4 | 24.6 | 62.6 | 119.0 | 222.4 |
| 2000..... | 20.2 | * | * | * | * | 0.5 | 2.1 | 8.2 | 25.5 | 64.0 | 125.8 | 237.3 |
| 1999..... | 20.3 | * | * | * | 0.1 | 0.5 | 2.2 | 8.3 | 26.6 | 64.3 | 126.0 | 239.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 17.9 | * | * | * | * | 0.5 | 2.2 | 8.6 | 25.4 | 64.1 | 130.5 | 224.1 |
| 2003..... | 19.1 | * | * | * | * | 0.6 | 2.3 | 8.9 | 26.7 | 72.1 | 139.8 | 240.0 |
| 2002..... | 19.5 | * | * | * | * | 0.7 | 2.6 | 9.4 | 29.6 | 74.9 | 140.8 | 241.9 |
| 2001..... | 19.7 | * | * | * | * | 0.7 | 2.5 | 9.5 | 29.7 | 77.0 | 143.6 | 256.3 |
| 2000..... | 20.1 | * | * | * | * | 0.6 | 2.4 | 9.4 | 30.3 | 78.4 | 152.3 | 274.5 |
| 1999..... | 20.3 | * | * | * | * | 0.5 | 2.4 | 9.5 | 31.9 | 79.0 | 154.6 | 285.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 17.9 | * | * | * | * | 0.5 | 2.0 | 6.1 | 17.5 | 42.4 | 95.4 | 185.8 |
| 2003..... | 18.9 | * | * | * | * | 0.5 | 1.9 | 6.6 | 18.5 | 46.4 | 99.6 | 197.0 |
| 2002..... | 19.3 | * | * | * | * | 0.5 | 2.0 | 7.4 | 19.3 | 48.1 | 102.3 | 200.0 |
| 2001..... | 19.7 | * | * | * | * | 0.5 | 2.1 | 7.4 | 19.8 | 50.5 | 102.7 | 208.3 |
| 2000..... | 20.4 | * | * | * | * | 0.4 | 1.8 | 7.1 | 21.1 | 51.9 | 108.4 | 222.2 |
| 1999..... | 20.2 | * | * | * | * | 0.5 | 2.0 | 7.1 | 21.6 | 52.0 | 107.4 | 220.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Colon (C18) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 15.2 | * | * | * | * | 0.6 | 3.2 | 13.5 | 38.7 | 84.5 | 149.1 | 228.5 |
| 2003..... | 16.1 | * | * | * | * | 0.7 | 4.1 | 15.7 | 40.8 | 88.9 | 153.9 | 250.2 |
| 2002..... | 16.2 | * | * | * | * | 0.6 | 3.9 | 15.9 | 42.5 | 87.3 | 163.8 | 247.4 |
| 2001..... | 16.7 | * | * | * | * | 0.9 | 4.1 | 16.1 | 43.5 | 93.3 | 167.8 | 252.2 |
| 2000..... | 16.8 | * | * | * | * | 0.8 | 4.2 | 16.7 | 43.3 | 91.1 | 173.2 | 282.4 |
| 1999..... | 16.8 | * | * | * | * | 0.9 | 4.0 | 15.3 | 45.9 | 93.0 | 173.0 | 276.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 15.1 | * | * | * | * | * | 2.8 | 15.4 | 45.5 | 105.2 | 191.4 | 241.8 |
| 2003..... | 16.0 | * | * | * | * | * | 4.1 | 17.7 | 49.2 | 110.4 | 191.6 | 273.8 |
| 2002..... | 16.4 | * | * | * | * | * | 4.5 | 17.5 | 52.2 | 106.1 | 209.8 | 296.5 |
| 2001..... | 16.2 | * | * | * | * | 0.9 | 4.3 | 17.7 | 49.0 | 111.5 | 204.0 | 311.8 |
| 2000..... | 16.8 | * | * | * | * | * | 4.7 | 18.4 | 52.8 | 110.3 | 218.8 | 351.1 |
| 1999..... | 16.3 | * | * | * | * | 0.9 | 3.7 | 16.7 | 55.6 | 109.6 | 215.8 | 306.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 15.3 | * | * | * | * | * | 3.5 | 11.8 | 33.2 | 69.8 | 125.3 | 223.3 |
| 2003..... | 16.3 | * | * | * | * | * | 4.1 | 14.0 | 34.1 | 73.6 | 132.6 | 241.0 |
| 2002..... | 16.1 | * | * | * | * | * | 3.5 | 14.6 | 34.7 | 73.9 | 137.8 | 228.7 |
| 2001..... | 17.1 | * | * | * | * | 0.9 | 3.9 | 14.7 | 39.1 | 80.5 | 147.4 | 230.1 |
| 2000..... | 16.9 | * | * | * | * | 0.8 | 3.7 | 15.2 | 35.7 | 77.5 | 147.6 | 256.9 |
| 1999..... | 17.3 | * | * | * | * | 0.9 | 4.3 | 14.0 | 38.0 | 81.3 | 149.0 | 265.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rectosigmoid junction and rectum (C19-C20) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 0.2 | 0.8 | 2.1 | 5.5 | 10.3 | 18.1 | 29.1 |
| 2003..... | 2.9 | * | * | * | * | 0.1 | 0.6 | 2.2 | 5.7 | 10.9 | 18.0 | 29.9 |
| 2002..... | 3.0 | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.9 | 11.7 | 18.0 | 29.0 |
| 2001..... | 3.0 | * | * | * | * | 0.2 | 0.7 | 2.5 | 5.5 | 11.2 | 18.3 | 32.6 |
| 2000..... | 3.0 | * | * | * | * | 0.1 | 0.6 | 2.1 | 5.8 | 11.8 | 18.8 | 32.6 |
| 1999..... | 3.0 | * | * | * | * | 0.2 | 0.6 | 2.0 | 5.8 | 11.7 | 18.8 | 32.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.2 | 0.9 | 2.6 | 7.4 | 14.3 | 24.5 | 33.5 |
| 2003..... | 3.3 | * | * | * | * | 0.2 | 0.7 | 2.7 | 7.6 | 14.6 | 23.7 | 36.4 |
| 2002..... | 3.4 | * | * | * | * | 0.2 | 0.8 | 2.9 | 8.1 | 15.3 | 24.2 | 33.6 |
| 2001..... | 3.4 | * | * | * | * | 0.2 | 0.8 | 3.0 | 7.4 | 15.5 | 24.4 | 40.7 |
| 2000..... | 3.4 | * | * | * | * | 0.2 | 0.7 | 2.6 | 7.8 | 16.3 | 24.8 | 38.9 |
| 1999..... | 3.3 | * | * | * | * | 0.2 | 0.6 | 2.5 | 7.6 | 15.4 | 25.1 | 40.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | 0.2 | 0.7 | 1.7 | 3.8 | 7.0 | 13.7 | 27.2 |
| 2003..... | 2.6 | * | * | * | * | 0.1 | 0.5 | 1.6 | 3.9 | 7.8 | 14.2 | 27.0 |
| 2002..... | 2.6 | * | * | * | * | 0.2 | 0.5 | 1.7 | 3.8 | 8.7 | 13.9 | 27.0 |
| 2001..... | 2.6 | * | * | * | * | 0.1 | 0.5 | 1.9 | 3.7 | 7.6 | 14.4 | 29.2 |
| 2000..... | 2.6 | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.0 | 8.1 | 14.8 | 30.1 |
| 1999..... | 2.7 | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.1 | 8.7 | 14.7 | 28.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.5 | 1.2 | 4.6 | 7.3 | 9.6 | 16.6 |
| 2003..... | 1.1 | * | * | * | * | * | 0.6 | 1.8 | 4.8 | 7.6 | 12.6 | 20.2 |
| 2002..... | 0.9 | * | * | * | * | * | * | 1.6 | 4.5 | 7.8 | 10.5 | 18.3 |
| 2001..... | 1.0 | * | * | * | * | * | 0.6 | 1.8 | 4.0 | 8.1 | 14.1 | 21.3 |
| 2000..... | 1.0 | * | * | * | * | * | 0.4 | 1.7 | 4.7 | 6.9 | 14.0 | 19.9 |
| 1999..... | 0.9 | * | * | * | * | * | * | 1.2 | 4.3 | 9.1 | 12.5 | 21.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.6 | 1.6 | 6.3 | 8.8 | 15.4 | * |
| 2003..... | 1.3 | * | * | * | * | * | 0.8 | 2.4 | 6.9 | 11.4 | 15.7 | * |
| 2002..... | 1.1 | * | * | * | * | * | * | 2.0 | 7.2 | 10.1 | 12.3 | * |
| 2001..... | 1.2 | * | * | * | * | * | 0.8 | 2.2 | 6.0 | 11.2 | 20.1 | * |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.8 | 6.8 | 10.6 | 14.8 | * |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.7 | 4.7 | 13.8 | 15.6 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 3.0 | 6.1 | 5.7 | 15.1 |
| 2003..... | 0.9 | * | * | * | * | * | * | 1.3 | 3.0 | 4.6 | 10.5 | 15.6 |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.0 | 5.9 | 9.3 | 17.4 |
| 2001..... | 0.8 | * | * | * | * | * | * | 1.3 | 2.2 | 5.7 | 10.0 | * |
| 2000..... | 0.9 | * | * | * | * | * | * | 1.5 | 2.9 | 4.0 | 13.5 | * |
| 1999..... | 0.8 | * | * | * | * | * | * | * | 4.0 | 5.3 | 10.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rectosigmoid junction and rectum (C19-C20) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | 0.2 | 0.8 | 2.2 | 5.6 | 10.5 | 18.5 | 29.7 |
| 2003..... | 3.2 | * | * | * | * | 0.2 | 0.6 | 2.2 | 5.7 | 11.1 | 18.3 | 30.3 |
| 2002..... | 3.3 | * | * | * | * | 0.2 | 0.7 | 2.3 | 6.0 | 11.9 | 18.3 | 29.3 |
| 2001..... | 3.3 | * | * | * | * | 0.2 | 0.7 | 2.5 | 5.6 | 11.4 | 18.5 | 33.0 |
| 2000..... | 3.3 | * | * | * | * | 0.1 | 0.6 | 2.2 | 5.9 | 12.1 | 18.9 | 33.0 |
| 1999..... | 3.2 | * | * | * | * | 0.2 | 0.6 | 2.1 | 5.9 | 11.9 | 19.0 | 32.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | 0.3 | 0.9 | 2.7 | 7.5 | 14.6 | 24.9 | 34.2 |
| 2003..... | 3.6 | * | * | * | * | 0.2 | 0.7 | 2.8 | 7.6 | 14.8 | 24.1 | 36.6 |
| 2002..... | 3.8 | * | * | * | * | 0.3 | 0.9 | 2.9 | 8.2 | 15.6 | 24.7 | 34.2 |
| 2001..... | 3.7 | * | * | * | * | 0.2 | 0.8 | 3.1 | 7.5 | 15.7 | 24.6 | 40.9 |
| 2000..... | 3.7 | * | * | * | * | 0.2 | 0.8 | 2.7 | 7.8 | 16.6 | 25.1 | 39.3 |
| 1999..... | 3.6 | * | * | * | * | 0.2 | 0.7 | 2.6 | 7.8 | 15.5 | 25.4 | 41.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 0.1 | 0.8 | 1.8 | 3.8 | 7.1 | 14.1 | 27.6 |
| 2003..... | 2.8 | * | * | * | * | 0.1 | 0.5 | 1.7 | 3.9 | 8.0 | 14.4 | 27.5 |
| 2002..... | 2.9 | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.0 | 8.8 | 14.1 | 27.2 |
| 2001..... | 2.9 | * | * | * | * | 0.2 | 0.5 | 2.0 | 3.8 | 7.8 | 14.5 | 29.7 |
| 2000..... | 2.9 | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.1 | 8.3 | 14.8 | 30.5 |
| 1999..... | 2.9 | * | * | * | * | * | 0.6 | 1.7 | 4.1 | 8.9 | 14.8 | 29.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.5 | 10.7 | 18.7 | 29.7 |
| 2003..... | 3.5 | * | * | * | * | 0.1 | 0.6 | 2.1 | 5.6 | 11.0 | 18.3 | 30.5 |
| 2002..... | 3.6 | * | * | * | * | 0.2 | 0.7 | 2.3 | 5.9 | 12.0 | 18.1 | 29.3 |
| 2001..... | 3.6 | * | * | * | * | 0.2 | 0.7 | 2.5 | 5.6 | 11.4 | 18.5 | 33.0 |
| 2000..... | 3.5 | * | * | * | * | 0.1 | 0.6 | 2.1 | 5.7 | 11.9 | 19.0 | 33.5 |
| 1999..... | 3.5 | * | * | * | * | 0.2 | 0.6 | 2.1 | 5.8 | 11.9 | 19.0 | 32.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | 0.2 | 0.8 | 2.7 | 7.4 | 14.8 | 25.2 | 34.3 |
| 2003..... | 3.9 | * | * | * | * | 0.2 | 0.7 | 2.6 | 7.5 | 14.8 | 24.2 | 36.1 |
| 2002..... | 4.0 | * | * | * | * | 0.2 | 0.8 | 2.8 | 8.0 | 15.8 | 24.4 | 34.3 |
| 2001..... | 4.0 | * | * | * | * | 0.2 | 0.8 | 3.1 | 7.3 | 15.6 | 25.0 | 41.1 |
| 2000..... | 4.0 | * | * | * | * | * | 0.7 | 2.6 | 7.6 | 16.2 | 25.4 | 40.3 |
| 1999..... | 3.9 | * | * | * | * | 0.2 | 0.6 | 2.4 | 7.6 | 15.4 | 25.3 | 41.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 0.7 | 1.7 | 3.7 | 7.2 | 14.4 | 27.6 |
| 2003..... | 3.0 | * | * | * | * | * | 0.5 | 1.6 | 3.8 | 7.8 | 14.2 | 28.0 |
| 2002..... | 3.1 | * | * | * | * | 0.2 | 0.5 | 1.7 | 3.8 | 8.8 | 13.9 | 27.2 |
| 2001..... | 3.1 | * | * | * | * | 0.2 | 0.6 | 1.8 | 3.9 | 7.9 | 14.3 | 29.6 |
| 2000..... | 3.1 | * | * | * | * | * | 0.4 | 1.6 | 4.0 | 8.3 | 14.9 | 30.7 |
| 1999..... | 3.2 | * | * | * | * | * | 0.6 | 1.7 | 4.1 | 8.9 | 14.9 | 29.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rectosigmoid junction and rectum (C19-C20) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 1.2 | 3.1 | 6.8 | 11.3 | 17.9 | 32.2 |
| 2003..... | 2.5 | * | * | * | * | * | 1.0 | 3.2 | 7.6 | 12.5 | 19.9 | 32.1 |
| 2002..... | 2.5 | * | * | * | * | * | 1.0 | 2.9 | 7.6 | 12.8 | 21.5 | 30.2 |
| 2001..... | 2.4 | * | * | * | * | * | 0.8 | 3.1 | 6.8 | 12.1 | 20.9 | 35.9 |
| 2000..... | 2.5 | * | * | * | * | * | 0.9 | 3.0 | 7.4 | 14.7 | 18.3 | 28.9 |
| 1999..... | 2.4 | * | * | * | * | * | 0.7 | 2.5 | 7.7 | 13.4 | 20.8 | 32.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | 1.6 | 3.7 | 8.9 | 16.8 | 27.1 | 32.5 |
| 2003..... | 2.9 | * | * | * | * | * | 1.1 | 4.3 | 10.5 | 16.9 | 24.5 | 46.9 |
| 2002..... | 2.8 | * | * | * | * | * | 1.2 | 4.0 | 10.2 | 16.4 | 30.2 | 35.0 |
| 2001..... | 2.8 | * | * | * | * | * | 1.3 | 3.5 | 9.9 | 18.7 | 24.9 | 39.3 |
| 2000..... | 2.9 | * | * | * | * | * | 1.3 | 3.9 | 10.5 | 22.8 | 24.4 | 29.4 |
| 1999..... | 2.8 | * | * | * | * | * | 0.8 | 3.5 | 11.1 | 17.7 | 29.7 | 45.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | 0.9 | 2.6 | 5.2 | 7.4 | 12.7 | 32.1 |
| 2003..... | 2.2 | * | * | * | * | * | 0.9 | 2.2 | 5.3 | 9.4 | 17.3 | 26.4 |
| 2002..... | 2.2 | * | * | * | * | * | 0.7 | 2.0 | 5.5 | 10.2 | 16.6 | 28.4 |
| 2001..... | 2.1 | * | * | * | * | * | * | 2.8 | 4.3 | 7.5 | 18.7 | 34.6 |
| 2000..... | 2.0 | * | * | * | * | * | * | 2.2 | 4.9 | 9.1 | 14.9 | 28.8 |
| 1999..... | 2.0 | * | * | * | * | * | * | 1.6 | 4.9 | 10.3 | 15.8 | 27.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Anus and anus canal (C21) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.0 | 1.2 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.0 | 1.4 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.9 | 1.2 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.9 | 1.5 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.8 | 1.2 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.8 | 1.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.6 | 0.7 | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.5 | * |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.3 | 0.5 | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.7 | * |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 0.8 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.7 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.4 | 0.4 | 0.7 | 1.2 | 1.1 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.3 | 1.6 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.0 | 1.1 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 1.1 | 1.7 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 0.9 | 1.3 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.0 | 1.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.0 | 1.1 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.9 | 1.4 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.9 | 1.2 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.9 | 1.5 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.8 | 1.2 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.9 | 1.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 0.8 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.4 | 0.5 | 0.5 | * |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.3 | 0.3 | 0.5 | 0.6 | * |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.7 | * |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.8 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.4 | 0.5 | 0.7 | 1.2 | 1.1 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.4 | 0.5 | 0.6 | 1.2 | 1.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.1 | 1.2 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.1 | 1.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 0.7 | 0.9 | 1.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.0 | 1.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of liver and intrahepatic bile ducts (C22) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.2 | * | 0.1 | * | 0.1 | 0.2 | 0.9 | 5.6 | 10.7 | 19.9 | 31.3 | 32.4 |
| 2003..... | 5.1 | * | 0.1 | * | 0.1 | 0.2 | 0.9 | 5.4 | 9.9 | 19.8 | 30.7 | 33.4 |
| 2002..... | 4.9 | * | 0.2 | * | 0.1 | 0.3 | 0.9 | 4.9 | 9.6 | 19.6 | 29.3 | 35.0 |
| 2001..... | 4.7 | * | * | * | 0.1 | 0.2 | 0.9 | 4.8 | 9.3 | 18.8 | 29.0 | 34.1 |
| 2000..... | 4.6 | * | 0.1 | 0.0 | 0.1 | 0.2 | 1.0 | 4.5 | 9.3 | 18.7 | 28.8 | 32.7 |
| 1999..... | 4.4 | * | 0.2 | * | 0.1 | 0.2 | 1.0 | 4.1 | 8.8 | 19.0 | 28.0 | 30.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | * | 0.1 | 0.3 | 1.3 | 9.2 | 16.6 | 28.3 | 42.7 | 41.5 |
| 2003..... | 6.6 | * | * | * | 0.1 | 0.3 | 1.3 | 8.9 | 14.9 | 27.9 | 44.6 | 47.5 |
| 2002..... | 6.4 | * | * | * | * | 0.3 | 1.4 | 8.2 | 14.5 | 28.5 | 41.6 | 46.2 |
| 2001..... | 6.1 | * | * | * | * | 0.2 | 1.3 | 7.7 | 13.9 | 26.8 | 41.0 | 50.3 |
| 2000..... | 5.9 | * | * | * | 0.1 | 0.3 | 1.3 | 7.2 | 13.8 | 27.0 | 42.1 | 44.7 |
| 1999..... | 5.6 | * | 0.3 | * | 0.1 | 0.2 | 1.4 | 6.5 | 12.9 | 27.5 | 39.0 | 40.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | 0.1 | 0.4 | 2.2 | 5.3 | 12.8 | 23.6 | 28.3 |
| 2003..... | 3.5 | * | * | * | * | 0.2 | 0.5 | 2.1 | 5.3 | 13.0 | 21.4 | 27.2 |
| 2002..... | 3.4 | * | * | * | * | 0.2 | 0.5 | 1.8 | 5.1 | 12.2 | 21.1 | 30.2 |
| 2001..... | 3.4 | * | * | * | * | 0.2 | 0.4 | 1.9 | 5.0 | 12.1 | 21.1 | 27.2 |
| 2000..... | 3.3 | * | * | * | * | 0.1 | 0.6 | 1.9 | 5.1 | 11.8 | 20.1 | 27.8 |
| 1999..... | 3.3 | * | * | * | * | 0.2 | 0.6 | 1.8 | 5.1 | 11.9 | 20.9 | 26.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 1.0 | 7.2 | 16.6 | 30.8 | 48.6 | 50.9 |
| 2003..... | 3.8 | * | * | * | * | * | 0.8 | 7.5 | 16.1 | 32.5 | 49.8 | 51.8 |
| 2002..... | 3.5 | * | * | * | * | * | 0.8 | 5.9 | 15.7 | 34.1 | 44.2 | 59.9 |
| 2001..... | 3.4 | * | * | * | * | * | 0.8 | 6.7 | 14.1 | 28.8 | 52.1 | 52.5 |
| 2000..... | 3.3 | * | * | * | * | * | 1.2 | 5.6 | 13.7 | 31.1 | 51.8 | 58.4 |
| 1999..... | 3.2 | * | * | * | * | * | 0.9 | 5.7 | 13.1 | 31.3 | 49.8 | 53.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | 1.6 | 11.2 | 27.5 | 42.3 | 62.9 | 71.4 |
| 2003..... | 4.8 | * | * | * | * | * | 1.2 | 12.5 | 24.3 | 45.2 | 66.0 | 55.4 |
| 2002..... | 4.5 | * | * | * | * | * | 1.1 | 9.8 | 24.1 | 48.9 | 59.9 | 75.3 |
| 2001..... | 4.3 | * | * | * | * | * | 1.3 | 11.1 | 21.5 | 41.1 | 68.0 | 64.8 |
| 2000..... | 4.2 | * | * | * | * | * | 1.7 | 9.1 | 20.5 | 43.9 | 74.7 | 72.6 |
| 1999..... | 3.9 | * | * | * | * | * | 1.1 | 9.6 | 20.7 | 42.8 | 63.1 | 66.9 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | * | 3.1 | 6.7 | 21.6 | 38.8 | 40.3 |
| 2003..... | 2.7 | * | * | * | * | * | * | 2.5 | 8.8 | 22.4 | 38.7 | 50.0 |
| 2002..... | 2.5 | * | * | * | * | * | * | 2.1 | 8.2 | 22.4 | 33.4 | 52.2 |
| 2001..... | 2.5 | * | * | * | * | * | * | 2.3 | 7.6 | 19.1 | 41.4 | 46.5 |
| 2000..... | 2.4 | * | * | * | * | * | * | 2.2 | 7.6 | 21.1 | 36.3 | 51.4 |
| 1999..... | 2.4 | * | * | * | * | * | * | 1.9 | 6.3 | 22.2 | 40.9 | 46.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of liver and intrahepatic bile ducts (C22) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.4 | * | 0.2 | * | 0.1 | 0.2 | 0.8 | 5.5 | 10.2 | 19.0 | 30.3 | 31.5 |
| 2003..... | 5.2 | * | * | * | 0.1 | 0.2 | 0.9 | 5.2 | 9.4 | 18.8 | 29.7 | 32.6 |
| 2002..... | 5.1 | * | 0.2 | * | 0.1 | 0.3 | 0.9 | 4.8 | 9.1 | 18.6 | 28.4 | 33.9 |
| 2001..... | 4.9 | * | * | * | 0.1 | 0.2 | 0.9 | 4.6 | 8.9 | 18.1 | 27.9 | 33.2 |
| 2000..... | 4.8 | * | 0.2 | * | 0.1 | 0.2 | 0.9 | 4.3 | 8.9 | 17.9 | 27.7 | 31.7 |
| 1999..... | 4.6 | * | 0.2 | * | 0.1 | 0.2 | 1.0 | 3.9 | 8.4 | 18.2 | 27.1 | 29.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | * | 0.1 | 0.3 | 1.3 | 9.0 | 15.6 | 27.2 | 41.4 | 40.0 |
| 2003..... | 6.9 | * | * | * | * | 0.3 | 1.3 | 8.4 | 14.1 | 26.6 | 43.5 | 47.1 |
| 2002..... | 6.6 | * | * | * | * | 0.4 | 1.4 | 8.0 | 13.7 | 27.1 | 40.5 | 44.8 |
| 2001..... | 6.3 | * | * | * | * | 0.2 | 1.3 | 7.4 | 13.2 | 25.8 | 39.7 | 49.7 |
| 2000..... | 6.2 | * | * | * | * | 0.3 | 1.3 | 6.9 | 13.3 | 25.9 | 40.5 | 43.4 |
| 1999..... | 5.8 | * | * | * | 0.1 | 0.3 | 1.5 | 6.2 | 12.2 | 26.5 | 37.9 | 38.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | 0.2 | 0.4 | 2.1 | 5.2 | 12.1 | 22.7 | 27.7 |
| 2003..... | 3.6 | * | * | * | * | 0.2 | 0.5 | 2.1 | 5.0 | 12.3 | 20.4 | 26.2 |
| 2002..... | 3.6 | * | * | * | * | 0.2 | 0.5 | 1.7 | 4.8 | 11.4 | 20.5 | 29.2 |
| 2001..... | 3.5 | * | * | * | * | 0.2 | 0.4 | 1.9 | 4.8 | 11.7 | 20.2 | 26.4 |
| 2000..... | 3.4 | * | * | * | * | 0.1 | 0.6 | 1.9 | 4.9 | 11.2 | 19.4 | 27.0 |
| 1999..... | 3.4 | * | * | * | * | 0.2 | 0.5 | 1.8 | 4.9 | 11.3 | 20.0 | 26.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | 0.1 | 0.1 | 0.6 | 4.5 | 8.6 | 17.1 | 28.8 | 30.1 |
| 2003..... | 5.1 | * | * | * | 0.1 | 0.1 | 0.7 | 4.2 | 8.0 | 17.0 | 28.2 | 31.1 |
| 2002..... | 5.0 | * | * | * | * | 0.2 | 0.7 | 3.9 | 7.7 | 17.0 | 27.4 | 32.6 |
| 2001..... | 4.8 | * | * | * | 0.1 | 0.1 | 0.6 | 3.7 | 7.6 | 16.6 | 26.5 | 32.4 |
| 2000..... | 4.6 | * | * | * | * | 0.1 | 0.6 | 3.4 | 7.5 | 16.2 | 26.5 | 30.8 |
| 1999..... | 4.5 | * | 0.2 | * | 0.1 | 0.1 | 0.7 | 3.1 | 7.1 | 16.8 | 26.1 | 28.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | * | * | 0.2 | 0.9 | 7.4 | 12.8 | 24.4 | 40.0 | 38.0 |
| 2003..... | 6.7 | * | * | * | * | * | 1.0 | 6.7 | 11.9 | 24.3 | 41.6 | 45.0 |
| 2002..... | 6.4 | * | * | * | * | 0.2 | 1.1 | 6.4 | 11.5 | 24.6 | 38.7 | 43.6 |
| 2001..... | 6.1 | * | * | * | * | * | 0.9 | 5.9 | 11.2 | 24.2 | 38.0 | 48.9 |
| 2000..... | 5.9 | * | * | * | * | 0.2 | 0.8 | 5.3 | 11.0 | 23.6 | 39.2 | 41.6 |
| 1999..... | 5.7 | * | * | * | * | * | 1.1 | 4.8 | 10.2 | 24.6 | 36.7 | 39.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 0.4 | 1.7 | 4.7 | 10.9 | 21.2 | 26.5 |
| 2003..... | 3.6 | * | * | * | * | * | 0.5 | 1.7 | 4.2 | 10.7 | 19.2 | 25.0 |
| 2002..... | 3.6 | * | * | * | * | 0.2 | 0.4 | 1.4 | 4.2 | 10.4 | 19.9 | 27.8 |
| 2001..... | 3.4 | * | * | * | * | * | 0.3 | 1.5 | 4.1 | 10.2 | 18.9 | 25.5 |
| 2000..... | 3.4 | * | * | * | * | * | 0.4 | 1.6 | 4.3 | 10.1 | 18.2 | 26.5 |
| 1999..... | 3.4 | * | * | * | * | * | 0.4 | 1.6 | 4.2 | 10.2 | 19.1 | 24.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of liver and intrahepatic bile ducts (C22) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | 0.4 | 1.4 | 9.8 | 17.7 | 26.1 | 34.7 | 36.6 |
| 2003..... | 5.1 | * | * | * | * | 0.5 | 1.4 | 10.0 | 16.5 | 23.1 | 31.7 | 41.9 |
| 2002..... | 4.7 | * | * | * | * | 0.5 | 1.5 | 8.9 | 15.0 | 23.3 | 30.9 | 38.1 |
| 2001..... | 4.6 | * | * | * | * | * | 1.7 | 8.9 | 15.1 | 21.9 | 32.3 | 35.3 |
| 2000..... | 4.5 | * | * | * | * | 0.4 | 1.8 | 8.8 | 14.5 | 23.7 | 29.1 | 31.8 |
| 1999..... | 4.3 | * | * | * | * | 0.4 | 1.7 | 7.4 | 13.4 | 24.6 | 29.7 | 34.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | * | * | 2.3 | 16.9 | 30.5 | 41.3 | 44.2 | 47.8 |
| 2003..... | 7.0 | * | * | * | * | * | 2.2 | 17.3 | 26.4 | 33.8 | 44.7 | 63.9 |
| 2002..... | 6.6 | * | * | * | * | * | 2.3 | 15.4 | 24.4 | 38.2 | 46.3 | 41.6 |
| 2001..... | 6.1 | * | * | * | * | * | 2.7 | 15.1 | 24.2 | 30.3 | 42.6 | 42.7 |
| 2000..... | 6.3 | * | * | * | * | * | 2.5 | 15.0 | 24.1 | 37.3 | 40.4 | 48.1 |
| 1999..... | 5.7 | * | * | * | * | * | 2.5 | 13.1 | 20.3 | 36.8 | 41.6 | 30.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | * | 3.7 | 7.4 | 15.2 | 29.3 | 32.1 |
| 2003..... | 3.3 | * | * | * | * | * | * | 3.6 | 8.6 | 15.5 | 24.3 | 33.4 |
| 2002..... | 3.0 | * | * | * | * | * | 0.8 | 3.3 | 7.4 | 12.8 | 22.2 | 36.8 |
| 2001..... | 3.3 | * | * | * | * | * | 0.8 | 3.6 | 7.8 | 16.0 | 26.4 | 32.5 |
| 2000..... | 2.9 | * | * | * | * | * | 1.2 | 3.4 | 6.7 | 14.1 | 22.8 | 25.7 |
| 1999..... | 3.1 | * | * | * | * | * | 0.9 | 2.5 | 7.9 | 15.9 | 23.1 | 36.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Liver (C22.0,C22.2-C22.9) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | 0.1 | * | 0.1 | 0.2 | 0.6 | 4.9 | 8.5 | 15.2 | 22.9 | 23.2 |
| 2003..... | 3.9 | * | 0.1 | * | 0.1 | 0.2 | 0.7 | 4.6 | 7.8 | 15.3 | 23.0 | 24.3 |
| 2002..... | 3.8 | * | 0.2 | * | 0.1 | 0.2 | 0.7 | 4.3 | 7.6 | 15.7 | 22.2 | 24.7 |
| 2001..... | 3.7 | * | * | * | 0.1 | 0.2 | 0.7 | 4.2 | 7.5 | 14.7 | 21.9 | 24.2 |
| 2000..... | 3.6 | * | 0.1 | 0.0 | 0.1 | 0.2 | 0.8 | 3.8 | 7.5 | 14.8 | 22.1 | 23.4 |
| 1999..... | 3.5 | * | 0.2 | * | 0.1 | 0.2 | 0.8 | 3.5 | 7.0 | 15.2 | 21.5 | 22.5 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 0.1 | 0.2 | 1.1 | 8.4 | 13.9 | 23.2 | 33.1 | 30.6 |
| 2003..... | 5.5 | * | * | * | * | 0.2 | 1.1 | 8.1 | 12.4 | 22.5 | 35.4 | 35.2 |
| 2002..... | 5.3 | * | * | * | * | 0.3 | 1.2 | 7.4 | 12.3 | 24.0 | 32.7 | 34.6 |
| 2001..... | 5.1 | * | * | * | * | 0.2 | 1.1 | 7.1 | 12.0 | 22.0 | 32.2 | 38.5 |
| 2000..... | 5.0 | * | * | * | * | 0.2 | 1.2 | 6.4 | 11.9 | 22.4 | 33.7 | 33.9 |
| 1999..... | 4.7 | * | 0.3 | * | 0.1 | 0.2 | 1.2 | 5.9 | 10.9 | 23.1 | 31.5 | 31.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | * | 0.2 | 1.5 | 3.4 | 8.6 | 16.1 | 19.8 |
| 2003..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.5 | 9.3 | 14.7 | 19.5 |
| 2002..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.2 | 8.7 | 15.2 | 20.4 |
| 2001..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.4 | 3.4 | 8.6 | 15.1 | 18.2 |
| 2000..... | 2.3 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.5 | 8.5 | 14.4 | 19.1 |
| 1999..... | 2.4 | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.5 | 8.6 | 15.0 | 18.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | 0.8 | 6.3 | 14.4 | 24.6 | 39.6 | 41.4 |
| 2003..... | 3.2 | * | * | * | * | * | 0.6 | 6.7 | 13.7 | 26.3 | 41.1 | 40.9 |
| 2002..... | 3.0 | * | * | * | * | * | 0.7 | 5.6 | 13.2 | 29.1 | 37.4 | 49.9 |
| 2001..... | 2.9 | * | * | * | * | * | 0.7 | 6.2 | 12.6 | 24.5 | 42.2 | 40.6 |
| 2000..... | 2.8 | * | * | * | * | * | 1.0 | 4.9 | 11.6 | 25.7 | 43.9 | 51.1 |
| 1999..... | 2.7 | * | * | * | * | * | 0.6 | 5.1 | 11.6 | 26.5 | 40.8 | 45.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | * | 1.3 | 10.5 | 24.8 | 35.9 | 53.3 | 57.4 |
| 2003..... | 4.2 | * | * | * | * | * | 1.1 | 11.6 | 22.0 | 38.8 | 57.1 | 43.1 |
| 2002..... | 4.0 | * | * | * | * | * | 1.0 | 9.5 | 21.3 | 43.6 | 52.9 | 63.6 |
| 2001..... | 3.8 | * | * | * | * | * | 1.1 | 10.6 | 20.4 | 35.9 | 57.2 | 59.0 |
| 2000..... | 3.8 | * | * | * | * | * | 1.6 | 8.4 | 18.7 | 38.0 | 65.9 | 66.5 |
| 1999..... | 3.6 | * | * | * | * | * | 0.9 | 9.3 | 19.5 | 38.7 | 53.7 | 58.2 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | 2.0 | 5.1 | 15.6 | 30.1 | 33.1 |
| 2003..... | 2.0 | * | * | * | * | * | * | 1.7 | 6.2 | 16.3 | 30.0 | 39.8 |
| 2002..... | 2.0 | * | * | * | * | * | * | 1.8 | 5.9 | 17.6 | 26.7 | 43.1 |
| 2001..... | 1.9 | * | * | * | * | * | * | 1.9 | 5.6 | 15.4 | 32.0 | 31.6 |
| 2000..... | 1.8 | * | * | * | * | * | * | 1.5 | 5.3 | 16.1 | 29.1 | 43.5 |
| 1999..... | 1.8 | * | * | * | * | * | * | * | 4.6 | 17.0 | 32.3 | 39.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Liver (C22.0,C22.2-C22.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | 0.1 | 0.2 | 0.6 | 4.7 | 7.9 | 14.5 | 22.0 | 22.3 |
| 2003..... | 4.0 | * | * | * | 0.1 | 0.2 | 0.7 | 4.4 | 7.3 | 14.5 | 22.0 | 23.6 |
| 2002..... | 3.9 | * | 0.2 | * | 0.1 | 0.2 | 0.7 | 4.1 | 7.1 | 14.7 | 21.4 | 23.6 |
| 2001..... | 3.8 | * | * | * | 0.1 | 0.2 | 0.7 | 4.0 | 7.1 | 14.0 | 20.9 | 23.5 |
| 2000..... | 3.7 | * | 0.2 | * | 0.1 | 0.2 | 0.8 | 3.7 | 7.2 | 14.1 | 21.1 | 22.3 |
| 1999..... | 3.6 | * | 0.2 | * | 0.1 | 0.2 | 0.8 | 3.4 | 6.7 | 14.4 | 20.7 | 21.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | 0.1 | 0.2 | 1.0 | 8.1 | 13.0 | 22.2 | 31.9 | 29.2 |
| 2003..... | 5.7 | * | * | * | * | 0.2 | 1.1 | 7.6 | 11.6 | 21.3 | 34.2 | 34.7 |
| 2002..... | 5.5 | * | * | * | * | 0.3 | 1.2 | 7.2 | 11.5 | 22.6 | 31.6 | 33.2 |
| 2001..... | 5.2 | * | * | * | * | 0.2 | 1.1 | 6.7 | 11.3 | 21.0 | 31.0 | 37.6 |
| 2000..... | 5.1 | * | * | * | * | 0.2 | 1.1 | 6.2 | 11.4 | 21.4 | 32.2 | 32.4 |
| 1999..... | 4.9 | * | * | * | 0.1 | 0.2 | 1.3 | 5.6 | 10.2 | 22.1 | 30.5 | 29.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | 0.2 | 1.5 | 3.3 | 8.0 | 15.3 | 19.2 |
| 2003..... | 2.5 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.3 | 8.7 | 13.9 | 18.7 |
| 2002..... | 2.5 | * | * | * | * | 0.2 | 0.3 | 1.2 | 3.0 | 8.1 | 14.6 | 19.4 |
| 2001..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.2 | 8.1 | 14.3 | 17.6 |
| 2000..... | 2.4 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.3 | 8.0 | 13.8 | 18.2 |
| 1999..... | 2.4 | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.4 | 8.1 | 14.3 | 18.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | 0.1 | * | 0.4 | 3.8 | 6.3 | 12.7 | 20.5 | 20.9 |
| 2003..... | 3.8 | * | * | * | * | * | 0.5 | 3.4 | 5.9 | 12.8 | 20.5 | 22.1 |
| 2002..... | 3.8 | * | * | * | * | 0.1 | 0.5 | 3.3 | 5.8 | 13.1 | 20.4 | 22.4 |
| 2001..... | 3.6 | * | * | * | * | 0.1 | 0.4 | 3.1 | 5.8 | 12.6 | 19.7 | 22.7 |
| 2000..... | 3.5 | * | * | * | * | 0.1 | 0.5 | 2.8 | 5.9 | 12.4 | 19.8 | 21.3 |
| 1999..... | 3.4 | * | 0.2 | * | 0.1 | * | 0.6 | 2.6 | 5.4 | 13.0 | 19.7 | 20.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | * | 0.6 | 6.6 | 10.2 | 19.5 | 30.2 | 27.1 |
| 2003..... | 5.3 | * | * | * | * | * | 0.7 | 6.0 | 9.5 | 19.3 | 32.2 | 32.5 |
| 2002..... | 5.2 | * | * | * | * | * | 0.8 | 5.6 | 9.4 | 20.3 | 30.1 | 32.0 |
| 2001..... | 5.0 | * | * | * | * | * | 0.7 | 5.2 | 9.4 | 19.5 | 29.5 | 36.8 |
| 2000..... | 4.8 | * | * | * | * | * | 0.7 | 4.6 | 9.2 | 19.0 | 30.8 | 30.1 |
| 1999..... | 4.6 | * | * | * | * | * | 0.9 | 4.1 | 8.2 | 20.1 | 29.4 | 30.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | * | 0.2 | 1.1 | 2.7 | 6.8 | 13.9 | 18.0 |
| 2003..... | 2.3 | * | * | * | * | * | 0.2 | 1.0 | 2.5 | 7.3 | 12.6 | 17.5 |
| 2002..... | 2.4 | * | * | * | * | * | 0.2 | 0.9 | 2.5 | 7.1 | 14.0 | 18.3 |
| 2001..... | 2.3 | * | * | * | * | * | 0.2 | 1.0 | 2.5 | 6.7 | 13.2 | 16.8 |
| 2000..... | 2.3 | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 6.8 | 12.6 | 17.7 |
| 1999..... | 2.3 | * | * | * | * | * | 0.3 | 1.0 | 2.7 | 7.0 | 13.3 | 16.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Liver (C22.0,C22.2-C22.9) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | 0.4 | 1.1 | 9.0 | 16.0 | 21.6 | 27.9 | 29.4 |
| 2003..... | 4.3 | * | * | * | * | 0.4 | 1.3 | 8.8 | 14.5 | 18.7 | 25.5 | 33.0 |
| 2002..... | 4.1 | * | * | * | * | 0.4 | 1.3 | 8.3 | 13.2 | 20.0 | 24.4 | 28.7 |
| 2001..... | 4.0 | * | * | * | * | * | 1.6 | 8.1 | 13.1 | 18.5 | 26.3 | 27.2 |
| 2000..... | 3.9 | * | * | * | * | * | 1.6 | 8.1 | 12.4 | 20.4 | 24.3 | 24.8 |
| 1999..... | 3.8 | * | * | * | * | * | 1.4 | 7.0 | 11.8 | 21.3 | 24.6 | 28.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | * | * | * | 2.1 | 16.1 | 28.4 | 36.1 | 38.2 | 37.6 |
| 2003..... | 6.2 | * | * | * | * | * | 2.2 | 16.2 | 23.9 | 27.9 | 37.5 | 51.1 |
| 2002..... | 6.0 | * | * | * | * | * | 2.1 | 14.8 | 22.7 | 33.2 | 37.2 | 35.0 |
| 2001..... | 5.5 | * | * | * | * | * | 2.6 | 14.4 | 21.5 | 26.0 | 35.7 | 34.6 |
| 2000..... | 5.7 | * | * | * | * | * | 2.2 | 14.3 | 22.1 | 33.2 | 34.9 | 42.3 |
| 1999..... | 5.2 | * | * | * | * | * | 2.3 | 12.8 | 19.1 | 32.8 | 35.0 | 25.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | * | 2.8 | 6.0 | 11.4 | 22.1 | 26.1 |
| 2003..... | 2.6 | * | * | * | * | * | * | 2.5 | 6.9 | 12.2 | 18.7 | 26.0 |
| 2002..... | 2.3 | * | * | * | * | * | * | 2.7 | 5.5 | 10.6 | 17.1 | 26.3 |
| 2001..... | 2.6 | * | * | * | * | * | * | 2.7 | 6.4 | 13.3 | 21.1 | 24.4 |
| 2000..... | 2.3 | * | * | * | * | * | 1.1 | 2.7 | 4.7 | 11.4 | 18.4 | 18.3 |
| 1999..... | 2.5 | * | * | * | * | * | * | 2.1 | 6.0 | 13.1 | 18.7 | 30.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intrahepatic bile duct carcinoma (C22.1) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.3 | 4.7 | 8.3 | 9.2 |
| 2003..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.1 | 4.5 | 7.7 | 9.1 |
| 2002..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.0 | 3.9 | 7.1 | 10.3 |
| 2001..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 4.1 | 7.1 | 9.9 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 3.9 | 6.7 | 9.3 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.8 | 6.5 | 8.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.2 | 0.9 | 2.6 | 5.1 | 9.5 | 10.9 |
| 2003..... | 1.2 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 5.4 | 9.2 | 12.4 |
| 2002..... | 1.0 | * | * | * | * | * | 0.2 | 0.8 | 2.2 | 4.5 | 8.8 | 11.6 |
| 2001..... | 1.0 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 4.8 | 8.8 | 11.9 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 4.6 | 8.3 | 10.8 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 4.4 | 7.5 | 8.9 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 4.3 | 7.5 | 8.5 |
| 2003..... | 1.1 | * | * | * | * | * | 0.2 | 0.8 | 1.8 | 3.7 | 6.6 | 7.6 |
| 2002..... | 1.0 | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 3.4 | 5.9 | 9.8 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 3.6 | 6.0 | 9.0 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.6 | 1.7 | 3.3 | 5.7 | 8.8 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 3.3 | 5.8 | 7.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.9 | 2.2 | 6.1 | 9.1 | 9.5 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.8 | 2.4 | 6.2 | 8.7 | 10.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 2.5 | 5.0 | 6.8 | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 1.6 | 4.3 | 9.9 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.7 | 2.1 | 5.4 | 7.9 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 1.5 | 4.7 | 9.0 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 2.7 | 6.4 | 9.7 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 2.3 | 6.5 | 8.8 | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 2.8 | 5.4 | * | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 5.2 | 10.7 | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 5.9 | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 1.1 | * | 6.0 | 8.6 | * |
| 2003..... | 0.7 | * | * | * | * | * | * | * | 2.6 | 6.0 | 8.7 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 2.3 | 4.7 | 6.7 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.6 | 9.4 | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 2.3 | 5.0 | 7.3 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 5.2 | 8.7 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intrahepatic bile duct carcinoma (C22.1) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 4.5 | 8.3 | 9.2 |
| 2003..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.1 | 4.4 | 7.6 | 9.0 |
| 2002..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.0 | 3.9 | 7.1 | 10.3 |
| 2001..... | 1.1 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 4.1 | 7.0 | 9.8 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 3.8 | 6.7 | 9.4 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 3.7 | 6.4 | 8.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.2 | 0.9 | 2.6 | 5.0 | 9.5 | 10.7 |
| 2003..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 5.3 | 9.3 | 12.4 |
| 2002..... | 1.1 | * | * | * | * | * | 0.2 | 0.8 | 2.2 | 4.5 | 8.8 | 11.6 |
| 2001..... | 1.1 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.7 | 8.6 | 12.1 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 4.5 | 8.3 | 11.0 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 4.4 | 7.4 | 8.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 4.1 | 7.5 | 8.5 |
| 2003..... | 1.2 | * | * | * | * | * | 0.2 | 0.8 | 1.7 | 3.6 | 6.5 | 7.5 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 3.3 | 5.9 | 9.8 |
| 2001..... | 1.1 | * | * | * | * | * | 0.1 | 0.6 | 1.6 | 3.5 | 5.9 | 8.8 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.6 | 1.6 | 3.2 | 5.6 | 8.8 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 3.2 | 5.7 | 7.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 4.4 | 8.3 | 9.2 |
| 2003..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.0 | 4.2 | 7.7 | 9.1 |
| 2002..... | 1.2 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 3.8 | 7.0 | 10.1 |
| 2001..... | 1.2 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.0 | 6.9 | 9.7 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.7 | 1.7 | 3.8 | 6.7 | 9.6 |
| 1999..... | 1.1 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.8 | 6.4 | 8.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 0.9 | 2.6 | 4.9 | 9.7 | 10.9 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 5.1 | 9.4 | 12.4 |
| 2002..... | 1.2 | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 4.4 | 8.6 | 11.6 |
| 2001..... | 1.2 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.6 | 8.6 | 12.1 |
| 2000..... | 1.1 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.6 | 8.4 | 11.5 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 4.5 | 7.2 | 9.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 4.0 | 7.2 | 8.5 |
| 2003..... | 1.2 | * | * | * | * | * | 0.2 | 0.8 | 1.7 | 3.5 | 6.6 | 7.6 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 3.4 | 5.9 | 9.5 |
| 2001..... | 1.1 | * | * | * | * | * | * | 0.5 | 1.6 | 3.5 | 5.8 | 8.7 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.6 | 1.6 | 3.2 | 5.6 | 8.8 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 3.2 | 5.8 | 7.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intrahepatic bile duct carcinoma (C22.1) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.8 | 1.7 | 4.4 | 6.8 | 7.2 |
| 2003..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.0 | 4.4 | 6.1 | 8.9 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.6 | 1.8 | 3.3 | 6.6 | 9.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.7 | 2.0 | 3.4 | 5.9 | 8.1 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.7 | 2.0 | 3.3 | 4.8 | 7.0 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 1.6 | 3.3 | 5.2 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | 2.1 | 5.3 | 6.0 | * |
| 2003..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.5 | 5.9 | 7.2 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 1.8 | 5.0 | 9.1 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 2.7 | 4.3 | 6.9 | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 2.1 | 4.0 | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 4.1 | 6.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.8 | 1.5 | 3.9 | 7.2 | * |
| 2003..... | 0.7 | * | * | * | * | * | * | 1.1 | 1.7 | 3.3 | 5.5 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 1.9 | 2.2 | 5.1 | 10.5 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 2.7 | 5.4 | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 2.0 | 2.7 | 4.3 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 1.9 | 2.8 | 4.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of gallbladder and extrahepatic bile ducts (C23-C24) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 4.7 | 8.7 | 13.3 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.6 | 1.7 | 4.2 | 9.3 | 14.3 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.6 | 1.7 | 4.6 | 8.9 | 14.5 |
| 2001..... | 1.3 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 4.8 | 9.9 | 15.8 |
| 2000..... | 1.3 | * | * | * | * | * | 0.1 | 0.6 | 2.0 | 5.3 | 10.0 | 16.0 |
| 1999..... | 1.3 | * | * | * | * | * | 0.1 | 0.6 | 2.0 | 5.1 | 10.0 | 16.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.3 | 1.4 | 4.1 | 8.6 | 12.9 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 4.2 | 8.5 | 14.1 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 4.3 | 8.5 | 15.0 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 4.0 | 9.9 | 15.8 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 5.0 | 9.5 | 15.8 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.5 | 9.3 | 15.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.1 | 0.5 | 2.1 | 5.2 | 8.8 | 13.5 |
| 2003..... | 1.4 | * | * | * | * | * | 0.1 | 0.7 | 2.0 | 4.3 | 9.8 | 14.4 |
| 2002..... | 1.4 | * | * | * | * | * | 0.2 | 0.8 | 1.9 | 4.9 | 9.2 | 14.3 |
| 2001..... | 1.6 | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 5.5 | 9.9 | 15.8 |
| 2000..... | 1.6 | * | * | * | * | * | 0.1 | 0.8 | 2.3 | 5.5 | 10.4 | 16.0 |
| 1999..... | 1.6 | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 5.5 | 10.5 | 16.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.7 | 2.1 | 5.8 | 10.7 | 14.3 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.6 | 2.3 | 5.6 | 11.9 | 13.0 |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.0 | 2.0 | 8.0 | 12.5 | 13.3 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.9 | 3.1 | 8.5 | 12.3 | 20.6 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.9 | 3.4 | 6.5 | 16.4 | 18.6 |
| 1999..... | 0.9 | * | * | * | * | * | * | 1.1 | 3.7 | 8.3 | 14.2 | 21.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 2.0 | 4.8 | 10.0 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 5.2 | 10.7 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 6.5 | 10.7 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 2.3 | 7.0 | 13.4 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 5.5 | 12.8 | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | 10.9 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.2 | 6.7 | 11.1 | 15.1 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | 3.0 | 5.9 | 12.6 | 16.4 |
| 2002..... | 1.0 | * | * | * | * | * | * | 1.4 | 2.8 | 9.1 | 13.8 | * |
| 2001..... | 1.1 | * | * | * | * | * | * | 1.3 | 3.9 | 9.8 | 11.5 | 25.1 |
| 2000..... | 1.1 | * | * | * | * | * | * | 1.4 | 4.9 | 7.3 | 18.8 | * |
| 1999..... | 1.3 | * | * | * | * | * | * | 1.7 | 5.1 | 11.7 | 16.3 | 22.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of gallbladder and extrahepatic bile ducts (C23-C24) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.4 | 1.7 | 4.6 | 8.6 | 13.3 |
| 2003..... | 1.3 | * | * | * | * | * | 0.1 | 0.6 | 1.7 | 4.1 | 9.1 | 14.3 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.6 | 1.6 | 4.4 | 8.7 | 14.5 |
| 2001..... | 1.3 | * | * | * | * | * | 0.1 | 0.6 | 1.7 | 4.6 | 9.8 | 15.6 |
| 2000..... | 1.4 | * | * | * | * | * | 0.1 | 0.6 | 1.8 | 5.2 | 9.7 | 15.8 |
| 1999..... | 1.3 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 4.9 | 9.8 | 16.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 0.3 | 1.4 | 4.0 | 8.6 | 12.9 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.5 | 1.4 | 4.1 | 8.3 | 14.4 |
| 2002..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 4.2 | 8.4 | 14.7 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 3.8 | 9.7 | 16.0 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.9 | 9.3 | 15.6 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.6 | 9.2 | 15.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.1 | 0.4 | 2.1 | 5.0 | 8.7 | 13.4 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.7 | 1.9 | 4.1 | 9.6 | 14.3 |
| 2002..... | 1.5 | * | * | * | * | * | 0.1 | 0.7 | 1.8 | 4.6 | 9.0 | 14.4 |
| 2001..... | 1.6 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 5.2 | 9.8 | 15.4 |
| 2000..... | 1.7 | * | * | * | * | * | 0.1 | 0.8 | 2.1 | 5.4 | 10.0 | 15.9 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.6 | 2.2 | 5.1 | 10.2 | 16.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 4.4 | 8.5 | 13.4 |
| 2003..... | 1.4 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 4.1 | 9.0 | 14.4 |
| 2002..... | 1.3 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.3 | 8.6 | 14.5 |
| 2001..... | 1.4 | * | * | * | * | * | 0.1 | 0.6 | 1.6 | 4.4 | 9.8 | 15.5 |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 0.6 | 1.8 | 5.1 | 9.8 | 16.2 |
| 1999..... | 1.5 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 4.8 | 9.6 | 16.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 0.4 | 1.3 | 3.8 | 8.4 | 13.2 |
| 2003..... | 1.1 | * | * | * | * | * | * | 0.4 | 1.4 | 4.2 | 8.2 | 14.8 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.3 | 1.4 | 4.1 | 8.3 | 15.0 |
| 2001..... | 1.1 | * | * | * | * | * | * | 0.5 | 1.4 | 3.7 | 9.6 | 15.6 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.9 | 9.5 | 16.1 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 4.4 | 9.3 | 15.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 0.4 | 2.0 | 4.8 | 8.5 | 13.5 |
| 2003..... | 1.6 | * | * | * | * | * | * | 0.6 | 1.9 | 4.1 | 9.5 | 14.3 |
| 2002..... | 1.6 | * | * | * | * | * | 0.1 | 0.7 | 1.8 | 4.5 | 8.8 | 14.4 |
| 2001..... | 1.7 | * | * | * | * | * | 0.1 | 0.7 | 1.9 | 4.9 | 9.9 | 15.5 |
| 2000..... | 1.8 | * | * | * | * | * | 0.1 | 0.7 | 2.0 | 5.3 | 10.1 | 16.2 |
| 1999..... | 1.8 | * | * | * | * | * | * | 0.6 | 2.1 | 5.1 | 9.9 | 16.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of gallbladder and extrahepatic bile ducts (C23-C24) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 2.3 | 5.5 | 8.9 | 11.2 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.8 | 1.5 | 3.9 | 9.3 | 12.5 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.9 | 2.3 | 5.2 | 9.5 | 13.9 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.7 | 2.1 | 5.8 | 8.8 | 15.9 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.8 | 2.0 | 5.9 | 7.8 | 11.4 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.7 | 2.2 | 5.3 | 10.1 | 12.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 1.7 | 4.6 | 9.4 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 3.3 | 7.8 | * |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 1.9 | 4.5 | 7.9 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 4.6 | 9.3 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 4.8 | 6.8 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 5.3 | 9.1 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | 2.8 | 6.2 | 8.5 | 14.1 |
| 2003..... | 1.0 | * | * | * | * | * | * | 1.0 | 1.6 | 4.3 | 10.1 | 14.4 |
| 2002..... | 1.2 | * | * | * | * | * | * | 1.2 | 2.7 | 5.7 | 10.4 | 15.9 |
| 2001..... | 1.2 | * | * | * | * | * | * | 1.0 | 2.6 | 6.6 | 8.6 | 16.3 |
| 2000..... | 1.1 | * | * | * | * | * | * | 1.3 | 2.5 | 6.6 | 8.3 | 12.2 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | 2.8 | 5.3 | 10.7 | 15.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of pancreas (C25) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 10.8 | * | * | * | * | 0.2 | 1.2 | 6.3 | 19.8 | 44.3 | 76.0 | 97.7 |
| 2003..... | 10.6 | * | * | * | * | 0.2 | 1.2 | 6.2 | 19.1 | 44.2 | 75.4 | 95.2 |
| 2002..... | 10.5 | * | * | * | * | 0.1 | 1.3 | 6.1 | 19.2 | 44.8 | 75.0 | 94.5 |
| 2001..... | 10.5 | * | * | * | * | 0.1 | 1.3 | 6.1 | 19.2 | 44.5 | 74.1 | 100.1 |
| 2000..... | 10.4 | * | * | * | * | 0.1 | 1.2 | 6.3 | 19.4 | 44.1 | 76.5 | 96.4 |
| 1999..... | 10.4 | * | * | * | * | 0.1 | 1.3 | 6.4 | 19.8 | 45.0 | 73.1 | 99.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 10.9 | * | * | * | * | 0.2 | 1.5 | 8.1 | 24.1 | 52.6 | 84.6 | 102.0 |
| 2003..... | 10.5 | * | * | * | * | 0.2 | 1.5 | 7.7 | 23.1 | 50.9 | 85.4 | 95.6 |
| 2002..... | 10.5 | * | * | * | * | 0.1 | 1.6 | 7.8 | 23.6 | 51.4 | 84.0 | 100.9 |
| 2001..... | 10.3 | * | * | * | * | 0.1 | 1.4 | 7.6 | 23.6 | 50.8 | 84.0 | 105.1 |
| 2000..... | 10.3 | * | * | * | * | 0.2 | 1.4 | 8.0 | 23.6 | 50.4 | 85.9 | 102.9 |
| 1999..... | 10.4 | * | * | * | * | 0.2 | 1.6 | 8.3 | 23.9 | 52.4 | 81.8 | 107.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 10.7 | * | * | * | * | 0.1 | 0.9 | 4.7 | 15.8 | 37.3 | 70.1 | 95.8 |
| 2003..... | 10.6 | * | * | * | * | 0.1 | 1.0 | 4.8 | 15.3 | 38.6 | 68.8 | 95.0 |
| 2002..... | 10.5 | * | * | * | * | * | 1.0 | 4.4 | 15.1 | 39.3 | 69.1 | 91.7 |
| 2001..... | 10.6 | * | * | * | * | 0.1 | 1.1 | 4.7 | 15.2 | 39.4 | 67.6 | 98.1 |
| 2000..... | 10.5 | * | * | * | * | 0.1 | 1.0 | 4.6 | 15.5 | 38.8 | 70.4 | 93.8 |
| 1999..... | 10.5 | * | * | * | * | 0.1 | 1.0 | 4.7 | 16.0 | 39.0 | 67.4 | 96.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 0.7 | 4.1 | 14.4 | 35.3 | 62.8 | 69.9 |
| 2003..... | 3.7 | * | * | * | * | * | 0.8 | 4.1 | 14.4 | 35.1 | 60.3 | 79.8 |
| 2002..... | 3.5 | * | * | * | * | * | 0.9 | 4.0 | 13.8 | 34.7 | 55.5 | 79.3 |
| 2001..... | 3.6 | * | * | * | * | * | 0.8 | 3.7 | 14.2 | 36.9 | 61.3 | 83.8 |
| 2000..... | 3.5 | * | * | * | * | * | 0.7 | 4.1 | 13.7 | 36.9 | 61.8 | 82.9 |
| 1999..... | 3.3 | * | * | * | * | * | 0.9 | 4.6 | 13.3 | 32.6 | 56.7 | 84.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | * | 1.0 | 4.7 | 17.4 | 42.4 | 68.3 | 57.4 |
| 2003..... | 3.6 | * | * | * | * | * | 1.0 | 5.0 | 17.8 | 40.0 | 67.1 | 70.8 |
| 2002..... | 3.3 | * | * | * | * | * | 1.2 | 5.3 | 15.7 | 40.9 | 55.8 | 60.2 |
| 2001..... | 3.4 | * | * | * | * | * | 0.9 | 4.3 | 16.2 | 42.3 | 68.9 | 62.9 |
| 2000..... | 3.4 | * | * | * | * | * | 0.8 | 5.2 | 16.6 | 45.2 | 62.9 | 80.6 |
| 1999..... | 3.2 | * | * | * | * | * | 1.2 | 5.7 | 15.2 | 40.0 | 56.3 | 86.3 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | * | * | 3.5 | 11.8 | 29.5 | 59.0 | 76.3 |
| 2003..... | 3.8 | * | * | * | * | * | * | 3.2 | 11.3 | 31.2 | 55.6 | 84.4 |
| 2002..... | 3.7 | * | * | * | * | * | * | 2.8 | 12.1 | 29.8 | 55.4 | 88.7 |
| 2001..... | 3.9 | * | * | * | * | * | 0.8 | 3.2 | 12.5 | 32.6 | 56.2 | 94.0 |
| 2000..... | 3.6 | * | * | * | * | * | * | 3.0 | 11.1 | 30.4 | 61.1 | 84.1 |
| 1999..... | 3.5 | * | * | * | * | * | * | 3.5 | 11.6 | 26.8 | 56.9 | 83.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of pancreas (C25) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 11.9 | * | * | * | * | 0.2 | 1.3 | 6.6 | 20.2 | 44.9 | 76.5 | 98.9 |
| 2003..... | 11.7 | * | * | * | * | 0.2 | 1.3 | 6.4 | 19.4 | 44.8 | 76.1 | 95.7 |
| 2002..... | 11.6 | * | * | * | * | 0.1 | 1.4 | 6.3 | 19.6 | 45.4 | 75.8 | 94.9 |
| 2001..... | 11.5 | * | * | * | * | 0.1 | 1.3 | 6.4 | 19.6 | 44.9 | 74.6 | 100.6 |
| 2000..... | 11.4 | * | * | * | * | 0.1 | 1.3 | 6.5 | 19.8 | 44.4 | 77.0 | 96.6 |
| 1999..... | 11.4 | * | * | * | * | 0.1 | 1.4 | 6.6 | 20.2 | 45.6 | 73.5 | 100.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 12.1 | * | * | * | * | 0.2 | 1.6 | 8.4 | 24.6 | 53.2 | 85.4 | 104.1 |
| 2003..... | 11.7 | * | * | * | * | 0.2 | 1.6 | 8.0 | 23.5 | 51.6 | 86.2 | 96.5 |
| 2002..... | 11.7 | * | * | * | * | 0.1 | 1.7 | 8.0 | 24.1 | 52.1 | 85.3 | 102.4 |
| 2001..... | 11.4 | * | * | * | * | 0.2 | 1.5 | 7.9 | 24.2 | 51.2 | 84.6 | 106.4 |
| 2000..... | 11.3 | * | * | * | * | 0.2 | 1.5 | 8.3 | 24.1 | 50.7 | 86.7 | 103.5 |
| 1999..... | 11.4 | * | * | * | * | 0.2 | 1.7 | 8.4 | 24.4 | 52.9 | 82.7 | 108.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 11.8 | * | * | * | * | * | 1.0 | 4.8 | 16.1 | 37.9 | 70.6 | 96.5 |
| 2003..... | 11.6 | * | * | * | * | 0.1 | 1.0 | 4.9 | 15.6 | 39.1 | 69.3 | 95.3 |
| 2002..... | 11.5 | * | * | * | * | * | 1.1 | 4.6 | 15.4 | 39.8 | 69.5 | 91.7 |
| 2001..... | 11.5 | * | * | * | * | 0.1 | 1.2 | 4.9 | 15.4 | 39.8 | 68.0 | 98.2 |
| 2000..... | 11.4 | * | * | * | * | * | 1.0 | 4.7 | 15.8 | 39.2 | 70.6 | 93.8 |
| 1999..... | 11.4 | * | * | * | * | * | 1.0 | 4.8 | 16.4 | 39.6 | 67.6 | 96.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 12.8 | * | * | * | * | 0.2 | 1.2 | 6.3 | 19.5 | 44.0 | 76.1 | 98.1 |
| 2003..... | 12.5 | * | * | * | * | 0.2 | 1.2 | 6.0 | 18.7 | 44.1 | 75.4 | 94.9 |
| 2002..... | 12.4 | * | * | * | * | 0.1 | 1.4 | 6.0 | 19.0 | 44.3 | 75.1 | 94.6 |
| 2001..... | 12.3 | * | * | * | * | 0.1 | 1.3 | 6.1 | 19.3 | 44.1 | 73.7 | 99.4 |
| 2000..... | 12.2 | * | * | * | * | 0.2 | 1.2 | 6.3 | 19.2 | 43.4 | 75.7 | 94.7 |
| 1999..... | 12.1 | * | * | * | * | 0.1 | 1.3 | 6.3 | 19.5 | 44.7 | 72.1 | 98.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 13.1 | * | * | * | * | 0.2 | 1.5 | 8.1 | 23.8 | 52.0 | 85.8 | 105.0 |
| 2003..... | 12.6 | * | * | * | * | 0.2 | 1.5 | 7.5 | 22.8 | 51.4 | 86.0 | 96.8 |
| 2002..... | 12.6 | * | * | * | * | * | 1.7 | 7.7 | 23.5 | 51.2 | 85.4 | 102.9 |
| 2001..... | 12.4 | * | * | * | * | 0.2 | 1.5 | 7.7 | 24.0 | 50.3 | 84.5 | 108.0 |
| 2000..... | 12.2 | * | * | * | * | 0.2 | 1.5 | 8.0 | 23.4 | 50.0 | 86.1 | 103.0 |
| 1999..... | 12.1 | * | * | * | * | 0.2 | 1.6 | 8.0 | 23.4 | 52.1 | 81.6 | 108.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 12.6 | * | * | * | * | * | 0.9 | 4.5 | 15.4 | 37.1 | 69.5 | 95.1 |
| 2003..... | 12.4 | * | * | * | * | * | 1.0 | 4.6 | 14.8 | 38.0 | 68.1 | 94.0 |
| 2002..... | 12.2 | * | * | * | * | * | 1.1 | 4.3 | 14.7 | 38.3 | 68.1 | 91.0 |
| 2001..... | 12.3 | * | * | * | * | * | 1.2 | 4.6 | 14.9 | 38.8 | 66.6 | 95.8 |
| 2000..... | 12.2 | * | * | * | * | * | 1.0 | 4.5 | 15.3 | 37.9 | 68.9 | 91.4 |
| 1999..... | 12.1 | * | * | * | * | * | 1.0 | 4.7 | 15.7 | 38.5 | 65.9 | 95.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of pancreas (C25) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 9.8 | * | * | * | * | * | 1.7 | 9.5 | 28.8 | 59.0 | 89.1 | 115.1 |
| 2003..... | 9.8 | * | * | * | * | * | 1.9 | 10.3 | 27.7 | 57.5 | 94.0 | 113.3 |
| 2002..... | 9.6 | * | * | * | * | * | 1.8 | 8.9 | 28.3 | 61.7 | 90.9 | 108.3 |
| 2001..... | 9.4 | * | * | * | * | * | 1.6 | 9.2 | 25.9 | 59.0 | 93.3 | 122.0 |
| 2000..... | 9.7 | * | * | * | * | * | 1.7 | 9.2 | 27.9 | 60.0 | 98.9 | 121.2 |
| 1999..... | 9.7 | * | * | * | * | * | 2.0 | 9.9 | 28.8 | 60.4 | 95.0 | 119.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 9.8 | * | * | * | * | * | 2.1 | 11.6 | 36.5 | 73.8 | 91.3 | 102.6 |
| 2003..... | 9.6 | * | * | * | * | * | 2.5 | 13.6 | 33.7 | 64.8 | 99.0 | 106.5 |
| 2002..... | 9.3 | * | * | * | * | * | 2.2 | 11.6 | 34.9 | 69.5 | 97.3 | 102.8 |
| 2001..... | 8.9 | * | * | * | * | * | 1.8 | 11.4 | 30.8 | 68.2 | 100.5 | 100.5 |
| 2000..... | 9.3 | * | * | * | * | * | 2.1 | 12.1 | 34.6 | 66.9 | 108.0 | 110.4 |
| 1999..... | 9.7 | * | * | * | * | * | 2.4 | 13.6 | 37.4 | 68.7 | 107.6 | 115.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 9.9 | * | * | * | * | * | 1.2 | 7.7 | 22.6 | 48.5 | 87.9 | 120.1 |
| 2003..... | 10.0 | * | * | * | * | * | 1.3 | 7.4 | 22.8 | 52.4 | 91.3 | 116.0 |
| 2002..... | 9.8 | * | * | * | * | * | 1.5 | 6.5 | 23.0 | 56.2 | 87.2 | 110.4 |
| 2001..... | 9.9 | * | * | * | * | * | 1.4 | 7.3 | 21.9 | 52.4 | 89.3 | 130.0 |
| 2000..... | 10.0 | * | * | * | * | * | 1.5 | 6.8 | 22.5 | 55.1 | 93.8 | 125.2 |
| 1999..... | 9.7 | * | * | * | * | * | 1.7 | 6.7 | 21.9 | 54.6 | 87.9 | 120.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other and ill-defined digestive organs (C26) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 2.6 | 5.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.4 | 7.0 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 3.7 | 7.0 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 | 3.7 | 7.8 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 1.7 | 3.5 | 7.9 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 4.0 | 9.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 3.1 | 5.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 4.1 | 7.4 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 4.8 | 6.5 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.9 | 2.1 | 4.2 | 8.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 1.0 | 2.3 | 4.1 | 7.5 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.3 | 5.1 | 10.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.3 | 5.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 2.9 | 6.8 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 3.0 | 7.3 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 3.3 | 7.4 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 3.1 | 8.0 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 3.3 | 8.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 2.7 | 5.4 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.3 | 7.0 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 3.8 | 7.1 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 3.6 | 7.7 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.8 | 1.6 | 3.4 | 7.9 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.0 | 4.1 | 9.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 3.0 | 5.5 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 4.0 | 7.1 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 4.8 | 6.3 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.9 | 2.1 | 4.2 | 8.5 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.0 | 2.1 | 4.2 | 7.6 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 5.2 | 10.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 5.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 1.3 | 2.8 | 7.0 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 0.5 | 1.0 | 3.0 | 7.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.5 | 3.2 | 7.4 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.0 | 8.0 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 3.4 | 8.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of respiratory and intrathoracic organs (C30-C39) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 55.4 | * | * | * | 0.1 | 0.4 | 5.8 | 31.3 | 110.7 | 272.3 | 382.0 | 306.3 |
| 2003..... | 56.0 | * | * | * | 0.1 | 0.4 | 5.9 | 31.6 | 115.0 | 276.6 | 386.7 | 307.9 |
| 2002..... | 56.3 | * | * | * | 0.1 | 0.5 | 6.3 | 31.7 | 119.3 | 282.1 | 386.9 | 305.1 |
| 2001..... | 56.5 | * | * | * | 0.1 | 0.5 | 6.6 | 32.2 | 121.9 | 287.3 | 380.6 | 312.2 |
| 2000..... | 57.0 | * | * | * | 0.1 | 0.6 | 6.4 | 33.0 | 126.7 | 291.9 | 380.7 | 310.8 |
| 1999..... | 56.2 | * | * | * | 0.1 | 0.5 | 6.4 | 32.7 | 127.9 | 289.4 | 371.2 | 306.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 64.4 | * | * | * | 0.1 | 0.5 | 6.2 | 37.9 | 137.8 | 349.2 | 532.5 | 485.7 |
| 2003..... | 65.4 | * | * | * | 0.1 | 0.5 | 6.5 | 38.6 | 143.3 | 358.8 | 541.5 | 493.3 |
| 2002..... | 66.1 | * | * | * | 0.2 | 0.6 | 6.6 | 38.8 | 150.7 | 367.5 | 544.9 | 498.8 |
| 2001..... | 67.2 | * | * | * | 0.1 | 0.4 | 7.2 | 39.7 | 154.0 | 382.6 | 546.1 | 534.1 |
| 2000..... | 68.1 | * | * | * | 0.1 | 0.6 | 7.3 | 40.7 | 161.1 | 390.8 | 550.4 | 541.7 |
| 1999..... | 68.0 | * | * | * | * | 0.6 | 7.1 | 40.6 | 165.3 | 390.0 | 544.0 | 550.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 46.7 | * | * | * | * | 0.3 | 5.4 | 24.9 | 85.5 | 207.8 | 280.7 | 225.6 |
| 2003..... | 46.9 | * | * | * | * | 0.4 | 5.3 | 24.9 | 88.8 | 207.9 | 283.2 | 225.9 |
| 2002..... | 46.8 | * | * | * | * | 0.4 | 6.0 | 24.9 | 90.3 | 211.1 | 282.0 | 221.1 |
| 2001..... | 46.1 | * | * | * | * | 0.5 | 5.9 | 25.0 | 92.3 | 208.3 | 271.8 | 219.3 |
| 2000..... | 46.2 | * | * | * | * | 0.5 | 5.5 | 25.6 | 94.9 | 210.4 | 270.0 | 216.8 |
| 1999..... | 45.0 | * | * | * | * | 0.5 | 5.7 | 25.1 | 93.4 | 207.0 | 258.9 | 208.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 10.8 | * | * | * | * | 0.4 | 1.5 | 10.2 | 37.7 | 108.2 | 187.3 | 202.2 |
| 2003..... | 10.9 | * | * | * | * | 0.3 | 1.7 | 11.0 | 41.0 | 110.1 | 191.1 | 198.0 |
| 2002..... | 11.0 | * | * | * | * | 0.4 | 1.8 | 10.8 | 43.3 | 115.7 | 191.0 | 199.0 |
| 2001..... | 10.9 | * | * | * | * | * | 2.1 | 11.6 | 43.6 | 114.7 | 190.1 | 210.7 |
| 2000..... | 10.9 | * | * | * | * | 0.3 | 1.9 | 11.6 | 41.7 | 116.0 | 206.6 | 222.3 |
| 1999..... | 11.1 | * | * | * | * | 0.4 | 2.2 | 11.5 | 46.5 | 118.8 | 197.7 | 240.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 13.7 | * | * | * | * | * | 1.9 | 13.9 | 54.4 | 163.6 | 296.9 | 305.4 |
| 2003..... | 13.7 | * | * | * | * | * | 1.8 | 14.0 | 60.7 | 170.5 | 291.1 | 312.5 |
| 2002..... | 14.0 | * | * | * | * | * | 1.9 | 14.0 | 65.3 | 170.1 | 312.7 | 321.3 |
| 2001..... | 14.1 | * | * | * | * | * | 2.7 | 15.7 | 64.0 | 175.9 | 307.6 | 356.2 |
| 2000..... | 14.1 | * | * | * | * | * | 2.3 | 16.1 | 63.1 | 176.4 | 343.2 | 391.0 |
| 1999..... | 14.6 | * | * | * | * | * | 2.6 | 16.2 | 72.2 | 186.5 | 333.2 | 396.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.7 | * | * | * | * | * | 1.1 | 6.4 | 22.6 | 63.7 | 111.6 | 149.1 |
| 2003..... | 7.9 | * | * | * | * | * | 1.5 | 8.0 | 23.2 | 61.9 | 122.5 | 139.8 |
| 2002..... | 7.9 | * | * | * | * | * | 1.8 | 7.6 | 23.7 | 72.4 | 107.6 | 138.4 |
| 2001..... | 7.6 | * | * | * | * | * | 1.6 | 7.6 | 25.6 | 66.5 | 110.6 | 139.6 |
| 2000..... | 7.4 | * | * | * | * | * | 1.4 | 7.3 | 22.7 | 68.5 | 114.9 | 139.5 |
| 1999..... | 7.4 | * | * | * | * | * | 1.8 | 7.0 | 23.7 | 65.7 | 107.5 | 165.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of respiratory and intrathoracic organs (C30-C39) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 62.6 | * | * | * | 0.1 | 0.4 | 6.4 | 33.5 | 116.4 | 283.8 | 392.2 | 310.4 |
| 2003..... | 63.0 | * | * | * | 0.1 | 0.5 | 6.5 | 33.7 | 120.7 | 287.8 | 396.1 | 312.0 |
| 2002..... | 63.2 | * | * | * | 0.1 | 0.5 | 7.0 | 33.7 | 125.1 | 292.9 | 395.6 | 308.8 |
| 2001..... | 63.1 | * | * | * | 0.1 | 0.5 | 7.1 | 34.0 | 127.7 | 298.1 | 388.6 | 315.2 |
| 2000..... | 63.4 | * | * | * | 0.1 | 0.6 | 7.0 | 34.8 | 132.7 | 302.0 | 387.4 | 313.2 |
| 1999..... | 62.3 | * | * | * | 0.1 | 0.5 | 6.9 | 34.4 | 133.5 | 298.7 | 377.3 | 308.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 73.0 | * | * | * | * | 0.5 | 6.9 | 40.4 | 144.3 | 361.6 | 544.9 | 493.7 |
| 2003..... | 73.9 | * | * | * | 0.1 | 0.5 | 7.3 | 41.1 | 149.5 | 370.8 | 553.6 | 501.2 |
| 2002..... | 74.5 | * | * | * | 0.2 | 0.6 | 7.4 | 41.2 | 157.0 | 379.8 | 555.2 | 505.7 |
| 2001..... | 75.4 | * | * | * | * | 0.5 | 7.8 | 41.8 | 160.3 | 395.1 | 556.1 | 540.1 |
| 2000..... | 76.1 | * | * | * | 0.1 | 0.7 | 8.0 | 42.8 | 167.8 | 402.7 | 558.2 | 546.6 |
| 1999..... | 75.5 | * | * | * | * | 0.6 | 7.6 | 42.6 | 171.6 | 400.6 | 551.2 | 555.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 52.6 | * | * | * | * | 0.3 | 6.0 | 26.7 | 90.6 | 218.3 | 289.5 | 228.5 |
| 2003..... | 52.7 | * | * | * | * | 0.4 | 5.8 | 26.5 | 94.0 | 218.2 | 291.1 | 228.9 |
| 2002..... | 52.4 | * | * | * | * | 0.4 | 6.6 | 26.5 | 95.5 | 220.4 | 289.8 | 223.9 |
| 2001..... | 51.5 | * | * | * | * | 0.5 | 6.5 | 26.5 | 97.5 | 217.4 | 278.7 | 221.7 |
| 2000..... | 51.3 | * | * | * | * | 0.5 | 6.0 | 27.2 | 100.3 | 218.9 | 276.1 | 218.8 |
| 1999..... | 49.8 | * | * | * | * | 0.5 | 6.1 | 26.6 | 98.4 | 215.1 | 264.4 | 209.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 68.9 | * | * | * | 0.1 | 0.4 | 6.4 | 32.2 | 116.1 | 289.7 | 397.6 | 315.0 |
| 2003..... | 69.3 | * | * | * | 0.1 | 0.5 | 6.4 | 32.2 | 120.8 | 293.6 | 401.1 | 315.5 |
| 2002..... | 69.5 | * | * | * | 0.1 | 0.5 | 6.8 | 32.4 | 124.3 | 298.9 | 401.4 | 312.8 |
| 2001..... | 69.2 | * | * | * | * | 0.5 | 7.0 | 32.7 | 128.0 | 302.4 | 391.9 | 317.9 |
| 2000..... | 69.3 | * | * | * | 0.1 | 0.6 | 6.5 | 33.2 | 132.9 | 305.4 | 391.3 | 315.7 |
| 1999..... | 67.8 | * | * | * | * | 0.6 | 6.5 | 32.8 | 133.0 | 301.2 | 379.3 | 309.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 79.1 | * | * | * | * | 0.5 | 6.6 | 37.7 | 139.8 | 362.2 | 545.0 | 498.7 |
| 2003..... | 80.0 | * | * | * | * | 0.6 | 7.0 | 38.4 | 145.4 | 371.0 | 552.4 | 504.2 |
| 2002..... | 80.6 | * | * | * | * | 0.6 | 6.9 | 38.4 | 151.7 | 380.6 | 554.8 | 510.5 |
| 2001..... | 81.2 | * | * | * | * | 0.5 | 7.4 | 38.9 | 156.7 | 392.7 | 552.1 | 541.6 |
| 2000..... | 81.8 | * | * | * | * | 0.7 | 7.1 | 39.5 | 163.8 | 399.3 | 555.4 | 548.0 |
| 1999..... | 80.9 | * | * | * | * | 0.6 | 7.1 | 39.2 | 166.8 | 396.5 | 547.1 | 556.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 59.0 | * | * | * | * | 0.3 | 6.1 | 26.9 | 93.6 | 227.4 | 297.2 | 232.8 |
| 2003..... | 59.0 | * | * | * | * | 0.4 | 5.9 | 26.1 | 97.4 | 227.3 | 299.0 | 232.5 |
| 2002..... | 58.8 | * | * | * | * | 0.4 | 6.7 | 26.5 | 98.4 | 229.3 | 298.6 | 227.5 |
| 2001..... | 57.6 | * | * | * | * | 0.5 | 6.5 | 26.7 | 100.9 | 225.7 | 285.7 | 224.8 |
| 2000..... | 57.3 | * | * | * | * | 0.5 | 5.9 | 27.1 | 103.7 | 226.2 | 283.5 | 222.1 |
| 1999..... | 55.2 | * | * | * | * | 0.5 | 6.0 | 26.5 | 101.2 | 221.5 | 269.5 | 211.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of respiratory and intrathoracic organs (C30-C39) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 46.2 | * | * | * | * | 0.5 | 8.4 | 49.3 | 151.5 | 297.0 | 403.6 | 291.6 |
| 2003..... | 46.7 | * | * | * | * | 0.5 | 8.4 | 51.4 | 155.8 | 306.8 | 403.8 | 296.0 |
| 2002..... | 46.9 | * | * | * | * | 0.6 | 9.7 | 51.3 | 164.0 | 311.6 | 400.6 | 280.7 |
| 2001..... | 46.8 | * | * | * | * | * | 9.9 | 50.9 | 158.3 | 324.7 | 404.8 | 291.2 |
| 2000..... | 47.8 | * | * | * | * | 0.7 | 11.1 | 54.9 | 165.3 | 334.4 | 395.7 | 298.9 |
| 1999..... | 47.8 | * | * | * | * | 0.5 | 10.5 | 55.0 | 170.9 | 330.9 | 402.3 | 303.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 59.6 | * | * | * | * | * | 9.6 | 69.1 | 224.7 | 441.2 | 647.7 | 515.2 |
| 2003..... | 60.7 | * | * | * | * | * | 10.3 | 70.2 | 230.1 | 461.4 | 665.4 | 538.0 |
| 2002..... | 61.6 | * | * | * | * | 0.8 | 12.0 | 71.6 | 244.1 | 473.3 | 665.6 | 504.3 |
| 2001..... | 62.7 | * | * | * | * | * | 12.1 | 73.2 | 234.1 | 509.3 | 698.0 | 574.0 |
| 2000..... | 64.2 | * | * | * | * | * | 14.8 | 78.2 | 246.3 | 521.9 | 680.3 | 589.4 |
| 1999..... | 64.5 | * | * | * | * | * | 13.0 | 79.8 | 259.6 | 514.6 | 689.2 | 606.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 34.0 | * | * | * | * | * | 7.3 | 32.3 | 92.8 | 194.2 | 265.8 | 203.2 |
| 2003..... | 34.0 | * | * | * | * | * | 6.8 | 35.3 | 96.1 | 196.9 | 256.0 | 202.2 |
| 2002..... | 33.5 | * | * | * | * | * | 7.7 | 33.9 | 99.7 | 196.8 | 250.7 | 195.2 |
| 2001..... | 32.3 | * | * | * | * | * | 7.9 | 31.7 | 97.4 | 194.5 | 240.2 | 186.5 |
| 2000..... | 32.9 | * | * | * | * | * | 7.9 | 35.0 | 100.2 | 202.4 | 235.5 | 191.0 |
| 1999..... | 32.7 | * | * | * | * | * | 8.2 | 33.7 | 99.8 | 201.2 | 241.6 | 190.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of nasal cavity, middle ear and accessory sinuses (C30-C31) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 0.9 | 1.4 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 0.9 | 1.2 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.2 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.1 | 1.6 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.8 | 1.2 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 1.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.2 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.4 | * |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.0 | 1.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.3 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 1.1 | * |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.2 | 1.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 0.7 | 1.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 0.9 | 1.4 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 1.7 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.8 | 1.3 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 1.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.2 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.3 | * |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.1 | 1.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.3 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.2 | * |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.3 | 0.4 | 0.7 | 1.2 | 1.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 0.8 | 1.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of larynx (C32) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 1.1 | 3.3 | 5.7 | 6.6 | 6.5 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 1.1 | 3.5 | 6.3 | 6.9 | 6.5 |
| 2002..... | 1.4 | * | * | * | * | * | 0.2 | 1.1 | 3.5 | 6.0 | 7.7 | 5.4 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 1.1 | 3.7 | 6.4 | 7.3 | 6.7 |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 1.2 | 3.7 | 6.7 | 8.1 | 6.5 |
| 1999..... | 1.5 | * | * | * | * | * | 0.2 | 1.1 | 3.9 | 6.8 | 7.2 | 6.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | 0.2 | 1.9 | 5.5 | 10.0 | 12.8 | 15.6 |
| 2003..... | 2.3 | * | * | * | * | * | 0.3 | 1.8 | 5.9 | 10.9 | 13.7 | 15.0 |
| 2002..... | 2.3 | * | * | * | * | * | 0.3 | 1.9 | 5.9 | 10.1 | 15.0 | 13.2 |
| 2001..... | 2.4 | * | * | * | * | * | 0.3 | 1.9 | 6.1 | 11.1 | 14.3 | 18.4 |
| 2000..... | 2.4 | * | * | * | * | * | 0.2 | 1.9 | 6.4 | 11.5 | 15.7 | 17.3 |
| 1999..... | 2.4 | * | * | * | * | * | 0.3 | 1.9 | 6.6 | 11.4 | 14.0 | 17.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.4 | 1.3 | 2.1 | 2.4 | 2.5 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.3 | 2.5 | 2.4 | 2.7 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.4 | 1.3 | 2.6 | 2.9 | 2.1 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 2.6 | 2.7 | 1.8 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.2 | 2.8 | 3.1 | 2.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.4 | 3.0 | 2.8 | 2.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.1 | 1.0 | 3.0 | 5.4 | 6.3 | 6.4 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.9 | 3.2 | 6.0 | 6.6 | 6.1 |
| 2002..... | 1.4 | * | * | * | * | * | 0.1 | 0.9 | 3.1 | 5.8 | 7.3 | 5.5 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 0.9 | 3.3 | 6.0 | 6.9 | 6.5 |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 1.0 | 3.2 | 6.2 | 7.7 | 6.5 |
| 1999..... | 1.5 | * | * | * | * | * | 0.1 | 0.9 | 3.5 | 6.4 | 7.1 | 6.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | 0.2 | 1.6 | 4.9 | 9.4 | 12.1 | 15.0 |
| 2003..... | 2.3 | * | * | * | * | * | 0.3 | 1.5 | 5.2 | 9.9 | 12.8 | 14.1 |
| 2002..... | 2.3 | * | * | * | * | * | 0.2 | 1.5 | 5.1 | 9.5 | 14.2 | 13.3 |
| 2001..... | 2.3 | * | * | * | * | * | 0.3 | 1.5 | 5.5 | 10.2 | 13.3 | 18.3 |
| 2000..... | 2.4 | * | * | * | * | * | 0.2 | 1.5 | 5.4 | 10.5 | 15.0 | 17.4 |
| 1999..... | 2.3 | * | * | * | * | * | 0.2 | 1.6 | 5.8 | 10.7 | 13.6 | 16.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.4 | 1.3 | 2.0 | 2.4 | 2.5 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.2 | 2.5 | 2.4 | 2.6 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.3 | 2.6 | 2.7 | 2.2 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.3 | 1.2 | 2.4 | 2.7 | 1.6 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.4 | 1.1 | 2.6 | 3.0 | 2.1 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.3 | 1.3 | 2.9 | 2.9 | 2.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of larynx (C32) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | 2.3 | 6.6 | 10.3 | 11.4 | 8.1 |
| 2003..... | 1.9 | * | * | * | * | * | * | 2.6 | 7.7 | 11.6 | 11.9 | 11.3 |
| 2002..... | 1.9 | * | * | * | * | * | 0.5 | 2.7 | 7.7 | 10.6 | 13.9 | * |
| 2001..... | 1.9 | * | * | * | * | * | 0.4 | 2.9 | 7.9 | 11.5 | 12.0 | 8.4 |
| 2000..... | 2.0 | * | * | * | * | * | * | 2.9 | 9.2 | 12.9 | 14.3 | 7.0 |
| 1999..... | 1.9 | * | * | * | * | * | 0.6 | 3.0 | 8.7 | 11.7 | 10.0 | 8.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | * | * | 4.3 | 12.6 | 20.2 | 25.4 | 23.4 |
| 2003..... | 3.3 | * | * | * | * | * | * | 4.6 | 14.0 | 24.0 | 28.3 | 27.7 |
| 2002..... | 3.1 | * | * | * | * | * | 0.8 | 4.6 | 15.1 | 19.8 | 29.6 | * |
| 2001..... | 3.1 | * | * | * | * | * | * | 5.2 | 14.2 | 21.4 | 28.2 | * |
| 2000..... | 3.4 | * | * | * | * | * | * | 5.2 | 17.5 | 24.6 | 29.0 | * |
| 1999..... | 3.2 | * | * | * | * | * | 1.0 | 5.3 | 16.3 | 22.1 | 22.5 | 25.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 1.7 | 3.1 | 3.5 | * |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.8 | 2.6 | 2.8 | * | * |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.1 | 1.8 | 4.1 | 5.0 | * |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 2.9 | 4.6 | * | * |
| 2000..... | 0.8 | * | * | * | * | * | * | 1.0 | 2.4 | 4.7 | 6.1 | * |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.9 | 2.7 | 4.3 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of trachea, bronchus and lung (C33-C34) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 53.8 | * | * | * | 0.1 | 0.4 | 5.5 | 29.9 | 106.9 | 265.5 | 373.6 | 297.5 |
| 2003..... | 54.4 | * | * | * | 0.1 | 0.4 | 5.6 | 30.3 | 111.0 | 269.3 | 377.8 | 298.9 |
| 2002..... | 54.7 | * | * | * | 0.1 | 0.4 | 6.0 | 30.3 | 115.3 | 275.0 | 377.6 | 297.2 |
| 2001..... | 54.8 | * | * | * | 0.1 | 0.4 | 6.2 | 30.7 | 117.7 | 279.7 | 371.4 | 302.7 |
| 2000..... | 55.3 | * | * | * | 0.1 | 0.5 | 6.1 | 31.6 | 122.4 | 284.2 | 370.8 | 302.1 |
| 1999..... | 54.5 | * | * | * | 0.1 | 0.4 | 6.0 | 31.3 | 123.4 | 281.5 | 362.1 | 297.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 62.0 | * | * | * | * | 0.4 | 5.8 | 35.8 | 131.7 | 337.6 | 517.3 | 466.8 |
| 2003..... | 62.9 | * | * | * | * | 0.4 | 6.1 | 36.5 | 136.7 | 346.6 | 525.1 | 475.1 |
| 2002..... | 63.7 | * | * | * | * | 0.4 | 6.2 | 36.6 | 144.0 | 355.9 | 527.9 | 482.2 |
| 2001..... | 64.7 | * | * | * | * | 0.4 | 6.7 | 37.4 | 147.1 | 370.0 | 529.5 | 512.4 |
| 2000..... | 65.5 | * | * | * | * | 0.5 | 6.9 | 38.5 | 154.0 | 377.9 | 532.2 | 521.2 |
| 1999..... | 65.4 | * | * | * | * | 0.5 | 6.6 | 38.2 | 158.0 | 377.2 | 527.6 | 529.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 45.9 | * | * | * | * | 0.3 | 5.2 | 24.2 | 83.9 | 205.0 | 277.0 | 221.3 |
| 2003..... | 46.1 | * | * | * | * | 0.4 | 5.1 | 24.4 | 87.1 | 204.8 | 279.4 | 221.0 |
| 2002..... | 46.0 | * | * | * | * | 0.4 | 5.8 | 24.3 | 88.8 | 207.7 | 277.8 | 217.0 |
| 2001..... | 45.3 | * | * | * | * | 0.4 | 5.7 | 24.3 | 90.5 | 204.9 | 267.5 | 215.0 |
| 2000..... | 45.4 | * | * | * | * | 0.5 | 5.3 | 25.0 | 93.3 | 206.9 | 265.6 | 212.8 |
| 1999..... | 44.1 | * | * | * | * | 0.4 | 5.5 | 24.6 | 91.5 | 203.1 | 254.5 | 203.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 10.2 | * | * | * | * | 0.3 | 1.4 | 9.4 | 35.4 | 102.6 | 178.7 | 193.1 |
| 2003..... | 10.4 | * | * | * | * | 0.3 | 1.6 | 10.2 | 38.1 | 105.8 | 183.8 | 189.2 |
| 2002..... | 10.4 | * | * | * | * | 0.4 | 1.5 | 9.9 | 40.9 | 110.4 | 183.7 | 189.0 |
| 2001..... | 10.3 | * | * | * | * | * | 1.9 | 10.7 | 41.0 | 108.3 | 182.3 | 200.7 |
| 2000..... | 10.3 | * | * | * | * | * | 1.8 | 10.9 | 39.3 | 110.6 | 198.7 | 213.7 |
| 1999..... | 10.4 | * | * | * | * | * | 2.0 | 10.4 | 44.2 | 112.9 | 187.7 | 227.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 12.7 | * | * | * | * | * | 1.7 | 12.6 | 50.0 | 151.7 | 279.7 | 287.2 |
| 2003..... | 12.8 | * | * | * | * | * | 1.7 | 12.6 | 55.4 | 161.8 | 277.3 | 290.9 |
| 2002..... | 13.1 | * | * | * | * | * | 1.5 | 12.6 | 60.7 | 161.7 | 299.1 | 307.9 |
| 2001..... | 13.1 | * | * | * | * | * | 2.4 | 14.2 | 59.5 | 163.7 | 293.8 | 335.2 |
| 2000..... | 13.3 | * | * | * | * | * | 2.1 | 14.8 | 58.6 | 167.3 | 327.5 | 368.8 |
| 1999..... | 13.6 | * | * | * | * | * | 2.3 | 14.2 | 68.1 | 175.3 | 316.0 | 362.3 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | * | * | 1.0 | 6.2 | 22.1 | 63.1 | 108.9 | 144.8 |
| 2003..... | 7.8 | * | * | * | * | * | 1.4 | 7.9 | 22.4 | 61.2 | 119.6 | 137.5 |
| 2002..... | 7.6 | * | * | * | * | * | 1.6 | 7.3 | 23.2 | 69.5 | 104.6 | 130.1 |
| 2001..... | 7.4 | * | * | * | * | * | 1.4 | 7.3 | 24.6 | 64.6 | 107.0 | 135.0 |
| 2000..... | 7.2 | * | * | * | * | * | 1.4 | 7.1 | 22.2 | 66.0 | 112.3 | 137.5 |
| 1999..... | 7.1 | * | * | * | * | * | 1.7 | 6.7 | 23.1 | 64.0 | 102.3 | 162.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of trachea, bronchus and lung (C33-C34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 60.9 | * | * | * | 0.1 | 0.4 | 6.2 | 32.0 | 112.6 | 277.0 | 383.9 | 301.6 |
| 2003..... | 61.2 | * | * | * | 0.1 | 0.4 | 6.2 | 32.3 | 116.6 | 280.3 | 387.2 | 303.0 |
| 2002..... | 61.4 | * | * | * | * | 0.4 | 6.7 | 32.3 | 121.0 | 285.7 | 386.2 | 301.0 |
| 2001..... | 61.3 | * | * | * | * | 0.4 | 6.8 | 32.5 | 123.3 | 290.4 | 379.4 | 305.8 |
| 2000..... | 61.5 | * | * | * | * | 0.5 | 6.6 | 33.4 | 128.4 | 294.2 | 377.4 | 304.5 |
| 1999..... | 60.5 | * | * | * | * | 0.5 | 6.5 | 33.0 | 128.9 | 290.7 | 368.2 | 298.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 70.4 | * | * | * | * | 0.4 | 6.5 | 38.2 | 138.1 | 350.1 | 529.8 | 474.8 |
| 2003..... | 71.1 | * | * | * | * | 0.4 | 6.8 | 38.9 | 142.9 | 358.3 | 537.2 | 483.1 |
| 2002..... | 71.8 | * | * | * | * | 0.5 | 6.9 | 38.9 | 150.3 | 368.1 | 538.0 | 488.9 |
| 2001..... | 72.6 | * | * | * | * | 0.4 | 7.3 | 39.5 | 153.2 | 382.4 | 539.5 | 518.4 |
| 2000..... | 73.2 | * | * | * | * | 0.6 | 7.5 | 40.5 | 160.6 | 389.7 | 539.9 | 526.2 |
| 1999..... | 72.7 | * | * | * | * | 0.5 | 7.1 | 40.1 | 164.0 | 387.8 | 534.9 | 534.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 51.8 | * | * | * | * | 0.3 | 5.8 | 26.1 | 88.9 | 215.4 | 285.8 | 224.2 |
| 2003..... | 51.8 | * | * | * | * | 0.4 | 5.6 | 26.0 | 92.2 | 214.9 | 287.2 | 223.9 |
| 2002..... | 51.6 | * | * | * | * | 0.4 | 6.4 | 25.9 | 93.9 | 216.9 | 285.5 | 220.0 |
| 2001..... | 50.5 | * | * | * | * | 0.5 | 6.2 | 25.8 | 95.6 | 213.9 | 274.3 | 217.4 |
| 2000..... | 50.4 | * | * | * | * | 0.5 | 5.8 | 26.5 | 98.6 | 215.4 | 271.6 | 214.8 |
| 1999..... | 48.8 | * | * | * | * | 0.4 | 6.0 | 26.1 | 96.5 | 211.0 | 260.0 | 204.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 67.1 | * | * | * | * | 0.3 | 6.1 | 31.0 | 112.5 | 283.2 | 389.6 | 306.3 |
| 2003..... | 67.5 | * | * | * | * | 0.4 | 6.1 | 31.0 | 117.0 | 286.5 | 392.6 | 306.7 |
| 2002..... | 67.7 | * | * | * | * | 0.4 | 6.5 | 31.2 | 120.6 | 292.0 | 392.2 | 304.8 |
| 2001..... | 67.3 | * | * | * | * | 0.5 | 6.6 | 31.5 | 124.0 | 295.2 | 383.0 | 308.6 |
| 2000..... | 67.4 | * | * | * | * | 0.5 | 6.2 | 32.0 | 129.1 | 298.0 | 381.7 | 306.8 |
| 1999..... | 65.9 | * | * | * | * | 0.5 | 6.2 | 31.5 | 128.9 | 293.5 | 370.3 | 300.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 76.5 | * | * | * | * | 0.4 | 6.3 | 35.8 | 134.2 | 351.4 | 530.6 | 480.3 |
| 2003..... | 77.3 | * | * | * | * | 0.4 | 6.5 | 36.5 | 139.4 | 359.6 | 537.0 | 486.8 |
| 2002..... | 77.8 | * | * | * | * | 0.5 | 6.5 | 36.5 | 145.7 | 369.5 | 538.3 | 493.3 |
| 2001..... | 78.4 | * | * | * | * | 0.4 | 6.9 | 37.0 | 150.3 | 381.1 | 536.5 | 520.0 |
| 2000..... | 78.9 | * | * | * | * | 0.6 | 6.7 | 37.7 | 157.7 | 387.3 | 537.7 | 527.3 |
| 1999..... | 78.1 | * | * | * | * | 0.5 | 6.6 | 37.1 | 160.1 | 384.4 | 531.1 | 536.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 58.1 | * | * | * | * | 0.3 | 5.9 | 26.2 | 92.0 | 224.6 | 293.5 | 228.4 |
| 2003..... | 58.1 | * | * | * | * | 0.4 | 5.7 | 25.6 | 95.7 | 223.9 | 295.1 | 227.4 |
| 2002..... | 57.9 | * | * | * | * | 0.3 | 6.5 | 26.0 | 96.8 | 225.9 | 294.3 | 223.4 |
| 2001..... | 56.6 | * | * | * | * | 0.5 | 6.4 | 26.0 | 99.2 | 222.3 | 281.3 | 220.6 |
| 2000..... | 56.4 | * | * | * | * | 0.5 | 5.7 | 26.4 | 102.2 | 222.9 | 279.2 | 218.0 |
| 1999..... | 54.3 | * | * | * | * | 0.5 | 5.8 | 26.1 | 99.5 | 217.5 | 265.0 | 206.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of trachea, bronchus and lung (C33-C34) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 44.2 | * | * | * | * | 0.5 | 8.0 | 46.5 | 144.2 | 285.3 | 390.1 | 281.5 |
| 2003..... | 44.5 | * | * | * | * | * | 8.0 | 48.5 | 147.4 | 293.9 | 389.4 | 283.5 |
| 2002..... | 44.7 | * | * | * | * | 0.4 | 9.0 | 48.1 | 155.6 | 299.5 | 385.6 | 274.9 |
| 2001..... | 44.6 | * | * | * | * | * | 9.2 | 47.5 | 149.4 | 311.4 | 390.9 | 281.6 |
| 2000..... | 45.4 | * | * | * | * | 0.6 | 10.4 | 51.5 | 155.0 | 320.2 | 379.4 | 291.0 |
| 1999..... | 45.6 | * | * | * | * | 0.4 | 9.6 | 51.5 | 161.1 | 317.7 | 390.9 | 293.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 56.3 | * | * | * | * | * | 9.0 | 64.3 | 211.1 | 419.0 | 619.8 | 488.8 |
| 2003..... | 57.0 | * | * | * | * | * | 9.7 | 65.3 | 215.1 | 434.9 | 632.5 | 509.3 |
| 2002..... | 58.1 | * | * | * | * | * | 11.0 | 66.3 | 228.0 | 451.4 | 633.9 | 490.1 |
| 2001..... | 59.2 | * | * | * | * | * | 11.4 | 67.5 | 218.3 | 484.4 | 668.0 | 550.9 |
| 2000..... | 60.3 | * | * | * | * | * | 13.8 | 72.4 | 227.1 | 495.4 | 649.5 | 567.1 |
| 1999..... | 60.9 | * | * | * | * | * | 11.7 | 73.7 | 241.9 | 491.0 | 665.1 | 580.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 33.2 | * | * | * | * | * | 7.0 | 31.3 | 90.5 | 190.0 | 260.5 | 199.6 |
| 2003..... | 33.1 | * | * | * | * | * | 6.5 | 34.1 | 93.1 | 193.5 | 252.1 | 196.0 |
| 2002..... | 32.6 | * | * | * | * | * | 7.3 | 32.5 | 97.5 | 191.8 | 245.1 | 192.7 |
| 2001..... | 31.3 | * | * | * | * | * | 7.3 | 30.4 | 94.0 | 189.4 | 235.3 | 181.8 |
| 2000..... | 31.9 | * | * | * | * | * | 7.4 | 33.6 | 97.0 | 196.9 | 227.3 | 188.4 |
| 1999..... | 31.8 | * | * | * | * | * | 7.9 | 32.5 | 96.4 | 195.5 | 237.4 | 186.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of thymus, heart, mediastinum and pleura (C37-C38) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 0.9 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 1.3 |
| 2002..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.6 | 0.8 | 1.1 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.0 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.0 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 0.9 | 1.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 1.4 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 1.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.0 | 1.7 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 0.9 | * |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | * |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.0 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.5 | 0.7 | 0.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.7 | 0.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | * |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.8 | 0.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.8 | 0.9 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.8 | 1.4 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 0.8 | 1.2 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.0 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 1.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 1.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.0 | 1.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 0.9 | * |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | * |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.6 | * |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.6 | 1.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.5 | 0.7 | 0.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.6 | 0.9 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | * |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 0.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of bone and articular cartilage (C40-C41) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | 0.2 | 0.5 | 0.2 | 0.2 | 0.3 | 0.6 | 1.0 | 1.8 | 2.8 |
| 2003..... | 0.5 | * | * | 0.2 | 0.5 | 0.2 | 0.2 | 0.3 | 0.5 | 0.9 | 1.9 | 2.9 |
| 2002..... | 0.4 | * | * | 0.2 | 0.4 | 0.3 | 0.2 | 0.3 | 0.6 | 0.8 | 1.6 | 2.9 |
| 2001..... | 0.5 | * | * | 0.2 | 0.5 | 0.3 | 0.2 | 0.3 | 0.5 | 1.1 | 1.9 | 3.5 |
| 2000..... | 0.4 | * | * | 0.2 | 0.4 | 0.2 | 0.2 | 0.3 | 0.6 | 0.9 | 1.9 | 3.4 |
| 1999..... | 0.5 | * | * | 0.2 | 0.5 | 0.2 | 0.2 | 0.4 | 0.4 | 1.0 | 1.8 | 2.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | 0.2 | 0.6 | 0.3 | 0.3 | 0.4 | 0.8 | 1.2 | 2.0 | 3.3 |
| 2003..... | 0.5 | * | * | 0.2 | 0.6 | 0.2 | 0.3 | 0.4 | 0.6 | 1.1 | 2.4 | 3.6 |
| 2002..... | 0.5 | * | * | 0.2 | 0.5 | 0.3 | 0.3 | 0.4 | 0.6 | 0.9 | 2.1 | 3.6 |
| 2001..... | 0.5 | * | * | 0.2 | 0.7 | 0.3 | 0.3 | 0.4 | 0.6 | 1.4 | 2.4 | 3.7 |
| 2000..... | 0.5 | * | * | 0.2 | 0.6 | 0.3 | 0.2 | 0.4 | 0.7 | 1.2 | 2.4 | 4.2 |
| 1999..... | 0.5 | * | * | 0.2 | 0.6 | 0.2 | 0.2 | 0.4 | 0.6 | 1.3 | 1.9 | 4.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.1 | 0.3 | 0.2 | 0.2 | 0.2 | 0.5 | 0.9 | 1.6 | 2.5 |
| 2003..... | 0.4 | * | * | 0.1 | 0.3 | 0.2 | 0.1 | 0.2 | 0.3 | 0.8 | 1.5 | 2.5 |
| 2002..... | 0.4 | * | * | 0.2 | 0.3 | 0.2 | 0.1 | 0.2 | 0.5 | 0.8 | 1.3 | 2.6 |
| 2001..... | 0.4 | * | * | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.5 | 0.8 | 1.5 | 3.4 |
| 2000..... | 0.4 | * | * | 0.2 | 0.3 | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.0 |
| 1999..... | 0.4 | * | * | 0.2 | 0.4 | 0.2 | 0.2 | 0.3 | 0.3 | 0.8 | 1.7 | 2.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | 0.2 | 0.5 | 0.3 | 0.2 | 0.3 | 0.6 | 1.1 | 1.8 | 2.9 |
| 2003..... | 0.5 | * | * | 0.2 | 0.5 | 0.2 | 0.2 | 0.3 | 0.5 | 0.9 | 1.9 | 2.9 |
| 2002..... | 0.5 | * | * | 0.2 | 0.5 | 0.3 | 0.2 | 0.3 | 0.6 | 0.8 | 1.7 | 3.0 |
| 2001..... | 0.5 | * | * | 0.2 | 0.6 | 0.3 | 0.2 | 0.3 | 0.5 | 1.0 | 2.0 | 3.6 |
| 2000..... | 0.5 | * | * | 0.2 | 0.4 | 0.2 | 0.2 | 0.3 | 0.5 | 1.0 | 1.9 | 3.5 |
| 1999..... | 0.5 | * | * | 0.2 | 0.5 | 0.2 | 0.2 | 0.4 | 0.4 | 1.0 | 1.9 | 3.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | 0.2 | 0.6 | 0.3 | 0.2 | 0.4 | 0.7 | 1.2 | 2.1 | 3.4 |
| 2003..... | 0.6 | * | * | 0.3 | 0.6 | 0.3 | 0.3 | 0.4 | 0.6 | 1.1 | 2.5 | 3.5 |
| 2002..... | 0.5 | * | * | 0.2 | 0.5 | 0.4 | 0.3 | 0.4 | 0.6 | 0.9 | 2.1 | 3.8 |
| 2001..... | 0.6 | * | * | 0.2 | 0.8 | 0.3 | 0.3 | 0.4 | 0.6 | 1.4 | 2.5 | 3.6 |
| 2000..... | 0.6 | * | * | 0.2 | 0.5 | 0.4 | 0.2 | 0.4 | 0.7 | 1.2 | 2.4 | 4.5 |
| 1999..... | 0.5 | * | * | 0.2 | 0.6 | 0.3 | 0.2 | 0.4 | 0.5 | 1.3 | 2.0 | 4.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.3 | 0.2 | 0.1 | 0.2 | 0.5 | 0.9 | 1.6 | 2.6 |
| 2003..... | 0.4 | * | * | * | 0.4 | 0.2 | 0.2 | 0.3 | 0.3 | 0.8 | 1.4 | 2.7 |
| 2002..... | 0.4 | * | * | 0.2 | 0.4 | 0.2 | * | 0.1 | 0.5 | 0.8 | 1.4 | 2.7 |
| 2001..... | 0.4 | * | * | 0.2 | 0.3 | 0.3 | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.5 |
| 2000..... | 0.4 | * | * | 0.2 | 0.3 | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.6 | 3.1 |
| 1999..... | 0.4 | * | * | * | 0.5 | 0.2 | 0.2 | 0.3 | 0.3 | 0.8 | 1.8 | 2.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Melanoma and other malignant neoplasms of skin (C43-C44) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | 0.1 | 0.6 | 1.4 | 3.5 | 7.0 | 12.4 | 20.6 | 31.1 |
| 2003..... | 4.0 | * | * | * | 0.1 | 0.6 | 1.6 | 3.6 | 7.2 | 11.8 | 20.4 | 31.4 |
| 2002..... | 3.9 | * | * | * | 0.1 | 0.6 | 1.6 | 3.6 | 6.7 | 12.1 | 20.0 | 30.4 |
| 2001..... | 3.9 | * | * | * | 0.1 | 0.6 | 1.6 | 3.9 | 7.2 | 12.6 | 19.2 | 31.8 |
| 2000..... | 3.8 | * | * | * | 0.1 | 0.6 | 1.6 | 3.9 | 7.0 | 12.3 | 18.5 | 32.5 |
| 1999..... | 3.8 | * | * | * | 0.1 | 0.6 | 1.7 | 3.8 | 7.0 | 12.2 | 19.2 | 29.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | 0.7 | 1.8 | 4.7 | 10.4 | 18.9 | 33.8 | 56.1 |
| 2003..... | 5.3 | * | * | * | 0.1 | 0.8 | 1.9 | 4.9 | 10.4 | 17.6 | 33.2 | 54.7 |
| 2002..... | 5.1 | * | * | * | * | 0.7 | 2.0 | 5.0 | 9.8 | 18.3 | 32.0 | 51.8 |
| 2001..... | 5.2 | * | * | * | 0.1 | 0.7 | 2.0 | 5.2 | 10.4 | 19.5 | 30.9 | 55.4 |
| 2000..... | 5.0 | * | * | * | 0.1 | 0.7 | 1.9 | 5.2 | 9.9 | 18.5 | 30.3 | 57.0 |
| 1999..... | 5.0 | * | * | * | 0.1 | 0.7 | 2.2 | 5.2 | 10.2 | 18.5 | 30.7 | 51.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 0.1 | 0.5 | 1.1 | 2.3 | 3.9 | 6.9 | 11.7 | 19.9 |
| 2003..... | 2.7 | * | * | * | * | 0.4 | 1.2 | 2.4 | 4.2 | 7.0 | 11.8 | 21.1 |
| 2002..... | 2.7 | * | * | * | * | 0.5 | 1.2 | 2.2 | 3.9 | 7.0 | 12.0 | 21.2 |
| 2001..... | 2.7 | * | * | * | * | 0.5 | 1.2 | 2.6 | 4.2 | 7.0 | 11.5 | 22.0 |
| 2000..... | 2.7 | * | * | * | * | 0.5 | 1.3 | 2.6 | 4.3 | 7.2 | 10.8 | 22.6 |
| 1999..... | 2.6 | * | * | * | * | 0.5 | 1.1 | 2.5 | 4.0 | 7.0 | 11.8 | 20.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | 0.1 | 0.8 | 1.8 | 4.1 | 8.0 | 14.0 | 22.6 | 33.8 |
| 2003..... | 4.8 | * | * | * | 0.1 | 0.8 | 1.9 | 4.2 | 8.2 | 13.3 | 22.3 | 34.0 |
| 2002..... | 4.7 | * | * | * | 0.1 | 0.8 | 2.0 | 4.1 | 7.6 | 13.4 | 21.9 | 32.9 |
| 2001..... | 4.8 | * | * | * | 0.2 | 0.7 | 2.0 | 4.5 | 8.1 | 14.0 | 20.9 | 34.4 |
| 2000..... | 4.6 | * | * | * | 0.2 | 0.7 | 1.9 | 4.5 | 7.8 | 13.6 | 20.0 | 35.0 |
| 1999..... | 4.6 | * | * | * | 0.1 | 0.7 | 2.0 | 4.4 | 7.9 | 13.5 | 20.8 | 32.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.6 | * | * | * | * | 0.9 | 2.1 | 5.5 | 11.6 | 21.3 | 36.9 | 61.5 |
| 2003..... | 6.4 | * | * | * | 0.2 | 1.0 | 2.3 | 5.6 | 11.7 | 19.7 | 36.3 | 59.6 |
| 2002..... | 6.2 | * | * | * | * | 0.9 | 2.4 | 5.7 | 10.8 | 20.0 | 35.0 | 56.3 |
| 2001..... | 6.3 | * | * | * | 0.2 | 0.8 | 2.4 | 5.9 | 11.6 | 21.4 | 33.7 | 60.4 |
| 2000..... | 6.0 | * | * | * | 0.2 | 0.8 | 2.2 | 6.0 | 11.0 | 20.3 | 32.7 | 62.5 |
| 1999..... | 6.0 | * | * | * | * | 0.8 | 2.6 | 5.8 | 11.4 | 20.5 | 33.1 | 56.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.6 | 1.4 | 2.8 | 4.5 | 7.8 | 12.8 | 21.4 |
| 2003..... | 3.3 | * | * | * | * | 0.6 | 1.4 | 2.8 | 4.7 | 7.9 | 12.8 | 22.7 |
| 2002..... | 3.3 | * | * | * | * | 0.6 | 1.5 | 2.6 | 4.5 | 7.8 | 13.1 | 22.8 |
| 2001..... | 3.3 | * | * | * | * | 0.6 | 1.5 | 3.1 | 4.8 | 7.7 | 12.4 | 23.6 |
| 2000..... | 3.3 | * | * | * | * | 0.6 | 1.6 | 3.1 | 4.8 | 7.9 | 11.7 | 24.0 |
| 1999..... | 3.2 | * | * | * | * | 0.6 | 1.4 | 3.0 | 4.6 | 7.8 | 12.7 | 22.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant melanoma of skin (C43) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | 0.1 | 0.6 | 1.3 | 3.0 | 5.6 | 9.8 | 15.3 | 18.3 |
| 2003..... | 3.0 | * | * | * | 0.1 | 0.6 | 1.4 | 3.1 | 5.9 | 9.1 | 15.1 | 18.1 |
| 2002..... | 2.9 | * | * | * | 0.1 | 0.6 | 1.5 | 3.0 | 5.4 | 9.3 | 14.1 | 18.2 |
| 2001..... | 3.0 | * | * | * | 0.1 | 0.5 | 1.5 | 3.3 | 5.6 | 9.7 | 13.7 | 18.3 |
| 2000..... | 3.0 | * | * | * | 0.1 | 0.5 | 1.5 | 3.4 | 5.7 | 9.3 | 13.5 | 19.3 |
| 1999..... | 2.9 | * | * | * | 0.1 | 0.6 | 1.5 | 3.3 | 5.6 | 9.3 | 13.7 | 16.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 0.7 | 1.6 | 3.9 | 8.0 | 14.5 | 24.8 | 32.7 |
| 2003..... | 4.0 | * | * | * | 0.1 | 0.7 | 1.7 | 4.0 | 8.2 | 13.2 | 24.2 | 30.9 |
| 2002..... | 3.8 | * | * | * | * | 0.7 | 1.8 | 4.1 | 7.6 | 13.7 | 22.0 | 30.2 |
| 2001..... | 3.9 | * | * | * | 0.1 | 0.6 | 1.9 | 4.2 | 7.8 | 14.6 | 22.0 | 31.0 |
| 2000..... | 3.7 | * | * | * | 0.1 | 0.6 | 1.7 | 4.4 | 7.8 | 13.4 | 21.3 | 32.6 |
| 1999..... | 3.7 | * | * | * | 0.1 | 0.6 | 2.0 | 4.2 | 7.8 | 13.7 | 21.5 | 28.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | 0.5 | 1.1 | 2.1 | 3.4 | 5.8 | 9.0 | 11.9 |
| 2003..... | 2.2 | * | * | * | * | 0.4 | 1.1 | 2.2 | 3.6 | 5.7 | 9.0 | 12.4 |
| 2002..... | 2.1 | * | * | * | * | 0.4 | 1.2 | 2.0 | 3.3 | 5.7 | 8.9 | 12.9 |
| 2001..... | 2.1 | * | * | * | * | 0.5 | 1.2 | 2.4 | 3.5 | 5.6 | 8.3 | 12.9 |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 1.3 | 2.4 | 3.7 | 5.9 | 8.4 | 13.9 |
| 1999..... | 2.1 | * | * | * | * | 0.5 | 1.1 | 2.3 | 3.5 | 5.7 | 8.7 | 12.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.1 | 0.8 | 1.7 | 3.6 | 6.6 | 11.2 | 16.9 | 19.9 |
| 2003..... | 3.8 | * | * | * | 0.1 | 0.7 | 1.8 | 3.7 | 6.8 | 10.4 | 16.6 | 19.6 |
| 2002..... | 3.6 | * | * | * | 0.1 | 0.7 | 1.8 | 3.6 | 6.2 | 10.5 | 15.6 | 19.6 |
| 2001..... | 3.7 | * | * | * | 0.2 | 0.7 | 1.9 | 4.0 | 6.5 | 10.8 | 15.0 | 19.9 |
| 2000..... | 3.6 | * | * | * | 0.2 | 0.7 | 1.8 | 4.0 | 6.5 | 10.4 | 14.7 | 20.8 |
| 1999..... | 3.5 | * | * | * | 0.1 | 0.7 | 1.9 | 3.9 | 6.4 | 10.5 | 14.9 | 17.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | * | 0.9 | 2.0 | 4.7 | 9.3 | 16.6 | 27.2 | 35.8 |
| 2003..... | 4.9 | * | * | * | 0.2 | 0.9 | 2.1 | 4.8 | 9.4 | 14.9 | 26.5 | 33.8 |
| 2002..... | 4.7 | * | * | * | * | 0.9 | 2.2 | 4.8 | 8.6 | 15.3 | 24.1 | 32.9 |
| 2001..... | 4.8 | * | * | * | 0.2 | 0.8 | 2.3 | 5.0 | 9.0 | 16.3 | 24.1 | 34.1 |
| 2000..... | 4.6 | * | * | * | 0.2 | 0.8 | 2.0 | 5.2 | 8.9 | 15.0 | 23.2 | 35.7 |
| 1999..... | 4.5 | * | * | * | * | 0.8 | 2.4 | 5.0 | 8.9 | 15.4 | 23.3 | 30.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | 0.6 | 1.4 | 2.5 | 4.0 | 6.6 | 9.9 | 12.8 |
| 2003..... | 2.6 | * | * | * | * | 0.5 | 1.4 | 2.6 | 4.2 | 6.4 | 9.8 | 13.3 |
| 2002..... | 2.6 | * | * | * | * | 0.6 | 1.4 | 2.4 | 3.9 | 6.5 | 9.8 | 13.8 |
| 2001..... | 2.6 | * | * | * | * | 0.6 | 1.5 | 2.9 | 4.1 | 6.2 | 9.0 | 13.9 |
| 2000..... | 2.7 | * | * | * | * | 0.6 | 1.6 | 2.8 | 4.2 | 6.6 | 9.1 | 14.7 |
| 1999..... | 2.5 | * | * | * | * | 0.6 | 1.4 | 2.8 | 4.1 | 6.4 | 9.4 | 12.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other malignant neoplasms of skin (C44) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 2.6 | 5.2 | 12.8 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 2.7 | 5.3 | 13.3 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 2.8 | 5.9 | 12.3 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 3.0 | 5.5 | 13.5 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 3.0 | 5.0 | 13.2 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 2.9 | 5.5 | 12.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 4.4 | 9.0 | 23.4 |
| 2003..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.2 | 4.5 | 9.0 | 23.8 |
| 2002..... | 1.3 | * | * | * | * | * | 0.2 | 0.9 | 2.2 | 4.6 | 10.0 | 21.6 |
| 2001..... | 1.3 | * | * | * | * | * | 0.1 | 0.9 | 2.6 | 4.9 | 8.9 | 24.4 |
| 2000..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 5.0 | 8.9 | 24.4 |
| 1999..... | 1.3 | * | * | * | * | * | 0.3 | 0.9 | 2.4 | 4.8 | 9.2 | 23.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.7 | 8.0 | |
| 2003..... | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.8 | 8.7 | |
| 2002..... | 0.6 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.1 | 8.2 | |
| 2001..... | 0.6 | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 3.3 | 9.0 | |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 8.6 | |
| 1999..... | 0.6 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.1 | 8.6 | |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 2.8 | 5.7 | 13.9 |
| 2003..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 3.0 | 5.7 | 14.4 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 2.9 | 6.4 | 13.3 |
| 2001..... | 1.1 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 3.1 | 5.9 | 14.5 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 3.1 | 5.3 | 14.3 |
| 1999..... | 1.1 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.1 | 5.9 | 14.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 4.7 | 9.7 | 25.7 |
| 2003..... | 1.5 | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 4.8 | 9.7 | 25.8 |
| 2002..... | 1.5 | * | * | * | * | * | 0.2 | 0.9 | 2.2 | 4.7 | 10.9 | 23.4 |
| 2001..... | 1.5 | * | * | * | * | * | 0.1 | 0.9 | 2.7 | 5.0 | 9.6 | 26.3 |
| 2000..... | 1.4 | * | * | * | * | * | 0.1 | 0.8 | 2.1 | 5.3 | 9.5 | 26.8 |
| 1999..... | 1.5 | * | * | * | * | * | 0.3 | 0.9 | 2.5 | 5.1 | 9.8 | 26.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.3 | 0.5 | 1.2 | 2.9 | 8.6 | |
| 2003..... | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.0 | 9.4 | |
| 2002..... | 0.7 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.3 | 8.9 | |
| 2001..... | 0.7 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.5 | 9.6 | |
| 2000..... | 0.6 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.6 | 9.3 | |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.3 | 9.3 | |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of mesothelial and soft tissue (C45-C49) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.8 | 1.6 | 4.2 | 8.7 | 16.1 | 15.7 |
| 2003..... | 2.4 | * | * | 0.2 | 0.4 | 0.5 | 0.7 | 1.7 | 4.1 | 8.8 | 15.7 | 14.9 |
| 2002..... | 2.4 | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 1.6 | 4.3 | 9.1 | 14.9 | 15.7 |
| 2001..... | 2.4 | * | * | 0.1 | 0.3 | 0.5 | 0.8 | 1.8 | 4.2 | 9.1 | 15.1 | 14.9 |
| 2000..... | 2.4 | * | 0.1 | 0.1 | 0.4 | 0.5 | 0.8 | 1.8 | 4.4 | 9.6 | 15.1 | 15.7 |
| 1999..... | 2.4 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.8 | 4.4 | 9.3 | 14.8 | 14.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.6 | 4.8 | 11.3 | 24.7 | 27.9 |
| 2003..... | 2.8 | * | * | 0.2 | 0.4 | 0.5 | 0.8 | 1.8 | 4.6 | 11.6 | 24.3 | 25.1 |
| 2002..... | 2.8 | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 1.6 | 5.0 | 12.3 | 22.5 | 28.1 |
| 2001..... | 2.8 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 2.0 | 4.8 | 12.5 | 22.7 | 25.9 |
| 2000..... | 2.9 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 1.8 | 5.3 | 12.9 | 23.5 | 28.4 |
| 1999..... | 2.8 | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 1.9 | 5.5 | 12.8 | 22.6 | 24.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 1.6 | 3.6 | 6.5 | 10.3 | 10.2 |
| 2003..... | 2.0 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.6 | 3.7 | 6.4 | 9.9 | 10.3 |
| 2002..... | 1.9 | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 1.5 | 3.7 | 6.4 | 9.8 | 10.4 |
| 2001..... | 2.0 | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.6 | 3.7 | 6.3 | 10.1 | 10.3 |
| 2000..... | 2.0 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.7 | 3.6 | 6.9 | 9.6 | 10.5 |
| 1999..... | 1.9 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.7 | 3.5 | 6.4 | 9.6 | 10.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.6 | 4.6 | 9.6 | * |
| 2003..... | 0.9 | * | * | * | 0.4 | 0.3 | 0.6 | 1.3 | 2.7 | 6.1 | 10.3 | 13.0 |
| 2002..... | 0.9 | * | * | * | 0.3 | 0.3 | 0.7 | 1.1 | 3.0 | 6.1 | 11.2 | * |
| 2001..... | 0.9 | * | * | * | 0.3 | * | 0.6 | 1.5 | 3.0 | 6.0 | 9.2 | * |
| 2000..... | 0.8 | * | * | * | * | 0.4 | * | 1.3 | 2.1 | 6.3 | 11.3 | * |
| 1999..... | 0.9 | * | * | * | 0.4 | 0.4 | 0.7 | 1.4 | 3.6 | 5.9 | 8.1 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.6 | 1.2 | 2.4 | 6.0 | 17.2 | * |
| 2003..... | 1.0 | * | * | * | * | * | * | 1.3 | 3.0 | 7.2 | 17.6 | * |
| 2002..... | 1.0 | * | * | * | * | * | 0.8 | * | 3.9 | 7.1 | 19.3 | * |
| 2001..... | 0.9 | * | * | * | * | * | * | 1.2 | 3.6 | 7.0 | 14.3 | * |
| 2000..... | 0.9 | * | * | * | * | * | * | * | 3.0 | 7.6 | 18.2 | * |
| 1999..... | 1.0 | * | * | * | * | * | 0.8 | 1.4 | 4.0 | 8.3 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 1.4 | 2.7 | 3.4 | * | * |
| 2003..... | 0.9 | * | * | * | * | * | * | 1.2 | 2.4 | 5.3 | 5.3 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.5 | 2.2 | 5.3 | 5.6 | * |
| 2001..... | 0.8 | * | * | * | * | * | * | 1.8 | 2.6 | 5.2 | * | * |
| 2000..... | 0.7 | * | * | * | * | * | * | 1.5 | * | 5.3 | 6.6 | * |
| 1999..... | 0.9 | * | * | * | * | * | * | 1.5 | 3.2 | 4.0 | 7.6 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of mesothelial and soft tissue (C45-C49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | 0.2 | 0.4 | 0.5 | 0.8 | 1.6 | 4.3 | 9.0 | 16.5 | 16.0 |
| 2003..... | 2.6 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.7 | 4.3 | 8.9 | 15.9 | 14.9 |
| 2002..... | 2.6 | * | 0.2 | 0.2 | 0.3 | 0.5 | 0.8 | 1.6 | 4.4 | 9.2 | 15.0 | 15.9 |
| 2001..... | 2.6 | * | * | 0.1 | 0.3 | 0.5 | 0.8 | 1.8 | 4.3 | 9.3 | 15.3 | 15.1 |
| 2000..... | 2.6 | * | * | 0.1 | 0.4 | 0.5 | 0.8 | 1.8 | 4.6 | 9.8 | 15.2 | 16.1 |
| 1999..... | 2.6 | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 1.9 | 4.5 | 9.5 | 15.0 | 15.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | 0.2 | 0.4 | 0.6 | 0.9 | 1.6 | 5.0 | 11.7 | 25.1 | 28.7 |
| 2003..... | 3.1 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 1.9 | 4.7 | 11.9 | 24.7 | 25.7 |
| 2002..... | 3.1 | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 1.7 | 5.1 | 12.6 | 22.6 | 28.7 |
| 2001..... | 3.1 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 2.1 | 4.8 | 12.8 | 23.1 | 26.2 |
| 2000..... | 3.1 | * | * | * | 0.4 | 0.6 | 0.9 | 1.9 | 5.5 | 13.1 | 23.7 | 29.3 |
| 1999..... | 3.1 | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 2.0 | 5.5 | 13.1 | 23.1 | 25.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | 0.2 | 0.4 | 0.4 | 0.8 | 1.6 | 3.6 | 6.8 | 10.7 | 10.4 |
| 2003..... | 2.1 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.6 | 3.8 | 6.5 | 10.1 | 10.2 |
| 2002..... | 2.1 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.5 | 3.8 | 6.5 | 9.9 | 10.4 |
| 2001..... | 2.1 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.6 | 3.8 | 6.3 | 10.3 | 10.4 |
| 2000..... | 2.2 | * | * | * | 0.3 | 0.5 | 0.8 | 1.7 | 3.7 | 7.0 | 9.7 | 10.8 |
| 1999..... | 2.1 | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.7 | 3.5 | 6.5 | 9.7 | 10.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.6 | 4.5 | 9.6 | 17.4 | 16.9 |
| 2003..... | 2.9 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.7 | 4.4 | 9.5 | 16.8 | 15.6 |
| 2002..... | 2.9 | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 1.6 | 4.5 | 9.7 | 15.7 | 16.7 |
| 2001..... | 2.9 | * | * | 0.1 | 0.3 | 0.5 | 0.8 | 1.8 | 4.5 | 9.9 | 16.1 | 15.6 |
| 2000..... | 3.0 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 1.8 | 4.7 | 10.2 | 16.1 | 16.6 |
| 1999..... | 2.8 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.9 | 4.6 | 9.9 | 15.5 | 15.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.6 | 5.3 | 12.5 | 26.6 | 30.5 |
| 2003..... | 3.5 | * | * | * | 0.4 | 0.6 | 0.8 | 1.8 | 4.9 | 12.6 | 26.2 | 26.7 |
| 2002..... | 3.5 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.8 | 5.3 | 13.4 | 23.9 | 30.5 |
| 2001..... | 3.5 | * | * | * | 0.4 | 0.6 | 0.8 | 2.1 | 5.1 | 13.9 | 24.3 | 27.4 |
| 2000..... | 3.6 | * | * | * | 0.4 | 0.6 | 0.8 | 2.0 | 5.7 | 13.8 | 25.0 | 30.5 |
| 1999..... | 3.5 | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 2.0 | 5.8 | 13.7 | 24.2 | 26.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 1.6 | 3.7 | 7.1 | 11.2 | 10.8 |
| 2003..... | 2.4 | * | * | * | 0.3 | 0.4 | 0.7 | 1.6 | 4.0 | 6.8 | 10.4 | 10.7 |
| 2002..... | 2.3 | * | * | * | 0.3 | 0.4 | 0.7 | 1.5 | 3.8 | 6.6 | 10.3 | 10.7 |
| 2001..... | 2.3 | * | * | * | 0.3 | 0.5 | 0.7 | 1.5 | 3.9 | 6.6 | 10.7 | 10.7 |
| 2000..... | 2.4 | * | * | * | 0.3 | 0.5 | 0.8 | 1.6 | 3.8 | 7.2 | 10.2 | 11.0 |
| 1999..... | 2.2 | * | * | * | 0.3 | 0.4 | 0.6 | 1.7 | 3.5 | 6.7 | 9.8 | 10.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of mesothelial and soft tissue (C45-C49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.6 | 0.6 | 1.1 | 2.1 | 3.5 | 6.2 | 10.6 | 9.2 |
| 2003..... | 1.6 | * | * | * | 0.4 | 0.8 | 0.9 | 2.3 | 4.0 | 6.4 | 10.3 | 10.1 |
| 2002..... | 1.6 | * | * | * | 0.4 | 0.7 | 1.1 | 1.8 | 4.2 | 7.0 | 10.4 | 7.6 |
| 2001..... | 1.5 | * | * | * | 0.4 | 0.6 | 0.9 | 2.3 | 3.8 | 6.0 | 8.5 | 10.6 |
| 2000..... | 1.6 | * | * | * | 0.5 | 0.6 | 1.2 | 2.2 | 4.3 | 8.3 | 8.4 | 11.1 |
| 1999..... | 1.7 | * | * | * | 0.6 | 0.7 | 1.4 | 1.9 | 4.3 | 6.7 | 11.7 | 10.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 1.1 | 2.1 | 3.6 | 7.3 | 15.4 | * |
| 2003..... | 1.7 | * | * | * | * | 0.8 | 1.1 | 2.8 | 4.6 | 6.9 | 13.0 | * |
| 2002..... | 1.6 | * | * | * | * | 0.9 | 1.4 | 2.1 | 4.4 | 7.9 | 13.2 | * |
| 2001..... | 1.5 | * | * | * | 0.7 | 0.8 | 1.1 | 2.2 | 3.9 | 6.4 | 11.1 | * |
| 2000..... | 1.7 | * | * | * | * | * | 1.5 | 1.7 | 5.1 | 10.2 | 12.6 | * |
| 1999..... | 1.8 | * | * | * | 0.8 | 0.8 | 1.7 | 1.9 | 5.2 | 8.1 | 14.7 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 1.1 | 2.1 | 3.5 | 5.4 | 7.9 | * |
| 2003..... | 1.5 | * | * | * | * | 0.8 | 0.8 | 1.8 | 3.4 | 6.1 | 8.8 | * |
| 2002..... | 1.5 | * | * | * | * | * | 0.9 | 1.6 | 4.1 | 6.4 | 8.8 | * |
| 2001..... | 1.4 | * | * | * | * | * | 0.7 | 2.4 | 3.7 | 5.6 | 7.1 | 8.6 |
| 2000..... | 1.6 | * | * | * | * | * | 1.1 | 2.7 | 3.7 | 7.0 | 6.1 | 8.7 |
| 1999..... | 1.6 | * | * | * | * | * | 1.1 | 2.0 | 3.6 | 5.7 | 10.0 | 10.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mesothelioma (C45) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 1.4 | 3.6 | 8.2 | 5.9 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.8 | 7.6 | 5.7 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.1 | 7.2 | 6.1 |
| 2001..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 4.0 | 7.2 | 5.6 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 1.5 | 4.4 | 7.0 | 5.9 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 4.3 | 7.0 | 4.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 0.4 | 2.3 | 6.4 | 16.7 | 15.0 |
| 2003..... | 1.6 | * | * | * | * | * | 0.1 | 0.6 | 2.3 | 7.0 | 15.7 | 14.4 |
| 2002..... | 1.6 | * | * | * | * | * | 0.1 | 0.4 | 2.4 | 7.8 | 14.9 | 16.4 |
| 2001..... | 1.5 | * | * | * | * | * | * | 0.6 | 2.3 | 7.5 | 14.6 | 14.8 |
| 2000..... | 1.6 | * | * | * | * | * | * | 0.5 | 2.6 | 8.0 | 14.4 | 16.1 |
| 1999..... | 1.5 | * | * | * | * | * | * | 0.6 | 2.6 | 7.9 | 14.4 | 13.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.4 | 1.8 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 1.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 2.2 | 1.6 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.3 | 1.8 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.1 | 1.8 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.7 | 1.3 | 2.1 | 1.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 1.5 | 4.0 | 8.9 | 6.3 |
| 2003..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 1.5 | 4.2 | 8.2 | 6.0 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 1.6 | 4.5 | 7.8 | 6.6 |
| 2001..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 4.5 | 7.8 | 6.1 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 1.6 | 4.7 | 7.5 | 6.3 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 4.7 | 7.3 | 5.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | 0.5 | 2.6 | 7.1 | 17.8 | 16.2 |
| 2003..... | 1.9 | * | * | * | * | * | * | 0.6 | 2.5 | 7.6 | 16.8 | 15.2 |
| 2002..... | 1.9 | * | * | * | * | * | 0.1 | 0.4 | 2.7 | 8.5 | 15.8 | 17.8 |
| 2001..... | 1.8 | * | * | * | * | * | * | 0.6 | 2.6 | 8.4 | 15.7 | 16.0 |
| 2000..... | 1.8 | * | * | * | * | * | * | 0.5 | 2.8 | 8.5 | 15.5 | 17.3 |
| 1999..... | 1.8 | * | * | * | * | * | * | 0.6 | 2.8 | 8.7 | 15.2 | 14.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.7 | 1.9 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 2.0 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.4 | 1.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.5 | 2.0 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 2.3 | 1.9 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.2 | 1.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other mesothelial and soft tissue (C47-C49) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.7 | 1.3 | 2.8 | 5.2 | 8.1 | 9.6 |
| 2003..... | 1.5 | * | * | 0.2 | 0.4 | 0.4 | 0.6 | 1.3 | 2.8 | 5.0 | 8.1 | 8.8 |
| 2002..... | 1.5 | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.7 | 1.3 | 2.9 | 5.1 | 7.6 | 9.3 |
| 2001..... | 1.5 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.4 | 2.8 | 5.2 | 8.0 | 9.1 |
| 2000..... | 1.6 | * | 0.1 | 0.1 | 0.4 | 0.5 | 0.7 | 1.4 | 2.9 | 5.3 | 8.1 | 9.6 |
| 1999..... | 1.5 | * | * | 0.2 | 0.4 | 0.5 | 0.8 | 1.5 | 2.9 | 5.1 | 7.9 | 9.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.2 | 2.6 | 5.1 | 8.3 | 12.7 |
| 2003..... | 1.4 | * | * | 0.2 | 0.4 | 0.5 | 0.7 | 1.3 | 2.4 | 4.7 | 8.7 | 10.7 |
| 2002..... | 1.4 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.3 | 2.6 | 4.7 | 7.7 | 11.7 |
| 2001..... | 1.4 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.4 | 2.5 | 5.2 | 8.2 | 10.7 |
| 2000..... | 1.4 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.4 | 2.7 | 5.0 | 9.2 | 12.5 |
| 1999..... | 1.5 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.4 | 2.9 | 5.1 | 8.5 | 11.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 1.4 | 3.1 | 5.3 | 8.0 | 8.2 |
| 2003..... | 1.6 | * | * | 0.2 | 0.3 | 0.4 | 0.6 | 1.4 | 3.2 | 5.3 | 7.6 | 8.0 |
| 2002..... | 1.6 | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 1.3 | 3.2 | 5.4 | 7.6 | 8.3 |
| 2001..... | 1.6 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 3.2 | 5.1 | 7.8 | 8.4 |
| 2000..... | 1.7 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.5 | 3.1 | 5.6 | 7.4 | 8.4 |
| 1999..... | 1.6 | * | * | 0.2 | 0.3 | 0.4 | 0.6 | 1.5 | 2.8 | 5.1 | 7.5 | 8.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.5 | 1.1 | 1.8 | 2.9 | 5.6 | * |
| 2003..... | 0.7 | * | * | * | 0.4 | * | 0.6 | 1.0 | 1.9 | 3.4 | 4.2 | * |
| 2002..... | 0.7 | * | * | * | 0.3 | 0.3 | 0.6 | 1.0 | 2.2 | 3.6 | 5.5 | * |
| 2001..... | 0.7 | * | * | * | 0.3 | * | 0.5 | 1.3 | 2.5 | 4.1 | 4.7 | * |
| 2000..... | 0.6 | * | * | * | * | 0.4 | * | 1.0 | 1.4 | 4.6 | 7.7 | * |
| 1999..... | 0.7 | * | * | * | 0.4 | 0.3 | 0.6 | 1.1 | 2.8 | 3.6 | 5.6 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 1.0 | * | * | 7.5 | * |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 2.5 | * | 8.2 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 2.8 | 4.0 | * | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | * | 10.8 | * |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 2.7 | 4.4 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 1.2 | 2.2 | 3.0 | * | * |
| 2003..... | 0.7 | * | * | * | * | * | * | 1.1 | 2.1 | 4.3 | * | * |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.4 | 2.0 | 4.3 | * | * |
| 2001..... | 0.7 | * | * | * | * | * | * | 1.6 | 2.2 | 4.3 | * | * |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 5.1 | * | * |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 2.9 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other mesothelial and soft tissue (C47-C49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.2 | 0.4 | 0.5 | 0.7 | 1.3 | 2.9 | 5.4 | 8.3 | 9.7 |
| 2003..... | 1.6 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.4 | 2.9 | 5.1 | 8.3 | 8.9 |
| 2002..... | 1.6 | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.7 | 1.3 | 3.0 | 5.1 | 7.7 | 9.5 |
| 2001..... | 1.6 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.4 | 2.9 | 5.2 | 8.1 | 9.1 |
| 2000..... | 1.7 | * | * | 0.1 | 0.4 | 0.5 | 0.8 | 1.5 | 3.0 | 5.4 | 8.1 | 9.9 |
| 1999..... | 1.6 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.5 | 2.9 | 5.2 | 8.0 | 9.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.2 | 0.4 | 0.6 | 0.8 | 1.2 | 2.7 | 5.2 | 8.4 | 12.9 |
| 2003..... | 1.5 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.3 | 2.4 | 4.8 | 8.9 | 10.9 |
| 2002..... | 1.5 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.3 | 2.6 | 4.8 | 7.6 | 11.8 |
| 2001..... | 1.5 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 1.5 | 2.5 | 5.3 | 8.4 | 10.7 |
| 2000..... | 1.6 | * | * | * | 0.4 | 0.5 | 0.8 | 1.4 | 2.8 | 5.1 | 9.1 | 12.8 |
| 1999..... | 1.6 | * | * | 0.2 | 0.4 | 0.6 | 1.0 | 1.4 | 2.9 | 5.1 | 8.6 | 11.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | 0.2 | 0.4 | 0.3 | 0.7 | 1.4 | 3.1 | 5.5 | 8.2 | 8.3 |
| 2003..... | 1.8 | * | * | 0.2 | 0.3 | 0.4 | 0.6 | 1.4 | 3.3 | 5.3 | 7.8 | 8.0 |
| 2002..... | 1.8 | * | * | 0.1 | 0.3 | 0.3 | 0.6 | 1.3 | 3.3 | 5.4 | 7.7 | 8.4 |
| 2001..... | 1.8 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 3.2 | 5.2 | 7.9 | 8.4 |
| 2000..... | 1.8 | * | * | * | 0.3 | 0.5 | 0.7 | 1.5 | 3.2 | 5.6 | 7.5 | 8.7 |
| 1999..... | 1.7 | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 1.6 | 2.8 | 5.2 | 7.6 | 9.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.3 | 3.0 | 5.6 | 8.5 | 10.1 |
| 2003..... | 1.8 | * | * | 0.1 | 0.4 | 0.4 | 0.6 | 1.3 | 2.9 | 5.3 | 8.5 | 9.2 |
| 2002..... | 1.8 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.3 | 2.9 | 5.2 | 7.9 | 9.7 |
| 2001..... | 1.8 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.4 | 2.9 | 5.4 | 8.3 | 9.2 |
| 2000..... | 1.8 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.5 | 3.1 | 5.5 | 8.4 | 10.0 |
| 1999..... | 1.7 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.5 | 2.9 | 5.2 | 8.0 | 9.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.2 | 0.3 | 0.6 | 0.8 | 1.1 | 2.7 | 5.4 | 8.7 | 13.4 |
| 2003..... | 1.6 | * | * | * | 0.4 | 0.5 | 0.7 | 1.2 | 2.4 | 5.0 | 9.3 | 11.2 |
| 2002..... | 1.6 | * | * | 0.2 | 0.4 | 0.5 | 0.8 | 1.3 | 2.6 | 4.9 | 8.0 | 12.1 |
| 2001..... | 1.6 | * | * | * | 0.4 | 0.6 | 0.7 | 1.5 | 2.5 | 5.5 | 8.5 | 10.7 |
| 2000..... | 1.7 | * | * | * | 0.4 | 0.6 | 0.7 | 1.5 | 2.9 | 5.2 | 9.3 | 12.8 |
| 1999..... | 1.7 | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 1.4 | 3.0 | 5.1 | 8.7 | 12.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 1.4 | 3.1 | 5.7 | 8.4 | 8.6 |
| 2003..... | 1.9 | * | * | * | 0.3 | 0.3 | 0.6 | 1.4 | 3.4 | 5.5 | 7.9 | 8.3 |
| 2002..... | 1.9 | * | * | * | 0.3 | 0.3 | 0.6 | 1.3 | 3.2 | 5.4 | 7.9 | 8.7 |
| 2001..... | 1.9 | * | * | * | 0.3 | 0.5 | 0.7 | 1.3 | 3.3 | 5.3 | 8.1 | 8.5 |
| 2000..... | 1.9 | * | * | * | 0.3 | 0.5 | 0.7 | 1.4 | 3.3 | 5.7 | 7.8 | 8.8 |
| 1999..... | 1.8 | * | * | * | 0.3 | 0.4 | 0.5 | 1.5 | 2.8 | 5.3 | 7.6 | 9.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other mesothelial and soft tissue (C47-C49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | 0.6 | 0.5 | 1.0 | 1.9 | 2.9 | 4.9 | 7.5 | 6.9 |
| 2003..... | 1.3 | * | * | * | 0.3 | 0.7 | 0.9 | 1.9 | 3.2 | 5.1 | 7.3 | 7.1 |
| 2002..... | 1.3 | * | * | * | 0.4 | 0.6 | 1.0 | 1.7 | 3.6 | 5.4 | 7.1 | 6.0 |
| 2001..... | 1.3 | * | * | * | 0.4 | 0.6 | 0.8 | 2.0 | 3.3 | 4.8 | 6.8 | 9.4 |
| 2000..... | 1.3 | * | * | * | 0.5 | 0.4 | 1.1 | 2.0 | 3.3 | 5.5 | 5.9 | 9.2 |
| 1999..... | 1.4 | * | * | * | 0.6 | 0.7 | 1.3 | 1.6 | 3.4 | 5.3 | 8.0 | 8.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.9 | 1.7 | 2.5 | 5.0 | 6.8 | * |
| 2003..... | 1.2 | * | * | * | * | * | 1.0 | 2.1 | 3.1 | 4.0 | 6.3 | * |
| 2002..... | 1.2 | * | * | * | * | * | 1.1 | 1.8 | 3.3 | 4.5 | 5.9 | * |
| 2001..... | 1.2 | * | * | * | 0.7 | * | 1.0 | 1.7 | 2.9 | 4.3 | 7.2 | * |
| 2000..... | 1.1 | * | * | * | * | * | 1.3 | 1.2 | 3.2 | 4.6 | 6.8 | * |
| 1999..... | 1.3 | * | * | * | 0.8 | * | 1.5 | 1.4 | 3.4 | 5.7 | 7.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 1.1 | 2.1 | 3.3 | 4.8 | 7.9 | * |
| 2003..... | 1.4 | * | * | * | * | 0.8 | 0.8 | 1.8 | 3.2 | 5.9 | 7.8 | * |
| 2002..... | 1.4 | * | * | * | * | * | 0.8 | 1.6 | 3.9 | 6.0 | 7.8 | * |
| 2001..... | 1.4 | * | * | * | * | * | 0.7 | 2.2 | 3.6 | 5.2 | 6.6 | 8.6 |
| 2000..... | 1.5 | * | * | * | * | * | 1.0 | 2.6 | 3.4 | 6.1 | 5.4 | * |
| 1999..... | 1.4 | * | * | * | * | * | 1.0 | 1.8 | 3.4 | 5.0 | 8.1 | 9.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasm of breast (C50) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 14.1 | * | * | * | * | 1.0 | 5.7 | 15.0 | 29.1 | 44.9 | 72.2 | 124.2 |
| 2003..... | 14.4 | * | * | * | * | 1.0 | 6.1 | 15.6 | 29.6 | 45.5 | 74.9 | 132.5 |
| 2002..... | 14.5 | * | * | * | * | 1.1 | 6.1 | 16.1 | 29.5 | 46.5 | 76.5 | 134.6 |
| 2001..... | 14.7 | * | * | * | * | 1.2 | 6.3 | 16.8 | 30.2 | 47.6 | 76.8 | 134.2 |
| 2000..... | 15.0 | * | * | * | * | 1.2 | 6.3 | 16.9 | 31.2 | 49.1 | 78.9 | 147.4 |
| 1999..... | 14.9 | * | * | * | * | 1.1 | 6.1 | 16.9 | 31.1 | 49.5 | 78.9 | 144.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.1 | 3.3 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.0 | 3.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.2 | 3.4 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.3 | 3.5 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.2 | 4.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.0 | 4.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 27.5 | * | * | * | * | 2.0 | 11.3 | 29.3 | 55.8 | 81.6 | 119.5 | 178.6 |
| 2003..... | 28.2 | * | * | * | * | 2.1 | 12.2 | 30.4 | 56.6 | 82.6 | 123.7 | 189.4 |
| 2002..... | 28.3 | * | * | * | * | 2.1 | 12.0 | 31.4 | 56.2 | 84.4 | 125.9 | 191.5 |
| 2001..... | 28.6 | * | * | * | * | 2.4 | 12.4 | 32.8 | 57.5 | 85.8 | 125.8 | 188.9 |
| 2000..... | 29.2 | * | * | * | * | 2.3 | 12.4 | 33.0 | 59.3 | 88.3 | 128.9 | 205.7 |
| 1999..... | 28.9 | * | * | * | * | 2.2 | 12.0 | 32.9 | 59.2 | 88.9 | 128.9 | 200.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | * | 0.6 | 3.7 | 9.9 | 20.2 | 29.4 | 41.2 | 72.3 |
| 2003..... | 4.7 | * | * | * | * | 0.7 | 4.3 | 10.5 | 20.5 | 27.1 | 42.0 | 77.7 |
| 2002..... | 4.5 | * | * | * | * | 0.7 | 3.8 | 11.0 | 17.6 | 27.4 | 43.7 | 71.0 |
| 2001..... | 4.8 | * | * | * | * | 0.8 | 4.5 | 12.0 | 22.0 | 28.0 | 39.5 | 68.8 |
| 2000..... | 4.8 | * | * | * | * | 0.8 | 4.3 | 12.1 | 21.0 | 31.3 | 45.6 | 71.7 |
| 1999..... | 4.6 | * | * | * | * | 0.7 | 4.0 | 12.2 | 19.6 | 28.0 | 43.5 | 86.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | * | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 9.6 | * | * | * | * | 1.4 | 7.7 | 19.5 | 38.2 | 52.1 | 69.1 | 108.8 |
| 2003..... | 9.8 | * | * | * | * | 1.6 | 9.1 | 21.0 | 38.8 | 48.4 | 70.8 | 114.8 |
| 2002..... | 9.2 | * | * | * | * | 1.6 | 7.8 | 21.6 | 33.5 | 48.7 | 73.1 | 105.3 |
| 2001..... | 9.7 | * | * | * | * | 1.7 | 9.2 | 23.8 | 41.5 | 49.3 | 64.7 | 102.4 |
| 2000..... | 9.7 | * | * | * | * | 1.8 | 8.7 | 23.9 | 39.1 | 54.9 | 74.9 | 105.8 |
| 1999..... | 9.3 | * | * | * | * | 1.6 | 8.3 | 23.9 | 36.6 | 49.1 | 71.8 | 127.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasm of breast (C50) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.6 | * | * | * | * | 1.1 | 6.0 | 15.5 | 29.9 | 45.9 | 73.9 | 126.3 |
| 2003..... | 15.9 | * | * | * | * | 1.1 | 6.4 | 16.1 | 30.3 | 46.7 | 76.4 | 134.7 |
| 2002..... | 16.0 | * | * | * | * | 1.1 | 6.4 | 16.6 | 30.3 | 47.7 | 78.0 | 136.8 |
| 2001..... | 16.1 | * | * | * | * | 1.2 | 6.5 | 17.2 | 30.7 | 48.8 | 78.4 | 136.4 |
| 2000..... | 16.5 | * | * | * | * | 1.2 | 6.5 | 17.3 | 31.9 | 50.1 | 80.1 | 149.7 |
| 1999..... | 16.3 | * | * | * | * | 1.2 | 6.3 | 17.3 | 31.9 | 50.6 | 80.2 | 146.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 3.3 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.1 | 3.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 2.3 | 3.5 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.5 | 2.3 | 3.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.2 | 4.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.1 | 4.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 30.2 | * | * | * | * | 2.1 | 11.9 | 30.3 | 57.2 | 83.7 | 122.1 | 181.3 |
| 2003..... | 30.8 | * | * | * | * | 2.1 | 12.6 | 31.3 | 57.9 | 84.9 | 126.0 | 192.2 |
| 2002..... | 31.0 | * | * | * | * | 2.2 | 12.6 | 32.3 | 57.9 | 86.7 | 128.2 | 194.4 |
| 2001..... | 31.1 | * | * | * | * | 2.5 | 12.8 | 33.6 | 58.6 | 88.1 | 128.3 | 191.6 |
| 2000..... | 31.8 | * | * | * | * | 2.4 | 12.8 | 33.8 | 60.7 | 90.3 | 130.8 | 208.6 |
| 1999..... | 31.4 | * | * | * | * | 2.3 | 12.4 | 33.6 | 60.7 | 91.2 | 130.9 | 202.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 16.4 | * | * | * | * | 0.9 | 5.2 | 14.0 | 28.7 | 45.8 | 74.7 | 126.8 |
| 2003..... | 16.8 | * | * | * | * | 0.9 | 5.7 | 14.8 | 29.0 | 46.3 | 76.7 | 134.7 |
| 2002..... | 16.9 | * | * | * | * | 0.9 | 5.6 | 15.2 | 29.4 | 47.4 | 78.0 | 137.7 |
| 2001..... | 17.0 | * | * | * | * | 1.0 | 5.8 | 16.0 | 29.6 | 48.2 | 78.7 | 137.1 |
| 2000..... | 17.4 | * | * | * | * | 1.1 | 5.8 | 16.1 | 30.8 | 50.3 | 80.6 | 149.6 |
| 1999..... | 17.1 | * | * | * | * | 1.0 | 5.6 | 16.0 | 30.6 | 50.2 | 80.3 | 147.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 2.2 | 2.9 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 3.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.2 | 3.4 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.2 | 3.5 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.2 | 4.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.0 | 4.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 32.0 | * | * | * | * | 1.8 | 10.4 | 27.7 | 55.6 | 84.4 | 124.2 | 182.4 |
| 2003..... | 32.6 | * | * | * | * | 1.8 | 11.3 | 29.1 | 56.0 | 85.1 | 127.1 | 192.3 |
| 2002..... | 32.9 | * | * | * | * | 1.9 | 11.1 | 30.0 | 56.7 | 87.0 | 128.9 | 195.8 |
| 2001..... | 32.9 | * | * | * | * | 2.0 | 11.5 | 31.6 | 57.0 | 88.0 | 129.4 | 192.6 |
| 2000..... | 33.8 | * | * | * | * | 2.1 | 11.6 | 31.7 | 59.2 | 91.4 | 132.2 | 208.3 |
| 1999..... | 33.3 | * | * | * | * | 2.0 | 11.1 | 31.5 | 58.9 | 91.2 | 131.5 | 204.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasm of breast (C50) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 15.0 | * | * | * | * | 2.2 | 11.3 | 26.7 | 44.7 | 57.4 | 83.1 | 145.6 |
| 2003..... | 15.5 | * | * | * | * | 2.3 | 11.4 | 27.1 | 46.1 | 60.1 | 91.3 | 163.3 |
| 2002..... | 15.4 | * | * | * | * | 2.4 | 12.0 | 27.7 | 43.7 | 61.1 | 95.3 | 154.0 |
| 2001..... | 15.5 | * | * | * | * | 2.7 | 11.5 | 27.9 | 45.6 | 64.6 | 93.0 | 153.9 |
| 2000..... | 15.3 | * | * | * | * | 2.3 | 11.5 | 28.4 | 46.4 | 59.4 | 91.7 | 178.7 |
| 1999..... | 15.5 | * | * | * | * | 2.4 | 12.0 | 29.2 | 47.1 | 64.5 | 95.6 | 149.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 28.4 | * | * | * | * | 4.2 | 21.2 | 49.3 | 79.8 | 96.8 | 128.5 | 199.6 |
| 2003..... | 29.4 | * | * | * | * | 4.3 | 21.4 | 50.1 | 82.4 | 101.6 | 141.8 | 224.5 |
| 2002..... | 29.1 | * | * | * | * | 4.5 | 22.5 | 51.0 | 77.8 | 102.7 | 147.1 | 210.7 |
| 2001..... | 29.2 | * | * | * | * | 5.1 | 21.5 | 51.3 | 81.6 | 108.4 | 143.0 | 208.3 |
| 2000..... | 28.9 | * | * | * | * | 4.3 | 21.5 | 52.5 | 82.5 | 100.1 | 141.5 | 241.6 |
| 1999..... | 29.2 | * | * | * | * | 4.5 | 22.3 | 53.7 | 83.7 | 108.2 | 147.2 | 201.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of female genital organs (C51-C58) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.2 | * | * | * | 0.1 | 0.8 | 2.9 | 7.7 | 17.5 | 34.8 | 54.8 | 72.7 |
| 2003..... | 9.3 | * | * | * | 0.1 | 0.8 | 2.9 | 7.8 | 18.5 | 34.3 | 55.7 | 75.2 |
| 2002..... | 9.4 | * | * | * | 0.1 | 0.9 | 3.0 | 8.2 | 18.3 | 34.9 | 56.9 | 77.5 |
| 2001..... | 9.4 | * | * | * | 0.1 | 0.9 | 3.0 | 8.4 | 18.2 | 34.7 | 58.2 | 79.9 |
| 2000..... | 9.4 | * | * | * | 0.1 | 0.9 | 3.0 | 8.3 | 18.5 | 34.8 | 57.7 | 82.3 |
| 1999..... | 9.3 | * | * | * | 0.1 | 0.8 | 3.1 | 8.3 | 18.5 | 35.0 | 56.9 | 79.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 18.1 | * | * | * | 0.3 | 1.7 | 5.9 | 15.0 | 33.8 | 64.1 | 91.7 | 105.4 |
| 2003..... | 18.3 | * | * | * | 0.3 | 1.7 | 5.8 | 15.3 | 35.7 | 63.0 | 93.0 | 108.5 |
| 2002..... | 18.5 | * | * | * | 0.3 | 1.7 | 5.9 | 16.2 | 35.3 | 63.9 | 94.6 | 111.2 |
| 2001..... | 18.5 | * | * | * | 0.2 | 1.8 | 6.0 | 16.4 | 35.1 | 63.4 | 96.4 | 113.4 |
| 2000..... | 18.4 | * | * | * | 0.2 | 1.9 | 6.0 | 16.3 | 35.6 | 63.5 | 95.3 | 115.8 |
| 1999..... | 18.2 | * | * | * | 0.2 | 1.6 | 6.1 | 16.4 | 35.6 | 63.6 | 93.8 | 110.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | 0.6 | 2.7 | 6.2 | 14.8 | 26.5 | 39.9 | 48.5 | 63.2 |
| 2003..... | 3.6 | * | * | * | 0.5 | 2.7 | 6.9 | 13.1 | 24.1 | 40.7 | 61.0 | 75.6 |
| 2002..... | 3.6 | * | * | * | 0.6 | 2.7 | 6.8 | 12.9 | 26.8 | 44.5 | 66.9 | 82.3 |
| 2001..... | 3.8 | * | * | * | 0.8 | 2.9 | 7.2 | 15.6 | 26.6 | 41.5 | 59.6 | 75.6 |
| 2000..... | 3.8 | * | * | * | 0.7 | 3.2 | 7.0 | 15.0 | 28.6 | 47.4 | 75.6 | 91.2 |
| 1999..... | 3.5 | * | * | * | 0.7 | 2.7 | 7.3 | 14.1 | 26.6 | 42.3 | 59.6 | 75.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | 1.4 | 5.7 | 12.4 | 28.3 | 47.7 | 67.4 | 73.5 | 91.2 |
| 2003..... | 7.4 | * | * | * | 1.2 | 5.7 | 13.8 | 24.9 | 43.4 | 68.7 | 95.3 | 115.8 |
| 2002..... | 7.5 | * | * | * | 1.4 | 5.6 | 13.6 | 24.5 | 48.2 | 75.0 | 91.2 | 115.8 |
| 2001..... | 7.8 | * | * | * | 1.8 | 6.1 | 14.3 | 29.4 | 47.6 | 69.5 | 99.6 | 122.8 |
| 2000..... | 7.9 | * | * | * | 1.6 | 6.6 | 13.8 | 28.2 | 51.1 | 79.2 | 112.8 | 145.8 |
| 1999..... | 7.2 | * | * | * | 1.5 | 5.5 | 14.3 | 26.6 | 47.5 | 70.4 | 88.4 | 115.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of female genital organs (C51-C58) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 10.1 | * | * | * | 0.1 | 0.9 | 3.0 | 7.8 | 17.7 | 35.4 | 55.6 | 73.6 |
| 2003..... | 10.2 | * | * | * | 0.1 | 0.9 | 2.9 | 7.8 | 18.9 | 34.9 | 56.4 | 75.7 |
| 2002..... | 10.3 | * | * | * | 0.1 | 0.9 | 3.0 | 8.4 | 18.7 | 35.3 | 57.4 | 78.1 |
| 2001..... | 10.2 | * | * | * | 0.1 | 0.9 | 3.0 | 8.4 | 18.4 | 35.1 | 58.9 | 80.2 |
| 2000..... | 10.2 | * | * | * | 0.1 | 1.0 | 3.0 | 8.4 | 18.7 | 35.1 | 58.0 | 82.4 |
| 1999..... | 10.0 | * | * | * | 0.1 | 0.8 | 3.1 | 8.4 | 18.8 | 35.4 | 57.2 | 79.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 19.7 | * | * | * | 0.3 | 1.8 | 5.9 | 15.3 | 34.2 | 65.2 | 93.0 | 106.6 |
| 2003..... | 19.9 | * | * | * | 0.3 | 1.7 | 5.8 | 15.4 | 36.5 | 64.2 | 94.1 | 109.0 |
| 2002..... | 20.0 | * | * | * | 0.3 | 1.8 | 6.0 | 16.4 | 36.0 | 64.8 | 95.4 | 111.8 |
| 2001..... | 20.0 | * | * | * | 0.2 | 1.8 | 6.0 | 16.6 | 35.4 | 64.3 | 97.5 | 113.5 |
| 2000..... | 19.8 | * | * | * | 0.2 | 1.9 | 5.9 | 16.4 | 36.1 | 64.1 | 95.8 | 115.7 |
| 1999..... | 19.6 | * | * | * | 0.2 | 1.7 | 6.1 | 16.5 | 36.1 | 64.4 | 94.4 | 111.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 10.8 | * | * | * | 0.1 | 0.8 | 2.9 | 7.5 | 17.3 | 34.1 | 55.8 | 74.0 |
| 2003..... | 10.9 | * | * | * | 0.1 | 0.8 | 2.8 | 7.7 | 18.6 | 34.1 | 56.0 | 75.6 |
| 2002..... | 10.9 | * | * | * | 0.1 | 0.9 | 2.9 | 8.2 | 18.3 | 34.2 | 56.8 | 77.3 |
| 2001..... | 10.9 | * | * | * | 0.1 | 0.9 | 2.9 | 8.3 | 17.9 | 34.2 | 58.6 | 79.8 |
| 2000..... | 10.8 | * | * | * | 0.1 | 0.9 | 2.8 | 8.2 | 18.4 | 33.9 | 57.5 | 82.1 |
| 1999..... | 10.7 | * | * | * | 0.1 | 0.8 | 3.0 | 8.4 | 18.4 | 34.2 | 56.6 | 79.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 21.2 | * | * | * | 0.3 | 1.7 | 5.7 | 14.9 | 33.8 | 63.4 | 93.8 | 107.2 |
| 2003..... | 21.3 | * | * | * | 0.2 | 1.6 | 5.5 | 15.2 | 36.3 | 63.3 | 93.8 | 108.9 |
| 2002..... | 21.4 | * | * | * | 0.3 | 1.8 | 5.8 | 16.2 | 35.7 | 63.4 | 94.9 | 110.7 |
| 2001..... | 21.4 | * | * | * | 0.2 | 1.7 | 5.8 | 16.5 | 34.7 | 63.2 | 97.4 | 113.1 |
| 2000..... | 21.1 | * | * | * | 0.2 | 1.9 | 5.5 | 16.3 | 35.7 | 62.4 | 95.2 | 115.2 |
| 1999..... | 21.0 | * | * | * | 0.2 | 1.7 | 6.0 | 16.7 | 35.7 | 62.8 | 93.7 | 111.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of female genital organs (C51-C58) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 8.9 | * | * | * | * | 1.3 | 4.0 | 10.3 | 23.4 | 53.2 | 66.4 | 81.2 |
| 2003..... | 8.8 | * | * | * | * | 1.4 | 3.9 | 9.9 | 24.5 | 48.6 | 70.5 | 90.4 |
| 2002..... | 8.9 | * | * | * | * | 1.3 | 4.2 | 10.0 | 23.9 | 50.7 | 71.6 | 99.8 |
| 2001..... | 8.8 | * | * | * | * | 1.4 | 4.2 | 9.6 | 24.4 | 50.0 | 72.1 | 95.5 |
| 2000..... | 8.9 | * | * | * | * | 1.3 | 4.4 | 9.8 | 24.3 | 52.6 | 74.3 | 97.0 |
| 1999..... | 8.7 | * | * | * | * | 1.0 | 4.0 | 9.1 | 25.4 | 53.2 | 72.3 | 92.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 16.9 | * | * | * | * | 2.5 | 7.5 | 19.1 | 42.2 | 91.1 | 103.9 | 113.2 |
| 2003..... | 16.8 | * | * | * | * | 2.7 | 7.4 | 18.4 | 44.1 | 83.2 | 110.3 | 125.5 |
| 2002..... | 17.0 | * | * | * | * | 2.4 | 7.9 | 18.6 | 43.1 | 86.7 | 112.1 | 138.0 |
| 2001..... | 16.7 | * | * | * | * | 2.6 | 7.9 | 17.8 | 44.0 | 85.3 | 112.5 | 130.9 |
| 2000..... | 17.0 | * | * | * | * | 2.5 | 8.2 | 18.1 | 43.9 | 89.5 | 116.2 | 133.0 |
| 1999..... | 16.5 | * | * | * | * | 2.0 | 7.5 | 16.8 | 45.8 | 90.8 | 112.8 | 127.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of vulva and vagina (C51-C52) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.9 | 7.0 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.9 | 7.2 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.0 | 7.7 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.1 | 7.3 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 2.9 | 7.7 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.0 | 8.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.3 | 4.9 | 10.1 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.2 | 4.8 | 10.3 |
| 2002..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.9 | 11.0 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 2.1 | 5.2 | 10.3 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 4.8 | 10.8 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.2 | 5.0 | 11.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.0 | 7.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 7.5 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.0 | 8.0 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.2 | 7.3 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 3.0 | 7.9 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.0 | 9.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.3 | 5.1 | 10.5 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.4 | 0.9 | 2.3 | 5.1 | 10.7 |
| 2002..... | 1.0 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.1 | 5.1 | 11.5 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.1 | 5.3 | 10.4 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.5 | 5.0 | 11.1 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.2 | 5.0 | 12.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of vulva and vagina (C51-C52) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.9 | 1.3 | 2.4 | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | 2.2 | 6.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 2.4 | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.9 | 1.5 | 3.5 | 6.6 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | 2.7 | 7.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 1.4 | 3.1 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 1.5 | 2.2 | 3.7 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 3.4 | 8.7 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 3.8 | * |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 1.6 | 2.6 | 5.4 | 9.0 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | 4.2 | 9.6 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 2.3 | 4.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of cervix uteri (C53) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.5 | 1.4 | 2.0 | 2.8 | 3.3 | 3.7 | 5.1 |
| 2003..... | 1.3 | * | * | * | * | 0.5 | 1.4 | 2.2 | 2.8 | 3.1 | 4.3 | 5.7 |
| 2002..... | 1.4 | * | * | * | * | 0.5 | 1.5 | 2.2 | 2.7 | 3.4 | 4.1 | 6.2 |
| 2001..... | 1.4 | * | * | * | * | 0.5 | 1.6 | 2.4 | 2.9 | 3.2 | 4.6 | 7.2 |
| 2000..... | 1.5 | * | * | * | * | 0.6 | 1.5 | 2.4 | 3.1 | 3.7 | 4.8 | 7.3 |
| 1999..... | 1.5 | * | * | * | * | 0.5 | 1.6 | 2.5 | 3.2 | 3.8 | 4.6 | 7.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | 1.1 | 2.8 | 4.0 | 5.4 | 6.0 | 6.2 | 7.3 |
| 2003..... | 2.7 | * | * | * | * | 1.0 | 2.8 | 4.3 | 5.5 | 5.6 | 7.2 | 8.3 |
| 2002..... | 2.7 | * | * | * | * | 1.1 | 3.0 | 4.4 | 5.2 | 6.3 | 6.9 | 8.9 |
| 2001..... | 2.8 | * | * | * | * | 1.1 | 3.1 | 4.8 | 5.5 | 5.8 | 7.7 | 10.1 |
| 2000..... | 2.9 | * | * | * | * | 1.2 | 3.1 | 4.8 | 5.9 | 6.8 | 8.0 | 10.3 |
| 1999..... | 3.0 | * | * | * | * | 1.1 | 3.1 | 4.9 | 6.2 | 6.9 | 7.7 | 11.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.4 | 1.6 | 2.4 | 4.3 | 4.3 | 5.7 | * |
| 2003..... | 1.0 | * | * | * | * | 0.4 | 1.4 | 2.9 | 3.1 | 3.5 | 6.6 | * |
| 2002..... | 1.0 | * | * | * | * | 0.4 | 1.5 | 2.4 | 3.6 | 4.7 | 6.7 | * |
| 2001..... | 1.1 | * | * | * | * | 0.5 | 1.9 | 3.0 | 4.1 | 3.4 | 7.0 | * |
| 2000..... | 1.2 | * | * | * | * | 0.5 | 2.1 | 2.8 | 4.6 | 5.6 | 8.5 | 13.9 |
| 1999..... | 1.1 | * | * | * | * | 0.5 | 1.8 | 3.3 | 3.9 | 4.9 | 6.5 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | 0.9 | 3.3 | 4.8 | 8.2 | 7.7 | 9.6 | * |
| 2003..... | 2.0 | * | * | * | * | 0.9 | 3.0 | 5.7 | 5.9 | 6.3 | 11.1 | * |
| 2002..... | 2.1 | * | * | * | * | 0.9 | 3.1 | 4.7 | 6.8 | 8.4 | 11.2 | * |
| 2001..... | 2.3 | * | * | * | * | 1.1 | 3.9 | 6.0 | 7.7 | 6.1 | 11.8 | * |
| 2000..... | 2.5 | * | * | * | * | 1.1 | 4.4 | 5.5 | 8.6 | 10.0 | 14.2 | 20.8 |
| 1999..... | 2.3 | * | * | * | * | 1.1 | 3.6 | 6.6 | 7.4 | 8.8 | 10.8 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of cervix uteri (C53) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | 0.6 | 1.4 | 2.0 | 2.7 | 3.2 | 3.6 | 5.0 |
| 2003..... | 1.4 | * | * | * | * | 0.5 | 1.4 | 2.1 | 2.8 | 3.0 | 4.2 | 5.7 |
| 2002..... | 1.4 | * | * | * | * | 0.5 | 1.5 | 2.2 | 2.6 | 3.4 | 4.0 | 6.1 |
| 2001..... | 1.5 | * | * | * | * | 0.5 | 1.5 | 2.4 | 2.8 | 3.2 | 4.5 | 7.0 |
| 2000..... | 1.5 | * | * | * | * | 0.6 | 1.5 | 2.4 | 3.0 | 3.6 | 4.7 | 7.1 |
| 1999..... | 1.6 | * | * | * | * | 0.5 | 1.5 | 2.4 | 3.2 | 3.7 | 4.6 | 7.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | 1.1 | 2.7 | 3.9 | 5.2 | 5.9 | 6.0 | 7.2 |
| 2003..... | 2.7 | * | * | * | * | 1.0 | 2.8 | 4.1 | 5.4 | 5.6 | 7.0 | 8.2 |
| 2002..... | 2.8 | * | * | * | * | 1.1 | 2.9 | 4.3 | 5.1 | 6.2 | 6.7 | 8.7 |
| 2001..... | 2.9 | * | * | * | * | 1.0 | 3.0 | 4.6 | 5.3 | 5.8 | 7.5 | 10.0 |
| 2000..... | 3.0 | * | * | * | * | 1.2 | 2.9 | 4.7 | 5.7 | 6.6 | 7.7 | 9.9 |
| 1999..... | 3.0 | * | * | * | * | 1.1 | 3.0 | 4.7 | 6.1 | 6.7 | 7.5 | 11.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | 0.5 | 1.2 | 1.8 | 2.4 | 2.6 | 3.1 | 3.9 |
| 2003..... | 1.3 | * | * | * | * | 0.4 | 1.3 | 1.9 | 2.5 | 2.5 | 3.4 | 4.8 |
| 2002..... | 1.3 | * | * | * | * | 0.5 | 1.3 | 1.9 | 2.3 | 2.7 | 3.4 | 4.9 |
| 2001..... | 1.4 | * | * | * | * | 0.5 | 1.4 | 2.1 | 2.4 | 2.6 | 3.8 | 6.1 |
| 2000..... | 1.4 | * | * | * | * | 0.5 | 1.3 | 2.1 | 2.5 | 2.9 | 4.1 | 6.0 |
| 1999..... | 1.4 | * | * | * | * | 0.6 | 1.4 | 2.3 | 2.7 | 3.0 | 3.9 | 6.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | 1.1 | 2.5 | 3.5 | 4.6 | 4.9 | 5.2 | 5.7 |
| 2003..... | 2.5 | * | * | * | * | 0.9 | 2.6 | 3.7 | 4.9 | 4.7 | 5.7 | 7.0 |
| 2002..... | 2.5 | * | * | * | * | 1.0 | 2.7 | 3.8 | 4.5 | 5.0 | 5.7 | 7.0 |
| 2001..... | 2.7 | * | * | * | * | 1.0 | 2.8 | 4.2 | 4.7 | 4.9 | 6.3 | 8.6 |
| 2000..... | 2.7 | * | * | * | * | 1.1 | 2.5 | 4.2 | 4.8 | 5.3 | 6.8 | 8.4 |
| 1999..... | 2.8 | * | * | * | * | 1.1 | 2.8 | 4.5 | 5.2 | 5.5 | 6.5 | 9.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of cervix uteri (C53) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | 0.8 | 2.3 | 3.8 | 5.4 | 7.7 | 9.1 | 16.4 |
| 2003..... | 2.2 | * | * | * | * | 0.9 | 2.3 | 3.7 | 5.7 | 7.5 | 11.6 | 14.9 |
| 2002..... | 2.3 | * | * | * | * | 0.8 | 2.6 | 4.2 | 5.1 | 9.2 | 10.7 | 17.5 |
| 2001..... | 2.2 | * | * | * | * | 0.8 | 2.5 | 3.8 | 5.3 | 7.9 | 12.0 | 19.0 |
| 2000..... | 2.5 | * | * | * | * | 1.0 | 2.6 | 4.3 | 6.4 | 9.8 | 12.0 | 19.7 |
| 1999..... | 2.4 | * | * | * | * | 0.7 | 2.5 | 3.6 | 7.2 | 10.8 | 11.3 | 21.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | 1.6 | 4.3 | 7.1 | 9.8 | 13.2 | 14.2 | 22.9 |
| 2003..... | 4.1 | * | * | * | * | 1.8 | 4.3 | 6.9 | 10.2 | 12.8 | 18.1 | 20.6 |
| 2002..... | 4.3 | * | * | * | * | 1.6 | 4.9 | 7.8 | 9.2 | 15.8 | 16.7 | 24.2 |
| 2001..... | 4.2 | * | * | * | * | 1.6 | 4.6 | 7.1 | 9.6 | 13.5 | 18.7 | 26.1 |
| 2000..... | 4.7 | * | * | * | * | 1.9 | 5.0 | 8.0 | 11.5 | 16.6 | 18.8 | 27.0 |
| 1999..... | 4.5 | * | * | * | * | 1.2 | 4.8 | 6.6 | 13.0 | 18.3 | 17.7 | 29.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of corpus uteri and uterus, part unspecified (C54-C55) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | 0.0 | 0.4 | 1.3 | 4.4 | 10.1 | 15.5 | 22.3 |
| 2003..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.7 | 10.1 | 15.2 | 22.8 |
| 2002..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.5 | 10.0 | 15.4 | 24.6 |
| 2001..... | 2.4 | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.4 | 9.7 | 16.4 | 24.7 |
| 2000..... | 2.3 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.6 | 9.7 | 15.8 | 24.7 |
| 1999..... | 2.3 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.3 | 9.8 | 16.4 | 24.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | * | 0.1 | 0.8 | 2.6 | 8.6 | 18.5 | 25.9 | 32.3 |
| 2003..... | 4.7 | * | * | * | * | 0.1 | 0.7 | 2.6 | 9.0 | 18.5 | 25.3 | 33.0 |
| 2002..... | 4.7 | * | * | * | * | 0.1 | 0.6 | 2.8 | 8.7 | 18.4 | 25.6 | 35.2 |
| 2001..... | 4.7 | * | * | * | * | 0.2 | 0.6 | 2.8 | 8.4 | 17.8 | 27.2 | 35.0 |
| 2000..... | 4.6 | * | * | * | * | 0.1 | 0.6 | 2.6 | 8.9 | 17.7 | 26.2 | 34.8 |
| 1999..... | 4.5 | * | * | * | * | 0.1 | 0.6 | 2.5 | 8.2 | 17.8 | 27.1 | 33.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.9 | 3.5 | 7.9 | 10.5 | 9.5 |
| 2003..... | 0.8 | * | * | * | * | * | 0.4 | 1.2 | 2.9 | 7.0 | 9.8 | 20.7 |
| 2002..... | 0.9 | * | * | * | * | * | 0.3 | 1.1 | 3.2 | 9.0 | 11.0 | 20.5 |
| 2001..... | 0.9 | * | * | * | * | * | * | 1.4 | 3.8 | 7.5 | 11.7 | 18.8 |
| 2000..... | 0.8 | * | * | * | * | * | * | 1.1 | 3.6 | 7.0 | 10.7 | 19.9 |
| 1999..... | 0.8 | * | * | * | * | * | * | 1.1 | 3.7 | 8.3 | 10.6 | 15.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 1.7 | 6.7 | 14.2 | 17.8 | 14.4 |
| 2003..... | 1.7 | * | * | * | * | * | 0.9 | 2.3 | 5.6 | 12.5 | 16.6 | 31.3 |
| 2002..... | 1.8 | * | * | * | * | * | 0.7 | 2.1 | 6.2 | 16.1 | 18.5 | 30.7 |
| 2001..... | 1.8 | * | * | * | * | * | * | 2.8 | 7.3 | 13.4 | 19.6 | 27.9 |
| 2000..... | 1.6 | * | * | * | * | * | * | 2.2 | 6.8 | 12.4 | 17.8 | 29.7 |
| 1999..... | 1.6 | * | * | * | * | * | * | 2.2 | 7.0 | 14.8 | 17.7 | 22.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of corpus uteri and uterus, part unspecified (C54-C55) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | 0.4 | 1.4 | 4.5 | 10.2 | 15.8 | 22.8 |
| 2003..... | 2.6 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.8 | 10.3 | 15.4 | 22.9 |
| 2002..... | 2.6 | * | * | * | * | * | 0.3 | 1.4 | 4.6 | 10.1 | 15.5 | 24.6 |
| 2001..... | 2.6 | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.4 | 9.9 | 16.6 | 24.8 |
| 2000..... | 2.6 | * | * | * | * | 0.1 | 0.3 | 1.4 | 4.7 | 9.8 | 16.0 | 24.9 |
| 1999..... | 2.5 | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.3 | 9.9 | 16.6 | 24.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.2 | * | * | * | * | * | 0.8 | 2.7 | 8.7 | 18.8 | 26.4 | 33.0 |
| 2003..... | 5.1 | * | * | * | * | 0.1 | 0.7 | 2.6 | 9.2 | 18.9 | 25.6 | 33.0 |
| 2002..... | 5.1 | * | * | * | * | * | 0.6 | 2.8 | 8.8 | 18.5 | 25.8 | 35.3 |
| 2001..... | 5.1 | * | * | * | * | 0.2 | 0.6 | 2.8 | 8.5 | 18.1 | 27.5 | 35.1 |
| 2000..... | 5.0 | * | * | * | * | 0.1 | 0.6 | 2.7 | 9.0 | 17.9 | 26.5 | 34.9 |
| 1999..... | 4.9 | * | * | * | * | 0.1 | 0.6 | 2.5 | 8.3 | 18.0 | 27.4 | 34.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | 0.4 | 1.3 | 4.1 | 8.9 | 15.2 | 23.0 |
| 2003..... | 2.7 | * | * | * | * | * | 0.3 | 1.3 | 4.4 | 9.3 | 14.5 | 22.8 |
| 2002..... | 2.6 | * | * | * | * | * | 0.3 | 1.3 | 4.1 | 8.9 | 14.6 | 23.8 |
| 2001..... | 2.7 | * | * | * | * | * | 0.3 | 1.3 | 4.0 | 8.9 | 15.9 | 24.5 |
| 2000..... | 2.6 | * | * | * | * | * | 0.3 | 1.3 | 4.3 | 8.6 | 15.0 | 24.5 |
| 1999..... | 2.6 | * | * | * | * | * | 0.3 | 1.3 | 4.0 | 8.9 | 16.0 | 23.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | * | * | 0.8 | 2.6 | 8.0 | 16.6 | 25.5 | 33.3 |
| 2003..... | 5.2 | * | * | * | * | * | 0.6 | 2.5 | 8.5 | 17.2 | 24.4 | 32.8 |
| 2002..... | 5.1 | * | * | * | * | * | 0.6 | 2.7 | 8.0 | 16.5 | 24.4 | 34.1 |
| 2001..... | 5.2 | * | * | * | * | * | 0.6 | 2.7 | 7.8 | 16.4 | 26.5 | 34.7 |
| 2000..... | 5.1 | * | * | * | * | * | 0.5 | 2.7 | 8.4 | 15.9 | 24.8 | 34.3 |
| 1999..... | 5.1 | * | * | * | * | * | 0.6 | 2.5 | 7.7 | 16.3 | 26.4 | 33.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of corpus uteri and uterus, part unspecified (C54-C55) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | * | 0.5 | 2.0 | 8.1 | 23.4 | 26.6 | 26.5 |
| 2003..... | 3.0 | * | * | * | * | * | 0.5 | 1.7 | 9.1 | 21.4 | 28.0 | 30.3 |
| 2002..... | 3.1 | * | * | * | * | * | 0.4 | 2.0 | 8.6 | 22.7 | 28.4 | 39.6 |
| 2001..... | 2.9 | * | * | * | * | * | 0.4 | 2.1 | 7.7 | 21.6 | 28.6 | 32.8 |
| 2000..... | 3.0 | * | * | * | * | * | 0.4 | 1.7 | 8.4 | 22.9 | 31.1 | 33.7 |
| 1999..... | 2.8 | * | * | * | * | * | 0.5 | 1.4 | 8.1 | 21.5 | 27.5 | 35.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 5.7 | * | * | * | * | * | 1.0 | 3.6 | 14.7 | 40.1 | 41.5 | 36.9 |
| 2003..... | 5.7 | * | * | * | * | * | 1.0 | 3.2 | 16.5 | 36.7 | 43.8 | 42.1 |
| 2002..... | 5.8 | * | * | * | * | * | 0.7 | 3.6 | 15.5 | 38.7 | 44.4 | 54.8 |
| 2001..... | 5.5 | * | * | * | * | * | 0.7 | 4.0 | 13.9 | 36.8 | 44.6 | 44.9 |
| 2000..... | 5.7 | * | * | * | * | * | 0.7 | 3.1 | 15.3 | 39.1 | 48.6 | 46.2 |
| 1999..... | 5.4 | * | * | * | * | * | 0.9 | 2.7 | 14.7 | 36.7 | 42.9 | 49.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of ovary (C56) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | 0.1 | 0.2 | 1.0 | 4.0 | 9.6 | 20.0 | 32.1 | 37.7 |
| 2003..... | 5.0 | * | * | * | 0.1 | 0.2 | 1.0 | 4.0 | 10.4 | 19.5 | 32.6 | 38.4 |
| 2002..... | 5.1 | * | * | * | 0.1 | 0.2 | 1.0 | 4.3 | 10.3 | 19.7 | 33.4 | 38.1 |
| 2001..... | 5.1 | * | * | * | 0.1 | 0.3 | 1.0 | 4.1 | 10.3 | 20.0 | 33.2 | 39.6 |
| 2000..... | 5.0 | * | * | * | 0.1 | 0.3 | 1.0 | 4.2 | 10.1 | 19.6 | 33.2 | 40.9 |
| 1999..... | 4.9 | * | * | * | 0.1 | 0.2 | 1.1 | 4.2 | 10.2 | 19.6 | 31.7 | 37.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 9.9 | * | * | * | 0.1 | 0.5 | 2.1 | 7.8 | 18.5 | 36.8 | 53.7 | 54.6 |
| 2003..... | 9.9 | * | * | * | 0.1 | 0.4 | 2.0 | 7.8 | 20.0 | 35.7 | 54.4 | 55.4 |
| 2002..... | 10.0 | * | * | * | 0.2 | 0.5 | 2.1 | 8.4 | 19.9 | 36.1 | 55.6 | 54.6 |
| 2001..... | 9.9 | * | * | * | 0.1 | 0.5 | 2.1 | 8.1 | 19.9 | 36.6 | 55.0 | 56.1 |
| 2000..... | 9.8 | * | * | * | 0.1 | 0.5 | 2.1 | 8.3 | 19.4 | 35.7 | 54.8 | 57.6 |
| 1999..... | 9.6 | * | * | * | 0.1 | 0.4 | 2.2 | 8.3 | 19.6 | 35.7 | 52.3 | 52.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.8 | 2.8 | 6.5 | 12.9 | 20.9 | 28.1 |
| 2003..... | 1.6 | * | * | * | * | * | 0.8 | 2.7 | 6.4 | 12.7 | 22.0 | 28.5 |
| 2002..... | 1.6 | * | * | * | * | * | 0.8 | 3.2 | 5.6 | 12.3 | 24.2 | 23.8 |
| 2001..... | 1.6 | * | * | * | * | * | 0.8 | 2.7 | 7.1 | 14.4 | 20.0 | 30.6 |
| 2000..... | 1.6 | * | * | * | * | * | 0.6 | 3.0 | 6.5 | 14.3 | 24.9 | 32.5 |
| 1999..... | 1.4 | * | * | * | * | * | 0.6 | 2.7 | 6.1 | 12.2 | 20.8 | 23.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 1.7 | 5.5 | 12.5 | 23.3 | 35.3 | 42.5 |
| 2003..... | 3.3 | * | * | * | * | * | 1.7 | 5.5 | 12.2 | 22.8 | 37.1 | 43.0 |
| 2002..... | 3.3 | * | * | * | * | * | 1.7 | 6.3 | 10.7 | 22.1 | 40.8 | 35.6 |
| 2001..... | 3.4 | * | * | * | * | * | 1.6 | 5.3 | 13.4 | 25.8 | 33.5 | 45.6 |
| 2000..... | 3.4 | * | * | * | * | * | 1.2 | 5.9 | 12.2 | 25.5 | 41.6 | 48.5 |
| 1999..... | 2.9 | * | * | * | * | * | 1.2 | 5.3 | 11.5 | 21.7 | 34.7 | 35.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of ovary (C56) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 0.1 | 0.3 | 1.1 | 4.1 | 9.8 | 20.5 | 32.7 | 38.1 |
| 2003..... | 5.6 | * | * | * | 0.1 | 0.2 | 1.0 | 4.1 | 10.7 | 19.9 | 33.1 | 38.8 |
| 2002..... | 5.6 | * | * | * | 0.1 | 0.3 | 1.1 | 4.4 | 10.7 | 20.2 | 33.8 | 38.7 |
| 2001..... | 5.6 | * | * | * | 0.1 | 0.3 | 1.1 | 4.3 | 10.5 | 20.3 | 33.7 | 39.8 |
| 2000..... | 5.5 | * | * | * | * | 0.3 | 1.1 | 4.3 | 10.3 | 19.9 | 33.4 | 41.2 |
| 1999..... | 5.3 | * | * | * | 0.1 | 0.2 | 1.2 | 4.4 | 10.5 | 20.0 | 32.0 | 38.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 10.8 | * | * | * | 0.1 | 0.5 | 2.1 | 8.0 | 19.0 | 37.7 | 54.7 | 55.1 |
| 2003..... | 10.9 | * | * | * | 0.1 | 0.5 | 2.1 | 8.1 | 20.5 | 36.6 | 55.2 | 55.9 |
| 2002..... | 11.0 | * | * | * | 0.2 | 0.5 | 2.1 | 8.6 | 20.6 | 37.0 | 56.3 | 55.3 |
| 2001..... | 10.9 | * | * | * | 0.1 | 0.5 | 2.1 | 8.4 | 20.3 | 37.3 | 55.8 | 56.4 |
| 2000..... | 10.7 | * | * | * | * | 0.5 | 2.2 | 8.4 | 19.9 | 36.3 | 55.2 | 57.8 |
| 1999..... | 10.4 | * | * | * | 0.2 | 0.4 | 2.3 | 8.6 | 20.1 | 36.4 | 52.7 | 53.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | * | 0.2 | 1.1 | 4.2 | 10.2 | 21.0 | 33.9 | 39.1 |
| 2003..... | 6.3 | * | * | * | * | 0.3 | 1.1 | 4.2 | 11.1 | 20.5 | 34.2 | 39.5 |
| 2002..... | 6.3 | * | * | * | 0.1 | 0.3 | 1.1 | 4.6 | 11.1 | 20.9 | 34.8 | 39.6 |
| 2001..... | 6.2 | * | * | * | * | 0.3 | 1.1 | 4.5 | 10.8 | 21.0 | 34.8 | 40.6 |
| 2000..... | 6.2 | * | * | * | * | 0.3 | 1.1 | 4.5 | 10.8 | 20.5 | 34.5 | 42.2 |
| 1999..... | 6.0 | * | * | * | 0.1 | 0.2 | 1.2 | 4.5 | 10.9 | 20.5 | 32.7 | 39.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 12.2 | * | * | * | * | 0.5 | 2.2 | 8.3 | 19.8 | 39.1 | 57.0 | 56.5 |
| 2003..... | 12.3 | * | * | * | * | 0.5 | 2.2 | 8.4 | 21.6 | 38.1 | 57.2 | 56.8 |
| 2002..... | 12.4 | * | * | * | 0.2 | 0.5 | 2.2 | 9.1 | 21.7 | 38.7 | 58.0 | 56.7 |
| 2001..... | 12.2 | * | * | * | * | 0.5 | 2.2 | 9.0 | 21.0 | 38.8 | 57.8 | 57.5 |
| 2000..... | 12.0 | * | * | * | * | 0.6 | 2.3 | 8.9 | 21.0 | 37.7 | 57.2 | 59.2 |
| 1999..... | 11.7 | * | * | * | 0.2 | 0.4 | 2.4 | 9.0 | 21.2 | 37.7 | 54.1 | 54.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of ovary (C56) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 0.9 | 4.0 | 8.8 | 20.5 | 28.0 | 32.5 |
| 2003..... | 3.2 | * | * | * | * | * | 1.0 | 3.9 | 8.8 | 17.8 | 27.8 | 37.2 |
| 2002..... | 3.2 | * | * | * | * | * | 1.1 | 3.5 | 9.2 | 17.4 | 29.3 | 35.4 |
| 2001..... | 3.2 | * | * | * | * | * | 1.2 | 3.1 | 10.2 | 18.5 | 27.2 | 36.5 |
| 2000..... | 3.0 | * | * | * | * | * | 1.1 | 3.4 | 8.6 | 18.2 | 27.1 | 34.7 |
| 1999..... | 3.1 | * | * | * | * | * | 0.9 | 3.6 | 9.1 | 18.7 | 28.7 | 29.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | 1.8 | 7.5 | 15.9 | 35.2 | 43.8 | 45.4 |
| 2003..... | 6.2 | * | * | * | * | * | 1.9 | 7.2 | 15.9 | 30.5 | 43.5 | 51.6 |
| 2002..... | 6.1 | * | * | * | * | * | 2.0 | 6.5 | 16.6 | 29.8 | 45.9 | 48.9 |
| 2001..... | 6.1 | * | * | * | * | * | 2.2 | 5.7 | 18.4 | 31.5 | 42.5 | 50.0 |
| 2000..... | 5.8 | * | * | * | * | * | 2.0 | 6.3 | 15.6 | 31.1 | 42.4 | 47.5 |
| 1999..... | 5.8 | * | * | * | * | * | 1.6 | 6.7 | 16.4 | 31.9 | 44.8 | 40.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other and unspecified female genital organs (C57) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 1.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 1.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 0.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 1.1 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.4 | 1.6 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 1.7 | 1.5 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.5 | 1.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.5 | 2.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.1 | 1.7 | 1.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 1.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 1.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.0 | 0.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.2 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 1.5 | 1.5 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 1.7 | 1.5 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 1.6 | 1.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.4 | 2.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.7 | 1.1 | 1.7 | 1.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of urinary tract (C64-C68) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 8.8 | * | * | 0.1 | * | 0.2 | 0.9 | 4.4 | 13.9 | 32.3 | 65.9 | 102.7 |
| 2003..... | 8.7 | * | * | 0.1 | 0.1 | 0.2 | 1.0 | 4.3 | 14.2 | 32.5 | 65.3 | 101.4 |
| 2002..... | 8.8 | * | * | 0.1 | 0.1 | 0.2 | 1.0 | 4.6 | 14.0 | 34.6 | 65.1 | 102.0 |
| 2001..... | 8.7 | * | 0.1 | 0.1 | 0.1 | 0.2 | 1.1 | 4.6 | 14.4 | 33.8 | 63.8 | 103.9 |
| 2000..... | 8.7 | * | * | 0.1 | 0.1 | 0.1 | 1.0 | 4.5 | 14.3 | 33.5 | 64.0 | 106.0 |
| 1999..... | 8.5 | * | * | 0.1 | * | 0.2 | 1.0 | 4.5 | 14.4 | 33.5 | 61.8 | 103.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 11.7 | * | * | * | * | 0.2 | 1.2 | 6.4 | 20.9 | 49.0 | 105.3 | 182.8 |
| 2003..... | 11.6 | * | * | * | * | 0.2 | 1.3 | 6.4 | 21.4 | 48.5 | 105.0 | 178.8 |
| 2002..... | 11.6 | * | * | 0.1 | * | 0.2 | 1.3 | 6.7 | 21.1 | 51.4 | 103.3 | 183.6 |
| 2001..... | 11.6 | * | * | * | * | 0.2 | 1.4 | 6.8 | 21.6 | 51.5 | 102.5 | 192.4 |
| 2000..... | 11.4 | * | * | * | * | 0.1 | 1.3 | 6.6 | 21.3 | 50.7 | 103.3 | 198.9 |
| 1999..... | 11.2 | * | * | * | * | 0.2 | 1.4 | 6.6 | 21.5 | 50.6 | 100.8 | 188.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.0 | * | * | * | * | 0.1 | 0.6 | 2.4 | 7.5 | 18.4 | 39.4 | 66.7 |
| 2003..... | 6.0 | * | * | 0.1 | * | 0.1 | 0.6 | 2.3 | 7.5 | 19.2 | 38.7 | 67.2 |
| 2002..... | 6.1 | * | * | * | * | 0.1 | 0.6 | 2.6 | 7.5 | 20.5 | 39.7 | 66.6 |
| 2001..... | 6.0 | * | * | 0.1 | * | 0.2 | 0.8 | 2.6 | 7.9 | 19.2 | 38.3 | 66.9 |
| 2000..... | 6.0 | * | * | * | * | 0.1 | 0.7 | 2.6 | 7.9 | 19.4 | 38.5 | 68.1 |
| 1999..... | 5.9 | * | * | * | * | 0.1 | 0.7 | 2.5 | 7.9 | 19.5 | 36.5 | 68.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | * | 0.8 | 3.7 | 11.0 | 21.7 | 45.4 | 61.4 |
| 2003..... | 2.8 | * | * | * | * | * | 0.6 | 2.9 | 10.7 | 24.3 | 45.0 | 70.0 |
| 2002..... | 2.7 | * | * | * | * | * | 0.7 | 3.5 | 11.0 | 20.9 | 43.9 | 82.0 |
| 2001..... | 2.7 | * | * | * | * | * | 0.8 | 3.9 | 9.2 | 22.5 | 43.1 | 81.9 |
| 2000..... | 2.5 | * | * | * | * | * | 0.8 | 3.4 | 10.9 | 19.1 | 43.5 | 73.7 |
| 1999..... | 2.5 | * | * | * | * | * | 0.7 | 3.6 | 11.4 | 22.6 | 39.8 | 68.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.9 | 5.5 | 15.5 | 34.8 | 64.4 | 102.3 |
| 2003..... | 3.4 | * | * | * | * | * | 0.8 | 3.7 | 15.0 | 35.6 | 66.3 | 117.0 |
| 2002..... | 3.4 | * | * | * | * | * | 0.9 | 4.9 | 15.5 | 31.9 | 68.5 | 142.2 |
| 2001..... | 3.5 | * | * | * | * | * | 1.2 | 5.5 | 14.3 | 33.3 | 67.5 | 150.5 |
| 2000..... | 3.3 | * | * | * | * | * | 1.2 | 4.6 | 16.2 | 28.9 | 73.8 | 125.0 |
| 1999..... | 3.3 | * | * | * | * | * | 1.1 | 5.4 | 17.8 | 36.7 | 58.4 | 118.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | 0.7 | 1.9 | 7.0 | 11.2 | 32.4 | 40.3 |
| 2003..... | 2.1 | * | * | * | * | * | * | 2.0 | 6.8 | 15.3 | 30.3 | 46.1 |
| 2002..... | 2.0 | * | * | * | * | * | * | 2.1 | 6.9 | 12.2 | 27.0 | 52.2 |
| 2001..... | 1.9 | * | * | * | * | * | * | 2.5 | 4.7 | 13.9 | 26.6 | 48.4 |
| 2000..... | 1.7 | * | * | * | * | * | * | 2.2 | 6.3 | 11.4 | 23.1 | 48.5 |
| 1999..... | 1.7 | * | * | * | * | * | * | 1.9 | 5.6 | 11.5 | 27.4 | 43.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of urinary tract (C64-C68) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.8 | * | * | 0.1 | * | 0.2 | 0.9 | 4.5 | 14.1 | 33.1 | 67.0 | 104.4 |
| 2003..... | 9.7 | * | * | 0.1 | * | 0.2 | 1.0 | 4.5 | 14.4 | 33.1 | 66.2 | 102.6 |
| 2002..... | 9.7 | * | * | 0.1 | 0.1 | 0.2 | 1.0 | 4.7 | 14.2 | 35.4 | 66.0 | 102.6 |
| 2001..... | 9.6 | * | * | 0.1 | 0.1 | 0.2 | 1.1 | 4.7 | 14.8 | 34.5 | 64.6 | 104.5 |
| 2000..... | 9.5 | * | * | 0.1 | 0.1 | 0.1 | 1.0 | 4.6 | 14.5 | 34.3 | 64.7 | 106.8 |
| 1999..... | 9.3 | * | * | 0.1 | * | 0.2 | 1.0 | 4.5 | 14.6 | 34.1 | 62.6 | 104.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 13.1 | * | * | * | * | 0.2 | 1.2 | 6.5 | 21.3 | 49.9 | 107.5 | 186.6 |
| 2003..... | 12.9 | * | * | * | * | 0.2 | 1.4 | 6.7 | 21.8 | 49.2 | 106.8 | 181.5 |
| 2002..... | 12.9 | * | * | * | * | 0.2 | 1.4 | 6.9 | 21.4 | 52.6 | 104.9 | 185.0 |
| 2001..... | 12.8 | * | * | * | * | 0.2 | 1.4 | 6.9 | 22.1 | 52.5 | 103.9 | 193.6 |
| 2000..... | 12.6 | * | * | * | * | 0.1 | 1.3 | 6.7 | 21.6 | 51.8 | 104.2 | 201.3 |
| 1999..... | 12.3 | * | * | * | * | 0.3 | 1.4 | 6.6 | 21.7 | 51.3 | 102.3 | 190.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.6 | * | * | * | * | * | 0.6 | 2.5 | 7.5 | 18.9 | 39.7 | 67.7 |
| 2003..... | 6.6 | * | * | * | * | 0.1 | 0.6 | 2.3 | 7.5 | 19.5 | 39.2 | 67.9 |
| 2002..... | 6.7 | * | * | * | * | 0.1 | 0.6 | 2.7 | 7.5 | 21.1 | 40.2 | 67.1 |
| 2001..... | 6.6 | * | * | 0.2 | * | 0.2 | 0.9 | 2.5 | 8.1 | 19.5 | 38.8 | 67.4 |
| 2000..... | 6.6 | * | * | * | * | 0.1 | 0.7 | 2.6 | 8.0 | 19.9 | 39.0 | 68.6 |
| 1999..... | 6.4 | * | * | * | * | 0.1 | 0.7 | 2.5 | 8.1 | 19.9 | 36.8 | 69.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 11.1 | * | * | 0.1 | * | 0.1 | 0.9 | 4.6 | 14.8 | 34.5 | 69.5 | 107.3 |
| 2003..... | 10.9 | * | * | 0.1 | * | 0.1 | 1.0 | 4.5 | 14.9 | 34.5 | 68.2 | 105.4 |
| 2002..... | 11.0 | * | * | 0.1 | * | 0.1 | 1.0 | 4.8 | 14.7 | 36.8 | 67.7 | 104.5 |
| 2001..... | 10.8 | * | * | * | * | 0.1 | 1.1 | 4.8 | 15.2 | 36.0 | 66.1 | 106.9 |
| 2000..... | 10.7 | * | * | * | * | 0.1 | 1.0 | 4.8 | 15.1 | 35.5 | 66.1 | 110.2 |
| 1999..... | 10.4 | * | * | 0.1 | * | 0.1 | 1.0 | 4.5 | 15.0 | 35.0 | 64.0 | 105.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 15.1 | * | * | * | * | * | 1.1 | 6.7 | 22.3 | 52.3 | 112.3 | 194.9 |
| 2003..... | 14.7 | * | * | * | * | * | 1.4 | 6.8 | 22.6 | 51.3 | 110.5 | 189.6 |
| 2002..... | 14.7 | * | * | * | * | 0.2 | 1.4 | 7.1 | 22.3 | 54.9 | 108.1 | 192.4 |
| 2001..... | 14.5 | * | * | * | * | * | 1.4 | 7.1 | 22.7 | 55.0 | 106.9 | 202.1 |
| 2000..... | 14.3 | * | * | * | * | * | 1.4 | 7.0 | 22.4 | 53.5 | 107.0 | 209.7 |
| 1999..... | 13.8 | * | * | * | * | 0.2 | 1.3 | 6.6 | 22.4 | 53.0 | 105.5 | 197.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.4 | * | * | * | * | * | 0.6 | 2.5 | 7.7 | 19.2 | 40.4 | 68.0 |
| 2003..... | 7.3 | * | * | * | * | * | 0.6 | 2.3 | 7.5 | 20.1 | 39.6 | 68.3 |
| 2002..... | 7.4 | * | * | * | * | * | 0.6 | 2.6 | 7.5 | 21.4 | 40.6 | 66.5 |
| 2001..... | 7.2 | * | * | * | * | * | 0.8 | 2.5 | 8.1 | 19.8 | 38.9 | 67.4 |
| 2000..... | 7.3 | * | * | * | * | * | 0.7 | 2.7 | 8.2 | 20.2 | 39.1 | 70.1 |
| 1999..... | 7.0 | * | * | * | * | * | 0.7 | 2.6 | 8.1 | 20.0 | 36.8 | 68.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of urinary tract (C64-C68) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | * | 0.4 | 1.3 | 5.0 | 12.9 | 28.6 | 54.7 | 90.7 |
| 2003..... | 5.7 | * | * | * | * | * | 1.5 | 5.1 | 13.9 | 28.2 | 58.6 | 89.8 |
| 2002..... | 5.7 | * | * | * | * | 0.4 | 1.0 | 5.0 | 14.1 | 31.0 | 59.0 | 94.1 |
| 2001..... | 5.7 | * | * | * | * | 0.4 | 1.7 | 5.1 | 15.3 | 28.3 | 59.1 | 87.4 |
| 2000..... | 5.5 | * | * | * | * | * | 1.2 | 4.6 | 13.4 | 31.2 | 61.9 | 80.1 |
| 1999..... | 5.7 | * | * | * | * | 0.5 | 1.5 | 5.6 | 14.8 | 32.3 | 54.3 | 94.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | 1.8 | 7.0 | 18.7 | 40.2 | 79.0 | 122.9 |
| 2003..... | 6.8 | * | * | * | * | * | 2.2 | 7.3 | 19.7 | 41.1 | 88.6 | 126.8 |
| 2002..... | 6.5 | * | * | * | * | * | 1.5 | 6.4 | 20.3 | 41.0 | 86.2 | 135.6 |
| 2001..... | 6.7 | * | * | * | * | * | 2.0 | 7.4 | 22.2 | 40.0 | 86.4 | 127.0 |
| 2000..... | 6.6 | * | * | * | * | * | 1.2 | 6.4 | 19.5 | 45.5 | 93.2 | 139.7 |
| 1999..... | 6.8 | * | * | * | * | * | 2.1 | 8.6 | 21.1 | 43.6 | 80.0 | 139.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 0.8 | 3.2 | 8.1 | 20.4 | 41.1 | 77.9 |
| 2003..... | 4.6 | * | * | * | * | * | 1.0 | 3.3 | 9.3 | 19.0 | 41.7 | 75.5 |
| 2002..... | 4.9 | * | * | * | * | * | 0.7 | 3.8 | 9.1 | 24.0 | 43.6 | 78.2 |
| 2001..... | 4.8 | * | * | * | * | * | 1.5 | 3.2 | 9.8 | 20.0 | 43.8 | 72.7 |
| 2000..... | 4.4 | * | * | * | * | * | 1.1 | 3.1 | 8.5 | 21.1 | 44.3 | 58.0 |
| 1999..... | 4.7 | * | * | * | * | * | 0.9 | 3.1 | 9.7 | 24.4 | 39.9 | 77.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of kidney and renal pelvis (C64-C65) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | 0.1 | * | 0.1 | 0.6 | 3.0 | 8.7 | 16.7 | 26.8 | 32.7 |
| 2003..... | 4.2 | * | * | 0.1 | 0.1 | 0.1 | 0.7 | 2.9 | 9.0 | 16.9 | 27.0 | 33.2 |
| 2002..... | 4.2 | * | * | 0.1 | 0.0 | 0.1 | 0.7 | 3.3 | 8.8 | 17.7 | 26.3 | 32.5 |
| 2001..... | 4.2 | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.8 | 3.3 | 9.2 | 17.6 | 26.3 | 32.3 |
| 2000..... | 4.2 | * | * | 0.1 | 0.1 | 0.1 | 0.7 | 3.3 | 9.0 | 17.1 | 26.6 | 33.8 |
| 1999..... | 4.0 | * | * | 0.1 | * | 0.1 | 0.8 | 3.1 | 8.9 | 16.7 | 24.6 | 31.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | * | 0.2 | 0.8 | 4.3 | 12.8 | 23.9 | 38.2 | 47.0 |
| 2003..... | 5.4 | * | * | * | * | 0.2 | 1.0 | 4.3 | 13.3 | 23.7 | 39.5 | 48.7 |
| 2002..... | 5.4 | * | * | * | * | 0.2 | 1.0 | 4.8 | 13.0 | 25.1 | 37.9 | 50.1 |
| 2001..... | 5.4 | * | * | * | * | 0.1 | 1.1 | 4.9 | 13.4 | 25.2 | 37.4 | 52.9 |
| 2000..... | 5.3 | * | * | * | * | 0.1 | 1.0 | 4.8 | 13.1 | 24.2 | 38.2 | 55.1 |
| 1999..... | 5.0 | * | * | * | * | 0.2 | 1.1 | 4.5 | 12.9 | 23.6 | 35.7 | 49.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 0.4 | 1.7 | 5.0 | 10.6 | 19.1 | 26.3 |
| 2003..... | 3.1 | * | * | 0.1 | * | 0.1 | 0.4 | 1.6 | 4.9 | 11.2 | 18.7 | 26.4 |
| 2002..... | 3.1 | * | * | * | * | * | 0.4 | 1.8 | 4.9 | 11.6 | 18.7 | 24.9 |
| 2001..... | 3.1 | * | * | 0.1 | * | 0.1 | 0.5 | 1.9 | 5.3 | 11.3 | 19.0 | 23.7 |
| 2000..... | 3.1 | * | * | * | * | * | 0.5 | 1.9 | 5.3 | 11.3 | 18.9 | 25.1 |
| 1999..... | 3.0 | * | * | * | * | * | 0.5 | 1.8 | 5.1 | 11.0 | 17.4 | 24.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | 0.7 | 3.1 | 8.2 | 14.0 | 23.4 | 28.1 |
| 2003..... | 1.7 | * | * | * | * | * | 0.5 | 2.1 | 8.1 | 16.2 | 23.6 | 26.9 |
| 2002..... | 1.7 | * | * | * | * | * | 0.6 | 2.8 | 8.1 | 12.9 | 24.4 | 26.1 |
| 2001..... | 1.7 | * | * | * | * | * | 0.7 | 3.3 | 6.9 | 14.4 | 23.3 | 31.9 |
| 2000..... | 1.6 | * | * | * | * | * | 0.6 | 2.8 | 7.9 | 12.3 | 24.7 | 25.9 |
| 1999..... | 1.6 | * | * | * | * | * | 0.6 | 2.8 | 7.9 | 13.9 | 22.5 | 28.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 0.8 | 4.4 | 11.2 | 23.3 | 34.7 | 43.4 |
| 2003..... | 2.1 | * | * | * | * | * | 0.7 | 2.8 | 11.0 | 22.3 | 32.2 | 38.5 |
| 2002..... | 2.1 | * | * | * | * | * | 0.9 | 3.9 | 11.3 | 18.5 | 39.0 | 40.2 |
| 2001..... | 2.2 | * | * | * | * | * | 1.2 | 4.7 | 9.9 | 21.4 | 35.3 | 57.1 |
| 2000..... | 2.0 | * | * | * | * | * | 1.0 | 3.9 | 11.4 | 17.9 | 38.4 | 40.3 |
| 1999..... | 2.0 | * | * | * | * | * | 0.9 | 4.3 | 11.8 | 21.1 | 29.7 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.7 | 5.5 | 6.5 | 15.6 | 20.2 |
| 2003..... | 1.4 | * | * | * | * | * | * | 1.5 | 5.6 | 11.3 | 17.6 | 21.1 |
| 2002..... | 1.2 | * | * | * | * | * | * | 1.7 | 5.2 | 8.4 | 14.3 | 19.1 |
| 2001..... | 1.2 | * | * | * | * | * | * | 1.9 | 4.2 | 8.8 | 15.1 | 19.5 |
| 2000..... | 1.1 | * | * | * | * | * | * | 1.8 | 4.7 | 7.8 | 15.5 | * |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.4 | 4.4 | 8.3 | 17.7 | 23.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of kidney and renal pelvis (C64-C65) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | 0.1 | * | 0.1 | 0.6 | 3.0 | 8.7 | 16.8 | 26.9 | 32.9 |
| 2003..... | 4.6 | * | * | 0.1 | * | 0.2 | 0.8 | 3.0 | 9.0 | 16.9 | 27.2 | 33.4 |
| 2002..... | 4.6 | * | * | 0.1 | * | 0.1 | 0.7 | 3.3 | 8.8 | 18.0 | 26.4 | 32.7 |
| 2001..... | 4.6 | * | * | 0.1 | 0.1 | 0.1 | 0.8 | 3.3 | 9.4 | 17.8 | 26.4 | 32.3 |
| 2000..... | 4.5 | * | * | 0.1 | * | 0.1 | 0.8 | 3.3 | 9.1 | 17.3 | 26.6 | 33.9 |
| 1999..... | 4.3 | * | * | 0.1 | * | 0.1 | 0.8 | 3.2 | 8.9 | 16.8 | 24.6 | 31.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | 0.2 | 0.8 | 4.3 | 12.9 | 23.9 | 38.4 | 47.1 |
| 2003..... | 5.9 | * | * | * | * | 0.2 | 1.1 | 4.4 | 13.5 | 23.7 | 39.8 | 49.0 |
| 2002..... | 5.9 | * | * | * | * | 0.2 | 1.0 | 4.9 | 13.0 | 25.5 | 37.7 | 50.4 |
| 2001..... | 5.9 | * | * | * | * | 0.1 | 1.0 | 4.9 | 13.7 | 25.4 | 37.5 | 52.6 |
| 2000..... | 5.7 | * | * | * | * | 0.1 | 1.0 | 4.8 | 13.2 | 24.4 | 38.1 | 55.4 |
| 1999..... | 5.4 | * | * | * | * | 0.2 | 1.1 | 4.6 | 13.0 | 23.7 | 35.9 | 49.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 0.4 | 1.7 | 4.9 | 10.9 | 19.3 | 26.5 |
| 2003..... | 3.4 | * | * | * | * | 0.1 | 0.4 | 1.7 | 4.9 | 11.1 | 18.7 | 26.6 |
| 2002..... | 3.4 | * | * | * | * | * | 0.4 | 1.8 | 4.9 | 11.8 | 18.8 | 25.1 |
| 2001..... | 3.4 | * | * | 0.1 | * | 0.1 | 0.6 | 1.8 | 5.4 | 11.4 | 19.1 | 23.8 |
| 2000..... | 3.4 | * | * | * | * | * | 0.5 | 1.9 | 5.3 | 11.4 | 19.1 | 25.2 |
| 1999..... | 3.2 | * | * | * | * | * | 0.5 | 1.8 | 5.2 | 11.2 | 17.3 | 24.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | 0.1 | * | 0.1 | 0.6 | 3.1 | 9.1 | 17.3 | 27.5 | 33.8 |
| 2003..... | 5.1 | * | * | 0.1 | * | 0.1 | 0.7 | 3.1 | 9.2 | 17.4 | 27.8 | 34.0 |
| 2002..... | 5.1 | * | * | 0.1 | * | 0.1 | 0.7 | 3.4 | 9.1 | 18.7 | 26.8 | 33.5 |
| 2001..... | 5.1 | * | * | * | * | 0.1 | 0.7 | 3.4 | 9.6 | 18.4 | 26.8 | 32.8 |
| 2000..... | 5.0 | * | * | * | * | 0.1 | 0.7 | 3.4 | 9.3 | 17.7 | 26.9 | 34.6 |
| 1999..... | 4.7 | * | * | 0.1 | * | 0.1 | 0.7 | 3.1 | 9.2 | 17.2 | 24.9 | 31.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | 0.8 | 4.5 | 13.3 | 24.4 | 39.1 | 48.6 |
| 2003..... | 6.6 | * | * | * | * | * | 1.0 | 4.5 | 13.7 | 24.3 | 40.7 | 50.4 |
| 2002..... | 6.5 | * | * | * | * | * | 1.0 | 5.0 | 13.3 | 26.3 | 38.4 | 52.4 |
| 2001..... | 6.5 | * | * | * | * | * | 1.0 | 4.9 | 13.8 | 26.2 | 37.7 | 54.3 |
| 2000..... | 6.3 | * | * | * | * | * | 1.0 | 4.9 | 13.5 | 24.6 | 38.5 | 56.9 |
| 1999..... | 6.0 | * | * | * | * | * | 1.0 | 4.5 | 13.3 | 24.0 | 36.5 | 50.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 0.3 | 1.8 | 5.1 | 11.2 | 19.6 | 27.2 |
| 2003..... | 3.8 | * | * | * | * | * | 0.4 | 1.7 | 4.8 | 11.6 | 19.1 | 26.8 |
| 2002..... | 3.8 | * | * | * | * | * | 0.4 | 1.8 | 5.1 | 12.2 | 19.0 | 25.3 |
| 2001..... | 3.8 | * | * | * | * | * | 0.5 | 1.8 | 5.5 | 11.8 | 19.5 | 23.8 |
| 2000..... | 3.8 | * | * | * | * | * | 0.5 | 1.9 | 5.4 | 11.8 | 19.3 | 25.7 |
| 1999..... | 3.5 | * | * | * | * | * | 0.5 | 1.8 | 5.3 | 11.5 | 17.3 | 24.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of kidney and renal pelvis (C64-C65) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | * | 0.8 | 3.2 | 8.2 | 16.0 | 26.3 | 28.8 |
| 2003..... | 3.0 | * | * | * | * | * | 1.1 | 3.3 | 9.7 | 14.8 | 24.2 | 34.2 |
| 2002..... | 2.9 | * | * | * | * | * | 0.8 | 3.6 | 8.8 | 15.8 | 26.5 | 28.1 |
| 2001..... | 3.1 | * | * | * | * | * | 1.3 | 3.8 | 9.7 | 15.3 | 26.0 | 30.0 |
| 2000..... | 2.9 | * | * | * | * | * | 0.9 | 3.5 | 9.0 | 16.8 | 25.9 | 29.6 |
| 1999..... | 3.0 | * | * | * | * | 0.4 | 1.1 | 3.9 | 8.7 | 16.4 | 23.9 | 31.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | * | 1.1 | 4.6 | 12.4 | 24.1 | 38.2 | 39.6 |
| 2003..... | 3.8 | * | * | * | * | * | 1.7 | 5.0 | 14.1 | 22.3 | 37.2 | 44.7 |
| 2002..... | 3.6 | * | * | * | * | * | 1.2 | 5.0 | 14.4 | 22.1 | 36.6 | 39.4 |
| 2001..... | 4.0 | * | * | * | * | * | 1.5 | 5.5 | 15.3 | 22.7 | 40.8 | 46.2 |
| 2000..... | 3.7 | * | * | * | * | * | 1.0 | 5.0 | 13.4 | 26.8 | 38.3 | 49.3 |
| 1999..... | 3.7 | * | * | * | * | * | 1.6 | 6.0 | 13.3 | 23.9 | 32.5 | 49.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | * | 1.9 | 4.9 | 10.3 | 19.6 | 24.5 |
| 2003..... | 2.3 | * | * | * | * | * | 0.7 | 1.8 | 6.2 | 9.5 | 16.8 | 30.1 |
| 2002..... | 2.3 | * | * | * | * | * | * | 2.5 | 4.4 | 11.3 | 20.7 | 23.8 |
| 2001..... | 2.3 | * | * | * | * | * | 1.1 | 2.2 | 5.3 | 10.0 | 17.7 | 24.0 |
| 2000..... | 2.2 | * | * | * | * | * | 0.8 | 2.2 | 5.4 | 9.7 | 18.9 | 22.2 |
| 1999..... | 2.2 | * | * | * | * | * | * | 2.1 | 5.0 | 11.0 | 19.1 | 24.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of bladder (C67) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | * | * | 0.3 | 1.4 | 5.2 | 15.4 | 38.2 | 69.4 |
| 2003..... | 4.8 | * | * | * | * | * | 0.2 | 1.3 | 5.2 | 15.4 | 37.1 | 65.6 |
| 2002..... | 4.9 | * | * | * | * | * | 0.3 | 1.3 | 5.1 | 16.4 | 37.8 | 66.9 |
| 2001..... | 4.8 | * | * | * | * | * | 0.3 | 1.3 | 5.2 | 15.8 | 36.4 | 69.6 |
| 2000..... | 4.7 | * | * | * | * | * | 0.3 | 1.2 | 5.2 | 16.1 | 36.3 | 70.3 |
| 1999..... | 4.7 | * | * | * | * | * | 0.2 | 1.3 | 5.4 | 16.3 | 36.0 | 69.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | * | * | * | 0.4 | 2.1 | 8.1 | 24.9 | 66.6 | 136.5 |
| 2003..... | 6.8 | * | * | * | * | * | 0.3 | 2.1 | 8.1 | 24.7 | 64.5 | 127.2 |
| 2002..... | 6.8 | * | * | * | * | * | 0.4 | 1.9 | 7.9 | 26.0 | 64.7 | 129.8 |
| 2001..... | 6.7 | * | * | * | * | * | 0.4 | 1.9 | 8.1 | 26.0 | 63.8 | 136.4 |
| 2000..... | 6.6 | * | * | * | * | * | 0.3 | 1.8 | 8.1 | 26.1 | 63.6 | 141.3 |
| 1999..... | 6.6 | * | * | * | * | * | 0.3 | 2.0 | 8.4 | 26.3 | 63.8 | 136.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 7.5 | 19.1 | 39.4 |
| 2003..... | 2.9 | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 7.6 | 18.8 | 38.6 |
| 2002..... | 3.1 | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 8.5 | 20.0 | 39.8 |
| 2001..... | 3.0 | * | * | * | * | * | 0.3 | 0.6 | 2.5 | 7.4 | 18.3 | 41.9 |
| 2000..... | 3.0 | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 7.8 | 18.5 | 41.6 |
| 1999..... | 3.0 | * | * | * | * | * | 0.2 | 0.6 | 2.6 | 8.1 | 17.9 | 42.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | * | 0.3 | 1.4 | 5.5 | 16.4 | 40.1 | 71.1 |
| 2003..... | 5.5 | * | * | * | * | * | 0.2 | 1.3 | 5.5 | 16.3 | 38.3 | 67.6 |
| 2002..... | 5.6 | * | * | * | * | * | 0.3 | 1.4 | 5.3 | 17.1 | 39.1 | 68.0 |
| 2001..... | 5.5 | * | * | * | * | * | 0.3 | 1.3 | 5.4 | 16.6 | 37.4 | 71.5 |
| 2000..... | 5.4 | * | * | * | * | * | 0.3 | 1.3 | 5.5 | 16.8 | 37.1 | 72.8 |
| 1999..... | 5.4 | * | * | * | * | * | 0.2 | 1.3 | 5.5 | 16.8 | 37.1 | 70.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 8.3 | * | * | * | * | * | 0.3 | 2.2 | 8.7 | 26.7 | 70.5 | 143.1 |
| 2003..... | 7.9 | * | * | * | * | * | 0.3 | 2.2 | 8.6 | 26.1 | 67.2 | 133.6 |
| 2002..... | 7.9 | * | * | * | * | * | 0.4 | 2.0 | 8.5 | 27.4 | 67.2 | 135.2 |
| 2001..... | 7.8 | * | * | * | * | * | 0.4 | 2.0 | 8.6 | 27.6 | 66.6 | 143.0 |
| 2000..... | 7.7 | * | * | * | * | * | 0.3 | 2.0 | 8.6 | 27.5 | 65.9 | 148.0 |
| 1999..... | 7.6 | * | * | * | * | * | 0.3 | 2.0 | 8.7 | 27.6 | 66.2 | 143.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 7.5 | 19.4 | 38.9 |
| 2003..... | 3.3 | * | * | * | * | * | 0.2 | 0.5 | 2.5 | 7.8 | 18.7 | 38.6 |
| 2002..... | 3.4 | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 8.4 | 20.2 | 39.0 |
| 2001..... | 3.2 | * | * | * | * | * | 0.3 | 0.6 | 2.4 | 7.4 | 18.1 | 41.7 |
| 2000..... | 3.3 | * | * | * | * | * | 0.2 | 0.7 | 2.6 | 7.8 | 18.2 | 42.5 |
| 1999..... | 3.2 | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 7.8 | 18.0 | 41.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of bladder (C67) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 0.5 | 1.6 | 4.2 | 11.9 | 27.6 | 61.3 |
| 2003..... | 2.5 | * | * | * | * | * | 0.4 | 1.7 | 4.1 | 12.3 | 32.8 | 53.8 |
| 2002..... | 2.6 | * | * | * | * | * | * | 1.1 | 4.5 | 14.7 | 31.2 | 64.1 |
| 2001..... | 2.4 | * | * | * | * | * | 0.4 | 1.3 | 5.1 | 12.1 | 31.4 | 55.6 |
| 2000..... | 2.4 | * | * | * | * | * | * | 1.1 | 4.2 | 13.5 | 34.9 | 49.6 |
| 1999..... | 2.6 | * | * | * | * | * | * | 1.5 | 5.6 | 15.2 | 29.4 | 60.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | * | 2.3 | 6.0 | 15.0 | 40.2 | 83.3 |
| 2003..... | 2.9 | * | * | * | * | * | * | 2.2 | 5.5 | 17.8 | 49.6 | 79.9 |
| 2002..... | 2.8 | * | * | * | * | * | * | 1.3 | 5.5 | 18.7 | 48.1 | 91.9 |
| 2001..... | 2.6 | * | * | * | * | * | * | 1.8 | 6.5 | 16.5 | 43.2 | 78.5 |
| 2000..... | 2.8 | * | * | * | * | * | * | 1.4 | 5.9 | 17.8 | 52.8 | 88.1 |
| 1999..... | 3.0 | * | * | * | * | * | * | 2.6 | 7.4 | 19.2 | 46.6 | 86.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | * | 1.0 | 2.7 | 9.7 | 20.4 | 52.6 |
| 2003..... | 2.1 | * | * | * | * | * | * | 1.2 | 2.9 | 8.3 | 23.3 | 43.7 |
| 2002..... | 2.4 | * | * | * | * | * | * | 0.9 | 3.7 | 11.8 | 21.7 | 53.5 |
| 2001..... | 2.3 | * | * | * | * | * | * | * | 3.9 | 9.0 | 24.8 | 47.1 |
| 2000..... | 2.1 | * | * | * | * | * | * | * | 2.7 | 10.4 | 24.8 | 35.3 |
| 1999..... | 2.3 | * | * | * | * | * | * | * | 4.2 | 12.3 | 19.8 | 51.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other and unspecified sites within the urinary tract (C66,C68) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 1.8 | 2.2 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.0 | 3.5 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 1.8 | 3.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 1.8 | 2.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 1.9 | 2.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 1.9 | 3.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.2 | 2.5 | 2.9 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 2.5 | 5.3 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 2.5 | 4.8 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.2 | 2.6 | 4.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.3 | 2.5 | 4.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.3 | 2.6 | 4.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 1.8 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 2.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.3 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 1.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 2.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.9 | 2.3 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.1 | 3.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 1.8 | 3.0 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 1.9 | 2.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.0 | 2.8 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.0 | 3.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.2 | 2.7 | 3.2 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.6 | 5.7 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.5 | 1.2 | 2.6 | 4.8 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 1.2 | 2.7 | 4.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.4 | 2.6 | 4.8 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.4 | 2.8 | 4.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 1.9 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 2.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.9 | 1.4 | 2.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 2.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 3.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasm of eye and adnexa (C69) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.3 | 0.5 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.3 | * | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.6 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.2 | 0.3 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.2 | * | 0.4 | 0.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 0.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 0.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.3 | * | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.6 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.5 | 0.5 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.4 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 0.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.4 | * | 0.6 | 0.8 | 0.5 | 0.9 | 2.0 | 4.6 | 9.5 | 15.7 | 19.6 | 16.6 |
| 2003..... | 4.4 | * | 0.7 | 0.8 | 0.6 | 1.0 | 2.4 | 4.8 | 9.7 | 15.5 | 19.5 | 15.2 |
| 2002..... | 4.4 | * | 0.7 | 0.8 | 0.5 | 1.0 | 2.2 | 5.1 | 9.6 | 16.2 | 19.1 | 16.4 |
| 2001..... | 4.4 | * | 0.7 | 0.7 | 0.6 | 1.1 | 2.3 | 5.0 | 9.7 | 15.4 | 20.4 | 16.4 |
| 2000..... | 4.5 | 0.5 | 0.7 | 0.8 | 0.6 | 1.0 | 2.4 | 5.1 | 9.6 | 16.5 | 20.0 | 16.9 |
| 1999..... | 4.6 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.4 | 5.1 | 10.3 | 16.9 | 20.4 | 16.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.9 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.5 | 5.8 | 11.8 | 18.4 | 23.7 | 19.4 |
| 2003..... | 5.0 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.9 | 6.2 | 11.4 | 18.6 | 24.6 | 20.0 |
| 2002..... | 5.0 | * | 0.8 | 0.9 | 0.7 | 1.1 | 2.8 | 6.2 | 11.7 | 20.2 | 23.8 | 20.1 |
| 2001..... | 5.0 | * | 0.7 | 0.7 | 0.6 | 1.3 | 2.8 | 6.1 | 12.1 | 19.2 | 25.8 | 20.1 |
| 2000..... | 5.1 | * | 0.7 | 0.8 | 0.7 | 1.3 | 3.1 | 6.2 | 11.8 | 20.2 | 24.3 | 22.4 |
| 1999..... | 5.1 | * | 0.7 | 0.7 | 0.7 | 1.4 | 2.8 | 6.2 | 12.4 | 20.8 | 25.5 | 20.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | 0.6 | 0.8 | 0.4 | 0.7 | 1.6 | 3.4 | 7.3 | 13.4 | 16.8 | 15.4 |
| 2003..... | 3.9 | * | 0.6 | 0.8 | 0.5 | 0.9 | 2.0 | 3.5 | 8.1 | 12.8 | 16.1 | 13.1 |
| 2002..... | 3.9 | * | 0.7 | 0.7 | 0.4 | 0.8 | 1.7 | 4.1 | 7.7 | 12.8 | 16.0 | 14.8 |
| 2001..... | 3.9 | * | 0.6 | 0.8 | 0.5 | 0.8 | 1.7 | 3.9 | 7.6 | 12.3 | 16.8 | 14.8 |
| 2000..... | 3.9 | * | 0.6 | 0.7 | 0.5 | 0.7 | 1.7 | 4.0 | 7.6 | 13.4 | 17.3 | 14.7 |
| 1999..... | 4.1 | * | 0.8 | 0.8 | 0.5 | 0.8 | 1.9 | 4.1 | 8.4 | 13.7 | 17.1 | 14.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.7 | 0.5 | 0.4 | 1.1 | 2.4 | 5.9 | 9.0 | 12.4 | 14.3 |
| 2003..... | 1.7 | * | 0.6 | 0.8 | 0.3 | 0.6 | 1.1 | 2.5 | 5.9 | 9.5 | 15.3 | * |
| 2002..... | 1.7 | * | 0.7 | 0.7 | 0.4 | 0.5 | 0.9 | 3.4 | 5.8 | 10.0 | 13.7 | 13.3 |
| 2001..... | 1.7 | * | 0.7 | 0.5 | 0.4 | 0.6 | 1.4 | 2.6 | 5.9 | 10.3 | 14.8 | * |
| 2000..... | 1.8 | * | * | 0.9 | 0.4 | 0.7 | 1.5 | 2.9 | 6.2 | 9.8 | 14.2 | 15.9 |
| 1999..... | 1.7 | * | 0.9 | 0.7 | * | 0.5 | 1.3 | 3.0 | 6.0 | 10.0 | 14.0 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.6 | 0.6 | * | 1.4 | 2.9 | 7.2 | 10.1 | 15.4 | * |
| 2003..... | 1.8 | * | * | 0.9 | * | 0.6 | 1.2 | 3.2 | 6.3 | 11.8 | 16.5 | * |
| 2002..... | 1.8 | * | * | 0.7 | * | * | 1.0 | 4.4 | 6.6 | 12.4 | 16.0 | * |
| 2001..... | 1.8 | * | * | * | * | 0.6 | 1.4 | 3.0 | 6.8 | 14.4 | 17.9 | * |
| 2000..... | 1.9 | * | * | 1.2 | * | 0.8 | 1.5 | 2.6 | 7.7 | 12.5 | 18.7 | * |
| 1999..... | 1.7 | * | * | 0.7 | * | * | 1.8 | 2.9 | 7.8 | 10.5 | 16.2 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | 0.9 | * | * | 0.8 | 1.8 | 4.7 | 8.1 | 10.4 | 15.1 |
| 2003..... | 1.6 | * | * | 0.7 | * | * | 0.9 | 1.9 | 5.6 | 7.6 | 14.5 | * |
| 2002..... | 1.5 | * | * | 0.7 | * | * | 0.9 | 2.3 | 5.1 | 8.1 | 12.1 | * |
| 2001..... | 1.6 | * | * | 0.6 | * | * | 1.4 | 2.2 | 5.1 | 7.1 | 12.7 | * |
| 2000..... | 1.7 | * | * | 0.7 | * | * | 1.5 | 3.2 | 4.9 | 7.6 | 11.2 | * |
| 1999..... | 1.7 | * | * | 0.8 | * | * | 0.8 | 3.1 | 4.5 | 9.6 | 12.5 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.8 | * | 0.7 | 0.8 | 0.5 | 1.0 | 2.2 | 4.8 | 9.7 | 16.2 | 20.0 | 16.7 |
| 2003..... | 4.9 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.6 | 5.0 | 10.0 | 15.9 | 19.7 | 15.4 |
| 2002..... | 4.9 | * | 0.7 | 0.8 | 0.5 | 1.1 | 2.4 | 5.3 | 9.9 | 16.6 | 19.4 | 16.5 |
| 2001..... | 4.8 | * | 0.6 | 0.8 | 0.6 | 1.2 | 2.4 | 5.2 | 10.0 | 15.7 | 20.6 | 16.5 |
| 2000..... | 4.9 | * | 0.7 | 0.7 | 0.6 | 1.1 | 2.5 | 5.3 | 9.8 | 16.9 | 20.2 | 16.9 |
| 1999..... | 5.0 | * | 0.7 | 0.8 | 0.6 | 1.2 | 2.5 | 5.3 | 10.6 | 17.3 | 20.6 | 16.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.4 | * | 0.7 | 0.9 | 0.6 | 1.2 | 2.7 | 6.1 | 12.2 | 19.0 | 24.2 | 19.7 |
| 2003..... | 5.5 | * | 0.7 | 0.8 | 0.7 | 1.3 | 3.1 | 6.5 | 11.8 | 19.1 | 25.0 | 20.1 |
| 2002..... | 5.5 | * | 0.8 | 0.9 | 0.7 | 1.3 | 3.0 | 6.4 | 12.0 | 20.7 | 24.1 | 20.2 |
| 2001..... | 5.5 | * | 0.6 | 0.7 | 0.7 | 1.5 | 3.0 | 6.4 | 12.5 | 19.4 | 26.1 | 20.5 |
| 2000..... | 5.5 | * | 0.7 | 0.8 | 0.7 | 1.4 | 3.3 | 6.5 | 12.1 | 20.6 | 24.5 | 22.3 |
| 1999..... | 5.6 | * | 0.7 | 0.8 | 0.7 | 1.5 | 3.0 | 6.5 | 12.7 | 21.4 | 25.8 | 21.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.2 | * | 0.6 | 0.7 | 0.4 | 0.8 | 1.7 | 3.5 | 7.5 | 13.8 | 17.2 | 15.4 |
| 2003..... | 4.3 | * | 0.6 | 0.8 | 0.5 | 1.0 | 2.1 | 3.7 | 8.3 | 13.2 | 16.2 | 13.4 |
| 2002..... | 4.2 | * | 0.7 | 0.7 | 0.4 | 0.9 | 1.8 | 4.2 | 7.9 | 13.1 | 16.2 | 14.9 |
| 2001..... | 4.2 | * | 0.6 | 0.8 | 0.5 | 0.8 | 1.8 | 4.1 | 7.7 | 12.7 | 17.0 | 14.9 |
| 2000..... | 4.2 | * | 0.7 | 0.7 | 0.5 | 0.7 | 1.8 | 4.1 | 7.8 | 13.8 | 17.5 | 14.7 |
| 1999..... | 4.4 | * | 0.7 | 0.8 | 0.5 | 0.9 | 2.0 | 4.2 | 8.6 | 14.0 | 17.2 | 14.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.6 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.5 | 5.3 | 10.7 | 17.6 | 21.3 | 17.4 |
| 2003..... | 5.6 | * | 0.7 | 0.8 | 0.7 | 1.3 | 3.0 | 5.5 | 10.8 | 17.1 | 20.9 | 15.9 |
| 2002..... | 5.6 | * | 0.6 | 0.8 | 0.6 | 1.2 | 2.8 | 5.9 | 10.7 | 17.9 | 20.4 | 17.3 |
| 2001..... | 5.6 | * | 0.7 | 0.8 | 0.6 | 1.3 | 2.7 | 5.7 | 10.8 | 16.8 | 21.9 | 17.2 |
| 2000..... | 5.6 | * | 0.7 | 0.8 | 0.6 | 1.2 | 2.8 | 5.9 | 10.7 | 18.3 | 21.2 | 17.5 |
| 1999..... | 5.7 | * | 0.7 | 0.8 | 0.6 | 1.4 | 2.8 | 5.8 | 11.5 | 18.7 | 21.7 | 16.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.3 | * | 0.8 | 0.9 | 0.7 | 1.4 | 3.0 | 6.6 | 13.2 | 20.8 | 25.4 | 21.0 |
| 2003..... | 6.3 | * | 0.8 | 0.8 | 0.7 | 1.5 | 3.5 | 7.0 | 12.8 | 20.4 | 26.3 | 21.0 |
| 2002..... | 6.4 | * | 0.7 | 0.9 | 0.8 | 1.4 | 3.5 | 7.1 | 12.8 | 22.3 | 25.2 | 21.4 |
| 2001..... | 6.3 | * | 0.7 | 0.7 | 0.7 | 1.7 | 3.5 | 7.0 | 13.4 | 20.9 | 27.5 | 20.9 |
| 2000..... | 6.3 | * | 0.7 | 0.8 | 0.7 | 1.5 | 3.7 | 7.1 | 13.1 | 22.2 | 25.7 | 23.5 |
| 1999..... | 6.4 | * | 0.8 | 0.8 | 0.7 | 1.8 | 3.3 | 7.1 | 13.7 | 23.0 | 27.2 | 22.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.9 | * | 0.7 | 0.7 | 0.4 | 0.9 | 1.9 | 3.9 | 8.3 | 14.9 | 18.5 | 15.8 |
| 2003..... | 4.9 | * | 0.6 | 0.9 | 0.6 | 1.1 | 2.5 | 4.1 | 9.0 | 14.3 | 17.3 | 13.7 |
| 2002..... | 4.9 | * | 0.6 | 0.8 | 0.4 | 1.1 | 2.1 | 4.7 | 8.7 | 14.2 | 17.1 | 15.5 |
| 2001..... | 4.8 | * | 0.6 | 0.9 | 0.5 | 0.9 | 2.0 | 4.5 | 8.4 | 13.4 | 18.2 | 15.7 |
| 2000..... | 4.9 | * | 0.7 | 0.7 | 0.5 | 0.9 | 1.9 | 4.6 | 8.4 | 15.0 | 18.3 | 15.1 |
| 1999..... | 5.0 | * | 0.7 | 0.7 | 0.5 | 1.1 | 2.3 | 4.6 | 9.5 | 15.1 | 18.1 | 14.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | 0.8 | 0.5 | 0.7 | 1.3 | 2.9 | 5.0 | 7.9 | 10.1 | 10.6 |
| 2003..... | 2.2 | * | * | 0.7 | 0.6 | 0.6 | 1.5 | 3.1 | 5.6 | 9.0 | 10.6 | 10.7 |
| 2002..... | 2.1 | * | 1.1 | 0.8 | 0.4 | 0.6 | 1.2 | 2.8 | 5.6 | 8.6 | 11.9 | 9.7 |
| 2001..... | 2.1 | * | * | 0.7 | 0.5 | 0.5 | 1.1 | 3.1 | 5.9 | 9.4 | 9.9 | 11.9 |
| 2000..... | 2.2 | * | * | 0.8 | 0.6 | 0.7 | 1.6 | 2.9 | 5.8 | 8.2 | 12.1 | 10.5 |
| 1999..... | 2.2 | * | * | 0.8 | 0.7 | 0.7 | 1.4 | 3.3 | 6.1 | 8.2 | 11.7 | 10.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | 0.7 | * | * | 1.9 | 4.0 | 6.9 | 8.1 | 14.5 | * |
| 2003..... | 2.5 | * | * | 0.8 | 0.7 | * | 2.1 | 4.1 | 6.8 | 11.9 | 15.6 | * |
| 2002..... | 2.3 | * | * | 1.1 | * | 0.9 | 1.3 | 2.7 | 8.1 | 10.2 | 16.1 | * |
| 2001..... | 2.2 | * | * | 0.8 | * | * | 1.3 | 3.7 | 7.9 | 10.4 | 12.0 | * |
| 2000..... | 2.4 | * | * | 0.9 | 0.8 | 1.1 | 2.0 | 3.7 | 7.2 | 9.9 | 13.3 | * |
| 1999..... | 2.4 | * | * | 0.7 | 0.9 | 0.9 | 1.8 | 4.5 | 7.4 | 10.2 | 12.5 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | 0.8 | * | * | 0.8 | 1.9 | 3.5 | 7.8 | 7.6 | 11.6 |
| 2003..... | 1.9 | * | * | 0.7 | * | * | 0.9 | 2.3 | 4.7 | 6.9 | 7.8 | 12.0 |
| 2002..... | 1.9 | * | * | * | * | * | 1.1 | 2.8 | 3.7 | 7.5 | 9.5 | 10.5 |
| 2001..... | 1.9 | * | * | * | * | * | 1.0 | 2.5 | 4.2 | 8.7 | 8.8 | 9.8 |
| 2000..... | 1.9 | * | * | 0.7 | * | * | 1.2 | 2.2 | 4.7 | 7.1 | 11.5 | 10.5 |
| 1999..... | 1.9 | * | * | 0.9 | * | * | 1.0 | 2.2 | 5.1 | 6.8 | 11.2 | 9.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of brain (C71) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | * | 0.6 | 0.8 | 0.5 | 0.9 | 2.0 | 4.5 | 9.3 | 15.5 | 19.3 | 16.1 |
| 2003..... | 4.4 | * | 0.6 | 0.8 | 0.6 | 1.0 | 2.4 | 4.7 | 9.5 | 15.2 | 19.1 | 14.6 |
| 2002..... | 4.4 | * | 0.7 | 0.8 | 0.5 | 0.9 | 2.2 | 5.0 | 9.5 | 15.8 | 18.7 | 16.0 |
| 2001..... | 4.3 | * | 0.6 | 0.7 | 0.5 | 1.0 | 2.3 | 4.9 | 9.6 | 15.1 | 20.0 | 16.0 |
| 2000..... | 4.4 | 0.5 | 0.7 | 0.8 | 0.6 | 1.0 | 2.4 | 5.0 | 9.5 | 16.2 | 19.7 | 16.3 |
| 1999..... | 4.5 | * | 0.7 | 0.7 | 0.5 | 1.1 | 2.3 | 5.0 | 10.1 | 16.6 | 19.9 | 15.5 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.8 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.5 | 5.7 | 11.6 | 18.2 | 23.3 | 19.1 |
| 2003..... | 4.9 | * | 0.7 | 0.8 | 0.6 | 1.1 | 2.9 | 6.1 | 11.2 | 18.4 | 24.3 | 19.2 |
| 2002..... | 4.9 | * | 0.8 | 0.9 | 0.6 | 1.1 | 2.7 | 6.1 | 11.5 | 19.7 | 23.1 | 19.7 |
| 2001..... | 4.9 | * | 0.6 | 0.7 | 0.6 | 1.3 | 2.8 | 6.0 | 11.9 | 18.8 | 25.3 | 19.7 |
| 2000..... | 5.0 | * | 0.7 | 0.8 | 0.6 | 1.3 | 3.1 | 6.1 | 11.7 | 19.9 | 23.9 | 21.8 |
| 1999..... | 5.0 | * | 0.7 | 0.7 | 0.6 | 1.3 | 2.7 | 6.1 | 12.2 | 20.5 | 25.0 | 20.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | 0.5 | 0.7 | 0.4 | 0.7 | 1.5 | 3.3 | 7.2 | 13.2 | 16.6 | 14.7 |
| 2003..... | 3.8 | * | 0.6 | 0.8 | 0.5 | 0.8 | 1.9 | 3.5 | 8.0 | 12.6 | 15.6 | 12.6 |
| 2002..... | 3.8 | * | 0.6 | 0.7 | 0.4 | 0.8 | 1.7 | 4.0 | 7.6 | 12.5 | 15.8 | 14.4 |
| 2001..... | 3.8 | * | 0.6 | 0.8 | 0.5 | 0.8 | 1.7 | 3.8 | 7.5 | 12.1 | 16.4 | 14.4 |
| 2000..... | 3.9 | * | 0.6 | 0.7 | 0.5 | 0.7 | 1.7 | 4.0 | 7.4 | 13.2 | 16.9 | 14.1 |
| 1999..... | 4.0 | * | 0.7 | 0.7 | 0.4 | 0.8 | 1.9 | 4.0 | 8.2 | 13.5 | 16.6 | 13.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.7 | 0.5 | 0.4 | 1.1 | 2.4 | 5.8 | 8.9 | 12.0 | 13.8 |
| 2003..... | 1.7 | * | * | 0.8 | 0.3 | 0.6 | 1.0 | 2.5 | 5.9 | 9.4 | 15.1 | * |
| 2002..... | 1.6 | * | 0.7 | 0.7 | 0.3 | 0.5 | 0.9 | 3.3 | 5.8 | 9.7 | 13.5 | 12.2 |
| 2001..... | 1.6 | * | * | 0.5 | 0.4 | 0.5 | 1.4 | 2.5 | 5.9 | 10.3 | 14.4 | * |
| 2000..... | 1.8 | * | * | 0.9 | 0.4 | 0.7 | 1.5 | 2.8 | 6.1 | 9.8 | 14.2 | 15.9 |
| 1999..... | 1.6 | * | 0.9 | 0.7 | * | 0.5 | 1.3 | 2.9 | 5.8 | 10.0 | 13.7 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.6 | 0.6 | * | 1.3 | 2.9 | 7.1 | 10.1 | 14.7 | * |
| 2003..... | 1.8 | * | * | 0.9 | * | 0.6 | 1.2 | 3.1 | 6.3 | 11.8 | 16.1 | * |
| 2002..... | 1.8 | * | * | 0.7 | * | * | 1.0 | 4.3 | 6.6 | 12.2 | 16.0 | * |
| 2001..... | 1.7 | * | * | * | * | 0.5 | 1.4 | 3.0 | 6.8 | 14.4 | 17.9 | * |
| 2000..... | 1.9 | * | * | 1.2 | * | 0.7 | 1.5 | 2.5 | 7.7 | 12.5 | 18.7 | * |
| 1999..... | 1.7 | * | * | 0.7 | * | * | 1.7 | 2.9 | 7.7 | 10.5 | 16.2 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | 0.8 | * | * | 0.8 | 1.8 | 4.6 | 7.9 | 10.1 | 14.4 |
| 2003..... | 1.5 | * | * | 0.7 | * | * | 0.8 | 1.8 | 5.6 | 7.5 | 14.5 | * |
| 2002..... | 1.5 | * | * | 0.7 | * | * | 0.9 | 2.3 | 5.0 | 7.8 | 11.8 | * |
| 2001..... | 1.5 | * | * | 0.6 | * | * | 1.4 | 2.1 | 5.1 | 7.1 | 12.1 | * |
| 2000..... | 1.6 | * | * | 0.7 | * | * | 1.5 | 3.0 | 4.6 | 7.6 | 11.2 | * |
| 1999..... | 1.6 | * | * | 0.8 | * | * | 0.8 | 3.0 | 4.1 | 9.6 | 12.1 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of brain (C71) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.7 | * | 0.6 | 0.8 | 0.5 | 1.0 | 2.2 | 4.7 | 9.6 | 16.0 | 19.7 | 16.2 |
| 2003..... | 4.8 | * | 0.6 | 0.8 | 0.6 | 1.1 | 2.6 | 5.0 | 9.8 | 15.6 | 19.3 | 14.8 |
| 2002..... | 4.8 | * | 0.7 | 0.8 | 0.5 | 1.1 | 2.4 | 5.2 | 9.7 | 16.2 | 18.9 | 16.1 |
| 2001..... | 4.7 | * | 0.6 | 0.8 | 0.6 | 1.1 | 2.4 | 5.1 | 9.8 | 15.4 | 20.2 | 16.2 |
| 2000..... | 4.8 | * | 0.7 | 0.7 | 0.6 | 1.0 | 2.5 | 5.2 | 9.7 | 16.6 | 19.9 | 16.3 |
| 1999..... | 4.9 | * | 0.7 | 0.7 | 0.6 | 1.2 | 2.4 | 5.2 | 10.4 | 17.0 | 20.1 | 15.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.4 | * | 0.7 | 0.9 | 0.6 | 1.2 | 2.7 | 6.1 | 11.9 | 18.8 | 23.8 | 19.4 |
| 2003..... | 5.4 | * | 0.7 | 0.8 | 0.7 | 1.3 | 3.1 | 6.4 | 11.6 | 18.8 | 24.7 | 19.3 |
| 2002..... | 5.4 | * | 0.8 | 0.9 | 0.7 | 1.2 | 3.0 | 6.3 | 11.8 | 20.2 | 23.4 | 19.8 |
| 2001..... | 5.4 | * | 0.6 | 0.7 | 0.6 | 1.5 | 3.0 | 6.3 | 12.2 | 19.0 | 25.6 | 20.2 |
| 2000..... | 5.4 | * | 0.7 | 0.8 | 0.7 | 1.4 | 3.3 | 6.4 | 12.0 | 20.4 | 24.1 | 21.7 |
| 1999..... | 5.5 | * | 0.7 | 0.7 | 0.7 | 1.5 | 2.9 | 6.4 | 12.5 | 21.0 | 25.2 | 20.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | 0.6 | 0.7 | 0.4 | 0.8 | 1.7 | 3.5 | 7.4 | 13.6 | 16.9 | 14.7 |
| 2003..... | 4.2 | * | 0.6 | 0.8 | 0.5 | 0.9 | 2.1 | 3.6 | 8.2 | 12.9 | 15.6 | 12.8 |
| 2002..... | 4.1 | * | 0.6 | 0.7 | 0.4 | 0.9 | 1.8 | 4.1 | 7.8 | 12.8 | 15.9 | 14.5 |
| 2001..... | 4.1 | * | 0.6 | 0.8 | 0.5 | 0.8 | 1.8 | 4.0 | 7.6 | 12.4 | 16.6 | 14.5 |
| 2000..... | 4.1 | * | 0.7 | 0.7 | 0.5 | 0.7 | 1.7 | 4.0 | 7.6 | 13.5 | 17.1 | 14.1 |
| 1999..... | 4.3 | * | 0.7 | 0.7 | 0.5 | 0.9 | 2.0 | 4.0 | 8.5 | 13.7 | 16.8 | 13.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | * | 0.7 | 0.8 | 0.5 | 1.1 | 2.4 | 5.2 | 10.5 | 17.4 | 20.9 | 16.8 |
| 2003..... | 5.5 | * | 0.7 | 0.8 | 0.7 | 1.2 | 2.9 | 5.5 | 10.7 | 16.9 | 20.5 | 15.2 |
| 2002..... | 5.5 | * | 0.6 | 0.8 | 0.6 | 1.2 | 2.7 | 5.8 | 10.6 | 17.5 | 20.0 | 16.9 |
| 2001..... | 5.5 | * | 0.6 | 0.8 | 0.6 | 1.3 | 2.7 | 5.7 | 10.6 | 16.5 | 21.4 | 16.8 |
| 2000..... | 5.5 | * | 0.7 | 0.7 | 0.6 | 1.2 | 2.7 | 5.7 | 10.5 | 18.0 | 20.9 | 16.9 |
| 1999..... | 5.6 | * | 0.7 | 0.7 | 0.6 | 1.4 | 2.7 | 5.7 | 11.3 | 18.4 | 21.2 | 16.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.2 | * | 0.8 | 0.8 | 0.7 | 1.4 | 2.9 | 6.5 | 13.0 | 20.5 | 25.0 | 20.7 |
| 2003..... | 6.2 | * | 0.8 | 0.8 | 0.7 | 1.5 | 3.5 | 7.0 | 12.6 | 20.1 | 26.0 | 20.0 |
| 2002..... | 6.2 | * | 0.7 | 0.9 | 0.7 | 1.4 | 3.4 | 7.0 | 12.7 | 21.7 | 24.5 | 21.1 |
| 2001..... | 6.2 | * | 0.7 | 0.7 | 0.6 | 1.7 | 3.4 | 6.9 | 13.1 | 20.4 | 27.0 | 20.6 |
| 2000..... | 6.2 | * | 0.7 | 0.7 | 0.7 | 1.5 | 3.6 | 7.0 | 13.0 | 21.9 | 25.3 | 22.9 |
| 1999..... | 6.3 | * | 0.8 | 0.8 | 0.7 | 1.7 | 3.2 | 6.9 | 13.5 | 22.6 | 26.6 | 21.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.8 | * | 0.6 | 0.7 | 0.4 | 0.9 | 1.9 | 3.9 | 8.2 | 14.7 | 18.2 | 15.1 |
| 2003..... | 4.8 | * | 0.6 | 0.8 | 0.6 | 1.0 | 2.4 | 4.0 | 8.9 | 14.1 | 16.8 | 13.2 |
| 2002..... | 4.8 | * | 0.5 | 0.8 | 0.4 | 1.0 | 2.1 | 4.6 | 8.6 | 13.9 | 16.9 | 15.0 |
| 2001..... | 4.7 | * | 0.6 | 0.9 | 0.5 | 0.9 | 2.0 | 4.4 | 8.3 | 13.2 | 17.7 | 15.2 |
| 2000..... | 4.8 | * | 0.7 | 0.7 | 0.5 | 0.9 | 1.9 | 4.5 | 8.2 | 14.7 | 17.9 | 14.5 |
| 1999..... | 4.9 | * | 0.7 | 0.7 | 0.5 | 1.0 | 2.2 | 4.5 | 9.3 | 14.8 | 17.7 | 14.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of brain (C71) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | 0.7 | 0.5 | 0.7 | 1.2 | 2.7 | 4.8 | 7.8 | 9.9 | 10.6 |
| 2003..... | 2.1 | * | * | 0.7 | 0.5 | 0.6 | 1.5 | 3.1 | 5.4 | 8.5 | 10.2 | 10.4 |
| 2002..... | 2.0 | * | 1.1 | 0.8 | 0.4 | 0.6 | 1.2 | 2.7 | 5.3 | 8.1 | 11.3 | 9.4 |
| 2001..... | 2.0 | * | * | 0.7 | 0.5 | 0.5 | 1.1 | 3.0 | 5.7 | 9.3 | 9.7 | 11.9 |
| 2000..... | 2.1 | * | * | 0.8 | 0.6 | 0.7 | 1.5 | 2.8 | 5.6 | 8.1 | 11.7 | 9.9 |
| 1999..... | 2.1 | * | * | 0.8 | 0.6 | 0.6 | 1.3 | 3.1 | 5.8 | 8.0 | 11.0 | 9.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | 0.7 | * | * | 1.8 | 3.9 | 6.6 | 7.8 | 14.0 | * |
| 2003..... | 2.5 | * | * | 0.8 | 0.7 | * | 2.1 | 4.1 | 6.5 | 11.4 | 15.6 | * |
| 2002..... | 2.2 | * | * | 1.0 | * | 0.9 | 1.3 | 2.6 | 7.6 | 9.5 | 15.2 | * |
| 2001..... | 2.2 | * | * | 0.8 | * | * | 1.2 | 3.6 | 7.6 | 10.4 | 12.0 | * |
| 2000..... | 2.4 | * | * | 0.9 | 0.8 | 1.1 | 2.0 | 3.7 | 7.0 | 9.9 | 12.3 | * |
| 1999..... | 2.3 | * | * | 0.7 | 0.8 | 0.8 | 1.7 | 4.3 | 7.2 | 9.8 | 11.9 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.7 | * | * | 0.7 | 1.8 | 3.3 | 7.7 | 7.6 | 11.6 |
| 2003..... | 1.8 | * | * | 0.7 | * | * | 0.9 | 2.3 | 4.5 | 6.5 | 7.2 | 11.6 |
| 2002..... | 1.9 | * | * | * | * | * | 1.1 | 2.8 | 3.4 | 7.2 | 9.1 | 10.5 |
| 2001..... | 1.9 | * | * | * | * | * | 0.9 | 2.5 | 4.2 | 8.6 | 8.4 | 9.8 |
| 2000..... | 1.9 | * | * | 0.7 | * | * | 1.1 | 2.1 | 4.6 | 6.8 | 11.3 | 10.0 |
| 1999..... | 1.9 | * | * | 0.9 | * | * | 1.0 | 2.2 | 4.7 | 6.8 | 10.5 | 8.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other parts of central nervous system (C70,C72) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.5 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 0.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.4 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 0.7 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 0.7 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.3 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.2 | 0.5 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.4 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.6 |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 0.7 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.5 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 0.7 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.4 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.3 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of thyroid and other endocrine glands (C73-C75) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | 0.5 | 0.3 | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 2.4 | 4.0 | 6.1 |
| 2003..... | 0.7 | * | 0.4 | 0.3 | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.1 | 3.8 | 5.1 |
| 2002..... | 0.8 | * | 0.5 | 0.3 | 0.1 | 0.1 | 0.2 | 0.5 | 1.1 | 2.3 | 4.2 | 5.3 |
| 2001..... | 0.8 | * | 0.5 | 0.2 | 0.1 | 0.1 | 0.2 | 0.6 | 1.3 | 2.6 | 4.1 | 5.3 |
| 2000..... | 0.8 | * | 0.5 | 0.2 | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 2.6 | 4.1 | 6.1 |
| 1999..... | 0.8 | * | 0.6 | 0.3 | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 2.2 | 4.0 | 5.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | 0.5 | 0.3 | 0.1 | * | 0.2 | 0.4 | 1.2 | 2.7 | 3.7 | 5.6 |
| 2003..... | 0.6 | * | 0.5 | 0.3 | * | * | 0.2 | 0.6 | 1.2 | 2.2 | 3.6 | 4.3 |
| 2002..... | 0.7 | * | 0.5 | 0.3 | * | * | 0.2 | 0.6 | 1.2 | 2.2 | 4.2 | 4.2 |
| 2001..... | 0.7 | * | 0.6 | 0.2 | * | 0.1 | 0.2 | 0.6 | 1.4 | 2.9 | 4.0 | 4.5 |
| 2000..... | 0.7 | * | 0.5 | 0.2 | * | * | 0.2 | 0.5 | 1.4 | 2.8 | 4.3 | 5.0 |
| 1999..... | 0.7 | * | 0.7 | 0.3 | * | * | 0.3 | 0.6 | 1.3 | 2.5 | 4.1 | 5.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | 0.5 | 0.2 | * | * | 0.2 | 0.4 | 1.1 | 2.1 | 4.2 | 6.3 |
| 2003..... | 0.8 | * | 0.4 | 0.2 | * | * | 0.2 | 0.4 | 1.2 | 2.0 | 4.0 | 5.4 |
| 2002..... | 0.8 | * | 0.6 | 0.3 | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.3 | 4.2 | 5.8 |
| 2001..... | 0.8 | * | 0.4 | 0.2 | * | * | 0.2 | 0.6 | 1.2 | 2.4 | 4.2 | 5.6 |
| 2000..... | 0.8 | * | 0.5 | 0.2 | 0.1 | * | 0.2 | 0.4 | 1.2 | 2.4 | 4.0 | 6.5 |
| 1999..... | 0.8 | * | 0.5 | 0.3 | * | * | 0.2 | 0.5 | 1.3 | 2.0 | 3.9 | 5.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | 0.6 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 2.4 | 4.0 | 6.0 |
| 2003..... | 0.8 | * | 0.5 | 0.2 | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.2 | 3.8 | 5.2 |
| 2002..... | 0.8 | * | 0.5 | 0.3 | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.2 | 4.2 | 5.4 |
| 2001..... | 0.9 | * | 0.6 | 0.2 | * | 0.1 | 0.2 | 0.6 | 1.3 | 2.8 | 4.0 | 5.3 |
| 2000..... | 0.8 | * | 0.5 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 1.3 | 2.5 | 4.2 | 6.1 |
| 1999..... | 0.8 | * | 0.7 | 0.3 | * | 0.1 | 0.2 | 0.5 | 1.3 | 2.2 | 4.0 | 5.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | 0.6 | 0.3 | * | * | 0.2 | 0.4 | 1.2 | 2.8 | 3.8 | 5.5 |
| 2003..... | 0.7 | * | 0.4 | 0.3 | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 3.7 | 4.4 |
| 2002..... | 0.7 | * | 0.5 | 0.3 | * | * | 0.2 | 0.6 | 1.3 | 2.3 | 4.4 | 4.2 |
| 2001..... | 0.8 | * | 0.7 | 0.2 | * | * | 0.2 | 0.6 | 1.3 | 3.2 | 4.1 | 4.4 |
| 2000..... | 0.8 | * | 0.5 | 0.3 | * | * | 0.2 | 0.5 | 1.4 | 2.9 | 4.5 | 5.3 |
| 1999..... | 0.8 | * | 0.8 | 0.3 | * | * | 0.3 | 0.6 | 1.3 | 2.6 | 4.3 | 5.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | 0.6 | 0.2 | * | * | 0.2 | 0.4 | 1.1 | 2.1 | 4.2 | 6.2 |
| 2003..... | 0.9 | * | 0.5 | 0.2 | * | * | 0.2 | 0.5 | 1.2 | 2.0 | 3.9 | 5.6 |
| 2002..... | 0.9 | * | 0.5 | 0.3 | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 4.1 | 5.9 |
| 2001..... | 0.9 | * | 0.6 | 0.2 | * | * | 0.2 | 0.5 | 1.2 | 2.4 | 4.0 | 5.7 |
| 2000..... | 0.9 | * | * | 0.2 | * | * | 0.2 | 0.4 | 1.1 | 2.2 | 3.9 | 6.5 |
| 1999..... | 0.9 | * | 0.5 | 0.3 | * | * | 0.2 | 0.5 | 1.3 | 2.0 | 3.8 | 6.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of thyroid and other endocrine glands (C73-C75) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | 0.4 | * | * | * | * | 1.1 | 2.3 | 3.0 | 6.9 |
| 2003..... | 0.5 | * | * | 0.4 | * | * | * | * | 0.9 | 1.7 | 4.3 | * |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.1 | 1.9 | 4.7 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.7 | 1.4 | 1.8 | 5.1 | * |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.8 | 1.4 | 2.7 | 3.7 | * |
| 1999..... | 0.6 | * | * | 0.3 | * | * | 0.4 | * | 1.7 | 2.5 | 3.3 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 2.8 | 3.4 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 2.2 | 4.9 | * |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | * | 6.0 | * |
| 2001..... | 0.7 | * | * | * | * | * | * | 1.0 | * | 2.2 | 6.1 | * |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 2.9 | 4.5 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 1.6 | 2.3 | 3.7 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other malignant neoplasms of other and unspecified sites (C76-C80,C97) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 13.5 | * | * | 0.1 | 0.3 | 0.6 | 2.2 | 8.1 | 22.3 | 50.7 | 94.4 | 141.2 |
| 2003..... | 13.8 | * | * | 0.1 | 0.2 | 0.5 | 2.2 | 8.7 | 23.1 | 51.9 | 96.4 | 147.9 |
| 2002..... | 14.1 | * | 0.1 | 0.1 | 0.3 | 0.6 | 2.4 | 8.7 | 24.1 | 53.7 | 96.7 | 154.5 |
| 2001..... | 14.1 | * | * | 0.1 | 0.2 | 0.6 | 2.4 | 8.6 | 24.5 | 53.9 | 97.4 | 155.2 |
| 2000..... | 14.5 | * | 0.1 | 0.1 | 0.2 | 0.6 | 2.5 | 9.0 | 25.1 | 56.4 | 101.9 | 159.0 |
| 1999..... | 15.7 | * | * | 0.1 | 0.3 | 0.7 | 2.8 | 9.4 | 28.3 | 63.0 | 109.3 | 168.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 14.1 | * | * | 0.1 | 0.3 | 0.6 | 2.2 | 9.5 | 27.0 | 61.8 | 117.1 | 170.3 |
| 2003..... | 14.3 | * | * | * | 0.3 | 0.6 | 2.2 | 10.1 | 28.1 | 62.6 | 119.2 | 181.3 |
| 2002..... | 14.7 | * | * | * | 0.4 | 0.7 | 2.4 | 10.4 | 29.0 | 66.0 | 118.5 | 196.9 |
| 2001..... | 14.5 | * | * | 0.1 | 0.3 | 0.7 | 2.5 | 10.3 | 29.4 | 65.1 | 119.1 | 196.5 |
| 2000..... | 14.8 | * | * | 0.1 | 0.2 | 0.6 | 2.6 | 10.5 | 30.0 | 68.4 | 124.6 | 200.1 |
| 1999..... | 16.2 | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 11.1 | 33.9 | 76.6 | 133.5 | 218.9 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 13.0 | * | * | * | 0.2 | 0.6 | 2.1 | 6.7 | 17.9 | 41.3 | 79.1 | 128.1 |
| 2003..... | 13.4 | * | * | * | 0.2 | 0.5 | 2.2 | 7.4 | 18.5 | 43.0 | 81.2 | 133.2 |
| 2002..... | 13.5 | * | * | * | 0.2 | 0.6 | 2.4 | 7.1 | 19.6 | 43.4 | 82.3 | 136.0 |
| 2001..... | 13.6 | * | * | * | 0.2 | 0.6 | 2.3 | 7.1 | 19.9 | 44.6 | 83.1 | 137.9 |
| 2000..... | 14.1 | * | * | * | 0.2 | 0.5 | 2.4 | 7.5 | 20.6 | 46.6 | 87.0 | 142.3 |
| 1999..... | 15.3 | * | * | * | 0.2 | 0.6 | 2.7 | 7.9 | 23.1 | 51.8 | 93.6 | 148.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 0.5 | 1.2 | 4.9 | 13.5 | 32.0 | 67.2 | 93.7 |
| 2003..... | 4.5 | * | * | * | 0.3 | 0.5 | 1.6 | 5.8 | 13.7 | 36.7 | 69.9 | 105.2 |
| 2002..... | 4.4 | * | * | * | 0.3 | 0.5 | 1.3 | 5.3 | 15.6 | 35.5 | 69.2 | 109.8 |
| 2001..... | 4.3 | * | * | * | * | 0.4 | 1.7 | 5.3 | 14.2 | 36.3 | 69.2 | 121.9 |
| 2000..... | 4.4 | * | * | * | * | 0.3 | 1.4 | 5.2 | 15.4 | 39.1 | 77.8 | 132.0 |
| 1999..... | 4.5 | * | * | * | * | 0.5 | 1.6 | 5.8 | 15.7 | 40.0 | 75.6 | 127.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | 1.1 | 5.7 | 16.1 | 37.7 | 83.7 | 109.3 |
| 2003..... | 4.6 | * | * | * | * | 0.7 | 1.5 | 6.5 | 17.4 | 47.1 | 80.5 | 135.5 |
| 2002..... | 4.4 | * | * | * | * | 0.5 | 1.2 | 6.2 | 18.0 | 43.6 | 88.2 | 137.2 |
| 2001..... | 4.3 | * | * | * | * | * | 1.6 | 6.2 | 17.8 | 43.1 | 90.3 | 131.4 |
| 2000..... | 4.4 | * | * | * | * | * | 1.3 | 5.7 | 17.8 | 48.3 | 98.3 | 167.3 |
| 1999..... | 4.7 | * | * | * | * | * | 1.5 | 6.4 | 19.1 | 53.6 | 100.1 | 153.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | 1.2 | 4.1 | 11.1 | 27.4 | 55.8 | 85.7 |
| 2003..... | 4.3 | * | * | * | * | * | 1.7 | 5.0 | 10.5 | 28.4 | 62.7 | 89.8 |
| 2002..... | 4.3 | * | * | * | * | 0.6 | 1.4 | 4.5 | 13.6 | 29.0 | 56.2 | 96.2 |
| 2001..... | 4.2 | * | * | * | * | * | 1.8 | 4.5 | 10.9 | 30.9 | 55.0 | 117.3 |
| 2000..... | 4.4 | * | * | * | * | * | 1.5 | 4.7 | 13.2 | 31.9 | 64.1 | 114.7 |
| 1999..... | 4.3 | * | * | * | * | * | 1.8 | 5.1 | 12.7 | 29.3 | 59.3 | 115.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other malignant neoplasms of other and unspecified sites (C76-C80,C97) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.0 | * | * | 0.1 | 0.3 | 0.7 | 2.4 | 8.4 | 22.9 | 51.9 | 95.7 | 143.2 |
| 2003..... | 15.3 | * | * | 0.1 | 0.2 | 0.5 | 2.3 | 9.0 | 23.8 | 52.9 | 97.6 | 149.4 |
| 2002..... | 15.6 | * | * | 0.1 | 0.3 | 0.7 | 2.6 | 9.0 | 24.8 | 54.8 | 97.9 | 155.8 |
| 2001..... | 15.5 | * | * | 0.1 | 0.3 | 0.7 | 2.5 | 8.9 | 25.2 | 54.8 | 98.4 | 156.0 |
| 2000..... | 15.9 | * | * | 0.1 | 0.2 | 0.6 | 2.6 | 9.2 | 25.8 | 57.3 | 102.5 | 159.7 |
| 1999..... | 17.2 | * | * | 0.1 | 0.3 | 0.8 | 2.9 | 9.7 | 29.1 | 64.2 | 110.4 | 169.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 15.8 | * | * | * | 0.3 | 0.7 | 2.4 | 9.9 | 27.8 | 63.4 | 118.9 | 173.1 |
| 2003..... | 15.9 | * | * | * | 0.3 | 0.5 | 2.4 | 10.5 | 28.8 | 63.5 | 121.0 | 183.2 |
| 2002..... | 16.3 | * | * | * | 0.4 | 0.7 | 2.6 | 10.8 | 29.8 | 67.3 | 119.7 | 198.9 |
| 2001..... | 16.0 | * | * | * | 0.3 | 0.7 | 2.7 | 10.6 | 30.2 | 66.2 | 120.2 | 198.7 |
| 2000..... | 16.3 | * | * | * | 0.2 | 0.7 | 2.7 | 10.9 | 30.8 | 69.4 | 125.3 | 201.0 |
| 1999..... | 17.8 | * | * | 0.1 | 0.3 | 0.9 | 3.0 | 11.4 | 34.9 | 77.7 | 134.4 | 221.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 14.3 | * | * | * | 0.2 | 0.7 | 2.3 | 6.9 | 18.4 | 42.2 | 80.2 | 129.8 |
| 2003..... | 14.7 | * | * | * | 0.2 | 0.5 | 2.2 | 7.6 | 19.1 | 44.0 | 82.1 | 134.6 |
| 2002..... | 14.9 | * | * | * | 0.2 | 0.6 | 2.5 | 7.4 | 20.1 | 44.3 | 83.4 | 137.1 |
| 2001..... | 14.9 | * | * | * | 0.2 | 0.6 | 2.4 | 7.3 | 20.6 | 45.4 | 84.1 | 138.2 |
| 2000..... | 15.4 | * | * | * | 0.2 | 0.6 | 2.5 | 7.6 | 21.2 | 47.4 | 87.7 | 143.1 |
| 1999..... | 16.7 | * | * | * | 0.2 | 0.6 | 2.8 | 8.1 | 23.8 | 53.1 | 94.8 | 148.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 16.3 | * | * | * | 0.3 | 0.7 | 2.3 | 7.7 | 22.1 | 51.5 | 95.9 | 145.3 |
| 2003..... | 16.6 | * | * | * | 0.2 | 0.5 | 2.1 | 8.4 | 22.9 | 52.4 | 98.5 | 151.0 |
| 2002..... | 16.8 | * | * | * | 0.3 | 0.6 | 2.4 | 8.5 | 24.0 | 54.1 | 97.6 | 156.2 |
| 2001..... | 16.6 | * | * | 0.1 | 0.2 | 0.6 | 2.3 | 8.3 | 24.1 | 53.7 | 98.0 | 157.8 |
| 2000..... | 17.0 | * | * | * | 0.3 | 0.6 | 2.4 | 8.6 | 25.0 | 56.6 | 101.8 | 161.0 |
| 1999..... | 18.5 | * | * | 0.1 | 0.3 | 0.8 | 2.7 | 9.1 | 27.8 | 63.5 | 109.9 | 169.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 16.9 | * | * | * | 0.3 | 0.7 | 2.3 | 8.9 | 26.3 | 62.1 | 119.4 | 175.1 |
| 2003..... | 17.1 | * | * | * | 0.3 | 0.5 | 2.2 | 9.6 | 27.5 | 62.4 | 121.6 | 186.4 |
| 2002..... | 17.5 | * | * | * | 0.4 | 0.7 | 2.3 | 10.1 | 28.5 | 66.6 | 118.9 | 201.0 |
| 2001..... | 17.0 | * | * | * | 0.3 | 0.7 | 2.4 | 9.7 | 28.3 | 64.3 | 119.2 | 201.1 |
| 2000..... | 17.3 | * | * | * | 0.3 | 0.7 | 2.4 | 9.9 | 29.0 | 68.3 | 123.5 | 202.1 |
| 1999..... | 18.9 | * | * | * | 0.3 | 0.9 | 2.8 | 10.4 | 33.1 | 76.3 | 133.8 | 221.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 15.7 | * | * | * | 0.2 | 0.6 | 2.3 | 6.7 | 18.1 | 42.5 | 79.9 | 132.0 |
| 2003..... | 16.1 | * | * | * | 0.2 | 0.5 | 2.1 | 7.2 | 18.6 | 43.9 | 82.8 | 135.4 |
| 2002..... | 16.1 | * | * | * | 0.2 | 0.5 | 2.5 | 7.0 | 19.7 | 43.4 | 83.3 | 136.9 |
| 2001..... | 16.2 | * | * | * | 0.2 | 0.6 | 2.3 | 7.0 | 20.2 | 44.6 | 84.0 | 139.8 |
| 2000..... | 16.8 | * | * | * | 0.3 | 0.6 | 2.4 | 7.2 | 21.1 | 46.8 | 87.6 | 144.4 |
| 1999..... | 18.0 | * | * | * | 0.2 | 0.6 | 2.6 | 7.8 | 22.9 | 52.7 | 94.3 | 148.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other malignant neoplasms of other and unspecified sites (C76-C80,C97) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 12.3 | * | * | * | * | 0.7 | 3.4 | 13.8 | 34.4 | 66.8 | 110.4 | 137.9 |
| 2003..... | 12.5 | * | * | * | * | 0.8 | 3.5 | 14.7 | 36.3 | 67.8 | 106.9 | 149.6 |
| 2002..... | 13.0 | * | * | * | * | 0.9 | 3.8 | 13.8 | 36.7 | 71.7 | 117.5 | 169.7 |
| 2001..... | 13.0 | * | * | * | 0.3 | 1.0 | 3.7 | 14.9 | 38.2 | 74.4 | 115.4 | 154.2 |
| 2000..... | 13.1 | * | * | * | * | 0.6 | 4.4 | 15.6 | 37.5 | 72.5 | 123.1 | 155.8 |
| 1999..... | 14.4 | * | * | * | * | 0.8 | 4.4 | 16.2 | 44.7 | 80.1 | 129.0 | 185.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 13.9 | * | * | * | * | * | 3.9 | 18.7 | 46.8 | 91.3 | 139.5 | 180.9 |
| 2003..... | 13.6 | * | * | * | * | * | 3.6 | 18.6 | 47.7 | 88.8 | 141.4 | 171.5 |
| 2002..... | 13.9 | * | * | * | * | 0.8 | 4.5 | 17.0 | 48.0 | 89.0 | 153.9 | 210.0 |
| 2001..... | 14.5 | * | * | * | * | 0.9 | 4.0 | 19.7 | 51.9 | 97.6 | 151.2 | 212.5 |
| 2000..... | 14.6 | * | * | * | * | * | 5.1 | 20.4 | 52.0 | 92.4 | 167.2 | 204.3 |
| 1999..... | 15.5 | * | * | * | * | 0.9 | 4.4 | 21.2 | 58.8 | 105.4 | 160.7 | 242.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 10.9 | * | * | * | * | 0.8 | 3.0 | 9.6 | 24.3 | 49.3 | 94.0 | 120.9 |
| 2003..... | 11.6 | * | * | * | * | 0.8 | 3.4 | 11.4 | 27.1 | 52.9 | 87.3 | 141.1 |
| 2002..... | 12.3 | * | * | * | * | 1.1 | 3.2 | 11.1 | 27.6 | 59.4 | 97.0 | 154.3 |
| 2001..... | 11.7 | * | * | * | * | 1.0 | 3.4 | 10.8 | 27.1 | 58.0 | 95.4 | 132.6 |
| 2000..... | 11.8 | * | * | * | * | * | 3.7 | 11.4 | 25.9 | 58.5 | 98.3 | 137.8 |
| 1999..... | 13.3 | * | * | * | * | * | 4.4 | 11.9 | 33.3 | 62.2 | 111.2 | 164.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 18.6 | 0.6 | 0.8 | 0.9 | 1.5 | 2.1 | 3.4 | 9.0 | 25.8 | 68.2 | 139.8 | 189.4 |
| 2003..... | 19.1 | 0.8 | 0.9 | 1.0 | 1.5 | 2.2 | 3.9 | 9.3 | 26.8 | 70.3 | 143.3 | 198.0 |
| 2002..... | 19.5 | 0.7 | 0.8 | 0.9 | 1.7 | 2.3 | 3.9 | 9.9 | 28.2 | 73.9 | 143.6 | 197.5 |
| 2001..... | 19.8 | 0.7 | 1.0 | 0.9 | 1.8 | 2.5 | 4.1 | 10.4 | 29.5 | 74.6 | 145.2 | 199.9 |
| 2000..... | 20.1 | 1.0 | 0.9 | 0.9 | 1.8 | 2.3 | 4.1 | 10.7 | 30.0 | 76.3 | 149.2 | 204.6 |
| 1999..... | 20.2 | 0.8 | 0.9 | 0.9 | 1.9 | 2.4 | 4.2 | 10.8 | 31.8 | 77.8 | 148.0 | 199.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 20.4 | * | 0.9 | 1.0 | 1.9 | 2.4 | 4.2 | 11.0 | 32.1 | 87.4 | 181.3 | 254.9 |
| 2003..... | 21.0 | * | 1.0 | 1.1 | 1.8 | 2.6 | 4.7 | 11.4 | 33.9 | 90.4 | 184.8 | 268.7 |
| 2002..... | 21.4 | * | 0.8 | 1.0 | 2.0 | 2.7 | 4.6 | 11.9 | 35.0 | 95.2 | 188.0 | 269.1 |
| 2001..... | 21.5 | * | 1.0 | 1.0 | 2.1 | 2.9 | 5.1 | 12.7 | 36.3 | 94.7 | 187.0 | 275.3 |
| 2000..... | 21.6 | 1.1 | 1.0 | 1.1 | 2.3 | 2.7 | 4.8 | 13.0 | 36.1 | 96.8 | 192.6 | 281.8 |
| 1999..... | 21.7 | * | 1.0 | 1.0 | 2.2 | 2.8 | 5.1 | 13.2 | 38.8 | 97.3 | 189.0 | 279.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 16.8 | * | 0.6 | 0.7 | 1.2 | 1.7 | 2.7 | 7.0 | 19.9 | 52.0 | 111.8 | 160.0 |
| 2003..... | 17.3 | 1.1 | 0.8 | 0.8 | 1.2 | 1.7 | 3.1 | 7.2 | 20.1 | 53.4 | 115.6 | 166.8 |
| 2002..... | 17.7 | * | 0.7 | 0.8 | 1.4 | 1.9 | 3.2 | 7.9 | 21.9 | 56.2 | 114.1 | 166.5 |
| 2001..... | 18.1 | * | 1.0 | 0.7 | 1.4 | 2.1 | 3.2 | 8.1 | 23.1 | 57.9 | 117.7 | 168.4 |
| 2000..... | 18.6 | * | 0.9 | 0.7 | 1.4 | 1.9 | 3.4 | 8.5 | 24.4 | 59.5 | 120.9 | 173.1 |
| 1999..... | 18.7 | * | 0.8 | 0.7 | 1.5 | 2.0 | 3.4 | 8.5 | 25.3 | 61.9 | 121.3 | 167.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 6.9 | * | 0.8 | 1.4 | 2.0 | 2.0 | 3.2 | 7.7 | 19.8 | 48.3 | 95.4 | 103.2 |
| 2003..... | 7.1 | * | 1.3 | 1.3 | 1.9 | 2.0 | 3.4 | 8.5 | 20.4 | 47.1 | 99.1 | 129.6 |
| 2002..... | 7.3 | * | 0.9 | 1.3 | 2.1 | 1.9 | 3.9 | 7.7 | 22.1 | 53.7 | 100.8 | 132.5 |
| 2001..... | 7.4 | * | 1.3 | 1.2 | 2.2 | 2.2 | 3.9 | 8.2 | 23.9 | 52.6 | 103.3 | 134.4 |
| 2000..... | 7.5 | * | 1.0 | 1.1 | 2.0 | 1.9 | 3.9 | 8.9 | 25.0 | 58.8 | 103.3 | 143.3 |
| 1999..... | 7.5 | * | 1.0 | 1.2 | 2.0 | 2.1 | 3.9 | 10.1 | 24.2 | 58.5 | 101.0 | 131.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | 1.8 | 2.6 | 2.2 | 4.1 | 9.9 | 23.9 | 60.8 | 117.0 | 119.1 |
| 2003..... | 7.4 | * | 1.5 | 1.5 | 2.4 | 2.4 | 4.2 | 9.5 | 24.3 | 57.4 | 114.7 | 149.3 |
| 2002..... | 7.6 | * | * | 1.3 | 2.7 | 2.3 | 4.4 | 8.9 | 26.2 | 63.3 | 119.0 | 180.7 |
| 2001..... | 8.3 | * | 1.4 | 1.3 | 2.7 | 2.7 | 5.0 | 9.7 | 28.9 | 67.5 | 142.2 | 180.9 |
| 2000..... | 8.0 | * | * | 1.4 | 2.3 | 2.4 | 4.3 | 11.1 | 30.7 | 71.3 | 126.9 | 197.5 |
| 1999..... | 7.6 | * | * | 1.3 | 2.1 | 2.2 | 4.2 | 12.0 | 29.4 | 68.5 | 117.8 | 135.9 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | 1.1 | 1.2 | 1.6 | 2.2 | 5.6 | 16.1 | 38.2 | 80.5 | 95.1 |
| 2003..... | 6.8 | * | * | 1.1 | 1.3 | 1.5 | 2.6 | 7.4 | 16.8 | 39.0 | 88.5 | 119.5 |
| 2002..... | 7.0 | * | * | 1.3 | 1.5 | 1.4 | 3.3 | 6.5 | 18.5 | 46.1 | 88.3 | 108.6 |
| 2001..... | 6.5 | * | * | 1.1 | 1.5 | 1.7 | 2.6 | 6.7 | 19.5 | 40.8 | 77.1 | 111.7 |
| 2000..... | 6.9 | * | * | 0.8 | 1.5 | 1.4 | 3.4 | 6.8 | 19.8 | 48.9 | 87.5 | 116.7 |
| 1999..... | 7.4 | * | * | 1.0 | 1.9 | 2.1 | 3.4 | 8.3 | 19.6 | 50.6 | 89.9 | 130.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.5 | * | 0.8 | 0.7 | 1.4 | 2.1 | 3.5 | 9.1 | 26.2 | 69.6 | 142.1 | 193.0 |
| 2003..... | 21.0 | 0.8 | 0.8 | 0.9 | 1.4 | 2.2 | 4.0 | 9.3 | 27.2 | 71.8 | 145.4 | 200.5 |
| 2002..... | 21.3 | 0.8 | 0.7 | 0.8 | 1.6 | 2.4 | 3.9 | 10.1 | 28.6 | 75.1 | 145.3 | 199.8 |
| 2001..... | 21.6 | 0.6 | 0.9 | 0.8 | 1.7 | 2.5 | 4.1 | 10.5 | 29.8 | 75.8 | 146.8 | 202.1 |
| 2000..... | 21.8 | 1.0 | 0.9 | 0.9 | 1.8 | 2.4 | 4.1 | 10.8 | 30.3 | 77.3 | 150.9 | 206.5 |
| 1999..... | 21.9 | 0.7 | 0.9 | 0.8 | 1.8 | 2.4 | 4.3 | 10.8 | 32.3 | 78.8 | 149.5 | 201.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 22.6 | * | 0.9 | 0.8 | 1.7 | 2.4 | 4.2 | 11.1 | 32.7 | 89.3 | 184.8 | 261.1 |
| 2003..... | 23.3 | * | 0.8 | 1.0 | 1.6 | 2.7 | 4.8 | 11.6 | 34.6 | 92.6 | 188.2 | 273.6 |
| 2002..... | 23.6 | * | 0.8 | 1.0 | 1.8 | 2.8 | 4.7 | 12.1 | 35.6 | 97.0 | 190.9 | 272.8 |
| 2001..... | 23.5 | * | 1.0 | 1.0 | 1.9 | 2.9 | 5.0 | 12.9 | 36.8 | 96.3 | 188.6 | 278.8 |
| 2000..... | 23.7 | * | 1.0 | 1.0 | 2.2 | 2.7 | 4.8 | 13.1 | 36.4 | 98.1 | 195.2 | 285.0 |
| 1999..... | 23.7 | * | 0.9 | 0.9 | 2.2 | 2.9 | 5.2 | 13.3 | 39.5 | 98.7 | 191.5 | 284.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 18.5 | * | 0.6 | 0.6 | 1.2 | 1.7 | 2.8 | 7.2 | 20.1 | 52.9 | 113.3 | 162.6 |
| 2003..... | 18.9 | * | 0.7 | 0.8 | 1.1 | 1.7 | 3.2 | 7.1 | 20.4 | 54.3 | 116.8 | 168.4 |
| 2002..... | 19.2 | * | 0.7 | 0.7 | 1.4 | 2.0 | 3.1 | 8.0 | 22.2 | 56.8 | 115.1 | 168.3 |
| 2001..... | 19.7 | * | 0.9 | 0.7 | 1.4 | 2.2 | 3.3 | 8.2 | 23.3 | 58.9 | 119.3 | 170.2 |
| 2000..... | 20.1 | * | 0.9 | 0.7 | 1.3 | 2.0 | 3.4 | 8.6 | 24.7 | 60.1 | 122.1 | 174.8 |
| 1999..... | 20.1 | * | 0.8 | 0.7 | 1.4 | 2.0 | 3.4 | 8.5 | 25.6 | 62.5 | 122.3 | 168.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 22.7 | * | 0.8 | 0.7 | 1.4 | 2.0 | 3.4 | 8.8 | 26.1 | 70.4 | 145.9 | 198.8 |
| 2003..... | 23.4 | 1.0 | 0.8 | 0.9 | 1.4 | 2.1 | 3.8 | 9.0 | 27.4 | 73.2 | 149.3 | 207.2 |
| 2002..... | 23.7 | * | 0.8 | 0.8 | 1.7 | 2.3 | 3.8 | 9.7 | 28.4 | 76.7 | 149.5 | 206.5 |
| 2001..... | 23.8 | * | 1.1 | 0.8 | 1.7 | 2.4 | 4.0 | 10.3 | 29.5 | 77.2 | 150.5 | 208.2 |
| 2000..... | 24.2 | 1.3 | 1.0 | 0.9 | 1.7 | 2.3 | 4.0 | 10.5 | 30.3 | 79.0 | 155.1 | 210.7 |
| 1999..... | 24.1 | * | 1.0 | 0.8 | 1.8 | 2.4 | 4.2 | 10.6 | 32.0 | 80.4 | 153.0 | 206.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 25.1 | * | 0.9 | 0.9 | 1.6 | 2.4 | 4.1 | 10.7 | 32.8 | 90.3 | 190.1 | 270.9 |
| 2003..... | 26.0 | * | 0.9 | 1.0 | 1.6 | 2.6 | 4.7 | 11.2 | 34.9 | 95.2 | 194.8 | 284.6 |
| 2002..... | 26.2 | * | 0.9 | 1.0 | 1.9 | 2.7 | 4.5 | 11.8 | 35.4 | 99.3 | 196.8 | 283.5 |
| 2001..... | 25.9 | * | 1.0 | 0.8 | 2.0 | 2.8 | 4.7 | 12.8 | 36.4 | 97.9 | 193.6 | 288.7 |
| 2000..... | 26.2 | * | 1.1 | 1.0 | 2.1 | 2.6 | 4.8 | 12.7 | 36.3 | 100.1 | 201.0 | 293.8 |
| 1999..... | 26.1 | * | 1.0 | 0.9 | 2.2 | 2.7 | 5.0 | 13.1 | 39.2 | 100.7 | 196.1 | 291.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 20.4 | * | 0.6 | 0.6 | 1.2 | 1.5 | 2.6 | 7.0 | 19.8 | 53.4 | 115.8 | 166.5 |
| 2003..... | 20.8 | * | 0.7 | 0.8 | 1.1 | 1.6 | 3.0 | 6.8 | 20.3 | 54.3 | 118.5 | 173.1 |
| 2002..... | 21.2 | * | 0.7 | 0.7 | 1.4 | 1.9 | 3.1 | 7.6 | 21.7 | 57.5 | 117.8 | 173.2 |
| 2001..... | 21.8 | * | 1.1 | 0.7 | 1.4 | 2.0 | 3.2 | 7.9 | 23.0 | 59.5 | 122.0 | 174.6 |
| 2000..... | 22.3 | * | 0.9 | 0.8 | 1.2 | 1.9 | 3.2 | 8.4 | 24.6 | 61.2 | 125.0 | 177.2 |
| 1999..... | 22.2 | * | 0.9 | 0.6 | 1.3 | 2.0 | 3.3 | 8.3 | 25.2 | 63.4 | 124.7 | 172.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 14.0 | * | * | 0.7 | 1.7 | 3.1 | 4.6 | 12.8 | 32.9 | 76.0 | 125.2 | 156.3 |
| 2003..... | 13.9 | * | * | 0.9 | 1.5 | 3.0 | 5.4 | 13.4 | 31.8 | 74.3 | 127.4 | 152.3 |
| 2002..... | 14.1 | * | * | 0.8 | 1.5 | 3.0 | 4.8 | 13.9 | 36.4 | 77.0 | 123.5 | 153.7 |
| 2001..... | 14.5 | * | * | 1.0 | 1.8 | 3.5 | 5.8 | 13.5 | 37.1 | 79.0 | 127.6 | 157.3 |
| 2000..... | 14.3 | * | * | 0.7 | 2.3 | 3.0 | 5.4 | 15.0 | 36.5 | 76.2 | 122.3 | 177.1 |
| 1999..... | 14.5 | * | * | 0.9 | 2.4 | 3.0 | 5.6 | 14.1 | 39.5 | 77.5 | 128.2 | 162.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 14.8 | * | * | 0.7 | 2.0 | 3.4 | 5.4 | 16.0 | 40.3 | 99.5 | 152.9 | 197.1 |
| 2003..... | 14.5 | * | * | 0.9 | 1.7 | 3.5 | 6.6 | 17.1 | 41.1 | 88.6 | 150.3 | 188.6 |
| 2002..... | 15.1 | * | * | 1.0 | 1.5 | 3.8 | 6.2 | 16.5 | 45.2 | 99.4 | 155.3 | 203.5 |
| 2001..... | 15.6 | * | * | 1.3 | 2.0 | 4.0 | 7.6 | 16.3 | 46.6 | 100.7 | 161.4 | 206.7 |
| 2000..... | 15.2 | * | * | 0.8 | 2.8 | 3.7 | 5.7 | 18.2 | 45.3 | 97.8 | 157.4 | 226.6 |
| 1999..... | 15.6 | * | * | 1.2 | 2.6 | 3.9 | 7.0 | 17.5 | 48.4 | 96.9 | 162.9 | 221.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 13.2 | * | * | 0.7 | 1.4 | 2.8 | 3.9 | 10.0 | 27.0 | 59.2 | 109.6 | 140.1 |
| 2003..... | 13.3 | * | * | 0.8 | 1.2 | 2.5 | 4.4 | 10.2 | 24.3 | 64.1 | 114.4 | 138.3 |
| 2002..... | 13.1 | * | * | * | 1.5 | 2.3 | 3.5 | 11.6 | 29.4 | 61.2 | 105.4 | 134.6 |
| 2001..... | 13.5 | * | * | * | 1.7 | 3.1 | 4.2 | 11.1 | 29.5 | 63.7 | 108.7 | 139.0 |
| 2000..... | 13.5 | * | * | * | 1.8 | 2.5 | 5.2 | 12.2 | 29.4 | 61.0 | 102.6 | 158.7 |
| 1999..... | 13.6 | * | * | 0.7 | 2.1 | 2.2 | 4.3 | 11.1 | 32.3 | 63.8 | 108.7 | 140.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hodgkin's disease (C81) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.7 | 1.2 | 1.9 | 2.1 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.4 | 0.4 | 0.7 | 1.2 | 2.2 | 2.3 |
| 2002..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.4 | 0.4 | 0.6 | 1.3 | 1.9 | 2.8 |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.4 | 0.5 | 0.5 | 1.1 | 2.2 | 2.2 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.4 | 0.4 | 0.6 | 1.3 | 1.9 | 2.4 |
| 1999..... | 0.5 | * | * | * | 0.3 | 0.4 | 0.4 | 0.4 | 0.7 | 1.5 | 2.1 | 2.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.5 | 1.0 | 1.6 | 2.5 | 3.1 |
| 2003..... | 0.5 | * | * | * | 0.1 | 0.5 | 0.5 | 0.5 | 0.9 | 1.5 | 2.5 | 2.7 |
| 2002..... | 0.6 | * | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.8 | 1.8 | 2.4 | 3.7 |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.5 | 0.6 | 0.7 | 1.5 | 2.6 | 2.5 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.7 | 1.6 | 2.3 | 3.7 |
| 1999..... | 0.6 | * | * | * | 0.3 | 0.5 | 0.5 | 0.5 | 1.0 | 1.9 | 2.3 | 3.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.5 | 0.9 | 1.5 | 1.6 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.6 | 0.9 | 1.9 | 2.2 |
| 2002..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.5 | 0.8 | 1.5 | 2.4 |
| 2001..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.5 | 0.8 | 2.0 | 2.0 |
| 2000..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.6 | 1.0 | 1.7 | 2.0 |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 | 1.2 | 1.9 | 1.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.7 | 1.2 | 2.0 | 2.1 |
| 2003..... | 0.6 | * | * | * | 0.2 | 0.5 | 0.4 | 0.4 | 0.7 | 1.2 | 2.3 | 2.5 |
| 2002..... | 0.6 | * | * | * | 0.2 | 0.5 | 0.4 | 0.5 | 0.6 | 1.4 | 2.0 | 2.9 |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.4 | 0.5 | 0.6 | 1.2 | 2.3 | 2.3 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.4 | 0.4 | 0.6 | 1.3 | 2.1 | 2.6 |
| 1999..... | 0.6 | * | * | * | 0.3 | 0.4 | 0.4 | 0.5 | 0.8 | 1.5 | 2.2 | 2.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | 0.2 | 0.4 | 0.3 | 0.5 | 1.0 | 1.6 | 2.6 | 3.3 |
| 2003..... | 0.6 | * | * | * | * | 0.5 | 0.5 | 0.5 | 0.9 | 1.5 | 2.6 | 2.7 |
| 2002..... | 0.6 | * | * | * | 0.2 | 0.5 | 0.4 | 0.5 | 0.8 | 1.9 | 2.6 | 4.0 |
| 2001..... | 0.6 | * | * | * | 0.3 | 0.5 | 0.5 | 0.6 | 0.7 | 1.6 | 2.7 | 2.6 |
| 2000..... | 0.6 | * | * | * | 0.2 | 0.5 | 0.4 | 0.5 | 0.7 | 1.6 | 2.5 | 3.9 |
| 1999..... | 0.7 | * | * | * | 0.3 | 0.5 | 0.5 | 0.5 | 1.1 | 2.0 | 2.5 | 3.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.3 | 0.3 | 0.5 | 0.9 | 1.6 | 1.6 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.6 | 1.0 | 2.0 | 2.4 |
| 2002..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.5 | 0.9 | 1.6 | 2.5 |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.5 | 0.9 | 2.0 | 2.3 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.6 | 1.1 | 1.9 | 2.1 |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.5 | 1.2 | 2.0 | 1.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other lymphoid, hematopoietic and related tissue (C82-C85,C96) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.2 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.7 | 9.9 | 26.2 | 54.8 | 75.3 |
| 2003..... | 7.4 | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.8 | 10.8 | 26.9 | 56.2 | 79.0 |
| 2002..... | 7.6 | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 4.0 | 11.5 | 28.9 | 57.4 | 77.6 |
| 2001..... | 7.9 | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.4 | 12.3 | 29.4 | 58.8 | 80.1 |
| 2000..... | 8.1 | * | * | 0.1 | 0.5 | 0.7 | 1.7 | 4.7 | 13.1 | 30.6 | 61.2 | 80.7 |
| 1999..... | 8.2 | * | * | 0.1 | 0.4 | 0.7 | 1.8 | 4.8 | 13.3 | 31.8 | 62.0 | 78.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.7 | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.8 | 12.4 | 32.5 | 67.5 | 96.3 |
| 2003..... | 8.0 | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 4.9 | 13.8 | 34.0 | 69.3 | 104.0 |
| 2002..... | 8.2 | * | * | 0.2 | 0.5 | 0.9 | 2.0 | 5.0 | 14.4 | 36.2 | 71.0 | 100.2 |
| 2001..... | 8.4 | * | * | 0.1 | 0.5 | 0.9 | 2.2 | 5.8 | 15.1 | 36.7 | 71.6 | 107.1 |
| 2000..... | 8.6 | * | * | 0.1 | 0.6 | 0.9 | 2.2 | 6.0 | 16.0 | 37.5 | 74.8 | 106.6 |
| 1999..... | 8.6 | * | * | 0.1 | 0.5 | 0.9 | 2.3 | 6.2 | 16.3 | 38.4 | 75.0 | 104.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | 0.3 | 0.5 | 0.8 | 2.6 | 7.6 | 20.8 | 46.2 | 65.8 |
| 2003..... | 6.8 | * | * | 0.1 | 0.3 | 0.4 | 1.0 | 2.8 | 8.0 | 21.0 | 47.4 | 68.0 |
| 2002..... | 7.1 | * | * | * | 0.2 | 0.5 | 1.1 | 3.0 | 8.8 | 22.9 | 48.4 | 67.8 |
| 2001..... | 7.3 | * | * | * | 0.3 | 0.6 | 1.1 | 3.1 | 9.7 | 23.3 | 50.4 | 68.8 |
| 2000..... | 7.6 | * | * | * | 0.3 | 0.6 | 1.2 | 3.3 | 10.5 | 24.8 | 52.3 | 70.2 |
| 1999..... | 7.8 | * | * | * | 0.3 | 0.6 | 1.2 | 3.4 | 10.6 | 26.5 | 53.6 | 67.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 0.4 | 0.6 | 1.2 | 3.3 | 7.5 | 18.8 | 40.6 | 43.8 |
| 2003..... | 2.7 | * | * | * | 0.4 | 0.6 | 1.4 | 3.0 | 8.4 | 19.1 | 40.4 | 54.4 |
| 2002..... | 2.8 | * | * | * | 0.4 | 0.6 | 1.3 | 3.0 | 9.4 | 22.8 | 43.7 | 50.4 |
| 2001..... | 2.7 | * | * | * | 0.3 | 0.5 | 1.4 | 3.6 | 10.9 | 18.5 | 43.3 | 59.4 |
| 2000..... | 2.9 | * | * | * | 0.3 | 0.6 | 1.6 | 3.6 | 11.2 | 25.9 | 41.7 | 60.4 |
| 1999..... | 2.8 | * | * | * | * | 0.6 | 1.4 | 4.4 | 9.3 | 25.3 | 43.1 | 56.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 0.7 | 1.8 | 4.9 | 8.8 | 24.2 | 47.2 | 50.4 |
| 2003..... | 2.8 | * | * | * | * | 0.8 | 1.8 | 3.4 | 10.5 | 23.8 | 45.3 | 57.0 |
| 2002..... | 2.8 | * | * | * | 0.5 | 0.6 | 1.5 | 3.6 | 11.7 | 26.0 | 45.5 | 60.2 |
| 2001..... | 3.0 | * | * | * | * | 0.6 | 2.0 | 4.4 | 13.2 | 24.6 | 58.6 | 68.6 |
| 2000..... | 3.1 | * | * | * | * | 0.7 | 1.9 | 4.7 | 14.1 | 30.2 | 48.2 | 84.6 |
| 1999..... | 2.8 | * | * | * | * | * | 1.7 | 5.7 | 11.8 | 27.5 | 48.5 | 60.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | * | 1.6 | 6.4 | 14.5 | 36.1 | 40.3 |
| 2003..... | 2.5 | * | * | * | * | * | 0.9 | 2.7 | 6.6 | 15.3 | 37.1 | 53.1 |
| 2002..... | 2.8 | * | * | * | * | * | 1.2 | 2.3 | 7.4 | 20.2 | 42.4 | 45.6 |
| 2001..... | 2.4 | * | * | * | * | * | 0.8 | 2.8 | 8.8 | 13.7 | 32.9 | 54.9 |
| 2000..... | 2.7 | * | * | * | * | * | 1.3 | 2.6 | 8.6 | 22.6 | 37.3 | 48.5 |
| 1999..... | 2.8 | * | * | * | * | * | 1.0 | 3.1 | 7.1 | 23.6 | 39.5 | 54.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other lymphoid, hematopoietic and related tissue (C82-C85,C96) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.9 | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 3.7 | 10.1 | 26.7 | 55.5 | 76.6 |
| 2003..... | 8.1 | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.9 | 10.9 | 27.4 | 56.9 | 79.9 |
| 2002..... | 8.4 | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 4.0 | 11.6 | 29.3 | 57.9 | 78.6 |
| 2001..... | 8.6 | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.5 | 12.4 | 30.0 | 59.4 | 80.8 |
| 2000..... | 8.8 | * | * | 0.1 | 0.5 | 0.8 | 1.7 | 4.7 | 13.2 | 30.8 | 61.9 | 81.3 |
| 1999..... | 8.9 | * | * | 0.1 | 0.4 | 0.7 | 1.8 | 4.8 | 13.6 | 32.2 | 62.6 | 78.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 8.5 | * | * | 0.1 | 0.4 | 0.9 | 1.7 | 4.8 | 12.6 | 33.1 | 68.6 | 98.3 |
| 2003..... | 8.9 | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 5.0 | 14.0 | 34.7 | 70.5 | 105.9 |
| 2002..... | 9.0 | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 5.1 | 14.6 | 36.7 | 72.0 | 102.0 |
| 2001..... | 9.2 | * | * | 0.1 | 0.5 | 0.9 | 2.3 | 5.9 | 15.2 | 37.4 | 72.0 | 108.6 |
| 2000..... | 9.4 | * | * | 0.2 | 0.7 | 0.9 | 2.2 | 6.1 | 16.0 | 37.9 | 75.9 | 107.4 |
| 1999..... | 9.5 | * | * | 0.2 | 0.5 | 0.9 | 2.3 | 6.2 | 16.6 | 38.9 | 75.8 | 106.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.3 | * | * | * | 0.3 | 0.4 | 0.8 | 2.7 | 7.7 | 21.2 | 46.7 | 66.9 |
| 2003..... | 7.5 | * | * | * | 0.3 | 0.5 | 1.0 | 2.8 | 8.1 | 21.4 | 47.9 | 68.5 |
| 2002..... | 7.7 | * | * | * | 0.3 | 0.5 | 1.1 | 3.0 | 8.9 | 23.0 | 48.5 | 68.5 |
| 2001..... | 8.0 | * | * | * | 0.3 | 0.6 | 1.2 | 3.1 | 9.7 | 23.9 | 51.1 | 69.2 |
| 2000..... | 8.3 | * | * | * | 0.3 | 0.6 | 1.2 | 3.4 | 10.6 | 24.9 | 52.8 | 70.8 |
| 1999..... | 8.4 | * | * | * | 0.3 | 0.5 | 1.3 | 3.4 | 10.8 | 26.6 | 54.0 | 67.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 9.0 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.7 | 10.4 | 28.1 | 58.7 | 80.5 |
| 2003..... | 9.3 | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 3.8 | 11.4 | 28.9 | 60.1 | 84.6 |
| 2002..... | 9.5 | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 4.0 | 12.1 | 30.7 | 61.1 | 83.5 |
| 2001..... | 9.8 | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.5 | 12.7 | 31.6 | 62.5 | 84.6 |
| 2000..... | 10.1 | * | * | 0.2 | 0.4 | 0.7 | 1.7 | 4.8 | 13.9 | 32.5 | 65.4 | 85.6 |
| 1999..... | 10.2 | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 4.9 | 14.2 | 34.0 | 65.8 | 82.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 9.7 | * | * | * | 0.4 | 0.8 | 1.7 | 4.7 | 13.0 | 34.6 | 72.3 | 104.0 |
| 2003..... | 10.1 | * | * | * | 0.4 | 0.7 | 1.8 | 4.9 | 14.6 | 36.8 | 74.6 | 112.4 |
| 2002..... | 10.3 | * | * | 0.2 | 0.4 | 0.8 | 1.9 | 5.2 | 15.0 | 38.5 | 75.8 | 108.1 |
| 2001..... | 10.4 | * | * | * | 0.4 | 0.9 | 2.2 | 6.0 | 15.5 | 39.1 | 75.5 | 113.6 |
| 2000..... | 10.7 | * | * | 0.2 | 0.6 | 0.8 | 2.2 | 6.1 | 16.6 | 39.7 | 80.0 | 113.6 |
| 1999..... | 10.7 | * | * | 0.2 | 0.5 | 0.8 | 2.2 | 6.4 | 17.0 | 40.9 | 79.2 | 110.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 8.4 | * | * | * | 0.3 | 0.4 | 0.8 | 2.7 | 8.0 | 22.6 | 49.4 | 69.9 |
| 2003..... | 8.5 | * | * | * | 0.3 | 0.4 | 1.0 | 2.8 | 8.4 | 22.1 | 50.3 | 72.4 |
| 2002..... | 8.8 | * | * | * | 0.2 | 0.5 | 1.1 | 2.9 | 9.2 | 24.2 | 51.2 | 72.8 |
| 2001..... | 9.2 | * | * | * | 0.3 | 0.5 | 1.2 | 3.1 | 10.1 | 25.2 | 53.9 | 72.5 |
| 2000..... | 9.5 | * | * | * | 0.3 | 0.6 | 1.1 | 3.5 | 11.3 | 26.4 | 55.7 | 74.2 |
| 1999..... | 9.7 | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 11.5 | 28.2 | 57.0 | 71.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other lymphoid, hematopoietic and related tissue (C82-C85,C96) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.5 | 1.1 | 1.5 | 4.9 | 9.2 | 18.4 | 28.5 | 40.0 |
| 2003..... | 3.7 | * | * | * | 0.4 | 1.1 | 2.0 | 5.0 | 9.1 | 18.5 | 28.6 | 31.5 |
| 2002..... | 4.0 | * | * | * | 0.4 | 1.2 | 2.0 | 4.8 | 10.8 | 21.7 | 30.4 | 31.8 |
| 2001..... | 4.2 | * | * | * | 0.6 | 1.2 | 2.3 | 5.1 | 11.0 | 21.5 | 31.1 | 40.3 |
| 2000..... | 3.9 | * | * | * | 0.6 | 1.1 | 2.1 | 5.1 | 10.8 | 19.7 | 27.4 | 37.2 |
| 1999..... | 4.1 | * | * | * | 0.6 | 1.1 | 2.5 | 4.8 | 11.2 | 20.5 | 32.2 | 40.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | 0.6 | 1.4 | 1.9 | 6.6 | 12.0 | 23.5 | 33.4 | 43.7 |
| 2003..... | 3.9 | * | * | * | * | 1.4 | 2.4 | 6.7 | 11.5 | 20.7 | 31.2 | 35.2 |
| 2002..... | 4.5 | * | * | * | * | 1.6 | 2.9 | 5.7 | 13.9 | 28.4 | 36.0 | 43.8 |
| 2001..... | 4.7 | * | * | * | 0.8 | 1.4 | 3.4 | 6.7 | 13.8 | 28.3 | 37.8 | 56.6 |
| 2000..... | 4.4 | * | * | * | 0.7 | 1.2 | 2.6 | 7.1 | 14.9 | 25.3 | 32.1 | 45.8 |
| 1999..... | 4.7 | * | * | * | * | 1.6 | 3.3 | 6.7 | 14.9 | 26.2 | 41.3 | 59.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | 0.8 | 1.2 | 3.4 | 7.0 | 14.8 | 25.8 | 38.6 |
| 2003..... | 3.6 | * | * | * | * | 0.9 | 1.5 | 3.4 | 7.1 | 17.0 | 27.2 | 30.1 |
| 2002..... | 3.6 | * | * | * | * | 0.8 | 1.2 | 4.0 | 8.3 | 16.9 | 27.2 | 27.2 |
| 2001..... | 3.7 | * | * | * | * | 1.1 | 1.4 | 3.7 | 8.8 | 16.6 | 27.3 | 34.2 |
| 2000..... | 3.5 | * | * | * | * | 0.9 | 1.7 | 3.3 | 7.5 | 15.7 | 24.8 | 34.0 |
| 1999..... | 3.6 | * | * | * | * | * | 1.7 | 3.3 | 8.2 | 16.5 | 27.1 | 33.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Non-Hodgkin's lymphoma (C82-C85) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.7 | 9.9 | 26.1 | 54.6 | 75.0 |
| 2003..... | 7.4 | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.8 | 10.8 | 26.8 | 56.1 | 78.8 |
| 2002..... | 7.6 | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.9 | 11.5 | 28.8 | 57.2 | 77.3 |
| 2001..... | 7.8 | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.4 | 12.2 | 29.3 | 58.6 | 79.6 |
| 2000..... | 8.1 | * | * | 0.1 | 0.5 | 0.7 | 1.7 | 4.6 | 13.1 | 30.4 | 61.0 | 80.2 |
| 1999..... | 8.2 | * | * | 0.1 | 0.4 | 0.7 | 1.8 | 4.8 | 13.3 | 31.7 | 61.8 | 77.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.8 | 12.3 | 32.5 | 67.3 | 95.8 |
| 2003..... | 8.0 | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 4.8 | 13.8 | 33.9 | 69.1 | 103.6 |
| 2002..... | 8.1 | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 5.0 | 14.4 | 36.0 | 70.7 | 99.7 |
| 2001..... | 8.4 | * | * | 0.1 | 0.5 | 0.9 | 2.2 | 5.8 | 15.0 | 36.5 | 71.3 | 106.6 |
| 2000..... | 8.6 | * | * | 0.1 | 0.6 | 0.9 | 2.1 | 6.0 | 15.9 | 37.4 | 74.5 | 105.9 |
| 1999..... | 8.6 | * | * | 0.1 | 0.5 | 0.9 | 2.3 | 6.2 | 16.2 | 38.3 | 74.7 | 104.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.6 | * | * | * | 0.3 | 0.5 | 0.8 | 2.6 | 7.6 | 20.7 | 46.0 | 65.7 |
| 2003..... | 6.8 | * | * | * | 0.3 | 0.4 | 1.0 | 2.8 | 8.0 | 20.9 | 47.4 | 67.8 |
| 2002..... | 7.1 | * | * | * | 0.2 | 0.5 | 1.1 | 3.0 | 8.8 | 22.8 | 48.2 | 67.6 |
| 2001..... | 7.3 | * | * | * | 0.3 | 0.6 | 1.1 | 3.1 | 9.7 | 23.3 | 50.2 | 68.3 |
| 2000..... | 7.6 | * | * | * | 0.3 | 0.6 | 1.2 | 3.3 | 10.5 | 24.7 | 52.1 | 69.8 |
| 1999..... | 7.7 | * | * | * | 0.3 | 0.6 | 1.2 | 3.4 | 10.6 | 26.4 | 53.4 | 66.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 0.4 | 0.6 | 1.2 | 3.3 | 7.5 | 18.7 | 40.6 | 43.8 |
| 2003..... | 2.6 | * | * | * | 0.4 | 0.6 | 1.4 | 3.0 | 8.4 | 19.1 | 40.4 | 54.4 |
| 2002..... | 2.8 | * | * | * | 0.4 | 0.6 | 1.3 | 3.0 | 9.4 | 22.7 | 43.7 | 50.4 |
| 2001..... | 2.7 | * | * | * | 0.3 | 0.5 | 1.4 | 3.6 | 10.9 | 18.3 | 43.3 | 59.4 |
| 2000..... | 2.9 | * | * | * | 0.3 | 0.6 | 1.6 | 3.6 | 11.1 | 25.7 | 41.5 | 59.7 |
| 1999..... | 2.8 | * | * | * | * | 0.6 | 1.4 | 4.4 | 9.3 | 25.2 | 42.7 | 56.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 0.7 | 1.8 | 4.9 | 8.8 | 24.0 | 47.2 | 50.4 |
| 2003..... | 2.8 | * | * | * | * | 0.8 | 1.8 | 3.4 | 10.5 | 23.8 | 45.3 | 57.0 |
| 2002..... | 2.7 | * | * | * | * | 0.6 | 1.5 | 3.6 | 11.7 | 26.0 | 45.5 | 60.2 |
| 2001..... | 3.0 | * | * | * | * | 0.6 | 2.0 | 4.4 | 13.2 | 24.4 | 58.6 | 68.6 |
| 2000..... | 3.1 | * | * | * | * | 0.7 | 1.9 | 4.6 | 13.9 | 30.2 | 48.2 | 82.6 |
| 1999..... | 2.8 | * | * | * | * | * | 1.7 | 5.7 | 11.8 | 27.5 | 48.0 | 60.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | * | 1.6 | 6.4 | 14.5 | 36.1 | 40.3 |
| 2003..... | 2.5 | * | * | * | * | * | 0.9 | 2.7 | 6.6 | 15.3 | 37.1 | 53.1 |
| 2002..... | 2.8 | * | * | * | * | * | 1.2 | 2.3 | 7.4 | 20.1 | 42.4 | 45.6 |
| 2001..... | 2.4 | * | * | * | * | * | 0.8 | 2.8 | 8.8 | 13.5 | 32.9 | 54.9 |
| 2000..... | 2.7 | * | * | * | * | * | 1.3 | 2.6 | 8.6 | 22.2 | 37.0 | 48.5 |
| 1999..... | 2.8 | * | * | * | * | * | 1.0 | 3.1 | 7.1 | 23.4 | 39.2 | 54.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Non-Hodgkin's lymphoma (C82-C85) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.9 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.7 | 10.0 | 26.6 | 55.3 | 76.3 |
| 2003..... | 8.1 | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.9 | 10.9 | 27.3 | 56.8 | 79.6 |
| 2002..... | 8.3 | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 4.0 | 11.6 | 29.2 | 57.7 | 78.2 |
| 2001..... | 8.6 | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 4.5 | 12.3 | 29.9 | 59.2 | 80.3 |
| 2000..... | 8.8 | * | * | 0.1 | 0.5 | 0.8 | 1.7 | 4.7 | 13.2 | 30.7 | 61.7 | 80.9 |
| 1999..... | 8.9 | * | * | 0.1 | 0.4 | 0.7 | 1.8 | 4.8 | 13.5 | 32.1 | 62.4 | 78.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 8.5 | * | * | 0.1 | 0.4 | 0.8 | 1.6 | 4.8 | 12.6 | 33.1 | 68.4 | 97.8 |
| 2003..... | 8.8 | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 5.0 | 14.0 | 34.5 | 70.3 | 105.6 |
| 2002..... | 9.0 | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 5.1 | 14.6 | 36.5 | 71.8 | 101.4 |
| 2001..... | 9.2 | * | * | 0.1 | 0.5 | 0.9 | 2.3 | 5.9 | 15.1 | 37.2 | 71.7 | 108.1 |
| 2000..... | 9.4 | * | * | 0.2 | 0.7 | 0.9 | 2.2 | 6.1 | 15.9 | 37.7 | 75.6 | 106.7 |
| 1999..... | 9.4 | * | * | 0.2 | 0.5 | 0.9 | 2.3 | 6.2 | 16.5 | 38.8 | 75.5 | 106.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.3 | * | * | * | 0.3 | 0.4 | 0.8 | 2.7 | 7.7 | 21.2 | 46.5 | 66.7 |
| 2003..... | 7.4 | * | * | * | 0.3 | 0.5 | 1.0 | 2.8 | 8.1 | 21.3 | 47.8 | 68.2 |
| 2002..... | 7.7 | * | * | * | 0.3 | 0.5 | 1.1 | 3.0 | 8.9 | 23.0 | 48.3 | 68.2 |
| 2001..... | 8.0 | * | * | * | 0.3 | 0.6 | 1.2 | 3.1 | 9.7 | 23.8 | 50.9 | 68.7 |
| 2000..... | 8.3 | * | * | * | 0.3 | 0.6 | 1.2 | 3.4 | 10.6 | 24.8 | 52.6 | 70.4 |
| 1999..... | 8.4 | * | * | * | 0.3 | 0.5 | 1.3 | 3.4 | 10.8 | 26.5 | 53.9 | 67.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 9.0 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.7 | 10.4 | 28.1 | 58.5 | 80.2 |
| 2003..... | 9.3 | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 3.8 | 11.4 | 28.8 | 60.0 | 84.3 |
| 2002..... | 9.5 | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 4.0 | 12.0 | 30.6 | 60.9 | 83.1 |
| 2001..... | 9.7 | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.5 | 12.7 | 31.4 | 62.3 | 84.1 |
| 2000..... | 10.1 | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 4.8 | 13.8 | 32.3 | 65.1 | 85.1 |
| 1999..... | 10.1 | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 4.9 | 14.1 | 33.9 | 65.6 | 82.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 9.6 | * | * | * | 0.4 | 0.8 | 1.7 | 4.7 | 12.9 | 34.6 | 72.0 | 103.5 |
| 2003..... | 10.1 | * | * | * | 0.4 | 0.7 | 1.8 | 4.9 | 14.6 | 36.6 | 74.4 | 112.0 |
| 2002..... | 10.2 | * | * | 0.2 | 0.4 | 0.8 | 1.9 | 5.1 | 15.0 | 38.3 | 75.6 | 107.6 |
| 2001..... | 10.4 | * | * | * | 0.4 | 0.9 | 2.1 | 6.0 | 15.5 | 38.8 | 75.2 | 113.1 |
| 2000..... | 10.6 | * | * | 0.2 | 0.6 | 0.8 | 2.2 | 6.1 | 16.5 | 39.5 | 79.6 | 113.0 |
| 1999..... | 10.6 | * | * | 0.2 | 0.5 | 0.8 | 2.2 | 6.3 | 17.0 | 40.8 | 78.9 | 110.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 8.4 | * | * | * | 0.3 | 0.4 | 0.8 | 2.7 | 8.0 | 22.5 | 49.2 | 69.8 |
| 2003..... | 8.5 | * | * | * | 0.3 | 0.4 | 1.0 | 2.8 | 8.4 | 22.1 | 50.3 | 72.2 |
| 2002..... | 8.8 | * | * | * | 0.2 | 0.5 | 1.1 | 2.9 | 9.2 | 24.1 | 51.0 | 72.6 |
| 2001..... | 9.2 | * | * | * | 0.3 | 0.5 | 1.2 | 3.1 | 10.1 | 25.2 | 53.7 | 72.0 |
| 2000..... | 9.5 | * | * | * | 0.3 | 0.6 | 1.1 | 3.4 | 11.3 | 26.3 | 55.6 | 73.8 |
| 1999..... | 9.6 | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 11.4 | 28.1 | 56.9 | 70.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Non-Hodgkin's lymphoma (C82-C85) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.5 | 1.1 | 1.5 | 4.8 | 9.1 | 18.4 | 28.2 | 39.4 |
| 2003..... | 3.7 | * | * | * | 0.4 | 1.1 | 2.0 | 4.9 | 9.1 | 18.4 | 28.6 | 31.2 |
| 2002..... | 4.0 | * | * | * | 0.4 | 1.2 | 2.0 | 4.8 | 10.8 | 21.6 | 30.1 | 31.2 |
| 2001..... | 4.1 | * | * | * | 0.6 | 1.2 | 2.3 | 5.1 | 10.9 | 21.3 | 30.6 | 40.0 |
| 2000..... | 3.9 | * | * | * | 0.6 | 1.1 | 2.1 | 5.0 | 10.8 | 19.7 | 27.1 | 36.6 |
| 1999..... | 4.1 | * | * | * | 0.6 | 1.1 | 2.4 | 4.8 | 11.2 | 20.5 | 32.1 | 40.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | 0.6 | 1.4 | 1.9 | 6.6 | 12.0 | 23.5 | 33.4 | 42.7 |
| 2003..... | 3.9 | * | * | * | * | 1.4 | 2.4 | 6.7 | 11.5 | 20.7 | 31.2 | 35.2 |
| 2002..... | 4.5 | * | * | * | * | 1.6 | 2.9 | 5.7 | 13.9 | 28.3 | 35.8 | 43.8 |
| 2001..... | 4.7 | * | * | * | 0.8 | 1.3 | 3.4 | 6.7 | 13.7 | 28.2 | 37.2 | 55.4 |
| 2000..... | 4.3 | * | * | * | * | 1.2 | 2.6 | 7.0 | 14.8 | 25.3 | 31.8 | 43.4 |
| 1999..... | 4.7 | * | * | * | * | 1.6 | 3.3 | 6.6 | 14.8 | 26.2 | 41.0 | 59.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | 0.8 | 1.2 | 3.4 | 6.9 | 14.8 | 25.3 | 38.1 |
| 2003..... | 3.5 | * | * | * | * | 0.9 | 1.5 | 3.4 | 7.1 | 16.9 | 27.2 | 29.7 |
| 2002..... | 3.6 | * | * | * | * | 0.8 | 1.2 | 4.0 | 8.3 | 16.9 | 26.9 | 26.3 |
| 2001..... | 3.7 | * | * | * | * | 1.1 | 1.4 | 3.7 | 8.6 | 16.5 | 27.0 | 34.2 |
| 2000..... | 3.5 | * | * | * | * | 0.9 | 1.7 | 3.3 | 7.5 | 15.7 | 24.5 | 34.0 |
| 1999..... | 3.6 | * | * | * | * | * | 1.7 | 3.2 | 8.2 | 16.5 | 27.1 | 33.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malignant immunoproliferative diseases (C88) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 1.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 2.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.4 | 2.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.7 | 1.8 | 3.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 2.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.6 | 3.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.8 | 3.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.9 | 4.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.9 | 1.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 2.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 1.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.6 | 1.3 | 2.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 2.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 3.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.5 | 2.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.8 | 1.9 | 4.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.3 | 3.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 3.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.7 | 1.9 | 3.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.0 | 4.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 1.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 2.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Multiple myeloma and malignant plasma cell neoplasms (C90) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.3 | 1.6 | 5.8 | 15.0 | 29.7 | 31.6 |
| 2003..... | 3.7 | * | * | * | * | * | 0.3 | 1.7 | 5.8 | 16.0 | 29.9 | 34.1 |
| 2002..... | 3.8 | * | * | * | * | * | 0.3 | 1.8 | 6.1 | 17.0 | 29.8 | 34.7 |
| 2001..... | 3.8 | * | * | * | * | * | 0.3 | 1.7 | 6.4 | 16.9 | 29.8 | 34.2 |
| 2000..... | 3.8 | * | * | * | * | * | 0.3 | 1.8 | 6.2 | 16.9 | 30.3 | 35.7 |
| 1999..... | 3.8 | * | * | * | * | 0.1 | 0.3 | 1.8 | 6.9 | 16.7 | 29.6 | 33.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 0.3 | 1.8 | 6.8 | 18.3 | 36.8 | 42.1 |
| 2003..... | 3.9 | * | * | * | * | * | 0.4 | 2.1 | 7.3 | 19.1 | 36.0 | 46.8 |
| 2002..... | 4.0 | * | * | * | * | * | 0.3 | 2.2 | 7.3 | 20.4 | 37.5 | 48.0 |
| 2001..... | 4.0 | * | * | * | * | * | 0.4 | 2.1 | 7.6 | 20.2 | 37.1 | 45.1 |
| 2000..... | 3.9 | * | * | * | * | * | 0.4 | 2.1 | 6.8 | 19.8 | 37.9 | 48.3 |
| 1999..... | 3.9 | * | * | * | * | * | 0.4 | 2.1 | 8.2 | 19.4 | 36.5 | 47.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 0.3 | 1.3 | 4.9 | 12.2 | 24.9 | 26.9 |
| 2003..... | 3.6 | * | * | * | * | * | 0.2 | 1.4 | 4.5 | 13.5 | 25.9 | 28.5 |
| 2002..... | 3.6 | * | * | * | * | * | 0.3 | 1.4 | 5.0 | 14.1 | 24.7 | 29.0 |
| 2001..... | 3.6 | * | * | * | * | * | 0.3 | 1.3 | 5.4 | 14.2 | 25.0 | 29.7 |
| 2000..... | 3.7 | * | * | * | * | * | 0.3 | 1.6 | 5.6 | 14.4 | 25.4 | 30.6 |
| 1999..... | 3.7 | * | * | * | * | * | 0.3 | 1.5 | 5.7 | 14.5 | 25.2 | 28.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.2 | 5.5 | 12.2 | 23.7 | 18.1 |
| 2003..... | 1.3 | * | * | * | * | * | * | 1.4 | 5.1 | 12.5 | 23.3 | 24.4 |
| 2002..... | 1.4 | * | * | * | * | * | * | 1.8 | 4.9 | 14.1 | 26.2 | 32.2 |
| 2001..... | 1.4 | * | * | * | * | * | * | 1.3 | 5.8 | 14.7 | 24.3 | 29.4 |
| 2000..... | 1.3 | * | * | * | * | * | * | 1.4 | 5.1 | 14.9 | 24.1 | 30.5 |
| 1999..... | 1.4 | * | * | * | * | * | * | 1.7 | 6.8 | 13.2 | 22.5 | 27.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.2 | 6.3 | 13.6 | 30.0 | * |
| 2003..... | 1.2 | * | * | * | * | * | * | 1.6 | 5.6 | 13.1 | 24.9 | * |
| 2002..... | 1.5 | * | * | * | * | * | * | 2.0 | 6.5 | 17.2 | 30.8 | 51.9 |
| 2001..... | 1.5 | * | * | * | * | * | * | 1.4 | 6.0 | 20.0 | 35.3 | 38.1 |
| 2000..... | 1.3 | * | * | * | * | * | * | 1.5 | 5.2 | 18.1 | 30.0 | * |
| 1999..... | 1.3 | * | * | * | * | * | * | 1.8 | 7.8 | 14.3 | 24.0 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.2 | 4.7 | 11.1 | 19.3 | 18.7 |
| 2003..... | 1.4 | * | * | * | * | * | * | 1.2 | 4.7 | 12.1 | 22.1 | 21.9 |
| 2002..... | 1.4 | * | * | * | * | * | * | 1.5 | 3.5 | 11.6 | 23.0 | 22.4 |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.2 | 5.6 | 10.6 | 16.9 | 25.1 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | 5.1 | 12.3 | 20.1 | 29.7 |
| 1999..... | 1.5 | * | * | * | * | * | * | 1.5 | 5.9 | 12.4 | 21.5 | 30.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Multiple myeloma and malignant plasma cell neoplasms (C90) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | * | 0.3 | 1.6 | 5.9 | 15.2 | 30.0 | 32.2 |
| 2003..... | 4.1 | * | * | * | * | * | 0.3 | 1.7 | 5.9 | 16.2 | 30.2 | 34.4 |
| 2002..... | 4.2 | * | * | * | * | * | 0.3 | 1.8 | 6.2 | 17.1 | 29.9 | 34.7 |
| 2001..... | 4.1 | * | * | * | * | * | 0.3 | 1.7 | 6.5 | 17.0 | 29.9 | 34.4 |
| 2000..... | 4.1 | * | * | * | * | * | 0.3 | 1.9 | 6.2 | 16.9 | 30.5 | 35.9 |
| 1999..... | 4.1 | * | * | * | * | * | 0.4 | 1.8 | 6.9 | 16.9 | 29.8 | 34.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | * | 0.3 | 1.9 | 6.9 | 18.6 | 37.2 | 43.3 |
| 2003..... | 4.4 | * | * | * | * | * | 0.5 | 2.1 | 7.4 | 19.5 | 36.4 | 47.5 |
| 2002..... | 4.4 | * | * | * | * | * | 0.3 | 2.2 | 7.4 | 20.5 | 37.8 | 47.7 |
| 2001..... | 4.3 | * | * | * | * | * | 0.4 | 2.1 | 7.7 | 20.1 | 37.0 | 45.3 |
| 2000..... | 4.3 | * | * | * | * | * | 0.4 | 2.1 | 6.8 | 19.9 | 38.1 | 49.0 |
| 1999..... | 4.3 | * | * | * | * | * | 0.4 | 2.1 | 8.2 | 19.6 | 36.9 | 48.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 0.3 | 1.3 | 4.9 | 12.3 | 25.2 | 27.2 |
| 2003..... | 3.9 | * | * | * | * | * | 0.2 | 1.4 | 4.5 | 13.5 | 26.0 | 28.7 |
| 2002..... | 3.9 | * | * | * | * | * | 0.3 | 1.4 | 5.1 | 14.3 | 24.7 | 29.1 |
| 2001..... | 4.0 | * | * | * | * | * | 0.3 | 1.3 | 5.4 | 14.4 | 25.2 | 29.8 |
| 2000..... | 4.0 | * | * | * | * | * | 0.3 | 1.6 | 5.6 | 14.5 | 25.6 | 30.5 |
| 1999..... | 4.0 | * | * | * | * | * | 0.3 | 1.5 | 5.7 | 14.6 | 25.2 | 28.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | 0.3 | 1.4 | 5.2 | 13.8 | 28.5 | 30.8 |
| 2003..... | 4.2 | * | * | * | * | * | 0.3 | 1.4 | 5.3 | 15.0 | 28.5 | 32.8 |
| 2002..... | 4.3 | * | * | * | * | * | 0.2 | 1.5 | 5.5 | 15.9 | 28.8 | 33.6 |
| 2001..... | 4.2 | * | * | * | * | * | 0.3 | 1.5 | 5.6 | 15.7 | 28.4 | 32.9 |
| 2000..... | 4.2 | * | * | * | * | * | 0.3 | 1.6 | 5.4 | 15.7 | 28.9 | 33.0 |
| 1999..... | 4.1 | * | * | * | * | * | 0.3 | 1.5 | 5.8 | 15.6 | 28.2 | 32.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | * | 0.3 | 1.6 | 6.2 | 17.3 | 35.7 | 41.9 |
| 2003..... | 4.5 | * | * | * | * | * | 0.4 | 1.7 | 6.7 | 18.5 | 35.1 | 46.3 |
| 2002..... | 4.6 | * | * | * | * | * | 0.3 | 1.9 | 6.6 | 19.5 | 36.8 | 47.1 |
| 2001..... | 4.4 | * | * | * | * | * | 0.3 | 1.9 | 6.7 | 18.8 | 35.6 | 44.1 |
| 2000..... | 4.4 | * | * | * | * | * | 0.4 | 1.8 | 6.1 | 18.5 | 36.6 | 47.3 |
| 1999..... | 4.4 | * | * | * | * | * | 0.3 | 1.9 | 7.2 | 18.4 | 35.4 | 46.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 0.2 | 1.1 | 4.2 | 10.8 | 23.5 | 25.8 |
| 2003..... | 3.9 | * | * | * | * | * | 0.2 | 1.1 | 3.9 | 12.0 | 24.1 | 26.9 |
| 2002..... | 3.9 | * | * | * | * | * | 0.2 | 1.1 | 4.4 | 12.9 | 23.4 | 27.7 |
| 2001..... | 4.0 | * | * | * | * | * | 0.2 | 1.1 | 4.5 | 13.1 | 23.6 | 28.2 |
| 2000..... | 4.0 | * | * | * | * | * | 0.2 | 1.4 | 4.8 | 13.3 | 23.8 | 27.2 |
| 1999..... | 3.9 | * | * | * | * | * | 0.2 | 1.2 | 4.5 | 13.3 | 23.5 | 26.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Multiple myeloma and malignant plasma cell neoplasms (C90) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | 0.7 | 3.5 | 12.6 | 30.9 | 53.5 | 54.7 |
| 2003..... | 5.0 | * | * | * | * | * | 0.8 | 4.2 | 12.0 | 31.2 | 55.1 | 60.4 |
| 2002..... | 4.8 | * | * | * | * | * | 0.7 | 4.2 | 13.2 | 31.7 | 48.6 | 54.4 |
| 2001..... | 5.0 | * | * | * | * | * | 0.9 | 3.6 | 15.1 | 31.6 | 52.9 | 58.1 |
| 2000..... | 5.1 | * | * | * | * | * | 0.8 | 4.5 | 14.1 | 32.3 | 54.9 | 71.2 |
| 1999..... | 5.1 | * | * | * | * | * | 0.9 | 4.2 | 16.5 | 30.6 | 53.0 | 62.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.8 | * | * | * | * | * | 0.8 | 4.0 | 14.4 | 36.8 | 64.2 | 72.1 |
| 2003..... | 4.9 | * | * | * | * | * | 0.9 | 5.2 | 15.6 | 34.4 | 63.5 | 74.6 |
| 2002..... | 4.8 | * | * | * | * | * | 0.8 | 4.8 | 15.8 | 37.2 | 59.2 | 70.0 |
| 2001..... | 5.0 | * | * | * | * | * | 1.1 | 4.3 | 18.3 | 36.9 | 65.7 | 70.5 |
| 2000..... | 4.9 | * | * | * | * | * | * 4.9 | 4.9 | 15.3 | 39.1 | 67.3 | 77.5 |
| 1999..... | 4.9 | * | * | * | * | * | 1.1 | 4.3 | 18.5 | 35.3 | 64.1 | 80.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | * | 3.1 | 11.1 | 26.7 | 47.5 | 47.8 |
| 2003..... | 5.0 | * | * | * | * | * | * | 3.4 | 9.2 | 29.0 | 50.4 | 54.9 |
| 2002..... | 4.8 | * | * | * | * | * | * | 3.8 | 11.2 | 27.7 | 42.6 | 48.5 |
| 2001..... | 5.0 | * | * | * | * | * | 0.8 | 3.0 | 12.5 | 27.8 | 45.7 | 53.5 |
| 2000..... | 5.4 | * | * | * | * | * | 0.9 | 4.1 | 13.1 | 27.6 | 47.9 | 68.9 |
| 1999..... | 5.2 | * | * | * | * | * | * | 4.2 | 15.0 | 27.2 | 46.7 | 56.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Leukemia (C91-C95) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.3 | 0.6 | 0.8 | 0.8 | 1.0 | 1.0 | 1.6 | 3.3 | 9.2 | 25.4 | 52.4 | 78.8 |
| 2003..... | 7.4 | 0.7 | 0.8 | 0.8 | 1.0 | 1.1 | 1.8 | 3.3 | 9.3 | 25.7 | 53.9 | 80.3 |
| 2002..... | 7.5 | 0.5 | 0.7 | 0.8 | 1.2 | 1.2 | 1.7 | 3.6 | 9.8 | 26.4 | 53.1 | 80.5 |
| 2001..... | 7.5 | 0.6 | 1.0 | 0.8 | 1.2 | 1.3 | 1.7 | 3.8 | 10.0 | 26.7 | 53.3 | 81.4 |
| 2000..... | 7.6 | 1.0 | 0.9 | 0.8 | 1.2 | 1.2 | 1.8 | 3.7 | 9.9 | 27.2 | 54.5 | 83.6 |
| 1999..... | 7.5 | 0.7 | 0.8 | 0.7 | 1.2 | 1.2 | 1.8 | 3.7 | 10.6 | 27.2 | 52.8 | 82.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 8.3 | * | 0.9 | 0.9 | 1.3 | 1.2 | 1.8 | 3.8 | 11.9 | 34.4 | 73.2 | 110.7 |
| 2003..... | 8.4 | * | 0.9 | 0.9 | 1.2 | 1.4 | 2.0 | 4.0 | 11.9 | 35.1 | 75.3 | 111.8 |
| 2002..... | 8.5 | * | 0.7 | 0.8 | 1.4 | 1.4 | 1.9 | 4.2 | 12.2 | 36.3 | 74.9 | 114.5 |
| 2001..... | 8.5 | * | 1.0 | 0.9 | 1.4 | 1.5 | 1.9 | 4.2 | 12.8 | 35.7 | 74.2 | 117.5 |
| 2000..... | 8.5 | 1.1 | 0.9 | 0.9 | 1.5 | 1.4 | 1.9 | 4.3 | 12.5 | 37.2 | 75.7 | 119.8 |
| 1999..... | 8.4 | * | 0.8 | 0.8 | 1.5 | 1.4 | 1.9 | 4.3 | 13.1 | 36.8 | 73.2 | 119.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.3 | * | 0.6 | 0.7 | 0.8 | 0.9 | 1.4 | 2.8 | 6.8 | 17.8 | 38.4 | 64.5 |
| 2003..... | 6.4 | 1.1 | 0.8 | 0.7 | 0.7 | 0.8 | 1.6 | 2.7 | 7.0 | 17.8 | 39.6 | 66.4 |
| 2002..... | 6.5 | * | 0.6 | 0.7 | 1.0 | 1.0 | 1.5 | 3.1 | 7.6 | 18.1 | 38.7 | 65.7 |
| 2001..... | 6.6 | * | 1.0 | 0.7 | 1.0 | 1.1 | 1.5 | 3.3 | 7.5 | 19.2 | 39.5 | 66.3 |
| 2000..... | 6.7 | * | 0.9 | 0.6 | 0.9 | 1.0 | 1.7 | 3.2 | 7.6 | 19.0 | 40.6 | 68.8 |
| 1999..... | 6.7 | * | 0.7 | 0.7 | 1.0 | 1.1 | 1.6 | 3.2 | 8.3 | 19.3 | 39.6 | 67.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.8 | * | 0.8 | 1.3 | 1.5 | 1.1 | 1.5 | 3.0 | 6.3 | 15.7 | 28.6 | 36.2 |
| 2003..... | 2.9 | * | 1.2 | 1.1 | 1.4 | 1.1 | 1.7 | 3.6 | 6.3 | 13.9 | 32.9 | 47.2 |
| 2002..... | 2.8 | * | 0.8 | 1.1 | 1.6 | 1.1 | 2.1 | 2.5 | 7.2 | 15.5 | 27.4 | 47.1 |
| 2001..... | 3.0 | * | 1.3 | 1.1 | 1.7 | 1.5 | 1.9 | 2.9 | 6.5 | 17.3 | 33.4 | 42.5 |
| 2000..... | 2.9 | * | 0.9 | 1.1 | 1.5 | 1.1 | 1.9 | 3.5 | 7.8 | 16.9 | 32.6 | 49.1 |
| 1999..... | 2.9 | * | 0.9 | 1.1 | 1.6 | 1.4 | 2.0 | 3.4 | 7.0 | 17.3 | 31.0 | 44.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | 1.6 | 2.0 | 1.3 | 1.8 | 3.3 | 8.1 | 20.2 | 37.9 | 43.4 |
| 2003..... | 3.2 | * | 1.4 | 1.3 | 1.7 | 1.3 | 1.9 | 4.1 | 7.6 | 18.3 | 40.6 | 61.6 |
| 2002..... | 3.0 | * | * | 1.1 | 2.0 | 1.4 | 2.3 | 2.8 | 7.2 | 18.5 | 38.6 | 65.3 |
| 2001..... | 3.4 | * | 1.4 | 1.1 | 2.2 | 1.8 | 2.3 | 3.3 | 8.4 | 19.6 | 45.6 | 68.6 |
| 2000..... | 3.3 | * | * | 1.3 | 1.7 | 1.5 | 2.0 | 4.1 | 10.2 | 20.7 | 42.8 | 76.6 |
| 1999..... | 3.1 | * | * | 1.1 | 1.7 | 1.4 | 1.9 | 3.6 | 8.6 | 23.5 | 39.1 | 49.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | 1.0 | 0.9 | 0.9 | 1.3 | 2.6 | 4.6 | 12.1 | 22.2 | 32.4 |
| 2003..... | 2.6 | * | * | 1.0 | 1.0 | 0.9 | 1.5 | 3.2 | 5.2 | 10.4 | 27.7 | 39.8 |
| 2002..... | 2.6 | * | * | 1.2 | 1.1 | 0.7 | 1.9 | 2.3 | 7.3 | 13.1 | 19.7 | 38.1 |
| 2001..... | 2.6 | * | * | 1.1 | 1.1 | 1.1 | 1.4 | 2.6 | 4.8 | 15.6 | 25.1 | 29.8 |
| 2000..... | 2.6 | * | * | 0.8 | 1.3 | * | 1.8 | 2.9 | 5.6 | 13.9 | 25.8 | 35.6 |
| 1999..... | 2.8 | * | * | 1.0 | 1.5 | 1.3 | 2.1 | 3.2 | 5.6 | 12.4 | 25.7 | 41.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Leukemia (C91-C95) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 8.0 | * | 0.8 | 0.6 | 0.9 | 1.0 | 1.6 | 3.3 | 9.5 | 26.1 | 53.7 | 80.6 |
| 2003..... | 8.1 | 0.8 | 0.7 | 0.8 | 0.9 | 1.1 | 1.8 | 3.3 | 9.6 | 26.4 | 54.9 | 81.6 |
| 2002..... | 8.2 | * | 0.6 | 0.7 | 1.1 | 1.2 | 1.6 | 3.7 | 10.0 | 27.0 | 54.3 | 81.7 |
| 2001..... | 8.2 | * | 0.9 | 0.7 | 1.1 | 1.2 | 1.7 | 3.8 | 10.3 | 27.2 | 54.1 | 82.7 |
| 2000..... | 8.2 | 1.0 | 0.9 | 0.7 | 1.1 | 1.2 | 1.7 | 3.8 | 10.1 | 27.8 | 55.3 | 84.7 |
| 1999..... | 8.2 | * | 0.7 | 0.7 | 1.2 | 1.2 | 1.7 | 3.8 | 10.8 | 27.8 | 53.6 | 83.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 9.2 | * | 0.9 | 0.7 | 1.1 | 1.2 | 1.8 | 3.9 | 12.2 | 35.4 | 75.1 | 113.9 |
| 2003..... | 9.3 | * | 0.8 | 0.8 | 1.1 | 1.4 | 2.0 | 4.0 | 12.2 | 36.2 | 77.0 | 113.7 |
| 2002..... | 9.4 | * | 0.7 | 0.8 | 1.2 | 1.4 | 1.9 | 4.3 | 12.6 | 37.3 | 76.5 | 116.5 |
| 2001..... | 9.3 | * | 0.9 | 0.8 | 1.2 | 1.4 | 1.9 | 4.3 | 13.1 | 36.6 | 75.4 | 119.3 |
| 2000..... | 9.3 | * | 0.9 | 0.8 | 1.4 | 1.4 | 1.9 | 4.3 | 12.6 | 38.2 | 77.0 | 121.5 |
| 1999..... | 9.2 | * | 0.8 | 0.8 | 1.4 | 1.3 | 1.9 | 4.4 | 13.4 | 37.5 | 74.6 | 122.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.9 | * | 0.6 | 0.6 | 0.8 | 0.9 | 1.4 | 2.8 | 7.0 | 18.2 | 39.3 | 65.7 |
| 2003..... | 7.0 | * | 0.7 | 0.7 | 0.8 | 0.8 | 1.6 | 2.7 | 7.1 | 18.3 | 40.2 | 67.4 |
| 2002..... | 7.0 | * | 0.5 | 0.6 | 0.9 | 1.0 | 1.4 | 3.2 | 7.6 | 18.4 | 39.5 | 66.6 |
| 2001..... | 7.2 | * | 0.9 | 0.6 | 0.9 | 1.1 | 1.5 | 3.4 | 7.7 | 19.4 | 40.1 | 67.5 |
| 2000..... | 7.2 | * | 0.9 | 0.6 | 0.8 | 1.0 | 1.6 | 3.3 | 7.7 | 19.3 | 41.1 | 69.8 |
| 1999..... | 7.2 | * | 0.7 | 0.6 | 0.9 | 1.1 | 1.5 | 3.2 | 8.5 | 19.8 | 40.0 | 68.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 9.0 | * | 0.7 | 0.7 | 0.9 | 1.0 | 1.6 | 3.3 | 9.7 | 26.8 | 55.7 | 83.6 |
| 2003..... | 9.1 | 1.0 | 0.8 | 0.8 | 0.9 | 1.1 | 1.8 | 3.3 | 9.8 | 27.5 | 57.1 | 84.7 |
| 2002..... | 9.1 | * | 0.7 | 0.7 | 1.1 | 1.2 | 1.7 | 3.7 | 10.0 | 28.2 | 56.2 | 84.4 |
| 2001..... | 9.1 | * | 1.0 | 0.7 | 1.1 | 1.2 | 1.7 | 3.8 | 10.5 | 28.1 | 56.0 | 86.1 |
| 2000..... | 9.2 | 1.3 | 0.9 | 0.7 | 1.1 | 1.1 | 1.7 | 3.7 | 10.2 | 29.0 | 57.5 | 87.3 |
| 1999..... | 9.1 | * | 0.8 | 0.7 | 1.1 | 1.2 | 1.7 | 3.7 | 11.0 | 28.6 | 55.3 | 86.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 10.3 | * | 0.8 | 0.7 | 1.1 | 1.2 | 1.7 | 3.8 | 12.5 | 36.1 | 78.0 | 118.8 |
| 2003..... | 10.5 | * | 0.8 | 0.9 | 1.1 | 1.3 | 1.9 | 4.0 | 12.5 | 37.7 | 80.6 | 119.0 |
| 2002..... | 10.5 | * | 0.8 | 0.7 | 1.3 | 1.3 | 1.9 | 4.2 | 12.7 | 38.7 | 79.2 | 121.3 |
| 2001..... | 10.3 | * | 1.0 | 0.7 | 1.3 | 1.4 | 1.8 | 4.3 | 13.3 | 37.7 | 78.0 | 125.3 |
| 2000..... | 10.4 | * | 1.0 | 0.8 | 1.3 | 1.3 | 1.9 | 4.2 | 12.7 | 39.5 | 79.9 | 125.5 |
| 1999..... | 10.2 | * | 0.8 | 0.7 | 1.4 | 1.3 | 1.9 | 4.3 | 13.6 | 38.6 | 77.0 | 127.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.7 | * | 0.6 | 0.6 | 0.8 | 0.8 | 1.4 | 2.8 | 7.0 | 18.8 | 40.4 | 67.8 |
| 2003..... | 7.8 | * | 0.7 | 0.6 | 0.7 | 0.8 | 1.6 | 2.6 | 7.3 | 18.8 | 41.3 | 69.6 |
| 2002..... | 7.8 | * | 0.5 | 0.6 | 0.9 | 1.0 | 1.5 | 3.2 | 7.5 | 19.3 | 40.7 | 68.4 |
| 2001..... | 8.0 | * | 1.1 | 0.6 | 0.9 | 1.0 | 1.6 | 3.3 | 7.8 | 19.9 | 41.4 | 69.8 |
| 2000..... | 8.1 | * | 0.9 | 0.6 | 0.8 | 0.9 | 1.5 | 3.2 | 7.8 | 20.1 | 42.7 | 72.0 |
| 1999..... | 8.0 | * | 0.9 | 0.6 | 0.8 | 1.1 | 1.5 | 3.2 | 8.5 | 20.3 | 41.1 | 70.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Leukemia (C91-C95) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | 0.6 | 1.0 | 1.3 | 2.1 | 4.0 | 10.4 | 25.4 | 41.7 | 60.7 |
| 2003..... | 4.8 | * | * | 0.7 | 0.9 | 1.2 | 2.4 | 3.8 | 9.8 | 23.4 | 42.1 | 60.4 |
| 2002..... | 4.8 | * | * | 0.6 | 0.9 | 1.3 | 1.6 | 4.4 | 11.5 | 22.6 | 42.9 | 65.3 |
| 2001..... | 4.9 | * | * | 0.9 | 1.0 | 1.5 | 2.1 | 4.3 | 10.1 | 25.0 | 42.1 | 57.1 |
| 2000..... | 4.9 | * | * | 0.6 | 1.4 | 1.7 | 2.1 | 5.0 | 10.8 | 23.1 | 38.8 | 66.1 |
| 1999..... | 4.9 | * | * | 0.8 | 1.5 | 1.3 | 1.9 | 4.4 | 10.8 | 25.2 | 41.6 | 55.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | 0.7 | 1.1 | 1.4 | 2.3 | 4.9 | 12.8 | 37.3 | 52.8 | 81.3 |
| 2003..... | 5.2 | * | * | 0.7 | 1.2 | 1.6 | 2.8 | 4.6 | 12.6 | 31.5 | 54.0 | 78.8 |
| 2002..... | 5.3 | * | * | 0.7 | 1.0 | 1.5 | 1.8 | 5.2 | 14.1 | 32.2 | 58.0 | 87.5 |
| 2001..... | 5.4 | * | * | 1.2 | 1.1 | 1.8 | 2.6 | 4.6 | 13.4 | 34.4 | 56.1 | 75.1 |
| 2000..... | 5.5 | * | * | 0.7 | 1.9 | 2.0 | 2.0 | 5.6 | 14.0 | 32.3 | 55.8 | 98.6 |
| 1999..... | 5.4 | * | * | 1.0 | 1.6 | 1.5 | 2.1 | 5.7 | 13.7 | 33.4 | 55.7 | 78.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | 0.9 | 1.2 | 1.8 | 3.2 | 8.5 | 17.0 | 35.4 | 52.6 |
| 2003..... | 4.4 | * | * | 0.8 | 0.7 | 0.9 | 1.9 | 3.1 | 7.5 | 17.6 | 35.4 | 53.2 |
| 2002..... | 4.4 | * | * | * | 0.8 | 1.1 | 1.4 | 3.6 | 9.4 | 15.8 | 34.3 | 56.9 |
| 2001..... | 4.4 | * | * | * | 1.0 | 1.1 | 1.7 | 4.0 | 7.4 | 18.4 | 34.2 | 50.5 |
| 2000..... | 4.4 | * | * | * | 0.9 | 1.4 | 2.2 | 4.4 | 8.3 | 16.5 | 29.2 | 54.1 |
| 1999..... | 4.4 | * | * | 0.6 | 1.3 | 1.1 | 1.7 | 3.3 | 8.5 | 19.4 | 33.8 | 47.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Lymphoid leukemia (C91) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | 0.8 | 2.4 | 6.5 | 14.1 | 28.9 |
| 2003..... | 2.2 | * | 0.4 | 0.4 | 0.4 | 0.3 | 0.4 | 0.7 | 2.3 | 6.8 | 15.4 | 29.6 |
| 2002..... | 2.2 | * | 0.2 | 0.4 | 0.5 | 0.3 | 0.4 | 0.9 | 2.5 | 6.7 | 14.9 | 30.1 |
| 2001..... | 2.2 | * | 0.3 | 0.4 | 0.5 | 0.4 | 0.4 | 0.8 | 2.5 | 7.0 | 15.1 | 31.2 |
| 2000..... | 2.2 | * | 0.3 | 0.4 | 0.5 | 0.3 | 0.3 | 0.8 | 2.4 | 7.1 | 15.7 | 32.0 |
| 1999..... | 2.2 | * | 0.3 | 0.4 | 0.5 | 0.3 | 0.3 | 0.8 | 2.8 | 7.6 | 15.9 | 30.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.5 | * | 0.4 | 0.5 | 0.6 | 0.4 | 0.4 | 1.0 | 3.5 | 9.1 | 20.1 | 41.7 |
| 2003..... | 2.6 | * | 0.4 | 0.6 | 0.6 | 0.4 | 0.5 | 1.0 | 3.3 | 10.2 | 22.1 | 41.7 |
| 2002..... | 2.6 | * | 0.5 | 0.7 | 0.7 | 0.4 | 0.6 | 1.1 | 3.3 | 10.0 | 21.3 | 43.0 |
| 2001..... | 2.5 | * | 0.3 | 0.5 | 0.7 | 0.4 | 0.5 | 1.1 | 3.6 | 10.0 | 21.2 | 43.8 |
| 2000..... | 2.6 | * | 0.3 | 0.5 | 0.7 | 0.4 | 0.4 | 1.0 | 3.5 | 10.4 | 22.4 | 43.8 |
| 1999..... | 2.6 | * | 0.3 | 0.4 | 0.7 | 0.4 | 0.4 | 1.1 | 3.9 | 10.7 | 22.7 | 45.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.3 | 0.3 | 0.2 | 0.3 | 0.6 | 1.3 | 4.3 | 10.1 | 23.1 |
| 2003..... | 1.8 | * | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | 0.5 | 1.4 | 3.9 | 10.8 | 24.3 |
| 2002..... | 1.8 | * | * | 0.4 | 0.4 | 0.3 | 0.3 | 0.6 | 1.7 | 4.0 | 10.6 | 24.5 |
| 2001..... | 1.8 | * | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.6 | 1.5 | 4.4 | 11.0 | 25.9 |
| 2000..... | 1.8 | * | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.6 | 1.4 | 4.5 | 11.2 | 27.3 |
| 1999..... | 1.9 | * | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | 0.5 | 1.7 | 5.0 | 11.5 | 24.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | 0.7 | 0.8 | 0.4 | 0.5 | 0.5 | 1.4 | 3.7 | 6.3 | * |
| 2003..... | 0.9 | * | 0.7 | 0.7 | 0.9 | 0.4 | 0.6 | 0.8 | 1.6 | 2.6 | 6.6 | 15.0 |
| 2002..... | 0.8 | * | * | 0.7 | 0.8 | 0.3 | 0.7 | 0.7 | 1.6 | 2.4 | 5.7 | * |
| 2001..... | 0.9 | * | * | 0.7 | 1.0 | 0.5 | 0.5 | * | 1.5 | 4.0 | 6.0 | * |
| 2000..... | 0.9 | * | * | 0.6 | 0.9 | 0.4 | * | 1.1 | 1.2 | 3.5 | 6.5 | 13.3 |
| 1999..... | 0.9 | * | * | 0.5 | 0.9 | 0.5 | 0.5 | 0.7 | 1.8 | 2.8 | 7.5 | 15.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | 0.9 | 1.1 | 0.6 | * | * | * | 3.5 | 7.9 | * |
| 2003..... | 1.1 | * | * | 0.9 | 1.1 | 0.5 | 0.8 | * | * | 4.4 | 8.4 | * |
| 2002..... | 0.9 | * | * | 0.6 | 0.9 | * | 0.9 | * | * | * | * | * |
| 2001..... | 1.0 | * | * | 0.7 | 1.3 | 0.6 | * | * | * | 4.2 | * | * |
| 2000..... | 1.0 | * | * | 0.9 | 1.0 | 0.6 | * | * | * | 4.9 | * | * |
| 1999..... | 1.0 | * | * | 0.9 | 0.9 | 0.6 | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | 0.6 | * | * | * | * | * | 3.8 | 5.2 | * |
| 2003..... | 0.7 | * | * | * | 0.6 | * | * | * | * | * | 5.3 | * |
| 2002..... | 0.8 | * | * | 0.8 | * | * | * | * | * | * | * | * |
| 2001..... | 0.7 | * | * | 0.7 | 0.6 | * | * | * | * | 3.8 | * | * |
| 2000..... | 0.7 | * | * | * | 0.7 | * | * | * | * | * | * | * |
| 1999..... | 0.8 | * | * | * | 0.9 | * | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Lymphoid leukemia (C91) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | * | 0.2 | 0.3 | 0.4 | 0.3 | 0.3 | 0.8 | 2.4 | 6.7 | 14.5 | 29.7 |
| 2003..... | 2.4 | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | 0.7 | 2.4 | 7.0 | 15.8 | 30.2 |
| 2002..... | 2.4 | * | 0.2 | 0.4 | 0.5 | 0.3 | 0.4 | 0.9 | 2.5 | 7.0 | 15.3 | 30.8 |
| 2001..... | 2.4 | * | 0.2 | 0.3 | 0.4 | 0.3 | 0.4 | 0.8 | 2.6 | 7.1 | 15.5 | 31.9 |
| 2000..... | 2.4 | * | 0.3 | 0.4 | 0.4 | 0.2 | 0.3 | 0.8 | 2.5 | 7.4 | 16.0 | 32.6 |
| 1999..... | 2.4 | * | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.8 | 2.8 | 7.8 | 16.2 | 31.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | 0.4 | 0.3 | 0.5 | 0.4 | 0.4 | 1.0 | 3.7 | 9.5 | 20.8 | 43.1 |
| 2003..... | 2.8 | * | 0.3 | 0.5 | 0.5 | 0.4 | 0.4 | 1.0 | 3.5 | 10.5 | 22.8 | 42.6 |
| 2002..... | 2.8 | * | * | 0.4 | 0.6 | 0.4 | 0.5 | 1.1 | 3.4 | 10.4 | 22.0 | 44.3 |
| 2001..... | 2.8 | * | * | 0.4 | 0.6 | 0.4 | 0.4 | 1.1 | 3.7 | 10.4 | 21.8 | 44.8 |
| 2000..... | 2.8 | * | * | 0.5 | 0.6 | 0.3 | 0.4 | 1.0 | 3.6 | 10.7 | 23.0 | 44.8 |
| 1999..... | 2.9 | * | * | 0.4 | 0.6 | 0.4 | 0.4 | 1.1 | 4.0 | 11.1 | 23.3 | 46.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | 0.3 | 0.3 | 0.2 | 0.3 | 0.6 | 1.3 | 4.3 | 10.3 | 23.7 |
| 2003..... | 1.9 | * | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.4 | 4.1 | 11.1 | 24.7 |
| 2002..... | 2.0 | * | * | 0.3 | 0.4 | 0.3 | 0.2 | 0.6 | 1.7 | 4.1 | 10.9 | 25.0 |
| 2001..... | 2.0 | * | * | 0.3 | 0.3 | 0.3 | 0.3 | 0.6 | 1.5 | 4.5 | 11.3 | 26.6 |
| 2000..... | 2.0 | * | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.6 | 1.4 | 4.6 | 11.4 | 27.8 |
| 1999..... | 2.0 | * | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.5 | 1.7 | 5.1 | 11.7 | 25.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.8 | 2.5 | 6.9 | 15.0 | 31.2 |
| 2003..... | 2.7 | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | 0.7 | 2.5 | 7.3 | 16.6 | 31.6 |
| 2002..... | 2.7 | * | * | 0.4 | 0.5 | 0.3 | 0.4 | 0.9 | 2.5 | 7.3 | 15.8 | 32.5 |
| 2001..... | 2.7 | * | 0.2 | 0.3 | 0.5 | 0.3 | 0.4 | 0.8 | 2.6 | 7.3 | 16.0 | 33.7 |
| 2000..... | 2.7 | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.7 | 2.6 | 7.6 | 16.7 | 33.8 |
| 1999..... | 2.7 | * | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.8 | 2.8 | 8.1 | 16.8 | 32.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | 0.4 | 0.5 | 0.4 | 0.4 | 1.0 | 3.7 | 9.6 | 21.5 | 45.2 |
| 2003..... | 3.2 | * | * | 0.5 | 0.5 | 0.4 | 0.4 | 1.0 | 3.6 | 10.9 | 24.0 | 44.6 |
| 2002..... | 3.2 | * | * | 0.4 | 0.6 | 0.4 | 0.6 | 1.2 | 3.4 | 10.9 | 22.7 | 46.5 |
| 2001..... | 3.1 | * | * | 0.4 | 0.6 | 0.4 | 0.4 | 1.1 | 3.7 | 10.6 | 22.6 | 47.5 |
| 2000..... | 3.1 | * | * | 0.5 | 0.6 | 0.3 | 0.5 | 0.9 | 3.7 | 11.0 | 23.9 | 46.2 |
| 1999..... | 3.2 | * | * | 0.4 | 0.6 | 0.4 | 0.4 | 1.1 | 4.0 | 11.4 | 24.2 | 48.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | 0.3 | 0.3 | 0.2 | 0.2 | 0.6 | 1.3 | 4.6 | 10.6 | 24.9 |
| 2003..... | 2.2 | * | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.4 | 4.1 | 11.5 | 25.9 |
| 2002..... | 2.2 | * | * | 0.4 | 0.4 | 0.3 | 0.3 | 0.6 | 1.6 | 4.2 | 11.2 | 26.4 |
| 2001..... | 2.2 | * | * | 0.3 | 0.3 | 0.3 | 0.3 | 0.5 | 1.5 | 4.5 | 11.7 | 28.0 |
| 2000..... | 2.3 | * | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.5 | 1.5 | 4.8 | 11.9 | 28.9 |
| 1999..... | 2.3 | * | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.5 | 1.7 | 5.4 | 12.0 | 26.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Lymphoid leukemia (C91) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.4 | 1.0 | 3.0 | 7.4 | 13.1 | 19.9 |
| 2003..... | 1.4 | * | * | * | 0.3 | * | 0.4 | 0.9 | 2.4 | 7.4 | 11.9 | 20.2 |
| 2002..... | 1.4 | * | * | * | 0.4 | * | * | 0.8 | 3.4 | 6.4 | 14.1 | 18.5 |
| 2001..... | 1.5 | * | * | 0.4 | 0.4 | * | * | 1.3 | 3.2 | 7.7 | 13.4 | 18.4 |
| 2000..... | 1.3 | * | * | * | 0.4 | * | * | 1.1 | 2.4 | 6.7 | 11.9 | 25.4 |
| 1999..... | 1.4 | * | * | 0.3 | 0.5 | * | 0.4 | 0.9 | 3.4 | 7.5 | 12.6 | 22.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | * | 1.3 | 4.4 | 12.5 | 18.5 | 31.5 |
| 2003..... | 1.6 | * | * | * | * | * | * | 1.4 | 3.4 | 10.1 | 16.2 | 26.6 |
| 2002..... | 1.6 | * | * | * | * | * | * | 1.0 | 4.5 | 9.1 | 20.8 | 31.7 |
| 2001..... | 1.7 | * | * | * | * | * | * | 1.5 | 4.7 | 11.1 | 19.2 | 25.4 |
| 2000..... | 1.6 | * | * | * | * | * | * | 1.5 | 4.1 | 10.3 | 18.5 | 41.1 |
| 1999..... | 1.8 | * | * | * | * | * | * | 1.3 | 4.7 | 11.6 | 17.8 | 33.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | 1.9 | 3.7 | 10.0 | 15.3 |
| 2003..... | 1.1 | * | * | * | * | * | * | * | 1.5 | 5.6 | 9.5 | 17.7 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | 2.6 | 4.6 | 10.3 | 13.4 |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.1 | 2.0 | 5.3 | 10.1 | 15.8 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | * | 4.2 | 8.2 | 19.6 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 2.4 | 4.7 | 9.6 | 17.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Myeloid leukemia (C92) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | * | 0.4 | 0.2 | 0.4 | 0.6 | 0.9 | 1.9 | 5.0 | 12.9 | 23.9 | 25.6 |
| 2003..... | 3.4 | * | 0.3 | 0.3 | 0.4 | 0.6 | 1.1 | 2.0 | 5.1 | 13.0 | 23.7 | 26.1 |
| 2002..... | 3.4 | * | 0.3 | 0.2 | 0.4 | 0.6 | 1.0 | 2.0 | 5.4 | 13.4 | 23.8 | 25.7 |
| 2001..... | 3.5 | * | 0.5 | 0.2 | 0.5 | 0.7 | 1.1 | 2.3 | 5.4 | 13.6 | 23.6 | 25.1 |
| 2000..... | 3.5 | * | 0.4 | 0.2 | 0.5 | 0.7 | 1.1 | 2.2 | 5.4 | 13.4 | 23.8 | 25.3 |
| 1999..... | 3.3 | * | 0.4 | 0.2 | 0.5 | 0.7 | 1.1 | 2.2 | 5.5 | 12.9 | 21.5 | 25.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.8 | * | 0.4 | 0.3 | 0.5 | 0.5 | 1.0 | 2.1 | 5.9 | 17.3 | 32.7 | 36.1 |
| 2003..... | 3.8 | * | 0.3 | 0.2 | 0.4 | 0.7 | 1.1 | 2.2 | 6.2 | 17.1 | 32.9 | 37.9 |
| 2002..... | 3.9 | * | 0.4 | 0.2 | 0.4 | 0.7 | 1.0 | 2.2 | 6.6 | 17.8 | 33.7 | 37.6 |
| 2001..... | 3.9 | * | 0.5 | 0.2 | 0.5 | 0.8 | 1.1 | 2.4 | 6.7 | 17.5 | 32.6 | 37.2 |
| 2000..... | 3.9 | * | 0.4 | 0.2 | 0.6 | 0.8 | 1.1 | 2.5 | 6.4 | 17.8 | 33.2 | 39.1 |
| 1999..... | 3.7 | * | 0.4 | 0.2 | 0.5 | 0.7 | 1.1 | 2.4 | 6.4 | 17.0 | 29.4 | 37.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.0 | * | 0.3 | 0.2 | 0.4 | 0.6 | 0.9 | 1.8 | 4.0 | 9.2 | 18.0 | 20.8 |
| 2003..... | 3.0 | * | * | 0.3 | 0.3 | 0.5 | 1.0 | 1.7 | 4.0 | 9.6 | 17.6 | 20.9 |
| 2002..... | 3.0 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.8 | 4.3 | 9.7 | 17.3 | 20.5 |
| 2001..... | 3.1 | * | 0.5 | 0.2 | 0.5 | 0.6 | 1.0 | 2.2 | 4.3 | 10.4 | 17.6 | 20.1 |
| 2000..... | 3.1 | * | 0.3 | 0.2 | 0.4 | 0.6 | 1.2 | 1.9 | 4.5 | 9.9 | 17.7 | 19.7 |
| 1999..... | 3.0 | * | 0.3 | 0.2 | 0.5 | 0.6 | 1.0 | 2.0 | 4.6 | 9.5 | 16.3 | 20.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | 0.3 | 0.5 | 0.5 | 0.8 | 1.9 | 3.3 | 8.4 | 12.3 | 12.4 |
| 2003..... | 1.3 | * | * | 0.4 | 0.3 | 0.5 | 0.7 | 2.1 | 3.4 | 7.5 | 14.1 | 17.1 |
| 2002..... | 1.2 | * | * | * | 0.5 | 0.5 | 1.0 | 1.4 | 3.5 | 8.6 | 12.0 | 15.5 |
| 2001..... | 1.4 | * | * | * | 0.4 | 0.7 | 1.1 | 1.8 | 3.6 | 8.8 | 15.0 | 16.3 |
| 2000..... | 1.4 | * | * | * | 0.5 | 0.5 | 1.2 | 1.8 | 4.9 | 9.2 | 15.6 | 16.6 |
| 1999..... | 1.4 | * | * | 0.4 | 0.4 | 0.6 | 1.1 | 2.1 | 3.7 | 9.7 | 15.0 | 14.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 0.6 | * | 1.0 | 2.0 | 4.3 | 11.8 | 15.0 | * |
| 2003..... | 1.4 | * | * | * | * | 0.5 | 0.6 | 2.4 | 4.0 | 10.1 | 19.6 | * |
| 2002..... | 1.3 | * | * | * | 0.6 | 0.6 | 1.1 | 1.4 | 3.5 | 10.9 | 16.4 | * |
| 2001..... | 1.5 | * | * | * | 0.6 | 0.9 | 1.2 | 2.0 | 5.3 | 10.0 | 17.0 | * |
| 2000..... | 1.5 | * | * | * | * | 0.7 | 1.1 | 2.4 | 6.0 | 12.0 | 20.2 | * |
| 1999..... | 1.4 | * | * | * | 0.6 | 0.7 | 0.9 | 2.0 | 4.0 | 13.6 | 18.8 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.6 | 0.7 | 1.7 | 2.4 | 5.7 | 10.4 | * |
| 2003..... | 1.2 | * | * | * | * | * | 0.8 | 1.8 | 2.7 | 5.4 | 10.3 | * |
| 2002..... | 1.1 | * | * | * | * | * | 0.9 | 1.5 | 3.5 | 6.7 | 9.0 | * |
| 2001..... | 1.2 | * | * | * | * | * | 0.9 | 1.6 | 2.1 | 7.9 | 13.6 | * |
| 2000..... | 1.2 | * | * | * | * | * | 1.2 | * | 3.9 | 7.0 | 12.5 | * |
| 1999..... | 1.4 | * | * | * | * | * | 1.4 | 2.2 | 3.3 | 6.7 | 12.5 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Myeloid leukemia (C92) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | 0.4 | 0.2 | 0.4 | 0.6 | 0.9 | 1.9 | 5.1 | 13.2 | 24.6 | 26.2 |
| 2003..... | 3.7 | * | 0.2 | 0.2 | 0.4 | 0.6 | 1.1 | 1.9 | 5.2 | 13.4 | 24.2 | 26.4 |
| 2002..... | 3.8 | * | 0.3 | 0.2 | 0.4 | 0.7 | 1.0 | 2.1 | 5.5 | 13.7 | 24.3 | 26.0 |
| 2001..... | 3.8 | * | 0.5 | 0.2 | 0.5 | 0.7 | 1.1 | 2.3 | 5.6 | 13.9 | 23.9 | 25.4 |
| 2000..... | 3.8 | * | 0.4 | 0.2 | 0.5 | 0.7 | 1.1 | 2.3 | 5.4 | 13.7 | 24.2 | 25.5 |
| 1999..... | 3.6 | * | 0.3 | 0.2 | 0.5 | 0.7 | 1.1 | 2.2 | 5.6 | 13.1 | 21.7 | 26.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.2 | * | 0.4 | 0.2 | 0.4 | 0.6 | 1.0 | 2.1 | 6.1 | 17.7 | 33.7 | 37.2 |
| 2003..... | 4.3 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.2 | 6.3 | 17.6 | 33.6 | 38.3 |
| 2002..... | 4.3 | * | 0.4 | 0.2 | 0.4 | 0.8 | 1.0 | 2.3 | 6.8 | 18.3 | 34.4 | 37.9 |
| 2001..... | 4.2 | * | 0.4 | 0.2 | 0.4 | 0.8 | 1.1 | 2.4 | 6.8 | 18.0 | 33.3 | 37.5 |
| 2000..... | 4.2 | * | 0.4 | 0.2 | 0.6 | 0.8 | 1.1 | 2.5 | 6.4 | 18.1 | 33.7 | 39.4 |
| 1999..... | 4.0 | * | 0.3 | 0.2 | 0.5 | 0.7 | 1.1 | 2.5 | 6.6 | 17.2 | 29.8 | 38.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | 0.2 | 0.4 | 0.6 | 0.9 | 1.8 | 4.2 | 9.5 | 18.4 | 21.2 |
| 2003..... | 3.2 | * | * | 0.2 | 0.4 | 0.4 | 1.0 | 1.7 | 4.2 | 9.9 | 18.0 | 21.2 |
| 2002..... | 3.2 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.9 | 4.3 | 9.9 | 17.7 | 20.9 |
| 2001..... | 3.4 | * | 0.5 | 0.2 | 0.5 | 0.6 | 1.0 | 2.2 | 4.4 | 10.5 | 17.8 | 20.4 |
| 2000..... | 3.3 | * | 0.3 | 0.2 | 0.3 | 0.7 | 1.2 | 2.0 | 4.6 | 10.1 | 17.9 | 19.9 |
| 1999..... | 3.2 | * | 0.4 | 0.2 | 0.5 | 0.6 | 1.0 | 2.0 | 4.7 | 9.7 | 16.4 | 20.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | 0.3 | 0.2 | 0.4 | 0.6 | 0.9 | 1.9 | 5.2 | 13.8 | 25.5 | 27.4 |
| 2003..... | 4.2 | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 2.0 | 5.4 | 14.1 | 25.3 | 27.2 |
| 2002..... | 4.2 | * | 0.3 | 0.2 | 0.4 | 0.6 | 1.0 | 2.1 | 5.6 | 14.4 | 25.3 | 26.9 |
| 2001..... | 4.2 | * | 0.5 | 0.2 | 0.5 | 0.7 | 1.0 | 2.3 | 5.8 | 14.5 | 24.8 | 26.3 |
| 2000..... | 4.2 | * | 0.4 | 0.2 | 0.4 | 0.6 | 1.1 | 2.2 | 5.4 | 14.4 | 25.2 | 26.3 |
| 1999..... | 4.0 | * | 0.4 | 0.2 | 0.4 | 0.6 | 1.1 | 2.2 | 5.8 | 13.5 | 22.4 | 27.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | 0.2 | 0.4 | 0.6 | 1.0 | 2.1 | 6.3 | 18.4 | 34.9 | 39.6 |
| 2003..... | 4.8 | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 2.2 | 6.5 | 18.4 | 35.3 | 39.9 |
| 2002..... | 4.8 | * | * | 0.2 | 0.4 | 0.7 | 1.0 | 2.3 | 6.9 | 19.1 | 35.8 | 39.0 |
| 2001..... | 4.7 | * | 0.4 | 0.2 | 0.4 | 0.7 | 1.0 | 2.4 | 7.0 | 18.5 | 34.7 | 39.4 |
| 2000..... | 4.7 | * | 0.4 | 0.2 | 0.5 | 0.7 | 1.0 | 2.5 | 6.4 | 18.9 | 35.4 | 41.4 |
| 1999..... | 4.4 | * | * | 0.2 | 0.5 | 0.6 | 1.1 | 2.4 | 6.7 | 17.7 | 30.8 | 40.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.8 | 4.2 | 9.8 | 19.1 | 21.9 |
| 2003..... | 3.6 | * | * | 0.2 | 0.4 | 0.5 | 1.0 | 1.7 | 4.3 | 10.4 | 18.5 | 21.6 |
| 2002..... | 3.6 | * | * | 0.2 | 0.4 | 0.5 | 0.9 | 1.9 | 4.3 | 10.4 | 18.3 | 21.7 |
| 2001..... | 3.8 | * | 0.6 | 0.2 | 0.5 | 0.6 | 1.0 | 2.2 | 4.6 | 11.0 | 18.3 | 20.8 |
| 2000..... | 3.6 | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.0 | 4.5 | 10.6 | 18.6 | 20.2 |
| 1999..... | 3.5 | * | 0.4 | 0.2 | 0.4 | 0.7 | 1.0 | 2.0 | 4.8 | 10.0 | 16.9 | 21.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Myeloid leukemia (C92) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | 0.5 | 0.8 | 1.1 | 2.3 | 5.1 | 10.3 | 17.3 | 17.8 |
| 2003..... | 2.2 | * | * | 0.3 | 0.5 | 0.7 | 1.4 | 2.2 | 4.9 | 10.4 | 16.9 | 21.4 |
| 2002..... | 2.2 | * | * | * | 0.4 | 0.8 | 0.9 | 2.2 | 5.7 | 10.3 | 16.6 | 19.7 |
| 2001..... | 2.2 | * | * | * | 0.4 | 0.8 | 1.4 | 2.2 | 4.5 | 11.5 | 16.8 | 17.5 |
| 2000..... | 2.4 | * | * | * | 0.8 | 1.1 | 1.5 | 2.9 | 6.1 | 10.0 | 14.6 | 19.7 |
| 1999..... | 2.2 | * | * | 0.3 | 0.8 | 0.7 | 1.2 | 2.4 | 5.0 | 11.1 | 15.8 | 14.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 0.8 | 1.2 | 2.7 | 5.7 | 13.7 | 21.1 | 20.3 |
| 2003..... | 2.4 | * | * | * | * | 0.8 | 1.6 | 2.2 | 5.9 | 14.3 | 21.9 | 32.0 |
| 2002..... | 2.4 | * | * | * | * | 0.9 | 1.1 | 2.6 | 7.0 | 14.2 | 21.4 | 32.8 |
| 2001..... | 2.4 | * | * | * | * | 1.0 | 1.6 | 2.2 | 5.9 | 15.9 | 20.7 | * |
| 2000..... | 2.5 | * | * | * | 1.0 | 1.4 | 1.4 | 3.2 | 7.2 | 13.2 | 18.5 | 23.5 |
| 1999..... | 2.4 | * | * | * | 0.8 | 0.9 | 1.2 | 3.1 | 6.5 | 14.4 | 20.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | 0.8 | 1.0 | 2.0 | 4.7 | 7.8 | 15.1 | 16.9 |
| 2003..... | 2.0 | * | * | * | * | * | 1.2 | 2.2 | 4.0 | 7.6 | 14.0 | 17.3 |
| 2002..... | 2.0 | * | * | * | * | 0.7 | 0.8 | 1.9 | 4.7 | 7.6 | 13.9 | 14.6 |
| 2001..... | 2.0 | * | * | * | * | * | 1.2 | 2.2 | 3.3 | 8.4 | 14.7 | 16.7 |
| 2000..... | 2.2 | * | * | * | * | 0.9 | 1.5 | 2.7 | 5.3 | 7.7 | 12.3 | 18.3 |
| 1999..... | 2.0 | * | * | * | 0.8 | * | 1.1 | 1.8 | 3.8 | 8.7 | 13.1 | 12.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Monocytic leukemia (C93) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | 0.1 | 0.5 | 0.5 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.5 | 0.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 0.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified leukemia (C94-C95) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | 1.9 | 5.8 | 13.9 | 23.8 |
| 2003..... | 1.8 | * | 0.2 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.9 | 5.7 | 14.4 | 24.0 |
| 2002..... | 1.8 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.9 | 6.1 | 14.0 | 24.1 |
| 2001..... | 1.8 | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.7 | 2.0 | 5.9 | 14.2 | 24.4 |
| 2000..... | 1.9 | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 2.0 | 6.5 | 14.6 | 25.6 |
| 1999..... | 1.9 | * | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 2.3 | 6.5 | 14.9 | 25.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | 0.2 | 0.2 | 0.2 | 0.4 | 0.7 | 2.4 | 7.8 | 19.7 | 31.7 |
| 2003..... | 1.9 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.8 | 2.3 | 7.6 | 19.6 | 31.1 |
| 2002..... | 2.0 | * | * | 0.2 | 0.3 | 0.3 | 0.3 | 0.8 | 2.3 | 8.2 | 19.3 | 32.6 |
| 2001..... | 2.0 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.3 | 0.8 | 2.4 | 8.0 | 19.8 | 35.3 |
| 2000..... | 2.0 | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 2.5 | 8.8 | 19.5 | 35.7 |
| 1999..... | 2.1 | * | * | 0.2 | 0.3 | 0.2 | 0.4 | 0.8 | 2.7 | 8.8 | 20.3 | 34.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.4 | 1.4 | 4.2 | 10.0 | 20.3 |
| 2003..... | 1.6 | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.5 | 4.2 | 10.9 | 20.8 |
| 2002..... | 1.6 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | 1.6 | 4.4 | 10.5 | 20.4 |
| 2001..... | 1.6 | * | * | 0.1 | 0.2 | 0.2 | 0.2 | 0.5 | 1.7 | 4.3 | 10.5 | 19.8 |
| 2000..... | 1.7 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.6 | 4.5 | 11.4 | 21.4 |
| 1999..... | 1.8 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | 1.9 | 4.6 | 11.5 | 21.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.5 | 1.5 | 3.5 | 10.1 | 13.8 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.4 | 1.3 | 3.7 | 11.9 | 15.0 |
| 2002..... | 0.7 | * | * | * | 0.4 | 0.3 | * | * | 2.1 | 4.3 | 9.3 | 21.1 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.7 | 1.4 | 4.5 | 12.3 | 16.9 |
| 2000..... | 0.7 | * | * | * | * | * | 0.4 | 0.6 | 1.6 | 4.0 | 10.3 | 19.2 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 1.6 | 4.6 | 8.1 | 14.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 2.1 | 4.8 | 15.0 | * |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | 3.7 | 11.9 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | * | 2.1 | 4.6 | 14.0 | * |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 5.2 | 20.1 | * |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 13.8 | * |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 6.2 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 6.7 | * |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | 3.7 | 11.9 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 2.0 | 4.1 | 6.2 | 19.9 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 3.9 | 7.0 | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 4.3 | 7.9 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 3.4 | 6.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified leukemia (C94-C95) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.6 | 1.9 | 6.0 | 14.1 | 24.2 |
| 2003..... | 2.0 | * | 0.2 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.9 | 5.9 | 14.5 | 24.4 |
| 2002..... | 2.0 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.9 | 6.2 | 14.3 | 24.2 |
| 2001..... | 2.0 | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 2.1 | 6.0 | 14.2 | 24.6 |
| 2000..... | 2.0 | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 2.1 | 6.6 | 14.8 | 25.8 |
| 1999..... | 2.1 | * | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | 2.3 | 6.6 | 15.2 | 25.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.7 | 2.4 | 8.0 | 19.9 | 32.4 |
| 2003..... | 2.2 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.8 | 2.3 | 7.9 | 20.0 | 31.7 |
| 2002..... | 2.2 | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.8 | 2.3 | 8.4 | 19.6 | 33.0 |
| 2001..... | 2.2 | * | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.8 | 2.5 | 8.1 | 19.6 | 35.8 |
| 2000..... | 2.2 | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.7 | 2.5 | 9.1 | 19.7 | 36.0 |
| 1999..... | 2.3 | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.8 | 2.7 | 8.9 | 20.7 | 35.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.1 | * | 0.3 | 0.4 | 1.5 | 4.3 | 10.2 | 20.5 |
| 2003..... | 1.8 | * | * | 0.2 | * | 0.1 | 0.3 | 0.5 | 1.5 | 4.2 | 10.9 | 21.1 |
| 2002..... | 1.8 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.6 | 4.3 | 10.7 | 20.4 |
| 2001..... | 1.7 | * | * | * | 0.2 | 0.2 | 0.2 | 0.5 | 1.7 | 4.3 | 10.7 | 20.0 |
| 2000..... | 1.9 | * | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.7 | 4.5 | 11.6 | 21.7 |
| 1999..... | 1.9 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | 2.0 | 4.7 | 11.6 | 21.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | 0.1 | 0.2 | 0.1 | 0.3 | 0.6 | 1.9 | 6.0 | 14.6 | 24.5 |
| 2003..... | 2.2 | * | 0.2 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.9 | 6.0 | 14.9 | 25.2 |
| 2002..... | 2.2 | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.9 | 6.4 | 14.7 | 24.4 |
| 2001..... | 2.2 | * | 0.3 | 0.1 | 0.2 | 0.2 | 0.2 | 0.7 | 2.0 | 6.1 | 14.7 | 25.3 |
| 2000..... | 2.3 | * | 0.3 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 2.1 | 6.8 | 15.2 | 26.5 |
| 1999..... | 2.3 | * | * | 0.1 | 0.3 | 0.2 | 0.3 | 0.7 | 2.4 | 6.7 | 15.5 | 26.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | 0.2 | 0.2 | 0.4 | 0.7 | 2.4 | 7.9 | 20.8 | 32.9 |
| 2003..... | 2.4 | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 2.3 | 8.1 | 20.6 | 33.3 |
| 2002..... | 2.4 | * | * | * | 0.3 | 0.2 | 0.3 | 0.7 | 2.3 | 8.5 | 20.2 | 34.5 |
| 2001..... | 2.4 | * | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.8 | 2.5 | 8.3 | 20.1 | 36.9 |
| 2000..... | 2.4 | * | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 2.5 | 9.4 | 20.1 | 36.6 |
| 1999..... | 2.5 | * | * | * | 0.3 | 0.2 | 0.4 | 0.7 | 2.8 | 9.2 | 21.2 | 36.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.3 | 0.4 | 1.4 | 4.3 | 10.4 | 20.7 |
| 2003..... | 2.0 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.2 | 10.9 | 21.6 |
| 2002..... | 2.0 | * | * | * | 0.2 | 0.2 | 0.3 | 0.6 | 1.5 | 4.6 | 10.9 | 20.0 |
| 2001..... | 1.9 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.3 | 11.1 | 20.5 |
| 2000..... | 2.1 | * | * | * | 0.3 | 0.2 | 0.3 | 0.7 | 1.7 | 4.6 | 11.9 | 22.4 |
| 1999..... | 2.1 | * | * | * | 0.2 | 0.2 | 0.3 | 0.6 | 2.0 | 4.7 | 11.8 | 22.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified leukemia (C94-C95) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.5 | 0.7 | 2.3 | 7.8 | 11.2 | 22.7 |
| 2003..... | 1.2 | * | * | * | * | * | 0.5 | 0.8 | 2.5 | 5.3 | 13.2 | 18.7 |
| 2002..... | 1.3 | * | * | * | * | * | * | 1.3 | 2.3 | 5.8 | 12.0 | 26.3 |
| 2001..... | 1.2 | * | * | * | * | * | 0.4 | 0.8 | 2.4 | 5.8 | 11.7 | 20.9 |
| 2000..... | 1.2 | * | * | * | * | * | 0.4 | 0.9 | 2.2 | 6.2 | 12.2 | 20.4 |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.1 | 2.4 | 6.3 | 12.8 | 19.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.8 | 11.1 | 12.8 | 28.5 |
| 2003..... | 1.2 | * | * | * | * | * | * | 1.0 | 3.2 | 6.8 | 15.9 | * |
| 2002..... | 1.3 | * | * | * | * | * | * | 1.5 | 2.6 | 8.9 | 15.5 | 21.9 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | 2.7 | 7.4 | 15.9 | 30.0 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | 2.7 | 8.8 | 18.5 | 32.9 |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.2 | 2.5 | 7.2 | 16.6 | 25.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.0 | 5.4 | 10.3 | 20.5 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | 1.9 | 4.3 | 11.7 | 18.2 |
| 2002..... | 1.3 | * | * | * | * | * | * | 1.1 | 2.1 | 3.5 | 9.9 | 28.0 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | 2.0 | 4.7 | 9.3 | 17.5 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | 1.8 | 4.4 | 8.7 | 15.7 |
| 1999..... | 1.3 | * | * | * | * | * | * | 1.0 | 2.2 | 5.7 | 10.7 | 17.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Benign neoplasms (D10-D36) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.3 | 3.7 | 7.9 |
| 2003..... | 0.6 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 3.7 | 7.4 |
| 2002..... | 0.6 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.6 | 3.6 | 7.8 |
| 2001..... | 0.6 | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.7 | 1.5 | 3.6 | 8.7 |
| 2000..... | 0.6 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 4.2 | 8.9 |
| 1999..... | 0.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.6 | 4.1 | 8.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.4 | 7.5 |
| 2003..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.7 | 7.5 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.6 | 3.3 | 6.2 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 4.0 | 9.1 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.5 | 3.8 | 8.8 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 3.7 | 9.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 3.9 | 8.1 |
| 2003..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 3.7 | 7.3 |
| 2002..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.6 | 3.9 | 8.6 |
| 2001..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.5 | 3.4 | 8.5 |
| 2000..... | 0.7 | * | * | * | * | * | 0.3 | 0.3 | 0.7 | 1.5 | 4.4 | 8.9 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 4.3 | 8.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 3.5 | 7.8 |
| 2003..... | 0.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.2 | 3.7 | 7.3 |
| 2002..... | 0.6 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.5 | 3.5 | 7.4 |
| 2001..... | 0.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 3.5 | 8.2 |
| 2000..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.1 | 8.8 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 3.9 | 8.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.1 | 7.4 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.6 | 7.5 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.6 | 3.2 | 5.6 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 3.8 | 8.6 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.4 | 1.3 | 3.8 | 8.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 3.5 | 9.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 3.8 | 8.0 |
| 2003..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 3.8 | 7.2 |
| 2002..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.5 | 3.7 | 8.2 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.3 | 8.0 |
| 2000..... | 0.8 | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.5 | 4.3 | 8.8 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 4.2 | 8.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Benign neoplasms (D10-D36) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.1 | 2.7 | 5.9 | 10.4 |
| 2003..... | 0.6 | * | * | * | * | * | 0.5 | 0.6 | 0.9 | 2.9 | 4.2 | 8.0 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.7 | 1.1 | 2.7 | 5.5 | 11.5 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.6 | 1.8 | 2.5 | 5.6 | 15.3 |
| 2000..... | 0.6 | * | * | * | * | * | 0.5 | 0.6 | 0.8 | 2.7 | 5.7 | 10.2 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.7 | 1.8 | 2.8 | 6.6 | 8.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 7.4 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 3.6 | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 2.1 | * | 6.3 | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 3.0 | * | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 2.1 | * | 7.5 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 3.3 | 5.0 | 10.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | 2.5 | 3.6 | 8.7 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.9 | * | 2.8 | 5.5 | 11.3 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 1.5 | 2.3 | 5.2 | 14.5 |
| 2000..... | 0.8 | * | * | * | * | * | 0.7 | * | * | 2.5 | 6.4 | 10.0 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 3.2 | 6.1 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of eye, brain and other parts of central nervous system (D31-D33) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.5 | 5.6 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.5 | 5.5 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.1 | 2.5 | 5.1 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.3 | 6.0 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.9 | 6.2 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.1 | 2.8 | 5.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 2.2 | 5.2 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.4 | 5.4 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.0 | 3.2 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.2 | 5.6 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.9 | 2.5 | 5.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.4 | 5.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.7 | 5.8 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.5 | 5.5 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.8 | 6.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.4 | 6.1 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 3.1 | 6.4 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 3.0 | 5.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.4 | 5.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 5.4 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.0 | 2.3 | 5.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 5.6 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 6.3 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 5.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | * | 0.6 | 2.0 | 5.3 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 5.5 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 1.8 | 2.9 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.9 | 2.1 | 5.1 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | * | 0.8 | 2.5 | 6.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.2 | 5.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.6 | 5.9 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.6 | 5.3 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.7 | 5.9 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.3 | 5.8 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.5 | 1.0 | 3.0 | 6.4 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.5 | 1.0 | 2.9 | 5.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of eye, brain and other parts of central nervous system (D31-D33) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.7 | 2.1 | 4.0 | 6.0 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | 2.3 | 2.7 | 6.2 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 1.6 | 3.9 | 6.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 1.2 | 1.8 | 3.7 | 10.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 1.3 | 4.0 | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 1.1 | 1.9 | 4.5 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | 2.5 | 3.5 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 2.1 | * | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 4.1 | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | 3.9 | 10.7 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | 4.9 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 2.7 | 4.2 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Of other and unspecified sites (D10-D24,D29-D30,D34-D36) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.4 | 1.1 | 2.2 |
| 2003..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 1.1 | 1.8 |
| 2002..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.5 | 1.2 | 2.6 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.3 | 2.5 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 2.5 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.2 | 2.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.4 | 1.2 | 2.3 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.3 | 2.1 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 3.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 1.8 | 3.5 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.3 | 2.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 2.2 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 1.0 | 1.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 2.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.1 | 2.0 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 1.8 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 2.3 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.2 | 2.4 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 2.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.2 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 1.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 2.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.4 | 1.7 | 3.5 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.3 | 2.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 4.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.1 | 2.0 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 1.1 | 1.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.3 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.2 | 2.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Neoplasms of uncertain or unknown behavior of specified sites (D37-D47) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | 0.8 | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 3.5 | 10.8 | 30.8 | 65.6 |
| 2003..... | 3.9 | 0.9 | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 3.6 | 11.0 | 31.6 | 69.6 |
| 2002..... | 3.8 | 0.9 | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.3 | 3.6 | 10.8 | 31.1 | 67.0 |
| 2001..... | 4.0 | 0.5 | 0.3 | 0.2 | 0.2 | 0.2 | 0.6 | 1.4 | 4.0 | 12.0 | 32.2 | 70.1 |
| 2000..... | 4.0 | 0.7 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 4.2 | 11.8 | 32.0 | 72.0 |
| 1999..... | 4.0 | 0.8 | 0.4 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 3.8 | 12.5 | 31.4 | 70.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 4.5 | 13.9 | 40.8 | 87.9 |
| 2003..... | 4.1 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.7 | 1.4 | 4.3 | 14.1 | 41.3 | 91.6 |
| 2002..... | 4.0 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 4.5 | 13.7 | 41.3 | 91.9 |
| 2001..... | 4.2 | * | 0.3 | 0.2 | 0.1 | 0.3 | 0.7 | 1.6 | 4.6 | 15.3 | 43.1 | 96.6 |
| 2000..... | 4.2 | * | 0.4 | 0.2 | 0.2 | 0.3 | 0.7 | 1.6 | 4.9 | 15.1 | 42.4 | 97.6 |
| 1999..... | 4.2 | * | 0.4 | 0.2 | 0.2 | 0.4 | 0.7 | 1.6 | 4.8 | 16.2 | 41.4 | 97.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | 0.2 | 0.2 | 0.2 | 0.4 | 1.0 | 2.7 | 8.2 | 24.0 | 55.5 |
| 2003..... | 3.8 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 3.0 | 8.4 | 25.0 | 59.9 |
| 2002..... | 3.7 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.1 | 2.8 | 8.3 | 24.3 | 56.2 |
| 2001..... | 3.9 | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.6 | 1.2 | 3.4 | 9.3 | 25.0 | 59.0 |
| 2000..... | 3.9 | * | * | 0.2 | 0.1 | 0.3 | 0.5 | 1.2 | 3.5 | 9.2 | 25.2 | 61.5 |
| 1999..... | 3.8 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 2.9 | 9.5 | 24.9 | 60.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 7.2 | 20.0 | 41.4 |
| 2003..... | 1.2 | * | * | * | * | 0.3 | * | 0.7 | 2.3 | 7.2 | 25.0 | 38.9 |
| 2002..... | 1.1 | * | * | 0.3 | * | * | 0.4 | 1.0 | 2.2 | 7.4 | 21.2 | 40.5 |
| 2001..... | 1.2 | * | * | * | * | * | 0.6 | 1.1 | 2.5 | 7.2 | 20.4 | 47.5 |
| 2000..... | 1.1 | * | * | * | * | * | 0.5 | 0.8 | 2.7 | 7.2 | 19.2 | 59.1 |
| 1999..... | 1.2 | * | * | * | * | 0.3 | 0.4 | 1.0 | 2.6 | 9.0 | 20.2 | 46.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 1.2 | 3.0 | 7.8 | 23.2 | 51.8 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | 2.8 | 8.7 | 31.1 | 49.3 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | 2.9 | 9.0 | 26.7 | 50.2 |
| 2001..... | 1.2 | * | * | * | * | * | * | 1.4 | 2.9 | 8.6 | 27.3 | 62.9 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | 2.7 | 7.8 | 21.6 | 82.6 |
| 1999..... | 1.1 | * | * | * | * | * | * | 1.5 | 2.7 | 9.0 | 22.9 | 49.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | * | 2.3 | 6.8 | 17.8 | 36.0 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | * | 6.0 | 20.8 | 33.6 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | * | 6.1 | 17.4 | 35.6 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | 2.2 | 6.1 | 15.7 | 40.0 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 2.8 | 6.6 | 17.5 | 47.5 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | 2.5 | 8.9 | 18.4 | 44.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Neoplasms of uncertain or unknown behavior of specified sites (D37-D47) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | 0.7 | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 3.6 | 11.1 | 31.3 | 66.7 |
| 2003..... | 4.4 | 0.9 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 1.3 | 3.7 | 11.2 | 31.9 | 70.8 |
| 2002..... | 4.2 | 0.8 | 0.3 | 0.1 | 0.2 | 0.3 | 0.5 | 1.3 | 3.7 | 11.0 | 31.5 | 68.0 |
| 2001..... | 4.4 | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.6 | 1.4 | 4.1 | 12.3 | 32.6 | 70.8 |
| 2000..... | 4.4 | 0.8 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 1.5 | 4.3 | 12.1 | 32.5 | 72.2 |
| 1999..... | 4.4 | 0.9 | 0.4 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 3.9 | 12.7 | 31.8 | 71.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.6 | * | 0.4 | 0.2 | 0.2 | 0.4 | 0.6 | 1.4 | 4.6 | 14.3 | 41.7 | 89.7 |
| 2003..... | 4.5 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.7 | 1.5 | 4.4 | 14.4 | 41.8 | 93.5 |
| 2002..... | 4.5 | * | * | 0.2 | 0.2 | 0.3 | 0.6 | 1.5 | 4.6 | 14.0 | 41.9 | 93.6 |
| 2001..... | 4.6 | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.6 | 4.7 | 15.7 | 43.7 | 97.9 |
| 2000..... | 4.6 | * | 0.4 | 0.2 | 0.2 | 0.3 | 0.7 | 1.7 | 5.1 | 15.5 | 43.2 | 97.8 |
| 1999..... | 4.6 | * | 0.4 | 0.2 | 0.2 | 0.4 | 0.7 | 1.7 | 4.9 | 16.6 | 42.1 | 99.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | 0.2 | 0.2 | 0.2 | 0.5 | 1.0 | 2.7 | 8.3 | 24.3 | 56.3 |
| 2003..... | 4.2 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 3.1 | 8.6 | 25.2 | 60.9 |
| 2002..... | 4.0 | * | 0.3 | 0.1 | 0.2 | 0.3 | 0.5 | 1.1 | 2.9 | 8.4 | 24.6 | 57.0 |
| 2001..... | 4.3 | * | * | 0.2 | 0.2 | 0.2 | 0.6 | 1.2 | 3.5 | 9.5 | 25.3 | 59.6 |
| 2000..... | 4.3 | * | * | 0.2 | 0.1 | 0.3 | 0.5 | 1.2 | 3.5 | 9.4 | 25.5 | 61.9 |
| 1999..... | 4.1 | * | 0.3 | 0.2 | 0.2 | 0.2 | 0.5 | 1.3 | 2.9 | 9.5 | 25.1 | 60.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.8 | * | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 3.6 | 11.5 | 32.4 | 69.4 |
| 2003..... | 5.0 | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 1.3 | 3.7 | 11.6 | 33.1 | 73.4 |
| 2002..... | 4.8 | 0.9 | 0.3 | 0.1 | 0.2 | 0.3 | 0.5 | 1.3 | 3.8 | 11.4 | 32.5 | 70.3 |
| 2001..... | 5.0 | * | 0.3 | 0.2 | 0.1 | 0.3 | 0.6 | 1.5 | 4.2 | 12.8 | 33.7 | 72.5 |
| 2000..... | 5.0 | * | 0.4 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 4.4 | 12.4 | 33.7 | 74.5 |
| 1999..... | 4.9 | * | 0.4 | 0.2 | 0.2 | 0.3 | 0.6 | 1.4 | 3.9 | 13.1 | 32.8 | 74.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 5.2 | * | * | 0.2 | 0.2 | 0.4 | 0.6 | 1.4 | 4.5 | 14.8 | 43.3 | 94.2 |
| 2003..... | 5.2 | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.5 | 4.3 | 15.0 | 43.5 | 97.7 |
| 2002..... | 5.1 | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 4.6 | 14.6 | 43.1 | 97.2 |
| 2001..... | 5.3 | * | * | 0.2 | * | 0.3 | 0.7 | 1.7 | 4.8 | 16.4 | 45.0 | 100.7 |
| 2000..... | 5.2 | * | 0.4 | 0.2 | 0.2 | 0.3 | 0.7 | 1.7 | 5.1 | 15.7 | 44.8 | 101.4 |
| 1999..... | 5.2 | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.7 | 5.0 | 17.1 | 43.6 | 104.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | 0.2 | 0.2 | 0.2 | 0.4 | 1.0 | 2.7 | 8.6 | 25.0 | 58.3 |
| 2003..... | 4.8 | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.2 | 8.8 | 26.0 | 62.8 |
| 2002..... | 4.6 | * | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 3.0 | 8.7 | 25.4 | 58.7 |
| 2001..... | 4.8 | * | * | 0.2 | 0.2 | 0.2 | 0.6 | 1.2 | 3.6 | 9.7 | 26.2 | 60.8 |
| 2000..... | 4.8 | * | * | 0.2 | * | 0.3 | 0.5 | 1.2 | 3.6 | 9.6 | 26.4 | 63.7 |
| 1999..... | 4.7 | * | * | 0.2 | 0.2 | 0.3 | 0.5 | 1.2 | 2.9 | 9.7 | 25.7 | 61.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Neoplasms of uncertain or unknown behavior of specified sites (D37-D47) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 0.4 | 0.7 | 1.5 | 4.7 | 9.7 | 24.6 | 42.9 |
| 2003..... | 2.3 | * | * | * | * | * | 0.7 | 1.8 | 4.1 | 9.8 | 24.4 | 47.6 |
| 2002..... | 2.3 | * | * | * | * | 0.4 | 0.7 | 1.5 | 4.4 | 9.5 | 25.2 | 50.5 |
| 2001..... | 2.4 | * | * | 0.4 | * | * | 0.8 | 1.3 | 3.9 | 10.6 | 25.4 | 54.9 |
| 2000..... | 2.4 | * | * | * | * | 0.4 | 0.5 | 2.0 | 4.3 | 11.6 | 23.6 | 51.8 |
| 1999..... | 2.5 | * | * | * | * | * | 0.8 | 1.7 | 4.2 | 12.4 | 25.1 | 49.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 0.8 | 2.0 | 6.5 | 13.0 | 30.5 | 48.8 |
| 2003..... | 2.3 | * | * | * | * | * | 0.8 | 1.8 | 5.4 | 12.6 | 30.3 | 57.5 |
| 2002..... | 2.3 | * | * | * | * | * | 1.7 | 5.5 | 11.3 | 33.1 | 67.8 | |
| 2001..... | 2.3 | * | * | * | * | * | 0.9 | 1.3 | 4.4 | 13.2 | 35.1 | 70.5 |
| 2000..... | 2.5 | * | * | * | * | * | 0.8 | 2.2 | 5.5 | 15.1 | 29.9 | 64.6 |
| 1999..... | 2.4 | * | * | * | * | * | 0.8 | 1.7 | 4.7 | 16.1 | 29.4 | 53.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | 0.7 | 1.2 | 3.3 | 7.3 | 21.3 | 40.6 |
| 2003..... | 2.3 | * | * | * | * | * | * | 1.8 | 3.2 | 7.8 | 21.0 | 43.7 |
| 2002..... | 2.4 | * | * | * | * | * | 0.8 | 1.4 | 3.4 | 8.3 | 20.7 | 43.9 |
| 2001..... | 2.4 | * | * | * | * | * | 0.8 | 1.3 | 3.5 | 8.8 | 19.9 | 49.2 |
| 2000..... | 2.3 | * | * | * | * | * | * | 1.8 | 3.3 | 9.1 | 20.0 | 47.1 |
| 1999..... | 2.5 | * | * | * | * | * | 0.8 | 1.7 | 3.8 | 9.8 | 22.8 | 48.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Neoplasms of uncertain or unknown behavior of unspecified sites (D48) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 4.3 |
| 2003..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 3.5 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 3.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 3.8 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.4 | 4.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.3 | 3.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.8 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | * | 0.2 | 0.5 | 1.1 | 3.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.1 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.4 | 1.3 | 2.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 4.5 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.5 | 4.1 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 4.6 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.5 | 4.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 4.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 4.3 |
| 2003..... | 0.2 | * | * | * | * | 0.1 | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.6 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.4 | 3.9 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.2 | 4.0 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.4 | 4.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.3 | 3.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | * | 0.5 | 1.2 | 1.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.3 | 2.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.5 | 4.5 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 4.0 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.6 | 4.2 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 4.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 4.7 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.2 | 3.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.2 | 2.3 | 0.4 | 0.2 | 0.5 | 0.8 | 1.4 | 2.2 | 3.5 | 7.0 | 17.3 | 52.9 |
| 2003..... | 3.3 | 2.6 | 0.5 | 0.2 | 0.4 | 0.8 | 1.4 | 2.3 | 3.5 | 7.4 | 17.6 | 56.7 |
| 2002..... | 3.4 | 2.0 | 0.5 | 0.2 | 0.5 | 0.8 | 1.5 | 2.4 | 3.6 | 7.6 | 18.4 | 56.8 |
| 2001..... | 3.3 | 2.6 | 0.5 | 0.2 | 0.4 | 0.8 | 1.5 | 2.3 | 3.6 | 7.3 | 18.9 | 58.5 |
| 2000..... | 3.3 | 1.8 | 0.5 | 0.2 | 0.4 | 0.8 | 1.4 | 2.3 | 3.7 | 7.6 | 18.8 | 58.9 |
| 1999..... | 3.2 | 2.5 | 0.5 | 0.2 | 0.5 | 0.8 | 1.4 | 2.1 | 3.3 | 7.3 | 18.2 | 60.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.8 | 2.8 | 0.5 | 0.2 | 0.5 | 0.8 | 1.5 | 2.2 | 3.7 | 7.3 | 18.1 | 52.1 |
| 2003..... | 2.9 | 2.9 | 0.5 | 0.2 | 0.4 | 0.8 | 1.5 | 2.4 | 3.4 | 7.9 | 18.8 | 55.1 |
| 2002..... | 2.9 | 2.5 | 0.5 | 0.2 | 0.5 | 0.9 | 1.4 | 2.3 | 3.7 | 7.7 | 18.9 | 57.2 |
| 2001..... | 2.8 | 2.9 | 0.6 | 0.2 | 0.4 | 0.9 | 1.5 | 2.2 | 3.3 | 7.2 | 20.0 | 58.1 |
| 2000..... | 2.8 | 2.1 | 0.5 | 0.3 | 0.4 | 0.8 | 1.4 | 2.3 | 3.5 | 8.1 | 19.1 | 60.4 |
| 1999..... | 2.8 | 3.0 | 0.6 | 0.2 | 0.5 | 0.9 | 1.5 | 2.1 | 3.1 | 7.5 | 19.3 | 63.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.6 | 1.8 | 0.4 | 0.2 | 0.4 | 0.8 | 1.3 | 2.1 | 3.3 | 6.8 | 16.8 | 53.3 |
| 2003..... | 3.7 | 2.3 | 0.4 | 0.2 | 0.4 | 0.8 | 1.3 | 2.2 | 3.6 | 6.9 | 16.7 | 57.5 |
| 2002..... | 3.8 | 1.5 | 0.5 | 0.2 | 0.4 | 0.7 | 1.5 | 2.5 | 3.5 | 7.5 | 18.1 | 56.6 |
| 2001..... | 3.9 | 2.4 | 0.3 | 0.2 | 0.4 | 0.7 | 1.5 | 2.3 | 3.8 | 7.5 | 18.2 | 58.7 |
| 2000..... | 3.8 | 1.6 | 0.6 | 0.2 | 0.4 | 0.8 | 1.5 | 2.2 | 3.8 | 7.2 | 18.6 | 58.3 |
| 1999..... | 3.7 | 1.9 | 0.4 | 0.2 | 0.5 | 0.8 | 1.4 | 2.1 | 3.5 | 7.1 | 17.4 | 59.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 0.4 | 0.3 | 0.6 | 1.4 | 2.6 | 4.6 | 11.8 | 37.6 |
| 2003..... | 1.2 | 2.9 | * | * | * | 0.4 | 0.7 | 1.6 | 2.6 | 5.7 | 13.4 | 38.9 |
| 2002..... | 1.3 | 2.5 | * | * | 0.4 | 0.4 | 0.6 | 1.5 | 2.7 | 8.3 | 15.0 | 46.6 |
| 2001..... | 1.3 | 2.4 | * | * | * | 0.5 | 0.8 | 1.1 | 3.1 | 5.8 | 18.9 | 41.9 |
| 2000..... | 1.2 | * | * | * | * | 0.4 | 0.9 | 1.6 | 3.0 | 6.4 | 15.0 | 27.9 |
| 1999..... | 1.2 | * | * | * | * | 0.4 | 0.9 | 1.5 | 2.8 | 6.8 | 13.7 | 47.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.7 | 1.7 | 2.9 | 4.2 | 13.6 | 30.8 |
| 2003..... | 1.2 | * | * | * | * | * | 0.7 | 1.9 | 3.0 | 5.0 | 13.8 | 41.6 |
| 2002..... | 1.3 | * | * | * | * | * | 0.7 | 1.6 | 3.1 | 8.0 | 16.4 | 73.6 |
| 2001..... | 1.2 | * | * | * | * | * | 1.1 | * | 3.5 | 5.6 | 23.3 | * |
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | 1.9 | * | 7.6 | 16.2 | * |
| 1999..... | 1.2 | * | * | * | * | * | 1.2 | 1.4 | * | 9.0 | 15.1 | 56.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | 1.0 | 2.2 | 5.0 | 10.6 | 41.1 |
| 2003..... | 1.3 | * | * | * | * | * | 0.7 | 1.2 | 2.2 | 6.3 | 13.2 | 37.5 |
| 2002..... | 1.3 | * | * | * | * | * | * | 1.3 | 2.4 | 8.5 | 14.1 | 33.2 |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.4 | 2.9 | 6.0 | 16.0 | 44.7 |
| 2000..... | 1.3 | * | * | * | * | * | 1.0 | 1.3 | 4.1 | 5.5 | 14.2 | 28.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.5 | 3.1 | 5.0 | 12.8 | 43.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | 2.5 | 0.4 | 0.2 | 0.5 | 0.9 | 1.5 | 2.3 | 3.6 | 7.2 | 17.6 | 53.6 |
| 2003..... | 3.6 | 2.5 | 0.5 | 0.2 | 0.5 | 0.9 | 1.5 | 2.3 | 3.6 | 7.5 | 17.8 | 57.4 |
| 2002..... | 3.7 | 1.8 | 0.5 | 0.2 | 0.5 | 0.9 | 1.6 | 2.5 | 3.7 | 7.5 | 18.5 | 57.1 |
| 2001..... | 3.6 | 2.6 | 0.5 | 0.2 | 0.4 | 0.9 | 1.6 | 2.4 | 3.6 | 7.4 | 18.8 | 59.0 |
| 2000..... | 3.6 | 1.9 | 0.6 | 0.2 | 0.4 | 0.9 | 1.5 | 2.3 | 3.7 | 7.6 | 18.9 | 59.9 |
| 1999..... | 3.5 | 2.5 | 0.5 | 0.3 | 0.6 | 0.9 | 1.5 | 2.1 | 3.4 | 7.3 | 18.3 | 60.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.1 | 3.1 | 0.5 | 0.2 | 0.5 | 0.9 | 1.6 | 2.3 | 3.8 | 7.5 | 18.3 | 53.2 |
| 2003..... | 3.1 | 2.6 | 0.5 | 0.2 | 0.5 | 0.8 | 1.6 | 2.4 | 3.5 | 8.1 | 19.1 | 55.6 |
| 2002..... | 3.1 | 2.3 | 0.5 | 0.2 | 0.6 | 1.1 | 1.5 | 2.4 | 3.7 | 7.6 | 18.9 | 56.5 |
| 2001..... | 3.1 | 2.8 | 0.6 | 0.2 | 0.5 | 1.0 | 1.5 | 2.4 | 3.3 | 7.3 | 19.8 | 58.9 |
| 2000..... | 3.1 | 2.2 | 0.6 | 0.3 | 0.4 | 0.8 | 1.5 | 2.3 | 3.6 | 8.0 | 19.2 | 61.8 |
| 1999..... | 3.0 | 3.1 | 0.6 | 0.3 | 0.6 | 0.9 | 1.5 | 2.2 | 3.2 | 7.4 | 19.5 | 63.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.0 | 1.8 | 0.4 | 0.2 | 0.4 | 0.8 | 1.4 | 2.2 | 3.4 | 6.9 | 17.1 | 53.8 |
| 2003..... | 4.1 | 2.4 | 0.4 | 0.2 | 0.5 | 0.9 | 1.4 | 2.3 | 3.7 | 7.0 | 16.9 | 58.2 |
| 2002..... | 4.2 | 1.4 | 0.5 | 0.2 | 0.4 | 0.8 | 1.7 | 2.6 | 3.6 | 7.4 | 18.2 | 57.4 |
| 2001..... | 4.2 | 2.4 | 0.3 | 0.2 | 0.4 | 0.8 | 1.7 | 2.4 | 3.9 | 7.5 | 18.2 | 59.0 |
| 2000..... | 4.1 | 1.6 | 0.6 | 0.2 | 0.4 | 0.9 | 1.5 | 2.3 | 3.8 | 7.3 | 18.7 | 59.2 |
| 1999..... | 4.0 | 1.9 | 0.4 | 0.2 | 0.5 | 0.9 | 1.4 | 2.1 | 3.5 | 7.2 | 17.5 | 59.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | 2.0 | 0.3 | 0.2 | 0.2 | 0.4 | 0.7 | 1.4 | 2.6 | 6.6 | 17.4 | 53.2 |
| 2003..... | 3.3 | 2.0 | 0.3 | 0.2 | 0.2 | 0.4 | 0.7 | 1.3 | 2.5 | 6.7 | 17.3 | 57.0 |
| 2002..... | 3.4 | 1.5 | 0.4 | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 2.8 | 6.8 | 17.9 | 56.4 |
| 2001..... | 3.4 | 2.1 | 0.4 | 0.2 | 0.2 | 0.4 | 0.7 | 1.3 | 2.8 | 6.8 | 18.2 | 58.7 |
| 2000..... | 3.4 | 1.7 | 0.4 | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 2.9 | 7.0 | 18.5 | 59.8 |
| 1999..... | 3.2 | 1.7 | 0.3 | 0.2 | 0.3 | 0.3 | 0.6 | 1.3 | 2.5 | 6.6 | 18.0 | 59.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.9 | 2.2 | * | 0.2 | 0.3 | 0.4 | 0.8 | 1.5 | 2.9 | 7.0 | 18.0 | 53.6 |
| 2003..... | 2.8 | 2.3 | * | 0.2 | 0.2 | 0.3 | 0.8 | 1.4 | 2.5 | 7.4 | 18.7 | 55.5 |
| 2002..... | 2.8 | 2.2 | * | * | 0.3 | 0.5 | 0.8 | 1.4 | 3.0 | 7.0 | 18.4 | 55.3 |
| 2001..... | 2.8 | 2.1 | 0.5 | 0.2 | 0.2 | 0.4 | 0.7 | 1.4 | 2.7 | 6.7 | 19.3 | 59.7 |
| 2000..... | 2.8 | 1.8 | 0.5 | 0.2 | 0.2 | 0.3 | 0.8 | 1.4 | 3.0 | 7.6 | 18.8 | 61.2 |
| 1999..... | 2.7 | 2.2 | * | 0.2 | 0.3 | 0.3 | 0.7 | 1.4 | 2.4 | 6.8 | 19.0 | 63.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.7 | 1.8 | * | * | 0.2 | 0.4 | 0.6 | 1.2 | 2.4 | 6.2 | 17.0 | 53.0 |
| 2003..... | 3.8 | * | * | * | 0.2 | 0.4 | 0.6 | 1.2 | 2.6 | 6.1 | 16.3 | 57.7 |
| 2002..... | 3.9 | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.4 | 2.6 | 6.7 | 17.6 | 56.9 |
| 2001..... | 3.9 | 2.1 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 3.0 | 6.9 | 17.5 | 58.3 |
| 2000..... | 3.9 | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 2.8 | 6.5 | 18.3 | 59.3 |
| 1999..... | 3.7 | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 1.1 | 2.6 | 6.5 | 17.3 | 58.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.9 | 3.6 | 0.9 | 0.5 | 1.7 | 3.6 | 6.2 | 8.9 | 12.1 | 14.2 | 22.4 | 66.5 |
| 2003..... | 6.2 | * | 0.9 | 0.5 | 1.6 | 3.5 | 6.6 | 9.8 | 12.3 | 15.3 | 26.2 | 70.8 |
| 2002..... | 6.1 | * | * | 0.5 | 1.6 | 3.6 | 6.7 | 9.9 | 11.5 | 15.1 | 28.3 | 72.9 |
| 2001..... | 6.1 | 3.8 | * | 0.5 | 1.6 | 3.7 | 7.2 | 10.3 | 10.3 | 14.4 | 28.3 | 65.6 |
| 2000..... | 6.0 | * | 1.2 | 0.6 | 1.4 | 4.1 | 6.5 | 10.0 | 11.2 | 14.0 | 26.9 | 67.7 |
| 1999..... | 5.9 | 5.5 | 1.4 | 0.6 | 1.9 | 4.2 | 6.8 | 8.7 | 10.9 | 13.9 | 24.9 | 71.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | 1.8 | 4.0 | 6.5 | 8.5 | 12.6 | 15.0 | 23.4 | 63.0 |
| 2003..... | 5.7 | * | * | * | 1.6 | 3.6 | 6.9 | 10.2 | 12.5 | 15.8 | 28.6 | 68.2 |
| 2002..... | 5.6 | * | * | * | 2.0 | 4.3 | 6.3 | 9.6 | 11.5 | 15.5 | 28.7 | 79.9 |
| 2001..... | 5.4 | * | * | * | 1.8 | 4.5 | 6.9 | 9.8 | 9.2 | 14.5 | 29.4 | 53.1 |
| 2000..... | 5.2 | * | * | 0.7 | 1.3 | 3.9 | 6.2 | 10.1 | 10.0 | 13.2 | 25.6 | 75.1 |
| 1999..... | 5.5 | * | * | 0.7 | 2.1 | 4.5 | 6.9 | 8.5 | 10.5 | 14.0 | 27.2 | 63.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | 1.6 | 3.3 | 6.0 | 9.2 | 11.7 | 13.7 | 21.9 | 67.9 |
| 2003..... | 6.5 | * | * | * | 1.6 | 3.5 | 6.3 | 9.4 | 12.1 | 15.0 | 24.9 | 71.8 |
| 2002..... | 6.6 | * | * | * | 1.2 | 2.9 | 7.1 | 10.2 | 11.5 | 14.9 | 28.0 | 70.2 |
| 2001..... | 6.6 | * | * | * | 1.4 | 3.0 | 7.4 | 10.8 | 11.1 | 14.3 | 27.6 | 70.2 |
| 2000..... | 6.6 | * | * | * | 1.5 | 4.2 | 6.8 | 10.0 | 12.1 | 14.5 | 27.6 | 65.0 |
| 1999..... | 6.3 | * | * | * | 1.6 | 3.8 | 6.7 | 8.9 | 11.2 | 13.8 | 23.6 | 74.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Anemias (D50-D64) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | 0.2 | 0.1 | 0.3 | 0.5 | 0.5 | 0.6 | 1.1 | 2.6 | 8.7 | 38.5 |
| 2003..... | 1.8 | 0.6 | * | 0.1 | 0.3 | 0.5 | 0.4 | 0.6 | 1.0 | 2.7 | 8.9 | 41.2 |
| 2002..... | 1.8 | * | * | 0.1 | 0.3 | 0.5 | 0.5 | 0.7 | 1.0 | 2.7 | 9.2 | 40.5 |
| 2001..... | 1.8 | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.6 | 1.0 | 2.7 | 9.8 | 42.5 |
| 2000..... | 1.8 | * | 0.3 | 0.1 | 0.2 | 0.4 | 0.5 | 0.6 | 1.1 | 2.8 | 9.5 | 43.2 |
| 1999..... | 1.8 | * | 0.2 | 0.1 | 0.4 | 0.4 | 0.5 | 0.7 | 1.0 | 2.6 | 9.6 | 44.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 0.4 | 0.5 | 0.5 | 0.6 | 1.2 | 2.8 | 9.0 | 37.1 |
| 2003..... | 1.4 | * | * | 0.1 | 0.3 | 0.5 | 0.5 | 0.7 | 1.2 | 3.1 | 9.4 | 38.3 |
| 2002..... | 1.4 | * | * | 0.1 | 0.4 | 0.6 | 0.4 | 0.6 | 1.0 | 2.9 | 9.0 | 37.6 |
| 2001..... | 1.4 | * | * | 0.1 | 0.3 | 0.5 | 0.4 | 0.6 | 0.9 | 2.9 | 10.0 | 40.8 |
| 2000..... | 1.3 | * | * | 0.1 | 0.2 | 0.4 | 0.4 | 0.6 | 1.2 | 2.8 | 9.6 | 41.0 |
| 1999..... | 1.4 | * | * | 0.1 | 0.4 | 0.5 | 0.5 | 0.7 | 1.0 | 2.6 | 10.1 | 43.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.6 | 1.0 | 2.5 | 8.5 | 39.1 |
| 2003..... | 2.1 | * | * | 0.1 | 0.3 | 0.4 | 0.4 | 0.6 | 0.8 | 2.3 | 8.6 | 42.4 |
| 2002..... | 2.1 | * | * | 0.2 | 0.2 | 0.4 | 0.5 | 0.7 | 1.0 | 2.6 | 9.3 | 41.8 |
| 2001..... | 2.2 | * | * | 0.2 | 0.3 | 0.4 | 0.6 | 0.6 | 1.0 | 2.6 | 9.7 | 43.1 |
| 2000..... | 2.2 | * | * | * | 0.3 | 0.4 | 0.5 | 0.6 | 1.0 | 2.8 | 9.5 | 44.1 |
| 1999..... | 2.1 | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.6 | 1.0 | 2.6 | 9.3 | 44.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.7 | 2.5 | 8.6 | 38.1 |
| 2003..... | 1.6 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 2.4 | 8.6 | 41.0 |
| 2002..... | 1.6 | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.7 | 2.4 | 8.7 | 39.9 |
| 2001..... | 1.7 | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.8 | 2.5 | 9.4 | 42.1 |
| 2000..... | 1.7 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 9.3 | 42.9 |
| 1999..... | 1.6 | * | * | * | 0.1 | * | 0.1 | 0.3 | 0.6 | 2.4 | 9.4 | 43.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.7 | 8.6 | 37.3 |
| 2003..... | 1.3 | * | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 9.2 | 38.1 |
| 2002..... | 1.2 | * | * | * | * | 0.2 | * | 0.3 | 0.7 | 2.5 | 8.5 | 36.3 |
| 2001..... | 1.3 | * | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 9.8 | 41.2 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.7 | 9.3 | 39.9 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.3 | 0.5 | 2.4 | 9.9 | 43.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 8.6 | 38.4 |
| 2003..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 | 8.2 | 42.2 |
| 2002..... | 2.1 | * | * | * | * | * | * | 0.3 | 0.7 | 2.4 | 8.8 | 41.4 |
| 2001..... | 2.1 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.4 | 9.2 | 42.4 |
| 2000..... | 2.1 | * | * | * | * | * | * | 0.3 | 0.7 | 2.4 | 9.4 | 44.1 |
| 1999..... | 2.0 | * | * | * | * | * | * | 0.3 | 0.7 | 2.4 | 9.1 | 43.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Anemias (D50-D64) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | 0.4 | 1.5 | 2.4 | 2.5 | 2.8 | 4.1 | 4.8 | 11.8 | 48.9 |
| 2003..... | 2.8 | * | * | 0.4 | 1.3 | 2.2 | 2.4 | 3.1 | 4.3 | 5.6 | 14.5 | 50.3 |
| 2002..... | 2.9 | * | * | 0.4 | 1.3 | 2.5 | 2.5 | 3.1 | 3.7 | 6.1 | 16.2 | 52.9 |
| 2001..... | 2.8 | * | * | 0.4 | 1.2 | 2.3 | 2.8 | 3.1 | 3.0 | 5.5 | 16.1 | 49.3 |
| 2000..... | 2.7 | * | * | 0.5 | 1.1 | 2.4 | 2.5 | 3.1 | 3.5 | 5.5 | 13.4 | 51.8 |
| 1999..... | 2.9 | * | 0.9 | 0.5 | 1.5 | 2.7 | 2.6 | 3.2 | 4.3 | 5.0 | 13.4 | 53.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 1.6 | 2.8 | 2.9 | 2.5 | 4.9 | 5.0 | 14.0 | 43.7 |
| 2003..... | 2.8 | * | * | * | 1.3 | 2.3 | 2.6 | 3.4 | 5.6 | 8.0 | 15.3 | 50.1 |
| 2002..... | 2.7 | * | * | * | 1.7 | 2.9 | 2.5 | 2.6 | 4.0 | 7.5 | 17.0 | 58.0 |
| 2001..... | 2.6 | * | * | * | 1.4 | 3.0 | 2.5 | 3.6 | 3.2 | 6.8 | 15.3 | 42.7 |
| 2000..... | 2.4 | * | * | * | 1.0 | 2.2 | 2.1 | 3.4 | 3.5 | 4.6 | 13.9 | 56.4 |
| 1999..... | 2.7 | * | * | * | 1.8 | 2.9 | 2.8 | 3.4 | 5.1 | 5.6 | 12.8 | 44.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | 1.3 | 2.0 | 2.2 | 3.0 | 3.3 | 4.7 | 10.6 | 51.0 |
| 2003..... | 2.9 | * | * | * | 1.3 | 2.1 | 2.2 | 2.8 | 3.3 | 4.0 | 14.0 | 50.3 |
| 2002..... | 3.0 | * | * | * | 1.0 | 2.0 | 2.6 | 3.5 | 3.4 | 5.1 | 15.8 | 51.0 |
| 2001..... | 2.9 | * | * | * | 1.1 | 1.6 | 3.0 | 2.7 | 2.9 | 4.6 | 16.5 | 51.8 |
| 2000..... | 3.1 | * | * | * | 1.1 | 2.5 | 2.8 | 2.9 | 3.5 | 6.1 | 13.2 | 50.2 |
| 1999..... | 3.1 | * | * | * | 1.2 | 2.4 | 2.4 | 3.0 | 3.6 | 4.7 | 13.7 | 56.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nutritional anemias (D50-D53) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 3.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 3.5 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 4.4 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.7 | 4.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.2 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 5.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 3.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 4.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 4.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.5 | 3.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 3.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 3.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 4.4 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 3.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 3.1 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 5.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 4.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.7 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.4 | 4.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 3.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.6 | 3.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 4.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 3.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hemolytic anemias (D55-D59) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | 0.1 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.7 | 1.9 |
| 2003..... | 0.3 | * | * | 0.1 | 0.2 | 0.4 | 0.3 | 0.3 | 0.3 | 0.2 | 0.8 | 1.9 |
| 2002..... | 0.3 | * | * | 0.1 | 0.2 | 0.4 | 0.4 | 0.4 | 0.2 | 0.4 | 0.8 | 2.2 |
| 2001..... | 0.3 | * | * | 0.1 | 0.2 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.9 | 2.3 |
| 2000..... | 0.3 | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | 0.9 | 2.4 |
| 1999..... | 0.3 | * | 0.2 | 0.1 | 0.2 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 0.9 | 2.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.3 | 0.5 | 0.4 | 0.3 | 0.3 | * | 0.9 | 2.2 |
| 2003..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.3 | * | 0.9 | 1.4 |
| 2002..... | 0.3 | * | * | * | 0.3 | 0.5 | 0.3 | 0.3 | 0.2 | 0.4 | 0.5 | 2.3 |
| 2001..... | 0.3 | * | * | * | 0.3 | 0.4 | 0.3 | 0.4 | 0.2 | 0.3 | 1.0 | 2.2 |
| 2000..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.2 | 0.3 | 0.9 | 2.6 |
| 1999..... | 0.3 | * | * | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 1.1 | 1.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.3 | 0.4 | 0.5 | 1.8 |
| 2003..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.8 | 2.1 |
| 2002..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.4 | 0.4 | 0.2 | 0.4 | 1.0 | 2.2 |
| 2001..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.4 | 0.3 | 0.3 | 0.4 | 0.9 | 2.3 |
| 2000..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.9 | 2.3 |
| 1999..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 0.8 | 2.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.9 | 2.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.8 | 2.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.0 | 2.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.0 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 2.5 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.1 | 2.3 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 2.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.1 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.8 | 2.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hemolytic anemias (D55-D59) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 1.4 | 2.3 | 2.2 | 2.2 | 2.0 | 1.2 | * | * |
| 2003..... | 1.4 | * | * | 0.3 | 1.1 | 2.1 | 2.2 | 2.3 | 1.9 | * | * | * |
| 2002..... | 1.5 | * | * | 0.3 | 1.2 | 2.3 | 2.3 | 2.5 | 1.4 | * | * | * |
| 2001..... | 1.4 | * | * | * | 1.1 | 2.0 | 2.4 | 2.4 | 1.5 | * | * | * |
| 2000..... | 1.4 | * | * | 0.4 | 1.0 | 2.1 | 2.2 | 2.4 | 1.7 | * | * | * |
| 1999..... | 1.5 | * | * | 0.4 | 1.3 | 2.5 | 2.3 | 2.3 | 2.0 | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 1.5 | 2.7 | 2.5 | 1.9 | 2.2 | * | * | * |
| 2003..... | 1.5 | * | * | * | 1.0 | 2.2 | 2.3 | 2.3 | 2.3 | * | * | * |
| 2002..... | 1.5 | * | * | * | 1.5 | 2.7 | 2.3 | 2.1 | * | * | * | * |
| 2001..... | 1.5 | * | * | * | 1.3 | 2.7 | 2.2 | 2.7 | * | * | * | * |
| 2000..... | 1.3 | * | * | * | 0.9 | 1.9 | 2.0 | 2.5 | * | * | * | * |
| 1999..... | 1.7 | * | * | * | 1.7 | 2.7 | 2.6 | 2.3 | 2.0 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 1.2 | 2.0 | 2.0 | 2.4 | 1.7 | * | * | * |
| 2003..... | 1.3 | * | * | * | 1.2 | 2.0 | 2.0 | 2.2 | 1.6 | * | * | * |
| 2002..... | 1.4 | * | * | * | 0.9 | 2.0 | 2.3 | 2.8 | 1.5 | * | * | * |
| 2001..... | 1.3 | * | * | * | 0.8 | 1.4 | 2.5 | 2.2 | 1.5 | * | * | * |
| 2000..... | 1.5 | * | * | * | 1.0 | 2.2 | 2.4 | 2.4 | 2.0 | * | * | * |
| 1999..... | 1.4 | * | * | * | 0.9 | 2.3 | 2.0 | 2.3 | 1.9 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Aplastic anemias (D60-D61) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.1 | * | 0.1 | 0.1 | 0.4 | 0.9 | 2.0 | 5.7 |
| 2003..... | 0.4 | * | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.9 | 2.2 | 5.8 |
| 2002..... | 0.4 | * | * | 0.1 | 0.1 | * | 0.1 | 0.1 | 0.3 | 0.8 | 2.4 | 5.3 |
| 2001..... | 0.4 | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 6.8 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 7.6 |
| 1999..... | 0.4 | * | * | * | 0.1 | * | 0.1 | 0.2 | 0.3 | 1.0 | 2.7 | 7.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.3 | 6.3 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 2.6 | 7.6 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.8 | 6.8 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.9 | 7.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.9 | 7.8 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.2 | 10.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.7 | 5.4 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.9 | 5.1 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 4.6 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.5 | 6.3 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.4 | 7.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.3 | 6.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.9 | 2.0 | 5.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 5.9 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 5.2 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.7 | 7.1 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.6 | 7.6 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 2.7 | 8.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.1 | 2.3 | 6.3 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 2.8 | 7.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.9 | 6.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 3.0 | 8.6 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.4 | 1.1 | 2.7 | 7.3 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.2 | 10.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.8 | 5.4 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.9 | 5.1 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 | 4.6 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.5 | 6.4 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.5 | 7.7 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 7.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute posthemorrhagic and other anemias (D62,D64) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.3 | 5.5 | 28.0 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 5.3 | 29.8 |
| 2002..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 5.6 | 29.6 |
| 2001..... | 0.9 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.3 | 5.6 | 29.8 |
| 2000..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 5.5 | 28.8 |
| 1999..... | 0.9 | * | * | * | * | * | 0.0 | 0.2 | 0.3 | 1.2 | 5.3 | 29.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 5.3 | 25.8 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 5.2 | 26.5 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 5.4 | 25.7 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.5 | 5.3 | 27.5 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 5.4 | 25.5 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 5.1 | 27.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 5.7 | 29.0 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 5.4 | 31.3 |
| 2002..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 5.7 | 31.2 |
| 2001..... | 1.2 | * | * | * | * | * | * | * | 0.3 | 1.1 | 5.7 | 30.8 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 0.3 | 1.4 | 5.5 | 30.2 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.3 | 1.2 | 5.4 | 30.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 5.3 | 27.6 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.2 | 1.1 | 5.0 | 29.4 |
| 2002..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 5.0 | 28.8 |
| 2001..... | 1.0 | * | * | * | * | * | * | * | 0.2 | 1.2 | 5.1 | 29.0 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 5.2 | 28.3 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 5.1 | 28.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.9 | 25.8 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | 0.3 | 1.2 | 4.9 | 26.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.8 | 24.3 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 0.2 | 1.3 | 5.0 | 26.9 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.3 | 5.2 | 24.8 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 1.0 | 5.0 | 27.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 0.2 | 1.1 | 5.5 | 28.4 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | * | 1.1 | 5.1 | 30.9 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | 0.3 | 1.1 | 5.2 | 30.7 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | 0.2 | 1.0 | 5.2 | 29.9 |
| 2000..... | 1.3 | * | * | * | * | * | * | * | 0.3 | 1.2 | 5.2 | 29.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 0.3 | 1.0 | 5.2 | 29.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute posthemorrhagic and other anemias (D62,D64) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.5 | 1.5 | 2.6 | 9.2 | 38.3 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.5 | 1.7 | 3.2 | 10.5 | 40.5 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | 1.5 | 3.3 | 13.0 | 42.0 |
| 2001..... | 1.0 | * | * | * | * | * | * | 0.5 | 0.8 | 2.9 | 12.0 | 41.8 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | 1.3 | 3.4 | 9.6 | 37.8 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.7 | 1.2 | 2.8 | 8.9 | 39.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 2.0 | 3.1 | 10.0 | 34.5 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 2.4 | 4.6 | 12.4 | 38.4 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | 1.9 | 4.5 | 13.8 | 45.9 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 3.7 | 11.1 | 38.1 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 8.9 | 34.1 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | * | 8.4 | 30.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | * | 2.1 | 8.7 | 39.8 |
| 2003..... | 1.1 | * | * | * | * | * | * | * | * | 2.2 | 9.5 | 41.3 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | * | 2.4 | 12.6 | 40.6 |
| 2001..... | 1.2 | * | * | * | * | * | * | * | * | 2.4 | 12.5 | 43.2 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | * | 3.8 | 9.9 | 39.2 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | * | 3.0 | 9.1 | 43.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Coagulation defects, purpura and other hemorrhagic conditions (D65-D69) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | 1.0 | * | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.5 | 6.1 | 10.2 |
| 2003..... | 1.0 | 0.9 | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.2 | 2.8 | 5.9 | 10.6 |
| 2002..... | 1.0 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.1 | 2.7 | 6.0 | 11.3 |
| 2001..... | 1.0 | 0.9 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 | 2.7 | 5.9 | 11.0 |
| 2000..... | 1.0 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 2.9 | 6.0 | 11.7 |
| 1999..... | 1.0 | 1.1 | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.2 | 2.9 | 6.0 | 11.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | 1.4 | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.8 | 6.7 | 11.2 |
| 2003..... | 0.9 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 3.1 | 6.6 | 11.9 |
| 2002..... | 0.9 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.2 | 2.7 | 6.6 | 13.8 |
| 2001..... | 0.9 | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.2 | 2.8 | 6.5 | 12.3 |
| 2000..... | 0.9 | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 3.2 | 6.2 | 14.9 |
| 1999..... | 0.9 | 1.3 | * | * | * | 0.1 | 0.4 | 0.7 | 1.1 | 3.0 | 6.2 | 14.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.0 | 2.3 | 5.7 | 9.7 |
| 2003..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.1 | 2.5 | 5.5 | 10.0 |
| 2002..... | 1.1 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.1 | 2.8 | 5.6 | 10.2 |
| 2001..... | 1.1 | * | * | * | * | 0.3 | 0.4 | 0.7 | 1.3 | 2.6 | 5.5 | 10.5 |
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.7 | 5.9 | 10.4 |
| 1999..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.3 | 2.7 | 5.8 | 10.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | 0.9 | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.4 | 6.2 | 10.1 |
| 2003..... | 1.0 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.0 | 2.7 | 5.8 | 10.3 |
| 2002..... | 1.0 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 2.6 | 6.0 | 11.4 |
| 2001..... | 1.0 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.6 | 5.8 | 11.0 |
| 2000..... | 1.0 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 2.8 | 5.9 | 11.9 |
| 1999..... | 1.0 | 0.9 | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 2.6 | 5.8 | 11.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.7 | 6.7 | 11.3 |
| 2003..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 3.0 | 6.5 | 11.7 |
| 2002..... | 0.9 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.6 | 6.5 | 14.0 |
| 2001..... | 0.9 | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.7 | 6.3 | 12.4 |
| 2000..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.0 | 6.2 | 15.3 |
| 1999..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 2.8 | 5.8 | 14.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.9 | 2.1 | 5.8 | 9.6 |
| 2003..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.4 | 5.3 | 9.7 |
| 2002..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 2.6 | 5.6 | 10.2 |
| 2001..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.2 | 2.6 | 5.4 | 10.4 |
| 2000..... | 1.1 | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 2.6 | 5.7 | 10.6 |
| 1999..... | 1.1 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.5 | 5.8 | 10.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Coagulation defects, purpura and other hemorrhagic conditions (D65-D69) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.9 | 1.6 | 2.3 | 4.0 | 6.8 | 13.0 |
| 2003..... | 1.1 | * | * | * | * | * | 0.6 | 1.8 | 2.3 | 4.3 | 7.7 | 14.0 |
| 2002..... | 1.0 | * | * | * | * | * | 0.4 | 1.6 | 2.1 | 4.6 | 7.5 | 13.3 |
| 2001..... | 1.1 | * | * | * | * | 0.5 | 0.9 | 1.9 | 2.1 | 3.6 | 8.1 | 12.2 |
| 2000..... | 1.1 | * | * | * | * | 0.4 | 0.6 | 1.7 | 2.6 | 4.5 | 8.2 | 11.4 |
| 1999..... | 1.2 | * | * | * | * | * | 1.0 | 1.5 | 2.7 | 5.0 | 8.3 | 13.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 1.7 | 2.6 | 4.7 | 7.4 | * |
| 2003..... | 1.0 | * | * | * | * | * | * | 1.8 | 2.5 | 4.9 | 8.1 | * |
| 2002..... | 0.9 | * | * | * | * | * | * | 1.6 | 2.0 | 4.2 | 8.2 | * |
| 2001..... | 1.0 | * | * | * | * | * | 0.8 | 1.9 | 2.0 | 4.0 | 9.0 | * |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.8 | 2.4 | 5.4 | 6.8 | * |
| 1999..... | 1.1 | * | * | * | * | * | 0.8 | 1.4 | 2.5 | 5.4 | 10.9 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 1.1 | 1.4 | 2.1 | 3.6 | 6.4 | 12.4 |
| 2003..... | 1.2 | * | * | * | * | * | 0.7 | 1.8 | 2.1 | 3.9 | 7.5 | 14.0 |
| 2002..... | 1.2 | * | * | * | * | * | * | 1.6 | 2.2 | 4.9 | 7.1 | 12.5 |
| 2001..... | 1.3 | * | * | * | * | * | 1.0 | 1.8 | 2.3 | 3.2 | 7.6 | 13.7 |
| 2000..... | 1.3 | * | * | * | * | * | * | 1.6 | 2.8 | 3.9 | 9.0 | 10.5 |
| 1999..... | 1.3 | * | * | * | * | * | 1.1 | 1.5 | 2.8 | 4.8 | 6.8 | 13.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of blood and blood-forming organs (D70-D76) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 0.6 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 1.6 | 3.1 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.7 | 4.0 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.0 | 3.4 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.9 | 3.9 |
| 2000..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 1.8 | 3.4 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.7 | 3.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 1.9 | 3.4 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.1 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 3.5 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 3.9 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 2.2 | 4.2 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.0 | 2.2 | 4.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 3.0 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 4.0 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.8 | 3.4 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 1.7 | 3.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.1 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.4 | 3.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 1.7 | 3.3 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.7 | 4.2 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.0 | 3.4 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.9 | 3.9 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.9 | 3.5 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.8 | 3.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 | 3.6 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.5 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.3 | 3.4 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 4.0 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 2.1 | 4.3 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.5 | 1.0 | 2.2 | 2.2 | 4.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 3.2 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.6 | 4.1 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.8 | 3.4 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.4 | 0.8 | 1.7 | 3.9 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.7 | 3.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 3.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Certain disorders involving the immune mechanism (D80-D89) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.0 | 1.2 | 1.2 | 1.8 |
| 2003..... | 0.5 | * | * | * | * | 0.2 | 0.7 | 0.8 | 1.0 | 1.2 | 1.2 | 1.6 |
| 2002..... | 0.6 | * | * | * | * | 0.2 | 0.7 | 0.9 | 1.1 | 1.1 | 1.3 | 1.9 |
| 2001..... | 0.5 | 0.7 | * | * | * | 0.2 | 0.6 | 0.9 | 1.0 | 1.2 | 1.3 | 1.6 |
| 2000..... | 0.5 | * | * | * | * | 0.2 | 0.6 | 0.8 | 0.9 | 0.9 | 1.5 | 1.5 |
| 1999..... | 0.4 | * | 0.2 | * | * | 0.2 | 0.5 | 0.7 | 0.7 | 0.9 | 1.0 | 1.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | 0.2 | 0.6 | 0.8 | 0.8 | 1.0 | 0.8 | 1.5 |
| 2003..... | 0.5 | * | * | * | * | 0.2 | 0.7 | 0.8 | 0.7 | 1.1 | 1.2 | * |
| 2002..... | 0.5 | * | * | * | * | 0.2 | 0.7 | 0.9 | 1.0 | 0.9 | 1.1 | 1.6 |
| 2001..... | 0.5 | * | * | * | * | 0.3 | 0.7 | 0.8 | 0.7 | 0.8 | 1.2 | 1.9 |
| 2000..... | 0.5 | * | * | * | * | 0.3 | 0.7 | 0.8 | 0.8 | 0.9 | 1.3 | * |
| 1999..... | 0.4 | * | * | * | * | 0.3 | 0.6 | 0.7 | 0.6 | 0.7 | 1.0 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.1 | 1.4 | 1.4 | 1.9 |
| 2003..... | 0.6 | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.3 | 1.3 | 1.3 | 1.8 |
| 2002..... | 0.6 | * | * | * | * | 0.1 | 0.7 | 0.9 | 1.1 | 1.3 | 1.4 | 2.0 |
| 2001..... | 0.6 | * | * | * | * | 0.1 | 0.6 | 1.0 | 1.2 | 1.5 | 1.3 | 1.5 |
| 2000..... | 0.5 | * | * | * | * | 0.2 | 0.6 | 0.9 | 0.9 | 0.9 | 1.7 | 1.5 |
| 1999..... | 0.5 | * | * | * | * | 0.2 | 0.5 | 0.7 | 0.8 | 1.1 | 1.1 | 1.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.0 | 1.8 |
| 2003..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 1.1 | 1.5 |
| 2002..... | 0.3 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 0.9 | 1.2 | 1.8 |
| 2001..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 1.2 | 1.7 |
| 2000..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.4 | 1.5 |
| 1999..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.0 | 1.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.3 | 0.4 | 0.5 | 0.7 | 0.8 | * |
| 2003..... | 0.3 | * | * | * | * | * | 0.3 | 0.3 | 0.4 | 1.0 | 1.0 | * |
| 2002..... | 0.3 | * | * | * | * | * | 0.3 | 0.4 | 0.7 | 0.8 | 1.1 | * |
| 2001..... | 0.3 | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.4 | 0.6 | 1.0 | 2.0 |
| 2000..... | 0.3 | * | * | * | * | * | 0.3 | 0.3 | 0.6 | 0.8 | 1.2 | * |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 1.0 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.2 | 1.9 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 0.8 | 1.2 | 1.7 |
| 2002..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.3 | 1.9 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.1 | 1.3 | 1.6 |
| 2000..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.5 | 1.4 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.0 | 1.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Certain disorders involving the immune mechanism (D80-D89) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.9 | 2.7 | 4.3 | 5.1 | 4.6 | 3.2 | * |
| 2003..... | 2.0 | * | * | * | * | 0.9 | 3.5 | 4.6 | 5.3 | 4.7 | 2.5 | * |
| 2002..... | 1.9 | * | * | * | * | 0.7 | 3.6 | 4.9 | 5.1 | 3.7 | 2.5 | * |
| 2001..... | 1.9 | * | * | * | * | 0.9 | 3.4 | 5.1 | 4.6 | 4.3 | 2.6 | * |
| 2000..... | 1.8 | * | * | * | * | 1.1 | 3.2 | 4.9 | 4.5 | 2.8 | 3.6 | * |
| 1999..... | 1.5 | * | * | * | * | 1.1 | 3.0 | 3.8 | 3.0 | 2.8 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 1.0 | 2.8 | 4.1 | 4.3 | 4.3 | * | * |
| 2003..... | 1.7 | * | * | * | * | 0.8 | 3.7 | 4.7 | 3.8 | * | * | * |
| 2002..... | 1.7 | * | * | * | * | 1.0 | 3.2 | 5.0 | 4.6 | * | * | * |
| 2001..... | 1.6 | * | * | * | * | 1.2 | 3.5 | 3.9 | 3.5 | * | * | * |
| 2000..... | 1.6 | * | * | * | * | 1.1 | 3.3 | 4.5 | 3.4 | * | * | * |
| 1999..... | 1.3 | * | * | * | * | 1.1 | 2.9 | 3.4 | 2.1 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | 0.9 | 2.5 | 4.5 | 5.7 | 4.8 | 4.3 | * |
| 2003..... | 2.2 | * | * | * | * | 1.1 | 3.3 | 4.6 | 6.4 | 6.4 | * | * |
| 2002..... | 2.1 | * | * | * | * | * | 4.0 | 4.9 | 5.6 | 4.6 | 3.3 | * |
| 2001..... | 2.2 | * | * | * | * | * | 3.3 | 6.1 | 5.5 | 5.5 | * | * |
| 2000..... | 2.0 | * | * | * | * | 1.0 | 3.1 | 5.1 | 5.4 | 3.5 | 4.2 | * |
| 1999..... | 1.7 | * | * | * | * | 1.0 | 3.2 | 4.1 | 3.7 | 3.4 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| IV. Endocrine, nutritional and metabolic diseases (E00-E88) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 33.7 | 5.7 | 0.9 | 0.5 | 1.2 | 3.0 | 7.1 | 18.8 | 47.1 | 107.4 | 226.8 | 470.6 |
| 2003..... | 34.5 | 6.2 | 1.2 | 0.5 | 1.3 | 3.1 | 7.2 | 19.2 | 48.4 | 111.0 | 232.1 | 492.8 |
| 2002..... | 34.5 | 6.6 | 0.9 | 0.5 | 1.3 | 3.1 | 7.4 | 18.7 | 47.9 | 111.1 | 235.3 | 504.8 |
| 2001..... | 34.0 | 7.2 | 1.2 | 0.5 | 1.2 | 2.8 | 6.7 | 18.6 | 46.9 | 110.8 | 232.5 | 519.2 |
| 2000..... | 33.5 | 6.8 | 1.0 | 0.5 | 1.3 | 2.9 | 6.6 | 17.8 | 46.9 | 109.9 | 232.0 | 521.0 |
| 1999..... | 33.3 | 6.9 | 1.0 | 0.5 | 1.4 | 2.6 | 6.5 | 17.2 | 46.5 | 110.5 | 229.3 | 529.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 32.2 | 7.1 | 1.0 | 0.6 | 1.3 | 3.2 | 8.4 | 22.7 | 55.5 | 123.3 | 254.9 | 481.0 |
| 2003..... | 32.5 | 6.8 | 1.4 | 0.6 | 1.4 | 3.6 | 8.5 | 22.8 | 56.5 | 127.3 | 259.6 | 490.8 |
| 2002..... | 31.9 | 7.4 | 0.8 | 0.6 | 1.4 | 3.6 | 8.9 | 22.1 | 56.1 | 123.3 | 255.6 | 504.5 |
| 2001..... | 30.9 | 8.0 | 1.2 | 0.6 | 1.2 | 3.0 | 7.8 | 22.0 | 53.6 | 124.1 | 251.9 | 507.7 |
| 2000..... | 30.2 | 7.4 | 1.1 | 0.6 | 1.4 | 3.4 | 7.7 | 20.8 | 53.1 | 120.8 | 251.5 | 517.4 |
| 1999..... | 29.8 | 7.3 | 1.1 | 0.5 | 1.3 | 2.9 | 7.6 | 20.3 | 52.3 | 120.9 | 250.5 | 521.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 35.2 | 4.2 | 0.8 | 0.5 | 1.2 | 2.7 | 5.8 | 14.9 | 39.2 | 94.0 | 207.8 | 465.9 |
| 2003..... | 36.5 | 5.5 | 1.1 | 0.4 | 1.2 | 2.7 | 5.9 | 15.8 | 40.9 | 97.4 | 213.7 | 493.7 |
| 2002..... | 37.0 | 5.8 | 1.0 | 0.4 | 1.2 | 2.6 | 5.9 | 15.5 | 40.2 | 100.9 | 221.9 | 504.9 |
| 2001..... | 37.0 | 6.4 | 1.2 | 0.5 | 1.3 | 2.5 | 5.5 | 15.3 | 40.6 | 99.7 | 219.8 | 524.0 |
| 2000..... | 36.7 | 6.2 | 1.0 | 0.5 | 1.2 | 2.3 | 5.6 | 14.8 | 41.2 | 100.8 | 219.4 | 522.5 |
| 1999..... | 36.6 | 6.5 | 0.9 | 0.5 | 1.4 | 2.4 | 5.4 | 14.2 | 41.2 | 102.0 | 215.4 | 533.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 17.1 | 5.7 | 1.1 | 0.4 | 0.5 | 1.5 | 4.4 | 15.6 | 53.6 | 136.1 | 290.0 | 472.8 |
| 2003..... | 18.3 | 6.0 | 1.2 | 0.4 | 0.7 | 1.6 | 4.7 | 18.4 | 59.7 | 143.6 | 306.6 | 539.5 |
| 2002..... | 18.0 | 5.4 | 0.8 | 0.4 | 0.7 | 1.4 | 4.9 | 16.8 | 56.7 | 147.9 | 324.3 | 554.3 |
| 2001..... | 18.0 | 5.5 | 0.9 | 0.4 | 0.7 | 1.5 | 4.4 | 17.3 | 58.5 | 153.5 | 315.2 | 620.3 |
| 2000..... | 17.6 | 5.2 | 0.8 | 0.4 | 0.6 | 1.6 | 4.3 | 16.7 | 60.3 | 161.0 | 313.3 | 595.9 |
| 1999..... | 17.8 | 5.7 | 0.7 | 0.3 | 0.6 | 1.2 | 4.3 | 16.3 | 61.6 | 161.8 | 321.2 | 648.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 15.7 | 6.5 | 1.2 | * | 0.6 | 1.7 | 4.9 | 18.8 | 62.5 | 147.3 | 311.5 | 414.7 |
| 2003..... | 16.9 | 6.1 | 1.3 | 0.5 | 0.7 | 1.7 | 5.3 | 20.8 | 69.5 | 158.3 | 331.7 | 528.0 |
| 2002..... | 16.3 | 4.9 | * | 0.6 | 0.8 | 1.6 | 5.7 | 19.3 | 62.8 | 158.6 | 346.3 | 543.8 |
| 2001..... | 16.0 | 5.3 | * | * | 0.7 | 1.8 | 5.4 | 20.9 | 64.8 | 164.3 | 324.2 | 558.0 |
| 2000..... | 16.0 | * | * | * | 0.6 | 2.2 | 4.9 | 18.4 | 70.1 | 176.2 | 330.4 | 560.3 |
| 1999..... | 15.6 | 7.0 | * | * | * | 1.2 | 4.8 | 19.9 | 67.8 | 168.9 | 334.7 | 590.9 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 18.6 | 5.0 | * | * | * | 1.3 | 3.9 | 12.3 | 45.5 | 127.1 | 275.1 | 502.7 |
| 2003..... | 19.7 | 5.9 | * | * | 0.8 | 1.5 | 4.1 | 16.0 | 50.9 | 131.9 | 289.4 | 545.3 |
| 2002..... | 19.7 | 5.9 | * | * | * | 1.3 | 4.1 | 14.3 | 51.1 | 139.3 | 309.2 | 559.5 |
| 2001..... | 20.0 | 5.7 | * | * | 0.7 | 1.2 | 3.3 | 13.8 | 53.0 | 145.0 | 309.2 | 650.7 |
| 2000..... | 19.4 | 6.1 | * | * | * | 0.9 | 3.6 | 15.0 | 51.7 | 149.0 | 301.8 | 613.3 |
| 1999..... | 20.1 | * | * | * | * | 1.1 | 3.8 | 12.7 | 56.2 | 156.3 | 312.2 | 675.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| IV. Endocrine, nutritional and metabolic diseases (E00-E88) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 36.4 | 5.7 | 0.8 | 0.6 | 1.4 | 3.3 | 7.5 | 19.0 | 46.4 | 105.0 | 222.7 | 469.4 |
| 2003..... | 37.0 | 6.2 | 1.2 | 0.5 | 1.4 | 3.5 | 7.6 | 19.3 | 47.3 | 108.5 | 227.6 | 489.6 |
| 2002..... | 36.9 | 6.9 | 0.9 | 0.6 | 1.4 | 3.5 | 7.8 | 18.9 | 47.0 | 108.2 | 230.1 | 501.4 |
| 2001..... | 36.3 | 7.6 | 1.2 | 0.6 | 1.3 | 3.0 | 7.0 | 18.6 | 45.8 | 107.5 | 228.1 | 514.2 |
| 2000..... | 35.7 | 7.2 | 1.1 | 0.5 | 1.4 | 3.1 | 6.9 | 17.8 | 45.7 | 106.3 | 227.9 | 516.7 |
| 1999..... | 35.3 | 7.1 | 1.0 | 0.6 | 1.5 | 2.9 | 6.7 | 17.2 | 45.2 | 107.2 | 224.9 | 524.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 34.9 | 7.3 | 0.9 | 0.6 | 1.4 | 3.6 | 9.0 | 23.1 | 54.7 | 121.3 | 251.1 | 483.3 |
| 2003..... | 35.1 | 7.0 | 1.4 | 0.6 | 1.5 | 4.0 | 9.0 | 22.9 | 55.3 | 124.8 | 255.1 | 487.5 |
| 2002..... | 34.3 | 8.0 | 0.8 | 0.6 | 1.5 | 4.1 | 9.4 | 22.3 | 55.4 | 120.5 | 250.1 | 501.0 |
| 2001..... | 33.1 | 8.7 | 1.2 | 0.6 | 1.3 | 3.3 | 8.2 | 22.0 | 52.6 | 121.0 | 247.8 | 504.3 |
| 2000..... | 32.3 | 8.0 | 1.1 | 0.6 | 1.5 | 3.6 | 8.0 | 21.0 | 51.7 | 117.1 | 247.1 | 514.0 |
| 1999..... | 31.8 | 7.2 | 1.3 | 0.6 | 1.4 | 3.2 | 7.9 | 20.2 | 51.0 | 117.8 | 246.4 | 516.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 37.7 | 4.0 | 0.8 | 0.6 | 1.3 | 3.0 | 6.1 | 15.2 | 38.6 | 91.3 | 203.7 | 463.2 |
| 2003..... | 38.9 | 5.3 | 1.0 | 0.4 | 1.3 | 2.9 | 6.2 | 15.8 | 39.9 | 94.7 | 209.3 | 490.6 |
| 2002..... | 39.4 | 5.8 | 1.1 | 0.5 | 1.4 | 2.9 | 6.2 | 15.6 | 39.2 | 97.9 | 216.9 | 501.6 |
| 2001..... | 39.3 | 6.5 | 1.3 | 0.5 | 1.3 | 2.7 | 5.8 | 15.4 | 39.6 | 96.3 | 215.1 | 518.4 |
| 2000..... | 38.9 | 6.2 | 1.1 | 0.5 | 1.3 | 2.6 | 5.9 | 14.7 | 40.2 | 97.5 | 215.4 | 517.7 |
| 1999..... | 38.7 | 6.9 | 0.8 | 0.6 | 1.6 | 2.6 | 5.6 | 14.3 | 39.9 | 98.5 | 210.9 | 527.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 36.6 | 5.8 | 0.9 | 0.6 | 1.4 | 2.9 | 6.5 | 16.7 | 40.7 | 93.9 | 208.1 | 450.9 |
| 2003..... | 37.3 | 5.1 | 1.2 | 0.6 | 1.4 | 3.0 | 6.7 | 16.7 | 41.4 | 97.2 | 212.4 | 470.6 |
| 2002..... | 37.1 | 6.5 | 1.0 | 0.6 | 1.4 | 3.1 | 6.9 | 16.3 | 40.8 | 96.0 | 215.1 | 482.1 |
| 2001..... | 36.5 | 6.9 | 1.3 | 0.6 | 1.3 | 2.6 | 6.2 | 16.1 | 39.7 | 96.2 | 213.0 | 495.7 |
| 2000..... | 35.9 | 6.3 | 1.2 | 0.6 | 1.4 | 2.7 | 6.2 | 15.3 | 39.8 | 95.0 | 212.0 | 497.5 |
| 1999..... | 35.3 | 6.0 | 1.1 | 0.6 | 1.5 | 2.6 | 5.9 | 14.6 | 38.1 | 95.0 | 209.5 | 504.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 35.6 | 7.5 | 0.9 | 0.6 | 1.4 | 3.1 | 8.0 | 20.8 | 49.2 | 111.3 | 240.0 | 471.9 |
| 2003..... | 35.8 | 6.3 | 1.4 | 0.7 | 1.5 | 3.5 | 8.1 | 20.5 | 49.7 | 115.0 | 243.4 | 479.4 |
| 2002..... | 34.9 | 8.4 | 0.8 | 0.7 | 1.4 | 3.6 | 8.5 | 19.8 | 49.3 | 110.6 | 239.8 | 493.9 |
| 2001..... | 33.7 | 7.6 | 1.3 | 0.6 | 1.3 | 2.8 | 7.4 | 19.3 | 46.8 | 112.1 | 235.5 | 495.6 |
| 2000..... | 32.9 | 7.2 | 1.3 | 0.7 | 1.5 | 3.2 | 7.3 | 18.7 | 46.0 | 108.0 | 235.9 | 505.5 |
| 1999..... | 32.1 | 5.9 | 1.4 | 0.6 | 1.5 | 2.9 | 7.0 | 17.7 | 44.6 | 108.0 | 235.2 | 499.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 37.6 | 4.0 | 0.8 | 0.6 | 1.3 | 2.7 | 5.1 | 12.7 | 32.6 | 79.0 | 186.3 | 441.4 |
| 2003..... | 38.7 | 3.9 | 1.0 | 0.4 | 1.3 | 2.5 | 5.2 | 13.0 | 33.5 | 82.1 | 191.5 | 466.8 |
| 2002..... | 39.2 | 4.4 | 1.2 | 0.4 | 1.4 | 2.6 | 5.3 | 12.9 | 32.6 | 83.5 | 198.6 | 477.0 |
| 2001..... | 39.2 | 6.2 | 1.3 | 0.6 | 1.3 | 2.5 | 5.0 | 12.9 | 33.0 | 82.7 | 198.1 | 495.7 |
| 2000..... | 38.7 | 5.4 | 1.1 | 0.5 | 1.3 | 2.1 | 5.1 | 11.9 | 34.0 | 83.9 | 196.3 | 494.3 |
| 1999..... | 38.3 | 6.0 | 0.9 | 0.6 | 1.6 | 2.4 | 4.7 | 11.5 | 32.0 | 84.0 | 192.7 | 506.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| IV. Endocrine, nutritional and metabolic diseases (E00-E88) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 43.2 | 5.7 | * | 0.5 | 1.8 | 6.2 | 14.9 | 37.8 | 97.9 | 214.5 | 402.5 | 725.6 |
| 2003..... | 43.8 | 10.0 | 1.0 | 0.4 | 1.7 | 6.7 | 14.3 | 39.4 | 102.6 | 217.5 | 411.2 | 743.7 |
| 2002..... | 44.1 | 8.3 | 0.9 | 0.6 | 2.0 | 6.5 | 14.4 | 39.7 | 103.7 | 223.2 | 414.2 | 765.8 |
| 2001..... | 43.0 | 11.0 | 0.9 | 0.5 | 1.8 | 5.7 | 13.2 | 39.1 | 100.6 | 219.9 | 413.8 | 770.4 |
| 2000..... | 42.8 | 9.5 | 0.8 | 0.4 | 1.8 | 6.3 | 12.7 | 38.7 | 98.3 | 220.2 | 429.6 | 772.4 |
| 1999..... | 42.8 | 12.0 | 0.9 | 0.6 | 1.6 | 5.2 | 13.4 | 38.7 | 106.0 | 225.6 | 409.7 | 766.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 39.0 | 7.1 | * | 0.6 | 1.8 | 7.0 | 17.3 | 43.0 | 109.9 | 236.7 | 416.4 | 688.9 |
| 2003..... | 38.5 | 9.3 | * | * | 1.6 | 8.0 | 16.0 | 43.7 | 113.3 | 234.2 | 426.5 | 626.5 |
| 2002..... | 37.9 | 7.3 | * | * | 2.4 | 7.8 | 16.8 | 44.3 | 116.8 | 226.4 | 408.5 | 618.1 |
| 2001..... | 37.1 | 12.4 | * | * | 1.7 | 6.8 | 14.9 | 44.1 | 110.0 | 224.8 | 428.5 | 664.1 |
| 2000..... | 35.7 | 10.6 | * | * | 2.0 | 7.1 | 14.0 | 42.0 | 106.6 | 219.8 | 417.1 | 662.2 |
| 1999..... | 36.4 | 13.2 | * | 0.6 | 1.6 | 5.9 | 15.2 | 42.7 | 111.5 | 227.0 | 400.2 | 748.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 47.0 | * | * | * | 1.7 | 5.4 | 12.7 | 33.4 | 88.2 | 198.7 | 394.6 | 740.1 |
| 2003..... | 48.7 | 10.6 | * | * | 1.8 | 5.5 | 12.8 | 35.7 | 94.1 | 205.6 | 402.5 | 789.1 |
| 2002..... | 49.7 | 9.5 | * | 0.7 | 1.6 | 5.4 | 12.4 | 35.7 | 93.1 | 220.9 | 417.3 | 822.3 |
| 2001..... | 48.4 | 9.4 | * | * | 1.9 | 4.7 | 11.7 | 34.9 | 93.1 | 216.5 | 405.5 | 809.7 |
| 2000..... | 49.2 | 8.5 | * | * | 1.7 | 5.6 | 11.5 | 35.9 | 91.6 | 220.5 | 436.6 | 813.3 |
| 1999..... | 48.7 | 10.7 | * | * | 1.6 | 4.7 | 11.9 | 35.4 | 101.6 | 224.7 | 414.9 | 773.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Endocrine diseases (E00-E34) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 25.8 | 0.6 | 0.1 | 0.1 | 0.5 | 1.6 | 4.8 | 13.8 | 37.8 | 88.8 | 182.3 | 330.4 |
| 2003..... | 26.5 | 0.7 | 0.1 | 0.1 | 0.5 | 1.8 | 4.8 | 14.2 | 39.3 | 92.5 | 187.3 | 342.7 |
| 2002..... | 26.4 | 1.0 | * | 0.1 | 0.5 | 1.7 | 5.0 | 14.1 | 38.4 | 93.2 | 188.8 | 347.6 |
| 2001..... | 26.1 | 1.1 | * | 0.1 | 0.4 | 1.6 | 4.6 | 14.0 | 38.5 | 93.0 | 187.7 | 350.3 |
| 2000..... | 25.6 | 1.2 | 0.1 | 0.1 | 0.5 | 1.7 | 4.5 | 13.5 | 38.6 | 92.3 | 185.5 | 345.2 |
| 1999..... | 25.5 | 1.0 | 0.1 | 0.1 | 0.5 | 1.6 | 4.5 | 13.4 | 39.0 | 93.7 | 183.9 | 343.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 25.0 | * | * | 0.1 | 0.5 | 1.8 | 5.9 | 16.8 | 44.6 | 101.5 | 206.9 | 349.1 |
| 2003..... | 25.3 | * | * | 0.1 | 0.5 | 2.0 | 5.9 | 17.0 | 45.6 | 105.4 | 211.6 | 354.4 |
| 2002..... | 24.8 | 1.1 | * | 0.1 | 0.5 | 2.0 | 6.1 | 16.7 | 45.1 | 102.8 | 206.6 | 365.2 |
| 2001..... | 24.1 | 1.3 | * | 0.1 | 0.5 | 1.8 | 5.5 | 16.7 | 43.7 | 103.2 | 203.5 | 362.2 |
| 2000..... | 23.4 | 1.3 | * | * | 0.5 | 2.0 | 5.3 | 15.5 | 43.0 | 100.4 | 203.5 | 360.9 |
| 1999..... | 23.3 | * | * | * | 0.5 | 1.7 | 5.3 | 15.9 | 43.2 | 101.3 | 203.1 | 361.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 26.7 | * | * | 0.1 | 0.5 | 1.4 | 3.7 | 10.8 | 31.5 | 78.0 | 165.8 | 322.0 |
| 2003..... | 27.7 | * | * | 0.1 | 0.4 | 1.6 | 3.8 | 11.6 | 33.4 | 81.8 | 171.0 | 337.6 |
| 2002..... | 28.0 | * | * | * | 0.5 | 1.4 | 3.9 | 11.6 | 32.2 | 85.3 | 177.1 | 339.9 |
| 2001..... | 28.0 | * | * | * | 0.4 | 1.4 | 3.7 | 11.4 | 33.7 | 84.6 | 177.3 | 345.3 |
| 2000..... | 27.7 | 1.1 | * | 0.1 | 0.4 | 1.4 | 3.7 | 11.5 | 34.5 | 85.6 | 173.8 | 338.8 |
| 1999..... | 27.5 | 1.3 | * | 0.1 | 0.5 | 1.5 | 3.8 | 11.0 | 35.0 | 87.4 | 171.3 | 336.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 14.8 | * | * | * | * | 0.9 | 3.1 | 13.4 | 48.7 | 125.8 | 261.4 | 382.9 |
| 2003..... | 15.8 | * | * | * | 0.3 | 0.9 | 3.5 | 15.8 | 54.4 | 133.2 | 277.4 | 444.6 |
| 2002..... | 15.6 | * | * | * | * | 0.8 | 3.7 | 14.3 | 52.6 | 135.8 | 289.2 | 459.0 |
| 2001..... | 15.6 | * | * | * | 0.3 | 0.8 | 3.1 | 15.2 | 53.6 | 142.8 | 283.3 | 510.8 |
| 2000..... | 15.4 | * | * | * | * | 1.1 | 3.1 | 14.3 | 56.1 | 149.0 | 282.3 | 481.1 |
| 1999..... | 15.6 | * | * | * | * | 0.7 | 3.2 | 14.2 | 57.2 | 151.6 | 290.6 | 515.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 13.6 | * | * | * | * | 1.0 | 3.5 | 16.1 | 57.1 | 135.1 | 282.2 | 337.6 |
| 2003..... | 14.7 | * | * | * | * | 0.9 | 4.0 | 18.2 | 63.0 | 147.1 | 303.0 | 435.6 |
| 2002..... | 14.1 | * | * | * | * | 1.0 | 4.2 | 16.9 | 59.2 | 144.5 | 309.0 | 441.7 |
| 2001..... | 13.8 | * | * | * | * | 0.9 | 3.9 | 18.8 | 58.8 | 152.8 | 288.4 | 436.1 |
| 2000..... | 14.0 | * | * | * | * | 1.4 | 3.7 | 15.7 | 65.2 | 161.4 | 301.4 | 463.6 |
| 1999..... | 13.6 | * | * | * | * | 0.7 | 3.3 | 17.4 | 62.7 | 157.3 | 299.3 | 478.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 16.1 | * | * | * | * | 0.8 | 2.7 | 10.8 | 41.1 | 118.3 | 247.0 | 406.2 |
| 2003..... | 17.1 | * | * | * | * | 1.0 | 3.0 | 13.5 | 46.7 | 122.1 | 259.9 | 449.2 |
| 2002..... | 17.1 | * | * | * | * | 0.7 | 3.1 | 11.7 | 46.7 | 128.8 | 275.7 | 467.5 |
| 2001..... | 17.5 | * | * | * | * | 0.7 | 2.3 | 11.7 | 49.0 | 135.0 | 279.9 | 547.3 |
| 2000..... | 16.8 | * | * | * | * | * | 2.5 | 12.9 | 48.1 | 139.2 | 269.4 | 489.7 |
| 1999..... | 17.7 | * | * | * | * | 0.8 | 3.1 | 11.1 | 52.4 | 147.2 | 284.8 | 533.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Endocrine diseases (E00-E34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 27.6 | 0.7 | * | 0.2 | 0.5 | 1.8 | 5.1 | 13.7 | 36.8 | 85.8 | 177.5 | 327.1 |
| 2003..... | 28.2 | 0.9 | * | 0.1 | 0.5 | 2.0 | 5.0 | 14.0 | 38.0 | 89.5 | 182.1 | 337.6 |
| 2002..... | 28.0 | 1.1 | * | 0.1 | 0.5 | 1.9 | 5.2 | 14.0 | 37.1 | 90.0 | 183.2 | 342.2 |
| 2001..... | 27.6 | 1.1 | * | 0.1 | 0.4 | 1.8 | 4.8 | 13.8 | 37.2 | 89.4 | 182.8 | 343.4 |
| 2000..... | 27.0 | 1.2 | 0.2 | 0.1 | 0.5 | 1.8 | 4.7 | 13.4 | 37.1 | 88.5 | 180.9 | 339.1 |
| 1999..... | 26.8 | 1.1 | 0.2 | 0.1 | 0.5 | 1.7 | 4.7 | 13.3 | 37.5 | 90.0 | 178.9 | 336.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 26.9 | * | * | 0.2 | 0.5 | 2.0 | 6.3 | 16.8 | 43.4 | 98.9 | 202.1 | 348.8 |
| 2003..... | 27.0 | * | * | 0.1 | 0.6 | 2.3 | 6.2 | 16.8 | 44.1 | 102.3 | 206.1 | 349.8 |
| 2002..... | 26.4 | * | * | 0.1 | 0.6 | 2.3 | 6.3 | 16.6 | 43.8 | 99.7 | 200.7 | 360.7 |
| 2001..... | 25.6 | 1.5 | * | 0.2 | 0.5 | 2.0 | 5.7 | 16.5 | 42.4 | 99.5 | 199.0 | 358.3 |
| 2000..... | 24.8 | * | * | * | 0.6 | 2.1 | 5.5 | 15.5 | 41.2 | 96.3 | 198.6 | 355.4 |
| 1999..... | 24.7 | * | * | * | 0.5 | 1.9 | 5.5 | 15.7 | 41.6 | 97.8 | 198.6 | 355.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 28.2 | * | * | 0.1 | 0.5 | 1.6 | 3.9 | 10.8 | 30.6 | 74.8 | 161.0 | 317.4 |
| 2003..... | 29.2 | * | * | * | 0.4 | 1.7 | 3.9 | 11.3 | 32.2 | 78.7 | 166.0 | 332.2 |
| 2002..... | 29.5 | * | * | * | 0.5 | 1.6 | 4.0 | 11.5 | 31.0 | 82.0 | 171.7 | 334.3 |
| 2001..... | 29.4 | * | * | * | 0.4 | 1.5 | 3.9 | 11.3 | 32.4 | 80.9 | 172.1 | 337.2 |
| 2000..... | 29.1 | * | * | * | 0.4 | 1.5 | 3.9 | 11.3 | 33.3 | 82.0 | 169.4 | 332.4 |
| 1999..... | 28.8 | 1.4 | * | 0.1 | 0.5 | 1.6 | 3.8 | 10.9 | 33.6 | 83.6 | 166.2 | 328.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 27.2 | * | * | 0.1 | 0.4 | 1.5 | 4.3 | 11.6 | 31.3 | 75.1 | 163.2 | 308.6 |
| 2003..... | 27.7 | * | * | 0.1 | 0.4 | 1.6 | 4.3 | 11.6 | 32.1 | 78.4 | 167.8 | 319.7 |
| 2002..... | 27.6 | 1.2 | * | 0.1 | 0.4 | 1.7 | 4.5 | 11.6 | 31.3 | 78.5 | 169.4 | 324.5 |
| 2001..... | 27.2 | 1.1 | * | 0.1 | 0.3 | 1.4 | 4.1 | 11.6 | 31.4 | 78.5 | 168.9 | 326.7 |
| 2000..... | 26.5 | 0.9 | * | * | 0.4 | 1.5 | 4.1 | 11.1 | 31.6 | 77.6 | 165.9 | 321.5 |
| 1999..... | 26.1 | 1.0 | * | 0.1 | 0.5 | 1.5 | 4.0 | 10.9 | 30.6 | 78.5 | 164.8 | 317.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 27.0 | * | * | * | 0.4 | 1.7 | 5.5 | 14.6 | 37.9 | 89.0 | 191.4 | 336.8 |
| 2003..... | 27.2 | * | * | * | 0.5 | 1.8 | 5.4 | 14.4 | 38.5 | 92.7 | 195.8 | 341.1 |
| 2002..... | 26.6 | * | * | * | 0.5 | 2.0 | 5.6 | 14.1 | 38.1 | 90.2 | 191.5 | 355.7 |
| 2001..... | 25.8 | * | * | * | 0.4 | 1.5 | 5.0 | 14.0 | 37.0 | 91.1 | 187.9 | 351.8 |
| 2000..... | 24.9 | * | * | * | 0.5 | 1.8 | 4.9 | 13.2 | 36.0 | 87.7 | 188.4 | 346.0 |
| 1999..... | 24.6 | * | * | * | 0.5 | 1.6 | 4.8 | 13.3 | 35.5 | 88.8 | 189.1 | 342.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 27.4 | * | * | * | 0.4 | 1.3 | 3.2 | 8.7 | 25.1 | 63.2 | 144.0 | 295.9 |
| 2003..... | 28.3 | * | * | * | 0.3 | 1.4 | 3.2 | 8.8 | 26.2 | 66.3 | 148.9 | 310.3 |
| 2002..... | 28.6 | * | * | * | 0.4 | 1.3 | 3.4 | 9.2 | 24.8 | 68.6 | 154.6 | 311.1 |
| 2001..... | 28.6 | * | * | * | 0.2 | 1.3 | 3.2 | 9.2 | 26.1 | 67.8 | 156.3 | 316.3 |
| 2000..... | 28.1 | * | * | * | 0.3 | 1.2 | 3.3 | 8.9 | 27.5 | 69.0 | 151.1 | 311.7 |
| 1999..... | 27.6 | * | * | * | 0.4 | 1.3 | 3.2 | 8.5 | 26.1 | 69.9 | 148.8 | 307.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Endocrine diseases (E00-E34) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 35.4 | * | * | 0.3 | 1.1 | 3.8 | 10.3 | 29.7 | 83.5 | 186.5 | 344.9 | 559.8 |
| 2003..... | 36.0 | * | * | * | 0.9 | 4.4 | 10.2 | 31.6 | 89.2 | 191.8 | 347.6 | 558.1 |
| 2002..... | 35.7 | * | * | * | 1.1 | 4.0 | 10.4 | 32.0 | 87.6 | 194.7 | 346.4 | 566.8 |
| 2001..... | 35.1 | * | * | * | 1.0 | 4.0 | 9.6 | 30.9 | 87.9 | 193.9 | 347.0 | 560.0 |
| 2000..... | 35.0 | * | * | * | 1.0 | 4.1 | 8.9 | 31.5 | 85.0 | 194.5 | 364.4 | 556.2 |
| 1999..... | 35.1 | * | * | * | 0.9 | 3.4 | 9.7 | 32.1 | 94.0 | 197.7 | 345.0 | 554.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 32.1 | * | * | * | 1.1 | 4.4 | 12.6 | 34.9 | 95.0 | 205.9 | 352.0 | 528.4 |
| 2003..... | 31.7 | * | * | * | 1.0 | 5.2 | 12.2 | 35.5 | 99.3 | 203.2 | 350.9 | 477.3 |
| 2002..... | 30.6 | * | * | * | 1.2 | 4.9 | 12.5 | 36.8 | 98.2 | 196.5 | 336.7 | 434.3 |
| 2001..... | 30.2 | * | * | * | 0.9 | 5.2 | 11.1 | 35.8 | 94.9 | 193.1 | 354.1 | 472.4 |
| 2000..... | 29.0 | * | * | * | 1.0 | 4.4 | 10.1 | 34.7 | 90.9 | 190.3 | 345.9 | 479.1 |
| 1999..... | 29.5 | * | * | * | 0.8 | 3.6 | 11.2 | 36.2 | 97.6 | 194.3 | 326.8 | 524.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 38.4 | * | * | * | 1.1 | 3.2 | 8.2 | 25.3 | 74.3 | 172.6 | 340.9 | 572.2 |
| 2003..... | 39.9 | * | * | * | 0.8 | 3.6 | 8.5 | 28.3 | 81.1 | 183.7 | 345.8 | 589.3 |
| 2002..... | 40.4 | * | * | * | 1.0 | 3.3 | 8.6 | 27.9 | 79.2 | 193.4 | 351.8 | 617.5 |
| 2001..... | 39.6 | * | * | * | 1.1 | 3.0 | 8.2 | 26.8 | 82.3 | 194.4 | 343.0 | 592.4 |
| 2000..... | 40.4 | * | * | * | 1.1 | 3.8 | 7.9 | 28.8 | 80.2 | 197.4 | 374.8 | 584.8 |
| 1999..... | 40.3 | * | * | * | 1.0 | 3.2 | 8.3 | 28.6 | 91.0 | 200.1 | 355.2 | 565.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of thyroid gland (E00-E07) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 3.6 | 19.3 |
| 2003..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.9 | 4.2 | 21.4 |
| 2002..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 1.0 | 4.0 | 22.8 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 4.2 | 23.4 |
| 2000..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.4 | 0.9 | 4.0 | 21.4 |
| 1999..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 1.0 | 4.0 | 22.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.3 | 11.1 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 10.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 12.6 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 2.4 | 14.2 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 12.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.6 | 13.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 4.5 | 22.9 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 5.6 | 26.1 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 5.2 | 27.2 |
| 2001..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 5.4 | 27.2 |
| 2000..... | 1.1 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 5.3 | 25.1 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.2 | 4.9 | 26.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 3.5 | 19.9 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 4.3 | 22.0 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 4.1 | 23.6 |
| 2001..... | 0.8 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 4.1 | 24.0 |
| 2000..... | 0.8 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 4.0 | 21.9 |
| 1999..... | 0.8 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 3.9 | 23.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.6 | 2.2 | 11.5 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | 0.6 | 2.1 | 11.1 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 0.6 | 2.2 | 13.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.3 | 14.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.5 | 2.1 | 12.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.5 | 0.5 | 2.4 | 13.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 4.4 | 23.7 |
| 2003..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 5.7 | 26.8 |
| 2002..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 5.4 | 28.2 |
| 2001..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 5.3 | 27.9 |
| 2000..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 5.2 | 25.9 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.3 | 1.2 | 4.9 | 27.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of thyroid gland (E00-E07) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 1.9 | 5.4 | 15.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.4 | * | 1.4 | 4.1 | 19.6 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.5 | 0.8 | 1.3 | 3.2 | 16.9 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 1.4 | 5.8 | 19.7 |
| 2000..... | 0.6 | * | * | * | * | * | 0.5 | * | * | 1.9 | 5.0 | 19.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.6 | 0.9 | 1.5 | 5.3 | 18.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 2.2 | 6.3 | 18.1 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 4.4 | 23.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | * | 4.0 | 20.1 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | * | 6.2 | 22.2 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | * | 6.9 | 20.1 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | * | 5.8 | 21.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diabetes mellitus (E10-E14) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 24.9 | * | * | 0.1 | 0.4 | 1.5 | 4.6 | 13.4 | 37.1 | 87.2 | 176.9 | 307.0 |
| 2003..... | 25.5 | * | * | 0.1 | 0.4 | 1.6 | 4.6 | 13.9 | 38.5 | 90.8 | 181.1 | 317.5 |
| 2002..... | 25.4 | * | * | 0.1 | 0.4 | 1.6 | 4.8 | 13.7 | 37.7 | 91.4 | 182.8 | 320.6 |
| 2001..... | 25.1 | * | * | 0.1 | 0.4 | 1.5 | 4.3 | 13.6 | 37.8 | 91.4 | 181.4 | 321.8 |
| 2000..... | 24.6 | * | * | 0.1 | 0.4 | 1.6 | 4.3 | 13.1 | 37.8 | 90.7 | 179.5 | 319.7 |
| 1999..... | 24.5 | * | * | 0.1 | 0.4 | 1.4 | 4.3 | 12.9 | 38.3 | 91.8 | 178.0 | 317.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 24.4 | * | * | * | 0.4 | 1.7 | 5.7 | 16.5 | 44.0 | 100.0 | 202.5 | 333.9 |
| 2003..... | 24.8 | * | * | * | 0.5 | 1.9 | 5.7 | 16.7 | 45.0 | 104.0 | 207.2 | 340.7 |
| 2002..... | 24.2 | * | * | 0.1 | 0.5 | 1.9 | 5.9 | 16.4 | 44.5 | 101.4 | 202.6 | 348.5 |
| 2001..... | 23.5 | * | * | 0.1 | 0.4 | 1.7 | 5.3 | 16.4 | 43.1 | 101.7 | 198.9 | 343.1 |
| 2000..... | 22.9 | * | * | * | 0.5 | 1.9 | 5.1 | 15.3 | 42.4 | 99.1 | 199.6 | 345.1 |
| 1999..... | 22.8 | * | * | * | 0.4 | 1.6 | 5.1 | 15.6 | 42.6 | 99.8 | 198.2 | 345.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 25.4 | * | * | 0.1 | 0.4 | 1.3 | 3.5 | 10.4 | 30.7 | 76.4 | 159.6 | 294.9 |
| 2003..... | 26.2 | * | * | * | 0.3 | 1.4 | 3.5 | 11.2 | 32.4 | 79.8 | 163.6 | 307.2 |
| 2002..... | 26.5 | * | * | * | 0.4 | 1.3 | 3.7 | 11.1 | 31.4 | 83.1 | 169.7 | 308.4 |
| 2001..... | 26.6 | * | * | * | 0.3 | 1.3 | 3.4 | 11.0 | 32.9 | 82.8 | 169.9 | 312.9 |
| 2000..... | 26.3 | * | * | * | 0.4 | 1.2 | 3.4 | 11.1 | 33.6 | 83.7 | 166.3 | 309.4 |
| 1999..... | 26.2 | * | * | * | 0.4 | 1.3 | 3.5 | 10.4 | 34.3 | 85.2 | 164.9 | 305.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 14.5 | * | * | * | * | 0.8 | 3.0 | 13.3 | 48.3 | 124.9 | 258.4 | 365.8 |
| 2003..... | 15.5 | * | * | * | * | 0.9 | 3.3 | 15.5 | 53.7 | 131.8 | 272.1 | 428.1 |
| 2002..... | 15.3 | * | * | * | * | 0.8 | 3.5 | 14.0 | 52.1 | 134.7 | 285.6 | 437.9 |
| 2001..... | 15.3 | * | * | * | * | 0.7 | 3.1 | 14.9 | 53.2 | 141.2 | 279.2 | 489.6 |
| 2000..... | 15.1 | * | * | * | * | 1.0 | 3.0 | 14.0 | 55.5 | 147.9 | 278.5 | 465.1 |
| 1999..... | 15.3 | * | * | * | * | 0.7 | 3.2 | 13.9 | 56.6 | 149.7 | 285.2 | 493.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 13.4 | * | * | * | * | 0.9 | 3.4 | 16.0 | 56.9 | 134.2 | 279.3 | 325.0 |
| 2003..... | 14.4 | * | * | * | * | 0.8 | 3.9 | 18.1 | 62.4 | 146.3 | 297.2 | 426.4 |
| 2002..... | 13.9 | * | * | * | * | 0.9 | 4.1 | 16.7 | 58.9 | 143.7 | 306.9 | 425.0 |
| 2001..... | 13.6 | * | * | * | * | 0.8 | 3.9 | 18.5 | 58.7 | 151.4 | 285.3 | 422.8 |
| 2000..... | 13.8 | * | * | * | * | 1.4 | 3.6 | 15.5 | 64.7 | 160.8 | 299.4 | 449.4 |
| 1999..... | 13.4 | * | * | * | * | 0.6 | 3.3 | 17.2 | 62.5 | 155.8 | 293.0 | 468.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 15.7 | * | * | * | * | 0.8 | 2.6 | 10.6 | 40.5 | 117.5 | 244.0 | 386.8 |
| 2003..... | 16.6 | * | * | * | * | 0.9 | 2.7 | 13.0 | 45.8 | 120.3 | 254.9 | 428.9 |
| 2002..... | 16.7 | * | * | * | * | 0.7 | 2.9 | 11.4 | 46.0 | 127.6 | 270.9 | 444.3 |
| 2001..... | 17.1 | * | * | * | * | 0.7 | 2.2 | 11.4 | 48.3 | 133.2 | 275.0 | 522.2 |
| 2000..... | 16.5 | * | * | * | * | * | 2.4 | 12.6 | 47.4 | 137.7 | 264.5 | 472.8 |
| 1999..... | 17.3 | * | * | * | * | 0.8 | 3.1 | 10.7 | 51.4 | 144.9 | 280.0 | 506.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diabetes mellitus (E10-E14) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 26.5 | * | * | 0.1 | 0.4 | 1.6 | 4.8 | 13.3 | 36.0 | 84.2 | 172.0 | 303.5 |
| 2003..... | 27.0 | * | * | 0.1 | 0.4 | 1.8 | 4.8 | 13.6 | 37.1 | 87.7 | 175.8 | 312.0 |
| 2002..... | 26.9 | * | * | 0.1 | 0.5 | 1.8 | 5.0 | 13.6 | 36.4 | 88.2 | 177.2 | 315.0 |
| 2001..... | 26.4 | * | * | 0.1 | 0.4 | 1.7 | 4.5 | 13.4 | 36.5 | 87.7 | 176.4 | 314.8 |
| 2000..... | 25.9 | * | * | 0.1 | 0.5 | 1.7 | 4.4 | 13.0 | 36.4 | 86.8 | 174.8 | 313.4 |
| 1999..... | 25.7 | * | * | 0.1 | 0.4 | 1.6 | 4.4 | 12.8 | 36.8 | 88.1 | 173.1 | 310.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 26.2 | * | * | * | 0.4 | 1.9 | 6.1 | 16.5 | 42.7 | 97.3 | 197.7 | 333.5 |
| 2003..... | 26.4 | * | * | * | 0.5 | 2.1 | 6.0 | 16.5 | 43.5 | 100.9 | 201.8 | 335.9 |
| 2002..... | 25.8 | * | * | 0.1 | 0.5 | 2.2 | 6.2 | 16.3 | 43.1 | 98.2 | 196.6 | 344.0 |
| 2001..... | 25.0 | * | * | 0.1 | 0.4 | 1.9 | 5.5 | 16.1 | 41.8 | 98.0 | 194.3 | 338.9 |
| 2000..... | 24.2 | * | * | * | 0.5 | 2.0 | 5.3 | 15.2 | 40.7 | 95.0 | 194.6 | 339.7 |
| 1999..... | 24.1 | * | * | * | 0.4 | 1.8 | 5.3 | 15.3 | 41.0 | 96.3 | 193.8 | 339.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 26.8 | * | * | 0.1 | 0.4 | 1.4 | 3.6 | 10.3 | 29.8 | 73.1 | 154.7 | 290.1 |
| 2003..... | 27.6 | * | * | * | 0.3 | 1.5 | 3.6 | 10.9 | 31.2 | 76.7 | 158.5 | 301.5 |
| 2002..... | 27.9 | * | * | * | 0.4 | 1.4 | 3.8 | 11.1 | 30.1 | 79.8 | 164.3 | 302.5 |
| 2001..... | 27.8 | * | * | * | 0.3 | 1.4 | 3.6 | 10.9 | 31.6 | 79.1 | 164.6 | 304.7 |
| 2000..... | 27.6 | * | * | * | 0.4 | 1.4 | 3.5 | 11.0 | 32.4 | 80.1 | 161.8 | 302.8 |
| 1999..... | 27.3 | * | * | * | 0.4 | 1.4 | 3.6 | 10.4 | 32.8 | 81.4 | 159.6 | 298.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 26.1 | * | * | * | 0.3 | 1.4 | 4.1 | 11.3 | 30.7 | 73.7 | 157.8 | 284.5 |
| 2003..... | 26.6 | * | * | * | 0.3 | 1.5 | 4.1 | 11.3 | 31.4 | 76.8 | 161.6 | 293.8 |
| 2002..... | 26.4 | * | * | * | 0.4 | 1.5 | 4.3 | 11.3 | 30.6 | 76.8 | 163.3 | 296.4 |
| 2001..... | 26.0 | * | * | * | 0.3 | 1.3 | 3.9 | 11.3 | 30.7 | 77.0 | 162.7 | 297.4 |
| 2000..... | 25.4 | * | * | * | 0.4 | 1.4 | 3.9 | 10.7 | 30.9 | 76.1 | 159.9 | 295.5 |
| 1999..... | 25.0 | * | * | * | 0.4 | 1.4 | 3.8 | 10.5 | 30.1 | 76.8 | 159.1 | 290.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 26.3 | * | * | * | 0.3 | 1.6 | 5.3 | 14.3 | 37.3 | 87.7 | 187.1 | 321.7 |
| 2003..... | 26.6 | * | * | * | 0.4 | 1.7 | 5.2 | 14.1 | 37.9 | 91.3 | 191.7 | 327.3 |
| 2002..... | 26.0 | * | * | * | 0.4 | 1.9 | 5.5 | 13.8 | 37.5 | 88.9 | 187.4 | 338.4 |
| 2001..... | 25.1 | * | * | * | 0.3 | 1.5 | 4.8 | 13.7 | 36.4 | 89.8 | 183.5 | 332.1 |
| 2000..... | 24.3 | * | * | * | 0.5 | 1.7 | 4.8 | 13.0 | 35.4 | 86.6 | 184.5 | 330.4 |
| 1999..... | 24.0 | * | * | * | 0.4 | 1.5 | 4.6 | 13.0 | 35.0 | 87.7 | 184.6 | 326.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 25.9 | * | * | * | 0.3 | 1.2 | 3.0 | 8.3 | 24.3 | 61.7 | 137.8 | 267.8 |
| 2003..... | 26.5 | * | * | * | 0.3 | 1.3 | 2.9 | 8.5 | 25.2 | 64.4 | 141.3 | 279.0 |
| 2002..... | 26.8 | * | * | * | 0.4 | 1.2 | 3.2 | 8.8 | 24.1 | 66.5 | 147.0 | 278.3 |
| 2001..... | 26.9 | * | * | * | 0.2 | 1.2 | 2.9 | 8.8 | 25.3 | 66.0 | 148.9 | 283.0 |
| 2000..... | 26.5 | * | * | * | 0.3 | 1.1 | 3.1 | 8.6 | 26.6 | 67.3 | 143.8 | 281.4 |
| 1999..... | 26.0 | * | * | * | 0.3 | 1.2 | 3.0 | 8.1 | 25.4 | 67.7 | 142.4 | 275.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diabetes mellitus (E10-E14) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 34.4 | * | * | * | 1.0 | 3.4 | 9.8 | 28.8 | 81.9 | 182.6 | 336.4 | 536.5 |
| 2003..... | 35.0 | * | * | * | 0.7 | 4.0 | 9.9 | 30.8 | 87.5 | 188.9 | 340.0 | 532.2 |
| 2002..... | 34.8 | * | * | * | 1.0 | 3.8 | 9.9 | 30.9 | 85.9 | 191.5 | 340.2 | 544.7 |
| 2001..... | 34.1 | * | * | * | 1.0 | 3.8 | 9.1 | 30.1 | 86.5 | 190.8 | 337.9 | 533.8 |
| 2000..... | 34.0 | * | * | * | 1.0 | 3.6 | 8.2 | 31.0 | 83.9 | 190.6 | 354.8 | 530.1 |
| 1999..... | 34.1 | * | * | * | 0.8 | 3.1 | 9.1 | 31.0 | 92.0 | 194.0 | 336.1 | 530.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 31.3 | * | * | * | 1.0 | 4.0 | 12.2 | 34.0 | 93.4 | 201.9 | 344.3 | 508.1 |
| 2003..... | 30.9 | * | * | * | 0.9 | 4.9 | 11.9 | 34.6 | 97.8 | 201.2 | 342.8 | 457.1 |
| 2002..... | 30.0 | * | * | * | 1.1 | 4.6 | 12.1 | 35.8 | 96.6 | 193.3 | 332.6 | 423.4 |
| 2001..... | 29.4 | * | * | * | 0.9 | 5.0 | 10.5 | 35.0 | 93.8 | 189.5 | 345.1 | 452.7 |
| 2000..... | 28.3 | * | * | * | 1.0 | 4.0 | 9.7 | 34.2 | 89.7 | 186.9 | 339.4 | 456.8 |
| 1999..... | 28.6 | * | * | * | * | 3.4 | 10.7 | 35.3 | 95.4 | 189.7 | 317.7 | 506.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 37.2 | * | * | * | 1.0 | 2.9 | 7.8 | 24.3 | 72.6 | 168.7 | 332.0 | 547.7 |
| 2003..... | 38.7 | * | * | * | * | 3.2 | 8.1 | 27.6 | 79.2 | 180.2 | 338.5 | 561.3 |
| 2002..... | 39.2 | * | * | * | 0.8 | 3.0 | 8.0 | 26.8 | 77.3 | 190.3 | 344.5 | 591.1 |
| 2001..... | 38.4 | * | * | * | 1.0 | 2.8 | 7.8 | 25.9 | 80.6 | 191.7 | 333.9 | 563.8 |
| 2000..... | 39.1 | * | * | * | 1.0 | 3.3 | 7.0 | 28.2 | 79.1 | 193.3 | 363.5 | 557.3 |
| 1999..... | 39.0 | * | * | * | 0.9 | 2.7 | 7.7 | 27.2 | 89.2 | 197.0 | 346.5 | 540.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of pituitary gland (E22-E23) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.5 | 1.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.4 | 1.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 1.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.3 | 1.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of adrenal glands (E24-E27) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.8 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.7 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.5 | 0.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.5 | 1.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.5 | 0.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 1.2 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 1.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | * | 0.7 | 1.2 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.5 | 1.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other endocrine diseases (E15-E21,E28-E31,E34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.0 | 2.1 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 |
| 2002..... | 0.2 | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.2 |
| 2001..... | 0.2 | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 2.5 |
| 2000..... | 0.2 | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.9 | 2.2 |
| 1999..... | 0.2 | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 1.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.6 |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 1.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.3 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | * | * | 0.9 | 1.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.2 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.4 | 0.9 | 2.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 2.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.5 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.8 | 2.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.9 |
| 2002..... | 0.2 | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.7 | 2.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.7 | 2.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 1.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.8 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.2 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.4 | 1.0 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | * | 0.8 | 2.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nutritional diseases (E40-E68) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | 0.1 | 0.8 | 1.5 | 2.9 | 4.0 | 5.3 | 9.9 | 32.3 |
| 2003..... | 2.5 | * | * | * | 0.2 | 0.7 | 1.5 | 2.8 | 3.9 | 5.4 | 10.2 | 37.5 |
| 2002..... | 2.6 | * | * | * | 0.2 | 0.8 | 1.6 | 2.6 | 3.9 | 4.9 | 11.3 | 42.9 |
| 2001..... | 2.4 | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 3.3 | 4.6 | 10.8 | 44.5 |
| 2000..... | 2.6 | * | * | * | 0.1 | 0.6 | 1.3 | 2.3 | 3.2 | 4.9 | 12.6 | 51.7 |
| 1999..... | 2.5 | * | * | * | 0.2 | 0.5 | 1.2 | 2.0 | 2.9 | 4.7 | 12.3 | 55.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | 0.2 | 0.8 | 1.6 | 3.0 | 3.9 | 4.9 | 9.5 | 28.8 |
| 2003..... | 2.1 | * | * | * | 0.2 | 0.9 | 1.5 | 2.8 | 3.9 | 5.1 | 9.7 | 31.5 |
| 2002..... | 2.2 | * | * | * | 0.2 | 0.9 | 1.8 | 2.6 | 3.7 | 4.5 | 10.3 | 37.8 |
| 2001..... | 1.9 | * | * | * | 0.2 | 0.7 | 1.4 | 2.4 | 3.1 | 4.3 | 9.8 | 35.8 |
| 2000..... | 2.0 | * | * | * | 0.2 | 0.7 | 1.5 | 2.4 | 3.2 | 4.6 | 11.2 | 44.6 |
| 1999..... | 2.0 | * | * | * | 0.2 | 0.6 | 1.4 | 2.0 | 3.1 | 4.6 | 11.6 | 48.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 0.7 | 1.4 | 2.9 | 4.0 | 5.6 | 10.1 | 33.9 |
| 2003..... | 2.9 | * | * | * | 0.2 | 0.5 | 1.5 | 2.8 | 3.9 | 5.6 | 10.6 | 40.1 |
| 2002..... | 3.0 | * | * | * | 0.2 | 0.6 | 1.4 | 2.6 | 4.0 | 5.1 | 12.0 | 45.1 |
| 2001..... | 2.9 | * | * | * | 0.2 | 0.5 | 1.2 | 2.5 | 3.5 | 4.9 | 11.4 | 48.1 |
| 2000..... | 3.1 | * | * | * | 0.1 | 0.4 | 1.2 | 2.2 | 3.2 | 5.2 | 13.6 | 54.6 |
| 1999..... | 2.9 | * | * | * | 0.2 | 0.4 | 1.0 | 2.0 | 2.6 | 4.7 | 12.8 | 58.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | 0.4 | 1.0 | 1.2 | 2.1 | 2.7 | 7.3 | 21.4 |
| 2003..... | 0.9 | * | * | * | * | 0.5 | 0.9 | 1.5 | 2.0 | 3.3 | 7.5 | 31.6 |
| 2002..... | 0.9 | * | * | * | * | 0.4 | 0.9 | 1.4 | 1.7 | 3.5 | 9.0 | 29.9 |
| 2001..... | 0.8 | * | * | * | * | 0.5 | 0.9 | 1.2 | 1.6 | 2.4 | 9.6 | 41.9 |
| 2000..... | 0.8 | * | * | * | * | 0.4 | 0.9 | 1.4 | 1.3 | 4.0 | 9.7 | 41.8 |
| 1999..... | 0.8 | * | * | * | * | 0.3 | 0.8 | 1.0 | 1.9 | 2.5 | 7.5 | 52.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.5 | 1.1 | 1.5 | 1.9 | * | * | * |
| 2003..... | 0.8 | * | * | * | * | 0.6 | 0.9 | 1.4 | * | * | * | * |
| 2002..... | 0.7 | * | * | * | * | * | 1.0 | 1.1 | * | * | * | * |
| 2001..... | 0.7 | * | * | * | * | 0.6 | 1.0 | * | * | * | 10.3 | 45.7 |
| 2000..... | 0.7 | * | * | * | * | 0.6 | 1.0 | 1.4 | * | * | * | * |
| 1999..... | 0.7 | * | * | * | * | * | 1.2 | * | * | * | * | 51.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.9 | 1.0 | 2.3 | * | 8.6 | 24.5 |
| 2003..... | 1.0 | * | * | * | * | * | 0.8 | 1.6 | 2.1 | 3.4 | 8.7 | 33.6 |
| 2002..... | 1.0 | * | * | * | * | * | 0.8 | 1.7 | 2.0 | 3.5 | 9.8 | 29.8 |
| 2001..... | 0.9 | * | * | * | * | * | 0.8 | 1.3 | * | * | 9.1 | 40.0 |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.4 | * | 4.3 | 10.6 | 44.5 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | * | 7.6 | 53.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nutritional diseases (E40-E68) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 0.1 | 0.8 | 1.5 | 3.1 | 4.1 | 5.5 | 10.0 | 32.8 |
| 2003..... | 2.8 | * | * | * | 0.2 | 0.8 | 1.6 | 2.9 | 4.0 | 5.5 | 10.3 | 37.7 |
| 2002..... | 2.9 | * | * | * | 0.2 | 0.8 | 1.7 | 2.7 | 4.0 | 4.9 | 11.4 | 43.3 |
| 2001..... | 2.6 | * | * | * | 0.2 | 0.6 | 1.3 | 2.5 | 3.4 | 4.8 | 10.8 | 44.5 |
| 2000..... | 2.8 | * | * | * | 0.1 | 0.6 | 1.4 | 2.4 | 3.4 | 5.0 | 12.7 | 51.9 |
| 1999..... | 2.7 | * | * | * | 0.2 | 0.5 | 1.3 | 2.1 | 2.9 | 4.8 | 12.5 | 55.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | 0.2 | 0.8 | 1.6 | 3.1 | 4.0 | 5.0 | 9.7 | 29.4 |
| 2003..... | 2.4 | * | * | * | 0.2 | 1.0 | 1.6 | 2.9 | 4.1 | 5.2 | 9.8 | 31.6 |
| 2002..... | 2.4 | * | * | * | 0.2 | 0.9 | 1.9 | 2.8 | 3.9 | 4.6 | 10.3 | 38.0 |
| 2001..... | 2.0 | * | * | * | 0.2 | 0.7 | 1.4 | 2.5 | 3.2 | 4.4 | 9.8 | 35.4 |
| 2000..... | 2.2 | * | * | * | 0.2 | 0.7 | 1.5 | 2.5 | 3.4 | 4.7 | 11.2 | 44.8 |
| 1999..... | 2.2 | * | * | * | 0.2 | 0.7 | 1.4 | 2.1 | 3.1 | 4.7 | 11.8 | 48.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | 0.8 | 1.5 | 3.1 | 4.2 | 5.9 | 10.2 | 34.3 |
| 2003..... | 3.2 | * | * | * | 0.2 | 0.6 | 1.6 | 2.9 | 4.0 | 5.8 | 10.7 | 40.3 |
| 2002..... | 3.3 | * | * | * | 0.2 | 0.7 | 1.4 | 2.7 | 4.2 | 5.2 | 12.1 | 45.5 |
| 2001..... | 3.2 | * | * | * | 0.2 | 0.5 | 1.3 | 2.6 | 3.6 | 5.0 | 11.5 | 48.3 |
| 2000..... | 3.3 | * | * | * | * | 0.4 | 1.3 | 2.3 | 3.3 | 5.2 | 13.7 | 54.8 |
| 1999..... | 3.2 | * | * | * | 0.2 | 0.4 | 1.1 | 2.1 | 2.7 | 4.8 | 13.0 | 58.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 0.1 | 0.7 | 1.3 | 2.9 | 4.0 | 5.1 | 9.6 | 31.4 |
| 2003..... | 2.9 | * | * | * | 0.2 | 0.6 | 1.5 | 2.8 | 4.0 | 5.5 | 9.9 | 36.2 |
| 2002..... | 2.9 | * | * | * | 0.1 | 0.6 | 1.5 | 2.6 | 3.8 | 4.6 | 10.8 | 42.0 |
| 2001..... | 2.7 | * | * | * | 0.1 | 0.5 | 1.1 | 2.3 | 3.3 | 4.4 | 10.1 | 43.7 |
| 2000..... | 2.9 | * | * | * | 0.1 | 0.4 | 1.3 | 2.2 | 3.1 | 4.6 | 12.1 | 50.7 |
| 1999..... | 2.7 | * | * | * | 0.1 | 0.4 | 1.1 | 1.9 | 2.7 | 4.3 | 11.7 | 54.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | 0.1 | 0.6 | 1.5 | 3.1 | 4.0 | 4.8 | 9.3 | 28.4 |
| 2003..... | 2.4 | * | * | * | 0.2 | 0.9 | 1.5 | 2.9 | 4.1 | 5.2 | 9.2 | 30.5 |
| 2002..... | 2.4 | * | * | * | * | 0.7 | 1.8 | 2.8 | 3.8 | 4.4 | 9.8 | 37.4 |
| 2001..... | 2.0 | * | * | * | * | 0.6 | 1.2 | 2.3 | 3.1 | 4.1 | 9.2 | 33.9 |
| 2000..... | 2.3 | * | * | * | 0.2 | 0.6 | 1.4 | 2.5 | 3.2 | 4.3 | 10.6 | 43.8 |
| 1999..... | 2.2 | * | * | * | * | 0.5 | 1.3 | 2.1 | 2.9 | 4.2 | 10.8 | 46.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | 0.7 | 1.1 | 2.7 | 3.9 | 5.4 | 9.8 | 32.8 |
| 2003..... | 3.3 | * | * | * | * | 0.4 | 1.4 | 2.7 | 3.8 | 5.7 | 10.4 | 38.7 |
| 2002..... | 3.3 | * | * | * | * | 0.6 | 1.2 | 2.3 | 3.8 | 4.8 | 11.5 | 44.0 |
| 2001..... | 3.3 | * | * | * | * | 0.4 | 1.0 | 2.3 | 3.5 | 4.7 | 10.7 | 47.8 |
| 2000..... | 3.4 | * | * | * | * | 0.3 | 1.1 | 1.9 | 3.1 | 4.9 | 13.1 | 53.5 |
| 1999..... | 3.3 | * | * | * | * | 0.3 | 0.9 | 1.8 | 2.5 | 4.4 | 12.3 | 57.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nutritional diseases (E40-E68) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | 0.3 | 2.0 | 3.3 | 5.3 | 6.8 | 10.0 | 16.1 | 51.2 |
| 2003..... | 3.1 | * | * | * | 0.4 | 1.8 | 2.9 | 4.4 | 6.2 | 7.6 | 17.3 | 58.9 |
| 2002..... | 3.4 | * | * | * | 0.5 | 2.0 | 2.9 | 4.5 | 7.5 | 8.9 | 20.0 | 61.7 |
| 2001..... | 3.2 | * | * | * | 0.5 | 1.3 | 2.9 | 4.9 | 5.6 | 8.7 | 20.1 | 56.8 |
| 2000..... | 3.2 | * | * | * | 0.4 | 1.6 | 2.6 | 4.4 | 6.0 | 9.0 | 20.8 | 69.0 |
| 1999..... | 3.1 | * | * | * | 0.4 | 1.3 | 2.4 | 3.6 | 5.5 | 9.9 | 23.3 | 72.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | 2.1 | 3.1 | 4.1 | 5.6 | 8.7 | 16.5 | 45.7 |
| 2003..... | 2.5 | * | * | * | * | 2.1 | 2.4 | 4.0 | 4.9 | 7.4 | 19.9 | 49.0 |
| 2002..... | 2.7 | * | * | * | * | 2.4 | 3.1 | 3.3 | 6.5 | 7.8 | 19.0 | 47.0 |
| 2001..... | 2.6 | * | * | * | * | 1.3 | 3.0 | 4.2 | 5.1 | 8.3 | 18.3 | 58.9 |
| 2000..... | 2.6 | * | * | * | * | 1.8 | 2.6 | 3.4 | 6.2 | 9.1 | 20.7 | 66.9 |
| 1999..... | 2.7 | * | * | * | * | 1.6 | 2.4 | 2.9 | 5.9 | 10.2 | 24.7 | 76.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 1.9 | 3.5 | 6.3 | 7.8 | 11.0 | 15.8 | 53.4 |
| 2003..... | 3.7 | * | * | * | * | 1.5 | 3.2 | 4.8 | 7.3 | 7.7 | 15.8 | 62.7 |
| 2002..... | 4.1 | * | * | * | * | 1.7 | 2.8 | 5.6 | 8.4 | 9.7 | 20.6 | 67.3 |
| 2001..... | 3.7 | * | * | * | * | 1.3 | 2.9 | 5.5 | 6.1 | 9.0 | 21.1 | 56.0 |
| 2000..... | 3.7 | * | * | * | * | 1.4 | 2.6 | 5.2 | 5.8 | 9.0 | 20.8 | 69.8 |
| 1999..... | 3.6 | * | * | * | * | 1.1 | 2.4 | 4.2 | 5.1 | 9.6 | 22.6 | 71.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malnutrition (E40-E46) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.6 | 6.8 | 28.6 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 7.5 | 33.7 |
| 2002..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 8.3 | 38.7 |
| 2001..... | 1.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.0 | 8.2 | 40.2 |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.3 | 9.7 | 46.5 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 9.6 | 50.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 6.7 | 25.8 |
| 2003..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 7.5 | 28.4 |
| 2002..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.1 | 8.0 | 35.3 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.3 | 0.6 | 2.0 | 8.0 | 33.2 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 8.8 | 40.9 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 9.5 | 44.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.5 | 6.8 | 29.9 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 7.5 | 36.0 |
| 2002..... | 1.7 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.8 | 8.6 | 40.1 |
| 2001..... | 1.7 | * | * | * | * | * | * | 0.3 | 0.6 | 1.9 | 8.3 | 43.2 |
| 2000..... | 2.0 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.2 | 10.3 | 48.7 |
| 1999..... | 2.0 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 2.0 | 9.7 | 52.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 6.4 | 27.3 |
| 2003..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 7.2 | 32.3 |
| 2002..... | 1.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 8.0 | 37.4 |
| 2001..... | 1.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 7.6 | 39.4 |
| 2000..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 9.3 | 45.3 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.9 | 8.9 | 48.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 6.2 | 24.6 |
| 2003..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.4 | 1.6 | 6.9 | 27.3 |
| 2002..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.5 | 34.5 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 7.4 | 31.7 |
| 2000..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 8.3 | 39.8 |
| 1999..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 8.6 | 42.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 6.6 | 28.5 |
| 2003..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 7.4 | 34.5 |
| 2002..... | 1.8 | * | * | * | * | * | * | 0.3 | 0.5 | 1.7 | 8.3 | 38.6 |
| 2001..... | 1.9 | * | * | * | * | * | * | 0.3 | 0.6 | 1.9 | 7.7 | 42.7 |
| 2000..... | 2.2 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.0 | 10.0 | 47.4 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.8 | 9.2 | 51.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Malnutrition (E40-E46) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 0.6 | 1.1 | 3.3 | 10.5 | 46.3 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.7 | 1.3 | 3.1 | 13.0 | 54.1 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | 1.7 | 3.6 | 14.2 | 57.2 |
| 2001..... | 1.2 | * | * | * | * | * | * | 0.6 | 1.1 | 3.4 | 15.0 | 52.1 |
| 2000..... | 1.4 | * | * | * | * | * | * | 0.6 | 1.6 | 4.8 | 15.1 | 64.2 |
| 1999..... | 1.6 | * | * | * | * | * | * | 0.6 | 1.8 | 5.5 | 18.1 | 66.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | 1.6 | 4.4 | 12.0 | 43.7 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | 1.8 | 4.3 | 17.3 | 46.9 |
| 2002..... | 1.0 | * | * | * | * | * | * | * | 2.7 | 4.0 | 15.2 | 44.9 |
| 2001..... | 1.1 | * | * | * | * | * | * | 1.0 | * | 5.0 | 15.9 | 56.6 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 2.5 | 6.4 | 17.3 | 62.2 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | 2.6 | 8.0 | 20.6 | 74.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | * | * | 2.5 | 9.7 | 47.4 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | * | 2.3 | 10.6 | 57.0 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | * | 3.4 | 13.6 | 61.9 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | * | 2.3 | 14.5 | 50.5 |
| 2000..... | 1.6 | * | * | * | * | * | * | * | * | 3.6 | 13.9 | 65.0 |
| 1999..... | 1.7 | * | * | * | * | * | * | * | * | 3.7 | 16.6 | 62.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nutritional marasmus (E41) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.3 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.5 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.5 | 2.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.3 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified malnutrition (E40,E42-E46) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.5 | 6.5 | 27.4 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 7.1 | 32.1 |
| 2002..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 8.0 | 36.3 |
| 2001..... | 1.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 | 7.7 | 37.8 |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.2 | 9.2 | 43.7 |
| 1999..... | 1.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.1 | 9.1 | 47.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 6.4 | 24.8 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 7.2 | 27.2 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.0 | 7.6 | 33.8 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 7.5 | 31.1 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 8.4 | 38.1 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 9.0 | 41.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 6.5 | 28.5 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 7.1 | 34.2 |
| 2002..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.7 | 8.2 | 37.4 |
| 2001..... | 1.7 | * | * | * | * | * | * | 0.3 | 0.6 | 1.8 | 7.9 | 40.6 |
| 2000..... | 1.9 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.1 | 9.8 | 46.0 |
| 1999..... | 1.9 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.9 | 9.2 | 50.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 6.1 | 26.1 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 6.8 | 30.7 |
| 2002..... | 1.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.7 | 7.6 | 35.0 |
| 2001..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 7.2 | 37.0 |
| 2000..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 2.0 | 8.8 | 42.5 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 8.5 | 46.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 5.9 | 23.7 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.4 | 1.6 | 6.5 | 26.1 |
| 2002..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.2 | 33.0 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 6.9 | 29.7 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 | 7.9 | 37.1 |
| 1999..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 8.1 | 39.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 6.3 | 27.1 |
| 2003..... | 1.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 7.0 | 32.7 |
| 2002..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 7.9 | 35.9 |
| 2001..... | 1.8 | * | * | * | * | * | * | 0.3 | 0.6 | 1.8 | 7.4 | 40.0 |
| 2000..... | 2.0 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.9 | 9.5 | 44.7 |
| 1999..... | 2.1 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.7 | 8.7 | 49.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified malnutrition (E40,E42-E46) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 0.6 | 1.1 | 3.1 | 10.5 | 45.2 |
| 2003..... | 1.1 | * | * | * | * | * | * | 0.6 | 1.3 | 2.9 | 12.2 | 53.2 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | 1.6 | 3.5 | 13.2 | 55.4 |
| 2001..... | 1.2 | * | * | * | * | * | * | 0.6 | 0.9 | 3.3 | 14.5 | 49.6 |
| 2000..... | 1.4 | * | * | * | * | * | * | 0.5 | 1.5 | 4.6 | 14.8 | 60.1 |
| 1999..... | 1.5 | * | * | * | * | * | * | 0.6 | 1.7 | 5.2 | 17.1 | 62.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | * | 4.3 | 12.0 | 42.7 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | 1.8 | 4.0 | 16.4 | 45.8 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | 2.5 | 4.0 | 14.1 | 43.8 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | * | 4.9 | 15.6 | 54.3 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | 2.4 | 6.1 | 16.4 | 56.4 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | 2.4 | 7.7 | 20.0 | 70.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | * | * | 2.3 | 9.7 | 46.2 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | * | 2.2 | 9.8 | 56.1 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | * | 3.2 | 12.8 | 59.8 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | * | 2.2 | 13.8 | 47.9 |
| 2000..... | 1.6 | * | * | * | * | * | * | * | * | 3.5 | 13.9 | 61.5 |
| 1999..... | 1.6 | * | * | * | * | * | * | * | * | 3.4 | 15.4 | 60.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other nutritional deficiencies (E50-E64) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | | 2.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.3 | 2.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 3.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.7 | 3.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 2.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 1.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 3.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 3.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.1 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.6 | 4.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 4.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 2.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.6 | 3.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.7 | 3.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 3.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 4.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 3.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 3.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 4.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.6 | 3.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Obesity and other hyperalimentation (E65-E68) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.1 | 0.8 | 1.4 | 2.8 | 3.5 | 3.8 | 3.0 | 2.0 |
| 2003..... | 1.5 | * | * | * | 0.2 | 0.8 | 1.5 | 2.7 | 3.4 | 3.7 | 2.6 | 1.9 |
| 2002..... | 1.4 | * | * | * | 0.2 | 0.8 | 1.5 | 2.4 | 3.4 | 2.8 | 2.5 | 1.4 |
| 2001..... | 1.2 | * | * | * | 0.2 | 0.6 | 1.2 | 2.2 | 2.8 | 2.7 | 2.0 | 1.5 |
| 2000..... | 1.1 | * | * | * | 0.1 | 0.5 | 1.3 | 2.1 | 2.6 | 2.5 | 2.3 | 1.7 |
| 1999..... | 1.0 | * | * | * | 0.1 | 0.5 | 1.1 | 1.8 | 2.2 | 2.4 | 2.2 | 1.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 0.2 | 0.8 | 1.6 | 2.9 | 3.4 | 3.3 | 2.8 | 1.4 |
| 2003..... | 1.4 | * | * | * | 0.2 | 1.0 | 1.5 | 2.7 | 3.5 | 3.2 | 2.0 | * |
| 2002..... | 1.3 | * | * | * | 0.2 | 0.9 | 1.8 | 2.5 | 3.1 | 2.4 | 1.8 | * |
| 2001..... | 1.1 | * | * | * | 0.1 | 0.7 | 1.3 | 2.2 | 2.5 | 2.2 | 1.3 | * |
| 2000..... | 1.1 | * | * | * | 0.2 | 0.7 | 1.4 | 2.3 | 2.5 | 2.0 | 1.6 | * |
| 1999..... | 1.0 | * | * | * | 0.1 | 0.6 | 1.3 | 1.8 | 2.2 | 2.0 | 1.4 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.8 | 1.3 | 2.7 | 3.7 | 4.3 | 3.1 | 2.2 |
| 2003..... | 1.6 | * | * | * | 0.2 | 0.6 | 1.5 | 2.6 | 3.4 | 4.0 | 2.9 | 2.4 |
| 2002..... | 1.5 | * | * | * | 0.2 | 0.7 | 1.3 | 2.4 | 3.6 | 3.2 | 3.0 | 1.7 |
| 2001..... | 1.3 | * | * | * | 0.2 | 0.5 | 1.2 | 2.3 | 3.0 | 3.0 | 2.5 | 1.9 |
| 2000..... | 1.2 | * | * | * | * | 0.4 | 1.1 | 2.0 | 2.7 | 2.8 | 2.7 | 2.0 |
| 1999..... | 1.1 | * | * | * | 0.2 | 0.4 | 0.9 | 1.7 | 2.1 | 2.7 | 2.6 | 1.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 0.1 | 0.6 | 1.2 | 2.6 | 3.5 | 3.7 | 2.9 | 1.9 |
| 2003..... | 1.5 | * | * | * | 0.1 | 0.6 | 1.4 | 2.6 | 3.4 | 3.7 | 2.5 | 1.8 |
| 2002..... | 1.4 | * | * | * | 0.1 | 0.6 | 1.4 | 2.3 | 3.2 | 2.8 | 2.3 | 1.5 |
| 2001..... | 1.1 | * | * | * | 0.1 | 0.5 | 1.1 | 2.1 | 2.7 | 2.5 | 1.9 | 1.5 |
| 2000..... | 1.1 | * | * | * | 0.1 | 0.4 | 1.1 | 2.0 | 2.5 | 2.4 | 2.2 | 1.7 |
| 1999..... | 1.0 | * | * | * | 0.1 | 0.4 | 1.0 | 1.7 | 2.1 | 2.3 | 2.1 | 1.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.6 | 1.5 | 2.9 | 3.5 | 3.3 | 2.7 | 1.5 |
| 2003..... | 1.5 | * | * | * | 0.2 | 0.8 | 1.4 | 2.7 | 3.7 | 3.4 | 2.0 | * |
| 2002..... | 1.4 | * | * | * | * | 0.7 | 1.7 | 2.6 | 3.2 | 2.4 | 1.8 | * |
| 2001..... | 1.1 | * | * | * | * | 0.6 | 1.2 | 2.1 | 2.6 | 2.2 | 1.3 | * |
| 2000..... | 1.1 | * | * | * | * | 0.6 | 1.4 | 2.3 | 2.5 | 2.0 | 1.6 | * |
| 1999..... | 1.0 | * | * | * | * | 0.5 | 1.2 | 1.8 | 2.2 | 2.1 | 1.4 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.6 | 1.0 | 2.4 | 3.4 | 4.0 | 2.9 | 2.0 |
| 2003..... | 1.5 | * | * | * | * | 0.4 | 1.3 | 2.5 | 3.1 | 4.0 | 2.8 | 2.2 |
| 2002..... | 1.3 | * | * | * | * | 0.5 | 1.1 | 2.0 | 3.2 | 3.0 | 2.6 | 1.7 |
| 2001..... | 1.2 | * | * | * | * | 0.4 | 1.0 | 2.0 | 3.2 | 2.7 | 2.4 | 1.9 |
| 2000..... | 1.1 | * | * | * | * | 0.3 | 0.9 | 1.7 | 2.5 | 2.6 | 2.6 | 2.1 |
| 1999..... | 1.0 | * | * | * | * | 0.3 | 0.7 | 1.6 | 2.0 | 2.5 | 2.5 | 1.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Obesity and other hyperalimentation (E65-E68) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 1.9 | 3.1 | 4.6 | 5.6 | 6.6 | 5.4 | * |
| 2003..... | 1.9 | * | * | * | 0.3 | 1.8 | 2.8 | 3.7 | 4.8 | 4.3 | 4.1 | * |
| 2002..... | 2.0 | * | * | * | 0.5 | 2.0 | 2.6 | 4.1 | 5.7 | 4.6 | 5.3 | * |
| 2001..... | 1.8 | * | * | * | 0.5 | 1.2 | 2.7 | 4.2 | 4.5 | 5.0 | 4.0 | * |
| 2000..... | 1.7 | * | * | * | * | 1.4 | 2.3 | 3.8 | 4.2 | 4.2 | 4.4 | * |
| 1999..... | 1.5 | * | * | * | 0.4 | 1.3 | 2.1 | 2.9 | 3.4 | 4.3 | 3.7 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 2.0 | 2.8 | 3.4 | 3.9 | 4.1 | * | * |
| 2003..... | 1.5 | * | * | * | * | 2.1 | 2.4 | 3.2 | 3.0 | 2.9 | * | * |
| 2002..... | 1.6 | * | * | * | * | 2.3 | 2.8 | 2.6 | 3.6 | 3.2 | * | * |
| 2001..... | 1.4 | * | * | * | * | 1.3 | 2.6 | 3.0 | 3.2 | * | * | * |
| 2000..... | 1.4 | * | * | * | * | 1.6 | 2.4 | 2.7 | 3.4 | * | * | * |
| 1999..... | 1.2 | * | * | * | * | 1.5 | 2.1 | 2.2 | 3.0 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 1.8 | 3.4 | 5.7 | 7.0 | 8.3 | 6.0 | * |
| 2003..... | 2.3 | * | * | * | * | 1.5 | 3.2 | 4.2 | 6.3 | 5.3 | 5.2 | * |
| 2002..... | 2.5 | * | * | * | * | 1.6 | 2.5 | 5.3 | 7.4 | 5.6 | 6.8 | * |
| 2001..... | 2.3 | * | * | * | * | 1.1 | 2.7 | 5.2 | 5.5 | 6.6 | 5.2 | * |
| 2000..... | 2.0 | * | * | * | * | 1.2 | 2.3 | 4.7 | 4.9 | 5.4 | 5.4 | * |
| 1999..... | 1.7 | * | * | * | * | 1.1 | 2.2 | 3.5 | 3.8 | 5.9 | 4.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Metabolic disorders (E70-E88) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.4 | 4.9 | 0.8 | 0.4 | 0.6 | 0.6 | 0.8 | 2.1 | 5.3 | 13.3 | 34.6 | 107.9 |
| 2003..... | 5.5 | 5.2 | 1.1 | 0.4 | 0.7 | 0.6 | 0.9 | 2.2 | 5.2 | 13.1 | 34.6 | 112.6 |
| 2002..... | 5.5 | 5.5 | 0.8 | 0.4 | 0.6 | 0.6 | 0.8 | 2.0 | 5.6 | 13.0 | 35.2 | 114.3 |
| 2001..... | 5.5 | 6.0 | 1.0 | 0.4 | 0.6 | 0.6 | 0.8 | 2.1 | 5.0 | 13.1 | 34.1 | 124.4 |
| 2000..... | 5.4 | 5.3 | 0.9 | 0.4 | 0.7 | 0.6 | 0.8 | 2.0 | 5.1 | 12.6 | 33.9 | 124.1 |
| 1999..... | 5.3 | 5.6 | 0.8 | 0.4 | 0.7 | 0.6 | 0.8 | 1.8 | 4.7 | 12.2 | 33.1 | 130.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.1 | 6.0 | 0.8 | 0.4 | 0.6 | 0.6 | 0.9 | 3.0 | 7.0 | 16.9 | 38.5 | 103.1 |
| 2003..... | 5.1 | 5.8 | 1.1 | 0.5 | 0.7 | 0.6 | 1.1 | 3.0 | 6.9 | 16.8 | 38.4 | 104.8 |
| 2002..... | 4.9 | 6.1 | 0.8 | 0.5 | 0.6 | 0.7 | 1.0 | 2.8 | 7.3 | 16.0 | 38.8 | 101.5 |
| 2001..... | 4.9 | 6.5 | 1.1 | 0.4 | 0.6 | 0.6 | 1.0 | 2.8 | 6.8 | 16.7 | 38.6 | 109.8 |
| 2000..... | 4.8 | 5.5 | 0.9 | 0.5 | 0.7 | 0.7 | 0.9 | 2.9 | 6.8 | 15.8 | 36.7 | 112.0 |
| 1999..... | 4.5 | 6.3 | 0.9 | 0.4 | 0.7 | 0.6 | 0.9 | 2.4 | 6.1 | 15.1 | 35.8 | 110.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.8 | 3.8 | 0.7 | 0.4 | 0.6 | 0.5 | 0.7 | 1.3 | 3.7 | 10.3 | 31.9 | 110.1 |
| 2003..... | 5.9 | 4.5 | 1.0 | 0.3 | 0.6 | 0.5 | 0.7 | 1.5 | 3.6 | 10.0 | 32.1 | 116.1 |
| 2002..... | 6.0 | 4.9 | 0.9 | 0.4 | 0.6 | 0.5 | 0.7 | 1.3 | 4.0 | 10.5 | 32.8 | 119.9 |
| 2001..... | 6.1 | 5.5 | 1.0 | 0.4 | 0.7 | 0.5 | 0.6 | 1.5 | 3.4 | 10.2 | 31.1 | 130.6 |
| 2000..... | 6.0 | 5.0 | 0.8 | 0.3 | 0.7 | 0.5 | 0.7 | 1.1 | 3.5 | 10.0 | 32.0 | 129.1 |
| 1999..... | 6.1 | 4.9 | 0.7 | 0.4 | 0.8 | 0.6 | 0.6 | 1.2 | 3.5 | 9.8 | 31.3 | 139.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | 5.1 | 0.9 | 0.3 | * | * | * | 0.9 | 2.8 | 7.6 | 21.3 | 68.5 |
| 2003..... | 1.6 | 5.4 | 1.0 | 0.4 | * | * | 0.3 | 1.1 | 3.3 | 7.2 | 21.7 | 63.2 |
| 2002..... | 1.6 | 4.4 | 0.7 | 0.4 | * | * | 0.3 | 1.1 | 2.4 | 8.6 | 26.0 | 65.4 |
| 2001..... | 1.5 | 4.5 | 0.8 | 0.3 | * | * | 0.4 | 0.9 | 3.4 | 8.4 | 22.4 | 67.5 |
| 2000..... | 1.4 | 4.2 | 0.7 | 0.4 | * | * | * | 1.1 | 2.9 | 8.0 | 21.3 | 73.0 |
| 1999..... | 1.4 | 4.9 | * | * | * | * | * | 1.0 | 2.5 | 7.7 | 23.1 | 80.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.5 | 5.8 | * | * | * | * | * | 1.2 | 3.5 | 9.5 | 24.0 | 61.6 |
| 2003..... | 1.5 | 5.7 | * | * | * | * | * | 1.3 | 4.5 | 8.1 | 23.0 | 64.7 |
| 2002..... | 1.5 | * | * | 0.6 | * | * | * | 1.3 | 2.4 | 10.7 | 29.5 | 71.9 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 4.6 | 10.0 | 25.5 | 76.2 |
| 2000..... | 1.3 | * | * | * | * | * | * | 1.4 | 3.7 | 11.2 | 20.7 | 60.5 |
| 1999..... | 1.3 | 6.2 | * | * | * | * | * | * | 3.1 | 9.4 | 28.2 | 60.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | * | 2.1 | 6.1 | 19.5 | 72.0 |
| 2003..... | 1.6 | 5.2 | * | * | * | * | * | * | 2.1 | 6.5 | 20.8 | 62.5 |
| 2002..... | 1.6 | 5.2 | * | * | * | * | * | * | 2.4 | 7.0 | 23.6 | 62.2 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 2.2 | 7.1 | 20.2 | 63.3 |
| 2000..... | 1.6 | 5.3 | * | * | * | * | * | * | 2.2 | 5.5 | 21.8 | 79.1 |
| 1999..... | 1.6 | * | * | * | * | * | * | * | * | 6.4 | 19.8 | 89.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex; All ages 1/; Under 1 year 2/; 1-4 years; 5-14 years; 15-24 years; 25-34 years; 35-44 years; 45-54 years; 55-64 years; 65-74 years; 75-84 years; 85 years and over. Rows include Metabolic disorders (E70-E88) for Non-Hispanic 4/, both sexes; Non-Hispanic 4/, male; Non-Hispanic 4/, female; Non-Hispanic white, both sexes; Non-Hispanic white, male; Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Metabolic disorders (E70-E88) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.4 | 4.9 | * | * | * | 0.5 | 1.3 | 2.8 | 7.5 | 18.0 | 41.6 | 114.6 |
| 2003..... | 4.7 | 7.9 | 0.8 | * | 0.4 | 0.5 | 1.2 | 3.3 | 7.2 | 18.1 | 46.2 | 126.7 |
| 2002..... | 5.0 | 7.6 | 0.8 | 0.3 | 0.4 | 0.4 | 1.1 | 3.2 | 8.5 | 19.6 | 47.8 | 137.3 |
| 2001..... | 4.7 | 9.3 | * | * | * | 0.4 | 0.7 | 3.3 | 7.1 | 17.4 | 46.7 | 153.6 |
| 2000..... | 4.6 | 6.6 | * | * | 0.4 | 0.7 | 1.1 | 2.8 | 7.4 | 16.7 | 44.4 | 147.2 |
| 1999..... | 4.5 | 9.7 | * | 0.4 | * | 0.5 | 1.4 | 3.0 | 6.5 | 18.0 | 41.3 | 139.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | * | 1.6 | 4.0 | 9.3 | 22.1 | 47.9 | 114.8 |
| 2003..... | 4.3 | 7.1 | * | * | * | * | 1.4 | 4.2 | 9.1 | 23.6 | 55.7 | 100.2 |
| 2002..... | 4.6 | 7.0 | * | * | 0.7 | * | 1.3 | 4.3 | 12.2 | 22.1 | 52.8 | 136.7 |
| 2001..... | 4.4 | 9.6 | * | * | * | * | 0.8 | 4.1 | 10.0 | 23.4 | 56.1 | 132.8 |
| 2000..... | 4.1 | 7.8 | * | * | * | 0.9 | 1.3 | 4.0 | 9.5 | 20.4 | 50.6 | 116.2 |
| 1999..... | 4.2 | 10.7 | * | * | * | * | 1.6 | 3.6 | 7.9 | 22.4 | 48.8 | 148.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 0.9 | 1.8 | 6.2 | 15.1 | 38.0 | 114.4 |
| 2003..... | 5.1 | 8.7 | * | * | * | * | 1.1 | 2.5 | 5.6 | 14.2 | 40.9 | 137.0 |
| 2002..... | 5.3 | 8.2 | * | * | * | * | 0.9 | 2.3 | 5.5 | 17.9 | 44.9 | 137.5 |
| 2001..... | 5.1 | 9.1 | * | * | * | * | * | 2.6 | 4.8 | 13.1 | 41.4 | 161.3 |
| 2000..... | 5.0 | * | * | * | * | * | 0.9 | 1.9 | 5.6 | 14.1 | 41.0 | 158.7 |
| 1999..... | 4.8 | 8.6 | * | * | * | * | 1.2 | 2.5 | 5.4 | 14.9 | 37.1 | 136.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cystic fibrosis (E84) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | 0.1 | 0.5 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| 2003..... | 0.2 | * | * | 0.1 | 0.4 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| 2002..... | 0.2 | * | * | 0.1 | 0.4 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| 2001..... | 0.2 | * | * | 0.1 | 0.5 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| 2000..... | 0.2 | * | * | 0.1 | 0.5 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| 1999..... | 0.2 | * | * | 0.1 | 0.5 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.4 | 0.4 | 0.1 | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.4 | 0.4 | 0.2 | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | 0.4 | 0.5 | 0.2 | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.4 | 0.4 | 0.2 | 0.1 | * | * | * | * |
| 2000..... | 0.2 | * | * | * | 0.4 | 0.4 | 0.2 | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.5 | 0.4 | 0.2 | * | * | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.5 | 0.4 | 0.2 | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.5 | 0.3 | 0.2 | * | * | * | * | * |
| 2002..... | 0.2 | * | * | 0.1 | 0.5 | 0.3 | 0.2 | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.6 | 0.4 | 0.1 | * | * | * | * | * |
| 2000..... | 0.2 | * | * | 0.1 | 0.5 | 0.4 | 0.3 | * | * | * | * | * |
| 1999..... | 0.2 | * | * | 0.2 | 0.6 | 0.4 | 0.1 | * | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | 0.1 | 0.6 | 0.5 | 0.2 | 0.1 | * | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.5 | 0.5 | 0.3 | 0.1 | * | * | * | * |
| 2002..... | 0.2 | * | * | 0.1 | 0.6 | 0.5 | 0.2 | 0.1 | * | * | * | * |
| 2001..... | 0.2 | * | * | 0.1 | 0.6 | 0.5 | 0.2 | 0.1 | * | * | * | * |
| 2000..... | 0.2 | * | * | 0.1 | 0.6 | 0.5 | 0.3 | 0.1 | * | * | * | * |
| 1999..... | 0.2 | * | * | 0.2 | 0.7 | 0.5 | 0.2 | 0.1 | * | * | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.6 | 0.5 | 0.2 | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.5 | 0.5 | 0.3 | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | 0.5 | 0.6 | 0.3 | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.5 | 0.5 | 0.3 | 0.1 | * | * | * | * |
| 2000..... | 0.2 | * | * | * | 0.6 | 0.5 | 0.3 | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.6 | 0.5 | 0.2 | * | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.6 | 0.5 | 0.3 | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.6 | 0.4 | 0.2 | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | 0.7 | 0.4 | 0.2 | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.7 | 0.4 | 0.2 | * | * | * | * | * |
| 2000..... | 0.2 | * | * | 0.2 | 0.7 | 0.5 | 0.3 | * | * | * | * | * |
| 1999..... | 0.2 | * | * | 0.2 | 0.8 | 0.5 | 0.2 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | 1.6 | * | * | 0.0 | 0.1 | 0.2 | 0.5 | 1.0 | 3.2 | 13.7 | 70.7 |
| 2003..... | 2.4 | 1.9 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.1 | 3.4 | 15.1 | 77.3 |
| 2002..... | 2.5 | 1.8 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 3.8 | 16.0 | 83.2 |
| 2001..... | 2.7 | 2.2 | 0.2 | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 4.1 | 16.8 | 93.9 |
| 2000..... | 2.8 | 1.9 | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.3 | 4.0 | 17.4 | 96.8 |
| 1999..... | 2.9 | 1.9 | 0.1 | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 4.2 | 18.2 | 106.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | 1.9 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 3.5 | 13.3 | 59.8 |
| 2003..... | 1.7 | 2.2 | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 3.4 | 15.0 | 65.6 |
| 2002..... | 1.7 | 2.1 | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.8 | 15.3 | 67.9 |
| 2001..... | 1.8 | 2.2 | * | * | * | * | 0.2 | 0.6 | 1.3 | 4.1 | 16.8 | 76.5 |
| 2000..... | 2.0 | 1.8 | * | * | * | * | 0.3 | 0.6 | 1.5 | 4.5 | 17.7 | 84.3 |
| 1999..... | 2.0 | 2.3 | * | * | * | * | 0.3 | 0.7 | 1.2 | 4.5 | 18.4 | 85.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.8 | 1.3 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.9 | 13.9 | 75.7 |
| 2003..... | 3.1 | 1.7 | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.3 | 15.2 | 82.5 |
| 2002..... | 3.3 | 1.5 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.7 | 16.5 | 89.8 |
| 2001..... | 3.6 | 2.2 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 4.1 | 16.8 | 101.2 |
| 2000..... | 3.5 | 2.0 | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.6 | 17.2 | 101.9 |
| 1999..... | 3.8 | 1.5 | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.9 | 18.1 | 115.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 2.3 | 7.6 | 40.4 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 2.0 | 8.4 | 35.2 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 2.5 | 9.7 | 33.3 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 3.1 | 9.0 | 38.8 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 2.1 | 9.7 | 46.4 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 2.1 | 12.3 | 54.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 7.9 | 35.0 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.8 | 38.5 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 10.7 | 33.5 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | 9.8 | 45.7 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | 11.3 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 16.7 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | * | 7.4 | 43.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.2 | 33.6 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | * | 9.0 | 33.2 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 4.3 | 8.5 | 35.4 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 8.6 | 51.4 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | * | 9.4 | 63.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.5 | 1.5 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.3 | 14.0 | 71.9 |
| 2003..... | 2.7 | 2.0 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.1 | 3.5 | 15.5 | 78.8 |
| 2002..... | 2.9 | 2.0 | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.8 | 16.3 | 84.9 |
| 2001..... | 3.1 | 2.4 | 0.2 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 4.1 | 17.1 | 95.8 |
| 2000..... | 3.1 | 2.1 | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 4.1 | 17.7 | 98.4 |
| 1999..... | 3.2 | 2.0 | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 4.3 | 18.4 | 108.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.8 | 1.8 | * | * | * | * | 0.2 | 0.6 | 1.2 | 3.5 | 13.6 | 61.1 |
| 2003..... | 1.9 | 2.4 | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 3.5 | 15.3 | 66.4 |
| 2002..... | 1.9 | 2.6 | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.9 | 15.5 | 69.0 |
| 2001..... | 2.1 | 2.4 | * | * | * | * | 0.3 | 0.6 | 1.3 | 4.2 | 17.1 | 77.4 |
| 2000..... | 2.2 | 2.2 | * | * | * | * | 0.3 | 0.6 | 1.5 | 4.6 | 18.0 | 86.0 |
| 1999..... | 2.2 | 2.3 | * | * | * | * | 0.3 | 0.7 | 1.2 | 4.7 | 18.5 | 86.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.2 | 1.3 | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.0 | 14.2 | 76.8 |
| 2003..... | 3.5 | 1.7 | * | * | * | * | 0.3 | 0.5 | 1.1 | 3.4 | 15.6 | 84.3 |
| 2002..... | 3.7 | 1.4 | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.8 | 16.8 | 91.8 |
| 2001..... | 4.0 | 2.4 | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 4.0 | 17.1 | 103.5 |
| 2000..... | 3.9 | 2.0 | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.7 | 17.6 | 103.3 |
| 1999..... | 4.2 | 1.8 | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.0 | 18.4 | 116.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | 1.2 | * | * | * | * | 0.2 | 0.5 | 0.9 | 3.0 | 13.7 | 72.0 |
| 2003..... | 2.9 | 1.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 3.2 | 14.8 | 78.2 |
| 2002..... | 3.0 | 1.5 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.5 | 15.4 | 84.2 |
| 2001..... | 3.3 | 1.7 | 0.2 | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 3.7 | 16.3 | 94.9 |
| 2000..... | 3.3 | 1.5 | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.8 | 17.2 | 97.7 |
| 1999..... | 3.5 | 1.3 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 3.9 | 18.0 | 108.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.2 | 0.6 | 0.9 | 3.3 | 13.3 | 60.1 |
| 2003..... | 2.0 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.1 | 14.3 | 66.3 |
| 2002..... | 2.0 | 2.2 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.4 | 14.5 | 67.2 |
| 2001..... | 2.1 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.7 | 16.0 | 76.7 |
| 2000..... | 2.3 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.2 | 17.1 | 86.8 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 3.9 | 17.4 | 84.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.7 | 14.1 | 77.3 |
| 2003..... | 3.8 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 3.2 | 15.1 | 83.4 |
| 2002..... | 4.1 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.5 | 16.0 | 91.5 |
| 2001..... | 4.4 | 1.9 | * | * | * | * | 0.3 | 0.4 | 1.0 | 3.7 | 16.6 | 102.5 |
| 2000..... | 4.3 | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.4 | 17.3 | 102.1 |
| 1999..... | 4.7 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.8 | 18.3 | 117.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | 0.6 | 0.9 | 2.7 | 6.5 | 20.0 | 85.5 |
| 2003..... | 2.5 | 4.7 | * | * | * | * | 0.6 | 1.4 | 2.4 | 7.1 | 26.8 | 97.6 |
| 2002..... | 2.6 | 3.6 | * | * | * | * | 0.6 | 1.3 | 2.6 | 8.1 | 28.8 | 106.2 |
| 2001..... | 2.9 | 5.4 | * | * | * | * | 0.4 | 1.5 | 2.9 | 8.5 | 29.6 | 121.1 |
| 2000..... | 2.8 | 3.5 | * | * | * | * | 0.6 | 1.4 | 2.8 | 8.4 | 27.6 | 119.9 |
| 1999..... | 2.8 | 5.1 | * | * | * | * | 0.8 | 1.4 | 2.9 | 9.4 | 26.5 | 115.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | 1.2 | 3.9 | 7.1 | 22.8 | 88.4 |
| 2003..... | 2.1 | * | * | * | * | * | * | 1.9 | 2.4 | 8.5 | 33.5 | 78.8 |
| 2002..... | 2.2 | * | * | * | * | * | * | 1.8 | 3.4 | 8.8 | 30.5 | 101.7 |
| 2001..... | 2.3 | * | * | * | * | * | * | 1.7 | 3.8 | 10.4 | 35.4 | 99.3 |
| 2000..... | 2.3 | * | * | * | * | * | 0.8 | 1.8 | 3.8 | 10.0 | 34.2 | 98.6 |
| 1999..... | 2.6 | * | * | * | * | * | 1.2 | 1.7 | 3.5 | 13.2 | 33.1 | 126.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | * | 1.8 | 6.0 | 18.4 | 84.3 |
| 2003..... | 2.9 | * | * | * | * | * | * | 1.0 | 2.4 | 6.2 | 23.0 | 104.8 |
| 2002..... | 3.1 | * | * | * | * | * | * | 0.9 | 1.9 | 7.7 | 27.9 | 107.9 |
| 2001..... | 3.3 | * | * | * | * | * | * | 1.4 | 2.2 | 7.1 | 26.3 | 129.2 |
| 2000..... | 3.2 | * | * | * | * | * | * | 1.1 | 2.0 | 7.3 | 23.8 | 127.8 |
| 1999..... | 3.0 | * | * | * | * | * | * | 1.2 | 2.5 | 6.8 | 22.8 | 111.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other metabolic disorders (E70-E83,E85,E88) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | 3.1 | 0.7 | 0.3 | 0.2 | 0.2 | 0.5 | 1.5 | 4.2 | 10.1 | 20.9 | 37.1 |
| 2003..... | 2.9 | 3.1 | 0.8 | 0.3 | 0.2 | 0.2 | 0.5 | 1.6 | 4.1 | 9.7 | 19.4 | 35.2 |
| 2002..... | 2.8 | 3.4 | 0.7 | 0.3 | 0.2 | 0.2 | 0.5 | 1.5 | 4.3 | 9.2 | 19.1 | 31.1 |
| 2001..... | 2.6 | 3.7 | 0.8 | 0.3 | 0.2 | 0.1 | 0.4 | 1.5 | 3.8 | 9.0 | 17.2 | 30.5 |
| 2000..... | 2.4 | 3.2 | 0.8 | 0.3 | 0.2 | 0.2 | 0.3 | 1.4 | 3.8 | 8.6 | 16.4 | 27.3 |
| 1999..... | 2.2 | 3.4 | 0.7 | 0.3 | 0.2 | 0.1 | 0.4 | 1.2 | 3.6 | 8.0 | 14.8 | 24.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.3 | 3.8 | 0.7 | 0.3 | 0.2 | 0.2 | 0.6 | 2.2 | 5.8 | 13.4 | 25.2 | 43.2 |
| 2003..... | 3.2 | 3.5 | 1.0 | 0.4 | 0.3 | 0.2 | 0.7 | 2.3 | 5.7 | 13.3 | 23.3 | 39.2 |
| 2002..... | 3.0 | 3.6 | 0.7 | 0.4 | 0.3 | 0.2 | 0.6 | 2.2 | 6.0 | 12.1 | 23.4 | 33.6 |
| 2001..... | 2.9 | 4.1 | 0.9 | 0.3 | 0.2 | 0.2 | 0.5 | 2.2 | 5.5 | 12.6 | 21.7 | 33.3 |
| 2000..... | 2.6 | 3.6 | 0.8 | 0.3 | 0.2 | 0.3 | 0.4 | 2.2 | 5.3 | 11.3 | 19.0 | 27.7 |
| 1999..... | 2.4 | 3.7 | 0.8 | 0.3 | 0.2 | 0.1 | 0.5 | 1.7 | 4.8 | 10.5 | 17.3 | 25.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.8 | 2.4 | 0.6 | 0.2 | 0.2 | 0.1 | 0.3 | 0.8 | 2.7 | 7.3 | 17.9 | 34.4 |
| 2003..... | 2.6 | 2.8 | 0.7 | 0.2 | 0.2 | 0.2 | 0.3 | 0.9 | 2.5 | 6.7 | 16.8 | 33.4 |
| 2002..... | 2.5 | 3.2 | 0.7 | 0.2 | 0.1 | 0.1 | 0.3 | 0.8 | 2.8 | 6.7 | 16.3 | 30.0 |
| 2001..... | 2.3 | 3.2 | 0.7 | 0.3 | 0.1 | 0.1 | 0.3 | 0.9 | 2.3 | 6.1 | 14.3 | 29.3 |
| 2000..... | 2.3 | 2.8 | 0.7 | 0.2 | 0.2 | * | 0.2 | 0.6 | 2.3 | 6.3 | 14.8 | 27.1 |
| 1999..... | 2.1 | 3.1 | 0.6 | 0.3 | 0.2 | 0.1 | 0.3 | 0.8 | 2.4 | 5.9 | 13.2 | 23.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | 3.3 | 0.8 | * | * | * | * | * | 2.1 | 5.4 | 13.7 | 28.1 |
| 2003..... | 1.0 | 3.7 | 0.9 | 0.3 | * | * | * | 0.8 | 2.6 | 5.2 | 13.3 | 28.0 |
| 2002..... | 1.0 | 2.9 | 0.7 | * | * | * | * | 0.7 | 2.0 | 6.2 | 16.3 | 32.2 |
| 2001..... | 0.9 | 2.9 | 0.7 | * | * | * | * | * | 2.8 | 5.3 | 13.2 | 28.8 |
| 2000..... | 0.9 | 3.0 | 0.7 | * | * | * | * | 0.8 | 2.0 | 5.9 | 11.7 | 26.5 |
| 1999..... | 0.8 | 3.4 | * | * | * | * | * | * | 2.0 | 5.6 | 10.8 | 25.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | 2.6 | 6.7 | 16.1 | * |
| 2003..... | 1.0 | * | * | * | * | * | * | * | 3.4 | 5.5 | 14.2 | * |
| 2002..... | 1.0 | * | * | * | * | * | * | * | 2.1 | 8.0 | 18.9 | 38.5 |
| 2001..... | 1.0 | * | * | * | * | * | * | * | 4.0 | 8.4 | 15.2 | * |
| 2000..... | 0.9 | * | * | * | * | * | * | * | 2.9 | 8.9 | * | * |
| 1999..... | 0.8 | * | * | * | * | * | * | * | 2.6 | 7.0 | 11.5 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | * | 4.3 | 12.1 | 28.8 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | 1.9 | 4.9 | 12.6 | 28.9 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | 4.7 | 14.6 | 29.0 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | * | 11.8 | 27.9 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | 3.5 | 13.2 | 27.7 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 4.5 | 10.4 | 26.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other metabolic disorders (E70-E83,E85,E88) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | 3.1 | 0.6 | 0.3 | 0.2 | 0.2 | 0.5 | 1.6 | 4.4 | 10.4 | 21.2 | 37.5 |
| 2003..... | 3.2 | 2.9 | 0.8 | 0.3 | 0.2 | 0.2 | 0.5 | 1.7 | 4.2 | 10.0 | 19.7 | 35.4 |
| 2002..... | 3.0 | 3.6 | 0.7 | 0.3 | 0.2 | 0.2 | 0.5 | 1.6 | 4.5 | 9.4 | 19.2 | 30.9 |
| 2001..... | 2.9 | 3.8 | 0.8 | 0.3 | 0.2 | 0.1 | 0.4 | 1.6 | 3.9 | 9.3 | 17.3 | 30.4 |
| 2000..... | 2.7 | 3.3 | 0.8 | 0.3 | 0.2 | 0.2 | 0.4 | 1.5 | 3.9 | 8.7 | 16.6 | 27.2 |
| 1999..... | 2.4 | 3.3 | 0.7 | 0.3 | 0.2 | 0.2 | 0.4 | 1.3 | 3.7 | 8.1 | 15.0 | 24.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.7 | 3.8 | 0.7 | 0.3 | 0.2 | 0.2 | 0.7 | 2.4 | 6.1 | 13.8 | 25.6 | 44.0 |
| 2003..... | 3.6 | 3.3 | 1.0 | 0.4 | 0.3 | 0.2 | 0.8 | 2.4 | 5.9 | 13.8 | 23.8 | 39.6 |
| 2002..... | 3.3 | 3.8 | 0.6 | 0.4 | 0.3 | 0.2 | 0.7 | 2.3 | 6.3 | 12.3 | 23.5 | 33.2 |
| 2001..... | 3.2 | 4.4 | 0.9 | 0.3 | 0.2 | 0.2 | 0.6 | 2.3 | 5.6 | 12.8 | 21.9 | 33.2 |
| 2000..... | 2.9 | 3.9 | 0.8 | 0.3 | 0.3 | 0.3 | 0.5 | 2.3 | 5.5 | 11.4 | 19.4 | 27.7 |
| 1999..... | 2.6 | 3.5 | 0.9 | 0.3 | 0.2 | 0.2 | 0.5 | 1.8 | 5.0 | 10.7 | 17.5 | 25.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.1 | 2.3 | 0.6 | 0.3 | 0.2 | 0.2 | 0.4 | 0.8 | 2.8 | 7.6 | 18.3 | 34.6 |
| 2003..... | 2.9 | 2.5 | 0.7 | 0.2 | 0.2 | 0.2 | 0.3 | 0.9 | 2.6 | 6.8 | 17.0 | 33.6 |
| 2002..... | 2.7 | 3.3 | 0.7 | 0.2 | 0.1 | 0.2 | 0.3 | 0.9 | 2.8 | 6.9 | 16.3 | 29.9 |
| 2001..... | 2.5 | 3.3 | 0.7 | 0.3 | 0.1 | 0.1 | 0.3 | 0.9 | 2.4 | 6.3 | 14.3 | 29.3 |
| 2000..... | 2.4 | 2.6 | 0.7 | 0.2 | 0.2 | * | 0.2 | 0.7 | 2.4 | 6.5 | 14.8 | 27.0 |
| 1999..... | 2.3 | 3.0 | 0.5 | 0.3 | 0.2 | 0.2 | 0.3 | 0.8 | 2.5 | 6.0 | 13.3 | 23.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | 3.4 | 0.7 | 0.3 | 0.2 | 0.2 | 0.5 | 1.6 | 4.5 | 10.6 | 21.5 | 38.8 |
| 2003..... | 3.6 | 2.9 | 0.9 | 0.4 | 0.2 | 0.2 | 0.5 | 1.7 | 4.2 | 10.1 | 19.9 | 36.4 |
| 2002..... | 3.3 | 3.5 | 0.7 | 0.3 | 0.2 | 0.2 | 0.5 | 1.6 | 4.5 | 9.3 | 19.4 | 31.3 |
| 2001..... | 3.2 | 3.8 | 0.9 | 0.3 | 0.2 | 0.2 | 0.5 | 1.7 | 4.0 | 9.5 | 17.6 | 30.3 |
| 2000..... | 2.9 | 3.4 | 0.8 | 0.3 | 0.2 | 0.2 | 0.4 | 1.5 | 3.9 | 8.9 | 16.8 | 27.4 |
| 1999..... | 2.7 | 2.9 | 0.8 | 0.3 | 0.2 | 0.2 | 0.4 | 1.3 | 3.8 | 8.3 | 15.1 | 24.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.2 | 4.5 | 0.7 | 0.3 | 0.3 | 0.2 | 0.7 | 2.4 | 6.4 | 14.1 | 26.0 | 46.4 |
| 2003..... | 4.0 | 3.6 | 1.0 | 0.5 | 0.3 | 0.2 | 0.8 | 2.5 | 6.0 | 13.9 | 24.0 | 41.4 |
| 2002..... | 3.7 | 4.3 | 0.6 | 0.4 | 0.3 | 0.2 | 0.7 | 2.3 | 6.2 | 12.5 | 23.8 | 33.6 |
| 2001..... | 3.5 | 4.2 | 1.0 | 0.3 | 0.3 | 0.2 | 0.6 | 2.4 | 5.6 | 13.1 | 22.4 | 33.2 |
| 2000..... | 3.2 | 4.0 | 0.9 | 0.4 | 0.2 | 0.2 | 0.5 | 2.4 | 5.6 | 11.7 | 19.8 | 29.0 |
| 1999..... | 2.9 | 3.2 | 1.0 | 0.3 | 0.2 | * | 0.5 | 1.8 | 5.2 | 11.0 | 17.7 | 25.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.4 | 2.3 | 0.6 | 0.3 | 0.2 | 0.2 | 0.4 | 0.8 | 2.7 | 7.7 | 18.4 | 35.4 |
| 2003..... | 3.2 | 2.1 | 0.7 | 0.2 | 0.2 | 0.2 | 0.3 | 0.9 | 2.5 | 6.9 | 17.0 | 34.1 |
| 2002..... | 3.0 | 2.6 | 0.8 | 0.2 | * | 0.2 | 0.3 | 0.8 | 2.8 | 6.6 | 16.4 | 30.3 |
| 2001..... | 2.8 | 3.3 | 0.7 | 0.3 | * | * | 0.3 | 0.9 | 2.4 | 6.5 | 14.4 | 29.1 |
| 2000..... | 2.7 | 2.7 | 0.7 | 0.2 | 0.2 | * | 0.2 | 0.7 | 2.3 | 6.6 | 14.8 | 26.8 |
| 1999..... | 2.5 | 2.5 | 0.6 | 0.3 | 0.2 | 0.2 | 0.2 | 0.8 | 2.5 | 5.9 | 13.3 | 23.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other metabolic disorders (E70-E83,E85,E88) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | 0.6 | 1.9 | 4.8 | 11.6 | 21.6 | 29.1 |
| 2003..... | 2.1 | 3.2 | * | * | * | * | 0.6 | 1.9 | 4.7 | 10.9 | 19.4 | 29.2 |
| 2002..... | 2.2 | 3.9 | * | * | * | * | 0.5 | 1.9 | 5.8 | 11.4 | 19.0 | 31.2 |
| 2001..... | 1.9 | 3.9 | * | * | * | * | * | 1.7 | 4.2 | 8.9 | 17.2 | 32.5 |
| 2000..... | 1.8 | * | * | * | * | * | 0.4 | 1.3 | 4.5 | 8.3 | 16.9 | 27.3 |
| 1999..... | 1.7 | 4.6 | * | 0.3 | * | * | 0.5 | 1.5 | 3.6 | 8.6 | 14.8 | 23.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | 0.9 | 2.7 | 5.3 | 15.0 | 25.1 | 26.4 |
| 2003..... | 2.2 | * | * | * | * | * | 0.8 | 2.4 | 6.6 | 15.0 | 22.2 | 21.3 |
| 2002..... | 2.3 | * | * | * | * | * | * | 2.5 | 8.7 | 13.2 | 22.3 | 35.0 |
| 2001..... | 2.0 | * | * | * | * | * | * | 2.3 | 6.2 | 13.1 | 20.7 | 33.5 |
| 2000..... | 1.8 | * | * | * | * | * | * | 2.1 | 5.7 | 10.3 | 16.4 | * |
| 1999..... | 1.6 | * | * | * | * | * | * | 1.9 | 4.5 | 9.2 | 15.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | * | 1.1 | 4.4 | 9.1 | 19.6 | 30.1 |
| 2003..... | 2.0 | * | * | * | * | * | * | 1.4 | 3.2 | 8.0 | 17.8 | 32.2 |
| 2002..... | 2.1 | * | * | * | * | * | * | 1.4 | 3.6 | 10.2 | 17.1 | 29.7 |
| 2001..... | 1.7 | * | * | * | * | * | * | 1.2 | 2.6 | 6.0 | 15.2 | 32.1 |
| 2000..... | 1.8 | * | * | * | * | * | * | * | 3.6 | 6.8 | 17.2 | 31.0 |
| 1999..... | 1.8 | * | * | * | * | * | 0.7 | 1.2 | 2.9 | 8.2 | 14.4 | 24.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| V. Mental and behavioral disorders (F01-F99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 21.0 | * | * | 0.1 | 0.7 | 1.9 | 5.0 | 9.0 | 9.8 | 20.6 | 117.1 | 671.4 |
| 2003..... | 20.7 | * | * | 0.1 | 0.7 | 1.8 | 5.2 | 8.7 | 9.2 | 20.0 | 115.9 | 679.3 |
| 2002..... | 19.3 | * | * | 0.1 | 0.6 | 1.7 | 5.3 | 8.1 | 8.8 | 19.0 | 107.6 | 646.7 |
| 2001..... | 17.8 | * | * | 0.1 | 0.6 | 1.8 | 4.9 | 7.8 | 8.4 | 18.2 | 99.0 | 604.8 |
| 2000..... | 16.4 | * | * | 0.1 | 0.6 | 1.8 | 4.9 | 7.3 | 8.1 | 16.9 | 89.9 | 561.6 |
| 1999..... | 15.1 | * | * | 0.1 | 0.6 | 1.8 | 5.1 | 7.1 | 8.0 | 16.0 | 81.3 | 515.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 16.2 | * | * | * | 0.9 | 2.7 | 7.2 | 13.6 | 15.1 | 25.6 | 114.3 | 529.7 |
| 2003..... | 15.7 | * | * | * | 0.9 | 2.6 | 7.5 | 13.3 | 13.7 | 24.7 | 114.0 | 523.8 |
| 2002..... | 14.8 | * | * | * | 0.8 | 2.3 | 7.5 | 12.7 | 13.5 | 23.9 | 106.9 | 506.0 |
| 2001..... | 13.8 | * | * | * | 0.8 | 2.6 | 7.0 | 12.2 | 12.8 | 23.3 | 97.3 | 488.0 |
| 2000..... | 12.8 | * | * | * | 0.9 | 2.6 | 7.2 | 11.5 | 12.4 | 21.9 | 89.9 | 451.0 |
| 1999..... | 12.3 | * | * | 0.1 | 0.8 | 2.6 | 7.7 | 11.3 | 12.6 | 21.2 | 85.5 | 415.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 25.6 | * | * | * | 0.5 | 1.0 | 2.7 | 4.6 | 4.9 | 16.4 | 119.0 | 735.1 |
| 2003..... | 25.5 | * | * | * | 0.4 | 1.0 | 2.9 | 4.2 | 5.0 | 16.1 | 117.1 | 748.1 |
| 2002..... | 23.7 | * | * | * | 0.4 | 1.0 | 3.0 | 3.7 | 4.5 | 14.9 | 108.0 | 707.8 |
| 2001..... | 21.7 | * | * | * | 0.3 | 1.0 | 2.8 | 3.6 | 4.3 | 13.9 | 100.1 | 653.7 |
| 2000..... | 19.8 | * | * | * | 0.4 | 1.0 | 2.5 | 3.3 | 4.2 | 12.9 | 90.0 | 606.6 |
| 1999..... | 17.8 | * | * | * | 0.3 | 1.0 | 2.5 | 3.1 | 3.8 | 11.8 | 78.6 | 555.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | 0.7 | 1.6 | 4.3 | 7.2 | 7.1 | 13.1 | 67.1 | 355.8 |
| 2003..... | 5.4 | * | * | * | 0.8 | 1.7 | 4.3 | 8.2 | 7.4 | 13.1 | 64.6 | 347.7 |
| 2002..... | 5.1 | * | * | * | 0.6 | 1.6 | 4.6 | 7.1 | 8.2 | 13.2 | 59.2 | 332.6 |
| 2001..... | 4.9 | * | * | * | 0.7 | 1.5 | 4.2 | 6.9 | 8.4 | 12.6 | 54.6 | 371.4 |
| 2000..... | 4.6 | * | * | * | 0.7 | 1.9 | 4.8 | 6.6 | 8.1 | 11.1 | 50.0 | 307.2 |
| 1999..... | 4.8 | * | * | * | 0.7 | 2.3 | 4.7 | 7.8 | 8.3 | 11.3 | 47.7 | 343.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | 1.1 | 2.6 | 6.9 | 11.8 | 12.0 | 18.7 | 69.4 | 285.8 |
| 2003..... | 5.9 | * | * | * | 1.1 | 2.6 | 6.7 | 13.0 | 13.2 | 19.0 | 69.8 | 283.2 |
| 2002..... | 6.0 | * | * | * | 1.0 | 2.5 | 7.3 | 12.3 | 13.9 | 21.6 | 68.1 | 331.3 |
| 2001..... | 5.4 | * | * | * | 1.0 | 2.1 | 6.6 | 11.9 | 15.1 | 19.6 | 57.2 | 327.6 |
| 2000..... | 5.5 | * | * | * | 1.1 | 2.9 | 8.0 | 11.1 | 14.4 | 20.0 | 48.2 | 298.3 |
| 1999..... | 5.8 | * | * | * | 1.2 | 3.7 | 8.0 | 13.3 | 15.4 | 17.8 | 51.6 | 325.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | * | * | 1.5 | 2.6 | 2.6 | 8.7 | 65.4 | 391.8 |
| 2003..... | 4.9 | * | * | * | * | 0.6 | 1.7 | 3.3 | 2.1 | 8.4 | 61.1 | 380.5 |
| 2002..... | 4.2 | * | * | * | * | 0.7 | 1.7 | 1.9 | 3.0 | 6.5 | 53.1 | 333.2 |
| 2001..... | 4.4 | * | * | * | * | 0.8 | 1.6 | 2.0 | 2.5 | 7.1 | 52.9 | 392.8 |
| 2000..... | 3.6 | * | * | * | * | 0.8 | 1.3 | 2.1 | 2.4 | 4.0 | 51.2 | 311.6 |
| 1999..... | 3.8 | * | * | * | * | 0.8 | 1.2 | 2.5 | * | 6.2 | 45.1 | 351.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| V. Mental and behavioral disorders (F01-F99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 23.4 | * | * | 0.1 | 0.7 | 1.9 | 5.0 | 9.2 | 10.0 | 21.1 | 119.7 | 684.6 |
| 2003..... | 23.0 | * | * | 0.1 | 0.7 | 1.8 | 5.3 | 8.7 | 9.3 | 20.4 | 118.3 | 692.3 |
| 2002..... | 21.5 | * | * | 0.1 | 0.6 | 1.6 | 5.3 | 8.1 | 8.8 | 19.3 | 109.5 | 658.0 |
| 2001..... | 19.7 | * | * | 0.1 | 0.6 | 1.9 | 4.9 | 7.8 | 8.3 | 18.5 | 100.9 | 612.1 |
| 2000..... | 18.0 | * | * | 0.1 | 0.6 | 1.8 | 4.8 | 7.3 | 8.1 | 17.2 | 91.4 | 569.7 |
| 1999..... | 16.4 | * | * | 0.1 | 0.5 | 1.7 | 5.0 | 7.0 | 7.9 | 16.2 | 82.5 | 520.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 17.9 | * | * | * | 0.9 | 2.7 | 7.2 | 13.7 | 15.2 | 26.0 | 116.7 | 541.0 |
| 2003..... | 17.2 | * | * | * | 0.9 | 2.6 | 7.5 | 13.2 | 13.6 | 25.0 | 116.2 | 534.1 |
| 2002..... | 16.2 | * | * | * | 0.8 | 2.2 | 7.4 | 12.5 | 13.4 | 23.9 | 108.5 | 512.8 |
| 2001..... | 15.1 | * | * | * | 0.8 | 2.8 | 6.9 | 12.1 | 12.5 | 23.5 | 99.0 | 493.5 |
| 2000..... | 13.8 | * | * | * | 0.8 | 2.5 | 7.0 | 11.3 | 12.1 | 21.8 | 91.4 | 456.4 |
| 1999..... | 13.1 | * | * | * | 0.7 | 2.4 | 7.5 | 11.0 | 12.2 | 21.2 | 86.7 | 418.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 28.7 | * | * | * | 0.6 | 1.2 | 2.9 | 4.8 | 5.1 | 16.9 | 121.7 | 748.8 |
| 2003..... | 28.5 | * | * | * | 0.4 | 1.1 | 3.1 | 4.2 | 5.3 | 16.6 | 119.7 | 761.7 |
| 2002..... | 26.5 | * | * | * | 0.4 | 1.1 | 3.2 | 3.8 | 4.6 | 15.5 | 110.2 | 720.6 |
| 2001..... | 24.1 | * | * | * | 0.3 | 1.0 | 2.9 | 3.7 | 4.4 | 14.3 | 102.1 | 661.4 |
| 2000..... | 21.9 | * | * | * | 0.4 | 1.0 | 2.6 | 3.4 | 4.3 | 13.4 | 91.4 | 615.5 |
| 1999..... | 19.6 | * | * | * | 0.3 | 1.0 | 2.6 | 3.2 | 3.9 | 12.1 | 79.8 | 560.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 26.3 | * | * | * | 0.8 | 1.9 | 5.0 | 8.7 | 9.5 | 20.5 | 121.1 | 702.1 |
| 2003..... | 25.9 | * | * | * | 0.7 | 1.8 | 5.1 | 8.1 | 8.7 | 20.3 | 119.5 | 711.5 |
| 2002..... | 24.1 | * | * | * | 0.6 | 1.7 | 5.1 | 7.5 | 8.1 | 18.9 | 111.0 | 675.7 |
| 2001..... | 22.0 | * | * | * | 0.6 | 1.9 | 4.5 | 6.8 | 7.6 | 18.1 | 102.8 | 629.4 |
| 2000..... | 19.9 | * | * | * | 0.6 | 1.7 | 4.5 | 6.5 | 7.3 | 16.7 | 92.3 | 586.2 |
| 1999..... | 18.2 | * | * | 0.1 | 0.5 | 1.7 | 4.5 | 6.0 | 7.1 | 15.7 | 83.6 | 540.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 19.4 | * | * | * | 1.0 | 2.6 | 7.3 | 13.0 | 14.2 | 24.6 | 115.8 | 551.0 |
| 2003..... | 18.7 | * | * | * | 0.9 | 2.6 | 7.3 | 12.2 | 12.4 | 24.3 | 115.6 | 547.0 |
| 2002..... | 17.4 | * | * | * | 0.8 | 2.2 | 7.2 | 11.5 | 11.9 | 22.4 | 108.2 | 522.2 |
| 2001..... | 16.1 | * | * | * | 0.8 | 2.7 | 6.4 | 10.5 | 11.0 | 22.3 | 99.7 | 504.2 |
| 2000..... | 14.6 | * | * | * | 0.8 | 2.4 | 6.5 | 10.0 | 10.7 | 20.5 | 91.2 | 464.8 |
| 1999..... | 13.8 | * | * | * | 0.7 | 2.5 | 6.8 | 9.5 | 10.7 | 20.0 | 86.1 | 432.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 33.0 | * | * | * | 0.6 | 1.2 | 2.8 | 4.5 | 5.1 | 16.9 | 124.7 | 769.8 |
| 2003..... | 32.8 | * | * | * | 0.4 | 1.0 | 2.8 | 4.0 | 5.2 | 16.9 | 122.1 | 783.9 |
| 2002..... | 30.5 | * | * | * | 0.4 | 1.1 | 2.9 | 3.6 | 4.5 | 16.0 | 112.8 | 742.0 |
| 2001..... | 27.6 | * | * | * | 0.3 | 1.0 | 2.6 | 3.2 | 4.4 | 14.5 | 104.8 | 681.5 |
| 2000..... | 25.0 | * | * | * | 0.5 | 1.0 | 2.4 | 3.0 | 4.2 | 13.5 | 93.0 | 635.1 |
| 1999..... | 22.5 | * | * | * | 0.3 | 1.0 | 2.2 | 2.6 | 3.7 | 12.1 | 81.9 | 583.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| V. Mental and behavioral disorders (F01-F99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 14.9 | * | * | * | 0.4 | 2.2 | 5.6 | 14.1 | 15.8 | 31.9 | 135.5 | 598.7 |
| 2003..... | 14.4 | * | * | * | 0.5 | 1.9 | 7.1 | 13.9 | 16.0 | 25.7 | 133.0 | 584.5 |
| 2002..... | 13.7 | * | * | * | 0.5 | 1.9 | 7.4 | 13.9 | 16.4 | 26.7 | 117.0 | 548.7 |
| 2001..... | 12.9 | * | * | * | 0.6 | 2.0 | 7.8 | 15.9 | 15.7 | 26.6 | 102.1 | 492.3 |
| 2000..... | 12.2 | * | * | * | 0.7 | 2.4 | 7.4 | 14.6 | 15.4 | 25.6 | 99.2 | 448.1 |
| 1999..... | 11.0 | * | * | * | 0.6 | 1.9 | 8.8 | 14.8 | 16.1 | 24.4 | 85.6 | 350.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 14.5 | * | * | * | * | 3.7 | 7.9 | 21.5 | 28.4 | 46.9 | 159.7 | 520.3 |
| 2003..... | 13.8 | * | * | * | 0.7 | 2.5 | 9.8 | 22.9 | 27.1 | 37.9 | 150.3 | 488.0 |
| 2002..... | 13.5 | * | * | * | * | 2.5 | 10.0 | 22.7 | 28.8 | 42.6 | 137.5 | 461.6 |
| 2001..... | 13.4 | * | * | * | 0.9 | 3.1 | 11.1 | 26.1 | 28.3 | 40.6 | 118.2 | 427.3 |
| 2000..... | 12.9 | * | * | * | 1.1 | 3.5 | 10.9 | 24.2 | 26.5 | 40.1 | 114.8 | 421.5 |
| 1999..... | 12.2 | * | * | * | 0.8 | 2.6 | 12.7 | 23.4 | 28.2 | 38.0 | 112.3 | 312.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 15.3 | * | * | * | * | 0.8 | 3.5 | 7.8 | 5.8 | 21.2 | 121.9 | 629.7 |
| 2003..... | 15.0 | * | * | * | * | 1.4 | 4.7 | 6.2 | 7.1 | 17.1 | 123.2 | 621.9 |
| 2002..... | 13.8 | * | * | * | * | 1.3 | 5.0 | 6.3 | 6.5 | 15.4 | 105.4 | 581.9 |
| 2001..... | 12.5 | * | * | * | * | 1.0 | 4.8 | 7.1 | 5.6 | 16.6 | 93.0 | 516.3 |
| 2000..... | 11.6 | * | * | * | * | 1.5 | 4.3 | 6.4 | 6.5 | 15.4 | 90.5 | 457.9 |
| 1999..... | 9.9 | * | * | * | * | 1.3 | 5.3 | 7.4 | 6.4 | 14.8 | 70.7 | 364.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Organic dementia (F01,F03) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.9 | * | * | * | * | * | * | 0.2 | 1.4 | 12.2 | 103.5 | 628.2 |
| 2003..... | 15.7 | * | * | * | * | * | * | 0.1 | 1.3 | 11.8 | 102.4 | 631.7 |
| 2002..... | 14.5 | * | * | * | * | * | * | 0.1 | 1.1 | 10.9 | 94.3 | 598.3 |
| 2001..... | 13.1 | * | * | * | * | * | * | 0.1 | 1.1 | 10.5 | 84.9 | 554.0 |
| 2000..... | 11.7 | * | * | * | * | * | * | 0.1 | 1.0 | 9.3 | 76.0 | 505.6 |
| 1999..... | 10.3 | * | * | * | * | * | * | 0.1 | 0.9 | 8.3 | 65.8 | 454.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 9.5 | * | * | * | * | * | * | 0.2 | 1.5 | 12.6 | 97.3 | 487.8 |
| 2003..... | 9.3 | * | * | * | * | * | * | 0.1 | 1.4 | 12.9 | 96.9 | 480.4 |
| 2002..... | 8.7 | * | * | * | * | * | * | 0.1 | 1.1 | 12.0 | 91.4 | 465.7 |
| 2001..... | 7.8 | * | * | * | * | * | * | 0.2 | 1.2 | 11.5 | 81.1 | 442.4 |
| 2000..... | 6.9 | * | * | * | * | * | * | 0.1 | 1.1 | 10.4 | 73.7 | 398.9 |
| 1999..... | 6.2 | * | * | * | * | * | * | 0.1 | 1.1 | 9.8 | 66.4 | 361.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 22.1 | * | * | * | * | * | * | 0.1 | 1.2 | 11.7 | 107.7 | 691.3 |
| 2003..... | 21.9 | * | * | * | * | * | * | 0.1 | 1.2 | 11.0 | 106.0 | 698.5 |
| 2002..... | 20.1 | * | * | * | * | * | * | 0.1 | 1.0 | 10.0 | 96.3 | 655.9 |
| 2001..... | 18.2 | * | * | * | * | * | * | 0.1 | 1.0 | 9.6 | 87.3 | 600.8 |
| 2000..... | 16.3 | * | * | * | * | * | * | 0.1 | 0.9 | 8.3 | 77.5 | 549.1 |
| 1999..... | 14.2 | * | * | * | * | * | * | * | 0.7 | 7.1 | 65.5 | 491.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | * | * | * | * | 7.9 | 59.5 | 329.7 |
| 2003..... | 2.7 | * | * | * | * | * | * | * | * | 6.4 | 54.2 | 319.7 |
| 2002..... | 2.3 | * | * | * | * | * | * | * | * | 5.8 | 48.5 | 293.8 |
| 2001..... | 2.4 | * | * | * | * | * | * | * | * | 6.0 | 44.7 | 337.6 |
| 2000..... | 1.9 | * | * | * | * | * | * | * | * | 4.6 | 40.9 | 268.1 |
| 1999..... | 1.9 | * | * | * | * | * | * | * | * | 4.5 | 35.2 | 295.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | * | * | 8.7 | 59.7 | 257.8 |
| 2003..... | 1.7 | * | * | * | * | * | * | * | * | 6.6 | 53.7 | 264.8 |
| 2002..... | 1.7 | * | * | * | * | * | * | * | * | 7.6 | 50.5 | 291.1 |
| 2001..... | 1.6 | * | * | * | * | * | * | * | * | 7.8 | 41.1 | 300.9 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 7.2 | 38.4 | 266.0 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 4.8 | 34.4 | 280.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | * | * | * | * | 7.2 | 59.3 | 366.6 |
| 2003..... | 3.6 | * | * | * | * | * | * | * | * | 6.2 | 54.5 | 347.7 |
| 2002..... | 3.0 | * | * | * | * | * | * | * | * | 4.4 | 47.2 | 295.1 |
| 2001..... | 3.2 | * | * | * | * | * | * | * | * | 4.6 | 47.1 | 355.6 |
| 2000..... | 2.5 | * | * | * | * | * | * | * | * | * | 42.6 | 269.1 |
| 1999..... | 2.6 | * | * | * | * | * | * | * | * | 4.3 | 35.7 | 302.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Organic dementia (F01,F03) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 18.0 | * | * | * | * | * | * | 0.2 | 1.4 | 12.4 | 105.8 | 640.6 |
| 2003..... | 17.7 | * | * | * | * | * | * | 0.1 | 1.4 | 12.2 | 104.8 | 643.9 |
| 2002..... | 16.4 | * | * | * | * | * | * | 0.1 | 1.1 | 11.2 | 96.3 | 609.3 |
| 2001..... | 14.7 | * | * | * | * | * | * | 0.1 | 1.1 | 10.7 | 86.6 | 560.9 |
| 2000..... | 13.0 | * | * | * | * | * | * | 0.1 | 1.0 | 9.5 | 77.3 | 513.2 |
| 1999..... | 11.4 | * | * | * | * | * | * | 0.1 | 0.9 | 8.5 | 66.9 | 458.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 10.8 | * | * | * | * | * | * | 0.2 | 1.6 | 12.9 | 99.3 | 498.4 |
| 2003..... | 10.5 | * | * | * | * | * | * | 0.1 | 1.5 | 13.3 | 99.1 | 489.8 |
| 2002..... | 9.8 | * | * | * | * | * | * | 0.1 | 1.1 | 12.3 | 93.2 | 472.6 |
| 2001..... | 8.8 | * | * | * | * | * | * | 0.1 | 1.2 | 11.7 | 82.8 | 447.3 |
| 2000..... | 7.7 | * | * | * | * | * | * | 0.1 | 1.1 | 10.6 | 75.0 | 403.5 |
| 1999..... | 6.9 | * | * | * | * | * | * | 0.1 | 1.1 | 10.0 | 67.6 | 364.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 24.8 | * | * | * | * | * | * | 0.2 | 1.3 | 12.0 | 110.2 | 704.3 |
| 2003..... | 24.6 | * | * | * | * | * | * | 0.1 | 1.2 | 11.3 | 108.5 | 711.6 |
| 2002..... | 22.6 | * | * | * | * | * | * | 0.1 | 1.1 | 10.4 | 98.3 | 668.3 |
| 2001..... | 20.3 | * | * | * | * | * | * | 0.1 | 1.0 | 9.9 | 89.0 | 608.1 |
| 2000..... | 18.1 | * | * | * | * | * | * | 0.1 | 0.9 | 8.7 | 78.8 | 557.6 |
| 1999..... | 15.7 | * | * | * | * | * | * | * | 0.7 | 7.3 | 66.5 | 496.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 20.7 | * | * | * | * | * | * | 0.2 | 1.4 | 12.2 | 106.9 | 656.1 |
| 2003..... | 20.4 | * | * | * | * | * | * | 0.1 | 1.3 | 12.1 | 105.6 | 661.1 |
| 2002..... | 18.8 | * | * | * | * | * | * | 0.1 | 1.1 | 11.2 | 97.5 | 625.6 |
| 2001..... | 16.9 | * | * | * | * | * | * | 0.1 | 1.1 | 10.6 | 88.2 | 576.4 |
| 2000..... | 14.9 | * | * | * | * | * | * | 0.1 | 0.9 | 9.3 | 78.0 | 527.7 |
| 1999..... | 13.2 | * | * | * | * | * | * | 0.1 | 0.9 | 8.4 | 67.9 | 476.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 12.2 | * | * | * | * | * | * | 0.2 | 1.6 | 12.5 | 98.2 | 507.8 |
| 2003..... | 11.9 | * | * | * | * | * | * | * | 1.4 | 12.9 | 98.4 | 501.1 |
| 2002..... | 11.1 | * | * | * | * | * | * | * | 1.2 | 11.9 | 93.0 | 481.1 |
| 2001..... | 9.9 | * | * | * | * | * | * | * | 1.2 | 11.3 | 83.6 | 457.4 |
| 2000..... | 8.6 | * | * | * | * | * | * | * | 1.0 | 10.0 | 74.9 | 410.6 |
| 1999..... | 7.8 | * | * | * | * | * | * | * | 1.1 | 9.8 | 67.5 | 376.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 28.8 | * | * | * | * | * | * | 0.2 | 1.3 | 11.9 | 112.8 | 722.6 |
| 2003..... | 28.5 | * | * | * | * | * | * | * | 1.3 | 11.5 | 110.5 | 731.6 |
| 2002..... | 26.3 | * | * | * | * | * | * | 0.2 | 1.1 | 10.6 | 100.4 | 688.0 |
| 2001..... | 23.5 | * | * | * | * | * | * | 0.1 | 1.1 | 10.0 | 91.2 | 626.0 |
| 2000..... | 20.9 | * | * | * | * | * | * | * | 0.9 | 8.6 | 79.9 | 574.9 |
| 1999..... | 18.3 | * | * | * | * | * | * | * | 0.7 | 7.3 | 68.1 | 516.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Organic dementia (F01,F03) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 9.6 | * | * | * | * | * | * | * | 1.8 | 18.5 | 122.7 | 569.0 |
| 2003..... | 9.1 | * | * | * | * | * | * | * | 1.9 | 15.7 | 119.3 | 551.2 |
| 2002..... | 8.2 | * | * | * | * | * | * | * | 1.3 | 14.6 | 103.6 | 510.3 |
| 2001..... | 7.2 | * | * | * | * | * | * | * | 1.5 | 15.0 | 87.9 | 455.7 |
| 2000..... | 6.6 | * | * | * | * | * | * | * | 1.6 | 14.0 | 84.3 | 408.0 |
| 1999..... | 5.1 | * | * | * | * | * | * | * | 1.1 | 11.4 | 68.1 | 310.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | * | * | 2.4 | 21.5 | 141.5 | 477.6 |
| 2003..... | 6.0 | * | * | * | * | * | * | * | 2.5 | 20.2 | 130.4 | 450.7 |
| 2002..... | 5.5 | * | * | * | * | * | * | * | * | 19.0 | 117.5 | 428.8 |
| 2001..... | 4.8 | * | * | * | * | * | * | * | 2.0 | 19.4 | 96.3 | 385.8 |
| 2000..... | 4.6 | * | * | * | * | * | * | * | 2.0 | 18.4 | 92.9 | 378.1 |
| 1999..... | 3.7 | * | * | * | * | * | * | * | * | 14.3 | 81.9 | 268.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 12.4 | * | * | * | * | * | * | * | 1.3 | 16.4 | 112.1 | 605.2 |
| 2003..... | 11.9 | * | * | * | * | * | * | * | 1.3 | 12.5 | 113.1 | 590.2 |
| 2002..... | 10.6 | * | * | * | * | * | * | * | 1.4 | 11.4 | 95.7 | 541.4 |
| 2001..... | 9.4 | * | * | * | * | * | * | * | * | 11.8 | 83.2 | 481.6 |
| 2000..... | 8.4 | * | * | * | * | * | * | * | * | 10.8 | 79.5 | 419.1 |
| 1999..... | 6.5 | * | * | * | * | * | * | * | * | 9.3 | 60.4 | 326.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified organic mental disorders (F04-F09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.1 | 0.6 | 3.8 | 22.5 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | 0.2 | 0.6 | 4.1 | 24.7 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 4.0 | 26.7 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 0.1 | 0.8 | 5.0 | 29.4 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | 0.2 | 0.8 | 5.0 | 33.7 |
| 1999..... | 1.0 | * | * | * | * | * | * | * | 0.2 | 0.8 | 5.5 | 37.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.2 | 19.8 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.6 | 21.2 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.0 | 4.1 | 19.9 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.0 | 4.8 | 23.9 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 1.0 | 4.7 | 28.7 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 0.9 | 5.6 | 29.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | * | 0.5 | 3.6 | 23.7 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | * | 0.6 | 3.7 | 26.2 |
| 2002..... | 1.0 | * | * | * | * | * | * | * | * | 0.7 | 4.0 | 29.6 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | * | 0.7 | 5.1 | 31.7 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 0.2 | 0.8 | 5.2 | 35.7 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 0.7 | 5.4 | 40.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 0.1 | 0.6 | 4.0 | 23.6 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | 0.1 | 0.6 | 4.2 | 25.3 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | 0.2 | 0.8 | 4.2 | 27.5 |
| 2001..... | 1.0 | * | * | * | * | * | * | * | 0.2 | 0.8 | 5.1 | 30.3 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | 0.2 | 0.9 | 5.1 | 34.7 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.8 | 5.5 | 38.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 0.7 | 4.4 | 20.6 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.6 | 21.8 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 0.9 | 4.2 | 20.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 1.0 | 4.8 | 24.3 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 0.9 | 4.8 | 28.8 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 0.2 | 0.8 | 5.4 | 30.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | * | 0.5 | 3.7 | 24.9 |
| 2003..... | 1.1 | * | * | * | * | * | * | * | * | 0.6 | 3.9 | 26.9 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | * | 0.6 | 4.1 | 30.5 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | * | 0.7 | 5.2 | 32.7 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 0.8 | 5.3 | 37.1 |
| 1999..... | 1.5 | * | * | * | * | * | * | * | * | 0.7 | 5.5 | 42.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified organic mental disorders (F04-F09) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | 3.4 | 17.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | 4.3 | 19.9 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 3.4 | 23.3 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 1.2 | 5.5 | 24.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | 5.6 | 25.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 1.4 | 6.7 | 28.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.3 | 23.1 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 25.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 9.7 | 29.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | 3.5 | 16.9 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | 3.6 | 20.2 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 3.6 | 25.5 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | * | 5.1 | 25.2 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | * | 5.7 | 25.7 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | * | 5.1 | 28.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to psychoactive substance use (F10-F19) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | 0.6 | 1.7 | 4.6 | 8.3 | 7.5 | 6.4 | 5.4 | 4.6 |
| 2003..... | 3.5 | * | * | * | 0.5 | 1.6 | 4.8 | 7.9 | 7.0 | 6.0 | 4.7 | 4.3 |
| 2002..... | 3.3 | * | * | * | 0.5 | 1.5 | 4.9 | 7.4 | 6.8 | 5.7 | 4.4 | 3.7 |
| 2001..... | 3.2 | * | * | * | 0.5 | 1.6 | 4.5 | 7.2 | 6.5 | 5.4 | 4.6 | 4.0 |
| 2000..... | 3.1 | * | * | * | 0.5 | 1.6 | 4.6 | 6.8 | 6.2 | 5.5 | 4.7 | 4.1 |
| 1999..... | 3.1 | * | * | * | 0.4 | 1.6 | 4.7 | 6.6 | 6.4 | 5.4 | 4.9 | 3.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 0.8 | 2.5 | 6.8 | 12.7 | 12.6 | 10.5 | 8.9 | 8.2 |
| 2003..... | 5.3 | * | * | * | 0.8 | 2.4 | 7.1 | 12.5 | 11.3 | 9.7 | 8.0 | 7.3 |
| 2002..... | 5.1 | * | * | * | 0.7 | 2.1 | 7.1 | 11.9 | 11.4 | 9.3 | 7.2 | 7.0 |
| 2001..... | 4.9 | * | * | * | 0.7 | 2.4 | 6.6 | 11.5 | 10.7 | 9.2 | 7.2 | 7.5 |
| 2000..... | 4.8 | * | * | * | 0.7 | 2.4 | 6.9 | 10.9 | 10.3 | 9.1 | 8.0 | 7.3 |
| 1999..... | 4.8 | * | * | * | 0.7 | 2.4 | 7.3 | 10.6 | 10.6 | 8.9 | 8.8 | 7.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | 0.3 | 0.8 | 2.3 | 4.0 | 2.8 | 3.0 | 3.0 | 3.0 |
| 2003..... | 1.7 | * | * | * | 0.2 | 0.8 | 2.6 | 3.5 | 3.0 | 2.9 | 2.5 | 3.0 |
| 2002..... | 1.6 | * | * | * | 0.2 | 0.8 | 2.7 | 3.1 | 2.6 | 2.7 | 2.5 | 2.2 |
| 2001..... | 1.5 | * | * | * | 0.2 | 0.8 | 2.4 | 3.1 | 2.6 | 2.3 | 2.9 | 2.5 |
| 2000..... | 1.5 | * | * | * | 0.2 | 0.8 | 2.2 | 2.8 | 2.5 | 2.5 | 2.5 | 2.8 |
| 1999..... | 1.4 | * | * | * | 0.2 | 0.8 | 2.2 | 2.7 | 2.5 | 2.6 | 2.4 | 2.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | 0.6 | 1.5 | 4.1 | 6.9 | 6.1 | 4.4 | 2.9 | * |
| 2003..... | 2.4 | * | * | * | 0.6 | 1.6 | 4.2 | 7.8 | 6.5 | 5.2 | 4.8 | * |
| 2002..... | 2.3 | * | * | * | 0.6 | 1.6 | 4.5 | 6.8 | 6.8 | 6.0 | 4.3 | * |
| 2001..... | 2.2 | * | * | * | 0.6 | 1.4 | 4.1 | 6.6 | 7.2 | 4.9 | 3.8 | * |
| 2000..... | 2.3 | * | * | * | 0.7 | 1.8 | 4.7 | 6.3 | 6.6 | 5.5 | 4.0 | * |
| 1999..... | 2.4 | * | * | * | 0.7 | 2.1 | 4.6 | 7.5 | 7.0 | 5.5 | 4.2 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | 1.1 | 2.5 | 6.5 | 11.4 | 11.1 | 8.5 | * | * |
| 2003..... | 3.9 | * | * | * | 1.0 | 2.5 | 6.6 | 12.6 | 12.5 | 10.7 | 10.4 | * |
| 2002..... | 3.9 | * | * | * | 0.9 | 2.4 | 7.2 | 12.0 | 12.6 | 11.7 | 9.0 | * |
| 2001..... | 3.5 | * | * | * | 0.9 | 2.0 | 6.5 | 11.5 | 13.6 | 9.8 | * | * |
| 2000..... | 3.8 | * | * | * | 1.0 | 2.7 | 7.8 | 10.8 | 12.7 | 11.0 | * | * |
| 1999..... | 4.1 | * | * | * | 1.2 | 3.5 | 7.8 | 13.0 | 13.4 | 11.2 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 1.4 | 2.4 | * | * | * | * |
| 2003..... | 0.8 | * | * | * | * | * | 1.6 | 3.0 | * | * | * | * |
| 2002..... | 0.7 | * | * | * | * | * | 1.6 | 1.7 | * | * | * | * |
| 2001..... | 0.7 | * | * | * | * | 0.7 | 1.5 | 1.8 | * | * | * | * |
| 2000..... | 0.7 | * | * | * | * | 0.8 | 1.3 | 2.0 | * | * | * | * |
| 1999..... | 0.6 | * | * | * | * | * | 1.2 | 2.1 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to psychoactive substance use (F10-F19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.6 | 1.7 | 4.6 | 8.3 | 7.5 | 6.5 | 5.5 | 4.7 |
| 2003..... | 3.6 | * | * | * | 0.5 | 1.6 | 4.9 | 7.9 | 7.0 | 6.0 | 4.7 | 4.4 |
| 2002..... | 3.4 | * | * | * | 0.4 | 1.4 | 4.9 | 7.4 | 6.8 | 5.6 | 4.3 | 3.6 |
| 2001..... | 3.3 | * | * | * | 0.4 | 1.6 | 4.5 | 7.2 | 6.3 | 5.4 | 4.6 | 4.1 |
| 2000..... | 3.2 | * | * | * | 0.4 | 1.6 | 4.5 | 6.7 | 6.2 | 5.4 | 4.6 | 4.1 |
| 1999..... | 3.1 | * | * | * | 0.4 | 1.5 | 4.7 | 6.4 | 6.2 | 5.3 | 4.9 | 3.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | 0.8 | 2.5 | 6.8 | 12.7 | 12.6 | 10.6 | 9.1 | 8.3 |
| 2003..... | 5.5 | * | * | * | 0.7 | 2.3 | 7.1 | 12.4 | 11.1 | 9.5 | 7.8 | 7.5 |
| 2002..... | 5.2 | * | * | * | 0.6 | 2.0 | 7.0 | 11.7 | 11.2 | 9.0 | 7.1 | 6.8 |
| 2001..... | 5.0 | * | * | * | 0.7 | 2.4 | 6.5 | 11.3 | 10.3 | 9.1 | 7.1 | 7.7 |
| 2000..... | 4.9 | * | * | * | 0.7 | 2.3 | 6.7 | 10.7 | 10.0 | 8.8 | 7.8 | 7.4 |
| 1999..... | 4.8 | * | * | * | 0.6 | 2.1 | 7.1 | 10.2 | 10.1 | 8.6 | 8.7 | 7.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | 0.3 | 1.0 | 2.5 | 4.1 | 2.8 | 3.1 | 3.1 | 3.1 |
| 2003..... | 1.8 | * | * | * | 0.2 | 0.8 | 2.7 | 3.6 | 3.1 | 3.0 | 2.5 | 3.0 |
| 2002..... | 1.7 | * | * | * | 0.2 | 0.8 | 2.8 | 3.2 | 2.7 | 2.8 | 2.5 | 2.3 |
| 2001..... | 1.6 | * | * | * | 0.1 | 0.8 | 2.6 | 3.2 | 2.7 | 2.4 | 2.9 | 2.5 |
| 2000..... | 1.5 | * | * | * | 0.2 | 0.8 | 2.4 | 2.8 | 2.6 | 2.5 | 2.6 | 2.8 |
| 1999..... | 1.5 | * | * | * | 0.2 | 0.8 | 2.3 | 2.7 | 2.6 | 2.7 | 2.4 | 2.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | 0.7 | 1.7 | 4.7 | 7.8 | 7.1 | 6.1 | 5.6 | 4.7 |
| 2003..... | 3.6 | * | * | * | 0.5 | 1.6 | 4.7 | 7.3 | 6.4 | 5.9 | 4.7 | 4.6 |
| 2002..... | 3.3 | * | * | * | 0.5 | 1.5 | 4.7 | 6.8 | 6.0 | 5.3 | 4.3 | 3.5 |
| 2001..... | 3.1 | * | * | * | 0.5 | 1.6 | 4.1 | 6.3 | 5.7 | 5.1 | 4.6 | 4.0 |
| 2000..... | 3.0 | * | * | * | 0.5 | 1.5 | 4.1 | 5.8 | 5.4 | 5.0 | 4.6 | 4.1 |
| 1999..... | 2.9 | * | * | * | 0.4 | 1.5 | 4.1 | 5.4 | 5.4 | 4.9 | 4.8 | 3.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | 0.9 | 2.5 | 6.9 | 11.9 | 11.6 | 9.5 | 9.1 | 8.2 |
| 2003..... | 5.4 | * | * | * | 0.8 | 2.4 | 6.9 | 11.3 | 9.9 | 9.2 | 7.7 | 7.5 |
| 2002..... | 5.0 | * | * | * | 0.7 | 2.1 | 6.8 | 10.7 | 9.6 | 8.1 | 6.8 | 6.7 |
| 2001..... | 4.7 | * | * | * | 0.7 | 2.4 | 5.9 | 9.9 | 8.9 | 8.3 | 6.9 | 7.5 |
| 2000..... | 4.6 | * | * | * | 0.7 | 2.2 | 6.2 | 9.4 | 8.7 | 8.0 | 7.6 | 7.3 |
| 1999..... | 4.5 | * | * | * | 0.6 | 2.1 | 6.3 | 8.8 | 8.7 | 7.6 | 8.3 | 7.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | 0.4 | 1.0 | 2.4 | 3.8 | 2.8 | 3.2 | 3.2 | 3.2 |
| 2003..... | 1.8 | * | * | * | 0.2 | 0.8 | 2.5 | 3.4 | 3.0 | 3.0 | 2.7 | 3.3 |
| 2002..... | 1.7 | * | * | * | 0.2 | 0.8 | 2.5 | 2.9 | 2.5 | 3.0 | 2.6 | 2.2 |
| 2001..... | 1.6 | * | * | * | 0.2 | 0.8 | 2.3 | 2.7 | 2.6 | 2.4 | 3.0 | 2.6 |
| 2000..... | 1.5 | * | * | * | 0.2 | 0.8 | 2.0 | 2.4 | 2.4 | 2.6 | 2.7 | 2.9 |
| 1999..... | 1.4 | * | * | * | 0.2 | 0.8 | 1.9 | 2.1 | 2.4 | 2.7 | 2.5 | 2.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to psychoactive substance use (F10-F19) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | 1.8 | 5.0 | 13.3 | 12.9 | 11.1 | 5.2 | * |
| 2003..... | 4.4 | * | * | * | * | 1.7 | 6.6 | 12.8 | 13.1 | 8.0 | 5.0 | * |
| 2002..... | 4.6 | * | * | * | * | 1.6 | 6.8 | 13.1 | 14.1 | 9.1 | 5.5 | * |
| 2001..... | 4.7 | * | * | * | 0.4 | 1.7 | 7.2 | 14.9 | 13.0 | 8.8 | 5.8 | * |
| 2000..... | 4.6 | * | * | * | 0.5 | 2.2 | 6.9 | 14.0 | 12.9 | 9.4 | 5.9 | * |
| 1999..... | 4.8 | * | * | * | 0.4 | 1.7 | 8.3 | 13.9 | 13.8 | 10.3 | 7.4 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | * | * | 3.0 | 7.3 | 20.6 | 24.7 | 22.7 | 10.8 | * |
| 2003..... | 6.9 | * | * | * | * | 2.2 | 9.2 | 21.6 | 23.3 | 14.9 | 11.0 | * |
| 2002..... | 7.2 | * | * | * | * | 2.3 | 9.2 | 21.7 | 26.3 | 19.0 | 11.7 | * |
| 2001..... | 7.6 | * | * | * | 0.7 | 2.6 | 10.4 | 24.5 | 24.6 | 18.2 | 12.0 | * |
| 2000..... | 7.4 | * | * | * | 0.8 | 3.2 | 10.0 | 23.4 | 23.5 | 18.7 | 13.3 | * |
| 1999..... | 7.6 | * | * | * | * | 2.3 | 12.2 | 22.1 | 25.5 | 20.6 | 15.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | 3.0 | 7.1 | 3.5 | 2.8 | * | * |
| 2003..... | 2.1 | * | * | * | * | 1.3 | 4.3 | 5.2 | 4.9 | 3.1 | * | * |
| 2002..... | 2.2 | * | * | * | * | 1.1 | 4.7 | 5.8 | 4.2 | * | * | * |
| 2001..... | 2.1 | * | * | * | * | 0.9 | 4.3 | 6.7 | 3.7 | 2.2 | * | * |
| 2000..... | 2.2 | * | * | * | * | 1.3 | 4.2 | 5.9 | 4.5 | 2.8 | * | * |
| 1999..... | 2.4 | * | * | * | * | 1.1 | 4.9 | 6.9 | 4.5 | 3.1 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to use of alcohol (F10) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 0.2 | 0.7 | 2.8 | 5.9 | 6.2 | 5.0 | 3.3 | 2.0 |
| 2003..... | 2.5 | * | * | * | 0.2 | 0.7 | 3.1 | 5.9 | 5.8 | 4.5 | 2.9 | 2.0 |
| 2002..... | 2.5 | * | * | * | 0.1 | 0.6 | 3.1 | 5.6 | 5.8 | 4.5 | 3.0 | 1.9 |
| 2001..... | 2.4 | * | * | * | 0.1 | 0.7 | 3.0 | 5.5 | 5.6 | 4.6 | 3.2 | 2.2 |
| 2000..... | 2.3 | * | * | * | 0.1 | 0.7 | 2.9 | 5.1 | 5.3 | 4.5 | 3.1 | 2.5 |
| 1999..... | 2.3 | * | * | * | 0.1 | 0.7 | 2.9 | 5.0 | 5.4 | 4.3 | 3.4 | 2.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | 0.2 | 1.0 | 4.4 | 9.4 | 10.6 | 8.6 | 6.0 | 3.7 |
| 2003..... | 4.0 | * | * | * | 0.2 | 1.1 | 4.7 | 9.3 | 9.5 | 7.5 | 5.4 | 3.6 |
| 2002..... | 4.0 | * | * | * | 0.2 | 0.9 | 4.7 | 9.1 | 9.8 | 7.7 | 5.6 | 4.1 |
| 2001..... | 3.8 | * | * | * | 0.3 | 1.1 | 4.5 | 8.8 | 9.2 | 8.0 | 5.5 | 4.9 |
| 2000..... | 3.7 | * | * | * | 0.2 | 1.1 | 4.5 | 8.3 | 9.0 | 7.8 | 5.8 | 5.3 |
| 1999..... | 3.6 | * | * | * | 0.2 | 0.9 | 4.5 | 8.1 | 9.0 | 7.3 | 6.7 | 4.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.4 | 1.4 | 2.6 | 2.2 | 1.9 | 1.4 | 1.2 |
| 2003..... | 1.2 | * | * | * | * | 0.3 | 1.6 | 2.5 | 2.4 | 1.9 | 1.2 | 1.3 |
| 2002..... | 1.1 | * | * | * | * | 0.3 | 1.6 | 2.3 | 2.0 | 1.8 | 1.3 | 1.0 |
| 2001..... | 1.1 | * | * | * | * | 0.4 | 1.5 | 2.3 | 2.2 | 1.7 | 1.6 | 1.1 |
| 2000..... | 1.0 | * | * | * | * | 0.3 | 1.3 | 2.0 | 2.0 | 1.7 | 1.4 | 1.4 |
| 1999..... | 1.0 | * | * | * | * | 0.4 | 1.4 | 1.9 | 2.1 | 1.8 | 1.3 | 1.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 0.2 | 0.7 | 3.0 | 6.0 | 6.0 | 4.4 | 3.3 | 2.0 |
| 2003..... | 2.6 | * | * | * | 0.1 | 0.7 | 3.2 | 5.8 | 5.5 | 4.3 | 2.8 | 2.1 |
| 2002..... | 2.5 | * | * | * | 0.1 | 0.6 | 3.2 | 5.6 | 5.2 | 4.1 | 2.9 | 1.9 |
| 2001..... | 2.4 | * | * | * | 0.1 | 0.7 | 2.9 | 5.1 | 5.0 | 4.2 | 3.1 | 2.1 |
| 2000..... | 2.2 | * | * | * | 0.1 | 0.7 | 2.8 | 4.7 | 4.7 | 4.1 | 3.0 | 2.5 |
| 1999..... | 2.2 | * | * | * | 0.1 | 0.7 | 2.7 | 4.5 | 4.8 | 3.9 | 3.2 | 2.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | 0.2 | 0.9 | 4.5 | 9.4 | 10.0 | 7.5 | 6.0 | 3.8 |
| 2003..... | 4.0 | * | * | * | 0.2 | 1.1 | 4.8 | 9.0 | 8.7 | 7.2 | 5.1 | 3.5 |
| 2002..... | 3.9 | * | * | * | 0.2 | 0.9 | 4.8 | 8.9 | 8.6 | 6.8 | 5.3 | 3.8 |
| 2001..... | 3.7 | * | * | * | 0.2 | 1.0 | 4.2 | 8.1 | 8.1 | 7.2 | 5.3 | 4.6 |
| 2000..... | 3.6 | * | * | * | * | 1.0 | 4.4 | 7.6 | 7.8 | 6.9 | 5.4 | 5.2 |
| 1999..... | 3.4 | * | * | * | 0.2 | 1.0 | 4.2 | 7.4 | 7.7 | 6.3 | 6.3 | 4.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | 0.4 | 1.4 | 2.7 | 2.2 | 1.8 | 1.4 | 1.2 |
| 2003..... | 1.2 | * | * | * | * | 0.3 | 1.6 | 2.7 | 2.4 | 1.9 | 1.2 | 1.4 |
| 2002..... | 1.1 | * | * | * | * | 0.3 | 1.5 | 2.3 | 1.9 | 1.9 | 1.3 | 1.0 |
| 2001..... | 1.1 | * | * | * | * | 0.3 | 1.5 | 2.1 | 2.1 | 1.7 | 1.7 | 1.1 |
| 2000..... | 1.0 | * | * | * | * | 0.4 | 1.2 | 1.8 | 1.9 | 1.7 | 1.5 | 1.4 |
| 1999..... | 0.9 | * | * | * | * | 0.4 | 1.2 | 1.7 | 2.0 | 1.8 | 1.2 | 1.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to use of alcohol (F10) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | 0.7 | 2.1 | 6.7 | 9.2 | 9.7 | 3.7 | * |
| 2003..... | 2.6 | * | * | * | * | 0.6 | 2.9 | 7.1 | 9.5 | 6.7 | 4.5 | * |
| 2002..... | 2.8 | * | * | * | * | 0.6 | 3.2 | 7.3 | 11.1 | 8.1 | 4.8 | * |
| 2001..... | 3.1 | * | * | * | * | 0.6 | 3.7 | 9.0 | 11.1 | 8.1 | 4.7 | * |
| 2000..... | 3.0 | * | * | * | * | 0.9 | 3.2 | 8.8 | 10.9 | 8.6 | 5.0 | * |
| 1999..... | 3.1 | * | * | * | * | 0.6 | 4.1 | 8.5 | 11.5 | 9.1 | 6.6 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | 1.2 | 3.5 | 11.6 | 18.1 | 19.7 | 7.7 | * |
| 2003..... | 4.4 | * | * | * | * | 1.1 | 4.4 | 12.9 | 17.6 | 12.6 | 10.1 | * |
| 2002..... | 4.8 | * | * | * | * | 1.1 | 4.8 | 12.5 | 21.1 | 17.3 | 10.0 | * |
| 2001..... | 5.3 | * | * | * | * | 1.0 | 6.1 | 15.3 | 21.2 | 16.9 | 10.2 | * |
| 2000..... | 5.0 | * | * | * | * | 1.5 | 4.8 | 15.0 | 20.5 | 17.7 | 12.3 | * |
| 1999..... | 5.0 | * | * | * | * | * | 6.2 | 14.0 | 21.6 | 18.6 | 13.4 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.9 | 2.5 | 2.1 | 2.6 | * | * |
| 2003..... | 1.0 | * | * | * | * | * | 1.5 | 2.2 | 3.1 | 2.6 | * | * |
| 2002..... | 1.1 | * | * | * | * | * | 1.8 | 2.8 | 3.2 | * | * | * |
| 2001..... | 1.1 | * | * | * | * | * | 1.6 | 3.6 | 2.9 | * | * | * |
| 2000..... | 1.1 | * | * | * | * | * | 1.8 | 3.4 | 3.2 | 2.2 | * | * |
| 1999..... | 1.3 | * | * | * | * | * | 2.3 | 3.7 | 3.5 | 2.4 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to other psychoactive substance use (F11-F19) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.4 | 1.0 | 1.8 | 2.4 | 1.4 | 1.5 | 2.2 | 2.7 |
| 2003..... | 1.1 | * | * | * | 0.3 | 0.9 | 1.8 | 2.1 | 1.2 | 1.5 | 1.8 | 2.3 |
| 2002..... | 0.9 | * | * | * | 0.3 | 0.8 | 1.8 | 1.8 | 1.0 | 1.1 | 1.3 | 1.7 |
| 2001..... | 0.9 | * | * | * | 0.3 | 0.8 | 1.6 | 1.8 | 0.8 | 0.8 | 1.4 | 1.8 |
| 2000..... | 0.9 | * | * | * | 0.3 | 0.9 | 1.6 | 1.6 | 0.8 | 0.9 | 1.5 | 1.6 |
| 1999..... | 0.9 | * | * | * | 0.3 | 0.9 | 1.8 | 1.6 | 0.9 | 1.0 | 1.4 | 1.5 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.6 | 1.4 | 2.5 | 3.3 | 2.1 | 2.0 | 3.0 | 4.6 |
| 2003..... | 1.5 | * | * | * | 0.5 | 1.3 | 2.5 | 3.2 | 1.7 | 2.0 | 2.5 | 3.8 |
| 2002..... | 1.3 | * | * | * | 0.4 | 1.1 | 2.4 | 2.8 | 1.4 | 1.3 | 1.5 | 2.6 |
| 2001..... | 1.2 | * | * | * | 0.4 | 1.2 | 2.1 | 2.7 | 1.2 | 1.1 | 1.6 | 2.8 |
| 2000..... | 1.2 | * | * | * | 0.5 | 1.2 | 2.3 | 2.5 | 1.1 | 1.0 | 1.9 | 2.1 |
| 1999..... | 1.3 | * | * | * | 0.4 | 1.3 | 2.7 | 2.3 | 1.3 | 1.3 | 2.0 | 2.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.2 | 0.5 | 1.1 | 1.5 | 0.7 | 1.1 | 1.6 | 1.8 |
| 2003..... | 0.7 | * | * | * | 0.2 | 0.5 | 1.1 | 1.1 | 0.7 | 1.0 | 1.3 | 1.7 |
| 2002..... | 0.6 | * | * | * | 0.2 | 0.5 | 1.2 | 0.9 | 0.6 | 1.0 | 1.2 | 1.2 |
| 2001..... | 0.5 | * | * | * | 0.1 | 0.4 | 1.0 | 0.9 | 0.5 | 0.6 | 1.3 | 1.4 |
| 2000..... | 0.6 | * | * | * | 0.2 | 0.6 | 1.0 | 0.8 | 0.6 | 0.8 | 1.2 | 1.3 |
| 1999..... | 0.5 | * | * | * | 0.1 | 0.4 | 0.9 | 0.8 | 0.4 | 0.8 | 1.1 | 1.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.3 | 0.7 | 1.8 | 2.4 | 1.9 | * | * | * |
| 2003..... | 0.9 | * | * | * | * | 0.9 | 2.0 | 2.6 | 1.2 | * | * | * |
| 2002..... | 0.8 | * | * | * | * | 0.7 | 1.9 | 2.4 | 1.1 | * | * | * |
| 2001..... | 0.7 | * | * | * | * | 0.5 | 1.8 | 2.0 | 1.1 | * | * | * |
| 2000..... | 0.7 | * | * | * | 0.3 | 1.0 | 1.9 | 1.7 | * | * | * | * |
| 1999..... | 0.9 | * | * | * | 0.4 | 1.1 | 2.0 | 2.6 | 1.2 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 1.1 | 2.6 | 3.5 | 3.3 | * | * | * |
| 2003..... | 1.3 | * | * | * | * | 1.3 | 2.9 | 4.0 | 2.2 | * | * | * |
| 2002..... | 1.2 | * | * | * | * | 1.1 | 3.0 | 4.0 | * | * | * | * |
| 2001..... | 1.0 | * | * | * | * | 0.7 | 2.6 | 3.4 | * | * | * | * |
| 2000..... | 1.1 | * | * | * | * | 1.3 | 2.9 | 2.6 | * | * | * | * |
| 1999..... | 1.5 | * | * | * | 0.6 | 1.8 | 3.1 | 4.5 | 2.6 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 1.0 | 1.2 | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | 0.9 | 1.3 | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | 0.8 | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | 0.9 | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | 0.7 | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | 0.8 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to other psychoactive substance use (F11-F19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.4 | 1.0 | 1.8 | 2.4 | 1.3 | 1.6 | 2.2 | 2.7 |
| 2003..... | 1.1 | * | * | * | 0.3 | 0.9 | 1.8 | 2.0 | 1.1 | 1.5 | 1.8 | 2.3 |
| 2002..... | 0.9 | * | * | * | 0.3 | 0.8 | 1.7 | 1.7 | 1.0 | 1.1 | 1.3 | 1.7 |
| 2001..... | 0.9 | * | * | * | 0.3 | 0.9 | 1.5 | 1.7 | 0.8 | 0.8 | 1.4 | 1.8 |
| 2000..... | 0.9 | * | * | * | 0.3 | 0.9 | 1.6 | 1.6 | 0.8 | 0.9 | 1.5 | 1.6 |
| 1999..... | 0.9 | * | * | * | 0.3 | 0.8 | 1.7 | 1.4 | 0.8 | 1.0 | 1.5 | 1.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.6 | 1.5 | 2.5 | 3.3 | 2.0 | 2.0 | 3.1 | 4.6 |
| 2003..... | 1.5 | * | * | * | 0.5 | 1.2 | 2.4 | 3.0 | 1.6 | 2.0 | 2.4 | 3.8 |
| 2002..... | 1.3 | * | * | * | 0.4 | 1.1 | 2.2 | 2.6 | 1.4 | 1.3 | 1.5 | 2.7 |
| 2001..... | 1.2 | * | * | * | 0.5 | 1.4 | 2.0 | 2.6 | 1.1 | 1.1 | 1.6 | 2.8 |
| 2000..... | 1.2 | * | * | * | 0.5 | 1.2 | 2.2 | 2.4 | 1.1 | 1.0 | 2.0 | 2.1 |
| 1999..... | 1.2 | * | * | * | 0.4 | 1.2 | 2.5 | 2.1 | 1.2 | 1.3 | 2.0 | 2.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.2 | 0.5 | 1.1 | 1.6 | 0.7 | 1.2 | 1.7 | 1.8 |
| 2003..... | 0.7 | * | * | * | 0.2 | 0.5 | 1.2 | 1.1 | 0.7 | 1.1 | 1.4 | 1.7 |
| 2002..... | 0.6 | * | * | * | 0.2 | 0.5 | 1.2 | 0.9 | 0.6 | 1.0 | 1.2 | 1.3 |
| 2001..... | 0.6 | * | * | * | * | 0.5 | 1.0 | 0.9 | 0.5 | 0.7 | 1.3 | 1.4 |
| 2000..... | 0.6 | * | * | * | 0.2 | 0.5 | 1.0 | 0.8 | 0.6 | 0.8 | 1.2 | 1.4 |
| 1999..... | 0.5 | * | * | * | 0.1 | 0.5 | 0.9 | 0.8 | 0.5 | 0.8 | 1.1 | 1.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.5 | 1.1 | 1.7 | 1.9 | 1.1 | 1.7 | 2.3 | 2.7 |
| 2003..... | 1.0 | * | * | * | 0.4 | 0.9 | 1.5 | 1.6 | 0.9 | 1.6 | 2.0 | 2.5 |
| 2002..... | 0.9 | * | * | * | 0.3 | 0.9 | 1.5 | 1.2 | 0.8 | 1.2 | 1.4 | 1.7 |
| 2001..... | 0.8 | * | * | * | 0.3 | 0.9 | 1.2 | 1.2 | 0.7 | 0.9 | 1.5 | 1.9 |
| 2000..... | 0.8 | * | * | * | 0.4 | 0.8 | 1.3 | 1.1 | 0.7 | 1.0 | 1.6 | 1.7 |
| 1999..... | 0.8 | * | * | * | 0.3 | 0.8 | 1.4 | 0.9 | 0.6 | 1.1 | 1.6 | 1.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.7 | 1.5 | 2.4 | 2.6 | 1.6 | 2.0 | 3.1 | 4.4 |
| 2003..... | 1.4 | * | * | * | 0.6 | 1.3 | 2.1 | 2.4 | 1.2 | 2.0 | 2.6 | 4.0 |
| 2002..... | 1.1 | * | * | * | 0.5 | 1.2 | 2.0 | 1.8 | 1.0 | 1.3 | 1.5 | 2.8 |
| 2001..... | 1.1 | * | * | * | 0.5 | 1.4 | 1.7 | 1.8 | 0.9 | 1.1 | 1.6 | 2.8 |
| 2000..... | 1.1 | * | * | * | 0.5 | 1.2 | 1.8 | 1.7 | 0.9 | 1.0 | 2.1 | 2.1 |
| 1999..... | 1.1 | * | * | * | 0.4 | 1.2 | 2.1 | 1.4 | 0.9 | 1.3 | 2.0 | 2.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.3 | 0.6 | 1.0 | 1.1 | 0.6 | 1.4 | 1.8 | 2.0 |
| 2003..... | 0.6 | * | * | * | 0.2 | 0.4 | 0.9 | 0.8 | 0.6 | 1.2 | 1.5 | 1.9 |
| 2002..... | 0.6 | * | * | * | 0.2 | 0.5 | 1.0 | 0.6 | 0.6 | 1.1 | 1.3 | 1.2 |
| 2001..... | 0.5 | * | * | * | * | 0.4 | 0.8 | 0.6 | 0.5 | 0.7 | 1.4 | 1.5 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.8 | 0.5 | 0.5 | 0.9 | 1.2 | 1.5 |
| 1999..... | 0.5 | * | * | * | * | 0.4 | 0.6 | 0.5 | 0.4 | 0.9 | 1.3 | 1.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental and behavioral disorders due to other psychoactive substance use (F11-F19) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 1.0 | 2.9 | 6.6 | 3.7 | 1.4 | * | * |
| 2003..... | 1.8 | * | * | * | * | 1.1 | 3.7 | 5.6 | 3.5 | 1.3 | * | * |
| 2002..... | 1.7 | * | * | * | * | 1.0 | 3.6 | 5.8 | 2.9 | * | * | * |
| 2001..... | 1.7 | * | * | * | * | 1.1 | 3.5 | 5.9 | 1.9 | * | * | * |
| 2000..... | 1.7 | * | * | * | * | 1.3 | 3.7 | 5.2 | 2.1 | * | * | * |
| 1999..... | 1.8 | * | * | * | * | 1.1 | 4.2 | 5.5 | 2.3 | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | 1.8 | 3.8 | 9.0 | 6.6 | 3.0 | * | * |
| 2003..... | 2.5 | * | * | * | * | 1.2 | 4.7 | 8.7 | 5.7 | * | * | * |
| 2002..... | 2.5 | * | * | * | * | 1.1 | 4.4 | 9.2 | 5.3 | * | * | * |
| 2001..... | 2.4 | * | * | * | * | 1.6 | 4.4 | 9.1 | 3.4 | * | * | * |
| 2000..... | 2.4 | * | * | * | * | 1.7 | 5.2 | 8.3 | 3.0 | * | * | * |
| 1999..... | 2.5 | * | * | * | * | 1.6 | 6.0 | 8.1 | 3.9 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 2.0 | 4.6 | 1.3 | * | * | * |
| 2003..... | 1.1 | * | * | * | * | 1.1 | 2.7 | 3.0 | 1.8 | * | * | * |
| 2002..... | 1.1 | * | * | * | * | 0.9 | 2.9 | 2.9 | * | * | * | * |
| 2001..... | 1.0 | * | * | * | * | * | 2.7 | 3.1 | * | * | * | * |
| 2000..... | 1.0 | * | * | * | * | 1.0 | 2.5 | 2.6 | * | * | * | * |
| 1999..... | 1.1 | * | * | * | * | 0.8 | 2.7 | 3.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Schizophrenia, schizotypal and delusional disorders (F20-F29) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.1 | 2.8 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.6 | 1.1 | 2.7 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.2 | 3.4 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.2 | 2.7 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 3.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.5 | 5.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 2.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.2 | * | 0.4 | 0.9 | 1.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | * |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.4 | 1.2 | 4.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 4.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.3 | 3.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 3.8 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.6 | 5.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.7 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.6 | 1.2 | 2.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.4 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.2 | 2.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.4 | 5.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.2 | * | 0.5 | 0.9 | 2.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.8 | 2.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.0 | * |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.1 | 4.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.1 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | 0.7 | 1.4 | 3.3 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 4.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 1.4 | 3.4 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 3.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.7 | 5.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mood (affective) disorders (F30-F39) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 2.0 | 8.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 2.1 | 10.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 2.3 | 9.3 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.5 | 2.2 | 9.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.9 | 10.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.1 | 0.4 | 2.4 | 9.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.5 | 7.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.9 | 8.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.6 | 6.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.5 | 1.8 | 8.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.5 | 9.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 2.1 | 9.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.4 | 2.3 | 9.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.1 | 0.4 | 2.3 | 12.0 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.8 | 10.4 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 0.5 | 2.5 | 10.4 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 0.4 | 2.3 | 10.3 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 0.5 | 2.6 | 9.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 2.1 | 9.4 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 2.2 | 11.7 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 2.4 | 10.0 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | * | 0.1 | 0.5 | 2.4 | 10.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 2.1 | 10.9 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.1 | 0.5 | 2.6 | 10.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.6 | 7.9 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 2.0 | 8.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.7 | 7.4 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | 0.6 | 1.9 | 8.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.5 | 10.5 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 2.2 | 9.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | 0.4 | 2.4 | 10.1 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 0.4 | 2.4 | 12.9 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 0.6 | 2.9 | 11.2 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 0.5 | 2.7 | 11.0 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 0.5 | 2.4 | 11.0 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 0.5 | 2.9 | 10.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Mental retardation (F70-F79) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 | * |
| 2003..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 |
| 2002..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.4 | 0.4 | 0.5 |
| 2001..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | * |
| 2000..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 |
| 1999..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | 0.4 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.4 | 0.5 | 0.5 | * |
| 2003..... | 0.2 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.4 | 0.6 | * |
| 2002..... | 0.2 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.4 | 0.5 | * |
| 2001..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.4 | 0.7 | * |
| 2000..... | 0.2 | * | * | * | 0.1 | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | * |
| 1999..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.5 | 0.6 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.3 | 0.4 | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | * |
| 2000..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.2 | 0.1 | 0.3 | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.5 | * |
| 2003..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.5 | 0.5 |
| 2002..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.4 | 0.4 | 0.6 |
| 2001..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | * |
| 2000..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | 0.4 | 0.6 |
| 1999..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | 0.4 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.2 | 0.4 | 0.4 | 0.6 | 0.5 | * |
| 2003..... | 0.2 | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 0.4 | 0.7 | * |
| 2002..... | 0.2 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.5 | * |
| 2001..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.7 | * |
| 2000..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.3 | 0.5 | * |
| 1999..... | 0.2 | * | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.5 | 0.7 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.5 | * |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.3 | * |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.4 | 0.4 | * |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.3 | * |
| 2000..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.2 | * | 0.2 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified mental and behavioral disorders (F50-F69,F80-F99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.7 | 4.0 |
| 2003..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.9 | 4.4 |
| 2002..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.1 | * | 0.2 | 0.8 | 4.4 |
| 2001..... | 0.2 | * | * | * | * | 0.1 | * | * | * | 0.2 | 0.7 | 4.1 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.9 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | * | 0.3 | 0.6 | 4.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.7 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.0 | 4.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 3.5 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.5 |
| 2000..... | 0.1 | * | * | * | * | *.1 | * | * | * | * | 0.6 | 3.2 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 2.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | * | 0.1 | 0.1 | 0.2 | * | 0.8 | 4.1 |
| 2003..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.1 | * | * | 0.3 | 0.9 | 4.4 |
| 2002..... | 0.2 | * | * | * | * | 0.2 | 0.1 | 0.1 | * | 0.2 | 0.7 | 4.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 4.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.8 | 4.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | * | 0.1 | 0.1 | 0.1 | * | 0.8 | 4.1 |
| 2003..... | 0.2 | * | * | * | 0.1 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.9 | 4.4 |
| 2002..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.1 | * | 0.2 | 0.8 | 4.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.0 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | * | * | 0.3 | 0.7 | 4.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.5 |
| 2003..... | 0.1 | * | * | * | * | *.1 | * | * | * | * | 1.0 | 4.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 3.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 3.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 3.2 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 2.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.2 | * | * | * | 0.2 | * | 0.8 | 4.4 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | * | * | 0.3 | 0.8 | 4.5 |
| 2002..... | 0.3 | * | * | * | * | 0.2 | 0.1 | * | * | * | 0.7 | 4.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.8 | 4.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.8 | 4.3 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | 0.8 | 5.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| VI. Diseases of the nervous system (G00-G98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 38.1 | 9.2 | 2.0 | 1.3 | 2.0 | 2.3 | 4.1 | 9.2 | 20.1 | 61.3 | 282.8 | 1,014.7 |
| 2003..... | 37.7 | 10.0 | 1.8 | 1.3 | 2.2 | 2.3 | 4.2 | 9.3 | 20.3 | 64.2 | 280.2 | 1,006.7 |
| 2002..... | 35.7 | 9.5 | 2.1 | 1.2 | 2.0 | 2.2 | 4.2 | 9.1 | 19.7 | 62.0 | 270.2 | 951.0 |
| 2001..... | 33.9 | 8.9 | 2.1 | 1.2 | 2.1 | 2.2 | 4.1 | 8.7 | 19.4 | 60.2 | 258.9 | 906.2 |
| 2000..... | 32.3 | 10.2 | 2.1 | 1.3 | 1.9 | 2.1 | 4.1 | 9.1 | 19.2 | 59.1 | 248.5 | 864.9 |
| 1999..... | 30.1 | 11.6 | 2.4 | 1.3 | 2.0 | 2.1 | 3.9 | 8.5 | 17.9 | 57.1 | 234.2 | 790.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 30.0 | 10.5 | 2.0 | 1.5 | 2.6 | 2.7 | 4.6 | 10.0 | 21.5 | 68.8 | 304.9 | 910.7 |
| 2003..... | 29.9 | 11.0 | 1.8 | 1.5 | 2.8 | 2.7 | 4.8 | 10.3 | 21.9 | 71.7 | 305.4 | 907.9 |
| 2002..... | 28.5 | 11.0 | 2.4 | 1.4 | 2.6 | 2.7 | 4.9 | 9.8 | 21.2 | 70.3 | 293.9 | 867.4 |
| 2001..... | 27.4 | 10.0 | 2.2 | 1.3 | 2.8 | 2.4 | 4.7 | 9.2 | 20.3 | 68.7 | 289.2 | 861.6 |
| 2000..... | 26.2 | 11.5 | 2.1 | 1.3 | 2.4 | 2.5 | 4.6 | 9.9 | 20.6 | 68.3 | 277.0 | 826.2 |
| 1999..... | 25.0 | 12.2 | 2.6 | 1.4 | 2.8 | 2.4 | 4.4 | 9.0 | 19.7 | 65.6 | 268.8 | 772.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 45.9 | 8.0 | 2.0 | 1.1 | 1.4 | 1.9 | 3.6 | 8.3 | 18.7 | 54.9 | 267.9 | 1,061.5 |
| 2003..... | 45.3 | 9.0 | 1.8 | 1.1 | 1.5 | 2.0 | 3.5 | 8.3 | 18.8 | 58.0 | 263.4 | 1,050.4 |
| 2002..... | 42.8 | 7.8 | 1.7 | 1.1 | 1.4 | 1.8 | 3.5 | 8.4 | 18.3 | 55.0 | 254.4 | 987.3 |
| 2001..... | 40.2 | 7.8 | 1.9 | 1.1 | 1.4 | 1.9 | 3.5 | 8.2 | 18.4 | 53.2 | 239.0 | 924.9 |
| 2000..... | 38.2 | 8.8 | 2.0 | 1.2 | 1.3 | 1.8 | 3.6 | 8.2 | 18.0 | 51.6 | 229.9 | 880.7 |
| 1999..... | 35.1 | 11.0 | 2.1 | 1.1 | 1.3 | 1.7 | 3.4 | 8.1 | 16.2 | 50.2 | 211.8 | 798.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 9.6 | 7.8 | 1.3 | 1.1 | 1.6 | 1.4 | 2.3 | 4.7 | 11.9 | 36.1 | 173.4 | 628.4 |
| 2003..... | 9.7 | 8.0 | 2.1 | 1.1 | 1.8 | 1.5 | 2.1 | 5.2 | 13.0 | 40.3 | 174.7 | 635.9 |
| 2002..... | 8.6 | 6.6 | 1.6 | 1.1 | 1.5 | 1.4 | 1.9 | 5.0 | 12.1 | 40.3 | 154.0 | 564.9 |
| 2001..... | 7.8 | 5.6 | 1.7 | 0.9 | 1.4 | 1.1 | 2.2 | 4.4 | 10.6 | 35.4 | 154.2 | 527.1 |
| 2000..... | 7.8 | 8.6 | 1.9 | 0.9 | 1.2 | 1.3 | 2.3 | 4.9 | 11.2 | 38.3 | 150.7 | 510.3 |
| 1999..... | 7.3 | 8.1 | 2.1 | 0.9 | 1.4 | 1.2 | 2.3 | 4.1 | 9.8 | 36.0 | 142.7 | 480.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 8.2 | 8.9 | 1.5 | 1.3 | 2.1 | 1.5 | 3.0 | 5.4 | 13.8 | 40.7 | 189.9 | 543.6 |
| 2003..... | 8.7 | 10.2 | 2.1 | 1.2 | 2.2 | 1.8 | 2.3 | 5.7 | 14.8 | 47.8 | 199.0 | 614.2 |
| 2002..... | 7.4 | 7.7 | 1.8 | 1.3 | 1.9 | 1.4 | 2.3 | 5.7 | 13.5 | 47.2 | 162.5 | 507.0 |
| 2001..... | 7.1 | 6.7 | 2.2 | 1.0 | 2.1 | 1.3 | 2.7 | 5.1 | 11.4 | 42.1 | 176.2 | 493.3 |
| 2000..... | 7.2 | 9.6 | 1.9 | 1.0 | 1.6 | 1.5 | 2.5 | 6.3 | 13.7 | 47.5 | 179.0 | 483.7 |
| 1999..... | 6.5 | 9.6 | 2.3 | 1.0 | 1.6 | 1.3 | 2.4 | 4.9 | 11.6 | 43.1 | 162.7 | 440.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 11.1 | 6.8 | * | 0.8 | 1.1 | 1.3 | 1.5 | 4.0 | 10.1 | 32.4 | 162.0 | 672.0 |
| 2003..... | 10.9 | 5.7 | 2.1 | 0.9 | 1.4 | 1.2 | 1.8 | 4.7 | 11.3 | 34.4 | 158.0 | 646.9 |
| 2002..... | 9.9 | 5.4 | 1.4 | 0.9 | 1.0 | 1.3 | 1.5 | 4.3 | 10.9 | 34.8 | 148.1 | 593.5 |
| 2001..... | 8.7 | * | * | 0.9 | 0.7 | 0.9 | 1.6 | 3.7 | 9.8 | 30.1 | 139.3 | 543.6 |
| 2000..... | 8.4 | 7.4 | 1.9 | 0.8 | 0.7 | 1.0 | 2.1 | 3.6 | 8.9 | 31.0 | 131.7 | 523.3 |
| 1999..... | 8.2 | 6.5 | 2.0 | 0.8 | 1.2 | 1.0 | 2.1 | 3.3 | 8.3 | 30.5 | 129.4 | 500.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| VI. Diseases of the nervous system (G00-G98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 42.6 | 9.6 | 2.2 | 1.3 | 2.1 | 2.5 | 4.4 | 9.6 | 20.7 | 63.0 | 288.5 | 1,030.8 |
| 2003..... | 42.1 | 10.5 | 1.7 | 1.4 | 2.2 | 2.5 | 4.5 | 9.6 | 20.9 | 65.8 | 285.2 | 1,020.6 |
| 2002..... | 39.8 | 10.2 | 2.2 | 1.2 | 2.1 | 2.4 | 4.5 | 9.4 | 20.2 | 63.3 | 275.2 | 964.4 |
| 2001..... | 37.7 | 9.7 | 2.1 | 1.2 | 2.2 | 2.4 | 4.4 | 9.1 | 20.0 | 61.7 | 263.2 | 918.3 |
| 2000..... | 35.8 | 10.6 | 2.1 | 1.3 | 2.0 | 2.3 | 4.3 | 9.4 | 19.8 | 60.3 | 252.2 | 875.9 |
| 1999..... | 33.2 | 12.3 | 2.4 | 1.3 | 2.2 | 2.2 | 4.1 | 8.9 | 18.4 | 58.3 | 237.5 | 800.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 33.7 | 10.8 | 2.2 | 1.5 | 2.7 | 3.0 | 4.9 | 10.5 | 22.1 | 70.8 | 310.9 | 927.8 |
| 2003..... | 33.4 | 11.0 | 1.7 | 1.6 | 2.9 | 2.9 | 5.2 | 10.7 | 22.5 | 73.2 | 310.4 | 919.8 |
| 2002..... | 31.9 | 11.9 | 2.6 | 1.4 | 2.7 | 3.0 | 5.2 | 10.1 | 21.7 | 71.7 | 299.8 | 881.9 |
| 2001..... | 30.5 | 10.7 | 2.2 | 1.3 | 2.9 | 2.6 | 4.9 | 9.6 | 20.9 | 70.3 | 293.9 | 874.7 |
| 2000..... | 29.1 | 12.0 | 2.2 | 1.4 | 2.5 | 2.6 | 4.8 | 10.2 | 21.1 | 69.3 | 280.7 | 838.8 |
| 1999..... | 27.6 | 12.8 | 2.6 | 1.4 | 3.0 | 2.6 | 4.7 | 9.3 | 20.2 | 66.8 | 272.8 | 784.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 51.2 | 8.3 | 2.3 | 1.2 | 1.5 | 2.1 | 3.9 | 8.8 | 19.4 | 56.5 | 273.4 | 1,076.9 |
| 2003..... | 50.3 | 9.9 | 1.7 | 1.2 | 1.5 | 2.2 | 3.8 | 8.6 | 19.4 | 59.6 | 268.4 | 1,064.9 |
| 2002..... | 47.4 | 8.4 | 1.8 | 1.1 | 1.5 | 1.8 | 3.8 | 8.7 | 18.9 | 56.3 | 258.9 | 1,000.0 |
| 2001..... | 44.5 | 8.6 | 2.0 | 1.2 | 1.5 | 2.2 | 3.8 | 8.6 | 19.1 | 54.6 | 243.0 | 936.5 |
| 2000..... | 42.1 | 9.1 | 2.0 | 1.2 | 1.5 | 1.9 | 3.8 | 8.6 | 18.6 | 52.8 | 233.6 | 891.0 |
| 1999..... | 38.5 | 11.9 | 2.2 | 1.2 | 1.3 | 1.8 | 3.5 | 8.5 | 16.8 | 51.3 | 214.6 | 806.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 48.9 | 9.5 | 1.9 | 1.2 | 1.9 | 2.3 | 4.3 | 9.5 | 20.9 | 65.5 | 299.7 | 1,068.8 |
| 2003..... | 48.3 | 9.6 | 1.4 | 1.2 | 2.2 | 2.4 | 4.3 | 9.4 | 21.0 | 68.2 | 297.2 | 1,062.7 |
| 2002..... | 45.8 | 10.1 | 1.9 | 1.1 | 1.9 | 2.3 | 4.4 | 9.2 | 20.5 | 65.6 | 287.5 | 1,004.6 |
| 2001..... | 43.1 | 8.6 | 1.9 | 1.2 | 2.1 | 2.2 | 4.2 | 8.9 | 20.1 | 64.1 | 273.5 | 955.6 |
| 2000..... | 40.8 | 9.5 | 1.7 | 1.2 | 2.0 | 2.2 | 4.1 | 9.2 | 20.0 | 62.1 | 262.3 | 912.9 |
| 1999..... | 37.8 | 11.0 | 2.1 | 1.2 | 2.0 | 2.1 | 3.9 | 8.6 | 18.4 | 59.9 | 246.9 | 835.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 38.4 | 11.1 | 1.9 | 1.4 | 2.5 | 2.9 | 4.9 | 10.4 | 22.2 | 73.5 | 323.2 | 964.4 |
| 2003..... | 38.0 | 11.3 | 1.3 | 1.5 | 3.0 | 2.7 | 5.1 | 10.5 | 22.5 | 75.6 | 322.7 | 957.6 |
| 2002..... | 36.2 | 11.9 | 2.2 | 1.2 | 2.6 | 2.9 | 5.1 | 9.8 | 22.0 | 73.8 | 312.5 | 921.0 |
| 2001..... | 34.6 | 9.3 | 2.1 | 1.3 | 2.8 | 2.3 | 4.7 | 9.3 | 21.1 | 72.6 | 305.4 | 911.0 |
| 2000..... | 32.8 | 10.6 | 1.9 | 1.3 | 2.6 | 2.6 | 4.5 | 10.1 | 21.0 | 71.5 | 291.5 | 873.6 |
| 1999..... | 31.2 | 12.2 | 2.5 | 1.2 | 2.9 | 2.5 | 4.4 | 8.9 | 20.2 | 68.7 | 284.2 | 819.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 59.1 | 7.8 | 1.9 | 1.0 | 1.3 | 1.7 | 3.7 | 8.6 | 19.6 | 58.7 | 283.6 | 1,115.5 |
| 2003..... | 58.3 | 7.9 | 1.5 | 1.0 | 1.4 | 2.1 | 3.5 | 8.3 | 19.7 | 61.8 | 280.0 | 1,108.9 |
| 2002..... | 55.0 | 8.1 | 1.5 | 1.0 | 1.3 | 1.6 | 3.6 | 8.6 | 19.1 | 58.6 | 270.7 | 1,040.8 |
| 2001..... | 51.2 | 7.8 | 1.6 | 1.1 | 1.4 | 2.0 | 3.6 | 8.5 | 19.1 | 56.9 | 252.3 | 974.2 |
| 2000..... | 48.5 | 8.4 | 1.5 | 1.1 | 1.3 | 1.7 | 3.6 | 8.2 | 19.0 | 54.2 | 243.1 | 928.7 |
| 1999..... | 44.1 | 9.7 | 1.7 | 1.2 | 1.1 | 1.7 | 3.4 | 8.2 | 16.7 | 52.4 | 222.5 | 842.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| VI. Diseases of the nervous system (G00-G98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 22.7 | 10.7 | 3.3 | 1.9 | 3.1 | 4.3 | 6.2 | 13.2 | 25.7 | 58.3 | 235.5 | 786.6 |
| 2003..... | 21.6 | 14.5 | 2.8 | 2.1 | 2.7 | 4.1 | 7.0 | 14.2 | 25.5 | 61.9 | 217.0 | 708.0 |
| 2002..... | 20.2 | 12.1 | 3.4 | 1.7 | 3.0 | 4.0 | 6.8 | 14.0 | 23.2 | 58.5 | 199.3 | 660.9 |
| 2001..... | 19.6 | 15.1 | 3.3 | 1.7 | 3.0 | 4.2 | 6.6 | 12.9 | 24.4 | 55.2 | 202.6 | 631.8 |
| 2000..... | 18.5 | 14.7 | 3.5 | 1.9 | 2.6 | 3.6 | 6.8 | 13.8 | 23.2 | 56.8 | 185.8 | 567.3 |
| 1999..... | 17.2 | 18.3 | 3.5 | 1.9 | 3.1 | 3.2 | 6.3 | 13.8 | 22.5 | 55.8 | 168.9 | 487.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 18.0 | 10.8 | 3.3 | 1.8 | 3.8 | 4.1 | 6.6 | 14.8 | 27.8 | 64.8 | 243.3 | 671.7 |
| 2003..... | 18.1 | 12.4 | 3.2 | 2.2 | 3.2 | 4.9 | 7.7 | 15.6 | 28.9 | 70.4 | 238.9 | 629.7 |
| 2002..... | 17.1 | 12.4 | 3.9 | 1.9 | 3.7 | 4.7 | 7.9 | 15.5 | 25.1 | 70.4 | 219.8 | 571.0 |
| 2001..... | 16.7 | 17.6 | 3.0 | 1.5 | 3.8 | 4.9 | 7.7 | 14.0 | 25.7 | 64.1 | 223.0 | 599.4 |
| 2000..... | 15.9 | 18.1 | 3.3 | 2.0 | 2.7 | 3.8 | 7.9 | 13.9 | 27.4 | 65.3 | 206.7 | 544.8 |
| 1999..... | 15.1 | 16.6 | 3.4 | 2.4 | 4.1 | 3.7 | 7.3 | 15.3 | 24.2 | 61.8 | 181.7 | 486.1 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 27.0 | 10.6 | 3.4 | 2.0 | 2.4 | 4.5 | 5.9 | 11.8 | 24.1 | 53.6 | 231.0 | 832.0 |
| 2003..... | 24.8 | 16.8 | 2.3 | 1.9 | 2.3 | 3.4 | 6.4 | 13.0 | 22.8 | 55.8 | 204.7 | 738.3 |
| 2002..... | 22.9 | 11.7 | 2.8 | 1.5 | 2.3 | 3.3 | 5.8 | 12.7 | 21.7 | 50.1 | 187.7 | 695.2 |
| 2001..... | 22.3 | 12.4 | 3.6 | 1.9 | 2.2 | 3.6 | 5.6 | 11.8 | 23.4 | 48.9 | 191.2 | 643.8 |
| 2000..... | 20.9 | 11.3 | 3.6 | 1.8 | 2.5 | 3.5 | 5.8 | 13.8 | 19.9 | 50.9 | 174.0 | 575.6 |
| 1999..... | 19.1 | 20.0 | 3.6 | 1.5 | 2.1 | 2.8 | 5.5 | 12.5 | 21.1 | 51.6 | 161.8 | 488.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Meningitis (G00,G03) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | 1.8 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.5 | 0.7 | 0.9 |
| 2003..... | 0.3 | 1.9 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.7 | 1.1 |
| 2002..... | 0.3 | 2.1 | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.3 | 0.6 | 0.6 | 1.2 |
| 2001..... | 0.3 | 2.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.5 | 0.7 | 0.9 |
| 2000..... | 0.3 | 2.7 | 0.3 | 0.1 | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 1.0 | 1.2 |
| 1999..... | 0.3 | 3.2 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 1.0 | 1.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | 1.8 | * | * | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.7 | * |
| 2003..... | 0.3 | 1.6 | * | * | * | 0.1 | 0.2 | 0.4 | 0.5 | 0.6 | 0.8 | * |
| 2002..... | 0.3 | 2.1 | * | * | * | * | 0.2 | 0.4 | 0.4 | 0.7 | 0.8 | * |
| 2001..... | 0.3 | 2.1 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.6 | 0.9 | * |
| 2000..... | 0.3 | 2.7 | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.2 | * |
| 1999..... | 0.3 | 3.1 | 0.3 | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.4 | 0.7 | 1.1 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | 1.9 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.6 | 0.7 |
| 2003..... | 0.2 | 2.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.0 |
| 2002..... | 0.2 | 2.0 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.5 | 1.2 |
| 2001..... | 0.2 | 2.3 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.4 | 0.6 | 0.7 |
| 2000..... | 0.3 | 2.7 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.1 |
| 1999..... | 0.3 | 3.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.5 | 1.0 | 1.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | 1.6 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 0.8 |
| 2003..... | 0.2 | 1.5 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.0 |
| 2002..... | 0.2 | 1.9 | * | 0.1 | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.5 | 1.2 |
| 2001..... | 0.2 | 1.6 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.5 | 0.7 | 0.7 |
| 2000..... | 0.2 | 2.1 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.1 |
| 1999..... | 0.3 | 2.2 | 0.3 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 1.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | 1.9 | * | * | * | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.7 | * |
| 2003..... | 0.3 | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.8 | * |
| 2002..... | 0.2 | 1.9 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 0.7 | * |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.4 | 0.5 | 0.8 | * |
| 2000..... | 0.3 | 2.3 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.1 | * |
| 1999..... | 0.3 | 2.8 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.9 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 0.5 | * |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.6 | 0.8 |
| 2002..... | 0.2 | 1.8 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.4 | 1.3 |
| 2001..... | 0.2 | 1.9 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.6 | * |
| 2000..... | 0.2 | 1.8 | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.8 | 1.0 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other inflammatory diseases of central nervous system (G04-G09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 1.2 |
| 2003..... | 0.3 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.7 | 1.2 | 1.0 |
| 2002..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.6 | 1.3 | 1.2 |
| 2001..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.0 | 1.0 |
| 2000..... | 0.2 | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 1.0 |
| 1999..... | 0.2 | * | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.5 | 1.6 |
| 2003..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.9 | 1.4 | * |
| 2002..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 1.7 | * |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.2 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.5 | * |
| 1999..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.6 | 1.2 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 1.0 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 0.9 |
| 2002..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 1.0 | 1.2 |
| 2001..... | 0.2 | * | * | * | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 0.7 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.6 | 0.6 | 0.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 1.2 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.2 | 1.1 |
| 2002..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 1.2 | 1.4 |
| 2001..... | 0.2 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 1.0 | 1.0 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 1.0 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 1.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.5 | * |
| 2003..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.4 | * |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.6 | * |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 1.9 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.5 | * |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.5 | 1.2 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 1.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 1.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 1.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | * |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | * | 0.6 | 0.6 | 0.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Parkinson's disease (G20-G21) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.1 | * | * | * | * | * | * | 0.2 | 1.2 | 12.0 | 67.5 | 135.8 |
| 2003..... | 6.2 | * | * | * | * | * | * | 0.2 | 1.3 | 12.7 | 67.8 | 138.2 |
| 2002..... | 5.9 | * | * | * | * | * | * | 0.1 | 1.2 | 12.2 | 63.9 | 135.2 |
| 2001..... | 5.8 | * | * | * | * | * | * | 0.1 | 1.2 | 11.7 | 64.6 | 134.2 |
| 2000..... | 5.6 | * | * | * | * | * | * | 0.1 | 1.1 | 11.5 | 61.9 | 131.9 |
| 1999..... | 5.2 | * | * | * | * | * | * | 0.1 | 1.0 | 11.0 | 58.2 | 124.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.2 | * | * | * | * | * | * | 0.2 | 1.6 | 17.3 | 101.7 | 220.0 |
| 2003..... | 7.1 | * | * | * | * | * | * | 0.2 | 1.6 | 18.2 | 100.7 | 222.4 |
| 2002..... | 6.8 | * | * | * | * | * | * | 0.2 | 1.5 | 17.9 | 94.7 | 220.1 |
| 2001..... | 6.7 | * | * | * | * | * | * | 0.2 | 1.8 | 16.8 | 97.0 | 224.4 |
| 2000..... | 6.4 | * | * | * | * | * | * | 0.2 | 1.5 | 16.1 | 92.8 | 222.3 |
| 1999..... | 6.0 | * | * | * | * | * | * | 0.2 | 1.4 | 15.7 | 89.5 | 205.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | * | * | * | * | 0.8 | 7.5 | 44.4 | 97.9 |
| 2003..... | 5.3 | * | * | * | * | * | * | 0.1 | 0.9 | 8.1 | 45.8 | 101.0 |
| 2002..... | 5.0 | * | * | * | * | * | * | * | 0.9 | 7.4 | 43.5 | 98.4 |
| 2001..... | 4.9 | * | * | * | * | * | * | * | 0.7 | 7.5 | 43.3 | 96.4 |
| 2000..... | 4.8 | * | * | * | * | * | * | 0.1 | 0.8 | 7.6 | 41.7 | 95.0 |
| 1999..... | 4.5 | * | * | * | * | * | * | * | 0.7 | 7.1 | 37.8 | 91.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | * | 1.0 | 6.8 | 44.1 | 85.1 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | * | 7.9 | 40.3 | 77.7 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | 1.1 | 8.6 | 37.2 | 76.5 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | * | 8.6 | 41.5 | 86.3 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 7.8 | 37.7 | 86.3 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 8.1 | 38.3 | 77.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | * | * | 9.5 | 66.5 | 126.1 |
| 2003..... | 1.5 | * | * | * | * | * | * | * | * | 12.4 | 58.3 | 129.3 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | * | 13.6 | 50.1 | 113.8 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | * | 11.2 | 63.0 | 127.6 |
| 2000..... | 1.3 | * | * | * | * | * | * | * | * | 9.7 | 55.1 | 143.1 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 11.4 | 58.9 | 105.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | * | * | 4.7 | 28.6 | 64.1 |
| 2003..... | 1.1 | * | * | * | * | * | * | * | * | 4.3 | 27.9 | 51.6 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | * | 4.7 | 28.4 | 58.0 |
| 2001..... | 1.2 | * | * | * | * | * | * | * | * | 6.6 | 26.9 | 66.1 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | * | 6.3 | 26.1 | 58.4 |
| 1999..... | 1.0 | * | * | * | * | * | * | * | * | 5.5 | 24.6 | 63.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Parkinson's disease (G20-G21) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | * | * | * | * | 0.2 | 1.2 | 12.4 | 68.6 | 137.9 |
| 2003..... | 7.0 | * | * | * | * | * | * | 0.2 | 1.3 | 13.0 | 69.1 | 140.6 |
| 2002..... | 6.6 | * | * | * | * | * | * | 0.1 | 1.2 | 12.4 | 65.1 | 137.4 |
| 2001..... | 6.5 | * | * | * | * | * | * | 0.1 | 1.2 | 11.9 | 65.5 | 135.6 |
| 2000..... | 6.2 | * | * | * | * | * | * | 0.1 | 1.1 | 11.7 | 62.8 | 133.3 |
| 1999..... | 5.8 | * | * | * | * | * | * | 0.1 | 1.1 | 11.1 | 58.9 | 125.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 8.1 | * | * | * | * | * | * | 0.3 | 1.6 | 17.8 | 103.4 | 224.4 |
| 2003..... | 8.0 | * | * | * | * | * | * | 0.2 | 1.7 | 18.5 | 102.7 | 226.5 |
| 2002..... | 7.6 | * | * | * | * | * | * | 0.2 | 1.5 | 18.2 | 96.7 | 224.4 |
| 2001..... | 7.5 | * | * | * | * | * | * | 0.2 | 1.8 | 17.1 | 98.3 | 227.8 |
| 2000..... | 7.1 | * | * | * | * | * | * | 0.2 | 1.5 | 16.5 | 94.2 | 225.2 |
| 1999..... | 6.7 | * | * | * | * | * | * | 0.2 | 1.5 | 15.9 | 90.6 | 209.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.7 | * | * | * | * | * | * | * | 0.8 | 7.8 | 45.3 | 99.2 |
| 2003..... | 5.9 | * | * | * | * | * | * | 0.2 | 0.9 | 8.4 | 46.6 | 102.8 |
| 2002..... | 5.6 | * | * | * | * | * | * | * | 0.9 | 7.6 | 44.2 | 99.8 |
| 2001..... | 5.4 | * | * | * | * | * | * | * | 0.7 | 7.6 | 44.0 | 97.3 |
| 2000..... | 5.3 | * | * | * | * | * | * | * | 0.8 | 7.7 | 42.3 | 96.1 |
| 1999..... | 4.9 | * | * | * | * | * | * | * | 0.7 | 7.1 | 38.3 | 92.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 8.2 | * | * | * | * | * | * | 0.2 | 1.3 | 13.5 | 73.3 | 145.5 |
| 2003..... | 8.3 | * | * | * | * | * | * | 0.2 | 1.3 | 14.0 | 73.7 | 148.8 |
| 2002..... | 7.9 | * | * | * | * | * | * | 0.2 | 1.3 | 13.3 | 69.6 | 145.8 |
| 2001..... | 7.7 | * | * | * | * | * | * | 0.1 | 1.3 | 12.8 | 69.9 | 143.5 |
| 2000..... | 7.3 | * | * | * | * | * | * | 0.1 | 1.2 | 12.6 | 66.7 | 141.2 |
| 1999..... | 6.8 | * | * | * | * | * | * | 0.1 | 1.1 | 11.9 | 62.4 | 132.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 9.7 | * | * | * | * | * | * | 0.3 | 1.7 | 19.2 | 110.0 | 236.3 |
| 2003..... | 9.5 | * | * | * | * | * | * | 0.2 | 1.7 | 19.8 | 109.0 | 237.9 |
| 2002..... | 9.1 | * | * | * | * | * | * | 0.2 | 1.6 | 19.3 | 102.9 | 237.8 |
| 2001..... | 9.0 | * | * | * | * | * | * | 0.2 | 2.0 | 18.3 | 104.5 | 241.7 |
| 2000..... | 8.4 | * | * | * | * | * | * | 0.2 | 1.6 | 17.6 | 99.2 | 238.6 |
| 1999..... | 7.9 | * | * | * | * | * | * | 0.2 | 1.6 | 17.1 | 95.7 | 221.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | * | * | * | * | * | 0.9 | 8.5 | 48.3 | 104.8 |
| 2003..... | 7.1 | * | * | * | * | * | * | 0.2 | 1.0 | 9.0 | 49.8 | 109.6 |
| 2002..... | 6.7 | * | * | * | * | * | * | * | 1.0 | 8.1 | 47.3 | 106.0 |
| 2001..... | 6.5 | * | * | * | * | * | * | * | 0.7 | 8.2 | 46.9 | 102.7 |
| 2000..... | 6.3 | * | * | * | * | * | * | * | 0.9 | 8.3 | 45.3 | 101.9 |
| 1999..... | 5.8 | * | * | * | * | * | * | * | 0.7 | 7.6 | 40.5 | 97.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Parkinson's disease (G20-G21) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | * | 0.7 | 5.5 | 29.8 | 61.6 |
| 2003..... | 1.7 | * | * | * | * | * | * | * | 1.2 | 7.2 | 29.4 | 53.5 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | * | 6.7 | 25.7 | 51.7 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 0.8 | 5.6 | 25.7 | 59.0 |
| 2000..... | 1.5 | * | * | * | * | * | * | * | * | 5.8 | 27.3 | 52.5 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | 0.9 | 5.9 | 24.8 | 51.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | * | * | 8.6 | 44.5 | 106.7 |
| 2003..... | 2.0 | * | * | * | * | * | * | * | 1.8 | 10.6 | 45.6 | 99.1 |
| 2002..... | 1.8 | * | * | * | * | * | * | * | * | 11.1 | 40.4 | 87.5 |
| 2001..... | 1.7 | * | * | * | * | * | * | * | * | 8.8 | 38.1 | 99.3 |
| 2000..... | 1.8 | * | * | * | * | * | * | * | * | 9.0 | 47.5 | 88.1 |
| 1999..... | 1.6 | * | * | * | * | * | * | * | * | 8.7 | 38.5 | 87.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | * | * | 3.3 | 21.6 | 43.8 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | * | 4.8 | 20.2 | 35.9 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | * | 3.6 | 17.4 | 38.0 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | * | 3.3 | 18.7 | 44.1 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 3.5 | 16.0 | 39.2 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 3.9 | 17.2 | 38.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Alzheimer's disease (G30) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 22.5 | * | * | * | * | * | * | 0.2 | 1.9 | 19.7 | 168.7 | 818.8 |
| 2003..... | 21.8 | * | * | * | * | * | * | 0.2 | 2.0 | 20.9 | 164.4 | 802.4 |
| 2002..... | 20.4 | * | * | * | * | * | * | 0.1 | 1.9 | 19.7 | 158.1 | 752.3 |
| 2001..... | 18.9 | * | * | * | * | * | * | 0.2 | 2.1 | 18.7 | 147.5 | 710.3 |
| 2000..... | 17.6 | * | * | * | * | * | * | 0.2 | 2.0 | 18.7 | 139.6 | 667.7 |
| 1999..... | 16.0 | * | * | * | * | * | * | 0.2 | 1.9 | 17.4 | 129.5 | 601.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 13.1 | * | * | * | * | * | * | 0.2 | 1.8 | 18.5 | 149.0 | 619.6 |
| 2003..... | 12.8 | * | * | * | * | * | * | 0.2 | 1.8 | 20.2 | 147.6 | 606.5 |
| 2002..... | 12.0 | * | * | * | * | * | * | 0.1 | 1.8 | 18.6 | 142.2 | 572.0 |
| 2001..... | 11.3 | * | * | * | * | * | * | 0.2 | 1.9 | 17.7 | 136.3 | 556.1 |
| 2000..... | 10.5 | * | * | * | * | * | * | 0.2 | 1.8 | 18.7 | 127.6 | 522.1 |
| 1999..... | 9.8 | * | * | * | * | * | * | 0.2 | 1.9 | 17.3 | 122.6 | 485.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 31.5 | * | * | * | * | * | * | 0.3 | 1.9 | 20.8 | 181.9 | 908.4 |
| 2003..... | 30.5 | * | * | * | * | * | * | 0.2 | 2.1 | 21.5 | 175.7 | 889.0 |
| 2002..... | 28.5 | * | * | * | * | * | * | 0.1 | 2.0 | 20.6 | 168.7 | 830.5 |
| 2001..... | 26.3 | * | * | * | * | * | * | 0.2 | 2.3 | 19.5 | 154.8 | 774.9 |
| 2000..... | 24.5 | * | * | * | * | * | * | 0.2 | 2.2 | 18.6 | 147.4 | 727.0 |
| 1999..... | 21.9 | * | * | * | * | * | * | 0.2 | 1.8 | 17.5 | 134.0 | 648.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | * | * | 1.0 | 12.0 | 99.5 | 490.9 |
| 2003..... | 4.6 | * | * | * | * | * | * | * | 1.3 | 14.6 | 103.7 | 493.4 |
| 2002..... | 3.8 | * | * | * | * | * | * | * | 1.0 | 10.8 | 89.2 | 438.5 |
| 2001..... | 3.4 | * | * | * | * | * | * | * | * | 9.9 | 83.9 | 399.5 |
| 2000..... | 3.2 | * | * | * | * | * | * | * | * | 11.9 | 79.2 | 380.2 |
| 1999..... | 2.9 | * | * | * | * | * | * | * | * | 9.7 | 75.4 | 351.5 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | * | * | * | 11.5 | 93.0 | 374.1 |
| 2003..... | 3.1 | * | * | * | * | * | * | * | * | 13.8 | 101.2 | 426.4 |
| 2002..... | 2.3 | * | * | * | * | * | * | * | * | 10.5 | 80.4 | 341.3 |
| 2001..... | 2.1 | * | * | * | * | * | * | * | * | 8.8 | 83.2 | 306.6 |
| 2000..... | 2.1 | * | * | * | * | * | * | * | * | 13.3 | 81.6 | 284.2 |
| 1999..... | 1.8 | * | * | * | * | * | * | * | * | 11.2 | 66.7 | 278.2 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | * | * | * | 12.5 | 104.0 | 551.0 |
| 2003..... | 6.2 | * | * | * | * | * | * | * | * | 15.3 | 105.3 | 527.3 |
| 2002..... | 5.4 | * | * | * | * | * | * | * | * | 10.9 | 95.3 | 486.6 |
| 2001..... | 4.7 | * | * | * | * | * | * | * | * | 10.7 | 84.3 | 444.9 |
| 2000..... | 4.3 | * | * | * | * | * | * | * | * | 10.8 | 77.6 | 427.3 |
| 1999..... | 4.0 | * | * | * | * | * | * | * | * | 8.4 | 81.2 | 386.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Alzheimer's disease (G30) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 25.4 | * | * | * | * | * | * | 0.2 | 1.9 | 20.2 | 172.3 | 832.6 |
| 2003..... | 24.5 | * | * | * | * | * | * | 0.2 | 2.0 | 21.3 | 167.3 | 814.0 |
| 2002..... | 22.9 | * | * | * | * | * | * | 0.1 | 2.0 | 20.3 | 161.1 | 763.2 |
| 2001..... | 21.2 | * | * | * | * | * | * | 0.2 | 2.2 | 19.2 | 150.2 | 720.5 |
| 2000..... | 19.6 | * | * | * | * | * | * | 0.2 | 2.1 | 19.0 | 142.0 | 676.7 |
| 1999..... | 17.7 | * | * | * | * | * | * | 0.2 | 2.0 | 17.8 | 131.5 | 608.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 14.9 | * | * | * | * | * | * | 0.2 | 1.9 | 19.0 | 152.0 | 631.2 |
| 2003..... | 14.4 | * | * | * | * | * | * | 0.2 | 1.9 | 20.5 | 149.8 | 613.6 |
| 2002..... | 13.6 | * | * | * | * | * | * | 0.2 | 1.9 | 19.1 | 145.0 | 581.4 |
| 2001..... | 12.7 | * | * | * | * | * | * | 0.2 | 2.0 | 18.3 | 138.6 | 565.2 |
| 2000..... | 11.7 | * | * | * | * | * | * | 0.2 | 1.8 | 19.0 | 129.6 | 531.0 |
| 1999..... | 10.9 | * | * | * | * | * | * | 0.2 | 2.0 | 17.7 | 124.8 | 492.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 35.3 | * | * | * | * | * | * | 0.3 | 2.0 | 21.3 | 186.0 | 922.7 |
| 2003..... | 34.1 | * | * | * | * | * | * | 0.2 | 2.2 | 22.0 | 179.0 | 902.1 |
| 2002..... | 31.9 | * | * | * | * | * | * | 0.1 | 2.1 | 21.2 | 171.8 | 841.6 |
| 2001..... | 29.3 | * | * | * | * | * | * | 0.2 | 2.4 | 20.0 | 157.8 | 785.1 |
| 2000..... | 27.2 | * | * | * | * | * | * | 0.2 | 2.3 | 19.1 | 150.0 | 735.6 |
| 1999..... | 24.2 | * | * | * | * | * | * | 0.2 | 1.9 | 18.0 | 135.9 | 655.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 29.6 | * | * | * | * | * | * | 0.3 | 2.1 | 21.1 | 178.1 | 861.9 |
| 2003..... | 28.7 | * | * | * | * | * | * | 0.2 | 2.2 | 22.2 | 173.9 | 846.9 |
| 2002..... | 26.9 | * | * | * | * | * | * | 0.2 | 2.1 | 21.3 | 168.2 | 793.9 |
| 2001..... | 24.8 | * | * | * | * | * | * | 0.2 | 2.4 | 20.3 | 155.6 | 749.2 |
| 2000..... | 22.9 | * | * | * | * | * | * | 0.2 | 2.2 | 19.9 | 147.4 | 705.1 |
| 1999..... | 20.7 | * | * | * | * | * | * | 0.2 | 2.1 | 18.6 | 136.7 | 636.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 17.3 | * | * | * | * | * | * | 0.2 | 2.0 | 19.6 | 157.0 | 653.3 |
| 2003..... | 16.8 | * | * | * | * | * | * | 0.2 | 2.0 | 21.2 | 154.8 | 638.1 |
| 2002..... | 15.8 | * | * | * | * | * | * | 0.2 | 2.0 | 19.9 | 150.5 | 605.4 |
| 2001..... | 14.7 | * | * | * | * | * | * | 0.2 | 2.1 | 19.1 | 143.0 | 587.2 |
| 2000..... | 13.6 | * | * | * | * | * | * | 0.2 | 2.0 | 19.8 | 134.6 | 551.7 |
| 1999..... | 12.7 | * | * | * | * | * | * | 0.2 | 2.2 | 18.3 | 129.8 | 515.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 41.4 | * | * | * | * | * | * | 0.4 | 2.2 | 22.3 | 192.5 | 955.3 |
| 2003..... | 40.2 | * | * | * | * | * | * | 0.2 | 2.5 | 23.0 | 186.8 | 938.8 |
| 2002..... | 37.6 | * | * | * | * | * | * | 0.2 | 2.3 | 22.5 | 180.1 | 875.3 |
| 2001..... | 34.4 | * | * | * | * | * | * | 0.2 | 2.7 | 21.3 | 163.9 | 816.6 |
| 2000..... | 31.9 | * | * | * | * | * | * | 0.2 | 2.5 | 20.0 | 155.8 | 767.0 |
| 1999..... | 28.4 | * | * | * | * | * | * | 0.2 | 2.0 | 18.8 | 141.2 | 685.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Alzheimer's disease (G30) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 11.4 | * | * | * | * | * | * | * | 1.3 | 19.4 | 155.4 | 666.9 |
| 2003..... | 10.1 | * | * | * | * | * | * | * | 1.1 | 19.6 | 140.3 | 591.1 |
| 2002..... | 9.1 | * | * | * | * | * | * | * | 1.2 | 16.8 | 124.2 | 548.7 |
| 2001..... | 8.7 | * | * | * | * | * | * | * | 1.1 | 15.5 | 126.9 | 513.2 |
| 2000..... | 7.7 | * | * | * | * | * | * | * | 1.3 | 16.1 | 111.6 | 450.9 |
| 1999..... | 6.7 | * | * | * | * | * | * | * | 1.2 | 15.5 | 99.1 | 374.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | * | * | * | 19.4 | 138.6 | 507.0 |
| 2003..... | 6.2 | * | * | * | * | * | * | * | * | 20.5 | 138.8 | 465.6 |
| 2002..... | 5.5 | * | * | * | * | * | * | * | * | 17.9 | 122.5 | 414.6 |
| 2001..... | 5.3 | * | * | * | * | * | * | * | * | 16.3 | 123.6 | 420.4 |
| 2000..... | 4.6 | * | * | * | * | * | * | * | * | 16.9 | 98.7 | 378.1 |
| 1999..... | 4.2 | * | * | * | * | * | * | * | * | 16.8 | 90.7 | 320.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 15.8 | * | * | * | * | * | * | * | 1.4 | 19.4 | 164.9 | 730.0 |
| 2003..... | 13.7 | * | * | * | * | * | * | * | * | 18.9 | 141.1 | 639.7 |
| 2002..... | 12.5 | * | * | * | * | * | * | * | * | 16.1 | 125.2 | 599.9 |
| 2001..... | 11.8 | * | * | * | * | * | * | * | * | 15.0 | 128.7 | 547.5 |
| 2000..... | 10.6 | * | * | * | * | * | * | * | * | 15.6 | 118.8 | 478.0 |
| 1999..... | 9.0 | * | * | * | * | * | * | * | * | 14.5 | 103.8 | 394.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Multiple sclerosis (G35) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.1 | 0.6 | 2.0 | 3.6 | 4.0 | 3.7 | 2.7 |
| 2003..... | 1.2 | * | * | * | * | 0.1 | 0.7 | 2.1 | 3.3 | 3.8 | 3.7 | 2.3 |
| 2002..... | 1.2 | * | * | * | * | 0.1 | 0.7 | 2.1 | 3.4 | 3.6 | 3.5 | 2.3 |
| 2001..... | 1.2 | * | * | * | * | 0.1 | 0.7 | 2.1 | 3.3 | 3.5 | 3.6 | 2.1 |
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.7 | 2.0 | 3.2 | 3.2 | 3.2 | 1.8 |
| 1999..... | 1.1 | * | * | * | * | 0.1 | 0.7 | 2.0 | 2.7 | 3.0 | 3.1 | 2.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.4 | 1.4 | 2.5 | 3.1 | 2.8 | 1.7 |
| 2003..... | 0.9 | * | * | * | * | * | 0.5 | 1.5 | 2.6 | 2.9 | 2.7 | 1.5 |
| 2002..... | 0.9 | * | * | * | * | * | 0.5 | 1.4 | 2.7 | 2.6 | 2.7 | 2.0 |
| 2001..... | 0.8 | * | * | * | * | * | 0.5 | 1.5 | 2.4 | 2.8 | 2.8 | 1.6 |
| 2000..... | 0.8 | * | * | * | * | * | 0.6 | 1.5 | 2.3 | 2.4 | 2.4 | 1.9 |
| 1999..... | 0.7 | * | * | * | * | * | 0.5 | 1.3 | 1.9 | 2.5 | 2.3 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.2 | 0.8 | 2.6 | 4.5 | 4.8 | 4.4 | 3.2 |
| 2003..... | 1.6 | * | * | * | * | 0.2 | 0.8 | 2.6 | 4.0 | 4.5 | 4.4 | 2.6 |
| 2002..... | 1.6 | * | * | * | * | * | 0.9 | 2.8 | 4.0 | 4.4 | 4.0 | 2.4 |
| 2001..... | 1.5 | * | * | * | * | 0.2 | 1.0 | 2.6 | 4.2 | 4.1 | 4.1 | 2.4 |
| 2000..... | 1.4 | * | * | * | * | 0.2 | 0.8 | 2.5 | 4.0 | 3.8 | 3.7 | 1.8 |
| 1999..... | 1.4 | * | * | * | * | 0.2 | 0.9 | 2.7 | 3.4 | 3.4 | 3.6 | 2.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.1 | 0.6 | 2.1 | 3.7 | 4.4 | 4.1 | 3.0 |
| 2003..... | 1.4 | * | * | * | * | 0.1 | 0.6 | 2.1 | 3.6 | 4.0 | 3.9 | 2.4 |
| 2002..... | 1.4 | * | * | * | * | * | 0.7 | 2.2 | 3.6 | 3.9 | 3.7 | 2.5 |
| 2001..... | 1.3 | * | * | * | * | 0.1 | 0.7 | 2.2 | 3.5 | 3.8 | 3.9 | 2.3 |
| 2000..... | 1.2 | * | * | * | * | 0.1 | 0.7 | 2.1 | 3.3 | 3.4 | 3.4 | 2.0 |
| 1999..... | 1.2 | * | * | * | * | 0.1 | 0.7 | 2.1 | 2.9 | 3.2 | 3.4 | 2.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.4 | 1.5 | 2.7 | 3.3 | 2.8 | 1.9 |
| 2003..... | 1.0 | * | * | * | * | * | 0.5 | 1.5 | 2.8 | 3.1 | 2.9 | * |
| 2002..... | 1.0 | * | * | * | * | * | 0.5 | 1.5 | 3.0 | 2.8 | 2.9 | 2.2 |
| 2001..... | 1.0 | * | * | * | * | * | 0.5 | 1.6 | 2.6 | 3.1 | 3.0 | 1.8 |
| 2000..... | 0.9 | * | * | * | * | * | 0.5 | 1.6 | 2.4 | 2.7 | 2.6 | 2.1 |
| 1999..... | 0.8 | * | * | * | * | * | 0.5 | 1.4 | 2.1 | 2.7 | 2.5 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.8 | 2.7 | 4.7 | 5.3 | 4.9 | 3.4 |
| 2003..... | 1.8 | * | * | * | * | * | 0.8 | 2.7 | 4.3 | 4.8 | 4.6 | 2.8 |
| 2002..... | 1.7 | * | * | * | * | * | 0.8 | 2.8 | 4.2 | 4.8 | 4.3 | 2.6 |
| 2001..... | 1.7 | * | * | * | * | 0.2 | 0.9 | 2.7 | 4.4 | 4.4 | 4.4 | 2.5 |
| 2000..... | 1.6 | * | * | * | * | 0.2 | 0.8 | 2.6 | 4.2 | 4.1 | 4.0 | 1.9 |
| 1999..... | 1.5 | * | * | * | * | * | 0.9 | 2.8 | 3.7 | 3.6 | 4.0 | 2.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Multiple sclerosis (G35) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.9 | 2.0 | 3.8 | 2.6 | * | * |
| 2003..... | 1.0 | * | * | * | * | 0.4 | 1.0 | 2.5 | 2.7 | 3.1 | 2.5 | * |
| 2002..... | 0.9 | * | * | * | * | * | 1.1 | 2.7 | 2.9 | 2.5 | * | * |
| 2001..... | 0.8 | * | * | * | * | * | 1.1 | 2.2 | 2.8 | 2.2 | * | * |
| 2000..... | 0.9 | * | * | * | * | * | 1.2 | 2.3 | 3.4 | 2.0 | * | * |
| 1999..... | 0.7 | * | * | * | * | * | 1.0 | 2.0 | 1.9 | 2.4 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 1.3 | 1.8 | 2.9 | * | * |
| 2003..... | 0.7 | * | * | * | * | * | * | 1.8 | 2.0 | * | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | 1.6 | 1.8 | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | 1.4 | * | * | * | * |
| 2000..... | 0.6 | * | * | * | * | * | 1.0 | 1.3 | 2.9 | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | 1.5 | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 1.2 | 2.6 | 5.4 | 2.3 | * | * |
| 2003..... | 1.2 | * | * | * | * | * | 1.5 | 3.1 | 3.2 | 3.8 | * | * |
| 2002..... | 1.3 | * | * | * | * | * | 1.6 | 3.6 | 3.9 | 3.4 | * | * |
| 2001..... | 1.2 | * | * | * | * | * | 1.6 | 2.9 | 3.8 | 2.7 | * | * |
| 2000..... | 1.1 | * | * | * | * | * | 1.4 | 3.2 | 3.7 | 2.7 | * | * |
| 1999..... | 0.9 | * | * | * | * | * | 1.4 | 2.3 | 2.2 | 3.2 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Epilepsy (G40-G41) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | 0.2 | 0.1 | 0.2 | 0.5 | 0.5 | 0.6 | 0.6 | 0.7 | 1.1 | 1.5 |
| 2003..... | 0.5 | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.7 | 0.6 | 0.8 | 1.2 | 1.5 |
| 2002..... | 0.5 | * | * | 0.1 | 0.3 | 0.4 | 0.6 | 0.6 | 0.5 | 0.8 | 1.3 | 1.7 |
| 2001..... | 0.5 | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.5 | 0.6 | 0.7 | 1.2 | 1.6 |
| 2000..... | 0.5 | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.6 | 0.6 | 0.8 | 1.2 | 2.0 |
| 1999..... | 0.5 | * | 0.2 | 0.1 | 0.3 | 0.5 | 0.6 | 0.6 | 0.6 | 0.9 | 1.2 | 1.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.3 | 0.6 | 0.7 | 0.8 | 0.6 | 0.6 | 1.1 | 1.9 |
| 2003..... | 0.5 | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.8 | 0.7 | 0.8 | 1.4 | * |
| 2002..... | 0.6 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 0.8 | 0.6 | 0.8 | 1.3 | 2.3 |
| 2001..... | 0.5 | * | * | 0.1 | 0.3 | 0.5 | 0.8 | 0.6 | 0.7 | 0.8 | 0.9 | 2.0 |
| 2000..... | 0.5 | * | * | * | 0.3 | 0.5 | 0.6 | 0.7 | 0.8 | 0.9 | 0.9 | 2.4 |
| 1999..... | 0.6 | * | * | 0.1 | 0.4 | 0.6 | 0.8 | 0.7 | 0.6 | 1.0 | 1.0 | 2.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.6 | 0.7 | 1.1 | 1.4 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.5 | 0.5 | 0.6 | 0.8 | 1.1 | 1.7 |
| 2002..... | 0.5 | * | * | * | 0.3 | 0.3 | 0.4 | 0.5 | 0.5 | 0.8 | 1.4 | 1.5 |
| 2001..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.4 | 0.4 | 0.5 | 0.7 | 1.4 | 1.5 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.5 | 0.4 | 0.5 | 0.8 | 1.4 | 1.9 |
| 1999..... | 0.4 | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | 0.8 | 1.3 | 1.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.1 | 0.2 | 0.4 | 0.5 | 0.6 | 0.5 | 0.6 | 1.0 | 1.4 |
| 2003..... | 0.5 | * | * | 0.1 | 0.3 | 0.4 | 0.5 | 0.6 | 0.5 | 0.6 | 1.0 | 1.4 |
| 2002..... | 0.5 | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.5 | 0.5 | 0.6 | 1.2 | 1.6 |
| 2001..... | 0.4 | * | * | 0.1 | 0.2 | 0.4 | 0.5 | 0.5 | 0.5 | 0.6 | 0.9 | 1.6 |
| 2000..... | 0.4 | * | * | 0.1 | 0.2 | 0.4 | 0.5 | 0.5 | 0.5 | 0.7 | 1.0 | 1.7 |
| 1999..... | 0.4 | * | 0.2 | 0.1 | 0.2 | 0.4 | 0.5 | 0.5 | 0.4 | 0.7 | 1.1 | 1.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.2 | 0.6 | 0.6 | 0.8 | 0.5 | 0.5 | 0.9 | 1.8 |
| 2003..... | 0.5 | * | * | * | 0.3 | 0.5 | 0.6 | 0.7 | 0.5 | 0.5 | 1.0 | * |
| 2002..... | 0.5 | * | * | * | 0.2 | 0.5 | 0.7 | 0.6 | 0.5 | 0.7 | 1.0 | 2.2 |
| 2001..... | 0.5 | * | * | * | 0.3 | 0.4 | 0.6 | 0.6 | 0.6 | 0.6 | 0.7 | 2.0 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.6 | 0.7 | 0.6 | 0.7 | 0.6 | * |
| 1999..... | 0.5 | * | * | * | 0.3 | 0.5 | 0.6 | 0.6 | 0.4 | 0.8 | 0.9 | 2.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 1.0 | 1.2 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | 0.7 | 1.0 | 1.5 |
| 2002..... | 0.4 | * | * | * | 0.3 | 0.3 | 0.4 | 0.4 | 0.5 | 0.6 | 1.3 | 1.3 |
| 2001..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.4 | 0.4 | 0.4 | 0.6 | 1.1 | 1.4 |
| 2000..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.5 | 0.3 | 0.4 | 0.7 | 1.3 | 1.7 |
| 1999..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.4 | 0.6 | 1.2 | 1.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Epilepsy (G40-G41) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.4 | 0.9 | 0.9 | 1.0 | 1.7 | 1.5 | 2.9 | * |
| 2003..... | 0.9 | * | * | * | 0.3 | 0.7 | 1.0 | 1.5 | 1.8 | 2.1 | 3.5 | * |
| 2002..... | 0.9 | * | * | * | 0.5 | 0.7 | 0.9 | 1.5 | 1.3 | 2.2 | 3.6 | * |
| 2001..... | 0.8 | * | * | * | 0.4 | 0.8 | 1.2 | 0.9 | 1.7 | 1.6 | 3.9 | * |
| 2000..... | 0.9 | * | * | * | 0.5 | 0.6 | 0.8 | 1.3 | 1.7 | 2.2 | 4.0 | * |
| 1999..... | 1.0 | * | * | * | 0.4 | 0.8 | 1.1 | 1.4 | 1.7 | 2.7 | 3.5 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | 0.6 | 0.9 | 1.1 | 1.5 | 2.1 | * | * | * |
| 2003..... | 1.1 | * | * | * | * | 0.8 | 1.2 | 2.1 | 2.1 | 3.3 | * | * |
| 2002..... | 1.0 | * | * | * | * | 1.0 | 1.2 | 2.1 | * | * | * | * |
| 2001..... | 1.0 | * | * | * | * | 1.0 | 1.7 | 1.2 | 2.0 | * | * | * |
| 2000..... | 0.9 | * | * | * | * | * | 1.0 | 1.5 | 2.3 | * | * | * |
| 1999..... | 1.1 | * | * | * | * | 1.1 | 1.7 | 1.9 | 2.3 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.8 | 0.7 | * | 1.4 | * | * | * |
| 2003..... | 0.7 | * | * | * | * | * | 0.8 | 0.9 | 1.5 | * | * | * |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.1 | * | 2.1 | * | * |
| 2001..... | 0.7 | * | * | * | * | * | 0.7 | * | * | * | 4.0 | * |
| 2000..... | 0.8 | * | * | * | * | * | * | 1.2 | * | * | 3.5 | * |
| 1999..... | 0.8 | * | * | * | * | * | * | 1.0 | * | 2.7 | 3.5 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Transient cerebral ischemic attacks and related syndromes (G45) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 3.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.5 | 3.9 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.7 | 3.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.7 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 4.3 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 3.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 3.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 4.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.4 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 3.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 4.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 5.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 4.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 3.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 4.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 4.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 5.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 3.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 4.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 3.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.0 | 4.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 4.5 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 3.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 4.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 5.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Infantile cerebral palsy (G80) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | 0.5 | 0.6 | 0.8 | 0.5 | 0.3 | 0.4 | 0.5 | 0.6 | 0.5 | * |
| 2003..... | 0.5 | * | 0.5 | 0.6 | 0.7 | 0.5 | 0.2 | 0.4 | 0.5 | 0.4 | 0.5 | * |
| 2002..... | 0.4 | * | 0.6 | 0.5 | 0.7 | 0.4 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | * |
| 2001..... | 0.4 | * | 0.5 | 0.5 | 0.7 | 0.4 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | * |
| 2000..... | 0.4 | * | 0.4 | 0.6 | 0.6 | 0.4 | 0.2 | 0.3 | 0.3 | 0.5 | 0.5 | * |
| 1999..... | 0.4 | * | 0.5 | 0.5 | 0.5 | 0.4 | 0.2 | 0.2 | 0.3 | 0.4 | 0.4 | 0.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | 0.4 | 0.7 | 1.0 | 0.6 | 0.3 | 0.4 | 0.6 | 0.7 | 0.8 | * |
| 2003..... | 0.6 | * | 0.6 | 0.7 | 0.8 | 0.5 | 0.3 | 0.5 | 0.5 | 0.5 | 0.8 | * |
| 2002..... | 0.5 | * | 0.8 | 0.5 | 0.9 | 0.4 | 0.3 | 0.3 | 0.5 | 0.5 | 0.7 | * |
| 2001..... | 0.5 | * | 0.5 | 0.5 | 0.8 | 0.4 | 0.3 | 0.3 | 0.4 | 0.7 | 0.7 | * |
| 2000..... | 0.5 | * | * | 0.7 | 0.7 | 0.4 | 0.2 | 0.3 | 0.3 | 0.7 | 0.5 | * |
| 1999..... | 0.4 | * | 0.6 | 0.5 | 0.6 | 0.4 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | 0.6 | 0.5 | 0.6 | 0.4 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 | * |
| 2003..... | 0.4 | * | 0.4 | 0.5 | 0.6 | 0.5 | 0.2 | 0.3 | 0.4 | 0.3 | 0.4 | * |
| 2002..... | 0.4 | * | 0.4 | 0.5 | 0.5 | 0.3 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 | * |
| 2001..... | 0.4 | * | 0.4 | 0.5 | 0.6 | 0.3 | 0.2 | 0.3 | 0.4 | 0.3 | 0.4 | * |
| 2000..... | 0.4 | * | 0.4 | 0.6 | 0.5 | 0.3 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | * |
| 1999..... | 0.3 | * | 0.4 | 0.4 | 0.4 | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | 0.3 | 0.5 | 0.7 | 0.5 | 0.3 | 0.4 | 0.5 | 0.6 | 0.6 | * |
| 2003..... | 0.4 | * | 0.3 | 0.5 | 0.6 | 0.5 | 0.2 | 0.4 | 0.5 | 0.5 | 0.6 | * |
| 2002..... | 0.4 | * | 0.5 | 0.4 | 0.6 | 0.3 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | * |
| 2001..... | 0.4 | * | 0.4 | 0.4 | 0.6 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.6 | * |
| 2000..... | 0.4 | * | * | 0.5 | 0.6 | 0.4 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | * |
| 1999..... | 0.4 | * | 0.4 | 0.4 | 0.5 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | 0.6 | 0.8 | 0.6 | 0.3 | 0.4 | 0.6 | 0.8 | 0.8 | * |
| 2003..... | 0.5 | * | 0.4 | 0.6 | 0.8 | 0.5 | 0.3 | 0.5 | 0.6 | 0.6 | 0.8 | * |
| 2002..... | 0.5 | * | 0.7 | 0.4 | 0.8 | 0.4 | 0.3 | 0.3 | 0.5 | 0.6 | 0.7 | * |
| 2001..... | 0.5 | * | 0.5 | 0.4 | 0.7 | 0.4 | 0.3 | 0.3 | 0.4 | 0.7 | 0.8 | * |
| 2000..... | 0.4 | * | * | 0.6 | 0.7 | 0.4 | 0.2 | 0.3 | 0.4 | 0.7 | * | * |
| 1999..... | 0.4 | * | 0.5 | 0.4 | 0.6 | 0.4 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.4 | 0.5 | 0.4 | 0.3 | 0.4 | 0.4 | 0.5 | 0.4 | * |
| 2003..... | 0.4 | * | * | 0.4 | 0.5 | 0.5 | 0.2 | 0.3 | 0.4 | 0.4 | 0.4 | * |
| 2002..... | 0.3 | * | * | 0.4 | 0.4 | 0.3 | 0.2 | 0.3 | 0.4 | 0.4 | 0.4 | * |
| 2001..... | 0.3 | * | * | 0.4 | 0.5 | 0.3 | 0.2 | 0.2 | 0.4 | 0.3 | 0.4 | * |
| 2000..... | 0.3 | * | * | 0.5 | 0.5 | 0.3 | 0.2 | 0.2 | 0.4 | 0.3 | 0.5 | * |
| 1999..... | 0.3 | * | * | 0.4 | 0.3 | 0.4 | 0.3 | 0.2 | * | 0.3 | 0.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.9 | 6.7 | 1.1 | 0.5 | 0.9 | 1.1 | 2.4 | 5.5 | 11.9 | 23.5 | 39.2 | 51.4 |
| 2003..... | 7.1 | 7.3 | 0.9 | 0.5 | 1.0 | 1.1 | 2.4 | 5.4 | 12.1 | 24.5 | 40.4 | 56.0 |
| 2002..... | 6.9 | 6.9 | 1.1 | 0.5 | 0.9 | 1.2 | 2.5 | 5.4 | 11.7 | 24.1 | 40.5 | 53.8 |
| 2001..... | 6.7 | 6.1 | 1.2 | 0.5 | 1.0 | 1.1 | 2.3 | 5.2 | 11.2 | 24.0 | 39.4 | 51.9 |
| 2000..... | 6.7 | 7.1 | 1.2 | 0.5 | 0.9 | 1.1 | 2.4 | 5.5 | 11.4 | 23.3 | 39.4 | 55.4 |
| 1999..... | 6.5 | 7.7 | 1.2 | 0.6 | 1.1 | 1.0 | 2.1 | 5.0 | 10.9 | 23.3 | 39.3 | 53.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.3 | 7.8 | 1.1 | 0.6 | 1.3 | 1.3 | 2.9 | 6.6 | 13.8 | 27.6 | 47.2 | 62.4 |
| 2003..... | 7.5 | 8.3 | 0.9 | 0.6 | 1.5 | 1.4 | 3.0 | 6.6 | 14.0 | 27.8 | 49.8 | 70.0 |
| 2002..... | 7.3 | 8.4 | 1.3 | 0.6 | 1.3 | 1.5 | 3.0 | 6.4 | 13.4 | 28.2 | 49.3 | 66.1 |
| 2001..... | 7.1 | 7.1 | 1.2 | 0.6 | 1.5 | 1.2 | 2.9 | 5.9 | 12.5 | 28.5 | 49.0 | 69.7 |
| 2000..... | 7.2 | 8.1 | 1.3 | 0.5 | 1.3 | 1.4 | 2.8 | 6.5 | 13.2 | 27.9 | 49.1 | 72.0 |
| 1999..... | 7.0 | 8.6 | 1.2 | 0.6 | 1.6 | 1.2 | 2.6 | 5.8 | 12.6 | 27.4 | 49.7 | 70.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.5 | 5.6 | 1.1 | 0.4 | 0.4 | 0.8 | 2.0 | 4.5 | 10.2 | 20.0 | 33.9 | 46.5 |
| 2003..... | 6.6 | 6.2 | 1.0 | 0.4 | 0.5 | 0.9 | 1.8 | 4.3 | 10.3 | 21.8 | 34.1 | 49.9 |
| 2002..... | 6.5 | 5.4 | 0.9 | 0.3 | 0.5 | 0.8 | 1.9 | 4.4 | 10.1 | 20.6 | 34.6 | 48.5 |
| 2001..... | 6.3 | 5.0 | 1.2 | 0.5 | 0.5 | 0.9 | 1.8 | 4.4 | 10.0 | 20.2 | 33.1 | 44.5 |
| 2000..... | 6.3 | 6.0 | 1.1 | 0.5 | 0.5 | 0.8 | 1.9 | 4.6 | 9.7 | 19.5 | 33.0 | 48.7 |
| 1999..... | 6.1 | 6.8 | 1.2 | 0.5 | 0.5 | 0.7 | 1.7 | 4.3 | 9.2 | 20.0 | 32.6 | 46.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.5 | 6.1 | 0.7 | 0.4 | 0.6 | 0.4 | 1.2 | 3.1 | 7.7 | 14.8 | 26.0 | 49.0 |
| 2003..... | 2.6 | 5.2 | 1.3 | 0.3 | 0.8 | 0.6 | 0.9 | 3.0 | 8.4 | 15.3 | 26.9 | 59.1 |
| 2002..... | 2.4 | 5.2 | 0.7 | 0.4 | 0.7 | 0.6 | 1.0 | 3.1 | 7.3 | 17.4 | 23.5 | 44.9 |
| 2001..... | 2.1 | 3.5 | 0.9 | 0.4 | 0.6 | 0.5 | 1.2 | 2.6 | 7.0 | 13.6 | 24.7 | 38.1 |
| 2000..... | 2.4 | 6.9 | 0.9 | 0.4 | 0.6 | 0.6 | 1.3 | 2.8 | 6.7 | 15.2 | 30.4 | 39.1 |
| 1999..... | 2.2 | 5.0 | 0.9 | 0.4 | 0.6 | 0.5 | 1.2 | 2.4 | 6.8 | 15.4 | 24.6 | 42.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.6 | 7.1 | * | 0.5 | 0.9 | 0.6 | 1.7 | 3.9 | 8.8 | 17.7 | 27.5 | 40.6 |
| 2003..... | 2.7 | 6.8 | 1.2 | * | 1.1 | 0.7 | 1.0 | 3.3 | 10.4 | 19.2 | 35.3 | 55.4 |
| 2002..... | 2.5 | 6.1 | * | * | 1.0 | 0.8 | 1.2 | 3.9 | 8.0 | 18.5 | 27.5 | 46.9 |
| 2001..... | 2.4 | * | * | * | 1.0 | 0.6 | 1.4 | 3.3 | 8.1 | 18.0 | 25.9 | 55.2 |
| 2000..... | 2.6 | 7.3 | * | * | 0.8 | 0.8 | 1.4 | 3.9 | 8.3 | 20.3 | 36.4 | 52.4 |
| 1999..... | 2.4 | 5.7 | * | * | 0.9 | 0.6 | 1.3 | 3.0 | 8.6 | 17.8 | 32.8 | 49.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.4 | 5.0 | * | * | * | * | 0.8 | 2.3 | 6.6 | 12.5 | 24.9 | 53.3 |
| 2003..... | 2.4 | * | 1.3 | * | * | * | 0.8 | 2.7 | 6.6 | 12.2 | 21.1 | 60.9 |
| 2002..... | 2.3 | * | * | * | * | * | 0.8 | 2.2 | 6.6 | 16.6 | 20.8 | 43.9 |
| 2001..... | 1.9 | * | * | * | * | * | 0.9 | 2.0 | 6.1 | 10.1 | 23.9 | 49.8 |
| 2000..... | 2.1 | 6.4 | * | * | * | * | 1.1 | 1.7 | 5.2 | 11.3 | 26.4 | 32.6 |
| 1999..... | 2.0 | * | * | * | * | * | 1.1 | 1.8 | 5.2 | 13.6 | 19.1 | 39.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.6 | 6.8 | 1.2 | 0.5 | 0.9 | 1.2 | 2.6 | 5.8 | 12.2 | 24.1 | 39.9 | 51.4 |
| 2003..... | 7.8 | 7.8 | 0.8 | 0.6 | 1.1 | 1.2 | 2.6 | 5.7 | 12.4 | 25.1 | 41.0 | 55.8 |
| 2002..... | 7.6 | 7.4 | 1.2 | 0.5 | 1.0 | 1.3 | 2.7 | 5.6 | 12.0 | 24.5 | 41.2 | 53.9 |
| 2001..... | 7.3 | 6.7 | 1.3 | 0.6 | 1.1 | 1.2 | 2.5 | 5.4 | 11.5 | 24.6 | 40.0 | 52.2 |
| 2000..... | 7.3 | 7.1 | 1.3 | 0.5 | 1.0 | 1.2 | 2.5 | 5.7 | 11.7 | 23.7 | 39.7 | 55.9 |
| 1999..... | 7.1 | 8.3 | 1.2 | 0.6 | 1.1 | 1.1 | 2.2 | 5.3 | 11.1 | 23.7 | 39.8 | 53.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 8.1 | 7.9 | 1.2 | 0.6 | 1.3 | 1.5 | 3.1 | 6.9 | 14.1 | 28.3 | 48.3 | 63.3 |
| 2003..... | 8.3 | 8.6 | 0.8 | 0.6 | 1.6 | 1.5 | 3.4 | 7.0 | 14.3 | 28.3 | 50.5 | 70.5 |
| 2002..... | 8.1 | 9.0 | 1.4 | 0.6 | 1.4 | 1.7 | 3.3 | 6.6 | 13.8 | 28.8 | 50.3 | 66.7 |
| 2001..... | 7.8 | 7.7 | 1.3 | 0.6 | 1.6 | 1.4 | 3.1 | 6.1 | 12.8 | 29.1 | 49.9 | 69.9 |
| 2000..... | 7.8 | 8.3 | 1.4 | 0.5 | 1.4 | 1.5 | 3.0 | 6.7 | 13.5 | 28.3 | 49.6 | 72.5 |
| 1999..... | 7.6 | 9.2 | 1.3 | 0.6 | 1.7 | 1.3 | 2.8 | 6.1 | 12.9 | 27.8 | 50.3 | 70.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.1 | 5.7 | 1.2 | 0.4 | 0.5 | 0.9 | 2.2 | 4.7 | 10.5 | 20.5 | 34.4 | 46.2 |
| 2003..... | 7.2 | 7.0 | 0.9 | 0.5 | 0.6 | 1.0 | 1.9 | 4.4 | 10.6 | 22.4 | 34.7 | 49.3 |
| 2002..... | 7.0 | 5.8 | 1.0 | 0.3 | 0.5 | 0.8 | 2.0 | 4.5 | 10.4 | 20.8 | 35.2 | 48.4 |
| 2001..... | 6.9 | 5.6 | 1.3 | 0.5 | 0.6 | 1.0 | 1.9 | 4.6 | 10.3 | 20.9 | 33.4 | 44.9 |
| 2000..... | 6.8 | 5.9 | 1.2 | 0.4 | 0.6 | 0.8 | 2.0 | 4.8 | 10.0 | 20.0 | 33.2 | 49.1 |
| 1999..... | 6.6 | 7.3 | 1.2 | 0.5 | 0.5 | 0.8 | 1.7 | 4.5 | 9.5 | 20.3 | 33.0 | 46.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 8.2 | 7.0 | 1.1 | 0.5 | 0.9 | 1.2 | 2.7 | 5.6 | 12.1 | 24.4 | 40.5 | 52.3 |
| 2003..... | 8.4 | 7.4 | 0.7 | 0.5 | 1.2 | 1.3 | 2.6 | 5.5 | 12.2 | 25.5 | 41.7 | 56.6 |
| 2002..... | 8.2 | 7.4 | 1.1 | 0.5 | 1.0 | 1.3 | 2.7 | 5.4 | 12.0 | 24.8 | 41.9 | 54.9 |
| 2001..... | 7.9 | 6.4 | 1.1 | 0.6 | 1.1 | 1.1 | 2.4 | 5.2 | 11.2 | 24.9 | 40.4 | 52.8 |
| 2000..... | 7.9 | 6.8 | 1.2 | 0.5 | 1.1 | 1.2 | 2.4 | 5.6 | 11.6 | 23.8 | 40.5 | 56.7 |
| 1999..... | 7.6 | 8.1 | 1.1 | 0.5 | 1.2 | 1.1 | 2.2 | 5.0 | 10.9 | 23.8 | 40.5 | 54.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 8.8 | 8.0 | 1.1 | 0.6 | 1.4 | 1.5 | 3.3 | 6.7 | 14.0 | 28.9 | 49.1 | 65.6 |
| 2003..... | 9.0 | 8.6 | 0.6 | 0.6 | 1.7 | 1.6 | 3.3 | 6.9 | 14.1 | 28.9 | 51.5 | 72.6 |
| 2002..... | 8.8 | 9.2 | 1.2 | 0.6 | 1.4 | 1.8 | 3.3 | 6.5 | 13.8 | 28.9 | 51.5 | 68.2 |
| 2001..... | 8.4 | 7.2 | 1.2 | 0.6 | 1.7 | 1.2 | 3.0 | 5.9 | 12.6 | 29.5 | 50.7 | 70.4 |
| 2000..... | 8.4 | 7.4 | 1.3 | 0.5 | 1.6 | 1.5 | 2.8 | 6.7 | 13.3 | 28.5 | 50.6 | 73.8 |
| 1999..... | 8.2 | 9.0 | 1.2 | 0.5 | 1.9 | 1.4 | 2.7 | 5.9 | 12.8 | 28.1 | 51.6 | 72.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.6 | 5.9 | 1.1 | 0.4 | 0.4 | 0.8 | 2.1 | 4.4 | 10.4 | 20.4 | 34.7 | 46.3 |
| 2003..... | 7.8 | 6.1 | 0.8 | 0.4 | 0.6 | 1.0 | 1.8 | 4.2 | 10.4 | 22.7 | 35.1 | 49.6 |
| 2002..... | 7.6 | 5.5 | 0.8 | 0.3 | 0.5 | 0.8 | 2.0 | 4.4 | 10.3 | 21.2 | 35.5 | 49.1 |
| 2001..... | 7.3 | 5.6 | 1.0 | 0.5 | 0.5 | 0.9 | 1.9 | 4.5 | 9.8 | 21.0 | 33.6 | 45.5 |
| 2000..... | 7.3 | 6.1 | 1.1 | 0.4 | 0.5 | 0.8 | 1.9 | 4.5 | 10.1 | 19.8 | 33.8 | 49.8 |
| 1999..... | 7.1 | 7.2 | 1.1 | 0.5 | 0.5 | 0.8 | 1.6 | 4.1 | 9.2 | 20.3 | 33.3 | 47.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 6.4 | 6.9 | 1.8 | 0.5 | 1.0 | 1.9 | 3.3 | 8.5 | 16.5 | 27.0 | 42.1 | 50.1 |
| 2003..... | 6.4 | 9.8 | 1.5 | 0.7 | 0.8 | 1.6 | 3.8 | 8.3 | 16.3 | 27.7 | 38.7 | 52.9 |
| 2002..... | 6.1 | 8.2 | 1.7 | 0.5 | 1.1 | 1.7 | 3.6 | 7.9 | 15.4 | 27.5 | 40.0 | 50.8 |
| 2001..... | 6.3 | 8.3 | 2.1 | 0.7 | 1.2 | 2.0 | 3.3 | 8.0 | 16.3 | 27.5 | 41.5 | 49.0 |
| 2000..... | 6.0 | 7.6 | 1.7 | 0.5 | 1.0 | 1.7 | 3.7 | 8.7 | 14.6 | 27.4 | 37.4 | 50.9 |
| 1999..... | 5.9 | 9.3 | 1.4 | 0.9 | 1.2 | 1.3 | 3.1 | 8.6 | 14.9 | 26.6 | 36.8 | 50.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.4 | 7.7 | 2.0 | * | 1.2 | 1.8 | 3.5 | 10.0 | 19.5 | 29.8 | 48.8 | 50.8 |
| 2003..... | 6.5 | 9.9 | 1.8 | 0.8 | 1.2 | 1.9 | 4.6 | 9.4 | 18.9 | 31.1 | 45.0 | 51.1 |
| 2002..... | 6.5 | 8.2 | 2.0 | 0.6 | 1.4 | 2.0 | 4.5 | 9.3 | 17.8 | 34.3 | 45.1 | 58.0 |
| 2001..... | 6.5 | 10.5 | 1.9 | * | 1.6 | 2.5 | 4.1 | 9.4 | 17.4 | 31.9 | 51.9 | 62.4 |
| 2000..... | 6.4 | 10.9 | 1.7 | * | 1.0 | 2.1 | 4.4 | 9.3 | 18.1 | 32.2 | 49.7 | 62.2 |
| 1999..... | 5.9 | 10.7 | * | 1.0 | 1.6 | 1.5 | 3.6 | 9.4 | 15.9 | 28.6 | 43.5 | 65.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | * | 0.7 | 1.9 | 3.2 | 7.1 | 14.0 | 25.1 | 38.3 | 49.8 |
| 2003..... | 6.2 | 9.7 | * | 0.7 | * | 1.3 | 3.2 | 7.3 | 14.2 | 25.3 | 35.2 | 53.7 |
| 2002..... | 5.8 | 8.2 | * | * | 0.7 | 1.4 | 2.7 | 6.7 | 13.6 | 22.6 | 37.1 | 48.1 |
| 2001..... | 6.1 | * | 2.2 | 0.8 | 0.9 | 1.6 | 2.6 | 6.9 | 15.5 | 24.5 | 35.7 | 44.1 |
| 2000..... | 5.7 | * | * | * | 1.0 | 1.4 | 3.1 | 8.2 | 11.8 | 24.0 | 30.6 | 46.7 |
| 1999..... | 5.9 | 7.9 | * | 0.7 | * | 1.2 | 2.7 | 8.0 | 14.2 | 25.2 | 33.1 | 45.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

VIII. Diseases of the ear and mastoid process (H60-H93)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, both sexes.

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, male.

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, female.

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, both sexes.

Non-Hispanic white, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, male.

Non-Hispanic white, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| IX. Diseases of the circulatory system (I00-I99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 294.9 | 14.5 | 1.5 | 0.8 | 3.3 | 10.3 | 37.3 | 112.2 | 270.3 | 701.3 | 2,044.6 | 6,639.9 |
| 2003..... | 311.9 | 14.8 | 1.5 | 0.9 | 3.6 | 10.6 | 38.9 | 114.8 | 287.0 | 752.4 | 2,179.9 | 7,162.1 |
| 2002..... | 320.2 | 16.5 | 1.5 | 0.9 | 3.2 | 10.3 | 38.4 | 115.6 | 297.1 | 792.2 | 2,266.8 | 7,414.0 |
| 2001..... | 325.5 | 15.4 | 1.9 | 0.9 | 3.3 | 10.5 | 37.7 | 114.6 | 303.4 | 815.0 | 2,331.6 | 7,699.8 |
| 2000..... | 334.6 | 17.4 | 1.5 | 0.9 | 3.4 | 9.8 | 37.5 | 117.0 | 321.3 | 853.6 | 2,405.6 | 8,055.8 |
| 1999..... | 342.0 | 17.6 | 1.5 | 0.9 | 3.5 | 9.9 | 38.3 | 117.3 | 329.2 | 891.0 | 2,484.0 | 8,220.5 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 282.7 | 15.6 | 1.5 | 0.8 | 4.1 | 13.1 | 49.8 | 158.3 | 374.1 | 907.8 | 2,472.0 | 6,837.7 |
| 2003..... | 297.7 | 16.2 | 1.5 | 1.0 | 4.3 | 13.2 | 51.8 | 162.5 | 395.7 | 980.4 | 2,644.5 | 7,348.4 |
| 2002..... | 304.7 | 17.3 | 1.5 | 0.9 | 4.2 | 13.1 | 51.5 | 163.7 | 409.5 | 1,031.7 | 2,740.6 | 7,659.3 |
| 2001..... | 307.6 | 15.8 | 2.0 | 0.9 | 4.0 | 13.0 | 50.6 | 161.8 | 416.5 | 1,062.5 | 2,833.1 | 7,958.0 |
| 2000..... | 317.2 | 18.3 | 1.7 | 1.0 | 4.0 | 12.1 | 50.3 | 166.7 | 443.8 | 1,118.8 | 2,936.2 | 8,470.3 |
| 1999..... | 324.2 | 18.5 | 1.6 | 1.0 | 4.1 | 12.3 | 51.9 | 166.9 | 454.0 | 1,172.7 | 3,033.6 | 8,624.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 306.7 | 13.5 | 1.5 | 0.8 | 2.4 | 7.5 | 24.8 | 67.7 | 174.0 | 527.9 | 1,756.9 | 6,551.0 |
| 2003..... | 325.8 | 13.3 | 1.5 | 0.7 | 2.9 | 7.9 | 26.1 | 68.7 | 186.1 | 561.7 | 1,869.4 | 7,079.7 |
| 2002..... | 335.2 | 15.7 | 1.4 | 0.8 | 2.3 | 7.5 | 25.5 | 69.2 | 193.1 | 592.8 | 1,952.2 | 7,307.6 |
| 2001..... | 342.8 | 15.0 | 1.8 | 0.9 | 2.5 | 7.9 | 24.9 | 69.1 | 198.9 | 609.9 | 2,002.1 | 7,591.7 |
| 2000..... | 351.3 | 16.5 | 1.4 | 0.8 | 2.9 | 7.5 | 24.9 | 69.2 | 208.3 | 635.2 | 2,059.6 | 7,887.0 |
| 1999..... | 359.1 | 16.6 | 1.4 | 0.8 | 2.8 | 7.4 | 24.9 | 69.5 | 214.3 | 660.4 | 2,127.0 | 8,057.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 89.8 | 12.0 | 1.3 | 0.6 | 2.6 | 5.8 | 21.5 | 72.5 | 203.9 | 557.7 | 1,572.8 | 4,601.3 |
| 2003..... | 94.0 | 12.2 | 1.4 | 0.6 | 2.6 | 6.6 | 20.9 | 73.0 | 213.6 | 597.5 | 1,689.9 | 5,194.2 |
| 2002..... | 94.8 | 13.9 | 1.2 | 0.7 | 2.7 | 6.3 | 20.7 | 74.6 | 227.2 | 624.9 | 1,734.8 | 5,388.0 |
| 2001..... | 96.8 | 12.6 | 1.5 | 0.8 | 2.5 | 6.5 | 20.1 | 76.3 | 217.3 | 649.1 | 1,835.8 | 6,068.8 |
| 2000..... | 97.1 | 14.5 | 1.3 | 0.8 | 2.6 | 6.6 | 22.0 | 76.0 | 237.6 | 664.0 | 1,861.5 | 6,192.8 |
| 1999..... | 99.7 | 15.5 | 1.4 | 0.7 | 2.6 | 6.6 | 21.7 | 81.6 | 244.1 | 694.1 | 1,940.9 | 6,360.5 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 91.1 | 12.3 | 1.6 | 0.5 | 3.5 | 8.0 | 29.6 | 102.5 | 284.4 | 732.9 | 1,932.8 | 4,606.3 |
| 2003..... | 93.1 | 13.4 | 1.6 | 0.7 | 3.3 | 8.0 | 28.5 | 104.2 | 297.3 | 782.8 | 2,014.1 | 5,093.6 |
| 2002..... | 94.6 | 13.4 | * | 0.6 | 3.4 | 8.3 | 27.7 | 103.8 | 318.2 | 825.7 | 2,074.3 | 5,503.2 |
| 2001..... | 95.9 | 13.7 | 1.4 | 0.8 | 3.1 | 8.4 | 27.1 | 108.5 | 298.3 | 856.4 | 2,229.5 | 6,186.1 |
| 2000..... | 96.7 | 14.4 | 1.4 | 0.9 | 3.1 | 8.3 | 28.6 | 109.0 | 332.3 | 873.4 | 2,283.4 | 6,330.5 |
| 1999..... | 99.7 | 12.7 | 1.4 | 0.9 | 3.1 | 8.7 | 30.0 | 117.3 | 334.6 | 929.0 | 2,362.9 | 6,621.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 88.5 | 11.7 | * | 0.7 | 1.5 | 3.1 | 12.5 | 42.3 | 130.7 | 417.1 | 1,324.2 | 4,598.8 |
| 2003..... | 94.9 | 11.1 | * | * | 1.8 | 4.9 | 12.6 | 41.8 | 138.1 | 449.9 | 1,467.3 | 5,245.3 |
| 2002..... | 95.1 | 14.5 | 1.4 | 0.8 | 1.8 | 3.9 | 13.2 | 45.4 | 145.5 | 465.2 | 1,502.2 | 5,330.9 |
| 2001..... | 97.7 | 11.5 | 1.6 | 0.8 | 1.8 | 4.3 | 12.6 | 44.6 | 145.3 | 485.7 | 1,569.7 | 6,011.4 |
| 2000..... | 97.6 | 14.6 | * | * | 2.0 | 4.6 | 14.8 | 43.8 | 153.7 | 499.4 | 1,578.2 | 6,125.2 |
| 1999..... | 99.8 | 18.4 | 1.4 | * | 2.0 | 4.0 | 12.8 | 46.9 | 164.1 | 510.0 | 1,660.0 | 6,234.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| IX. Diseases of the circulatory system (I00-I99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 327.7 | 15.2 | 1.5 | 0.9 | 3.4 | 11.4 | 39.7 | 116.1 | 274.8 | 709.7 | 2,066.7 | 6,721.6 |
| 2003..... | 345.6 | 15.3 | 1.5 | 0.9 | 3.8 | 11.5 | 41.5 | 118.6 | 291.6 | 760.8 | 2,200.0 | 7,231.1 |
| 2002..... | 354.0 | 17.1 | 1.5 | 0.9 | 3.4 | 11.2 | 40.9 | 119.2 | 301.4 | 800.5 | 2,286.3 | 7,476.1 |
| 2001..... | 358.5 | 16.0 | 2.0 | 0.9 | 3.4 | 11.3 | 39.9 | 117.7 | 308.7 | 823.2 | 2,347.5 | 7,741.0 |
| 2000..... | 367.4 | 18.0 | 1.6 | 0.9 | 3.6 | 10.4 | 39.3 | 120.1 | 326.0 | 862.1 | 2,421.6 | 8,102.0 |
| 1999..... | 374.3 | 18.1 | 1.5 | 0.9 | 3.6 | 10.5 | 40.2 | 119.8 | 333.7 | 899.1 | 2,498.5 | 8,263.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 315.0 | 16.5 | 1.4 | 0.9 | 4.3 | 14.4 | 53.1 | 163.8 | 379.8 | 917.1 | 2,496.9 | 6,937.9 |
| 2003..... | 331.0 | 16.9 | 1.4 | 1.0 | 4.5 | 14.4 | 55.4 | 167.9 | 401.5 | 990.5 | 2,670.8 | 7,438.9 |
| 2002..... | 338.0 | 18.3 | 1.7 | 1.0 | 4.3 | 14.1 | 54.9 | 169.0 | 414.7 | 1,041.1 | 2,765.3 | 7,734.9 |
| 2001..... | 339.7 | 16.2 | 2.1 | 0.9 | 4.2 | 14.0 | 53.7 | 166.1 | 423.2 | 1,071.8 | 2,852.4 | 8,012.2 |
| 2000..... | 349.2 | 19.0 | 1.7 | 1.0 | 4.2 | 12.9 | 52.8 | 171.0 | 449.6 | 1,128.8 | 2,955.6 | 8,535.2 |
| 1999..... | 355.6 | 20.0 | 1.7 | 1.0 | 4.3 | 13.0 | 54.4 | 170.4 | 459.9 | 1,181.4 | 3,051.5 | 8,678.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 339.8 | 13.8 | 1.6 | 0.9 | 2.5 | 8.4 | 26.6 | 70.1 | 177.2 | 534.9 | 1,777.5 | 6,624.9 |
| 2003..... | 359.6 | 13.7 | 1.6 | 0.8 | 3.1 | 8.5 | 27.9 | 71.2 | 189.4 | 568.1 | 1,885.9 | 7,139.8 |
| 2002..... | 369.3 | 15.9 | 1.4 | 0.8 | 2.4 | 8.3 | 27.2 | 71.3 | 196.2 | 599.7 | 1,968.8 | 7,364.4 |
| 2001..... | 376.3 | 15.9 | 1.8 | 0.9 | 2.7 | 8.6 | 26.5 | 71.0 | 202.5 | 616.6 | 2,016.2 | 7,628.2 |
| 2000..... | 384.6 | 16.9 | 1.5 | 0.8 | 3.0 | 8.0 | 26.0 | 71.2 | 211.6 | 642.0 | 2,073.7 | 7,926.7 |
| 1999..... | 392.0 | 16.0 | 1.4 | 0.8 | 3.0 | 8.0 | 26.3 | 71.2 | 217.3 | 667.3 | 2,139.6 | 8,096.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 354.9 | 12.7 | 1.2 | 0.8 | 2.8 | 9.1 | 34.5 | 100.8 | 249.1 | 674.0 | 2,045.6 | 6,813.3 |
| 2003..... | 374.2 | 11.7 | 1.2 | 0.8 | 3.1 | 9.3 | 35.5 | 102.7 | 264.3 | 721.7 | 2,178.4 | 7,328.5 |
| 2002..... | 383.7 | 14.0 | 1.4 | 0.8 | 2.8 | 8.8 | 35.5 | 104.1 | 274.0 | 764.3 | 2,261.8 | 7,570.6 |
| 2001..... | 387.6 | 12.6 | 1.6 | 0.8 | 2.7 | 9.2 | 33.8 | 102.4 | 280.5 | 783.7 | 2,322.1 | 7,837.2 |
| 2000..... | 397.3 | 15.1 | 1.4 | 0.8 | 2.9 | 8.4 | 33.4 | 105.1 | 297.2 | 824.2 | 2,394.0 | 8,204.4 |
| 1999..... | 404.3 | 14.9 | 1.3 | 0.8 | 3.0 | 8.5 | 33.9 | 104.9 | 304.3 | 860.2 | 2,471.9 | 8,365.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 338.6 | 14.0 | 1.1 | 0.8 | 3.5 | 11.8 | 47.7 | 146.7 | 348.9 | 880.1 | 2,485.7 | 7,065.8 |
| 2003..... | 355.6 | 12.3 | 1.0 | 0.9 | 3.7 | 11.9 | 49.2 | 149.8 | 370.5 | 949.4 | 2,656.2 | 7,580.3 |
| 2002..... | 363.7 | 15.4 | 1.6 | 0.9 | 3.6 | 11.4 | 49.3 | 152.4 | 383.2 | 1,004.9 | 2,753.4 | 7,866.2 |
| 2001..... | 364.6 | 13.4 | 1.7 | 0.8 | 3.4 | 11.6 | 47.2 | 149.3 | 392.0 | 1,031.7 | 2,836.9 | 8,137.5 |
| 2000..... | 375.2 | 16.3 | 1.6 | 0.9 | 3.4 | 10.7 | 46.9 | 154.6 | 417.4 | 1,092.7 | 2,940.1 | 8,672.2 |
| 1999..... | 381.1 | 16.9 | 1.5 | 0.8 | 3.5 | 10.8 | 47.5 | 153.6 | 426.4 | 1,143.7 | 3,039.5 | 8,796.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 370.6 | 11.3 | 1.3 | 0.7 | 2.1 | 6.4 | 21.2 | 55.7 | 154.3 | 496.8 | 1,745.8 | 6,700.2 |
| 2003..... | 392.1 | 11.1 | 1.4 | 0.6 | 2.5 | 6.6 | 21.7 | 56.3 | 163.4 | 526.7 | 1,855.7 | 7,217.7 |
| 2002..... | 402.9 | 12.5 | 1.1 | 0.7 | 1.9 | 6.3 | 21.7 | 56.5 | 170.7 | 559.2 | 1,932.4 | 7,442.9 |
| 2001..... | 409.7 | 11.7 | 1.5 | 0.8 | 2.1 | 6.8 | 20.4 | 56.2 | 175.2 | 573.4 | 1,980.8 | 7,712.2 |
| 2000..... | 418.5 | 14.0 | 1.2 | 0.6 | 2.3 | 6.0 | 20.0 | 56.4 | 183.9 | 598.2 | 2,035.1 | 8,015.8 |
| 1999..... | 426.5 | 12.9 | 1.1 | 0.7 | 2.5 | 6.2 | 20.2 | 57.0 | 189.3 | 623.1 | 2,100.2 | 8,194.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| IX. Diseases of the circulatory system (I00-I99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 271.4 | 23.5 | 3.0 | 1.6 | 6.2 | 24.7 | 78.8 | 239.2 | 539.8 | 1,147.5 | 2,620.1 | 6,465.6 |
| 2003..... | 284.4 | 27.5 | 2.9 | 1.5 | 7.4 | 24.7 | 83.3 | 247.2 | 569.9 | 1,231.8 | 2,761.1 | 6,957.5 |
| 2002..... | 287.3 | 29.4 | 2.1 | 1.5 | 6.2 | 25.3 | 79.9 | 245.2 | 583.0 | 1,250.0 | 2,868.8 | 7,166.9 |
| 2001..... | 291.7 | 30.9 | 3.8 | 1.3 | 7.1 | 24.3 | 82.2 | 245.5 | 592.2 | 1,301.4 | 2,930.1 | 7,396.3 |
| 2000..... | 296.3 | 29.6 | 2.6 | 1.5 | 7.3 | 23.3 | 80.7 | 249.2 | 614.2 | 1,342.1 | 3,009.6 | 7,606.4 |
| 1999..... | 300.2 | 30.8 | 2.6 | 1.4 | 7.2 | 22.2 | 85.1 | 249.2 | 626.7 | 1,389.5 | 3,058.9 | 7,652.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 268.0 | 25.7 | 2.9 | 1.5 | 8.1 | 30.1 | 97.7 | 315.9 | 735.1 | 1,460.8 | 3,086.6 | 6,306.0 |
| 2003..... | 279.6 | 33.2 | 3.3 | 1.6 | 9.0 | 30.1 | 102.8 | 328.2 | 759.1 | 1,570.8 | 3,317.7 | 6,658.9 |
| 2002..... | 279.3 | 29.7 | 2.0 | 1.5 | 7.6 | 31.2 | 99.3 | 322.4 | 781.1 | 1,584.6 | 3,336.9 | 7,014.4 |
| 2001..... | 283.2 | 30.6 | 4.0 | 1.2 | 8.5 | 29.6 | 101.8 | 321.5 | 778.6 | 1,648.8 | 3,477.5 | 7,442.7 |
| 2000..... | 286.3 | 30.7 | 2.3 | 1.5 | 8.1 | 27.4 | 99.4 | 327.5 | 812.8 | 1,667.2 | 3,541.0 | 7,786.1 |
| 1999..... | 292.8 | 32.6 | 2.8 | 1.6 | 8.8 | 27.2 | 107.1 | 332.5 | 835.8 | 1,747.0 | 3,559.7 | 7,974.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 274.5 | 21.2 | 3.0 | 1.7 | 4.4 | 19.9 | 62.0 | 173.4 | 382.9 | 924.1 | 2,356.8 | 6,528.7 |
| 2003..... | 288.8 | 21.6 | 2.4 | 1.3 | 5.9 | 19.8 | 66.1 | 177.7 | 418.1 | 990.8 | 2,446.8 | 7,073.1 |
| 2002..... | 294.5 | 29.0 | 2.2 | 1.4 | 4.8 | 19.9 | 62.8 | 179.2 | 424.0 | 1,012.7 | 2,603.9 | 7,225.2 |
| 2001..... | 299.5 | 31.2 | 3.6 | 1.4 | 5.7 | 19.5 | 64.9 | 180.3 | 442.2 | 1,056.3 | 2,622.8 | 7,379.1 |
| 2000..... | 305.4 | 28.5 | 2.9 | 1.6 | 6.5 | 19.6 | 64.1 | 182.0 | 454.3 | 1,113.4 | 2,710.4 | 7,539.6 |
| 1999..... | 307.0 | 28.9 | 2.4 | 1.3 | 5.6 | 17.7 | 65.8 | 178.0 | 459.0 | 1,137.4 | 2,778.5 | 7,532.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.3 | 8.8 | 17.1 |
| 2003..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.8 | 9.6 | 20.5 |
| 2002..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 3.8 | 9.6 | 17.8 |
| 2001..... | 1.2 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.1 | 9.4 | 17.7 |
| 2000..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.5 | 9.5 | 16.8 |
| 1999..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.6 | 10.0 | 18.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.4 | 7.3 | 11.5 |
| 2003..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.9 | 7.5 | 16.2 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 6.7 | 11.8 |
| 2001..... | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.8 | 6.6 | 12.8 |
| 2000..... | 0.8 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 2.9 | 6.6 | 12.2 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 6.8 | 10.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 4.1 | 9.9 | 19.5 |
| 2003..... | 1.7 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 4.5 | 11.0 | 22.3 |
| 2002..... | 1.7 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.6 | 11.6 | 20.4 |
| 2001..... | 1.7 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 5.1 | 11.3 | 19.8 |
| 2000..... | 1.8 | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.0 | 5.8 | 11.4 | 18.7 |
| 1999..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 5.6 | 12.2 | 21.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 1.3 | 3.5 | 6.3 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.5 | 1.0 | 3.3 | 6.7 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.7 | 1.8 | 4.6 | 7.0 | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 1.5 | 4.2 | 5.4 | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 1.8 | 4.4 | 5.3 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.8 | 1.6 | 4.2 | 7.3 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 2.1 | 3.7 | 6.4 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 4.0 | 8.4 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | * | 2.1 | 5.9 | 10.7 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 2.0 | 5.2 | * | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 5.6 | 6.9 | * |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 5.9 | 10.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.3 | 9.0 | 17.7 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.8 | 9.7 | 20.9 |
| 2002..... | 1.4 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 3.8 | 9.7 | 18.1 |
| 2001..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.1 | 9.6 | 18.0 |
| 2000..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.5 | 9.7 | 17.1 |
| 1999..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.6 | 10.1 | 18.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.3 | 7.3 | 11.9 |
| 2003..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.9 | 7.6 | 16.6 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 7.0 | 12.0 |
| 2001..... | 0.8 | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.8 | 6.7 | 13.1 |
| 2000..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 2.9 | 6.7 | 12.5 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.6 | 6.9 | 10.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.1 | 10.0 | 20.2 |
| 2003..... | 1.8 | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 4.6 | 11.1 | 22.8 |
| 2002..... | 1.8 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.5 | 11.6 | 20.8 |
| 2001..... | 1.8 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 5.1 | 11.5 | 20.0 |
| 2000..... | 1.9 | * | * | * | * | * | 0.4 | 0.8 | 2.0 | 5.8 | 11.6 | 19.0 |
| 1999..... | 2.0 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 5.5 | 12.2 | 21.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 3.4 | 9.5 | 18.9 |
| 2003..... | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.9 | 10.4 | 22.3 |
| 2002..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.9 | 10.2 | 19.3 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.0 | 10.2 | 19.0 |
| 2000..... | 1.6 | * | * | * | * | * | 0.3 | 0.6 | 1.7 | 4.6 | 10.3 | 18.0 |
| 1999..... | 1.6 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 4.8 | 10.6 | 19.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.4 | 0.7 | 2.4 | 7.8 | 12.8 |
| 2003..... | 1.0 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.0 | 8.3 | 17.7 |
| 2002..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.0 | 7.3 | 12.9 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.7 | 7.0 | 13.7 |
| 2000..... | 0.9 | * | * | * | * | * | 0.3 | 0.5 | 1.5 | 2.9 | 7.2 | 13.0 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.8 | 7.2 | 11.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.2 | 10.6 | 21.6 |
| 2003..... | 2.1 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.6 | 11.8 | 24.4 |
| 2002..... | 2.1 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 4.6 | 12.2 | 22.0 |
| 2001..... | 2.0 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 5.1 | 12.2 | 21.2 |
| 2000..... | 2.1 | * | * | * | * | * | 0.3 | 0.7 | 1.9 | 6.0 | 12.4 | 20.0 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.7 | 12.9 | 22.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.6 | 1.0 | 1.6 | 2.7 | 4.1 | 6.0 |
| 2003..... | 0.8 | * | * | * | * | * | 0.5 | 1.4 | 2.7 | 4.0 | 4.2 | 7.1 |
| 2002..... | 0.7 | * | * | * | * | * | 0.4 | 1.0 | 2.4 | 2.9 | 4.4 | * |
| 2001..... | 0.8 | * | * | * | * | * | 0.7 | 1.4 | 2.1 | 4.2 | 4.3 | 6.9 |
| 2000..... | 0.7 | * | * | * | * | * | 0.6 | 1.0 | 2.0 | 3.6 | 4.0 | 8.6 |
| 1999..... | 0.8 | * | * | * | * | * | 0.5 | 1.2 | 2.2 | 3.4 | 4.4 | 8.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.9 | * | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.9 | 2.8 | 3.0 | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 1.8 | 2.9 | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | 1.0 | * | * | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 1.1 | 1.7 | 3.4 | 5.2 | * |
| 2003..... | 1.0 | * | * | * | * | * | * | 1.7 | 2.6 | 4.6 | 5.2 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.9 | 2.9 | 5.1 | * |
| 2001..... | 1.2 | * | * | * | * | * | 0.9 | 1.7 | 2.6 | 5.4 | 5.4 | * |
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | 1.4 | 2.7 | 4.4 | 4.9 | * |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.5 | 2.7 | 4.1 | 4.4 | 11.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic rheumatic heart diseases (I05-I09) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.3 | 8.8 | 17.0 |
| 2003..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.7 | 9.5 | 20.3 |
| 2002..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 3.8 | 9.5 | 17.5 |
| 2001..... | 1.2 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.0 | 9.3 | 17.2 |
| 2000..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.4 | 9.5 | 16.7 |
| 1999..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.6 | 10.0 | 17.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.3 | 7.2 | 11.4 |
| 2003..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.8 | 7.4 | 16.1 |
| 2002..... | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.9 | 6.6 | 11.4 |
| 2001..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.7 | 6.5 | 12.3 |
| 2000..... | 0.8 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 2.8 | 6.5 | 12.1 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.5 | 6.7 | 10.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 4.0 | 9.8 | 19.5 |
| 2003..... | 1.7 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 4.5 | 10.9 | 22.2 |
| 2002..... | 1.7 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.6 | 11.4 | 20.2 |
| 2001..... | 1.7 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 5.0 | 11.2 | 19.3 |
| 2000..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 5.7 | 11.4 | 18.5 |
| 1999..... | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 5.5 | 12.1 | 21.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 1.3 | 3.5 | 6.3 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.5 | 1.0 | 3.3 | 6.7 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.7 | 1.8 | 4.5 | 7.0 | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 1.5 | 4.1 | 5.4 | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 1.8 | 4.3 | 5.3 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.8 | 1.6 | 4.2 | 7.3 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 2.0 | 3.7 | 6.4 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 4.0 | 8.4 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 2.1 | 5.8 | 10.7 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 2.0 | 5.2 | * | * |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 5.6 | 6.9 | * |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 5.9 | 10.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic rheumatic heart diseases (I05-I09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.3 | 8.9 | 17.6 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.8 | 9.6 | 20.8 |
| 2002..... | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 3.8 | 9.6 | 17.9 |
| 2001..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.0 | 9.5 | 17.5 |
| 2000..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.4 | 9.6 | 17.0 |
| 1999..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.7 | 4.6 | 10.1 | 18.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.3 | 7.3 | 11.8 |
| 2003..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.9 | 7.6 | 16.4 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.9 | 6.9 | 11.7 |
| 2001..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.7 | 6.6 | 12.6 |
| 2000..... | 0.8 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 2.8 | 6.7 | 12.4 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.5 | 6.8 | 10.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.1 | 9.9 | 20.1 |
| 2003..... | 1.8 | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 4.5 | 11.0 | 22.7 |
| 2002..... | 1.8 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 4.5 | 11.4 | 20.6 |
| 2001..... | 1.8 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 5.0 | 11.4 | 19.5 |
| 2000..... | 1.9 | * | * | * | * | * | 0.3 | 0.8 | 2.0 | 5.7 | 11.6 | 18.8 |
| 1999..... | 2.0 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 5.5 | 12.2 | 21.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.3 | 9.4 | 18.8 |
| 2003..... | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.8 | 10.3 | 22.2 |
| 2002..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.9 | 10.1 | 19.0 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.0 | 10.0 | 18.6 |
| 2000..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.6 | 4.5 | 10.3 | 17.9 |
| 1999..... | 1.6 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 4.8 | 10.6 | 19.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.4 | 0.7 | 2.4 | 7.8 | 12.7 |
| 2003..... | 1.0 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.9 | 8.2 | 17.7 |
| 2002..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.0 | 7.2 | 12.5 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.7 | 6.9 | 13.2 |
| 2000..... | 0.9 | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 2.8 | 7.1 | 12.9 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.7 | 7.1 | 11.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.1 | 10.5 | 21.5 |
| 2003..... | 2.1 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.6 | 11.7 | 24.3 |
| 2002..... | 2.1 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 4.6 | 12.0 | 21.8 |
| 2001..... | 2.0 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 5.1 | 12.1 | 20.8 |
| 2000..... | 2.1 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 6.0 | 12.4 | 19.9 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.6 | 12.8 | 22.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic rheumatic heart diseases (I05-I09) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.6 | 1.0 | 1.5 | 2.6 | 4.0 | 6.0 |
| 2003..... | 0.8 | * | * | * | * | * | 0.5 | 1.3 | 2.6 | 3.9 | 3.9 | 7.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.0 | 2.4 | 2.9 | 4.2 | * |
| 2001..... | 0.8 | * | * | * | * | * | 0.7 | 1.4 | 2.1 | 4.0 | 4.3 | * |
| 2000..... | 0.7 | * | * | * | * | * | 0.6 | 1.0 | 2.0 | 3.5 | 4.0 | 8.3 |
| 1999..... | 0.8 | * | * | * | * | * | 0.5 | 1.2 | 2.1 | 3.4 | 4.3 | 7.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.9 | * | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 2.7 | 3.0 | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 1.8 | 2.9 | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | 1.0 | * | * | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 1.1 | 1.7 | 3.3 | 5.0 | * |
| 2003..... | 1.0 | * | * | * | * | * | * | 1.7 | 2.6 | 4.5 | 4.7 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.1 | 2.9 | 2.9 | 4.8 | * |
| 2001..... | 1.1 | * | * | * | * | * | 0.9 | 1.7 | 2.6 | 5.1 | 5.4 | * |
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | 1.3 | 2.7 | 4.3 | 4.9 | * |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.5 | 2.6 | 4.1 | 4.4 | 10.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rheumatic mitral valve diseases (I05) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.6 | 4.0 | 6.9 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 4.8 | 7.8 |
| 2002..... | 0.7 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.0 | 5.0 | 7.5 |
| 2001..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.3 | 5.2 | 8.6 |
| 2000..... | 0.8 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 5.5 | 8.7 |
| 1999..... | 0.8 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.6 | 5.7 | 9.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.0 | 4.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.5 | 5.9 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 3.1 | 4.3 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 3.3 | 6.4 |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.7 | 3.9 | 6.2 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 | 3.7 | 5.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.0 | 4.6 | 7.8 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.4 | 5.6 | 8.6 |
| 2002..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 2.3 | 6.3 | 8.9 |
| 2001..... | 1.0 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.9 | 6.4 | 9.5 |
| 2000..... | 1.1 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 6.5 | 9.7 |
| 1999..... | 1.1 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.2 | 7.0 | 10.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.7 | 4.1 | 7.4 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 5.0 | 8.3 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.0 | 5.3 | 8.0 |
| 2001..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.3 | 5.5 | 9.2 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.8 | 5.8 | 9.3 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 6.0 | 9.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 1.2 | 3.2 | 5.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 3.8 | 6.3 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 3.3 | 4.6 |
| 2001..... | 0.5 | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.5 | 3.5 | 6.7 |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.7 | 4.1 | 6.5 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.6 | 2.0 | 3.8 | 5.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 4.7 | 8.4 |
| 2003..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.5 | 5.9 | 9.1 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.3 | 6.7 | 9.5 |
| 2001..... | 1.1 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.9 | 6.8 | 10.2 |
| 2000..... | 1.2 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.7 | 7.0 | 10.5 |
| 1999..... | 1.3 | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.3 | 7.5 | 11.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rheumatic mitral valve diseases (I05) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.4 | 0.6 | 0.7 | 1.2 | 2.4 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.8 | 1.6 | 1.7 | 2.4 | * |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.6 | 1.6 | 1.3 | * | * |
| 2001..... | 0.5 | * | * | * | * | * | 0.4 | 0.9 | 1.2 | 2.4 | 2.3 | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 1.1 | 1.8 | 2.2 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.9 | 1.2 | 2.5 | 2.5 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 2.1 | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.5 | * | * | * | * | * | * | 1.0 | * | 2.1 | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 1.9 | * | * | * |
| 2001..... | 0.7 | * | * | * | * | * | * | 1.1 | 1.7 | 2.9 | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 1.5 | 2.3 | * | * |
| 1999..... | 0.6 | * | * | * | * | * | * | 1.2 | 1.6 | 3.1 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex; All ages 1/; Under 1 year 2/; 1-4 years; 5-14 years; 15-24 years; 25-34 years; 35-44 years; 45-54 years; 55-64 years; 65-74 years; 75-84 years; 85 years and over. Rows include: Disorders of both mitral and aortic valves (I08.0); Non-Hispanic 4/, both sexes; Non-Hispanic 4/, male; Non-Hispanic 4/, female; Non-Hispanic white, both sexes; Non-Hispanic white, male; Non-Hispanic white, female. Data points range from 0.1 to 7.3.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.6 | 4.5 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.1 | 2.5 | 5.5 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.2 | 2.7 | 5.7 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.2 | 3.1 | 6.0 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 3.2 | 5.9 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.6 | 1.5 | 3.4 | 6.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 2.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 3.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.0 | 3.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.0 | 3.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.0 | 4.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.2 | 2.3 | 3.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.2 | 5.4 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.0 | 6.4 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 3.1 | 6.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 3.7 | 6.9 |
| 2000..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 | 4.0 | 6.7 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 4.2 | 8.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.8 | 4.8 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.7 | 5.8 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 2.8 | 6.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.2 | 6.4 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.4 | 3.4 | 6.1 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 3.6 | 7.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 2.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.9 | 3.6 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 | 4.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.4 | 0.7 | 2.1 | 3.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.2 | 4.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 1.2 | 2.4 | 4.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.4 | 5.7 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 3.2 | 6.8 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 0.6 | 1.7 | 3.2 | 6.8 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.6 | 3.9 | 7.4 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 4.2 | 6.9 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.4 | 8.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive diseases (I10-I13) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 18.6 | * | * | * | 0.2 | 1.1 | 4.0 | 11.1 | 20.6 | 40.8 | 108.5 | 416.1 |
| 2003..... | 18.4 | * | * | * | 0.2 | 1.1 | 4.0 | 10.9 | 20.3 | 41.2 | 109.7 | 409.5 |
| 2002..... | 17.2 | * | * | * | 0.1 | 1.0 | 3.7 | 9.9 | 19.2 | 39.5 | 104.6 | 390.8 |
| 2001..... | 16.4 | * | * | * | 0.1 | 1.0 | 3.4 | 9.0 | 17.7 | 37.5 | 103.1 | 384.7 |
| 2000..... | 15.9 | * | * | * | 0.1 | 0.8 | 3.1 | 8.3 | 17.8 | 37.0 | 100.7 | 380.3 |
| 1999..... | 15.4 | * | * | * | 0.1 | 0.8 | 3.1 | 7.9 | 17.3 | 36.8 | 100.1 | 366.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 16.0 | * | * | * | 0.2 | 1.5 | 5.4 | 14.8 | 27.0 | 47.8 | 108.5 | 335.7 |
| 2003..... | 15.4 | * | * | * | 0.3 | 1.4 | 5.2 | 14.7 | 25.7 | 47.8 | 107.3 | 322.4 |
| 2002..... | 14.5 | * | * | * | 0.2 | 1.4 | 5.0 | 13.2 | 24.8 | 44.6 | 103.7 | 311.1 |
| 2001..... | 13.5 | * | * | * | 0.2 | 1.2 | 4.5 | 12.0 | 21.9 | 42.7 | 101.0 | 309.4 |
| 2000..... | 13.0 | * | * | * | 0.2 | 1.0 | 4.0 | 11.2 | 22.4 | 42.0 | 98.8 | 309.3 |
| 1999..... | 12.6 | * | * | * | 0.2 | 1.1 | 4.1 | 10.3 | 21.4 | 41.1 | 99.3 | 297.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 21.2 | * | * | * | 0.1 | 0.7 | 2.6 | 7.5 | 14.7 | 34.9 | 108.5 | 452.3 |
| 2003..... | 21.2 | * | * | * | 0.1 | 0.8 | 2.8 | 7.3 | 15.3 | 35.7 | 111.3 | 448.0 |
| 2002..... | 19.9 | * | * | * | * | 0.7 | 2.4 | 6.7 | 14.1 | 35.2 | 105.2 | 425.4 |
| 2001..... | 19.2 | * | * | * | * | 0.7 | 2.3 | 6.1 | 13.8 | 33.2 | 104.5 | 416.3 |
| 2000..... | 18.6 | * | * | * | 0.1 | 0.6 | 2.2 | 5.5 | 13.5 | 32.9 | 101.9 | 409.3 |
| 1999..... | 18.1 | * | * | * | * | 0.5 | 2.1 | 5.6 | 13.6 | 33.3 | 100.7 | 393.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | * | * | 0.5 | 2.3 | 7.4 | 17.7 | 40.1 | 100.2 | 342.5 |
| 2003..... | 7.0 | * | * | * | * | 0.5 | 2.3 | 8.0 | 18.5 | 42.3 | 104.6 | 375.7 |
| 2002..... | 6.5 | * | * | * | * | 0.6 | 2.3 | 6.3 | 19.7 | 40.7 | 107.1 | 337.0 |
| 2001..... | 6.1 | * | * | * | * | 0.5 | 2.2 | 6.9 | 15.6 | 40.5 | 103.2 | 338.3 |
| 2000..... | 5.5 | * | * | * | * | 0.5 | 1.9 | 6.0 | 15.5 | 35.9 | 98.2 | 316.5 |
| 1999..... | 5.4 | * | * | * | * | 0.4 | 1.8 | 5.9 | 15.1 | 35.1 | 101.2 | 305.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | 0.7 | 3.2 | 9.5 | 23.9 | 48.1 | 114.5 | 302.6 |
| 2003..... | 6.7 | * | * | * | * | 0.7 | 3.5 | 11.4 | 25.4 | 48.5 | 115.4 | 314.0 |
| 2002..... | 6.2 | * | * | * | * | 0.8 | 3.2 | 8.9 | 26.0 | 47.2 | 115.7 | 294.5 |
| 2001..... | 5.7 | * | * | * | * | 0.6 | 3.4 | 9.7 | 19.3 | 46.5 | 106.9 | 304.7 |
| 2000..... | 5.2 | * | * | * | * | 0.7 | 2.5 | 8.7 | 19.4 | 41.6 | 112.1 | 276.1 |
| 1999..... | 4.9 | * | * | * | * | 0.6 | 2.6 | 7.5 | 19.3 | 42.4 | 101.7 | 267.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.0 | * | * | * | * | * | 1.2 | 5.4 | 12.1 | 33.8 | 90.4 | 363.0 |
| 2003..... | 7.3 | * | * | * | * | * | 1.0 | 4.5 | 12.3 | 37.4 | 97.2 | 407.0 |
| 2002..... | 6.9 | * | * | * | * | * | 1.2 | 3.7 | 14.1 | 35.6 | 101.2 | 358.1 |
| 2001..... | 6.5 | * | * | * | * | * | 1.0 | 4.2 | 12.3 | 35.7 | 100.6 | 354.7 |
| 2000..... | 5.8 | * | * | * | * | * | 1.3 | 3.3 | 12.0 | 31.5 | 88.8 | 336.3 |
| 1999..... | 5.9 | * | * | * | * | * | 1.0 | 4.3 | 11.4 | 29.4 | 101.0 | 323.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive diseases (I10-I13) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.5 | * | * | * | 0.2 | 1.2 | 4.2 | 11.4 | 20.7 | 40.6 | 108.7 | 418.6 |
| 2003..... | 20.1 | * | * | * | 0.2 | 1.2 | 4.2 | 11.2 | 20.4 | 40.9 | 109.7 | 410.0 |
| 2002..... | 18.8 | * | * | * | 0.1 | 1.1 | 3.8 | 10.2 | 19.1 | 39.1 | 104.1 | 391.7 |
| 2001..... | 17.9 | * | * | * | 0.1 | 1.0 | 3.5 | 9.1 | 17.7 | 37.1 | 102.8 | 385.4 |
| 2000..... | 17.3 | * | * | * | 0.2 | 0.8 | 3.2 | 8.4 | 17.9 | 36.8 | 100.4 | 381.4 |
| 1999..... | 16.7 | * | * | * | 0.1 | 0.8 | 3.2 | 8.0 | 17.3 | 36.7 | 99.7 | 366.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 17.5 | * | * | * | 0.2 | 1.7 | 5.7 | 15.3 | 27.0 | 47.4 | 108.0 | 336.7 |
| 2003..... | 16.8 | * | * | * | 0.2 | 1.5 | 5.5 | 14.9 | 25.6 | 47.4 | 106.6 | 322.3 |
| 2002..... | 15.7 | * | * | * | 0.2 | 1.5 | 5.2 | 13.5 | 24.5 | 44.0 | 102.6 | 310.5 |
| 2001..... | 14.6 | * | * | * | 0.2 | 1.3 | 4.6 | 12.1 | 21.8 | 42.2 | 100.4 | 308.4 |
| 2000..... | 14.1 | * | * | * | 0.2 | 1.0 | 4.2 | 11.3 | 22.5 | 41.6 | 97.7 | 309.5 |
| 1999..... | 13.6 | * | * | * | 0.2 | 1.1 | 4.3 | 10.4 | 21.3 | 40.8 | 98.7 | 297.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 23.3 | * | * | * | 0.1 | 0.8 | 2.8 | 7.7 | 14.9 | 34.8 | 109.2 | 455.2 |
| 2003..... | 23.2 | * | * | * | 0.1 | 0.9 | 3.0 | 7.6 | 15.5 | 35.4 | 111.8 | 448.5 |
| 2002..... | 21.7 | * | * | * | * | 0.7 | 2.5 | 7.0 | 14.1 | 35.0 | 105.1 | 426.7 |
| 2001..... | 21.0 | * | * | * | * | 0.7 | 2.4 | 6.3 | 13.9 | 32.9 | 104.4 | 417.5 |
| 2000..... | 20.3 | * | * | * | 0.1 | 0.6 | 2.3 | 5.6 | 13.6 | 32.9 | 102.2 | 410.5 |
| 1999..... | 19.7 | * | * | * | * | 0.5 | 2.2 | 5.7 | 13.6 | 33.4 | 100.4 | 394.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 19.1 | * | * | * | 0.1 | 0.7 | 2.7 | 7.4 | 14.4 | 31.4 | 94.7 | 396.0 |
| 2003..... | 18.6 | * | * | * | 0.1 | 0.7 | 2.6 | 7.1 | 14.1 | 30.7 | 96.4 | 389.1 |
| 2002..... | 17.3 | * | * | * | * | 0.6 | 2.3 | 6.7 | 13.0 | 29.8 | 90.0 | 368.4 |
| 2001..... | 16.5 | * | * | * | * | 0.6 | 2.0 | 5.6 | 12.0 | 27.9 | 89.6 | 363.9 |
| 2000..... | 15.7 | * | * | * | * | 0.4 | 2.0 | 5.1 | 11.7 | 27.6 | 86.6 | 357.0 |
| 1999..... | 15.1 | * | * | * | * | 0.5 | 1.7 | 4.8 | 11.4 | 27.3 | 85.1 | 343.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 15.6 | * | * | * | 0.2 | 1.0 | 3.9 | 10.6 | 19.3 | 37.3 | 93.7 | 318.3 |
| 2003..... | 14.9 | * | * | * | * | 1.0 | 3.8 | 10.1 | 18.5 | 36.4 | 93.7 | 299.9 |
| 2002..... | 13.8 | * | * | * | * | 0.8 | 3.4 | 9.4 | 17.3 | 34.0 | 88.1 | 289.1 |
| 2001..... | 12.8 | * | * | * | * | 0.8 | 2.9 | 7.8 | 15.5 | 32.3 | 87.0 | 286.3 |
| 2000..... | 12.2 | * | * | * | * | 0.6 | 2.8 | 7.4 | 15.4 | 32.0 | 84.4 | 285.6 |
| 1999..... | 11.7 | * | * | * | * | 0.8 | 2.5 | 6.6 | 14.6 | 30.7 | 84.1 | 272.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 22.4 | * | * | * | * | 0.3 | 1.5 | 4.2 | 9.8 | 26.4 | 95.4 | 430.8 |
| 2003..... | 22.2 | * | * | * | * | 0.4 | 1.5 | 4.1 | 10.1 | 25.9 | 98.2 | 428.3 |
| 2002..... | 20.7 | * | * | * | * | 0.3 | 1.3 | 4.0 | 8.9 | 26.2 | 91.2 | 402.6 |
| 2001..... | 20.0 | * | * | * | * | 0.3 | 1.1 | 3.4 | 8.7 | 24.0 | 91.3 | 396.2 |
| 2000..... | 19.1 | * | * | * | * | 0.2 | 1.2 | 2.9 | 8.3 | 23.9 | 88.1 | 385.8 |
| 1999..... | 18.4 | * | * | * | * | 0.2 | 0.9 | 3.0 | 8.3 | 24.5 | 85.8 | 371.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive diseases (I10-I13) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 33.3 | * | * | * | 0.5 | 4.1 | 14.0 | 40.1 | 75.2 | 126.8 | 274.0 | 707.5 |
| 2003..... | 33.2 | * | * | * | 0.6 | 4.0 | 14.2 | 41.0 | 75.4 | 134.7 | 269.3 | 693.1 |
| 2002..... | 31.7 | * | * | * | 0.4 | 4.1 | 13.2 | 36.7 | 72.8 | 126.6 | 273.9 | 691.7 |
| 2001..... | 30.1 | * | * | * | 0.4 | 3.7 | 12.8 | 35.5 | 66.9 | 124.6 | 259.7 | 670.8 |
| 2000..... | 29.9 | * | * | * | 0.6 | 3.2 | 11.4 | 33.7 | 71.1 | 127.1 | 264.1 | 680.8 |
| 1999..... | 29.5 | * | * | * | 0.6 | 2.8 | 12.8 | 33.0 | 68.1 | 127.6 | 272.8 | 646.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 32.3 | * | * | * | * | 5.1 | 18.0 | 50.9 | 100.5 | 153.9 | 296.6 | 582.2 |
| 2003..... | 31.8 | * | * | * | 0.9 | 4.9 | 16.9 | 52.6 | 94.7 | 163.8 | 279.0 | 623.3 |
| 2002..... | 30.2 | * | * | * | * | 5.6 | 17.0 | 47.0 | 93.5 | 146.9 | 290.7 | 603.9 |
| 2001..... | 28.5 | * | * | * | * | 4.5 | 15.9 | 46.8 | 81.9 | 147.3 | 275.8 | 611.0 |
| 2000..... | 27.6 | * | * | * | * | 3.8 | 14.0 | 43.4 | 89.7 | 144.5 | 270.0 | 609.4 |
| 1999..... | 27.6 | * | * | * | 0.8 | 3.3 | 15.9 | 42.1 | 83.7 | 146.7 | 288.3 | 595.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 34.2 | * | * | * | * | 3.3 | 10.4 | 30.8 | 54.9 | 107.5 | 261.2 | 757.0 |
| 2003..... | 34.4 | * | * | * | * | 3.1 | 11.8 | 31.1 | 59.8 | 114.0 | 263.8 | 720.1 |
| 2002..... | 33.0 | * | * | * | * | 2.7 | 9.7 | 27.9 | 56.1 | 112.3 | 264.5 | 725.3 |
| 2001..... | 31.5 | * | * | * | * | 3.0 | 10.1 | 25.8 | 54.8 | 108.6 | 250.7 | 693.0 |
| 2000..... | 31.9 | * | * | * | * | 2.7 | 9.2 | 25.4 | 56.2 | 114.8 | 260.8 | 707.3 |
| 1999..... | 31.3 | * | * | * | * | 2.4 | 10.0 | 25.2 | 55.5 | 114.2 | 264.2 | 665.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Essential (primary) hypertension (I10) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | 0.1 | 0.3 | 1.6 | 3.6 | 9.1 | 28.5 | 127.4 |
| 2003..... | 4.4 | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.6 | 9.2 | 28.5 | 121.9 |
| 2002..... | 4.0 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.2 | 8.6 | 26.3 | 112.0 |
| 2001..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.4 | 8.4 | 25.9 | 107.4 |
| 2000..... | 3.7 | * | * | * | * | 0.1 | 0.3 | 1.4 | 3.4 | 8.2 | 24.1 | 101.1 |
| 1999..... | 3.6 | * | * | * | * | 0.1 | 0.3 | 1.2 | 3.3 | 8.1 | 24.0 | 98.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | 0.5 | 2.1 | 4.4 | 10.0 | 26.6 | 86.5 |
| 2003..... | 3.0 | * | * | * | * | * | 0.4 | 1.9 | 4.5 | 10.4 | 24.3 | 77.4 |
| 2002..... | 2.8 | * | * | * | * | * | 0.5 | 1.7 | 4.0 | 9.4 | 23.3 | 74.2 |
| 2001..... | 2.8 | * | * | * | * | 0.1 | 0.4 | 1.7 | 4.2 | 9.3 | 23.2 | 74.8 |
| 2000..... | 2.6 | * | * | * | * | * | 0.4 | 1.9 | 4.2 | 8.9 | 22.0 | 70.3 |
| 1999..... | 2.5 | * | * | * | * | * | 0.4 | 1.5 | 4.1 | 9.2 | 21.9 | 68.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | * | * | 0.2 | 1.1 | 2.9 | 8.3 | 29.8 | 145.8 |
| 2003..... | 5.8 | * | * | * | * | * | 0.4 | 1.0 | 2.9 | 8.3 | 31.3 | 141.5 |
| 2002..... | 5.2 | * | * | * | * | * | 0.3 | 1.0 | 2.5 | 8.0 | 28.2 | 128.3 |
| 2001..... | 5.0 | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 7.7 | 27.6 | 121.1 |
| 2000..... | 4.6 | * | * | * | * | * | 0.3 | 0.9 | 2.6 | 7.5 | 25.5 | 113.6 |
| 1999..... | 4.5 | * | * | * | * | * | 0.2 | 1.0 | 2.7 | 7.3 | 25.4 | 110.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.0 | 2.0 | 5.7 | 20.0 | 102.3 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.9 | 2.2 | 6.6 | 16.9 | 104.2 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.5 | 2.2 | 5.6 | 20.4 | 80.9 |
| 2001..... | 1.1 | * | * | * | * | * | * | 0.8 | 1.5 | 6.2 | 24.2 | 90.0 |
| 2000..... | 1.1 | * | * | * | * | * | * | 0.8 | 2.9 | 6.1 | 21.3 | 80.3 |
| 1999..... | 1.0 | * | * | * | * | * | * | 0.8 | 2.2 | 5.6 | 21.2 | 72.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 1.3 | 2.3 | 6.0 | 21.1 | 65.8 |
| 2003..... | 0.8 | * | * | * | * | * | * | 1.3 | 2.5 | 6.5 | 13.8 | 67.7 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 2.6 | 5.2 | 14.8 | 58.6 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 6.2 | 24.6 | 53.3 |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.3 | 3.7 | 6.3 | 25.1 | 72.6 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 7.0 | 21.9 | 51.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | * | 1.7 | 5.4 | 19.3 | 121.0 |
| 2003..... | 1.6 | * | * | * | * | * | * | * | 1.9 | 6.8 | 19.0 | 122.7 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 1.9 | 5.9 | 24.2 | 92.0 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | * | 6.1 | 23.9 | 108.0 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 6.0 | 18.8 | 84.1 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | * | 4.5 | 20.8 | 82.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Essential (primary) hypertension (I10) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | * | 0.1 | 0.4 | 1.6 | 3.7 | 9.3 | 29.0 | 128.4 |
| 2003..... | 4.9 | * | * | * | * | 0.1 | 0.4 | 1.5 | 3.7 | 9.4 | 29.0 | 122.4 |
| 2002..... | 4.5 | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.3 | 8.8 | 26.5 | 112.7 |
| 2001..... | 4.3 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.5 | 8.5 | 25.9 | 107.9 |
| 2000..... | 4.0 | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.4 | 8.3 | 24.2 | 101.4 |
| 1999..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.4 | 8.3 | 24.0 | 98.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.5 | 2.1 | 4.5 | 10.2 | 26.9 | 87.4 |
| 2003..... | 3.4 | * | * | * | * | * | 0.5 | 1.9 | 4.6 | 10.6 | 24.8 | 77.8 |
| 2002..... | 3.1 | * | * | * | * | * | 0.5 | 1.8 | 4.1 | 9.6 | 23.6 | 74.5 |
| 2001..... | 3.0 | * | * | * | * | * | 0.4 | 1.7 | 4.3 | 9.5 | 23.0 | 75.4 |
| 2000..... | 2.9 | * | * | * | * | * | 0.4 | 1.9 | 4.2 | 9.0 | 21.7 | 70.0 |
| 1999..... | 2.8 | * | * | * | * | * | 0.4 | 1.5 | 4.2 | 9.3 | 21.7 | 68.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | 0.2 | 1.2 | 2.9 | 8.5 | 30.3 | 146.7 |
| 2003..... | 6.4 | * | * | * | * | * | 0.4 | 1.0 | 3.0 | 8.3 | 31.8 | 142.0 |
| 2002..... | 5.8 | * | * | * | * | * | 0.3 | 1.0 | 2.6 | 8.1 | 28.3 | 129.2 |
| 2001..... | 5.5 | * | * | * | * | * | 0.3 | 0.9 | 2.8 | 7.8 | 27.7 | 121.3 |
| 2000..... | 5.1 | * | * | * | * | * | 0.3 | 0.9 | 2.6 | 7.6 | 25.8 | 114.2 |
| 1999..... | 5.0 | * | * | * | * | * | 0.2 | 1.0 | 2.7 | 7.4 | 25.5 | 111.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | * | * | 0.3 | 1.2 | 3.0 | 8.1 | 27.0 | 124.2 |
| 2003..... | 5.0 | * | * | * | * | * | 0.3 | 1.2 | 3.0 | 7.7 | 26.5 | 120.3 |
| 2002..... | 4.6 | * | * | * | * | * | 0.3 | 1.0 | 2.8 | 7.6 | 24.3 | 109.1 |
| 2001..... | 4.3 | * | * | * | * | * | 0.2 | 0.9 | 2.8 | 7.1 | 23.9 | 104.8 |
| 2000..... | 4.0 | * | * | * | * | * | 0.3 | 1.0 | 2.7 | 7.2 | 22.2 | 97.7 |
| 1999..... | 3.9 | * | * | * | * | * | 0.2 | 0.9 | 2.7 | 7.2 | 21.9 | 94.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.4 | 1.7 | 3.8 | 9.0 | 24.7 | 84.7 |
| 2003..... | 3.3 | * | * | * | * | * | 0.4 | 1.6 | 3.7 | 9.0 | 21.8 | 74.4 |
| 2002..... | 3.1 | * | * | * | * | * | 0.4 | 1.4 | 3.6 | 8.4 | 21.2 | 71.3 |
| 2001..... | 2.9 | * | * | * | * | * | 0.3 | 1.3 | 3.5 | 8.0 | 20.5 | 71.6 |
| 2000..... | 2.7 | * | * | * | * | * | 0.3 | 1.4 | 3.3 | 8.0 | 19.5 | 65.7 |
| 1999..... | 2.7 | * | * | * | * | * | 0.3 | 1.2 | 3.5 | 8.1 | 19.5 | 62.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 7.4 | 28.5 | 141.9 |
| 2003..... | 6.7 | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 6.5 | 29.7 | 140.5 |
| 2002..... | 6.0 | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 6.9 | 26.4 | 125.5 |
| 2001..... | 5.7 | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 6.4 | 26.1 | 118.7 |
| 2000..... | 5.3 | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 6.6 | 23.9 | 110.5 |
| 1999..... | 5.1 | * | * | * | * | * | * | 0.6 | 2.1 | 6.5 | 23.5 | 107.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Essential (primary) hypertension (I10) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.7 | * | * | * | * | * | 1.0 | 4.7 | 9.9 | 20.5 | 53.9 | 185.9 |
| 2003..... | 5.7 | * | * | * | * | * | 1.1 | 3.9 | 10.9 | 25.5 | 60.6 | 159.4 |
| 2002..... | 5.1 | * | * | * | * | * | 1.2 | 4.1 | 8.0 | 21.0 | 53.8 | 160.3 |
| 2001..... | 5.1 | * | * | * | * | * | 1.1 | 4.1 | 10.1 | 22.0 | 50.8 | 152.3 |
| 2000..... | 4.9 | * | * | * | * | * | 1.0 | 4.5 | 9.9 | 19.1 | 48.3 | 150.7 |
| 1999..... | 4.8 | * | * | * | * | * | 1.1 | 4.1 | 9.5 | 19.3 | 49.3 | 145.2 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 1.3 | 5.4 | 11.9 | 22.7 | 58.2 | 127.0 |
| 2003..... | 4.9 | * | * | * | * | * | 1.1 | 5.1 | 13.6 | 28.2 | 64.1 | 122.5 |
| 2002..... | 4.2 | * | * | * | * | * | 1.5 | 4.9 | 9.6 | 23.0 | 56.3 | 119.2 |
| 2001..... | 4.5 | * | * | * | * | * | 1.3 | 5.3 | 12.6 | 25.5 | 56.4 | 132.8 |
| 2000..... | 4.2 | * | * | * | * | * | 1.0 | 5.9 | 13.3 | 21.1 | 49.7 | 124.5 |
| 1999..... | 4.0 | * | * | * | * | * | 1.2 | 4.4 | 11.3 | 22.7 | 50.7 | 126.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | * | 0.7 | 4.0 | 8.3 | 19.0 | 51.5 | 209.2 |
| 2003..... | 6.5 | * | * | * | * | * | 1.1 | 2.8 | 8.7 | 23.5 | 58.7 | 173.7 |
| 2002..... | 6.0 | * | * | * | * | * | 0.9 | 3.4 | 6.7 | 19.6 | 52.4 | 176.0 |
| 2001..... | 5.7 | * | * | * | * | * | 1.0 | 3.1 | 8.1 | 19.6 | 47.7 | 159.5 |
| 2000..... | 5.5 | * | * | * | * | * | 1.0 | 3.3 | 7.1 | 17.7 | 47.6 | 160.5 |
| 1999..... | 5.5 | * | * | * | * | * | 1.0 | 3.8 | 8.1 | 16.9 | 48.5 | 152.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive heart disease (I11) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.7 | * | * | * | 0.1 | 0.8 | 3.0 | 7.8 | 13.4 | 21.7 | 49.3 | 192.8 |
| 2003..... | 9.7 | * | * | * | 0.1 | 0.8 | 2.9 | 7.9 | 13.1 | 22.1 | 51.0 | 195.0 |
| 2002..... | 9.2 | * | * | * | 0.1 | 0.7 | 2.7 | 7.2 | 12.5 | 21.3 | 49.6 | 187.0 |
| 2001..... | 8.7 | * | * | * | 0.1 | 0.6 | 2.4 | 6.3 | 10.9 | 19.8 | 48.7 | 189.5 |
| 2000..... | 8.4 | * | * | * | 0.1 | 0.5 | 2.2 | 5.6 | 11.0 | 19.6 | 48.2 | 193.1 |
| 1999..... | 8.1 | * | * | * | 0.1 | 0.5 | 2.1 | 5.3 | 10.7 | 19.1 | 47.7 | 184.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 8.9 | * | * | * | 0.1 | 1.1 | 4.1 | 10.7 | 18.5 | 27.3 | 48.0 | 145.4 |
| 2003..... | 8.8 | * | * | * | 0.2 | 1.0 | 4.0 | 10.8 | 17.2 | 27.1 | 51.0 | 142.2 |
| 2002..... | 8.2 | * | * | * | 0.1 | 1.1 | 3.8 | 9.8 | 16.9 | 25.2 | 49.4 | 134.6 |
| 2001..... | 7.5 | * | * | * | * | 0.8 | 3.3 | 8.6 | 14.4 | 23.6 | 48.4 | 134.4 |
| 2000..... | 7.2 | * | * | * | 0.1 | 0.7 | 2.9 | 7.8 | 14.5 | 23.7 | 46.5 | 139.1 |
| 1999..... | 6.8 | * | * | * | 0.1 | 0.7 | 2.9 | 7.0 | 13.7 | 21.9 | 46.8 | 133.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 10.5 | * | * | * | * | 0.4 | 1.9 | 5.1 | 8.6 | 16.9 | 50.2 | 214.1 |
| 2003..... | 10.7 | * | * | * | * | 0.5 | 1.9 | 5.2 | 9.3 | 17.8 | 51.0 | 218.3 |
| 2002..... | 10.2 | * | * | * | * | 0.4 | 1.6 | 4.6 | 8.4 | 18.1 | 49.8 | 209.7 |
| 2001..... | 9.8 | * | * | * | * | 0.4 | 1.6 | 4.1 | 7.8 | 16.6 | 48.8 | 212.6 |
| 2000..... | 9.7 | * | * | * | * | 0.4 | 1.5 | 3.5 | 7.8 | 16.2 | 49.4 | 215.1 |
| 1999..... | 9.4 | * | * | * | * | 0.3 | 1.3 | 3.6 | 7.9 | 16.8 | 48.3 | 205.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | 0.3 | 1.7 | 4.8 | 11.0 | 22.2 | 47.0 | 150.3 |
| 2003..... | 4.0 | * | * | * | * | 0.4 | 1.8 | 5.7 | 11.8 | 22.9 | 54.2 | 179.3 |
| 2002..... | 3.7 | * | * | * | * | 0.4 | 1.7 | 4.9 | 12.2 | 22.4 | 51.4 | 165.7 |
| 2001..... | 3.3 | * | * | * | * | 0.3 | 1.7 | 4.9 | 10.0 | 21.0 | 46.9 | 151.9 |
| 2000..... | 2.8 | * | * | * | * | 0.3 | 1.2 | 3.8 | 8.2 | 18.4 | 46.0 | 143.3 |
| 1999..... | 2.7 | * | * | * | * | * | 1.2 | 3.3 | 8.9 | 16.8 | 44.8 | 152.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | 2.7 | 6.1 | 16.0 | 27.8 | 55.4 | 131.7 |
| 2003..... | 4.3 | * | * | * | * | 0.5 | 2.9 | 8.3 | 17.0 | 30.8 | 63.7 | 144.7 |
| 2002..... | 3.9 | * | * | * | * | 0.6 | 2.6 | 7.4 | 17.0 | 28.5 | 61.1 | 137.2 |
| 2001..... | 3.4 | * | * | * | * | * | 2.7 | 7.0 | 13.8 | 26.4 | 48.3 | 144.7 |
| 2000..... | 2.7 | * | * | * | * | * | 1.6 | 5.4 | 10.7 | 22.4 | 50.6 | 112.9 |
| 1999..... | 2.6 | * | * | * | * | * | 1.8 | 4.7 | 12.5 | 20.9 | 42.8 | 107.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | * | 3.4 | 6.4 | 17.7 | 41.2 | 159.9 |
| 2003..... | 3.7 | * | * | * | * | * | * | 3.1 | 7.2 | 16.6 | 47.7 | 196.9 |
| 2002..... | 3.4 | * | * | * | * | * | * | 2.5 | 7.9 | 17.5 | 44.7 | 179.9 |
| 2001..... | 3.1 | * | * | * | * | * | * | 2.7 | 6.5 | 16.8 | 45.9 | 155.5 |
| 2000..... | 2.9 | * | * | * | * | * | 0.8 | 2.1 | 6.1 | 15.3 | 42.9 | 158.3 |
| 1999..... | 2.9 | * | * | * | * | * | * | 2.0 | 5.8 | 13.6 | 46.1 | 173.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive heart disease (I11) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 10.7 | * | * | * | 0.1 | 0.9 | 3.2 | 8.1 | 13.5 | 21.5 | 49.3 | 194.3 |
| 2003..... | 10.6 | * | * | * | 0.1 | 0.9 | 3.1 | 8.1 | 13.1 | 21.9 | 50.7 | 195.0 |
| 2002..... | 10.0 | * | * | * | 0.1 | 0.8 | 2.8 | 7.4 | 12.4 | 21.0 | 49.3 | 187.1 |
| 2001..... | 9.4 | * | * | * | * | 0.7 | 2.5 | 6.3 | 10.9 | 19.6 | 48.6 | 190.3 |
| 2000..... | 9.2 | * | * | * | 0.1 | 0.6 | 2.3 | 5.7 | 11.2 | 19.5 | 48.1 | 194.3 |
| 1999..... | 8.8 | * | * | * | 0.1 | 0.6 | 2.2 | 5.4 | 10.7 | 19.1 | 47.6 | 185.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 9.8 | * | * | * | 0.2 | 1.3 | 4.3 | 11.1 | 18.5 | 27.1 | 47.4 | 145.6 |
| 2003..... | 9.5 | * | * | * | 0.2 | 1.1 | 4.1 | 10.9 | 17.1 | 26.7 | 50.1 | 141.7 |
| 2002..... | 8.9 | * | * | * | 0.1 | 1.2 | 4.0 | 10.0 | 16.8 | 24.6 | 48.5 | 133.7 |
| 2001..... | 8.1 | * | * | * | * | 0.9 | 3.4 | 8.6 | 14.2 | 23.3 | 48.3 | 133.4 |
| 2000..... | 7.8 | * | * | * | 0.1 | 0.7 | 3.0 | 7.9 | 14.6 | 23.5 | 46.0 | 139.6 |
| 1999..... | 7.4 | * | * | * | 0.1 | 0.8 | 3.1 | 7.2 | 13.6 | 21.7 | 46.7 | 134.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 11.6 | * | * | * | * | 0.5 | 2.1 | 5.2 | 8.8 | 16.7 | 50.6 | 216.0 |
| 2003..... | 11.7 | * | * | * | * | 0.6 | 2.1 | 5.4 | 9.4 | 17.8 | 51.1 | 218.5 |
| 2002..... | 11.1 | * | * | * | * | 0.5 | 1.7 | 4.8 | 8.4 | 18.0 | 49.9 | 210.2 |
| 2001..... | 10.7 | * | * | * | * | 0.4 | 1.7 | 4.2 | 7.8 | 16.5 | 48.8 | 214.0 |
| 2000..... | 10.6 | * | * | * | * | 0.4 | 1.6 | 3.6 | 8.0 | 16.2 | 49.5 | 216.5 |
| 1999..... | 10.2 | * | * | * | * | 0.3 | 1.4 | 3.7 | 8.0 | 16.9 | 48.2 | 205.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 9.8 | * | * | * | * | 0.5 | 2.2 | 5.6 | 9.8 | 17.0 | 42.7 | 184.6 |
| 2003..... | 9.7 | * | * | * | * | 0.5 | 2.0 | 5.3 | 9.5 | 16.7 | 44.3 | 184.9 |
| 2002..... | 9.1 | * | * | * | * | 0.4 | 1.9 | 5.1 | 8.8 | 16.1 | 42.2 | 177.0 |
| 2001..... | 8.6 | * | * | * | * | 0.4 | 1.6 | 4.1 | 7.6 | 14.8 | 42.2 | 180.4 |
| 2000..... | 8.3 | * | * | * | * | 0.3 | 1.5 | 3.6 | 7.5 | 14.5 | 41.2 | 182.5 |
| 1999..... | 7.9 | * | * | * | * | 0.3 | 1.3 | 3.4 | 7.3 | 14.1 | 40.6 | 175.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 8.6 | * | * | * | * | 0.8 | 3.2 | 8.3 | 13.6 | 21.6 | 40.9 | 135.9 |
| 2003..... | 8.2 | * | * | * | * | 0.8 | 3.0 | 7.8 | 13.0 | 20.4 | 43.5 | 130.8 |
| 2002..... | 7.6 | * | * | * | * | 0.7 | 2.8 | 7.3 | 12.2 | 19.0 | 41.2 | 123.5 |
| 2001..... | 6.9 | * | * | * | * | 0.6 | 2.3 | 5.9 | 10.5 | 18.1 | 41.5 | 122.0 |
| 2000..... | 6.6 | * | * | * | * | 0.4 | 2.1 | 5.4 | 10.3 | 18.0 | 38.8 | 129.5 |
| 1999..... | 6.2 | * | * | * | * | 0.6 | 2.0 | 4.9 | 9.6 | 16.2 | 39.4 | 123.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 11.0 | * | * | * | * | * | 1.2 | 3.0 | 6.2 | 13.0 | 43.9 | 206.5 |
| 2003..... | 11.1 | * | * | * | * | 0.3 | 1.1 | 2.8 | 6.3 | 13.6 | 44.9 | 208.7 |
| 2002..... | 10.5 | * | * | * | * | 0.2 | 1.0 | 2.9 | 5.5 | 13.6 | 42.9 | 200.1 |
| 2001..... | 10.2 | * | * | * | * | 0.2 | 0.8 | 2.3 | 4.9 | 11.9 | 42.6 | 204.7 |
| 2000..... | 9.9 | * | * | * | * | 0.2 | 0.8 | 1.9 | 4.9 | 11.6 | 42.7 | 203.9 |
| 1999..... | 9.6 | * | * | * | * | * | 0.6 | 2.0 | 5.0 | 12.4 | 41.4 | 195.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive heart disease (I11) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 18.2 | * | * | * | 0.3 | 2.8 | 9.5 | 26.7 | 45.8 | 65.4 | 129.1 | 318.6 |
| 2003..... | 18.5 | * | * | * | 0.4 | 2.6 | 9.8 | 28.5 | 45.1 | 70.3 | 128.4 | 334.9 |
| 2002..... | 17.7 | * | * | * | * | 2.9 | 8.8 | 24.8 | 45.1 | 68.3 | 134.2 | 322.1 |
| 2001..... | 16.5 | * | * | * | * | 2.2 | 8.5 | 23.2 | 39.4 | 66.1 | 125.6 | 328.7 |
| 2000..... | 16.6 | * | * | * | * | 2.2 | 7.5 | 21.6 | 43.1 | 67.6 | 132.7 | 344.1 |
| 1999..... | 16.0 | * | * | * | 0.4 | 1.9 | 8.1 | 21.3 | 40.1 | 67.4 | 132.7 | 308.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 19.0 | * | * | * | * | 3.7 | 12.1 | 34.0 | 65.4 | 87.5 | 133.8 | 271.3 |
| 2003..... | 18.7 | * | * | * | * | 3.2 | 11.5 | 35.6 | 57.5 | 93.4 | 139.1 | 291.9 |
| 2002..... | 18.0 | * | * | * | * | 4.1 | 11.7 | 31.9 | 61.9 | 84.0 | 141.8 | 264.7 |
| 2001..... | 16.3 | * | * | * | * | 2.7 | 10.5 | 30.4 | 50.1 | 79.8 | 137.4 | 288.7 |
| 2000..... | 16.1 | * | * | * | * | 2.7 | 9.2 | 28.4 | 56.0 | 81.9 | 139.5 | 275.9 |
| 1999..... | 15.6 | * | * | * | * | 2.2 | 10.3 | 26.8 | 51.1 | 80.6 | 143.2 | 266.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 17.5 | * | * | * | * | 2.0 | 7.3 | 20.4 | 30.1 | 49.6 | 126.4 | 337.3 |
| 2003..... | 18.4 | * | * | * | * | 2.1 | 8.2 | 22.4 | 35.1 | 53.9 | 122.4 | 351.6 |
| 2002..... | 17.4 | * | * | * | * | 1.8 | 6.2 | 18.7 | 31.7 | 57.1 | 129.8 | 344.1 |
| 2001..... | 16.7 | * | * | * | * | 1.7 | 6.7 | 17.1 | 30.8 | 56.5 | 118.9 | 343.5 |
| 2000..... | 17.1 | * | * | * | * | 1.8 | 5.9 | 15.8 | 32.7 | 57.5 | 128.9 | 369.4 |
| 1999..... | 16.4 | * | * | * | * | 1.6 | 6.1 | 16.5 | 31.3 | 58.1 | 126.8 | 324.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive renal disease (I12) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.7 | 8.0 | 24.0 | 71.1 |
| 2003..... | 3.1 | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.6 | 7.7 | 23.2 | 67.0 |
| 2002..... | 3.0 | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.5 | 7.3 | 21.9 | 68.4 |
| 2001..... | 2.9 | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.4 | 7.0 | 21.8 | 64.5 |
| 2000..... | 2.8 | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.6 | 7.0 | 21.4 | 61.8 |
| 1999..... | 2.5 | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.2 | 7.1 | 19.6 | 53.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.0 | 8.2 | 26.7 | 79.5 |
| 2003..... | 2.7 | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.9 | 8.0 | 24.7 | 76.1 |
| 2002..... | 2.6 | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.7 | 7.7 | 24.1 | 79.2 |
| 2001..... | 2.4 | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.3 | 7.3 | 22.6 | 76.0 |
| 2000..... | 2.4 | * | * | * | * | 0.1 | 0.5 | 1.1 | 2.8 | 6.8 | 23.2 | 72.0 |
| 1999..... | 2.2 | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.4 | 7.4 | 21.1 | 64.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.5 | 7.8 | 22.3 | 67.4 |
| 2003..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.5 | 22.2 | 63.0 |
| 2002..... | 3.4 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 7.1 | 20.6 | 63.7 |
| 2001..... | 3.3 | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.5 | 6.8 | 21.3 | 59.7 |
| 2000..... | 3.1 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.1 | 20.2 | 57.7 |
| 1999..... | 2.8 | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 6.8 | 18.6 | 49.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 1.2 | 3.5 | 10.5 | 24.3 | 69.9 |
| 2003..... | 1.4 | * | * | * | * | * | * | 1.1 | 3.6 | 10.1 | 27.6 | 66.9 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 4.0 | 9.9 | 26.9 | 67.6 |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.0 | 3.0 | 10.3 | 24.7 | 70.0 |
| 2000..... | 1.2 | * | * | * | * | * | * | 0.9 | 3.5 | 8.9 | 24.5 | 67.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.3 | 2.8 | 9.7 | 27.1 | 57.5 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 1.5 | 4.2 | 12.4 | 27.5 | 81.3 |
| 2003..... | 1.3 | * | * | * | * | * | * | 1.5 | 4.8 | 9.0 | 30.7 | 66.2 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | 4.5 | 10.1 | 31.6 | 73.6 |
| 2001..... | 1.1 | * | * | * | * | * | * | 1.2 | * | 10.6 | 26.4 | 76.2 |
| 2000..... | 1.1 | * | * | * | * | * | * | 1.5 | 4.4 | 8.9 | 27.5 | 68.5 |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.4 | 3.0 | 11.4 | 29.7 | 79.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | * | 2.8 | 8.9 | 22.0 | 64.1 |
| 2003..... | 1.6 | * | * | * | * | * | * | * | 2.5 | 10.9 | 25.5 | 67.2 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | 3.6 | 9.7 | 23.6 | 64.7 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 3.8 | 10.1 | 23.6 | 67.0 |
| 2000..... | 1.3 | * | * | * | * | * | * | * | 2.8 | 9.0 | 22.5 | 67.3 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | 2.6 | 8.3 | 25.3 | 46.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive renal disease (I12) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.7 | 7.8 | 24.0 | 71.0 |
| 2003..... | 3.4 | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.6 | 7.5 | 22.9 | 66.9 |
| 2002..... | 3.2 | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.4 | 7.1 | 21.7 | 68.4 |
| 2001..... | 3.1 | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.3 | 6.8 | 21.6 | 64.2 |
| 2000..... | 3.0 | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.5 | 6.8 | 21.2 | 61.4 |
| 1999..... | 2.7 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.2 | 6.9 | 19.2 | 53.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.9 | 7.9 | 26.6 | 79.3 |
| 2003..... | 2.9 | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.7 | 7.8 | 24.3 | 76.5 |
| 2002..... | 2.8 | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.5 | 7.5 | 23.6 | 79.4 |
| 2001..... | 2.6 | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.2 | 7.1 | 22.3 | 75.7 |
| 2000..... | 2.5 | * | * | * | * | * | 0.6 | 1.1 | 2.7 | 6.7 | 22.9 | 72.0 |
| 1999..... | 2.4 | * | * | * | * | 0.1 | 0.5 | 1.2 | 2.3 | 7.1 | 20.6 | 63.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 7.7 | 22.2 | 67.3 |
| 2003..... | 3.8 | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.4 | 7.2 | 22.0 | 62.7 |
| 2002..... | 3.6 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 6.8 | 20.4 | 63.6 |
| 2001..... | 3.5 | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.3 | 6.6 | 21.1 | 59.4 |
| 2000..... | 3.4 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 7.0 | 20.0 | 57.2 |
| 1999..... | 3.0 | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 6.7 | 18.3 | 49.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 5.0 | 19.7 | 64.1 |
| 2003..... | 2.9 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 4.9 | 19.6 | 60.7 |
| 2002..... | 2.8 | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 4.9 | 17.8 | 61.4 |
| 2001..... | 2.7 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 4.6 | 17.9 | 58.0 |
| 2000..... | 2.5 | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 4.3 | 17.7 | 55.0 |
| 1999..... | 2.2 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 4.6 | 15.4 | 47.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 5.2 | 22.2 | 74.8 |
| 2003..... | 2.5 | * | * | * | * | * | 0.3 | 0.5 | 1.2 | 5.3 | 21.8 | 71.0 |
| 2002..... | 2.4 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 5.2 | 19.8 | 74.1 |
| 2001..... | 2.2 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 4.8 | 19.1 | 70.5 |
| 2000..... | 2.2 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 4.4 | 20.1 | 66.0 |
| 1999..... | 1.9 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 4.9 | 17.0 | 57.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | * | 0.4 | 1.0 | 4.9 | 18.1 | 59.3 |
| 2003..... | 3.3 | * | * | * | * | * | * | 0.4 | 1.2 | 4.6 | 18.2 | 56.1 |
| 2002..... | 3.1 | * | * | * | * | * | * | 0.4 | 0.9 | 4.6 | 16.5 | 56.0 |
| 2001..... | 3.1 | * | * | * | * | * | * | 0.4 | 1.1 | 4.5 | 17.1 | 52.8 |
| 2000..... | 2.9 | * | * | * | * | * | * | 0.3 | 1.0 | 4.3 | 16.2 | 50.6 |
| 1999..... | 2.5 | * | * | * | * | * | * | 0.3 | 0.8 | 4.3 | 14.3 | 43.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive renal disease (I12) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | * | * | 0.7 | 2.2 | 5.8 | 14.4 | 32.3 | 71.0 | 153.7 |
| 2003..... | 6.4 | * | * | * | * | 0.8 | 2.0 | 5.6 | 13.9 | 29.7 | 60.4 | 144.9 |
| 2002..... | 6.4 | * | * | * | * | 0.6 | 2.2 | 5.1 | 13.4 | 27.3 | 66.2 | 152.1 |
| 2001..... | 6.1 | * | * | * | * | 0.8 | 2.0 | 6.0 | 12.3 | 26.4 | 63.1 | 139.2 |
| 2000..... | 6.1 | * | * | * | * | 0.5 | 2.2 | 5.4 | 13.5 | 30.4 | 60.6 | 135.1 |
| 1999..... | 5.9 | * | * | * | * | 0.4 | 2.1 | 5.3 | 12.3 | 28.2 | 63.6 | 127.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | 0.8 | 2.7 | 7.8 | 16.9 | 34.9 | 81.0 | 141.2 |
| 2003..... | 5.9 | * | * | * | * | 1.0 | 2.4 | 7.8 | 16.4 | 33.1 | 56.8 | 157.7 |
| 2002..... | 5.7 | * | * | * | * | * | 2.8 | 6.4 | 14.4 | 29.6 | 73.3 | 157.5 |
| 2001..... | 5.5 | * | * | * | * | 0.9 | 2.6 | 7.8 | 13.3 | 30.0 | 63.3 | 144.4 |
| 2000..... | 5.3 | * | * | * | * | * | 2.8 | 6.2 | 15.1 | 30.4 | 60.8 | 144.4 |
| 1999..... | 5.4 | * | * | * | * | * | 2.5 | 7.7 | 13.5 | 29.6 | 68.5 | 142.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | * | * | 1.8 | 4.0 | 12.4 | 30.4 | 65.4 | 158.6 |
| 2003..... | 6.9 | * | * | * | * | * | 1.7 | 3.7 | 11.8 | 27.2 | 62.4 | 139.9 |
| 2002..... | 7.0 | * | * | * | * | * | 1.7 | 4.0 | 12.7 | 25.7 | 62.2 | 150.1 |
| 2001..... | 6.7 | * | * | * | * | 0.7 | 1.6 | 4.5 | 11.6 | 23.9 | 63.0 | 137.3 |
| 2000..... | 6.9 | * | * | * | * | * | 1.7 | 4.7 | 12.1 | 30.4 | 60.4 | 131.7 |
| 1999..... | 6.3 | * | * | * | * | * | 1.6 | 3.3 | 11.3 | 27.2 | 60.9 | 121.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive heart and renal disease (I13) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.1 | 6.6 | 24.8 |
| 2003..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.2 | 7.0 | 25.6 |
| 2002..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.2 | 6.8 | 23.5 |
| 2001..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.3 | 6.8 | 23.3 |
| 2000..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 7.0 | 24.3 |
| 1999..... | 1.2 | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.1 | 2.6 | 8.9 | 29.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.2 | 7.2 | 24.3 |
| 2003..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.3 | 7.4 | 26.7 |
| 2002..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.4 | 7.0 | 23.1 |
| 2001..... | 0.8 | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.5 | 6.8 | 24.2 |
| 2000..... | 0.8 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.5 | 7.2 | 27.8 |
| 1999..... | 1.0 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.7 | 9.6 | 31.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.9 | 6.2 | 25.0 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 6.8 | 25.2 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 6.7 | 23.7 |
| 2001..... | 1.1 | * | * | * | * | * | 0.2 | 0.2 | 0.9 | 2.1 | 6.8 | 22.9 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 6.9 | 22.9 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.4 | 8.5 | 28.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 1.3 | 1.8 | 8.9 | 20.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 2.7 | 5.9 | 25.4 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 1.3 | 2.9 | 8.5 | 22.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 1.1 | 3.0 | 7.4 | 26.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 2.5 | 6.3 | 25.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 3.1 | 8.1 | 23.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | 10.4 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 35.4 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 8.2 | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 7.9 | 18.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 3.1 | * | 20.3 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.7 | 21.6 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | 7.3 | 24.2 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 26.7 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.7 | 20.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive heart and renal disease (I13) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.1 | 6.5 | 24.9 |
| 2003..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.2 | 7.1 | 25.6 |
| 2002..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.1 | 6.7 | 23.5 |
| 2001..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.2 | 6.7 | 23.1 |
| 2000..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.2 | 7.0 | 24.2 |
| 1999..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.1 | 2.5 | 8.9 | 29.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.2 | 7.0 | 24.4 |
| 2003..... | 1.0 | * | * | * | * | * | 0.3 | 0.7 | 1.2 | 2.3 | 7.4 | 26.3 |
| 2002..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 6.9 | 22.9 |
| 2001..... | 0.9 | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.4 | 6.7 | 23.8 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.4 | 7.0 | 27.9 |
| 1999..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.7 | 9.6 | 31.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.9 | 6.1 | 25.2 |
| 2003..... | 1.3 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.1 | 6.8 | 25.3 |
| 2002..... | 1.2 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.0 | 6.5 | 23.7 |
| 2001..... | 1.2 | * | * | * | * | * | 0.2 | 0.2 | 0.9 | 2.0 | 6.7 | 22.8 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 6.9 | 22.8 |
| 1999..... | 1.5 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.4 | 8.4 | 28.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 5.3 | 23.0 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 5.9 | 23.2 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 5.6 | 20.8 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 5.6 | 20.7 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.5 | 5.6 | 21.8 |
| 1999..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 7.3 | 26.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 5.8 | 22.8 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 6.6 | 23.6 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 0.5 | 1.4 | 5.9 | 20.2 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.4 | 1.5 | 5.8 | 22.2 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.4 | 1.6 | 6.1 | 24.3 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 8.3 | 28.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | 0.3 | 1.1 | 4.9 | 23.1 |
| 2003..... | 1.1 | * | * | * | * | * | * | * | 0.3 | 1.1 | 5.5 | 23.0 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | 0.3 | 1.1 | 5.4 | 21.0 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | 0.5 | 1.3 | 5.4 | 20.1 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | 0.3 | 1.4 | 5.2 | 20.8 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 0.4 | 1.3 | 6.6 | 25.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hypertensive heart and renal disease (I13) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 1.2 | 2.9 | 5.0 | 8.7 | 20.0 | 49.2 |
| 2003..... | 2.5 | * | * | * | * | 0.4 | 1.3 | 3.1 | 5.5 | 9.3 | 19.9 | 53.8 |
| 2002..... | 2.5 | * | * | * | * | * | 1.0 | 2.8 | 6.2 | 10.0 | 19.8 | 57.2 |
| 2001..... | 2.3 | * | * | * | * | 0.4 | 1.2 | 2.1 | 5.1 | 10.1 | 20.2 | 50.6 |
| 2000..... | 2.3 | * | * | * | * | * | 0.8 | 2.2 | 4.7 | 10.0 | 22.6 | 50.9 |
| 1999..... | 2.8 | * | * | * | * | * | 1.5 | 2.3 | 6.2 | 12.7 | 27.3 | 66.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | 1.9 | 3.6 | 6.3 | 8.8 | 23.7 | 42.7 |
| 2003..... | 2.4 | * | * | * | * | * | 1.8 | 4.1 | 7.2 | 9.1 | 19.0 | 51.1 |
| 2002..... | 2.3 | * | * | * | * | * | 1.0 | 3.9 | 7.7 | 10.2 | 19.3 | 62.4 |
| 2001..... | 2.2 | * | * | * | * | * | 1.6 | 3.2 | 6.0 | 12.0 | 18.6 | 45.0 |
| 2000..... | 2.1 | * | * | * | * | * | 1.0 | 2.9 | 5.4 | 11.1 | 20.1 | 64.6 |
| 1999..... | 2.6 | * | * | * | * | * | 1.8 | 3.3 | 7.8 | 13.8 | 26.0 | 60.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 2.3 | 4.0 | 8.5 | 17.9 | 51.8 |
| 2003..... | 2.6 | * | * | * | * | * | 0.8 | 2.2 | 4.2 | 9.4 | 20.4 | 54.9 |
| 2002..... | 2.6 | * | * | * | * | * | 0.9 | 1.8 | 5.0 | 9.9 | 20.1 | 55.2 |
| 2001..... | 2.4 | * | * | * | * | * | 0.8 | 1.2 | 4.3 | 8.7 | 21.1 | 52.6 |
| 2000..... | 2.4 | * | * | * | * | * | * | 1.6 | 4.2 | 9.2 | 24.0 | 45.8 |
| 1999..... | 3.1 | * | * | * | * | * | 1.2 | 1.5 | 4.8 | 12.0 | 28.0 | 68.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 153.7 | * | * | * | 0.3 | 2.7 | 16.2 | 59.9 | 158.2 | 395.9 | 1,072.6 | 3,287.4 |
| 2003..... | 165.1 | 0.5 | * | 0.1 | 0.4 | 2.7 | 17.0 | 62.2 | 169.8 | 430.2 | 1,160.7 | 3,610.2 |
| 2002..... | 171.4 | 0.6 | * | 0.0 | 0.3 | 2.5 | 17.1 | 63.8 | 177.7 | 458.9 | 1,220.2 | 3,775.0 |
| 2001..... | 176.3 | 0.6 | * | 0.0 | 0.4 | 2.8 | 16.7 | 64.0 | 184.0 | 477.6 | 1,271.3 | 3,958.2 |
| 2000..... | 183.1 | 0.8 | * | * | 0.3 | 2.3 | 16.7 | 66.1 | 196.7 | 505.1 | 1,324.2 | 4,182.5 |
| 1999..... | 189.8 | 0.7 | * | 0.1 | 0.4 | 2.5 | 17.2 | 67.7 | 203.5 | 536.6 | 1,384.8 | 4,318.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 161.6 | * | * | * | 0.4 | 4.0 | 24.4 | 92.8 | 234.8 | 549.1 | 1,410.3 | 3,680.3 |
| 2003..... | 172.2 | * | * | * | 0.5 | 4.0 | 25.7 | 97.2 | 251.5 | 598.8 | 1,524.4 | 4,005.2 |
| 2002..... | 178.4 | * | * | * | 0.4 | 3.7 | 26.1 | 99.4 | 263.3 | 640.1 | 1,598.9 | 4,205.3 |
| 2001..... | 181.7 | * | * | * | 0.6 | 4.1 | 25.5 | 99.5 | 271.6 | 663.7 | 1,663.5 | 4,395.8 |
| 2000..... | 188.7 | * | * | * | 0.4 | 3.5 | 25.8 | 103.5 | 290.2 | 705.2 | 1,736.8 | 4,701.7 |
| 1999..... | 195.4 | * | * | * | 0.5 | 3.7 | 26.6 | 106.2 | 299.3 | 749.8 | 1,813.7 | 4,837.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 146.1 | * | * | * | 0.1 | 1.4 | 8.0 | 28.0 | 87.1 | 267.3 | 845.3 | 3,110.6 |
| 2003..... | 158.1 | * | * | * | 0.3 | 1.4 | 8.4 | 28.3 | 94.0 | 289.2 | 917.7 | 3,435.5 |
| 2002..... | 164.7 | * | * | * | 0.2 | 1.3 | 8.2 | 29.5 | 98.5 | 308.1 | 968.9 | 3,588.3 |
| 2001..... | 171.2 | * | * | * | 0.2 | 1.5 | 8.1 | 29.7 | 103.1 | 323.4 | 1,013.6 | 3,775.1 |
| 2000..... | 177.6 | * | * | * | 0.2 | 1.2 | 7.6 | 29.9 | 110.4 | 340.3 | 1,055.2 | 3,971.0 |
| 1999..... | 184.5 | * | * | * | 0.2 | 1.4 | 7.9 | 30.7 | 115.1 | 362.0 | 1,106.3 | 4,108.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 49.6 | * | * | * | * | 1.2 | 7.7 | 35.3 | 117.4 | 326.5 | 906.9 | 2,613.4 |
| 2003..... | 52.1 | * | * | * | 0.3 | 1.4 | 7.5 | 35.4 | 122.0 | 355.0 | 973.0 | 2,970.0 |
| 2002..... | 54.0 | * | * | * | * | 1.2 | 7.6 | 38.0 | 135.9 | 382.0 | 1,010.8 | 3,182.9 |
| 2001..... | 55.9 | * | * | * | * | 1.6 | 6.8 | 38.7 | 130.2 | 400.9 | 1,099.1 | 3,620.2 |
| 2000..... | 55.9 | * | * | * | * | 1.1 | 7.6 | 39.5 | 143.6 | 405.3 | 1,111.3 | 3,697.2 |
| 1999..... | 58.7 | * | * | * | * | 1.4 | 7.4 | 44.8 | 150.4 | 433.5 | 1,178.0 | 3,876.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 52.7 | * | * | * | * | 1.9 | 11.9 | 55.2 | 175.1 | 455.7 | 1,175.3 | 2,695.4 |
| 2003..... | 53.7 | * | * | * | 0.5 | 2.1 | 11.0 | 55.8 | 179.7 | 494.6 | 1,214.9 | 2,933.9 |
| 2002..... | 56.5 | * | * | * | * | 1.9 | 11.4 | 58.0 | 204.0 | 530.4 | 1,271.7 | 3,374.9 |
| 2001..... | 57.9 | * | * | * | * | 2.4 | 10.4 | 60.3 | 191.9 | 559.9 | 1,393.3 | 3,828.2 |
| 2000..... | 58.2 | * | * | * | * | 1.6 | 12.1 | 62.7 | 214.8 | 561.5 | 1,420.0 | 3,865.6 |
| 1999..... | 60.8 | * | * | * | * | 2.1 | 12.2 | 71.3 | 218.3 | 608.5 | 1,495.9 | 3,987.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 46.2 | * | * | * | * | * | 3.1 | 15.4 | 65.0 | 222.8 | 721.6 | 2,571.3 |
| 2003..... | 50.4 | * | * | * | * | 0.6 | 3.7 | 15.0 | 69.8 | 243.7 | 806.9 | 2,988.3 |
| 2002..... | 51.4 | * | * | * | * | * | 3.5 | 18.1 | 74.6 | 263.9 | 832.2 | 3,087.8 |
| 2001..... | 53.7 | * | * | * | * | * | 2.8 | 17.4 | 75.4 | 275.4 | 900.3 | 3,518.6 |
| 2000..... | 53.6 | * | * | * | * | * | 2.8 | 16.8 | 80.6 | 282.5 | 904.0 | 3,614.6 |
| 1999..... | 56.6 | * | * | * | * | * | 2.3 | 19.0 | 90.5 | 296.4 | 966.5 | 3,822.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 170.3 | * | * | * | 0.3 | 3.1 | 17.5 | 62.3 | 160.9 | 399.6 | 1,079.3 | 3,312.1 |
| 2003..... | 182.5 | 0.6 | * | 0.1 | 0.4 | 3.0 | 18.4 | 64.7 | 172.8 | 433.9 | 1,167.1 | 3,629.3 |
| 2002..... | 189.0 | * | * | * | 0.3 | 2.8 | 18.4 | 66.1 | 180.1 | 462.1 | 1,226.5 | 3,787.0 |
| 2001..... | 193.6 | 0.7 | * | * | 0.4 | 3.1 | 18.0 | 66.1 | 187.3 | 480.9 | 1,274.9 | 3,959.4 |
| 2000..... | 200.5 | 0.8 | * | * | 0.3 | 2.6 | 17.7 | 68.1 | 199.5 | 509.0 | 1,328.8 | 4,187.0 |
| 1999..... | 207.1 | 0.8 | * | 0.1 | 0.4 | 2.7 | 18.3 | 69.4 | 206.3 | 540.3 | 1,388.1 | 4,319.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 179.9 | * | * | * | 0.5 | 4.6 | 26.4 | 96.6 | 238.7 | 553.4 | 1,419.8 | 3,722.6 |
| 2003..... | 191.5 | * | * | * | 0.5 | 4.5 | 27.9 | 101.1 | 255.9 | 603.6 | 1,536.0 | 4,045.5 |
| 2002..... | 197.6 | * | * | * | 0.5 | 4.1 | 28.3 | 103.1 | 266.5 | 644.3 | 1,609.7 | 4,228.4 |
| 2001..... | 200.4 | * | * | * | 0.6 | 4.5 | 27.5 | 102.8 | 276.1 | 667.7 | 1,670.1 | 4,408.5 |
| 2000..... | 207.6 | * | * | * | 0.4 | 3.9 | 27.4 | 106.7 | 294.0 | 710.4 | 1,744.6 | 4,721.1 |
| 1999..... | 214.1 | * | * | * | 0.5 | 4.0 | 28.3 | 108.8 | 303.4 | 754.2 | 1,820.2 | 4,854.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 161.1 | * | * | * | 0.2 | 1.6 | 8.7 | 29.3 | 88.6 | 269.9 | 850.4 | 3,128.5 |
| 2003..... | 173.9 | * | * | * | 0.3 | 1.5 | 9.0 | 29.6 | 95.6 | 291.5 | 920.9 | 3,446.4 |
| 2002..... | 180.7 | * | * | * | 0.2 | 1.5 | 8.8 | 30.5 | 100.0 | 309.9 | 972.5 | 3,596.5 |
| 2001..... | 187.2 | * | * | * | 0.2 | 1.6 | 8.7 | 30.8 | 105.0 | 325.6 | 1,015.5 | 3,772.7 |
| 2000..... | 193.8 | * | * | * | 0.2 | 1.3 | 8.2 | 31.0 | 112.1 | 342.8 | 1,057.9 | 3,971.0 |
| 1999..... | 200.6 | * | * | * | 0.3 | 1.5 | 8.5 | 31.5 | 116.6 | 364.6 | 1,107.8 | 4,105.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 187.7 | * | * | * | 0.2 | 2.8 | 17.2 | 59.1 | 153.5 | 388.1 | 1,079.6 | 3,369.7 |
| 2003..... | 200.8 | * | * | * | 0.4 | 2.8 | 17.9 | 61.1 | 164.5 | 421.6 | 1,164.6 | 3,691.6 |
| 2002..... | 208.0 | * | * | * | 0.3 | 2.6 | 17.9 | 62.8 | 171.2 | 450.8 | 1,223.1 | 3,848.2 |
| 2001..... | 212.6 | * | * | * | 0.3 | 2.9 | 17.3 | 62.6 | 177.8 | 467.8 | 1,271.3 | 4,022.8 |
| 2000..... | 220.2 | * | * | * | 0.3 | 2.4 | 16.9 | 64.6 | 189.5 | 496.4 | 1,324.5 | 4,261.3 |
| 1999..... | 227.2 | * | * | * | 0.3 | 2.5 | 17.3 | 65.8 | 196.4 | 527.7 | 1,386.5 | 4,390.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 198.0 | * | * | * | 0.3 | 4.3 | 26.4 | 93.4 | 230.4 | 543.1 | 1,431.6 | 3,814.0 |
| 2003..... | 210.2 | * | * | * | 0.4 | 4.3 | 27.5 | 97.2 | 247.3 | 591.8 | 1,541.1 | 4,149.5 |
| 2002..... | 217.2 | * | * | * | 0.4 | 4.0 | 28.0 | 100.0 | 256.8 | 636.0 | 1,618.3 | 4,324.5 |
| 2001..... | 219.6 | * | * | * | 0.5 | 4.3 | 26.9 | 99.5 | 266.3 | 656.5 | 1,678.7 | 4,506.9 |
| 2000..... | 227.6 | * | * | * | 0.4 | 3.7 | 26.8 | 103.5 | 283.5 | 701.7 | 1,752.3 | 4,833.4 |
| 1999..... | 234.2 | * | * | * | 0.5 | 3.6 | 27.2 | 105.1 | 292.3 | 745.2 | 1,834.9 | 4,954.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 177.7 | * | * | * | * | 1.3 | 7.9 | 25.4 | 80.4 | 254.7 | 839.7 | 3,170.6 |
| 2003..... | 191.8 | * | * | * | 0.3 | 1.4 | 8.2 | 25.5 | 85.9 | 275.9 | 910.4 | 3,490.0 |
| 2002..... | 199.1 | * | * | * | 0.2 | 1.3 | 7.8 | 26.1 | 90.2 | 293.0 | 958.3 | 3,642.5 |
| 2001..... | 205.8 | * | * | * | 0.2 | 1.4 | 7.6 | 26.4 | 94.2 | 307.7 | 1,001.1 | 3,821.4 |
| 2000..... | 213.1 | * | * | * | 0.2 | 1.0 | 6.9 | 26.4 | 100.9 | 323.6 | 1,043.3 | 4,030.7 |
| 1999..... | 220.5 | * | * | * | * | 1.4 | 7.4 | 27.2 | 106.1 | 345.6 | 1,092.9 | 4,165.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 124.8 | * | * | * | 0.7 | 4.9 | 24.0 | 97.9 | 259.1 | 576.3 | 1,255.5 | 3,038.5 |
| 2003..... | 133.2 | * | * | * | 0.7 | 4.4 | 25.5 | 101.8 | 277.3 | 622.3 | 1,375.4 | 3,328.7 |
| 2002..... | 137.0 | * | * | * | 0.5 | 4.1 | 25.8 | 103.6 | 292.4 | 643.9 | 1,433.9 | 3,481.1 |
| 2001..... | 141.2 | * | * | * | 0.9 | 4.8 | 26.4 | 104.6 | 302.1 | 681.4 | 1,487.0 | 3,628.7 |
| 2000..... | 144.4 | * | * | * | 0.4 | 4.4 | 26.1 | 107.7 | 318.9 | 710.1 | 1,534.5 | 3,701.7 |
| 1999..... | 147.9 | * | * | * | 0.8 | 4.1 | 28.2 | 110.8 | 324.7 | 741.6 | 1,559.0 | 3,817.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 129.2 | * | * | * | 1.2 | 6.7 | 32.9 | 141.2 | 368.2 | 771.0 | 1,551.8 | 3,133.7 |
| 2003..... | 137.6 | * | * | * | 0.9 | 6.4 | 36.2 | 148.3 | 387.0 | 836.1 | 1,758.6 | 3,309.2 |
| 2002..... | 139.5 | * | * | * | * | 5.3 | 36.2 | 146.9 | 413.2 | 850.8 | 1,762.3 | 3,572.8 |
| 2001..... | 143.2 | * | * | * | 1.2 | 6.8 | 37.1 | 149.0 | 418.0 | 903.1 | 1,834.9 | 3,791.8 |
| 2000..... | 145.7 | * | * | * | * | 6.1 | 36.3 | 152.2 | 444.2 | 916.5 | 1,887.6 | 3,911.2 |
| 1999..... | 150.0 | * | * | * | 0.9 | 6.0 | 40.3 | 160.0 | 458.4 | 967.5 | 1,873.3 | 4,082.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 120.8 | * | * | * | * | 3.2 | 16.0 | 60.7 | 171.5 | 437.6 | 1,088.3 | 3,000.9 |
| 2003..... | 129.1 | * | * | * | * | 2.6 | 16.0 | 62.0 | 189.2 | 470.4 | 1,159.1 | 3,336.2 |
| 2002..... | 134.8 | * | * | * | * | 2.9 | 16.7 | 66.5 | 195.4 | 497.1 | 1,248.1 | 3,446.0 |
| 2001..... | 139.4 | * | * | * | * | 2.9 | 16.9 | 66.5 | 208.9 | 525.0 | 1,291.6 | 3,568.3 |
| 2000..... | 143.2 | * | * | * | * | 2.9 | 17.1 | 69.4 | 218.1 | 565.0 | 1,335.7 | 3,623.9 |
| 1999..... | 146.0 | * | * | * | 0.7 | 2.4 | 17.4 | 68.7 | 217.5 | 582.3 | 1,383.0 | 3,719.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute myocardial infarction (I21-I22) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 53.4 | * | * | * | 0.1 | 1.1 | 6.4 | 24.2 | 64.5 | 153.4 | 370.3 | 992.9 |
| 2003..... | 58.7 | * | * | * | 0.2 | 1.0 | 7.0 | 25.5 | 70.9 | 168.6 | 412.7 | 1,118.8 |
| 2002..... | 62.3 | * | * | * | 0.2 | 1.0 | 7.0 | 26.5 | 74.9 | 186.1 | 444.4 | 1,191.3 |
| 2001..... | 64.9 | * | * | * | 0.2 | 1.2 | 7.1 | 27.6 | 78.4 | 193.2 | 471.2 | 1,265.4 |
| 2000..... | 68.5 | 0.5 | * | * | 0.1 | 0.9 | 6.9 | 28.9 | 86.6 | 209.1 | 503.3 | 1,338.5 |
| 1999..... | 71.5 | * | * | * | 0.2 | 1.1 | 7.2 | 29.6 | 90.0 | 225.0 | 529.3 | 1,379.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 57.4 | * | * | * | 0.2 | 1.6 | 9.6 | 37.3 | 95.3 | 209.5 | 477.0 | 1,120.5 |
| 2003..... | 62.6 | * | * | * | 0.2 | 1.5 | 10.4 | 39.8 | 104.1 | 231.0 | 530.6 | 1,263.5 |
| 2002..... | 66.2 | * | * | * | 0.2 | 1.4 | 10.5 | 40.6 | 109.2 | 256.2 | 569.2 | 1,366.4 |
| 2001..... | 68.6 | * | * | * | 0.2 | 1.7 | 10.6 | 42.4 | 114.0 | 264.7 | 604.1 | 1,461.3 |
| 2000..... | 72.7 | * | * | * | 0.1 | 1.3 | 10.6 | 44.7 | 125.7 | 286.9 | 648.3 | 1,568.3 |
| 1999..... | 76.1 | * | * | * | 0.2 | 1.5 | 11.0 | 46.3 | 131.8 | 310.0 | 684.4 | 1,622.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 49.6 | * | * | * | * | 0.6 | 3.3 | 11.6 | 35.9 | 106.3 | 298.5 | 935.5 |
| 2003..... | 54.8 | * | * | * | 0.1 | 0.6 | 3.6 | 11.7 | 40.0 | 116.5 | 333.9 | 1,054.8 |
| 2002..... | 58.4 | * | * | * | 0.1 | 0.5 | 3.6 | 12.9 | 43.2 | 127.7 | 361.5 | 1,115.3 |
| 2001..... | 61.3 | * | * | * | * | 0.7 | 3.5 | 13.3 | 45.4 | 134.0 | 383.9 | 1,183.5 |
| 2000..... | 64.6 | * | * | * | 0.1 | 0.5 | 3.3 | 13.6 | 50.5 | 145.1 | 408.7 | 1,244.8 |
| 1999..... | 67.0 | * | * | * | 0.1 | 0.6 | 3.4 | 13.5 | 51.5 | 155.4 | 428.6 | 1,281.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 17.3 | * | * | * | * | 0.5 | 2.9 | 12.9 | 44.2 | 120.4 | 315.7 | 816.8 |
| 2003..... | 18.5 | * | * | * | * | 0.5 | 3.0 | 13.3 | 44.5 | 137.7 | 341.6 | 952.0 |
| 2002..... | 19.7 | * | * | * | * | 0.5 | 3.0 | 14.2 | 55.1 | 146.2 | 364.3 | 1,048.2 |
| 2001..... | 20.0 | * | * | * | * | 0.7 | 2.5 | 14.8 | 49.5 | 149.8 | 394.9 | 1,185.5 |
| 2000..... | 20.1 | * | * | * | * | 0.5 | 3.0 | 15.5 | 55.0 | 154.3 | 404.7 | 1,177.1 |
| 1999..... | 21.1 | * | * | * | * | 0.5 | 2.9 | 17.8 | 56.1 | 164.0 | 433.7 | 1,227.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 18.4 | * | * | * | * | 0.6 | 4.3 | 20.1 | 65.4 | 164.1 | 403.4 | 840.6 |
| 2003..... | 18.9 | * | * | * | * | 0.7 | 4.1 | 20.6 | 64.1 | 187.5 | 409.2 | 963.6 |
| 2002..... | 20.6 | * | * | * | * | 0.7 | 4.2 | 21.0 | 80.8 | 199.3 | 460.0 | 1,107.7 |
| 2001..... | 20.9 | * | * | * | * | 1.0 | 3.9 | 22.7 | 69.4 | 206.9 | 488.7 | 1,331.3 |
| 2000..... | 21.1 | * | * | * | * | 0.7 | 4.7 | 23.4 | 80.0 | 207.6 | 514.8 | 1,273.8 |
| 1999..... | 22.0 | * | * | * | * | 0.8 | 4.6 | 26.7 | 79.0 | 225.8 | 558.9 | 1,276.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 16.0 | * | * | * | * | * | 1.3 | 5.6 | 24.9 | 85.4 | 255.1 | 804.5 |
| 2003..... | 18.0 | * | * | * | * | * | 1.7 | 6.0 | 26.8 | 98.1 | 295.2 | 946.1 |
| 2002..... | 18.7 | * | * | * | * | * | 1.8 | 7.4 | 32.1 | 103.9 | 298.8 | 1,018.8 |
| 2001..... | 19.1 | * | * | * | * | * | 1.1 | 7.0 | 31.8 | 104.7 | 331.5 | 1,114.2 |
| 2000..... | 19.1 | * | * | * | * | * | 1.2 | 7.8 | 32.9 | 112.3 | 330.8 | 1,129.7 |
| 1999..... | 20.1 | * | * | * | * | * | 1.1 | 9.1 | 35.9 | 115.5 | 350.4 | 1,204.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute myocardial infarction (I21-I22) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 59.2 | * | * | * | 0.1 | 1.3 | 7.0 | 25.4 | 66.0 | 155.4 | 372.7 | 999.2 |
| 2003..... | 64.9 | * | * | * | 0.2 | 1.2 | 7.6 | 26.7 | 72.8 | 170.4 | 415.5 | 1,123.8 |
| 2002..... | 68.7 | * | * | * | 0.2 | 1.1 | 7.6 | 27.7 | 76.3 | 188.2 | 447.3 | 1,194.0 |
| 2001..... | 71.4 | * | * | * | 0.2 | 1.3 | 7.6 | 28.8 | 80.3 | 195.6 | 473.6 | 1,265.3 |
| 2000..... | 75.3 | * | * | * | 0.1 | 1.0 | 7.4 | 30.0 | 88.6 | 211.9 | 506.3 | 1,340.8 |
| 1999..... | 78.2 | * | * | * | 0.2 | 1.2 | 7.7 | 30.5 | 92.2 | 227.9 | 531.9 | 1,381.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 64.0 | * | * | * | 0.2 | 1.8 | 10.5 | 39.1 | 97.5 | 212.1 | 480.2 | 1,132.5 |
| 2003..... | 69.7 | * | * | * | 0.2 | 1.7 | 11.4 | 41.8 | 106.9 | 233.4 | 535.7 | 1,274.5 |
| 2002..... | 73.5 | * | * | * | 0.2 | 1.6 | 11.5 | 42.6 | 111.3 | 259.2 | 573.4 | 1,374.2 |
| 2001..... | 75.9 | * | * | * | 0.2 | 1.8 | 11.6 | 44.2 | 117.0 | 267.7 | 607.9 | 1,463.9 |
| 2000..... | 80.3 | * | * | * | 0.2 | 1.5 | 11.4 | 46.5 | 128.7 | 290.9 | 652.6 | 1,576.2 |
| 1999..... | 83.8 | * | * | * | 0.2 | 1.7 | 11.8 | 47.8 | 135.1 | 313.8 | 687.7 | 1,631.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 54.7 | * | * | * | * | 0.7 | 3.6 | 12.2 | 36.7 | 107.7 | 300.4 | 939.6 |
| 2003..... | 60.3 | * | * | * | 0.2 | 0.6 | 3.8 | 12.3 | 41.0 | 117.5 | 335.3 | 1,057.5 |
| 2002..... | 64.1 | * | * | * | * | 0.6 | 3.8 | 13.3 | 44.0 | 128.9 | 363.8 | 1,116.3 |
| 2001..... | 67.1 | * | * | * | * | 0.8 | 3.8 | 13.9 | 46.3 | 135.7 | 385.4 | 1,182.7 |
| 2000..... | 70.6 | * | * | * | 0.1 | 0.6 | 3.5 | 14.1 | 51.6 | 146.8 | 411.0 | 1,245.6 |
| 1999..... | 72.9 | * | * | * | * | 0.7 | 3.7 | 13.9 | 52.5 | 157.3 | 430.7 | 1,280.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 65.3 | * | * | * | 0.1 | 1.2 | 7.0 | 24.8 | 64.9 | 152.9 | 374.2 | 1,010.1 |
| 2003..... | 71.5 | * | * | * | 0.2 | 1.1 | 7.5 | 25.9 | 71.4 | 168.3 | 415.1 | 1,137.4 |
| 2002..... | 75.5 | * | * | * | 0.1 | 1.1 | 7.5 | 27.0 | 74.7 | 186.1 | 445.9 | 1,202.8 |
| 2001..... | 78.4 | * | * | * | 0.1 | 1.2 | 7.5 | 28.1 | 78.7 | 193.0 | 473.0 | 1,276.8 |
| 2000..... | 82.7 | * | * | * | 0.1 | 0.9 | 7.3 | 29.3 | 86.9 | 209.3 | 505.3 | 1,355.3 |
| 1999..... | 86.0 | * | * | * | 0.1 | 1.1 | 7.5 | 30.0 | 90.7 | 226.1 | 532.4 | 1,394.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 70.9 | * | * | * | * | 1.9 | 10.7 | 38.8 | 97.3 | 210.8 | 487.4 | 1,156.5 |
| 2003..... | 77.0 | * | * | * | 0.2 | 1.6 | 11.5 | 41.2 | 106.5 | 232.9 | 537.8 | 1,300.9 |
| 2002..... | 81.2 | * | * | * | 0.2 | 1.6 | 11.5 | 42.5 | 110.8 | 260.2 | 575.8 | 1,393.0 |
| 2001..... | 83.7 | * | * | * | 0.2 | 1.7 | 11.6 | 44.3 | 116.8 | 267.7 | 612.5 | 1,483.2 |
| 2000..... | 88.5 | * | * | * | 0.2 | 1.4 | 11.4 | 46.7 | 128.0 | 291.0 | 656.7 | 1,600.7 |
| 1999..... | 92.4 | * | * | * | 0.2 | 1.5 | 11.8 | 48.1 | 134.6 | 315.0 | 695.7 | 1,653.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 59.9 | * | * | * | * | 0.6 | 3.3 | 10.9 | 34.2 | 103.1 | 297.1 | 944.5 |
| 2003..... | 66.2 | * | * | * | * | 0.6 | 3.5 | 10.8 | 38.0 | 113.0 | 332.2 | 1,065.3 |
| 2002..... | 70.1 | * | * | * | * | 0.5 | 3.6 | 11.8 | 40.6 | 122.9 | 358.8 | 1,120.7 |
| 2001..... | 73.4 | * | * | * | * | 0.8 | 3.4 | 12.2 | 42.7 | 129.6 | 380.6 | 1,190.9 |
| 2000..... | 77.2 | * | * | * | * | 0.5 | 3.1 | 12.3 | 48.2 | 140.5 | 405.9 | 1,256.3 |
| 1999..... | 79.8 | * | * | * | * | 0.7 | 3.3 | 12.3 | 49.3 | 151.8 | 425.5 | 1,290.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute myocardial infarction (I21-I22) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 43.4 | * | * | * | 0.3 | 1.8 | 9.1 | 36.2 | 93.0 | 208.2 | 424.3 | 991.3 |
| 2003..... | 47.3 | * | * | * | * | 1.7 | 10.4 | 38.6 | 100.6 | 223.1 | 486.5 | 1,103.6 |
| 2002..... | 50.2 | * | * | * | * | 1.7 | 10.0 | 39.0 | 109.1 | 242.6 | 523.0 | 1,212.9 |
| 2001..... | 51.6 | * | * | * | 0.4 | 1.9 | 10.2 | 40.2 | 111.6 | 256.0 | 543.7 | 1,247.0 |
| 2000..... | 54.0 | * | * | * | * | 1.8 | 10.1 | 41.9 | 122.3 | 273.0 | 576.6 | 1,281.8 |
| 1999..... | 55.1 | * | * | * | * | 1.7 | 10.5 | 42.0 | 123.2 | 282.4 | 589.9 | 1,328.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 43.9 | * | * | * | * | 2.1 | 12.0 | 51.6 | 126.2 | 269.9 | 491.7 | 1,020.2 |
| 2003..... | 48.2 | * | * | * | * | 2.4 | 14.3 | 54.7 | 135.9 | 287.1 | 606.9 | 1,121.9 |
| 2002..... | 50.1 | * | * | * | * | 2.2 | 13.7 | 53.4 | 144.5 | 305.4 | 631.3 | 1,296.3 |
| 2001..... | 50.9 | * | * | * | * | 2.5 | 13.5 | 53.4 | 146.1 | 320.8 | 646.6 | 1,389.4 |
| 2000..... | 53.7 | * | * | * | * | 2.2 | 13.6 | 56.3 | 163.1 | 341.2 | 693.3 | 1,426.6 |
| 1999..... | 55.2 | * | * | * | * | 2.5 | 14.5 | 58.1 | 168.7 | 357.0 | 691.4 | 1,509.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 43.0 | * | * | * | * | 1.5 | 6.5 | 23.0 | 66.2 | 164.3 | 386.3 | 979.8 |
| 2003..... | 46.5 | * | * | * | * | 1.0 | 6.9 | 24.7 | 72.2 | 177.6 | 418.5 | 1,096.5 |
| 2002..... | 50.4 | * | * | * | * | 1.3 | 6.7 | 26.7 | 80.7 | 198.0 | 461.8 | 1,181.0 |
| 2001..... | 52.3 | * | * | * | * | 1.3 | 7.4 | 28.8 | 83.8 | 210.3 | 485.9 | 1,194.3 |
| 2000..... | 54.2 | * | * | * | * | 1.4 | 7.0 | 29.6 | 89.5 | 225.0 | 510.9 | 1,228.0 |
| 1999..... | 54.9 | * | * | * | * | 1.0 | 7.0 | 28.2 | 86.8 | 229.7 | 533.1 | 1,261.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other acute ischemic heart diseases (I24) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.3 | 5.4 | 15.3 |
| 2003..... | 1.2 | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 3.3 | 6.7 | 20.6 |
| 2002..... | 1.3 | * | * | * | * | * | 0.2 | 0.9 | 1.8 | 3.6 | 7.3 | 21.8 |
| 2001..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 1.9 | 3.6 | 7.1 | 22.0 |
| 2000..... | 1.3 | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 3.6 | 7.5 | 22.2 |
| 1999..... | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 3.8 | 7.2 | 22.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.3 | 0.8 | 1.9 | 3.2 | 6.5 | 17.0 |
| 2003..... | 1.3 | * | * | * | * | * | 0.4 | 1.1 | 2.2 | 4.5 | 8.4 | 20.2 |
| 2002..... | 1.4 | * | * | * | * | * | 0.3 | 1.4 | 2.6 | 4.9 | 9.0 | 22.1 |
| 2001..... | 1.4 | * | * | * | * | * | 0.3 | 1.3 | 2.7 | 4.8 | 8.7 | 20.9 |
| 2000..... | 1.4 | * | * | * | * | * | 0.3 | 1.3 | 3.1 | 4.9 | 9.3 | 21.0 |
| 1999..... | 1.4 | * | * | * | * | * | 0.3 | 1.1 | 2.9 | 5.2 | 9.0 | 22.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 4.6 | 14.6 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.3 | 5.7 | 20.7 |
| 2002..... | 1.3 | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 2.5 | 6.1 | 21.6 |
| 2001..... | 1.3 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.6 | 6.0 | 22.4 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 2.5 | 6.3 | 22.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.4 | 1.1 | 2.7 | 6.0 | 21.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.1 | 5.1 | 15.6 |
| 2003..... | 1.3 | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 3.1 | 6.6 | 20.9 |
| 2002..... | 1.4 | * | * | * | * | * | 0.2 | 0.8 | 1.7 | 3.5 | 7.2 | 22.0 |
| 2001..... | 1.4 | * | * | * | * | * | 0.2 | 0.8 | 1.7 | 3.4 | 7.0 | 22.0 |
| 2000..... | 1.4 | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 3.4 | 7.3 | 22.5 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 3.6 | 7.0 | 22.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 2.8 | 6.3 | 16.9 |
| 2003..... | 1.3 | * | * | * | * | * | 0.4 | 1.0 | 2.0 | 4.4 | 7.9 | 20.4 |
| 2002..... | 1.5 | * | * | * | * | * | 0.3 | 1.2 | 2.5 | 4.6 | 9.0 | 22.7 |
| 2001..... | 1.4 | * | * | * | * | * | 0.3 | 1.2 | 2.5 | 4.5 | 8.7 | 20.9 |
| 2000..... | 1.5 | * | * | * | * | * | 0.3 | 1.2 | 2.8 | 4.6 | 9.0 | 21.3 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.9 | 2.6 | 4.9 | 8.7 | 23.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 4.3 | 15.0 |
| 2003..... | 1.3 | * | * | * | * | * | * | 0.4 | 0.8 | 2.1 | 5.7 | 21.1 |
| 2002..... | 1.4 | * | * | * | * | * | * | 0.5 | 1.0 | 2.5 | 6.0 | 21.8 |
| 2001..... | 1.4 | * | * | * | * | * | * | 0.3 | 1.0 | 2.5 | 5.9 | 22.4 |
| 2000..... | 1.3 | * | * | * | * | * | * | 0.3 | 1.0 | 2.4 | 6.1 | 23.0 |
| 1999..... | 1.3 | * | * | * | * | * | * | 0.3 | 1.1 | 2.6 | 5.8 | 22.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other acute ischemic heart diseases (I24) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 1.4 | 3.0 | 4.2 | 9.1 | 15.5 |
| 2003..... | 1.2 | * | * | * | * | * | 0.5 | 1.3 | 3.0 | 5.5 | 10.1 | 20.2 |
| 2002..... | 1.2 | * | * | * | * | * | * | 1.7 | 3.0 | 6.1 | 9.4 | 21.5 |
| 2001..... | 1.3 | * | * | * | * | * | 0.5 | 1.6 | 4.0 | 5.8 | 8.5 | 24.3 |
| 2000..... | 1.3 | * | * | * | * | * | 0.4 | 1.5 | 4.3 | 5.6 | 10.8 | 21.9 |
| 1999..... | 1.2 | * | * | * | * | * | 0.5 | 1.9 | 3.6 | 5.8 | 10.9 | 17.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.9 | 4.7 | 7.1 | 12.0 | 23.4 |
| 2003..... | 1.5 | * | * | * | * | * | * | 2.0 | 4.9 | 7.7 | 16.4 | 24.5 |
| 2002..... | 1.4 | * | * | * | * | * | * | 2.8 | 4.4 | 9.4 | 11.7 | 21.9 |
| 2001..... | 1.5 | * | * | * | * | * | * | 2.5 | 5.6 | 8.8 | 10.5 | 24.3 |
| 2000..... | 1.5 | * | * | * | * | * | * | 2.3 | 6.3 | 8.7 | 14.2 | * |
| 1999..... | 1.5 | * | * | * | * | * | 0.8 | 2.7 | 6.1 | 8.4 | 12.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.9 | 1.7 | 2.0 | 7.4 | 12.4 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 1.5 | 3.9 | 6.5 | 18.6 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.8 | 2.0 | 3.7 | 8.1 | 21.3 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | 2.6 | 3.8 | 7.4 | 24.4 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | 2.7 | 3.5 | 8.9 | 22.2 |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.1 | 1.6 | 3.9 | 9.8 | 16.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other forms of chronic ischemic heart disease (I20,I25) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 99.5 | * | * | * | 0.2 | 1.6 | 9.6 | 35.1 | 92.5 | 240.3 | 697.1 | 2,279.7 |
| 2003..... | 105.3 | * | * | * | 0.2 | 1.7 | 9.8 | 35.9 | 97.5 | 258.4 | 741.5 | 2,471.4 |
| 2002..... | 108.0 | * | * | * | 0.2 | 1.4 | 9.9 | 36.5 | 101.0 | 269.4 | 768.8 | 2,562.5 |
| 2001..... | 110.3 | * | * | * | 0.2 | 1.6 | 9.5 | 35.6 | 103.8 | 280.9 | 793.1 | 2,671.3 |
| 2000..... | 113.3 | * | * | * | 0.2 | 1.4 | 9.5 | 36.4 | 108.1 | 292.4 | 813.6 | 2,822.1 |
| 1999..... | 117.2 | * | * | * | 0.2 | 1.4 | 9.8 | 37.5 | 111.5 | 308.0 | 848.5 | 2,917.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 103.3 | * | * | * | 0.2 | 2.4 | 14.5 | 54.8 | 137.7 | 336.5 | 926.9 | 2,543.6 |
| 2003..... | 108.5 | * | * | * | 0.3 | 2.5 | 14.9 | 56.4 | 145.4 | 363.5 | 985.8 | 2,722.2 |
| 2002..... | 110.9 | * | * | * | 0.2 | 2.2 | 15.4 | 57.5 | 151.5 | 379.3 | 1,021.0 | 2,817.4 |
| 2001..... | 111.9 | * | * | * | 0.3 | 2.4 | 14.6 | 55.9 | 154.9 | 394.3 | 1,050.9 | 2,913.8 |
| 2000..... | 114.8 | * | * | * | 0.2 | 2.1 | 14.8 | 57.7 | 161.6 | 413.5 | 1,079.4 | 3,112.7 |
| 1999..... | 118.0 | * | * | * | 0.3 | 2.1 | 15.3 | 59.0 | 164.7 | 434.9 | 1,120.5 | 3,193.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 95.7 | * | * | * | * | 0.7 | 4.6 | 16.2 | 50.5 | 159.6 | 542.4 | 2,161.0 |
| 2003..... | 102.3 | * | * | * | 0.1 | 0.8 | 4.7 | 16.2 | 53.1 | 170.5 | 578.3 | 2,360.5 |
| 2002..... | 105.2 | * | * | * | * | 0.7 | 4.4 | 16.2 | 54.3 | 178.0 | 601.4 | 2,451.9 |
| 2001..... | 108.8 | * | * | * | 0.1 | 0.7 | 4.4 | 16.0 | 56.7 | 186.9 | 623.7 | 2,569.8 |
| 2000..... | 111.9 | * | * | * | * | 0.6 | 4.3 | 16.0 | 58.8 | 192.8 | 640.3 | 2,703.8 |
| 1999..... | 116.4 | * | * | * | 0.1 | 0.7 | 4.3 | 16.7 | 62.6 | 204.0 | 671.8 | 2,805.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 32.2 | * | * | * | * | 0.8 | 4.8 | 22.4 | 72.9 | 205.5 | 589.6 | 1,795.7 |
| 2003..... | 33.5 | * | * | * | * | 0.9 | 4.6 | 22.0 | 77.2 | 216.4 | 629.5 | 2,012.8 |
| 2002..... | 34.2 | * | * | * | * | 0.7 | 4.5 | 23.6 | 80.1 | 234.6 | 644.0 | 2,129.7 |
| 2001..... | 35.7 | * | * | * | * | 0.9 | 4.2 | 23.8 | 80.0 | 249.4 | 701.7 | 2,426.6 |
| 2000..... | 35.6 | * | * | * | * | 0.6 | 4.6 | 23.9 | 88.0 | 249.7 | 703.2 | 2,508.8 |
| 1999..... | 37.5 | * | * | * | * | 0.9 | 4.5 | 26.8 | 93.5 | 268.8 | 741.8 | 2,640.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 34.2 | * | * | * | * | 1.3 | 7.5 | 35.0 | 109.3 | 291.0 | 769.7 | 1,853.5 |
| 2003..... | 34.6 | * | * | * | * | 1.4 | 6.9 | 35.1 | 115.3 | 306.1 | 804.2 | 1,965.7 |
| 2002..... | 35.7 | * | * | * | * | 1.2 | 7.2 | 36.6 | 122.3 | 329.1 | 808.8 | 2,262.2 |
| 2001..... | 36.8 | * | * | * | * | 1.4 | 6.5 | 37.5 | 121.4 | 350.2 | 901.5 | 2,479.8 |
| 2000..... | 36.9 | * | * | * | * | 0.9 | 7.4 | 39.1 | 134.2 | 351.6 | 900.8 | 2,577.7 |
| 1999..... | 38.7 | * | * | * | * | 1.3 | 7.6 | 44.3 | 138.5 | 381.8 | 932.8 | 2,706.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 30.1 | * | * | * | * | * | 1.8 | 9.8 | 39.9 | 136.9 | 465.3 | 1,766.0 |
| 2003..... | 32.3 | * | * | * | * | * | 2.0 | 8.9 | 42.8 | 144.9 | 509.6 | 2,036.7 |
| 2002..... | 32.6 | * | * | * | * | * | 1.6 | 10.5 | 42.2 | 159.4 | 531.2 | 2,064.1 |
| 2001..... | 34.5 | * | * | * | * | * | 1.7 | 10.3 | 43.3 | 169.9 | 566.7 | 2,400.7 |
| 2000..... | 34.2 | * | * | * | * | * | 1.7 | 8.9 | 47.2 | 169.6 | 570.5 | 2,475.0 |
| 1999..... | 36.3 | * | * | * | * | * | 1.3 | 9.7 | 53.8 | 180.2 | 614.8 | 2,608.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other forms of chronic ischemic heart disease (I20,I25) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 110.1 | * | * | * | 0.2 | 1.8 | 10.3 | 36.3 | 93.6 | 241.8 | 701.2 | 2,297.5 |
| 2003..... | 116.3 | * | * | * | 0.2 | 1.8 | 10.6 | 37.2 | 98.5 | 260.2 | 744.8 | 2,484.9 |
| 2002..... | 118.9 | * | * | * | 0.2 | 1.6 | 10.6 | 37.5 | 102.0 | 270.3 | 771.9 | 2,571.2 |
| 2001..... | 120.9 | * | * | * | 0.2 | 1.7 | 10.2 | 36.5 | 105.1 | 281.6 | 794.2 | 2,672.2 |
| 2000..... | 123.9 | * | * | * | 0.2 | 1.5 | 10.1 | 37.3 | 108.8 | 293.5 | 815.0 | 2,824.0 |
| 1999..... | 127.6 | * | * | * | 0.2 | 1.5 | 10.4 | 38.1 | 112.1 | 308.6 | 849.1 | 2,916.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 114.9 | * | * | * | 0.2 | 2.7 | 15.6 | 56.7 | 139.3 | 338.1 | 933.0 | 2,573.1 |
| 2003..... | 120.4 | * | * | * | 0.3 | 2.8 | 16.2 | 58.3 | 146.7 | 365.6 | 991.9 | 2,750.7 |
| 2002..... | 122.7 | * | * | * | 0.3 | 2.4 | 16.6 | 59.2 | 152.6 | 380.2 | 1,027.3 | 2,832.2 |
| 2001..... | 123.1 | * | * | * | 0.3 | 2.6 | 15.7 | 57.3 | 156.4 | 395.1 | 1,053.5 | 2,923.7 |
| 2000..... | 126.0 | * | * | * | 0.2 | 2.3 | 15.7 | 58.9 | 162.3 | 414.7 | 1,082.7 | 3,123.9 |
| 1999..... | 128.9 | * | * | * | 0.3 | 2.3 | 16.2 | 59.9 | 165.4 | 435.2 | 1,123.5 | 3,200.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 105.6 | * | * | * | 0.8 | 5.0 | 16.7 | 51.1 | 160.7 | 545.5 | 2,174.3 | |
| 2003..... | 112.4 | * | * | * | 0.1 | 0.9 | 5.1 | 16.9 | 53.7 | 171.7 | 580.0 | 2,368.1 |
| 2002..... | 115.4 | * | * | * | 0.8 | 4.8 | 16.7 | 55.0 | 178.4 | 602.6 | 2,458.6 | |
| 2001..... | 118.8 | * | * | * | 0.1 | 0.8 | 4.8 | 16.5 | 57.5 | 187.3 | 624.1 | 2,567.6 |
| 2000..... | 122.0 | * | * | * | 0.7 | 4.6 | 16.5 | 59.3 | 193.5 | 640.6 | 2,702.7 | |
| 1999..... | 126.4 | * | * | * | 0.1 | 0.8 | 4.7 | 17.3 | 62.9 | 204.6 | 671.1 | 2,802.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 121.4 | * | * | * | 0.1 | 1.6 | 10.0 | 33.9 | 87.4 | 233.1 | 700.2 | 2,343.9 |
| 2003..... | 128.0 | * | * | * | 0.2 | 1.7 | 10.2 | 34.6 | 91.7 | 250.2 | 743.0 | 2,533.4 |
| 2002..... | 131.0 | * | * | * | 0.2 | 1.5 | 10.2 | 34.9 | 94.8 | 261.3 | 770.0 | 2,623.4 |
| 2001..... | 132.7 | * | * | * | 0.2 | 1.6 | 9.6 | 33.8 | 97.4 | 271.4 | 791.2 | 2,724.1 |
| 2000..... | 136.1 | * | * | * | 0.2 | 1.4 | 9.4 | 34.6 | 100.7 | 283.7 | 811.9 | 2,883.6 |
| 1999..... | 139.9 | * | * | * | 0.2 | 1.4 | 9.6 | 35.2 | 103.9 | 297.9 | 847.2 | 2,973.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 126.1 | * | * | * | 0.2 | 2.4 | 15.4 | 53.9 | 131.5 | 329.5 | 937.9 | 2,640.6 |
| 2003..... | 131.9 | * | * | * | 0.2 | 2.6 | 15.7 | 55.1 | 138.9 | 354.5 | 995.5 | 2,828.2 |
| 2002..... | 134.5 | * | * | * | 0.2 | 2.3 | 16.2 | 56.4 | 143.5 | 371.2 | 1,033.6 | 2,908.8 |
| 2001..... | 134.4 | * | * | * | 0.3 | 2.5 | 15.0 | 54.1 | 147.0 | 384.3 | 1,057.5 | 3,002.8 |
| 2000..... | 137.7 | * | * | * | 0.2 | 2.2 | 15.1 | 55.7 | 152.7 | 406.1 | 1,086.5 | 3,211.3 |
| 1999..... | 140.4 | * | * | * | 0.2 | 2.1 | 15.2 | 56.1 | 155.1 | 425.3 | 1,130.5 | 3,276.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 116.9 | * | * | * | 0.7 | 4.5 | 14.2 | 45.6 | 150.1 | 538.3 | 2,211.1 | |
| 2003..... | 124.3 | * | * | * | 0.8 | 4.6 | 14.4 | 47.0 | 160.9 | 572.4 | 2,403.6 | |
| 2002..... | 127.7 | * | * | * | 0.7 | 4.2 | 13.8 | 48.7 | 167.6 | 593.4 | 2,500.1 | |
| 2001..... | 131.1 | * | * | * | 0.7 | 4.1 | 13.9 | 50.5 | 175.6 | 614.6 | 2,608.1 | |
| 2000..... | 134.6 | * | * | * | 0.5 | 3.8 | 13.8 | 51.8 | 180.6 | 631.3 | 2,751.4 | |
| 1999..... | 139.3 | * | * | * | 0.7 | 4.0 | 14.6 | 55.7 | 191.3 | 661.7 | 2,852.2 | |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other forms of chronic ischemic heart disease (I20,I25) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 80.3 | * | * | * | 0.4 | 3.0 | 14.6 | 60.3 | 163.1 | 363.9 | 822.2 | 2,031.7 |
| 2003..... | 84.7 | * | * | * | 0.4 | 2.7 | 14.6 | 61.9 | 173.6 | 393.8 | 878.9 | 2,204.8 |
| 2002..... | 85.5 | * | * | * | * | 2.2 | 15.5 | 62.9 | 180.3 | 395.2 | 901.5 | 2,246.7 |
| 2001..... | 88.2 | * | * | * | 0.5 | 2.9 | 15.6 | 62.9 | 186.6 | 419.6 | 934.8 | 2,357.3 |
| 2000..... | 89.1 | * | * | * | * | 2.5 | 15.6 | 64.3 | 192.3 | 431.5 | 947.1 | 2,398.0 |
| 1999..... | 91.6 | * | * | * | 0.4 | 2.4 | 17.2 | 66.9 | 197.9 | 453.5 | 958.2 | 2,471.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 83.9 | * | * | * | * | 4.5 | 20.4 | 87.7 | 237.3 | 493.9 | 1,048.2 | 2,090.1 |
| 2003..... | 87.9 | * | * | * | * | 4.0 | 21.3 | 91.6 | 246.2 | 541.3 | 1,135.2 | 2,162.8 |
| 2002..... | 88.0 | * | * | * | * | 3.0 | 22.2 | 90.8 | 264.4 | 536.0 | 1,119.3 | 2,254.6 |
| 2001..... | 90.8 | * | * | * | * | 4.3 | 22.9 | 93.1 | 266.3 | 573.5 | 1,177.8 | 2,378.1 |
| 2000..... | 90.4 | * | * | * | * | 3.7 | 22.1 | 93.6 | 274.7 | 566.6 | 1,180.1 | 2,463.5 |
| 1999..... | 93.3 | * | * | * | * | 3.5 | 25.1 | 99.2 | 283.6 | 602.1 | 1,169.2 | 2,552.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 76.9 | * | * | * | * | 1.6 | 9.4 | 36.8 | 103.6 | 271.3 | 694.6 | 2,008.6 |
| 2003..... | 81.7 | * | * | * | * | 1.5 | 8.7 | 36.5 | 115.4 | 288.8 | 734.1 | 2,221.1 |
| 2002..... | 83.3 | * | * | * | * | 1.6 | 9.6 | 39.0 | 112.8 | 295.4 | 778.3 | 2,243.7 |
| 2001..... | 86.0 | * | * | * | * | 1.6 | 9.2 | 36.9 | 122.4 | 310.9 | 798.4 | 2,349.6 |
| 2000..... | 88.0 | * | * | * | * | 1.5 | 10.0 | 39.1 | 125.9 | 336.5 | 815.9 | 2,373.7 |
| 1999..... | 90.1 | * | * | * | * | 1.3 | 10.2 | 39.4 | 129.2 | 348.6 | 840.0 | 2,441.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Atherosclerotic cardiovascular disease, so described (I25.0) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 21.7 | * | * | * | * | 0.5 | 3.3 | 13.6 | 32.0 | 58.7 | 129.0 | 400.4 |
| 2003..... | 23.1 | * | * | * | * | 0.5 | 3.4 | 13.9 | 32.7 | 62.7 | 142.5 | 442.9 |
| 2002..... | 23.6 | * | * | * | * | 0.5 | 3.4 | 13.4 | 33.1 | 64.7 | 146.2 | 474.5 |
| 2001..... | 23.9 | * | * | * | 0.1 | 0.5 | 3.2 | 12.9 | 33.3 | 65.6 | 152.7 | 493.6 |
| 2000..... | 24.7 | * | * | * | * | 0.4 | 3.0 | 13.2 | 34.0 | 68.3 | 158.1 | 534.6 |
| 1999..... | 25.8 | * | * | * | * | 0.4 | 3.3 | 13.0 | 35.3 | 72.7 | 166.2 | 563.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 23.9 | * | * | * | * | 0.8 | 5.2 | 22.0 | 49.0 | 83.6 | 164.9 | 414.0 |
| 2003..... | 25.2 | * | * | * | * | 0.9 | 5.2 | 22.7 | 49.9 | 88.9 | 181.1 | 462.1 |
| 2002..... | 25.4 | * | * | * | * | 0.7 | 5.4 | 21.5 | 51.2 | 91.8 | 185.3 | 491.8 |
| 2001..... | 25.3 | * | * | * | * | 0.7 | 5.0 | 21.0 | 51.4 | 92.4 | 193.9 | 502.0 |
| 2000..... | 26.0 | * | * | * | * | 0.6 | 4.9 | 21.5 | 52.6 | 97.2 | 199.3 | 550.8 |
| 1999..... | 26.8 | * | * | * | * | 0.6 | 5.3 | 20.9 | 53.7 | 103.0 | 207.0 | 578.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 19.5 | * | * | * | * | 0.2 | 1.4 | 5.5 | 16.1 | 37.7 | 104.8 | 394.3 |
| 2003..... | 21.1 | * | * | * | * | 0.2 | 1.5 | 5.5 | 16.7 | 40.8 | 116.7 | 434.4 |
| 2002..... | 21.9 | * | * | * | * | 0.2 | 1.3 | 5.5 | 16.2 | 42.1 | 120.2 | 467.1 |
| 2001..... | 22.5 | * | * | * | * | 0.2 | 1.4 | 5.1 | 16.7 | 43.3 | 125.5 | 490.1 |
| 2000..... | 23.5 | * | * | * | * | 0.1 | 1.2 | 5.2 | 16.8 | 44.6 | 131.3 | 528.0 |
| 1999..... | 24.8 | * | * | * | * | 0.2 | 1.3 | 5.3 | 18.3 | 47.9 | 139.8 | 557.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.9 | * | * | * | * | * | 1.7 | 9.7 | 29.3 | 55.8 | 118.9 | 279.7 |
| 2003..... | 8.3 | * | * | * | * | 0.3 | 1.6 | 9.8 | 29.4 | 57.3 | 124.6 | 357.1 |
| 2002..... | 8.2 | * | * | * | * | 0.4 | 1.9 | 8.9 | 29.7 | 61.1 | 124.1 | 358.6 |
| 2001..... | 8.5 | * | * | * | * | 0.4 | 1.5 | 10.1 | 29.7 | 65.9 | 132.5 | 417.0 |
| 2000..... | 8.4 | * | * | * | * | * | 1.6 | 9.5 | 30.5 | 65.7 | 128.2 | 452.5 |
| 1999..... | 9.0 | * | * | * | * | * | 1.5 | 10.4 | 33.0 | 72.1 | 147.5 | 469.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 9.8 | * | * | * | * | * | 2.7 | 16.6 | 47.1 | 83.3 | 169.5 | 297.0 |
| 2003..... | 10.2 | * | * | * | * | 0.5 | 2.6 | 17.1 | 48.7 | 88.6 | 169.5 | 397.1 |
| 2002..... | 9.7 | * | * | * | * | 0.6 | 2.9 | 14.8 | 49.3 | 90.8 | 153.9 | 415.0 |
| 2001..... | 10.1 | * | * | * | * | 0.6 | 2.2 | 17.1 | 47.9 | 100.2 | 181.1 | 440.0 |
| 2000..... | 9.9 | * | * | * | * | * | 2.7 | 16.6 | 50.5 | 92.4 | 180.9 | 493.8 |
| 1999..... | 10.6 | * | * | * | * | * | 2.5 | 17.8 | 50.0 | 115.8 | 194.0 | 485.3 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | * | * | * | 2.8 | 13.2 | 33.8 | 84.0 | 270.8 |
| 2003..... | 6.2 | * | * | * | * | * | * | 2.5 | 12.0 | 32.4 | 93.7 | 336.7 |
| 2002..... | 6.5 | * | * | * | * | * | * | 3.1 | 12.0 | 37.6 | 103.7 | 330.7 |
| 2001..... | 6.8 | * | * | * | * | * | * | 3.2 | 13.5 | 38.7 | 99.7 | 405.8 |
| 2000..... | 6.8 | * | * | * | * | * | * | 2.6 | 12.8 | 44.6 | 92.8 | 432.3 |
| 1999..... | 7.4 | * | * | * | * | * | * | 3.2 | 18.0 | 37.9 | 116.6 | 461.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Atherosclerotic cardiovascular disease, so described (I25.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 23.8 | * | * | * | * | 0.5 | 3.6 | 14.0 | 32.0 | 58.3 | 128.9 | 404.7 |
| 2003..... | 25.3 | * | * | * | * | 0.6 | 3.6 | 14.2 | 32.6 | 62.6 | 142.6 | 445.3 |
| 2002..... | 25.9 | * | * | * | * | 0.5 | 3.6 | 13.7 | 32.9 | 64.3 | 146.4 | 477.8 |
| 2001..... | 26.0 | * | * | * | * | 0.5 | 3.5 | 13.0 | 33.3 | 65.0 | 152.6 | 494.8 |
| 2000..... | 26.9 | * | * | * | * | 0.4 | 3.2 | 13.4 | 33.8 | 67.9 | 158.3 | 535.9 |
| 1999..... | 27.9 | * | * | * | * | 0.4 | 3.5 | 13.1 | 35.0 | 72.0 | 165.7 | 564.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 26.2 | * | * | * | * | 0.9 | 5.6 | 22.5 | 48.9 | 82.8 | 163.7 | 418.3 |
| 2003..... | 27.5 | * | * | * | * | 1.0 | 5.6 | 23.1 | 49.3 | 88.2 | 180.4 | 463.3 |
| 2002..... | 27.8 | * | * | * | * | 0.7 | 5.8 | 22.0 | 50.7 | 91.0 | 185.6 | 493.3 |
| 2001..... | 27.5 | * | * | * | * | 0.7 | 5.4 | 21.2 | 51.1 | 91.0 | 193.2 | 503.0 |
| 2000..... | 28.2 | * | * | * | * | 0.7 | 5.1 | 21.7 | 52.1 | 96.4 | 198.5 | 551.0 |
| 1999..... | 28.9 | * | * | * | * | 0.6 | 5.6 | 21.0 | 53.3 | 101.1 | 205.6 | 580.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 21.5 | * | * | * | * | 0.2 | 1.6 | 5.8 | 16.3 | 37.7 | 105.5 | 398.6 |
| 2003..... | 23.3 | * | * | * | * | 0.2 | 1.6 | 5.7 | 16.9 | 41.2 | 117.4 | 437.4 |
| 2002..... | 24.0 | * | * | * | * | 0.2 | 1.4 | 5.7 | 16.5 | 42.0 | 120.5 | 471.1 |
| 2001..... | 24.6 | * | * | * | * | 0.3 | 1.5 | 5.2 | 16.8 | 43.3 | 126.0 | 491.4 |
| 2000..... | 25.6 | * | * | * | * | 0.2 | 1.3 | 5.4 | 16.9 | 44.4 | 132.2 | 529.8 |
| 1999..... | 27.0 | * | * | * | * | 0.2 | 1.4 | 5.4 | 18.2 | 48.2 | 139.8 | 558.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 24.6 | * | * | * | * | 0.5 | 3.5 | 12.6 | 28.4 | 52.4 | 121.7 | 399.9 |
| 2003..... | 26.2 | * | * | * | * | 0.6 | 3.4 | 12.8 | 28.6 | 55.9 | 135.1 | 441.5 |
| 2002..... | 26.8 | * | * | * | * | 0.4 | 3.3 | 12.4 | 28.9 | 57.2 | 138.2 | 473.5 |
| 2001..... | 26.8 | * | * | * | * | 0.4 | 3.2 | 11.6 | 28.4 | 57.1 | 144.2 | 490.7 |
| 2000..... | 27.6 | * | * | * | * | 0.4 | 2.9 | 11.9 | 28.9 | 60.2 | 149.0 | 530.2 |
| 1999..... | 28.6 | * | * | * | * | 0.4 | 3.1 | 11.3 | 29.8 | 63.6 | 156.2 | 558.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 26.7 | * | * | * | * | 0.7 | 5.5 | 20.8 | 43.4 | 75.2 | 154.3 | 412.6 |
| 2003..... | 28.0 | * | * | * | * | 0.9 | 5.3 | 21.0 | 43.9 | 78.5 | 170.6 | 459.5 |
| 2002..... | 28.2 | * | * | * | * | 0.7 | 5.4 | 20.3 | 44.8 | 81.3 | 174.4 | 489.2 |
| 2001..... | 27.8 | * | * | * | * | 0.6 | 5.0 | 19.1 | 44.3 | 80.4 | 182.9 | 499.4 |
| 2000..... | 28.4 | * | * | * | * | 0.6 | 4.8 | 19.6 | 45.2 | 86.3 | 186.1 | 544.7 |
| 1999..... | 29.0 | * | * | * | * | 0.6 | 5.2 | 18.4 | 46.0 | 90.0 | 193.5 | 572.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 22.6 | * | * | * | * | 0.2 | 1.4 | 4.5 | 14.1 | 32.9 | 99.4 | 394.2 |
| 2003..... | 24.6 | * | * | * | * | 0.2 | 1.5 | 4.6 | 14.2 | 36.5 | 111.0 | 433.5 |
| 2002..... | 25.3 | * | * | * | * | 0.2 | 1.1 | 4.6 | 13.8 | 36.6 | 113.9 | 466.8 |
| 2001..... | 25.8 | * | * | * | * | 0.2 | 1.3 | 4.1 | 13.3 | 37.3 | 118.6 | 487.0 |
| 2000..... | 26.9 | * | * | * | * | * | 1.1 | 4.3 | 13.4 | 38.2 | 124.6 | 524.4 |
| 1999..... | 28.2 | * | * | * | * | 0.2 | 1.1 | 4.2 | 14.6 | 41.6 | 131.8 | 553.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Atherosclerotic cardiovascular disease, so described (I25.0) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 25.7 | * | * | * | * | 0.9 | 4.9 | 26.1 | 67.3 | 120.8 | 229.1 | 509.2 |
| 2003..... | 26.7 | * | * | * | * | 0.9 | 5.1 | 26.1 | 70.1 | 129.6 | 244.1 | 546.7 |
| 2002..... | 27.7 | * | * | * | * | 0.8 | 6.0 | 25.7 | 71.9 | 137.2 | 261.1 | 580.4 |
| 2001..... | 28.5 | * | * | * | * | 0.7 | 5.8 | 25.7 | 78.1 | 144.8 | 266.7 | 595.9 |
| 2000..... | 29.3 | * | * | * | * | 0.7 | 5.3 | 27.3 | 78.9 | 147.2 | 280.2 | 652.2 |
| 1999..... | 30.7 | * | * | * | * | 0.8 | 6.5 | 28.1 | 82.0 | 158.7 | 291.0 | 670.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 29.7 | * | * | * | * | 1.6 | 7.1 | 38.5 | 105.6 | 173.1 | 308.9 | 540.6 |
| 2003..... | 31.2 | * | * | * | * | 1.6 | 7.8 | 41.1 | 106.0 | 191.8 | 334.2 | 588.1 |
| 2002..... | 31.9 | * | * | * | * | 1.1 | 8.9 | 38.8 | 111.8 | 198.7 | 358.4 | 603.9 |
| 2001..... | 32.3 | * | * | * | * | 1.0 | 8.8 | 39.9 | 118.1 | 209.4 | 352.3 | 609.8 |
| 2000..... | 33.2 | * | * | * | * | 1.1 | 7.9 | 42.3 | 120.8 | 209.3 | 371.5 | 711.6 |
| 1999..... | 34.5 | * | * | * | * | 1.1 | 9.8 | 43.6 | 123.7 | 228.2 | 378.4 | 714.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 22.1 | * | * | * | * | * | 3.1 | 15.4 | 36.5 | 83.4 | 184.0 | 496.7 |
| 2003..... | 22.7 | * | * | * | * | * | 2.7 | 13.3 | 41.2 | 85.3 | 193.3 | 530.7 |
| 2002..... | 24.0 | * | * | * | * | * | 3.4 | 14.4 | 39.8 | 93.6 | 206.1 | 571.5 |
| 2001..... | 25.0 | * | * | * | * | * | 3.2 | 13.6 | 45.9 | 99.3 | 218.7 | 590.7 |
| 2000..... | 25.8 | * | * | * | * | * | 2.9 | 14.4 | 45.1 | 103.4 | 228.9 | 630.2 |
| 1999..... | 27.1 | * | * | * | * | * | 3.6 | 14.7 | 48.5 | 109.7 | 242.1 | 653.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other forms of chronic ischemic heart disease (I20,I25.1-I25.9) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 77.8 | * | * | * | 0.1 | 1.1 | 6.2 | 21.5 | 60.5 | 181.7 | 568.1 | 1,879.3 |
| 2003..... | 82.2 | * | * | * | 0.2 | 1.1 | 6.5 | 22.0 | 64.9 | 195.7 | 599.0 | 2,028.5 |
| 2002..... | 84.4 | * | * | * | 0.1 | 1.0 | 6.5 | 23.1 | 68.0 | 204.7 | 622.6 | 2,088.0 |
| 2001..... | 86.4 | * | * | * | 0.2 | 1.1 | 6.2 | 22.7 | 70.5 | 215.3 | 640.5 | 2,177.7 |
| 2000..... | 88.6 | * | * | * | 0.1 | 1.0 | 6.5 | 23.2 | 74.1 | 224.1 | 655.5 | 2,287.5 |
| 1999..... | 91.4 | * | * | * | 0.2 | 1.0 | 6.5 | 24.5 | 76.3 | 235.2 | 682.3 | 2,353.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 79.4 | * | * | * | 0.2 | 1.7 | 9.3 | 32.8 | 88.7 | 252.9 | 762.0 | 2,129.6 |
| 2003..... | 83.3 | * | * | * | 0.2 | 1.6 | 9.7 | 33.7 | 95.5 | 274.6 | 804.7 | 2,260.1 |
| 2002..... | 85.5 | * | * | * | 0.2 | 1.5 | 10.0 | 35.9 | 100.3 | 287.5 | 835.7 | 2,325.6 |
| 2001..... | 86.5 | * | * | * | 0.2 | 1.8 | 9.5 | 34.9 | 103.5 | 301.9 | 856.9 | 2,411.8 |
| 2000..... | 88.8 | * | * | * | 0.2 | 1.5 | 9.9 | 36.2 | 109.0 | 316.3 | 880.1 | 2,561.9 |
| 1999..... | 91.2 | * | * | * | 0.2 | 1.5 | 10.0 | 38.1 | 111.0 | 331.9 | 913.6 | 2,614.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 76.2 | * | * | * | * | 0.5 | 3.2 | 10.6 | 34.3 | 121.9 | 437.6 | 1,766.7 |
| 2003..... | 81.1 | * | * | * | 0.1 | 0.6 | 3.2 | 10.8 | 36.4 | 129.7 | 461.6 | 1,926.2 |
| 2002..... | 83.3 | * | * | * | * | 0.5 | 3.1 | 10.7 | 38.0 | 135.9 | 481.1 | 1,984.9 |
| 2001..... | 86.2 | * | * | * | 0.1 | 0.5 | 3.0 | 11.0 | 40.0 | 143.6 | 498.2 | 2,079.8 |
| 2000..... | 88.4 | * | * | * | * | 0.5 | 3.0 | 10.7 | 42.0 | 148.2 | 509.0 | 2,175.8 |
| 1999..... | 91.6 | * | * | * | 0.1 | 0.5 | 3.0 | 11.4 | 44.3 | 156.1 | 532.1 | 2,248.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 24.3 | * | * | * | * | 0.6 | 3.2 | 12.7 | 43.6 | 149.7 | 470.7 | 1,516.0 |
| 2003..... | 25.2 | * | * | * | * | 0.6 | 3.0 | 12.2 | 47.8 | 159.1 | 504.9 | 1,655.7 |
| 2002..... | 26.0 | * | * | * | * | 0.4 | 2.7 | 14.6 | 50.4 | 173.4 | 519.9 | 1,771.1 |
| 2001..... | 27.2 | * | * | * | * | 0.6 | 2.7 | 13.7 | 50.4 | 183.6 | 569.1 | 2,009.6 |
| 2000..... | 27.2 | * | * | * | * | 0.5 | 3.0 | 14.3 | 57.5 | 184.0 | 575.0 | 2,056.3 |
| 1999..... | 28.5 | * | * | * | * | 0.6 | 3.0 | 16.4 | 60.5 | 196.7 | 594.3 | 2,171.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 24.3 | * | * | * | * | 0.9 | 4.8 | 18.3 | 62.2 | 207.8 | 600.2 | 1,556.5 |
| 2003..... | 24.4 | * | * | * | * | 0.9 | 4.3 | 18.0 | 66.6 | 217.6 | 634.7 | 1,568.6 |
| 2002..... | 25.9 | * | * | * | * | 0.6 | 4.3 | 21.8 | 73.0 | 238.3 | 654.9 | 1,847.2 |
| 2001..... | 26.7 | * | * | * | * | 0.9 | 4.3 | 20.4 | 73.6 | 250.0 | 720.4 | 2,039.8 |
| 2000..... | 27.0 | * | * | * | * | 0.7 | 4.7 | 22.6 | 83.6 | 259.1 | 719.8 | 2,084.0 |
| 1999..... | 28.1 | * | * | * | * | 1.0 | 5.1 | 26.5 | 88.5 | 266.0 | 738.8 | 2,221.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 24.2 | * | * | * | * | * | 1.3 | 7.0 | 26.7 | 103.2 | 381.3 | 1,495.2 |
| 2003..... | 26.1 | * | * | * | * | * | 1.6 | 6.4 | 30.7 | 112.5 | 415.8 | 1,700.0 |
| 2002..... | 26.1 | * | * | * | * | * | 0.9 | 7.4 | 30.2 | 121.8 | 427.5 | 1,733.3 |
| 2001..... | 27.7 | * | * | * | * | * | 1.0 | 7.1 | 29.8 | 131.2 | 466.9 | 1,994.8 |
| 2000..... | 27.5 | * | * | * | * | * | 1.2 | 6.2 | 34.4 | 124.9 | 477.7 | 2,042.7 |
| 1999..... | 28.9 | * | * | * | * | * | 0.8 | 6.5 | 35.8 | 142.3 | 498.2 | 2,147.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other forms of chronic ischemic heart disease (I20,I25.1-I25.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 86.3 | * | * | * | 0.2 | 1.2 | 6.7 | 22.4 | 61.6 | 183.5 | 572.4 | 1,892.8 |
| 2003..... | 91.0 | * | * | * | 0.2 | 1.2 | 7.0 | 22.9 | 65.9 | 197.6 | 602.3 | 2,039.6 |
| 2002..... | 93.1 | * | * | * | 0.1 | 1.1 | 7.1 | 23.8 | 69.0 | 206.0 | 625.4 | 2,093.4 |
| 2001..... | 94.9 | * | * | * | 0.2 | 1.2 | 6.7 | 23.5 | 71.8 | 216.7 | 641.6 | 2,177.4 |
| 2000..... | 97.0 | * | * | * | 0.1 | 1.1 | 6.9 | 23.9 | 75.0 | 225.6 | 656.7 | 2,288.1 |
| 1999..... | 99.7 | * | * | * | 0.2 | 1.1 | 6.9 | 25.1 | 77.1 | 236.5 | 683.4 | 2,351.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 88.7 | * | * | * | 0.2 | 1.8 | 10.0 | 34.2 | 90.4 | 255.3 | 769.4 | 2,154.8 |
| 2003..... | 92.9 | * | * | * | 0.2 | 1.8 | 10.6 | 35.2 | 97.4 | 277.5 | 811.5 | 2,287.4 |
| 2002..... | 94.9 | * | * | * | 0.2 | 1.6 | 10.8 | 37.2 | 102.0 | 289.3 | 841.7 | 2,338.9 |
| 2001..... | 95.6 | * | * | * | 0.2 | 2.0 | 10.3 | 36.1 | 105.3 | 304.1 | 860.3 | 2,420.7 |
| 2000..... | 97.8 | * | * | * | 0.2 | 1.7 | 10.5 | 37.2 | 110.2 | 318.2 | 884.2 | 2,572.9 |
| 1999..... | 100.0 | * | * | * | 0.2 | 1.6 | 10.6 | 38.9 | 112.1 | 334.1 | 917.9 | 2,619.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 84.1 | * | * | * | * | 0.6 | 3.4 | 11.0 | 34.9 | 123.0 | 439.9 | 1,775.7 |
| 2003..... | 89.2 | * | * | * | 0.1 | 0.7 | 3.4 | 11.2 | 36.7 | 130.6 | 462.6 | 1,930.7 |
| 2002..... | 91.4 | * | * | * | * | 0.6 | 3.4 | 11.0 | 38.5 | 136.4 | 482.1 | 1,987.5 |
| 2001..... | 94.2 | * | * | * | * | 0.5 | 3.2 | 11.3 | 40.7 | 143.9 | 498.0 | 2,076.2 |
| 2000..... | 96.3 | * | * | * | * | 0.5 | 3.3 | 11.1 | 42.4 | 149.1 | 508.5 | 2,172.9 |
| 1999..... | 99.4 | * | * | * | * | 0.6 | 3.3 | 11.8 | 44.8 | 156.4 | 531.2 | 2,243.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 96.8 | * | * | * | 0.1 | 1.1 | 6.5 | 21.3 | 59.1 | 180.6 | 578.5 | 1,944.1 |
| 2003..... | 101.8 | * | * | * | 0.1 | 1.1 | 6.7 | 21.8 | 63.1 | 194.3 | 607.9 | 2,091.9 |
| 2002..... | 104.3 | * | * | * | 0.1 | 1.1 | 6.9 | 22.5 | 65.9 | 204.1 | 631.9 | 2,149.8 |
| 2001..... | 106.0 | * | * | * | 0.1 | 1.1 | 6.4 | 22.2 | 69.0 | 214.3 | 647.0 | 2,233.4 |
| 2000..... | 108.4 | * | * | * | 0.1 | 1.0 | 6.5 | 22.7 | 71.9 | 223.5 | 662.9 | 2,353.3 |
| 1999..... | 111.3 | * | * | * | 0.1 | 1.0 | 6.5 | 23.9 | 74.1 | 234.2 | 690.9 | 2,414.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 99.3 | * | * | * | 0.2 | 1.7 | 9.9 | 33.1 | 88.1 | 254.3 | 783.6 | 2,228.0 |
| 2003..... | 103.9 | * | * | * | 0.2 | 1.7 | 10.4 | 34.1 | 94.9 | 276.0 | 824.8 | 2,368.6 |
| 2002..... | 106.2 | * | * | * | 0.2 | 1.6 | 10.7 | 36.1 | 98.7 | 289.9 | 859.2 | 2,419.7 |
| 2001..... | 106.6 | * | * | * | 0.2 | 1.8 | 10.0 | 34.9 | 102.7 | 303.9 | 874.6 | 2,503.4 |
| 2000..... | 109.2 | * | * | * | 0.2 | 1.6 | 10.3 | 36.1 | 107.5 | 319.8 | 900.4 | 2,666.6 |
| 1999..... | 111.4 | * | * | * | 0.2 | 1.5 | 10.1 | 37.7 | 109.1 | 335.3 | 936.9 | 2,703.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 94.3 | * | * | * | * | 0.5 | 3.1 | 9.7 | 31.5 | 117.3 | 438.8 | 1,816.9 |
| 2003..... | 99.8 | * | * | * | * | 0.6 | 3.1 | 9.7 | 32.8 | 124.3 | 461.4 | 1,970.1 |
| 2002..... | 102.4 | * | * | * | * | 0.5 | 3.1 | 9.2 | 34.9 | 131.0 | 479.6 | 2,033.3 |
| 2001..... | 105.3 | * | * | * | * | 0.5 | 2.9 | 9.8 | 37.2 | 138.2 | 496.0 | 2,121.0 |
| 2000..... | 107.7 | * | * | * | * | 0.4 | 2.7 | 9.5 | 38.3 | 142.4 | 506.8 | 2,227.0 |
| 1999..... | 111.1 | * | * | * | * | 0.5 | 2.8 | 10.4 | 41.1 | 149.7 | 529.9 | 2,299.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other forms of chronic ischemic heart disease (I20,I25.1-I25.9) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 54.6 | * | * | * | 0.4 | 2.1 | 9.6 | 34.2 | 95.9 | 243.2 | 593.1 | 1,522.6 |
| 2003..... | 57.9 | * | * | * | * | 1.8 | 9.5 | 35.8 | 103.6 | 264.2 | 634.8 | 1,658.1 |
| 2002..... | 57.8 | * | * | * | * | 1.5 | 9.6 | 37.2 | 108.4 | 258.0 | 640.3 | 1,666.3 |
| 2001..... | 59.7 | * | * | * | 0.4 | 2.1 | 9.8 | 37.1 | 108.4 | 274.7 | 668.1 | 1,761.4 |
| 2000..... | 59.8 | * | * | * | * | 1.8 | 10.4 | 37.0 | 113.4 | 284.4 | 666.8 | 1,745.8 |
| 1999..... | 61.0 | * | * | * | * | 1.6 | 10.7 | 38.9 | 115.9 | 294.7 | 667.1 | 1,801.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 54.2 | * | * | * | * | 3.0 | 13.4 | 49.2 | 131.6 | 320.8 | 739.3 | 1,549.6 |
| 2003..... | 56.7 | * | * | * | * | 2.4 | 13.4 | 50.5 | 140.2 | 349.5 | 801.1 | 1,574.7 |
| 2002..... | 56.1 | * | * | * | * | 1.9 | 13.3 | 52.0 | 152.5 | 337.3 | 760.8 | 1,650.8 |
| 2001..... | 58.4 | * | * | * | * | 3.3 | 14.1 | 53.2 | 148.2 | 364.1 | 825.5 | 1,768.3 |
| 2000..... | 57.3 | * | * | * | * | 2.6 | 14.1 | 51.3 | 153.9 | 357.2 | 808.7 | 1,751.9 |
| 1999..... | 58.8 | * | * | * | * | 2.4 | 15.3 | 55.5 | 160.0 | 374.0 | 790.8 | 1,838.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 54.9 | * | * | * | * | 1.2 | 6.3 | 21.3 | 67.1 | 187.8 | 510.6 | 1,511.9 |
| 2003..... | 59.1 | * | * | * | * | 1.2 | 6.1 | 23.2 | 74.2 | 203.5 | 540.8 | 1,690.4 |
| 2002..... | 59.3 | * | * | * | * | 1.1 | 6.2 | 24.5 | 73.0 | 201.8 | 572.2 | 1,672.2 |
| 2001..... | 60.9 | * | * | * | * | 1.0 | 6.0 | 23.3 | 76.5 | 211.7 | 579.7 | 1,758.9 |
| 2000..... | 62.1 | * | * | * | * | 1.1 | 7.1 | 24.7 | 80.7 | 233.1 | 587.0 | 1,743.5 |
| 1999..... | 63.0 | * | * | * | * | 0.8 | 6.6 | 24.6 | 80.7 | 238.9 | 597.9 | 1,787.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other heart diseases (I26-I51) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 56.6 | 10.0 | 1.1 | 0.5 | 2.1 | 4.3 | 9.7 | 21.5 | 45.2 | 118.6 | 368.9 | 1,373.9 |
| 2003..... | 58.5 | 10.4 | 1.1 | 0.6 | 2.2 | 4.5 | 10.3 | 21.4 | 48.0 | 126.8 | 382.8 | 1,427.2 |
| 2002..... | 58.8 | 11.7 | 1.0 | 0.6 | 2.1 | 4.5 | 10.3 | 21.7 | 48.7 | 129.6 | 390.9 | 1,443.5 |
| 2001..... | 58.6 | 11.1 | 1.4 | 0.6 | 2.0 | 4.4 | 9.9 | 21.6 | 49.5 | 131.4 | 389.5 | 1,475.5 |
| 2000..... | 58.8 | 12.0 | 1.1 | 0.6 | 2.2 | 4.4 | 9.9 | 21.5 | 50.9 | 134.2 | 391.3 | 1,509.4 |
| 1999..... | 59.4 | 12.9 | 1.2 | 0.6 | 2.3 | 4.4 | 10.3 | 21.6 | 52.9 | 138.8 | 398.5 | 1,512.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 50.6 | 10.5 | 1.1 | 0.5 | 2.6 | 5.3 | 12.0 | 27.7 | 57.7 | 142.8 | 420.8 | 1,377.7 |
| 2003..... | 52.3 | 11.4 | 1.0 | 0.7 | 2.6 | 5.3 | 12.7 | 27.2 | 60.7 | 154.2 | 440.0 | 1,431.3 |
| 2002..... | 52.4 | 12.1 | 1.0 | 0.6 | 2.7 | 5.6 | 12.8 | 28.2 | 60.8 | 156.5 | 448.2 | 1,448.6 |
| 2001..... | 51.8 | 10.8 | 1.5 | 0.6 | 2.4 | 5.2 | 12.4 | 27.6 | 61.7 | 158.7 | 452.0 | 1,473.4 |
| 2000..... | 52.2 | 12.2 | 1.2 | 0.7 | 2.6 | 5.2 | 12.2 | 27.8 | 64.7 | 164.0 | 451.0 | 1,549.1 |
| 1999..... | 53.1 | 13.1 | 1.3 | 0.7 | 2.7 | 5.3 | 12.9 | 27.7 | 67.5 | 172.7 | 459.7 | 1,548.9 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 62.4 | 9.4 | 1.1 | 0.6 | 1.5 | 3.3 | 7.4 | 15.5 | 33.6 | 98.4 | 334.0 | 1,372.1 |
| 2003..... | 64.5 | 9.3 | 1.2 | 0.5 | 1.7 | 3.7 | 7.9 | 15.7 | 36.3 | 103.8 | 344.5 | 1,425.4 |
| 2002..... | 65.0 | 11.4 | 1.0 | 0.5 | 1.4 | 3.4 | 7.8 | 15.5 | 37.5 | 107.2 | 352.8 | 1,441.3 |
| 2001..... | 65.2 | 11.5 | 1.3 | 0.6 | 1.5 | 3.6 | 7.5 | 15.9 | 38.3 | 108.7 | 348.4 | 1,476.4 |
| 2000..... | 65.1 | 11.8 | 0.9 | 0.5 | 1.8 | 3.5 | 7.5 | 15.3 | 38.2 | 109.6 | 352.3 | 1,493.2 |
| 1999..... | 65.5 | 12.7 | 1.0 | 0.6 | 1.8 | 3.6 | 7.7 | 15.6 | 39.4 | 111.1 | 358.7 | 1,498.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 13.2 | 8.5 | 1.0 | 0.4 | 1.5 | 2.3 | 5.3 | 11.4 | 24.7 | 70.3 | 208.5 | 659.3 |
| 2003..... | 14.1 | 8.3 | 1.0 | 0.3 | 1.4 | 2.8 | 5.3 | 11.3 | 29.7 | 77.3 | 221.2 | 749.9 |
| 2002..... | 13.3 | 9.7 | 0.7 | 0.4 | 1.5 | 2.6 | 5.1 | 11.4 | 25.7 | 72.6 | 219.8 | 740.0 |
| 2001..... | 13.3 | 9.2 | 1.0 | 0.5 | 1.3 | 2.6 | 4.9 | 10.9 | 27.9 | 75.4 | 217.5 | 798.5 |
| 2000..... | 13.6 | 10.2 | 1.0 | 0.5 | 1.7 | 2.8 | 5.5 | 11.6 | 28.8 | 75.9 | 227.0 | 829.4 |
| 1999..... | 13.8 | 12.2 | 1.2 | 0.5 | 1.6 | 2.8 | 6.3 | 11.6 | 30.6 | 78.7 | 229.6 | 827.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 13.0 | 8.4 | 1.3 | * | 2.2 | 3.3 | 7.1 | 15.5 | 31.6 | 83.6 | 241.8 | 641.6 |
| 2003..... | 13.6 | 9.1 | * | * | 1.8 | 3.4 | 7.0 | 14.7 | 37.0 | 95.0 | 253.5 | 752.7 |
| 2002..... | 12.9 | 9.1 | * | * | 2.1 | 3.4 | 6.4 | 14.3 | 31.7 | 92.5 | 256.9 | 758.0 |
| 2001..... | 12.6 | 9.6 | * | * | 1.6 | 3.2 | 6.5 | 15.0 | 34.1 | 91.1 | 248.6 | 775.2 |
| 2000..... | 13.2 | 9.9 | 1.3 | 0.7 | 2.1 | 3.5 | 6.7 | 15.1 | 37.2 | 94.1 | 250.8 | 890.8 |
| 1999..... | 13.7 | 10.3 | 1.4 | 0.6 | 1.8 | 3.8 | 8.2 | 15.4 | 38.8 | 102.4 | 267.5 | 888.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 13.5 | 8.6 | * | * | 0.8 | 1.1 | 3.3 | 7.3 | 18.5 | 59.6 | 185.5 | 668.4 |
| 2003..... | 14.5 | 7.6 | * | * | 0.9 | 2.2 | 3.3 | 8.0 | 23.2 | 63.2 | 199.1 | 748.4 |
| 2002..... | 13.7 | 10.3 | * | * | 0.8 | 1.6 | 3.7 | 8.5 | 20.3 | 56.7 | 194.5 | 731.1 |
| 2001..... | 13.9 | 8.7 | * | * | 0.9 | 1.9 | 3.2 | 6.9 | 22.5 | 63.0 | 196.4 | 809.8 |
| 2000..... | 14.0 | 10.6 | * | * | 1.1 | 1.9 | 4.3 | 8.1 | 21.4 | 61.6 | 211.0 | 799.3 |
| 1999..... | 13.9 | 14.1 | * | * | 1.3 | 1.6 | 4.2 | 8.0 | 23.4 | 60.1 | 204.3 | 798.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other heart diseases (I26-I51) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 63.6 | 10.3 | 1.1 | 0.6 | 2.1 | 4.7 | 10.4 | 22.5 | 46.7 | 122.0 | 377.3 | 1,404.5 |
| 2003..... | 65.4 | 10.9 | 1.1 | 0.6 | 2.3 | 4.9 | 11.0 | 22.3 | 49.3 | 130.0 | 390.5 | 1,453.7 |
| 2002..... | 65.7 | 12.2 | 1.1 | 0.6 | 2.2 | 4.9 | 11.0 | 22.7 | 50.4 | 133.2 | 398.4 | 1,469.2 |
| 2001..... | 65.2 | 11.6 | 1.5 | 0.6 | 2.1 | 4.7 | 10.6 | 22.6 | 51.1 | 134.7 | 396.5 | 1,497.8 |
| 2000..... | 65.1 | 12.4 | 1.1 | 0.6 | 2.3 | 4.7 | 10.4 | 22.3 | 52.4 | 137.5 | 397.5 | 1,530.9 |
| 1999..... | 65.6 | 13.1 | 1.1 | 0.6 | 2.4 | 4.7 | 10.8 | 22.4 | 54.3 | 142.0 | 404.4 | 1,533.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 57.0 | 11.0 | 1.0 | 0.6 | 2.7 | 5.8 | 12.8 | 28.9 | 59.6 | 146.8 | 430.2 | 1,412.5 |
| 2003..... | 58.6 | 12.0 | 1.0 | 0.7 | 2.8 | 5.8 | 13.5 | 28.4 | 62.2 | 158.0 | 448.9 | 1,460.5 |
| 2002..... | 58.7 | 12.8 | 1.1 | 0.7 | 2.8 | 6.2 | 13.7 | 29.5 | 62.9 | 160.4 | 456.6 | 1,477.3 |
| 2001..... | 57.8 | 11.0 | 1.6 | 0.6 | 2.6 | 5.6 | 13.2 | 28.7 | 63.6 | 162.5 | 460.6 | 1,499.0 |
| 2000..... | 58.0 | 12.7 | 1.2 | 0.7 | 2.7 | 5.5 | 12.9 | 28.9 | 66.4 | 167.8 | 458.9 | 1,573.6 |
| 1999..... | 58.6 | 13.8 | 1.3 | 0.7 | 2.9 | 5.5 | 13.5 | 28.7 | 69.3 | 176.2 | 466.4 | 1,571.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 69.9 | 9.6 | 1.2 | 0.6 | 1.6 | 3.7 | 8.0 | 16.4 | 34.8 | 101.1 | 341.7 | 1,400.9 |
| 2003..... | 71.8 | 9.7 | 1.2 | 0.5 | 1.9 | 4.0 | 8.5 | 16.4 | 37.3 | 106.5 | 351.5 | 1,450.7 |
| 2002..... | 72.3 | 11.6 | 1.0 | 0.5 | 1.5 | 3.7 | 8.4 | 16.1 | 38.7 | 110.4 | 359.7 | 1,465.7 |
| 2001..... | 72.3 | 12.2 | 1.3 | 0.6 | 1.6 | 3.9 | 8.0 | 16.7 | 39.4 | 111.6 | 354.4 | 1,497.3 |
| 2000..... | 71.8 | 12.0 | 1.0 | 0.5 | 1.9 | 3.8 | 7.9 | 15.9 | 39.4 | 112.5 | 357.5 | 1,513.7 |
| 1999..... | 72.2 | 12.3 | 1.0 | 0.6 | 1.9 | 3.9 | 8.1 | 16.2 | 40.5 | 113.9 | 364.2 | 1,519.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 69.3 | 8.7 | 0.8 | 0.5 | 1.8 | 3.8 | 8.6 | 19.0 | 41.9 | 116.5 | 377.3 | 1,442.1 |
| 2003..... | 71.0 | 8.1 | 0.8 | 0.5 | 1.9 | 3.8 | 9.1 | 18.8 | 44.1 | 123.1 | 391.4 | 1,490.1 |
| 2002..... | 71.3 | 10.0 | 0.9 | 0.5 | 1.8 | 3.8 | 9.3 | 19.0 | 45.3 | 127.6 | 398.1 | 1,503.1 |
| 2001..... | 70.6 | 9.1 | 1.1 | 0.5 | 1.7 | 3.7 | 8.5 | 19.0 | 45.8 | 128.7 | 394.8 | 1,533.2 |
| 2000..... | 70.3 | 10.5 | 0.9 | 0.5 | 1.7 | 3.7 | 8.4 | 18.7 | 47.4 | 131.2 | 395.8 | 1,564.2 |
| 1999..... | 70.4 | 10.5 | 0.9 | 0.5 | 2.0 | 3.7 | 8.7 | 18.7 | 48.3 | 135.2 | 401.1 | 1,565.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 60.9 | 9.3 | 0.7 | 0.5 | 2.3 | 4.6 | 10.8 | 24.7 | 53.6 | 140.9 | 428.9 | 1,445.4 |
| 2003..... | 62.6 | 8.7 | 0.7 | 0.6 | 2.3 | 4.6 | 11.4 | 24.3 | 56.0 | 150.6 | 450.4 | 1,499.3 |
| 2002..... | 62.7 | 10.7 | 1.1 | 0.6 | 2.3 | 4.7 | 11.8 | 25.2 | 57.1 | 154.1 | 457.1 | 1,512.9 |
| 2001..... | 61.4 | 9.4 | 1.2 | 0.6 | 2.0 | 4.4 | 10.9 | 24.8 | 57.5 | 155.8 | 457.4 | 1,530.7 |
| 2000..... | 61.6 | 10.9 | 1.1 | 0.6 | 2.0 | 4.4 | 10.7 | 24.6 | 60.7 | 161.0 | 457.6 | 1,606.4 |
| 1999..... | 61.7 | 11.6 | 1.1 | 0.5 | 2.3 | 4.4 | 11.2 | 24.5 | 62.1 | 168.5 | 462.9 | 1,595.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 77.3 | 8.2 | 0.9 | 0.4 | 1.3 | 2.9 | 6.4 | 13.3 | 30.7 | 95.6 | 342.1 | 1,440.6 |
| 2003..... | 79.1 | 7.5 | 1.0 | 0.4 | 1.6 | 3.0 | 6.8 | 13.5 | 32.7 | 99.6 | 351.5 | 1,486.1 |
| 2002..... | 79.6 | 9.3 | 0.8 | 0.5 | 1.2 | 2.8 | 6.8 | 13.0 | 34.1 | 105.0 | 358.5 | 1,498.8 |
| 2001..... | 79.4 | 8.8 | 1.0 | 0.5 | 1.3 | 3.1 | 6.1 | 13.4 | 34.8 | 105.7 | 353.3 | 1,534.2 |
| 2000..... | 78.7 | 10.1 | 0.8 | 0.4 | 1.4 | 2.9 | 6.2 | 12.9 | 34.8 | 106.0 | 355.3 | 1,547.2 |
| 1999..... | 78.8 | 9.4 | 0.8 | 0.5 | 1.7 | 3.0 | 6.2 | 13.1 | 35.3 | 107.3 | 360.6 | 1,554.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other heart diseases (I26-I51) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 52.7 | 15.3 | 2.4 | 1.1 | 3.8 | 10.4 | 23.0 | 51.4 | 98.5 | 199.4 | 458.1 | 1,197.9 |
| 2003..... | 54.4 | 19.8 | 2.0 | 1.0 | 4.5 | 11.3 | 24.4 | 51.1 | 104.2 | 219.8 | 455.6 | 1,253.2 |
| 2002..... | 54.3 | 20.9 | 1.6 | 0.9 | 4.2 | 11.8 | 23.4 | 52.6 | 104.1 | 211.2 | 474.7 | 1,285.5 |
| 2001..... | 54.7 | 21.3 | 3.0 | 1.0 | 4.4 | 11.0 | 24.5 | 52.2 | 105.7 | 216.1 | 486.4 | 1,292.9 |
| 2000..... | 54.6 | 19.6 | 2.0 | 1.1 | 4.9 | 10.8 | 23.7 | 52.8 | 104.5 | 223.6 | 484.3 | 1,319.4 |
| 1999..... | 56.2 | 23.2 | 2.0 | 1.0 | 4.7 | 11.0 | 25.3 | 53.5 | 115.3 | 232.9 | 499.9 | 1,314.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 52.3 | 16.4 | 2.5 | 1.0 | 4.8 | 12.6 | 27.7 | 65.3 | 128.9 | 242.3 | 550.5 | 1,244.7 |
| 2003..... | 53.4 | 23.6 | 2.2 | 1.2 | 5.8 | 13.0 | 29.6 | 65.1 | 134.7 | 269.3 | 530.4 | 1,239.1 |
| 2002..... | 52.6 | 21.8 | * | 0.9 | 5.1 | 14.8 | 28.2 | 67.5 | 130.4 | 259.4 | 543.4 | 1,258.0 |
| 2001..... | 53.5 | 18.8 | 3.3 | 0.9 | 5.5 | 13.4 | 29.3 | 63.9 | 134.9 | 261.9 | 593.5 | 1,362.9 |
| 2000..... | 53.3 | 19.1 | 2.0 | 1.2 | 5.7 | 12.7 | 28.7 | 68.3 | 132.9 | 269.8 | 565.5 | 1,416.1 |
| 1999..... | 56.2 | 22.9 | 2.3 | 1.2 | 6.0 | 13.4 | 30.9 | 68.7 | 149.5 | 291.0 | 584.7 | 1,466.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 53.1 | 14.2 | 2.3 | 1.2 | 2.8 | 8.4 | 19.0 | 39.4 | 74.0 | 168.8 | 405.9 | 1,179.4 |
| 2003..... | 55.4 | 15.8 | 1.8 | 0.9 | 3.3 | 9.7 | 19.8 | 39.2 | 79.6 | 184.6 | 413.4 | 1,258.7 |
| 2002..... | 55.9 | 19.9 | 1.9 | 0.8 | 3.2 | 9.0 | 19.2 | 39.9 | 83.0 | 177.0 | 435.9 | 1,296.0 |
| 2001..... | 55.8 | 23.8 | 2.7 | 1.0 | 3.2 | 8.9 | 20.3 | 42.2 | 82.3 | 183.7 | 426.2 | 1,267.0 |
| 2000..... | 55.8 | 20.1 | 1.9 | 1.0 | 4.2 | 9.0 | 19.2 | 39.6 | 81.6 | 191.1 | 438.5 | 1,283.4 |
| 1999..... | 56.3 | 23.6 | 1.8 | 0.9 | 3.4 | 8.9 | 20.4 | 40.5 | 87.9 | 192.0 | 452.4 | 1,257.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pulmonary heart disease and diseases of pulmonary circulation (I26-I28) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.5 | 3.8 | 0.2 | 0.1 | 0.3 | 0.8 | 1.7 | 3.3 | 5.8 | 13.4 | 27.5 | 54.8 |
| 2003..... | 4.8 | 3.3 | 0.2 | 0.1 | 0.3 | 0.9 | 1.9 | 3.3 | 6.5 | 14.4 | 30.4 | 58.0 |
| 2002..... | 4.8 | 4.6 | 0.2 | 0.1 | 0.3 | 0.8 | 1.7 | 3.4 | 6.9 | 14.9 | 29.8 | 56.3 |
| 2001..... | 4.7 | 4.2 | 0.1 | 0.1 | 0.3 | 0.9 | 1.7 | 3.4 | 6.9 | 14.6 | 28.5 | 57.5 |
| 2000..... | 4.6 | 4.9 | 0.2 | 0.1 | 0.3 | 0.9 | 1.7 | 3.5 | 6.7 | 14.5 | 28.3 | 57.4 |
| 1999..... | 4.7 | 5.7 | * | 0.1 | 0.4 | 0.9 | 1.8 | 3.3 | 6.7 | 14.8 | 30.0 | 59.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.6 | 4.1 | * | * | 0.2 | 0.6 | 1.4 | 3.1 | 5.8 | 12.5 | 25.2 | 48.3 |
| 2003..... | 3.9 | 3.7 | * | * | 0.2 | 0.6 | 1.6 | 3.3 | 6.2 | 14.2 | 29.9 | 54.4 |
| 2002..... | 3.8 | 4.7 | * | * | 0.2 | 0.7 | 1.4 | 3.1 | 6.5 | 14.1 | 27.7 | 52.1 |
| 2001..... | 3.8 | 3.9 | * | * | 0.2 | 0.6 | 1.5 | 3.0 | 6.5 | 14.5 | 27.9 | 55.6 |
| 2000..... | 3.8 | 5.1 | * | * | 0.2 | 0.7 | 1.4 | 3.3 | 6.4 | 13.9 | 28.3 | 54.8 |
| 1999..... | 3.9 | 5.9 | * | * | 0.3 | 0.6 | 1.5 | 3.0 | 6.8 | 15.1 | 29.1 | 60.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.3 | 3.5 | * | * | 0.4 | 0.9 | 1.9 | 3.4 | 5.7 | 14.3 | 29.1 | 57.7 |
| 2003..... | 5.7 | 2.9 | * | * | 0.5 | 1.1 | 2.2 | 3.4 | 6.9 | 14.6 | 30.8 | 59.6 |
| 2002..... | 5.7 | 4.5 | * | * | 0.4 | 1.0 | 1.9 | 3.6 | 7.2 | 15.6 | 31.2 | 58.1 |
| 2001..... | 5.5 | 4.6 | * | * | 0.4 | 1.1 | 2.0 | 3.7 | 7.3 | 14.6 | 28.9 | 58.2 |
| 2000..... | 5.5 | 4.6 | * | * | 0.5 | 1.0 | 2.1 | 3.7 | 7.0 | 15.0 | 28.4 | 58.5 |
| 1999..... | 5.5 | 5.6 | * | * | 0.5 | 1.1 | 2.0 | 3.6 | 6.6 | 14.6 | 30.5 | 59.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | 2.2 | * | * | * | 0.4 | 0.9 | 1.9 | 2.8 | 6.8 | 12.4 | 31.9 |
| 2003..... | 1.2 | * | * | * | * | 0.5 | 0.9 | 1.3 | 3.4 | 6.0 | 15.9 | 29.5 |
| 2002..... | 1.3 | 3.1 | * | * | * | 0.4 | 0.9 | 1.8 | 3.1 | 7.5 | 16.8 | 30.5 |
| 2001..... | 1.3 | 3.7 | * | * | * | 0.5 | 0.9 | 1.4 | 3.5 | 6.9 | 14.6 | 32.5 |
| 2000..... | 1.2 | 4.3 | * | * | * | 0.5 | 1.1 | 2.0 | 3.0 | 6.2 | 14.4 | 27.9 |
| 1999..... | 1.4 | 5.4 | * | * | 0.3 | 0.5 | 1.0 | 1.6 | 3.4 | 7.1 | 16.2 | 37.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 1.0 | 1.7 | 2.6 | 5.8 | 8.9 | * |
| 2003..... | 0.8 | * | * | * | * | * | 0.8 | * | 3.1 | 4.2 | 15.0 | * |
| 2002..... | 0.9 | * | * | * | * | * | 0.7 | 1.6 | 2.8 | 8.6 | 11.9 | * |
| 2001..... | 1.1 | * | * | * | * | * | 0.9 | 1.4 | 2.9 | 7.2 | 15.2 | 43.8 |
| 2000..... | 0.9 | * | * | * | * | * | 0.8 | 2.0 | 2.5 | 5.7 | 12.3 | * |
| 1999..... | 1.1 | 5.2 | * | * | * | * | 0.8 | * | 3.3 | 6.8 | 15.1 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.9 | 2.1 | 3.0 | 7.7 | 14.8 | 36.0 |
| 2003..... | 1.6 | * | * | * | * | 0.8 | 1.1 | 1.7 | 3.7 | 7.4 | 16.6 | 33.6 |
| 2002..... | 1.6 | * | * | * | * | * | 1.3 | 2.0 | 3.5 | 6.5 | 20.2 | 35.6 |
| 2001..... | 1.5 | * | * | * | * | 0.8 | 1.0 | 1.5 | 4.1 | 6.6 | 14.2 | 27.0 |
| 2000..... | 1.5 | * | * | * | * | 0.7 | 1.4 | 2.0 | 3.4 | 6.6 | 15.8 | 32.6 |
| 1999..... | 1.7 | 5.7 | * | * | * | 0.7 | 1.3 | 2.0 | 3.5 | 7.4 | 17.0 | 35.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pulmonary heart disease and diseases of pulmonary circulation (I26-I28) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.0 | 4.2 | * | 0.1 | 0.3 | 0.9 | 1.8 | 3.4 | 6.0 | 13.9 | 28.3 | 55.8 |
| 2003..... | 5.4 | 3.8 | 0.2 | 0.1 | 0.4 | 0.9 | 2.0 | 3.5 | 6.8 | 15.0 | 31.1 | 59.1 |
| 2002..... | 5.3 | 5.0 | 0.2 | 0.1 | 0.3 | 0.9 | 1.8 | 3.5 | 7.1 | 15.4 | 30.4 | 57.3 |
| 2001..... | 5.2 | 4.3 | * | 0.1 | 0.3 | 1.0 | 1.8 | 3.5 | 7.1 | 15.0 | 29.1 | 58.3 |
| 2000..... | 5.1 | 5.0 | 0.2 | 0.1 | 0.4 | 0.9 | 1.8 | 3.6 | 6.9 | 14.9 | 28.9 | 58.4 |
| 1999..... | 5.2 | 5.8 | * | 0.1 | 0.4 | 0.9 | 1.9 | 3.5 | 6.9 | 15.2 | 30.4 | 60.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.0 | 4.6 | * | * | 0.2 | 0.7 | 1.5 | 3.3 | 6.1 | 12.9 | 26.0 | 49.6 |
| 2003..... | 4.4 | 4.4 | * | * | 0.3 | 0.7 | 1.7 | 3.5 | 6.4 | 14.8 | 30.6 | 55.9 |
| 2002..... | 4.2 | 5.4 | * | * | 0.2 | 0.8 | 1.6 | 3.3 | 6.7 | 14.5 | 28.4 | 53.5 |
| 2001..... | 4.2 | 4.0 | * | * | 0.3 | 0.7 | 1.6 | 3.1 | 6.7 | 14.9 | 28.3 | 55.8 |
| 2000..... | 4.2 | 5.3 | * | * | 0.2 | 0.8 | 1.5 | 3.4 | 6.7 | 14.3 | 28.9 | 56.3 |
| 1999..... | 4.3 | 6.0 | * | * | 0.3 | 0.7 | 1.6 | 3.2 | 7.0 | 15.5 | 29.6 | 60.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.9 | 3.8 | * | * | 0.4 | 1.1 | 2.0 | 3.5 | 5.9 | 14.8 | 29.8 | 58.6 |
| 2003..... | 6.2 | 3.1 | * | * | 0.5 | 1.2 | 2.4 | 3.5 | 7.1 | 15.1 | 31.5 | 60.5 |
| 2002..... | 6.3 | 4.5 | * | * | 0.4 | 1.1 | 2.0 | 3.7 | 7.5 | 16.2 | 31.7 | 58.9 |
| 2001..... | 6.1 | 4.7 | * | * | 0.4 | 1.2 | 2.1 | 3.9 | 7.6 | 15.2 | 29.6 | 59.3 |
| 2000..... | 6.0 | 4.7 | * | * | 0.5 | 1.1 | 2.1 | 3.8 | 7.2 | 15.4 | 28.9 | 59.2 |
| 1999..... | 6.0 | 5.5 | * | * | 0.6 | 1.1 | 2.1 | 3.7 | 6.8 | 15.0 | 31.0 | 59.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.1 | 3.6 | * | * | 0.3 | 0.7 | 1.5 | 3.0 | 5.5 | 13.1 | 27.4 | 56.4 |
| 2003..... | 5.5 | 2.9 | * | * | 0.3 | 0.8 | 1.6 | 3.2 | 6.2 | 13.8 | 30.8 | 58.9 |
| 2002..... | 5.4 | 4.1 | * | 0.1 | 0.2 | 0.7 | 1.6 | 3.1 | 6.6 | 14.7 | 29.7 | 57.1 |
| 2001..... | 5.3 | 3.5 | * | * | 0.3 | 0.8 | 1.5 | 3.2 | 6.6 | 14.3 | 28.4 | 57.6 |
| 2000..... | 5.2 | 4.0 | * | * | 0.3 | 0.7 | 1.5 | 3.2 | 6.5 | 14.3 | 28.2 | 57.4 |
| 1999..... | 5.3 | 4.9 | * | 0.1 | 0.4 | 0.7 | 1.5 | 3.0 | 6.1 | 14.1 | 29.4 | 59.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.1 | 4.0 | * | * | 0.2 | 0.5 | 1.3 | 3.0 | 5.6 | 12.4 | 24.6 | 49.3 |
| 2003..... | 4.5 | 3.6 | * | * | 0.2 | 0.6 | 1.3 | 3.1 | 5.9 | 13.9 | 30.5 | 55.6 |
| 2002..... | 4.4 | 4.6 | * | * | 0.2 | 0.6 | 1.4 | 3.1 | 6.4 | 14.0 | 27.9 | 53.4 |
| 2001..... | 4.3 | 3.3 | * | * | 0.2 | 0.6 | 1.4 | 2.9 | 6.1 | 14.4 | 27.5 | 53.8 |
| 2000..... | 4.3 | 4.4 | * | * | 0.2 | 0.7 | 1.3 | 3.1 | 6.3 | 14.1 | 28.4 | 54.4 |
| 1999..... | 4.4 | 5.5 | * | * | 0.2 | 0.5 | 1.4 | 2.9 | 6.4 | 14.5 | 28.5 | 60.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.1 | 3.2 | * | * | 0.4 | 0.9 | 1.7 | 3.0 | 5.4 | 13.7 | 29.2 | 59.5 |
| 2003..... | 6.4 | 2.1 | * | * | 0.4 | 1.0 | 1.9 | 3.2 | 6.4 | 13.6 | 31.0 | 60.4 |
| 2002..... | 6.4 | 3.6 | * | * | 0.3 | 0.7 | 1.7 | 3.2 | 6.8 | 15.3 | 30.9 | 58.7 |
| 2001..... | 6.3 | 3.8 | * | * | 0.3 | 0.9 | 1.7 | 3.4 | 7.0 | 14.3 | 29.0 | 59.2 |
| 2000..... | 6.1 | 3.6 | * | * | 0.4 | 0.8 | 1.7 | 3.3 | 6.7 | 14.5 | 28.2 | 58.6 |
| 1999..... | 6.1 | 4.4 | * | * | 0.5 | 0.8 | 1.6 | 3.2 | 5.9 | 13.8 | 29.9 | 59.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pulmonary heart disease and diseases of pulmonary circulation (I26-I28) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.8 | 6.2 | * | * | 0.7 | 2.0 | 4.0 | 6.8 | 11.8 | 25.2 | 45.3 | 62.2 |
| 2003..... | 6.3 | 6.5 | * | * | 0.8 | 2.0 | 4.7 | 6.9 | 13.4 | 29.9 | 42.9 | 73.2 |
| 2002..... | 6.1 | 8.5 | * | * | 0.6 | 2.5 | 3.6 | 7.0 | 14.0 | 26.4 | 46.5 | 69.6 |
| 2001..... | 6.0 | 7.0 | * | * | 0.6 | 2.2 | 4.0 | 7.0 | 13.8 | 26.1 | 42.6 | 78.0 |
| 2000..... | 5.9 | 8.1 | * | * | 0.8 | 2.2 | 4.1 | 7.6 | 12.2 | 25.1 | 42.9 | 77.6 |
| 1999..... | 6.4 | 10.0 | * | * | 0.7 | 2.3 | 4.1 | 7.4 | 15.0 | 29.2 | 49.7 | 72.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.8 | 6.8 | * | * | * | 1.6 | 3.4 | 6.0 | 11.9 | 21.4 | 50.5 | 64.0 |
| 2003..... | 5.4 | 7.4 | * | * | 0.7 | 1.6 | 4.2 | 7.0 | 13.1 | 28.6 | 43.0 | 77.8 |
| 2002..... | 4.7 | 9.4 | * | * | * | 2.1 | 3.0 | 5.7 | 12.0 | 23.4 | 43.1 | 66.7 |
| 2001..... | 4.9 | 7.3 | * | * | * | 1.6 | 3.4 | 5.5 | 13.8 | 24.6 | 44.7 | 94.7 |
| 2000..... | 4.8 | 7.8 | * | * | * | 1.7 | 3.3 | 6.6 | 12.2 | 21.6 | 43.8 | 93.9 |
| 1999..... | 5.3 | 9.4 | * | * | * | 2.0 | 3.3 | 6.3 | 14.4 | 30.5 | 50.7 | 71.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | * | 0.8 | 2.3 | 4.5 | 7.5 | 11.7 | 27.9 | 42.3 | 61.4 |
| 2003..... | 7.1 | * | * | * | 0.8 | 2.3 | 5.1 | 6.7 | 13.7 | 30.8 | 42.9 | 71.4 |
| 2002..... | 7.4 | 7.6 | * | * | 0.8 | 2.8 | 4.2 | 8.0 | 15.6 | 28.6 | 48.4 | 70.7 |
| 2001..... | 7.0 | 6.7 | * | * | * | 2.8 | 4.5 | 8.2 | 13.8 | 27.2 | 41.4 | 71.9 |
| 2000..... | 7.0 | 8.5 | * | * | 1.2 | 2.5 | 4.8 | 8.3 | 12.3 | 27.6 | 42.4 | 71.5 |
| 1999..... | 7.4 | 10.7 | * | * | 0.8 | 2.7 | 4.9 | 8.2 | 15.5 | 28.3 | 49.2 | 72.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pulmonary embolism (I26) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 3.6 | 8.2 | 17.1 | 33.4 |
| 2003..... | 3.0 | * | * | * | 0.2 | 0.6 | 1.3 | 2.3 | 4.2 | 8.7 | 18.9 | 35.5 |
| 2002..... | 3.0 | * | * | * | 0.2 | 0.5 | 1.2 | 2.2 | 4.5 | 9.5 | 18.5 | 35.4 |
| 2001..... | 3.0 | * | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 4.7 | 9.5 | 18.4 | 37.8 |
| 2000..... | 3.0 | * | * | * | 0.2 | 0.5 | 1.2 | 2.3 | 4.5 | 9.3 | 18.4 | 38.4 |
| 1999..... | 3.2 | * | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 4.7 | 10.1 | 21.1 | 42.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | 0.1 | 0.5 | 1.0 | 2.1 | 4.0 | 8.4 | 16.8 | 30.6 |
| 2003..... | 2.6 | * | * | * | 0.2 | 0.4 | 1.1 | 2.4 | 4.5 | 9.4 | 19.6 | 34.1 |
| 2002..... | 2.5 | * | * | * | 0.1 | 0.5 | 1.0 | 2.2 | 4.5 | 9.4 | 17.5 | 34.1 |
| 2001..... | 2.6 | * | * | * | 0.1 | 0.5 | 1.1 | 2.1 | 4.5 | 10.0 | 19.0 | 38.0 |
| 2000..... | 2.6 | * | * | * | 0.1 | 0.5 | 1.0 | 2.4 | 4.6 | 9.5 | 19.3 | 37.5 |
| 1999..... | 2.7 | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 5.0 | 10.9 | 20.4 | 42.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | 0.2 | 0.6 | 1.3 | 2.2 | 3.3 | 8.0 | 17.3 | 34.7 |
| 2003..... | 3.4 | * | * | * | 0.3 | 0.7 | 1.4 | 2.2 | 4.0 | 8.2 | 18.4 | 36.0 |
| 2002..... | 3.5 | * | * | * | 0.2 | 0.6 | 1.3 | 2.2 | 4.5 | 9.5 | 19.2 | 36.0 |
| 2001..... | 3.5 | * | * | * | 0.2 | 0.7 | 1.3 | 2.5 | 4.9 | 9.1 | 18.0 | 37.7 |
| 2000..... | 3.4 | * | * | * | 0.3 | 0.6 | 1.3 | 2.2 | 4.4 | 9.1 | 17.8 | 38.8 |
| 1999..... | 3.7 | * | * | * | 0.3 | 0.7 | 1.3 | 2.4 | 4.4 | 9.6 | 21.5 | 42.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.5 | 0.7 | 1.6 | 3.5 | 8.0 | 20.5 |
| 2003..... | 0.6 | * | * | * | * | * | 0.4 | 0.7 | 1.8 | 3.0 | 8.7 | 18.7 |
| 2002..... | 0.7 | * | * | * | * | * | 0.4 | 0.8 | 1.7 | 4.6 | 9.8 | 21.1 |
| 2001..... | 0.7 | * | * | * | * | * | 0.5 | 0.6 | 2.2 | 4.4 | 9.2 | 22.5 |
| 2000..... | 0.6 | * | * | * | * | * | * | 1.0 | 1.7 | 3.7 | 7.7 | 14.6 |
| 1999..... | 0.7 | * | * | * | * | * | * | 1.0 | 2.3 | 5.0 | 10.2 | 28.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 1.9 | 3.5 | * | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 2.1 | * | 10.4 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 5.9 | * | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 5.4 | 10.7 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 4.2 | * | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 4.8 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 3.4 | 9.9 | 23.0 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | 2.9 | 7.6 | 21.1 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | 3.5 | 12.1 | 23.2 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 2.5 | 3.6 | 8.2 | 19.5 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 3.3 | 7.9 | * |
| 1999..... | 0.9 | * | * | * | * | * | * | * | 2.3 | 5.2 | 11.1 | 26.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pulmonary embolism (I26) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 3.8 | 8.5 | 17.6 | 34.0 |
| 2003..... | 3.4 | * | * | * | 0.3 | 0.6 | 1.4 | 2.4 | 4.4 | 9.1 | 19.3 | 36.1 |
| 2002..... | 3.3 | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 4.7 | 9.8 | 18.9 | 36.0 |
| 2001..... | 3.4 | * | * | * | 0.2 | 0.7 | 1.2 | 2.4 | 4.9 | 9.8 | 18.8 | 38.3 |
| 2000..... | 3.3 | * | * | * | 0.2 | 0.6 | 1.3 | 2.5 | 4.7 | 9.6 | 18.8 | 39.2 |
| 1999..... | 3.6 | * | * | * | 0.3 | 0.6 | 1.3 | 2.4 | 4.9 | 10.4 | 21.4 | 42.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 0.2 | 0.5 | 1.1 | 2.3 | 4.2 | 8.7 | 17.4 | 31.4 |
| 2003..... | 3.0 | * | * | * | 0.2 | 0.5 | 1.2 | 2.5 | 4.6 | 9.8 | 20.0 | 35.1 |
| 2002..... | 2.8 | * | * | * | 0.1 | 0.5 | 1.1 | 2.3 | 4.8 | 9.6 | 18.0 | 34.9 |
| 2001..... | 2.8 | * | * | * | 0.1 | 0.5 | 1.1 | 2.2 | 4.7 | 10.2 | 19.3 | 38.2 |
| 2000..... | 2.9 | * | * | * | 0.1 | 0.6 | 1.1 | 2.6 | 4.8 | 9.8 | 19.8 | 38.6 |
| 1999..... | 3.0 | * | * | * | 0.2 | 0.5 | 1.1 | 2.3 | 5.2 | 11.2 | 20.8 | 42.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | 0.3 | 0.7 | 1.4 | 2.4 | 3.4 | 8.3 | 17.7 | 35.1 |
| 2003..... | 3.8 | * | * | * | 0.3 | 0.8 | 1.6 | 2.3 | 4.2 | 8.5 | 18.9 | 36.6 |
| 2002..... | 3.9 | * | * | * | 0.3 | 0.7 | 1.4 | 2.4 | 4.7 | 9.9 | 19.5 | 36.4 |
| 2001..... | 3.9 | * | * | * | 0.2 | 0.8 | 1.3 | 2.6 | 5.0 | 9.5 | 18.4 | 38.4 |
| 2000..... | 3.8 | * | * | * | 0.3 | 0.7 | 1.4 | 2.3 | 4.6 | 9.5 | 18.2 | 39.5 |
| 1999..... | 4.1 | * | * | * | 0.4 | 0.7 | 1.4 | 2.5 | 4.5 | 9.8 | 21.8 | 42.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | 0.2 | 0.5 | 1.0 | 2.1 | 3.6 | 7.9 | 16.9 | 33.9 |
| 2003..... | 3.4 | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 4.1 | 8.3 | 19.1 | 35.5 |
| 2002..... | 3.4 | * | * | * | 0.2 | 0.5 | 1.2 | 2.2 | 4.4 | 9.3 | 18.3 | 35.3 |
| 2001..... | 3.5 | * | * | * | 0.2 | 0.5 | 1.0 | 2.2 | 4.6 | 9.4 | 18.3 | 37.5 |
| 2000..... | 3.4 | * | * | * | 0.2 | 0.5 | 1.0 | 2.2 | 4.4 | 9.2 | 18.2 | 38.2 |
| 1999..... | 3.6 | * | * | * | 0.2 | 0.5 | 1.1 | 2.1 | 4.3 | 9.7 | 20.6 | 42.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | 0.4 | 0.9 | 2.1 | 3.9 | 8.3 | 16.2 | 30.8 |
| 2003..... | 3.0 | * | * | * | * | 0.4 | 1.0 | 2.3 | 4.2 | 9.2 | 19.6 | 34.1 |
| 2002..... | 2.9 | * | * | * | * | 0.4 | 1.1 | 2.2 | 4.5 | 9.3 | 17.5 | 34.1 |
| 2001..... | 2.9 | * | * | * | * | 0.4 | 1.0 | 2.1 | 4.3 | 9.9 | 18.6 | 36.3 |
| 2000..... | 2.9 | * | * | * | * | 0.5 | 0.9 | 2.3 | 4.5 | 9.6 | 19.1 | 36.9 |
| 1999..... | 3.0 | * | * | * | * | 0.4 | 1.1 | 2.1 | 4.7 | 10.4 | 19.7 | 41.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | 0.2 | 0.6 | 1.2 | 2.1 | 3.3 | 7.6 | 17.4 | 35.4 |
| 2003..... | 3.9 | * | * | * | 0.2 | 0.6 | 1.3 | 2.2 | 3.9 | 7.6 | 18.7 | 36.1 |
| 2002..... | 4.0 | * | * | * | 0.2 | 0.5 | 1.2 | 2.1 | 4.3 | 9.3 | 18.8 | 35.9 |
| 2001..... | 4.0 | * | * | * | 0.2 | 0.7 | 1.1 | 2.3 | 4.8 | 9.0 | 18.0 | 38.0 |
| 2000..... | 3.9 | * | * | * | 0.2 | 0.6 | 1.2 | 2.0 | 4.4 | 8.8 | 17.6 | 38.8 |
| 1999..... | 4.1 | * | * | * | 0.3 | 0.5 | 1.1 | 2.2 | 4.0 | 9.0 | 21.2 | 42.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pulmonary embolism (I26) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | 0.4 | 1.2 | 2.8 | 4.7 | 7.0 | 16.1 | 29.3 | 42.9 |
| 2003..... | 4.1 | * | * | * | 0.6 | 1.5 | 3.2 | 4.7 | 8.5 | 19.3 | 28.1 | 51.8 |
| 2002..... | 3.9 | * | * | * | 0.4 | 1.6 | 2.3 | 4.6 | 9.0 | 17.0 | 30.8 | 49.9 |
| 2001..... | 4.0 | * | * | * | * | 1.7 | 2.8 | 4.8 | 8.9 | 16.7 | 28.9 | 55.9 |
| 2000..... | 4.0 | * | * | * | 0.6 | 1.3 | 2.9 | 5.3 | 8.2 | 16.8 | 30.9 | 57.2 |
| 1999..... | 4.4 | * | * | * | 0.5 | 1.7 | 2.6 | 5.2 | 10.5 | 20.9 | 36.6 | 53.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 1.3 | 2.5 | 4.2 | 8.1 | 15.4 | 37.6 | 45.7 |
| 2003..... | 3.8 | * | * | * | * | 1.3 | 2.9 | 5.2 | 9.6 | 19.7 | 32.9 | 58.6 |
| 2002..... | 3.2 | * | * | * | * | 1.6 | 1.9 | 4.4 | 8.8 | 15.5 | 29.0 | 52.5 |
| 2001..... | 3.4 | * | * | * | * | 1.4 | 2.6 | 3.8 | 10.1 | 16.2 | 33.3 | 71.6 |
| 2000..... | 3.4 | * | * | * | * | 1.0 | 2.6 | 5.4 | 9.3 | 14.8 | 33.9 | 72.8 |
| 1999..... | 3.9 | * | * | * | * | 1.6 | 2.1 | 4.8 | 11.0 | 23.6 | 41.0 | 60.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 1.2 | 3.1 | 5.1 | 6.2 | 16.7 | 24.6 | 41.8 |
| 2003..... | 4.4 | * | * | * | 0.7 | 1.7 | 3.5 | 4.3 | 7.6 | 19.1 | 25.4 | 49.1 |
| 2002..... | 4.6 | * | * | * | * | 1.6 | 2.7 | 4.8 | 9.1 | 18.0 | 31.8 | 48.9 |
| 2001..... | 4.4 | * | * | * | * | 1.9 | 3.1 | 5.6 | 7.9 | 17.1 | 26.4 | 50.0 |
| 2000..... | 4.5 | * | * | * | 0.9 | 1.6 | 3.1 | 5.1 | 7.4 | 18.1 | 29.2 | 51.5 |
| 1999..... | 4.9 | * | * | * | * | 1.9 | 3.2 | 5.5 | 10.0 | 19.0 | 34.1 | 50.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | 3.7 | 0.1 | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.1 | 5.3 | 10.4 | 21.4 |
| 2003..... | 1.8 | 3.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 1.0 | 2.3 | 5.6 | 11.6 | 22.6 |
| 2002..... | 1.8 | 4.6 | 0.2 | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 2.4 | 5.5 | 11.3 | 20.9 |
| 2001..... | 1.6 | 4.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 2.2 | 5.1 | 10.1 | 19.6 |
| 2000..... | 1.6 | 4.7 | 0.2 | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 2.2 | 5.2 | 9.9 | 19.0 |
| 1999..... | 1.5 | 5.7 | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.0 | 2.0 | 4.6 | 8.9 | 17.5 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.2 | 4.0 | * | * | * | 0.2 | 0.5 | 1.0 | 1.8 | 4.1 | 8.4 | 17.7 |
| 2003..... | 1.3 | 3.6 | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 4.8 | 10.3 | 20.3 |
| 2002..... | 1.3 | 4.7 | * | * | * | 0.2 | 0.4 | 0.9 | 1.9 | 4.7 | 10.3 | 18.0 |
| 2001..... | 1.2 | 3.9 | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 | 4.5 | 8.9 | 17.6 |
| 2000..... | 1.2 | 4.9 | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 4.4 | 9.0 | 17.3 |
| 1999..... | 1.2 | 5.9 | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 1.7 | 4.2 | 8.7 | 18.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.2 | 3.5 | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.4 | 6.3 | 11.8 | 23.1 |
| 2003..... | 2.3 | 2.8 | * | * | 0.2 | 0.4 | 0.8 | 1.2 | 2.8 | 6.4 | 12.4 | 23.6 |
| 2002..... | 2.2 | 4.4 | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 2.8 | 6.1 | 12.0 | 22.1 |
| 2001..... | 2.0 | 4.5 | * | * | 0.1 | 0.4 | 0.7 | 1.3 | 2.5 | 5.5 | 11.0 | 20.5 |
| 2000..... | 2.1 | 4.5 | * | * | 0.2 | 0.4 | 0.7 | 1.4 | 2.5 | 5.8 | 10.6 | 19.7 |
| 1999..... | 1.8 | 5.5 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.2 | 5.0 | 9.0 | 17.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | 2.2 | * | * | * | * | 0.5 | 1.2 | 1.2 | 3.4 | 4.4 | 11.4 |
| 2003..... | 0.6 | * | * | * | * | 0.3 | 0.5 | 0.6 | 1.6 | 2.9 | 7.2 | 10.9 |
| 2002..... | 0.6 | 3.0 | * | * | * | * | 0.5 | 0.9 | 1.4 | 2.9 | 7.0 | * |
| 2001..... | 0.6 | 3.7 | * | * | * | 0.4 | 0.4 | 0.8 | 1.4 | 2.5 | 5.4 | * |
| 2000..... | 0.7 | 3.9 | * | * | * | 0.3 | 0.7 | 1.0 | 1.3 | 2.5 | 6.7 | 13.3 |
| 1999..... | 0.6 | 5.3 | * | * | * | * | 0.7 | * | * | 2.1 | 6.0 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 1.1 | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | 5.2 | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 1.3 | * | 4.3 | 4.9 | * |
| 2003..... | 0.9 | * | * | * | * | 0.6 | * | * | 2.2 | 4.4 | 9.0 | * |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.3 | * | 3.0 | 8.2 | * |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | * | 6.0 | * |
| 2000..... | 0.9 | * | * | * | * | * | 1.0 | * | * | 3.3 | 7.9 | * |
| 1999..... | 0.8 | 5.4 | * | * | * | * | 0.9 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | 4.1 | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.2 | 5.4 | 10.7 | 21.9 |
| 2003..... | 2.0 | 3.7 | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 2.4 | 5.8 | 11.8 | 23.0 |
| 2002..... | 1.9 | 4.9 | 0.2 | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 2.4 | 5.6 | 11.5 | 21.3 |
| 2001..... | 1.8 | 4.3 | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 2.3 | 5.2 | 10.3 | 20.0 |
| 2000..... | 1.8 | 4.9 | 0.2 | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.2 | 5.3 | 10.1 | 19.1 |
| 1999..... | 1.6 | 5.8 | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.1 | 2.1 | 4.8 | 9.0 | 17.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | 4.5 | * | * | * | 0.1 | 0.4 | 1.0 | 1.9 | 4.2 | 8.6 | 18.2 |
| 2003..... | 1.5 | 4.3 | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 5.0 | 10.6 | 20.9 |
| 2002..... | 1.5 | 5.3 | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.8 | 10.5 | 18.6 |
| 2001..... | 1.4 | 4.0 | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 | 4.7 | 9.0 | 17.6 |
| 2000..... | 1.3 | 5.2 | * | * | * | 0.2 | 0.5 | 0.9 | 1.9 | 4.5 | 9.1 | 17.7 |
| 1999..... | 1.3 | 6.0 | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 4.3 | 8.8 | 18.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.4 | 3.7 | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 2.5 | 6.4 | 12.1 | 23.5 |
| 2003..... | 2.5 | 3.0 | * | * | 0.2 | 0.4 | 0.8 | 1.2 | 2.9 | 6.5 | 12.6 | 23.9 |
| 2002..... | 2.4 | 4.5 | * | * | 0.1 | 0.4 | 0.6 | 1.3 | 2.8 | 6.3 | 12.2 | 22.5 |
| 2001..... | 2.2 | 4.5 | * | * | 0.1 | 0.4 | 0.7 | 1.3 | 2.5 | 5.6 | 11.2 | 21.0 |
| 2000..... | 2.2 | 4.6 | * | * | 0.2 | 0.4 | 0.7 | 1.4 | 2.6 | 6.0 | 10.7 | 19.7 |
| 1999..... | 2.0 | 5.5 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.3 | 5.2 | 9.2 | 17.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | 3.5 | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.9 | 5.2 | 10.4 | 22.4 |
| 2003..... | 2.1 | 2.8 | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 | 5.4 | 11.7 | 23.5 |
| 2002..... | 2.0 | 4.1 | * | * | * | 0.2 | 0.4 | 1.0 | 2.2 | 5.4 | 11.4 | 21.8 |
| 2001..... | 1.8 | 3.5 | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.0 | 4.9 | 10.2 | 20.1 |
| 2000..... | 1.8 | 4.0 | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 | 5.1 | 10.1 | 19.1 |
| 1999..... | 1.6 | 4.9 | * | 0.1 | 0.2 | 0.2 | 0.5 | 0.9 | 1.8 | 4.5 | 8.8 | 17.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | 3.8 | * | * | * | * | 0.4 | 0.9 | 1.7 | 4.2 | 8.4 | 18.6 |
| 2003..... | 1.6 | 3.6 | * | * | * | * | 0.4 | 0.9 | 1.7 | 4.8 | 10.8 | 21.4 |
| 2002..... | 1.5 | 4.5 | * | * | * | 0.2 | 0.3 | 0.9 | 1.9 | 4.7 | 10.4 | 19.3 |
| 2001..... | 1.4 | 3.3 | * | * | * | 0.2 | 0.4 | 0.8 | 1.9 | 4.4 | 8.9 | 17.5 |
| 2000..... | 1.4 | 4.4 | * | * | * | * | 0.4 | 0.8 | 1.8 | 4.5 | 9.3 | 17.5 |
| 1999..... | 1.3 | 5.5 | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 4.1 | 8.8 | 19.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.5 | 3.2 | * | * | * | 0.2 | 0.5 | 1.0 | 2.1 | 6.0 | 11.8 | 24.2 |
| 2003..... | 2.6 | 2.1 | * | * | 0.2 | 0.4 | 0.6 | 1.0 | 2.5 | 6.0 | 12.3 | 24.4 |
| 2002..... | 2.5 | 3.6 | * | * | * | 0.2 | 0.5 | 1.1 | 2.5 | 6.0 | 12.1 | 22.9 |
| 2001..... | 2.3 | 3.6 | * | * | * | 0.3 | 0.6 | 1.1 | 2.1 | 5.3 | 11.0 | 21.2 |
| 2000..... | 2.2 | 3.6 | * | * | * | 0.3 | 0.5 | 1.2 | 2.3 | 5.7 | 10.6 | 19.8 |
| 1999..... | 1.9 | 4.4 | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 1.9 | 4.8 | 8.7 | 17.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | 6.0 | * | * | 0.3 | 0.8 | 1.2 | 2.1 | 4.8 | 9.1 | 16.0 | 19.3 |
| 2003..... | 2.2 | 6.2 | * | * | * | 0.5 | 1.4 | 2.2 | 4.9 | 10.6 | 14.8 | 21.4 |
| 2002..... | 2.2 | 8.5 | * | * | * | 0.9 | 1.3 | 2.3 | 5.0 | 9.5 | 15.7 | 19.7 |
| 2001..... | 2.1 | 7.0 | * | * | * | 0.6 | 1.1 | 2.2 | 4.9 | 9.4 | 13.7 | 22.2 |
| 2000..... | 1.9 | 7.8 | * | * | * | 0.9 | 1.2 | 2.3 | 4.0 | 8.3 | 12.0 | 20.4 |
| 1999..... | 2.0 | 9.8 | * | * | * | 0.6 | 1.5 | 2.2 | 4.6 | 8.3 | 13.1 | 18.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.5 | 6.8 | * | * | * | * | 0.9 | 1.9 | 3.7 | 6.0 | 12.8 | * |
| 2003..... | 1.6 | 7.1 | * | * | * | * | 1.3 | 1.9 | 3.4 | 9.0 | 10.1 | * |
| 2002..... | 1.5 | 9.4 | * | * | * | * | 1.1 | 1.3 | 3.2 | 7.9 | 14.1 | * |
| 2001..... | 1.5 | 7.3 | * | * | * | * | 0.8 | 1.7 | 3.7 | 8.5 | 11.4 | 23.1 |
| 2000..... | 1.3 | 7.5 | * | * | * | * | 0.8 | 1.2 | 2.8 | 6.7 | 9.9 | * |
| 1999..... | 1.4 | 9.4 | * | * | * | * | 1.2 | 1.5 | 3.4 | 6.9 | 9.7 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | 1.1 | 1.5 | 2.4 | 5.6 | 11.3 | 17.7 | 19.7 |
| 2003..... | 2.7 | * | * | * | * | * | 1.6 | 2.4 | 6.1 | 11.7 | 17.4 | 22.3 |
| 2002..... | 2.9 | 7.6 | * | * | * | 1.2 | 1.5 | 3.2 | 6.5 | 10.6 | 16.6 | 21.7 |
| 2001..... | 2.6 | 6.7 | * | * | * | 0.9 | 1.5 | 2.6 | 5.9 | 10.1 | 15.0 | 21.8 |
| 2000..... | 2.5 | 8.1 | * | * | * | 1.0 | 1.7 | 3.2 | 4.9 | 9.5 | 13.2 | 20.1 |
| 1999..... | 2.5 | 10.4 | * | * | * | 0.8 | 1.7 | 2.8 | 5.5 | 9.3 | 15.1 | 21.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other non-pulmonary forms of heart disease (I30-I51) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 52.2 | 6.2 | 0.9 | 0.5 | 1.7 | 3.5 | 8.1 | 18.2 | 39.4 | 105.2 | 341.4 | 1,319.1 |
| 2003..... | 53.6 | 7.1 | 0.9 | 0.5 | 1.8 | 3.7 | 8.4 | 18.0 | 41.5 | 112.4 | 352.4 | 1,369.2 |
| 2002..... | 54.0 | 7.1 | 0.8 | 0.5 | 1.8 | 3.7 | 8.6 | 18.4 | 41.8 | 114.6 | 361.1 | 1,387.2 |
| 2001..... | 53.9 | 6.9 | 1.3 | 0.5 | 1.7 | 3.5 | 8.2 | 18.3 | 42.6 | 116.8 | 361.0 | 1,418.1 |
| 2000..... | 54.1 | 7.1 | 0.9 | 0.5 | 1.8 | 3.5 | 8.1 | 18.0 | 44.2 | 119.7 | 362.9 | 1,452.0 |
| 1999..... | 54.7 | 7.2 | 1.0 | 0.5 | 1.9 | 3.6 | 8.5 | 18.2 | 46.2 | 124.0 | 368.5 | 1,453.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 47.1 | 6.4 | 1.0 | 0.5 | 2.4 | 4.7 | 10.6 | 24.5 | 51.8 | 130.3 | 395.6 | 1,329.4 |
| 2003..... | 48.3 | 7.7 | 0.8 | 0.6 | 2.4 | 4.7 | 11.1 | 24.0 | 54.4 | 140.0 | 410.2 | 1,376.9 |
| 2002..... | 48.6 | 7.3 | 0.9 | 0.5 | 2.5 | 5.0 | 11.4 | 25.1 | 54.3 | 142.4 | 420.4 | 1,396.5 |
| 2001..... | 48.0 | 6.9 | 1.3 | 0.5 | 2.2 | 4.5 | 10.9 | 24.6 | 55.3 | 144.2 | 424.1 | 1,417.8 |
| 2000..... | 48.5 | 7.1 | 1.0 | 0.6 | 2.4 | 4.5 | 10.8 | 24.5 | 58.3 | 150.2 | 422.7 | 1,494.4 |
| 1999..... | 49.1 | 7.3 | 1.1 | 0.6 | 2.4 | 4.6 | 11.4 | 24.7 | 60.8 | 157.6 | 430.5 | 1,488.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 57.1 | 5.9 | 0.9 | 0.5 | 1.1 | 2.3 | 5.5 | 12.2 | 27.9 | 84.1 | 304.9 | 1,314.4 |
| 2003..... | 58.8 | 6.4 | 1.0 | 0.4 | 1.3 | 2.6 | 5.7 | 12.3 | 29.4 | 89.2 | 313.7 | 1,365.8 |
| 2002..... | 59.3 | 6.9 | 0.8 | 0.4 | 1.1 | 2.4 | 5.9 | 11.9 | 30.2 | 91.6 | 321.7 | 1,383.1 |
| 2001..... | 59.7 | 7.0 | 1.2 | 0.6 | 1.1 | 2.4 | 5.5 | 12.2 | 30.9 | 94.1 | 319.5 | 1,418.2 |
| 2000..... | 59.6 | 7.2 | 0.7 | 0.4 | 1.3 | 2.5 | 5.5 | 11.7 | 31.3 | 94.7 | 323.9 | 1,434.8 |
| 1999..... | 60.0 | 7.1 | 0.9 | 0.5 | 1.3 | 2.5 | 5.7 | 12.0 | 32.8 | 96.6 | 328.2 | 1,439.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 12.0 | 6.3 | 0.9 | 0.3 | 1.4 | 2.0 | 4.4 | 9.5 | 21.9 | 63.5 | 196.1 | 627.4 |
| 2003..... | 12.9 | 6.9 | 0.9 | 0.3 | 1.2 | 2.3 | 4.3 | 10.0 | 26.3 | 71.4 | 205.3 | 720.3 |
| 2002..... | 12.0 | 6.6 | 0.7 | 0.3 | 1.3 | 2.2 | 4.2 | 9.6 | 22.5 | 65.1 | 203.0 | 709.5 |
| 2001..... | 12.0 | 5.5 | 0.9 | 0.4 | 1.1 | 2.1 | 4.0 | 9.4 | 24.4 | 68.5 | 202.9 | 765.9 |
| 2000..... | 12.3 | 6.0 | 0.8 | 0.5 | 1.5 | 2.3 | 4.4 | 9.6 | 25.8 | 69.7 | 212.5 | 801.6 |
| 1999..... | 12.5 | 6.7 | 1.0 | 0.4 | 1.3 | 2.3 | 5.2 | 10.1 | 27.2 | 71.5 | 213.3 | 790.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 12.1 | 6.5 | 1.1 | * | 2.0 | 3.0 | 6.2 | 13.8 | 29.0 | 77.8 | 232.8 | 617.8 |
| 2003..... | 12.8 | 7.9 | * | * | 1.8 | 3.1 | 6.2 | 13.7 | 33.9 | 90.8 | 238.5 | 731.2 |
| 2002..... | 12.0 | 6.8 | * | * | 1.9 | 3.1 | 5.8 | 12.7 | 28.9 | 83.9 | 245.0 | 737.9 |
| 2001..... | 11.6 | 6.2 | * | * | 1.4 | 2.9 | 5.7 | 13.5 | 31.2 | 83.9 | 233.4 | 731.4 |
| 2000..... | 12.2 | 5.6 | * | 0.7 | 2.0 | 3.2 | 5.9 | 13.1 | 34.7 | 88.4 | 238.5 | 872.7 |
| 1999..... | 12.7 | 5.2 | * | * | 1.4 | 3.5 | 7.4 | 14.2 | 35.5 | 95.6 | 252.4 | 847.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 11.9 | 6.1 | * | * | 0.7 | 0.7 | 2.3 | 5.2 | 15.5 | 51.9 | 170.7 | 632.4 |
| 2003..... | 13.0 | 5.9 | * | * | 0.6 | 1.4 | 2.3 | 6.3 | 19.4 | 55.9 | 182.5 | 714.8 |
| 2002..... | 12.0 | 6.4 | * | * | 0.6 | 1.1 | 2.4 | 6.6 | 16.8 | 50.2 | 174.3 | 695.5 |
| 2001..... | 12.4 | * | * | * | 0.6 | 1.1 | 2.2 | 5.4 | 18.4 | 56.4 | 182.2 | 782.8 |
| 2000..... | 12.5 | 6.4 | * | * | 0.9 | 1.3 | 2.9 | 6.2 | 18.0 | 54.9 | 195.1 | 766.6 |
| 1999..... | 12.3 | 8.4 | * | * | 1.0 | 0.9 | 2.9 | 6.0 | 19.9 | 52.7 | 187.3 | 763.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other non-pulmonary forms of heart disease (I30-I51) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 58.6 | 6.1 | 0.9 | 0.5 | 1.8 | 3.9 | 8.6 | 19.1 | 40.7 | 108.1 | 349.0 | 1,348.6 |
| 2003..... | 60.0 | 7.1 | 0.9 | 0.5 | 2.0 | 4.0 | 9.0 | 18.8 | 42.5 | 115.0 | 359.4 | 1,394.6 |
| 2002..... | 60.4 | 7.2 | 0.9 | 0.5 | 1.9 | 4.0 | 9.2 | 19.2 | 43.2 | 117.8 | 368.0 | 1,411.9 |
| 2001..... | 60.0 | 7.2 | 1.3 | 0.6 | 1.8 | 3.8 | 8.7 | 19.0 | 43.9 | 119.7 | 367.5 | 1,439.5 |
| 2000..... | 60.0 | 7.4 | 0.9 | 0.5 | 1.9 | 3.7 | 8.5 | 18.7 | 45.5 | 122.5 | 368.6 | 1,472.5 |
| 1999..... | 60.4 | 7.3 | 1.0 | 0.5 | 2.0 | 3.8 | 8.9 | 18.9 | 47.4 | 126.8 | 374.0 | 1,473.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 53.0 | 6.3 | 0.9 | 0.5 | 2.4 | 5.1 | 11.3 | 25.6 | 53.5 | 133.8 | 404.2 | 1,363.0 |
| 2003..... | 54.1 | 7.7 | 0.8 | 0.7 | 2.5 | 5.1 | 11.8 | 24.9 | 55.8 | 143.1 | 418.3 | 1,404.6 |
| 2002..... | 54.5 | 7.4 | 0.9 | 0.6 | 2.6 | 5.4 | 12.1 | 26.2 | 56.2 | 145.9 | 428.2 | 1,423.8 |
| 2001..... | 53.6 | 7.0 | 1.5 | 0.5 | 2.3 | 4.9 | 11.6 | 25.5 | 56.9 | 147.6 | 432.3 | 1,443.2 |
| 2000..... | 53.8 | 7.4 | 1.0 | 0.6 | 2.4 | 4.8 | 11.4 | 25.5 | 59.8 | 153.5 | 430.0 | 1,517.3 |
| 1999..... | 54.3 | 7.8 | 1.1 | 0.6 | 2.6 | 4.8 | 11.8 | 25.5 | 62.3 | 160.7 | 436.8 | 1,510.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 64.0 | 5.8 | 1.0 | 0.5 | 1.2 | 2.7 | 6.0 | 12.9 | 28.9 | 86.4 | 311.9 | 1,342.2 |
| 2003..... | 65.6 | 6.6 | 1.0 | 0.4 | 1.4 | 2.8 | 6.2 | 12.9 | 30.2 | 91.4 | 320.0 | 1,390.2 |
| 2002..... | 66.0 | 7.0 | 0.8 | 0.4 | 1.1 | 2.7 | 6.4 | 12.4 | 31.2 | 94.3 | 328.1 | 1,406.8 |
| 2001..... | 66.2 | 7.5 | 1.2 | 0.6 | 1.2 | 2.7 | 6.0 | 12.8 | 31.9 | 96.4 | 324.9 | 1,438.0 |
| 2000..... | 65.9 | 7.4 | 0.8 | 0.4 | 1.3 | 2.7 | 5.8 | 12.1 | 32.2 | 97.0 | 328.6 | 1,454.4 |
| 1999..... | 66.1 | 6.7 | 0.9 | 0.5 | 1.4 | 2.8 | 6.0 | 12.5 | 33.6 | 99.0 | 333.2 | 1,459.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 64.1 | 5.1 | 0.7 | 0.4 | 1.5 | 3.1 | 7.1 | 15.9 | 36.4 | 103.4 | 349.9 | 1,385.7 |
| 2003..... | 65.5 | 5.2 | 0.7 | 0.5 | 1.6 | 3.1 | 7.5 | 15.6 | 37.9 | 109.3 | 360.6 | 1,431.2 |
| 2002..... | 65.9 | 5.9 | 0.8 | 0.4 | 1.5 | 3.1 | 7.7 | 15.9 | 38.7 | 112.9 | 368.4 | 1,446.0 |
| 2001..... | 65.2 | 5.6 | 1.0 | 0.5 | 1.4 | 3.0 | 7.0 | 15.9 | 39.3 | 114.4 | 366.4 | 1,475.6 |
| 2000..... | 65.1 | 6.5 | 0.7 | 0.4 | 1.4 | 2.9 | 6.9 | 15.5 | 40.8 | 116.8 | 367.6 | 1,506.8 |
| 1999..... | 65.2 | 5.6 | 0.8 | 0.4 | 1.6 | 3.0 | 7.2 | 15.7 | 42.1 | 121.0 | 371.7 | 1,506.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 56.8 | 5.3 | 0.6 | 0.4 | 2.1 | 4.0 | 9.6 | 21.7 | 48.0 | 128.4 | 404.3 | 1,396.0 |
| 2003..... | 58.0 | 5.1 | 0.6 | 0.6 | 2.0 | 4.1 | 10.1 | 21.1 | 50.1 | 136.6 | 420.0 | 1,443.7 |
| 2002..... | 58.3 | 6.1 | 0.9 | 0.5 | 2.2 | 4.1 | 10.4 | 22.2 | 50.7 | 140.0 | 429.2 | 1,459.5 |
| 2001..... | 57.1 | 6.1 | 1.1 | 0.5 | 1.8 | 3.8 | 9.6 | 21.9 | 51.3 | 141.5 | 429.8 | 1,476.9 |
| 2000..... | 57.3 | 6.5 | 0.9 | 0.5 | 1.8 | 3.8 | 9.4 | 21.5 | 54.3 | 146.9 | 429.2 | 1,551.9 |
| 1999..... | 57.4 | 6.1 | 0.9 | 0.5 | 2.1 | 3.9 | 9.7 | 21.6 | 55.7 | 154.0 | 434.3 | 1,534.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 71.2 | 5.0 | 0.7 | 0.4 | 1.0 | 2.1 | 4.7 | 10.2 | 25.3 | 81.9 | 312.9 | 1,381.1 |
| 2003..... | 72.6 | 5.4 | 0.8 | 0.3 | 1.2 | 2.0 | 4.9 | 10.3 | 26.3 | 86.0 | 320.5 | 1,425.7 |
| 2002..... | 73.1 | 5.7 | 0.6 | 0.4 | 0.9 | 2.1 | 5.0 | 9.7 | 27.3 | 89.8 | 327.6 | 1,440.1 |
| 2001..... | 73.1 | 5.0 | 0.9 | 0.5 | 1.0 | 2.1 | 4.4 | 10.0 | 27.9 | 91.4 | 324.3 | 1,475.0 |
| 2000..... | 72.6 | 6.4 | 0.6 | 0.3 | 1.0 | 2.1 | 4.5 | 9.6 | 28.1 | 91.5 | 327.1 | 1,488.7 |
| 1999..... | 72.7 | 4.9 | 0.7 | 0.4 | 1.1 | 2.2 | 4.6 | 9.9 | 29.4 | 93.5 | 330.7 | 1,494.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other non-pulmonary forms of heart disease (I30-I51) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 46.9 | 9.2 | 2.1 | 1.0 | 3.2 | 8.4 | 19.0 | 44.6 | 86.7 | 174.2 | 412.8 | 1,135.7 |
| 2003..... | 48.1 | 13.3 | 1.7 | 0.9 | 3.8 | 9.3 | 19.7 | 44.3 | 90.8 | 189.9 | 412.7 | 1,180.1 |
| 2002..... | 48.2 | 12.4 | 1.2 | 0.8 | 3.6 | 9.3 | 19.8 | 45.7 | 90.1 | 184.8 | 428.2 | 1,215.9 |
| 2001..... | 48.7 | 14.2 | 2.8 | 0.9 | 3.7 | 8.8 | 20.6 | 45.3 | 91.9 | 189.9 | 443.8 | 1,214.9 |
| 2000..... | 48.7 | 11.4 | 1.6 | 1.0 | 4.2 | 8.6 | 19.6 | 45.3 | 92.2 | 198.5 | 441.4 | 1,241.8 |
| 1999..... | 49.8 | 13.2 | 1.9 | 1.0 | 4.0 | 8.7 | 21.2 | 46.1 | 100.3 | 203.7 | 450.2 | 1,241.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 47.5 | 9.6 | 2.3 | 0.9 | 4.3 | 11.0 | 24.2 | 59.3 | 117.0 | 220.9 | 500.0 | 1,180.7 |
| 2003..... | 48.0 | 16.1 | 1.8 | 1.0 | 5.1 | 11.4 | 25.4 | 58.0 | 121.7 | 240.7 | 487.4 | 1,161.3 |
| 2002..... | 47.9 | 12.4 | * | 0.9 | 4.7 | 12.7 | 25.2 | 61.7 | 118.4 | 236.0 | 500.3 | 1,191.3 |
| 2001..... | 48.6 | 11.5 | 3.1 | 0.8 | 4.9 | 11.8 | 25.9 | 58.4 | 121.1 | 237.3 | 548.8 | 1,268.2 |
| 2000..... | 48.5 | 11.3 | * | 1.1 | 5.3 | 11.0 | 25.3 | 61.7 | 120.7 | 248.2 | 521.7 | 1,322.1 |
| 1999..... | 50.9 | 13.5 | 2.1 | 1.1 | 5.4 | 11.4 | 27.6 | 62.4 | 135.1 | 260.5 | 534.1 | 1,395.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 46.3 | 8.7 | 2.0 | 1.1 | 2.0 | 6.1 | 14.4 | 31.9 | 62.3 | 140.9 | 363.6 | 1,118.0 |
| 2003..... | 48.2 | 10.3 | * | 0.8 | 2.5 | 7.4 | 14.7 | 32.4 | 65.9 | 153.8 | 370.6 | 1,187.3 |
| 2002..... | 48.5 | 12.3 | * | 0.7 | 2.4 | 6.2 | 15.0 | 31.9 | 67.4 | 148.4 | 387.5 | 1,225.3 |
| 2001..... | 48.8 | 17.1 | 2.4 | 0.9 | 2.6 | 6.1 | 15.8 | 34.0 | 68.5 | 156.5 | 384.8 | 1,195.1 |
| 2000..... | 48.8 | 11.6 | * | 0.9 | 3.0 | 6.5 | 14.5 | 31.2 | 69.3 | 163.5 | 396.1 | 1,211.9 |
| 1999..... | 48.9 | 12.9 | * | 0.9 | 2.7 | 6.2 | 15.5 | 32.3 | 72.4 | 163.7 | 403.2 | 1,184.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute and subacute endocarditis (I33) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.5 | 1.3 | 2.5 | 3.2 |
| 2003..... | 0.5 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 1.3 | 2.5 | 3.0 |
| 2002..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.3 | 2.4 | 2.7 |
| 2001..... | 0.5 | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.6 | 1.4 | 2.5 | 2.6 |
| 2000..... | 0.4 | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.2 | 2.5 | 2.9 |
| 1999..... | 0.4 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.3 | 2.4 | 2.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.6 | 0.7 | 1.4 | 3.1 | 4.8 |
| 2003..... | 0.5 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 1.4 | 3.3 | 4.1 |
| 2002..... | 0.5 | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.7 | 1.6 | 3.0 | 3.6 |
| 2001..... | 0.5 | * | * | * | * | * | 0.3 | 0.7 | 0.8 | 1.6 | 3.6 | 3.9 |
| 2000..... | 0.5 | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.8 | 1.4 | 3.6 | 4.2 |
| 1999..... | 0.5 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 1.4 | 3.2 | 4.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.2 | 2.0 | 2.6 |
| 2003..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.9 | 2.5 |
| 2002..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.0 | 2.3 |
| 2001..... | 0.4 | * | * | * | * | * | 0.3 | 0.3 | 0.5 | 1.2 | 1.8 | 2.0 |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.0 | 1.8 | 2.3 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.2 | 1.8 | 2.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.2 | 2.4 | 3.3 |
| 2003..... | 0.5 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 2.5 | 3.2 |
| 2002..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.2 | 2.5 | 2.7 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 1.2 | 2.5 | 2.7 |
| 2000..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.1 | 2.4 | 2.8 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.4 | 3.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.4 | 3.1 | 4.9 |
| 2003..... | 0.5 | * | * | * | * | * | 0.3 | 0.4 | 0.8 | 1.3 | 3.4 | 4.5 |
| 2002..... | 0.5 | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 1.5 | 3.2 | 3.7 |
| 2001..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.4 | 3.6 | 4.0 |
| 2000..... | 0.5 | * | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.3 | 3.6 | 4.3 |
| 1999..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.3 | 3.3 | 5.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 1.9 | 2.5 |
| 2003..... | 0.4 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 1.9 | 2.7 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.0 | 2.3 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 1.7 | 2.1 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 1.7 | 2.2 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 1.8 | 2.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute and subacute endocarditis (I33) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 1.4 | 1.2 | 2.1 | 2.8 | * |
| 2003..... | 0.6 | * | * | * | * | * | 0.6 | 1.3 | 2.0 | 2.2 | 3.0 | * |
| 2002..... | 0.5 | * | * | * | * | * | 0.4 | 1.3 | 1.6 | 2.3 | * | * |
| 2001..... | 0.7 | * | * | * | * | * | 0.7 | 1.3 | 1.7 | 2.6 | 4.0 | * |
| 2000..... | 0.6 | * | * | * | * | * | 0.6 | 1.4 | 1.6 | 1.9 | 2.9 | * |
| 1999..... | 0.6 | * | * | * | * | * | 0.7 | 1.2 | 2.0 | 2.6 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 1.9 | * | * | * | * |
| 2003..... | 0.7 | * | * | * | * | * | * | 2.0 | 2.3 | * | * | * |
| 2002..... | 0.6 | * | * | * | * | * | * | 1.6 | 1.9 | * | * | * |
| 2001..... | 0.8 | * | * | * | * | * | 0.8 | 1.9 | 2.2 | 3.9 | * | * |
| 2000..... | 0.7 | * | * | * | * | * | 0.8 | 1.8 | * | * | * | * |
| 1999..... | 0.7 | * | * | * | * | * | 1.0 | 1.9 | 2.5 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 1.0 | * | * | * | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 1.7 | 2.1 | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | 1.0 | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | 0.7 | * | * | * | * | * |
| 2000..... | 0.6 | * | * | * | * | * | * | 1.0 | 1.6 | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 1.6 | 2.9 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of pericardium and acute myocarditis (I30-I31,I40) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 1.2 | 1.8 |
| 2003..... | 0.3 | 0.9 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 1.2 | 1.7 |
| 2002..... | 0.3 | 0.7 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 1.3 | 2.1 |
| 2001..... | 0.3 | 0.9 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.7 | 1.2 | 1.9 |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.1 | 1.8 |
| 1999..... | 0.3 | 0.8 | 0.2 | * | 0.1 | 0.1 | 0.3 | 0.3 | 0.6 | 0.8 | 1.3 | 2.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 1.3 | 2.4 |
| 2003..... | 0.3 | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.4 | 1.0 | 1.3 | 1.6 |
| 2002..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.5 | 0.9 | 1.2 | 2.3 |
| 2001..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 1.2 | 1.8 |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.4 | 0.9 | 1.4 | 2.1 |
| 1999..... | 0.3 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 0.9 | 1.4 | 2.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.2 | 1.6 |
| 2003..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.1 | 1.7 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 2.0 |
| 2001..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.6 | 1.2 | 1.9 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.0 | 1.7 |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.7 | 1.2 | 1.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.6 | 1.2 | 1.8 |
| 2003..... | 0.3 | * | * | * | 0.1 | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.2 | 1.7 |
| 2002..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.4 | 0.8 | 1.2 | 2.1 |
| 2001..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.7 | 1.2 | 1.9 |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | 1.1 | 1.8 |
| 1999..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.3 | 2.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 | 2.3 |
| 2003..... | 0.3 | * | * | * | 0.2 | * | 0.2 | 0.3 | 0.4 | 1.0 | 1.4 | 1.8 |
| 2002..... | 0.3 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.9 | 1.2 | 2.3 |
| 2001..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.2 | 2.0 |
| 2000..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.4 | 2.1 |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 0.8 | 1.4 | 2.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.6 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.1 | 1.7 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.2 | 2.1 |
| 2001..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.6 | 1.3 | 1.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.0 | 1.7 |
| 1999..... | 0.3 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 0.7 | 1.2 | 1.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of endocardium (I34-I38) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 3.6 | 12.0 | 46.7 | 200.3 |
| 2003..... | 6.9 | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.7 | 12.3 | 47.7 | 201.9 |
| 2002..... | 6.9 | * | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 3.7 | 12.0 | 49.3 | 203.9 |
| 2001..... | 6.9 | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 3.7 | 13.0 | 48.8 | 204.2 |
| 2000..... | 7.0 | * | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 4.0 | 13.2 | 49.8 | 213.2 |
| 1999..... | 7.0 | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.9 | 13.2 | 51.7 | 209.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 4.3 | 12.9 | 48.8 | 194.2 |
| 2003..... | 5.4 | * | * | * | 0.1 | 0.3 | 0.9 | 1.7 | 4.3 | 13.2 | 51.1 | 195.3 |
| 2002..... | 5.4 | * | * | * | 0.1 | 0.3 | 0.9 | 2.0 | 4.1 | 12.7 | 51.6 | 198.5 |
| 2001..... | 5.4 | * | * | * | * | 0.3 | 0.8 | 2.1 | 4.2 | 14.0 | 52.8 | 200.8 |
| 2000..... | 5.5 | * | * | * | 0.1 | 0.3 | 0.9 | 2.1 | 4.7 | 14.0 | 51.6 | 215.8 |
| 1999..... | 5.5 | * | * | * | 0.2 | 0.3 | 1.0 | 1.9 | 4.5 | 14.7 | 56.4 | 206.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 8.3 | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.1 | 11.3 | 45.3 | 203.0 |
| 2003..... | 8.3 | * | * | * | * | 0.3 | 0.6 | 1.3 | 3.1 | 11.5 | 45.4 | 204.8 |
| 2002..... | 8.4 | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.4 | 11.4 | 47.7 | 206.3 |
| 2001..... | 8.2 | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 3.2 | 12.1 | 46.2 | 205.6 |
| 2000..... | 8.5 | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 3.3 | 12.6 | 48.6 | 212.1 |
| 1999..... | 8.5 | * | * | * | 0.2 | 0.4 | 0.7 | 1.4 | 3.3 | 12.1 | 48.7 | 211.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 1.0 | 3.5 | 10.9 | 26.0 | 79.9 |
| 2003..... | 1.5 | * | * | * | * | * | 0.5 | 0.9 | 3.1 | 10.2 | 27.9 | 85.0 |
| 2002..... | 1.5 | * | * | * | * | * | 0.6 | 1.3 | 3.5 | 7.8 | 24.4 | 92.6 |
| 2001..... | 1.5 | * | * | * | * | * | * | 1.2 | 3.1 | 10.0 | 27.6 | 95.0 |
| 2000..... | 1.6 | * | * | * | * | * | 0.4 | 1.4 | 3.6 | 9.3 | 29.6 | 116.8 |
| 1999..... | 1.6 | * | * | * | * | * | 0.6 | 0.9 | 3.1 | 9.4 | 34.4 | 98.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 1.5 | 4.2 | 11.5 | 26.5 | 85.5 |
| 2003..... | 1.3 | * | * | * | * | * | 0.7 | * | 3.1 | 11.3 | 26.1 | 84.7 |
| 2002..... | 1.4 | * | * | * | * | * | 0.8 | 1.6 | 3.7 | 8.4 | 26.7 | 103.7 |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.5 | 3.9 | 10.4 | 33.1 | 81.9 |
| 2000..... | 1.4 | * | * | * | * | * | * | 1.9 | 4.4 | 9.3 | 32.0 | 120.9 |
| 1999..... | 1.5 | * | * | * | * | * | * | 1.4 | 4.6 | 10.5 | 37.0 | 110.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | * | 2.9 | 10.4 | 25.7 | 77.1 |
| 2003..... | 1.8 | * | * | * | * | * | * | * | 3.0 | 9.4 | 29.2 | 85.2 |
| 2002..... | 1.6 | * | * | * | * | * | * | * | 3.4 | 7.3 | 22.8 | 87.0 |
| 2001..... | 1.7 | * | * | * | * | * | * | * | 2.5 | 9.8 | 23.9 | 101.5 |
| 2000..... | 1.8 | * | * | * | * | * | * | * | 2.9 | 9.3 | 28.1 | 114.7 |
| 1999..... | 1.7 | * | * | * | * | * | * | * | * | 8.4 | 32.6 | 92.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of endocardium (I34-I38) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.8 | * | * | * | 0.1 | 0.3 | 0.8 | 1.8 | 3.6 | 12.1 | 47.8 | 205.5 |
| 2003..... | 7.7 | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 3.7 | 12.4 | 48.6 | 206.6 |
| 2002..... | 7.8 | * | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 3.7 | 12.2 | 50.3 | 208.2 |
| 2001..... | 7.7 | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 3.7 | 13.2 | 49.7 | 207.9 |
| 2000..... | 7.8 | * | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 4.0 | 13.5 | 50.6 | 216.2 |
| 1999..... | 7.8 | * | * | * | 0.2 | 0.4 | 0.9 | 1.7 | 4.0 | 13.4 | 52.3 | 213.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | 0.1 | 0.3 | 0.9 | 2.3 | 4.3 | 13.0 | 50.0 | 199.4 |
| 2003..... | 6.1 | * | * | * | 0.2 | 0.3 | 0.9 | 1.8 | 4.4 | 13.2 | 52.3 | 200.0 |
| 2002..... | 6.1 | * | * | * | 0.1 | 0.4 | 0.9 | 2.0 | 4.1 | 12.9 | 52.8 | 202.4 |
| 2001..... | 6.1 | * | * | * | * | 0.4 | 0.8 | 2.1 | 4.2 | 14.2 | 53.7 | 205.5 |
| 2000..... | 6.1 | * | * | * | 0.1 | 0.3 | 0.9 | 2.1 | 4.7 | 14.2 | 52.4 | 219.2 |
| 1999..... | 6.1 | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 | 4.5 | 14.9 | 57.0 | 209.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 9.3 | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.1 | 11.3 | 46.3 | 208.2 |
| 2003..... | 9.3 | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.1 | 11.7 | 46.2 | 209.5 |
| 2002..... | 9.4 | * | * | * | * | 0.3 | 0.7 | 1.3 | 3.4 | 11.7 | 48.7 | 210.7 |
| 2001..... | 9.2 | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.2 | 12.3 | 47.1 | 208.8 |
| 2000..... | 9.4 | * | * | * | * | 0.3 | 0.7 | 1.2 | 3.3 | 12.8 | 49.4 | 214.9 |
| 1999..... | 9.3 | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 3.4 | 12.2 | 49.2 | 214.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 9.0 | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.5 | 12.3 | 50.2 | 218.5 |
| 2003..... | 9.0 | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.5 | 12.6 | 51.0 | 219.8 |
| 2002..... | 9.0 | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 3.6 | 12.4 | 53.2 | 221.1 |
| 2001..... | 8.8 | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.5 | 13.3 | 52.2 | 220.7 |
| 2000..... | 8.9 | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.7 | 13.6 | 52.8 | 229.3 |
| 1999..... | 8.9 | * | * | * | 0.2 | 0.4 | 0.8 | 1.5 | 3.7 | 13.3 | 54.6 | 225.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 7.0 | * | * | * | * | 0.3 | 0.8 | 1.9 | 4.1 | 13.5 | 52.3 | 211.1 |
| 2003..... | 7.0 | * | * | * | * | 0.3 | 0.8 | 1.5 | 4.1 | 13.6 | 54.9 | 213.4 |
| 2002..... | 7.0 | * | * | * | * | 0.3 | 0.9 | 1.7 | 4.0 | 13.3 | 55.7 | 216.3 |
| 2001..... | 6.9 | * | * | * | * | 0.3 | 0.7 | 1.9 | 4.0 | 14.4 | 56.4 | 217.3 |
| 2000..... | 6.9 | * | * | * | * | 0.2 | 0.8 | 1.8 | 4.5 | 14.5 | 54.7 | 233.1 |
| 1999..... | 6.9 | * | * | * | 0.2 | 0.3 | 0.9 | 1.7 | 4.3 | 14.9 | 59.6 | 222.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 10.9 | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.9 | 11.4 | 48.9 | 221.9 |
| 2003..... | 10.8 | * | * | * | * | 0.3 | 0.5 | 1.2 | 2.9 | 11.7 | 48.4 | 222.6 |
| 2002..... | 11.0 | * | * | * | * | 0.3 | 0.5 | 1.0 | 3.2 | 11.7 | 51.5 | 223.2 |
| 2001..... | 10.7 | * | * | * | * | 0.3 | 0.5 | 1.0 | 3.0 | 12.4 | 49.5 | 222.1 |
| 2000..... | 10.9 | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.9 | 12.8 | 51.6 | 227.8 |
| 1999..... | 10.8 | * | * | * | * | 0.4 | 0.6 | 1.3 | 3.2 | 12.0 | 51.4 | 226.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of endocardium (I34-I38) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | 0.5 | 1.3 | 3.9 | 5.6 | 12.8 | 30.3 | 89.5 |
| 2003..... | 3.5 | * | * | * | * | 0.6 | 1.5 | 3.5 | 6.5 | 12.9 | 31.2 | 86.0 |
| 2002..... | 3.3 | * | * | * | * | 0.5 | 1.5 | 3.8 | 5.8 | 13.1 | 28.6 | 87.4 |
| 2001..... | 3.4 | * | * | * | * | 0.6 | 1.3 | 3.3 | 6.2 | 13.7 | 31.0 | 86.5 |
| 2000..... | 3.6 | * | * | * | * | 0.6 | 1.5 | 3.2 | 6.6 | 14.9 | 33.9 | 92.5 |
| 1999..... | 3.6 | * | * | * | * | 0.6 | 1.6 | 3.5 | 6.6 | 16.1 | 32.3 | 92.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | * | 1.6 | 5.1 | 6.7 | 11.8 | 31.1 | 86.4 |
| 2003..... | 3.0 | * | * | * | * | * | 1.8 | 4.3 | 8.1 | 12.7 | 32.3 | 71.4 |
| 2002..... | 2.7 | * | * | * | * | * | 1.4 | 4.3 | 6.7 | 12.3 | 28.7 | 67.8 |
| 2001..... | 3.0 | * | * | * | * | * | 1.5 | 4.3 | 7.5 | 14.5 | 31.2 | 87.8 |
| 2000..... | 3.1 | * | * | * | * | * | 1.6 | 4.3 | 7.2 | 14.8 | 34.9 | 85.7 |
| 1999..... | 3.1 | * | * | * | * | * | 1.7 | 4.0 | 7.4 | 17.0 | 34.1 | 77.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | * | 1.1 | 2.9 | 4.8 | 13.4 | 29.8 | 90.8 |
| 2003..... | 3.9 | * | * | * | * | 0.7 | 1.2 | 2.8 | 5.2 | 13.0 | 30.6 | 91.6 |
| 2002..... | 3.9 | * | * | * | * | * | 1.5 | 3.3 | 5.1 | 13.6 | 28.5 | 94.9 |
| 2001..... | 3.7 | * | * | * | * | * | 1.0 | 2.5 | 5.3 | 13.2 | 30.8 | 86.0 |
| 2000..... | 4.1 | * | * | * | * | * | 1.5 | 2.3 | 6.1 | 15.0 | 33.3 | 95.1 |
| 1999..... | 4.1 | * | * | * | * | * | 1.4 | 3.1 | 6.0 | 15.5 | 31.3 | 97.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nonrheumatic mitral valve disorders (I34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.8 | 6.4 | 20.7 |
| 2003..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.9 | 6.4 | 20.6 |
| 2002..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.2 | 7.2 | 22.7 |
| 2001..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.6 | 2.2 | 7.4 | 23.2 |
| 2000..... | 1.1 | * | * | * | * | 0.1 | 0.3 | 0.3 | 0.7 | 2.3 | 7.9 | 24.8 |
| 1999..... | 1.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.7 | 2.3 | 7.8 | 25.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.6 | 5.9 | 17.3 |
| 2003..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.0 | 16.9 |
| 2002..... | 0.7 | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.9 | 5.9 | 21.7 |
| 2001..... | 0.8 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 2.2 | 6.1 | 19.3 |
| 2000..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 2.0 | 7.2 | 23.4 |
| 1999..... | 0.8 | * | * | * | * | * | 0.3 | 0.3 | 0.6 | 2.2 | 7.7 | 22.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.3 | 0.6 | 2.0 | 6.8 | 22.3 |
| 2003..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.9 | 6.7 | 22.2 |
| 2002..... | 1.4 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.8 | 2.5 | 8.0 | 23.2 |
| 2001..... | 1.4 | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.6 | 2.2 | 8.2 | 24.8 |
| 2000..... | 1.4 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.8 | 2.4 | 8.3 | 25.3 |
| 1999..... | 1.5 | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 2.5 | 7.8 | 26.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.9 | 6.7 | 21.8 |
| 2003..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 2.0 | 6.7 | 21.9 |
| 2002..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.7 | 2.2 | 7.5 | 24.5 |
| 2001..... | 1.2 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.6 | 2.2 | 7.8 | 24.8 |
| 2000..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.3 | 0.7 | 2.3 | 8.3 | 26.0 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 2.3 | 8.1 | 27.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.7 | 6.0 | 18.2 |
| 2003..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 6.2 | 18.1 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.9 | 6.2 | 23.3 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 2.2 | 6.6 | 20.9 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.1 | 7.7 | 24.8 |
| 1999..... | 0.9 | * | * | * | * | * | 0.3 | 0.3 | 0.7 | 2.2 | 8.1 | 24.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.3 | 0.2 | 0.6 | 2.1 | 7.2 | 23.4 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 7.0 | 23.6 |
| 2002..... | 1.6 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.8 | 2.5 | 8.4 | 25.0 |
| 2001..... | 1.6 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.8 | 2.2 | 8.6 | 26.4 |
| 2000..... | 1.6 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.8 | 2.5 | 8.7 | 26.4 |
| 1999..... | 1.7 | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 2.5 | 8.1 | 28.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nonrheumatic mitral valve disorders (I34) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.5 | 1.0 | 2.0 | 4.4 | 10.1 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.5 | 1.1 | 1.6 | 4.3 | 7.7 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.6 | 0.9 | 2.2 | 4.0 | 7.0 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.6 | 0.9 | 2.7 | 3.6 | 7.8 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.6 | 1.0 | 2.2 | 4.4 | 12.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.6 | 1.0 | 2.5 | 4.5 | 10.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 2.2 | 4.5 | 11.2 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | * | 4.6 | 8.7 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 2.5 | 4.5 | * |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 2.8 | 4.5 | 9.8 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 2.7 | 5.0 | 13.5 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 2.4 | 5.4 | 12.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nonrheumatic aortic valve disorders (I35) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 6.1 | 28.8 | 142.5 |
| 2003..... | 4.3 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 6.4 | 29.2 | 144.9 |
| 2002..... | 4.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 6.3 | 30.2 | 144.8 |
| 2001..... | 4.3 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 6.8 | 30.0 | 144.1 |
| 2000..... | 4.4 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 7.1 | 30.9 | 151.3 |
| 1999..... | 4.4 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 6.9 | 32.4 | 147.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 7.2 | 31.1 | 141.9 |
| 2003..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 7.5 | 33.3 | 144.7 |
| 2002..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 7.4 | 34.4 | 144.9 |
| 2001..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.1 | 8.0 | 34.8 | 145.9 |
| 2000..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.5 | 8.2 | 33.3 | 158.2 |
| 1999..... | 3.5 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.3 | 8.6 | 37.2 | 150.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.2 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 5.3 | 27.2 | 142.7 |
| 2003..... | 5.1 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 5.4 | 26.5 | 145.0 |
| 2002..... | 5.2 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 5.3 | 27.4 | 144.8 |
| 2001..... | 5.1 | * | * | * | * | * | * | 0.4 | 1.3 | 5.8 | 26.9 | 143.4 |
| 2000..... | 5.3 | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 6.1 | 29.2 | 148.5 |
| 1999..... | 5.2 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 5.6 | 29.2 | 146.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | 1.6 | 5.4 | 15.3 | 59.5 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 1.5 | 5.3 | 15.5 | 63.2 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.6 | 1.5 | 4.8 | 15.5 | 69.3 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 1.6 | 4.7 | 16.6 | 70.7 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | 1.6 | 4.6 | 19.8 | 88.9 |
| 1999..... | 0.9 | * | * | * | * | * | * | * | 1.3 | 5.2 | 22.7 | 75.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 5.8 | 16.1 | 70.0 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | * | 6.1 | 15.7 | 64.7 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | 5.5 | 19.3 | 83.7 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 2.4 | 5.2 | 21.9 | 59.0 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | 2.6 | 4.6 | 24.1 | 96.7 |
| 1999..... | 0.9 | * | * | * | * | * | * | * | * | 6.6 | 27.6 | 82.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | * | 5.1 | 14.8 | 54.0 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | * | 4.7 | 15.3 | 62.5 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | 4.3 | 12.9 | 62.2 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 4.3 | 13.0 | 76.3 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | * | 4.6 | 16.8 | 85.1 |
| 1999..... | 0.9 | * | * | * | * | * | * | * | * | 4.1 | 19.4 | 72.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nonrheumatic aortic valve disorders (I35) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 6.2 | 29.5 | 146.1 |
| 2003..... | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 6.4 | 29.9 | 148.2 |
| 2002..... | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 6.3 | 30.8 | 147.7 |
| 2001..... | 4.8 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 6.9 | 30.6 | 146.6 |
| 2000..... | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 7.2 | 31.3 | 153.2 |
| 1999..... | 4.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 7.0 | 32.7 | 150.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.1 | 7.2 | 31.9 | 145.4 |
| 2003..... | 3.9 | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 7.6 | 34.1 | 148.0 |
| 2002..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 7.5 | 35.1 | 147.4 |
| 2001..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.1 | 8.1 | 35.3 | 149.3 |
| 2000..... | 3.9 | * | * | * | * | * | 0.3 | 1.0 | 2.5 | 8.5 | 33.7 | 160.3 |
| 1999..... | 3.9 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.3 | 8.7 | 37.5 | 152.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 5.3 | 27.9 | 146.4 |
| 2003..... | 5.8 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 5.4 | 27.0 | 148.3 |
| 2002..... | 5.8 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 5.3 | 28.0 | 147.8 |
| 2001..... | 5.6 | * | * | * | * | * | * | 0.4 | 1.3 | 5.9 | 27.5 | 145.4 |
| 2000..... | 5.8 | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 6.2 | 29.7 | 150.4 |
| 1999..... | 5.7 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 5.6 | 29.5 | 148.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 6.5 | 31.1 | 155.8 |
| 2003..... | 5.7 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 6.7 | 31.5 | 157.9 |
| 2002..... | 5.7 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 6.5 | 32.7 | 156.5 |
| 2001..... | 5.6 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 7.2 | 32.2 | 155.6 |
| 2000..... | 5.7 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 7.4 | 32.9 | 162.9 |
| 1999..... | 5.6 | * | * | * | * | * | 0.3 | 0.6 | 1.8 | 7.0 | 34.4 | 158.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | * | * | 0.4 | 0.9 | 2.1 | 7.6 | 33.6 | 154.1 |
| 2003..... | 4.6 | * | * | * | * | * | 0.3 | 0.6 | 2.1 | 7.9 | 36.1 | 158.2 |
| 2002..... | 4.6 | * | * | * | * | * | 0.3 | 0.8 | 2.1 | 7.9 | 37.2 | 157.4 |
| 2001..... | 4.5 | * | * | * | * | * | 0.3 | 1.0 | 2.1 | 8.4 | 37.0 | 157.3 |
| 2000..... | 4.5 | * | * | * | * | * | 0.3 | 0.9 | 2.5 | 8.6 | 35.4 | 171.0 |
| 1999..... | 4.5 | * | * | * | * | * | 0.4 | 0.8 | 2.3 | 8.8 | 39.3 | 162.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.0 | * | * | * | * | * | * | 0.4 | 1.3 | 5.5 | 29.5 | 156.5 |
| 2003..... | 6.8 | * | * | * | * | * | * | 0.4 | 1.0 | 5.6 | 28.5 | 157.7 |
| 2002..... | 6.8 | * | * | * | * | * | * | 0.3 | 1.4 | 5.4 | 29.7 | 156.2 |
| 2001..... | 6.7 | * | * | * | * | * | * | 0.4 | 1.4 | 6.1 | 29.1 | 154.9 |
| 2000..... | 6.9 | * | * | * | * | * | * | 0.3 | 1.1 | 6.3 | 31.3 | 159.7 |
| 1999..... | 6.7 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 5.6 | 31.1 | 157.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Nonrheumatic aortic valve disorders (I35) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 0.9 | 1.8 | 4.7 | 17.4 | 59.9 |
| 2003..... | 1.6 | * | * | * | * | * | 0.4 | 0.8 | 1.8 | 5.3 | 17.3 | 59.5 |
| 2002..... | 1.6 | * | * | * | * | * | 0.4 | 1.0 | 1.6 | 5.9 | 16.5 | 63.5 |
| 2001..... | 1.6 | * | * | * | * | * | * | 1.0 | 1.9 | 5.6 | 18.6 | 60.2 |
| 2000..... | 1.7 | * | * | * | * | * | 0.4 | 0.9 | 2.4 | 7.3 | 18.6 | 63.0 |
| 1999..... | 1.8 | * | * | * | * | * | 0.4 | 1.2 | 2.2 | 7.6 | 18.0 | 64.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 1.3 | 3.0 | 5.0 | 17.7 | 60.0 |
| 2003..... | 1.3 | * | * | * | * | * | * | 1.1 | 2.7 | 5.6 | 18.2 | 49.0 |
| 2002..... | 1.3 | * | * | * | * | * | * | 1.4 | 2.2 | 6.3 | 17.0 | 52.5 |
| 2001..... | 1.5 | * | * | * | * | * | * | 1.6 | 2.3 | 6.4 | 21.9 | 69.3 |
| 2000..... | 1.5 | * | * | * | * | * | * | 1.5 | 2.8 | 8.4 | 19.7 | 57.5 |
| 1999..... | 1.6 | * | * | * | * | * | * | 1.8 | 2.8 | 9.3 | 21.9 | 58.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | * | * | * | 4.6 | 17.2 | 59.8 |
| 2003..... | 1.8 | * | * | * | * | * | * | * | * | 5.1 | 16.8 | 63.6 |
| 2002..... | 1.9 | * | * | * | * | * | * | * | * | 5.6 | 16.2 | 67.7 |
| 2001..... | 1.7 | * | * | * | * | * | * | * | 1.5 | 5.1 | 16.7 | 56.9 |
| 2000..... | 2.0 | * | * | * | * | * | * | * | 2.0 | 6.5 | 17.9 | 65.0 |
| 1999..... | 1.9 | * | * | * | * | * | * | * | 1.8 | 6.5 | 15.8 | 67.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of endocardium (I36-I38) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 4.1 | 11.9 | 38.7 |
| 2003..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 4.1 | 12.3 | 37.9 |
| 2002..... | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 3.7 | 12.4 | 37.8 |
| 2001..... | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.0 | 11.7 | 38.1 |
| 2000..... | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 4.0 | 11.5 | 38.2 |
| 1999..... | 1.8 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 4.1 | 11.8 | 37.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.1 | 0.3 | 1.1 | 1.6 | 4.2 | 12.2 | 36.8 |
| 2003..... | 1.5 | * | * | * | * | * | 0.4 | 0.9 | 1.6 | 3.8 | 12.1 | 35.1 |
| 2002..... | 1.4 | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.5 | 3.5 | 11.8 | 33.3 |
| 2001..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.9 | 12.2 | 36.9 |
| 2000..... | 1.4 | * | * | * | * | * | 0.4 | 0.8 | 1.6 | 3.7 | 11.5 | 35.6 |
| 1999..... | 1.4 | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.6 | 4.0 | 11.8 | 34.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 4.0 | 11.6 | 39.5 |
| 2003..... | 2.3 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 4.3 | 12.4 | 39.1 |
| 2002..... | 2.2 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.9 | 12.7 | 39.7 |
| 2001..... | 2.1 | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 4.2 | 11.4 | 38.6 |
| 2000..... | 2.1 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 4.2 | 11.5 | 39.2 |
| 1999..... | 2.1 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 4.1 | 11.8 | 39.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 4.0 | 12.4 | 41.0 |
| 2003..... | 2.1 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.9 | 12.8 | 40.0 |
| 2002..... | 2.0 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 3.6 | 13.0 | 40.1 |
| 2001..... | 2.0 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.0 | 12.2 | 40.3 |
| 2000..... | 1.9 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.9 | 11.6 | 40.4 |
| 1999..... | 1.9 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 4.0 | 12.2 | 39.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.3 | 0.8 | 1.5 | 4.2 | 12.7 | 38.8 |
| 2003..... | 1.6 | * | * | * | * | * | 0.3 | 0.7 | 1.4 | 3.8 | 12.6 | 37.0 |
| 2002..... | 1.5 | * | * | * | * | * | 0.3 | 0.7 | 1.2 | 3.5 | 12.3 | 35.6 |
| 2001..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.8 | 12.9 | 39.1 |
| 2000..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.7 | 11.6 | 37.4 |
| 1999..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 4.0 | 12.2 | 36.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.8 | 12.2 | 42.0 |
| 2003..... | 2.6 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.1 | 13.0 | 41.3 |
| 2002..... | 2.5 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.7 | 13.5 | 42.1 |
| 2001..... | 2.4 | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 4.1 | 11.8 | 40.8 |
| 2000..... | 2.4 | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 4.1 | 11.7 | 41.6 |
| 1999..... | 2.4 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 4.0 | 12.2 | 41.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of endocardium (I36-I38) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.7 | 2.5 | 2.8 | 6.0 | 8.4 | 19.6 |
| 2003..... | 1.4 | * | * | * | * | * | 0.8 | 2.2 | 3.6 | 5.9 | 9.7 | 18.7 |
| 2002..... | 1.3 | * | * | * | * | * | 0.8 | 2.2 | 3.2 | 5.0 | 8.0 | 16.9 |
| 2001..... | 1.3 | * | * | * | * | * | 0.6 | 1.8 | 3.5 | 5.4 | 8.8 | 18.4 |
| 2000..... | 1.3 | * | * | * | * | * | 0.8 | 1.8 | 3.2 | 5.4 | 10.9 | 17.5 |
| 1999..... | 1.3 | * | * | * | * | * | 0.9 | 1.7 | 3.3 | 6.0 | 9.9 | 17.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.9 | 3.5 | 2.8 | 5.3 | 9.1 | * |
| 2003..... | 1.3 | * | * | * | * | * | 0.9 | 2.8 | 4.4 | 5.5 | 10.4 | * |
| 2002..... | 1.1 | * | * | * | * | * | 0.8 | 2.5 | 3.7 | 4.2 | 8.5 | * |
| 2001..... | 1.2 | * | * | * | * | * | * | 2.2 | 4.5 | 5.6 | 7.5 | * |
| 2000..... | 1.2 | * | * | * | * | * | 0.8 | 2.2 | 3.6 | 5.1 | 11.7 | * |
| 1999..... | 1.1 | * | * | * | * | * | 0.9 | 1.8 | 4.0 | 5.1 | 9.4 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 1.8 | 2.8 | 6.6 | 8.1 | 19.7 |
| 2003..... | 1.5 | * | * | * | * | * | * | 1.7 | 3.0 | 6.3 | 9.3 | 19.4 |
| 2002..... | 1.4 | * | * | * | * | * | 0.9 | 1.9 | 2.8 | 5.5 | 7.8 | 19.6 |
| 2001..... | 1.3 | * | * | * | * | * | * | 1.4 | 2.7 | 5.2 | 9.6 | 19.2 |
| 2000..... | 1.4 | * | * | * | * | * | 0.8 | 1.4 | 3.0 | 5.7 | 10.4 | 16.6 |
| 1999..... | 1.4 | * | * | * | * | * | 0.9 | 1.6 | 2.8 | 6.6 | 10.2 | 18.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cardiomyopathy (I42) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 8.7 | 2.8 | 0.2 | 0.2 | 0.6 | 1.4 | 3.1 | 6.1 | 10.7 | 22.8 | 54.6 | 127.6 |
| 2003..... | 9.2 | 3.1 | 0.4 | 0.1 | 0.7 | 1.5 | 3.3 | 6.3 | 11.5 | 25.5 | 57.0 | 137.8 |
| 2002..... | 9.6 | 2.9 | 0.2 | 0.2 | 0.7 | 1.6 | 3.5 | 6.5 | 11.9 | 27.0 | 61.1 | 142.0 |
| 2001..... | 9.4 | 3.4 | 0.4 | 0.2 | 0.7 | 1.4 | 3.3 | 6.3 | 12.3 | 27.2 | 60.3 | 138.3 |
| 2000..... | 9.7 | 3.1 | 0.3 | 0.2 | 0.7 | 1.5 | 3.3 | 6.7 | 13.0 | 28.7 | 61.4 | 142.3 |
| 1999..... | 9.8 | 3.5 | 0.2 | 0.2 | 0.7 | 1.5 | 3.6 | 6.7 | 14.0 | 30.4 | 60.4 | 139.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 10.3 | 2.7 | 0.3 | 0.2 | 0.9 | 2.1 | 4.4 | 8.6 | 15.1 | 31.6 | 76.2 | 174.6 |
| 2003..... | 10.8 | 3.4 | 0.3 | 0.2 | 0.9 | 2.1 | 4.7 | 8.9 | 16.1 | 35.2 | 81.0 | 178.8 |
| 2002..... | 11.3 | 2.7 | * | 0.2 | 1.0 | 2.3 | 4.8 | 9.4 | 16.7 | 37.8 | 87.2 | 189.7 |
| 2001..... | 11.0 | 2.9 | 0.5 | 0.2 | 0.9 | 2.0 | 4.8 | 8.9 | 17.3 | 37.4 | 84.9 | 184.7 |
| 2000..... | 11.4 | 2.8 | 0.3 | 0.3 | 0.9 | 2.1 | 4.8 | 9.6 | 18.2 | 40.0 | 87.6 | 195.3 |
| 1999..... | 11.5 | 3.1 | * | 0.2 | 0.9 | 2.0 | 5.2 | 9.6 | 19.6 | 43.0 | 84.3 | 188.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 7.2 | 3.0 | * | 0.2 | 0.3 | 0.7 | 1.9 | 3.6 | 6.7 | 15.4 | 40.1 | 106.4 |
| 2003..... | 7.7 | 2.9 | 0.4 | * | 0.5 | 0.9 | 1.9 | 3.7 | 7.2 | 17.4 | 40.9 | 119.7 |
| 2002..... | 8.0 | 3.1 | * | 0.1 | 0.4 | 0.9 | 2.2 | 3.6 | 7.5 | 18.1 | 43.7 | 121.3 |
| 2001..... | 8.0 | 3.9 | 0.3 | 0.2 | 0.4 | 0.8 | 1.9 | 3.9 | 7.7 | 18.8 | 44.2 | 118.9 |
| 2000..... | 8.0 | 3.4 | * | 0.1 | 0.4 | 1.0 | 1.9 | 4.0 | 8.2 | 19.4 | 44.3 | 120.7 |
| 1999..... | 8.1 | 3.8 | * | 0.1 | 0.4 | 1.0 | 2.0 | 4.0 | 8.9 | 20.0 | 44.9 | 119.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | 2.9 | * | * | 0.6 | 0.8 | 2.0 | 3.8 | 6.8 | 17.9 | 40.8 | 91.8 |
| 2003..... | 3.5 | 3.7 | * | * | 0.5 | 1.1 | 1.9 | 4.2 | 9.2 | 20.6 | 47.0 | 113.0 |
| 2002..... | 3.4 | 3.7 | * | * | 0.5 | 1.1 | 1.8 | 4.4 | 7.6 | 19.5 | 51.2 | 100.9 |
| 2001..... | 3.4 | 2.9 | * | * | 0.5 | 0.9 | 1.8 | 4.0 | 8.9 | 21.7 | 50.1 | 122.6 |
| 2000..... | 3.5 | 2.6 | * | * | 0.6 | 1.1 | 2.1 | 4.0 | 9.2 | 21.8 | 52.7 | 122.8 |
| 1999..... | 3.4 | 3.7 | * | * | 0.3 | 0.9 | 2.5 | 4.8 | 10.5 | 21.5 | 47.9 | 105.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.8 | 1.3 | 2.9 | 5.9 | 9.7 | 25.3 | 56.2 | 112.1 |
| 2003..... | 4.3 | * | * | * | 0.8 | 1.5 | 2.7 | 6.8 | 12.4 | 29.5 | 64.0 | 144.7 |
| 2002..... | 4.1 | * | * | * | 0.8 | 1.6 | 2.6 | 6.1 | 10.5 | 28.3 | 71.0 | 137.2 |
| 2001..... | 3.9 | * | * | * | 0.7 | 1.4 | 2.7 | 6.0 | 12.7 | 30.6 | 64.8 | 129.5 |
| 2000..... | 4.1 | * | * | * | 0.8 | 1.5 | 2.9 | 5.9 | 13.6 | 29.3 | 70.3 | 151.2 |
| 1999..... | 4.2 | * | * | * | * | 1.5 | 3.7 | 7.0 | 14.6 | 33.4 | 65.2 | 125.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 1.0 | 1.7 | 4.2 | 12.1 | 30.1 | 81.4 |
| 2003..... | 2.7 | * | * | * | * | * | 1.0 | 1.6 | 6.3 | 13.5 | 35.3 | 96.9 |
| 2002..... | 2.6 | * | * | * | * | * | 1.0 | 2.7 | 5.1 | 12.5 | 37.7 | 82.9 |
| 2001..... | 2.8 | * | * | * | * | * | 0.9 | 2.1 | 5.6 | 14.8 | 40.2 | 119.1 |
| 2000..... | 2.9 | * | * | * | * | * | 1.2 | 2.3 | 5.3 | 15.9 | 40.9 | 108.8 |
| 1999..... | 2.6 | 5.4 | * | * | * | * | 1.1 | 2.7 | 6.9 | 12.2 | 36.4 | 95.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cardiomyopathy (I42) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.6 | 2.8 | 0.2 | 0.2 | 0.7 | 1.6 | 3.3 | 6.3 | 11.0 | 23.1 | 55.2 | 129.0 |
| 2003..... | 10.1 | 3.0 | 0.4 | 0.1 | 0.7 | 1.6 | 3.5 | 6.5 | 11.7 | 25.8 | 57.4 | 138.6 |
| 2002..... | 10.6 | 2.7 | 0.2 | 0.2 | 0.7 | 1.7 | 3.7 | 6.7 | 12.2 | 27.5 | 61.4 | 143.5 |
| 2001..... | 10.3 | 3.5 | 0.4 | 0.2 | 0.7 | 1.5 | 3.5 | 6.5 | 12.6 | 27.5 | 60.7 | 138.6 |
| 2000..... | 10.5 | 3.2 | 0.2 | 0.2 | 0.7 | 1.6 | 3.5 | 6.9 | 13.3 | 29.0 | 61.7 | 142.7 |
| 1999..... | 10.6 | 3.4 | 0.2 | 0.2 | 0.7 | 1.6 | 3.7 | 6.9 | 14.2 | 30.8 | 60.9 | 140.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 11.4 | 2.7 | * | 0.2 | 1.0 | 2.2 | 4.7 | 8.9 | 15.5 | 32.0 | 77.1 | 177.5 |
| 2003..... | 11.9 | 3.2 | 0.4 | 0.2 | 0.9 | 2.3 | 5.0 | 9.1 | 16.4 | 35.5 | 81.8 | 180.3 |
| 2002..... | 12.5 | 2.5 | * | 0.2 | 1.0 | 2.5 | 5.1 | 9.7 | 17.2 | 38.4 | 87.8 | 192.0 |
| 2001..... | 12.0 | 2.9 | 0.5 | 0.2 | 0.9 | 2.1 | 5.1 | 9.1 | 17.6 | 37.7 | 85.7 | 186.3 |
| 2000..... | 12.4 | 2.8 | * | 0.2 | 1.0 | 2.2 | 5.0 | 9.9 | 18.5 | 40.5 | 88.2 | 196.7 |
| 1999..... | 12.5 | 3.4 | * | 0.2 | 1.0 | 2.1 | 5.3 | 9.8 | 19.8 | 43.4 | 84.9 | 190.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.9 | 2.9 | * | 0.2 | 0.3 | 0.9 | 2.0 | 3.7 | 6.9 | 15.6 | 40.5 | 107.3 |
| 2003..... | 8.5 | 2.8 | 0.4 | * | 0.5 | 1.0 | 2.1 | 3.9 | 7.3 | 17.6 | 41.1 | 120.3 |
| 2002..... | 8.7 | 2.9 | * | 0.1 | 0.4 | 1.0 | 2.4 | 3.7 | 7.7 | 18.5 | 43.9 | 122.6 |
| 2001..... | 8.7 | 4.1 | 0.3 | 0.1 | 0.4 | 0.9 | 2.0 | 4.1 | 7.9 | 19.0 | 44.3 | 118.7 |
| 2000..... | 8.7 | 3.6 | * | * | 0.4 | 1.0 | 2.0 | 4.1 | 8.4 | 19.6 | 44.4 | 120.8 |
| 1999..... | 8.8 | 3.4 | * | 0.1 | 0.4 | 1.1 | 2.1 | 4.1 | 9.0 | 20.5 | 45.3 | 120.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 9.7 | 2.6 | * | 0.2 | 0.5 | 1.1 | 2.5 | 4.9 | 9.4 | 21.1 | 54.2 | 130.7 |
| 2003..... | 10.2 | 2.8 | 0.3 | 0.1 | 0.5 | 1.1 | 2.8 | 5.1 | 9.7 | 23.7 | 56.1 | 140.7 |
| 2002..... | 10.7 | 2.5 | 0.2 | 0.2 | 0.5 | 1.2 | 2.9 | 5.3 | 10.5 | 25.7 | 60.1 | 143.2 |
| 2001..... | 10.3 | 2.7 | 0.3 | 0.1 | 0.5 | 1.1 | 2.6 | 5.1 | 10.6 | 25.6 | 58.8 | 139.3 |
| 2000..... | 10.5 | 3.0 | * | 0.1 | 0.5 | 1.1 | 2.6 | 5.3 | 11.3 | 26.6 | 59.6 | 142.6 |
| 1999..... | 10.5 | 2.8 | 0.2 | 0.1 | 0.5 | 1.2 | 2.7 | 5.3 | 11.7 | 28.5 | 58.9 | 139.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 11.4 | 2.5 | * | * | 0.8 | 1.7 | 3.6 | 7.2 | 13.1 | 29.6 | 76.4 | 181.3 |
| 2003..... | 11.9 | 2.5 | * | 0.2 | 0.7 | 1.6 | 4.0 | 7.4 | 13.8 | 32.7 | 80.7 | 184.1 |
| 2002..... | 12.5 | 2.0 | * | 0.2 | 0.7 | 1.7 | 4.0 | 7.8 | 14.9 | 35.9 | 86.9 | 193.1 |
| 2001..... | 12.0 | 2.5 | 0.4 | * | 0.7 | 1.5 | 3.8 | 7.3 | 15.0 | 35.4 | 83.7 | 188.4 |
| 2000..... | 12.3 | 2.7 | * | 0.2 | 0.6 | 1.5 | 3.7 | 7.8 | 15.9 | 37.4 | 85.7 | 198.7 |
| 1999..... | 12.2 | 2.8 | * | 0.2 | 0.7 | 1.6 | 3.9 | 7.6 | 16.4 | 40.1 | 83.0 | 189.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 8.1 | 2.7 | * | 0.2 | 0.2 | 0.6 | 1.4 | 2.7 | 5.8 | 13.7 | 39.1 | 108.1 |
| 2003..... | 8.6 | 3.1 | * | * | 0.4 | 0.6 | 1.5 | 2.9 | 5.8 | 15.9 | 39.5 | 121.5 |
| 2002..... | 8.9 | 3.0 | * | * | 0.2 | 0.6 | 1.7 | 2.8 | 6.3 | 17.1 | 42.1 | 121.7 |
| 2001..... | 8.7 | 2.8 | * | * | 0.3 | 0.6 | 1.3 | 2.9 | 6.4 | 17.3 | 42.4 | 118.8 |
| 2000..... | 8.8 | 3.4 | * | * | 0.2 | 0.7 | 1.4 | 2.9 | 7.0 | 17.6 | 42.4 | 119.9 |
| 1999..... | 8.9 | 2.9 | * | * | 0.3 | 0.8 | 1.4 | 3.0 | 7.3 | 18.7 | 43.2 | 119.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cardiomyopathy (I42) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 11.6 | * | * | * | 1.3 | 3.8 | 8.4 | 16.2 | 27.3 | 45.6 | 77.1 | 126.6 |
| 2003..... | 12.3 | 3.5 | * | * | 1.5 | 4.4 | 8.7 | 16.6 | 29.9 | 49.0 | 82.1 | 137.1 |
| 2002..... | 12.7 | 3.4 | * | * | 1.7 | 4.8 | 9.5 | 17.1 | 29.1 | 48.4 | 86.0 | 161.8 |
| 2001..... | 12.9 | 7.2 | 0.9 | 0.3 | 1.7 | 4.1 | 9.8 | 17.8 | 31.0 | 49.6 | 91.4 | 143.6 |
| 2000..... | 13.3 | 3.6 | * | 0.4 | 1.9 | 4.4 | 9.1 | 19.9 | 30.9 | 55.8 | 93.9 | 153.0 |
| 1999..... | 13.6 | 5.1 | * | 0.4 | 1.9 | 3.9 | 10.3 | 19.7 | 35.9 | 56.9 | 89.5 | 154.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 14.2 | * | * | * | 2.0 | 5.4 | 11.9 | 22.8 | 40.1 | 63.8 | 104.7 | 161.6 |
| 2003..... | 15.0 | * | * | * | 1.8 | 6.1 | 12.1 | 23.3 | 42.4 | 70.4 | 113.7 | 174.7 |
| 2002..... | 15.4 | * | * | * | 2.2 | 7.0 | 13.0 | 25.3 | 40.8 | 69.7 | 114.9 | 207.9 |
| 2001..... | 15.5 | * | * | * | 2.3 | 5.7 | 13.6 | 24.3 | 44.4 | 67.8 | 129.6 | 183.6 |
| 2000..... | 16.3 | * | * | * | 2.5 | 6.1 | 13.2 | 27.6 | 44.6 | 78.5 | 133.6 | 196.1 |
| 1999..... | 17.0 | * | * | * | 2.7 | 5.4 | 14.7 | 28.4 | 52.9 | 82.8 | 118.5 | 218.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 9.3 | * | * | * | 0.7 | 2.4 | 5.4 | 10.6 | 17.0 | 32.5 | 61.5 | 112.8 |
| 2003..... | 9.8 | * | * | * | 1.2 | 2.9 | 5.7 | 10.8 | 19.8 | 33.8 | 64.2 | 122.6 |
| 2002..... | 10.2 | * | * | * | 1.2 | 2.8 | 6.4 | 10.1 | 19.7 | 33.2 | 69.6 | 144.2 |
| 2001..... | 10.5 | 9.4 | * | * | 1.1 | 2.7 | 6.4 | 12.2 | 20.2 | 36.7 | 69.9 | 128.8 |
| 2000..... | 10.6 | * | * | * | 1.2 | 2.9 | 5.5 | 13.2 | 20.0 | 39.9 | 71.6 | 136.9 |
| 1999..... | 10.6 | * | * | * | 1.1 | 2.6 | 6.4 | 12.2 | 22.2 | 38.6 | 73.2 | 131.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Conduction disorders and cardiac dysrhythmias (I44-I49) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 12.8 | 0.8 | 0.1 | 0.1 | 0.4 | 0.8 | 1.9 | 5.0 | 11.9 | 28.8 | 84.5 | 295.3 |
| 2003..... | 13.4 | 0.8 | 0.1 | 0.1 | 0.4 | 0.8 | 1.9 | 5.0 | 12.5 | 31.3 | 90.0 | 313.7 |
| 2002..... | 13.6 | 1.4 | 0.1 | 0.1 | 0.5 | 0.8 | 2.0 | 4.9 | 12.6 | 31.5 | 93.4 | 323.2 |
| 2001..... | 13.5 | 0.9 | 0.2 | 0.1 | 0.4 | 0.7 | 1.7 | 5.0 | 12.6 | 32.6 | 92.4 | 328.2 |
| 2000..... | 13.6 | 1.0 | 0.2 | 0.1 | 0.5 | 0.7 | 1.9 | 4.8 | 13.1 | 33.3 | 92.2 | 340.2 |
| 1999..... | 14.3 | 1.0 | 0.3 | 0.1 | 0.5 | 0.8 | 1.9 | 5.2 | 14.5 | 35.7 | 97.1 | 353.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 11.5 | * | * | 0.1 | 0.5 | 0.9 | 2.3 | 6.5 | 16.2 | 36.2 | 93.8 | 278.5 |
| 2003..... | 12.0 | * | * | 0.2 | 0.5 | 0.9 | 2.4 | 6.6 | 16.6 | 40.1 | 99.9 | 292.7 |
| 2002..... | 12.2 | 1.6 | * | 0.1 | 0.6 | 0.9 | 2.5 | 6.6 | 16.8 | 39.0 | 103.8 | 310.5 |
| 2001..... | 12.0 | * | * | 0.1 | 0.6 | 0.9 | 2.2 | 6.7 | 16.5 | 39.8 | 104.9 | 311.6 |
| 2000..... | 12.1 | 1.1 | * | 0.1 | 0.5 | 0.8 | 2.3 | 6.6 | 17.5 | 42.9 | 101.9 | 333.2 |
| 1999..... | 12.9 | 1.2 | 0.3 | 0.2 | 0.6 | 1.0 | 2.3 | 7.1 | 19.2 | 45.4 | 110.8 | 353.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 14.1 | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 3.5 | 8.0 | 22.6 | 78.3 | 302.9 |
| 2003..... | 14.8 | * | * | * | 0.3 | 0.7 | 1.5 | 3.5 | 8.7 | 24.0 | 83.3 | 323.0 |
| 2002..... | 15.1 | 1.2 | * | * | 0.3 | 0.6 | 1.4 | 3.3 | 8.6 | 25.3 | 86.6 | 328.8 |
| 2001..... | 15.1 | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.4 | 9.0 | 26.7 | 84.2 | 335.2 |
| 2000..... | 15.1 | * | * | * | 0.4 | 0.6 | 1.5 | 3.0 | 8.9 | 25.4 | 85.8 | 343.0 |
| 1999..... | 15.7 | * | * | 0.1 | 0.4 | 0.6 | 1.4 | 3.4 | 10.2 | 27.8 | 88.2 | 353.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | 0.3 | 0.4 | 0.6 | 1.5 | 3.9 | 9.8 | 33.7 | 104.7 |
| 2003..... | 2.2 | * | * | * | * | 0.4 | 0.6 | 1.9 | 5.7 | 11.3 | 35.3 | 127.0 |
| 2002..... | 2.0 | * | * | * | * | 0.4 | 0.6 | 1.3 | 3.9 | 10.7 | 32.9 | 134.7 |
| 2001..... | 2.1 | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.8 | 12.6 | 36.4 | 145.1 |
| 2000..... | 2.2 | * | * | * | 0.4 | 0.4 | 0.8 | 1.7 | 4.6 | 13.1 | 39.3 | 142.7 |
| 1999..... | 2.3 | * | * | * | * | 0.3 | 0.6 | 1.5 | 5.5 | 14.0 | 40.6 | 164.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.6 | 0.9 | 1.8 | 4.4 | 12.0 | 37.6 | 95.3 |
| 2003..... | 2.1 | * | * | * | * | * | 0.9 | 2.3 | 7.0 | 14.2 | 42.6 | 118.5 |
| 2002..... | 2.0 | * | * | * | * | 0.6 | 0.8 | 1.8 | 5.0 | 13.8 | 39.0 | 130.5 |
| 2001..... | 1.9 | * | * | * | * | * | 0.8 | 1.9 | 4.8 | 14.8 | 38.9 | 137.1 |
| 2000..... | 2.2 | * | * | * | * | * | 0.9 | 2.6 | 5.6 | 18.4 | 39.8 | 155.2 |
| 1999..... | 2.3 | * | * | * | * | * | 0.8 | 2.3 | 7.2 | 17.1 | 45.9 | 198.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | * | 1.1 | 3.4 | 7.9 | 31.1 | 109.5 |
| 2003..... | 2.4 | * | * | * | * | * | * | 1.5 | 4.6 | 9.0 | 30.3 | 131.3 |
| 2002..... | 2.1 | * | * | * | * | * | * | * | 3.0 | 8.2 | 28.7 | 136.8 |
| 2001..... | 2.3 | * | * | * | * | * | * | * | 2.9 | 10.9 | 34.8 | 148.9 |
| 2000..... | 2.3 | * | * | * | * | * | * | * | 3.6 | 9.0 | 39.0 | 136.5 |
| 1999..... | 2.4 | * | * | * | * | * | * | * | 4.0 | 11.5 | 37.1 | 147.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Conduction disorders and cardiac dysrhythmias (I44-I49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 14.6 | 0.8 | * | 0.1 | 0.5 | 0.9 | 2.1 | 5.3 | 12.6 | 30.2 | 87.3 | 303.7 |
| 2003..... | 15.2 | 0.9 | * | 0.1 | 0.5 | 0.9 | 2.1 | 5.3 | 13.0 | 32.7 | 92.6 | 321.2 |
| 2002..... | 15.4 | 1.6 | * | 0.1 | 0.5 | 0.9 | 2.2 | 5.3 | 13.2 | 32.8 | 96.2 | 330.2 |
| 2001..... | 15.2 | 1.1 | 0.2 | 0.2 | 0.4 | 0.8 | 1.9 | 5.4 | 13.2 | 33.8 | 94.8 | 334.2 |
| 2000..... | 15.2 | 1.2 | 0.2 | 0.1 | 0.5 | 0.8 | 2.0 | 5.0 | 13.7 | 34.5 | 94.3 | 346.6 |
| 1999..... | 15.9 | 1.2 | 0.3 | 0.1 | 0.5 | 0.9 | 2.0 | 5.5 | 15.1 | 36.9 | 99.2 | 359.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 13.1 | * | * | 0.1 | 0.5 | 1.0 | 2.6 | 7.0 | 17.1 | 37.9 | 96.9 | 287.6 |
| 2003..... | 13.6 | * | * | 0.2 | 0.6 | 1.0 | 2.6 | 7.1 | 17.3 | 41.8 | 102.6 | 300.5 |
| 2002..... | 13.8 | 1.6 | * | 0.2 | 0.7 | 1.0 | 2.8 | 7.1 | 17.7 | 40.6 | 106.8 | 318.1 |
| 2001..... | 13.5 | * | * | 0.1 | 0.6 | 0.9 | 2.4 | 7.1 | 17.4 | 41.2 | 107.7 | 318.1 |
| 2000..... | 13.6 | 1.3 | * | 0.1 | 0.5 | 0.9 | 2.5 | 6.9 | 18.4 | 44.3 | 104.4 | 339.7 |
| 1999..... | 14.4 | 1.4 | 0.3 | 0.2 | 0.6 | 1.0 | 2.5 | 7.5 | 20.0 | 46.9 | 113.2 | 359.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 15.9 | * | * | * | 0.4 | 0.8 | 1.6 | 3.7 | 8.4 | 23.7 | 80.8 | 310.9 |
| 2003..... | 16.7 | * | * | * | 0.3 | 0.7 | 1.7 | 3.6 | 9.0 | 25.1 | 85.9 | 330.3 |
| 2002..... | 16.9 | 1.5 | * | * | 0.3 | 0.7 | 1.6 | 3.5 | 9.1 | 26.4 | 89.2 | 335.4 |
| 2001..... | 16.8 | * | * | 0.2 | 0.2 | 0.6 | 1.4 | 3.7 | 9.4 | 27.7 | 86.2 | 340.9 |
| 2000..... | 16.8 | * | * | * | 0.4 | 0.6 | 1.6 | 3.2 | 9.3 | 26.4 | 87.7 | 349.3 |
| 1999..... | 17.4 | * | * | * | 0.4 | 0.7 | 1.5 | 3.6 | 10.6 | 28.7 | 90.2 | 360.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 15.7 | * | * | 0.1 | 0.4 | 0.7 | 1.8 | 4.8 | 11.5 | 28.7 | 85.8 | 303.9 |
| 2003..... | 16.4 | * | * | 0.1 | 0.4 | 0.7 | 2.0 | 4.7 | 12.0 | 30.7 | 91.5 | 322.4 |
| 2002..... | 16.6 | 1.3 | * | 0.1 | 0.5 | 0.7 | 2.0 | 4.7 | 12.0 | 31.4 | 94.6 | 331.7 |
| 2001..... | 16.3 | 1.0 | * | 0.1 | 0.4 | 0.7 | 1.6 | 4.8 | 12.1 | 32.1 | 92.7 | 336.0 |
| 2000..... | 16.4 | 1.1 | * | 0.1 | 0.4 | 0.7 | 1.8 | 4.5 | 12.6 | 33.0 | 93.1 | 348.3 |
| 1999..... | 17.2 | 0.9 | 0.3 | 0.1 | 0.5 | 0.7 | 1.8 | 5.0 | 14.0 | 35.5 | 97.5 | 362.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 13.9 | * | * | * | 0.5 | 0.9 | 2.4 | 6.4 | 15.9 | 36.0 | 94.6 | 282.9 |
| 2003..... | 14.4 | * | * | 0.2 | 0.5 | 0.9 | 2.5 | 6.4 | 16.1 | 39.4 | 100.8 | 299.7 |
| 2002..... | 14.7 | 1.7 | * | 0.2 | 0.7 | 0.9 | 2.7 | 6.5 | 16.2 | 38.9 | 104.7 | 316.8 |
| 2001..... | 14.2 | * | * | 0.2 | 0.5 | 0.8 | 2.1 | 6.5 | 15.8 | 39.3 | 104.4 | 316.9 |
| 2000..... | 14.5 | * | * | * | 0.5 | 0.8 | 2.3 | 6.3 | 17.1 | 42.6 | 103.7 | 339.9 |
| 1999..... | 15.3 | * | * | 0.2 | 0.6 | 0.9 | 2.4 | 7.0 | 18.7 | 45.3 | 110.4 | 357.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 17.4 | * | * | * | 0.4 | 0.6 | 1.3 | 3.2 | 7.3 | 22.5 | 79.9 | 313.2 |
| 2003..... | 18.2 | * | * | * | 0.3 | 0.6 | 1.5 | 3.0 | 8.0 | 23.3 | 85.2 | 332.4 |
| 2002..... | 18.5 | * | * | * | 0.3 | 0.5 | 1.4 | 3.0 | 7.9 | 24.9 | 87.8 | 338.2 |
| 2001..... | 18.3 | * | * | * | 0.2 | 0.5 | 1.1 | 3.1 | 8.5 | 26.0 | 84.9 | 343.9 |
| 2000..... | 18.3 | * | * | * | 0.3 | 0.5 | 1.3 | 2.8 | 8.4 | 25.0 | 86.1 | 351.7 |
| 1999..... | 19.0 | * | * | * | 0.4 | 0.6 | 1.3 | 3.1 | 9.7 | 27.3 | 89.0 | 364.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Conduction disorders and cardiac dysrhythmias (I44-I49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 12.6 | * | * | * | 0.7 | 1.7 | 4.2 | 10.2 | 24.5 | 49.3 | 116.0 | 321.2 |
| 2003..... | 12.9 | * | * | * | 0.7 | 1.8 | 3.7 | 10.8 | 24.0 | 56.3 | 116.7 | 332.0 |
| 2002..... | 13.0 | * | * | * | 0.6 | 1.7 | 3.7 | 10.4 | 26.6 | 52.4 | 125.1 | 342.4 |
| 2001..... | 13.1 | * | * | * | 0.8 | 1.5 | 3.8 | 10.8 | 25.5 | 54.9 | 129.1 | 343.0 |
| 2000..... | 12.7 | * | * | * | 0.7 | 1.4 | 3.7 | 10.1 | 25.6 | 53.6 | 123.6 | 349.8 |
| 1999..... | 13.4 | * | * | * | 0.8 | 1.8 | 3.7 | 10.8 | 27.4 | 56.8 | 134.0 | 362.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 12.4 | * | * | * | 0.8 | 1.8 | 4.6 | 12.8 | 32.6 | 64.8 | 139.8 | 340.4 |
| 2003..... | 12.4 | * | * | * | 1.0 | 1.7 | 4.1 | 13.7 | 31.8 | 74.0 | 137.9 | 311.1 |
| 2002..... | 12.4 | * | * | * | 0.8 | 2.0 | 4.3 | 13.6 | 34.7 | 66.0 | 142.1 | 341.3 |
| 2001..... | 12.7 | * | * | * | 1.1 | 1.9 | 4.3 | 13.4 | 34.6 | 67.0 | 161.1 | 359.2 |
| 2000..... | 12.1 | * | * | * | 1.6 | 4.3 | 13.9 | 34.2 | 68.4 | 134.5 | 362.8 | |
| 1999..... | 13.3 | * | * | * | 1.0 | 2.0 | 4.1 | 13.8 | 37.0 | 72.0 | 165.4 | 413.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 12.7 | * | * | * | * | 1.7 | 3.8 | 7.9 | 17.9 | 38.3 | 102.6 | 313.6 |
| 2003..... | 13.3 | * | * | * | * | 1.8 | 3.3 | 8.4 | 17.8 | 43.8 | 104.8 | 340.1 |
| 2002..... | 13.5 | * | * | * | * | 1.5 | 3.2 | 7.7 | 20.1 | 42.8 | 115.4 | 342.8 |
| 2001..... | 13.4 | * | * | * | * | 1.2 | 3.3 | 8.5 | 18.3 | 46.3 | 111.2 | 337.1 |
| 2000..... | 13.3 | * | * | * | 0.8 | 1.2 | 3.3 | 6.8 | 18.7 | 43.2 | 117.4 | 345.0 |
| 1999..... | 13.6 | * | * | * | * | 1.6 | 3.4 | 8.2 | 19.8 | 46.1 | 116.4 | 342.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Heart failure (I50) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 19.5 | 0.7 | * | * | 0.1 | 0.2 | 0.7 | 2.5 | 8.4 | 32.2 | 131.2 | 621.2 |
| 2003..... | 19.8 | 0.9 | * | * | 0.1 | 0.2 | 0.7 | 2.4 | 8.8 | 33.5 | 133.2 | 641.1 |
| 2002..... | 19.6 | 0.6 | * | * | 0.1 | 0.3 | 0.6 | 2.5 | 8.8 | 33.8 | 132.8 | 645.0 |
| 2001..... | 20.0 | 0.6 | * | * | 0.1 | 0.2 | 0.7 | 2.4 | 9.4 | 34.3 | 136.3 | 674.6 |
| 2000..... | 19.8 | 0.8 | * | 0.1 | 0.1 | 0.2 | 0.6 | 2.2 | 9.1 | 34.8 | 136.1 | 684.9 |
| 1999..... | 19.7 | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 8.9 | 35.4 | 137.2 | 681.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 15.4 | * | * | * | * | 0.3 | 0.8 | 3.0 | 10.1 | 37.5 | 147.0 | 605.5 |
| 2003..... | 15.7 | 1.4 | * | * | 0.1 | 0.3 | 0.9 | 2.9 | 10.8 | 39.7 | 148.6 | 629.8 |
| 2002..... | 15.3 | * | * | * | * | 0.3 | 0.8 | 3.1 | 10.5 | 39.2 | 149.0 | 621.4 |
| 2001..... | 15.5 | * | * | * | * | 0.3 | 0.8 | 3.1 | 11.1 | 40.3 | 153.8 | 646.4 |
| 2000..... | 15.3 | 1.1 | * | * | 0.1 | 0.2 | 0.7 | 2.7 | 10.9 | 41.1 | 153.7 | 672.2 |
| 1999..... | 15.2 | * | * | * | 0.1 | 0.2 | 0.7 | 2.7 | 10.8 | 42.5 | 152.0 | 667.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 23.4 | * | * | * | * | 0.2 | 0.6 | 2.0 | 6.8 | 27.8 | 120.5 | 628.2 |
| 2003..... | 23.7 | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.0 | 28.3 | 122.9 | 646.1 |
| 2002..... | 23.7 | * | * | * | * | 0.2 | 0.5 | 1.8 | 7.2 | 29.2 | 122.0 | 655.3 |
| 2001..... | 24.3 | * | * | * | * | 0.2 | 0.6 | 1.8 | 7.8 | 29.3 | 124.7 | 686.4 |
| 2000..... | 24.1 | * | * | * | * | 0.1 | 0.5 | 1.6 | 7.5 | 29.6 | 124.6 | 690.0 |
| 1999..... | 24.0 | * | * | * | * | 0.2 | 0.5 | 1.6 | 7.1 | 29.6 | 127.6 | 687.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | * | 0.4 | 1.3 | 4.7 | 19.2 | 72.9 | 287.3 |
| 2003..... | 4.0 | * | * | * | * | * | * | 1.3 | 5.7 | 22.5 | 77.1 | 330.1 |
| 2002..... | 3.6 | * | * | * | * | * | * | 1.2 | 4.6 | 19.9 | 75.2 | 316.0 |
| 2001..... | 3.7 | * | * | * | * | * | 0.4 | 1.3 | 5.6 | 18.8 | 72.9 | 349.5 |
| 2000..... | 3.6 | * | * | * | * | * | * | 0.7 | 5.8 | 18.9 | 72.1 | 369.6 |
| 1999..... | 3.7 | * | * | * | * | * | * | 1.0 | 5.3 | 19.5 | 73.5 | 377.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | * | 1.7 | 6.0 | 22.1 | 84.1 | 263.4 |
| 2003..... | 3.5 | * | * | * | * | * | * | 1.3 | 7.8 | 26.0 | 92.8 | 312.5 |
| 2002..... | 3.0 | * | * | * | * | * | * | 1.1 | 5.8 | 23.1 | 85.8 | 296.2 |
| 2001..... | 3.0 | * | * | * | * | * | * | 1.8 | 6.0 | 21.8 | 75.6 | 333.3 |
| 2000..... | 2.9 | * | * | * | * | * | * | * | 7.7 | 23.0 | 74.2 | 380.9 |
| 1999..... | 3.0 | * | * | * | * | * | * | * | 6.1 | 24.6 | 82.9 | 358.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | * | * | 1.0 | 3.4 | 16.9 | 65.2 | 299.6 |
| 2003..... | 4.6 | * | * | * | * | * | * | 1.2 | 3.7 | 19.7 | 66.4 | 339.1 |
| 2002..... | 4.4 | * | * | * | * | * | * | 1.3 | 3.6 | 17.3 | 68.0 | 325.8 |
| 2001..... | 4.5 | * | * | * | * | * | * | * | 5.2 | 16.4 | 71.0 | 357.4 |
| 2000..... | 4.3 | * | * | * | * | * | * | * | 4.2 | 15.6 | 70.7 | 364.0 |
| 1999..... | 4.4 | * | * | * | * | * | * | * | 4.6 | 15.5 | 67.3 | 386.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Heart failure (I50) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 22.0 | 0.8 | * | * | 0.1 | 0.3 | 0.8 | 2.6 | 8.7 | 33.1 | 134.2 | 635.3 |
| 2003..... | 22.2 | 1.0 | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 9.1 | 34.1 | 135.9 | 653.2 |
| 2002..... | 22.0 | * | * | * | 0.1 | 0.3 | 0.7 | 2.6 | 9.1 | 34.6 | 135.3 | 656.8 |
| 2001..... | 22.4 | * | * | * | 0.1 | 0.3 | 0.7 | 2.5 | 9.6 | 35.2 | 138.8 | 685.6 |
| 2000..... | 22.1 | 0.8 | * | * | 0.1 | 0.2 | 0.6 | 2.3 | 9.3 | 35.7 | 138.5 | 694.8 |
| 1999..... | 21.8 | * | * | * | 0.1 | 0.2 | 0.6 | 2.2 | 9.1 | 36.3 | 139.4 | 691.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 17.5 | * | * | * | * | 0.3 | 0.9 | 3.1 | 10.4 | 38.6 | 150.3 | 621.3 |
| 2003..... | 17.7 | 1.4 | * | * | 0.1 | 0.3 | 1.0 | 3.1 | 11.0 | 40.5 | 151.2 | 643.2 |
| 2002..... | 17.3 | * | * | * | * | 0.4 | 0.9 | 3.3 | 10.9 | 40.2 | 151.8 | 634.7 |
| 2001..... | 17.4 | * | * | * | * | 0.3 | 0.9 | 3.2 | 11.4 | 41.4 | 157.2 | 658.0 |
| 2000..... | 17.2 | * | * | * | 0.1 | 0.2 | 0.8 | 2.9 | 11.0 | 42.1 | 156.9 | 683.3 |
| 1999..... | 16.9 | * | * | * | 0.1 | 0.2 | 0.8 | 2.8 | 11.0 | 43.5 | 154.5 | 677.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 26.3 | * | * | * | * | 0.2 | 0.6 | 2.1 | 7.1 | 28.6 | 123.3 | 641.6 |
| 2003..... | 26.5 | * | * | * | * | 0.2 | 0.6 | 2.0 | 7.3 | 28.8 | 125.6 | 657.6 |
| 2002..... | 26.5 | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.5 | 29.9 | 124.3 | 666.4 |
| 2001..... | 27.1 | * | * | * | * | 0.2 | 0.6 | 1.9 | 8.0 | 30.1 | 126.8 | 697.0 |
| 2000..... | 26.7 | * | * | * | * | 0.2 | 0.5 | 1.7 | 7.7 | 30.5 | 126.5 | 699.5 |
| 1999..... | 26.5 | * | * | * | * | 0.2 | 0.5 | 1.6 | 7.2 | 30.4 | 129.7 | 696.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 24.8 | * | * | * | * | 0.2 | 0.6 | 2.0 | 7.6 | 32.3 | 135.7 | 657.1 |
| 2003..... | 25.0 | * | * | * | * | 0.2 | 0.5 | 1.9 | 8.0 | 32.8 | 137.1 | 673.5 |
| 2002..... | 24.7 | * | * | * | * | 0.2 | 0.5 | 2.0 | 8.2 | 33.3 | 135.9 | 677.3 |
| 2001..... | 25.1 | * | * | * | * | 0.2 | 0.5 | 2.1 | 8.7 | 33.9 | 139.7 | 706.8 |
| 2000..... | 24.7 | * | * | * | * | 0.1 | 0.5 | 1.9 | 8.4 | 34.3 | 139.2 | 714.3 |
| 1999..... | 24.3 | * | * | * | * | 0.1 | 0.5 | 1.7 | 8.1 | 34.9 | 139.1 | 709.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 19.5 | * | * | * | * | 0.2 | 0.7 | 2.4 | 9.0 | 37.8 | 151.7 | 643.0 |
| 2003..... | 19.7 | * | * | * | * | 0.3 | 0.7 | 2.4 | 9.7 | 39.3 | 153.2 | 665.3 |
| 2002..... | 19.2 | * | * | * | * | 0.2 | 0.7 | 2.6 | 9.8 | 38.6 | 153.2 | 656.2 |
| 2001..... | 19.3 | * | * | * | * | 0.2 | 0.7 | 2.7 | 10.5 | 40.0 | 158.1 | 679.1 |
| 2000..... | 19.0 | * | * | * | * | 0.2 | 0.6 | 2.4 | 10.1 | 41.0 | 157.9 | 701.6 |
| 1999..... | 18.6 | * | * | * | * | * | 0.6 | 2.2 | 10.1 | 42.2 | 154.7 | 690.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 30.0 | * | * | * | * | * | 0.4 | 1.5 | 6.2 | 27.6 | 124.7 | 663.4 |
| 2003..... | 30.1 | * | * | * | * | * | 0.4 | 1.5 | 6.3 | 27.2 | 126.3 | 677.2 |
| 2002..... | 30.0 | * | * | * | * | 0.2 | 0.4 | 1.4 | 6.7 | 28.8 | 124.2 | 686.4 |
| 2001..... | 30.7 | * | * | * | * | 0.2 | 0.4 | 1.4 | 6.9 | 28.8 | 127.5 | 718.4 |
| 2000..... | 30.2 | * | * | * | * | * | 0.3 | 1.3 | 6.8 | 28.7 | 126.9 | 719.4 |
| 1999..... | 29.9 | * | * | * | * | * | 0.3 | 1.2 | 6.3 | 28.8 | 128.8 | 716.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Heart failure (I50) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 14.1 | * | * | * | * | 0.7 | 2.0 | 7.7 | 19.9 | 49.2 | 153.0 | 513.2 |
| 2003..... | 14.5 | * | * | * | * | 0.7 | 2.4 | 7.1 | 21.0 | 55.3 | 152.7 | 538.7 |
| 2002..... | 14.2 | * | * | * | * | 0.9 | 1.9 | 7.4 | 19.0 | 54.1 | 158.3 | 534.5 |
| 2001..... | 14.3 | * | * | * | * | 0.6 | 2.0 | 6.6 | 20.0 | 55.5 | 159.1 | 551.2 |
| 2000..... | 14.1 | * | * | * | * | 0.5 | 1.7 | 5.9 | 19.0 | 56.2 | 156.3 | 564.8 |
| 1999..... | 14.3 | * | * | * | * | 0.6 | 1.7 | 6.5 | 19.2 | 57.3 | 164.8 | 557.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 12.2 | * | * | * | * | 1.0 | 2.3 | 9.5 | 25.6 | 56.6 | 178.5 | 510.1 |
| 2003..... | 12.3 | * | * | * | * | 0.8 | 2.8 | 8.5 | 26.8 | 64.3 | 165.9 | 521.0 |
| 2002..... | 12.0 | * | * | * | * | 1.1 | 2.3 | 9.2 | 23.3 | 65.9 | 176.4 | 489.0 |
| 2001..... | 12.0 | * | * | * | * | 0.9 | 2.3 | 7.6 | 22.6 | 66.3 | 185.4 | 547.5 |
| 2000..... | 11.6 | * | * | * | * | * | 1.8 | 7.3 | 21.8 | 62.9 | 176.5 | 589.4 |
| 1999..... | 12.2 | * | * | * | * | * | 1.9 | 8.6 | 23.0 | 66.4 | 180.4 | 609.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 15.7 | * | * | * | * | * | 1.8 | 6.1 | 15.4 | 43.9 | 138.6 | 514.4 |
| 2003..... | 16.5 | * | * | * | * | * | 2.0 | 5.8 | 16.3 | 48.9 | 145.2 | 545.6 |
| 2002..... | 16.3 | * | * | * | * | * | 1.5 | 5.7 | 15.5 | 45.8 | 148.1 | 551.8 |
| 2001..... | 16.4 | * | * | * | * | * | 1.7 | 5.6 | 18.0 | 47.9 | 144.4 | 552.6 |
| 2000..... | 16.3 | * | * | * | * | * | 1.6 | 4.7 | 16.8 | 51.5 | 145.0 | 555.6 |
| 1999..... | 16.2 | * | * | * | * | * | 1.5 | 4.8 | 16.1 | 50.9 | 156.0 | 537.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Congestive heart failure (I50.0) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 18.1 | * | * | * | 0.0 | 0.2 | 0.6 | 2.1 | 7.2 | 29.1 | 122.3 | 588.9 |
| 2003..... | 18.4 | 0.6 | * | * | 0.1 | 0.2 | 0.6 | 2.0 | 7.7 | 30.3 | 124.4 | 608.2 |
| 2002..... | 18.2 | * | * | * | * | 0.2 | 0.5 | 2.0 | 7.6 | 30.3 | 123.7 | 609.0 |
| 2001..... | 18.5 | * | * | * | 0.1 | 0.2 | 0.5 | 2.1 | 8.0 | 31.1 | 126.6 | 636.1 |
| 2000..... | 18.3 | * | * | * | 0.1 | 0.1 | 0.5 | 1.8 | 7.9 | 31.0 | 126.4 | 644.4 |
| 1999..... | 18.2 | * | * | * | 0.1 | 0.1 | 0.5 | 1.7 | 7.4 | 31.9 | 127.3 | 642.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 14.2 | * | * | * | * | 0.2 | 0.7 | 2.5 | 8.7 | 33.6 | 136.4 | 571.8 |
| 2003..... | 14.5 | * | * | * | * | 0.2 | 0.7 | 2.4 | 9.3 | 35.8 | 138.3 | 594.6 |
| 2002..... | 14.0 | * | * | * | * | 0.3 | 0.6 | 2.5 | 8.9 | 35.2 | 138.1 | 584.6 |
| 2001..... | 14.2 | * | * | * | * | 0.2 | 0.6 | 2.6 | 9.3 | 36.2 | 141.2 | 609.3 |
| 2000..... | 14.0 | * | * | * | * | 0.1 | 0.5 | 2.2 | 9.3 | 36.4 | 142.4 | 632.2 |
| 1999..... | 13.9 | * | * | * | * | 0.1 | 0.6 | 2.2 | 8.9 | 38.0 | 140.5 | 628.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 21.9 | * | * | * | * | 0.1 | 0.5 | 1.7 | 5.8 | 25.3 | 112.8 | 596.6 |
| 2003..... | 22.3 | * | * | * | * | 0.2 | 0.5 | 1.6 | 6.2 | 25.7 | 115.2 | 614.2 |
| 2002..... | 22.2 | * | * | * | * | 0.2 | 0.4 | 1.5 | 6.3 | 26.3 | 114.1 | 619.6 |
| 2001..... | 22.8 | * | * | * | * | 0.1 | 0.4 | 1.6 | 6.8 | 26.8 | 117.0 | 647.3 |
| 2000..... | 22.4 | * | * | * | * | 0.1 | 0.4 | 1.4 | 6.6 | 26.6 | 116.1 | 649.4 |
| 1999..... | 22.4 | * | * | * | * | 0.1 | 0.4 | 1.3 | 6.1 | 26.9 | 118.7 | 648.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 0.3 | 1.3 | 4.0 | 17.0 | 67.9 | 271.1 |
| 2003..... | 3.7 | * | * | * | * | * | * | 1.2 | 5.2 | 20.1 | 71.7 | 315.1 |
| 2002..... | 3.3 | * | * | * | * | * | * | 0.9 | 4.3 | 17.2 | 69.2 | 296.0 |
| 2001..... | 3.4 | * | * | * | * | * | * | 1.2 | 4.9 | 17.3 | 68.0 | 322.6 |
| 2000..... | 3.3 | * | * | * | * | * | * | * | 5.4 | 16.9 | 67.0 | 343.7 |
| 1999..... | 3.4 | * | * | * | * | * | * | 0.7 | 4.9 | 17.9 | 68.7 | 359.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | * | 1.6 | 5.2 | 19.3 | 76.9 | 249.4 |
| 2003..... | 3.2 | * | * | * | * | * | * | 1.3 | 7.2 | 22.5 | 85.5 | 295.5 |
| 2002..... | 2.6 | * | * | * | * | * | * | * | 5.2 | 20.3 | 76.7 | 274.4 |
| 2001..... | 2.7 | * | * | * | * | * | * | 1.7 | 5.3 | 20.6 | 68.4 | 295.2 |
| 2000..... | 2.6 | * | * | * | * | * | * | * | 7.0 | 20.7 | 67.4 | 342.6 |
| 1999..... | 2.7 | * | * | * | * | * | * | * | 5.6 | 22.2 | 77.7 | 338.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | * | * | 2.8 | 15.2 | 61.7 | 282.3 |
| 2003..... | 4.4 | * | * | * | * | * | * | 1.1 | 3.4 | 18.2 | 62.1 | 325.0 |
| 2002..... | 4.0 | * | * | * | * | * | * | * | 3.4 | 14.8 | 64.1 | 306.7 |
| 2001..... | 4.1 | * | * | * | * | * | * | * | 4.5 | 14.6 | 67.7 | 336.0 |
| 2000..... | 4.0 | * | * | * | * | * | * | * | 4.0 | 13.9 | 66.7 | 344.2 |
| 1999..... | 4.1 | * | * | * | * | * | * | * | 4.4 | 14.6 | 62.8 | 369.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Congestive heart failure (I50.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.5 | * | * | * | * | 0.2 | 0.6 | 2.2 | 7.4 | 29.9 | 125.1 | 602.4 |
| 2003..... | 20.7 | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 7.9 | 31.0 | 126.9 | 619.6 |
| 2002..... | 20.4 | * | * | * | * | 0.2 | 0.6 | 2.1 | 7.8 | 31.1 | 126.1 | 620.2 |
| 2001..... | 20.8 | * | * | * | 0.1 | 0.2 | 0.6 | 2.1 | 8.2 | 31.9 | 129.0 | 646.7 |
| 2000..... | 20.4 | * | * | * | 0.1 | 0.1 | 0.5 | 1.9 | 8.1 | 31.8 | 128.7 | 653.9 |
| 1999..... | 20.2 | * | * | * | 0.1 | 0.1 | 0.5 | 1.8 | 7.6 | 32.6 | 129.4 | 651.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 16.1 | * | * | * | * | 0.3 | 0.7 | 2.6 | 8.9 | 34.5 | 139.5 | 586.8 |
| 2003..... | 16.3 | * | * | * | * | 0.2 | 0.7 | 2.5 | 9.4 | 36.6 | 140.7 | 607.4 |
| 2002..... | 15.9 | * | * | * | * | 0.3 | 0.7 | 2.6 | 9.2 | 36.1 | 140.8 | 597.3 |
| 2001..... | 15.9 | * | * | * | * | 0.2 | 0.7 | 2.6 | 9.6 | 37.1 | 144.4 | 621.0 |
| 2000..... | 15.7 | * | * | * | * | 0.2 | 0.6 | 2.3 | 9.4 | 37.2 | 145.4 | 643.2 |
| 1999..... | 15.5 | * | * | * | * | 0.1 | 0.6 | 2.3 | 9.0 | 38.8 | 142.7 | 637.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 24.6 | * | * | * | * | 0.1 | 0.5 | 1.8 | 6.0 | 26.0 | 115.4 | 609.4 |
| 2003..... | 24.9 | * | * | * | * | 0.2 | 0.5 | 1.7 | 6.4 | 26.2 | 117.7 | 625.0 |
| 2002..... | 24.8 | * | * | * | * | 0.2 | 0.4 | 1.6 | 6.6 | 27.0 | 116.3 | 630.1 |
| 2001..... | 25.4 | * | * | * | * | 0.2 | 0.5 | 1.7 | 6.9 | 27.6 | 118.9 | 657.3 |
| 2000..... | 24.9 | * | * | * | * | 0.1 | 0.4 | 1.4 | 6.8 | 27.3 | 117.8 | 658.3 |
| 1999..... | 24.7 | * | * | * | * | 0.1 | 0.4 | 1.3 | 6.2 | 27.6 | 120.7 | 656.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 23.3 | * | * | * | * | 0.2 | 0.5 | 1.7 | 6.5 | 29.3 | 126.8 | 624.1 |
| 2003..... | 23.4 | * | * | * | * | 0.1 | 0.4 | 1.6 | 6.9 | 29.9 | 128.5 | 640.9 |
| 2002..... | 23.1 | * | * | * | * | 0.1 | 0.4 | 1.6 | 7.0 | 30.1 | 127.0 | 641.1 |
| 2001..... | 23.4 | * | * | * | * | 0.2 | 0.4 | 1.7 | 7.4 | 30.8 | 130.2 | 667.8 |
| 2000..... | 23.0 | * | * | * | * | 0.1 | 0.4 | 1.5 | 7.3 | 30.7 | 129.7 | 673.7 |
| 1999..... | 22.6 | * | * | * | * | * | 0.4 | 1.3 | 6.8 | 31.4 | 129.4 | 669.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 18.1 | * | * | * | * | 0.2 | 0.6 | 2.0 | 7.7 | 34.1 | 141.3 | 608.3 |
| 2003..... | 18.3 | * | * | * | * | 0.2 | 0.5 | 2.0 | 8.3 | 35.7 | 143.1 | 630.0 |
| 2002..... | 17.7 | * | * | * | * | 0.2 | 0.5 | 2.0 | 8.2 | 34.6 | 142.7 | 619.6 |
| 2001..... | 17.8 | * | * | * | * | 0.2 | 0.5 | 2.2 | 8.9 | 36.0 | 145.9 | 641.8 |
| 2000..... | 17.5 | * | * | * | * | * | 0.5 | 1.9 | 8.7 | 36.5 | 146.8 | 662.2 |
| 1999..... | 17.0 | * | * | * | * | * | 0.5 | 1.7 | 8.2 | 37.8 | 143.1 | 652.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 28.2 | * | * | * | * | * | 0.3 | 1.3 | 5.4 | 25.2 | 117.0 | 631.3 |
| 2003..... | 28.4 | * | * | * | * | * | 0.3 | 1.3 | 5.6 | 24.9 | 118.6 | 645.7 |
| 2002..... | 28.2 | * | * | * | * | * | 0.3 | 1.1 | 5.8 | 26.2 | 116.5 | 650.4 |
| 2001..... | 28.8 | * | * | * | * | * | 0.3 | 1.3 | 6.0 | 26.4 | 119.8 | 678.6 |
| 2000..... | 28.2 | * | * | * | * | * | 0.3 | 1.1 | 6.0 | 25.7 | 118.4 | 678.4 |
| 1999..... | 27.9 | * | * | * | * | * | 0.3 | 0.9 | 5.4 | 26.1 | 120.4 | 676.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Congestive heart failure (I50.0) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 12.5 | * | * | * | * | 0.6 | 1.6 | 6.3 | 16.9 | 43.3 | 138.3 | 474.9 |
| 2003..... | 12.9 | * | * | * | * | 0.6 | 2.0 | 5.9 | 18.1 | 48.9 | 138.2 | 490.5 |
| 2002..... | 12.8 | * | * | * | * | 0.8 | 1.6 | 6.2 | 16.6 | 48.2 | 143.5 | 488.2 |
| 2001..... | 12.8 | * | * | * | * | 0.4 | 1.7 | 5.6 | 16.9 | 49.9 | 143.8 | 509.4 |
| 2000..... | 12.5 | * | * | * | * | * | 1.3 | 5.1 | 16.4 | 49.3 | 141.8 | 515.8 |
| 1999..... | 12.8 | * | * | * | * | 0.5 | 1.3 | 5.7 | 16.1 | 51.2 | 149.5 | 514.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 10.7 | * | * | * | * | 0.8 | 1.7 | 7.5 | 22.1 | 49.0 | 160.3 | 472.5 |
| 2003..... | 10.7 | * | * | * | * | * | 2.4 | 7.0 | 22.8 | 57.1 | 147.7 | 474.1 |
| 2002..... | 10.6 | * | * | * | * | 1.1 | 1.9 | 7.6 | 20.2 | 60.6 | 157.1 | 437.6 |
| 2001..... | 10.6 | * | * | * | * | * | 2.0 | 6.6 | 18.6 | 58.6 | 162.0 | 507.0 |
| 2000..... | 10.2 | * | * | * | * | * | 1.4 | 6.4 | 18.4 | 53.6 | 157.4 | 535.4 |
| 1999..... | 10.8 | * | * | * | * | * | 1.6 | 7.3 | 19.0 | 58.9 | 165.1 | 561.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 14.2 | * | * | * | * | * | 1.5 | 5.3 | 12.7 | 39.2 | 125.9 | 475.9 |
| 2003..... | 14.9 | * | * | * | * | * | 1.7 | 5.0 | 14.4 | 43.1 | 132.8 | 496.9 |
| 2002..... | 14.7 | * | * | * | * | * | 1.4 | 4.9 | 13.7 | 39.5 | 135.8 | 507.5 |
| 2001..... | 14.9 | * | * | * | * | * | 1.4 | 4.8 | 15.6 | 43.7 | 133.6 | 510.3 |
| 2000..... | 14.7 | * | * | * | * | * | 1.3 | 4.0 | 14.9 | 46.2 | 133.0 | 508.5 |
| 1999..... | 14.7 | * | * | * | * | * | 1.1 | 4.3 | 13.9 | 45.8 | 140.8 | 497.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified heart failure (I50.1-I50.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.2 | 9.1 | 32.9 |
| 2003..... | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.2 | 8.9 | 33.6 |
| 2002..... | 1.6 | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 3.5 | 9.2 | 36.6 |
| 2001..... | 1.6 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.3 | 9.9 | 38.9 |
| 2000..... | 1.6 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.9 | 9.8 | 40.9 |
| 1999..... | 1.6 | * | * | * | * | * | 0.1 | 0.4 | 1.5 | 3.6 | 10.1 | 39.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.0 | 10.8 | 34.5 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 3.8 | 10.5 | 35.8 |
| 2002..... | 1.4 | * | * | * | * | * | 0.2 | 0.7 | 1.7 | 4.1 | 11.0 | 37.4 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 4.2 | 12.9 | 37.0 |
| 2000..... | 1.4 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.9 | 11.5 | 40.1 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 4.6 | 11.8 | 39.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.5 | 7.9 | 32.2 |
| 2003..... | 1.6 | * | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 7.9 | 32.6 |
| 2002..... | 1.7 | * | * | * | * | * | * | 0.3 | 0.9 | 2.9 | 8.0 | 36.3 |
| 2001..... | 1.7 | * | * | * | * | * | 0.1 | 0.2 | 1.1 | 2.5 | 7.9 | 39.7 |
| 2000..... | 1.8 | * | * | * | * | * | * | 0.3 | 0.9 | 3.1 | 8.7 | 41.3 |
| 1999..... | 1.8 | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.8 | 9.0 | 39.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.0 | 8.8 | 33.0 |
| 2003..... | 1.5 | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 2.9 | 8.6 | 32.6 |
| 2002..... | 1.7 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.2 | 8.9 | 36.2 |
| 2001..... | 1.7 | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 3.1 | 9.5 | 39.0 |
| 2000..... | 1.7 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.7 | 9.5 | 40.6 |
| 1999..... | 1.7 | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 3.5 | 9.7 | 39.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.8 | 10.5 | 34.7 |
| 2003..... | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 3.6 | 10.1 | 35.3 |
| 2002..... | 1.5 | * | * | * | * | * | 0.1 | 0.6 | 1.6 | 4.0 | 10.6 | 36.6 |
| 2001..... | 1.5 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 4.0 | 12.2 | 37.3 |
| 2000..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.5 | 11.0 | 39.5 |
| 1999..... | 1.5 | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 4.4 | 11.6 | 38.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 7.7 | 32.2 |
| 2003..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.7 | 2.3 | 7.6 | 31.5 |
| 2002..... | 1.8 | * | * | * | * | * | * | 0.2 | 0.9 | 2.5 | 7.8 | 36.0 |
| 2001..... | 1.9 | * | * | * | * | * | * | 0.2 | 0.9 | 2.3 | 7.7 | 39.8 |
| 2000..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.8 | 3.0 | 8.5 | 41.1 |
| 1999..... | 1.9 | * | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 8.5 | 40.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified heart failure (I50.1-I50.9) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.4 | 1.4 | 3.1 | 5.9 | 14.7 | 38.3 |
| 2003..... | 1.6 | * | * | * | * | * | * | 1.2 | 2.9 | 6.4 | 14.5 | 48.2 |
| 2002..... | 1.5 | * | * | * | * | * | * | 1.2 | 2.4 | 5.9 | 14.8 | 46.3 |
| 2001..... | 1.5 | * | * | * | * | * | * | 0.9 | 3.1 | 5.6 | 15.3 | 41.8 |
| 2000..... | 1.5 | * | * | * | * | * | 0.4 | 0.8 | 2.6 | 6.9 | 14.6 | 49.0 |
| 1999..... | 1.5 | * | * | * | * | * | 0.4 | 0.8 | 3.1 | 6.1 | 15.3 | 43.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 2.0 | 3.5 | 7.6 | 18.3 | 37.6 |
| 2003..... | 1.5 | * | * | * | * | * | * | 1.6 | 4.0 | 7.2 | 18.2 | 46.9 |
| 2002..... | 1.4 | * | * | * | * | * | * | 1.7 | 3.1 | 5.3 | 19.3 | 51.4 |
| 2001..... | 1.5 | * | * | * | * | * | * | 1.0 | 3.9 | 7.7 | 23.4 | 40.4 |
| 2000..... | 1.5 | * | * | * | * | * | * | * | 3.4 | 9.3 | 19.1 | 54.0 |
| 1999..... | 1.3 | * | * | * | * | * | * | 1.3 | 4.1 | 7.5 | 15.3 | 48.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 0.8 | 2.7 | 4.7 | 12.7 | 38.6 |
| 2003..... | 1.6 | * | * | * | * | * | * | 0.8 | 1.9 | 5.9 | 12.4 | 48.7 |
| 2002..... | 1.6 | * | * | * | * | * | * | * | 1.8 | 6.3 | 12.3 | 44.3 |
| 2001..... | 1.5 | * | * | * | * | * | * | 0.9 | 2.4 | 4.2 | 10.8 | 42.3 |
| 2000..... | 1.5 | * | * | * | * | * | * | * | 2.0 | 5.3 | 12.0 | 47.1 |
| 1999..... | 1.6 | * | * | * | * | * | * | * | 2.2 | 5.1 | 15.2 | 40.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 2.1 |
| 2003..... | 0.2 | 0.8 | 0.2 | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.2 | 0.1 | 0.4 | 2.2 |
| 2002..... | 0.3 | 0.8 | * | * | 0.2 | 0.1 | 0.3 | 0.3 | 0.3 | 0.5 | 0.9 | 3.2 |
| 2001..... | 0.3 | 0.6 | 0.3 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.2 | 0.4 | 0.5 | 2.1 |
| 2000..... | 0.3 | * | 0.2 | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 | 0.3 | 0.6 | 2.5 |
| 1999..... | 0.3 | 0.7 | 0.2 | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.2 | 0.3 | 0.5 | 2.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.2 | 0.1 | 0.3 | 0.3 | 0.2 | 0.3 | 0.4 | 1.8 |
| 2003..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | * | 0.6 | 2.6 |
| 2002..... | 0.3 | * | * | * | 0.2 | 0.2 | 0.4 | 0.3 | 0.3 | 0.6 | 0.9 | 3.8 |
| 2001..... | 0.3 | * | 0.3 | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.5 | 0.8 | 2.2 |
| 2000..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 | 0.3 | 0.7 | 2.7 |
| 1999..... | 0.3 | * | * | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.2 | 0.4 | 0.6 | 2.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 | 0.3 | 2.3 |
| 2003..... | 0.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | * | 0.3 | 2.0 |
| 2002..... | 0.3 | * | * | * | 0.2 | * | 0.3 | 0.3 | 0.4 | 0.4 | 0.9 | 3.0 |
| 2001..... | 0.3 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.2 | 0.3 | 0.4 | 2.0 |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 2.4 |
| 1999..... | 0.3 | * | * | * | 0.2 | 0.1 | 0.2 | 0.3 | 0.2 | 0.2 | 0.4 | 2.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.1 | 0.3 | 0.3 | 2.2 |
| 2003..... | 0.2 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.4 | 2.3 |
| 2002..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.5 | 0.9 | 3.2 |
| 2001..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.2 | 0.4 | 0.5 | 2.1 |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 2.6 |
| 1999..... | 0.2 | * | * | * | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 2.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.2 | * | 0.2 | 0.2 | * | * | 0.5 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | * | 0.6 | 2.8 |
| 2002..... | 0.3 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.3 | 0.6 | 0.9 | 4.0 |
| 2001..... | 0.3 | * | * | * | * | * | 0.3 | 0.3 | 0.2 | 0.5 | 0.7 | 2.2 |
| 2000..... | 0.2 | * | * | * | * | 0.2 | 0.3 | * | 0.3 | 0.3 | 0.6 | 2.8 |
| 1999..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.3 | 0.3 | * | 0.4 | 0.6 | 2.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.2 | 0.3 | * | 0.3 | * | 2.3 |
| 2003..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | * | 0.3 | 2.0 |
| 2002..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.9 | 2.8 |
| 2001..... | 0.2 | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | 2.0 |
| 2000..... | 0.3 | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 2.5 |
| 1999..... | 0.2 | * | * | * | 0.2 | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 2.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | 0.6 | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 | 3.6 | 7.2 | 20.4 | 67.7 |
| 2003..... | 3.4 | 0.5 | * | 0.0 | 0.2 | 0.4 | 1.0 | 1.9 | 3.6 | 7.6 | 20.5 | 67.8 |
| 2002..... | 3.3 | 0.6 | * | * | 0.2 | 0.4 | 1.1 | 1.9 | 3.4 | 7.8 | 20.0 | 65.0 |
| 2001..... | 3.2 | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 1.8 | 3.4 | 7.3 | 18.9 | 66.3 |
| 2000..... | 3.1 | 0.8 | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 3.7 | 7.5 | 19.3 | 64.6 |
| 1999..... | 2.9 | 0.8 | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 3.5 | 6.9 | 17.8 | 61.5 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | 0.4 | 0.7 | 1.4 | 3.0 | 4.8 | 9.6 | 24.9 | 67.8 |
| 2003..... | 3.4 | * | * | * | 0.4 | 0.7 | 1.5 | 2.5 | 4.9 | 9.3 | 24.6 | 72.1 |
| 2002..... | 3.3 | * | * | * | 0.4 | 0.6 | 1.5 | 2.7 | 4.6 | 10.6 | 23.6 | 66.7 |
| 2001..... | 3.2 | * | * | * | 0.3 | 0.7 | 1.5 | 2.6 | 4.7 | 9.9 | 22.1 | 66.4 |
| 2000..... | 3.1 | * | * | * | 0.4 | 0.6 | 1.3 | 2.5 | 5.3 | 9.6 | 22.4 | 69.0 |
| 1999..... | 3.0 | * | * | * | 0.3 | 0.6 | 1.3 | 2.3 | 4.9 | 9.4 | 21.8 | 64.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | 0.1 | 0.2 | 0.7 | 1.1 | 2.4 | 5.1 | 17.3 | 67.7 |
| 2003..... | 3.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.3 | 6.1 | 17.8 | 66.0 |
| 2002..... | 3.2 | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.2 | 5.4 | 17.5 | 64.3 |
| 2001..... | 3.1 | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 5.2 | 16.8 | 66.3 |
| 2000..... | 3.1 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.2 | 5.7 | 17.3 | 62.7 |
| 1999..... | 2.8 | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.1 | 4.8 | 15.2 | 60.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.4 | 0.6 | 1.1 | 1.6 | 4.1 | 18.4 | 60.9 |
| 2003..... | 1.0 | * | * | * | * | * | 0.6 | 1.0 | 1.5 | 4.6 | 15.1 | 62.2 |
| 2002..... | 1.0 | * | * | * | * | * | 0.6 | 0.7 | 1.3 | 4.9 | 15.5 | 58.2 |
| 2001..... | 0.9 | * | * | * | * | * | 0.4 | 1.0 | 2.1 | 3.9 | 12.1 | 48.1 |
| 2000..... | 0.9 | * | * | * | * | * | 0.5 | 0.8 | 1.4 | 4.3 | 15.8 | 46.4 |
| 1999..... | 0.9 | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.0 | 4.8 | 12.5 | 39.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.6 | 0.8 | 1.7 | 2.5 | 5.3 | 24.7 | 57.4 |
| 2003..... | 1.1 | * | * | * | * | * | 1.0 | 1.5 | 2.2 | 6.6 | 11.5 | 66.2 |
| 2002..... | 1.1 | * | * | * | * | * | 0.9 | * | * | 7.5 | 19.3 | 60.2 |
| 2001..... | 1.0 | * | * | * | * | * | * | 1.6 | 3.0 | 4.4 | 17.0 | 43.8 |
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | * | * | 5.9 | 17.7 | 60.5 |
| 1999..... | 1.1 | * | * | * | * | 0.6 | 0.9 | 1.8 | * | 7.2 | 16.2 | 49.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | * | 3.1 | 14.1 | 62.7 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | * | 2.9 | 17.6 | 60.2 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | * | 12.9 | 57.2 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 3.5 | 8.8 | 50.3 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | * | 14.5 | 39.6 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | * | 10.1 | 34.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 | 3.7 | 7.4 | 20.5 | 67.9 |
| 2003..... | 3.7 | * | * | * | 0.3 | 0.5 | 1.1 | 1.9 | 3.7 | 7.7 | 20.8 | 68.0 |
| 2002..... | 3.6 | 0.6 | * | * | 0.2 | 0.4 | 1.1 | 2.0 | 3.5 | 8.0 | 20.1 | 65.2 |
| 2001..... | 3.5 | * | * | * | 0.2 | 0.5 | 1.1 | 1.9 | 3.5 | 7.5 | 19.2 | 66.8 |
| 2000..... | 3.4 | 0.9 | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 3.9 | 7.7 | 19.4 | 65.1 |
| 1999..... | 3.2 | 0.7 | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 3.6 | 7.0 | 18.0 | 62.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.4 | 0.8 | 1.5 | 3.2 | 5.0 | 9.9 | 24.9 | 68.2 |
| 2003..... | 3.8 | * | * | * | 0.4 | 0.8 | 1.6 | 2.6 | 5.1 | 9.5 | 25.2 | 72.2 |
| 2002..... | 3.7 | * | * | * | 0.4 | 0.6 | 1.6 | 2.8 | 4.8 | 10.8 | 23.8 | 66.8 |
| 2001..... | 3.5 | * | * | * | 0.3 | 0.7 | 1.6 | 2.7 | 4.8 | 10.2 | 22.3 | 67.4 |
| 2000..... | 3.5 | * | * | * | 0.4 | 0.7 | 1.3 | 2.6 | 5.5 | 9.8 | 22.5 | 69.2 |
| 1999..... | 3.2 | * | * | * | 0.3 | 0.6 | 1.3 | 2.3 | 5.1 | 9.5 | 22.0 | 64.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | 0.2 | 0.7 | 1.2 | 2.5 | 5.3 | 17.5 | 67.8 |
| 2003..... | 3.7 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 2.4 | 6.3 | 17.8 | 66.2 |
| 2002..... | 3.5 | * | * | * | * | 0.2 | 0.7 | 1.3 | 2.3 | 5.6 | 17.7 | 64.4 |
| 2001..... | 3.5 | * | * | * | * | 0.2 | 0.6 | 1.1 | 2.3 | 5.3 | 17.1 | 66.6 |
| 2000..... | 3.4 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.3 | 5.9 | 17.4 | 63.4 |
| 1999..... | 3.1 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.1 | 4.9 | 15.3 | 61.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | 0.2 | 0.4 | 1.0 | 2.0 | 3.5 | 6.9 | 20.0 | 68.2 |
| 2003..... | 4.0 | * | * | * | 0.2 | 0.4 | 1.0 | 1.7 | 3.5 | 7.4 | 20.8 | 67.6 |
| 2002..... | 3.8 | * | * | * | 0.2 | 0.3 | 1.0 | 1.7 | 3.2 | 7.6 | 20.1 | 64.6 |
| 2001..... | 3.6 | * | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 3.2 | 7.2 | 18.6 | 66.2 |
| 2000..... | 3.6 | * | * | * | 0.1 | 0.4 | 0.8 | 1.6 | 3.6 | 7.1 | 18.8 | 65.1 |
| 1999..... | 3.3 | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 3.1 | 6.6 | 17.4 | 62.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | 0.3 | 0.7 | 1.4 | 2.9 | 4.7 | 9.3 | 24.4 | 68.6 |
| 2003..... | 4.0 | * | * | * | 0.3 | 0.7 | 1.5 | 2.4 | 4.9 | 9.1 | 25.1 | 72.2 |
| 2002..... | 3.8 | * | * | * | 0.3 | 0.5 | 1.4 | 2.5 | 4.4 | 10.5 | 23.4 | 67.1 |
| 2001..... | 3.6 | * | * | * | 0.3 | 0.6 | 1.5 | 2.4 | 4.6 | 9.8 | 21.6 | 67.0 |
| 2000..... | 3.5 | * | * | * | 0.2 | 0.6 | 1.2 | 2.4 | 5.2 | 9.2 | 21.7 | 69.3 |
| 1999..... | 3.3 | * | * | * | 0.3 | 0.5 | 1.1 | 2.1 | 4.7 | 8.9 | 21.4 | 64.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | 0.2 | 0.6 | 1.0 | 2.3 | 4.8 | 17.1 | 68.0 |
| 2003..... | 3.9 | * | * | * | * | * | 0.5 | 1.0 | 2.2 | 5.9 | 17.8 | 65.6 |
| 2002..... | 3.8 | * | * | * | * | 0.2 | 0.6 | 1.0 | 2.1 | 5.3 | 17.8 | 63.5 |
| 2001..... | 3.7 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.0 | 4.9 | 16.7 | 65.8 |
| 2000..... | 3.6 | * | * | * | * | * | 0.4 | 0.8 | 2.1 | 5.4 | 17.0 | 63.5 |
| 1999..... | 3.3 | * | * | * | * | * | 0.4 | 0.7 | 1.7 | 4.6 | 14.9 | 60.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 0.4 | 1.0 | 2.0 | 3.9 | 7.0 | 13.8 | 30.7 | 77.4 |
| 2003..... | 3.6 | * | * | * | 0.6 | 1.0 | 2.0 | 4.0 | 6.6 | 13.1 | 25.3 | 81.5 |
| 2002..... | 3.6 | * | * | * | 0.5 | 0.8 | 1.9 | 4.6 | 6.8 | 12.8 | 25.1 | 81.4 |
| 2001..... | 3.6 | * | * | * | 0.4 | 1.0 | 2.1 | 4.2 | 6.4 | 12.1 | 27.2 | 84.9 |
| 2000..... | 3.6 | * | * | * | 0.7 | 0.9 | 2.0 | 3.8 | 7.3 | 14.8 | 28.7 | 74.7 |
| 1999..... | 3.4 | * | * | * | 0.4 | 0.9 | 2.1 | 3.5 | 8.0 | 12.4 | 25.6 | 69.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 1.7 | 2.6 | 5.4 | 9.5 | 19.0 | 40.2 | 76.2 |
| 2003..... | 3.8 | * | * | * | 0.9 | 1.3 | 2.9 | 5.1 | 9.1 | 15.9 | 32.9 | 82.0 |
| 2002..... | 3.9 | * | * | * | 0.8 | 1.3 | 2.5 | 6.0 | 9.5 | 17.1 | 33.1 | 77.7 |
| 2001..... | 3.7 | * | * | * | * | 1.5 | 2.7 | 5.2 | 8.5 | 15.6 | 33.0 | 85.5 |
| 2000..... | 4.0 | * | * | * | 1.1 | 1.3 | 2.6 | 5.5 | 10.2 | 19.5 | 36.4 | 81.0 |
| 1999..... | 3.7 | * | * | * | * | 1.5 | 2.9 | 4.5 | 10.8 | 17.7 | 32.2 | 73.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 1.4 | 2.7 | 4.9 | 10.2 | 25.3 | 77.9 |
| 2003..... | 3.4 | * | * | * | * | 0.8 | 1.1 | 3.0 | 4.5 | 11.1 | 21.0 | 81.3 |
| 2002..... | 3.3 | * | * | * | * | * | 1.3 | 3.3 | 4.6 | 9.7 | 20.6 | 82.8 |
| 2001..... | 3.5 | * | * | * | * | * | 1.6 | 3.2 | 4.6 | 9.6 | 23.9 | 84.7 |
| 2000..... | 3.3 | * | * | * | * | * | 1.5 | 2.3 | 5.0 | 11.6 | 24.3 | 72.4 |
| 1999..... | 3.1 | * | * | * | * | * | 1.4 | 2.7 | 5.7 | 8.6 | 21.9 | 68.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cerebrovascular diseases (I60-I69) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 51.1 | 3.1 | 0.3 | 0.2 | 0.5 | 1.4 | 5.4 | 14.9 | 34.3 | 107.8 | 386.2 | 1,245.9 |
| 2003..... | 54.2 | 2.5 | 0.3 | 0.2 | 0.5 | 1.5 | 5.5 | 15.0 | 35.6 | 112.9 | 410.7 | 1,370.1 |
| 2002..... | 56.4 | 2.9 | 0.3 | 0.2 | 0.4 | 1.4 | 5.4 | 15.1 | 37.2 | 120.3 | 431.0 | 1,445.9 |
| 2001..... | 57.4 | 2.7 | 0.4 | 0.2 | 0.5 | 1.5 | 5.5 | 15.1 | 38.0 | 123.4 | 443.9 | 1,500.2 |
| 2000..... | 59.6 | 3.3 | 0.3 | 0.2 | 0.5 | 1.5 | 5.8 | 16.0 | 41.0 | 128.6 | 461.3 | 1,589.2 |
| 1999..... | 60.0 | 2.7 | 0.3 | 0.2 | 0.5 | 1.4 | 5.7 | 15.2 | 40.6 | 130.8 | 469.8 | 1,614.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 40.7 | 3.4 | 0.3 | 0.2 | 0.5 | 1.4 | 5.6 | 16.7 | 39.5 | 121.1 | 402.9 | 1,118.1 |
| 2003..... | 42.9 | 2.8 | 0.3 | 0.2 | 0.5 | 1.6 | 5.8 | 16.7 | 40.8 | 127.8 | 431.4 | 1,236.0 |
| 2002..... | 44.2 | 3.2 | 0.4 | 0.2 | 0.5 | 1.4 | 5.3 | 16.7 | 42.7 | 135.0 | 445.9 | 1,317.9 |
| 2001..... | 45.2 | 3.1 | 0.3 | 0.2 | 0.5 | 1.6 | 5.7 | 16.7 | 43.4 | 140.4 | 467.3 | 1,380.2 |
| 2000..... | 46.9 | 3.8 | * | 0.2 | 0.5 | 1.5 | 5.8 | 17.5 | 47.2 | 145.0 | 490.8 | 1,484.3 |
| 1999..... | 47.1 | 3.3 | 0.3 | 0.2 | 0.5 | 1.5 | 5.8 | 16.7 | 46.5 | 147.3 | 500.4 | 1,512.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 61.2 | 2.8 | * | 0.2 | 0.5 | 1.4 | 5.1 | 13.1 | 29.5 | 96.6 | 374.9 | 1,303.4 |
| 2003..... | 65.1 | 2.2 | 0.3 | 0.1 | 0.5 | 1.4 | 5.3 | 13.4 | 30.9 | 100.5 | 396.8 | 1,429.4 |
| 2002..... | 68.2 | 2.5 | 0.3 | 0.2 | 0.3 | 1.4 | 5.5 | 13.6 | 32.1 | 108.1 | 421.2 | 1,501.5 |
| 2001..... | 69.2 | 2.3 | 0.4 | 0.2 | 0.5 | 1.5 | 5.4 | 13.6 | 32.9 | 109.3 | 428.6 | 1,550.4 |
| 2000..... | 71.8 | 2.7 | 0.4 | 0.2 | 0.5 | 1.5 | 5.7 | 14.5 | 35.3 | 115.1 | 442.1 | 1,632.0 |
| 1999..... | 72.3 | 2.1 | 0.3 | 0.2 | 0.5 | 1.4 | 5.6 | 13.8 | 35.1 | 117.2 | 449.8 | 1,656.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 16.4 | 2.2 | * | * | 0.5 | 1.2 | 4.9 | 14.6 | 34.9 | 97.0 | 293.1 | 813.0 |
| 2003..... | 16.7 | * | * | * | 0.4 | 1.2 | 4.5 | 14.9 | 35.3 | 98.4 | 316.8 | 880.5 |
| 2002..... | 16.6 | 2.4 | * | 0.3 | 0.4 | 1.3 | 4.4 | 15.3 | 35.9 | 102.9 | 319.9 | 898.0 |
| 2001..... | 17.4 | * | * | * | 0.6 | 1.4 | 5.0 | 15.9 | 35.4 | 106.2 | 341.2 | 1,042.9 |
| 2000..... | 17.5 | * | * | * | 0.5 | 1.7 | 5.2 | 15.2 | 39.8 | 113.2 | 344.1 | 1,082.9 |
| 1999..... | 17.4 | * | * | * | 0.5 | 1.6 | 4.9 | 15.7 | 37.7 | 116.9 | 348.3 | 1,077.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 15.0 | * | * | * | 0.5 | 1.5 | 5.9 | 17.5 | 42.9 | 114.4 | 323.3 | 778.9 |
| 2003..... | 14.9 | * | * | * | * | 1.3 | 5.2 | 18.1 | 43.5 | 113.9 | 337.1 | 837.4 |
| 2002..... | 15.0 | * | * | * | * | 1.5 | 5.1 | 18.6 | 45.0 | 124.6 | 338.5 | 856.7 |
| 2001..... | 15.7 | * | * | * | 0.7 | 1.6 | 5.3 | 18.7 | 43.5 | 127.2 | 386.3 | 1,005.6 |
| 2000..... | 15.8 | * | * | * | * | 1.9 | 5.5 | 18.1 | 48.8 | 136.1 | 392.9 | 1,029.9 |
| 1999..... | 16.1 | * | * | * | 0.6 | 1.8 | 5.7 | 19.5 | 46.1 | 140.8 | 395.2 | 1,136.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 17.9 | * | * | * | * | 0.9 | 3.9 | 11.8 | 27.7 | 83.0 | 272.2 | 830.4 |
| 2003..... | 18.6 | * | * | * | * | 1.1 | 3.7 | 11.7 | 27.8 | 86.0 | 302.8 | 902.3 |
| 2002..... | 18.4 | * | * | * | * | 1.0 | 3.7 | 12.0 | 27.6 | 85.6 | 307.2 | 918.5 |
| 2001..... | 19.1 | * | * | * | * | 1.2 | 4.7 | 13.1 | 28.2 | 89.6 | 310.7 | 1,061.2 |
| 2000..... | 19.4 | * | * | * | * | 1.4 | 4.9 | 12.4 | 31.9 | 95.2 | 311.3 | 1,108.9 |
| 1999..... | 18.8 | * | * | * | * | 1.4 | 4.0 | 11.9 | 30.3 | 98.1 | 317.1 | 1,049.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cerebrovascular diseases (I60-I69) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 56.7 | 3.3 | 0.3 | 0.2 | 0.5 | 1.5 | 5.4 | 14.8 | 34.1 | 108.4 | 390.8 | 1,263.8 |
| 2003..... | 60.1 | 2.6 | 0.3 | 0.2 | 0.6 | 1.5 | 5.7 | 15.0 | 35.6 | 113.7 | 415.0 | 1,388.4 |
| 2002..... | 62.4 | 3.0 | 0.3 | 0.2 | 0.4 | 1.5 | 5.5 | 15.0 | 37.2 | 121.3 | 435.5 | 1,465.0 |
| 2001..... | 63.2 | 2.8 | 0.3 | 0.2 | 0.5 | 1.5 | 5.6 | 14.9 | 38.0 | 124.3 | 447.7 | 1,514.0 |
| 2000..... | 65.4 | 3.6 | 0.3 | 0.2 | 0.5 | 1.5 | 5.8 | 15.9 | 40.9 | 129.2 | 465.0 | 1,604.5 |
| 1999..... | 65.7 | 2.9 | 0.3 | 0.2 | 0.5 | 1.4 | 5.8 | 15.1 | 40.6 | 131.2 | 473.8 | 1,630.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 45.0 | 3.7 | 0.3 | 0.2 | 0.5 | 1.4 | 5.6 | 16.5 | 39.0 | 121.4 | 406.7 | 1,133.7 |
| 2003..... | 47.6 | 2.8 | * | 0.2 | 0.6 | 1.6 | 5.9 | 16.5 | 40.4 | 128.4 | 435.7 | 1,252.9 |
| 2002..... | 48.9 | 3.5 | 0.4 | 0.2 | 0.5 | 1.4 | 5.3 | 16.4 | 42.3 | 135.4 | 450.1 | 1,336.0 |
| 2001..... | 49.7 | 3.3 | 0.3 | 0.2 | 0.5 | 1.5 | 5.7 | 16.4 | 43.2 | 140.9 | 469.8 | 1,392.9 |
| 2000..... | 51.5 | 4.2 | * | 0.2 | 0.5 | 1.5 | 5.8 | 17.3 | 46.8 | 145.1 | 493.7 | 1,499.9 |
| 1999..... | 51.5 | 3.7 | 0.3 | 0.2 | 0.5 | 1.4 | 5.8 | 16.4 | 46.3 | 147.3 | 503.6 | 1,524.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 67.8 | 3.0 | * | 0.2 | 0.5 | 1.5 | 5.2 | 13.2 | 29.5 | 97.4 | 380.0 | 1,321.9 |
| 2003..... | 72.0 | 2.4 | 0.3 | 0.1 | 0.5 | 1.4 | 5.5 | 13.5 | 31.0 | 101.4 | 401.1 | 1,448.0 |
| 2002..... | 75.3 | 2.5 | * | 0.2 | 0.3 | 1.5 | 5.8 | 13.7 | 32.4 | 109.5 | 425.8 | 1,520.7 |
| 2001..... | 76.1 | 2.4 | 0.4 | 0.2 | 0.5 | 1.5 | 5.5 | 13.5 | 33.2 | 110.4 | 433.2 | 1,564.4 |
| 2000..... | 78.7 | 2.8 | 0.4 | 0.2 | 0.5 | 1.5 | 5.8 | 14.6 | 35.5 | 116.1 | 446.4 | 1,646.8 |
| 1999..... | 79.2 | 2.0 | * | 0.2 | 0.5 | 1.4 | 5.8 | 13.9 | 35.4 | 118.1 | 454.4 | 1,672.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 60.6 | 2.6 | 0.3 | 0.2 | 0.4 | 1.1 | 4.2 | 10.9 | 27.8 | 98.7 | 381.8 | 1,276.4 |
| 2003..... | 64.2 | 2.0 | 0.2 | 0.1 | 0.5 | 1.2 | 4.1 | 10.9 | 28.7 | 103.6 | 406.6 | 1,401.1 |
| 2002..... | 67.0 | 2.2 | 0.3 | 0.2 | 0.4 | 1.2 | 4.1 | 11.1 | 30.8 | 110.8 | 427.5 | 1,479.2 |
| 2001..... | 67.6 | 1.9 | 0.3 | 0.2 | 0.4 | 1.3 | 4.2 | 10.8 | 31.2 | 113.0 | 439.5 | 1,525.0 |
| 2000..... | 70.0 | 2.6 | 0.3 | 0.2 | 0.5 | 1.2 | 4.3 | 12.0 | 34.2 | 119.3 | 455.7 | 1,616.8 |
| 1999..... | 70.4 | 2.4 | 0.3 | 0.2 | 0.4 | 1.1 | 4.3 | 11.2 | 33.4 | 120.7 | 465.1 | 1,648.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 47.0 | 2.9 | * | 0.2 | 0.5 | 1.1 | 4.4 | 12.1 | 31.1 | 110.0 | 396.9 | 1,145.3 |
| 2003..... | 49.7 | 1.9 | * | 0.2 | 0.5 | 1.3 | 4.2 | 12.1 | 32.1 | 116.9 | 426.0 | 1,264.2 |
| 2002..... | 51.3 | 2.7 | * | 0.2 | 0.5 | 1.1 | 4.0 | 12.1 | 34.5 | 123.2 | 441.1 | 1,345.9 |
| 2001..... | 52.0 | 2.2 | * | 0.2 | 0.4 | 1.3 | 4.3 | 11.9 | 35.1 | 128.0 | 460.5 | 1,399.0 |
| 2000..... | 53.9 | 3.3 | * | 0.2 | 0.5 | 1.2 | 4.3 | 13.0 | 38.7 | 133.1 | 482.3 | 1,505.9 |
| 1999..... | 53.9 | 3.1 | * | 0.2 | 0.4 | 1.1 | 4.4 | 12.0 | 38.0 | 135.0 | 492.8 | 1,535.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 73.7 | 2.2 | * | 0.2 | 0.4 | 1.2 | 4.0 | 9.8 | 24.7 | 89.0 | 371.6 | 1,335.1 |
| 2003..... | 78.2 | 2.1 | * | * | 0.4 | 1.1 | 4.0 | 9.7 | 25.5 | 92.1 | 393.6 | 1,461.3 |
| 2002..... | 82.1 | * | * | 0.2 | 0.3 | 1.2 | 4.3 | 10.1 | 27.2 | 100.2 | 418.4 | 1,536.7 |
| 2001..... | 82.6 | * | * | * | 0.3 | 1.3 | 4.1 | 9.8 | 27.4 | 100.3 | 425.6 | 1,577.4 |
| 2000..... | 85.5 | 1.9 | * | 0.2 | 0.4 | 1.2 | 4.3 | 10.9 | 29.9 | 107.6 | 438.3 | 1,661.6 |
| 1999..... | 86.1 | * | * | * | 0.4 | 1.1 | 4.3 | 10.5 | 29.1 | 108.8 | 447.0 | 1,693.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cerebrovascular diseases (I60-I69) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 48.7 | 6.2 | * | * | 0.8 | 3.3 | 12.7 | 40.0 | 86.0 | 199.4 | 519.1 | 1,226.1 |
| 2003..... | 51.1 | 4.9 | * | * | 1.0 | 3.2 | 14.2 | 42.0 | 91.3 | 204.9 | 539.1 | 1,366.0 |
| 2002..... | 51.7 | 5.6 | * | 0.4 | 0.6 | 3.2 | 13.3 | 41.6 | 89.7 | 218.7 | 558.8 | 1,398.0 |
| 2001..... | 52.9 | 6.9 | * | * | 0.9 | 3.0 | 13.6 | 43.5 | 92.8 | 226.0 | 567.2 | 1,465.5 |
| 2000..... | 54.3 | 8.0 | * | * | 0.7 | 3.0 | 14.9 | 44.3 | 95.2 | 226.3 | 593.8 | 1,545.8 |
| 1999..... | 53.9 | 4.7 | * | * | 0.7 | 2.9 | 14.7 | 42.2 | 97.7 | 235.7 | 596.4 | 1,501.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 43.1 | 7.1 | * | * | 1.0 | 3.1 | 13.3 | 46.0 | 110.2 | 240.1 | 558.7 | 1,077.1 |
| 2003..... | 44.9 | * | * | * | 0.8 | 3.5 | 14.9 | 48.1 | 114.1 | 241.8 | 597.6 | 1,200.7 |
| 2002..... | 45.1 | * | * | * | 0.7 | 2.9 | 12.7 | 47.5 | 112.7 | 267.3 | 595.2 | 1,266.8 |
| 2001..... | 46.2 | 8.3 | * | * | 0.7 | 3.1 | 13.5 | 49.9 | 113.9 | 273.5 | 620.5 | 1,359.4 |
| 2000..... | 47.7 | 9.2 | * | * | * | 2.8 | 15.1 | 50.4 | 116.9 | 272.1 | 664.9 | 1,479.5 |
| 1999..... | 47.5 | * | * | * | * | 2.9 | 15.0 | 49.6 | 119.5 | 279.1 | 656.7 | 1,455.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 53.8 | * | * | * | * | 3.5 | 12.2 | 34.8 | 66.6 | 170.4 | 496.7 | 1,285.0 |
| 2003..... | 56.7 | * | * | * | 1.1 | 3.0 | 13.7 | 36.8 | 73.0 | 178.6 | 506.0 | 1,430.0 |
| 2002..... | 57.7 | * | * | * | * | 3.5 | 13.8 | 36.6 | 71.3 | 184.2 | 538.2 | 1,448.1 |
| 2001..... | 59.0 | * | * | * | 1.1 | 2.9 | 13.6 | 38.1 | 75.9 | 192.5 | 537.2 | 1,504.8 |
| 2000..... | 60.3 | * | * | * | 0.8 | 3.2 | 14.8 | 39.2 | 77.8 | 194.1 | 553.8 | 1,570.4 |
| 1999..... | 59.8 | * | * | * | 0.8 | 2.8 | 14.4 | 35.9 | 80.2 | 205.1 | 562.7 | 1,518.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Subarachnoid hemorrhage (I60) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | 0.0 | 0.2 | 0.5 | 1.7 | 3.1 | 3.9 | 5.3 | 8.3 | 12.2 |
| 2003..... | 2.1 | * | * | * | 0.1 | 0.5 | 1.7 | 3.1 | 4.0 | 5.3 | 8.7 | 11.5 |
| 2002..... | 2.1 | * | * | 0.0 | 0.1 | 0.5 | 1.8 | 3.1 | 4.2 | 5.8 | 8.9 | 12.2 |
| 2001..... | 2.2 | * | * | * | 0.2 | 0.5 | 1.8 | 3.2 | 4.2 | 5.8 | 8.8 | 13.2 |
| 2000..... | 2.3 | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 3.5 | 4.9 | 5.8 | 9.4 | 14.1 |
| 1999..... | 2.3 | * | * | * | 0.2 | 0.6 | 2.1 | 3.5 | 4.7 | 6.2 | 9.4 | 13.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 0.2 | 0.5 | 1.3 | 2.7 | 3.0 | 3.6 | 6.0 | 9.3 |
| 2003..... | 1.5 | * | * | * | 0.2 | 0.5 | 1.4 | 2.4 | 3.1 | 4.0 | 6.2 | 9.1 |
| 2002..... | 1.5 | * | * | * | 0.2 | 0.4 | 1.4 | 2.3 | 3.3 | 4.4 | 5.8 | 9.1 |
| 2001..... | 1.5 | * | * | * | 0.2 | 0.5 | 1.4 | 2.6 | 3.2 | 4.3 | 6.4 | 9.5 |
| 2000..... | 1.6 | * | * | * | 0.2 | 0.6 | 1.5 | 2.7 | 3.5 | 4.2 | 6.8 | 11.2 |
| 1999..... | 1.6 | * | * | * | 0.2 | 0.6 | 1.8 | 2.8 | 3.4 | 4.3 | 6.7 | 9.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 0.2 | 0.5 | 2.0 | 3.5 | 4.8 | 6.7 | 9.9 | 13.5 |
| 2003..... | 2.6 | * | * | * | 0.1 | 0.5 | 1.9 | 3.7 | 4.8 | 6.5 | 10.4 | 12.5 |
| 2002..... | 2.8 | * | * | * | 0.1 | 0.6 | 2.1 | 3.8 | 5.1 | 6.9 | 11.0 | 13.6 |
| 2001..... | 2.8 | * | * | * | 0.2 | 0.5 | 2.1 | 3.7 | 5.2 | 7.0 | 10.3 | 14.8 |
| 2000..... | 3.0 | * | * | * | 0.2 | 0.6 | 2.3 | 4.3 | 6.2 | 7.0 | 11.2 | 15.3 |
| 1999..... | 3.0 | * | * | * | 0.2 | 0.6 | 2.4 | 4.1 | 5.9 | 7.7 | 11.1 | 14.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 0.5 | 1.7 | 3.2 | 4.7 | 7.9 | 9.8 | 17.1 | * |
| 2003..... | 1.4 | * | * | * | 0.5 | 1.5 | 4.0 | 5.2 | 6.8 | 9.1 | * | * |
| 2002..... | 1.3 | * | * | * | 0.6 | 1.6 | 3.3 | 5.0 | 6.6 | 9.3 | * | * |
| 2001..... | 1.4 | * | * | * | 0.6 | 1.6 | 3.6 | 5.4 | 7.0 | 9.7 | 16.3 | * |
| 2000..... | 1.5 | * | * | * | 0.8 | 1.8 | 4.0 | 5.0 | 6.8 | 11.1 | 15.9 | * |
| 1999..... | 1.6 | * | * | * | 0.6 | 2.0 | 4.1 | 5.1 | 7.3 | 12.5 | 23.2 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.7 | 1.6 | 3.1 | 4.1 | 6.2 | 8.2 | * | * |
| 2003..... | 1.1 | * | * | * | 0.5 | 1.5 | 3.4 | 4.3 | 5.5 | * | * | * |
| 2002..... | 1.1 | * | * | * | 0.6 | 1.7 | 3.0 | 4.4 | 4.8 | * | * | * |
| 2001..... | 1.1 | * | * | * | * | 1.3 | 3.1 | 4.3 | 5.4 | * | * | * |
| 2000..... | 1.1 | * | * | * | 0.9 | 1.5 | 2.9 | 4.1 | * | 10.8 | * | * |
| 1999..... | 1.2 | * | * | * | 0.8 | 2.0 | 3.9 | 4.3 | 4.4 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 1.9 | 3.4 | 5.3 | 9.2 | 10.9 | 19.4 | * |
| 2003..... | 1.6 | * | * | * | * | 1.5 | 4.5 | 5.9 | 7.8 | 11.9 | * | * |
| 2002..... | 1.6 | * | * | * | * | 1.5 | 3.5 | 5.5 | 8.1 | 11.8 | * | * |
| 2001..... | 1.8 | * | * | * | 0.6 | 1.8 | 4.1 | 6.3 | 8.3 | 11.5 | 20.5 | * |
| 2000..... | 1.9 | * | * | * | 0.7 | 2.1 | 5.0 | 5.7 | 9.0 | 11.2 | 19.8 | * |
| 1999..... | 1.9 | * | * | * | * | 2.0 | 4.3 | 5.9 | 9.6 | 15.6 | 32.2 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Subarachnoid hemorrhage (I60) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | 0.2 | 0.5 | 1.6 | 3.1 | 3.9 | 5.1 | 8.3 | 12.0 |
| 2003..... | 2.2 | * | * | * | 0.1 | 0.5 | 1.7 | 3.0 | 3.9 | 5.2 | 8.7 | 11.7 |
| 2002..... | 2.3 | * | * | * | 0.1 | 0.5 | 1.8 | 3.1 | 4.2 | 5.7 | 8.9 | 12.3 |
| 2001..... | 2.3 | * | * | * | 0.1 | 0.5 | 1.8 | 3.1 | 4.1 | 5.7 | 8.7 | 13.1 |
| 2000..... | 2.4 | * | * | * | 0.2 | 0.6 | 1.9 | 3.5 | 4.8 | 5.7 | 9.4 | 14.1 |
| 1999..... | 2.4 | * | * | * | 0.1 | 0.6 | 2.1 | 3.4 | 4.7 | 6.1 | 9.3 | 12.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 0.2 | 0.4 | 1.2 | 2.6 | 2.9 | 3.4 | 5.9 | 9.1 |
| 2003..... | 1.5 | * | * | * | 0.1 | 0.5 | 1.3 | 2.3 | 3.0 | 3.9 | 6.3 | 9.3 |
| 2002..... | 1.6 | * | * | * | 0.2 | 0.4 | 1.4 | 2.3 | 3.2 | 4.4 | 5.8 | 9.2 |
| 2001..... | 1.6 | * | * | * | 0.1 | 0.6 | 1.4 | 2.5 | 3.1 | 4.2 | 6.3 | 9.5 |
| 2000..... | 1.7 | * | * | * | 0.2 | 0.5 | 1.5 | 2.7 | 3.4 | 4.2 | 6.7 | 11.4 |
| 1999..... | 1.7 | * | * | * | 0.1 | 0.5 | 1.7 | 2.7 | 3.4 | 4.3 | 6.6 | 9.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | 0.2 | 0.5 | 2.1 | 3.5 | 4.8 | 6.5 | 9.9 | 13.2 |
| 2003..... | 2.8 | * | * | * | 0.1 | 0.5 | 2.0 | 3.6 | 4.7 | 6.3 | 10.3 | 12.7 |
| 2002..... | 2.9 | * | * | * | * | 0.6 | 2.2 | 3.8 | 5.0 | 6.8 | 10.9 | 13.6 |
| 2001..... | 2.9 | * | * | * | 0.2 | 0.5 | 2.1 | 3.7 | 5.1 | 6.9 | 10.3 | 14.5 |
| 2000..... | 3.1 | * | * | * | 0.2 | 0.6 | 2.3 | 4.2 | 6.2 | 6.9 | 11.1 | 15.1 |
| 1999..... | 3.1 | * | * | * | 0.2 | 0.6 | 2.5 | 4.1 | 5.9 | 7.6 | 11.0 | 14.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | 0.1 | 0.4 | 1.4 | 2.7 | 3.6 | 4.9 | 8.0 | 12.1 |
| 2003..... | 2.1 | * | * | * | 0.1 | 0.5 | 1.4 | 2.6 | 3.7 | 4.9 | 8.3 | 11.3 |
| 2002..... | 2.2 | * | * | * | 0.1 | 0.4 | 1.6 | 2.7 | 3.9 | 5.4 | 8.7 | 12.0 |
| 2001..... | 2.2 | * | * | * | 0.1 | 0.5 | 1.5 | 2.7 | 3.8 | 5.3 | 8.5 | 12.6 |
| 2000..... | 2.4 | * | * | * | 0.2 | 0.5 | 1.6 | 3.1 | 4.6 | 5.3 | 9.1 | 13.5 |
| 1999..... | 2.4 | * | * | * | 0.1 | 0.5 | 1.8 | 3.0 | 4.3 | 5.9 | 9.1 | 12.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.3 | 1.0 | 2.3 | 2.6 | 3.4 | 5.8 | 9.7 |
| 2003..... | 1.5 | * | * | * | 0.2 | 0.5 | 1.1 | 2.1 | 2.7 | 3.8 | 5.9 | 8.8 |
| 2002..... | 1.6 | * | * | * | 0.2 | 0.3 | 1.2 | 2.1 | 3.0 | 4.2 | 5.7 | 9.2 |
| 2001..... | 1.6 | * | * | * | * | 0.5 | 1.3 | 2.2 | 2.8 | 4.0 | 6.4 | 9.2 |
| 2000..... | 1.7 | * | * | * | 0.2 | 0.5 | 1.3 | 2.5 | 3.2 | 3.9 | 6.7 | 12.1 |
| 1999..... | 1.7 | * | * | * | * | 0.5 | 1.6 | 2.4 | 3.2 | 4.1 | 6.8 | 9.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 0.4 | 1.8 | 3.1 | 4.5 | 6.2 | 9.5 | 13.2 |
| 2003..... | 2.7 | * | * | * | * | 0.5 | 1.6 | 3.1 | 4.6 | 5.9 | 9.9 | 12.4 |
| 2002..... | 2.9 | * | * | * | * | 0.5 | 1.9 | 3.3 | 4.7 | 6.4 | 10.6 | 13.2 |
| 2001..... | 2.8 | * | * | * | * | 0.4 | 1.7 | 3.2 | 4.8 | 6.3 | 10.0 | 14.0 |
| 2000..... | 3.1 | * | * | * | * | 0.5 | 1.9 | 3.7 | 6.0 | 6.4 | 10.7 | 14.1 |
| 1999..... | 3.1 | * | * | * | 0.2 | 0.5 | 2.0 | 3.5 | 5.4 | 7.5 | 10.6 | 13.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Subarachnoid hemorrhage (I60) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 1.0 | 3.1 | 5.1 | 6.0 | 5.3 | 9.1 | 9.2 |
| 2003..... | 2.4 | * | * | * | * | 0.9 | 3.3 | 5.0 | 5.2 | 6.1 | 9.9 | 14.9 |
| 2002..... | 2.4 | * | * | * | * | 1.0 | 3.3 | 5.6 | 5.9 | 6.4 | 8.7 | 13.0 |
| 2001..... | 2.6 | * | * | * | * | 0.9 | 3.5 | 5.9 | 6.2 | 7.4 | 8.6 | 14.0 |
| 2000..... | 2.7 | * | * | * | * | 0.9 | 3.7 | 6.2 | 6.2 | 8.0 | 10.8 | 17.5 |
| 1999..... | 2.7 | * | * | * | * | 1.0 | 3.9 | 6.2 | 6.8 | 7.0 | 9.3 | 14.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 2.4 | 4.2 | 5.3 | 3.7 | 5.7 | * |
| 2003..... | 1.7 | * | * | * | * | * | 2.6 | 3.3 | 5.4 | 3.9 | 7.5 | * |
| 2002..... | 1.6 | * | * | * | * | * | 2.2 | 3.3 | 5.3 | 6.1 | 5.9 | * |
| 2001..... | 1.8 | * | * | * | * | 0.9 | 2.6 | 4.7 | 5.1 | 4.9 | 6.3 | * |
| 2000..... | 1.7 | * | * | * | * | * | 2.7 | 4.4 | 4.2 | 6.3 | 7.1 | * |
| 1999..... | 1.8 | * | * | * | * | 0.8 | 2.8 | 5.0 | 5.0 | 5.4 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 1.1 | 3.7 | 5.8 | 6.6 | 6.4 | 10.9 | 12.0 |
| 2003..... | 3.0 | * | * | * | * | 1.1 | 4.0 | 6.4 | 5.0 | 7.7 | 11.2 | 15.3 |
| 2002..... | 3.2 | * | * | * | * | 1.3 | 4.2 | 7.5 | 6.4 | 6.5 | 10.3 | 15.9 |
| 2001..... | 3.3 | * | * | * | * | 0.9 | 4.3 | 6.9 | 7.2 | 9.2 | 9.9 | 16.7 |
| 2000..... | 3.6 | * | * | * | * | 1.3 | 4.6 | 7.8 | 7.8 | 9.3 | 12.9 | 23.1 |
| 1999..... | 3.4 | * | * | * | * | 1.1 | 4.8 | 7.3 | 8.2 | 8.1 | 11.6 | 18.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intracerebral and other intracranial hemorrhage (I61-I62) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.4 | 1.3 | * | * | 0.2 | 0.5 | 2.1 | 6.2 | 11.6 | 27.1 | 71.3 | 124.3 |
| 2003..... | 9.4 | 0.8 | * | 0.1 | 0.2 | 0.5 | 2.2 | 6.3 | 11.4 | 27.2 | 71.1 | 126.6 |
| 2002..... | 9.4 | 1.0 | * | 0.1 | 0.2 | 0.5 | 2.1 | 6.0 | 11.7 | 28.1 | 72.3 | 127.2 |
| 2001..... | 9.4 | 1.0 | * | 0.1 | 0.2 | 0.6 | 2.3 | 6.2 | 11.8 | 27.9 | 71.7 | 127.9 |
| 2000..... | 9.7 | 1.1 | * | 0.1 | 0.2 | 0.5 | 2.3 | 6.5 | 12.9 | 29.3 | 74.1 | 137.9 |
| 1999..... | 9.1 | 1.1 | * | 0.1 | 0.2 | 0.5 | 2.2 | 5.9 | 12.3 | 28.3 | 71.9 | 122.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 8.9 | 1.4 | * | * | 0.2 | 0.6 | 2.6 | 7.4 | 13.9 | 30.7 | 77.4 | 134.4 |
| 2003..... | 8.9 | * | * | * | 0.2 | 0.5 | 2.5 | 7.6 | 13.3 | 31.1 | 77.8 | 137.4 |
| 2002..... | 8.7 | 1.1 | * | * | 0.2 | 0.5 | 2.3 | 7.3 | 13.7 | 32.3 | 77.2 | 135.0 |
| 2001..... | 8.8 | 1.2 | * | * | 0.3 | 0.6 | 2.6 | 7.5 | 13.9 | 32.0 | 78.6 | 134.2 |
| 2000..... | 9.1 | 1.3 | * | * | 0.2 | 0.6 | 2.7 | 7.8 | 15.9 | 33.1 | 79.0 | 149.6 |
| 1999..... | 8.5 | 1.3 | * | * | 0.2 | 0.5 | 2.6 | 7.0 | 14.7 | 32.0 | 75.9 | 130.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 9.8 | 1.3 | * | * | 0.2 | 0.5 | 1.6 | 5.1 | 9.4 | 24.2 | 67.1 | 119.8 |
| 2003..... | 9.8 | * | * | * | 0.2 | 0.5 | 1.9 | 5.1 | 9.6 | 24.0 | 66.6 | 121.8 |
| 2002..... | 10.0 | 1.0 | * | * | 0.1 | 0.4 | 1.8 | 4.8 | 9.8 | 24.6 | 69.0 | 123.8 |
| 2001..... | 9.9 | * | * | * | 0.2 | 0.5 | 1.9 | 5.1 | 10.0 | 24.6 | 67.1 | 125.2 |
| 2000..... | 10.4 | * | * | * | 0.2 | 0.4 | 2.0 | 5.3 | 10.2 | 26.1 | 70.9 | 133.1 |
| 1999..... | 9.8 | * | * | * | 0.1 | 0.4 | 1.8 | 4.9 | 10.1 | 25.2 | 69.2 | 118.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | 0.4 | 2.1 | 7.1 | 14.2 | 26.8 | 56.4 | 95.6 |
| 2003..... | 3.8 | * | * | * | * | 0.4 | 1.9 | 7.0 | 12.5 | 25.4 | 52.6 | 91.7 |
| 2002..... | 3.9 | * | * | * | * | 0.4 | 1.8 | 6.9 | 12.5 | 24.8 | 60.6 | 98.7 |
| 2001..... | 4.0 | * | * | * | * | 0.6 | 2.3 | 7.2 | 12.5 | 25.4 | 60.1 | 107.5 |
| 2000..... | 4.0 | * | * | * | * | 0.6 | 2.3 | 6.5 | 15.6 | 26.8 | 61.8 | 106.8 |
| 1999..... | 4.0 | * | * | * | * | 0.6 | 2.0 | 7.4 | 13.8 | 29.5 | 63.3 | 98.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | * | 2.9 | 8.8 | 19.2 | 31.1 | 60.1 | 109.3 |
| 2003..... | 3.9 | * | * | * | * | * | 2.5 | 10.0 | 16.2 | 27.7 | 51.8 | 104.7 |
| 2002..... | 4.0 | * | * | * | * | * | 2.3 | 9.7 | 16.5 | 30.4 | 60.3 | 90.4 |
| 2001..... | 4.2 | * | * | * | * | 0.9 | 2.9 | 9.5 | 16.5 | 29.0 | 65.7 | 112.4 |
| 2000..... | 4.2 | * | * | * | * | 0.8 | 3.1 | 9.0 | 21.8 | 31.7 | 62.0 | 112.9 |
| 1999..... | 4.4 | * | * | * | * | 0.7 | 2.7 | 10.3 | 19.0 | 37.3 | 67.8 | 107.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | * | 1.2 | 5.3 | 9.6 | 23.4 | 53.8 | 88.6 |
| 2003..... | 3.6 | * | * | * | * | * | 1.2 | 3.9 | 9.1 | 23.5 | 53.2 | 85.2 |
| 2002..... | 3.8 | * | * | * | * | * | 1.3 | 4.0 | 8.9 | 20.4 | 60.7 | 102.8 |
| 2001..... | 3.8 | * | * | * | * | * | 1.6 | 5.0 | 9.0 | 22.7 | 56.2 | 105.2 |
| 2000..... | 3.7 | * | * | * | * | * | 1.5 | 4.0 | 10.0 | 22.9 | 61.7 | 103.9 |
| 1999..... | 3.6 | * | * | * | * | * | 1.3 | 4.5 | 9.2 | 23.4 | 60.4 | 93.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intracerebral and other intracranial hemorrhage (I61-I62) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 10.2 | 1.5 | * | * | 0.2 | 0.5 | 2.1 | 6.1 | 11.3 | 27.1 | 72.0 | 125.5 |
| 2003..... | 10.2 | 0.9 | * | 0.1 | 0.2 | 0.5 | 2.3 | 6.2 | 11.3 | 27.3 | 72.0 | 127.9 |
| 2002..... | 10.2 | 0.9 | * | 0.1 | 0.1 | 0.5 | 2.1 | 5.9 | 11.6 | 28.3 | 72.7 | 128.1 |
| 2001..... | 10.1 | 0.9 | * | 0.1 | 0.2 | 0.6 | 2.3 | 6.1 | 11.7 | 28.0 | 72.1 | 128.3 |
| 2000..... | 10.5 | 1.2 | * | 0.1 | 0.2 | 0.5 | 2.3 | 6.5 | 12.7 | 29.4 | 74.4 | 138.9 |
| 1999..... | 9.8 | 0.9 | * | * | 0.2 | 0.4 | 2.2 | 5.7 | 12.1 | 28.1 | 72.1 | 122.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 9.7 | 1.5 | * | * | 0.2 | 0.6 | 2.5 | 7.2 | 13.4 | 30.6 | 78.3 | 135.7 |
| 2003..... | 9.7 | * | * | * | 0.2 | 0.6 | 2.5 | 7.3 | 13.1 | 31.2 | 79.1 | 138.6 |
| 2002..... | 9.5 | * | * | * | 0.2 | 0.6 | 2.3 | 7.0 | 13.5 | 32.3 | 77.9 | 136.6 |
| 2001..... | 9.5 | * | * | * | 0.2 | 0.6 | 2.5 | 7.2 | 13.6 | 32.2 | 79.2 | 134.9 |
| 2000..... | 9.8 | 1.5 | * | * | 0.2 | 0.6 | 2.6 | 7.7 | 15.3 | 33.1 | 79.5 | 151.0 |
| 1999..... | 9.0 | * | * | * | 0.2 | 0.5 | 2.5 | 6.6 | 14.3 | 31.6 | 76.3 | 131.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 10.7 | 1.5 | * | * | 0.2 | 0.5 | 1.6 | 5.0 | 9.4 | 24.2 | 67.7 | 120.9 |
| 2003..... | 10.8 | * | * | * | 0.2 | 0.5 | 2.0 | 5.2 | 9.6 | 24.0 | 67.2 | 123.1 |
| 2002..... | 10.8 | * | * | * | * | 0.5 | 1.9 | 4.9 | 9.9 | 24.9 | 69.2 | 124.4 |
| 2001..... | 10.7 | * | * | * | 0.2 | 0.6 | 2.0 | 5.0 | 10.0 | 24.6 | 67.5 | 125.6 |
| 2000..... | 11.2 | * | * | * | 0.2 | 0.4 | 2.1 | 5.4 | 10.2 | 26.3 | 71.1 | 134.0 |
| 1999..... | 10.5 | * | * | * | 0.1 | 0.4 | 1.8 | 4.9 | 10.1 | 25.2 | 69.4 | 119.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 10.4 | 1.1 | * | * | 0.2 | 0.4 | 1.5 | 4.0 | 9.0 | 25.5 | 71.7 | 127.7 |
| 2003..... | 10.3 | * | * | * | 0.2 | 0.4 | 1.5 | 3.9 | 8.7 | 25.5 | 71.6 | 129.0 |
| 2002..... | 10.4 | * | * | * | 0.1 | 0.4 | 1.3 | 3.8 | 9.6 | 26.6 | 72.8 | 130.3 |
| 2001..... | 10.3 | * | * | * | 0.2 | 0.4 | 1.5 | 3.9 | 9.5 | 26.1 | 72.1 | 129.3 |
| 2000..... | 10.7 | * | * | * | 0.2 | 0.4 | 1.5 | 4.4 | 10.3 | 28.0 | 74.2 | 140.4 |
| 1999..... | 10.0 | * | * | * | 0.1 | 0.3 | 1.4 | 3.7 | 9.8 | 26.9 | 71.9 | 123.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 9.7 | * | * | * | 0.2 | 0.5 | 1.8 | 4.8 | 10.4 | 28.6 | 78.4 | 138.8 |
| 2003..... | 9.5 | * | * | * | 0.2 | 0.4 | 1.7 | 4.6 | 9.9 | 29.2 | 79.0 | 140.1 |
| 2002..... | 9.5 | * | * | * | 0.2 | 0.4 | 1.4 | 4.5 | 11.0 | 30.5 | 78.6 | 138.7 |
| 2001..... | 9.4 | * | * | * | 0.2 | 0.4 | 1.7 | 4.6 | 11.2 | 29.9 | 79.2 | 135.6 |
| 2000..... | 9.7 | * | * | * | 0.2 | 0.4 | 1.6 | 5.2 | 12.2 | 31.2 | 79.5 | 151.7 |
| 1999..... | 8.9 | * | * | * | * | 0.3 | 1.6 | 4.3 | 11.6 | 30.4 | 75.5 | 132.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 11.1 | * | * | * | * | 0.3 | 1.1 | 3.1 | 7.7 | 22.8 | 67.1 | 122.8 |
| 2003..... | 11.1 | * | * | * | * | 0.4 | 1.3 | 3.1 | 7.6 | 22.4 | 66.7 | 124.1 |
| 2002..... | 11.3 | * | * | * | * | 0.3 | 1.2 | 3.2 | 8.2 | 23.3 | 68.9 | 126.7 |
| 2001..... | 11.1 | * | * | * | * | 0.4 | 1.4 | 3.1 | 8.0 | 22.8 | 67.4 | 126.7 |
| 2000..... | 11.7 | * | * | * | * | 0.3 | 1.3 | 3.5 | 8.5 | 25.3 | 70.8 | 135.8 |
| 1999..... | 11.0 | * | * | * | * | 0.3 | 1.2 | 3.2 | 8.1 | 23.9 | 69.5 | 120.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intracerebral and other intracranial hemorrhage (I61-I62) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 10.6 | * | * | * | 0.3 | 1.4 | 5.7 | 19.6 | 29.0 | 40.2 | 73.0 | 96.4 |
| 2003..... | 10.9 | * | * | * | 0.4 | 1.2 | 6.3 | 21.1 | 30.6 | 38.6 | 73.8 | 108.6 |
| 2002..... | 10.0 | * | * | * | * | 1.2 | 6.2 | 19.3 | 26.3 | 39.7 | 69.4 | 97.4 |
| 2001..... | 10.4 | * | * | * | 0.4 | 1.4 | 6.1 | 21.1 | 27.2 | 40.7 | 69.5 | 106.8 |
| 2000..... | 10.6 | * | * | * | * | 1.2 | 7.3 | 21.5 | 29.9 | 39.4 | 69.4 | 118.0 |
| 1999..... | 9.8 | * | * | * | * | 1.1 | 7.0 | 19.1 | 28.3 | 37.2 | 68.9 | 96.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 11.1 | * | * | * | * | 1.4 | 6.9 | 23.3 | 38.9 | 47.6 | 77.0 | 101.6 |
| 2003..... | 11.4 | * | * | * | * | 1.5 | 6.9 | 25.8 | 39.3 | 45.4 | 85.7 | 118.3 |
| 2002..... | 10.4 | * | * | * | * | 1.4 | 7.0 | 24.5 | 33.8 | 48.2 | 68.6 | 101.7 |
| 2001..... | 10.7 | * | * | * | * | 1.5 | 6.9 | 25.9 | 32.8 | 50.3 | 72.3 | 113.2 |
| 2000..... | 11.2 | * | * | * | * | 1.5 | 8.5 | 26.2 | 40.8 | 47.2 | 73.7 | 132.7 |
| 1999..... | 10.1 | * | * | * | * | 1.3 | 8.4 | 23.6 | 35.5 | 40.7 | 74.7 | 108.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 10.1 | * | * | * | * | 1.4 | 4.6 | 16.4 | 21.0 | 34.9 | 70.7 | 94.4 |
| 2003..... | 10.4 | * | * | * | * | 1.0 | 5.8 | 17.1 | 23.5 | 33.7 | 67.1 | 104.8 |
| 2002..... | 9.7 | * | * | * | * | 1.1 | 5.5 | 14.9 | 20.4 | 33.8 | 69.8 | 95.7 |
| 2001..... | 10.1 | * | * | * | * | 1.3 | 5.4 | 17.0 | 22.7 | 33.9 | 67.9 | 104.4 |
| 2000..... | 10.1 | * | * | * | * | 1.0 | 6.2 | 17.4 | 21.2 | 33.9 | 67.0 | 112.5 |
| 1999..... | 9.5 | * | * | * | * | 0.9 | 5.8 | 15.2 | 22.5 | 34.8 | 65.7 | 92.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cerebral infarction (I63) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 7.7 | 27.8 | 97.7 |
| 2003..... | 4.2 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.4 | 8.4 | 31.3 | 112.2 |
| 2002..... | 4.3 | * | * | * | * | 0.1 | 0.2 | 1.0 | 2.5 | 8.6 | 32.2 | 118.3 |
| 2001..... | 4.4 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.4 | 9.4 | 33.9 | 124.8 |
| 2000..... | 4.9 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.9 | 9.6 | 38.1 | 141.0 |
| 1999..... | 5.3 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.1 | 11.3 | 41.2 | 153.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.0 | 9.0 | 30.1 | 88.5 |
| 2003..... | 3.3 | * | * | * | * | * | 0.4 | 1.2 | 2.9 | 10.2 | 33.1 | 100.3 |
| 2002..... | 3.4 | * | * | * | * | 0.1 | 0.2 | 1.2 | 3.1 | 9.6 | 35.4 | 108.7 |
| 2001..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.1 | 11.1 | 37.8 | 111.2 |
| 2000..... | 3.9 | * | * | * | * | * | 0.4 | 1.1 | 3.6 | 11.4 | 42.3 | 133.3 |
| 1999..... | 4.2 | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.8 | 13.5 | 45.0 | 144.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.7 | 6.6 | 26.3 | 101.9 |
| 2003..... | 5.0 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 6.9 | 30.1 | 117.5 |
| 2002..... | 5.1 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.9 | 7.7 | 30.1 | 122.4 |
| 2001..... | 5.3 | * | * | * | * | * | 0.3 | 0.8 | 1.8 | 8.0 | 31.3 | 130.4 |
| 2000..... | 5.8 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 8.1 | 35.3 | 144.2 |
| 1999..... | 6.3 | * | * | * | * | * | 0.3 | 0.7 | 2.5 | 9.6 | 38.8 | 156.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 0.6 | 1.9 | 5.4 | 19.3 | 55.2 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.6 | 1.8 | 6.6 | 20.3 | 53.4 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.8 | 1.6 | 7.1 | 19.8 | 68.7 |
| 2001..... | 1.2 | * | * | * | * | * | * | 0.8 | 1.8 | 8.2 | 23.3 | 88.8 |
| 2000..... | 1.3 | * | * | * | * | * | * | 0.8 | 2.2 | 8.5 | 26.7 | 84.9 |
| 1999..... | 1.3 | * | * | * | * | * | * | 0.7 | 2.0 | 9.6 | 26.2 | 100.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 2.6 | 5.5 | 19.0 | 54.6 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 2.5 | 7.9 | 22.6 | 30.8 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | 2.8 | 7.5 | 20.1 | 71.9 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | 2.6 | 12.2 | 25.5 | 64.8 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 3.4 | 10.6 | 32.9 | 92.7 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | 2.6 | 11.9 | 27.1 | 105.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | * | * | 5.3 | 19.5 | 55.5 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | * | 5.6 | 18.7 | 64.8 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | * | 6.8 | 19.7 | 67.1 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | * | 5.0 | 21.8 | 100.5 |
| 2000..... | 1.3 | * | * | * | * | * | * | * | * | 7.0 | 22.5 | 81.1 |
| 1999..... | 1.5 | * | * | * | * | * | * | * | * | 7.7 | 25.7 | 97.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cerebral infarction (I63) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.8 | 28.2 | 99.5 |
| 2003..... | 4.6 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.4 | 8.5 | 31.8 | 114.5 |
| 2002..... | 4.8 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.5 | 8.7 | 32.8 | 120.1 |
| 2001..... | 4.9 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.5 | 9.4 | 34.2 | 126.0 |
| 2000..... | 5.4 | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.0 | 9.6 | 38.4 | 142.7 |
| 1999..... | 5.8 | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.2 | 11.4 | 41.8 | 154.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 9.2 | 30.6 | 90.2 |
| 2003..... | 3.7 | * | * | * | * | * | 0.4 | 1.2 | 3.0 | 10.3 | 33.6 | 103.6 |
| 2002..... | 3.8 | * | * | * | * | 0.1 | 0.2 | 1.3 | 3.1 | 9.7 | 36.1 | 110.1 |
| 2001..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.1 | 11.0 | 38.2 | 113.2 |
| 2000..... | 4.3 | * | * | * | * | * | 0.4 | 1.1 | 3.7 | 11.3 | 42.6 | 134.6 |
| 1999..... | 4.7 | * | * | * | * | * | 0.4 | 1.2 | 3.8 | 13.5 | 45.6 | 145.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | 0.3 | 0.6 | 1.7 | 6.7 | 26.6 | 103.7 |
| 2003..... | 5.5 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 7.0 | 30.6 | 119.3 |
| 2002..... | 5.7 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 7.8 | 30.6 | 124.4 |
| 2001..... | 5.9 | * | * | * | * | * | 0.3 | 0.8 | 1.9 | 8.1 | 31.7 | 131.3 |
| 2000..... | 6.4 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 8.2 | 35.7 | 146.0 |
| 1999..... | 7.0 | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 9.7 | 39.3 | 158.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 7.3 | 27.6 | 99.5 |
| 2003..... | 5.1 | * | * | * | * | * | 0.2 | 0.8 | 2.0 | 7.9 | 31.4 | 116.1 |
| 2002..... | 5.2 | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.3 | 8.1 | 32.2 | 121.0 |
| 2001..... | 5.3 | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.1 | 8.8 | 33.8 | 127.2 |
| 2000..... | 5.8 | * | * | * | * | * | 0.3 | 0.8 | 2.5 | 8.9 | 37.7 | 143.4 |
| 1999..... | 6.3 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.7 | 10.5 | 41.1 | 155.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.3 | 0.8 | 2.5 | 8.4 | 29.9 | 89.8 |
| 2003..... | 4.0 | * | * | * | * | * | 0.3 | 1.0 | 2.5 | 9.5 | 33.3 | 104.9 |
| 2002..... | 4.1 | * | * | * | * | * | 0.2 | 1.1 | 2.7 | 9.0 | 35.5 | 110.0 |
| 2001..... | 4.2 | * | * | * | * | * | 0.3 | 0.9 | 2.7 | 10.4 | 37.3 | 112.5 |
| 2000..... | 4.5 | * | * | * | * | * | 0.3 | 0.9 | 3.1 | 10.5 | 41.4 | 134.2 |
| 1999..... | 4.9 | * | * | * | * | * | 0.4 | 1.0 | 3.3 | 12.5 | 44.7 | 145.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 6.3 | 26.1 | 103.9 |
| 2003..... | 6.1 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 6.6 | 30.1 | 121.1 |
| 2002..... | 6.3 | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 7.2 | 29.9 | 125.8 |
| 2001..... | 6.5 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 7.4 | 31.4 | 133.3 |
| 2000..... | 7.1 | * | * | * | * | * | 0.2 | 0.7 | 2.0 | 7.6 | 35.3 | 147.1 |
| 1999..... | 7.7 | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 8.9 | 38.8 | 160.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cerebral infarction (I63) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | * | 0.7 | 1.8 | 5.6 | 14.1 | 38.2 | 105.1 |
| 2003..... | 3.6 | * | * | * | * | * | 0.8 | 2.1 | 6.0 | 14.5 | 41.0 | 109.2 |
| 2002..... | 3.6 | * | * | * | * | * | 0.5 | 2.2 | 5.0 | 15.2 | 42.7 | 113.7 |
| 2001..... | 3.8 | * | * | * | * | * | 0.7 | 2.2 | 5.8 | 16.7 | 42.8 | 123.0 |
| 2000..... | 4.2 | * | * | * | * | * | 0.8 | 2.1 | 6.6 | 16.9 | 49.6 | 144.1 |
| 1999..... | 4.5 | * | * | * | * | * | 0.7 | 2.2 | 6.7 | 20.8 | 53.6 | 152.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | * | 2.5 | 7.9 | 17.8 | 44.2 | 99.6 |
| 2003..... | 3.2 | * | * | * | * | * | 0.9 | 2.5 | 8.0 | 18.6 | 44.2 | 99.1 |
| 2002..... | 3.2 | * | * | * | * | * | * | 2.8 | 6.7 | 18.4 | 46.9 | 114.9 |
| 2001..... | 3.4 | * | * | * | * | * | * | 2.6 | 7.5 | 19.1 | 54.6 | 129.4 |
| 2000..... | 3.8 | * | * | * | * | * | 0.8 | 2.9 | 8.5 | 21.1 | 60.2 | 139.7 |
| 1999..... | 4.1 | * | * | * | * | * | * | 3.0 | 8.4 | 26.5 | 63.8 | 162.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | * | 0.7 | 1.2 | 3.8 | 11.4 | 34.8 | 107.2 |
| 2003..... | 4.0 | * | * | * | * | * | 0.7 | 1.8 | 4.5 | 11.5 | 39.3 | 113.1 |
| 2002..... | 4.0 | * | * | * | * | * | * | 1.6 | 3.7 | 13.0 | 40.3 | 113.3 |
| 2001..... | 4.1 | * | * | * | * | * | 0.8 | 1.8 | 4.5 | 15.1 | 36.2 | 120.6 |
| 2000..... | 4.7 | * | * | * | * | * | 0.8 | 1.5 | 5.0 | 14.0 | 43.6 | 145.7 |
| 1999..... | 4.9 | * | * | * | * | * | 0.8 | 1.6 | 5.4 | 16.8 | 48.0 | 147.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Stroke, not specified as hemorrhage or infarction (I64) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 26.0 | * | * | * | * | 0.2 | 0.9 | 3.7 | 12.7 | 51.7 | 204.5 | 713.4 |
| 2003..... | 28.4 | * | * | * | 0.1 | 0.3 | 1.0 | 3.7 | 14.1 | 54.8 | 223.0 | 799.7 |
| 2002..... | 30.1 | * | * | * | * | 0.2 | 0.9 | 4.0 | 15.0 | 60.5 | 237.6 | 859.4 |
| 2001..... | 31.1 | * | * | * | * | 0.2 | 0.9 | 3.8 | 15.6 | 62.8 | 249.4 | 906.2 |
| 2000..... | 32.1 | * | * | * | * | 0.2 | 0.9 | 3.8 | 16.2 | 66.4 | 257.6 | 954.7 |
| 1999..... | 32.6 | * | * | * | * | 0.2 | 0.8 | 3.8 | 16.3 | 66.5 | 263.9 | 982.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 19.7 | * | * | * | * | 0.2 | 1.1 | 4.4 | 14.9 | 58.9 | 209.0 | 615.5 |
| 2003..... | 21.5 | * | * | * | * | 0.3 | 1.2 | 4.4 | 16.9 | 62.5 | 231.4 | 699.0 |
| 2002..... | 22.6 | * | * | * | * | 0.2 | 1.0 | 4.6 | 18.0 | 68.8 | 241.0 | 762.8 |
| 2001..... | 23.5 | * | * | * | * | 0.1 | 0.9 | 4.5 | 18.7 | 72.1 | 256.3 | 819.7 |
| 2000..... | 24.2 | * | * | * | * | 0.2 | 0.9 | 4.7 | 19.4 | 75.1 | 270.6 | 863.2 |
| 1999..... | 24.7 | * | * | * | * | 0.2 | 0.8 | 4.6 | 19.4 | 75.9 | 280.4 | 896.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 32.2 | * | * | * | * | 0.2 | 0.8 | 2.9 | 10.6 | 45.7 | 201.4 | 757.4 |
| 2003..... | 35.0 | * | * | * | * | 0.2 | 0.8 | 3.0 | 11.5 | 48.3 | 217.4 | 844.2 |
| 2002..... | 37.4 | * | * | * | * | 0.2 | 0.9 | 3.3 | 12.1 | 53.6 | 235.4 | 901.3 |
| 2001..... | 38.5 | * | * | * | * | 0.2 | 0.8 | 3.0 | 12.7 | 55.1 | 244.9 | 942.4 |
| 2000..... | 39.7 | * | * | * | * | 0.2 | 0.8 | 3.0 | 13.1 | 59.2 | 249.0 | 992.0 |
| 1999..... | 40.3 | * | * | * | * | 0.2 | 0.7 | 3.0 | 13.5 | 58.9 | 253.3 | 1,017.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | * | * | * | 0.6 | 3.0 | 11.0 | 44.2 | 151.8 | 437.2 |
| 2003..... | 7.7 | * | * | * | * | * | 0.7 | 2.6 | 12.6 | 45.3 | 171.6 | 517.7 |
| 2002..... | 7.6 | * | * | * | * | * | 0.5 | 3.3 | 13.0 | 49.2 | 171.5 | 508.9 |
| 2001..... | 7.9 | * | * | * | * | * | 0.6 | 3.6 | 12.0 | 49.3 | 183.4 | 607.8 |
| 2000..... | 8.1 | * | * | * | * | * | 0.6 | 2.8 | 13.5 | 56.9 | 179.6 | 650.9 |
| 1999..... | 7.9 | * | * | * | * | * | 0.5 | 2.7 | 13.3 | 53.8 | 181.2 | 646.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.3 | * | * | * | * | * | 0.7 | 4.3 | 13.2 | 55.0 | 173.5 | 409.1 |
| 2003..... | 6.6 | * | * | * | * | * | 0.7 | 3.1 | 16.4 | 54.3 | 195.2 | 483.3 |
| 2002..... | 6.6 | * | * | * | * | * | * | 3.8 | 16.5 | 61.2 | 186.7 | 480.2 |
| 2001..... | 6.9 | * | * | * | * | * | * | 4.5 | 14.6 | 59.1 | 210.2 | 622.8 |
| 2000..... | 7.0 | * | * | * | * | * | * | 3.9 | 15.3 | 69.8 | 213.9 | 608.7 |
| 1999..... | 7.0 | * | * | * | * | * | * | 3.6 | 16.1 | 65.2 | 212.2 | 683.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 8.0 | * | * | * | * | * | * | 1.7 | 9.0 | 35.6 | 136.8 | 451.6 |
| 2003..... | 8.8 | * | * | * | * | * | * | 2.1 | 9.3 | 38.1 | 155.4 | 535.2 |
| 2002..... | 8.7 | * | * | * | * | * | * | 2.8 | 9.9 | 39.7 | 161.0 | 523.1 |
| 2001..... | 9.0 | * | * | * | * | * | * | 2.8 | 9.6 | 41.6 | 165.3 | 600.4 |
| 2000..... | 9.3 | * | * | * | * | * | * | 1.8 | 11.9 | 46.8 | 156.5 | 671.7 |
| 1999..... | 8.9 | * | * | * | * | * | * | 1.7 | 10.8 | 44.7 | 160.6 | 628.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Stroke, not specified as hemorrhage or infarction (I64) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 29.1 | * | * | * | * | 0.2 | 1.0 | 3.7 | 12.7 | 52.2 | 207.1 | 724.7 |
| 2003..... | 31.6 | * | * | * | 0.1 | 0.3 | 1.1 | 3.8 | 14.2 | 55.3 | 225.3 | 810.4 |
| 2002..... | 33.5 | * | * | * | * | 0.2 | 1.0 | 4.0 | 15.1 | 61.1 | 240.4 | 871.6 |
| 2001..... | 34.5 | * | * | * | * | 0.2 | 0.9 | 3.7 | 15.8 | 63.6 | 251.9 | 915.4 |
| 2000..... | 35.5 | * | * | * | * | 0.2 | 0.9 | 3.9 | 16.3 | 66.8 | 260.2 | 964.0 |
| 1999..... | 36.0 | * | * | * | * | 0.2 | 0.8 | 3.9 | 16.5 | 67.1 | 266.8 | 992.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 21.9 | * | * | * | * | 0.2 | 1.1 | 4.4 | 15.0 | 59.0 | 210.7 | 624.6 |
| 2003..... | 24.0 | * | * | * | * | 0.3 | 1.3 | 4.5 | 16.9 | 62.9 | 232.9 | 708.3 |
| 2002..... | 25.2 | * | * | * | * | 0.2 | 1.0 | 4.6 | 18.1 | 69.1 | 243.0 | 774.2 |
| 2001..... | 26.0 | * | * | * | * | * | 1.0 | 4.5 | 18.9 | 72.8 | 257.7 | 826.2 |
| 2000..... | 26.8 | * | * | * | * | 0.3 | 1.0 | 4.7 | 19.7 | 75.3 | 272.3 | 871.9 |
| 1999..... | 27.2 | * | * | * | * | 0.2 | 0.9 | 4.7 | 19.5 | 76.3 | 282.5 | 903.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 35.9 | * | * | * | * | 0.3 | 0.9 | 3.1 | 10.6 | 46.4 | 204.7 | 769.5 |
| 2003..... | 38.9 | * | * | * | * | 0.2 | 0.9 | 3.1 | 11.7 | 49.0 | 220.2 | 855.2 |
| 2002..... | 41.5 | * | * | * | * | 0.3 | 0.9 | 3.4 | 12.3 | 54.4 | 238.6 | 913.6 |
| 2001..... | 42.6 | * | * | * | * | 0.2 | 0.8 | 3.0 | 12.9 | 55.9 | 248.1 | 952.5 |
| 2000..... | 43.8 | * | * | * | * | 0.2 | 0.8 | 3.1 | 13.2 | 59.8 | 252.3 | 1,001.2 |
| 1999..... | 44.3 | * | * | * | * | 0.2 | 0.7 | 3.1 | 13.6 | 59.6 | 256.6 | 1,027.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 31.5 | * | * | * | * | 0.2 | 0.7 | 2.7 | 10.1 | 46.6 | 201.5 | 733.1 |
| 2003..... | 34.2 | * | * | * | * | 0.2 | 0.7 | 2.8 | 11.3 | 49.7 | 220.1 | 818.1 |
| 2002..... | 36.4 | * | * | * | * | 0.2 | 0.7 | 2.9 | 11.9 | 54.9 | 235.0 | 879.7 |
| 2001..... | 37.3 | * | * | * | * | 0.2 | 0.7 | 2.7 | 12.5 | 57.0 | 246.4 | 922.8 |
| 2000..... | 38.4 | * | * | * | * | 0.2 | 0.7 | 2.9 | 13.2 | 60.7 | 254.2 | 972.8 |
| 1999..... | 38.9 | * | * | * | * | 0.1 | 0.6 | 2.8 | 13.1 | 60.7 | 261.2 | 1,003.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 23.2 | * | * | * | * | * | 0.8 | 3.3 | 11.9 | 52.5 | 204.6 | 632.1 |
| 2003..... | 25.4 | * | * | * | * | 0.2 | 0.9 | 3.4 | 13.4 | 56.5 | 226.8 | 713.5 |
| 2002..... | 26.6 | * | * | * | * | 0.2 | 0.8 | 3.5 | 14.2 | 61.5 | 236.6 | 778.0 |
| 2001..... | 27.4 | * | * | * | * | * | 0.8 | 3.3 | 14.8 | 64.8 | 251.1 | 829.8 |
| 2000..... | 28.3 | * | * | * | * | 0.2 | 0.8 | 3.6 | 16.1 | 68.1 | 264.9 | 873.6 |
| 1999..... | 28.8 | * | * | * | * | * | 0.6 | 3.4 | 15.7 | 68.9 | 276.0 | 910.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 39.5 | * | * | * | * | 0.2 | 0.6 | 2.1 | 8.5 | 41.5 | 199.4 | 778.3 |
| 2003..... | 42.8 | * | * | * | * | 0.2 | 0.6 | 2.2 | 9.2 | 43.8 | 215.5 | 864.2 |
| 2002..... | 45.7 | * | * | * | * | 0.2 | 0.6 | 2.3 | 9.8 | 49.3 | 233.9 | 923.6 |
| 2001..... | 46.8 | * | * | * | * | 0.2 | 0.5 | 2.1 | 10.4 | 50.3 | 243.3 | 961.5 |
| 2000..... | 48.1 | * | * | * | * | 0.2 | 0.6 | 2.2 | 10.5 | 54.5 | 247.3 | 1,012.8 |
| 1999..... | 48.7 | * | * | * | * | 0.2 | 0.5 | 2.2 | 10.7 | 53.8 | 251.5 | 1,041.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Stroke, not specified as hemorrhage or infarction (I64) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 23.9 | * | * | * | * | 0.6 | 2.8 | 10.9 | 35.3 | 106.7 | 294.3 | 713.2 |
| 2003..... | 25.6 | * | * | * | * | 0.8 | 3.2 | 10.9 | 39.3 | 111.1 | 309.6 | 808.2 |
| 2002..... | 26.9 | * | * | * | * | 0.4 | 2.6 | 11.7 | 42.5 | 122.7 | 329.5 | 851.1 |
| 2001..... | 27.5 | * | * | * | * | 0.4 | 2.6 | 11.3 | 43.4 | 128.5 | 337.4 | 891.5 |
| 2000..... | 28.2 | * | * | * | * | 0.5 | 2.3 | 11.5 | 42.7 | 131.5 | 354.7 | 929.5 |
| 1999..... | 28.4 | * | * | * | * | 0.4 | 2.3 | 12.1 | 45.5 | 134.2 | 354.4 | 923.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 19.9 | * | * | * | * | * | 2.9 | 12.7 | 44.9 | 127.8 | 308.0 | 595.4 |
| 2003..... | 21.4 | * | * | * | * | 1.1 | 3.9 | 13.0 | 48.8 | 130.4 | 337.3 | 694.7 |
| 2002..... | 22.9 | * | * | * | * | * | 2.4 | 13.6 | 54.5 | 153.6 | 349.6 | 774.5 |
| 2001..... | 23.5 | * | * | * | * | * | 2.7 | 13.4 | 56.5 | 158.5 | 367.3 | 824.7 |
| 2000..... | 24.0 | * | * | * | * | * | 2.3 | 14.0 | 52.3 | 157.4 | 395.8 | 911.2 |
| 1999..... | 24.1 | * | * | * | * | * | 2.4 | 14.7 | 56.2 | 158.3 | 389.6 | 885.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 27.6 | * | * | * | * | * | 2.6 | 9.2 | 27.6 | 91.6 | 286.6 | 759.8 |
| 2003..... | 29.3 | * | * | * | * | * | 2.6 | 9.1 | 31.6 | 97.3 | 294.0 | 852.2 |
| 2002..... | 30.6 | * | * | * | * | * | 2.8 | 10.0 | 32.8 | 100.7 | 318.2 | 880.4 |
| 2001..... | 31.2 | * | * | * | * | * | 2.5 | 9.5 | 32.9 | 107.2 | 320.6 | 916.2 |
| 2000..... | 32.0 | * | * | * | * | * | 2.4 | 9.4 | 34.9 | 113.2 | 331.5 | 936.3 |
| 1999..... | 32.3 | * | * | * | * | * | 2.2 | 9.9 | 36.9 | 117.2 | 334.8 | 937.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other cerebrovascular diseases and their sequelae (I67,I69) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.9 | 1.4 | 0.2 | 0.1 | 0.1 | 0.1 | 0.4 | 1.1 | 3.8 | 16.0 | 74.3 | 298.3 |
| 2003..... | 10.3 | 1.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 1.0 | 3.8 | 17.2 | 76.6 | 320.1 |
| 2002..... | 10.5 | 1.4 | 0.2 | 0.1 | 0.0 | 0.1 | 0.4 | 1.1 | 3.8 | 17.4 | 80.0 | 328.9 |
| 2001..... | 10.3 | 1.2 | 0.2 | 0.1 | * | 0.1 | 0.3 | 1.0 | 3.9 | 17.5 | 80.2 | 328.2 |
| 2000..... | 10.5 | 1.6 | 0.2 | * | 0.1 | 0.1 | 0.3 | 1.1 | 4.1 | 17.6 | 82.1 | 341.4 |
| 1999..... | 10.6 | 1.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 1.1 | 4.1 | 18.5 | 83.3 | 343.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.6 | 1.5 | * | * | * | 0.1 | 0.4 | 1.1 | 4.6 | 19.0 | 80.4 | 270.4 |
| 2003..... | 7.8 | 1.2 | * | * | * | 0.1 | 0.3 | 1.1 | 4.6 | 20.0 | 82.9 | 290.2 |
| 2002..... | 7.9 | 1.7 | 0.3 | 0.1 | * | 0.1 | 0.3 | 1.2 | 4.5 | 20.0 | 86.4 | 302.3 |
| 2001..... | 7.9 | 1.5 | * | * | * | 0.1 | 0.3 | 1.1 | 4.6 | 20.8 | 88.2 | 305.5 |
| 2000..... | 8.1 | 1.8 | * | * | * | * | 0.3 | 1.2 | 4.7 | 21.2 | 92.0 | 326.9 |
| 1999..... | 8.1 | 1.4 | * | * | * | * | 0.3 | 1.1 | 5.2 | 21.6 | 92.4 | 332.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 12.1 | 1.4 | * | * | * | 0.1 | 0.3 | 1.0 | 3.0 | 13.4 | 70.2 | 310.8 |
| 2003..... | 12.7 | 1.2 | * | * | * | * | 0.4 | 0.9 | 3.1 | 14.8 | 72.4 | 333.3 |
| 2002..... | 13.0 | 1.1 | * | * | * | * | 0.4 | 1.0 | 3.2 | 15.3 | 75.7 | 340.4 |
| 2001..... | 12.7 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.2 | 14.7 | 74.9 | 337.6 |
| 2000..... | 12.8 | 1.4 | * | * | * | 0.1 | 0.3 | 1.1 | 3.6 | 14.6 | 75.6 | 347.4 |
| 1999..... | 12.9 | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.2 | 15.9 | 77.4 | 348.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | * | 0.8 | 3.1 | 12.7 | 55.8 | 207.9 |
| 2003..... | 2.8 | * | * | * | * | * | * | 0.8 | 3.2 | 14.3 | 63.2 | 211.4 |
| 2002..... | 2.8 | * | * | * | * | * | * | 1.1 | 3.7 | 15.2 | 58.7 | 212.3 |
| 2001..... | 2.8 | * | * | * | * | * | * | 0.6 | 3.8 | 16.2 | 64.7 | 222.6 |
| 2000..... | 2.7 | * | * | * | * | * | * | 1.1 | 3.6 | 14.2 | 65.0 | 224.3 |
| 1999..... | 2.6 | * | * | * | * | * | * | 0.8 | 3.4 | 16.7 | 65.0 | 209.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | * | * | 3.8 | 16.6 | 62.6 | 193.3 |
| 2003..... | 2.3 | * | * | * | * | * | * | * | 4.1 | 18.5 | 62.5 | 214.0 |
| 2002..... | 2.5 | * | * | * | * | * | * | 1.4 | 4.9 | 20.8 | 65.7 | 210.8 |
| 2001..... | 2.4 | * | * | * | * | * | * | * | 5.5 | 21.6 | 77.8 | 198.1 |
| 2000..... | 2.3 | * | * | * | * | * | * | 1.4 | 4.2 | 20.0 | 73.3 | 207.6 |
| 1999..... | 2.4 | * | * | * | * | * | * | * | 4.0 | 22.0 | 80.3 | 235.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | * | * | 2.4 | 9.5 | 51.1 | 215.4 |
| 2003..... | 3.3 | * | * | * | * | * | * | * | 2.3 | 11.0 | 63.7 | 210.2 |
| 2002..... | 3.1 | * | * | * | * | * | * | * | 2.7 | 10.6 | 54.0 | 213.0 |
| 2001..... | 3.1 | * | * | * | * | * | * | * | 2.2 | 12.0 | 55.9 | 234.6 |
| 2000..... | 3.1 | * | * | * | * | * | * | * | 3.1 | 9.6 | 59.4 | 232.5 |
| 1999..... | 2.8 | * | * | * | * | * | * | * | 2.9 | 12.6 | 54.8 | 197.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other cerebrovascular diseases and their sequelae (I67,I69) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 11.0 | 1.5 | 0.2 | 0.1 | 0.1 | 0.1 | 0.4 | 1.1 | 3.8 | 16.2 | 75.2 | 302.1 |
| 2003..... | 11.4 | 1.1 | * | * | 0.1 | 0.1 | 0.3 | 1.1 | 3.8 | 17.3 | 77.2 | 324.0 |
| 2002..... | 11.7 | 1.6 | 0.2 | 0.1 | 0.1 | 0.1 | 0.4 | 1.1 | 3.8 | 17.5 | 80.9 | 333.0 |
| 2001..... | 11.4 | 1.4 | 0.2 | * | * | 0.1 | 0.3 | 1.0 | 3.9 | 17.5 | 80.7 | 331.2 |
| 2000..... | 11.6 | 1.9 | 0.2 | * | 0.1 | 0.1 | 0.3 | 1.1 | 4.2 | 17.7 | 82.6 | 344.9 |
| 1999..... | 11.7 | 1.2 | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.1 | 4.2 | 18.5 | 83.8 | 348.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 8.5 | 1.8 | * | * | * | * | 0.4 | 1.2 | 4.6 | 19.1 | 81.3 | 274.1 |
| 2003..... | 8.7 | * | * | * | * | 0.1 | 0.3 | 1.1 | 4.6 | 20.1 | 83.8 | 293.1 |
| 2002..... | 8.8 | 2.0 | * | * | * | 0.1 | 0.3 | 1.2 | 4.4 | 19.9 | 87.3 | 305.9 |
| 2001..... | 8.7 | 1.6 | * | * | * | 0.1 | 0.4 | 1.1 | 4.5 | 20.8 | 88.4 | 309.0 |
| 2000..... | 9.0 | 2.2 | * | * | * | * | 0.4 | 1.1 | 4.8 | 21.2 | 92.7 | 331.0 |
| 1999..... | 9.0 | 1.7 | * | * | * | * | 0.3 | 1.2 | 5.3 | 21.6 | 92.5 | 335.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 13.4 | 1.3 | * | * | * | 0.1 | 0.4 | 1.0 | 3.1 | 13.7 | 71.1 | 314.5 |
| 2003..... | 14.1 | * | * | * | * | * | 0.4 | 1.0 | 3.2 | 15.0 | 72.8 | 337.6 |
| 2002..... | 14.4 | 1.3 | * | * | * | * | 0.5 | 1.0 | 3.2 | 15.6 | 76.6 | 344.6 |
| 2001..... | 14.0 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.3 | 14.9 | 75.7 | 340.5 |
| 2000..... | 14.1 | 1.6 | * | * | * | 0.1 | 0.3 | 1.1 | 3.6 | 14.9 | 76.1 | 350.5 |
| 1999..... | 14.2 | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.2 | 16.0 | 78.2 | 353.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 12.0 | 1.1 | * | 0.1 | * | 0.1 | 0.4 | 0.9 | 3.1 | 14.5 | 73.0 | 304.0 |
| 2003..... | 12.5 | * | * | * | * | 0.1 | 0.3 | 0.8 | 3.1 | 15.5 | 75.3 | 326.5 |
| 2002..... | 12.8 | 0.9 | * | * | * | 0.1 | 0.3 | 0.8 | 3.1 | 15.9 | 78.9 | 336.1 |
| 2001..... | 12.5 | 1.0 | * | * | * | 0.1 | 0.3 | 0.8 | 3.2 | 15.9 | 78.7 | 333.1 |
| 2000..... | 12.6 | 1.4 | * | * | * | 0.1 | 0.3 | 0.9 | 3.5 | 16.4 | 80.4 | 346.8 |
| 1999..... | 12.7 | 1.1 | * | * | * | 0.1 | 0.3 | 0.9 | 3.4 | 16.7 | 81.8 | 352.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 9.0 | * | * | * | * | * | 0.4 | 0.8 | 3.7 | 17.0 | 78.2 | 274.9 |
| 2003..... | 9.3 | * | * | * | * | * | 0.3 | 0.8 | 3.6 | 17.9 | 80.9 | 296.9 |
| 2002..... | 9.5 | * | * | * | * | * | 0.3 | 0.9 | 3.6 | 18.1 | 84.7 | 310.0 |
| 2001..... | 9.4 | * | * | * | * | * | 0.3 | 0.8 | 3.7 | 18.9 | 86.5 | 311.8 |
| 2000..... | 9.6 | 2.0 | * | * | * | * | 0.3 | 0.9 | 4.1 | 19.3 | 89.9 | 334.4 |
| 1999..... | 9.5 | * | * | * | * | * | 0.2 | 0.9 | 4.2 | 19.1 | 89.9 | 338.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 14.9 | * | * | * | * | * | 0.3 | 0.9 | 2.5 | 12.3 | 69.5 | 317.0 |
| 2003..... | 15.6 | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 13.4 | 71.4 | 339.6 |
| 2002..... | 15.9 | * | * | * | * | * | 0.4 | 0.7 | 2.7 | 14.0 | 75.0 | 347.4 |
| 2001..... | 15.4 | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 13.4 | 73.6 | 341.9 |
| 2000..... | 15.5 | * | * | * | * | * | 0.2 | 0.8 | 2.9 | 13.9 | 74.2 | 351.8 |
| 1999..... | 15.8 | * | * | * | * | * | 0.4 | 0.9 | 2.7 | 14.7 | 76.5 | 358.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other cerebrovascular diseases and their sequelae (I67,I69) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 8.4 | 3.2 | * | * | * | * | 0.5 | 2.6 | 10.1 | 33.2 | 104.6 | 302.2 |
| 2003..... | 8.7 | * | * | * | * | * | 0.6 | 2.8 | 10.2 | 34.7 | 104.7 | 325.1 |
| 2002..... | 8.7 | 3.6 | * | * | * | * | 0.7 | 2.9 | 10.0 | 34.7 | 108.5 | 322.7 |
| 2001..... | 8.6 | 3.3 | * | * | * | * | 0.7 | 3.0 | 10.2 | 32.7 | 108.9 | 330.2 |
| 2000..... | 8.5 | 4.2 | * | * | * | * | 0.8 | 2.9 | 9.8 | 30.5 | 109.3 | 336.8 |
| 1999..... | 8.5 | * | * | * | * | * | 0.8 | 2.6 | 10.4 | 36.6 | 110.1 | 314.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 7.3 | * | * | * | * | * | * | 3.2 | 13.2 | 43.2 | 123.8 | 278.4 |
| 2003..... | 7.2 | * | * | * | * | * | * | 3.5 | 12.6 | 43.5 | 122.9 | 274.9 |
| 2002..... | 7.0 | * | * | * | * | * | * | 3.2 | 12.4 | 41.0 | 124.3 | 270.2 |
| 2001..... | 6.8 | * | * | * | * | * | 0.8 | 3.2 | 12.0 | 40.6 | 120.0 | 285.3 |
| 2000..... | 6.8 | * | * | * | * | * | 0.8 | 2.8 | 11.0 | 40.1 | 128.0 | 293.5 |
| 1999..... | 7.3 | * | * | * | * | * | 0.8 | 3.4 | 14.3 | 48.2 | 123.2 | 294.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 9.4 | * | * | * | * | * | * | 2.1 | 7.6 | 26.0 | 93.7 | 311.6 |
| 2003..... | 10.0 | * | * | * | * | * | 0.7 | 2.2 | 8.3 | 28.4 | 94.4 | 344.6 |
| 2002..... | 10.2 | * | * | * | * | * | 0.8 | 2.6 | 8.1 | 30.2 | 99.6 | 342.8 |
| 2001..... | 10.2 | * | * | * | * | * | * | 2.9 | 8.7 | 27.1 | 102.6 | 346.9 |
| 2000..... | 10.0 | * | * | * | * | * | 0.7 | 3.1 | 8.9 | 23.7 | 98.8 | 352.8 |
| 1999..... | 9.7 | * | * | * | * | * | 0.8 | 1.9 | 7.2 | 28.4 | 102.8 | 322.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of arteries, arterioles and capillaries (I70-I78) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 12.1 | * | * | * | 0.2 | 0.4 | 1.1 | 3.1 | 8.8 | 31.2 | 90.6 | 276.3 |
| 2003..... | 13.0 | * | * | * | 0.2 | 0.4 | 1.1 | 3.4 | 9.8 | 33.5 | 97.4 | 300.6 |
| 2002..... | 13.4 | * | * | * | 0.2 | 0.4 | 1.1 | 3.1 | 10.4 | 36.2 | 100.6 | 316.5 |
| 2001..... | 13.8 | * | * | * | 0.2 | 0.4 | 1.2 | 3.1 | 10.5 | 36.5 | 104.9 | 337.3 |
| 2000..... | 14.4 | * | * | * | 0.2 | 0.4 | 1.1 | 3.2 | 11.1 | 40.0 | 109.2 | 352.0 |
| 1999..... | 14.6 | * | * | * | 0.1 | 0.4 | 1.2 | 3.2 | 11.3 | 39.5 | 111.6 | 367.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 11.6 | * | * | * | 0.2 | 0.6 | 1.5 | 4.3 | 11.9 | 40.8 | 112.5 | 294.7 |
| 2003..... | 12.6 | * | * | * | 0.2 | 0.5 | 1.5 | 4.7 | 13.5 | 44.8 | 125.5 | 318.3 |
| 2002..... | 13.0 | * | * | * | 0.2 | 0.5 | 1.4 | 4.3 | 14.3 | 48.6 | 127.8 | 344.0 |
| 2001..... | 13.3 | * | * | * | 0.2 | 0.6 | 1.6 | 4.2 | 14.3 | 49.3 | 133.0 | 364.5 |
| 2000..... | 14.1 | * | * | * | 0.2 | 0.5 | 1.4 | 4.4 | 15.7 | 55.1 | 142.5 | 391.4 |
| 1999..... | 14.1 | * | * | * | 0.2 | 0.5 | 1.5 | 4.3 | 16.0 | 54.0 | 144.2 | 396.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 12.6 | * | * | * | * | 0.3 | 0.8 | 1.9 | 6.0 | 23.2 | 75.8 | 268.0 |
| 2003..... | 13.3 | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 6.4 | 24.1 | 78.6 | 292.8 |
| 2002..... | 13.8 | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 6.8 | 25.8 | 82.5 | 304.6 |
| 2001..... | 14.4 | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 7.1 | 25.8 | 86.4 | 325.9 |
| 2000..... | 14.6 | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 6.8 | 27.5 | 87.5 | 336.0 |
| 1999..... | 15.2 | * | * | * | * | 0.2 | 0.9 | 2.0 | 7.0 | 27.7 | 90.5 | 355.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 0.3 | 0.7 | 2.1 | 6.2 | 18.3 | 53.0 | 151.7 |
| 2003..... | 3.2 | * | * | * | * | 0.3 | 0.7 | 1.8 | 5.7 | 18.7 | 61.2 | 192.3 |
| 2002..... | 3.3 | * | * | * | * | 0.3 | 0.8 | 1.9 | 6.9 | 19.1 | 63.7 | 201.2 |
| 2001..... | 3.2 | * | * | * | * | * | 0.7 | 2.6 | 5.8 | 17.8 | 62.0 | 225.7 |
| 2000..... | 3.5 | * | * | * | * | 0.4 | 0.8 | 2.1 | 6.7 | 26.0 | 67.0 | 235.6 |
| 1999..... | 3.4 | * | * | * | * | * | 0.7 | 1.8 | 7.3 | 21.6 | 71.0 | 239.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 1.0 | 3.1 | 8.3 | 25.6 | 67.2 | 177.9 |
| 2003..... | 3.4 | * | * | * | * | * | 1.0 | 2.4 | 9.1 | 25.1 | 82.8 | 234.0 |
| 2002..... | 3.2 | * | * | * | * | * | 1.1 | 2.6 | 8.3 | 25.0 | 83.3 | 204.1 |
| 2001..... | 3.1 | * | * | * | * | * | 0.9 | 3.3 | 7.6 | 24.8 | 80.5 | 236.2 |
| 2000..... | 3.4 | * | * | * | * | * | 0.9 | 2.6 | 9.0 | 34.0 | 93.9 | 235.8 |
| 1999..... | 3.4 | * | * | * | * | * | 0.9 | 2.2 | 9.8 | 29.0 | 93.3 | 314.9 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | * | 1.1 | 4.2 | 12.3 | 43.2 | 138.3 |
| 2003..... | 2.9 | * | * | * | * | * | * | 1.2 | 2.7 | 13.5 | 46.3 | 171.1 |
| 2002..... | 3.3 | * | * | * | * | * | * | 1.3 | 5.7 | 14.3 | 50.3 | 199.8 |
| 2001..... | 3.2 | * | * | * | * | * | * | 1.9 | 4.2 | 12.3 | 49.6 | 220.6 |
| 2000..... | 3.5 | * | * | * | * | * | * | 1.6 | 4.7 | 19.7 | 48.9 | 235.4 |
| 1999..... | 3.3 | * | * | * | * | * | * | 1.3 | 5.1 | 15.8 | 56.2 | 203.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of arteries, arterioles and capillaries (I70-I78) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 13.6 | * | * | * | 0.2 | 0.5 | 1.2 | 3.2 | 9.0 | 32.1 | 92.5 | 281.5 |
| 2003..... | 14.5 | * | * | * | 0.2 | 0.4 | 1.2 | 3.6 | 10.1 | 34.5 | 98.9 | 304.6 |
| 2002..... | 15.0 | * | * | * | 0.2 | 0.4 | 1.1 | 3.2 | 10.7 | 37.3 | 102.1 | 320.4 |
| 2001..... | 15.4 | * | * | * | 0.2 | 0.5 | 1.3 | 3.1 | 10.9 | 37.6 | 106.5 | 340.5 |
| 2000..... | 15.9 | * | * | * | 0.2 | 0.4 | 1.1 | 3.3 | 11.4 | 40.7 | 110.8 | 355.4 |
| 1999..... | 16.1 | * | * | * | 0.1 | 0.4 | 1.2 | 3.3 | 11.6 | 40.5 | 113.1 | 370.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 13.1 | * | * | * | 0.2 | 0.6 | 1.6 | 4.4 | 12.1 | 41.8 | 114.9 | 300.2 |
| 2003..... | 14.2 | * | * | * | 0.2 | 0.6 | 1.6 | 5.0 | 13.8 | 46.0 | 127.4 | 321.7 |
| 2002..... | 14.6 | * | * | * | 0.2 | 0.6 | 1.5 | 4.5 | 14.7 | 50.1 | 129.7 | 349.5 |
| 2001..... | 14.8 | * | * | * | 0.2 | 0.6 | 1.7 | 4.3 | 14.8 | 50.7 | 135.0 | 368.8 |
| 2000..... | 15.6 | * | * | * | 0.2 | 0.6 | 1.4 | 4.6 | 16.1 | 56.2 | 144.3 | 396.6 |
| 1999..... | 15.6 | * | * | * | 0.2 | 0.6 | 1.5 | 4.5 | 16.4 | 55.2 | 146.0 | 398.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 14.1 | * | * | * | * | 0.3 | 0.8 | 1.9 | 6.1 | 24.0 | 77.5 | 273.2 |
| 2003..... | 14.8 | * | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 6.7 | 24.8 | 80.0 | 297.1 |
| 2002..... | 15.3 | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 6.9 | 26.6 | 83.8 | 307.9 |
| 2001..... | 15.9 | * | * | * | * | 0.3 | 0.8 | 2.0 | 7.3 | 26.7 | 87.9 | 328.8 |
| 2000..... | 16.1 | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 7.0 | 28.0 | 88.9 | 338.7 |
| 1999..... | 16.7 | * | * | * | * | 0.2 | 0.9 | 2.1 | 7.2 | 28.4 | 91.7 | 359.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 15.2 | * | * | * | 0.2 | 0.4 | 1.0 | 2.8 | 8.5 | 32.4 | 93.8 | 287.0 |
| 2003..... | 16.2 | * | * | * | 0.2 | 0.4 | 1.0 | 3.1 | 9.8 | 34.9 | 100.2 | 310.5 |
| 2002..... | 16.7 | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 10.2 | 37.7 | 103.4 | 327.9 |
| 2001..... | 17.1 | * | * | * | 0.2 | 0.4 | 1.1 | 2.8 | 10.4 | 38.0 | 107.6 | 347.6 |
| 2000..... | 17.7 | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 10.9 | 41.3 | 111.8 | 361.6 |
| 1999..... | 18.0 | * | * | * | 0.1 | 0.4 | 1.1 | 2.9 | 11.4 | 41.1 | 114.4 | 376.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 14.6 | * | * | * | 0.2 | 0.5 | 1.4 | 4.0 | 11.4 | 42.6 | 117.2 | 309.8 |
| 2003..... | 15.7 | * | * | * | 0.2 | 0.5 | 1.4 | 4.4 | 13.5 | 46.8 | 128.8 | 331.0 |
| 2002..... | 16.3 | * | * | * | 0.2 | 0.5 | 1.3 | 3.9 | 14.1 | 51.0 | 132.2 | 360.3 |
| 2001..... | 16.4 | * | * | * | 0.2 | 0.6 | 1.4 | 3.7 | 14.3 | 51.6 | 136.9 | 379.8 |
| 2000..... | 17.4 | * | * | * | 0.2 | 0.5 | 1.2 | 4.0 | 15.7 | 57.6 | 146.9 | 406.3 |
| 1999..... | 17.4 | * | * | * | 0.2 | 0.6 | 1.4 | 4.1 | 16.3 | 56.7 | 148.3 | 407.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 15.7 | * | * | * | * | 0.2 | 0.7 | 1.6 | 5.7 | 23.7 | 78.0 | 276.7 |
| 2003..... | 16.6 | * | * | * | * | 0.2 | 0.6 | 1.8 | 6.3 | 24.6 | 81.0 | 301.4 |
| 2002..... | 17.2 | * | * | * | * | 0.3 | 0.7 | 1.6 | 6.5 | 26.4 | 84.0 | 313.9 |
| 2001..... | 17.8 | * | * | * | * | 0.3 | 0.7 | 1.8 | 6.6 | 26.4 | 88.2 | 334.2 |
| 2000..... | 17.9 | * | * | * | * | 0.2 | 0.7 | 1.7 | 6.3 | 27.5 | 88.8 | 343.5 |
| 1999..... | 18.6 | * | * | * | * | 0.2 | 0.8 | 1.8 | 6.7 | 28.1 | 92.1 | 364.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of arteries, arterioles and capillaries (I70-I78) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 9.1 | * | * | * | * | 0.9 | 2.5 | 5.9 | 15.2 | 36.1 | 95.5 | 260.5 |
| 2003..... | 9.5 | * | * | * | * | 0.8 | 2.5 | 7.2 | 14.8 | 38.7 | 102.0 | 275.8 |
| 2002..... | 9.7 | * | * | * | * | 0.9 | 2.2 | 7.1 | 16.6 | 39.8 | 106.1 | 275.5 |
| 2001..... | 10.0 | * | * | * | * | 0.8 | 2.6 | 6.0 | 17.5 | 41.2 | 111.2 | 301.8 |
| 2000..... | 10.3 | * | * | * | * | 0.7 | 2.4 | 6.8 | 17.9 | 42.5 | 114.5 | 320.2 |
| 1999..... | 10.0 | * | * | * | * | 0.6 | 2.2 | 6.1 | 15.8 | 41.5 | 112.3 | 335.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 8.6 | * | * | * | * | 1.2 | 3.3 | 8.0 | 21.2 | 44.6 | 111.2 | 239.8 |
| 2003..... | 9.2 | * | * | * | * | 1.2 | 3.3 | 9.7 | 20.4 | 48.4 | 131.9 | 247.2 |
| 2002..... | 9.3 | * | * | * | * | 1.3 | 2.9 | 9.7 | 23.8 | 49.5 | 123.7 | 277.9 |
| 2001..... | 9.3 | * | * | * | * | 1.0 | 3.7 | 8.5 | 21.8 | 51.3 | 134.4 | 281.8 |
| 2000..... | 9.6 | * | * | * | * | 0.9 | 3.1 | 9.1 | 22.7 | 52.9 | 137.3 | 329.9 |
| 1999..... | 9.3 | * | * | * | * | 1.0 | 2.6 | 8.3 | 20.4 | 52.0 | 135.1 | 350.6 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 9.5 | * | * | * | * | * | 1.9 | 4.0 | 10.3 | 30.0 | 86.6 | 268.6 |
| 2003..... | 9.8 | * | * | * | * | * | 1.8 | 5.1 | 10.3 | 31.8 | 85.1 | 286.8 |
| 2002..... | 10.0 | * | * | * | * | * | 1.5 | 4.8 | 10.9 | 32.9 | 96.2 | 274.7 |
| 2001..... | 10.6 | * | * | * | * | * | 1.7 | 3.8 | 14.0 | 34.0 | 98.2 | 309.3 |
| 2000..... | 10.9 | * | * | * | * | * | 1.8 | 4.8 | 14.0 | 35.2 | 101.6 | 316.6 |
| 1999..... | 10.7 | * | * | * | * | * | 1.8 | 4.3 | 12.0 | 34.0 | 99.6 | 330.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Atherosclerosis (I70) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 6.1 | 26.3 | 137.0 |
| 2003..... | 4.5 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 6.6 | 29.8 | 154.6 |
| 2002..... | 4.8 | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 7.1 | 31.9 | 168.1 |
| 2001..... | 4.9 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.3 | 33.1 | 179.1 |
| 2000..... | 5.1 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.7 | 34.6 | 188.8 |
| 1999..... | 5.4 | * | * | * | * | * | 0.1 | 0.5 | 2.0 | 8.0 | 35.9 | 202.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 0.1 | 0.5 | 2.0 | 7.3 | 29.7 | 128.2 |
| 2003..... | 3.4 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 7.8 | 33.6 | 138.6 |
| 2002..... | 3.7 | * | * | * | * | * | 0.1 | 0.6 | 2.4 | 8.8 | 35.5 | 156.6 |
| 2001..... | 3.7 | * | * | * | * | * | 0.1 | 0.6 | 2.4 | 9.0 | 35.9 | 167.2 |
| 2000..... | 3.9 | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 9.6 | 39.7 | 176.5 |
| 1999..... | 4.0 | * | * | * | * | * | 0.1 | 0.7 | 2.7 | 9.6 | 40.0 | 189.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | * | 0.2 | 1.2 | 5.1 | 24.0 | 141.0 |
| 2003..... | 5.5 | * | * | * | * | * | * | 0.3 | 1.2 | 5.6 | 27.3 | 161.7 |
| 2002..... | 5.9 | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 5.7 | 29.5 | 173.1 |
| 2001..... | 6.2 | * | * | * | * | * | * | 0.3 | 1.4 | 5.9 | 31.2 | 184.1 |
| 2000..... | 6.3 | * | * | * | * | * | * | 0.3 | 1.2 | 6.2 | 31.3 | 193.8 |
| 1999..... | 6.7 | * | * | * | * | * | * | 0.3 | 1.4 | 6.7 | 33.2 | 207.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | 1.0 | 4.3 | 18.6 | 74.7 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 1.1 | 4.6 | 18.9 | 86.0 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | 1.0 | 4.2 | 22.7 | 108.6 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | * | 4.0 | 23.1 | 130.7 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 5.9 | 24.5 | 132.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 1.5 | 5.2 | 27.1 | 129.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 5.1 | 22.9 | 88.3 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | * | 6.1 | 24.2 | 95.4 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | 5.2 | 26.7 | 102.1 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 4.8 | 27.3 | 127.6 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | * | 8.2 | 32.0 | 104.8 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 6.8 | 31.3 | 151.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | * | 3.7 | 15.6 | 67.7 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | * | 3.4 | 15.3 | 81.3 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | * | 3.5 | 20.0 | 111.9 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | * | 3.5 | 20.2 | 132.2 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 4.0 | 19.5 | 146.4 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | * | 4.0 | 24.3 | 118.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Atherosclerosis (I70) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 6.3 | 26.7 | 139.7 |
| 2003..... | 5.0 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 6.7 | 30.2 | 157.1 |
| 2002..... | 5.4 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.3 | 32.2 | 170.2 |
| 2001..... | 5.5 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.5 | 33.4 | 180.5 |
| 2000..... | 5.7 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 7.8 | 35.0 | 190.3 |
| 1999..... | 5.9 | * | * | * | * | * | 0.1 | 0.5 | 2.1 | 8.1 | 36.1 | 203.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | * | 0.1 | 0.5 | 2.0 | 7.5 | 30.0 | 130.1 |
| 2003..... | 3.8 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 7.9 | 34.0 | 140.3 |
| 2002..... | 4.1 | * | * | * | * | * | 0.1 | 0.6 | 2.5 | 9.0 | 35.8 | 158.7 |
| 2001..... | 4.1 | * | * | * | * | * | 0.1 | 0.6 | 2.4 | 9.2 | 36.1 | 168.2 |
| 2000..... | 4.3 | * | * | * | * | * | 0.2 | 0.7 | 2.6 | 9.7 | 39.9 | 178.7 |
| 1999..... | 4.4 | * | * | * | * | * | 0.2 | 0.7 | 2.7 | 9.7 | 40.3 | 189.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | * | * | * | 0.2 | 1.2 | 5.2 | 24.4 | 144.0 |
| 2003..... | 6.2 | * | * | * | * | * | * | 0.3 | 1.2 | 5.8 | 27.7 | 164.5 |
| 2002..... | 6.6 | * | * | * | * | * | * | 0.3 | 1.4 | 5.9 | 29.8 | 175.1 |
| 2001..... | 6.8 | * | * | * | * | * | * | 0.3 | 1.5 | 6.0 | 31.6 | 185.5 |
| 2000..... | 7.0 | * | * | * | * | * | * | 0.3 | 1.3 | 6.3 | 31.8 | 195.1 |
| 1999..... | 7.4 | * | * | * | * | * | * | 0.3 | 1.5 | 6.9 | 33.4 | 209.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 6.2 | 27.6 | 145.0 |
| 2003..... | 5.8 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 6.6 | 30.8 | 162.2 |
| 2002..... | 6.1 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 7.3 | 32.5 | 177.0 |
| 2001..... | 6.3 | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 7.5 | 34.1 | 187.4 |
| 2000..... | 6.5 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 7.8 | 35.5 | 196.8 |
| 1999..... | 6.7 | * | * | * | * | * | 0.1 | 0.5 | 2.0 | 8.2 | 36.7 | 209.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | * | 0.6 | 2.0 | 7.6 | 30.8 | 136.8 |
| 2003..... | 4.3 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 7.7 | 34.1 | 145.2 |
| 2002..... | 4.6 | * | * | * | * | * | 0.1 | 0.5 | 2.3 | 9.0 | 35.9 | 165.7 |
| 2001..... | 4.7 | * | * | * | * | * | 0.1 | 0.6 | 2.3 | 9.2 | 36.6 | 175.2 |
| 2000..... | 4.9 | * | * | * | * | * | 0.2 | 0.6 | 2.6 | 9.8 | 40.6 | 183.6 |
| 1999..... | 4.9 | * | * | * | * | * | 0.1 | 0.7 | 2.6 | 9.6 | 40.4 | 194.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | * | * | * | * | 0.2 | 1.2 | 5.1 | 25.4 | 148.7 |
| 2003..... | 7.2 | * | * | * | * | * | * | 0.3 | 1.1 | 5.8 | 28.5 | 169.6 |
| 2002..... | 7.6 | * | * | * | * | * | * | 0.3 | 1.3 | 5.9 | 30.2 | 181.9 |
| 2001..... | 7.9 | * | * | * | * | * | * | 0.3 | 1.3 | 6.1 | 32.5 | 192.6 |
| 2000..... | 8.0 | * | * | * | * | * | * | 0.2 | 1.1 | 6.2 | 32.1 | 202.1 |
| 1999..... | 8.5 | * | * | * | * | * | * | 0.3 | 1.4 | 6.9 | 34.3 | 215.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Atherosclerosis (I70) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | * | 0.5 | 2.2 | 7.5 | 24.3 | 103.6 |
| 2003..... | 2.7 | * | * | * | * | * | * | 0.8 | 2.7 | 9.1 | 30.8 | 122.9 |
| 2002..... | 2.8 | * | * | * | * | * | * | 1.0 | 3.0 | 8.1 | 37.0 | 118.0 |
| 2001..... | 2.7 | * | * | * | * | * | * | 0.7 | 3.4 | 8.9 | 30.7 | 126.1 |
| 2000..... | 3.0 | * | * | * | * | * | * | 1.1 | 3.3 | 9.1 | 35.6 | 140.2 |
| 1999..... | 3.1 | * | * | * | * | * | * | 0.7 | 3.3 | 9.3 | 35.0 | 155.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | * | 3.0 | 8.6 | 30.2 | 78.2 |
| 2003..... | 2.3 | * | * | * | * | * | * | 1.2 | 3.7 | 12.0 | 40.4 | 105.5 |
| 2002..... | 2.4 | * | * | * | * | * | * | 1.4 | 4.3 | 10.2 | 44.8 | 109.4 |
| 2001..... | 2.2 | * | * | * | * | * | * | 1.0 | 3.9 | 11.0 | 37.2 | 113.2 |
| 2000..... | 2.5 | * | * | * | * | * | * | 1.5 | 4.1 | 11.1 | 40.4 | 150.3 |
| 1999..... | 2.6 | * | * | * | * | * | * | * | 4.5 | 12.0 | 44.4 | 166.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | * | * | 1.7 | 6.7 | 20.9 | 113.6 |
| 2003..... | 3.0 | * | * | * | * | * | * | * | 1.9 | 7.1 | 25.4 | 129.6 |
| 2002..... | 3.2 | * | * | * | * | * | * | * | 2.0 | 6.6 | 32.5 | 121.2 |
| 2001..... | 3.2 | * | * | * | * | * | * | * | 3.0 | 7.4 | 27.1 | 130.9 |
| 2000..... | 3.5 | * | * | * | * | * | * | * | 2.7 | 7.7 | 32.8 | 136.5 |
| 1999..... | 3.5 | * | * | * | * | * | * | * | 2.3 | 7.4 | 29.8 | 151.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Aortic aneurysm and dissection (I71) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 4.5 | 15.9 | 39.6 | 64.8 |
| 2003..... | 5.1 | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 5.3 | 17.8 | 42.5 | 70.0 |
| 2002..... | 5.1 | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 5.5 | 19.1 | 43.3 | 69.4 |
| 2001..... | 5.3 | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 5.5 | 19.2 | 45.7 | 77.5 |
| 2000..... | 5.6 | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 6.2 | 21.8 | 47.2 | 79.2 |
| 1999..... | 5.7 | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 6.0 | 21.7 | 48.6 | 80.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 0.2 | 0.5 | 1.1 | 2.7 | 6.9 | 23.1 | 54.8 | 95.0 |
| 2003..... | 6.3 | * | * | * | 0.1 | 0.4 | 1.1 | 3.1 | 8.2 | 26.0 | 61.9 | 105.5 |
| 2002..... | 6.4 | * | * | * | 0.2 | 0.4 | 1.0 | 2.7 | 8.5 | 28.5 | 62.7 | 106.6 |
| 2001..... | 6.5 | * | * | * | 0.2 | 0.4 | 1.2 | 2.6 | 8.5 | 28.7 | 66.4 | 114.8 |
| 2000..... | 7.0 | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 9.5 | 32.9 | 70.6 | 127.3 |
| 1999..... | 7.0 | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 9.5 | 32.7 | 72.7 | 125.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.3 | 9.8 | 29.3 | 51.2 |
| 2003..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.6 | 11.0 | 29.5 | 54.2 |
| 2002..... | 4.0 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.7 | 11.2 | 30.5 | 53.3 |
| 2001..... | 4.2 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.8 | 11.3 | 32.1 | 61.9 |
| 2000..... | 4.3 | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.0 | 12.7 | 32.0 | 59.6 |
| 1999..... | 4.4 | * | * | * | * | * | 0.4 | 0.8 | 2.8 | 12.7 | 32.9 | 62.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.4 | 1.3 | 3.3 | 7.9 | 18.0 | 30.4 |
| 2003..... | 1.3 | * | * | * | * | * | 0.4 | 0.9 | 3.1 | 8.3 | 26.7 | 49.2 |
| 2002..... | 1.1 | * | * | * | * | * | 0.4 | 1.0 | 3.8 | 7.5 | 21.2 | 31.0 |
| 2001..... | 1.1 | * | * | * | * | * | 0.4 | 1.2 | 2.5 | 8.2 | 22.9 | 31.9 |
| 2000..... | 1.3 | * | * | * | * | * | 0.4 | 1.3 | 3.4 | 11.2 | 27.1 | 41.8 |
| 1999..... | 1.1 | * | * | * | * | * | 0.5 | 0.9 | 3.4 | 9.2 | 24.0 | 40.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.7 | 2.2 | 5.5 | 12.6 | 25.4 | 40.6 |
| 2003..... | 1.6 | * | * | * | * | * | * | 1.4 | 5.1 | 12.5 | 38.3 | 80.0 |
| 2002..... | 1.4 | * | * | * | * | * | 0.7 | 1.6 | 5.5 | 12.0 | 32.0 | 43.5 |
| 2001..... | 1.5 | * | * | * | * | * | 0.7 | 1.8 | 3.8 | 14.2 | 34.9 | 53.3 |
| 2000..... | 1.6 | * | * | * | * | * | * | 1.8 | 5.2 | 16.0 | 42.3 | 64.5 |
| 1999..... | 1.5 | * | * | * | * | * | * | 1.5 | 5.3 | 14.7 | 33.9 | 79.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 4.1 | 12.8 | 25.2 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | * | 5.0 | 18.7 | 33.6 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | 2.3 | 4.0 | 13.8 | 24.9 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 3.5 | 14.8 | 21.4 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | * | 7.5 | 16.8 | 30.7 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 4.8 | 17.3 | 21.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Aortic aneurysm and dissection (I71) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | 0.1 | 0.3 | 0.8 | 1.8 | 4.6 | 16.4 | 40.7 | 66.2 |
| 2003..... | 5.7 | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 5.5 | 18.5 | 43.2 | 70.7 |
| 2002..... | 5.7 | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 5.6 | 19.8 | 44.3 | 70.8 |
| 2001..... | 6.0 | * | * | * | 0.1 | 0.3 | 0.8 | 1.7 | 5.7 | 19.8 | 46.7 | 79.0 |
| 2000..... | 6.2 | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 6.3 | 22.4 | 47.9 | 80.3 |
| 1999..... | 6.3 | * | * | * | 0.1 | 0.3 | 0.8 | 1.8 | 6.2 | 22.4 | 49.5 | 81.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.3 | * | * | * | 0.2 | 0.5 | 1.2 | 2.7 | 7.0 | 23.8 | 56.4 | 97.5 |
| 2003..... | 7.0 | * | * | * | 0.2 | 0.4 | 1.2 | 3.2 | 8.5 | 26.9 | 63.0 | 106.6 |
| 2002..... | 7.2 | * | * | * | 0.2 | 0.4 | 1.1 | 2.9 | 8.7 | 29.6 | 64.1 | 109.1 |
| 2001..... | 7.3 | * | * | * | 0.2 | 0.5 | 1.2 | 2.7 | 8.8 | 29.5 | 67.7 | 117.0 |
| 2000..... | 7.8 | * | * | * | * | 0.4 | 1.0 | 2.7 | 9.8 | 33.8 | 71.6 | 129.6 |
| 1999..... | 7.8 | * | * | * | * | 0.4 | 1.1 | 2.8 | 9.8 | 33.6 | 74.3 | 127.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 10.2 | 30.2 | 52.3 |
| 2003..... | 4.4 | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.7 | 11.4 | 30.0 | 55.0 |
| 2002..... | 4.4 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.7 | 11.6 | 31.2 | 54.3 |
| 2001..... | 4.7 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.9 | 11.8 | 32.9 | 63.2 |
| 2000..... | 4.7 | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.1 | 13.0 | 32.5 | 60.4 |
| 1999..... | 4.8 | * | * | * | * | * | 0.4 | 0.9 | 2.9 | 13.2 | 33.4 | 63.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 4.3 | 17.2 | 41.9 | 68.4 |
| 2003..... | 6.4 | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 5.4 | 19.3 | 44.8 | 73.2 |
| 2002..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 5.5 | 20.7 | 46.0 | 72.8 |
| 2001..... | 6.7 | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 5.6 | 20.7 | 47.9 | 81.3 |
| 2000..... | 7.0 | * | * | * | * | 0.2 | 0.6 | 1.6 | 6.3 | 23.4 | 49.3 | 82.0 |
| 1999..... | 7.1 | * | * | * | 0.1 | 0.2 | 0.7 | 1.6 | 6.3 | 23.4 | 51.1 | 83.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 7.0 | * | * | * | 0.2 | 0.4 | 1.0 | 2.4 | 6.6 | 24.8 | 58.3 | 101.4 |
| 2003..... | 7.9 | * | * | * | 0.2 | 0.3 | 1.0 | 2.8 | 8.4 | 28.3 | 65.5 | 111.2 |
| 2002..... | 8.0 | * | * | * | * | 0.3 | 0.9 | 2.4 | 8.6 | 31.0 | 67.1 | 114.7 |
| 2001..... | 8.2 | * | * | * | 0.2 | 0.5 | 1.0 | 2.3 | 8.7 | 30.9 | 70.0 | 122.9 |
| 2000..... | 8.8 | * | * | * | * | 0.3 | 0.8 | 2.5 | 9.8 | 35.6 | 74.2 | 135.3 |
| 1999..... | 8.9 | * | * | * | * | 0.4 | 1.0 | 2.4 | 10.1 | 35.4 | 77.5 | 131.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | * | * | 0.3 | 0.7 | 2.2 | 10.6 | 30.7 | 53.6 |
| 2003..... | 4.9 | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 11.6 | 30.9 | 56.5 |
| 2002..... | 4.9 | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 11.9 | 31.9 | 54.7 |
| 2001..... | 5.2 | * | * | * | * | * | 0.3 | 0.6 | 2.7 | 12.0 | 33.3 | 64.0 |
| 2000..... | 5.2 | * | * | * | * | * | 0.3 | 0.7 | 2.9 | 13.1 | 32.9 | 60.5 |
| 1999..... | 5.3 | * | * | * | * | * | 0.4 | 0.7 | 2.7 | 13.3 | 33.9 | 64.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Aortic aneurysm and dissection (I71) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | 0.6 | 1.7 | 3.6 | 7.6 | 13.3 | 32.4 | 44.6 |
| 2003..... | 3.3 | * | * | * | * | 0.6 | 1.7 | 4.4 | 7.3 | 14.5 | 30.4 | 47.3 |
| 2002..... | 3.3 | * | * | * | * | 0.6 | 1.4 | 4.3 | 7.5 | 14.8 | 30.2 | 50.5 |
| 2001..... | 3.5 | * | * | * | * | 0.4 | 1.6 | 3.6 | 7.7 | 14.7 | 38.0 | 55.9 |
| 2000..... | 3.6 | * | * | * | * | 0.5 | 1.5 | 3.3 | 8.2 | 17.3 | 36.8 | 63.0 |
| 1999..... | 3.4 | * | * | * | * | 0.4 | 1.5 | 3.9 | 6.5 | 16.8 | 34.0 | 61.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | 1.0 | 2.6 | 5.5 | 11.8 | 18.5 | 37.4 | 57.9 |
| 2003..... | 3.9 | * | * | * | * | 1.1 | 2.5 | 6.5 | 11.0 | 17.9 | 39.8 | 52.2 |
| 2002..... | 3.9 | * | * | * | * | 1.0 | 2.3 | 6.6 | 12.1 | 19.3 | 34.9 | 49.2 |
| 2001..... | 3.9 | * | * | * | * | * | 2.6 | 5.4 | 11.3 | 19.9 | 47.4 | 49.7 |
| 2000..... | 4.0 | * | * | * | * | * | 2.3 | 4.7 | 11.6 | 22.6 | 47.2 | 70.5 |
| 1999..... | 3.7 | * | * | * | * | * | 2.0 | 6.0 | 9.3 | 22.1 | 38.8 | 77.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | 0.9 | 2.0 | 4.3 | 9.5 | 29.6 | 39.4 |
| 2003..... | 2.8 | * | * | * | * | * | 1.0 | 2.5 | 4.3 | 12.1 | 25.1 | 45.4 |
| 2002..... | 2.8 | * | * | * | * | * | * | 2.2 | 3.8 | 11.6 | 27.5 | 51.0 |
| 2001..... | 3.1 | * | * | * | * | * | 0.7 | 2.0 | 4.8 | 11.1 | 32.7 | 58.2 |
| 2000..... | 3.3 | * | * | * | * | * | 0.9 | 2.1 | 5.5 | 13.5 | 30.9 | 60.2 |
| 1999..... | 3.1 | * | * | * | * | * | 1.0 | 2.0 | 4.3 | 13.1 | 31.3 | 55.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of arteries, arterioles and capillaries (I72-I78) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.7 | 9.2 | 24.7 | 74.5 |
| 2003..... | 3.4 | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 2.8 | 9.1 | 25.0 | 76.0 |
| 2002..... | 3.5 | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 3.1 | 10.0 | 25.3 | 79.0 |
| 2001..... | 3.5 | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.2 | 10.0 | 26.1 | 80.7 |
| 2000..... | 3.6 | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 3.1 | 10.4 | 27.4 | 84.0 |
| 1999..... | 3.6 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.3 | 9.8 | 27.2 | 84.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.0 | 10.3 | 28.0 | 71.6 |
| 2003..... | 3.0 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.1 | 11.0 | 29.9 | 74.2 |
| 2002..... | 3.0 | * | * | * | * | * | 0.3 | 1.0 | 3.4 | 11.3 | 29.6 | 80.8 |
| 2001..... | 3.1 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.5 | 11.6 | 30.6 | 82.5 |
| 2000..... | 3.2 | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.6 | 12.5 | 32.3 | 87.6 |
| 1999..... | 3.1 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.8 | 11.7 | 31.5 | 82.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.4 | 8.3 | 22.5 | 75.8 |
| 2003..... | 3.8 | * | * | * | * | * | 0.3 | 0.9 | 2.6 | 7.5 | 21.8 | 76.9 |
| 2002..... | 4.0 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.8 | 8.9 | 22.5 | 78.2 |
| 2001..... | 4.0 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.9 | 8.6 | 23.1 | 79.9 |
| 2000..... | 4.0 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.6 | 8.7 | 24.2 | 82.6 |
| 1999..... | 4.1 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.8 | 8.3 | 24.5 | 86.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.7 | 1.8 | 6.1 | 16.5 | 46.6 |
| 2003..... | 0.9 | * | * | * | * | * | * | 0.7 | 1.5 | 5.7 | 15.6 | 57.0 |
| 2002..... | 1.1 | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 7.3 | 19.8 | 61.5 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.9 | 2.2 | 5.5 | 16.0 | 63.2 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | 2.4 | 8.9 | 15.4 | 61.0 |
| 1999..... | 1.0 | * | * | * | * | * | * | * | 2.4 | 7.2 | 20.0 | 70.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 8.0 | 19.0 | 49.0 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 2.2 | 6.5 | 20.3 | 58.5 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | 7.8 | 24.6 | 58.6 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 2.5 | 5.8 | 18.3 | 55.2 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | 2.7 | 9.7 | 19.7 | 66.5 |
| 1999..... | 0.9 | * | * | * | * | * | * | * | * | 7.5 | 28.2 | 84.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | 1.9 | 4.5 | 14.8 | 45.4 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | * | 5.1 | 12.4 | 56.3 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | 2.6 | 6.8 | 16.6 | 63.0 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | * | 5.4 | 14.5 | 67.0 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | * | 8.3 | 12.5 | 58.4 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | 2.4 | 7.1 | 14.6 | 63.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of arteries, arterioles and capillaries (I72-I78) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.8 | 9.4 | 25.1 | 75.6 |
| 2003..... | 3.8 | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 2.9 | 9.3 | 25.5 | 76.7 |
| 2002..... | 3.9 | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 3.2 | 10.2 | 25.6 | 79.5 |
| 2001..... | 3.9 | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.2 | 10.3 | 26.5 | 81.1 |
| 2000..... | 4.0 | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 3.1 | 10.5 | 27.9 | 84.7 |
| 1999..... | 3.9 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.3 | 9.9 | 27.5 | 85.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | 0.3 | 1.1 | 3.1 | 10.5 | 28.5 | 72.6 |
| 2003..... | 3.3 | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 11.3 | 30.3 | 74.9 |
| 2002..... | 3.4 | * | * | * | * | * | 0.3 | 1.0 | 3.5 | 11.5 | 29.8 | 81.7 |
| 2001..... | 3.4 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.5 | 12.0 | 31.1 | 83.5 |
| 2000..... | 3.5 | * | * | * | * | 0.1 | 0.2 | 1.2 | 3.6 | 12.7 | 32.8 | 88.2 |
| 1999..... | 3.4 | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.9 | 11.9 | 31.5 | 82.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | * | 0.4 | 0.8 | 2.5 | 8.5 | 22.9 | 76.9 |
| 2003..... | 4.2 | * | * | * | * | * | 0.4 | 0.9 | 2.8 | 7.6 | 22.3 | 77.6 |
| 2002..... | 4.3 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.8 | 9.0 | 22.8 | 78.5 |
| 2001..... | 4.4 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.9 | 8.8 | 23.5 | 80.1 |
| 2000..... | 4.4 | * | * | * | * | * | 0.4 | 0.9 | 2.6 | 8.7 | 24.6 | 83.3 |
| 1999..... | 4.5 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.8 | 8.3 | 24.8 | 86.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.5 | 9.0 | 24.4 | 73.6 |
| 2003..... | 4.0 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.8 | 8.9 | 24.6 | 75.1 |
| 2002..... | 4.1 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.9 | 9.7 | 24.9 | 78.2 |
| 2001..... | 4.2 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.9 | 9.8 | 25.6 | 78.8 |
| 2000..... | 4.3 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 10.1 | 27.1 | 82.8 |
| 1999..... | 4.2 | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.1 | 9.6 | 26.5 | 83.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | * | 0.3 | 1.1 | 2.8 | 10.2 | 28.1 | 71.6 |
| 2003..... | 3.5 | * | * | * | * | * | 0.2 | 0.9 | 2.9 | 10.8 | 29.2 | 74.6 |
| 2002..... | 3.6 | * | * | * | * | * | 0.2 | 0.9 | 3.2 | 11.0 | 29.3 | 79.9 |
| 2001..... | 3.6 | * | * | * | * | * | 0.2 | 0.8 | 3.3 | 11.5 | 30.3 | 81.8 |
| 2000..... | 3.8 | * | * | * | * | * | 0.2 | 1.0 | 3.3 | 12.3 | 32.1 | 87.4 |
| 1999..... | 3.6 | * | * | * | * | * | 0.3 | 1.0 | 3.6 | 11.6 | 30.4 | 81.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 0.3 | 0.7 | 2.3 | 8.0 | 21.9 | 74.4 |
| 2003..... | 4.5 | * | * | * | * | * | 0.3 | 0.7 | 2.7 | 7.3 | 21.6 | 75.4 |
| 2002..... | 4.7 | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 8.6 | 21.9 | 77.4 |
| 2001..... | 4.7 | * | * | * | * | * | 0.4 | 0.9 | 2.6 | 8.3 | 22.4 | 77.6 |
| 2000..... | 4.7 | * | * | * | * | * | 0.3 | 0.7 | 2.2 | 8.2 | 23.8 | 81.0 |
| 1999..... | 4.8 | * | * | * | * | * | 0.4 | 0.8 | 2.5 | 7.9 | 24.0 | 83.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of arteries, arterioles and capillaries (I72-I78) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | * | 0.6 | 1.7 | 5.3 | 15.3 | 38.8 | 112.2 |
| 2003..... | 3.5 | * | * | * | * | * | 0.7 | 2.0 | 4.8 | 15.0 | 40.7 | 105.6 |
| 2002..... | 3.6 | * | * | * | * | * | 0.6 | 1.8 | 6.2 | 16.8 | 39.0 | 107.1 |
| 2001..... | 3.8 | * | * | * | * | * | 0.9 | 1.7 | 6.4 | 17.5 | 42.5 | 119.9 |
| 2000..... | 3.7 | * | * | * | * | * | 0.6 | 2.4 | 6.3 | 16.1 | 42.1 | 117.0 |
| 1999..... | 3.6 | * | * | * | * | * | 0.6 | 1.5 | 5.9 | 15.3 | 43.3 | 119.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | * | * | 2.0 | 6.5 | 17.5 | 43.6 | 103.6 |
| 2003..... | 3.0 | * | * | * | * | * | * | 1.9 | 5.7 | 18.5 | 51.7 | 89.5 |
| 2002..... | 3.1 | * | * | * | * | * | * | 1.6 | 7.4 | 19.9 | 44.0 | 119.2 |
| 2001..... | 3.3 | * | * | * | * | * | 0.9 | 2.0 | 6.6 | 20.5 | 49.8 | 119.0 |
| 2000..... | 3.2 | * | * | * | * | * | * | 2.9 | 7.1 | 19.2 | 49.7 | 109.2 |
| 1999..... | 3.0 | * | * | * | * | * | * | 1.4 | 6.7 | 17.9 | 51.9 | 107.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | 0.8 | 1.5 | 4.4 | 13.7 | 36.1 | 115.7 |
| 2003..... | 4.0 | * | * | * | * | * | 0.8 | 2.1 | 4.1 | 12.5 | 34.5 | 111.8 |
| 2002..... | 4.0 | * | * | * | * | * | 0.8 | 1.9 | 5.1 | 14.6 | 36.1 | 102.4 |
| 2001..... | 4.4 | * | * | * | * | * | 0.9 | 1.4 | 6.3 | 15.5 | 38.4 | 120.2 |
| 2000..... | 4.2 | * | * | * | * | * | 0.7 | 1.9 | 5.7 | 14.0 | 37.9 | 119.9 |
| 1999..... | 4.2 | * | * | * | * | * | 0.8 | 1.6 | 5.4 | 13.5 | 38.5 | 123.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other disorders of circulatory system (I80-I99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | 0.7 | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 2.1 | 3.6 | 9.0 | 23.3 |
| 2003..... | 1.6 | 0.9 | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.0 | 4.1 | 9.0 | 24.0 |
| 2002..... | 1.6 | 0.7 | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 2.2 | 3.8 | 9.8 | 24.4 |
| 2001..... | 1.6 | 0.5 | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.2 | 4.6 | 9.4 | 26.1 |
| 2000..... | 1.6 | 0.7 | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.1 | 4.3 | 9.4 | 25.6 |
| 1999..... | 1.4 | 0.6 | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 1.9 | 3.8 | 9.1 | 23.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | 0.4 | 0.7 | 1.6 | 2.5 | 3.9 | 9.7 | 19.6 |
| 2003..... | 1.4 | 1.0 | * | * | 0.1 | 0.3 | 0.8 | 1.6 | 2.4 | 4.2 | 8.5 | 19.1 |
| 2002..... | 1.4 | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 2.4 | 3.9 | 9.5 | 20.5 |
| 2001..... | 1.4 | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.5 | 4.9 | 9.8 | 22.0 |
| 2000..... | 1.4 | * | * | * | * | 0.3 | 0.8 | 1.6 | 2.3 | 4.7 | 9.7 | 22.2 |
| 1999..... | 1.3 | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.0 | 4.2 | 9.6 | 20.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.3 | 0.7 | 1.1 | 1.8 | 3.4 | 8.5 | 25.0 |
| 2003..... | 1.9 | * | * | * | 0.1 | 0.4 | 0.8 | 1.2 | 1.7 | 4.0 | 9.4 | 26.2 |
| 2002..... | 1.9 | * | * | * | * | 0.4 | 0.7 | 1.2 | 2.1 | 3.8 | 10.1 | 26.2 |
| 2001..... | 1.9 | * | * | * | 0.1 | 0.4 | 0.6 | 1.0 | 2.0 | 4.4 | 9.2 | 27.8 |
| 2000..... | 1.8 | * | * | * | 0.1 | 0.4 | 0.6 | 1.1 | 2.0 | 4.0 | 9.2 | 26.9 |
| 1999..... | 1.6 | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.8 | 3.4 | 8.8 | 24.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.3 | 1.2 | 1.6 | 2.0 | 4.8 | 17.6 |
| 2003..... | 0.6 | * | * | * | * | * | 0.4 | 1.1 | 1.4 | 2.6 | 6.4 | 17.6 |
| 2002..... | 0.6 | * | * | * | * | * | 0.4 | 1.0 | 1.4 | 3.1 | 6.3 | 18.3 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.9 | * | 4.2 | 7.4 | 32.5 |
| 2000..... | 0.6 | * | * | * | * | * | 0.6 | 1.1 | 1.3 | 3.3 | 8.7 | 24.6 |
| 1999..... | 0.5 | * | * | * | * | * | * | 1.1 | 1.5 | 4.1 | 5.4 | 24.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 1.4 | 2.2 | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | * | 1.4 | * | * | * | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | 1.3 | * | 4.0 | 8.9 | * |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.8 | * | * | 10.8 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | * | 4.9 | 23.8 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | * | 6.6 | 19.5 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 3.2 | 6.2 | 23.2 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 4.4 | 6.3 | 33.5 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 7.3 | 23.7 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 4.3 | * | 23.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other disorders of circulatory system (I80-I99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | 0.8 | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.2 | 3.7 | 9.2 | 23.5 |
| 2003..... | 1.8 | 0.8 | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.1 | 4.1 | 9.1 | 24.3 |
| 2002..... | 1.8 | 0.6 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.3 | 3.9 | 10.0 | 24.7 |
| 2001..... | 1.8 | * | * | * | 0.1 | 0.4 | 0.7 | 1.2 | 2.3 | 4.6 | 9.5 | 25.8 |
| 2000..... | 1.8 | 0.7 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.2 | 4.4 | 9.4 | 25.6 |
| 1999..... | 1.6 | 0.7 | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.9 | 3.7 | 9.3 | 23.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.4 | 0.8 | 1.6 | 2.5 | 4.0 | 9.9 | 20.2 |
| 2003..... | 1.5 | * | * | * | * | 0.3 | 0.8 | 1.6 | 2.4 | 4.2 | 8.6 | 19.3 |
| 2002..... | 1.5 | * | * | * | * | 0.4 | 0.7 | 1.5 | 2.5 | 3.9 | 9.6 | 21.1 |
| 2001..... | 1.5 | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.6 | 4.9 | 9.8 | 21.6 |
| 2000..... | 1.5 | * | * | * | * | 0.3 | 0.8 | 1.6 | 2.3 | 4.8 | 9.6 | 22.1 |
| 1999..... | 1.4 | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.0 | 4.2 | 9.7 | 20.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.4 | 0.7 | 1.1 | 1.9 | 3.5 | 8.7 | 25.0 |
| 2003..... | 2.1 | * | * | * | 0.1 | 0.4 | 0.9 | 1.2 | 1.7 | 4.1 | 9.5 | 26.5 |
| 2002..... | 2.1 | * | * | * | * | 0.4 | 0.7 | 1.2 | 2.1 | 3.8 | 10.2 | 26.2 |
| 2001..... | 2.0 | * | * | * | 0.1 | 0.4 | 0.6 | 1.0 | 2.1 | 4.3 | 9.2 | 27.6 |
| 2000..... | 2.0 | * | * | * | * | 0.4 | 0.6 | 1.2 | 2.1 | 4.1 | 9.3 | 27.0 |
| 1999..... | 1.8 | * | * | * | * | 0.3 | 0.5 | 0.9 | 1.8 | 3.3 | 9.0 | 24.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.0 | 3.5 | 9.0 | 23.4 |
| 2003..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 1.9 | 3.9 | 8.7 | 23.8 |
| 2002..... | 1.8 | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.1 | 3.6 | 9.6 | 24.5 |
| 2001..... | 1.8 | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.1 | 4.3 | 9.2 | 25.7 |
| 2000..... | 1.8 | * | * | * | * | 0.3 | 0.6 | 1.2 | 1.9 | 4.0 | 9.2 | 25.5 |
| 1999..... | 1.6 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 3.5 | 9.0 | 23.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.4 | 3.8 | 9.6 | 20.3 |
| 2003..... | 1.5 | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.2 | 3.9 | 8.0 | 18.7 |
| 2002..... | 1.5 | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.2 | 3.7 | 9.1 | 20.5 |
| 2001..... | 1.5 | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.3 | 4.6 | 9.4 | 21.2 |
| 2000..... | 1.5 | * | * | * | * | 0.2 | 0.7 | 1.5 | 2.0 | 4.5 | 9.4 | 21.8 |
| 1999..... | 1.4 | * | * | * | * | 0.2 | 0.6 | 0.9 | 2.0 | 3.9 | 9.3 | 20.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | 0.3 | 0.6 | 0.9 | 1.6 | 3.3 | 8.5 | 24.8 |
| 2003..... | 2.1 | * | * | * | * | 0.3 | 0.6 | 1.2 | 1.6 | 3.9 | 9.3 | 26.1 |
| 2002..... | 2.2 | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.9 | 3.6 | 9.9 | 26.3 |
| 2001..... | 2.1 | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.9 | 4.1 | 9.0 | 27.6 |
| 2000..... | 2.0 | * | * | * | * | 0.3 | 0.5 | 1.0 | 1.9 | 3.6 | 9.0 | 27.0 |
| 1999..... | 1.8 | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.7 | 3.1 | 8.8 | 24.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other disorders of circulatory system (I80-I99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | 1.0 | 2.0 | 3.0 | 4.2 | 6.8 | 13.9 | 29.1 |
| 2003..... | 2.2 | * | * | * | * | 0.9 | 2.0 | 2.7 | 4.4 | 7.4 | 15.5 | 33.6 |
| 2002..... | 2.2 | * | * | * | * | 1.0 | 1.7 | 2.7 | 5.0 | 6.9 | 16.8 | 29.9 |
| 2001..... | 2.0 | * | * | * | * | 0.8 | 1.6 | 2.3 | 4.9 | 7.9 | 14.3 | 29.7 |
| 2000..... | 2.1 | * | * | * | * | 0.9 | 1.5 | 2.8 | 4.6 | 9.0 | 14.4 | 29.9 |
| 1999..... | 1.8 | * | * | * | * | 0.5 | 1.5 | 2.4 | 3.0 | 6.8 | 14.0 | 28.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | 1.1 | 2.0 | 3.5 | 4.5 | 7.3 | 15.4 | 25.4 |
| 2003..... | 2.1 | * | * | * | * | 0.9 | 1.4 | 3.5 | 5.4 | 8.4 | 17.9 | 33.0 |
| 2002..... | 2.1 | * | * | * | * | 1.1 | 1.8 | 3.0 | 5.6 | 7.8 | 18.5 | 31.7 |
| 2001..... | 2.0 | * | * | * | * | * | 1.8 | 2.4 | 6.5 | 9.2 | 15.9 | 28.9 |
| 2000..... | 2.0 | * | * | * | * | 0.9 | 1.8 | 3.5 | 5.3 | 9.1 | 13.3 | 30.5 |
| 1999..... | 1.7 | * | * | * | * | * | 1.9 | 2.9 | 2.7 | 8.4 | 17.2 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 0.8 | 1.9 | 2.6 | 3.9 | 6.5 | 13.0 | 30.5 |
| 2003..... | 2.3 | * | * | * | * | 1.0 | 2.6 | 1.9 | 3.6 | 6.8 | 14.2 | 33.8 |
| 2002..... | 2.3 | * | * | * | * | 1.0 | 1.6 | 2.4 | 4.5 | 6.2 | 15.9 | 29.3 |
| 2001..... | 2.1 | * | * | * | * | 0.9 | 1.4 | 2.1 | 3.7 | 7.0 | 13.5 | 29.9 |
| 2000..... | 2.3 | * | * | * | * | 1.0 | 1.3 | 2.3 | 4.0 | 8.8 | 15.1 | 29.7 |
| 1999..... | 1.8 | * | * | * | * | * | 1.1 | 2.0 | 3.2 | 5.6 | 12.3 | 30.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | 0.1 | 0.3 | 0.7 | 1.0 | 1.6 | 2.7 | 7.0 | 17.4 |
| 2003..... | 1.4 | * | * | * | 0.1 | 0.3 | 0.7 | 1.1 | 1.6 | 3.2 | 6.9 | 18.3 |
| 2002..... | 1.4 | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 1.8 | 2.9 | 7.5 | 18.8 |
| 2001..... | 1.4 | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 1.8 | 3.4 | 7.2 | 19.3 |
| 2000..... | 1.3 | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.5 | 3.3 | 7.0 | 18.6 |
| 1999..... | 1.0 | * | * | * | 0.1 | 0.2 | 0.5 | 0.7 | 1.2 | 2.4 | 5.8 | 14.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.4 | 0.7 | 1.1 | 1.7 | 2.8 | 7.2 | 14.7 |
| 2003..... | 1.1 | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.8 | 3.1 | 6.1 | 14.0 |
| 2002..... | 1.1 | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.8 | 2.9 | 6.8 | 15.3 |
| 2001..... | 1.1 | * | * | * | * | 0.3 | 0.6 | 0.9 | 1.9 | 3.5 | 6.9 | 14.8 |
| 2000..... | 1.1 | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.4 | 3.3 | 6.8 | 14.9 |
| 1999..... | 0.8 | * | * | * | * | 0.2 | 0.5 | 0.6 | 1.1 | 2.6 | 5.7 | 10.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.3 | 0.7 | 0.9 | 1.5 | 2.6 | 6.9 | 18.6 |
| 2003..... | 1.6 | * | * | * | 0.1 | 0.4 | 0.7 | 1.0 | 1.4 | 3.3 | 7.4 | 20.2 |
| 2002..... | 1.6 | * | * | * | * | 0.4 | 0.6 | 1.1 | 1.8 | 2.9 | 7.9 | 20.2 |
| 2001..... | 1.6 | * | * | * | * | 0.4 | 0.6 | 0.8 | 1.7 | 3.3 | 7.4 | 21.2 |
| 2000..... | 1.5 | * | * | * | * | 0.4 | 0.5 | 1.0 | 1.6 | 3.2 | 7.1 | 20.1 |
| 1999..... | 1.2 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.3 | 2.2 | 5.9 | 15.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.4 | 2.5 | 6.8 | 17.3 |
| 2003..... | 1.4 | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 1.4 | 3.0 | 6.6 | 17.9 |
| 2002..... | 1.4 | * | * | * | * | 0.3 | 0.5 | 0.9 | 1.6 | 2.7 | 7.2 | 18.6 |
| 2001..... | 1.4 | * | * | * | * | 0.3 | 0.4 | 0.8 | 1.6 | 3.2 | 7.0 | 19.3 |
| 2000..... | 1.3 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.4 | 3.0 | 6.8 | 18.7 |
| 1999..... | 1.0 | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.2 | 2.2 | 5.7 | 14.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.6 | 2.6 | 7.0 | 14.7 |
| 2003..... | 1.1 | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.6 | 2.8 | 5.7 | 13.5 |
| 2002..... | 1.1 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.6 | 2.7 | 6.5 | 14.9 |
| 2001..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.3 | 6.7 | 14.7 |
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.3 | 3.1 | 6.8 | 14.8 |
| 1999..... | 0.8 | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 2.4 | 5.5 | 11.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.2 | 2.4 | 6.7 | 18.4 |
| 2003..... | 1.7 | * | * | * | * | 0.3 | 0.5 | 1.0 | 1.2 | 3.1 | 7.2 | 19.9 |
| 2002..... | 1.7 | * | * | * | * | 0.3 | 0.5 | 1.0 | 1.6 | 2.8 | 7.6 | 20.2 |
| 2001..... | 1.6 | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.5 | 3.0 | 7.1 | 21.2 |
| 2000..... | 1.6 | * | * | * | * | 0.3 | 0.4 | 0.9 | 1.5 | 2.9 | 6.8 | 20.3 |
| 1999..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.3 | 2.1 | 5.7 | 15.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.9 | 1.7 | 2.4 | 3.6 | 4.9 | 10.8 | 23.0 |
| 2003..... | 1.8 | * | * | * | * | 0.9 | 1.7 | 2.1 | 3.7 | 5.9 | 11.9 | 26.8 |
| 2002..... | 1.8 | * | * | * | * | 1.0 | 1.5 | 2.2 | 3.8 | 5.3 | 13.0 | 23.9 |
| 2001..... | 1.7 | * | * | * | * | 0.8 | 1.3 | 1.8 | 4.0 | 6.4 | 10.9 | 22.8 |
| 2000..... | 1.7 | * | * | * | * | 0.9 | 1.3 | 2.2 | 3.4 | 7.0 | 10.9 | 21.6 |
| 1999..... | 1.2 | * | * | * | * | 0.5 | 1.2 | 1.6 | 2.0 | 4.7 | 9.4 | 18.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 1.1 | 1.7 | 2.5 | 3.6 | 5.1 | 11.1 | 20.3 |
| 2003..... | 1.6 | * | * | * | * | 0.8 | 1.2 | 2.7 | 4.1 | 6.4 | 13.0 | 25.6 |
| 2002..... | 1.6 | * | * | * | * | 1.0 | 1.5 | 2.3 | 4.3 | 5.6 | 12.3 | 23.0 |
| 2001..... | 1.5 | * | * | * | * | * | 1.4 | 1.8 | 5.0 | 6.8 | 11.1 | * |
| 2000..... | 1.5 | * | * | * | * | 0.8 | 1.5 | 2.4 | 3.6 | 6.7 | 8.3 | * |
| 1999..... | 1.1 | * | * | * | * | * | 1.4 | 1.5 | 1.9 | 5.9 | 10.3 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.7 | 1.8 | 2.3 | 3.5 | 4.8 | 10.6 | 24.1 |
| 2003..... | 2.0 | * | * | * | * | 0.9 | 2.2 | 1.6 | 3.3 | 5.6 | 11.2 | 27.2 |
| 2002..... | 1.9 | * | * | * | * | 1.0 | 1.5 | 2.1 | 3.4 | 5.1 | 13.4 | 24.2 |
| 2001..... | 1.8 | * | * | * | * | 0.9 | 1.2 | 1.8 | 3.2 | 6.2 | 10.8 | 23.5 |
| 2000..... | 1.9 | * | * | * | * | 1.0 | 1.1 | 2.0 | 3.3 | 7.2 | 12.3 | 21.8 |
| 1999..... | 1.4 | * | * | * | * | * | 1.1 | 1.7 | 2.1 | 3.9 | 8.9 | 21.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified disorders of circulatory system (I83-I99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.1 | 2.2 | 6.1 |
| 2003..... | 0.4 | 0.7 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.2 | 6.0 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.5 | 5.9 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.3 | 6.5 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.1 | 2.4 | 6.9 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 3.4 | 8.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.6 | 0.8 | 1.2 | 2.7 | 5.5 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.1 | 2.5 | 5.4 |
| 2002..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 2.9 | 5.7 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.4 | 2.9 | 6.7 |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.4 | 2.8 | 7.1 |
| 1999..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.6 | 4.0 | 9.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 6.4 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.1 | 6.2 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.3 | 6.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 | 6.4 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 6.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.1 | 8.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 2.1 | 6.1 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 2.1 | 5.9 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 2.5 | 5.9 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 2.2 | 6.5 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 2.4 | 6.8 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.3 | 8.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.5 | 0.8 | 1.2 | 2.6 | 5.6 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.1 | 2.3 | 5.2 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.0 | 2.6 | 5.6 |
| 2001..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.7 | 6.6 |
| 2000..... | 0.5 | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.4 | 2.7 | 7.0 |
| 1999..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 1.5 | 3.8 | 9.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 6.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.1 | 6.2 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.3 | 6.1 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 1.1 | 1.9 | 6.4 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.2 | 6.7 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.0 | 8.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| X. Diseases of the respiratory system (J00-J98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 77.3 | 15.8 | 1.9 | 0.7 | 1.3 | 2.0 | 5.6 | 17.2 | 63.6 | 229.1 | 627.0 | 1,497.1 |
| 2003..... | 81.1 | 17.5 | 2.2 | 0.8 | 1.5 | 2.3 | 5.9 | 18.2 | 66.4 | 239.5 | 651.9 | 1,625.4 |
| 2002..... | 81.4 | 15.8 | 1.9 | 0.8 | 1.3 | 2.3 | 6.0 | 17.6 | 65.7 | 240.3 | 662.2 | 1,663.7 |
| 2001..... | 80.8 | 17.3 | 1.8 | 0.7 | 1.3 | 2.3 | 6.0 | 17.6 | 67.3 | 245.9 | 651.9 | 1,671.8 |
| 2000..... | 82.1 | 17.1 | 1.7 | 0.8 | 1.4 | 2.5 | 6.2 | 17.9 | 68.8 | 250.2 | 667.5 | 1,734.0 |
| 1999..... | 82.4 | 18.1 | 2.1 | 0.8 | 1.4 | 2.5 | 6.0 | 17.3 | 70.8 | 255.0 | 671.7 | 1,733.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 75.2 | 17.6 | 1.8 | 0.8 | 1.5 | 2.3 | 5.9 | 19.3 | 70.6 | 267.2 | 779.3 | 1,915.3 |
| 2003..... | 78.2 | 20.6 | 2.5 | 0.9 | 1.6 | 2.7 | 6.1 | 20.3 | 74.0 | 279.9 | 808.5 | 2,057.2 |
| 2002..... | 78.9 | 18.8 | 2.2 | 0.8 | 1.5 | 2.4 | 6.5 | 19.5 | 73.0 | 282.2 | 832.8 | 2,150.0 |
| 2001..... | 78.2 | 19.7 | 1.9 | 0.8 | 1.5 | 2.4 | 6.4 | 19.7 | 74.6 | 289.4 | 824.1 | 2,187.2 |
| 2000..... | 80.2 | 18.9 | 1.7 | 0.8 | 1.6 | 2.7 | 6.8 | 19.9 | 77.1 | 297.1 | 851.4 | 2,344.4 |
| 1999..... | 81.3 | 19.6 | 2.2 | 0.9 | 1.5 | 2.6 | 6.4 | 19.1 | 79.5 | 310.4 | 870.5 | 2,356.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 79.4 | 14.1 | 1.9 | 0.6 | 1.1 | 1.7 | 5.4 | 15.3 | 57.2 | 197.1 | 524.4 | 1,309.0 |
| 2003..... | 84.0 | 14.2 | 1.8 | 0.8 | 1.3 | 2.0 | 5.8 | 16.2 | 59.3 | 205.8 | 547.3 | 1,434.5 |
| 2002..... | 83.7 | 12.6 | 1.5 | 0.8 | 1.1 | 2.1 | 5.6 | 15.8 | 58.9 | 205.5 | 548.9 | 1,452.7 |
| 2001..... | 83.2 | 14.7 | 1.8 | 0.7 | 1.2 | 2.3 | 5.6 | 15.5 | 60.6 | 209.9 | 538.8 | 1,456.2 |
| 2000..... | 84.0 | 15.2 | 1.7 | 0.7 | 1.2 | 2.3 | 5.7 | 16.0 | 61.1 | 211.7 | 547.6 | 1,485.4 |
| 1999..... | 83.4 | 16.6 | 1.9 | 0.7 | 1.2 | 2.4 | 5.7 | 15.6 | 62.9 | 209.7 | 542.6 | 1,481.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 19.7 | 14.6 | 1.7 | 0.5 | 1.0 | 1.3 | 3.4 | 9.8 | 31.4 | 117.8 | 382.8 | 1,092.7 |
| 2003..... | 20.2 | 17.2 | 2.2 | 0.7 | 1.1 | 1.6 | 3.3 | 10.4 | 34.2 | 120.6 | 388.4 | 1,210.1 |
| 2002..... | 20.0 | 12.7 | 1.5 | 0.6 | 0.8 | 1.4 | 3.4 | 10.0 | 34.7 | 119.9 | 406.5 | 1,273.8 |
| 2001..... | 20.1 | 13.8 | 2.1 | 0.4 | 0.7 | 1.0 | 3.2 | 11.0 | 33.9 | 123.3 | 401.4 | 1,442.5 |
| 2000..... | 20.1 | 15.8 | 1.6 | 0.7 | 0.8 | 1.7 | 4.1 | 10.5 | 37.2 | 122.6 | 413.2 | 1,406.0 |
| 1999..... | 19.8 | 15.9 | 1.4 | 0.6 | 0.9 | 1.3 | 4.1 | 10.9 | 35.3 | 126.3 | 402.3 | 1,420.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 19.1 | 17.1 | 1.6 | 0.7 | 1.3 | 1.5 | 3.9 | 11.1 | 37.1 | 146.6 | 476.0 | 1,291.7 |
| 2003..... | 20.0 | 19.9 | 2.7 | 0.8 | 1.3 | 1.6 | 3.5 | 12.2 | 41.9 | 151.5 | 497.4 | 1,477.7 |
| 2002..... | 19.9 | 14.3 | 1.9 | 0.6 | 1.0 | 1.6 | 3.7 | 11.7 | 44.4 | 155.8 | 525.7 | 1,551.1 |
| 2001..... | 19.7 | 14.9 | 2.0 | * | 0.9 | 1.1 | 3.6 | 12.5 | 40.9 | 156.4 | 515.1 | 1,839.8 |
| 2000..... | 20.1 | 16.0 | * | 0.7 | 0.9 | 1.8 | 4.6 | 12.8 | 47.5 | 156.4 | 562.0 | 1,751.4 |
| 1999..... | 19.7 | 18.1 | 1.5 | 0.7 | 0.8 | 1.3 | 4.8 | 11.8 | 41.5 | 161.7 | 538.1 | 1,882.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 20.3 | 12.0 | 1.8 | * | 0.8 | 1.2 | 2.9 | 8.4 | 26.3 | 94.7 | 318.3 | 990.3 |
| 2003..... | 20.5 | 14.4 | 1.6 | 0.6 | 1.0 | 1.5 | 3.0 | 8.6 | 27.3 | 96.0 | 313.6 | 1,074.2 |
| 2002..... | 20.1 | 11.0 | * | 0.6 | * | 1.2 | 3.1 | 8.4 | 26.1 | 91.4 | 324.9 | 1,136.5 |
| 2001..... | 20.5 | 12.7 | 2.2 | * | * | 0.8 | 2.8 | 9.5 | 27.7 | 97.3 | 324.6 | 1,248.3 |
| 2000..... | 20.0 | 15.7 | 1.9 | 0.7 | 0.7 | 1.6 | 3.5 | 8.2 | 28.0 | 96.1 | 313.3 | 1,236.5 |
| 1999..... | 19.8 | 13.6 | * | 0.6 | 1.0 | 1.3 | 3.4 | 10.0 | 29.8 | 98.6 | 311.9 | 1,197.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| X. Diseases of the respiratory system (J00-J98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 86.6 | 16.2 | 1.9 | 0.7 | 1.3 | 2.2 | 6.0 | 18.0 | 66.1 | 236.6 | 639.5 | 1,513.0 |
| 2003..... | 90.6 | 17.4 | 2.2 | 0.9 | 1.5 | 2.5 | 6.3 | 18.9 | 68.6 | 247.3 | 664.4 | 1,639.7 |
| 2002..... | 90.6 | 16.5 | 2.0 | 0.8 | 1.4 | 2.5 | 6.4 | 18.3 | 68.0 | 247.9 | 673.0 | 1,675.8 |
| 2001..... | 89.5 | 18.0 | 1.8 | 0.8 | 1.4 | 2.6 | 6.4 | 18.1 | 69.7 | 253.1 | 661.8 | 1,676.5 |
| 2000..... | 90.7 | 17.3 | 1.7 | 0.8 | 1.5 | 2.6 | 6.5 | 18.4 | 70.9 | 257.3 | 676.4 | 1,741.6 |
| 1999..... | 90.7 | 18.5 | 2.2 | 0.8 | 1.5 | 2.7 | 6.2 | 17.8 | 73.1 | 261.9 | 680.7 | 1,738.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 84.7 | 17.7 | 1.9 | 0.8 | 1.5 | 2.6 | 6.2 | 20.1 | 73.0 | 274.7 | 795.0 | 1,943.4 |
| 2003..... | 87.7 | 20.5 | 2.4 | 0.9 | 1.7 | 2.9 | 6.5 | 21.0 | 76.1 | 287.9 | 823.2 | 2,080.3 |
| 2002..... | 88.3 | 19.8 | 2.3 | 0.8 | 1.6 | 2.6 | 6.9 | 20.2 | 75.0 | 289.6 | 845.6 | 2,172.0 |
| 2001..... | 87.1 | 20.9 | 1.9 | 0.8 | 1.6 | 2.7 | 6.8 | 20.2 | 76.7 | 296.7 | 836.5 | 2,196.6 |
| 2000..... | 89.0 | 19.6 | 1.8 | 0.9 | 1.7 | 2.8 | 7.0 | 20.4 | 78.9 | 304.4 | 861.1 | 2,363.3 |
| 1999..... | 90.0 | 19.8 | 2.3 | 1.0 | 1.7 | 2.9 | 6.5 | 19.6 | 81.8 | 317.8 | 881.7 | 2,367.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 88.4 | 14.6 | 1.9 | 0.6 | 1.1 | 1.7 | 5.7 | 16.0 | 59.6 | 204.4 | 534.9 | 1,320.5 |
| 2003..... | 93.3 | 14.1 | 1.9 | 0.8 | 1.3 | 2.1 | 6.2 | 16.9 | 61.7 | 213.3 | 558.5 | 1,446.1 |
| 2002..... | 92.9 | 12.9 | 1.6 | 0.9 | 1.2 | 2.3 | 5.9 | 16.5 | 61.4 | 213.1 | 558.6 | 1,461.8 |
| 2001..... | 91.9 | 15.0 | 1.7 | 0.7 | 1.3 | 2.5 | 6.0 | 16.0 | 63.1 | 216.9 | 547.2 | 1,460.1 |
| 2000..... | 92.4 | 14.9 | 1.6 | 0.7 | 1.3 | 2.4 | 5.9 | 16.6 | 63.4 | 218.4 | 556.1 | 1,490.2 |
| 1999..... | 91.5 | 17.2 | 2.0 | 0.7 | 1.3 | 2.6 | 5.9 | 16.0 | 65.1 | 216.0 | 550.3 | 1,487.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 98.9 | 11.3 | 1.4 | 0.5 | 1.1 | 1.9 | 5.3 | 16.8 | 67.1 | 248.4 | 663.6 | 1,558.7 |
| 2003..... | 103.3 | 13.2 | 1.6 | 0.7 | 1.2 | 2.1 | 5.5 | 17.8 | 69.4 | 258.6 | 688.6 | 1,691.8 |
| 2002..... | 103.2 | 11.5 | 1.4 | 0.6 | 1.2 | 2.1 | 5.6 | 17.2 | 68.6 | 258.9 | 695.5 | 1,726.5 |
| 2001..... | 101.3 | 11.8 | 1.4 | 0.6 | 1.1 | 2.1 | 5.4 | 16.7 | 70.7 | 263.4 | 682.9 | 1,718.0 |
| 2000..... | 102.5 | 12.1 | 1.4 | 0.6 | 1.2 | 2.1 | 5.5 | 16.9 | 71.3 | 265.5 | 697.1 | 1,791.6 |
| 1999..... | 102.2 | 12.2 | 1.6 | 0.7 | 1.2 | 2.1 | 5.1 | 16.2 | 73.7 | 270.4 | 700.2 | 1,785.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 95.3 | 12.0 | 1.4 | 0.5 | 1.3 | 2.1 | 5.4 | 18.7 | 72.4 | 282.6 | 814.2 | 1,990.5 |
| 2003..... | 98.3 | 15.6 | 1.7 | 0.7 | 1.3 | 2.4 | 5.6 | 19.7 | 75.1 | 294.9 | 838.2 | 2,131.2 |
| 2002..... | 99.0 | 14.6 | 1.6 | 0.6 | 1.3 | 2.2 | 5.9 | 18.7 | 74.0 | 297.2 | 862.1 | 2,224.6 |
| 2001..... | 97.2 | 14.2 | 1.3 | 0.7 | 1.3 | 2.1 | 5.8 | 18.6 | 76.3 | 303.3 | 851.4 | 2,239.8 |
| 2000..... | 99.0 | 14.4 | 1.3 | 0.7 | 1.4 | 2.2 | 6.0 | 18.2 | 77.8 | 307.5 | 877.2 | 2,421.7 |
| 1999..... | 99.9 | 13.3 | 1.4 | 0.8 | 1.4 | 2.3 | 5.2 | 17.5 | 80.7 | 322.7 | 896.0 | 2,422.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 102.3 | 10.6 | 1.4 | 0.5 | 0.9 | 1.6 | 5.2 | 15.0 | 62.1 | 219.0 | 561.0 | 1,365.3 |
| 2003..... | 108.0 | 10.6 | 1.6 | 0.7 | 1.1 | 1.8 | 5.3 | 15.9 | 63.9 | 227.5 | 587.6 | 1,498.3 |
| 2002..... | 107.1 | 8.3 | 1.2 | 0.7 | 1.1 | 2.1 | 5.2 | 15.8 | 63.6 | 226.2 | 583.8 | 1,511.4 |
| 2001..... | 105.3 | 9.2 | 1.4 | 0.6 | 1.0 | 2.1 | 5.0 | 14.9 | 65.4 | 229.6 | 571.1 | 1,501.0 |
| 2000..... | 105.8 | 9.6 | 1.5 | 0.5 | 1.0 | 2.1 | 5.1 | 15.6 | 65.0 | 230.1 | 578.7 | 1,537.5 |
| 1999..... | 104.4 | 11.1 | 1.8 | 0.6 | 1.1 | 1.9 | 5.0 | 14.9 | 67.0 | 226.7 | 572.0 | 1,531.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| X. Diseases of the respiratory system (J00-J98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 45.8 | 34.7 | 3.7 | 1.6 | 2.5 | 4.3 | 11.4 | 30.2 | 77.0 | 197.3 | 488.0 | 1,095.4 |
| 2003..... | 47.4 | 33.4 | 3.5 | 1.6 | 3.0 | 5.3 | 12.3 | 31.3 | 80.0 | 209.7 | 505.7 | 1,144.1 |
| 2002..... | 48.2 | 35.2 | 4.2 | 1.8 | 2.8 | 4.7 | 12.8 | 29.7 | 81.1 | 209.0 | 526.5 | 1,214.4 |
| 2001..... | 48.6 | 41.2 | 3.3 | 1.4 | 2.9 | 5.8 | 13.3 | 31.9 | 78.7 | 215.1 | 518.4 | 1,266.4 |
| 2000..... | 49.6 | 39.7 | 3.2 | 1.6 | 3.1 | 5.4 | 13.0 | 33.7 | 85.0 | 233.6 | 522.4 | 1,247.8 |
| 1999..... | 50.7 | 43.6 | 4.1 | 1.6 | 2.9 | 6.5 | 13.9 | 33.2 | 86.2 | 233.8 | 537.4 | 1,301.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 49.6 | 38.7 | 3.7 | 2.0 | 2.7 | 5.7 | 13.0 | 35.2 | 99.4 | 274.2 | 706.8 | 1,521.1 |
| 2003..... | 51.7 | 38.8 | 4.5 | 2.0 | 3.6 | 6.8 | 12.5 | 35.2 | 103.9 | 290.1 | 763.0 | 1,590.7 |
| 2002..... | 51.9 | 40.3 | 4.8 | 1.8 | 3.4 | 5.3 | 14.6 | 34.8 | 104.5 | 282.9 | 774.6 | 1,705.5 |
| 2001..... | 52.1 | 45.3 | 3.9 | 1.5 | 3.2 | 6.5 | 14.0 | 37.0 | 99.0 | 295.7 | 778.7 | 1,792.5 |
| 2000..... | 54.1 | 41.3 | 4.1 | 1.9 | 3.5 | 6.9 | 14.9 | 40.2 | 106.7 | 330.3 | 763.9 | 1,822.3 |
| 1999..... | 55.7 | 46.8 | 5.4 | 2.0 | 3.4 | 6.7 | 15.2 | 40.2 | 111.0 | 327.8 | 813.9 | 1,907.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 42.3 | 30.6 | 3.7 | 1.1 | 2.3 | 3.1 | 10.1 | 25.8 | 59.0 | 142.4 | 364.5 | 927.2 |
| 2003..... | 43.6 | 27.7 | 2.5 | 1.1 | 2.4 | 3.9 | 12.0 | 28.0 | 60.8 | 152.5 | 360.5 | 971.1 |
| 2002..... | 44.8 | 30.0 | 3.6 | 1.9 | 2.2 | 4.2 | 11.2 | 25.2 | 62.3 | 156.6 | 386.2 | 1,026.7 |
| 2001..... | 45.5 | 36.9 | 2.6 | 1.3 | 2.6 | 5.2 | 12.7 | 27.6 | 62.5 | 158.3 | 372.3 | 1,071.5 |
| 2000..... | 45.4 | 38.1 | 2.2 | 1.3 | 2.6 | 4.0 | 11.3 | 28.1 | 67.6 | 165.6 | 386.4 | 1,034.4 |
| 1999..... | 46.2 | 40.3 | 2.9 | 1.2 | 2.4 | 6.3 | 12.8 | 27.1 | 66.4 | 167.5 | 382.5 | 1,074.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute upper respiratory infections (J00-J06) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 0.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.5 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.3 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.3 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.7 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 0.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.3 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.3 | 2.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 0.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.5 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.3 | 1.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 0.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.2 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute laryngitis and tracheitis (J04)

All origins 3/, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Acute laryngitis and tracheitis (J04) - All origins 3/, both sexes.

All origins 3/, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Acute laryngitis and tracheitis (J04) - All origins 3/, male.

All origins 3/, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Acute laryngitis and tracheitis (J04) - All origins 3/, female.

Hispanic, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Acute laryngitis and tracheitis (J04) - Hispanic, both sexes.

Hispanic, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Acute laryngitis and tracheitis (J04) - Hispanic, male.

Hispanic, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Acute laryngitis and tracheitis (J04) - Hispanic, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of the respiratory system (J10-J98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 77.3 | 15.7 | 1.8 | 0.7 | 1.3 | 2.0 | 5.6 | 17.2 | 63.6 | 229.0 | 626.8 | 1,496.3 |
| 2003..... | 81.1 | 17.1 | 2.1 | 0.8 | 1.4 | 2.3 | 5.9 | 18.2 | 66.3 | 239.5 | 651.8 | 1,623.9 |
| 2002..... | 81.3 | 15.5 | 1.8 | 0.8 | 1.3 | 2.3 | 6.0 | 17.6 | 65.6 | 240.2 | 662.0 | 1,662.6 |
| 2001..... | 80.7 | 17.0 | 1.7 | 0.7 | 1.3 | 2.3 | 6.0 | 17.5 | 67.3 | 245.9 | 651.7 | 1,670.6 |
| 2000..... | 82.0 | 16.9 | 1.7 | 0.8 | 1.4 | 2.4 | 6.2 | 17.9 | 68.7 | 250.1 | 667.2 | 1,732.2 |
| 1999..... | 82.3 | 17.8 | 2.0 | 0.8 | 1.4 | 2.5 | 6.0 | 17.3 | 70.8 | 255.0 | 671.3 | 1,731.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 75.1 | 17.4 | 1.8 | 0.8 | 1.4 | 2.3 | 5.9 | 19.2 | 70.6 | 267.1 | 779.2 | 1,914.6 |
| 2003..... | 78.1 | 20.1 | 2.4 | 0.9 | 1.6 | 2.7 | 6.0 | 20.3 | 73.9 | 279.9 | 808.3 | 2,055.5 |
| 2002..... | 78.9 | 18.6 | 2.1 | 0.8 | 1.5 | 2.4 | 6.5 | 19.4 | 72.9 | 282.1 | 832.7 | 2,149.3 |
| 2001..... | 78.1 | 19.4 | 1.8 | 0.7 | 1.5 | 2.4 | 6.4 | 19.7 | 74.5 | 289.4 | 824.0 | 2,186.6 |
| 2000..... | 80.1 | 18.7 | 1.7 | 0.8 | 1.6 | 2.6 | 6.8 | 19.9 | 77.1 | 296.9 | 851.2 | 2,343.3 |
| 1999..... | 81.3 | 19.2 | 2.1 | 0.9 | 1.5 | 2.6 | 6.3 | 19.1 | 79.5 | 310.3 | 870.1 | 2,355.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 79.4 | 14.0 | 1.8 | 0.5 | 1.1 | 1.7 | 5.4 | 15.3 | 57.1 | 197.0 | 524.2 | 1,308.1 |
| 2003..... | 83.9 | 13.9 | 1.8 | 0.8 | 1.3 | 2.0 | 5.7 | 16.2 | 59.2 | 205.7 | 547.2 | 1,433.1 |
| 2002..... | 83.7 | 12.4 | 1.4 | 0.8 | 1.1 | 2.1 | 5.6 | 15.8 | 58.9 | 205.4 | 548.8 | 1,451.5 |
| 2001..... | 83.2 | 14.6 | 1.7 | 0.6 | 1.1 | 2.3 | 5.6 | 15.5 | 60.6 | 209.8 | 538.5 | 1,454.6 |
| 2000..... | 83.9 | 15.1 | 1.6 | 0.7 | 1.2 | 2.2 | 5.6 | 16.0 | 61.1 | 211.5 | 547.3 | 1,483.4 |
| 1999..... | 83.3 | 16.4 | 1.9 | 0.7 | 1.2 | 2.4 | 5.7 | 15.6 | 62.8 | 209.6 | 542.2 | 1,479.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 19.6 | 14.2 | 1.6 | 0.5 | 1.0 | 1.3 | 3.4 | 9.8 | 31.4 | 117.8 | 382.8 | 1,092.7 |
| 2003..... | 20.2 | 16.9 | 2.2 | 0.7 | 1.1 | 1.5 | 3.3 | 10.4 | 34.2 | 120.6 | 388.3 | 1,209.0 |
| 2002..... | 20.0 | 12.5 | 1.3 | 0.6 | 0.8 | 1.4 | 3.4 | 10.0 | 34.7 | 119.9 | 406.0 | 1,273.3 |
| 2001..... | 20.1 | 13.5 | 2.0 | 0.4 | 0.7 | 1.0 | 3.2 | 11.0 | 33.8 | 123.3 | 401.4 | 1,442.5 |
| 2000..... | 20.0 | 15.8 | 1.6 | 0.7 | 0.8 | 1.7 | 4.1 | 10.4 | 37.1 | 122.4 | 413.0 | 1,406.0 |
| 1999..... | 19.7 | 15.5 | 1.3 | 0.6 | 0.9 | 1.3 | 4.0 | 10.9 | 35.3 | 126.3 | 402.3 | 1,419.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 19.0 | 16.4 | 1.5 | 0.7 | 1.2 | 1.4 | 3.8 | 11.1 | 37.1 | 146.6 | 476.0 | 1,291.7 |
| 2003..... | 19.9 | 19.7 | 2.7 | 0.8 | 1.3 | 1.6 | 3.5 | 12.2 | 41.8 | 151.5 | 497.0 | 1,477.7 |
| 2002..... | 19.9 | 14.1 | 1.7 | 0.6 | 1.0 | 1.6 | 3.7 | 11.6 | 44.4 | 155.8 | 525.2 | 1,551.1 |
| 2001..... | 19.7 | 14.4 | 2.0 | * | 0.9 | 1.1 | 3.6 | 12.5 | 40.9 | 156.4 | 515.1 | 1,839.8 |
| 2000..... | 20.1 | 16.0 | * | 0.7 | 0.9 | 1.7 | 4.6 | 12.6 | 47.4 | 156.2 | 562.0 | 1,751.4 |
| 1999..... | 19.7 | 17.3 | 1.4 | 0.7 | 0.8 | 1.2 | 4.6 | 11.8 | 41.5 | 161.7 | 538.1 | 1,880.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 20.3 | 12.0 | 1.7 | * | 0.7 | 1.2 | 2.9 | 8.4 | 26.3 | 94.7 | 318.3 | 990.3 |
| 2003..... | 20.5 | 13.9 | 1.6 | 0.6 | 1.0 | 1.5 | 3.0 | 8.6 | 27.3 | 96.0 | 313.6 | 1,072.7 |
| 2002..... | 20.1 | 10.8 | * | 0.6 | * | 1.2 | 3.1 | 8.4 | 26.0 | 91.4 | 324.3 | 1,135.6 |
| 2001..... | 20.5 | 12.5 | 2.1 | * | * | 0.8 | 2.8 | 9.5 | 27.5 | 97.2 | 324.6 | 1,248.3 |
| 2000..... | 20.0 | 15.7 | 1.9 | 0.7 | 0.7 | 1.6 | 3.5 | 8.1 | 28.0 | 95.9 | 313.0 | 1,236.5 |
| 1999..... | 19.8 | 13.6 | * | 0.6 | 1.0 | 1.3 | 3.4 | 9.9 | 29.8 | 98.6 | 311.9 | 1,196.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of the respiratory system (J10-J98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 86.5 | 16.1 | 1.8 | 0.7 | 1.3 | 2.2 | 5.9 | 18.0 | 66.0 | 236.5 | 639.3 | 1,512.1 |
| 2003..... | 90.5 | 17.0 | 2.0 | 0.9 | 1.5 | 2.5 | 6.3 | 18.9 | 68.6 | 247.2 | 664.2 | 1,638.2 |
| 2002..... | 90.6 | 16.3 | 1.9 | 0.8 | 1.4 | 2.4 | 6.4 | 18.2 | 67.9 | 247.8 | 672.9 | 1,674.7 |
| 2001..... | 89.5 | 17.8 | 1.6 | 0.7 | 1.4 | 2.6 | 6.4 | 18.1 | 69.6 | 253.1 | 661.6 | 1,675.1 |
| 2000..... | 90.6 | 17.1 | 1.7 | 0.8 | 1.5 | 2.6 | 6.4 | 18.4 | 70.8 | 257.1 | 676.1 | 1,739.8 |
| 1999..... | 90.7 | 18.3 | 2.1 | 0.8 | 1.5 | 2.7 | 6.2 | 17.7 | 73.1 | 261.8 | 680.4 | 1,737.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 84.6 | 17.6 | 1.8 | 0.8 | 1.5 | 2.6 | 6.2 | 20.0 | 73.0 | 274.7 | 794.8 | 1,942.7 |
| 2003..... | 87.6 | 20.0 | 2.3 | 0.9 | 1.7 | 2.9 | 6.4 | 21.0 | 76.0 | 287.8 | 823.0 | 2,078.5 |
| 2002..... | 88.3 | 19.6 | 2.2 | 0.8 | 1.6 | 2.6 | 6.8 | 20.1 | 74.9 | 289.5 | 845.4 | 2,171.2 |
| 2001..... | 87.1 | 20.6 | 1.7 | 0.8 | 1.6 | 2.7 | 6.8 | 20.2 | 76.7 | 296.7 | 836.3 | 2,196.0 |
| 2000..... | 88.9 | 19.2 | 1.8 | 0.9 | 1.7 | 2.8 | 7.0 | 20.3 | 78.9 | 304.2 | 860.9 | 2,362.1 |
| 1999..... | 89.9 | 19.6 | 2.2 | 0.9 | 1.7 | 2.8 | 6.5 | 19.6 | 81.8 | 317.7 | 881.3 | 2,366.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 88.4 | 14.5 | 1.8 | 0.6 | 1.1 | 1.7 | 5.7 | 16.0 | 59.6 | 204.4 | 534.7 | 1,319.6 |
| 2003..... | 93.3 | 13.9 | 1.8 | 0.8 | 1.3 | 2.1 | 6.1 | 16.9 | 61.6 | 213.2 | 558.3 | 1,444.7 |
| 2002..... | 92.8 | 12.7 | 1.5 | 0.9 | 1.2 | 2.3 | 5.9 | 16.4 | 61.4 | 213.0 | 558.5 | 1,460.5 |
| 2001..... | 91.8 | 14.9 | 1.5 | 0.7 | 1.2 | 2.5 | 6.0 | 16.0 | 63.0 | 216.8 | 546.9 | 1,458.5 |
| 2000..... | 92.3 | 14.9 | 1.5 | 0.7 | 1.2 | 2.3 | 5.9 | 16.6 | 63.4 | 218.3 | 555.7 | 1,488.2 |
| 1999..... | 91.4 | 16.9 | 2.0 | 0.7 | 1.3 | 2.6 | 5.9 | 16.0 | 65.1 | 215.9 | 550.0 | 1,485.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 98.8 | 11.2 | 1.3 | 0.5 | 1.1 | 1.8 | 5.3 | 16.8 | 67.1 | 248.4 | 663.4 | 1,557.9 |
| 2003..... | 103.2 | 12.8 | 1.5 | 0.7 | 1.2 | 2.1 | 5.5 | 17.7 | 69.3 | 258.5 | 688.4 | 1,690.3 |
| 2002..... | 103.1 | 11.5 | 1.3 | 0.6 | 1.1 | 2.1 | 5.5 | 17.2 | 68.6 | 258.8 | 695.4 | 1,725.4 |
| 2001..... | 101.3 | 11.7 | 1.2 | 0.6 | 1.1 | 2.1 | 5.4 | 16.7 | 70.7 | 263.4 | 682.6 | 1,716.6 |
| 2000..... | 102.4 | 11.8 | 1.3 | 0.6 | 1.2 | 2.1 | 5.5 | 16.8 | 71.2 | 265.4 | 696.8 | 1,789.7 |
| 1999..... | 102.1 | 12.0 | 1.5 | 0.7 | 1.2 | 2.1 | 5.1 | 16.2 | 73.6 | 270.4 | 699.8 | 1,783.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 95.3 | 11.9 | 1.3 | 0.5 | 1.3 | 2.1 | 5.4 | 18.6 | 72.4 | 282.5 | 814.0 | 1,989.8 |
| 2003..... | 98.3 | 15.0 | 1.6 | 0.7 | 1.3 | 2.4 | 5.6 | 19.7 | 75.0 | 294.8 | 838.0 | 2,129.6 |
| 2002..... | 99.0 | 14.6 | 1.5 | 0.6 | 1.2 | 2.2 | 5.9 | 18.7 | 74.0 | 297.2 | 862.0 | 2,224.0 |
| 2001..... | 97.2 | 14.1 | 1.2 | 0.6 | 1.3 | 2.0 | 5.8 | 18.5 | 76.3 | 303.2 | 851.2 | 2,239.0 |
| 2000..... | 98.9 | 13.9 | 1.3 | 0.7 | 1.4 | 2.1 | 5.9 | 18.1 | 77.8 | 307.4 | 877.0 | 2,420.6 |
| 1999..... | 99.8 | 13.1 | 1.3 | 0.8 | 1.4 | 2.3 | 5.2 | 17.4 | 80.7 | 322.6 | 895.6 | 2,421.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 102.2 | 10.5 | 1.3 | 0.5 | 0.9 | 1.6 | 5.2 | 15.0 | 62.0 | 219.0 | 560.7 | 1,364.4 |
| 2003..... | 107.9 | 10.5 | 1.4 | 0.7 | 1.1 | 1.8 | 5.3 | 15.8 | 63.9 | 227.4 | 587.4 | 1,496.9 |
| 2002..... | 107.0 | 8.2 | 1.2 | 0.6 | 1.0 | 2.1 | 5.2 | 15.7 | 63.5 | 226.1 | 583.7 | 1,510.0 |
| 2001..... | 105.2 | 9.2 | 1.3 | 0.6 | 1.0 | 2.1 | 5.0 | 14.9 | 65.3 | 229.6 | 570.8 | 1,499.3 |
| 2000..... | 105.7 | 9.6 | 1.4 | 0.5 | 1.0 | 2.1 | 5.0 | 15.5 | 65.0 | 230.0 | 578.3 | 1,535.3 |
| 1999..... | 104.3 | 10.9 | 1.8 | 0.6 | 1.1 | 1.9 | 5.0 | 14.9 | 67.0 | 226.7 | 571.6 | 1,529.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of the respiratory system (J10-J98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 45.7 | 34.7 | 3.7 | 1.6 | 2.5 | 4.3 | 11.3 | 30.1 | 77.0 | 197.1 | 487.9 | 1,094.6 |
| 2003..... | 47.4 | 32.9 | 3.4 | 1.6 | 3.0 | 5.3 | 12.2 | 31.2 | 79.9 | 209.6 | 505.7 | 1,142.6 |
| 2002..... | 48.1 | 34.6 | 4.1 | 1.8 | 2.8 | 4.7 | 12.8 | 29.6 | 81.1 | 208.9 | 526.3 | 1,214.4 |
| 2001..... | 48.6 | 40.7 | 3.1 | 1.4 | 2.9 | 5.8 | 13.2 | 31.9 | 78.6 | 215.1 | 518.3 | 1,265.4 |
| 2000..... | 49.5 | 39.5 | 3.1 | 1.6 | 3.0 | 5.4 | 13.0 | 33.6 | 85.0 | 233.5 | 522.3 | 1,247.2 |
| 1999..... | 50.6 | 43.1 | 4.0 | 1.6 | 2.9 | 6.4 | 13.9 | 33.1 | 86.1 | 233.6 | 537.3 | 1,299.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 49.5 | 38.7 | 3.7 | 2.0 | 2.7 | 5.7 | 12.8 | 35.2 | 99.3 | 274.0 | 706.5 | 1,520.1 |
| 2003..... | 51.6 | 38.5 | 4.4 | 2.0 | 3.6 | 6.8 | 12.3 | 35.0 | 103.7 | 289.8 | 763.0 | 1,587.5 |
| 2002..... | 51.8 | 39.4 | 4.8 | 1.8 | 3.4 | 5.3 | 14.6 | 34.7 | 104.5 | 282.7 | 774.3 | 1,705.5 |
| 2001..... | 52.0 | 44.4 | 3.6 | 1.5 | 3.2 | 6.5 | 13.9 | 37.0 | 98.8 | 295.7 | 778.7 | 1,792.5 |
| 2000..... | 54.0 | 41.3 | 4.0 | 1.9 | 3.5 | 6.9 | 14.9 | 40.2 | 106.7 | 330.0 | 763.6 | 1,820.0 |
| 1999..... | 55.6 | 46.4 | 5.1 | 1.9 | 3.3 | 6.5 | 15.2 | 40.2 | 110.8 | 327.7 | 813.9 | 1,902.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 42.3 | 30.6 | 3.7 | 1.1 | 2.3 | 3.1 | 10.0 | 25.8 | 59.0 | 142.3 | 364.5 | 926.4 |
| 2003..... | 43.5 | 27.1 | 2.4 | 1.1 | 2.4 | 3.9 | 12.0 | 27.9 | 60.8 | 152.5 | 360.5 | 970.2 |
| 2002..... | 44.8 | 29.7 | 3.4 | 1.9 | 2.1 | 4.2 | 11.2 | 25.2 | 62.2 | 156.5 | 386.0 | 1,026.7 |
| 2001..... | 45.4 | 36.9 | 2.5 | 1.2 | 2.6 | 5.2 | 12.7 | 27.5 | 62.4 | 158.2 | 372.1 | 1,070.2 |
| 2000..... | 45.4 | 37.7 | 2.2 | 1.3 | 2.5 | 4.0 | 11.3 | 27.9 | 67.6 | 165.6 | 386.4 | 1,034.4 |
| 1999..... | 46.1 | 39.6 | 2.9 | 1.2 | 2.4 | 6.3 | 12.7 | 27.1 | 66.3 | 167.3 | 382.4 | 1,074.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Influenza (J10-J11) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.3 | 11.8 |
| 2003..... | 0.7 | 0.7 | 0.4 | 0.1 | 0.1 | * | 0.1 | 0.2 | 0.3 | 1.1 | 3.6 | 17.5 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.1 | 0.3 | 1.4 | 9.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 2.3 |
| 2000..... | 0.7 | * | * | * | * | * | 0.0 | 0.1 | 0.4 | 1.1 | 4.1 | 20.5 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.7 | 21.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 11.6 |
| 2003..... | 0.5 | * | 0.4 | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.7 | 17.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.6 | 9.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.3 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 4.1 | 21.1 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.8 | 4.3 | 20.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.4 | 2.2 | 11.8 |
| 2003..... | 0.8 | * | 0.4 | 0.1 | * | * | * | 0.1 | 0.3 | 1.0 | 3.6 | 17.6 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 0.2 | 1.3 | 8.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.2 |
| 2000..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 4.1 | 20.3 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.4 | 22.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.5 | 12.7 |
| 2003..... | 0.8 | * | 0.3 | 0.1 | * | * | * | 0.2 | 0.3 | 1.1 | 3.9 | 18.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.1 | 0.3 | 1.6 | 9.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.4 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 4.3 | 21.7 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.9 | 23.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.8 | 2.5 | 12.7 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 1.2 | 4.0 | 18.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.7 | 10.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.4 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.4 | 4.3 | 22.0 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 0.9 | 4.6 | 22.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.4 | 12.8 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.9 | 18.7 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 1.4 | 9.5 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.4 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 4.3 | 21.6 |
| 1999..... | 1.0 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.5 | 23.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonia (J12-J18) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 19.9 | 6.3 | 0.6 | 0.2 | 0.4 | 0.7 | 2.0 | 4.5 | 10.7 | 34.0 | 137.0 | 571.2 |
| 2003..... | 21.8 | 7.3 | 0.7 | 0.2 | 0.5 | 0.9 | 2.2 | 5.1 | 10.9 | 36.3 | 147.5 | 649.1 |
| 2002..... | 22.5 | 6.3 | 0.7 | 0.2 | 0.4 | 0.9 | 2.1 | 4.7 | 11.1 | 37.2 | 155.5 | 687.8 |
| 2001..... | 21.7 | 7.2 | 0.7 | 0.2 | 0.4 | 0.8 | 2.2 | 4.6 | 10.6 | 36.2 | 148.1 | 683.4 |
| 2000..... | 22.6 | 7.4 | 0.6 | 0.2 | 0.4 | 0.9 | 2.3 | 4.6 | 11.5 | 38.0 | 156.4 | 723.9 |
| 1999..... | 22.2 | 8.1 | 0.8 | 0.2 | 0.4 | 0.8 | 2.3 | 4.6 | 10.7 | 36.4 | 153.3 | 730.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 18.3 | 7.5 | 0.5 | 0.2 | 0.5 | 0.9 | 2.2 | 5.5 | 12.8 | 42.1 | 170.3 | 666.1 |
| 2003..... | 19.6 | 8.5 | 0.7 | 0.2 | 0.5 | 1.1 | 2.5 | 6.2 | 12.9 | 45.4 | 183.2 | 740.9 |
| 2002..... | 20.2 | 7.6 | 0.8 | 0.2 | 0.4 | 0.9 | 2.4 | 5.7 | 13.4 | 46.0 | 193.1 | 796.9 |
| 2001..... | 19.5 | 7.8 | 0.7 | 0.2 | 0.5 | 0.9 | 2.5 | 5.7 | 13.0 | 44.9 | 186.0 | 806.2 |
| 2000..... | 20.3 | 8.1 | 0.6 | 0.2 | 0.5 | 1.0 | 2.7 | 5.6 | 14.0 | 47.1 | 195.8 | 871.3 |
| 1999..... | 19.8 | 8.5 | 0.7 | 0.2 | 0.4 | 0.9 | 2.6 | 5.8 | 12.5 | 46.0 | 192.3 | 874.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 21.5 | 5.0 | 0.8 | 0.2 | 0.3 | 0.5 | 1.8 | 3.5 | 8.6 | 27.3 | 114.6 | 528.5 |
| 2003..... | 23.9 | 6.0 | 0.6 | 0.3 | 0.4 | 0.7 | 1.9 | 4.0 | 9.1 | 28.6 | 123.7 | 608.5 |
| 2002..... | 24.8 | 5.0 | 0.6 | 0.2 | 0.3 | 0.8 | 1.9 | 3.8 | 9.0 | 29.8 | 130.6 | 640.5 |
| 2001..... | 23.8 | 6.7 | 0.7 | 0.2 | 0.4 | 0.7 | 1.9 | 3.5 | 8.4 | 29.0 | 123.2 | 632.0 |
| 2000..... | 24.8 | 6.6 | 0.6 | 0.2 | 0.4 | 0.8 | 1.9 | 3.6 | 9.1 | 30.6 | 130.6 | 663.9 |
| 1999..... | 24.6 | 7.7 | 0.8 | 0.2 | 0.4 | 0.7 | 2.0 | 3.4 | 9.1 | 28.4 | 128.0 | 672.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.0 | 5.2 | 0.6 | * | 0.4 | 0.6 | 1.3 | 3.7 | 9.6 | 32.9 | 124.9 | 479.5 |
| 2003..... | 7.2 | 6.5 | 0.7 | * | 0.5 | 0.7 | 1.2 | 3.9 | 10.4 | 34.8 | 132.4 | 511.0 |
| 2002..... | 7.2 | 5.5 | * | * | * | 0.6 | 1.3 | 3.4 | 10.5 | 33.6 | 142.1 | 563.2 |
| 2001..... | 7.3 | 7.0 | 1.0 | * | * | 0.3 | 1.3 | 3.5 | 9.1 | 35.3 | 136.3 | 660.3 |
| 2000..... | 7.3 | 7.5 | * | * | 0.3 | 0.7 | 1.6 | 3.8 | 11.0 | 37.2 | 134.3 | 633.0 |
| 1999..... | 6.5 | 7.9 | * | * | 0.4 | 0.4 | 1.5 | 3.4 | 9.2 | 30.9 | 116.7 | 607.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.5 | 7.3 | * | * | 0.6 | 0.7 | 1.5 | 4.9 | 11.5 | 38.5 | 151.3 | 526.8 |
| 2003..... | 6.7 | 7.5 | * | * | * | 0.8 | 1.3 | 5.2 | 11.4 | 42.8 | 163.0 | 571.1 |
| 2002..... | 6.9 | 7.3 | * | * | * | 0.6 | 1.7 | 4.7 | 13.9 | 43.6 | 172.8 | 654.2 |
| 2001..... | 6.8 | 7.4 | * | * | * | * | 1.7 | 5.0 | 11.4 | 40.7 | 172.6 | 765.6 |
| 2000..... | 7.1 | 7.6 | * | * | * | 0.7 | 2.1 | 5.2 | 14.6 | 45.6 | 178.0 | 751.8 |
| 1999..... | 6.1 | 10.3 | * | * | * | * | 2.0 | 4.7 | 9.9 | 39.5 | 147.0 | 718.2 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.5 | * | * | * | * | * | 1.2 | 2.5 | 7.9 | 28.4 | 106.7 | 455.2 |
| 2003..... | 7.7 | 5.4 | * | * | * | * | 1.1 | 2.6 | 9.6 | 28.4 | 111.4 | 480.5 |
| 2002..... | 7.5 | * | * | * | * | * | 0.9 | 2.1 | 7.4 | 25.7 | 121.1 | 518.1 |
| 2001..... | 7.8 | 6.5 | * | * | * | * | 0.9 | 2.1 | 7.1 | 31.0 | 111.8 | 608.8 |
| 2000..... | 7.6 | 7.4 | * | * | * | * | 1.0 | 2.3 | 7.9 | 30.7 | 105.0 | 574.7 |
| 1999..... | 6.8 | 5.4 | * | * | * | * | 0.8 | 2.2 | 8.5 | 24.1 | 96.4 | 554.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonia (J12-J18) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 22.0 | 6.6 | 0.6 | 0.2 | 0.4 | 0.8 | 2.1 | 4.6 | 10.7 | 33.9 | 137.4 | 574.2 |
| 2003..... | 24.0 | 7.4 | 0.6 | 0.3 | 0.5 | 0.9 | 2.3 | 5.2 | 10.9 | 36.2 | 148.0 | 653.6 |
| 2002..... | 24.8 | 6.5 | 0.7 | 0.2 | 0.4 | 0.9 | 2.3 | 4.9 | 11.1 | 37.3 | 155.6 | 691.2 |
| 2001..... | 23.8 | 7.2 | 0.6 | 0.2 | 0.5 | 0.9 | 2.3 | 4.6 | 10.6 | 36.0 | 148.2 | 682.5 |
| 2000..... | 24.7 | 7.3 | 0.6 | 0.2 | 0.5 | 0.9 | 2.4 | 4.6 | 11.4 | 37.9 | 156.6 | 725.2 |
| 1999..... | 24.3 | 8.1 | 0.8 | 0.2 | 0.4 | 0.9 | 2.4 | 4.6 | 10.7 | 36.5 | 154.2 | 732.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 20.3 | 7.5 | 0.5 | 0.2 | 0.5 | 1.0 | 2.3 | 5.6 | 12.8 | 42.0 | 171.0 | 671.6 |
| 2003..... | 21.7 | 8.7 | 0.7 | 0.2 | 0.5 | 1.1 | 2.7 | 6.3 | 12.9 | 45.3 | 183.9 | 747.1 |
| 2002..... | 22.3 | 7.6 | 0.8 | 0.2 | 0.5 | 1.0 | 2.5 | 5.8 | 13.4 | 45.9 | 193.5 | 801.5 |
| 2001..... | 21.4 | 7.8 | 0.6 | 0.2 | 0.5 | 1.0 | 2.6 | 5.7 | 13.0 | 44.8 | 186.1 | 805.9 |
| 2000..... | 22.2 | 8.2 | 0.7 | 0.2 | 0.5 | 1.0 | 2.8 | 5.6 | 13.9 | 46.8 | 195.7 | 873.8 |
| 1999..... | 21.7 | 8.0 | 0.8 | 0.2 | 0.5 | 1.1 | 2.6 | 5.8 | 12.5 | 46.2 | 193.4 | 877.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 23.7 | 5.6 | 0.8 | 0.2 | 0.3 | 0.6 | 1.8 | 3.6 | 8.7 | 27.1 | 114.8 | 530.7 |
| 2003..... | 26.3 | 6.1 | 0.6 | 0.3 | 0.4 | 0.7 | 2.0 | 4.1 | 9.0 | 28.5 | 124.1 | 612.4 |
| 2002..... | 27.2 | 5.3 | 0.6 | 0.2 | 0.4 | 0.8 | 2.0 | 4.0 | 9.1 | 30.1 | 130.6 | 643.7 |
| 2001..... | 26.0 | 6.6 | 0.6 | 0.2 | 0.4 | 0.8 | 2.0 | 3.6 | 8.5 | 28.8 | 123.3 | 631.3 |
| 2000..... | 27.1 | 6.4 | 0.5 | 0.1 | 0.4 | 0.8 | 2.0 | 3.7 | 9.2 | 30.4 | 131.2 | 665.2 |
| 1999..... | 26.8 | 8.2 | 0.9 | 0.2 | 0.4 | 0.8 | 2.2 | 3.5 | 9.1 | 28.5 | 128.8 | 674.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 24.3 | 4.6 | 0.5 | 0.1 | 0.4 | 0.7 | 1.8 | 4.0 | 9.9 | 32.6 | 136.3 | 581.1 |
| 2003..... | 26.6 | 5.7 | 0.6 | 0.3 | 0.5 | 0.8 | 2.0 | 4.5 | 9.8 | 34.6 | 148.0 | 665.7 |
| 2002..... | 27.6 | 4.7 | 0.5 | 0.2 | 0.4 | 0.8 | 2.0 | 4.4 | 10.2 | 36.4 | 155.3 | 703.1 |
| 2001..... | 26.1 | 4.6 | 0.5 | 0.2 | 0.4 | 0.8 | 1.9 | 4.0 | 9.8 | 34.9 | 146.7 | 691.5 |
| 2000..... | 27.1 | 4.8 | 0.5 | 0.2 | 0.4 | 0.8 | 2.1 | 3.9 | 10.3 | 35.6 | 155.9 | 738.1 |
| 1999..... | 26.8 | 5.2 | 0.6 | 0.2 | 0.4 | 0.7 | 1.9 | 3.9 | 9.7 | 35.2 | 154.8 | 743.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 22.0 | 5.3 | * | * | 0.5 | 0.9 | 2.0 | 4.8 | 11.8 | 39.5 | 168.2 | 678.4 |
| 2003..... | 23.5 | 6.8 | 0.6 | 0.2 | 0.5 | 1.0 | 2.3 | 5.4 | 11.1 | 42.5 | 181.5 | 755.2 |
| 2002..... | 24.3 | 6.0 | 0.5 | 0.2 | 0.5 | 0.9 | 2.2 | 5.1 | 11.9 | 44.5 | 191.5 | 810.6 |
| 2001..... | 23.0 | 5.3 | 0.6 | 0.2 | 0.5 | 0.9 | 2.1 | 4.8 | 11.6 | 42.6 | 182.0 | 811.5 |
| 2000..... | 23.8 | 5.4 | 0.4 | 0.2 | 0.5 | 0.8 | 2.4 | 4.6 | 12.2 | 43.3 | 193.2 | 886.1 |
| 1999..... | 23.4 | 5.4 | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.7 | 11.1 | 43.8 | 192.7 | 886.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 26.5 | 3.8 | 0.6 | * | 0.2 | 0.5 | 1.6 | 3.2 | 8.1 | 26.6 | 114.6 | 537.5 |
| 2003..... | 29.6 | 4.5 | 0.5 | 0.3 | 0.4 | 0.6 | 1.8 | 3.7 | 8.5 | 27.9 | 125.4 | 626.4 |
| 2002..... | 30.7 | 3.4 | 0.5 | 0.2 | 0.3 | 0.8 | 1.8 | 3.7 | 8.5 | 29.5 | 131.1 | 656.6 |
| 2001..... | 29.1 | 3.8 | 0.4 | 0.2 | 0.3 | 0.7 | 1.7 | 3.2 | 8.0 | 28.3 | 123.3 | 641.7 |
| 2000..... | 30.3 | 4.2 | 0.5 | * | 0.3 | 0.8 | 1.7 | 3.3 | 8.5 | 29.1 | 131.4 | 678.4 |
| 1999..... | 30.0 | 4.9 | 0.7 | 0.2 | 0.3 | 0.6 | 1.8 | 3.0 | 8.4 | 28.0 | 130.0 | 686.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonia (J12-J18) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 15.1 | 14.0 | 1.2 | * | 0.6 | 1.2 | 4.0 | 9.2 | 18.9 | 50.0 | 156.7 | 509.2 |
| 2003..... | 15.6 | 13.3 | 0.9 | * | 0.6 | 1.9 | 4.5 | 10.2 | 20.8 | 54.4 | 156.4 | 522.4 |
| 2002..... | 16.0 | 13.1 | 1.5 | * | 0.6 | 1.6 | 4.5 | 8.9 | 21.4 | 50.5 | 167.3 | 577.4 |
| 2001..... | 15.9 | 16.9 | 1.0 | * | 0.8 | 1.7 | 4.9 | 9.9 | 19.0 | 48.7 | 169.1 | 581.5 |
| 2000..... | 16.6 | 17.2 | 1.2 | * | 0.8 | 1.8 | 4.5 | 10.1 | 22.1 | 59.9 | 170.8 | 595.0 |
| 1999..... | 16.5 | 19.3 | 1.7 | * | 0.9 | 2.0 | 5.4 | 10.7 | 20.7 | 54.0 | 156.6 | 628.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 15.4 | 15.8 | * | * | * | 1.6 | 4.9 | 12.0 | 24.7 | 72.7 | 218.2 | 606.6 |
| 2003..... | 16.2 | 15.2 | * | * | 0.7 | 2.3 | 5.4 | 13.0 | 29.4 | 78.5 | 219.9 | 639.3 |
| 2002..... | 15.8 | 13.9 | 1.6 | * | * | 1.8 | 5.2 | 11.5 | 29.3 | 66.0 | 230.6 | 693.6 |
| 2001..... | 16.3 | 16.6 | * | * | 0.8 | 2.0 | 5.9 | 13.5 | 26.4 | 69.7 | 242.5 | 716.1 |
| 2000..... | 17.1 | 18.8 | 1.7 | * | 0.7 | 2.3 | 5.7 | 14.0 | 30.1 | 83.4 | 233.9 | 750.3 |
| 1999..... | 16.8 | 19.1 | 2.0 | * | 0.8 | 2.3 | 6.6 | 15.1 | 27.2 | 77.0 | 213.3 | 799.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 14.8 | 12.2 | 1.7 | * | 0.7 | 0.8 | 3.3 | 6.9 | 14.3 | 33.8 | 122.0 | 470.6 |
| 2003..... | 15.0 | 11.3 | * | * | * | 1.5 | 3.7 | 7.8 | 13.9 | 37.2 | 120.6 | 477.1 |
| 2002..... | 16.1 | 12.3 | * | * | * | 1.4 | 3.9 | 6.6 | 15.1 | 39.5 | 131.5 | 533.0 |
| 2001..... | 15.6 | 17.1 | * | * | 0.8 | 1.3 | 4.0 | 6.9 | 13.0 | 33.9 | 127.9 | 531.7 |
| 2000..... | 16.3 | 15.5 | * | * | 0.8 | 1.2 | 3.5 | 6.8 | 15.6 | 43.3 | 135.3 | 537.3 |
| 1999..... | 16.3 | 19.6 | * | * | 0.9 | 1.7 | 4.4 | 6.9 | 15.5 | 37.7 | 124.8 | 565.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Viral pneumonia, not elsewhere classified (J12) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.3 | * | * | * | * | * | * | * | * | 0.4 | 1.3 |
| 2003..... | 0.1 | 1.6 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 |
| 2002..... | 0.1 | 1.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.7 |
| 2001..... | 0.1 | 1.9 | * | * | * | * | * | * | * | 0.1 | 0.5 | 1.5 |
| 2000..... | 0.1 | 1.5 | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.9 |
| 1999..... | 0.2 | 2.0 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.6 | * | * | * | * | * | * | * | * | 0.4 | * |
| 2003..... | 0.1 | 1.7 | * | * | * | * | * | * | * | * | 0.6 | 2.1 |
| 2002..... | 0.1 | 1.8 | * | * | * | * | * | * | * | * | 0.7 | 2.0 |
| 2001..... | 0.1 | 2.3 | * | * | * | * | * | * | * | * | 0.8 | 1.9 |
| 2000..... | 0.1 | 1.7 | * | * | * | * | * | * | * | * | 0.8 | 2.5 |
| 1999..... | 0.1 | 2.1 | * | * | * | * | * | * | * | * | 1.3 | 1.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.4 |
| 2003..... | 0.1 | 1.5 | * | * | * | * | * | * | * | * | 0.4 | 2.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.7 |
| 2001..... | 0.1 | 1.5 | * | * | * | * | * | * | * | * | 0.3 | 1.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 3.0 |
| 1999..... | 0.2 | 1.9 | * | * | * | * | * | * | * | 0.3 | 0.5 | 2.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | 0.9 | * | * | * | * | * | * | * | * | 0.5 | 1.4 |
| 2003..... | 0.1 | 1.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.3 |
| 2002..... | 0.1 | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 |
| 2001..... | 0.1 | 1.3 | * | * | * | * | * | * | * | * | 0.5 | 1.6 |
| 2000..... | 0.2 | 1.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 3.1 |
| 1999..... | 0.2 | 1.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 2.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.9 | 2.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.2 | 2.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.5 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 3.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 2.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bacterial pneumonia (J13-J15) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | 0.8 | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 3.4 | 10.5 | 30.1 |
| 2003..... | 1.6 | 1.0 | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 | 0.8 | 1.4 | 3.7 | 11.1 | 34.9 |
| 2002..... | 1.6 | 1.1 | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 3.6 | 11.3 | 35.7 |
| 2001..... | 1.6 | 0.8 | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.5 | 11.0 | 34.8 |
| 2000..... | 1.6 | 0.8 | * | * | * | 0.1 | 0.4 | 0.7 | 1.6 | 3.7 | 11.0 | 35.7 |
| 1999..... | 1.4 | 1.3 | 0.1 | * | * | 0.1 | 0.4 | 0.5 | 1.3 | 3.3 | 9.8 | 30.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 4.2 | 14.1 | 41.1 |
| 2003..... | 1.7 | 1.2 | * | * | * | 0.1 | 0.4 | 0.9 | 1.7 | 4.9 | 14.5 | 45.7 |
| 2002..... | 1.6 | 1.2 | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 4.4 | 15.1 | 46.8 |
| 2001..... | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 4.5 | 15.1 | 46.8 |
| 2000..... | 1.7 | 1.1 | * | * | * | 0.1 | 0.4 | 0.8 | 1.8 | 4.8 | 15.6 | 52.3 |
| 1999..... | 1.4 | 1.2 | * | * | * | * | 0.4 | 0.7 | 1.4 | 4.2 | 12.5 | 40.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 2.7 | 8.0 | 25.2 |
| 2003..... | 1.6 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.7 | 8.8 | 30.2 |
| 2002..... | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.9 | 8.8 | 31.0 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.6 | 8.3 | 29.8 |
| 2000..... | 1.5 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 2.9 | 8.0 | 29.0 |
| 1999..... | 1.4 | 1.5 | * | * | * | * | 0.4 | 0.4 | 1.2 | 2.6 | 7.9 | 26.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.7 | 1.2 | 4.6 | 13.6 | 34.2 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.6 | 1.5 | 4.6 | 13.7 | 41.5 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.9 | 1.4 | 3.0 | 10.0 | 41.0 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 1.5 | 4.0 | 12.1 | 50.0 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.7 | 2.0 | 4.1 | 11.7 | 45.8 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 1.4 | 3.3 | 9.8 | 37.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 1.0 | * | 5.8 | 16.5 | 47.6 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | * | 5.7 | 16.5 | 47.7 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 4.2 | 9.0 | 51.9 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 6.4 | 15.2 | 49.5 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | * | 5.5 | 16.2 | 66.5 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 13.6 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 3.5 | 11.6 | 27.4 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | * | 3.7 | 11.9 | 38.3 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | * | 10.7 | 35.6 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | * | 10.0 | 50.3 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 8.6 | 35.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | * | 7.3 | 38.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bacterial pneumonia (J13-J15) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | 0.8 | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 3.3 | 10.3 | 29.8 |
| 2003..... | 1.8 | 0.9 | 0.2 | 0.1 | * | 0.1 | 0.3 | 0.8 | 1.4 | 3.7 | 11.0 | 34.6 |
| 2002..... | 1.7 | 1.0 | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.7 | 11.3 | 35.3 |
| 2001..... | 1.7 | 0.6 | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.4 | 10.9 | 34.1 |
| 2000..... | 1.7 | 0.9 | * | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 3.7 | 10.9 | 35.3 |
| 1999..... | 1.5 | 1.5 | * | * | * | 0.1 | 0.4 | 0.6 | 1.3 | 3.3 | 9.7 | 29.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 4.0 | 14.0 | 40.7 |
| 2003..... | 1.8 | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.6 | 4.9 | 14.4 | 45.4 |
| 2002..... | 1.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 4.4 | 15.3 | 46.4 |
| 2001..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.4 | 15.1 | 46.5 |
| 2000..... | 1.8 | * | * | * | * | * | 0.4 | 0.7 | 1.8 | 4.7 | 15.5 | 51.6 |
| 1999..... | 1.5 | * | * | * | * | * | 0.4 | 0.7 | 1.3 | 4.2 | 12.5 | 40.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 2.6 | 7.8 | 25.0 |
| 2003..... | 1.7 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.7 | 8.7 | 29.8 |
| 2002..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 3.0 | 8.7 | 30.6 |
| 2001..... | 1.6 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.6 | 8.2 | 29.0 |
| 2000..... | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 2.9 | 8.0 | 28.6 |
| 1999..... | 1.5 | 1.8 | * | * | * | * | 0.4 | 0.4 | 1.2 | 2.5 | 8.0 | 25.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.1 | 10.0 | 29.6 |
| 2003..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 3.6 | 10.9 | 34.6 |
| 2002..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.6 | 11.2 | 35.8 |
| 2001..... | 1.8 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 3.3 | 10.8 | 34.2 |
| 2000..... | 1.8 | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.4 | 3.5 | 10.7 | 35.0 |
| 1999..... | 1.6 | 1.0 | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 3.2 | 9.7 | 29.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.2 | 0.7 | 1.3 | 3.7 | 13.5 | 41.1 |
| 2003..... | 1.9 | * | * | * | * | * | 0.3 | 0.8 | 1.3 | 4.7 | 14.3 | 45.7 |
| 2002..... | 1.9 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 4.3 | 14.9 | 47.5 |
| 2001..... | 1.8 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 4.1 | 14.6 | 46.2 |
| 2000..... | 1.9 | * | * | * | * | * | 0.4 | 0.6 | 1.6 | 4.4 | 15.2 | 50.2 |
| 1999..... | 1.5 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 4.0 | 12.3 | 40.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 2.6 | 7.5 | 24.4 |
| 2003..... | 1.9 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.7 | 8.6 | 29.7 |
| 2002..... | 1.9 | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 3.0 | 8.7 | 30.8 |
| 2001..... | 1.8 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.6 | 8.3 | 29.2 |
| 2000..... | 1.7 | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 2.7 | 7.7 | 28.9 |
| 1999..... | 1.6 | * | * | * | * | * | 0.4 | 0.4 | 1.1 | 2.5 | 8.0 | 25.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bacterial pneumonia (J13-J15) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.4 | 1.3 | 2.5 | 4.7 | 13.1 | 35.4 |
| 2003..... | 1.4 | * | * | * | * | * | 0.6 | 1.3 | 2.8 | 4.1 | 11.8 | 34.2 |
| 2002..... | 1.2 | * | * | * | * | * | 0.7 | 1.0 | 2.3 | 4.0 | 11.4 | 28.1 |
| 2001..... | 1.3 | * | * | * | * | * | 0.6 | 1.0 | 2.2 | 4.7 | 12.5 | 33.1 |
| 2000..... | 1.4 | * | * | * | * | * | 0.5 | 1.2 | 2.4 | 5.8 | 12.3 | 35.0 |
| 1999..... | 1.2 | * | * | * | * | * | 0.8 | 1.0 | 2.4 | 4.7 | 9.7 | 24.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 1.6 | 3.7 | 7.6 | 18.3 | 42.7 |
| 2003..... | 1.5 | * | * | * | * | * | 0.9 | 1.7 | 4.3 | 5.8 | 15.6 | 40.5 |
| 2002..... | 1.3 | * | * | * | * | * | * | 1.2 | 3.3 | 5.0 | 17.9 | 31.7 |
| 2001..... | 1.5 | * | * | * | * | * | 0.9 | 1.5 | 2.9 | 7.4 | 20.1 | 50.8 |
| 2000..... | 1.5 | * | * | * | * | * | * | 1.3 | 3.4 | 7.9 | 17.9 | 61.1 |
| 1999..... | 1.3 | * | * | * | * | * | 0.9 | 1.6 | 2.7 | 6.6 | 14.4 | 39.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 1.0 | 1.5 | 2.7 | 10.1 | 32.5 |
| 2003..... | 1.2 | * | * | * | * | * | * | 1.0 | 1.5 | 2.9 | 9.6 | 31.8 |
| 2002..... | 1.2 | * | * | * | * | * | * | 0.9 | 1.4 | 3.3 | 7.8 | 26.8 |
| 2001..... | 1.0 | * | * | * | * | * | * | * | 1.7 | 2.7 | 8.3 | 26.5 |
| 2000..... | 1.2 | * | * | * | * | * | * | 1.1 | 1.6 | 4.3 | 9.2 | 25.3 |
| 1999..... | 1.1 | * | * | * | * | * | 0.8 | * | 2.2 | 3.4 | 7.0 | 19.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonia due to other or unspecified organisms (J16,J18) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 18.4 | 4.1 | 0.4 | 0.1 | 0.4 | 0.6 | 1.7 | 3.8 | 9.2 | 30.6 | 126.1 | 539.8 |
| 2003..... | 20.0 | 4.6 | 0.4 | 0.2 | 0.4 | 0.8 | 1.8 | 4.3 | 9.5 | 32.4 | 135.9 | 612.1 |
| 2002..... | 20.8 | 3.9 | 0.5 | 0.2 | 0.3 | 0.7 | 1.8 | 4.1 | 9.7 | 33.4 | 143.7 | 650.3 |
| 2001..... | 20.0 | 4.6 | 0.5 | 0.2 | 0.4 | 0.7 | 1.9 | 3.9 | 9.1 | 32.6 | 136.6 | 647.1 |
| 2000..... | 20.8 | 5.1 | 0.4 | 0.1 | 0.4 | 0.7 | 1.9 | 3.9 | 9.8 | 34.0 | 144.6 | 685.3 |
| 1999..... | 20.7 | 4.7 | 0.5 | 0.1 | 0.4 | 0.7 | 1.9 | 4.0 | 9.4 | 32.8 | 142.8 | 698.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 16.7 | 4.9 | 0.3 | 0.1 | 0.5 | 0.8 | 1.9 | 4.7 | 11.3 | 37.8 | 155.8 | 623.9 |
| 2003..... | 17.8 | 5.5 | 0.4 | 0.2 | 0.5 | 0.9 | 2.1 | 5.3 | 11.1 | 40.3 | 168.0 | 693.0 |
| 2002..... | 18.5 | 4.6 | 0.5 | 0.1 | 0.4 | 0.8 | 2.1 | 5.0 | 11.8 | 41.4 | 177.4 | 748.2 |
| 2001..... | 17.8 | 4.9 | 0.5 | 0.2 | 0.4 | 0.8 | 2.1 | 4.9 | 11.2 | 40.2 | 170.0 | 757.4 |
| 2000..... | 18.5 | 5.3 | 0.4 | 0.2 | 0.4 | 0.9 | 2.3 | 4.8 | 12.1 | 42.0 | 179.5 | 816.5 |
| 1999..... | 18.3 | 5.0 | 0.5 | 0.2 | 0.4 | 0.8 | 2.2 | 5.1 | 11.0 | 41.7 | 178.5 | 832.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 20.0 | 3.3 | 0.5 | 0.1 | 0.3 | 0.5 | 1.4 | 2.9 | 7.3 | 24.5 | 106.2 | 502.0 |
| 2003..... | 22.2 | 3.8 | 0.5 | 0.1 | 0.3 | 0.6 | 1.6 | 3.4 | 7.9 | 25.7 | 114.5 | 576.3 |
| 2002..... | 23.0 | 3.1 | 0.4 | 0.2 | 0.3 | 0.6 | 1.5 | 3.2 | 7.8 | 26.7 | 121.3 | 607.9 |
| 2001..... | 22.2 | 4.2 | 0.5 | 0.2 | 0.3 | 0.6 | 1.6 | 2.9 | 7.1 | 26.3 | 114.6 | 601.0 |
| 2000..... | 23.1 | 4.9 | 0.4 | 0.1 | 0.4 | 0.6 | 1.6 | 3.0 | 7.7 | 27.4 | 121.9 | 631.9 |
| 1999..... | 23.0 | 4.3 | 0.5 | 0.1 | 0.4 | 0.6 | 1.6 | 2.9 | 7.9 | 25.6 | 119.6 | 643.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 6.1 | 2.8 | * | * | 0.4 | 0.5 | 1.1 | 3.0 | 8.4 | 28.3 | 111.0 | 444.3 |
| 2003..... | 6.3 | 3.6 | * | * | 0.3 | 0.6 | 0.9 | 3.3 | 9.0 | 30.2 | 118.2 | 468.0 |
| 2002..... | 6.5 | 2.9 | * | * | * | 0.4 | 1.1 | 2.4 | 9.0 | 30.5 | 131.3 | 521.6 |
| 2001..... | 6.5 | 3.9 | 0.7 | * | * | 0.3 | 1.0 | 3.0 | 7.5 | 31.1 | 124.1 | 609.6 |
| 2000..... | 6.4 | 5.6 | * | * | * | 0.6 | 1.2 | 2.9 | 9.0 | 32.4 | 121.9 | 584.6 |
| 1999..... | 5.8 | 4.8 | * | * | 0.3 | 0.3 | 1.1 | 3.1 | 7.8 | 27.3 | 106.4 | 569.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 0.5 | 0.5 | 1.3 | 3.9 | 10.5 | 32.5 | 134.5 | 477.7 |
| 2003..... | 5.8 | * | * | * | * | 0.6 | 0.9 | 4.3 | 9.7 | 37.1 | 145.7 | 518.7 |
| 2002..... | 6.2 | * | * | * | * | 0.5 | 1.4 | 3.6 | 11.9 | 39.0 | 163.3 | 600.7 |
| 2001..... | 6.0 | 4.8 | * | * | * | * | 1.3 | 4.0 | 9.2 | 34.1 | 156.9 | 714.2 |
| 2000..... | 6.1 | 5.3 | * | * | * | 0.6 | 1.5 | 4.0 | 12.2 | 39.0 | 161.3 | 685.2 |
| 1999..... | 5.4 | 6.2 | * | * | * | * | 1.6 | 4.1 | 7.9 | 35.6 | 133.5 | 683.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | * | 0.9 | 2.1 | 6.5 | 24.8 | 94.8 | 427.1 |
| 2003..... | 6.8 | * | * | * | * | * | 1.0 | 2.3 | 8.3 | 24.7 | 99.3 | 442.2 |
| 2002..... | 6.8 | * | * | * | * | * | 0.8 | 1.3 | 6.4 | 23.7 | 109.3 | 482.4 |
| 2001..... | 7.1 | * | * | * | * | * | * | 1.9 | 5.9 | 28.7 | 101.8 | 558.5 |
| 2000..... | 6.8 | 5.8 | * | * | * | * | 0.8 | 1.9 | 6.2 | 27.2 | 95.4 | 535.2 |
| 1999..... | 6.2 | * | * | * | * | * | * | 2.1 | 7.6 | 20.8 | 88.5 | 514.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonia due to other or unspecified organisms (J16,J18) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.3 | 4.5 | 0.4 | 0.1 | 0.4 | 0.7 | 1.8 | 3.9 | 9.2 | 30.6 | 126.7 | 543.1 |
| 2003..... | 22.2 | 4.9 | 0.4 | 0.2 | 0.4 | 0.8 | 2.0 | 4.4 | 9.4 | 32.4 | 136.6 | 616.9 |
| 2002..... | 23.0 | 4.1 | 0.5 | 0.2 | 0.4 | 0.8 | 1.9 | 4.2 | 9.7 | 33.5 | 143.8 | 654.2 |
| 2001..... | 22.0 | 4.7 | 0.5 | 0.2 | 0.4 | 0.8 | 2.0 | 4.0 | 9.1 | 32.5 | 136.7 | 646.9 |
| 2000..... | 22.8 | 4.9 | 0.4 | 0.1 | 0.4 | 0.8 | 2.0 | 3.9 | 9.8 | 33.9 | 145.0 | 687.1 |
| 1999..... | 22.7 | 4.6 | 0.5 | 0.2 | 0.4 | 0.8 | 2.0 | 4.0 | 9.4 | 33.0 | 143.7 | 700.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 18.5 | 5.3 | * | 0.1 | 0.5 | 0.8 | 2.0 | 4.8 | 11.3 | 37.9 | 156.6 | 629.9 |
| 2003..... | 19.8 | 5.9 | 0.4 | 0.2 | 0.5 | 1.0 | 2.3 | 5.3 | 11.2 | 40.3 | 168.9 | 699.6 |
| 2002..... | 20.4 | 4.9 | 0.6 | 0.2 | 0.4 | 0.8 | 2.2 | 5.1 | 11.7 | 41.3 | 177.5 | 753.1 |
| 2001..... | 19.6 | 4.9 | 0.5 | 0.2 | 0.5 | 0.9 | 2.2 | 4.9 | 11.2 | 40.2 | 170.2 | 757.4 |
| 2000..... | 20.2 | 5.2 | 0.4 | 0.2 | 0.5 | 0.9 | 2.4 | 4.8 | 12.0 | 41.9 | 179.4 | 819.6 |
| 1999..... | 20.1 | 4.8 | 0.5 | 0.2 | 0.4 | 0.9 | 2.2 | 5.1 | 11.1 | 41.8 | 179.6 | 834.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 22.0 | 3.7 | 0.6 | * | 0.3 | 0.5 | 1.5 | 3.0 | 7.3 | 24.4 | 106.6 | 504.3 |
| 2003..... | 24.4 | 4.0 | 0.4 | 0.2 | 0.4 | 0.7 | 1.7 | 3.5 | 7.8 | 25.7 | 115.0 | 580.6 |
| 2002..... | 25.4 | 3.3 | 0.5 | 0.2 | 0.3 | 0.7 | 1.6 | 3.4 | 7.9 | 26.9 | 121.4 | 611.5 |
| 2001..... | 24.3 | 4.5 | 0.4 | 0.2 | 0.4 | 0.7 | 1.7 | 3.0 | 7.2 | 26.1 | 114.8 | 601.0 |
| 2000..... | 25.3 | 4.7 | 0.4 | 0.1 | 0.4 | 0.7 | 1.6 | 3.1 | 7.8 | 27.3 | 122.6 | 633.5 |
| 1999..... | 25.1 | 4.5 | 0.5 | 0.1 | 0.4 | 0.7 | 1.7 | 3.0 | 7.8 | 25.8 | 120.4 | 646.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 22.5 | 3.0 | 0.2 | 0.1 | 0.4 | 0.6 | 1.5 | 3.4 | 8.6 | 29.3 | 125.8 | 550.1 |
| 2003..... | 24.6 | 3.7 | 0.3 | 0.2 | 0.4 | 0.7 | 1.7 | 3.8 | 8.5 | 30.8 | 136.7 | 628.9 |
| 2002..... | 25.6 | 2.9 | 0.4 | 0.2 | 0.3 | 0.7 | 1.7 | 3.8 | 8.9 | 32.6 | 143.6 | 665.4 |
| 2001..... | 24.2 | 2.8 | 0.4 | 0.2 | 0.4 | 0.7 | 1.6 | 3.4 | 8.3 | 31.5 | 135.4 | 655.8 |
| 2000..... | 25.2 | 3.3 | 0.3 | 0.1 | 0.4 | 0.7 | 1.7 | 3.3 | 8.7 | 31.9 | 144.4 | 700.0 |
| 1999..... | 25.0 | 2.8 | 0.4 | 0.1 | 0.3 | 0.6 | 1.6 | 3.4 | 8.5 | 31.8 | 144.3 | 711.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 20.2 | 3.3 | * | * | 0.5 | 0.8 | 1.7 | 4.1 | 10.5 | 35.8 | 154.1 | 636.3 |
| 2003..... | 21.5 | 4.5 | * | 0.2 | 0.4 | 0.9 | 2.0 | 4.6 | 9.8 | 37.5 | 166.6 | 707.3 |
| 2002..... | 22.4 | 3.9 | * | * | 0.4 | 0.7 | 1.9 | 4.5 | 10.5 | 40.0 | 175.9 | 761.2 |
| 2001..... | 21.1 | 3.2 | * | 0.2 | 0.4 | 0.7 | 1.8 | 4.1 | 9.9 | 38.3 | 166.6 | 763.3 |
| 2000..... | 21.8 | 3.8 | * | 0.2 | 0.4 | 0.7 | 2.0 | 3.9 | 10.4 | 38.7 | 177.2 | 833.0 |
| 1999..... | 21.7 | 3.0 | * | 0.2 | 0.3 | 0.7 | 1.7 | 4.2 | 9.8 | 39.5 | 179.2 | 844.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 24.7 | 2.7 | * | * | 0.2 | 0.5 | 1.3 | 2.7 | 6.8 | 23.8 | 106.6 | 511.5 |
| 2003..... | 27.6 | 2.9 | * | * | 0.4 | 0.5 | 1.5 | 3.0 | 7.4 | 25.0 | 116.4 | 594.4 |
| 2002..... | 28.6 | 1.8 | * | 0.2 | 0.3 | 0.6 | 1.4 | 3.1 | 7.4 | 26.3 | 121.9 | 624.1 |
| 2001..... | 27.2 | 2.4 | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.6 | 6.7 | 25.7 | 114.7 | 611.1 |
| 2000..... | 28.3 | 2.7 | * | * | 0.3 | 0.6 | 1.4 | 2.7 | 7.1 | 26.1 | 122.9 | 646.3 |
| 1999..... | 28.2 | 2.5 | 0.5 | * | 0.3 | 0.5 | 1.4 | 2.6 | 7.2 | 25.3 | 121.5 | 658.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonia due to other or unspecified organisms (J16,J18) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 13.6 | 10.7 | 1.0 | * | 0.6 | 1.0 | 3.6 | 8.0 | 16.4 | 45.2 | 143.7 | 473.5 |
| 2003..... | 14.2 | 9.2 | * | * | 0.5 | 1.7 | 3.9 | 8.9 | 17.9 | 50.3 | 144.4 | 487.6 |
| 2002..... | 14.6 | 8.5 | 1.0 | * | 0.5 | 1.3 | 3.8 | 7.8 | 19.2 | 46.4 | 155.7 | 548.4 |
| 2001..... | 14.5 | 12.1 | * | * | 0.7 | 1.4 | 4.3 | 8.9 | 16.7 | 43.9 | 156.4 | 548.1 |
| 2000..... | 15.2 | 11.8 | 0.8 | * | 0.7 | 1.6 | 4.0 | 8.9 | 19.5 | 53.9 | 158.3 | 559.7 |
| 1999..... | 15.1 | 12.1 | 1.1 | * | 0.8 | 1.8 | 4.6 | 9.6 | 18.3 | 49.1 | 146.4 | 602.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 13.8 | 13.0 | * | * | * | 1.3 | 4.5 | 10.4 | 21.0 | 65.1 | 199.9 | 562.9 |
| 2003..... | 14.7 | 10.6 | * | * | 0.7 | 2.0 | 4.6 | 11.3 | 24.9 | 72.7 | 203.7 | 597.7 |
| 2002..... | 14.4 | 8.5 | * | * | * | 1.6 | 4.5 | 10.3 | 26.0 | 61.1 | 212.2 | 659.7 |
| 2001..... | 14.7 | 11.5 | * | * | 0.7 | 1.9 | 5.0 | 11.9 | 23.5 | 62.0 | 222.4 | 664.1 |
| 2000..... | 15.4 | 10.9 | * | * | * | 2.1 | 5.1 | 12.7 | 26.5 | 75.3 | 216.0 | 689.3 |
| 1999..... | 15.2 | 12.8 | * | * | 0.8 | 2.1 | 5.7 | 13.5 | 24.5 | 70.3 | 197.3 | 759.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 13.5 | 8.4 | * | * | * | * | 2.7 | 5.8 | 12.7 | 31.0 | 111.9 | 438.1 |
| 2003..... | 13.8 | 7.7 | * | * | * | 1.4 | 3.4 | 6.8 | 12.3 | 34.3 | 111.0 | 444.9 |
| 2002..... | 14.9 | 8.5 | * | * | * | 1.1 | 3.2 | 5.7 | 13.7 | 36.0 | 123.7 | 505.8 |
| 2001..... | 14.4 | 12.8 | * | * | 0.7 | 1.0 | 3.7 | 6.3 | 11.3 | 31.1 | 119.4 | 505.2 |
| 2000..... | 15.0 | 12.7 | * | * | * | 1.2 | 3.0 | 5.7 | 14.0 | 38.8 | 125.9 | 511.5 |
| 1999..... | 15.0 | 11.4 | * | * | 0.8 | 1.5 | 3.6 | 6.4 | 13.3 | 34.1 | 117.8 | 544.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other acute lower respiratory infections (J20-J22) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.4 | * | * | * | * | * | * | * | * | 0.4 | 2.6 |
| 2003..... | 0.2 | 1.2 | 0.2 | * | * | * | * | * | * | 0.2 | 0.6 | 3.6 |
| 2002..... | 0.1 | 1.3 | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 3.4 |
| 2001..... | 0.1 | 1.3 | * | * | * | * | * | * | * | 0.2 | 0.4 | 3.3 |
| 2000..... | 0.2 | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.5 |
| 1999..... | 0.2 | 1.7 | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 6.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.6 | * | * | * | * | * | * | * | * | * | 2.5 |
| 2003..... | 0.1 | 1.4 | * | * | * | * | * | * | * | * | 0.7 | 2.4 |
| 2002..... | 0.1 | 2.1 | * | * | * | * | * | * | * | * | 0.6 | 2.9 |
| 2001..... | 0.1 | 1.2 | * | * | * | * | * | * | * | * | * | 2.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 3.7 |
| 1999..... | 0.2 | 2.0 | * | * | * | * | * | * | * | * | 1.1 | 4.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.7 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.6 | 4.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 3.6 |
| 2001..... | 0.2 | 1.4 | * | * | * | * | * | * | * | * | 0.5 | 3.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 4.8 |
| 1999..... | 0.3 | 1.5 | * | * | * | * | * | * | * | * | 0.8 | 6.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.2 | * | * | * | * | * | * | * | * | 0.4 | 2.8 |
| 2003..... | 0.2 | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.8 |
| 2002..... | 0.2 | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.5 | 3.5 |
| 2001..... | 0.1 | 1.0 | * | * | * | * | * | * | * | 0.1 | 0.5 | 3.5 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 4.7 |
| 1999..... | 0.2 | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.9 | 6.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.7 |
| 2002..... | 0.1 | 1.8 | * | * | * | * | * | * | * | * | 0.6 | 3.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.5 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.1 | 4.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.6 | 4.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 3.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 3.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 5.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | 0.8 | 7.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Acute bronchitis and bronchiolitis (J20-J21) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.4 | * | * | * | * | * | * | * | * | 0.3 | 1.4 |
| 2003..... | 0.1 | 1.1 | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 2.2 |
| 2002..... | 0.1 | 1.3 | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 2.1 |
| 2001..... | 0.1 | 1.3 | * | * | * | * | * | * | * | 0.1 | 0.3 | 2.1 |
| 2000..... | 0.1 | 1.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 2.6 |
| 1999..... | 0.1 | 1.7 | 0.2 | * | * | * | * | * | * | 0.2 | 0.6 | 3.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.6 | * | * | * | * | * | * | * | * | * | 1.7 |
| 2003..... | 0.1 | 1.4 | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | 2.0 | * | * | * | * | * | * | * | * | * | 1.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 2.2 |
| 1999..... | 0.1 | 2.0 | * | * | * | * | * | * | * | * | 0.7 | 2.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.2 |
| 2001..... | 0.1 | 1.4 | * | * | * | * | * | * | * | * | 0.4 | 2.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.7 |
| 1999..... | 0.2 | 1.4 | * | * | * | * | * | * | * | * | 0.5 | 4.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | 1.2 | * | * | * | * | * | * | * | * | 0.2 | 1.5 |
| 2003..... | 0.1 | 1.0 | * | * | * | * | * | * | * | * | 0.4 | 2.3 |
| 2002..... | 0.1 | 1.0 | * | * | * | * | * | * | * | 0.1 | 0.3 | 2.1 |
| 2001..... | 0.1 | 1.0 | * | * | * | * | * | * | * | * | 0.3 | 2.2 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 2.7 |
| 1999..... | 0.2 | 1.1 | * | * | * | * | * | * | * | 0.1 | 0.6 | 3.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | 1.7 | * | * | * | * | * | * | * | * | * | 1.7 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.3 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 2.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 2.3 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.5 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 2.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.5 | 4.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified acute lower respiratory infection (J22) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 1.2 |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.3 | 1.4 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 1.3 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.2 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.2 | 1.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.5 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.3 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.3 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.1 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 1.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 1.5 |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | 0.2 | 1.4 |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | 1.3 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | 1.4 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 2.2 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.3 | 3.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of upper respiratory tract (J30-J39) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 2.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.1 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.9 | 2.3 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 2.2 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.0 | 2.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.1 | 3.2 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.6 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.1 | 3.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.5 | 1.2 | 3.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.9 | 2.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 2.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 2.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | * | 1.0 | 3.3 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.2 | 3.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 2.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.0 | 3.1 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.5 | 1.2 | 3.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.2 | 3.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 1.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic lower respiratory diseases (J40-J47) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 41.5 | 0.9 | 0.3 | 0.3 | 0.4 | 0.6 | 2.0 | 8.4 | 40.4 | 153.8 | 366.7 | 601.7 |
| 2003..... | 43.5 | 0.8 | 0.3 | 0.3 | 0.5 | 0.7 | 2.1 | 8.7 | 43.3 | 163.2 | 383.0 | 635.1 |
| 2002..... | 43.3 | 1.0 | 0.4 | 0.3 | 0.5 | 0.8 | 2.2 | 8.7 | 42.4 | 163.0 | 386.7 | 637.6 |
| 2001..... | 43.2 | 1.0 | 0.3 | 0.3 | 0.4 | 0.7 | 2.2 | 8.5 | 44.1 | 167.9 | 379.8 | 644.7 |
| 2000..... | 43.4 | 0.9 | 0.3 | 0.3 | 0.5 | 0.7 | 2.1 | 8.6 | 44.2 | 169.4 | 386.1 | 648.6 |
| 1999..... | 44.5 | 0.9 | 0.4 | 0.3 | 0.5 | 0.8 | 2.0 | 8.5 | 47.5 | 177.2 | 397.8 | 646.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 40.6 | 1.2 | 0.4 | 0.4 | 0.5 | 0.7 | 2.0 | 8.8 | 43.1 | 172.1 | 445.6 | 811.1 |
| 2003..... | 42.4 | 1.1 | 0.5 | 0.4 | 0.5 | 0.8 | 1.9 | 9.1 | 46.5 | 183.6 | 464.9 | 865.9 |
| 2002..... | 42.9 | 1.1 | 0.6 | 0.4 | 0.6 | 0.8 | 2.2 | 9.1 | 45.2 | 184.8 | 480.8 | 894.8 |
| 2001..... | 42.7 | 1.1 | 0.4 | 0.3 | 0.5 | 0.7 | 2.0 | 8.8 | 46.9 | 191.3 | 475.1 | 916.9 |
| 2000..... | 43.5 | 1.2 | 0.4 | 0.4 | 0.6 | 0.8 | 1.9 | 9.0 | 47.8 | 195.2 | 488.5 | 967.9 |
| 1999..... | 45.6 | * | 0.4 | 0.4 | 0.6 | 0.8 | 1.8 | 8.6 | 52.3 | 210.7 | 513.2 | 996.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 42.5 | * | * | 0.2 | 0.3 | 0.5 | 2.1 | 8.1 | 38.0 | 138.4 | 313.6 | 507.4 |
| 2003..... | 44.4 | * | * | 0.2 | 0.4 | 0.6 | 2.3 | 8.2 | 40.3 | 146.0 | 328.3 | 533.0 |
| 2002..... | 43.7 | * | 0.3 | 0.3 | 0.4 | 0.7 | 2.3 | 8.2 | 39.8 | 144.9 | 324.1 | 526.0 |
| 2001..... | 43.7 | * | * | 0.2 | 0.4 | 0.7 | 2.3 | 8.1 | 41.5 | 148.5 | 317.3 | 530.8 |
| 2000..... | 43.2 | * | 0.3 | 0.3 | 0.4 | 0.7 | 2.2 | 8.3 | 41.0 | 148.2 | 319.2 | 518.5 |
| 1999..... | 43.4 | * | 0.3 | 0.2 | 0.5 | 0.9 | 2.1 | 8.4 | 43.1 | 149.8 | 322.9 | 504.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.5 | * | * | * | * | 0.4 | 1.1 | 3.1 | 12.2 | 51.7 | 158.7 | 391.5 |
| 2003..... | 8.0 | * | * | * | 0.3 | 0.4 | 1.0 | 3.5 | 12.7 | 53.3 | 171.0 | 460.2 |
| 2002..... | 7.9 | * | * | * | * | 0.4 | 1.2 | 3.5 | 13.9 | 53.8 | 176.0 | 457.3 |
| 2001..... | 7.7 | * | * | * | * | 0.3 | 1.0 | 3.6 | 14.0 | 55.8 | 169.5 | 480.2 |
| 2000..... | 7.6 | * | * | * | * | 0.3 | 1.1 | 3.3 | 14.3 | 53.1 | 181.9 | 481.1 |
| 1999..... | 8.4 | * | * | * | * | 0.5 | 1.6 | 3.9 | 14.6 | 62.6 | 194.8 | 536.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | * | 0.5 | 1.4 | 2.8 | 13.7 | 68.4 | 206.4 | 518.4 |
| 2003..... | 8.2 | * | * | * | * | * | 1.0 | 3.2 | 16.6 | 68.1 | 231.2 | 646.5 |
| 2002..... | 8.1 | * | * | * | * | * | 1.0 | 3.8 | 17.5 | 69.2 | 243.3 | 602.4 |
| 2001..... | 7.8 | * | * | * | * | * | 0.7 | 3.2 | 16.1 | 75.5 | 224.0 | 676.1 |
| 2000..... | 8.0 | * | * | * | * | * | 0.9 | 3.4 | 18.2 | 72.4 | 250.3 | 671.1 |
| 1999..... | 8.9 | * | * | * | * | * | 1.4 | 3.6 | 17.6 | 81.9 | 272.7 | 836.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.4 | * | * | * | * | * | 0.7 | 3.4 | 10.9 | 38.3 | 125.7 | 326.3 |
| 2003..... | 7.7 | * | * | * | * | * | 1.0 | 3.8 | 9.3 | 41.5 | 129.6 | 365.6 |
| 2002..... | 7.6 | * | * | * | * | * | 1.4 | 3.1 | 10.6 | 41.5 | 129.8 | 385.5 |
| 2001..... | 7.5 | * | * | * | * | * | 1.2 | 4.1 | 12.1 | 40.3 | 132.7 | 384.4 |
| 2000..... | 7.2 | * | * | * | * | * | 1.3 | 3.3 | 10.8 | 38.0 | 136.0 | 387.8 |
| 1999..... | 8.0 | * | * | * | * | * | 1.8 | 4.2 | 12.0 | 47.5 | 142.9 | 391.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic lower respiratory diseases (J40-J47) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 47.0 | 0.8 | 0.3 | 0.3 | 0.5 | 0.7 | 2.2 | 9.0 | 42.7 | 160.9 | 377.8 | 610.3 |
| 2003..... | 49.0 | 0.8 | 0.3 | 0.3 | 0.5 | 0.8 | 2.3 | 9.1 | 45.6 | 170.6 | 393.3 | 641.2 |
| 2002..... | 48.6 | 1.1 | 0.4 | 0.4 | 0.5 | 0.8 | 2.4 | 9.2 | 44.6 | 170.1 | 396.0 | 643.5 |
| 2001..... | 48.4 | 1.1 | 0.3 | 0.3 | 0.5 | 0.8 | 2.3 | 8.9 | 46.3 | 174.8 | 388.6 | 649.4 |
| 2000..... | 48.3 | 1.0 | 0.3 | 0.4 | 0.5 | 0.8 | 2.2 | 9.0 | 46.3 | 176.1 | 393.8 | 653.2 |
| 1999..... | 49.4 | 1.0 | 0.4 | 0.4 | 0.6 | 0.9 | 2.0 | 8.9 | 49.8 | 183.5 | 405.1 | 648.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 46.2 | * | 0.4 | 0.4 | 0.6 | 0.8 | 2.1 | 9.5 | 45.3 | 178.9 | 458.4 | 824.6 |
| 2003..... | 48.1 | * | 0.5 | 0.4 | 0.6 | 0.9 | 2.1 | 9.7 | 48.7 | 191.1 | 476.2 | 874.6 |
| 2002..... | 48.4 | 1.2 | 0.6 | 0.4 | 0.6 | 0.9 | 2.4 | 9.6 | 47.3 | 191.9 | 491.2 | 906.0 |
| 2001..... | 48.0 | * | 0.4 | 0.3 | 0.6 | 0.8 | 2.2 | 9.3 | 49.0 | 198.1 | 485.7 | 924.8 |
| 2000..... | 48.7 | * | 0.4 | 0.4 | 0.6 | 0.9 | 2.1 | 9.4 | 49.7 | 201.9 | 497.6 | 978.0 |
| 1999..... | 50.8 | * | 0.4 | 0.5 | 0.7 | 0.8 | 1.9 | 9.0 | 54.5 | 217.4 | 521.9 | 1,000.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 47.8 | * | * | 0.2 | 0.4 | 0.6 | 2.3 | 8.5 | 40.2 | 145.6 | 323.6 | 514.5 |
| 2003..... | 49.9 | * | * | 0.2 | 0.4 | 0.7 | 2.5 | 8.6 | 42.7 | 153.3 | 338.0 | 538.6 |
| 2002..... | 48.9 | * | * | 0.3 | 0.4 | 0.8 | 2.4 | 8.7 | 42.0 | 151.8 | 332.9 | 530.3 |
| 2001..... | 48.7 | * | * | 0.2 | 0.4 | 0.8 | 2.4 | 8.5 | 43.8 | 155.4 | 324.9 | 534.8 |
| 2000..... | 48.0 | * | * | 0.3 | 0.4 | 0.7 | 2.3 | 8.7 | 43.2 | 154.8 | 326.2 | 521.9 |
| 1999..... | 48.0 | * | * | 0.2 | 0.5 | 0.9 | 2.1 | 8.7 | 45.4 | 155.7 | 329.3 | 507.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 54.9 | * | * | 0.2 | 0.3 | 0.5 | 2.0 | 8.8 | 44.8 | 173.3 | 399.6 | 640.0 |
| 2003..... | 57.0 | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 9.0 | 47.8 | 182.6 | 414.9 | 671.6 |
| 2002..... | 56.4 | 0.9 | 0.2 | 0.2 | 0.4 | 0.6 | 2.0 | 9.0 | 46.7 | 181.6 | 416.3 | 673.1 |
| 2001..... | 55.9 | * | * | 0.2 | 0.3 | 0.5 | 1.9 | 8.7 | 48.6 | 185.9 | 408.6 | 676.9 |
| 2000..... | 55.7 | * | * | 0.2 | 0.4 | 0.6 | 1.8 | 8.6 | 48.2 | 186.5 | 413.2 | 680.8 |
| 1999..... | 56.6 | * | 0.2 | 0.2 | 0.4 | 0.6 | 1.6 | 8.5 | 51.8 | 193.6 | 422.4 | 675.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 53.0 | * | * | 0.2 | 0.3 | 0.5 | 1.8 | 9.2 | 46.3 | 188.5 | 476.6 | 852.5 |
| 2003..... | 54.9 | * | * | 0.2 | 0.4 | 0.5 | 1.8 | 9.5 | 50.0 | 200.2 | 491.0 | 903.6 |
| 2002..... | 55.1 | * | * | 0.2 | 0.4 | 0.6 | 2.0 | 9.3 | 48.3 | 200.4 | 506.7 | 935.4 |
| 2001..... | 54.6 | * | * | 0.2 | 0.3 | 0.5 | 1.9 | 9.1 | 50.5 | 206.1 | 500.9 | 951.5 |
| 2000..... | 55.1 | * | * | 0.2 | 0.4 | 0.5 | 1.7 | 8.9 | 50.8 | 208.8 | 513.6 | 1,008.6 |
| 1999..... | 57.3 | * | * | 0.3 | 0.5 | 0.5 | 1.5 | 8.5 | 55.6 | 224.9 | 534.9 | 1,033.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 56.7 | * | * | * | 0.2 | 0.5 | 2.1 | 8.4 | 43.5 | 160.1 | 347.2 | 544.8 |
| 2003..... | 59.0 | * | * | * | 0.3 | 0.6 | 2.2 | 8.5 | 45.7 | 167.6 | 363.5 | 569.5 |
| 2002..... | 57.7 | * | * | * | 0.4 | 0.6 | 2.1 | 8.6 | 45.1 | 165.5 | 355.7 | 559.8 |
| 2001..... | 57.2 | * | * | * | 0.3 | 0.6 | 2.0 | 8.3 | 46.8 | 168.8 | 347.3 | 562.7 |
| 2000..... | 56.2 | * | * | 0.2 | 0.3 | 0.6 | 1.9 | 8.3 | 45.8 | 167.6 | 347.2 | 548.7 |
| 1999..... | 55.9 | * | * | * | 0.4 | 0.6 | 1.8 | 8.5 | 48.3 | 167.4 | 348.8 | 532.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic lower respiratory diseases (J40-J47) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 19.9 | * | 0.8 | 1.0 | 1.3 | 1.9 | 4.0 | 13.0 | 39.1 | 103.9 | 225.9 | 335.9 |
| 2003..... | 20.9 | * | 1.0 | 1.0 | 1.3 | 2.0 | 4.4 | 12.9 | 41.3 | 115.3 | 238.5 | 354.9 |
| 2002..... | 21.4 | * | 1.2 | 1.1 | 1.3 | 2.0 | 5.2 | 13.1 | 41.6 | 115.0 | 250.5 | 370.8 |
| 2001..... | 21.1 | * | 0.9 | 0.8 | 1.3 | 2.5 | 4.9 | 12.7 | 41.3 | 120.4 | 236.3 | 379.3 |
| 2000..... | 21.5 | * | 0.9 | 1.1 | 1.4 | 2.2 | 4.9 | 14.5 | 44.7 | 123.2 | 236.5 | 376.2 |
| 1999..... | 22.6 | * | * | 1.0 | 1.4 | 2.7 | 4.7 | 14.1 | 46.7 | 129.9 | 265.3 | 377.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 22.8 | * | * | 1.4 | 1.6 | 2.6 | 4.2 | 14.4 | 51.4 | 142.5 | 342.3 | 569.0 |
| 2003..... | 24.2 | * | * | 1.3 | 1.6 | 2.9 | 4.1 | 13.7 | 51.5 | 158.0 | 386.7 | 608.4 |
| 2002..... | 24.9 | * | 1.6 | 1.2 | 2.0 | 2.3 | 5.7 | 14.6 | 53.1 | 160.0 | 395.7 | 653.1 |
| 2001..... | 24.5 | * | * | 1.0 | 1.7 | 2.8 | 4.7 | 13.5 | 50.4 | 170.1 | 385.6 | 682.6 |
| 2000..... | 25.1 | * | * | 1.3 | 1.8 | 2.8 | 4.9 | 15.2 | 55.3 | 178.9 | 372.4 | 696.3 |
| 1999..... | 27.0 | * | * | 1.4 | 1.9 | 2.7 | 4.7 | 15.6 | 60.4 | 186.1 | 438.4 | 707.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 17.2 | * | * | 0.6 | 1.0 | 1.3 | 3.9 | 11.9 | 29.3 | 76.3 | 160.2 | 243.8 |
| 2003..... | 17.9 | * | * | * | 1.1 | 1.2 | 4.7 | 12.2 | 33.2 | 85.0 | 154.8 | 256.7 |
| 2002..... | 18.2 | * | * | 1.1 | 0.7 | 1.7 | 4.8 | 11.8 | 32.3 | 83.0 | 168.5 | 263.0 |
| 2001..... | 18.0 | * | * | 0.7 | 1.0 | 2.2 | 5.0 | 11.9 | 34.0 | 85.3 | 152.5 | 266.9 |
| 2000..... | 18.2 | * | * | 0.8 | 1.0 | 1.6 | 4.9 | 13.8 | 36.1 | 84.1 | 159.9 | 257.3 |
| 1999..... | 18.6 | * | * | 0.6 | 0.9 | 2.7 | 4.7 | 12.8 | 35.8 | 90.3 | 168.4 | 253.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 41.2 | 0.9 | 0.3 | 0.3 | 0.4 | 0.6 | 2.0 | 8.4 | 40.2 | 153.0 | 363.6 | 595.3 |
| 2003..... | 43.1 | 0.8 | 0.3 | 0.3 | 0.5 | 0.7 | 2.1 | 8.6 | 43.1 | 162.4 | 379.9 | 629.1 |
| 2002..... | 42.9 | 1.0 | 0.4 | 0.3 | 0.5 | 0.7 | 2.2 | 8.6 | 42.1 | 162.1 | 383.3 | 631.8 |
| 2001..... | 42.9 | 1.0 | 0.3 | 0.3 | 0.4 | 0.7 | 2.1 | 8.4 | 43.9 | 167.0 | 376.9 | 637.7 |
| 2000..... | 43.0 | 0.9 | 0.3 | 0.3 | 0.5 | 0.7 | 2.0 | 8.6 | 44.0 | 168.5 | 382.8 | 642.4 |
| 1999..... | 44.2 | 0.9 | 0.3 | 0.3 | 0.5 | 0.8 | 2.0 | 8.4 | 47.2 | 176.3 | 394.4 | 639.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 40.4 | 1.2 | 0.4 | 0.4 | 0.5 | 0.7 | 1.9 | 8.8 | 42.9 | 171.6 | 443.6 | 806.6 |
| 2003..... | 42.3 | 1.1 | 0.5 | 0.4 | 0.5 | 0.8 | 1.9 | 9.1 | 46.3 | 183.2 | 462.7 | 861.8 |
| 2002..... | 42.7 | 1.1 | 0.6 | 0.4 | 0.6 | 0.8 | 2.2 | 9.1 | 45.0 | 184.2 | 478.6 | 890.1 |
| 2001..... | 42.5 | 1.1 | 0.4 | 0.3 | 0.5 | 0.7 | 2.0 | 8.8 | 46.6 | 190.6 | 473.2 | 912.1 |
| 2000..... | 43.3 | 1.2 | 0.4 | 0.4 | 0.6 | 0.8 | 1.9 | 8.9 | 47.5 | 194.5 | 486.4 | 963.6 |
| 1999..... | 45.4 | * | 0.4 | 0.4 | 0.6 | 0.7 | 1.8 | 8.6 | 52.0 | 210.1 | 510.1 | 990.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 42.0 | * | * | 0.2 | 0.3 | 0.5 | 2.0 | 8.0 | 37.8 | 137.4 | 309.8 | 500.3 |
| 2003..... | 44.0 | * | * | 0.2 | 0.4 | 0.6 | 2.3 | 8.2 | 40.0 | 145.0 | 324.6 | 526.2 |
| 2002..... | 43.2 | * | 0.3 | 0.3 | 0.4 | 0.7 | 2.2 | 8.2 | 39.4 | 143.7 | 320.0 | 519.7 |
| 2001..... | 43.2 | * | * | 0.2 | 0.4 | 0.7 | 2.2 | 8.1 | 41.3 | 147.3 | 313.6 | 522.9 |
| 2000..... | 42.8 | * | 0.3 | 0.3 | 0.4 | 0.6 | 2.1 | 8.2 | 40.6 | 147.1 | 315.2 | 511.6 |
| 1999..... | 43.0 | * | 0.3 | 0.2 | 0.5 | 0.9 | 2.1 | 8.2 | 42.8 | 148.6 | 319.3 | 498.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.4 | * | * | * | * | 0.4 | 1.0 | 3.0 | 12.0 | 50.9 | 157.8 | 389.1 |
| 2003..... | 7.9 | * | * | * | * | 0.4 | 1.0 | 3.4 | 12.5 | 52.8 | 169.4 | 458.1 |
| 2002..... | 7.8 | * | * | * | * | 0.4 | 1.2 | 3.4 | 13.5 | 53.1 | 174.0 | 454.0 |
| 2001..... | 7.6 | * | * | * | * | 0.3 | 0.9 | 3.5 | 13.9 | 54.8 | 167.4 | 477.7 |
| 2000..... | 7.5 | * | * | * | * | * | 1.0 | 3.3 | 14.0 | 52.5 | 178.6 | 476.4 |
| 1999..... | 8.3 | * | * | * | * | 0.5 | 1.6 | 3.7 | 14.5 | 61.9 | 192.1 | 531.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | * | 0.5 | 1.3 | 2.8 | 13.4 | 67.9 | 206.0 | 515.6 |
| 2003..... | 8.2 | * | * | * | * | * | 1.0 | 3.0 | 16.2 | 67.9 | 229.7 | 646.5 |
| 2002..... | 8.1 | * | * | * | * | * | 1.0 | 3.8 | 17.3 | 68.6 | 241.7 | 599.0 |
| 2001..... | 7.7 | * | * | * | * | * | 0.7 | 3.1 | 16.1 | 74.1 | 223.1 | 672.3 |
| 2000..... | 7.9 | * | * | * | * | * | 0.8 | 3.4 | 17.9 | 72.2 | 247.3 | 669.1 |
| 1999..... | 8.8 | * | * | * | * | * | 1.4 | 3.6 | 17.6 | 81.7 | 269.0 | 836.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.3 | * | * | * | * | * | 0.7 | 3.2 | 10.7 | 37.3 | 124.5 | 324.1 |
| 2003..... | 7.6 | * | * | * | * | * | 1.0 | 3.8 | 9.3 | 40.7 | 128.0 | 362.5 |
| 2002..... | 7.5 | * | * | * | * | * | 1.4 | 3.1 | 10.1 | 40.8 | 127.6 | 382.1 |
| 2001..... | 7.4 | * | * | * | * | * | 1.1 | 4.0 | 12.0 | 39.5 | 129.7 | 382.6 |
| 2000..... | 7.0 | * | * | * | * | * | 1.2 | 3.2 | 10.5 | 37.0 | 132.4 | 381.8 |
| 1999..... | 7.8 | * | * | * | * | * | 1.8 | 3.9 | 11.7 | 46.3 | 140.9 | 384.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 46.7 | 0.8 | 0.3 | 0.3 | 0.5 | 0.7 | 2.1 | 8.9 | 42.5 | 160.1 | 374.5 | 603.8 |
| 2003..... | 48.6 | 0.8 | 0.3 | 0.3 | 0.5 | 0.8 | 2.3 | 9.1 | 45.3 | 169.7 | 390.2 | 635.0 |
| 2002..... | 48.3 | 1.1 | 0.4 | 0.4 | 0.5 | 0.8 | 2.4 | 9.1 | 44.3 | 169.1 | 392.6 | 637.6 |
| 2001..... | 48.0 | 1.0 | 0.3 | 0.3 | 0.5 | 0.8 | 2.3 | 8.8 | 46.0 | 173.9 | 385.7 | 642.2 |
| 2000..... | 48.0 | 1.0 | 0.3 | 0.4 | 0.5 | 0.8 | 2.2 | 9.0 | 46.0 | 175.1 | 390.6 | 647.0 |
| 1999..... | 49.0 | 1.0 | 0.4 | 0.3 | 0.6 | 0.8 | 2.0 | 8.8 | 49.5 | 182.6 | 401.6 | 641.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 46.0 | * | 0.4 | 0.4 | 0.6 | 0.8 | 2.0 | 9.4 | 45.2 | 178.4 | 456.2 | 820.0 |
| 2003..... | 47.9 | * | 0.5 | 0.4 | 0.6 | 0.9 | 2.0 | 9.6 | 48.5 | 190.6 | 473.9 | 870.2 |
| 2002..... | 48.2 | 1.2 | 0.6 | 0.4 | 0.6 | 0.9 | 2.4 | 9.6 | 47.1 | 191.3 | 489.0 | 901.2 |
| 2001..... | 47.8 | * | 0.4 | 0.3 | 0.5 | 0.8 | 2.2 | 9.3 | 48.7 | 197.5 | 483.8 | 919.9 |
| 2000..... | 48.5 | * | 0.4 | 0.4 | 0.6 | 0.9 | 2.1 | 9.3 | 49.5 | 201.1 | 495.4 | 973.6 |
| 1999..... | 50.6 | * | 0.4 | 0.5 | 0.7 | 0.8 | 1.9 | 9.0 | 54.3 | 216.7 | 518.9 | 993.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 47.3 | * | * | 0.2 | 0.4 | 0.6 | 2.2 | 8.5 | 40.0 | 144.6 | 319.6 | 507.1 |
| 2003..... | 49.3 | * | * | 0.2 | 0.4 | 0.7 | 2.5 | 8.6 | 42.4 | 152.2 | 334.3 | 531.7 |
| 2002..... | 48.3 | * | * | 0.3 | 0.4 | 0.7 | 2.4 | 8.6 | 41.7 | 150.7 | 328.7 | 523.9 |
| 2001..... | 48.1 | * | * | 0.2 | 0.4 | 0.8 | 2.4 | 8.4 | 43.5 | 154.2 | 321.2 | 526.7 |
| 2000..... | 47.5 | * | * | 0.3 | 0.4 | 0.7 | 2.2 | 8.6 | 42.8 | 153.7 | 322.3 | 515.0 |
| 1999..... | 47.5 | * | * | 0.2 | 0.5 | 0.9 | 2.1 | 8.6 | 45.0 | 154.6 | 325.6 | 501.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 54.5 | * | * | 0.2 | 0.3 | 0.5 | 1.9 | 8.7 | 44.7 | 172.4 | 396.1 | 633.1 |
| 2003..... | 56.6 | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 9.0 | 47.6 | 181.8 | 411.6 | 665.0 |
| 2002..... | 56.0 | 0.9 | 0.2 | 0.2 | 0.4 | 0.6 | 2.0 | 8.9 | 46.4 | 180.7 | 412.6 | 666.9 |
| 2001..... | 55.5 | * | * | 0.2 | 0.3 | 0.5 | 1.9 | 8.6 | 48.3 | 184.9 | 405.5 | 669.3 |
| 2000..... | 55.3 | * | * | 0.2 | 0.3 | 0.5 | 1.7 | 8.5 | 47.9 | 185.5 | 409.8 | 674.2 |
| 1999..... | 56.2 | * | 0.2 | 0.2 | 0.4 | 0.5 | 1.6 | 8.4 | 51.5 | 192.6 | 418.9 | 668.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 52.7 | * | * | 0.2 | 0.3 | 0.5 | 1.8 | 9.1 | 46.1 | 188.0 | 474.3 | 847.5 |
| 2003..... | 54.7 | * | * | 0.2 | 0.4 | 0.5 | 1.8 | 9.5 | 49.8 | 199.7 | 488.6 | 898.8 |
| 2002..... | 54.9 | * | * | 0.2 | 0.4 | 0.6 | 1.9 | 9.3 | 48.1 | 199.9 | 504.4 | 930.6 |
| 2001..... | 54.4 | * | * | 0.2 | 0.3 | 0.5 | 1.8 | 9.0 | 50.2 | 205.5 | 499.2 | 946.6 |
| 2000..... | 54.9 | * | * | 0.2 | 0.4 | 0.5 | 1.6 | 8.8 | 50.5 | 208.1 | 511.4 | 1,003.9 |
| 1999..... | 57.1 | * | * | 0.3 | 0.5 | 0.5 | 1.5 | 8.4 | 55.4 | 224.2 | 531.8 | 1,026.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 56.1 | * | * | * | 0.2 | 0.5 | 2.1 | 8.3 | 43.3 | 159.0 | 342.9 | 537.1 |
| 2003..... | 58.4 | * | * | * | 0.3 | 0.6 | 2.2 | 8.5 | 45.4 | 166.5 | 359.5 | 562.0 |
| 2002..... | 57.0 | * | * | * | 0.4 | 0.6 | 2.0 | 8.6 | 44.8 | 164.3 | 351.2 | 553.0 |
| 2001..... | 56.5 | * | * | * | 0.3 | 0.6 | 2.0 | 8.2 | 46.5 | 167.5 | 343.3 | 553.9 |
| 2000..... | 55.6 | * | * | 0.2 | 0.3 | 0.6 | 1.9 | 8.3 | 45.5 | 166.5 | 343.0 | 541.3 |
| 1999..... | 55.4 | * | * | * | 0.4 | 0.6 | 1.8 | 8.4 | 47.9 | 166.3 | 345.0 | 526.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 19.8 | * | 0.8 | 1.0 | 1.3 | 1.9 | 4.0 | 13.0 | 38.9 | 103.4 | 225.2 | 334.4 |
| 2003..... | 20.8 | * | 1.0 | 1.0 | 1.3 | 2.0 | 4.4 | 12.8 | 41.1 | 115.1 | 237.8 | 354.6 |
| 2002..... | 21.3 | * | 1.2 | 1.1 | 1.3 | 2.0 | 5.2 | 12.9 | 41.5 | 114.4 | 249.6 | 369.6 |
| 2001..... | 21.0 | * | 0.9 | 0.8 | 1.3 | 2.4 | 4.9 | 12.6 | 41.2 | 120.1 | 235.2 | 378.0 |
| 2000..... | 21.4 | * | 0.9 | 1.1 | 1.4 | 2.2 | 4.9 | 14.4 | 44.3 | 122.4 | 235.9 | 375.5 |
| 1999..... | 22.5 | * | * | 1.0 | 1.4 | 2.6 | 4.7 | 14.0 | 46.5 | 129.4 | 263.8 | 376.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 22.8 | * | * | 1.4 | 1.6 | 2.6 | 4.1 | 14.4 | 51.1 | 142.4 | 342.3 | 567.0 |
| 2003..... | 24.1 | * | * | 1.3 | 1.6 | 2.9 | 4.0 | 13.6 | 51.3 | 157.6 | 385.8 | 608.4 |
| 2002..... | 24.8 | * | 1.6 | 1.2 | 2.0 | 2.3 | 5.7 | 14.5 | 52.9 | 159.1 | 394.5 | 653.1 |
| 2001..... | 24.4 | * | * | 1.0 | 1.7 | 2.8 | 4.7 | 13.5 | 50.2 | 169.5 | 383.5 | 680.3 |
| 2000..... | 25.0 | * | * | 1.3 | 1.8 | 2.8 | 4.9 | 15.2 | 55.1 | 177.6 | 371.8 | 696.3 |
| 1999..... | 26.9 | * | * | 1.4 | 1.9 | 2.7 | 4.6 | 15.5 | 60.0 | 185.8 | 436.8 | 706.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 17.1 | * | * | * | 1.0 | 1.3 | 3.9 | 11.8 | 29.2 | 75.7 | 159.1 | 242.5 |
| 2003..... | 17.8 | * | * | * | 1.1 | 1.1 | 4.7 | 12.1 | 32.9 | 84.9 | 154.2 | 256.3 |
| 2002..... | 18.1 | * | * | 1.1 | 0.7 | 1.7 | 4.8 | 11.6 | 32.2 | 82.7 | 167.6 | 261.3 |
| 2001..... | 17.9 | * | * | 0.7 | 1.0 | 2.1 | 5.0 | 11.8 | 33.9 | 85.2 | 152.0 | 266.1 |
| 2000..... | 18.1 | * | * | 0.8 | 1.0 | 1.6 | 4.8 | 13.7 | 35.7 | 83.6 | 159.4 | 256.4 |
| 1999..... | 18.4 | * | * | 0.6 | 0.9 | 2.6 | 4.7 | 12.7 | 35.6 | 89.6 | 166.9 | 252.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bronchitis, not specified as acute or chronic (J40) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.7 | 3.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 4.6 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 5.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.1 | 5.3 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.5 | 7.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 6.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.6 | 3.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 4.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 5.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 2.0 | 6.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 5.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 3.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 4.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 5.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.1 | 5.3 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 0.3 | 1.2 | 7.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 6.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 3.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 4.6 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 5.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.1 | 5.6 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.6 | 7.5 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 6.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 4.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.0 | 3.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 5.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 5.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 2.1 | 7.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.5 | 6.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 3.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.8 | 5.0 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 5.4 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | 0.3 | 1.2 | 5.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 0.3 | 1.2 | 7.6 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 0.5 | 1.3 | 6.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic bronchitis (J41-J42) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.1 | 2.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.2 | 2.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 4.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 4.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.3 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.3 | 3.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.5 | 4.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.6 | 1.2 | 4.1 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.6 | 6.2 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.7 | 5.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.6 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 2.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 3.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 0.9 | 4.3 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 2.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 3.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 4.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 5.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 4.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.2 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 1.2 | 3.4 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.5 | 4.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.2 | 4.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.7 | 6.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 6.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.7 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 2.6 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 2.6 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.9 | 4.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 0.9 | 4.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 4.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Emphysema (J43) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 0.2 | 1.0 | 5.4 | 20.5 | 41.0 | 49.9 |
| 2003..... | 5.1 | * | * | * | * | * | 0.2 | 1.1 | 6.3 | 22.2 | 46.0 | 54.6 |
| 2002..... | 5.4 | * | * | * | * | * | 0.2 | 1.1 | 6.4 | 24.0 | 48.0 | 59.3 |
| 2001..... | 5.7 | * | * | * | * | * | 0.2 | 1.1 | 7.2 | 26.2 | 50.4 | 62.1 |
| 2000..... | 5.9 | * | * | * | * | * | 0.2 | 1.3 | 7.4 | 27.8 | 52.7 | 63.3 |
| 1999..... | 6.4 | * | * | * | * | * | 0.2 | 1.3 | 8.5 | 30.7 | 56.9 | 62.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | * | 0.2 | 1.2 | 6.2 | 24.3 | 52.2 | 74.4 |
| 2003..... | 5.3 | * | * | * | * | * | 0.2 | 1.4 | 6.9 | 25.6 | 58.1 | 79.2 |
| 2002..... | 5.7 | * | * | * | * | * | 0.3 | 1.4 | 7.2 | 28.1 | 63.5 | 90.0 |
| 2001..... | 6.0 | * | * | * | * | * | 0.2 | 1.4 | 7.9 | 30.8 | 65.3 | 101.2 |
| 2000..... | 6.3 | * | * | * | * | * | 0.2 | 1.5 | 8.1 | 33.0 | 69.1 | 104.2 |
| 1999..... | 6.9 | * | * | * | * | * | 0.2 | 1.5 | 9.5 | 37.8 | 76.4 | 105.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | * | * | 0.1 | 0.8 | 4.7 | 17.4 | 33.5 | 38.9 |
| 2003..... | 4.9 | * | * | * | * | * | 0.2 | 0.8 | 5.7 | 19.2 | 37.9 | 43.7 |
| 2002..... | 5.0 | * | * | * | * | * | 0.1 | 0.9 | 5.6 | 20.6 | 37.6 | 46.0 |
| 2001..... | 5.4 | * | * | * | * | * | 0.1 | 0.9 | 6.6 | 22.5 | 40.6 | 45.8 |
| 2000..... | 5.6 | * | * | * | * | * | 0.2 | 1.0 | 6.8 | 23.6 | 41.9 | 46.7 |
| 1999..... | 5.8 | * | * | * | * | * | 0.2 | 1.1 | 7.5 | 24.9 | 44.2 | 44.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | 0.9 | 5.0 | 16.1 | 30.0 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | 1.4 | 6.0 | 20.0 | 34.7 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | 1.5 | 6.7 | 18.8 | 36.6 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 1.4 | 6.6 | 20.4 | 39.4 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | 1.5 | 6.0 | 22.3 | 31.2 |
| 1999..... | 0.9 | * | * | * | * | * | * | * | 1.3 | 7.1 | 23.5 | 44.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 6.5 | 20.4 | 47.6 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 2.1 | 7.6 | 29.1 | 53.9 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | 9.2 | 26.7 | 53.5 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 9.2 | 34.0 | 60.9 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | * | 9.1 | 32.5 | 58.4 |
| 1999..... | 1.0 | * | * | * | * | * | * | * | * | 11.2 | 36.0 | 77.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 3.7 | 13.1 | 20.9 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | 4.7 | 13.7 | 25.0 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 4.7 | 13.5 | 28.2 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 4.6 | 11.2 | 28.9 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 3.6 | 15.5 | * |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 4.0 | 15.3 | 28.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Emphysema (J43) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | * | * | 0.2 | 1.1 | 5.8 | 21.7 | 42.4 | 50.7 |
| 2003..... | 5.8 | * | * | * | * | * | 0.2 | 1.2 | 6.6 | 23.3 | 47.3 | 55.4 |
| 2002..... | 6.1 | * | * | * | * | * | 0.2 | 1.2 | 6.8 | 25.1 | 49.3 | 60.1 |
| 2001..... | 6.4 | * | * | * | * | * | 0.2 | 1.2 | 7.6 | 27.5 | 51.7 | 62.8 |
| 2000..... | 6.7 | * | * | * | * | * | 0.2 | 1.3 | 7.9 | 29.1 | 53.8 | 64.3 |
| 1999..... | 7.1 | * | * | * | * | * | 0.2 | 1.3 | 9.0 | 32.0 | 58.1 | 62.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | * | * | 0.2 | 1.3 | 6.6 | 25.6 | 54.0 | 75.5 |
| 2003..... | 6.0 | * | * | * | * | * | 0.2 | 1.5 | 7.3 | 26.8 | 59.5 | 80.4 |
| 2002..... | 6.5 | * | * | * | * | * | 0.3 | 1.5 | 7.7 | 29.3 | 65.2 | 91.3 |
| 2001..... | 6.8 | * | * | * | * | * | 0.3 | 1.5 | 8.4 | 32.1 | 66.6 | 102.7 |
| 2000..... | 7.1 | * | * | * | * | * | 0.2 | 1.6 | 8.5 | 34.3 | 70.6 | 105.9 |
| 1999..... | 7.8 | * | * | * | * | * | 0.2 | 1.5 | 10.0 | 39.2 | 77.8 | 106.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | * | * | 0.2 | 0.9 | 5.1 | 18.4 | 34.6 | 39.6 |
| 2003..... | 5.6 | * | * | * | * | * | 0.2 | 0.9 | 6.0 | 20.3 | 39.1 | 44.5 |
| 2002..... | 5.7 | * | * | * | * | * | 0.1 | 1.0 | 6.0 | 21.7 | 38.8 | 46.6 |
| 2001..... | 6.1 | * | * | * | * | * | 0.1 | 1.0 | 7.0 | 23.6 | 41.9 | 46.3 |
| 2000..... | 6.3 | * | * | * | * | * | 0.2 | 1.1 | 7.3 | 24.8 | 42.9 | 47.5 |
| 1999..... | 6.5 | * | * | * | * | * | 0.2 | 1.1 | 8.1 | 26.1 | 45.3 | 45.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | * | * | 0.2 | 1.1 | 6.2 | 23.7 | 45.2 | 53.4 |
| 2003..... | 6.8 | * | * | * | * | * | 0.2 | 1.2 | 7.1 | 25.2 | 49.9 | 57.9 |
| 2002..... | 7.1 | * | * | * | * | * | 0.2 | 1.3 | 7.3 | 27.0 | 52.3 | 63.1 |
| 2001..... | 7.5 | * | * | * | * | * | 0.2 | 1.3 | 8.2 | 29.6 | 54.9 | 65.9 |
| 2000..... | 7.8 | * | * | * | * | * | 0.2 | 1.3 | 8.4 | 31.2 | 57.0 | 66.9 |
| 1999..... | 8.3 | * | * | * | * | * | 0.2 | 1.3 | 9.7 | 34.5 | 61.2 | 65.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | * | 0.2 | 1.2 | 6.8 | 27.2 | 56.5 | 78.2 |
| 2003..... | 6.9 | * | * | * | * | * | 0.2 | 1.5 | 7.6 | 28.5 | 61.3 | 81.6 |
| 2002..... | 7.4 | * | * | * | * | * | 0.3 | 1.5 | 8.0 | 30.6 | 67.7 | 94.3 |
| 2001..... | 7.8 | * | * | * | * | * | 0.3 | 1.5 | 8.7 | 33.8 | 69.6 | 106.6 |
| 2000..... | 8.1 | * | * | * | * | * | 0.2 | 1.4 | 8.9 | 35.8 | 73.9 | 107.3 |
| 1999..... | 8.9 | * | * | * | * | * | 0.2 | 1.4 | 10.4 | 41.1 | 80.6 | 110.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.0 | * | * | * | * | * | 0.2 | 0.9 | 5.6 | 20.6 | 37.6 | 42.3 |
| 2003..... | 6.7 | * | * | * | * | * | 0.2 | 1.0 | 6.7 | 22.5 | 42.3 | 47.4 |
| 2002..... | 6.8 | * | * | * | * | * | * | 1.1 | 6.7 | 24.0 | 41.9 | 49.7 |
| 2001..... | 7.2 | * | * | * | * | * | 0.1 | 1.0 | 7.7 | 26.1 | 45.2 | 48.9 |
| 2000..... | 7.4 | * | * | * | * | * | 0.2 | 1.1 | 8.0 | 27.3 | 45.8 | 50.6 |
| 1999..... | 7.7 | * | * | * | * | * | 0.2 | 1.2 | 9.0 | 29.0 | 48.5 | 47.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Emphysema (J43) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | 1.3 | 4.7 | 11.0 | 22.5 | 27.9 |
| 2003..... | 2.2 | * | * | * | * | * | * | 1.2 | 5.3 | 13.2 | 27.7 | 31.2 |
| 2002..... | 2.3 | * | * | * | * | * | 0.5 | 1.2 | 4.6 | 16.2 | 28.2 | 32.1 |
| 2001..... | 2.2 | * | * | * | * | * | * | 1.4 | 5.6 | 15.6 | 23.6 | 35.9 |
| 2000..... | 2.5 | * | * | * | * | * | * | 2.0 | 6.1 | 17.1 | 27.0 | 37.5 |
| 1999..... | 2.7 | * | * | * | * | * | 0.4 | 1.9 | 6.3 | 17.3 | 31.8 | 37.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | * | 2.0 | 7.1 | 17.4 | 37.1 | 57.9 |
| 2003..... | 2.8 | * | * | * | * | * | * | 1.7 | 7.5 | 18.6 | 46.7 | 71.4 |
| 2002..... | 3.1 | * | * | * | * | * | * | 1.8 | 6.7 | 25.3 | 49.5 | 65.6 |
| 2001..... | 2.8 | * | * | * | * | * | * | 1.6 | 7.9 | 23.0 | 40.2 | 69.3 |
| 2000..... | 3.3 | * | * | * | * | * | * | 3.0 | 8.2 | 27.4 | 41.3 | 89.2 |
| 1999..... | 3.6 | * | * | * | * | * | * | 3.0 | 9.1 | 29.3 | 55.7 | 73.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.7 | 6.4 | 14.3 | 16.1 |
| 2003..... | 1.6 | * | * | * | * | * | * | * | 3.5 | 9.4 | 16.9 | 15.7 |
| 2002..... | 1.6 | * | * | * | * | * | * | * | 2.8 | 9.8 | 16.1 | 19.2 |
| 2001..... | 1.7 | * | * | * | * | * | * | 1.1 | 3.7 | 10.4 | 14.3 | 23.5 |
| 2000..... | 1.8 | * | * | * | * | * | * | 1.3 | 4.5 | 9.8 | 18.9 | 18.3 |
| 1999..... | 1.8 | * | * | * | * | * | * | 1.0 | 4.0 | 8.9 | 18.4 | 24.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other chronic obstructive pulmonary disease (J44) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 35.0 | * | * | * | * | 0.1 | 0.9 | 5.9 | 32.8 | 129.2 | 315.8 | 524.8 |
| 2003..... | 36.3 | * | * | * | * | 0.1 | 0.9 | 5.9 | 34.6 | 136.7 | 326.2 | 552.9 |
| 2002..... | 35.8 | * | * | * | * | 0.1 | 0.9 | 5.9 | 33.5 | 134.2 | 326.7 | 549.7 |
| 2001..... | 35.3 | * | * | * | 0.1 | 0.1 | 0.9 | 5.6 | 34.2 | 136.5 | 318.0 | 550.3 |
| 2000..... | 35.1 | * | * | * | * | 0.1 | 0.8 | 5.6 | 34.0 | 135.8 | 320.6 | 550.6 |
| 1999..... | 35.7 | * | * | * | * | 0.1 | 0.7 | 5.4 | 35.9 | 140.5 | 327.2 | 549.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 34.3 | * | * | * | * | 0.1 | 1.0 | 6.4 | 35.1 | 144.5 | 386.3 | 714.1 |
| 2003..... | 35.7 | * | * | * | * | 0.1 | 0.9 | 6.4 | 37.6 | 154.9 | 397.8 | 764.9 |
| 2002..... | 35.5 | * | * | * | * | 0.1 | 1.0 | 6.4 | 36.0 | 152.8 | 407.6 | 780.4 |
| 2001..... | 35.1 | * | * | * | * | 0.1 | 0.9 | 6.2 | 36.9 | 156.3 | 401.1 | 789.6 |
| 2000..... | 35.4 | * | * | * | * | * | 0.8 | 6.0 | 37.5 | 157.5 | 408.7 | 833.7 |
| 1999..... | 36.9 | * | * | * | * | * | 0.7 | 5.9 | 40.3 | 167.9 | 424.8 | 860.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 35.7 | * | * | * | * | * | 0.9 | 5.5 | 30.7 | 116.3 | 268.4 | 439.6 |
| 2003..... | 36.9 | * | * | * | * | * | 1.0 | 5.4 | 31.8 | 121.6 | 278.3 | 459.2 |
| 2002..... | 36.0 | * | * | * | * | 0.1 | 0.9 | 5.4 | 31.2 | 118.7 | 273.0 | 449.5 |
| 2001..... | 35.5 | * | * | * | * | * | 0.9 | 5.1 | 31.7 | 120.2 | 263.4 | 450.2 |
| 2000..... | 34.7 | * | * | * | * | * | 0.8 | 5.2 | 30.8 | 117.9 | 263.1 | 435.3 |
| 1999..... | 34.5 | * | * | * | * | 0.1 | 0.7 | 4.9 | 31.8 | 118.0 | 263.8 | 424.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 6.0 | * | * | * | * | * | 0.4 | 1.6 | 9.1 | 42.4 | 134.5 | 343.0 |
| 2003..... | 6.2 | * | * | * | * | * | 0.4 | 1.6 | 8.9 | 43.2 | 141.9 | 401.6 |
| 2002..... | 6.2 | * | * | * | * | * | 0.4 | 2.0 | 10.0 | 42.4 | 145.8 | 394.1 |
| 2001..... | 6.0 | * | * | * | * | * | * | 1.8 | 10.0 | 44.1 | 137.4 | 416.4 |
| 2000..... | 5.8 | * | * | * | * | * | * | 1.6 | 9.6 | 40.0 | 145.8 | 418.0 |
| 1999..... | 6.4 | * | * | * | * | * | * | 1.7 | 10.7 | 48.5 | 155.4 | 458.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.3 | * | * | * | * | * | * | 1.7 | 11.0 | 58.7 | 180.6 | 452.5 |
| 2003..... | 6.5 | * | * | * | * | * | * | 1.3 | 11.8 | 57.6 | 192.9 | 574.2 |
| 2002..... | 6.6 | * | * | * | * | * | * | 2.2 | 13.8 | 57.3 | 207.2 | 532.1 |
| 2001..... | 6.2 | * | * | * | * | * | * | 2.0 | 12.2 | 61.1 | 180.6 | 601.8 |
| 2000..... | 6.1 | * | * | * | * | * | * | 1.6 | 13.3 | 56.3 | 206.0 | 582.5 |
| 1999..... | 7.0 | * | * | * | * | * | * | 1.6 | 13.9 | 65.7 | 225.2 | 735.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | * | * | 1.5 | 7.4 | 29.4 | 102.7 | 286.7 |
| 2003..... | 5.9 | * | * | * | * | * | * | 1.8 | 6.3 | 31.8 | 106.9 | 314.1 |
| 2002..... | 5.7 | * | * | * | * | * | * | 1.8 | 6.6 | 30.6 | 103.7 | 325.8 |
| 2001..... | 5.7 | * | * | * | * | * | * | 1.7 | 8.0 | 30.7 | 108.2 | 325.8 |
| 2000..... | 5.4 | * | * | * | * | * | * | 1.6 | 6.3 | 27.2 | 105.3 | 337.3 |
| 1999..... | 5.7 | * | * | * | * | * | * | 1.8 | 7.8 | 35.1 | 108.9 | 324.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other chronic obstructive pulmonary disease (J44) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 39.7 | * | * | * | * | 0.1 | 1.0 | 6.4 | 34.7 | 135.2 | 325.4 | 532.3 |
| 2003..... | 41.0 | * | * | * | * | 0.1 | 1.0 | 6.3 | 36.5 | 143.0 | 335.1 | 558.2 |
| 2002..... | 40.3 | * | * | * | * | 0.1 | 1.0 | 6.2 | 35.3 | 140.1 | 334.7 | 554.9 |
| 2001..... | 39.6 | * | * | * | 0.1 | 0.1 | 1.0 | 5.9 | 35.9 | 142.2 | 325.6 | 554.1 |
| 2000..... | 39.2 | * | * | * | * | 0.1 | 0.8 | 5.9 | 35.7 | 141.3 | 327.2 | 554.3 |
| 1999..... | 39.6 | * | * | * | * | 0.1 | 0.8 | 5.7 | 37.6 | 145.5 | 333.3 | 551.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 39.1 | * | * | * | * | 0.1 | 1.0 | 6.9 | 36.9 | 150.1 | 397.2 | 726.4 |
| 2003..... | 40.5 | * | * | * | * | * | 1.0 | 6.9 | 39.5 | 161.2 | 407.8 | 772.3 |
| 2002..... | 40.2 | * | * | * | * | 0.1 | 1.0 | 6.8 | 37.6 | 158.6 | 416.3 | 790.1 |
| 2001..... | 39.6 | * | * | * | * | 0.1 | 1.0 | 6.5 | 38.6 | 161.9 | 410.5 | 795.7 |
| 2000..... | 39.7 | * | * | * | * | * | 0.9 | 6.4 | 39.0 | 163.0 | 416.3 | 842.2 |
| 1999..... | 41.1 | * | * | * | * | * | 0.8 | 6.2 | 42.0 | 173.2 | 432.1 | 862.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 40.3 | * | * | * | * | * | 1.0 | 5.9 | 32.6 | 122.6 | 277.2 | 445.6 |
| 2003..... | 41.5 | * | * | * | * | * | 1.1 | 5.7 | 33.8 | 127.8 | 286.6 | 464.1 |
| 2002..... | 40.3 | * | * | * | * | * | 0.9 | 5.7 | 33.1 | 124.6 | 280.6 | 453.4 |
| 2001..... | 39.6 | * | * | * | * | * | 1.0 | 5.4 | 33.5 | 125.9 | 269.9 | 453.6 |
| 2000..... | 38.6 | * | * | * | * | * | 0.8 | 5.4 | 32.6 | 123.4 | 269.2 | 437.9 |
| 1999..... | 38.2 | * | * | * | * | 0.1 | 0.7 | 5.1 | 33.5 | 122.7 | 269.2 | 426.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 46.7 | * | * | * | * | 0.1 | 1.1 | 6.7 | 36.9 | 145.8 | 344.7 | 559.2 |
| 2003..... | 48.1 | * | * | * | * | 0.1 | 1.0 | 6.6 | 38.8 | 153.6 | 354.3 | 585.5 |
| 2002..... | 47.1 | * | * | * | * | 0.1 | 1.0 | 6.5 | 37.4 | 150.2 | 352.0 | 580.7 |
| 2001..... | 46.2 | * | * | * | * | 0.1 | 1.0 | 6.1 | 38.3 | 151.7 | 342.5 | 578.0 |
| 2000..... | 45.6 | * | * | * | * | * | 0.8 | 6.0 | 37.7 | 150.3 | 343.6 | 578.9 |
| 1999..... | 45.9 | * | * | * | * | 0.1 | 0.7 | 5.8 | 39.5 | 153.8 | 347.9 | 576.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 45.2 | * | * | * | * | * | 1.1 | 7.1 | 38.2 | 158.4 | 413.3 | 751.7 |
| 2003..... | 46.6 | * | * | * | * | * | 1.0 | 7.1 | 41.0 | 169.0 | 421.3 | 799.5 |
| 2002..... | 46.2 | * | * | * | * | * | 1.1 | 6.9 | 38.8 | 166.4 | 429.5 | 816.2 |
| 2001..... | 45.3 | * | * | * | * | * | 1.0 | 6.6 | 40.2 | 168.7 | 423.4 | 819.0 |
| 2000..... | 45.4 | * | * | * | * | * | 0.9 | 6.3 | 40.3 | 169.1 | 429.2 | 871.8 |
| 1999..... | 46.8 | * | * | * | * | * | 0.7 | 6.2 | 43.1 | 179.5 | 442.8 | 893.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 48.1 | * | * | * | * | * | 1.0 | 6.2 | 35.8 | 135.1 | 297.9 | 472.9 |
| 2003..... | 49.6 | * | * | * | * | * | 1.1 | 6.1 | 36.7 | 140.4 | 309.1 | 491.3 |
| 2002..... | 48.0 | * | * | * | * | * | 1.0 | 6.1 | 36.0 | 136.4 | 300.0 | 478.9 |
| 2001..... | 46.9 | * | * | * | * | * | 1.0 | 5.7 | 36.5 | 137.2 | 288.9 | 477.8 |
| 2000..... | 45.7 | * | * | * | * | * | 0.8 | 5.8 | 35.2 | 134.4 | 287.4 | 460.8 |
| 1999..... | 45.0 | * | * | * | * | * | 0.7 | 5.4 | 36.2 | 132.3 | 285.7 | 449.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other chronic obstructive pulmonary disease (J44) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 15.0 | * | * | * | * | * | 1.1 | 6.8 | 28.2 | 86.4 | 192.4 | 286.7 |
| 2003..... | 15.6 | * | * | * | * | * | 1.2 | 6.4 | 30.3 | 94.3 | 199.5 | 305.8 |
| 2002..... | 15.8 | * | * | * | * | * | 1.2 | 6.6 | 30.1 | 89.8 | 212.1 | 321.5 |
| 2001..... | 15.5 | * | * | * | * | * | 1.3 | 6.5 | 28.5 | 95.8 | 199.7 | 322.1 |
| 2000..... | 15.3 | * | * | * | * | * | 1.2 | 6.7 | 31.1 | 95.0 | 196.2 | 312.0 |
| 1999..... | 16.3 | * | * | * | * | * | 1.2 | 6.4 | 32.9 | 101.3 | 218.5 | 313.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 17.4 | * | * | * | * | * | 1.2 | 7.9 | 38.5 | 118.9 | 295.8 | 493.8 |
| 2003..... | 18.6 | * | * | * | * | * | 1.3 | 7.6 | 39.2 | 133.1 | 328.1 | 526.3 |
| 2002..... | 18.6 | * | * | * | * | * | 1.3 | 8.3 | 40.2 | 125.8 | 336.7 | 572.1 |
| 2001..... | 18.7 | * | * | * | * | * | 1.4 | 7.5 | 37.4 | 138.5 | 332.8 | 590.2 |
| 2000..... | 18.7 | * | * | * | * | * | 1.3 | 8.4 | 40.5 | 141.3 | 321.8 | 578.9 |
| 1999..... | 20.2 | * | * | * | * | * | 1.4 | 8.2 | 45.0 | 146.2 | 367.7 | 609.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 12.7 | * | * | * | * | * | 1.0 | 5.9 | 19.9 | 63.2 | 134.1 | 204.8 |
| 2003..... | 13.0 | * | * | * | * | * | 1.2 | 5.4 | 23.1 | 66.7 | 126.9 | 220.4 |
| 2002..... | 13.2 | * | * | * | * | * | 1.1 | 5.1 | 22.1 | 64.3 | 141.6 | 225.7 |
| 2001..... | 12.6 | * | * | * | * | * | 1.2 | 5.6 | 21.4 | 65.7 | 125.0 | 222.9 |
| 2000..... | 12.3 | * | * | * | * | * | 1.2 | 5.3 | 23.5 | 62.4 | 125.6 | 212.8 |
| 1999..... | 12.8 | * | * | * | * | * | 1.0 | 4.9 | 23.2 | 69.7 | 135.0 | 202.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Asthma (J45-J46) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.9 | 1.4 | 1.8 | 2.7 | 5.2 | 14.1 |
| 2003..... | 1.4 | * | 0.2 | 0.3 | 0.4 | 0.6 | 0.9 | 1.5 | 2.0 | 2.9 | 5.8 | 14.2 |
| 2002..... | 1.5 | * | 0.3 | 0.3 | 0.4 | 0.6 | 1.1 | 1.5 | 2.0 | 3.2 | 6.4 | 14.7 |
| 2001..... | 1.5 | * | 0.2 | 0.2 | 0.4 | 0.6 | 1.0 | 1.5 | 2.2 | 3.5 | 6.4 | 16.1 |
| 2000..... | 1.6 | * | 0.2 | 0.3 | 0.4 | 0.6 | 1.0 | 1.6 | 2.2 | 4.0 | 6.9 | 16.7 |
| 1999..... | 1.7 | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.0 | 1.7 | 2.5 | 4.2 | 7.6 | 17.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | 0.3 | 0.5 | 0.6 | 0.7 | 1.0 | 1.3 | 2.1 | 3.5 | 11.1 |
| 2003..... | 1.0 | * | 0.4 | 0.3 | 0.4 | 0.6 | 0.7 | 1.2 | 1.6 | 2.1 | 4.5 | 10.7 |
| 2002..... | 1.1 | * | 0.4 | 0.3 | 0.5 | 0.6 | 0.9 | 1.2 | 1.6 | 2.4 | 5.0 | 10.9 |
| 2001..... | 1.1 | * | 0.3 | 0.3 | 0.4 | 0.6 | 0.9 | 1.1 | 1.5 | 2.7 | 4.5 | 12.1 |
| 2000..... | 1.2 | * | 0.3 | 0.4 | 0.5 | 0.7 | 0.9 | 1.3 | 1.7 | 3.0 | 5.0 | 12.9 |
| 1999..... | 1.2 | * | * | 0.4 | 0.5 | 0.6 | 0.8 | 1.1 | 1.8 | 3.2 | 5.7 | 12.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.2 | 0.3 | 0.4 | 1.0 | 1.6 | 2.2 | 3.2 | 6.3 | 15.4 |
| 2003..... | 1.8 | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 1.9 | 2.4 | 3.6 | 6.7 | 15.8 |
| 2002..... | 1.8 | * | * | 0.3 | 0.3 | 0.5 | 1.2 | 1.8 | 2.4 | 3.9 | 7.3 | 16.3 |
| 2001..... | 1.9 | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 1.9 | 2.8 | 4.1 | 7.6 | 17.8 |
| 2000..... | 2.0 | * | * | 0.3 | 0.3 | 0.5 | 1.2 | 2.0 | 2.7 | 4.8 | 8.1 | 18.2 |
| 1999..... | 2.1 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.2 | 3.0 | 5.0 | 8.8 | 19.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.3 | 0.5 | 1.1 | 1.9 | 3.3 | 5.3 | 11.4 |
| 2003..... | 0.7 | * | * | * | * | 0.3 | 0.5 | 1.6 | 2.2 | 3.4 | 6.1 | 14.5 |
| 2002..... | 0.7 | * | * | * | * | 0.3 | 0.7 | 1.2 | 1.9 | 3.8 | 7.0 | 12.2 |
| 2001..... | 0.7 | * | * | * | * | * | 0.6 | 1.4 | 2.4 | 3.8 | 7.9 | 13.8 |
| 2000..... | 0.8 | * | * | * | * | * | 0.6 | 1.3 | 2.8 | 5.6 | 7.9 | 16.6 |
| 1999..... | 0.9 | * | * | * | * | 0.3 | 0.9 | 1.6 | 2.4 | 5.8 | 10.4 | 17.5 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.6 | * | * | * | * | * | * | 1.4 | 2.2 | * | * | * |
| 2002..... | 0.6 | * | * | * | * | * | * | 1.2 | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.7 | * | * | * | * | * | * | 1.3 | * | 5.3 | * | * |
| 1999..... | 0.7 | * | * | * | * | * | 0.9 | 1.4 | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 1.5 | 2.8 | 3.8 | 6.9 | * |
| 2003..... | 0.9 | * | * | * | * | * | * | 1.8 | 2.1 | 4.1 | 6.3 | 17.2 |
| 2002..... | 0.9 | * | * | * | * | * | 1.0 | 2.2 | 2.2 | 5.3 | 7.6 | * |
| 2001..... | 1.0 | * | * | * | * | * | * | 2.0 | 2.9 | 3.9 | 9.4 | 18.6 |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.3 | 3.2 | 5.8 | 9.2 | * |
| 1999..... | 1.2 | * | * | * | * | * | 1.0 | 1.9 | 3.1 | 7.1 | 13.2 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Asthma (J45-J46) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.9 | 1.4 | 1.8 | 2.7 | 5.2 | 14.2 |
| 2003..... | 1.5 | * | 0.2 | 0.3 | 0.4 | 0.6 | 1.0 | 1.5 | 2.0 | 2.9 | 5.8 | 14.1 |
| 2002..... | 1.6 | * | 0.3 | 0.3 | 0.5 | 0.6 | 1.1 | 1.5 | 2.0 | 3.1 | 6.3 | 14.7 |
| 2001..... | 1.6 | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.1 | 1.5 | 2.2 | 3.4 | 6.3 | 16.1 |
| 2000..... | 1.7 | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.1 | 1.6 | 2.2 | 3.9 | 6.8 | 16.6 |
| 1999..... | 1.8 | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.0 | 1.7 | 2.5 | 4.1 | 7.4 | 17.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | 0.3 | 0.5 | 0.6 | 0.7 | 1.1 | 1.4 | 2.1 | 3.5 | 11.1 |
| 2003..... | 1.1 | * | 0.4 | 0.4 | 0.5 | 0.7 | 0.8 | 1.1 | 1.5 | 2.0 | 4.5 | 10.7 |
| 2002..... | 1.2 | * | 0.4 | 0.3 | 0.6 | 0.7 | 0.9 | 1.2 | 1.6 | 2.4 | 4.9 | 11.1 |
| 2001..... | 1.1 | * | * | 0.3 | 0.4 | 0.6 | 0.9 | 1.2 | 1.5 | 2.6 | 4.4 | 12.2 |
| 2000..... | 1.2 | * | * | 0.4 | 0.6 | 0.8 | 0.9 | 1.3 | 1.6 | 2.9 | 4.9 | 12.7 |
| 1999..... | 1.3 | * | * | 0.4 | 0.6 | 0.7 | 0.8 | 1.1 | 1.8 | 3.1 | 5.6 | 12.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 1.7 | 2.2 | 3.2 | 6.3 | 15.5 |
| 2003..... | 1.9 | * | * | 0.2 | 0.4 | 0.5 | 1.2 | 1.9 | 2.4 | 3.6 | 6.7 | 15.6 |
| 2002..... | 2.0 | * | * | 0.3 | 0.3 | 0.6 | 1.2 | 1.9 | 2.4 | 3.8 | 7.2 | 16.3 |
| 2001..... | 2.1 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 1.9 | 2.8 | 4.1 | 7.5 | 17.7 |
| 2000..... | 2.1 | * | * | 0.3 | 0.4 | 0.6 | 1.2 | 2.0 | 2.6 | 4.7 | 8.0 | 18.2 |
| 1999..... | 2.2 | * | * | 0.2 | 0.4 | 0.7 | 1.2 | 2.2 | 3.0 | 4.9 | 8.6 | 19.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 0.9 | 1.3 | 2.3 | 4.6 | 13.8 |
| 2003..... | 1.3 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.0 | 1.5 | 2.4 | 5.3 | 14.1 |
| 2002..... | 1.4 | * | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.1 | 1.5 | 2.6 | 6.1 | 14.7 |
| 2001..... | 1.4 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.1 | 1.5 | 2.9 | 5.8 | 15.7 |
| 2000..... | 1.4 | * | * | 0.2 | 0.3 | 0.4 | 0.7 | 1.1 | 1.5 | 3.2 | 6.4 | 15.8 |
| 1999..... | 1.5 | * | * | 0.2 | 0.4 | 0.4 | 0.7 | 1.2 | 1.9 | 3.4 | 7.1 | 16.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.3 | 0.3 | 0.5 | 0.7 | 0.9 | 1.7 | 2.8 | 10.4 |
| 2003..... | 0.9 | * | * | * | 0.3 | 0.4 | 0.5 | 0.8 | 1.1 | 1.7 | 3.8 | 10.6 |
| 2002..... | 0.9 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.1 | 1.8 | 4.5 | 10.8 | |
| 2001..... | 0.9 | * | * | 0.2 | 0.2 | 0.3 | 0.6 | 0.8 | 1.0 | 2.1 | 3.9 | 11.4 |
| 2000..... | 1.0 | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.9 | 1.0 | 2.3 | 4.6 | 11.1 |
| 1999..... | 1.0 | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.7 | 1.4 | 2.4 | 5.2 | 10.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.2 | 0.4 | 0.8 | 1.1 | 1.7 | 2.9 | 5.9 | 15.3 |
| 2003..... | 1.7 | * | * | * | 0.2 | 0.5 | 0.9 | 1.3 | 2.0 | 2.9 | 6.3 | 15.7 |
| 2002..... | 1.8 | * | * | * | 0.3 | 0.5 | 0.9 | 1.3 | 1.8 | 3.2 | 7.1 | 16.4 |
| 2001..... | 1.9 | * | * | * | 0.2 | 0.4 | 0.8 | 1.4 | 2.1 | 3.6 | 7.1 | 17.5 |
| 2000..... | 1.9 | * | * | 0.2 | 0.3 | 0.4 | 0.9 | 1.3 | 2.0 | 4.0 | 7.6 | 17.7 |
| 1999..... | 2.1 | * | * | * | 0.3 | 0.5 | 0.9 | 1.6 | 2.3 | 4.2 | 8.3 | 18.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Asthma (J45-J46) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | 0.9 | 1.2 | 1.7 | 2.5 | 4.6 | 5.8 | 5.8 | 9.4 | 15.3 |
| 2003..... | 2.8 | * | 0.8 | 0.9 | 1.2 | 1.7 | 2.8 | 4.9 | 5.2 | 7.2 | 9.3 | 11.9 |
| 2002..... | 3.0 | * | 0.9 | 1.1 | 1.3 | 1.6 | 3.5 | 4.9 | 6.6 | 8.0 | 7.9 | 11.8 |
| 2001..... | 3.1 | * | * | 0.8 | 1.2 | 2.2 | 3.3 | 4.7 | 6.9 | 7.7 | 10.7 | 16.5 |
| 2000..... | 3.3 | * | * | 1.1 | 1.3 | 2.0 | 3.2 | 5.5 | 6.9 | 9.4 | 10.4 | 22.3 |
| 1999..... | 3.3 | * | * | 1.0 | 1.3 | 2.4 | 3.0 | 5.4 | 7.0 | 9.8 | 10.6 | 20.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | 1.2 | 1.5 | 2.4 | 2.3 | 4.0 | 5.3 | 5.7 | 8.8 | * |
| 2003..... | 2.5 | * | * | 1.3 | 1.4 | 2.5 | 2.4 | 4.0 | 4.4 | 5.5 | 8.9 | * |
| 2002..... | 2.8 | * | * | 1.1 | 1.9 | 2.0 | 3.7 | 4.0 | 5.9 | 7.5 | 6.7 | * |
| 2001..... | 2.7 | * | * | 0.9 | 1.4 | 2.5 | 3.0 | 4.1 | 4.7 | 6.8 | 8.7 | * |
| 2000..... | 2.8 | * | * | 1.3 | 1.6 | 2.5 | 3.0 | 3.7 | 6.1 | 7.9 | 6.5 | 24.7 |
| 1999..... | 2.9 | * | * | 1.4 | 1.7 | 2.7 | 2.8 | 4.0 | 5.5 | 9.2 | 8.4 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | 1.0 | 1.1 | 2.7 | 5.1 | 6.2 | 5.9 | 9.7 | 16.9 |
| 2003..... | 3.0 | * | * | * | 1.1 | 1.0 | 3.2 | 5.7 | 5.9 | 8.4 | 9.5 | 13.6 |
| 2002..... | 3.1 | * | * | 1.0 | * | 1.3 | 3.4 | 5.7 | 7.2 | 8.4 | 8.6 | 12.5 |
| 2001..... | 3.4 | * | * | 0.7 | 0.9 | 2.0 | 3.6 | 5.1 | 8.6 | 8.4 | 11.8 | 17.1 |
| 2000..... | 3.7 | * | * | 0.8 | 1.0 | 1.5 | 3.4 | 7.0 | 7.6 | 10.4 | 12.7 | 21.4 |
| 1999..... | 3.6 | * | * | 0.6 | 0.9 | 2.2 | 3.2 | 6.5 | 8.1 | 10.2 | 11.7 | 21.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Bronchiectasis (J47) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 3.2 | 6.5 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.1 | 6.1 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.4 | 5.9 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.0 | 7.2 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.3 | 6.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.4 | 6.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.2 | 4.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 2.3 | 4.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.2 | 4.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.9 | 4.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.8 | 2.1 | 4.4 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.1 | 6.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.2 | 1.0 | 4.0 | 7.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.1 | 3.7 | 6.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.2 | 4.2 | 6.4 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.2 | 3.7 | 8.1 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.1 | 4.0 | 6.9 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 1.2 | 3.7 | 6.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 3.5 | 6.9 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.3 | 6.6 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.7 | 6.2 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 7.6 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.4 | 6.6 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.5 | 6.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 2.3 | 4.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 2.3 | 4.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.4 | 4.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.8 | 4.9 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 4.7 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.0 | 7.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 1.1 | 4.3 | 7.8 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.1 | 4.0 | 7.5 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.3 | 4.6 | 6.8 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.3 | 4.0 | 8.8 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.1 | 4.2 | 7.4 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.2 | 3.8 | 6.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Lung diseases due to external agents (J60-J70) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.8 | 10.9 | 46.6 | 171.1 |
| 2003..... | 6.4 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 3.1 | 11.0 | 49.0 | 182.8 |
| 2002..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.9 | 11.2 | 50.3 | 192.9 |
| 2001..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.9 | 11.6 | 50.1 | 195.9 |
| 2000..... | 6.3 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 12.0 | 48.6 | 195.3 |
| 1999..... | 5.9 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.8 | 11.1 | 45.5 | 181.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.8 | 15.5 | 67.1 | 256.1 |
| 2003..... | 6.9 | * | * | * | 0.2 | 0.2 | 0.4 | 1.3 | 4.0 | 15.6 | 71.2 | 266.2 |
| 2002..... | 7.0 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.7 | 15.8 | 73.1 | 290.1 |
| 2001..... | 7.0 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.7 | 16.6 | 73.3 | 291.3 |
| 2000..... | 6.9 | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.5 | 17.7 | 72.1 | 301.5 |
| 1999..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 3.8 | 16.4 | 70.2 | 279.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 7.0 | 32.7 | 132.9 |
| 2003..... | 5.9 | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.2 | 7.0 | 34.2 | 145.9 |
| 2002..... | 6.0 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.1 | 7.4 | 35.2 | 150.7 |
| 2001..... | 6.1 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.1 | 7.4 | 34.9 | 156.0 |
| 2000..... | 5.8 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 7.2 | 33.3 | 152.1 |
| 1999..... | 5.3 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.8 | 29.5 | 142.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 0.5 | 1.7 | 5.5 | 28.3 | 104.7 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | 1.9 | 6.9 | 24.8 | 113.5 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | 2.1 | 6.5 | 30.2 | 136.4 |
| 2001..... | 1.6 | * | * | * | * | * | * | 0.6 | 1.9 | 6.7 | 28.7 | 165.1 |
| 2000..... | 1.5 | * | * | * | * | * | * | 0.7 | 2.0 | 6.4 | 30.4 | 140.7 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | 1.7 | 6.2 | 25.2 | 132.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | * | 2.6 | 8.3 | 35.4 | 130.3 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | 2.3 | 9.2 | 36.8 | 123.1 |
| 2002..... | 1.6 | * | * | * | * | * | * | * | 3.0 | 8.2 | 45.5 | 182.4 |
| 2001..... | 1.6 | * | * | * | * | * | * | * | 2.8 | 9.8 | 39.3 | 230.5 |
| 2000..... | 1.5 | * | * | * | * | * | * | * | 2.6 | 8.4 | 49.2 | 177.4 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | * | 8.1 | 35.5 | 161.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | * | * | 3.3 | 23.5 | 91.5 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | * | 5.0 | 16.6 | 108.6 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | * | 5.2 | 19.7 | 113.6 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | * | 4.3 | 21.5 | 133.1 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 4.8 | 17.8 | 122.7 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | * | 4.6 | 18.4 | 118.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Lung diseases due to external agents (J60-J70) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 2.9 | 11.2 | 47.5 | 173.8 |
| 2003..... | 7.2 | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.2 | 11.2 | 50.2 | 185.3 |
| 2002..... | 7.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.9 | 11.5 | 51.2 | 194.7 |
| 2001..... | 7.2 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 3.0 | 11.9 | 51.0 | 196.5 |
| 2000..... | 7.0 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.8 | 12.3 | 49.2 | 196.9 |
| 1999..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.8 | 11.4 | 46.2 | 183.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.8 | 16.0 | 68.8 | 262.1 |
| 2003..... | 7.8 | * | * | * | 0.2 | 0.2 | 0.5 | 1.3 | 4.2 | 16.0 | 72.9 | 272.4 |
| 2002..... | 7.9 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.7 | 16.2 | 74.2 | 294.1 |
| 2001..... | 7.8 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.8 | 17.0 | 74.7 | 293.0 |
| 2000..... | 7.7 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.6 | 18.3 | 72.8 | 305.9 |
| 1999..... | 7.2 | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 3.9 | 16.8 | 71.5 | 283.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.1 | 7.3 | 33.1 | 134.3 |
| 2003..... | 6.6 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 7.2 | 35.0 | 147.0 |
| 2002..... | 6.7 | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.2 | 7.5 | 35.9 | 151.9 |
| 2001..... | 6.7 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.2 | 7.6 | 35.4 | 156.4 |
| 2000..... | 6.4 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 7.4 | 33.9 | 152.8 |
| 1999..... | 5.8 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.9 | 6.9 | 29.9 | 143.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 7.7 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 11.2 | 48.1 | 177.0 |
| 2003..... | 8.0 | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.0 | 11.1 | 50.1 | 188.3 |
| 2002..... | 8.2 | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.8 | 11.3 | 51.6 | 198.7 |
| 2001..... | 8.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.9 | 11.7 | 51.3 | 198.1 |
| 2000..... | 7.8 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 11.9 | 49.3 | 200.8 |
| 1999..... | 7.2 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.7 | 11.0 | 46.1 | 185.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 8.5 | * | * | * | * | 0.3 | 0.5 | 1.4 | 3.5 | 15.7 | 69.7 | 267.6 |
| 2003..... | 8.7 | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.9 | 15.9 | 72.6 | 279.1 |
| 2002..... | 8.9 | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.6 | 15.9 | 74.9 | 301.6 |
| 2001..... | 8.7 | * | * | * | * | 0.3 | 0.6 | 1.3 | 3.7 | 16.9 | 75.6 | 299.5 |
| 2000..... | 8.6 | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 17.4 | 73.1 | 315.9 |
| 1999..... | 8.0 | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.6 | 16.1 | 71.0 | 289.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.0 | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 7.3 | 33.5 | 136.4 |
| 2003..... | 7.4 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.1 | 7.0 | 34.8 | 148.4 |
| 2002..... | 7.5 | * | * | * | * | 0.2 | 0.4 | 0.6 | 2.1 | 7.3 | 36.1 | 154.2 |
| 2001..... | 7.4 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.2 | 7.2 | 35.2 | 155.9 |
| 2000..... | 7.1 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.9 | 7.2 | 33.7 | 154.4 |
| 1999..... | 6.5 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.7 | 29.8 | 144.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Lung diseases due to external agents (J60-J70) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | * | 0.6 | 1.4 | 5.2 | 15.1 | 49.4 | 152.5 |
| 2003..... | 4.5 | * | * | * | * | * | 0.8 | 1.4 | 5.4 | 14.5 | 58.1 | 168.4 |
| 2002..... | 4.4 | * | * | * | * | * | 0.7 | 1.5 | 4.1 | 15.9 | 55.2 | 171.5 |
| 2001..... | 4.6 | * | * | * | * | * | 0.6 | 1.8 | 4.4 | 16.2 | 55.8 | 191.0 |
| 2000..... | 4.5 | * | * | * | * | * | 0.7 | 1.9 | 4.4 | 19.0 | 54.8 | 164.4 |
| 1999..... | 4.5 | * | * | * | * | * | 0.8 | 1.6 | 5.0 | 17.4 | 54.9 | 173.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 0.8 | 1.8 | 7.7 | 23.5 | 73.3 | 225.6 |
| 2003..... | 4.7 | * | * | * | * | * | * | 1.8 | 7.3 | 21.4 | 86.6 | 223.7 |
| 2002..... | 4.7 | * | * | * | * | * | 0.9 | 2.1 | 5.3 | 23.7 | 80.0 | 251.6 |
| 2001..... | 4.6 | * | * | * | * | * | 0.8 | 2.0 | 6.2 | 22.5 | 81.6 | 257.6 |
| 2000..... | 4.9 | * | * | * | * | * | 0.9 | 2.7 | 5.4 | 31.0 | 82.1 | 224.3 |
| 1999..... | 5.1 | * | * | * | * | * | 0.9 | 2.0 | 7.5 | 27.5 | 86.9 | 247.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | * | 1.1 | 3.1 | 9.0 | 35.9 | 123.7 |
| 2003..... | 4.3 | * | * | * | * | * | 0.9 | 1.0 | 3.9 | 9.7 | 42.0 | 146.9 |
| 2002..... | 4.1 | * | * | * | * | * | * | 0.9 | 3.1 | 10.3 | 41.1 | 140.9 |
| 2001..... | 4.6 | * | * | * | * | * | * | 1.5 | 2.9 | 11.7 | 41.3 | 166.4 |
| 2000..... | 4.1 | * | * | * | * | * | * | 1.2 | 3.6 | 10.5 | 39.4 | 142.2 |
| 1999..... | 4.0 | * | * | * | * | * | * | 1.2 | 2.9 | 10.3 | 36.9 | 146.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonitis due to solids and liquids (J69) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 9.6 | 42.9 | 164.9 |
| 2003..... | 6.0 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.8 | 9.6 | 45.0 | 176.2 |
| 2002..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.5 | 9.8 | 46.3 | 186.0 |
| 2001..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 10.0 | 45.8 | 189.4 |
| 2000..... | 5.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 10.3 | 44.5 | 187.6 |
| 1999..... | 5.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.5 | 9.5 | 41.1 | 175.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.2 | 12.9 | 58.6 | 237.2 |
| 2003..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.5 | 12.8 | 61.7 | 246.2 |
| 2002..... | 6.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.1 | 13.0 | 63.8 | 268.5 |
| 2001..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.2 | 13.3 | 63.1 | 270.3 |
| 2000..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.0 | 14.4 | 62.3 | 276.0 |
| 1999..... | 5.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 3.2 | 13.1 | 59.7 | 258.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.9 | 6.8 | 32.3 | 132.4 |
| 2003..... | 5.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 6.9 | 33.8 | 145.3 |
| 2002..... | 5.9 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 7.1 | 34.7 | 150.2 |
| 2001..... | 6.0 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.1 | 7.2 | 34.5 | 155.6 |
| 2000..... | 5.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.9 | 7.0 | 33.0 | 151.6 |
| 1999..... | 5.3 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.8 | 6.6 | 29.1 | 142.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 1.6 | 5.1 | 26.3 | 101.3 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | 1.7 | 6.5 | 22.8 | 113.0 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 1.8 | 5.3 | 29.0 | 131.4 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 1.8 | 6.1 | 27.4 | 161.3 |
| 2000..... | 1.4 | * | * | * | * | * | * | 0.6 | 1.9 | 5.9 | 27.9 | 137.4 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 1.6 | 5.5 | 24.6 | 130.5 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.4 | 7.4 | 31.8 | 121.9 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | 2.0 | 8.7 | 32.2 | 123.1 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 2.6 | 6.1 | 43.1 | 167.3 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 2.5 | 8.4 | 36.2 | 220.9 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | 2.5 | 7.4 | 42.8 | 169.3 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 6.8 | 33.9 | 159.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | * | 3.3 | 22.5 | 90.8 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | * | 4.9 | 16.3 | 107.8 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | * | 4.6 | 19.4 | 113.6 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | * | 4.3 | 21.5 | 132.2 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 4.8 | 17.8 | 121.7 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 4.5 | 18.4 | 116.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonitis due to solids and liquids (J69) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 9.9 | 43.7 | 167.4 |
| 2003..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.9 | 9.7 | 46.0 | 178.5 |
| 2002..... | 6.8 | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.6 | 10.1 | 47.1 | 187.8 |
| 2001..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 10.2 | 46.5 | 190.0 |
| 2000..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 10.6 | 45.1 | 189.0 |
| 1999..... | 6.0 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.5 | 9.7 | 41.7 | 176.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.3 | 13.2 | 60.1 | 242.7 |
| 2003..... | 6.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.6 | 13.0 | 63.1 | 251.5 |
| 2002..... | 7.0 | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 3.1 | 13.4 | 64.7 | 272.1 |
| 2001..... | 6.8 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.2 | 13.6 | 64.2 | 271.6 |
| 2000..... | 6.8 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.1 | 14.8 | 62.8 | 279.7 |
| 1999..... | 6.3 | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 3.3 | 13.3 | 60.6 | 261.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.1 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.0 | 7.1 | 32.8 | 133.8 |
| 2003..... | 6.5 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.1 | 7.0 | 34.6 | 146.5 |
| 2002..... | 6.6 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.1 | 7.3 | 35.4 | 151.4 |
| 2001..... | 6.6 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.2 | 7.4 | 34.9 | 156.0 |
| 2000..... | 6.3 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.0 | 7.1 | 33.6 | 152.3 |
| 1999..... | 5.8 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.8 | 29.4 | 142.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 7.2 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.4 | 9.7 | 44.1 | 170.4 |
| 2003..... | 7.5 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 9.5 | 45.7 | 181.0 |
| 2002..... | 7.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.5 | 9.7 | 47.3 | 191.3 |
| 2001..... | 7.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 9.9 | 46.6 | 191.2 |
| 2000..... | 7.3 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 10.0 | 45.0 | 192.5 |
| 1999..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 2.4 | 9.2 | 41.2 | 179.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 7.5 | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.9 | 12.7 | 60.2 | 247.4 |
| 2003..... | 7.6 | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.3 | 12.6 | 62.4 | 256.5 |
| 2002..... | 7.9 | * | * | * | * | 0.3 | 0.6 | 1.1 | 3.0 | 12.9 | 64.9 | 278.4 |
| 2001..... | 7.6 | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.1 | 13.2 | 64.4 | 276.9 |
| 2000..... | 7.5 | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.9 | 13.7 | 62.7 | 288.1 |
| 1999..... | 6.9 | * | * | * | * | 0.2 | 0.4 | 0.9 | 3.0 | 12.4 | 59.6 | 267.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 7.1 | 33.1 | 135.9 |
| 2003..... | 7.3 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.0 | 6.8 | 34.4 | 147.8 |
| 2002..... | 7.4 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 7.1 | 35.6 | 153.7 |
| 2001..... | 7.4 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.1 | 7.0 | 34.7 | 155.5 |
| 2000..... | 7.1 | * | * | * | * | * | 0.3 | 0.7 | 1.8 | 7.0 | 33.4 | 153.9 |
| 1999..... | 6.5 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.7 | 6.5 | 29.3 | 144.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonitis due to solids and liquids (J69) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | * | 0.6 | 1.4 | 5.0 | 14.2 | 47.6 | 147.4 |
| 2003..... | 4.4 | * | * | * | * | * | 0.7 | 1.3 | 5.3 | 13.9 | 55.5 | 165.7 |
| 2002..... | 4.2 | * | * | * | * | * | 0.7 | 1.4 | 3.8 | 15.2 | 52.8 | 167.3 |
| 2001..... | 4.4 | * | * | * | * | * | 0.6 | 1.6 | 4.1 | 15.4 | 53.6 | 187.3 |
| 2000..... | 4.3 | * | * | * | * | * | 0.7 | 1.9 | 4.3 | 17.9 | 52.0 | 160.6 |
| 1999..... | 4.3 | * | * | * | * | * | 0.7 | 1.5 | 4.5 | 16.4 | 52.6 | 168.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | * | 0.8 | 1.8 | 7.3 | 21.5 | 68.7 | 207.3 |
| 2003..... | 4.5 | * | * | * | * | * | * | 1.7 | 7.2 | 19.8 | 79.6 | 215.2 |
| 2002..... | 4.4 | * | * | * | * | * | 0.9 | 2.0 | 4.9 | 22.3 | 74.1 | 237.4 |
| 2001..... | 4.3 | * | * | * | * | * | 0.8 | 1.9 | 5.5 | 20.8 | 75.6 | 243.7 |
| 2000..... | 4.5 | * | * | * | * | * | 0.8 | 2.7 | 5.2 | 28.4 | 75.0 | 210.2 |
| 1999..... | 4.7 | * | * | * | * | * | 0.9 | 1.8 | 6.6 | 25.3 | 81.6 | 229.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | * | 1.0 | 3.1 | 8.9 | 35.7 | 123.7 |
| 2003..... | 4.3 | * | * | * | * | * | 0.8 | 1.0 | 3.7 | 9.7 | 41.9 | 146.5 |
| 2002..... | 4.1 | * | * | * | * | * | * | 0.9 | 3.0 | 10.2 | 40.8 | 140.5 |
| 2001..... | 4.6 | * | * | * | * | * | * | 1.4 | 2.9 | 11.6 | 41.3 | 166.4 |
| 2000..... | 4.1 | * | * | * | * | * | * | 1.2 | 3.6 | 10.4 | 39.1 | 142.2 |
| 1999..... | 4.0 | * | * | * | * | * | * | 1.2 | 2.9 | 10.2 | 36.4 | 145.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonitis due to food and vomit (J69.0) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 9.6 | 42.9 | 164.9 |
| 2003..... | 6.0 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.8 | 9.6 | 45.0 | 176.2 |
| 2002..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.5 | 9.8 | 46.3 | 186.0 |
| 2001..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 10.0 | 45.8 | 189.4 |
| 2000..... | 5.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 10.3 | 44.5 | 187.6 |
| 1999..... | 5.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.5 | 9.5 | 41.1 | 175.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.2 | 12.9 | 58.6 | 237.2 |
| 2003..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.5 | 12.8 | 61.7 | 246.2 |
| 2002..... | 6.3 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.1 | 13.0 | 63.8 | 268.5 |
| 2001..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.2 | 13.3 | 63.0 | 270.2 |
| 2000..... | 6.1 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.0 | 14.4 | 62.3 | 276.0 |
| 1999..... | 5.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 3.2 | 13.0 | 59.7 | 258.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.9 | 6.8 | 32.3 | 132.4 |
| 2003..... | 5.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 6.9 | 33.8 | 145.3 |
| 2002..... | 5.9 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 7.1 | 34.7 | 150.2 |
| 2001..... | 6.0 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.1 | 7.2 | 34.5 | 155.5 |
| 2000..... | 5.8 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.9 | 7.0 | 33.0 | 151.6 |
| 1999..... | 5.3 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.8 | 6.6 | 29.0 | 142.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 1.6 | 5.1 | 26.3 | 100.8 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | 1.7 | 6.5 | 22.8 | 113.0 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 1.8 | 5.3 | 29.0 | 131.4 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 1.8 | 6.1 | 27.4 | 161.3 |
| 2000..... | 1.4 | * | * | * | * | * | * | 0.6 | 1.9 | 5.9 | 27.9 | 137.4 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 1.6 | 5.5 | 24.6 | 130.5 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.4 | 7.4 | 31.8 | 121.9 |
| 2003..... | 1.3 | * | * | * | * | * | * | * | 2.0 | 8.7 | 32.2 | 123.1 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 2.6 | 6.1 | 43.1 | 167.3 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | 2.5 | 8.4 | 36.2 | 220.9 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | 2.5 | 7.4 | 42.8 | 169.3 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 6.8 | 33.9 | 159.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | * | 3.3 | 22.5 | 90.0 |
| 2003..... | 1.4 | * | * | * | * | * | * | * | * | 4.9 | 16.3 | 107.8 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | * | 4.6 | 19.4 | 113.6 |
| 2001..... | 1.5 | * | * | * | * | * | * | * | * | 4.3 | 21.5 | 132.2 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 4.8 | 17.8 | 121.7 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 4.5 | 18.4 | 116.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonitis due to food and vomit (J69.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 9.9 | 43.7 | 167.4 |
| 2003..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.9 | 9.7 | 46.0 | 178.5 |
| 2002..... | 6.8 | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.6 | 10.1 | 47.1 | 187.8 |
| 2001..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 10.2 | 46.5 | 189.9 |
| 2000..... | 6.5 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.5 | 10.6 | 45.1 | 189.0 |
| 1999..... | 6.0 | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.5 | 9.7 | 41.6 | 176.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.3 | 13.2 | 60.0 | 242.7 |
| 2003..... | 6.9 | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.6 | 13.0 | 63.1 | 251.5 |
| 2002..... | 7.0 | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 3.1 | 13.4 | 64.7 | 272.1 |
| 2001..... | 6.8 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.2 | 13.6 | 64.1 | 271.5 |
| 2000..... | 6.8 | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.1 | 14.8 | 62.8 | 279.6 |
| 1999..... | 6.3 | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 3.3 | 13.3 | 60.6 | 261.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.1 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.0 | 7.1 | 32.8 | 133.8 |
| 2003..... | 6.5 | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.1 | 7.0 | 34.6 | 146.4 |
| 2002..... | 6.6 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.1 | 7.3 | 35.4 | 151.4 |
| 2001..... | 6.6 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.2 | 7.4 | 34.9 | 156.0 |
| 2000..... | 6.3 | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.9 | 7.1 | 33.5 | 152.3 |
| 1999..... | 5.8 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 6.8 | 29.3 | 142.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 7.2 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.4 | 9.7 | 44.1 | 170.4 |
| 2003..... | 7.4 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.6 | 9.5 | 45.7 | 181.0 |
| 2002..... | 7.6 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.5 | 9.7 | 47.3 | 191.3 |
| 2001..... | 7.5 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 9.9 | 46.6 | 191.1 |
| 2000..... | 7.3 | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 10.0 | 45.0 | 192.4 |
| 1999..... | 6.7 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 2.4 | 9.2 | 41.2 | 179.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 7.5 | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.9 | 12.7 | 60.2 | 247.4 |
| 2003..... | 7.6 | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.3 | 12.6 | 62.4 | 256.5 |
| 2002..... | 7.9 | * | * | * | * | 0.3 | 0.6 | 1.1 | 3.0 | 12.9 | 64.9 | 278.4 |
| 2001..... | 7.6 | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.1 | 13.2 | 64.4 | 276.9 |
| 2000..... | 7.5 | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.9 | 13.7 | 62.7 | 288.0 |
| 1999..... | 6.9 | * | * | * | * | 0.2 | 0.4 | 0.9 | 3.0 | 12.4 | 59.6 | 267.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | * | * | * | 0.3 | 0.7 | 2.0 | 7.1 | 33.1 | 135.9 |
| 2003..... | 7.3 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.0 | 6.8 | 34.4 | 147.8 |
| 2002..... | 7.4 | * | * | * | * | 0.2 | 0.3 | 0.6 | 2.0 | 7.1 | 35.5 | 153.7 |
| 2001..... | 7.4 | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.1 | 7.0 | 34.7 | 155.4 |
| 2000..... | 7.1 | * | * | * | * | * | 0.3 | 0.7 | 1.8 | 7.0 | 33.4 | 153.9 |
| 1999..... | 6.4 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.7 | 6.5 | 29.2 | 144.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pneumonitis due to food and vomit (J69.0) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | * | 0.6 | 1.4 | 5.0 | 14.2 | 47.5 | 147.4 |
| 2003..... | 4.4 | * | * | * | * | * | 0.7 | 1.3 | 5.3 | 13.9 | 55.5 | 165.4 |
| 2002..... | 4.2 | * | * | * | * | * | 0.7 | 1.4 | 3.8 | 15.2 | 52.7 | 167.3 |
| 2001..... | 4.4 | * | * | * | * | * | 0.6 | 1.6 | 4.1 | 15.4 | 53.6 | 187.3 |
| 2000..... | 4.3 | * | * | * | * | * | 0.7 | 1.9 | 4.3 | 17.9 | 52.0 | 160.6 |
| 1999..... | 4.3 | * | * | * | * | * | 0.7 | 1.5 | 4.5 | 16.4 | 52.5 | 168.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | * | 0.8 | 1.8 | 7.3 | 21.5 | 68.5 | 207.3 |
| 2003..... | 4.5 | * | * | * | * | * | * | 1.7 | 7.2 | 19.8 | 79.6 | 215.2 |
| 2002..... | 4.4 | * | * | * | * | * | 0.9 | 2.0 | 4.9 | 22.3 | 74.1 | 237.4 |
| 2001..... | 4.3 | * | * | * | * | * | 0.8 | 1.9 | 5.5 | 20.8 | 75.6 | 243.7 |
| 2000..... | 4.5 | * | * | * | * | * | 0.8 | 2.7 | 5.2 | 28.4 | 75.0 | 210.2 |
| 1999..... | 4.7 | * | * | * | * | * | 0.9 | 1.8 | 6.6 | 25.1 | 81.6 | 229.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | * | * | 1.0 | 3.1 | 8.9 | 35.7 | 123.7 |
| 2003..... | 4.3 | * | * | * | * | * | 0.8 | 1.0 | 3.7 | 9.7 | 41.9 | 146.1 |
| 2002..... | 4.1 | * | * | * | * | * | * | 0.9 | 3.0 | 10.2 | 40.6 | 140.5 |
| 2001..... | 4.6 | * | * | * | * | * | * | 1.4 | 2.9 | 11.6 | 41.3 | 166.4 |
| 2000..... | 4.1 | * | * | * | * | * | * | 1.2 | 3.6 | 10.4 | 39.1 | 142.2 |
| 1999..... | 4.0 | * | * | * | * | * | * | 1.2 | 2.9 | 10.2 | 36.2 | 145.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other pneumonitis due to solids and liquids (J69.1-J69.8)

All origins 3/, both sexes

Table with 13 columns (Year 2004-1999) and 13 rows (Age groups) showing death rates for 'Other pneumonitis due to solids and liquids' for 'All origins 3/, both sexes'. All cells contain an asterisk (*).

All origins 3/, male

Table with 13 columns (Year 2004-1999) and 13 rows (Age groups) showing death rates for 'Other pneumonitis due to solids and liquids' for 'All origins 3/, male'. All cells contain an asterisk (*).

All origins 3/, female

Table with 13 columns (Year 2004-1999) and 13 rows (Age groups) showing death rates for 'Other pneumonitis due to solids and liquids' for 'All origins 3/, female'. All cells contain an asterisk (*).

Hispanic, both sexes

Table with 13 columns (Year 2004-1999) and 13 rows (Age groups) showing death rates for 'Other pneumonitis due to solids and liquids' for 'Hispanic, both sexes'. All cells contain an asterisk (*).

Hispanic, male

Table with 13 columns (Year 2004-1999) and 13 rows (Age groups) showing death rates for 'Other pneumonitis due to solids and liquids' for 'Hispanic, male'. All cells contain an asterisk (*).

Hispanic, female

Table with 13 columns (Year 2004-1999) and 13 rows (Age groups) showing death rates for 'Other pneumonitis due to solids and liquids' for 'Hispanic, female'. All cells contain an asterisk (*).

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Suppurative and necrotic conditions of lower respiratory tract (J85-J86) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.3 | 2.3 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 1.4 | 2.1 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.5 | 2.3 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 1.7 | 3.1 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.5 | 2.9 |
| 1999..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.0 | 1.5 | 2.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 1.7 | 4.0 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.4 | 0.5 | 1.1 | 1.7 | 3.3 |
| 2002..... | 0.3 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.0 | 2.3 | 3.5 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 2.3 | 4.9 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.2 | 5.8 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.5 | 2.2 | 3.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.2 | 1.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 1.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 1.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.2 | 2.2 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 1.3 | 2.1 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.4 | 2.2 |
| 2001..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.6 | 3.1 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.5 | 2.7 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.9 | 1.5 | 2.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 1.6 | 4.0 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.4 | 0.5 | 1.1 | 1.5 | 3.2 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.1 | 3.2 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 2.2 | 4.9 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.1 | 5.5 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.4 | 1.4 | 2.2 | 3.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.1 | 1.6 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.7 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 2.3 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.2 | 1.5 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pleural effusion and plaque (J90,J92) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.5 | 9.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 8.8 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.5 | 10.1 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.6 | 10.2 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.1 | 0.6 | 2.4 | 8.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 9.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.2 | 11.3 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.9 | 11.4 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.7 | 3.0 | 11.2 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | 0.9 | 2.9 | 12.4 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 0.7 | 2.8 | 11.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.0 | 11.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 7.9 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 0.7 | 2.2 | 7.7 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | 0.7 | 2.2 | 9.6 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.4 | 9.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 0.5 | 2.1 | 8.0 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.0 | 8.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.6 | 9.2 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 9.2 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 | 10.4 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.6 | 10.1 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.1 | 0.6 | 2.4 | 9.0 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.4 | 9.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.3 | 11.8 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 0.7 | 2.8 | 12.1 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.6 | 3.0 | 11.8 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 0.9 | 2.8 | 12.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 0.7 | 2.8 | 11.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | 0.9 | 3.0 | 11.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 8.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 0.7 | 2.3 | 7.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 0.7 | 2.2 | 9.8 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 0.5 | 2.4 | 9.1 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 0.5 | 2.1 | 8.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 | 8.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of respiratory system (J80-J84,J93-J98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 8.5 | 5.9 | 0.5 | 0.1 | 0.2 | 0.4 | 1.0 | 2.8 | 8.7 | 28.1 | 69.2 | 125.2 |
| 2003..... | 8.0 | 6.2 | 0.4 | 0.1 | 0.2 | 0.4 | 0.9 | 2.8 | 7.8 | 26.2 | 63.5 | 123.3 |
| 2002..... | 7.9 | 6.1 | 0.4 | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 8.3 | 26.6 | 63.0 | 118.1 |
| 2001..... | 8.3 | 6.5 | 0.4 | 0.1 | 0.3 | 0.4 | 1.0 | 3.0 | 8.8 | 28.1 | 67.6 | 125.7 |
| 2000..... | 8.3 | 6.4 | 0.4 | 0.2 | 0.3 | 0.5 | 1.1 | 3.1 | 9.2 | 27.5 | 66.8 | 125.8 |
| 1999..... | 8.1 | 5.8 | 0.4 | 0.1 | 0.2 | 0.5 | 1.0 | 2.8 | 8.6 | 27.1 | 65.3 | 131.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 8.5 | 5.9 | 0.6 | 0.1 | 0.2 | 0.3 | 1.0 | 3.0 | 9.7 | 34.6 | 87.5 | 149.7 |
| 2003..... | 7.8 | 7.1 | 0.4 | 0.1 | 0.3 | 0.4 | 0.9 | 2.9 | 9.1 | 31.7 | 79.1 | 145.5 |
| 2002..... | 7.7 | 7.3 | 0.5 | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 9.4 | 32.9 | 77.5 | 138.9 |
| 2001..... | 8.1 | 8.1 | 0.4 | 0.2 | 0.3 | 0.4 | 1.1 | 3.2 | 9.9 | 34.1 | 82.5 | 147.4 |
| 2000..... | 8.1 | 7.1 | 0.4 | 0.2 | 0.3 | 0.5 | 1.2 | 3.2 | 10.4 | 33.0 | 83.8 | 158.3 |
| 1999..... | 7.9 | 6.6 | 0.4 | 0.1 | 0.3 | 0.5 | 1.1 | 2.9 | 9.5 | 33.1 | 82.9 | 161.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 8.6 | 5.8 | 0.3 | * | 0.3 | 0.4 | 1.0 | 2.6 | 7.8 | 22.7 | 56.9 | 114.2 |
| 2003..... | 8.1 | 5.3 | 0.4 | * | 0.2 | 0.4 | 1.0 | 2.7 | 6.7 | 21.6 | 53.0 | 113.5 |
| 2002..... | 8.0 | 4.9 | 0.3 | 0.2 | 0.3 | 0.4 | 1.0 | 2.8 | 7.3 | 21.4 | 53.4 | 109.0 |
| 2001..... | 8.6 | 4.9 | 0.4 | 0.1 | 0.3 | 0.5 | 1.0 | 2.7 | 7.7 | 23.2 | 57.9 | 116.6 |
| 2000..... | 8.4 | 5.7 | 0.3 | 0.1 | 0.3 | 0.5 | 1.1 | 2.9 | 8.0 | 23.0 | 55.7 | 112.6 |
| 1999..... | 8.3 | 4.9 | 0.3 | 0.2 | 0.2 | 0.6 | 1.0 | 2.8 | 7.7 | 22.2 | 53.8 | 119.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | 4.5 | * | * | * | * | 0.8 | 2.1 | 7.5 | 26.2 | 67.3 | 105.1 |
| 2003..... | 3.2 | 5.4 | * | * | * | 0.3 | 0.7 | 2.1 | 8.0 | 24.6 | 56.4 | 106.2 |
| 2002..... | 3.0 | 4.2 | * | * | * | 0.3 | 0.6 | 2.3 | 7.6 | 24.5 | 54.2 | 104.2 |
| 2001..... | 3.2 | 4.4 | * | * | * | * | 0.7 | 2.6 | 8.2 | 24.0 | 61.0 | 125.7 |
| 2000..... | 3.2 | 4.9 | * | * | * | 0.4 | 1.0 | 2.4 | 8.9 | 24.0 | 60.4 | 124.1 |
| 1999..... | 3.1 | 3.7 | * | * | * | * | 0.8 | 2.7 | 8.7 | 23.9 | 61.2 | 122.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | 0.7 | 2.6 | 8.6 | 30.1 | 78.7 | 105.1 |
| 2003..... | 3.1 | 5.7 | * | * | * | * | 0.7 | 2.8 | 10.5 | 29.5 | 61.7 | 117.0 |
| 2002..... | 2.9 | * | * | * | * | * | 0.7 | 2.2 | 9.2 | 32.3 | 58.7 | 98.7 |
| 2001..... | 3.1 | * | * | * | * | * | 0.8 | 2.8 | 9.7 | 28.6 | 70.6 | 154.3 |
| 2000..... | 3.1 | * | * | * | * | * | 1.1 | 2.6 | 11.0 | 27.4 | 74.7 | 120.9 |
| 1999..... | 2.9 | * | * | * | * | * | 0.9 | 2.4 | 10.3 | 27.7 | 77.2 | 140.2 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.7 | 5.0 | * | * | * | * | 0.9 | 1.6 | 6.5 | 23.1 | 59.5 | 105.2 |
| 2003..... | 3.3 | 5.2 | * | * | * | * | * | 1.5 | 5.8 | 20.6 | 52.7 | 100.8 |
| 2002..... | 3.1 | * | * | * | * | * | * | 2.4 | 6.2 | 18.2 | 51.2 | 106.9 |
| 2001..... | 3.3 | * | * | * | * | * | * | 2.5 | 6.9 | 20.5 | 54.4 | 111.7 |
| 2000..... | 3.4 | * | * | * | * | * | 0.8 | 2.1 | 7.2 | 21.2 | 50.8 | 125.6 |
| 1999..... | 3.2 | * | * | * | * | * | * | 3.0 | 7.4 | 21.0 | 50.7 | 113.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of respiratory system (J80-J84,J93-J98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 9.3 | 6.2 | 0.5 | 0.1 | 0.2 | 0.4 | 1.1 | 2.9 | 8.8 | 28.2 | 69.2 | 126.1 |
| 2003..... | 8.7 | 6.4 | 0.4 | 0.1 | 0.3 | 0.5 | 1.0 | 2.9 | 7.8 | 26.2 | 63.7 | 123.8 |
| 2002..... | 8.6 | 6.6 | 0.4 | 0.1 | 0.3 | 0.4 | 1.1 | 2.8 | 8.4 | 26.7 | 63.3 | 118.4 |
| 2001..... | 9.1 | 7.0 | 0.4 | 0.2 | 0.3 | 0.5 | 1.1 | 3.0 | 8.8 | 28.3 | 67.7 | 125.4 |
| 2000..... | 8.9 | 6.7 | 0.4 | 0.2 | 0.3 | 0.5 | 1.2 | 3.1 | 9.2 | 27.6 | 66.8 | 125.6 |
| 1999..... | 8.8 | 6.2 | 0.4 | 0.2 | 0.2 | 0.6 | 1.1 | 2.8 | 8.5 | 27.3 | 65.3 | 131.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 9.4 | 6.5 | 0.6 | 0.1 | 0.2 | 0.4 | 1.1 | 3.0 | 9.8 | 34.8 | 87.8 | 151.8 |
| 2003..... | 8.6 | 7.5 | 0.4 | * | 0.3 | 0.5 | 0.9 | 2.9 | 8.9 | 31.7 | 79.9 | 146.7 |
| 2002..... | 8.5 | 8.0 | 0.5 | 0.1 | 0.3 | 0.4 | 1.1 | 2.7 | 9.4 | 32.9 | 78.3 | 140.4 |
| 2001..... | 8.8 | 9.0 | 0.4 | 0.2 | 0.3 | 0.4 | 1.1 | 3.3 | 9.9 | 34.3 | 82.8 | 147.0 |
| 2000..... | 8.9 | 7.6 | 0.4 | 0.2 | 0.3 | 0.5 | 1.2 | 3.3 | 10.4 | 33.2 | 84.0 | 159.5 |
| 1999..... | 8.6 | 7.3 | 0.5 | 0.2 | 0.3 | 0.5 | 1.1 | 2.9 | 9.4 | 33.3 | 82.8 | 162.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 9.3 | 5.9 | 0.4 | * | 0.3 | 0.4 | 1.1 | 2.7 | 7.9 | 22.6 | 56.7 | 114.5 |
| 2003..... | 8.8 | 5.3 | 0.4 | * | 0.2 | 0.4 | 1.0 | 2.8 | 6.7 | 21.6 | 53.0 | 113.7 |
| 2002..... | 8.7 | 5.1 | * | 0.2 | 0.3 | 0.5 | 1.0 | 2.8 | 7.4 | 21.5 | 53.4 | 108.9 |
| 2001..... | 9.3 | 4.9 | 0.5 | 0.1 | 0.3 | 0.6 | 1.1 | 2.8 | 7.8 | 23.3 | 57.9 | 116.4 |
| 2000..... | 9.0 | 5.8 | 0.3 | 0.1 | 0.3 | 0.6 | 1.1 | 2.9 | 8.0 | 23.0 | 55.7 | 111.9 |
| 1999..... | 8.9 | 5.2 | 0.4 | 0.2 | 0.2 | 0.6 | 1.1 | 2.7 | 7.7 | 22.3 | 53.9 | 119.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 10.4 | 4.1 | 0.4 | 0.1 | 0.2 | 0.3 | 0.9 | 2.5 | 8.6 | 29.1 | 71.8 | 130.7 |
| 2003..... | 9.7 | 4.3 | 0.3 | 0.1 | 0.2 | 0.4 | 0.8 | 2.5 | 7.6 | 27.0 | 66.2 | 128.4 |
| 2002..... | 9.6 | 4.2 | 0.3 | 0.1 | 0.2 | 0.4 | 0.9 | 2.4 | 8.0 | 27.3 | 65.4 | 122.5 |
| 2001..... | 10.0 | 4.4 | 0.3 | 0.2 | 0.2 | 0.4 | 0.9 | 2.6 | 8.5 | 28.9 | 69.9 | 128.7 |
| 2000..... | 9.9 | 4.8 | 0.3 | 0.2 | 0.2 | 0.5 | 1.0 | 2.8 | 8.9 | 28.2 | 68.5 | 129.7 |
| 1999..... | 9.7 | 4.2 | 0.4 | 0.1 | 0.3 | 0.5 | 0.9 | 2.5 | 8.3 | 27.5 | 66.7 | 134.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 10.4 | 4.2 | 0.5 | * | 0.2 | 0.3 | 0.8 | 2.7 | 9.6 | 35.8 | 90.7 | 156.9 |
| 2003..... | 9.6 | 4.6 | * | * | 0.2 | 0.5 | 0.8 | 2.7 | 8.7 | 32.7 | 82.6 | 151.6 |
| 2002..... | 9.4 | 5.2 | * | * | 0.2 | 0.3 | 0.9 | 2.5 | 9.0 | 33.7 | 80.8 | 145.5 |
| 2001..... | 9.8 | 5.7 | * | 0.2 | 0.2 | 0.3 | 1.0 | 2.9 | 9.6 | 35.4 | 85.8 | 151.0 |
| 2000..... | 9.8 | 5.9 | * | 0.2 | 0.3 | 0.4 | 1.0 | 2.9 | 10.1 | 33.8 | 86.0 | 164.6 |
| 1999..... | 9.5 | 4.9 | * | * | 0.3 | 0.5 | 0.9 | 2.6 | 9.2 | 34.0 | 84.8 | 166.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 10.4 | 3.9 | * | * | 0.2 | 0.3 | 0.9 | 2.3 | 7.6 | 23.3 | 58.8 | 119.0 |
| 2003..... | 9.9 | 4.0 | * | * | 0.2 | 0.4 | 0.7 | 2.4 | 6.6 | 22.2 | 55.1 | 118.2 |
| 2002..... | 9.7 | 3.2 | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 7.1 | 21.8 | 55.1 | 112.6 |
| 2001..... | 10.3 | 3.0 | * | * | 0.2 | 0.5 | 0.8 | 2.3 | 7.5 | 23.5 | 59.5 | 119.4 |
| 2000..... | 10.0 | 3.5 | * | * | 0.2 | 0.5 | 1.0 | 2.7 | 7.8 | 23.5 | 57.1 | 115.6 |
| 1999..... | 9.8 | 3.3 | * | * | 0.2 | 0.5 | 0.8 | 2.4 | 7.5 | 22.2 | 54.9 | 122.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of respiratory system (J80-J84,J93-J98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.7 | 14.5 | 0.8 | * | * | 0.8 | 2.4 | 5.8 | 12.5 | 25.6 | 48.6 | 80.6 |
| 2003..... | 5.3 | 14.4 | * | * | 0.6 | 0.9 | 2.2 | 5.5 | 10.6 | 22.5 | 44.8 | 78.5 |
| 2002..... | 5.5 | 15.5 | 0.9 | 0.3 | 0.6 | 0.9 | 2.0 | 5.3 | 12.2 | 24.4 | 45.6 | 77.4 |
| 2001..... | 6.0 | 17.5 | * | * | 0.5 | 1.1 | 2.4 | 6.4 | 12.9 | 26.6 | 50.5 | 90.8 |
| 2000..... | 5.8 | 15.9 | * | * | 0.5 | 0.9 | 2.4 | 6.2 | 12.4 | 26.8 | 53.0 | 83.6 |
| 1999..... | 5.8 | 15.1 | * | * | * | 1.1 | 2.4 | 5.7 | 11.3 | 28.1 | 53.0 | 95.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 5.8 | 15.5 | * | * | * | 0.8 | 2.7 | 6.0 | 13.7 | 32.1 | 62.2 | 102.6 |
| 2003..... | 5.3 | 17.7 | * | * | 0.8 | 0.9 | 1.9 | 5.0 | 13.2 | 28.2 | 59.7 | 100.2 |
| 2002..... | 5.4 | 18.5 | * | * | * | 0.8 | 2.2 | 5.2 | 14.1 | 29.6 | 56.9 | 88.6 |
| 2001..... | 5.6 | 22.3 | * | * | * | 1.1 | 1.9 | 6.5 | 14.2 | 29.5 | 58.2 | 108.6 |
| 2000..... | 5.8 | 15.7 | * | * | * | 0.9 | 2.6 | 7.0 | 13.9 | 31.0 | 67.0 | 106.9 |
| 1999..... | 5.5 | 17.7 | * | * | * | 0.8 | 2.2 | 6.1 | 12.2 | 31.6 | 66.9 | 115.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 5.7 | 13.5 | * | * | * | 0.8 | 2.1 | 5.5 | 11.5 | 20.9 | 40.9 | 71.9 |
| 2003..... | 5.2 | 11.0 | * | * | * | 0.9 | 2.5 | 6.0 | 8.5 | 18.4 | 36.3 | 70.2 |
| 2002..... | 5.6 | 12.3 | * | * | * | 0.9 | 1.8 | 5.4 | 10.7 | 20.7 | 39.3 | 73.2 |
| 2001..... | 6.4 | 12.4 | * | * | * | 1.1 | 2.9 | 6.3 | 11.8 | 24.6 | 46.2 | 84.3 |
| 2000..... | 5.9 | 16.2 | * | * | * | 0.9 | 2.1 | 5.5 | 11.3 | 23.9 | 45.2 | 75.0 |
| 1999..... | 6.1 | 12.5 | * | * | * | 1.4 | 2.6 | 5.4 | 10.5 | 25.7 | 45.2 | 87.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XI. Diseases of the digestive system (K00-K92) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 29.4 | 14.2 | 0.8 | 0.2 | 0.7 | 2.1 | 10.9 | 29.5 | 44.5 | 78.4 | 162.5 | 387.5 |
| 2003..... | 30.2 | 13.8 | 0.8 | 0.3 | 0.6 | 2.4 | 11.5 | 30.2 | 45.4 | 82.5 | 167.1 | 403.8 |
| 2002..... | 30.1 | 12.6 | 0.8 | 0.3 | 0.6 | 2.3 | 11.7 | 29.8 | 44.7 | 81.7 | 170.0 | 415.3 |
| 2001..... | 30.1 | 11.9 | 0.8 | 0.3 | 0.5 | 2.4 | 11.7 | 29.7 | 45.2 | 83.8 | 168.8 | 431.4 |
| 2000..... | 29.9 | 13.1 | 0.7 | 0.3 | 0.7 | 2.4 | 11.9 | 28.6 | 45.6 | 82.1 | 170.4 | 440.7 |
| 1999..... | 29.7 | 13.2 | 0.7 | 0.3 | 0.6 | 2.4 | 11.8 | 28.0 | 45.8 | 85.2 | 169.0 | 436.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 30.0 | 15.6 | 0.9 | 0.2 | 0.7 | 2.8 | 14.1 | 41.3 | 58.6 | 91.6 | 176.1 | 366.9 |
| 2003..... | 30.9 | 16.6 | 0.9 | 0.3 | 0.7 | 2.9 | 14.6 | 42.6 | 60.5 | 97.6 | 181.6 | 388.9 |
| 2002..... | 30.7 | 15.1 | 0.9 | 0.3 | 0.7 | 2.9 | 15.0 | 42.5 | 59.0 | 96.9 | 184.1 | 404.4 |
| 2001..... | 30.6 | 13.4 | 0.9 | 0.3 | 0.6 | 2.9 | 15.2 | 41.9 | 59.8 | 99.5 | 181.8 | 428.2 |
| 2000..... | 30.3 | 14.8 | 0.9 | 0.3 | 0.7 | 3.0 | 15.8 | 40.7 | 60.1 | 97.7 | 186.4 | 441.9 |
| 1999..... | 30.4 | 16.5 | 0.7 | 0.3 | 0.6 | 2.9 | 15.8 | 40.4 | 60.7 | 102.3 | 185.8 | 438.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 28.8 | 12.8 | 0.6 | 0.3 | 0.7 | 1.5 | 7.7 | 18.2 | 31.4 | 67.3 | 153.3 | 396.8 |
| 2003..... | 29.4 | 10.8 | 0.7 | 0.3 | 0.5 | 1.8 | 8.4 | 18.2 | 31.4 | 69.9 | 157.4 | 410.4 |
| 2002..... | 29.5 | 10.0 | 0.7 | 0.3 | 0.6 | 1.8 | 8.5 | 17.7 | 31.5 | 69.0 | 160.6 | 420.0 |
| 2001..... | 29.7 | 10.3 | 0.6 | 0.3 | 0.5 | 1.9 | 8.3 | 18.0 | 31.8 | 70.8 | 160.2 | 432.7 |
| 2000..... | 29.4 | 11.4 | 0.5 | 0.3 | 0.5 | 1.8 | 8.1 | 16.9 | 32.3 | 69.3 | 159.9 | 440.2 |
| 1999..... | 29.1 | 9.7 | 0.8 | 0.4 | 0.6 | 1.9 | 7.8 | 15.9 | 32.2 | 71.1 | 158.1 | 435.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 15.6 | 12.2 | 0.6 | * | 0.6 | 1.9 | 11.0 | 34.6 | 53.7 | 89.7 | 166.0 | 271.6 |
| 2003..... | 16.5 | 12.2 | 0.8 | 0.4 | 0.5 | 2.5 | 11.7 | 35.3 | 60.0 | 98.5 | 179.4 | 293.8 |
| 2002..... | 16.2 | 11.8 | 0.9 | 0.3 | 0.7 | 2.2 | 12.1 | 36.4 | 58.1 | 102.6 | 168.1 | 290.5 |
| 2001..... | 16.3 | 10.8 | 0.8 | * | 0.5 | 2.1 | 12.3 | 36.8 | 58.3 | 102.0 | 178.2 | 364.5 |
| 2000..... | 16.4 | 11.5 | 0.7 | * | 0.6 | 2.4 | 13.4 | 37.3 | 59.5 | 102.9 | 183.3 | 366.3 |
| 1999..... | 16.7 | 12.0 | 0.7 | * | 0.7 | 2.7 | 14.3 | 37.1 | 60.5 | 106.5 | 172.3 | 411.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 18.4 | 12.3 | * | * | 0.6 | 2.8 | 16.4 | 55.1 | 75.9 | 103.4 | 176.0 | 263.4 |
| 2003..... | 19.7 | 15.4 | * | * | 0.6 | 3.4 | 17.1 | 55.5 | 88.4 | 115.9 | 199.4 | 275.5 |
| 2002..... | 19.9 | 12.9 | * | * | 0.7 | 3.1 | 18.1 | 57.5 | 85.3 | 133.2 | 194.9 | 284.4 |
| 2001..... | 19.7 | 12.7 | * | * | * | 2.8 | 18.2 | 58.4 | 87.8 | 128.2 | 203.9 | 373.3 |
| 2000..... | 20.1 | 12.4 | * | * | 0.7 | 3.5 | 20.3 | 58.7 | 90.2 | 134.8 | 208.5 | 387.0 |
| 1999..... | 20.3 | 15.8 | * | * | 0.7 | 3.6 | 21.9 | 58.9 | 89.8 | 138.0 | 196.0 | 427.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 12.6 | 12.2 | * | * | * | 0.9 | 5.1 | 13.9 | 33.6 | 78.6 | 159.0 | 275.9 |
| 2003..... | 13.2 | 9.0 | * | * | * | 1.5 | 5.8 | 15.1 | 34.4 | 84.7 | 165.6 | 303.1 |
| 2002..... | 12.3 | 10.6 | * | * | * | 1.2 | 5.5 | 15.3 | 33.7 | 78.3 | 149.8 | 293.4 |
| 2001..... | 12.7 | 8.7 | * | * | * | 1.3 | 5.9 | 15.5 | 32.1 | 81.4 | 160.8 | 360.2 |
| 2000..... | 12.5 | 10.6 | * | * | * | 1.0 | 6.1 | 16.3 | 32.2 | 77.8 | 166.4 | 356.1 |
| 1999..... | 12.9 | 8.1 | * | * | * | 1.6 | 6.2 | 15.9 | 34.6 | 81.9 | 156.5 | 403.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XI. Diseases of the digestive system (K00-K92) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 31.6 | 14.7 | 0.8 | 0.2 | 0.7 | 2.2 | 10.8 | 28.8 | 43.6 | 77.3 | 162.0 | 392.0 |
| 2003..... | 32.3 | 14.0 | 0.8 | 0.3 | 0.6 | 2.3 | 11.4 | 29.5 | 44.1 | 81.2 | 166.1 | 407.8 |
| 2002..... | 32.1 | 12.6 | 0.8 | 0.3 | 0.6 | 2.4 | 11.6 | 29.1 | 43.4 | 79.9 | 169.6 | 419.6 |
| 2001..... | 32.1 | 11.9 | 0.8 | 0.3 | 0.5 | 2.4 | 11.6 | 28.9 | 44.0 | 82.3 | 167.9 | 433.0 |
| 2000..... | 31.7 | 13.3 | 0.7 | 0.3 | 0.7 | 2.4 | 11.7 | 27.7 | 44.4 | 80.5 | 169.4 | 442.5 |
| 1999..... | 31.4 | 13.3 | 0.7 | 0.3 | 0.6 | 2.4 | 11.4 | 27.0 | 44.6 | 83.6 | 168.4 | 436.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 31.9 | 16.5 | 1.0 | 0.2 | 0.7 | 2.7 | 13.6 | 39.6 | 56.9 | 90.4 | 175.9 | 371.5 |
| 2003..... | 32.7 | 16.7 | 0.9 | 0.3 | 0.7 | 2.8 | 14.1 | 41.0 | 58.0 | 96.1 | 180.3 | 393.5 |
| 2002..... | 32.3 | 15.6 | 0.9 | 0.3 | 0.7 | 2.8 | 14.4 | 40.7 | 56.5 | 94.0 | 183.0 | 408.8 |
| 2001..... | 32.2 | 13.4 | 0.9 | 0.4 | 0.6 | 2.9 | 14.8 | 40.1 | 57.2 | 97.2 | 180.3 | 430.0 |
| 2000..... | 31.7 | 15.1 | 1.0 | 0.3 | 0.8 | 2.8 | 15.1 | 38.8 | 57.5 | 94.9 | 185.0 | 443.4 |
| 1999..... | 31.7 | 16.4 | 0.7 | 0.3 | 0.6 | 2.8 | 15.0 | 38.5 | 58.3 | 99.9 | 184.9 | 437.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 31.3 | 12.9 | 0.6 | 0.3 | 0.7 | 1.6 | 8.0 | 18.5 | 31.1 | 66.4 | 152.6 | 401.3 |
| 2003..... | 31.8 | 11.2 | 0.8 | 0.2 | 0.6 | 1.9 | 8.7 | 18.4 | 31.1 | 68.7 | 156.7 | 414.0 |
| 2002..... | 31.9 | 9.5 | 0.7 | 0.3 | 0.6 | 1.9 | 8.9 | 17.8 | 31.1 | 68.2 | 160.8 | 424.2 |
| 2001..... | 32.0 | 10.4 | 0.6 | 0.3 | 0.5 | 1.9 | 8.5 | 18.2 | 31.7 | 70.0 | 159.8 | 434.3 |
| 2000..... | 31.6 | 11.4 | 0.4 | 0.3 | 0.5 | 1.9 | 8.3 | 16.9 | 32.2 | 68.5 | 159.3 | 442.1 |
| 1999..... | 31.1 | 10.0 | 0.8 | 0.4 | 0.6 | 2.0 | 8.0 | 15.9 | 31.9 | 70.3 | 157.7 | 435.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 34.3 | 10.0 | 0.6 | 0.2 | 0.5 | 2.0 | 10.6 | 28.2 | 42.1 | 77.3 | 164.2 | 401.2 |
| 2003..... | 35.0 | 10.4 | 0.7 | 0.2 | 0.5 | 2.1 | 11.2 | 28.4 | 42.4 | 81.3 | 168.6 | 416.6 |
| 2002..... | 34.9 | 9.2 | 0.6 | 0.2 | 0.5 | 2.1 | 11.5 | 27.9 | 42.0 | 80.0 | 171.8 | 430.9 |
| 2001..... | 34.5 | 7.7 | 0.6 | 0.3 | 0.4 | 2.2 | 11.1 | 27.4 | 42.0 | 82.1 | 169.3 | 443.2 |
| 2000..... | 33.9 | 9.2 | 0.4 | 0.3 | 0.6 | 2.0 | 11.0 | 25.9 | 42.4 | 79.8 | 170.5 | 451.0 |
| 1999..... | 33.6 | 9.1 | 0.6 | 0.3 | 0.5 | 2.2 | 10.7 | 25.2 | 42.5 | 82.6 | 169.3 | 444.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 34.0 | 11.1 | 0.7 | 0.2 | 0.5 | 2.5 | 13.4 | 38.6 | 54.5 | 89.4 | 176.6 | 379.8 |
| 2003..... | 34.8 | 13.0 | 0.8 | 0.3 | 0.7 | 2.5 | 14.0 | 39.6 | 55.4 | 95.2 | 181.2 | 400.6 |
| 2002..... | 34.5 | 11.5 | 0.7 | 0.3 | 0.6 | 2.5 | 14.4 | 39.1 | 54.4 | 93.5 | 184.2 | 417.2 |
| 2001..... | 33.9 | 7.9 | 0.8 | 0.3 | 0.5 | 2.6 | 14.0 | 38.0 | 54.2 | 95.9 | 180.8 | 436.1 |
| 2000..... | 33.4 | 10.5 | 0.6 | 0.3 | 0.6 | 2.5 | 14.3 | 36.5 | 54.9 | 93.6 | 185.2 | 449.5 |
| 1999..... | 33.3 | 11.0 | 0.5 | 0.3 | 0.5 | 2.5 | 14.1 | 36.0 | 55.6 | 98.8 | 185.6 | 441.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 34.5 | 8.9 | 0.5 | 0.2 | 0.5 | 1.5 | 7.7 | 17.8 | 30.3 | 66.9 | 155.7 | 410.8 |
| 2003..... | 35.1 | 7.6 | 0.6 | 0.2 | 0.4 | 1.7 | 8.4 | 17.4 | 30.1 | 69.4 | 160.1 | 423.6 |
| 2002..... | 35.2 | 6.8 | 0.6 | 0.2 | 0.5 | 1.6 | 8.6 | 16.8 | 30.2 | 68.6 | 163.5 | 436.9 |
| 2001..... | 35.2 | 7.4 | 0.5 | 0.3 | 0.3 | 1.8 | 8.2 | 17.0 | 30.5 | 70.3 | 161.7 | 446.1 |
| 2000..... | 34.4 | 7.8 | * | 0.3 | 0.5 | 1.5 | 7.7 | 15.3 | 30.6 | 68.2 | 160.9 | 451.6 |
| 1999..... | 33.8 | 7.2 | 0.8 | 0.3 | 0.4 | 1.8 | 7.2 | 14.6 | 30.2 | 69.1 | 158.7 | 445.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XI. Diseases of the digestive system (K00-K92) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 24.5 | 32.7 | 1.4 | 0.4 | 1.2 | 3.0 | 13.0 | 38.5 | 62.1 | 90.4 | 160.3 | 333.9 |
| 2003..... | 25.0 | 29.7 | 1.3 | 0.4 | 1.1 | 3.5 | 13.6 | 41.1 | 64.3 | 93.4 | 159.4 | 351.6 |
| 2002..... | 24.4 | 26.3 | 1.5 | 0.5 | 1.1 | 3.5 | 13.2 | 41.9 | 59.9 | 90.6 | 166.4 | 336.6 |
| 2001..... | 25.9 | 29.8 | 1.3 | 0.5 | 1.1 | 3.6 | 15.5 | 43.8 | 66.1 | 98.1 | 172.0 | 354.3 |
| 2000..... | 26.1 | 30.5 | 1.6 | 0.4 | 1.1 | 4.0 | 16.4 | 44.2 | 65.7 | 97.9 | 171.8 | 385.7 |
| 1999..... | 25.8 | 30.6 | 1.1 | 0.5 | 1.3 | 3.5 | 16.5 | 43.5 | 66.9 | 101.3 | 173.3 | 366.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 27.4 | 37.4 | 1.7 | * | 1.1 | 3.9 | 16.0 | 53.7 | 88.6 | 118.4 | 190.2 | 329.2 |
| 2003..... | 28.0 | 33.5 | * | * | 1.1 | 4.2 | 16.2 | 58.0 | 91.6 | 121.7 | 193.1 | 352.7 |
| 2002..... | 26.8 | 32.4 | 1.7 | * | 0.9 | 3.9 | 15.4 | 58.7 | 83.2 | 114.9 | 189.9 | 359.9 |
| 2001..... | 29.3 | 35.1 | * | * | 1.0 | 4.5 | 20.5 | 63.1 | 92.1 | 128.9 | 196.8 | 391.5 |
| 2000..... | 28.6 | 34.5 | 2.2 | * | 1.4 | 4.2 | 21.1 | 61.6 | 89.7 | 123.7 | 203.3 | 396.9 |
| 1999..... | 28.6 | 37.8 | * | * | 1.2 | 4.2 | 21.3 | 64.1 | 90.3 | 123.1 | 197.0 | 402.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 21.9 | 27.7 | * | * | 1.3 | 2.2 | 10.4 | 25.5 | 40.9 | 70.5 | 143.5 | 335.7 |
| 2003..... | 22.3 | 25.8 | * | * | 1.1 | 2.9 | 11.3 | 26.6 | 42.4 | 73.2 | 140.5 | 351.2 |
| 2002..... | 22.2 | 19.9 | * | * | 1.3 | 3.1 | 11.1 | 27.5 | 41.2 | 73.3 | 153.0 | 327.8 |
| 2001..... | 22.8 | 24.2 | * | * | 1.2 | 2.8 | 11.0 | 27.3 | 45.1 | 76.4 | 158.0 | 340.5 |
| 2000..... | 23.7 | 26.4 | * | * | 0.9 | 3.7 | 12.2 | 29.3 | 46.4 | 79.8 | 154.0 | 381.6 |
| 1999..... | 23.4 | 23.2 | * | * | 1.4 | 2.9 | 12.4 | 26.0 | 48.2 | 85.9 | 160.0 | 353.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of other parts of digestive system (K20-K92) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 29.3 | 14.1 | 0.8 | 0.2 | 0.7 | 2.1 | 10.9 | 29.5 | 44.4 | 78.3 | 162.0 | 385.7 |
| 2003..... | 30.1 | 13.7 | 0.8 | 0.3 | 0.6 | 2.4 | 11.5 | 30.2 | 45.4 | 82.4 | 166.9 | 402.6 |
| 2002..... | 30.0 | 12.5 | 0.8 | 0.3 | 0.6 | 2.3 | 11.7 | 29.8 | 44.6 | 81.6 | 169.7 | 413.8 |
| 2001..... | 30.1 | 11.8 | 0.8 | 0.3 | 0.5 | 2.4 | 11.7 | 29.7 | 45.2 | 83.8 | 168.3 | 430.2 |
| 2000..... | 29.8 | 13.0 | 0.7 | 0.3 | 0.7 | 2.4 | 11.9 | 28.6 | 45.6 | 82.1 | 170.1 | 439.5 |
| 1999..... | 29.7 | 13.1 | 0.7 | 0.3 | 0.6 | 2.4 | 11.8 | 27.9 | 45.8 | 85.0 | 168.8 | 435.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 29.9 | 15.5 | 0.9 | 0.2 | 0.7 | 2.7 | 14.1 | 41.3 | 58.5 | 91.5 | 175.6 | 365.5 |
| 2003..... | 30.9 | 16.5 | 0.9 | 0.3 | 0.7 | 2.9 | 14.6 | 42.6 | 60.5 | 97.5 | 181.4 | 388.0 |
| 2002..... | 30.6 | 15.0 | 0.9 | 0.3 | 0.7 | 2.9 | 15.0 | 42.4 | 58.9 | 96.7 | 183.8 | 403.1 |
| 2001..... | 30.5 | 13.3 | 0.9 | 0.3 | 0.6 | 2.8 | 15.2 | 41.8 | 59.7 | 99.5 | 181.4 | 427.0 |
| 2000..... | 30.3 | 14.8 | 0.9 | 0.3 | 0.7 | 3.0 | 15.8 | 40.6 | 60.1 | 97.6 | 186.2 | 440.8 |
| 1999..... | 30.3 | 16.4 | 0.7 | 0.3 | 0.6 | 2.9 | 15.8 | 40.4 | 60.6 | 102.2 | 185.6 | 437.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 28.7 | 12.6 | 0.6 | 0.3 | 0.7 | 1.5 | 7.7 | 18.1 | 31.3 | 67.2 | 152.9 | 394.7 |
| 2003..... | 29.4 | 10.8 | 0.7 | 0.3 | 0.5 | 1.8 | 8.3 | 18.2 | 31.4 | 69.8 | 157.2 | 409.1 |
| 2002..... | 29.4 | 9.9 | 0.7 | 0.3 | 0.6 | 1.8 | 8.5 | 17.6 | 31.4 | 68.9 | 160.3 | 418.4 |
| 2001..... | 29.7 | 10.2 | 0.6 | 0.3 | 0.5 | 1.8 | 8.3 | 18.0 | 31.8 | 70.8 | 159.8 | 431.5 |
| 2000..... | 29.4 | 11.2 | 0.5 | 0.3 | 0.5 | 1.8 | 8.1 | 16.9 | 32.3 | 69.3 | 159.7 | 439.0 |
| 1999..... | 29.0 | 9.7 | 0.8 | 0.4 | 0.6 | 1.9 | 7.8 | 15.9 | 32.2 | 71.0 | 157.8 | 434.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 15.6 | 12.1 | 0.6 | * | 0.6 | 1.9 | 11.0 | 34.5 | 53.7 | 89.6 | 165.5 | 271.1 |
| 2003..... | 16.5 | 12.1 | 0.8 | 0.4 | 0.5 | 2.5 | 11.7 | 35.3 | 59.9 | 98.5 | 179.1 | 292.3 |
| 2002..... | 16.2 | 11.8 | 0.9 | 0.3 | 0.7 | 2.2 | 12.0 | 36.3 | 57.9 | 102.6 | 168.1 | 289.9 |
| 2001..... | 16.3 | 10.8 | 0.8 | * | 0.5 | 2.1 | 12.3 | 36.8 | 58.3 | 102.0 | 178.0 | 363.3 |
| 2000..... | 16.4 | 11.4 | 0.7 | * | 0.6 | 2.4 | 13.4 | 37.3 | 59.5 | 102.9 | 183.1 | 365.6 |
| 1999..... | 16.7 | 12.0 | 0.7 | * | 0.7 | 2.7 | 14.3 | 37.1 | 60.5 | 106.3 | 172.3 | 411.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 18.4 | 12.3 | * | * | 0.6 | 2.8 | 16.4 | 55.1 | 75.8 | 103.4 | 175.3 | 262.0 |
| 2003..... | 19.6 | 15.2 | * | * | 0.6 | 3.4 | 17.0 | 55.4 | 88.3 | 115.7 | 198.6 | 274.0 |
| 2002..... | 19.8 | 12.9 | * | * | 0.7 | 3.1 | 18.0 | 57.5 | 85.1 | 133.2 | 194.9 | 284.4 |
| 2001..... | 19.7 | 12.7 | * | * | * | 2.8 | 18.2 | 58.4 | 87.7 | 128.2 | 203.9 | 369.5 |
| 2000..... | 20.1 | 12.4 | * | * | 0.7 | 3.5 | 20.3 | 58.7 | 90.2 | 134.8 | 208.5 | 387.0 |
| 1999..... | 20.2 | 15.8 | * | * | 0.7 | 3.5 | 21.9 | 58.8 | 89.8 | 138.0 | 196.0 | 427.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 12.6 | 12.0 | * | * | * | 0.9 | 5.1 | 13.8 | 33.6 | 78.5 | 158.8 | 275.9 |
| 2003..... | 13.2 | 9.0 | * | * | * | 1.5 | 5.8 | 15.1 | 34.3 | 84.7 | 165.6 | 301.6 |
| 2002..... | 12.3 | 10.6 | * | * | * | 1.2 | 5.5 | 15.3 | 33.6 | 78.2 | 149.8 | 292.6 |
| 2001..... | 12.7 | 8.7 | * | * | * | 1.3 | 5.9 | 15.5 | 32.1 | 81.4 | 160.5 | 360.2 |
| 2000..... | 12.5 | 10.4 | * | * | * | 1.0 | 6.1 | 16.3 | 32.2 | 77.8 | 166.1 | 355.1 |
| 1999..... | 12.9 | 8.1 | * | * | * | 1.6 | 6.2 | 15.9 | 34.6 | 81.6 | 156.5 | 403.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of other parts of digestive system (K20-K92) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 31.5 | 14.6 | 0.8 | 0.2 | 0.7 | 2.2 | 10.8 | 28.8 | 43.5 | 77.2 | 161.6 | 390.1 |
| 2003..... | 32.2 | 14.0 | 0.8 | 0.3 | 0.6 | 2.3 | 11.4 | 29.5 | 44.0 | 81.1 | 165.9 | 406.6 |
| 2002..... | 32.1 | 12.5 | 0.8 | 0.3 | 0.6 | 2.3 | 11.6 | 29.0 | 43.3 | 79.8 | 169.3 | 418.0 |
| 2001..... | 32.0 | 11.9 | 0.8 | 0.3 | 0.5 | 2.4 | 11.6 | 28.9 | 44.0 | 82.3 | 167.5 | 431.9 |
| 2000..... | 31.6 | 13.2 | 0.7 | 0.3 | 0.7 | 2.4 | 11.6 | 27.6 | 44.4 | 80.4 | 169.2 | 441.3 |
| 1999..... | 31.3 | 13.2 | 0.7 | 0.3 | 0.6 | 2.4 | 11.4 | 27.0 | 44.6 | 83.5 | 168.2 | 435.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 31.8 | 16.3 | 1.0 | 0.2 | 0.7 | 2.7 | 13.6 | 39.5 | 56.9 | 90.3 | 175.4 | 370.0 |
| 2003..... | 32.7 | 16.7 | 0.9 | 0.3 | 0.7 | 2.8 | 14.1 | 41.0 | 58.0 | 96.0 | 180.1 | 392.5 |
| 2002..... | 32.2 | 15.5 | 0.9 | 0.3 | 0.6 | 2.8 | 14.4 | 40.7 | 56.4 | 93.9 | 182.7 | 407.5 |
| 2001..... | 32.1 | 13.4 | 0.9 | 0.3 | 0.6 | 2.9 | 14.8 | 40.1 | 57.2 | 97.2 | 179.9 | 428.8 |
| 2000..... | 31.7 | 15.1 | 1.0 | 0.3 | 0.8 | 2.8 | 15.1 | 38.8 | 57.5 | 94.8 | 184.8 | 442.3 |
| 1999..... | 31.7 | 16.3 | 0.7 | 0.3 | 0.6 | 2.8 | 14.9 | 38.5 | 58.3 | 99.8 | 184.7 | 436.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 31.2 | 12.7 | 0.6 | 0.3 | 0.7 | 1.6 | 8.0 | 18.5 | 31.0 | 66.2 | 152.3 | 399.1 |
| 2003..... | 31.7 | 11.1 | 0.8 | 0.2 | 0.5 | 1.9 | 8.7 | 18.4 | 31.0 | 68.6 | 156.4 | 412.7 |
| 2002..... | 31.9 | 9.5 | 0.7 | 0.3 | 0.6 | 1.9 | 8.9 | 17.8 | 31.1 | 68.1 | 160.5 | 422.6 |
| 2001..... | 32.0 | 10.3 | 0.6 | 0.3 | 0.5 | 1.9 | 8.5 | 18.2 | 31.7 | 69.9 | 159.3 | 433.1 |
| 2000..... | 31.6 | 11.3 | 0.4 | 0.3 | 0.5 | 1.9 | 8.3 | 16.9 | 32.2 | 68.5 | 159.0 | 440.9 |
| 1999..... | 31.0 | 10.0 | 0.8 | 0.4 | 0.6 | 1.9 | 7.9 | 15.8 | 31.9 | 70.2 | 157.5 | 434.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 34.2 | 9.8 | 0.6 | 0.2 | 0.5 | 2.0 | 10.6 | 28.1 | 42.0 | 77.2 | 163.7 | 399.1 |
| 2003..... | 34.9 | 10.4 | 0.7 | 0.2 | 0.5 | 2.1 | 11.2 | 28.4 | 42.4 | 81.2 | 168.4 | 415.3 |
| 2002..... | 34.8 | 9.1 | 0.6 | 0.2 | 0.5 | 2.1 | 11.5 | 27.8 | 42.0 | 79.9 | 171.5 | 429.4 |
| 2001..... | 34.5 | 7.6 | 0.6 | 0.3 | 0.4 | 2.2 | 11.1 | 27.4 | 42.0 | 82.0 | 168.9 | 441.9 |
| 2000..... | 33.9 | 9.1 | 0.4 | 0.3 | 0.6 | 2.0 | 11.0 | 25.8 | 42.4 | 79.7 | 170.3 | 449.8 |
| 1999..... | 33.5 | 9.1 | 0.6 | 0.3 | 0.5 | 2.1 | 10.6 | 25.2 | 42.5 | 82.6 | 169.1 | 443.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 33.9 | 10.9 | 0.7 | 0.2 | 0.5 | 2.5 | 13.4 | 38.6 | 54.5 | 89.3 | 176.1 | 378.2 |
| 2003..... | 34.8 | 13.0 | 0.8 | 0.3 | 0.7 | 2.5 | 14.0 | 39.6 | 55.4 | 95.1 | 181.0 | 399.5 |
| 2002..... | 34.4 | 11.4 | 0.6 | 0.3 | 0.6 | 2.5 | 14.4 | 39.1 | 54.4 | 93.3 | 183.9 | 415.9 |
| 2001..... | 33.8 | 7.8 | 0.7 | 0.3 | 0.5 | 2.6 | 13.9 | 38.0 | 54.2 | 95.9 | 180.4 | 434.9 |
| 2000..... | 33.4 | 10.5 | 0.6 | 0.3 | 0.6 | 2.5 | 14.3 | 36.5 | 54.9 | 93.5 | 185.0 | 448.2 |
| 1999..... | 33.3 | 10.8 | 0.5 | 0.3 | 0.5 | 2.5 | 14.1 | 36.0 | 55.6 | 98.6 | 185.5 | 440.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 34.4 | 8.6 | 0.5 | 0.2 | 0.5 | 1.5 | 7.7 | 17.8 | 30.2 | 66.7 | 155.3 | 408.5 |
| 2003..... | 35.0 | 7.6 | 0.6 | 0.2 | 0.4 | 1.7 | 8.3 | 17.4 | 30.1 | 69.3 | 159.8 | 422.2 |
| 2002..... | 35.2 | 6.7 | 0.6 | 0.2 | 0.5 | 1.6 | 8.6 | 16.7 | 30.2 | 68.5 | 163.2 | 435.2 |
| 2001..... | 35.1 | 7.4 | 0.5 | 0.3 | 0.3 | 1.8 | 8.2 | 17.0 | 30.5 | 70.3 | 161.3 | 444.9 |
| 2000..... | 34.3 | 7.7 | * | 0.3 | 0.5 | 1.5 | 7.7 | 15.3 | 30.6 | 68.1 | 160.6 | 450.4 |
| 1999..... | 33.7 | 7.2 | 0.8 | 0.3 | 0.4 | 1.8 | 7.2 | 14.5 | 30.2 | 69.1 | 158.4 | 444.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of other parts of digestive system (K20-K92) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 24.4 | 32.7 | 1.4 | 0.4 | 1.2 | 2.9 | 13.0 | 38.4 | 62.1 | 90.2 | 160.1 | 333.6 |
| 2003..... | 25.0 | 29.6 | 1.3 | 0.4 | 1.1 | 3.5 | 13.6 | 41.1 | 64.2 | 93.2 | 159.4 | 351.6 |
| 2002..... | 24.4 | 26.3 | 1.5 | 0.5 | 1.1 | 3.4 | 13.1 | 41.9 | 59.9 | 90.5 | 166.1 | 335.7 |
| 2001..... | 25.8 | 29.8 | 1.3 | 0.5 | 1.1 | 3.6 | 15.5 | 43.8 | 66.1 | 98.1 | 171.1 | 353.3 |
| 2000..... | 26.0 | 30.3 | 1.6 | 0.4 | 1.1 | 3.9 | 16.3 | 44.2 | 65.7 | 97.8 | 171.5 | 385.4 |
| 1999..... | 25.8 | 30.6 | 1.1 | 0.5 | 1.3 | 3.5 | 16.5 | 43.5 | 66.9 | 101.2 | 172.8 | 366.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 27.3 | 37.4 | 1.7 | * | 1.1 | 3.8 | 16.0 | 53.5 | 88.6 | 118.1 | 190.0 | 329.2 |
| 2003..... | 28.0 | 33.5 | * | * | 1.0 | 4.2 | 16.2 | 58.0 | 91.5 | 121.6 | 193.1 | 352.7 |
| 2002..... | 26.7 | 32.4 | 1.7 | * | 0.9 | 3.9 | 15.4 | 58.6 | 83.2 | 114.8 | 189.6 | 359.9 |
| 2001..... | 29.2 | 35.1 | * | * | 1.0 | 4.5 | 20.5 | 63.1 | 92.1 | 128.9 | 195.6 | 390.4 |
| 2000..... | 28.6 | 34.5 | 2.2 | * | 1.4 | 4.2 | 21.1 | 61.6 | 89.7 | 123.5 | 203.3 | 396.9 |
| 1999..... | 28.5 | 37.8 | * | * | 1.2 | 4.2 | 21.2 | 64.1 | 90.2 | 123.1 | 196.7 | 402.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 21.8 | 27.7 | * | * | 1.3 | 2.1 | 10.3 | 25.4 | 40.8 | 70.2 | 143.3 | 335.3 |
| 2003..... | 22.3 | 25.5 | * | * | 1.1 | 2.9 | 11.3 | 26.6 | 42.3 | 73.1 | 140.5 | 351.2 |
| 2002..... | 22.2 | 19.9 | * | * | 1.3 | 3.1 | 11.1 | 27.5 | 41.2 | 73.3 | 152.9 | 326.5 |
| 2001..... | 22.7 | 24.2 | * | * | 1.2 | 2.8 | 11.0 | 27.3 | 45.1 | 76.4 | 157.3 | 339.6 |
| 2000..... | 23.7 | 26.1 | * | * | 0.9 | 3.7 | 12.2 | 29.3 | 46.4 | 79.7 | 153.5 | 381.1 |
| 1999..... | 23.3 | 23.2 | * | * | 1.4 | 2.9 | 12.3 | 26.0 | 48.2 | 85.8 | 159.5 | 352.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of esophagus (K20-K22) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 4.2 | 18.2 |
| 2003..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 4.5 | 19.5 |
| 2002..... | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.3 | 4.9 | 21.1 |
| 2001..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 5.2 | 21.5 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 5.2 | 23.7 |
| 1999..... | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 5.1 | 23.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 5.1 | 19.4 |
| 2003..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.6 | 5.0 | 23.5 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.4 | 0.4 | 1.6 | 5.8 | 25.7 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 6.0 | 24.1 |
| 2000..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.8 | 6.4 | 28.7 |
| 1999..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.0 | 6.6 | 28.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.6 | 17.7 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 4.1 | 17.8 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 4.4 | 19.1 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | 0.3 | 1.2 | 4.7 | 20.5 |
| 2000..... | 0.9 | * | * | * | * | * | * | 0.1 | 0.3 | 1.2 | 4.5 | 21.6 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.1 | 21.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 4.4 | 19.2 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 4.6 | 20.4 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.3 | 5.0 | 22.1 |
| 2001..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 5.3 | 22.6 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 5.4 | 24.8 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 5.3 | 24.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 5.2 | 20.4 |
| 2003..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.7 | 5.2 | 24.8 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.4 | 1.6 | 5.8 | 27.1 |
| 2001..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.7 | 6.0 | 25.3 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.8 | 6.6 | 30.1 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.0 | 6.7 | 29.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | 0.3 | 1.1 | 3.8 | 18.6 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 4.3 | 18.5 |
| 2002..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.5 | 20.0 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | 0.4 | 1.2 | 4.9 | 21.4 |
| 2000..... | 1.1 | * | * | * | * | * | * | * | 0.3 | 1.2 | 4.7 | 22.7 |
| 1999..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.4 | 22.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Peptic ulcer (K25-K28) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 0.7 | 1.3 | 3.2 | 8.9 | 24.6 |
| 2003..... | 1.5 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.4 | 3.2 | 9.9 | 26.7 |
| 2002..... | 1.6 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.7 | 10.2 | 28.6 |
| 2001..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 4.0 | 11.5 | 33.6 |
| 2000..... | 1.8 | * | * | * | * | * | 0.3 | 0.8 | 1.8 | 4.1 | 11.5 | 35.7 |
| 1999..... | 1.8 | * | * | * | * | * | 0.3 | 0.7 | 1.8 | 4.6 | 12.4 | 34.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.3 | 0.9 | 1.6 | 3.7 | 10.2 | 24.8 |
| 2003..... | 1.4 | * | * | * | * | * | 0.3 | 0.9 | 1.8 | 4.0 | 11.7 | 29.6 |
| 2002..... | 1.5 | * | * | * | * | * | 0.4 | 0.9 | 1.9 | 4.5 | 12.3 | 31.7 |
| 2001..... | 1.6 | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.2 | 4.8 | 12.8 | 35.5 |
| 2000..... | 1.8 | * | * | * | * | * | 0.4 | 1.2 | 2.4 | 5.4 | 13.9 | 41.0 |
| 1999..... | 1.7 | * | * | * | * | * | 0.4 | 1.0 | 2.3 | 6.1 | 14.8 | 37.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.8 | 8.1 | 24.6 |
| 2003..... | 1.5 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.5 | 8.7 | 25.4 |
| 2002..... | 1.6 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.0 | 8.8 | 27.3 |
| 2001..... | 1.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.3 | 10.6 | 32.8 |
| 2000..... | 1.8 | * | * | * | * | * | 0.1 | 0.5 | 1.3 | 3.1 | 9.9 | 33.5 |
| 1999..... | 1.9 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.3 | 10.8 | 33.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 3.1 | 9.0 | 25.4 |
| 2003..... | 1.6 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.2 | 9.8 | 27.3 |
| 2002..... | 1.7 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 3.6 | 10.4 | 29.4 |
| 2001..... | 1.9 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 3.9 | 11.6 | 34.7 |
| 2000..... | 1.9 | * | * | * | * | * | 0.2 | 0.7 | 1.8 | 4.0 | 11.6 | 36.3 |
| 1999..... | 2.0 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 4.5 | 12.4 | 35.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 0.2 | 0.8 | 1.5 | 3.6 | 10.0 | 25.7 |
| 2003..... | 1.5 | * | * | * | * | * | 0.2 | 0.8 | 1.6 | 4.0 | 11.3 | 29.8 |
| 2002..... | 1.6 | * | * | * | * | * | 0.3 | 0.8 | 1.7 | 4.4 | 12.4 | 32.6 |
| 2001..... | 1.7 | * | * | * | * | * | 0.3 | 0.8 | 2.0 | 4.5 | 12.7 | 36.3 |
| 2000..... | 1.9 | * | * | * | * | * | 0.3 | 1.1 | 2.3 | 5.2 | 13.7 | 41.0 |
| 1999..... | 1.8 | * | * | * | * | * | 0.3 | 0.8 | 2.1 | 5.9 | 14.9 | 38.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.7 | 8.3 | 25.3 |
| 2003..... | 1.7 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.6 | 8.8 | 26.2 |
| 2002..... | 1.8 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.0 | 9.0 | 27.9 |
| 2001..... | 2.1 | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 3.4 | 10.9 | 34.0 |
| 2000..... | 2.0 | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.1 | 10.2 | 34.4 |
| 1999..... | 2.1 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.2 | 10.8 | 34.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Peptic ulcer (K25-K28) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.5 | 1.3 | 2.1 | 4.3 | 7.8 | 16.1 |
| 2003..... | 1.0 | * | * | * | * | * | 0.4 | 1.5 | 2.2 | 3.2 | 11.1 | 17.3 |
| 2002..... | 1.0 | * | * | * | * | * | 0.6 | 1.2 | 2.6 | 4.2 | 7.9 | 18.8 |
| 2001..... | 1.1 | * | * | * | * | * | 0.6 | 1.4 | 2.8 | 4.5 | 9.0 | 20.9 |
| 2000..... | 1.2 | * | * | * | * | * | 0.5 | 1.7 | 2.6 | 4.9 | 10.1 | 26.7 |
| 1999..... | 1.3 | * | * | * | * | * | 0.8 | 1.5 | 2.8 | 5.7 | 11.0 | 25.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 1.8 | 3.0 | 4.7 | 11.1 | * |
| 2003..... | 1.3 | * | * | * | * | * | * | 2.3 | 3.4 | 5.2 | 16.4 | * |
| 2002..... | 1.2 | * | * | * | * | * | 0.8 | 2.0 | 3.9 | 6.0 | 11.1 | * |
| 2001..... | 1.4 | * | * | * | * | * | 1.0 | 2.3 | 4.3 | 7.0 | 11.7 | 24.3 |
| 2000..... | 1.4 | * | * | * | * | * | 0.8 | 2.4 | 3.8 | 7.2 | 13.9 | 31.7 |
| 1999..... | 1.5 | * | * | * | * | * | 1.2 | 2.4 | 4.5 | 8.3 | 12.2 | 28.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.9 | 1.3 | 4.1 | 6.0 | 16.9 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | * | * | 8.1 | 16.1 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | 1.6 | 2.9 | 6.1 | 18.4 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | 1.5 | 2.7 | 7.4 | 19.7 |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.1 | 1.7 | 3.3 | 8.0 | 24.9 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 3.9 | 10.3 | 24.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Gastric ulcer (K25) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.0 | 2.7 | 6.8 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 2.7 | 7.2 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.9 | 7.7 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 3.3 | 8.2 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 3.0 | 9.4 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 3.5 | 8.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.9 | 6.3 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.8 | 6.7 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 2.9 | 7.9 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 3.3 | 6.3 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.6 | 3.3 | 9.6 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.5 | 3.7 | 8.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.5 | 6.9 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.7 | 7.4 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 7.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.3 | 9.0 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.8 | 9.3 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.4 | 8.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 6.9 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.6 | 7.2 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 2.8 | 7.7 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.3 | 8.2 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 3.0 | 9.3 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.5 | 8.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.4 | 0.9 | 2.7 | 6.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 0.9 | 2.6 | 6.4 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.8 | 8.0 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.3 | 6.1 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.4 | 0.5 | 1.4 | 3.2 | 8.9 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 3.6 | 8.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.6 | 7.0 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.6 | 7.5 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.7 | 7.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.2 | 9.0 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 2.9 | 9.4 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.5 | 8.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Duodenal ulcer (K26) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.1 | 7.8 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.1 | 3.4 | 8.7 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.7 | 9.1 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.0 | 11.0 |
| 2000..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 4.2 | 10.2 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 3.8 | 10.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.5 | 3.6 | 9.2 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.6 | 4.6 | 10.5 |
| 2002..... | 0.6 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.7 | 4.7 | 10.3 |
| 2001..... | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.8 | 4.7 | 13.8 |
| 2000..... | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 4.9 | 12.5 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 5.1 | 12.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 7.2 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.6 | 7.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.0 | 8.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.5 | 9.8 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 3.7 | 9.2 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.9 | 9.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.2 | 8.2 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.2 | 3.4 | 9.0 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.8 | 9.7 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 4.1 | 11.7 |
| 2000..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.4 | 4.3 | 10.6 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.6 | 3.9 | 10.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.6 | 1.5 | 3.7 | 9.5 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.5 | 1.6 | 4.6 | 10.5 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.7 | 4.8 | 11.1 |
| 2001..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.8 | 4.8 | 14.6 |
| 2000..... | 0.7 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 1.8 | 5.0 | 12.9 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.3 | 1.0 | 2.4 | 5.3 | 12.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.9 | 7.7 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.6 | 8.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.2 | 9.1 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.7 | 10.5 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 3.8 | 9.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 3.0 | 9.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Peptic ulcer, site unspecified (K27) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.0 | 9.8 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 3.6 | 10.5 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.6 | 11.5 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 4.1 | 14.2 |
| 2000..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.2 | 15.7 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.7 | 4.9 | 15.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.5 | 9.0 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 4.2 | 12.0 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.4 | 1.4 | 4.5 | 13.0 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.8 | 1.6 | 4.6 | 15.2 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.8 | 1.8 | 5.4 | 18.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.8 | 16.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.7 | 10.1 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.8 | 3.2 | 9.9 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.9 | 10.8 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.8 | 13.8 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.4 | 14.5 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 0.4 | 1.2 | 4.3 | 15.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.0 | 10.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.6 | 10.8 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.6 | 11.7 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.1 | 14.6 |
| 2000..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.2 | 16.0 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 4.9 | 16.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.4 | 9.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.9 | 12.4 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.4 | 1.4 | 4.6 | 13.1 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 4.5 | 15.4 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.8 | 1.8 | 5.4 | 18.8 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 5.8 | 17.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.7 | 10.4 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 0.9 | 3.4 | 10.1 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.0 | 11.0 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.8 | 14.3 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | 0.2 | 1.0 | 3.4 | 14.9 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | 0.4 | 1.2 | 4.3 | 16.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Peptic ulcer, site unspecified (K27) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.7 | 1.2 | 2.7 | 8.3 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | 3.7 | 5.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.8 | 1.5 | 2.4 | 9.1 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 1.1 | 1.3 | 3.9 | 9.7 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 1.2 | 3.8 | 12.1 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.9 | 2.2 | 4.5 | 9.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | 7.5 | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 1.9 | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 3.3 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | 8.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | 8.4 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | 8.6 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | 3.6 | 11.3 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | 4.2 | 10.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Gastritis and duodenitis (K29) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.2 | 0.8 | 2.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 2.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.8 | 3.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 3.7 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 3.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 3.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.1 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.9 | 2.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 4.6 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.4 | 0.8 | 3.3 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.2 | * | 0.3 | 1.1 | 3.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.4 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.6 | 3.0 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 3.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 3.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 3.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 3.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 2.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 2.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.8 | 3.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 4.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.7 | 3.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 3.2 |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.2 | * | * | 0.8 | 2.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.0 | 3.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 4.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 3.3 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.2 | * | * | 0.9 | 3.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 2.9 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.6 | 2.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.7 | 3.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.9 | 3.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.6 | 4.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Dyspepsia and other diseases of stomach and duodenum (K30-K31) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.4 | 3.1 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.4 | 3.6 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.4 | 3.3 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 3.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 4.1 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.4 | 4.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 3.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 2.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.3 | 2.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.4 | 3.2 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.8 | 1.5 | 4.3 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.3 | 2.8 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.9 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 3.5 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.5 | 1.2 | 3.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.3 | 3.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 4.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 1.4 | 3.1 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 3.6 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.3 | 3.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.2 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.4 | 4.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.7 | 1.5 | 3.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 3.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.7 | 1.5 | 4.4 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 4.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.4 | 2.9 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.3 | 3.6 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | * | 0.5 | 1.2 | 3.9 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.3 | 4.1 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 4.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of appendix (K35-K38) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 1.8 |
| 2003..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.0 | 1.6 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 0.7 | 2.2 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.0 | 1.7 |
| 1999..... | 0.1 | * | * | * | * | * | 0.0 | 0.1 | 0.2 | 0.4 | 0.8 | 1.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 2.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 2.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.3 | 2.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.0 | 3.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 1.2 | 2.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.0 | 3.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.5 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 1.4 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.5 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 1.5 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 0.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 1.8 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 1.7 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 0.7 | 2.1 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 1.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 2.3 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 2.4 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.0 | 2.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.3 | 0.4 | 1.2 | 2.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 3.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | 1.6 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 1.3 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.4 | 2.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.6 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Hernia (K40-K46) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | 0.9 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.9 | 11.2 |
| 2003..... | 0.6 | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.6 | 12.1 |
| 2002..... | 0.6 | 0.8 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.8 | 13.0 |
| 2001..... | 0.6 | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.5 | 12.5 |
| 2000..... | 0.6 | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.7 | 13.1 |
| 1999..... | 0.6 | 1.3 | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.1 | 3.7 | 12.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 3.9 | 11.6 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 3.9 | 12.5 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.7 | 11.9 |
| 2001..... | 0.5 | 1.2 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 12.1 |
| 2000..... | 0.5 | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.9 | 12.8 |
| 1999..... | 0.5 | 1.5 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 4.0 | 13.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 4.0 | 11.0 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 3.3 | 12.0 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.9 | 13.4 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 12.7 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.6 | 13.2 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.5 | 12.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 4.1 | 11.4 |
| 2003..... | 0.7 | 1.1 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.6 | 12.4 |
| 2002..... | 0.7 | 0.9 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 4.0 | 13.6 |
| 2001..... | 0.6 | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 12.8 |
| 2000..... | 0.6 | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.7 | 13.3 |
| 1999..... | 0.6 | 1.4 | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.1 | 3.7 | 13.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.0 | 11.7 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 3.9 | 12.6 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.8 | 12.2 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.5 | 11.9 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 4.0 | 13.0 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 4.1 | 13.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 4.1 | 11.3 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 3.3 | 12.4 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 4.1 | 14.2 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.5 | 13.1 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.6 | 13.4 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 0.4 | 1.1 | 3.4 | 12.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Crohn's disease and ulcerative colitis (K50-K51) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 2.1 | 3.5 |
| 2003..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.0 | 2.0 | 3.1 |
| 2002..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.9 | 2.0 | 3.2 |
| 2001..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 0.9 | 2.0 | 3.8 |
| 2000..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.9 | 1.9 | 3.4 |
| 1999..... | 0.3 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 0.8 | 2.0 | 3.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.9 | 2.1 | 2.9 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.1 | 3.7 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 2.0 | 2.9 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.9 | 3.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.7 | 3.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.0 | 2.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.9 | 2.2 | 3.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.4 | 1.0 | 1.9 | 2.8 |
| 2002..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 2.0 | 3.3 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 2.0 | 3.9 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.0 | 3.5 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.8 | 2.1 | 3.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 2.3 | 3.8 |
| 2003..... | 0.4 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.0 | 2.1 | 3.3 |
| 2002..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.0 | 2.2 | 3.5 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.1 | 3.9 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.0 | 3.6 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.9 | 2.1 | 3.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.0 | 2.2 | 3.0 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.2 | 3.9 |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.0 | 2.1 | 3.1 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.1 | 3.6 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 3.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.1 | 3.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.5 | 0.9 | 2.4 | 4.1 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.1 | 3.0 |
| 2002..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.0 | 2.2 | 3.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.1 | 4.1 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.2 | 3.7 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.4 | 0.9 | 2.2 | 3.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Vascular disorders and obstruction of intestine without hernia (K55-K56) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | 6.9 | 0.2 | 0.0 | 0.1 | 0.1 | 0.4 | 1.1 | 3.5 | 12.1 | 38.2 | 106.9 |
| 2003..... | 5.1 | 6.8 | 0.1 | 0.1 | 0.1 | 0.1 | 0.4 | 1.2 | 3.7 | 13.3 | 39.2 | 111.5 |
| 2002..... | 5.0 | 5.8 | 0.1 | 0.1 | 0.1 | 0.1 | 0.4 | 1.1 | 3.7 | 12.8 | 39.4 | 113.3 |
| 2001..... | 5.0 | 5.8 | 0.1 | 0.1 | * | 0.1 | 0.3 | 1.2 | 3.9 | 13.8 | 38.9 | 113.5 |
| 2000..... | 5.1 | 6.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.4 | 1.3 | 3.9 | 13.4 | 40.1 | 116.2 |
| 1999..... | 5.0 | 6.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.4 | 1.1 | 4.0 | 13.5 | 38.9 | 116.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.5 | 7.7 | 0.2 | * | * | 0.1 | 0.5 | 1.1 | 3.3 | 10.9 | 35.0 | 88.8 |
| 2003..... | 3.7 | 8.2 | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.2 | 3.7 | 12.2 | 35.6 | 92.6 |
| 2002..... | 3.6 | 6.5 | * | * | 0.1 | 0.1 | 0.4 | 1.1 | 3.4 | 11.6 | 35.8 | 96.7 |
| 2001..... | 3.6 | 6.5 | * | * | * | 0.1 | 0.3 | 1.1 | 3.5 | 13.1 | 35.2 | 98.5 |
| 2000..... | 3.6 | 6.3 | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.2 | 3.8 | 12.5 | 36.8 | 98.5 |
| 1999..... | 3.5 | 7.6 | * | * | * | 0.1 | 0.4 | 1.1 | 3.8 | 12.5 | 35.6 | 95.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.3 | 6.0 | * | * | * | 0.2 | 0.4 | 1.1 | 3.7 | 13.1 | 40.5 | 115.1 |
| 2003..... | 6.5 | 5.3 | * | * | * | 0.1 | 0.4 | 1.3 | 3.7 | 14.2 | 41.6 | 119.8 |
| 2002..... | 6.4 | 5.0 | * | * | * | 0.1 | 0.3 | 1.1 | 3.8 | 13.8 | 41.7 | 120.5 |
| 2001..... | 6.4 | 5.1 | * | * | * | 0.1 | 0.3 | 1.2 | 4.1 | 14.4 | 41.3 | 119.8 |
| 2000..... | 6.5 | 5.8 | * | * | * | 0.1 | 0.4 | 1.3 | 4.0 | 14.0 | 42.1 | 123.4 |
| 1999..... | 6.4 | 4.4 | * | * | * | 0.2 | 0.4 | 1.1 | 4.2 | 14.3 | 41.1 | 124.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | 5.8 | * | * | * | * | * | 0.6 | 2.3 | 7.6 | 27.8 | 73.7 |
| 2003..... | 1.6 | 6.4 | * | * | * | * | * | 0.8 | 2.4 | 10.0 | 29.8 | 73.1 |
| 2002..... | 1.5 | 5.3 | * | * | * | * | * | 0.8 | 2.1 | 10.8 | 31.4 | 75.9 |
| 2001..... | 1.5 | 5.1 | * | * | * | * | * | 0.6 | 3.0 | 10.0 | 29.8 | 95.7 |
| 2000..... | 1.5 | 5.1 | * | * | * | * | * | 0.8 | 2.3 | 9.8 | 30.4 | 88.3 |
| 1999..... | 1.5 | 6.1 | * | * | * | * | * | 0.8 | 2.9 | 9.7 | 29.8 | 92.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.1 | 5.4 | * | * | * | * | * | * | 2.1 | 6.7 | 19.7 | 72.8 |
| 2003..... | 1.2 | 7.7 | * | * | * | * | * | * | 2.4 | 8.7 | 27.6 | 60.0 |
| 2002..... | 1.2 | 5.2 | * | * | * | * | * | * | * | 12.4 | 32.0 | 60.2 |
| 2001..... | 1.2 | 5.5 | * | * | * | * | * | * | 2.5 | 8.2 | 28.2 | 104.8 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 10.8 | 33.9 | 70.5 |
| 1999..... | 1.2 | 8.5 | * | * | * | * | * | * | * | 8.6 | 31.3 | 69.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.9 | 6.3 | * | * | * | * | * | * | 2.5 | 8.4 | 33.3 | 74.2 |
| 2003..... | 1.9 | 5.0 | * | * | * | * | * | * | 2.5 | 11.0 | 31.3 | 79.7 |
| 2002..... | 1.9 | 5.4 | * | * | * | * | * | * | 2.5 | 9.6 | 30.9 | 83.7 |
| 2001..... | 1.9 | * | * | * | * | * | * | * | 3.4 | 11.5 | 30.8 | 91.2 |
| 2000..... | 1.8 | 5.6 | * | * | * | * | * | * | 2.6 | 9.1 | 28.1 | 96.9 |
| 1999..... | 1.9 | * | * | * | * | * | * | * | 3.5 | 10.7 | 28.8 | 104.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Vascular disorders and obstruction of intestine without hernia (K55-K56) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | 7.2 | 0.2 | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.6 | 12.4 | 38.7 | 108.1 |
| 2003..... | 5.7 | 6.8 | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.3 | 3.8 | 13.5 | 39.6 | 113.0 |
| 2002..... | 5.6 | 5.9 | 0.2 | 0.1 | 0.1 | 0.1 | 0.4 | 1.2 | 3.8 | 12.9 | 39.7 | 114.6 |
| 2001..... | 5.5 | 5.8 | * | 0.1 | * | 0.1 | 0.3 | 1.2 | 3.9 | 14.0 | 39.2 | 113.9 |
| 2000..... | 5.6 | 6.1 | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.3 | 4.0 | 13.5 | 40.4 | 117.0 |
| 1999..... | 5.5 | 5.9 | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.1 | 4.1 | 13.7 | 39.2 | 116.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.9 | 8.4 | * | * | * | 0.1 | 0.5 | 1.2 | 3.4 | 11.2 | 35.7 | 89.2 |
| 2003..... | 4.1 | 8.2 | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.8 | 12.5 | 35.9 | 94.0 |
| 2002..... | 4.0 | 6.8 | * | * | * | 0.2 | 0.5 | 1.2 | 3.6 | 11.6 | 35.9 | 98.0 |
| 2001..... | 4.0 | 6.7 | * | * | * | 0.1 | 0.4 | 1.2 | 3.6 | 13.3 | 35.4 | 98.2 |
| 2000..... | 4.0 | 6.4 | * | 0.1 | 0.1 | 0.1 | 0.4 | 1.2 | 3.9 | 12.6 | 36.9 | 99.4 |
| 1999..... | 3.9 | 7.2 | * | * | * | 0.1 | 0.4 | 1.1 | 3.9 | 12.7 | 35.7 | 96.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.0 | 5.9 | * | * | * | 0.2 | 0.5 | 1.2 | 3.9 | 13.4 | 40.7 | 116.6 |
| 2003..... | 7.2 | 5.3 | * | * | * | * | 0.4 | 1.3 | 3.8 | 14.4 | 42.1 | 121.3 |
| 2002..... | 7.1 | 4.9 | * | * | * | 0.1 | 0.3 | 1.2 | 3.9 | 14.1 | 42.2 | 121.8 |
| 2001..... | 7.0 | 4.9 | * | * | * | 0.1 | 0.3 | 1.3 | 4.2 | 14.6 | 41.7 | 120.4 |
| 2000..... | 7.1 | 5.9 | * | * | * | 0.1 | 0.4 | 1.3 | 4.1 | 14.3 | 42.6 | 124.1 |
| 1999..... | 7.0 | 4.6 | * | * | * | 0.2 | 0.4 | 1.2 | 4.3 | 14.5 | 41.4 | 125.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 6.0 | 4.0 | * | * | * | 0.1 | 0.4 | 1.1 | 3.4 | 12.3 | 38.7 | 109.8 |
| 2003..... | 6.2 | 4.1 | * | * | 0.1 | 0.1 | 0.3 | 1.1 | 3.5 | 13.4 | 39.7 | 115.3 |
| 2002..... | 6.1 | 3.7 | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 3.6 | 12.5 | 39.8 | 117.3 |
| 2001..... | 6.1 | 3.5 | * | * | * | 0.1 | 0.3 | 1.1 | 3.6 | 13.8 | 39.2 | 116.5 |
| 2000..... | 6.0 | 3.3 | * | 0.1 | 0.1 | 0.1 | 0.3 | 1.1 | 3.6 | 13.3 | 40.1 | 118.8 |
| 1999..... | 6.0 | 3.5 | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 4.0 | 13.1 | 39.0 | 118.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.2 | 5.0 | * | * | * | * | 0.5 | 1.1 | 3.1 | 10.6 | 34.7 | 88.9 |
| 2003..... | 4.3 | 5.4 | * | * | * | * | 0.3 | 1.0 | 3.3 | 11.8 | 35.0 | 94.3 |
| 2002..... | 4.2 | 4.3 | * | * | * | * | 0.4 | 1.0 | 3.2 | 11.0 | 35.5 | 98.9 |
| 2001..... | 4.2 | 3.7 | * | * | * | * | 0.3 | 1.0 | 3.3 | 12.8 | 35.0 | 98.3 |
| 2000..... | 4.1 | 3.4 | * | * | * | * | 0.3 | 1.0 | 3.3 | 12.0 | 35.9 | 99.7 |
| 1999..... | 4.1 | 4.3 | * | * | * | * | 0.3 | 0.9 | 3.7 | 12.1 | 35.1 | 95.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.8 | 3.0 | * | * | * | * | 0.4 | 1.1 | 3.7 | 13.8 | 41.4 | 119.2 |
| 2003..... | 8.1 | 2.9 | * | * | * | * | 0.3 | 1.2 | 3.7 | 14.8 | 42.9 | 124.5 |
| 2002..... | 7.9 | 3.2 | * | * | * | * | 0.3 | 1.1 | 3.9 | 13.8 | 42.7 | 125.3 |
| 2001..... | 7.8 | 3.2 | * | * | * | * | 0.3 | 1.1 | 3.8 | 14.6 | 42.0 | 124.1 |
| 2000..... | 7.9 | 3.2 | * | * | * | * | 0.4 | 1.1 | 3.9 | 14.3 | 43.0 | 126.5 |
| 1999..... | 7.8 | 2.7 | * | * | * | 0.2 | 0.3 | 1.0 | 4.3 | 14.0 | 41.5 | 127.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Vascular disorders and obstruction of intestine without hernia (K55-K56) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | 19.6 | * | * | * | 0.4 | 0.9 | 2.1 | 6.2 | 15.4 | 45.9 | 107.1 |
| 2003..... | 4.3 | 18.7 | * | * | * | * | 0.9 | 2.5 | 6.8 | 17.4 | 45.2 | 103.8 |
| 2002..... | 4.2 | 14.5 | * | * | * | * | 0.7 | 2.4 | 5.9 | 18.8 | 45.1 | 101.0 |
| 2001..... | 4.3 | 16.5 | * | * | * | * | 0.6 | 2.8 | 7.5 | 19.6 | 45.7 | 97.7 |
| 2000..... | 4.5 | 17.2 | * | * | * | * | 0.8 | 3.2 | 7.8 | 18.1 | 48.1 | 112.9 |
| 1999..... | 4.3 | 16.0 | * | * | * | * | 1.0 | 2.4 | 5.8 | 20.8 | 47.6 | 110.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.8 | 22.3 | * | * | * | * | * | 2.1 | 6.9 | 18.2 | 53.3 | 109.7 |
| 2003..... | 4.0 | 21.1 | * | * | * | * | * | 2.9 | 8.5 | 21.4 | 52.5 | 103.3 |
| 2002..... | 3.6 | 17.3 | * | * | * | * | 0.9 | 2.6 | 7.0 | 19.0 | 46.0 | 106.1 |
| 2001..... | 3.9 | 19.8 | * | * | * | * | 0.8 | 2.8 | 7.2 | 22.7 | 48.3 | 122.4 |
| 2000..... | 4.1 | 18.4 | * | * | * | * | 1.0 | 3.6 | 9.9 | 19.9 | 55.2 | 111.5 |
| 1999..... | 3.6 | 18.7 | * | * | * | * | 1.1 | 2.8 | 6.3 | 19.8 | 50.0 | 111.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.7 | 16.7 | * | * | * | * | 1.0 | 2.0 | 5.6 | 13.4 | 41.7 | 106.0 |
| 2003..... | 4.5 | 16.1 | * | * | * | * | 1.0 | 2.0 | 5.4 | 14.6 | 41.1 | 104.0 |
| 2002..... | 4.7 | 11.7 | * | * | * | * | * | 2.2 | 5.1 | 18.6 | 44.6 | 99.1 |
| 2001..... | 4.7 | 13.1 | * | * | * | * | * | 2.8 | 7.8 | 17.5 | 44.3 | 88.5 |
| 2000..... | 4.8 | 15.9 | * | * | * | * | 0.7 | 2.9 | 6.1 | 16.7 | 44.1 | 113.4 |
| 1999..... | 5.0 | 13.2 | * | * | * | * | 0.9 | 2.1 | 5.4 | 21.5 | 46.2 | 109.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diverticular disease of intestine (K57) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.8 | 9.3 | 27.0 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.7 | 9.8 | 28.9 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.7 | 9.9 | 27.7 |
| 2001..... | 1.2 | * | * | * | * | * | 0.0 | 0.2 | 0.9 | 2.9 | 9.4 | 31.4 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 3.0 | 9.2 | 30.9 |
| 1999..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.1 | 9.3 | 33.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 7.8 | 18.8 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 7.6 | 22.3 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.4 | 8.0 | 19.9 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.8 | 2.4 | 8.2 | 23.4 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.3 | 0.7 | 2.7 | 7.3 | 22.3 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.8 | 2.7 | 7.4 | 25.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.8 | 3.1 | 10.3 | 30.6 |
| 2003..... | 1.6 | * | * | * | * | * | * | 0.2 | 0.9 | 3.0 | 11.3 | 31.9 |
| 2002..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.9 | 11.1 | 31.0 |
| 2001..... | 1.6 | * | * | * | * | * | * | 0.2 | 1.0 | 3.3 | 10.2 | 34.7 |
| 2000..... | 1.6 | * | * | * | * | * | * | 0.2 | 0.7 | 3.1 | 10.5 | 34.4 |
| 1999..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.9 | 3.4 | 10.5 | 36.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | 2.3 | 7.0 | 14.7 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 2.5 | 8.0 | 21.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 2.1 | 9.2 | 16.6 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 2.3 | 7.6 | 22.5 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 2.8 | 7.1 | 21.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 2.4 | 8.1 | 33.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 7.2 | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.9 | 17.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 9.7 | 25.0 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 10.4 | 17.4 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | 8.8 | 27.0 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | 9.2 | 22.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.0 | 35.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diverticular disease of intestine (K57) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.8 | 9.4 | 27.5 |
| 2003..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.8 | 9.9 | 29.2 |
| 2002..... | 1.3 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.7 | 9.9 | 28.0 |
| 2001..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.9 | 2.9 | 9.5 | 31.6 |
| 2000..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 3.0 | 9.3 | 31.2 |
| 1999..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.2 | 9.3 | 33.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 7.8 | 19.3 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 7.7 | 22.6 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.4 | 0.7 | 2.4 | 8.0 | 20.1 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 8.3 | 23.8 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.7 | 2.7 | 7.4 | 22.5 |
| 1999..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.9 | 2.7 | 7.3 | 25.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | 0.1 | 0.8 | 3.1 | 10.5 | 31.1 |
| 2003..... | 1.8 | * | * | * | * | * | * | 0.2 | 0.9 | 3.0 | 11.3 | 32.1 |
| 2002..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.7 | 2.9 | 11.1 | 31.4 |
| 2001..... | 1.8 | * | * | * | * | * | * | 0.2 | 1.0 | 3.4 | 10.2 | 34.9 |
| 2000..... | 1.7 | * | * | * | * | * | * | 0.2 | 0.7 | 3.1 | 10.5 | 34.7 |
| 1999..... | 1.8 | * | * | * | * | * | * | 0.2 | 0.9 | 3.5 | 10.6 | 36.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.8 | 9.8 | 28.5 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.8 | 2.8 | 10.2 | 30.6 |
| 2002..... | 1.5 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.8 | 10.0 | 28.9 |
| 2001..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.9 | 2.9 | 9.9 | 32.7 |
| 2000..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.7 | 3.0 | 9.6 | 32.6 |
| 1999..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.9 | 3.2 | 9.6 | 34.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 7.9 | 19.8 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 8.0 | 23.8 |
| 2002..... | 0.9 | * | * | * | * | * | * | 0.4 | 0.7 | 2.5 | 7.8 | 20.8 |
| 2001..... | 1.0 | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 8.5 | 24.8 |
| 2000..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.7 | 2.7 | 7.8 | 23.9 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.8 | 2.8 | 7.5 | 26.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | * | * | * | 0.8 | 3.2 | 11.0 | 32.4 |
| 2003..... | 2.1 | * | * | * | * | * | * | 0.2 | 0.8 | 3.1 | 11.8 | 33.5 |
| 2002..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.7 | 3.0 | 11.5 | 32.4 |
| 2001..... | 2.1 | * | * | * | * | * | * | 0.2 | 1.0 | 3.3 | 10.8 | 36.0 |
| 2000..... | 2.0 | * | * | * | * | * | * | 0.2 | 0.7 | 3.2 | 10.8 | 36.1 |
| 1999..... | 2.1 | * | * | * | * | * | * | 0.2 | 0.9 | 3.5 | 11.0 | 37.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diverticular disease of intestine (K57) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 3.1 | 7.4 | 22.2 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | 1.5 | 2.6 | 8.1 | 18.7 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 0.9 | 2.6 | 10.0 | 22.1 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 1.1 | 4.0 | 6.9 | 24.3 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | 1.3 | 3.8 | 7.2 | 20.4 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 1.5 | 3.5 | 7.3 | 23.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.0 | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.3 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 11.4 | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 3.0 | 7.2 | * |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | 3.7 | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 3.4 | 7.1 | 23.7 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | 1.9 | 3.0 | 9.1 | 20.6 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | 3.1 | 9.1 | 25.1 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 4.7 | 6.7 | 27.8 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | * | 3.8 | 8.7 | 24.9 |
| 1999..... | 1.0 | * | * | * | * | * | * | * | 1.6 | 4.2 | 7.7 | 27.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of intestines and peritoneum (K52,K58-K63) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | 2.0 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.4 | 4.4 | 12.8 | 33.3 |
| 2003..... | 1.7 | 2.4 | 0.1 | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.6 | 4.3 | 12.6 | 32.8 |
| 2002..... | 1.7 | 2.0 | 0.2 | 0.0 | 0.0 | 0.1 | 0.2 | 0.6 | 1.6 | 4.3 | 12.5 | 32.8 |
| 2001..... | 1.7 | 1.4 | 0.2 | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 4.5 | 12.8 | 34.1 |
| 2000..... | 1.6 | 1.7 | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 4.1 | 12.4 | 33.3 |
| 1999..... | 1.6 | 1.5 | 0.2 | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.4 | 12.1 | 32.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.3 | 2.3 | * | * | * | * | 0.1 | 0.6 | 1.2 | 4.1 | 12.5 | 26.4 |
| 2003..... | 1.3 | 2.9 | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 4.2 | 12.3 | 27.1 |
| 2002..... | 1.3 | 2.4 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.4 | 11.8 | 29.4 |
| 2001..... | 1.2 | 1.6 | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.2 | 12.1 | 29.9 |
| 2000..... | 1.2 | 1.8 | * | * | * | * | 0.2 | 0.4 | 1.7 | 4.0 | 11.4 | 28.7 |
| 1999..... | 1.2 | 2.0 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.2 | 11.9 | 28.9 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.1 | 1.7 | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.6 | 12.9 | 36.4 |
| 2003..... | 2.1 | 1.8 | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.5 | 12.8 | 35.2 |
| 2002..... | 2.0 | 1.5 | * | * | * | * | 0.2 | 0.5 | 1.7 | 4.3 | 13.0 | 34.3 |
| 2001..... | 2.1 | 1.1 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.7 | 13.3 | 35.8 |
| 2000..... | 2.0 | 1.6 | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.2 | 13.1 | 35.2 |
| 1999..... | 1.9 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.6 | 12.2 | 33.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | 2.8 | 8.8 | 20.9 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 1.0 | 2.9 | 8.9 | 20.2 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 1.3 | 2.3 | 8.3 | 14.4 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 3.2 | 10.8 | 25.6 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 2.1 | 10.1 | 21.2 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.7 | 1.4 | 3.3 | 8.1 | 21.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | 9.3 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 10.4 | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | 8.2 | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | 13.4 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | 8.4 | 20.9 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | 2.9 | 7.9 | 21.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.4 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | 9.1 | 28.9 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | 11.2 | 20.8 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | * | 9.4 | 27.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of intestines and peritoneum (K52,K58-K63) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | 2.1 | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 4.5 | 12.9 | 33.8 |
| 2003..... | 1.9 | 2.4 | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.6 | 4.4 | 12.8 | 33.2 |
| 2002..... | 1.9 | 1.9 | 0.2 | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.6 | 4.5 | 12.7 | 33.5 |
| 2001..... | 1.8 | 1.4 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.5 | 12.9 | 34.4 |
| 2000..... | 1.8 | 1.6 | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 4.2 | 12.5 | 33.7 |
| 1999..... | 1.7 | 1.3 | 0.2 | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.5 | 12.3 | 32.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.4 | 2.4 | * | * | * | * | 0.1 | 0.6 | 1.2 | 4.2 | 12.7 | 26.7 |
| 2003..... | 1.4 | 3.0 | * | * | * | * | 0.2 | 0.6 | 1.6 | 4.2 | 12.4 | 27.5 |
| 2002..... | 1.4 | 2.4 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.6 | 12.0 | 29.9 |
| 2001..... | 1.4 | 1.7 | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.2 | 12.1 | 30.4 |
| 2000..... | 1.3 | 1.8 | * | * | * | * | 0.2 | 0.5 | 1.7 | 4.1 | 11.4 | 28.9 |
| 1999..... | 1.3 | 1.7 | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.2 | 12.1 | 29.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.3 | 1.8 | * | * | * | * | 0.3 | 0.5 | 1.7 | 4.7 | 13.1 | 37.0 |
| 2003..... | 2.3 | 1.8 | * | * | * | * | 0.3 | 0.5 | 1.5 | 4.6 | 13.0 | 35.7 |
| 2002..... | 2.3 | 1.4 | * | * | * | * | 0.3 | 0.6 | 1.7 | 4.4 | 13.2 | 35.1 |
| 2001..... | 2.3 | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 4.8 | 13.5 | 36.0 |
| 2000..... | 2.2 | 1.5 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.3 | 13.2 | 35.6 |
| 1999..... | 2.1 | * | * | * | * | * | 0.2 | 0.4 | 1.7 | 4.7 | 12.3 | 33.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | 1.5 | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.5 | 13.3 | 35.0 |
| 2003..... | 2.1 | 2.0 | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.3 | 13.2 | 34.2 |
| 2002..... | 2.1 | 1.4 | * | * | * | * | 0.2 | 0.6 | 1.6 | 4.5 | 13.1 | 34.9 |
| 2001..... | 2.0 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.6 | 12.9 | 35.5 |
| 2000..... | 1.9 | 1.2 | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.3 | 12.7 | 34.1 |
| 1999..... | 1.9 | 0.9 | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.5 | 12.4 | 33.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.5 | 1.7 | * | * | * | * | * | 0.5 | 1.2 | 4.0 | 12.8 | 27.2 |
| 2003..... | 1.6 | 2.6 | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.1 | 12.6 | 27.9 |
| 2002..... | 1.6 | 2.2 | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.4 | 12.1 | 30.9 |
| 2001..... | 1.5 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 4.1 | 12.0 | 31.7 |
| 2000..... | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 1.5 | 4.2 | 11.4 | 28.3 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 4.3 | 12.1 | 28.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | * | 0.3 | 0.5 | 1.7 | 4.9 | 13.7 | 38.5 |
| 2003..... | 2.6 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.5 | 13.6 | 37.0 |
| 2002..... | 2.6 | * | * | * | * | * | 0.3 | 0.6 | 1.7 | 4.6 | 13.8 | 36.6 |
| 2001..... | 2.6 | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 5.1 | 13.4 | 37.0 |
| 2000..... | 2.5 | * | * | * | * | * | 0.2 | 0.6 | 1.5 | 4.3 | 13.6 | 36.5 |
| 1999..... | 2.4 | * | * | * | * | * | 0.2 | 0.4 | 1.7 | 4.7 | 12.6 | 35.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of intestines and peritoneum (K52,K58-K63) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | 3.8 | * | * | * | * | * | 0.7 | 2.2 | 5.5 | 11.3 | 24.5 |
| 2003..... | 1.3 | 3.8 | * | * | * | * | 0.4 | 1.1 | 2.6 | 6.1 | 12.2 | 25.9 |
| 2002..... | 1.2 | 3.7 | * | * | * | * | 0.4 | 0.7 | 2.4 | 5.4 | 10.8 | 23.6 |
| 2001..... | 1.4 | 4.1 | * | * | * | * | 0.4 | 1.0 | 2.5 | 5.0 | 16.0 | 27.5 |
| 2000..... | 1.2 | 3.6 | * | * | * | * | 0.4 | 0.8 | 2.6 | 4.8 | 10.9 | 29.9 |
| 1999..... | 1.2 | * | * | * | * | * | 0.4 | 1.0 | 1.9 | 5.2 | 11.8 | 23.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | 1.0 | 2.1 | 6.8 | 12.8 | 25.4 |
| 2003..... | 1.2 | * | * | * | * | * | * | 1.2 | 3.1 | 6.4 | 13.3 | 25.6 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | 2.5 | 7.6 | 12.3 | 25.2 |
| 2001..... | 1.2 | * | * | * | * | * | * | * | 3.1 | 7.0 | 16.5 | * |
| 2000..... | 1.1 | * | * | * | * | * | * | * | 3.2 | 4.0 | 13.6 | 29.4 |
| 1999..... | 1.1 | * | * | * | * | * | * | 1.3 | 1.9 | 4.7 | 12.8 | 40.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.2 | 4.5 | 10.5 | 24.1 |
| 2003..... | 1.5 | * | * | * | * | * | * | 1.0 | 2.2 | 6.0 | 11.6 | 26.0 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | 2.3 | 3.8 | 9.9 | 23.0 |
| 2001..... | 1.5 | * | * | * | * | * | * | 1.2 | 2.0 | 3.6 | 15.7 | 29.9 |
| 2000..... | 1.4 | * | * | * | * | * | * | 1.0 | 2.1 | 5.3 | 9.4 | 30.1 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | 1.9 | 5.5 | 11.2 | 16.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of peritoneum (K65-K66) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.8 | 4.2 | 8.7 |
| 2003..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 1.9 | 4.2 | 9.2 |
| 2002..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.8 | 1.8 | 4.9 | 10.5 |
| 2001..... | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 4.5 | 10.4 |
| 2000..... | 0.8 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.1 | 4.5 | 10.8 |
| 1999..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.2 | 4.5 | 9.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 2.0 | 4.5 | 7.9 |
| 2003..... | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.9 | 4.6 | 8.8 |
| 2002..... | 0.6 | * | * | * | * | 0.1 | 0.5 | 0.9 | 1.7 | 4.4 | 9.2 | |
| 2001..... | 0.6 | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.9 | 4.6 | 10.3 |
| 2000..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 2.1 | 4.6 | 11.0 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.6 | 1.0 | 2.4 | 5.0 | 9.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.6 | 3.9 | 9.1 |
| 2003..... | 0.8 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.8 | 3.9 | 9.4 |
| 2002..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.9 | 5.2 | 11.1 |
| 2001..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.7 | 4.3 | 10.4 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.1 | 4.3 | 10.7 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.0 | 4.2 | 9.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 4.2 | 8.9 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.8 | 4.2 | 9.1 |
| 2002..... | 0.8 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.7 | 4.9 | 10.8 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.7 | 4.3 | 10.4 |
| 2000..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.0 | 4.4 | 10.8 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.1 | 4.4 | 9.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.5 | 0.7 | 1.9 | 4.5 | 8.0 |
| 2003..... | 0.6 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.8 | 4.7 | 8.9 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.5 | 0.8 | 1.6 | 4.3 | 9.5 |
| 2001..... | 0.6 | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.8 | 4.6 | 9.9 |
| 2000..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 1.9 | 4.5 | 10.8 |
| 1999..... | 0.7 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.3 | 4.9 | 10.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.9 | 9.3 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.5 | 1.8 | 3.9 | 9.2 |
| 2002..... | 1.0 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.8 | 5.3 | 11.3 |
| 2001..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.6 | 1.6 | 4.2 | 10.6 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.1 | 4.3 | 10.7 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.9 | 4.1 | 9.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of peritoneum (K65-K66) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.5 | 0.7 | 1.3 | 2.8 | 5.2 | 8.1 |
| 2003..... | 0.7 | * | * | * | * | * | * | 1.1 | 2.0 | 2.7 | 4.5 | 12.2 |
| 2002..... | 0.7 | * | * | * | * | * | * | 1.1 | 1.5 | 2.9 | 5.5 | 10.6 |
| 2001..... | 0.8 | * | * | * | * | * | 0.5 | 1.2 | 2.1 | 2.9 | 6.1 | 11.9 |
| 2000..... | 0.8 | * | * | * | * | * | 0.5 | 1.1 | 2.2 | 3.2 | 6.3 | 11.1 |
| 1999..... | 0.8 | * | * | * | * | * | * | 1.0 | 2.1 | 3.9 | 5.9 | 12.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | * | * |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.9 | 2.0 | * | * | * |
| 2002..... | 0.6 | * | * | * | * | * | * | 1.3 | * | * | 6.2 | * |
| 2001..... | 0.7 | * | * | * | * | * | * | 1.1 | 2.5 | 3.4 | * | * |
| 2000..... | 0.8 | * | * | * | * | * | * | 1.3 | 2.5 | 3.9 | 7.4 | * |
| 1999..... | 0.7 | * | * | * | * | * | * | 1.3 | 2.6 | 4.2 | 7.2 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 2.5 | 5.3 | 8.0 |
| 2003..... | 0.8 | * | * | * | * | * | * | 1.2 | 1.9 | 2.7 | 3.9 | 12.8 |
| 2002..... | 0.8 | * | * | * | * | * | * | 1.0 | 1.4 | 3.3 | 5.1 | 11.3 |
| 2001..... | 0.9 | * | * | * | * | * | * | 1.2 | 1.8 | 2.6 | 6.4 | 10.7 |
| 2000..... | 0.9 | * | * | * | * | * | * | 1.0 | 2.0 | 2.6 | 5.7 | 10.0 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | 1.7 | 3.6 | 5.3 | 12.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of liver (K70-K76) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 12.3 | 2.5 | 0.2 | * | 0.2 | 1.2 | 8.1 | 23.0 | 29.4 | 36.9 | 40.0 | 32.8 |
| 2003..... | 12.6 | 1.5 | 0.2 | * | 0.2 | 1.4 | 8.7 | 23.4 | 30.0 | 38.7 | 41.0 | 33.6 |
| 2002..... | 12.4 | 2.2 | 0.2 | 0.1 | 0.2 | 1.3 | 8.9 | 23.0 | 29.1 | 38.1 | 41.9 | 35.7 |
| 2001..... | 12.2 | 1.9 | 0.2 | 0.1 | 0.2 | 1.3 | 9.0 | 22.9 | 28.8 | 38.1 | 41.4 | 36.1 |
| 2000..... | 12.0 | 2.0 | 0.2 | 0.1 | 0.2 | 1.4 | 9.2 | 21.8 | 29.3 | 37.9 | 41.2 | 36.7 |
| 1999..... | 12.0 | 2.3 | 0.1 | 0.1 | 0.2 | 1.4 | 9.0 | 21.4 | 29.4 | 39.4 | 42.2 | 37.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 15.7 | 2.5 | * | * | 0.2 | 1.6 | 10.8 | 33.3 | 41.7 | 47.5 | 49.5 | 40.7 |
| 2003..... | 16.1 | 2.2 | 0.2 | * | 0.2 | 1.7 | 11.4 | 34.6 | 42.9 | 50.3 | 49.6 | 44.4 |
| 2002..... | 15.8 | 3.1 | * | * | 0.2 | 1.6 | 11.6 | 33.9 | 41.2 | 49.5 | 51.7 | 47.3 |
| 2001..... | 15.7 | 2.3 | * | * | 0.2 | 1.6 | 11.9 | 33.7 | 40.9 | 50.2 | 50.9 | 49.4 |
| 2000..... | 15.5 | 2.7 | * | * | 0.2 | 1.8 | 12.5 | 32.5 | 41.3 | 49.7 | 51.0 | 52.2 |
| 1999..... | 15.6 | 2.9 | * | * | 0.2 | 1.8 | 12.5 | 32.4 | 42.1 | 52.5 | 52.0 | 52.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 9.0 | 2.4 | * | * | 0.2 | 0.8 | 5.5 | 13.0 | 18.1 | 27.9 | 33.7 | 29.3 |
| 2003..... | 9.1 | * | * | * | 0.2 | 1.0 | 6.1 | 12.7 | 18.1 | 29.0 | 35.2 | 28.9 |
| 2002..... | 9.0 | 1.3 | * | * | 0.2 | 1.1 | 6.2 | 12.5 | 17.9 | 28.5 | 35.4 | 30.7 |
| 2001..... | 8.9 | 1.6 | * | * | 0.2 | 1.0 | 6.2 | 12.5 | 17.5 | 28.1 | 35.1 | 30.5 |
| 2000..... | 8.7 | 1.3 | * | * | 0.2 | 1.0 | 5.9 | 11.4 | 18.3 | 28.3 | 34.7 | 30.4 |
| 1999..... | 8.6 | 1.6 | * | * | 0.2 | 1.1 | 5.6 | 10.7 | 17.7 | 28.8 | 35.8 | 31.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 10.2 | * | * | * | * | 1.3 | 9.2 | 30.2 | 42.1 | 60.1 | 73.5 | 38.5 |
| 2003..... | 10.7 | * | * | * | * | 1.8 | 9.7 | 29.9 | 48.3 | 63.5 | 81.2 | 48.2 |
| 2002..... | 10.7 | 2.5 | * | * | * | 1.6 | 10.2 | 31.8 | 46.1 | 67.9 | 67.6 | 57.1 |
| 2001..... | 10.7 | * | * | * | * | 1.4 | 10.7 | 32.4 | 46.5 | 66.5 | 77.5 | 55.6 |
| 2000..... | 10.7 | * | * | * | * | 1.6 | 11.5 | 32.0 | 47.4 | 69.6 | 77.6 | 67.0 |
| 1999..... | 10.7 | * | * | * | * | 1.7 | 12.1 | 31.8 | 47.2 | 68.2 | 73.3 | 73.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 13.5 | * | * | * | * | 1.9 | 13.9 | 49.2 | 62.0 | 73.2 | 82.3 | 42.0 |
| 2003..... | 14.2 | * | * | * | * | 2.4 | 14.3 | 48.3 | 74.8 | 75.5 | 92.0 | 63.1 |
| 2002..... | 14.5 | * | * | * | * | 2.1 | 15.7 | 51.6 | 70.0 | 92.7 | 82.9 | 61.9 |
| 2001..... | 14.7 | * | * | * | * | 2.0 | 16.3 | 53.1 | 72.9 | 91.7 | 96.6 | 68.6 |
| 2000..... | 14.7 | * | * | * | * | 2.5 | 17.6 | 52.0 | 74.7 | 95.8 | 96.4 | 88.7 |
| 1999..... | 14.6 | * | * | * | * | 2.4 | 18.9 | 52.1 | 74.3 | 94.5 | 86.6 | 99.2 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | 0.7 | 4.0 | 11.0 | 24.1 | 49.7 | 67.4 | 36.7 |
| 2003..... | 7.1 | * | * | * | * | 1.1 | 4.5 | 11.5 | 24.3 | 54.0 | 73.7 | 40.6 |
| 2002..... | 6.6 | * | * | * | * | 0.9 | 4.2 | 12.2 | 24.7 | 48.2 | 57.1 | 54.7 |
| 2001..... | 6.5 | * | * | * | * | 0.8 | 4.8 | 12.0 | 23.0 | 46.6 | 64.7 | 49.3 |
| 2000..... | 6.5 | * | * | * | * | * | 5.0 | 12.4 | 23.2 | 48.9 | 65.0 | 56.4 |
| 1999..... | 6.5 | * | * | * | * | 0.9 | 4.8 | 12.0 | 23.3 | 47.7 | 64.5 | 60.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of liver (K70-K76) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 12.6 | 2.6 | 0.2 | * | 0.2 | 1.2 | 8.0 | 22.1 | 28.3 | 35.0 | 38.1 | 32.6 |
| 2003..... | 12.8 | 1.6 | 0.2 | * | 0.2 | 1.3 | 8.6 | 22.6 | 28.4 | 36.9 | 38.8 | 33.0 |
| 2002..... | 12.6 | 2.1 | * | 0.1 | 0.2 | 1.3 | 8.7 | 22.0 | 27.5 | 35.8 | 40.5 | 34.8 |
| 2001..... | 12.4 | 1.8 | 0.2 | 0.1 | 0.2 | 1.3 | 8.8 | 21.9 | 27.2 | 36.0 | 39.6 | 35.3 |
| 2000..... | 12.2 | 2.2 | 0.2 | 0.1 | 0.2 | 1.3 | 8.8 | 20.7 | 27.8 | 35.8 | 39.5 | 35.5 |
| 1999..... | 12.2 | 2.2 | 0.2 | 0.1 | 0.2 | 1.4 | 8.6 | 20.3 | 28.0 | 37.6 | 40.8 | 35.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 16.0 | 2.7 | * | * | 0.2 | 1.5 | 10.2 | 31.3 | 39.8 | 45.5 | 47.6 | 40.5 |
| 2003..... | 16.4 | 2.2 | * | * | 0.2 | 1.5 | 10.8 | 32.9 | 40.1 | 48.4 | 47.3 | 43.5 |
| 2002..... | 15.9 | 3.0 | * | * | 0.2 | 1.5 | 10.9 | 32.0 | 38.6 | 46.4 | 50.0 | 46.6 |
| 2001..... | 15.8 | 2.1 | * | * | 0.2 | 1.5 | 11.3 | 31.7 | 38.2 | 47.3 | 48.6 | 48.5 |
| 2000..... | 15.5 | 3.0 | * | * | 0.2 | 1.6 | 11.7 | 30.5 | 38.5 | 46.6 | 49.0 | 50.5 |
| 1999..... | 15.7 | 3.1 | * | * | 0.2 | 1.6 | 11.6 | 30.4 | 39.6 | 49.8 | 50.5 | 50.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 9.3 | 2.6 | * | * | 0.2 | 0.8 | 5.7 | 13.2 | 17.5 | 26.2 | 31.8 | 29.0 |
| 2003..... | 9.4 | * | * | * | 0.3 | 1.0 | 6.3 | 12.8 | 17.6 | 27.2 | 33.1 | 28.3 |
| 2002..... | 9.4 | * | * | * | 0.2 | 1.1 | 6.5 | 12.5 | 17.3 | 27.0 | 34.3 | 29.7 |
| 2001..... | 9.2 | 1.5 | * | * | 0.2 | 1.1 | 6.3 | 12.5 | 17.0 | 26.7 | 33.6 | 29.8 |
| 2000..... | 9.0 | * | * | * | 0.2 | 1.1 | 6.0 | 11.3 | 17.9 | 26.9 | 33.3 | 29.4 |
| 1999..... | 8.9 | 1.3 | * | * | 0.2 | 1.1 | 5.7 | 10.6 | 17.3 | 27.6 | 34.5 | 30.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 13.4 | 1.8 | * | * | 0.1 | 1.1 | 8.0 | 21.9 | 27.7 | 35.8 | 39.4 | 33.1 |
| 2003..... | 13.7 | 1.2 | * | * | 0.1 | 1.2 | 8.7 | 22.3 | 27.9 | 37.7 | 40.3 | 33.5 |
| 2002..... | 13.4 | 1.6 | * | * | 0.2 | 1.2 | 8.8 | 21.5 | 27.1 | 36.8 | 41.7 | 35.7 |
| 2001..... | 13.1 | 1.3 | * | * | 0.1 | 1.2 | 8.6 | 21.2 | 26.6 | 36.6 | 40.6 | 36.1 |
| 2000..... | 12.7 | 1.7 | * | * | 0.1 | 1.2 | 8.5 | 19.8 | 27.1 | 36.2 | 40.5 | 36.1 |
| 1999..... | 12.7 | 1.4 | * | * | 0.1 | 1.2 | 8.1 | 19.3 | 26.8 | 37.9 | 41.7 | 36.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 17.0 | 1.8 | * | * | * | 1.5 | 10.2 | 31.1 | 38.8 | 46.2 | 49.2 | 42.5 |
| 2003..... | 17.4 | 1.7 | * | * | * | 1.5 | 11.2 | 32.4 | 39.0 | 49.2 | 49.0 | 44.0 |
| 2002..... | 17.0 | 2.4 | * | * | * | 1.4 | 11.2 | 31.3 | 37.9 | 47.4 | 51.4 | 47.9 |
| 2001..... | 16.5 | * | * | * | * | 1.4 | 10.9 | 30.7 | 36.9 | 47.7 | 49.6 | 49.7 |
| 2000..... | 16.2 | 2.1 | * | * | * | 1.4 | 11.4 | 29.4 | 37.7 | 47.1 | 50.3 | 51.3 |
| 1999..... | 16.3 | 1.9 | * | * | * | 1.5 | 11.1 | 29.0 | 38.2 | 50.0 | 51.8 | 51.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 9.9 | 1.9 | * | * | * | 0.8 | 5.7 | 12.9 | 17.1 | 26.8 | 32.8 | 28.9 |
| 2003..... | 10.0 | * | * | * | 0.2 | 0.9 | 6.2 | 12.3 | 17.3 | 27.8 | 34.5 | 28.8 |
| 2002..... | 10.0 | * | * | * | 0.2 | 0.9 | 6.5 | 11.8 | 16.9 | 27.8 | 35.3 | 30.4 |
| 2001..... | 9.8 | * | * | * | * | 1.0 | 6.3 | 11.8 | 16.8 | 27.2 | 34.6 | 30.4 |
| 2000..... | 9.4 | * | * | * | * | 0.9 | 5.6 | 10.4 | 17.1 | 27.1 | 34.1 | 30.0 |
| 1999..... | 9.2 | * | * | * | * | 0.9 | 5.2 | 9.7 | 16.2 | 27.7 | 35.2 | 30.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of liver (K70-K76) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 10.1 | 5.7 | * | * | 0.4 | 1.1 | 8.1 | 26.3 | 36.2 | 32.9 | 25.4 | 24.5 |
| 2003..... | 10.3 | * | * | * | 0.5 | 1.7 | 8.1 | 28.0 | 36.2 | 35.4 | 24.0 | 27.7 |
| 2002..... | 10.2 | 3.9 | * | * | 0.4 | 1.5 | 8.1 | 29.0 | 34.4 | 31.8 | 29.4 | 27.2 |
| 2001..... | 10.7 | 3.6 | * | * | 0.5 | 1.5 | 10.0 | 29.9 | 36.1 | 35.5 | 31.1 | 22.2 |
| 2000..... | 10.8 | 4.5 | * | * | 0.5 | 1.8 | 10.7 | 30.1 | 35.6 | 36.6 | 29.1 | 26.4 |
| 1999..... | 11.1 | 5.1 | * | * | 0.4 | 1.8 | 11.1 | 30.2 | 39.6 | 37.7 | 32.1 | 30.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 13.3 | * | * | * | * | 1.4 | 10.5 | 38.1 | 55.3 | 47.7 | 33.4 | 20.3 |
| 2003..... | 13.7 | * | * | * | * | 1.8 | 9.4 | 41.8 | 56.1 | 50.6 | 32.0 | 41.6 |
| 2002..... | 13.0 | * | * | * | * | 1.6 | 9.4 | 41.4 | 51.4 | 44.8 | 37.2 | 36.1 |
| 2001..... | 14.4 | * | * | * | * | 1.7 | 13.6 | 44.9 | 54.8 | 51.9 | 41.4 | 27.7 |
| 2000..... | 13.9 | 6.8 | * | * | * | 1.9 | 13.8 | 43.9 | 51.3 | 51.2 | 37.6 | 36.4 |
| 1999..... | 14.5 | 7.3 | * | * | * | 1.7 | 14.5 | 45.8 | 56.7 | 54.0 | 39.7 | 34.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 7.2 | * | * | * | * | 0.8 | 6.0 | 16.2 | 20.9 | 22.4 | 20.9 | 26.1 |
| 2003..... | 7.3 | * | * | * | * | 1.5 | 7.0 | 16.1 | 20.3 | 24.7 | 19.4 | 22.3 |
| 2002..... | 7.6 | * | * | * | * | 1.4 | 7.0 | 18.4 | 20.8 | 22.5 | 25.0 | 23.8 |
| 2001..... | 7.4 | * | * | * | * | 1.3 | 6.8 | 17.0 | 21.0 | 24.0 | 25.3 | 20.1 |
| 2000..... | 7.9 | * | * | * | * | 1.8 | 7.9 | 18.3 | 22.9 | 26.3 | 24.3 | 22.7 |
| 1999..... | 8.1 | * | * | * | * | 1.8 | 8.0 | 16.8 | 25.8 | 26.3 | 27.8 | 28.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Alcoholic liver disease (K70) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | 0.6 | 4.2 | 10.6 | 11.7 | 10.0 | 5.5 | 2.1 |
| 2003..... | 4.3 | * | * | * | * | 0.6 | 4.4 | 10.5 | 11.6 | 9.8 | 5.7 | 1.8 |
| 2002..... | 4.2 | * | * | * | * | 0.6 | 4.4 | 10.3 | 11.6 | 10.1 | 5.7 | 2.1 |
| 2001..... | 4.3 | * | * | * | * | 0.7 | 4.8 | 10.7 | 11.5 | 10.1 | 5.7 | 2.4 |
| 2000..... | 4.3 | * | * | * | * | 0.7 | 4.9 | 10.3 | 12.2 | 10.3 | 6.1 | 2.2 |
| 1999..... | 4.3 | * | * | * | * | 0.6 | 4.8 | 10.4 | 12.0 | 10.5 | 6.5 | 2.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.3 | * | * | * | * | 0.8 | 5.8 | 15.5 | 18.2 | 16.5 | 10.1 | 4.6 |
| 2003..... | 6.4 | * | * | * | * | 0.9 | 6.0 | 16.1 | 18.3 | 16.1 | 10.1 | 3.6 |
| 2002..... | 6.2 | * | * | * | * | 0.8 | 5.8 | 15.6 | 18.3 | 16.5 | 9.7 | 4.7 |
| 2001..... | 6.4 | * | * | * | * | 0.9 | 6.5 | 16.1 | 18.4 | 16.6 | 10.1 | 5.5 |
| 2000..... | 6.5 | * | * | * | * | 1.0 | 6.8 | 16.0 | 19.0 | 17.3 | 11.5 | 5.3 |
| 1999..... | 6.5 | * | * | * | * | 0.8 | 6.8 | 16.3 | 19.2 | 18.1 | 11.2 | 6.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 0.3 | 2.7 | 5.7 | 5.6 | 4.5 | 2.3 | 1.0 |
| 2003..... | 2.2 | * | * | * | * | 0.3 | 2.9 | 5.1 | 5.4 | 4.5 | 2.7 | 1.0 |
| 2002..... | 2.3 | * | * | * | * | 0.4 | 3.0 | 5.2 | 5.4 | 4.8 | 3.1 | 1.0 |
| 2001..... | 2.3 | * | * | * | * | 0.4 | 3.1 | 5.5 | 5.1 | 4.7 | 2.8 | 1.0 |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 3.1 | 4.8 | 6.0 | 4.5 | 2.5 | 1.0 |
| 1999..... | 2.1 | * | * | * | * | 0.5 | 2.9 | 4.7 | 5.4 | 4.3 | 3.4 | 1.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 0.7 | 5.3 | 15.2 | 17.2 | 16.1 | 14.6 | * |
| 2003..... | 4.3 | * | * | * | * | 0.9 | 5.6 | 14.4 | 21.4 | 17.7 | 14.1 | * |
| 2002..... | 4.3 | * | * | * | * | 0.7 | 5.5 | 16.1 | 19.9 | 20.8 | 10.5 | * |
| 2001..... | 4.7 | * | * | * | * | 0.9 | 6.3 | 17.1 | 21.4 | 21.1 | 17.0 | * |
| 2000..... | 4.8 | * | * | * | * | 1.0 | 6.9 | 16.8 | 23.4 | 22.9 | 15.4 | * |
| 1999..... | 4.6 | * | * | * | * | 0.8 | 6.9 | 16.8 | 21.7 | 20.6 | 14.4 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.6 | * | * | * | * | 1.2 | 8.2 | 26.6 | 29.7 | 28.6 | 28.3 | * |
| 2003..... | 6.9 | * | * | * | * | 1.3 | 8.8 | 25.5 | 38.4 | 29.0 | 26.5 | * |
| 2002..... | 7.1 | * | * | * | * | 1.2 | 9.0 | 27.0 | 36.3 | 40.3 | 18.1 | * |
| 2001..... | 7.7 | * | * | * | * | 1.5 | 10.0 | 29.3 | 38.9 | 41.9 | 30.4 | * |
| 2000..... | 7.9 | * | * | * | * | 1.8 | 10.6 | 29.3 | 42.8 | 46.6 | 33.9 | * |
| 1999..... | 7.4 | * | * | * | * | 1.2 | 10.8 | 29.1 | 40.0 | 39.5 | 27.1 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 2.0 | 3.6 | 5.9 | 6.1 | 5.2 | * |
| 2003..... | 1.5 | * | * | * | * | * | 2.1 | 3.4 | 6.1 | 8.7 | 5.5 | * |
| 2002..... | 1.4 | * | * | * | * | * | 1.8 | 5.1 | 5.1 | 5.3 | * | * |
| 2001..... | 1.5 | * | * | * | * | * | 2.2 | 5.1 | 5.9 | 4.7 | 7.9 | * |
| 2000..... | 1.4 | * | * | * | * | * | 2.9 | 4.5 | 6.2 | 4.3 | * | * |
| 1999..... | 1.5 | * | * | * | * | * | 2.8 | 4.9 | 5.6 | 5.7 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Alcoholic liver disease (K70) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | 0.5 | 4.1 | 10.0 | 11.2 | 9.5 | 4.9 | 2.1 |
| 2003..... | 4.2 | * | * | * | * | 0.5 | 4.2 | 10.1 | 10.8 | 9.1 | 5.2 | 1.7 |
| 2002..... | 4.2 | * | * | * | * | 0.6 | 4.2 | 9.7 | 10.8 | 9.3 | 5.5 | 1.9 |
| 2001..... | 4.2 | * | * | * | * | 0.6 | 4.6 | 10.1 | 10.7 | 9.3 | 5.2 | 2.2 |
| 2000..... | 4.2 | * | * | * | * | 0.6 | 4.7 | 9.7 | 11.3 | 9.5 | 5.7 | 2.2 |
| 1999..... | 4.2 | * | * | * | * | 0.6 | 4.6 | 9.8 | 11.3 | 9.9 | 6.1 | 2.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.3 | * | * | * | * | 0.7 | 5.3 | 14.2 | 17.2 | 15.6 | 9.1 | 4.5 |
| 2003..... | 6.2 | * | * | * | * | 0.8 | 5.5 | 15.0 | 16.6 | 15.1 | 9.2 | 3.4 |
| 2002..... | 6.0 | * | * | * | * | 0.7 | 5.3 | 14.4 | 16.7 | 14.8 | 9.2 | 4.3 |
| 2001..... | 6.2 | * | * | * | * | 0.8 | 5.9 | 14.8 | 16.7 | 14.9 | 9.2 | 5.1 |
| 2000..... | 6.2 | * | * | * | * | 0.8 | 6.3 | 14.7 | 17.1 | 15.4 | 10.5 | 5.0 |
| 1999..... | 6.4 | * | * | * | * | 0.7 | 6.2 | 15.1 | 17.7 | 16.7 | 10.5 | 5.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | 0.3 | 2.8 | 5.9 | 5.6 | 4.3 | 2.2 | 1.0 |
| 2003..... | 2.3 | * | * | * | * | 0.3 | 3.0 | 5.3 | 5.4 | 4.1 | 2.6 | 1.0 |
| 2002..... | 2.4 | * | * | * | * | 0.5 | 3.1 | 5.2 | 5.4 | 4.8 | 3.0 | 0.9 |
| 2001..... | 2.4 | * | * | * | * | 0.5 | 3.2 | 5.5 | 5.0 | 4.7 | 2.5 | 1.0 |
| 2000..... | 2.3 | * | * | * | * | 0.4 | 3.1 | 4.8 | 5.9 | 4.5 | 2.5 | 1.0 |
| 1999..... | 2.2 | * | * | * | * | 0.5 | 2.9 | 4.7 | 5.3 | 4.2 | 3.3 | 1.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | 0.5 | 4.2 | 10.3 | 11.1 | 9.7 | 5.1 | 2.2 |
| 2003..... | 4.5 | * | * | * | * | 0.5 | 4.4 | 10.1 | 10.7 | 9.3 | 5.4 | 1.7 |
| 2002..... | 4.4 | * | * | * | * | 0.5 | 4.3 | 9.7 | 10.8 | 9.5 | 5.5 | 1.9 |
| 2001..... | 4.4 | * | * | * | * | 0.6 | 4.5 | 9.9 | 10.5 | 9.3 | 5.3 | 2.1 |
| 2000..... | 4.3 | * | * | * | * | 0.6 | 4.5 | 9.4 | 11.1 | 9.5 | 5.6 | 2.2 |
| 1999..... | 4.3 | * | * | * | * | 0.6 | 4.4 | 9.5 | 10.8 | 9.7 | 6.0 | 2.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | * | 0.7 | 5.5 | 14.6 | 17.0 | 15.7 | 9.2 | 4.7 |
| 2003..... | 6.6 | * | * | * | * | 0.8 | 5.6 | 15.1 | 16.4 | 15.3 | 9.4 | 3.5 |
| 2002..... | 6.4 | * | * | * | * | 0.7 | 5.4 | 14.4 | 16.7 | 14.9 | 9.4 | 4.1 |
| 2001..... | 6.4 | * | * | * | * | 0.8 | 5.8 | 14.7 | 16.4 | 14.6 | 9.3 | 4.8 |
| 2000..... | 6.5 | * | * | * | * | 0.8 | 6.1 | 14.5 | 17.0 | 15.4 | 10.6 | 5.1 |
| 1999..... | 6.5 | * | * | * | * | 0.7 | 6.0 | 14.7 | 17.1 | 16.3 | 10.3 | 5.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | 0.3 | 3.0 | 6.0 | 5.5 | 4.5 | 2.3 | 1.0 |
| 2003..... | 2.4 | * | * | * | * | 0.3 | 3.1 | 5.3 | 5.2 | 4.1 | 2.8 | 1.0 |
| 2002..... | 2.4 | * | * | * | * | 0.4 | 3.2 | 5.1 | 5.2 | 4.9 | 3.0 | 1.0 |
| 2001..... | 2.4 | * | * | * | * | 0.4 | 3.2 | 5.3 | 5.0 | 4.8 | 2.5 | 1.0 |
| 2000..... | 2.2 | * | * | * | * | 0.4 | 3.0 | 4.4 | 5.5 | 4.6 | 2.4 | 1.0 |
| 1999..... | 2.2 | * | * | * | * | 0.5 | 2.8 | 4.3 | 4.9 | 4.2 | 3.2 | 1.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Alcoholic liver disease (K70) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 3.0 | 9.2 | 12.8 | 9.1 | 3.2 | * |
| 2003..... | 3.3 | * | * | * | * | * | 3.4 | 10.0 | 12.7 | 9.4 | 3.4 | * |
| 2002..... | 3.2 | * | * | * | * | 0.5 | 3.0 | 10.2 | 12.3 | 8.7 | 5.1 | * |
| 2001..... | 3.7 | * | * | * | * | 0.5 | 4.4 | 11.7 | 13.1 | 10.8 | 4.6 | * |
| 2000..... | 3.9 | * | * | * | * | 0.7 | 5.1 | 12.3 | 13.5 | 10.4 | 5.6 | * |
| 1999..... | 4.2 | * | * | * | * | 0.5 | 5.1 | 12.9 | 15.4 | 11.9 | 7.2 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.6 | * | * | * | * | * | 4.1 | 13.2 | 20.7 | 17.2 | * | * |
| 2003..... | 4.9 | * | * | * | * | * | 4.2 | 16.0 | 20.3 | 16.2 | 6.9 | * |
| 2002..... | 4.6 | * | * | * | * | * | 3.9 | 15.1 | 18.6 | 15.1 | 8.8 | * |
| 2001..... | 5.5 | * | * | * | * | * | 6.1 | 17.4 | 21.7 | 20.0 | 7.8 | * |
| 2000..... | 5.5 | * | * | * | * | * | 7.1 | 17.9 | 19.6 | 18.4 | 10.2 | * |
| 1999..... | 6.1 | * | * | * | * | * | 6.9 | 19.7 | 24.6 | 21.9 | 12.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | 2.0 | 5.7 | 6.4 | 3.4 | * | * |
| 2003..... | 1.9 | * | * | * | * | * | 2.6 | 4.8 | 6.6 | 4.6 | * | * |
| 2002..... | 2.1 | * | * | * | * | * | 2.2 | 6.1 | 7.2 | 4.2 | * | * |
| 2001..... | 2.2 | * | * | * | * | * | 2.8 | 6.8 | 6.2 | 4.3 | * | * |
| 2000..... | 2.5 | * | * | * | * | * | 3.4 | 7.5 | 8.6 | 4.8 | * | * |
| 1999..... | 2.4 | * | * | * | * | * | 3.5 | 7.0 | 8.1 | 4.8 | 4.0 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Chronic hepatitis, not elsewhere classified (K73) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | 0.1 | * | * | 0.2 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.2 | 0.2 | * |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.9 | 0.7 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 0.8 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 0.8 | 0.8 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 0.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | 0.1 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.2 | * | 0.4 | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.4 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.2 | * | 0.3 | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 1.2 | 0.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.0 | 0.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 1.1 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.1 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | 0.1 | * | * | 0.2 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.2 | 0.2 | * |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.0 | 0.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 0.8 | 0.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 0.8 | 0.8 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 0.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | 0.1 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.4 | 0.5 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | 0.2 | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.0 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.2 | 0.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.0 | 0.8 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.5 | 0.9 | 1.1 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.1 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Fibrosis and cirrhosis of liver (K74) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | * | 0.2 | 2.1 | 7.4 | 10.9 | 17.6 | 23.1 | 17.4 |
| 2003..... | 5.2 | * | * | * | * | 0.3 | 2.3 | 7.7 | 11.3 | 19.6 | 24.1 | 18.0 |
| 2002..... | 5.1 | * | * | * | * | 0.3 | 2.5 | 7.5 | 11.0 | 18.6 | 24.7 | 18.5 |
| 2001..... | 5.0 | * | * | * | * | 0.3 | 2.6 | 7.6 | 10.9 | 19.2 | 23.7 | 19.1 |
| 2000..... | 5.0 | * | * | * | 0.1 | 0.3 | 2.5 | 7.2 | 11.3 | 18.9 | 24.1 | 20.0 |
| 1999..... | 5.0 | * | * | * | * | 0.3 | 2.4 | 6.8 | 11.3 | 19.6 | 24.6 | 19.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | 0.3 | 2.9 | 10.6 | 14.5 | 20.4 | 26.5 | 20.9 |
| 2003..... | 6.1 | * | * | * | * | 0.3 | 3.2 | 11.3 | 15.4 | 22.8 | 26.7 | 23.7 |
| 2002..... | 6.0 | * | * | * | * | 0.3 | 3.4 | 11.1 | 14.7 | 21.7 | 28.1 | 23.2 |
| 2001..... | 6.0 | * | * | * | * | 0.4 | 3.3 | 11.3 | 14.6 | 23.2 | 26.2 | 25.7 |
| 2000..... | 5.9 | * | * | * | * | 0.4 | 3.4 | 10.7 | 15.3 | 22.5 | 26.5 | 28.4 |
| 1999..... | 5.9 | * | * | * | * | 0.4 | 3.5 | 10.4 | 15.2 | 23.3 | 27.9 | 26.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.0 | * | * | * | * | 0.1 | 1.3 | 4.2 | 7.5 | 15.3 | 20.9 | 15.8 |
| 2003..... | 4.2 | * | * | * | * | 0.2 | 1.5 | 4.2 | 7.5 | 16.8 | 22.4 | 15.5 |
| 2002..... | 4.2 | * | * | * | * | 0.2 | 1.7 | 4.1 | 7.6 | 16.0 | 22.4 | 16.5 |
| 2001..... | 4.2 | * | * | * | * | 0.2 | 1.8 | 4.1 | 7.5 | 15.9 | 22.1 | 16.4 |
| 2000..... | 4.1 | * | * | * | * | 0.2 | 1.5 | 3.8 | 7.6 | 15.9 | 22.5 | 16.6 |
| 1999..... | 4.1 | * | * | * | * | 0.2 | 1.3 | 3.4 | 7.8 | 16.5 | 22.5 | 17.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | * | 2.3 | 9.5 | 17.3 | 30.9 | 45.4 | 23.8 |
| 2003..... | 4.1 | * | * | * | * | 0.4 | 2.3 | 9.7 | 16.7 | 33.4 | 47.6 | 26.9 |
| 2002..... | 4.3 | * | * | * | * | 0.3 | 3.1 | 10.6 | 17.8 | 33.3 | 40.7 | 37.1 |
| 2001..... | 4.1 | * | * | * | * | * | 3.4 | 10.5 | 16.9 | 32.1 | 42.0 | 30.0 |
| 2000..... | 4.1 | * | * | * | * | 0.4 | 2.8 | 10.3 | 17.4 | 33.8 | 46.6 | 41.8 |
| 1999..... | 4.1 | * | * | * | * | 0.4 | 3.5 | 9.5 | 16.7 | 34.5 | 44.2 | 38.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | * | * | 3.4 | 14.0 | 22.6 | 30.2 | 42.6 | * |
| 2003..... | 4.6 | * | * | * | * | 0.6 | 3.4 | 14.3 | 23.4 | 32.3 | 44.5 | * |
| 2002..... | 5.0 | * | * | * | * | * | 4.5 | 16.8 | 23.6 | 37.1 | 46.0 | * |
| 2001..... | 4.9 | * | * | * | * | * | 5.0 | 16.9 | 24.7 | 34.7 | 45.2 | * |
| 2000..... | 4.7 | * | * | * | * | * | 4.3 | 16.2 | 23.0 | 36.5 | 46.2 | 46.4 |
| 1999..... | 4.8 | * | * | * | * | 0.6 | 5.7 | 15.0 | 22.4 | 39.3 | 43.3 | 45.3 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | * | * | * | 1.1 | 5.0 | 12.5 | 31.4 | 47.4 | 23.8 |
| 2003..... | 3.7 | * | * | * | * | * | 1.1 | 5.0 | 10.7 | 34.3 | 49.8 | 26.6 |
| 2002..... | 3.4 | * | * | * | * | * | 1.6 | 4.4 | 12.5 | 30.3 | 37.1 | 39.8 |
| 2001..... | 3.2 | * | * | * | * | * | 1.7 | 4.3 | 10.0 | 30.1 | 39.9 | 32.6 |
| 2000..... | 3.5 | * | * | * | * | * | 1.1 | 4.5 | 12.4 | 31.7 | 46.9 | 39.6 |
| 1999..... | 3.3 | * | * | * | * | * | 1.2 | 4.1 | 11.6 | 30.8 | 44.8 | 35.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Fibrosis and cirrhosis of liver (K74) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | * | 0.2 | 2.0 | 7.1 | 10.3 | 16.6 | 21.9 | 17.1 |
| 2003..... | 5.3 | * | * | * | * | 0.2 | 2.3 | 7.5 | 10.8 | 18.5 | 22.8 | 17.7 |
| 2002..... | 5.2 | * | * | * | * | 0.3 | 2.5 | 7.2 | 10.4 | 17.5 | 23.8 | 17.7 |
| 2001..... | 5.2 | * | * | * | * | 0.3 | 2.5 | 7.3 | 10.4 | 18.3 | 22.8 | 18.6 |
| 2000..... | 5.1 | * | * | * | 0.1 | 0.3 | 2.4 | 6.9 | 10.8 | 17.8 | 23.1 | 19.2 |
| 1999..... | 5.1 | * | * | * | * | 0.3 | 2.3 | 6.6 | 10.9 | 18.6 | 23.7 | 19.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.0 | * | * | * | * | 0.2 | 2.8 | 10.2 | 13.8 | 19.6 | 25.6 | 20.7 |
| 2003..... | 6.3 | * | * | * | * | 0.3 | 3.1 | 10.9 | 14.7 | 22.1 | 25.7 | 23.5 |
| 2002..... | 6.1 | * | * | * | * | 0.3 | 3.2 | 10.4 | 13.9 | 20.5 | 27.2 | 22.8 |
| 2001..... | 6.1 | * | * | * | * | 0.4 | 3.1 | 10.7 | 13.7 | 22.4 | 25.3 | 25.6 |
| 2000..... | 6.0 | * | * | * | * | 0.3 | 3.3 | 10.2 | 14.6 | 21.5 | 25.7 | 27.5 |
| 1999..... | 6.0 | * | * | * | * | 0.4 | 3.2 | 9.9 | 14.6 | 22.3 | 27.2 | 25.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | * | 0.1 | 1.3 | 4.1 | 7.1 | 14.1 | 19.4 | 15.5 |
| 2003..... | 4.3 | * | * | * | * | 0.2 | 1.5 | 4.2 | 7.3 | 15.5 | 20.9 | 15.1 |
| 2002..... | 4.3 | * | * | * | * | 0.2 | 1.7 | 4.0 | 7.2 | 15.0 | 21.6 | 15.6 |
| 2001..... | 4.3 | * | * | * | * | 0.2 | 1.8 | 4.0 | 7.3 | 14.8 | 21.2 | 15.7 |
| 2000..... | 4.2 | * | * | * | * | 0.3 | 1.6 | 3.7 | 7.2 | 14.8 | 21.4 | 15.8 |
| 1999..... | 4.2 | * | * | * | * | 0.2 | 1.4 | 3.3 | 7.5 | 15.6 | 21.5 | 16.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | * | 0.2 | 2.0 | 7.0 | 10.1 | 17.3 | 23.0 | 17.6 |
| 2003..... | 5.8 | * | * | * | * | 0.2 | 2.4 | 7.3 | 10.7 | 19.4 | 24.1 | 18.4 |
| 2002..... | 5.7 | * | * | * | * | 0.3 | 2.5 | 6.9 | 10.4 | 18.3 | 25.0 | 18.5 |
| 2001..... | 5.6 | * | * | * | * | 0.3 | 2.4 | 6.9 | 10.2 | 19.0 | 23.9 | 19.6 |
| 2000..... | 5.5 | * | * | * | * | 0.2 | 2.4 | 6.6 | 10.7 | 18.3 | 24.2 | 19.8 |
| 1999..... | 5.5 | * | * | * | * | 0.3 | 2.2 | 6.2 | 10.7 | 19.3 | 24.8 | 19.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | * | * | * | 0.2 | 2.8 | 10.0 | 13.5 | 20.2 | 27.0 | 21.6 |
| 2003..... | 6.9 | * | * | * | * | 0.2 | 3.3 | 10.8 | 14.3 | 23.2 | 26.9 | 24.3 |
| 2002..... | 6.6 | * | * | * | * | 0.3 | 3.4 | 10.2 | 13.6 | 21.3 | 28.7 | 24.0 |
| 2001..... | 6.5 | * | * | * | * | 0.3 | 3.0 | 10.0 | 13.2 | 23.2 | 26.5 | 27.2 |
| 2000..... | 6.5 | * | * | * | * | 0.3 | 3.3 | 9.8 | 14.5 | 22.1 | 26.8 | 28.4 |
| 1999..... | 6.5 | * | * | * | * | 0.4 | 3.1 | 9.3 | 14.2 | 23.1 | 28.5 | 27.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | * | * | 1.2 | 4.0 | 6.9 | 14.8 | 20.3 | 15.7 |
| 2003..... | 4.8 | * | * | * | * | 0.2 | 1.5 | 4.0 | 7.3 | 16.2 | 22.2 | 15.8 |
| 2002..... | 4.8 | * | * | * | * | 0.2 | 1.7 | 3.7 | 7.3 | 15.8 | 22.5 | 16.2 |
| 2001..... | 4.7 | * | * | * | * | 0.2 | 1.8 | 3.9 | 7.4 | 15.5 | 22.2 | 16.4 |
| 2000..... | 4.6 | * | * | * | * | 0.2 | 1.5 | 3.5 | 7.2 | 15.1 | 22.4 | 16.3 |
| 1999..... | 4.6 | * | * | * | * | 0.2 | 1.3 | 3.1 | 7.4 | 16.1 | 22.4 | 16.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Fibrosis and cirrhosis of liver (K74) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 2.6 | 9.4 | 13.0 | 12.8 | 11.2 | 11.2 |
| 2003..... | 3.7 | * | * | * | * | 0.4 | 2.4 | 10.0 | 13.4 | 14.4 | 10.6 | 11.0 |
| 2002..... | 3.6 | * | * | * | * | * | 2.4 | 10.6 | 12.7 | 12.5 | 12.8 | 9.1 |
| 2001..... | 3.8 | * | * | * | * | * | 3.0 | 11.0 | 13.0 | 13.9 | 12.7 | 7.5 |
| 2000..... | 3.7 | * | * | * | * | * | 2.8 | 10.1 | 12.5 | 15.4 | 12.2 | 11.8 |
| 1999..... | 3.7 | * | * | * | * | 0.4 | 3.0 | 9.8 | 13.9 | 14.2 | 13.8 | 11.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.8 | * | * | * | * | * | 3.5 | 14.3 | 19.1 | 17.2 | 15.1 | * |
| 2003..... | 4.8 | * | * | * | * | * | 3.0 | 14.9 | 20.8 | 17.2 | 13.3 | * |
| 2002..... | 4.6 | * | * | * | * | * | 2.9 | 15.2 | 19.6 | 17.1 | 12.9 | * |
| 2001..... | 5.2 | * | * | * | * | * | 4.2 | 17.5 | 20.2 | 19.4 | 14.7 | * |
| 2000..... | 4.7 | * | * | * | * | * | 3.5 | 15.3 | 18.8 | 19.6 | 14.8 | * |
| 1999..... | 4.8 | * | * | * | * | * | 4.4 | 15.8 | 19.3 | 18.8 | 16.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | 1.8 | 5.3 | 8.1 | 9.7 | 9.0 | 10.8 |
| 2003..... | 2.7 | * | * | * | * | * | 2.0 | 5.9 | 7.4 | 12.3 | 9.1 | 8.3 |
| 2002..... | 2.7 | * | * | * | * | * | 1.9 | 6.7 | 7.2 | 9.2 | 12.8 | 8.8 |
| 2001..... | 2.5 | * | * | * | * | * | 2.0 | 5.4 | 7.2 | 10.0 | 11.6 | * |
| 2000..... | 2.7 | * | * | * | * | * | 2.2 | 5.6 | 7.4 | 12.4 | 10.8 | 10.5 |
| 1999..... | 2.7 | * | * | * | * | * | 1.7 | 4.8 | 9.5 | 11.0 | 12.3 | 12.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of liver (K71-K72,K75-K76) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | 2.3 | 0.2 | * | 0.2 | 0.4 | 1.8 | 5.0 | 6.9 | 9.1 | 11.3 | 13.1 |
| 2003..... | 3.1 | 1.5 | 0.2 | * | 0.2 | 0.5 | 1.9 | 5.1 | 7.0 | 9.2 | 11.0 | 13.5 |
| 2002..... | 2.9 | 2.1 | 0.2 | 0.1 | 0.1 | 0.4 | 1.9 | 5.0 | 6.2 | 8.6 | 10.5 | 14.3 |
| 2001..... | 2.7 | 1.9 | 0.2 | 0.1 | 0.1 | 0.4 | 1.6 | 4.4 | 6.0 | 8.1 | 11.1 | 13.8 |
| 2000..... | 2.6 | 1.9 | 0.2 | 0.1 | 0.1 | 0.4 | 1.7 | 4.1 | 5.5 | 8.1 | 10.2 | 13.6 |
| 1999..... | 2.6 | 2.1 | 0.1 | 0.1 | 0.1 | 0.4 | 1.7 | 4.0 | 5.7 | 8.8 | 10.3 | 14.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.5 | 2.3 | * | * | 0.2 | 0.5 | 2.1 | 6.9 | 8.9 | 10.5 | 12.8 | 15.1 |
| 2003..... | 3.6 | 2.1 | 0.2 | * | 0.1 | 0.5 | 2.2 | 7.1 | 9.1 | 11.2 | 12.7 | 16.9 |
| 2002..... | 3.5 | 2.8 | * | * | 0.1 | 0.5 | 2.3 | 7.1 | 8.1 | 11.0 | 13.3 | 18.9 |
| 2001..... | 3.2 | 2.2 | * | * | 0.1 | 0.3 | 2.1 | 6.2 | 7.8 | 10.0 | 14.1 | 17.4 |
| 2000..... | 3.0 | 2.5 | * | * | 0.1 | 0.4 | 2.2 | 5.6 | 6.9 | 9.7 | 12.6 | 17.8 |
| 1999..... | 3.1 | 2.7 | * | * | 0.1 | 0.5 | 2.1 | 5.5 | 7.5 | 10.8 | 12.5 | 19.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.6 | 2.3 | * | * | 0.2 | 0.4 | 1.5 | 3.0 | 4.9 | 8.0 | 10.2 | 12.3 |
| 2003..... | 2.6 | * | * | * | 0.2 | 0.5 | 1.7 | 3.3 | 5.0 | 7.6 | 9.8 | 12.1 |
| 2002..... | 2.3 | 1.3 | * | * | 0.1 | 0.4 | 1.5 | 3.0 | 4.4 | 6.6 | 8.6 | 12.3 |
| 2001..... | 2.3 | 1.5 | * | * | 0.1 | 0.4 | 1.2 | 2.7 | 4.4 | 6.6 | 9.2 | 12.3 |
| 2000..... | 2.2 | 1.3 | * | * | * | 0.3 | 1.2 | 2.7 | 4.3 | 6.8 | 8.6 | 11.9 |
| 1999..... | 2.2 | 1.5 | * | * | 0.2 | 0.3 | 1.3 | 2.5 | 4.0 | 7.2 | 8.8 | 11.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | 0.4 | 1.6 | 5.4 | 7.5 | 12.8 | 13.3 | 12.4 |
| 2003..... | 2.3 | * | * | * | * | 0.5 | 1.8 | 5.7 | 9.9 | 12.4 | 19.2 | 17.1 |
| 2002..... | 1.9 | 2.4 | * | * | * | 0.5 | 1.4 | 5.0 | 7.7 | 11.8 | 14.0 | 11.6 |
| 2001..... | 1.8 | * | * | * | * | * | 1.0 | 4.4 | 7.4 | 12.1 | 17.1 | 18.8 |
| 2000..... | 1.7 | * | * | * | * | * | 1.7 | 4.7 | 6.0 | 11.2 | 13.8 | 18.6 |
| 1999..... | 1.9 | * | * | * | * | 0.5 | 1.6 | 5.2 | 8.2 | 12.0 | 14.0 | 21.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | 2.2 | 8.5 | 9.6 | 13.8 | 11.4 | * |
| 2003..... | 2.6 | * | * | * | * | * | 2.1 | 8.1 | 12.6 | 14.2 | 20.7 | * |
| 2002..... | 2.3 | * | * | * | * | 0.6 | 2.1 | 7.5 | 9.9 | 14.9 | 18.5 | * |
| 2001..... | 2.0 | * | * | * | * | * | 1.3 | 6.6 | 9.0 | 14.6 | 20.6 | * |
| 2000..... | 2.0 | * | * | * | * | * | 2.6 | 6.4 | 8.6 | 12.5 | 16.2 | * |
| 1999..... | 2.4 | * | * | * | * | 0.6 | 2.3 | 7.8 | 11.7 | 15.4 | 15.6 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 0.9 | 2.3 | 5.6 | 12.1 | 14.6 | * |
| 2003..... | 1.9 | * | * | * | * | * | 1.4 | 3.2 | 7.4 | 11.0 | 18.2 | * |
| 2002..... | 1.4 | * | * | * | * | * | * | 2.5 | 5.8 | 9.3 | 11.0 | * |
| 2001..... | 1.5 | * | * | * | * | * | 0.8 | 2.2 | 6.0 | 10.1 | 14.8 | * |
| 2000..... | 1.4 | * | * | * | * | * | 0.8 | 3.1 | 3.6 | 10.3 | 12.2 | * |
| 1999..... | 1.5 | * | * | * | * | * | * | 2.7 | 5.2 | 9.3 | 12.8 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of liver (K71-K72,K75-K76) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.2 | 2.4 | 0.2 | * | 0.2 | 0.4 | 1.8 | 4.9 | 6.8 | 8.8 | 11.1 | 13.2 |
| 2003..... | 3.2 | 1.5 | 0.2 | * | 0.2 | 0.5 | 2.0 | 5.1 | 6.7 | 9.0 | 10.6 | 13.4 |
| 2002..... | 3.1 | 1.9 | * | 0.1 | 0.1 | 0.4 | 2.0 | 5.0 | 6.0 | 8.3 | 10.3 | 14.4 |
| 2001..... | 2.9 | 1.7 | 0.2 | 0.1 | 0.1 | 0.4 | 1.7 | 4.4 | 5.9 | 7.8 | 10.8 | 13.6 |
| 2000..... | 2.7 | 2.1 | 0.2 | * | 0.1 | 0.4 | 1.7 | 4.1 | 5.5 | 7.9 | 10.0 | 13.4 |
| 1999..... | 2.7 | 2.1 | * | * | 0.1 | 0.4 | 1.7 | 3.8 | 5.5 | 8.6 | 10.1 | 13.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.7 | 2.4 | * | * | 0.1 | 0.5 | 2.1 | 6.8 | 8.8 | 10.2 | 12.8 | 15.2 |
| 2003..... | 3.8 | 2.1 | * | * | 0.1 | 0.5 | 2.2 | 6.9 | 8.8 | 11.0 | 12.3 | 16.4 |
| 2002..... | 3.7 | 2.7 | * | * | * | 0.4 | 2.4 | 7.0 | 7.9 | 10.7 | 13.0 | 18.9 |
| 2001..... | 3.4 | 2.1 | * | * | 0.1 | 0.4 | 2.2 | 6.1 | 7.6 | 9.6 | 13.8 | 17.0 |
| 2000..... | 3.1 | 2.8 | * | * | 0.1 | 0.4 | 2.1 | 5.5 | 6.7 | 9.5 | 12.5 | 17.4 |
| 1999..... | 3.2 | 2.8 | * | * | * | 0.5 | 2.1 | 5.2 | 7.2 | 10.5 | 12.3 | 18.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.8 | 2.4 | * | * | 0.2 | 0.4 | 1.6 | 3.1 | 4.9 | 7.7 | 10.0 | 12.3 |
| 2003..... | 2.7 | * | * | * | 0.2 | 0.5 | 1.7 | 3.3 | 4.8 | 7.3 | 9.4 | 12.0 |
| 2002..... | 2.5 | * | * | * | 0.1 | 0.4 | 1.6 | 3.1 | 4.2 | 6.4 | 8.5 | 12.4 |
| 2001..... | 2.4 | 1.4 | * | * | 0.1 | 0.4 | 1.2 | 2.7 | 4.3 | 6.3 | 8.9 | 12.2 |
| 2000..... | 2.3 | * | * | * | * | 0.4 | 1.3 | 2.6 | 4.3 | 6.6 | 8.4 | 11.8 |
| 1999..... | 2.3 | * | * | * | 0.1 | 0.3 | 1.3 | 2.5 | 4.0 | 7.0 | 8.6 | 11.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | 1.6 | * | * | 0.1 | 0.4 | 1.7 | 4.6 | 6.4 | 8.6 | 11.2 | 13.2 |
| 2003..... | 3.3 | 1.2 | * | * | 0.1 | 0.4 | 1.9 | 4.8 | 6.4 | 8.8 | 10.6 | 13.2 |
| 2002..... | 3.1 | 1.5 | * | * | 0.1 | 0.4 | 1.9 | 4.7 | 5.6 | 8.3 | 10.2 | 14.5 |
| 2001..... | 2.9 | 1.2 | * | * | 0.1 | 0.3 | 1.6 | 4.1 | 5.5 | 7.7 | 10.7 | 13.6 |
| 2000..... | 2.7 | 1.7 | * | * | 0.1 | 0.3 | 1.6 | 3.6 | 5.0 | 7.8 | 9.9 | 13.3 |
| 1999..... | 2.7 | 1.3 | * | * | 0.1 | 0.3 | 1.5 | 3.4 | 5.0 | 8.3 | 10.1 | 13.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | 0.5 | 2.0 | 6.3 | 8.2 | 10.1 | 12.8 | 16.1 |
| 2003..... | 3.8 | 1.7 | * | * | * | 0.4 | 2.2 | 6.5 | 8.2 | 10.6 | 12.5 | 16.1 |
| 2002..... | 3.8 | 2.1 | * | * | * | 0.4 | 2.3 | 6.6 | 7.4 | 10.7 | 12.7 | 19.2 |
| 2001..... | 3.5 | * | * | * | * | 0.3 | 2.0 | 5.8 | 7.2 | 9.6 | 13.3 | 16.9 |
| 2000..... | 3.1 | 2.1 | * | * | * | 0.3 | 2.0 | 4.9 | 6.1 | 9.3 | 12.5 | 17.2 |
| 1999..... | 3.2 | * | * | * | * | 0.4 | 1.9 | 4.7 | 6.6 | 10.3 | 12.5 | 19.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | 0.3 | 1.5 | 2.8 | 4.6 | 7.4 | 10.0 | 11.9 |
| 2003..... | 2.8 | * | * | * | * | 0.4 | 1.6 | 3.1 | 4.7 | 7.3 | 9.3 | 11.9 |
| 2002..... | 2.5 | * | * | * | * | 0.3 | 1.5 | 2.8 | 4.0 | 6.2 | 8.5 | 12.4 |
| 2001..... | 2.4 | * | * | * | * | 0.4 | 1.2 | 2.5 | 4.0 | 6.1 | 8.9 | 12.2 |
| 2000..... | 2.3 | * | * | * | * | 0.3 | 1.1 | 2.3 | 4.0 | 6.5 | 8.2 | 11.8 |
| 1999..... | 2.3 | * | * | * | * | 0.2 | 1.1 | 2.1 | 3.5 | 6.7 | 8.5 | 11.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of liver (K71-K72,K75-K76) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | 5.4 | * | * | 0.3 | 0.6 | 2.5 | 7.7 | 10.3 | 10.9 | 11.0 | 12.4 |
| 2003..... | 3.3 | * | * | * | 0.5 | 0.9 | 2.3 | 7.8 | 10.1 | 11.4 | 9.9 | 15.2 |
| 2002..... | 3.1 | 3.7 | * | * | * | 0.6 | 2.5 | 7.8 | 9.1 | 10.0 | 11.1 | 15.1 |
| 2001..... | 3.0 | 3.6 | * | * | * | 0.6 | 2.5 | 6.9 | 9.8 | 10.2 | 12.8 | 11.5 |
| 2000..... | 3.1 | 4.2 | * | * | * | 0.8 | 2.7 | 7.5 | 9.4 | 10.1 | 11.0 | 12.4 |
| 1999..... | 3.1 | 5.1 | * | * | * | 0.7 | 2.8 | 7.2 | 9.8 | 10.9 | 10.4 | 13.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | * | 0.8 | 2.8 | 10.5 | 15.3 | 13.3 | 12.8 | * |
| 2003..... | 4.0 | * | * | * | * | 0.8 | 2.1 | 10.9 | 14.9 | 16.9 | 11.8 | * |
| 2002..... | 3.7 | * | * | * | * | * | 2.5 | 10.8 | 13.2 | 12.4 | 15.2 | * |
| 2001..... | 3.7 | * | * | * | * | * | 3.2 | 9.8 | 12.8 | 12.2 | 18.9 | * |
| 2000..... | 3.7 | * | * | * | * | 0.8 | 3.1 | 10.5 | 12.8 | 13.0 | 12.6 | * |
| 1999..... | 3.5 | 7.3 | * | * | * | * | 3.2 | 10.0 | 12.9 | 12.9 | 9.7 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | 2.2 | 5.2 | 6.3 | 9.2 | 10.0 | 14.9 |
| 2003..... | 2.7 | * | * | * | * | 0.9 | 2.4 | 5.2 | 6.3 | 7.4 | 8.8 | 13.6 |
| 2002..... | 2.6 | * | * | * | * | * | 2.5 | 5.3 | 5.8 | 8.3 | 8.8 | 13.4 |
| 2001..... | 2.5 | * | * | * | * | * | 1.8 | 4.5 | 7.3 | 8.9 | 9.4 | 10.3 |
| 2000..... | 2.5 | * | * | * | * | * | 2.3 | 4.8 | 6.7 | 8.0 | 10.1 | 10.9 |
| 1999..... | 2.7 | * | * | * | * | 0.8 | 2.6 | 4.8 | 7.4 | 9.4 | 10.9 | 13.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cholelithiasis and other disorders of gallbladder (K80-K82) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 7.8 | 23.8 |
| 2003..... | 1.0 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 2.7 | 8.2 | 21.8 |
| 2002..... | 1.0 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 2.8 | 7.8 | 22.9 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.9 | 7.5 | 23.7 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.5 | 24.8 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.6 | 7.8 | 23.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 2.9 | 9.3 | 25.5 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 3.2 | 10.1 | 22.7 |
| 2002..... | 0.9 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.1 | 9.8 | 24.1 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 3.2 | 9.3 | 26.9 |
| 2000..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.9 | 9.4 | 27.6 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.0 | 9.3 | 27.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 6.8 | 23.0 |
| 2003..... | 1.1 | * | * | * | * | * | * | 0.3 | 0.5 | 2.2 | 6.9 | 21.4 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.5 | 6.4 | 22.4 |
| 2001..... | 1.1 | * | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 6.3 | 22.4 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.1 | 6.2 | 23.7 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.2 | 6.8 | 22.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 1.0 | 2.8 | 9.6 | 27.1 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 3.6 | 9.5 | 18.7 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 1.3 | 3.6 | 9.2 | 24.4 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 1.2 | 4.5 | 8.5 | 31.3 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 3.8 | 10.1 | 29.2 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 1.2 | 3.3 | 11.9 | 32.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 12.2 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 4.4 | 13.8 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 4.0 | 9.4 | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | 5.0 | 11.2 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 5.3 | 15.7 | * |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 15.1 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 2.8 | 7.9 | 31.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | 2.9 | 6.6 | 19.5 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 3.2 | 9.0 | 23.2 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 4.1 | 6.6 | 30.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | 31.7 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | * | 9.7 | 32.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cholelithiasis and other disorders of gallbladder (K80-K82) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.7 | 23.6 |
| 2003..... | 1.1 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 2.6 | 8.1 | 21.9 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.7 | 7.7 | 22.8 |
| 2001..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.8 | 7.5 | 23.4 |
| 2000..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.3 | 7.4 | 24.6 |
| 1999..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 7.6 | 23.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.9 | 9.1 | 25.8 |
| 2003..... | 1.0 | * | * | * | * | * | * | 0.3 | 0.8 | 3.1 | 9.9 | 23.0 |
| 2002..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 3.0 | 9.8 | 23.9 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.1 | 9.2 | 26.6 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.7 | 9.2 | 27.7 |
| 1999..... | 1.0 | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.0 | 9.0 | 27.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.0 | 6.7 | 22.6 |
| 2003..... | 1.2 | * | * | * | * | * | * | 0.3 | 0.5 | 2.2 | 6.9 | 21.4 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.4 | 6.3 | 22.4 |
| 2001..... | 1.2 | * | * | * | * | * | * | 0.3 | 0.8 | 2.5 | 6.3 | 22.1 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.0 | 6.2 | 23.4 |
| 1999..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 6.6 | 22.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.3 | 7.5 | 24.0 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.6 | 8.1 | 22.0 |
| 2002..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.6 | 7.5 | 23.4 |
| 2001..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.6 | 7.3 | 23.6 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.2 | 7.3 | 25.3 |
| 1999..... | 1.2 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.4 | 7.6 | 24.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 0.3 | 0.9 | 2.9 | 9.1 | 27.0 |
| 2003..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.7 | 3.1 | 9.8 | 23.3 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.3 | 0.9 | 3.0 | 9.8 | 24.5 |
| 2001..... | 1.1 | * | * | * | * | * | * | 0.4 | 1.0 | 2.9 | 9.2 | 26.4 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.8 | 2.6 | 9.3 | 28.9 |
| 1999..... | 1.1 | * | * | * | * | * | * | 0.3 | 0.9 | 2.9 | 9.3 | 27.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 6.5 | 22.7 |
| 2003..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.5 | 2.1 | 6.9 | 21.4 |
| 2002..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.6 | 2.3 | 6.1 | 22.9 |
| 2001..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 6.1 | 22.4 |
| 2000..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 6.0 | 23.9 |
| 1999..... | 1.3 | * | * | * | * | * | * | 0.3 | 0.6 | 1.9 | 6.5 | 22.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Cholelithiasis and other disorders of gallbladder (K80-K82) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.8 | 1.6 | 3.6 | 9.3 | 19.6 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.7 | 1.2 | 3.0 | 8.1 | 21.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.6 | 1.1 | 3.3 | 9.6 | 15.1 |
| 2001..... | 0.9 | * | * | * | * | * | * | 0.7 | 1.7 | 4.8 | 8.3 | 20.0 |
| 2000..... | 0.8 | * | * | * | * | * | * | 0.7 | 1.2 | 3.8 | 8.3 | 18.1 |
| 1999..... | 0.7 | * | * | * | * | * | 0.4 | 0.5 | 1.5 | 3.7 | 7.3 | 15.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 1.0 | 1.6 | 3.7 | 9.1 | * |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | 3.3 | 10.7 | 21.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | * | 10.8 | * |
| 2001..... | 0.8 | * | * | * | * | * | * | * | 2.2 | 5.0 | 6.3 | 27.7 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 3.4 | 8.0 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 3.8 | 7.2 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | 1.6 | 3.6 | 9.3 | 21.7 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.8 | * | 2.8 | 6.7 | 21.0 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | 3.8 | 9.0 | 16.3 |
| 2001..... | 1.0 | * | * | * | * | * | * | * | * | 4.6 | 9.4 | 17.1 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | * | 4.0 | 8.5 | 19.6 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 3.6 | 7.4 | 15.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of biliary tract and pancreas (K83-K86) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.2 | 4.7 | 9.4 | 20.8 |
| 2003..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.3 | 4.8 | 10.0 | 21.9 |
| 2002..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.5 | 4.8 | 9.7 | 21.7 |
| 2001..... | 1.7 | * | * | * | 0.1 | 0.4 | 0.9 | 1.6 | 2.5 | 5.1 | 9.4 | 22.9 |
| 2000..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.5 | 4.6 | 9.4 | 22.0 |
| 1999..... | 1.6 | * | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.4 | 4.5 | 8.5 | 22.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | 0.1 | 0.5 | 1.1 | 2.0 | 3.0 | 5.7 | 10.6 | 21.1 |
| 2003..... | 1.8 | * | * | * | * | 0.4 | 1.0 | 1.9 | 3.1 | 6.1 | 11.8 | 20.7 |
| 2002..... | 1.9 | * | * | * | 0.2 | 0.5 | 1.1 | 2.0 | 3.4 | 6.4 | 11.9 | 21.2 |
| 2001..... | 1.8 | * | * | * | 0.1 | 0.5 | 1.1 | 2.0 | 3.3 | 6.5 | 10.2 | 22.4 |
| 2000..... | 1.8 | * | * | * | 0.1 | 0.5 | 1.2 | 1.9 | 3.0 | 5.9 | 11.4 | 22.9 |
| 1999..... | 1.7 | * | * | * | 0.1 | 0.5 | 1.1 | 1.9 | 3.1 | 5.6 | 10.1 | 23.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | 0.1 | 0.5 | 0.9 | 1.5 | 3.8 | 8.6 | 20.7 |
| 2003..... | 1.6 | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.5 | 3.7 | 8.8 | 22.4 |
| 2002..... | 1.5 | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.6 | 3.5 | 8.3 | 21.9 |
| 2001..... | 1.7 | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.8 | 3.9 | 8.8 | 23.2 |
| 2000..... | 1.6 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.0 | 3.5 | 8.0 | 21.6 |
| 1999..... | 1.5 | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 1.7 | 3.6 | 7.4 | 21.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.6 | 3.7 | 9.6 | 18.1 |
| 2003..... | 0.7 | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.0 | 3.8 | 10.1 | 18.1 |
| 2002..... | 0.8 | * | * | * | * | 0.3 | 0.6 | 1.2 | 1.5 | 4.7 | 8.5 | 22.7 |
| 2001..... | 0.7 | * | * | * | * | 0.3 | 0.6 | 0.9 | 1.9 | 4.8 | 10.8 | 18.8 |
| 2000..... | 0.8 | * | * | * | * | * | 0.8 | 1.1 | 1.9 | 4.0 | 10.7 | 27.2 |
| 1999..... | 0.8 | * | * | * | * | 0.4 | 0.7 | 1.0 | 2.3 | 5.1 | 8.3 | 30.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | 0.5 | 0.8 | 1.3 | 2.4 | 4.4 | 10.7 | * |
| 2003..... | 0.9 | * | * | * | * | * | 0.8 | 1.5 | 3.0 | 5.2 | 15.0 | * |
| 2002..... | 0.9 | * | * | * | * | 0.5 | 0.7 | 1.4 | 2.6 | 5.9 | 13.1 | * |
| 2001..... | 0.7 | * | * | * | * | * | 0.8 | * | 2.4 | 5.4 | 9.4 | * |
| 2000..... | 0.9 | * | * | * | * | * | 1.3 | 1.5 | * | 4.6 | 14.3 | * |
| 1999..... | 0.9 | * | * | * | * | * | 0.8 | 1.7 | * | 6.8 | 10.9 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | 8.9 | 18.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | * | 6.8 | 21.1 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 3.6 | * | 25.7 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 4.3 | 11.8 | 24.2 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 3.5 | 8.3 | 23.7 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | 2.3 | 3.8 | * | 30.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of biliary tract and pancreas (K83-K86) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 2.3 | 4.7 | 9.4 | 20.9 |
| 2003..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.3 | 4.9 | 10.0 | 22.0 |
| 2002..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.6 | 4.8 | 9.8 | 21.6 |
| 2001..... | 1.9 | * | * | * | 0.1 | 0.4 | 0.9 | 1.6 | 2.6 | 5.1 | 9.3 | 23.0 |
| 2000..... | 1.8 | * | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.5 | 4.6 | 9.3 | 21.8 |
| 1999..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 2.4 | 4.4 | 8.4 | 21.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.5 | 1.2 | 2.0 | 3.1 | 5.8 | 10.6 | 21.2 |
| 2003..... | 1.9 | * | * | * | * | 0.4 | 1.1 | 1.9 | 3.1 | 6.2 | 11.6 | 20.9 |
| 2002..... | 2.0 | * | * | * | 0.1 | 0.5 | 1.1 | 2.1 | 3.5 | 6.4 | 11.8 | 21.3 |
| 2001..... | 2.0 | * | * | * | * | 0.5 | 1.2 | 2.1 | 3.3 | 6.5 | 10.2 | 22.9 |
| 2000..... | 1.9 | * | * | * | * | 0.5 | 1.2 | 1.9 | 3.1 | 5.9 | 11.3 | 22.4 |
| 1999..... | 1.8 | * | * | * | * | 0.5 | 1.1 | 1.9 | 3.2 | 5.5 | 10.0 | 22.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.1 | 0.5 | 1.0 | 1.5 | 3.8 | 8.6 | 20.8 |
| 2003..... | 1.8 | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.6 | 3.8 | 8.9 | 22.5 |
| 2002..... | 1.7 | * | * | * | * | 0.2 | 0.6 | 0.8 | 1.7 | 3.5 | 8.4 | 21.7 |
| 2001..... | 1.8 | * | * | * | * | 0.3 | 0.6 | 1.1 | 1.9 | 3.9 | 8.7 | 23.1 |
| 2000..... | 1.7 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.0 | 3.5 | 8.0 | 21.5 |
| 1999..... | 1.6 | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.7 | 3.6 | 7.4 | 21.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 2.0 | 4.6 | 9.4 | 21.1 |
| 2003..... | 1.9 | * | * | * | * | 0.3 | 0.7 | 1.2 | 2.1 | 4.6 | 10.2 | 22.0 |
| 2002..... | 1.9 | * | * | * | 0.1 | 0.3 | 0.8 | 1.2 | 2.4 | 4.8 | 9.9 | 21.8 |
| 2001..... | 2.0 | * | * | * | * | 0.3 | 0.8 | 1.4 | 2.4 | 5.0 | 9.3 | 23.1 |
| 2000..... | 1.8 | * | * | * | * | 0.3 | 0.8 | 1.3 | 2.3 | 4.3 | 9.2 | 21.7 |
| 1999..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 2.2 | 4.4 | 8.3 | 21.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | * | 0.4 | 1.1 | 1.8 | 2.7 | 5.7 | 10.6 | 21.5 |
| 2003..... | 2.0 | * | * | * | * | 0.4 | 0.8 | 1.7 | 2.7 | 5.9 | 11.7 | 20.7 |
| 2002..... | 2.1 | * | * | * | * | 0.4 | 1.0 | 1.8 | 3.2 | 6.3 | 12.0 | 21.2 |
| 2001..... | 2.0 | * | * | * | * | 0.4 | 1.1 | 1.8 | 3.1 | 6.5 | 10.3 | 22.8 |
| 2000..... | 1.9 | * | * | * | * | 0.4 | 1.0 | 1.7 | 2.8 | 5.6 | 11.2 | 21.5 |
| 1999..... | 1.8 | * | * | * | * | 0.4 | 1.0 | 1.6 | 2.8 | 5.6 | 9.9 | 21.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | 0.4 | 0.9 | 1.3 | 3.7 | 8.5 | 20.9 |
| 2003..... | 1.9 | * | * | * | * | * | 0.5 | 0.7 | 1.5 | 3.5 | 9.1 | 22.6 |
| 2002..... | 1.8 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.6 | 3.5 | 8.5 | 22.0 |
| 2001..... | 1.9 | * | * | * | * | 0.2 | 0.5 | 1.0 | 1.8 | 3.7 | 8.7 | 23.2 |
| 2000..... | 1.8 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 3.3 | 7.9 | 21.7 |
| 1999..... | 1.7 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.5 | 3.4 | 7.2 | 21.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of biliary tract and pancreas (K83-K86) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.6 | 1.6 | 3.0 | 4.9 | 6.5 | 10.7 | 20.7 |
| 2003..... | 1.9 | * | * | * | * | 0.6 | 1.9 | 3.2 | 4.6 | 7.5 | 8.9 | 23.8 |
| 2002..... | 1.8 | * | * | * | * | 0.6 | 1.5 | 3.1 | 4.6 | 6.1 | 8.8 | 21.8 |
| 2001..... | 2.0 | * | * | * | * | 0.8 | 2.0 | 3.8 | 4.2 | 6.7 | 9.5 | 22.2 |
| 2000..... | 1.9 | * | * | * | * | 0.8 | 1.5 | 3.0 | 4.6 | 7.4 | 10.1 | 23.8 |
| 1999..... | 1.8 | * | * | * | * | 0.7 | 1.5 | 3.4 | 4.8 | 5.5 | 10.0 | 23.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | 1.0 | 2.0 | 4.4 | 6.9 | 8.7 | 12.3 | * |
| 2003..... | 2.3 | * | * | * | * | * | 2.8 | 4.2 | 6.9 | 9.3 | 11.5 | 26.6 |
| 2002..... | 2.2 | * | * | * | * | 0.9 | 2.0 | 4.7 | 6.6 | 8.6 | 10.0 | 26.3 |
| 2001..... | 2.2 | * | * | * | * | 1.1 | 2.4 | 5.2 | 6.0 | 8.8 | 10.2 | * |
| 2000..... | 2.2 | * | * | * | * | 1.1 | 2.2 | 4.2 | 6.1 | 10.3 | 12.0 | 32.9 |
| 1999..... | 2.2 | * | * | * | * | 1.2 | 2.2 | 5.3 | 6.6 | 6.2 | 10.9 | 33.3 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | 1.2 | 1.8 | 3.3 | 4.9 | 9.8 | 21.3 |
| 2003..... | 1.6 | * | * | * | * | * | 1.1 | 2.3 | 2.8 | 6.3 | 7.3 | 22.7 |
| 2002..... | 1.4 | * | * | * | * | * | 1.0 | 1.7 | 3.0 | 4.3 | 8.1 | 20.1 |
| 2001..... | 1.7 | * | * | * | * | * | 1.6 | 2.5 | 2.7 | 5.3 | 9.1 | 22.7 |
| 2000..... | 1.6 | * | * | * | * | * | 0.8 | 2.0 | 3.4 | 5.3 | 9.0 | 20.5 |
| 1999..... | 1.5 | * | * | * | * | * | 0.9 | 1.7 | 3.3 | 5.0 | 9.5 | 19.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pancreatitis (K85,K86.0-K86.1) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 1.8 | 3.5 | 5.9 | 11.5 |
| 2003..... | 1.3 | * | * | * | * | 0.3 | 0.7 | 1.2 | 1.8 | 3.5 | 6.6 | 11.9 |
| 2002..... | 1.3 | * | * | * | 0.1 | 0.3 | 0.7 | 1.2 | 2.1 | 3.4 | 6.6 | 11.8 |
| 2001..... | 1.4 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.2 | 3.8 | 6.3 | 13.4 |
| 2000..... | 1.3 | * | * | * | 0.1 | 0.3 | 0.8 | 1.2 | 2.0 | 3.4 | 6.3 | 12.0 |
| 1999..... | 1.3 | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 1.9 | 3.4 | 5.8 | 12.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.5 | 1.1 | 1.9 | 2.5 | 4.4 | 6.5 | 10.4 |
| 2003..... | 1.4 | * | * | * | * | 0.4 | 1.0 | 1.7 | 2.5 | 4.5 | 7.6 | 10.7 |
| 2002..... | 1.5 | * | * | * | 0.1 | 0.4 | 1.0 | 1.9 | 2.9 | 4.6 | 7.7 | 11.1 |
| 2001..... | 1.5 | * | * | * | * | 0.4 | 1.1 | 1.8 | 2.8 | 4.9 | 6.9 | 13.3 |
| 2000..... | 1.4 | * | * | * | * | 0.4 | 1.1 | 1.6 | 2.5 | 4.6 | 7.3 | 11.4 |
| 1999..... | 1.4 | * | * | * | * | 0.5 | 1.0 | 1.7 | 2.5 | 4.4 | 6.8 | 12.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.1 | 0.4 | 0.8 | 1.2 | 2.7 | 5.4 | 11.9 |
| 2003..... | 1.2 | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.2 | 2.7 | 6.0 | 12.4 |
| 2002..... | 1.1 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.4 | 2.4 | 5.9 | 12.1 |
| 2001..... | 1.3 | * | * | * | * | 0.2 | 0.6 | 0.9 | 1.5 | 2.9 | 6.0 | 13.4 |
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.6 | 2.5 | 5.6 | 12.2 |
| 1999..... | 1.1 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.3 | 2.7 | 5.2 | 12.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | 0.3 | 0.7 | 1.2 | 1.6 | 3.4 | 5.8 | 11.4 |
| 2003..... | 1.3 | * | * | * | * | 0.2 | 0.6 | 1.0 | 1.7 | 3.3 | 6.8 | 11.7 |
| 2002..... | 1.4 | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.0 | 3.4 | 6.8 | 12.0 |
| 2001..... | 1.4 | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.0 | 3.8 | 6.4 | 13.4 |
| 2000..... | 1.3 | * | * | * | * | 0.3 | 0.7 | 1.1 | 1.8 | 3.3 | 6.3 | 12.0 |
| 1999..... | 1.3 | * | * | * | * | 0.3 | 0.7 | 1.0 | 1.7 | 3.4 | 5.7 | 12.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | 0.4 | 1.0 | 1.6 | 2.2 | 4.2 | 6.4 | 10.5 |
| 2003..... | 1.4 | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.2 | 4.2 | 7.6 | 10.4 |
| 2002..... | 1.5 | * | * | * | * | 0.3 | 0.9 | 1.6 | 2.6 | 4.6 | 7.8 | 10.8 |
| 2001..... | 1.5 | * | * | * | * | 0.3 | 0.9 | 1.5 | 2.6 | 5.0 | 6.9 | 13.4 |
| 2000..... | 1.4 | * | * | * | * | 0.4 | 1.0 | 1.4 | 2.3 | 4.3 | 7.2 | 10.9 |
| 1999..... | 1.3 | * | * | * | * | 0.4 | 0.9 | 1.3 | 2.3 | 4.3 | 6.8 | 11.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.3 | 0.7 | 1.0 | 2.7 | 5.5 | 11.8 |
| 2003..... | 1.2 | * | * | * | * | * | 0.4 | 0.6 | 1.1 | 2.5 | 6.2 | 12.3 |
| 2002..... | 1.2 | * | * | * | * | * | 0.5 | 0.5 | 1.3 | 2.4 | 6.1 | 12.4 |
| 2001..... | 1.3 | * | * | * | * | * | 0.5 | 0.8 | 1.5 | 2.8 | 6.1 | 13.4 |
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.4 | 2.4 | 5.6 | 12.5 |
| 1999..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.1 | 2.6 | 5.0 | 13.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pancreatitis (K85,K86.0-K86.1) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | 0.5 | 1.4 | 2.7 | 4.1 | 4.8 | 7.2 | 13.5 |
| 2003..... | 1.5 | * | * | * | * | 0.5 | 1.7 | 2.8 | 3.7 | 5.4 | 5.8 | 14.3 |
| 2002..... | 1.3 | * | * | * | * | 0.5 | 1.3 | 2.7 | 3.9 | 3.8 | 5.3 | 11.5 |
| 2001..... | 1.5 | * | * | * | * | 0.7 | 1.7 | 3.4 | 3.4 | 4.7 | 6.5 | 13.4 |
| 2000..... | 1.5 | * | * | * | * | 0.7 | 1.4 | 2.6 | 4.0 | 5.5 | 6.2 | 12.1 |
| 1999..... | 1.5 | * | * | * | * | 0.6 | 1.4 | 3.1 | 3.8 | 4.4 | 7.2 | 12.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.8 | 1.8 | 4.0 | 5.9 | 7.0 | 8.8 | * |
| 2003..... | 1.9 | * | * | * | * | * | 2.5 | 3.9 | 6.1 | 7.4 | 7.8 | * |
| 2002..... | 1.7 | * | * | * | * | * | 1.8 | 4.1 | 5.7 | 5.3 | 6.7 | * |
| 2001..... | 1.8 | * | * | * | * | 0.9 | 2.2 | 4.6 | 5.3 | 5.8 | 7.5 | * |
| 2000..... | 1.8 | * | * | * | * | 1.0 | 2.2 | 3.5 | 5.3 | 8.2 | 8.0 | * |
| 1999..... | 1.9 | * | * | * | * | 1.1 | 2.2 | 4.9 | 5.3 | 5.1 | 7.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 1.0 | 1.7 | 2.7 | 3.3 | 6.3 | 13.7 |
| 2003..... | 1.1 | * | * | * | * | * | 1.0 | 1.8 | 1.8 | 4.0 | 4.7 | 13.6 |
| 2002..... | 1.0 | * | * | * | * | * | 0.8 | 1.5 | 2.4 | 2.7 | 4.5 | 9.6 |
| 2001..... | 1.3 | * | * | * | * | * | 1.3 | 2.3 | 2.0 | 4.0 | 5.9 | 14.1 |
| 2000..... | 1.1 | * | * | * | * | * | 0.7 | 1.8 | 2.9 | 3.6 | 5.2 | 10.0 |
| 1999..... | 1.1 | * | * | * | * | * | 0.8 | 1.6 | 2.6 | 3.8 | 6.8 | 10.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other disorders of biliary tract and pancreas (K83,K86.2-K86.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 3.5 | 9.5 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 3.3 | 10.2 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 3.2 | 9.8 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.3 | 2.9 | 9.7 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 3.0 | 9.8 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.6 | 9.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.1 | 10.8 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.6 | 1.7 | 4.0 | 10.3 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 4.1 | 10.2 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.6 | 3.4 | 9.6 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.4 | 0.5 | 1.3 | 4.0 | 11.0 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.2 | 10.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.1 | 8.9 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 2.8 | 10.1 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 2.5 | 9.7 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.7 | 9.7 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.4 | 9.3 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.2 | 8.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.5 | 9.6 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 3.4 | 10.3 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.3 | 3.1 | 9.8 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.9 | 9.7 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.0 | 9.6 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.6 | 8.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.2 | 11.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.5 | 1.7 | 4.1 | 10.4 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 4.2 | 10.4 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.5 | 1.5 | 3.4 | 9.4 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 4.0 | 10.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 3.2 | 9.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 3.1 | 9.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 0.4 | 1.0 | 2.9 | 10.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.1 | 2.5 | 9.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.6 | 9.8 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.3 | 9.3 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.2 | 8.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of digestive system (K90,K92) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | 0.7 | * | * | * | 0.1 | 0.3 | 0.9 | 2.2 | 5.6 | 18.2 | 68.5 |
| 2003..... | 2.8 | 0.7 | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 6.1 | 19.1 | 74.2 |
| 2002..... | 2.9 | 0.6 | 0.1 | * | * | 0.1 | 0.3 | 1.0 | 2.1 | 6.1 | 19.7 | 76.2 |
| 2001..... | 2.9 | 0.6 | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 6.2 | 19.9 | 77.8 |
| 2000..... | 2.9 | 0.6 | * | * | * | 0.1 | 0.3 | 0.9 | 2.1 | 5.9 | 20.6 | 80.6 |
| 1999..... | 2.8 | 0.7 | 0.1 | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 6.2 | 19.6 | 77.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | * | 0.3 | 1.3 | 2.9 | 7.4 | 21.8 | 69.8 |
| 2003..... | 2.6 | * | * | * | * | * | 0.5 | 1.2 | 2.8 | 7.7 | 23.6 | 73.3 |
| 2002..... | 2.6 | * | * | * | * | 0.1 | 0.4 | 1.4 | 2.9 | 7.8 | 23.0 | 76.2 |
| 2001..... | 2.6 | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.1 | 7.9 | 23.5 | 81.4 |
| 2000..... | 2.6 | * | * | * | * | 0.1 | 0.4 | 1.3 | 2.8 | 7.1 | 25.1 | 83.1 |
| 1999..... | 2.5 | * | * | * | * | * | 0.4 | 1.2 | 2.9 | 7.9 | 23.7 | 80.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.2 | 15.7 | 67.9 |
| 2003..... | 3.1 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.7 | 16.0 | 74.6 |
| 2002..... | 3.1 | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 4.7 | 17.6 | 76.2 |
| 2001..... | 3.1 | * | * | * | * | * | 0.2 | 0.4 | 1.7 | 4.8 | 17.5 | 76.3 |
| 2000..... | 3.2 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 4.9 | 17.7 | 79.6 |
| 1999..... | 3.0 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.8 | 16.9 | 75.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.7 | 3.1 | 9.9 | 32.8 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.6 | 3.1 | 11.2 | 35.2 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 1.2 | 4.0 | 11.8 | 29.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 1.5 | 3.3 | 9.7 | 46.9 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 3.5 | 11.5 | 40.5 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 4.1 | 8.5 | 44.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | 1.9 | 4.8 | 12.9 | 32.2 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | 5.0 | 12.3 | * |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 2.1 | 5.7 | 14.4 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | * | 10.3 | 66.7 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | 4.2 | 11.8 | 42.3 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 5.9 | * | 56.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | * | 7.9 | 33.1 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 10.5 | 38.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | * | 10.1 | 31.5 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.5 | 9.4 | 37.2 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | * | 11.2 | 39.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | * | 8.7 | 38.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of digestive system (K90,K92) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | 0.7 | * | * | * | 0.1 | 0.3 | 1.0 | 2.2 | 5.8 | 18.6 | 70.0 |
| 2003..... | 3.2 | 0.7 | * | * | * | 0.1 | 0.4 | 0.9 | 2.2 | 6.3 | 19.4 | 75.6 |
| 2002..... | 3.2 | * | * | * | * | 0.1 | 0.3 | 1.0 | 2.2 | 6.2 | 20.1 | 77.9 |
| 2001..... | 3.2 | 0.7 | * | * | * | 0.1 | 0.3 | 0.9 | 2.4 | 6.4 | 20.3 | 78.8 |
| 2000..... | 3.2 | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.1 | 6.0 | 20.9 | 81.9 |
| 1999..... | 3.1 | 0.8 | * | * | * | * | 0.3 | 0.8 | 2.3 | 6.3 | 20.0 | 78.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | 0.3 | 1.4 | 2.9 | 7.5 | 22.3 | 71.6 |
| 2003..... | 2.9 | * | * | * | * | * | 0.5 | 1.2 | 2.9 | 7.9 | 24.2 | 75.3 |
| 2002..... | 2.9 | * | * | * | * | * | 0.4 | 1.4 | 3.0 | 7.9 | 23.4 | 78.3 |
| 2001..... | 2.9 | * | * | * | * | 0.1 | 0.5 | 1.3 | 3.1 | 8.2 | 24.1 | 82.0 |
| 2000..... | 2.9 | * | * | * | * | 0.1 | 0.4 | 1.3 | 2.9 | 7.3 | 25.6 | 84.8 |
| 1999..... | 2.8 | * | * | * | * | * | 0.4 | 1.2 | 3.0 | 8.0 | 24.3 | 81.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.4 | 16.1 | 69.3 |
| 2003..... | 3.4 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.9 | 16.3 | 75.8 |
| 2002..... | 3.5 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 4.8 | 17.8 | 77.8 |
| 2001..... | 3.5 | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 4.9 | 17.9 | 77.5 |
| 2000..... | 3.5 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 5.0 | 17.9 | 80.7 |
| 1999..... | 3.4 | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 4.9 | 17.2 | 76.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | * | * | * | * | * | 0.3 | 0.9 | 2.0 | 5.5 | 18.5 | 71.4 |
| 2003..... | 3.5 | * | * | * | * | * | 0.4 | 0.8 | 2.1 | 5.9 | 19.3 | 77.0 |
| 2002..... | 3.5 | * | * | * | * | * | 0.2 | 0.9 | 2.0 | 5.9 | 19.8 | 79.8 |
| 2001..... | 3.5 | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.1 | 6.1 | 20.2 | 80.2 |
| 2000..... | 3.5 | * | * | * | * | * | 0.2 | 0.8 | 1.8 | 5.7 | 20.5 | 83.1 |
| 1999..... | 3.4 | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 6.0 | 19.6 | 79.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.0 | * | * | * | * | * | 0.3 | 1.3 | 2.6 | 7.1 | 22.2 | 73.4 |
| 2003..... | 3.1 | * | * | * | * | * | 0.4 | 1.1 | 2.7 | 7.4 | 23.9 | 77.4 |
| 2002..... | 3.1 | * | * | * | * | * | 0.3 | 1.3 | 2.8 | 7.4 | 23.2 | 79.5 |
| 2001..... | 3.1 | * | * | * | * | * | 0.4 | 1.1 | 2.8 | 7.8 | 23.8 | 83.4 |
| 2000..... | 3.0 | * | * | * | * | * | 0.3 | 1.1 | 2.5 | 6.8 | 25.1 | 86.7 |
| 1999..... | 3.0 | * | * | * | * | * | 0.4 | 1.0 | 2.7 | 7.6 | 23.9 | 82.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.1 | 16.0 | 70.5 |
| 2003..... | 3.8 | * | * | * | * | * | 0.3 | 0.5 | 1.4 | 4.7 | 16.2 | 76.9 |
| 2002..... | 3.9 | * | * | * | * | * | 0.1 | 0.5 | 1.2 | 4.6 | 17.6 | 79.9 |
| 2001..... | 3.9 | * | * | * | * | * | 0.2 | 0.4 | 1.5 | 4.8 | 17.8 | 78.8 |
| 2000..... | 3.8 | * | * | * | * | * | 0.1 | 0.4 | 1.2 | 4.7 | 17.5 | 81.6 |
| 1999..... | 3.7 | * | * | * | * | * | 0.1 | 0.4 | 1.5 | 4.7 | 16.8 | 78.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other diseases of digestive system (K90,K92) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 1.6 | 4.1 | 10.0 | 24.8 | 63.0 |
| 2003..... | 2.4 | * | * | * | * | * | 0.6 | 1.4 | 4.2 | 10.6 | 24.9 | 71.1 |
| 2002..... | 2.5 | * | * | * | * | * | 0.6 | 1.8 | 4.4 | 10.0 | 26.7 | 65.9 |
| 2001..... | 2.5 | * | * | * | * | * | 0.6 | 1.7 | 5.2 | 9.9 | 26.0 | 73.4 |
| 2000..... | 2.6 | * | * | * | * | * | 0.7 | 2.0 | 5.0 | 10.0 | 28.2 | 79.2 |
| 1999..... | 2.3 | * | * | * | * | * | 0.4 | 1.7 | 4.3 | 10.0 | 26.7 | 66.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | * | 2.2 | 6.6 | 14.1 | 29.7 | 60.0 |
| 2003..... | 2.4 | * | * | * | * | * | 1.0 | 2.1 | 5.7 | 14.2 | 32.0 | 60.7 |
| 2002..... | 2.5 | * | * | * | * | * | 0.8 | 2.6 | 5.7 | 14.1 | 30.8 | 72.2 |
| 2001..... | 2.6 | * | * | * | * | * | 0.8 | 2.8 | 6.9 | 13.9 | 32.7 | 71.6 |
| 2000..... | 2.6 | * | * | * | * | * | 1.0 | 2.8 | 6.7 | 12.1 | 35.5 | 79.8 |
| 1999..... | 2.3 | * | * | * | * | * | * | 2.5 | 5.7 | 13.1 | 32.8 | 72.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | * | * | 1.1 | 2.1 | 7.1 | 22.1 | 64.3 |
| 2003..... | 2.5 | * | * | * | * | * | * | 0.9 | 3.1 | 8.1 | 20.9 | 75.1 |
| 2002..... | 2.4 | * | * | * | * | * | * | 1.1 | 3.3 | 7.1 | 24.4 | 63.5 |
| 2001..... | 2.5 | * | * | * | * | * | * | * | 3.8 | 7.0 | 22.2 | 74.0 |
| 2000..... | 2.7 | * | * | * | * | * | * | 1.3 | 3.7 | 8.4 | 24.1 | 78.9 |
| 1999..... | 2.4 | * | * | * | * | * | * | 1.0 | 3.1 | 7.8 | 23.3 | 64.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XII. Diseases of the skin and subcutaneous tissue (L00-L98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.4 | 3.4 | 9.5 | 28.8 |
| 2003..... | 1.4 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.4 | 3.3 | 9.2 | 30.5 |
| 2002..... | 1.4 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.2 | 3.1 | 8.7 | 32.8 |
| 2001..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.8 | 8.7 | 32.2 |
| 2000..... | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 3.0 | 8.9 | 34.7 |
| 1999..... | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.9 | 9.0 | 35.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 3.0 | 9.0 | 25.3 |
| 2003..... | 1.0 | * | * | * | * | * | 0.3 | 0.7 | 1.4 | 3.3 | 8.4 | 26.6 |
| 2002..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 3.2 | 8.6 | 28.6 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 3.0 | 8.3 | 24.9 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 2.8 | 8.4 | 29.7 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.8 | 8.4 | 31.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.7 | 9.8 | 30.4 |
| 2003..... | 1.8 | * | * | * | * | 0.3 | 0.3 | 0.7 | 1.4 | 3.4 | 9.7 | 32.2 |
| 2002..... | 1.7 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 3.0 | 8.8 | 34.7 |
| 2001..... | 1.7 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.6 | 9.1 | 35.2 |
| 2000..... | 1.7 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 3.1 | 9.2 | 36.7 |
| 1999..... | 1.7 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.9 | 9.4 | 36.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.7 | 1.2 | 2.6 | 7.5 | 21.9 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.8 | 1.3 | 2.7 | 7.5 | 23.8 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | 1.9 | 7.0 | 24.4 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.6 | 1.8 | 2.3 | 6.5 | 24.4 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 1.2 | 2.0 | 6.5 | 29.9 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 2.0 | 7.5 | 25.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | 7.2 | * |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | 9.2 | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | * | * | 7.7 | 22.3 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 6.3 | 24.2 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 6.7 | 24.0 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | 8.5 | 24.2 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | * | 7.3 | 30.7 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 9.0 | 29.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XII. Diseases of the skin and subcutaneous tissue (L00-L98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 3.5 | 9.6 | 29.0 |
| 2003..... | 1.6 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 3.4 | 9.2 | 30.7 |
| 2002..... | 1.5 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.1 | 8.8 | 33.1 |
| 2001..... | 1.4 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.8 | 8.8 | 32.4 |
| 2000..... | 1.5 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 3.0 | 9.0 | 34.7 |
| 1999..... | 1.4 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.9 | 9.1 | 35.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 3.1 | 9.1 | 25.4 |
| 2003..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.3 | 8.4 | 26.7 |
| 2002..... | 1.1 | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 3.2 | 8.6 | 28.6 |
| 2001..... | 1.0 | * | * | * | * | * | 0.2 | 0.6 | 1.1 | 3.0 | 8.4 | 24.9 |
| 2000..... | 1.0 | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 2.8 | 8.5 | 29.6 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.8 | 8.5 | 32.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.8 | 9.9 | 30.6 |
| 2003..... | 1.9 | * | * | * | * | 0.3 | 0.3 | 0.6 | 1.4 | 3.4 | 9.8 | 32.5 |
| 2002..... | 1.9 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 3.0 | 8.8 | 35.0 |
| 2001..... | 1.8 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.7 | 9.1 | 35.5 |
| 2000..... | 1.9 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 3.2 | 9.3 | 36.8 |
| 1999..... | 1.8 | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 3.0 | 9.4 | 36.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 2.8 | 8.6 | 25.9 |
| 2003..... | 1.5 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.7 | 8.1 | 27.4 |
| 2002..... | 1.4 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.5 | 7.4 | 28.6 |
| 2001..... | 1.3 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.4 | 7.5 | 28.2 |
| 2000..... | 1.3 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.4 | 7.6 | 29.4 |
| 1999..... | 1.3 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.3 | 7.4 | 29.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.5 | 8.3 | 23.1 |
| 2003..... | 1.1 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.5 | 7.3 | 24.2 |
| 2002..... | 1.1 | * | * | * | * | * | 0.2 | 0.5 | 0.9 | 2.6 | 7.7 | 26.3 |
| 2001..... | 0.9 | * | * | * | * | * | 0.1 | 0.5 | 1.0 | 2.6 | 7.0 | 21.6 |
| 2000..... | 0.9 | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 2.1 | 7.1 | 25.4 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.2 | 6.9 | 27.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.1 | 8.7 | 27.1 |
| 2003..... | 1.8 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.9 | 8.6 | 28.8 |
| 2002..... | 1.7 | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.5 | 7.3 | 29.5 |
| 2001..... | 1.7 | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 2.2 | 7.9 | 30.9 |
| 2000..... | 1.7 | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.6 | 7.9 | 31.0 |
| 1999..... | 1.6 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.4 | 7.8 | 29.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XII. Diseases of the skin and subcutaneous tissue (L00-L98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | 0.6 | 0.8 | 1.5 | 3.7 | 9.8 | 23.8 | 71.1 |
| 2003..... | 2.5 | * | * | * | * | 0.8 | 0.9 | 1.8 | 3.6 | 9.7 | 24.2 | 74.1 |
| 2002..... | 2.6 | * | * | * | * | 0.5 | 1.0 | 1.5 | 3.7 | 8.6 | 25.9 | 93.2 |
| 2001..... | 2.4 | * | * | * | * | 0.4 | 0.8 | 1.4 | 3.2 | 7.4 | 25.2 | 89.6 |
| 2000..... | 2.5 | * | * | * | * | 0.4 | 0.7 | 1.3 | 3.1 | 9.2 | 26.6 | 98.9 |
| 1999..... | 2.8 | * | * | * | * | * | 0.8 | 1.9 | 3.6 | 9.0 | 29.5 | 106.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | * | 1.5 | 3.0 | 9.3 | 22.2 | 58.9 |
| 2003..... | 2.0 | * | * | * | * | * | 0.8 | 1.6 | 3.9 | 11.6 | 24.0 | 67.1 |
| 2002..... | 1.7 | * | * | * | * | * | * | 1.3 | 3.0 | 10.2 | 24.0 | 61.3 |
| 2001..... | 1.8 | * | * | * | * | * | * | 1.6 | 2.6 | 7.4 | 28.8 | 72.8 |
| 2000..... | 1.8 | * | * | * | * | * | * | * | 3.0 | 10.3 | 28.4 | 83.4 |
| 1999..... | 2.0 | * | * | * | * | * | * | 2.1 | 3.3 | 9.3 | 30.0 | 86.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | 0.8 | 0.9 | 1.4 | 4.2 | 10.3 | 24.6 | 75.9 |
| 2003..... | 3.0 | * | * | * | * | 1.1 | 1.0 | 2.0 | 3.4 | 8.3 | 24.3 | 76.8 |
| 2002..... | 3.4 | * | * | * | * | * | 1.2 | 1.6 | 4.4 | 7.4 | 27.0 | 105.3 |
| 2001..... | 3.0 | * | * | * | * | * | 1.0 | 1.2 | 3.6 | 7.4 | 23.2 | 95.8 |
| 2000..... | 3.2 | * | * | * | * | * | 0.7 | 1.7 | 3.2 | 8.3 | 25.5 | 104.7 |
| 1999..... | 3.4 | * | * | * | * | * | 0.9 | 1.6 | 3.8 | 8.8 | 29.2 | 113.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Infections of skin and subcutaneous tissue (L00-L08) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 3.9 | 10.6 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.3 | 3.5 | 10.8 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.8 | 9.5 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.6 | 9.7 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.6 | 9.4 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.1 | 8.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.4 | 4.0 | 10.7 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 1.3 | 3.4 | 10.7 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.2 | 3.0 | 9.3 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.6 | 1.2 | 2.5 | 8.3 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.9 | 9.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.4 | 0.8 | 2.1 | 9.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.3 | 0.7 | 1.5 | 3.8 | 10.6 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.5 | 10.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.7 | 9.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.7 | 10.2 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 9.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.0 | 8.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 3.9 | 10.6 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 3.5 | 11.1 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.8 | 9.9 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.0 | 2.6 | 9.9 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.0 | 2.5 | 9.4 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.1 | 8.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.3 | 4.1 | 11.1 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.7 | 1.3 | 3.4 | 10.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 3.1 | 9.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.5 | 8.6 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.7 | 1.1 | 2.8 | 9.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.0 | 9.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.6 | 1.5 | 3.8 | 10.4 |
| 2003..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 3.6 | 11.1 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.6 | 10.0 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.7 | 10.4 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.4 | 9.5 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.2 | 8.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified diseases of skin and subcutaneous tissue (L10-L98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.7 | 2.0 | 5.7 | 18.4 |
| 2003..... | 0.9 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.0 | 5.7 | 19.9 |
| 2002..... | 0.9 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.0 | 5.9 | 23.5 |
| 2001..... | 0.9 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.8 | 6.1 | 22.7 |
| 2000..... | 0.9 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.9 | 6.4 | 25.4 |
| 1999..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 2.0 | 7.0 | 26.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 5.0 | 14.9 |
| 2003..... | 0.6 | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.9 | 4.9 | 16.2 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 2.0 | 5.7 | 19.4 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 5.8 | 16.8 |
| 2000..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 5.6 | 20.8 |
| 1999..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 2.0 | 6.3 | 22.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.2 | 6.1 | 20.0 |
| 2003..... | 1.2 | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 2.1 | 6.2 | 21.5 |
| 2002..... | 1.2 | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.8 | 2.0 | 6.1 | 25.3 |
| 2001..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 1.8 | 6.4 | 25.2 |
| 2000..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 2.2 | 6.9 | 27.3 |
| 1999..... | 1.3 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.0 | 7.4 | 28.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | 5.1 | 16.2 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | 4.8 | 18.7 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | 5.2 | 20.5 |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | 4.5 | 21.3 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | 4.7 | 23.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | 5.6 | 19.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | 5.7 | 15.8 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | * | * | 18.8 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 20.7 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.0 | 21.4 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 22.8 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.9 | 21.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified diseases of skin and subcutaneous tissue (L10-L98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 2.0 | 5.7 | 18.4 |
| 2003..... | 1.0 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 2.1 | 5.7 | 19.9 |
| 2002..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 2.1 | 6.0 | 23.6 |
| 2001..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.8 | 6.2 | 22.7 |
| 2000..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 2.0 | 6.4 | 25.3 |
| 1999..... | 1.1 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 2.0 | 7.0 | 26.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.7 | 5.1 | 14.8 |
| 2003..... | 0.6 | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.9 | 4.9 | 16.1 |
| 2002..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 2.1 | 5.6 | 19.3 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 5.9 | 16.6 |
| 2000..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 5.7 | 20.4 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 | 6.4 | 22.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 6.1 | 20.1 |
| 2003..... | 1.3 | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 2.1 | 6.2 | 21.6 |
| 2002..... | 1.3 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.9 | 2.0 | 6.2 | 25.4 |
| 2001..... | 1.3 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.7 | 1.8 | 6.4 | 25.3 |
| 2000..... | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.3 | 6.9 | 27.2 |
| 1999..... | 1.4 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.1 | 7.4 | 28.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 4.6 | 15.3 |
| 2003..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.6 | 16.3 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 4.7 | 18.6 |
| 2001..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.9 | 18.3 |
| 2000..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.4 | 5.0 | 20.0 |
| 1999..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.4 | 5.3 | 20.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 4.2 | 12.0 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.9 | 13.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 1.5 | 4.6 | 16.7 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 4.5 | 13.0 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 4.2 | 16.4 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 4.9 | 18.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.7 | 4.9 | 16.8 |
| 2003..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 5.0 | 17.7 |
| 2002..... | 1.1 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 4.7 | 19.5 |
| 2001..... | 1.1 | * | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 5.2 | 20.5 |
| 2000..... | 1.2 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 5.6 | 21.4 |
| 1999..... | 1.2 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.5 | 5.6 | 21.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified diseases of skin and subcutaneous tissue (L10-L98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | 0.5 | 0.6 | 1.0 | 2.6 | 7.6 | 19.3 | 58.1 |
| 2003..... | 2.0 | * | * | * | * | 0.7 | 0.7 | 1.2 | 2.7 | 7.4 | 20.2 | 64.6 |
| 2002..... | 2.2 | * | * | * | * | 0.4 | 0.7 | 1.1 | 2.8 | 6.6 | 22.1 | 86.2 |
| 2001..... | 2.0 | * | * | * | * | * | 0.6 | 0.9 | 2.1 | 6.3 | 22.1 | 80.2 |
| 2000..... | 2.2 | * | * | * | * | * | 0.5 | 1.0 | 2.4 | 7.7 | 23.4 | 88.7 |
| 1999..... | 2.4 | * | * | * | * | * | 0.7 | 1.2 | 2.5 | 7.9 | 27.8 | 97.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | * | 2.0 | 7.0 | 17.7 | 49.8 |
| 2003..... | 1.5 | * | * | * | * | * | * | * | 2.8 | 9.4 | 19.9 | 56.5 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 2.2 | 7.9 | 20.8 | 55.8 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | 1.8 | 6.4 | 25.8 | 67.0 |
| 2000..... | 1.5 | * | * | * | * | * | * | * | 2.4 | 8.2 | 25.0 | 74.0 |
| 1999..... | 1.7 | * | * | * | * | * | * | 1.1 | 2.2 | 8.9 | 26.9 | 74.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | * | 0.8 | 1.1 | 3.1 | 8.0 | 20.3 | 61.4 |
| 2003..... | 2.5 | * | * | * | * | 1.0 | 0.9 | 1.6 | 2.6 | 6.0 | 20.4 | 67.7 |
| 2002..... | 3.0 | * | * | * | * | * | 1.1 | 1.3 | 3.3 | 5.6 | 22.9 | 97.8 |
| 2001..... | 2.5 | * | * | * | * | * | 0.8 | 0.9 | 2.4 | 6.2 | 20.0 | 85.1 |
| 2000..... | 2.8 | * | * | * | * | * | * | 1.5 | 2.4 | 7.3 | 22.6 | 94.2 |
| 1999..... | 3.1 | * | * | * | * | * | 0.8 | 1.3 | 2.7 | 7.2 | 28.4 | 106.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | 0.1 | 0.4 | 0.7 | 1.3 | 2.7 | 5.4 | 12.1 | 31.4 | 85.9 |
| 2003..... | 5.0 | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.7 | 5.6 | 12.7 | 31.7 | 90.4 |
| 2002..... | 5.0 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.7 | 5.5 | 12.9 | 31.7 | 94.9 |
| 2001..... | 4.9 | * | * | 0.1 | 0.5 | 0.7 | 1.5 | 2.6 | 5.3 | 12.8 | 32.2 | 93.4 |
| 2000..... | 4.9 | * | * | 0.1 | 0.4 | 0.8 | 1.4 | 2.6 | 5.4 | 12.6 | 32.2 | 94.0 |
| 1999..... | 4.8 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.7 | 5.4 | 12.7 | 31.1 | 91.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | 0.3 | 0.3 | 0.8 | 2.1 | 4.1 | 9.0 | 24.2 | 59.1 |
| 2003..... | 3.0 | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 | 4.1 | 10.3 | 24.5 | 59.9 |
| 2002..... | 3.0 | * | * | * | 0.2 | 0.4 | 0.9 | 1.9 | 4.0 | 10.2 | 24.1 | 63.5 |
| 2001..... | 2.9 | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 | 3.9 | 9.6 | 24.9 | 63.2 |
| 2000..... | 2.9 | * | * | * | 0.3 | 0.5 | 0.9 | 1.7 | 4.0 | 9.8 | 25.2 | 67.5 |
| 1999..... | 2.8 | * | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 3.8 | 9.5 | 24.2 | 63.7 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | 0.1 | 0.5 | 1.1 | 1.9 | 3.4 | 6.7 | 14.6 | 36.3 | 98.0 |
| 2003..... | 6.9 | * | * | * | 0.6 | 1.2 | 1.8 | 3.5 | 6.9 | 14.6 | 36.5 | 103.8 |
| 2002..... | 7.0 | * | * | 0.1 | 0.6 | 1.0 | 1.8 | 3.3 | 6.9 | 15.1 | 36.8 | 108.5 |
| 2001..... | 6.9 | * | * | * | 0.7 | 1.1 | 2.0 | 3.2 | 6.6 | 15.6 | 37.0 | 106.0 |
| 2000..... | 6.8 | * | * | 0.1 | 0.5 | 1.1 | 1.9 | 3.5 | 6.6 | 15.0 | 36.8 | 104.8 |
| 1999..... | 6.7 | * | * | 0.1 | 0.6 | 1.1 | 1.9 | 3.7 | 6.9 | 15.3 | 35.5 | 102.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | 0.3 | 0.7 | 1.2 | 2.9 | 6.3 | 11.1 | 23.7 | 43.8 |
| 2003..... | 2.0 | * | * | * | 0.5 | 0.7 | 1.6 | 2.6 | 6.3 | 10.7 | 22.3 | 44.6 |
| 2002..... | 1.9 | * | * | * | 0.4 | 0.5 | 1.4 | 2.7 | 5.6 | 12.6 | 20.9 | 53.2 |
| 2001..... | 2.0 | * | * | * | 0.6 | 0.6 | 1.6 | 3.0 | 5.0 | 12.9 | 22.4 | 55.6 |
| 2000..... | 1.9 | * | * | * | * | 0.9 | 1.2 | 2.5 | 5.7 | 13.0 | 26.9 | 55.1 |
| 1999..... | 2.1 | * | * | * | 0.4 | 0.8 | 1.5 | 3.0 | 6.9 | 12.8 | 22.7 | 73.7 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.7 | 2.3 | 3.8 | 6.7 | 16.1 | 32.2 |
| 2003..... | 1.2 | * | * | * | * | * | 0.9 | 1.6 | 4.2 | 7.4 | 19.6 | * |
| 2002..... | 1.0 | * | * | * | * | * | 1.1 | 1.6 | 3.1 | 6.7 | 13.5 | 36.8 |
| 2001..... | 1.2 | * | * | * | * | * | 1.1 | 2.6 | 2.9 | 7.4 | 19.2 | 41.9 |
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | 1.5 | 4.5 | 7.4 | 14.3 | 40.3 |
| 1999..... | 1.1 | * | * | * | * | * | 0.9 | 1.6 | 4.3 | 8.6 | 16.7 | 47.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | * | 1.4 | 1.9 | 3.5 | 8.6 | 14.6 | 28.9 | 49.7 |
| 2003..... | 2.9 | * | * | * | 0.9 | 1.1 | 2.5 | 3.7 | 8.1 | 13.4 | 24.2 | 52.3 |
| 2002..... | 2.9 | * | * | * | 0.8 | 0.9 | 1.8 | 3.7 | 7.9 | 17.3 | 25.9 | 61.3 |
| 2001..... | 2.8 | * | * | * | 1.1 | 0.9 | 2.1 | 3.5 | 7.0 | 17.3 | 24.5 | 62.4 |
| 2000..... | 2.9 | * | * | * | * | 1.4 | 1.7 | 3.5 | 6.7 | 17.4 | 35.3 | 62.3 |
| 1999..... | 3.1 | * | * | * | 0.7 | 1.3 | 2.2 | 4.3 | 9.2 | 16.2 | 26.7 | 86.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.7 | 5.4 | 12.1 | 31.8 | 87.6 |
| 2003..... | 5.4 | * | * | 0.1 | 0.4 | 0.8 | 1.3 | 2.7 | 5.5 | 12.8 | 32.1 | 92.2 |
| 2002..... | 5.5 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.6 | 5.5 | 12.8 | 32.1 | 96.2 |
| 2001..... | 5.4 | * | * | 0.1 | 0.4 | 0.8 | 1.5 | 2.5 | 5.3 | 12.8 | 32.6 | 94.6 |
| 2000..... | 5.3 | * | * | 0.1 | 0.4 | 0.8 | 1.5 | 2.6 | 5.3 | 12.6 | 32.4 | 95.2 |
| 1999..... | 5.1 | * | * | 0.1 | 0.4 | 0.7 | 1.4 | 2.7 | 5.3 | 12.7 | 31.3 | 91.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | 0.3 | 0.4 | 0.8 | 2.0 | 4.1 | 9.2 | 24.6 | 60.1 |
| 2003..... | 3.3 | * | * | * | 0.2 | 0.4 | 1.0 | 1.9 | 4.0 | 10.5 | 24.7 | 61.4 |
| 2002..... | 3.3 | * | * | * | 0.2 | 0.5 | 0.8 | 2.0 | 4.1 | 10.4 | 24.6 | 64.7 |
| 2001..... | 3.2 | * | * | * | 0.2 | 0.4 | 1.0 | 1.8 | 3.9 | 9.7 | 25.0 | 64.0 |
| 2000..... | 3.2 | * | * | * | 0.3 | 0.5 | 0.9 | 1.7 | 3.9 | 9.9 | 25.6 | 68.4 |
| 1999..... | 3.0 | * | * | * | 0.2 | 0.4 | 0.9 | 1.8 | 3.8 | 9.6 | 24.5 | 63.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.3 | * | * | 0.1 | 0.5 | 1.0 | 1.9 | 3.3 | 6.5 | 14.6 | 36.7 | 99.9 |
| 2003..... | 7.5 | * | * | * | 0.6 | 1.3 | 1.7 | 3.5 | 6.8 | 14.7 | 37.0 | 105.7 |
| 2002..... | 7.5 | * | * | 0.2 | 0.6 | 1.0 | 1.9 | 3.3 | 6.8 | 14.9 | 37.2 | 109.9 |
| 2001..... | 7.4 | * | * | * | 0.6 | 1.1 | 1.9 | 3.2 | 6.5 | 15.4 | 37.6 | 107.3 |
| 2000..... | 7.3 | * | * | 0.1 | 0.5 | 1.1 | 1.9 | 3.5 | 6.6 | 14.8 | 36.7 | 106.1 |
| 1999..... | 7.1 | * | * | 0.1 | 0.6 | 1.0 | 1.9 | 3.6 | 6.6 | 15.3 | 35.8 | 102.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | 0.1 | 0.2 | 0.3 | 0.9 | 2.0 | 4.8 | 11.9 | 32.3 | 90.6 |
| 2003..... | 5.8 | * | * | * | 0.2 | 0.4 | 1.0 | 2.1 | 5.0 | 12.5 | 32.4 | 94.8 |
| 2002..... | 5.8 | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.1 | 5.0 | 12.7 | 32.6 | 99.7 |
| 2001..... | 5.7 | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.0 | 4.8 | 12.5 | 33.3 | 97.1 |
| 2000..... | 5.6 | * | * | * | 0.2 | 0.5 | 0.9 | 2.0 | 4.8 | 12.5 | 32.8 | 98.2 |
| 1999..... | 5.4 | * | * | * | 0.2 | 0.4 | 0.8 | 2.1 | 4.7 | 12.5 | 32.0 | 94.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | 0.2 | 0.2 | 0.6 | 1.5 | 3.8 | 8.8 | 24.6 | 60.8 |
| 2003..... | 3.4 | * | * | * | 0.2 | 0.3 | 0.8 | 1.6 | 3.6 | 9.9 | 24.2 | 61.5 |
| 2002..... | 3.4 | * | * | * | * | 0.3 | 0.6 | 1.7 | 3.7 | 10.0 | 24.5 | 65.5 |
| 2001..... | 3.3 | * | * | * | 0.2 | 0.3 | 0.8 | 1.5 | 3.5 | 9.4 | 25.3 | 65.1 |
| 2000..... | 3.3 | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.7 | 9.4 | 25.4 | 68.2 |
| 1999..... | 3.1 | * | * | * | * | 0.3 | 0.6 | 1.5 | 3.4 | 9.5 | 24.4 | 64.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.8 | * | * | * | 0.2 | 0.4 | 1.2 | 2.5 | 5.7 | 14.6 | 37.6 | 103.9 |
| 2003..... | 8.0 | * | * | * | 0.3 | 0.6 | 1.2 | 2.6 | 6.3 | 14.7 | 38.0 | 109.5 |
| 2002..... | 8.1 | * | * | * | 0.2 | 0.5 | 1.1 | 2.6 | 6.2 | 14.9 | 38.0 | 114.5 |
| 2001..... | 7.9 | * | * | * | 0.3 | 0.6 | 1.2 | 2.5 | 6.0 | 15.3 | 38.6 | 110.4 |
| 2000..... | 7.8 | * | * | * | 0.3 | 0.6 | 1.2 | 2.6 | 5.9 | 15.0 | 37.6 | 110.3 |
| 1999..... | 7.6 | * | * | * | 0.2 | 0.6 | 1.0 | 2.6 | 6.0 | 15.1 | 37.0 | 106.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | * | 1.1 | 2.6 | 3.9 | 7.4 | 10.4 | 16.0 | 31.2 | 66.2 |
| 2003..... | 5.1 | * | * | * | 1.2 | 2.6 | 3.5 | 7.3 | 10.1 | 17.2 | 32.3 | 69.3 |
| 2002..... | 5.0 | * | * | * | 1.2 | 2.6 | 4.3 | 6.4 | 10.2 | 16.2 | 31.3 | 68.4 |
| 2001..... | 4.9 | * | * | * | 1.2 | 2.5 | 4.4 | 6.5 | 9.6 | 16.7 | 29.7 | 76.2 |
| 2000..... | 5.0 | * | * | * | 1.3 | 2.5 | 4.7 | 7.3 | 10.0 | 15.5 | 31.1 | 68.7 |
| 1999..... | 4.8 | * | * | * | 1.2 | 2.2 | 4.9 | 7.2 | 9.8 | 16.4 | 27.3 | 60.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | 0.7 | 1.6 | 2.0 | 5.5 | 7.3 | 15.0 | 29.1 | 62.0 |
| 2003..... | 3.4 | * | * | * | * | 1.1 | 2.4 | 4.3 | 8.3 | 18.4 | 34.1 | 59.7 |
| 2002..... | 3.2 | * | * | * | * | 1.5 | 2.3 | 4.1 | 8.0 | 15.5 | 28.4 | 61.3 |
| 2001..... | 3.2 | * | * | * | * | 1.2 | 2.8 | 4.9 | 7.9 | 13.9 | 28.2 | 60.1 |
| 2000..... | 3.5 | * | * | * | 1.0 | 1.4 | 3.1 | 4.1 | 7.2 | 16.6 | 30.9 | 78.7 |
| 1999..... | 3.0 | * | * | * | * | 0.9 | 3.0 | 4.2 | 6.7 | 13.1 | 29.4 | 64.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | 1.6 | 3.6 | 5.6 | 9.0 | 12.9 | 16.7 | 32.4 | 67.9 |
| 2003..... | 6.6 | * | * | * | 2.1 | 4.0 | 4.5 | 10.0 | 11.6 | 16.3 | 31.3 | 73.0 |
| 2002..... | 6.6 | * | * | * | 1.9 | 3.6 | 6.2 | 8.3 | 12.1 | 16.6 | 33.0 | 71.1 |
| 2001..... | 6.5 | * | * | * | 2.0 | 3.7 | 5.7 | 7.8 | 11.0 | 18.7 | 30.5 | 82.1 |
| 2000..... | 6.4 | * | * | * | 1.6 | 3.5 | 6.1 | 10.0 | 12.4 | 14.6 | 31.3 | 65.0 |
| 1999..... | 6.4 | * | * | * | 1.9 | 3.4 | 6.6 | 9.8 | 12.3 | 18.7 | 26.1 | 59.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.1 | 8.0 | 11.9 |
| 2003..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.1 | 8.7 | 12.8 |
| 2002..... | 1.0 | * | * | * | * | * | 0.0 | 0.2 | 1.0 | 3.3 | 9.1 | 14.2 |
| 2001..... | 0.9 | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 3.3 | 8.8 | 12.7 |
| 2000..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 3.5 | 9.2 | 12.9 |
| 1999..... | 1.0 | * | * | * | * | * | 0.0 | 0.2 | 1.1 | 3.5 | 9.5 | 13.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.6 | 1.9 | 4.3 | 6.5 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.7 | 1.9 | 5.0 | 6.5 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 5.3 | 7.6 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.7 | 2.0 | 5.1 | 5.5 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.6 | 2.3 | 5.2 | 6.9 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.8 | 2.3 | 5.2 | 7.2 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 0.3 | 1.1 | 4.1 | 10.5 | 14.3 |
| 2003..... | 1.4 | * | * | * | * | * | * | 0.3 | 1.1 | 4.1 | 11.1 | 15.6 |
| 2002..... | 1.4 | * | * | * | * | * | * | 0.3 | 1.3 | 4.3 | 11.6 | 17.0 |
| 2001..... | 1.4 | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.4 | 11.2 | 15.7 |
| 2000..... | 1.4 | * | * | * | * | * | * | 0.4 | 1.3 | 4.5 | 11.7 | 15.3 |
| 1999..... | 1.5 | * | * | * | * | * | * | 0.3 | 1.5 | 4.4 | 12.2 | 16.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 1.0 | 2.3 | 6.6 | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 1.0 | 2.8 | 6.4 | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | 3.0 | 6.7 | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | 4.5 | 6.0 | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | 3.7 | 8.3 | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 1.4 | 3.3 | 7.3 | 20.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | 3.3 | 9.4 | * |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | 4.7 | 8.7 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | 4.3 | 10.1 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 7.2 | 7.3 | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | 5.1 | 11.9 | * |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 5.2 | 10.1 | 25.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.2 | 8.1 | 12.0 |
| 2003..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.2 | 8.8 | 13.0 |
| 2002..... | 1.1 | * | * | * | * | * | * | 0.2 | 1.0 | 3.3 | 9.2 | 14.3 |
| 2001..... | 1.0 | * | * | * | * | * | 0.1 | 0.2 | 1.0 | 3.2 | 8.9 | 12.8 |
| 2000..... | 1.0 | * | * | * | * | * | * | 0.3 | 1.0 | 3.5 | 9.2 | 13.2 |
| 1999..... | 1.1 | * | * | * | * | * | * | 0.2 | 1.1 | 3.5 | 9.5 | 13.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.6 | 1.9 | 4.4 | 6.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.7 | 2.0 | 5.1 | 6.6 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.5 | 7.7 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.7 | 2.1 | 5.1 | 5.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.6 | 2.4 | 5.3 | 7.1 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.8 | 2.4 | 5.3 | 7.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 0.3 | 1.1 | 4.2 | 10.5 | 14.5 |
| 2003..... | 1.5 | * | * | * | * | * | * | 0.3 | 1.1 | 4.1 | 11.2 | 15.8 |
| 2002..... | 1.6 | * | * | * | * | * | * | 0.3 | 1.3 | 4.3 | 11.6 | 17.1 |
| 2001..... | 1.5 | * | * | * | * | * | 0.1 | 0.3 | 1.3 | 4.2 | 11.4 | 15.8 |
| 2000..... | 1.6 | * | * | * | * | * | * | 0.4 | 1.3 | 4.4 | 11.7 | 15.6 |
| 1999..... | 1.6 | * | * | * | * | * | * | 0.3 | 1.4 | 4.4 | 12.3 | 16.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | 0.2 | 0.9 | 3.3 | 8.3 | 12.2 |
| 2003..... | 1.2 | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 3.3 | 9.0 | 13.3 |
| 2002..... | 1.2 | * | * | * | * | * | * | 0.2 | 1.0 | 3.5 | 9.5 | 14.9 |
| 2001..... | 1.2 | * | * | * | * | * | * | 0.2 | 1.1 | 3.4 | 9.3 | 13.0 |
| 2000..... | 1.2 | * | * | * | * | * | * | 0.3 | 1.0 | 3.7 | 9.5 | 13.7 |
| 1999..... | 1.2 | * | * | * | * | * | * | 0.3 | 1.2 | 3.7 | 9.9 | 14.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.6 | 1.9 | 4.6 | 6.4 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | 0.7 | 2.1 | 5.1 | 6.7 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 5.7 | 8.1 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | 0.8 | 2.2 | 5.3 | 5.8 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.7 | 2.5 | 5.4 | 7.5 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.9 | 2.5 | 5.4 | 7.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 0.3 | 1.2 | 4.5 | 10.8 | 14.9 |
| 2003..... | 1.7 | * | * | * | * | * | * | 0.3 | 1.2 | 4.3 | 11.6 | 16.2 |
| 2002..... | 1.8 | * | * | * | * | * | * | 0.3 | 1.3 | 4.5 | 12.0 | 17.8 |
| 2001..... | 1.8 | * | * | * | * | * | * | 0.3 | 1.3 | 4.5 | 12.0 | 16.0 |
| 2000..... | 1.8 | * | * | * | * | * | * | 0.4 | 1.3 | 4.7 | 12.2 | 16.1 |
| 1999..... | 1.9 | * | * | * | * | * | * | 0.3 | 1.5 | 4.7 | 12.8 | 17.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.7 | 2.5 | 6.5 | 9.5 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | * | 2.3 | 6.9 | 9.8 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 0.9 | 2.5 | 6.7 | 8.2 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.9 | 2.2 | 5.1 | 12.2 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.9 | 2.0 | 5.2 | 7.9 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 2.0 | 6.2 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 2.7 | 8.4 | 11.2 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | 2.9 | 8.0 | 11.6 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 2.7 | 8.1 | 10.0 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 2.7 | 5.4 | 15.0 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | * | 6.4 | 9.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 2.2 | 6.8 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Systemic lupus erythematosus (M32) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 0.8 | 0.8 | 1.3 | 1.1 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 0.8 | 0.9 | 1.4 | 1.1 |
| 2002..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.8 | 1.0 | 1.3 | 1.3 |
| 2001..... | 0.5 | * | * | * | 0.3 | 0.4 | 0.5 | 0.6 | 0.8 | 1.0 | 1.5 | 1.2 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.6 | 0.7 | 0.9 | 0.9 | 1.5 | 1.2 |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 0.8 | 1.1 | 1.3 | 1.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.4 | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | * |
| 2000..... | 0.1 | * | * | * | * | * | 0.2 | 0.1 | 0.3 | 0.4 | 0.7 | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.6 | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.3 | 0.7 | 0.8 | 1.0 | 1.3 | 1.3 | 1.8 | 1.3 |
| 2003..... | 0.8 | * | * | * | 0.4 | 0.6 | 0.8 | 1.1 | 1.4 | 1.3 | 2.1 | 1.3 |
| 2002..... | 0.8 | * | * | * | 0.4 | 0.5 | 0.9 | 1.0 | 1.4 | 1.5 | 1.8 | 1.7 |
| 2001..... | 0.8 | * | * | * | 0.5 | 0.7 | 0.8 | 1.1 | 1.3 | 1.6 | 2.1 | 1.3 |
| 2000..... | 0.8 | * | * | * | 0.3 | 0.6 | 1.0 | 1.1 | 1.4 | 1.3 | 2.0 | 1.4 |
| 1999..... | 0.8 | * | * | * | 0.4 | 0.6 | 0.9 | 1.2 | 1.3 | 1.8 | 1.8 | 1.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | 0.5 | 0.5 | 0.7 | 1.0 | * | * | * |
| 2003..... | 0.4 | * | * | * | 0.3 | 0.4 | 0.7 | 0.7 | 1.2 | * | * | * |
| 2002..... | 0.4 | * | * | * | * | 0.3 | 0.6 | 0.8 | 1.2 | 1.8 | * | * |
| 2001..... | 0.4 | * | * | * | 0.4 | * | 0.5 | 0.7 | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | 0.4 | 0.5 | 0.7 | 1.3 | * | * | * |
| 1999..... | 0.5 | * | * | * | * | 0.4 | 0.7 | 1.1 | 1.4 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | * | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 1.1 | 0.9 | 1.2 | 1.7 | * | * | * |
| 2003..... | 0.7 | * | * | * | 0.6 | 0.6 | 1.3 | 1.2 | 2.0 | * | * | * |
| 2002..... | 0.7 | * | * | * | * | * | 1.0 | 1.4 | 2.1 | * | * | * |
| 2001..... | 0.7 | * | * | * | 0.8 | * | 0.8 | 1.2 | * | * | * | * |
| 2000..... | 0.7 | * | * | * | * | 0.8 | 0.9 | 1.5 | 2.3 | * | * | * |
| 1999..... | 0.8 | * | * | * | * | 0.9 | 1.3 | 2.0 | 2.4 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Systemic lupus erythematosus (M32) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.8 | 0.8 | 1.3 | 1.1 |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.4 | 0.6 | 0.8 | 0.9 | 1.5 | 1.1 |
| 2002..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.8 | 0.9 | 1.3 | 1.4 |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 0.8 | 1.0 | 1.5 | 1.3 |
| 2000..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.6 | 0.6 | 0.9 | 0.9 | 1.4 | 1.2 |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.7 | 1.1 | 1.3 | 1.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.4 | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | * |
| 2000..... | 0.2 | * | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.6 | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.7 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.3 | 0.6 | 0.7 | 1.0 | 1.3 | 1.2 | 1.8 | 1.3 |
| 2003..... | 0.8 | * | * | * | 0.4 | 0.6 | 0.7 | 1.0 | 1.3 | 1.3 | 2.2 | 1.3 |
| 2002..... | 0.8 | * | * | * | 0.4 | 0.5 | 0.8 | 1.0 | 1.3 | 1.4 | 1.8 | 1.8 |
| 2001..... | 0.8 | * | * | * | 0.4 | 0.7 | 0.8 | 1.1 | 1.3 | 1.6 | 2.1 | 1.3 |
| 2000..... | 0.8 | * | * | * | 0.3 | 0.6 | 1.0 | 1.1 | 1.3 | 1.3 | 2.0 | 1.5 |
| 1999..... | 0.8 | * | * | * | 0.3 | 0.6 | 0.9 | 1.1 | 1.1 | 1.8 | 1.8 | 1.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.2 | 1.2 |
| 2003..... | 0.3 | * | * | * | * | 0.2 | 0.2 | 0.4 | 0.6 | 0.7 | 1.5 | 1.2 |
| 2002..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.7 | 1.3 | 1.4 |
| 2001..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.9 | 1.5 | 1.2 |
| 2000..... | 0.4 | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.7 | 0.8 | 1.4 | 1.4 |
| 1999..... | 0.3 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 0.9 | 1.4 | 1.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.2 | * | 0.5 | * |
| 2003..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.4 | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.6 | * |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.4 | 0.6 | * |
| 1999..... | 0.1 | * | * | * | * | * | * | * | 0.3 | 0.3 | 0.7 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.4 | 0.6 | 0.9 | 1.0 | 1.7 | 1.4 |
| 2003..... | 0.6 | * | * | * | * | 0.3 | 0.4 | 0.6 | 1.0 | 1.0 | 2.2 | 1.3 |
| 2002..... | 0.5 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 1.0 | 1.8 | 1.8 |
| 2001..... | 0.6 | * | * | * | * | 0.3 | 0.3 | 0.7 | 1.0 | 1.4 | 2.1 | 1.2 |
| 2000..... | 0.6 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.0 | 1.2 | 1.9 | 1.6 |
| 1999..... | 0.6 | * | * | * | * | 0.2 | 0.4 | 0.7 | 0.9 | 1.5 | 1.8 | 1.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Systemic lupus erythematosus (M32) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | 0.7 | 1.5 | 1.7 | 2.4 | 2.8 | 2.0 | 2.3 | * |
| 2003..... | 1.1 | * | * | * | 0.8 | 1.2 | 1.4 | 2.4 | 2.2 | 2.1 | * | * |
| 2002..... | 1.3 | * | * | * | 0.8 | 1.4 | 2.0 | 2.0 | 2.5 | 2.6 | * | * |
| 2001..... | 1.2 | * | * | * | 0.8 | 1.5 | 2.0 | 2.0 | 2.1 | 1.7 | * | * |
| 2000..... | 1.3 | * | * | * | 0.8 | 1.3 | 2.1 | 2.4 | 2.5 | 1.9 | * | * |
| 1999..... | 1.2 | * | * | * | 0.8 | 1.2 | 2.2 | 2.2 | 1.9 | 2.9 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.4 | * | * | * | * | * | 0.8 | * | * | * | * | * |
| 1999..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | 1.1 | 2.5 | 2.5 | 3.7 | 4.5 | 2.9 | * | * |
| 2003..... | 2.0 | * | * | * | 1.5 | 2.2 | 2.4 | 4.0 | 3.4 | 3.4 | * | * |
| 2002..... | 2.2 | * | * | * | 1.4 | 2.3 | 3.6 | 3.5 | 4.4 | 3.8 | * | * |
| 2001..... | 2.0 | * | * | * | 1.5 | 2.5 | 3.4 | 3.3 | 3.1 | 2.6 | * | * |
| 2000..... | 2.1 | * | * | * | 1.1 | 2.3 | 3.3 | 4.1 | 3.8 | 2.8 | * | * |
| 1999..... | 2.1 | * | * | * | 1.4 | 2.1 | 3.7 | 3.8 | 3.0 | 4.5 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.2 | 4.3 | 9.5 | 24.5 |
| 2003..... | 1.6 | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.2 | 4.6 | 9.4 | 25.9 |
| 2002..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.2 | 4.7 | 9.7 | 27.2 |
| 2001..... | 1.7 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 2.1 | 5.0 | 10.1 | 27.7 |
| 2000..... | 1.7 | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.2 | 5.0 | 10.5 | 28.0 |
| 1999..... | 1.7 | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 2.3 | 5.0 | 9.5 | 27.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.5 | 2.9 | 7.2 | 17.5 |
| 2003..... | 1.0 | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 3.5 | 7.7 | 19.4 |
| 2002..... | 1.0 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.3 | 7.6 | 18.6 |
| 2001..... | 1.0 | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.5 | 3.4 | 8.1 | 20.5 |
| 2000..... | 1.1 | * | * | * | * | 0.3 | 0.4 | 0.8 | 1.6 | 3.4 | 8.8 | 21.1 |
| 1999..... | 1.0 | * | * | * | * | 0.2 | 0.5 | 0.8 | 1.6 | 3.5 | 7.8 | 22.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.8 | 5.6 | 11.1 | 27.7 |
| 2003..... | 2.2 | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 2.9 | 5.6 | 10.5 | 28.8 |
| 2002..... | 2.3 | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 2.8 | 5.9 | 11.1 | 31.0 |
| 2001..... | 2.3 | * | * | * | 0.2 | 0.3 | 0.8 | 1.3 | 2.7 | 6.3 | 11.5 | 30.7 |
| 2000..... | 2.3 | * | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 2.8 | 6.3 | 11.6 | 30.8 |
| 1999..... | 2.3 | * | * | * | 0.2 | 0.3 | 0.7 | 1.7 | 3.0 | 6.3 | 10.6 | 29.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.5 | 1.0 | 2.3 | 4.7 | 7.3 | 12.4 |
| 2003..... | 0.7 | * | * | * | * | * | 0.7 | 1.2 | 2.8 | 3.5 | 8.3 | 14.0 |
| 2002..... | 0.7 | * | * | * | * | * | 0.6 | 1.0 | 2.3 | 4.8 | 6.7 | 12.7 |
| 2001..... | 0.8 | * | * | * | * | * | 0.7 | 1.2 | 2.7 | 5.2 | 7.4 | 18.8 |
| 2000..... | 0.7 | * | * | * | * | 0.3 | 0.4 | 1.1 | 2.3 | 4.6 | 9.3 | 19.9 |
| 1999..... | 0.7 | * | * | * | * | * | 0.5 | 1.2 | 2.6 | 5.5 | 6.9 | 21.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.5 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | 4.4 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.8 | 1.4 | 3.5 | 6.2 | 8.4 | 14.4 |
| 2003..... | 1.1 | * | * | * | * | * | 1.0 | 1.8 | 3.8 | 4.1 | 10.3 | 18.0 |
| 2002..... | 1.0 | * | * | * | * | * | * | 1.4 | 3.3 | 7.5 | 7.9 | * |
| 2001..... | 1.1 | * | * | * | * | * | 0.9 | 1.4 | 3.8 | 6.6 | 9.1 | 19.5 |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.3 | 2.5 | 6.8 | 11.9 | * |
| 1999..... | 1.1 | * | * | * | * | * | * | 1.7 | 3.6 | 6.4 | 6.9 | 27.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.2 | 4.3 | 9.7 | 25.0 |
| 2003..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.2 | 4.7 | 9.4 | 26.4 |
| 2002..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.2 | 4.7 | 9.8 | 27.7 |
| 2001..... | 1.8 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 2.1 | 5.0 | 10.3 | 27.9 |
| 2000..... | 1.8 | * | * | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.2 | 5.0 | 10.5 | 28.2 |
| 1999..... | 1.8 | * | * | * | 0.1 | 0.2 | 0.6 | 1.2 | 2.3 | 5.0 | 9.6 | 27.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.6 | 2.9 | 7.3 | 17.9 |
| 2003..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 3.5 | 7.8 | 20.0 |
| 2002..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 3.4 | 7.8 | 19.0 |
| 2001..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.9 | 1.5 | 3.4 | 8.2 | 20.6 |
| 2000..... | 1.1 | * | * | * | * | 0.3 | 0.4 | 0.7 | 1.6 | 3.4 | 8.9 | 20.7 |
| 1999..... | 1.1 | * | * | * | * | 0.1 | 0.5 | 0.8 | 1.6 | 3.5 | 7.8 | 22.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | 0.1 | 0.3 | 0.8 | 1.4 | 2.7 | 5.5 | 11.2 | 28.2 |
| 2003..... | 2.4 | * | * | * | 0.2 | 0.4 | 0.7 | 1.4 | 2.9 | 5.7 | 10.5 | 29.2 |
| 2002..... | 2.5 | * | * | * | 0.1 | 0.4 | 0.7 | 1.5 | 2.8 | 5.8 | 11.2 | 31.4 |
| 2001..... | 2.5 | * | * | * | 0.2 | 0.3 | 0.8 | 1.3 | 2.6 | 6.3 | 11.6 | 30.9 |
| 2000..... | 2.5 | * | * | * | 0.1 | 0.4 | 0.8 | 1.5 | 2.8 | 6.3 | 11.5 | 31.3 |
| 1999..... | 2.4 | * | * | * | 0.1 | 0.3 | 0.7 | 1.6 | 2.9 | 6.2 | 10.7 | 29.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 1.9 | 4.3 | 10.0 | 26.1 |
| 2003..... | 1.8 | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 2.0 | 4.6 | 9.6 | 26.9 |
| 2002..... | 1.9 | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.1 | 4.7 | 10.0 | 28.5 |
| 2001..... | 1.9 | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 1.9 | 4.8 | 10.6 | 28.8 |
| 2000..... | 1.9 | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.0 | 4.9 | 10.7 | 29.0 |
| 1999..... | 1.8 | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.0 | 4.9 | 9.8 | 28.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.8 | 7.5 | 18.4 |
| 2003..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 3.3 | 7.6 | 20.6 |
| 2002..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.3 | 7.9 | 19.6 |
| 2001..... | 1.2 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.3 | 8.4 | 21.2 |
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.4 | 3.3 | 8.8 | 20.2 |
| 1999..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.4 | 3.4 | 7.9 | 23.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | * | * | 0.5 | 1.1 | 2.4 | 5.5 | 11.6 | 29.5 |
| 2003..... | 2.5 | * | * | * | * | * | 0.5 | 1.0 | 2.6 | 5.7 | 10.8 | 29.6 |
| 2002..... | 2.7 | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.7 | 5.8 | 11.5 | 32.4 |
| 2001..... | 2.6 | * | * | * | * | 0.2 | 0.6 | 1.0 | 2.4 | 6.2 | 12.1 | 31.9 |
| 2000..... | 2.6 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.6 | 6.3 | 11.9 | 32.6 |
| 1999..... | 2.5 | * | * | * | * | 0.2 | 0.4 | 1.1 | 2.6 | 6.2 | 11.0 | 30.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | 0.8 | 1.6 | 3.2 | 4.0 | 5.4 | 7.7 | 17.0 |
| 2003..... | 1.9 | * | * | * | * | 1.0 | 1.6 | 3.3 | 4.2 | 6.1 | 9.2 | 20.8 |
| 2002..... | 1.8 | * | * | * | * | 0.9 | 1.8 | 2.8 | 3.9 | 5.7 | 8.9 | 22.1 |
| 2001..... | 1.8 | * | * | * | * | 0.8 | 1.8 | 3.2 | 3.2 | 6.1 | 8.6 | 22.5 |
| 2000..... | 1.8 | * | * | * | * | 0.9 | 2.0 | 3.2 | 4.0 | 5.8 | 9.3 | 21.9 |
| 1999..... | 1.9 | * | * | * | * | 0.8 | 2.0 | 3.8 | 4.8 | 6.1 | 7.9 | 20.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.8 | 2.4 | 2.9 | 4.0 | 6.0 | * |
| 2003..... | 1.3 | * | * | * | * | * | 1.3 | 1.9 | 3.3 | 6.1 | 10.4 | * |
| 2002..... | 1.2 | * | * | * | * | * | 1.3 | 2.1 | 3.9 | 5.0 | 6.4 | * |
| 2001..... | 1.2 | * | * | * | * | * | 1.4 | 2.7 | 2.4 | 4.3 | 7.5 | * |
| 2000..... | 1.3 | * | * | * | * | * | 1.5 | 1.8 | 3.2 | 5.1 | 10.8 | 27.0 |
| 1999..... | 1.3 | * | * | * | * | * | 1.8 | 2.4 | 3.6 | 4.7 | 7.8 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | * | 0.8 | 2.4 | 3.9 | 4.9 | 6.4 | 8.7 | 18.1 |
| 2003..... | 2.4 | * | * | * | * | 1.4 | 1.8 | 4.5 | 5.0 | 6.2 | 8.5 | 23.9 |
| 2002..... | 2.3 | * | * | * | * | 1.1 | 2.3 | 3.5 | 3.9 | 6.1 | 10.3 | 25.1 |
| 2001..... | 2.3 | * | * | * | * | 1.1 | 2.2 | 3.6 | 3.9 | 7.4 | 9.3 | 24.4 |
| 2000..... | 2.3 | * | * | * | * | 1.0 | 2.4 | 4.4 | 4.6 | 6.2 | 8.5 | 20.1 |
| 1999..... | 2.5 | * | * | * | * | 1.1 | 2.2 | 4.9 | 5.7 | 7.1 | 7.9 | 22.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Dorsopathies (M40-M54) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 7.2 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.7 | 5.7 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.3 | 6.2 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 2.4 | 7.5 |
| 2000..... | 0.3 | * | * | * | * | * | 0.0 | 0.1 | 0.3 | 0.8 | 2.4 | 6.1 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.8 | 1.9 | 5.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 6.8 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.9 | 4.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 5.6 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 6.3 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.4 | 6.5 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.3 | 5.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.5 | 7.5 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.6 | 6.1 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.2 | 6.4 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 7.9 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.6 | 2.4 | 6.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.8 | 5.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.8 | 7.6 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.8 | 6.0 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.3 | 6.4 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 2.4 | 7.9 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 6.4 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 2.0 | 5.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.7 | 7.0 |
| 2003..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.9 | 4.9 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.4 | 5.5 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.5 | 6.6 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 2.4 | 7.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.8 | 2.2 | 5.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.8 | 2.8 | 7.8 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 0.8 | 2.7 | 6.4 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 0.9 | 2.3 | 6.8 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 8.4 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.6 | 2.4 | 6.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.8 | 5.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Soft tissue disorders (M60-M79) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.6 | 1.2 | 2.7 | 5.2 |
| 2003..... | 0.5 | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.3 | 2.6 | 5.5 |
| 2002..... | 0.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.1 | 2.5 | 5.3 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.3 | 4.4 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.4 | 0.5 | 1.0 | 2.2 | 4.3 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 2.1 | 4.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.3 | 3.4 | 7.8 |
| 2003..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.5 | 3.4 | 7.2 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 1.4 | 2.9 | 8.1 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.4 | 3.2 | 6.3 |
| 2000..... | 0.4 | * | * | * | * | * | 0.2 | 0.4 | 0.5 | 1.2 | 2.9 | 6.9 |
| 1999..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 3.1 | 6.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.1 | 2.2 | 4.0 |
| 2003..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 2.1 | 4.8 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.1 | 4.2 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 0.8 | 1.7 | 3.6 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.8 | 1.7 | 3.2 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.5 | 3.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 2.6 | 4.9 |
| 2003..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.5 | 5.5 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.4 | 5.0 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 2.1 | 4.3 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 0.9 | 2.0 | 4.0 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.0 | 3.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 3.4 | 7.2 |
| 2003..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.4 | 3.3 | 7.1 |
| 2002..... | 0.5 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 2.9 | 7.9 |
| 2001..... | 0.4 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 3.1 | 6.1 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.8 | 6.5 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.8 | 6.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.4 | 1.1 | 2.1 | 3.8 |
| 2003..... | 0.5 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.1 | 1.9 | 4.8 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 0.8 | 2.1 | 3.8 |
| 2001..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.5 | 3.5 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.3 | 0.3 | 0.8 | 1.4 | 3.0 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.5 | 3.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 7.5 | 37.1 |
| 2003..... | 1.3 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 | 7.1 | 40.5 |
| 2002..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 7.1 | 41.4 |
| 2001..... | 1.3 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 7.2 | 40.7 |
| 2000..... | 1.2 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.4 | 6.7 | 42.2 |
| 1999..... | 1.1 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 6.8 | 39.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.3 | 0.6 | 1.9 | 6.1 | 20.6 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 2.1 | 5.2 | 22.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 5.4 | 23.7 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 5.4 | 23.8 |
| 2000..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 5.5 | 26.4 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 5.3 | 20.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 8.4 | 44.4 |
| 2003..... | 1.9 | * | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 8.4 | 48.6 |
| 2002..... | 1.9 | * | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 8.3 | 49.0 |
| 2001..... | 1.8 | * | * | * | * | * | * | 0.1 | 0.5 | 1.8 | 8.4 | 47.7 |
| 2000..... | 1.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.4 | 7.5 | 48.6 |
| 1999..... | 1.7 | * | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 7.8 | 46.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 7.5 | 38.6 |
| 2003..... | 1.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 7.1 | 42.1 |
| 2002..... | 1.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.0 | 43.4 |
| 2001..... | 1.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 7.3 | 42.0 |
| 2000..... | 1.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.3 | 6.8 | 43.7 |
| 1999..... | 1.3 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 6.9 | 40.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.6 | 1.8 | 5.9 | 21.0 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 4.9 | 21.4 |
| 2002..... | 0.8 | * | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 5.0 | 23.9 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.4 | 1.5 | 5.3 | 24.1 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 5.3 | 26.3 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 5.2 | 20.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 8.6 | 46.5 |
| 2003..... | 2.2 | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 8.7 | 51.2 |
| 2002..... | 2.2 | * | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 8.4 | 51.9 |
| 2001..... | 2.1 | * | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 8.7 | 49.4 |
| 2000..... | 2.0 | * | * | * | * | * | * | * | 0.3 | 1.4 | 7.7 | 50.8 |
| 1999..... | 2.0 | * | * | * | * | * | * | 0.1 | 0.5 | 1.4 | 8.0 | 48.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | 0.5 | 0.9 | 2.6 | 9.4 | 25.3 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.5 | 1.3 | 2.9 | 7.5 | 27.1 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | 1.2 | 2.9 | 8.8 | 24.2 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 1.0 | 3.4 | 6.8 | 30.0 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | 1.1 | 2.5 | 7.4 | 27.7 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 2.2 | 7.0 | 21.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 3.4 | 11.4 | 22.4 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | 1.8 | 5.2 | 10.1 | 27.7 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 4.1 | 10.0 | 25.2 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.9 | 7.5 | 26.6 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 4.0 | 8.3 | 32.9 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | * | 7.2 | 25.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 2.0 | 8.2 | 26.5 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 6.0 | 26.8 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | * | 8.1 | 23.8 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 3.0 | 6.4 | 31.2 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 6.9 | 25.7 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 2.1 | 6.8 | 19.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Osteoporosis (M80-M81) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 0.3 | 3.3 | 25.1 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | 0.4 | 3.3 | 27.7 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 0.4 | 3.5 | 29.6 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 0.4 | 3.4 | 27.4 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 0.4 | 3.5 | 29.0 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 0.4 | 3.6 | 27.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.3 | 7.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.0 | 8.2 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.1 | 10.0 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.0 | 9.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.3 | 10.3 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 1.3 | 7.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | * | 0.5 | 4.7 | 33.1 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | * | 0.6 | 4.9 | 36.2 |
| 2002..... | 1.3 | * | * | * | * | * | * | * | * | 0.6 | 5.0 | 38.0 |
| 2001..... | 1.2 | * | * | * | * | * | * | * | * | 0.6 | 4.9 | 34.8 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | 0.5 | 4.9 | 36.5 |
| 1999..... | 1.2 | * | * | * | * | * | * | * | * | 0.6 | 5.0 | 35.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | 0.4 | 3.6 | 27.2 |
| 2003..... | 0.9 | * | * | * | * | * | * | * | * | 0.5 | 3.6 | 30.0 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | 0.5 | 3.7 | 32.0 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 0.4 | 3.6 | 29.5 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | 0.4 | 3.8 | 31.3 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 0.4 | 3.9 | 29.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.3 | 8.0 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.1 | 8.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.2 | 10.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.1 | 10.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.4 | 10.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.4 | 7.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | * | * | 0.6 | 5.2 | 35.8 |
| 2003..... | 1.5 | * | * | * | * | * | * | * | * | 0.7 | 5.3 | 39.5 |
| 2002..... | 1.5 | * | * | * | * | * | * | * | * | 0.7 | 5.4 | 41.3 |
| 2001..... | 1.4 | * | * | * | * | * | * | * | * | 0.6 | 5.3 | 37.4 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | * | 0.6 | 5.4 | 39.5 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | * | 0.7 | 5.5 | 38.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.1 | 11.9 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 3.7 | 12.9 |
| 2002..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 3.6 | 11.8 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.4 | 3.9 | 13.3 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.1 | 3.2 | 13.2 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 3.3 | 11.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.3 | 0.5 | 1.7 | 4.9 | 13.2 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.7 | 1.9 | 4.1 | 13.9 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 4.3 | 13.7 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.4 | 14.2 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 4.1 | 16.1 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 4.0 | 13.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.6 | 11.3 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.5 | 12.4 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.2 | 11.0 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.3 | 3.5 | 12.9 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.6 | 12.1 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.4 | 0.9 | 2.8 | 10.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.8 | 11.5 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.6 | 12.1 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 3.3 | 11.4 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 | 3.7 | 12.5 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.0 | 12.5 |
| 1999..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 3.0 | 11.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.5 | 13.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 3.8 | 12.9 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.8 | 3.9 | 13.2 |
| 2001..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 4.2 | 13.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | * | 0.4 | 1.1 | 3.9 | 15.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 1.1 | 3.8 | 12.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 3.4 | 10.8 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.4 | 11.7 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.1 | 2.9 | 10.6 |
| 2001..... | 0.7 | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 | 3.3 | 12.0 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.4 | 11.2 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.6 | 10.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.5 | 0.9 | 2.6 | 8.9 | 20.4 |
| 2003..... | 0.7 | * | * | * | * | * | * | 0.4 | 1.3 | 2.9 | 7.0 | 23.8 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 1.2 | 2.7 | 7.7 | 19.7 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | 1.0 | 3.3 | 6.4 | 24.3 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | 1.0 | 2.4 | 6.9 | 23.8 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | * | 2.2 | 6.2 | 17.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | 3.4 | 10.8 | 21.3 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | 1.8 | 5.2 | 9.5 | 24.5 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | 4.1 | 9.7 | 24.1 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.9 | 7.5 | 24.3 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | 3.9 | 7.7 | 28.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | 6.9 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | * | 7.7 | 20.1 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 5.5 | 23.5 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | * | 6.6 | 18.0 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 2.8 | 5.7 | 24.4 |
| 2000..... | 0.6 | * | * | * | * | * | * | * | * | * | 6.4 | 22.2 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | 2.1 | 5.8 | 15.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XIV. Diseases of the genitourinary system (N00-N98) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.5 | 4.9 | 0.2 | 0.1 | 0.3 | 0.8 | 2.2 | 6.2 | 16.5 | 49.2 | 153.4 | 458.5 |
| 2003..... | 20.7 | 5.1 | 0.1 | 0.1 | 0.3 | 0.9 | 2.2 | 6.0 | 16.7 | 50.7 | 153.8 | 469.5 |
| 2002..... | 20.2 | 4.6 | * | 0.1 | 0.3 | 0.9 | 2.1 | 5.7 | 15.8 | 49.7 | 153.5 | 469.4 |
| 2001..... | 20.0 | 3.9 | 0.2 | * | 0.3 | 0.8 | 2.1 | 5.7 | 15.8 | 51.2 | 149.8 | 476.4 |
| 2000..... | 19.4 | 5.0 | 0.1 | 0.1 | 0.3 | 0.8 | 2.0 | 5.4 | 15.6 | 48.0 | 148.2 | 471.4 |
| 1999..... | 19.0 | 5.1 | 0.2 | 0.1 | 0.2 | 0.8 | 2.0 | 5.0 | 15.1 | 48.1 | 144.7 | 471.1 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 18.5 | 5.8 | * | * | 0.3 | 0.7 | 2.4 | 7.0 | 18.2 | 53.8 | 179.0 | 533.9 |
| 2003..... | 18.6 | 6.0 | * | * | 0.3 | 0.9 | 2.4 | 6.8 | 18.3 | 55.3 | 181.1 | 547.9 |
| 2002..... | 18.1 | 5.6 | * | * | 0.2 | 0.8 | 2.1 | 6.5 | 17.0 | 55.9 | 178.6 | 553.0 |
| 2001..... | 17.7 | 4.2 | * | * | 0.3 | 0.7 | 2.2 | 6.4 | 16.7 | 56.9 | 173.5 | 564.2 |
| 2000..... | 17.2 | 6.1 | * | * | 0.3 | 0.8 | 2.2 | 6.3 | 16.8 | 52.6 | 173.2 | 567.5 |
| 1999..... | 16.9 | 5.0 | * | 0.1 | 0.2 | 0.8 | 2.1 | 5.6 | 16.5 | 53.3 | 172.1 | 558.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 22.5 | 4.0 | * | * | 0.3 | 0.9 | 2.1 | 5.4 | 15.0 | 45.4 | 136.3 | 424.5 |
| 2003..... | 22.6 | 4.3 | * | * | 0.3 | 0.9 | 2.1 | 5.2 | 15.3 | 46.7 | 135.6 | 434.9 |
| 2002..... | 22.3 | 3.6 | * | * | 0.3 | 0.9 | 2.0 | 5.0 | 14.8 | 44.5 | 136.8 | 433.2 |
| 2001..... | 22.2 | 3.6 | * | * | 0.3 | 0.8 | 2.0 | 5.1 | 15.0 | 46.5 | 134.3 | 439.6 |
| 2000..... | 21.5 | 3.9 | * | * | 0.3 | 0.8 | 1.8 | 4.6 | 14.5 | 44.3 | 132.0 | 432.3 |
| 1999..... | 21.1 | 5.1 | * | 0.1 | 0.3 | 0.8 | 1.8 | 4.5 | 13.8 | 43.9 | 127.0 | 436.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.2 | 4.6 | * | * | * | 0.3 | 1.5 | 5.7 | 16.9 | 52.2 | 126.5 | 306.3 |
| 2003..... | 7.1 | 4.4 | * | * | 0.3 | 0.5 | 1.6 | 5.4 | 17.3 | 51.3 | 124.0 | 338.4 |
| 2002..... | 6.6 | 4.7 | * | * | * | 0.6 | 1.4 | 4.9 | 15.7 | 45.9 | 120.8 | 327.6 |
| 2001..... | 6.8 | 2.6 | * | * | * | 0.4 | 1.7 | 5.0 | 15.4 | 52.0 | 130.2 | 361.4 |
| 2000..... | 6.5 | 3.8 | * | * | * | 0.5 | 1.2 | 4.5 | 16.5 | 47.0 | 125.8 | 372.9 |
| 1999..... | 6.5 | 3.3 | * | * | * | 0.4 | 1.7 | 4.7 | 17.0 | 46.2 | 126.4 | 385.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.6 | 5.4 | * | * | * | * | 1.7 | 6.6 | 19.8 | 56.4 | 139.1 | 361.4 |
| 2003..... | 6.4 | 4.5 | * | * | * | * | 1.7 | 6.4 | 18.0 | 54.3 | 141.9 | 394.1 |
| 2002..... | 5.9 | 5.4 | * | * | * | * | 1.6 | 5.3 | 17.2 | 53.5 | 141.2 | 319.6 |
| 2001..... | 5.9 | * | * | * | * | * | 2.0 | 4.9 | 17.5 | 54.1 | 140.0 | 400.0 |
| 2000..... | 5.7 | 5.1 | * | * | * | * | 1.5 | 5.7 | 16.3 | 50.0 | 138.7 | 411.1 |
| 1999..... | 6.0 | * | * | * | * | * | 1.6 | 5.2 | 21.2 | 50.1 | 147.6 | 455.1 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.8 | * | * | * | * | * | 1.2 | 4.8 | 14.3 | 48.8 | 117.8 | 278.0 |
| 2003..... | 7.8 | * | * | * | * | * | 1.5 | 4.4 | 16.8 | 49.0 | 111.7 | 310.2 |
| 2002..... | 7.3 | * | * | * | * | 0.8 | 1.3 | 4.4 | 14.4 | 39.8 | 106.8 | 331.6 |
| 2001..... | 7.8 | * | * | * | * | * | 1.5 | 5.1 | 13.5 | 50.4 | 123.6 | 342.6 |
| 2000..... | 7.3 | * | * | * | * | 0.7 | 1.0 | 3.3 | 16.8 | 44.6 | 117.2 | 354.1 |
| 1999..... | 7.1 | * | * | * | * | * | 1.8 | 4.3 | 13.4 | 43.2 | 112.4 | 351.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XIV. Diseases of the genitourinary system (N00-N98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 22.7 | 5.0 | 0.2 | * | 0.3 | 0.9 | 2.4 | 6.2 | 16.5 | 48.9 | 154.6 | 464.6 |
| 2003..... | 22.7 | 5.3 | 0.2 | 0.1 | 0.3 | 1.0 | 2.3 | 6.0 | 16.6 | 50.5 | 155.0 | 473.8 |
| 2002..... | 22.3 | 4.6 | * | 0.1 | 0.3 | 0.9 | 2.2 | 5.8 | 15.8 | 49.8 | 154.7 | 474.0 |
| 2001..... | 21.9 | 4.2 | 0.2 | * | 0.3 | 0.8 | 2.2 | 5.7 | 15.8 | 51.0 | 150.3 | 479.6 |
| 2000..... | 21.2 | 5.2 | * | 0.1 | 0.3 | 0.8 | 2.1 | 5.5 | 15.4 | 48.0 | 148.8 | 474.3 |
| 1999..... | 20.7 | 5.5 | 0.2 | 0.1 | 0.3 | 0.9 | 2.0 | 5.1 | 14.9 | 48.1 | 145.2 | 472.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 20.6 | 5.9 | * | * | 0.3 | 0.8 | 2.6 | 7.0 | 18.0 | 53.5 | 180.8 | 541.5 |
| 2003..... | 20.6 | 6.4 | * | * | 0.3 | 1.0 | 2.5 | 6.8 | 18.2 | 55.2 | 182.8 | 553.6 |
| 2002..... | 20.1 | 5.7 | * | * | 0.3 | 0.9 | 2.2 | 6.6 | 16.9 | 55.8 | 179.9 | 562.0 |
| 2001..... | 19.5 | 4.3 | * | * | 0.3 | 0.8 | 2.3 | 6.4 | 16.6 | 56.9 | 174.6 | 569.8 |
| 2000..... | 18.9 | 6.1 | * | * | 0.3 | 0.9 | 2.3 | 6.3 | 16.8 | 52.5 | 174.3 | 573.3 |
| 1999..... | 18.4 | 5.1 | * | * | 0.2 | 0.8 | 2.2 | 5.6 | 16.1 | 53.3 | 172.8 | 560.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 24.7 | 4.0 | * | * | 0.3 | 1.0 | 2.2 | 5.4 | 15.0 | 45.0 | 136.9 | 430.2 |
| 2003..... | 24.8 | 4.1 | * | * | 0.3 | 1.0 | 2.1 | 5.3 | 15.1 | 46.5 | 136.5 | 438.8 |
| 2002..... | 24.4 | 3.5 | * | * | 0.3 | 1.0 | 2.1 | 5.1 | 14.8 | 44.8 | 138.0 | 436.0 |
| 2001..... | 24.2 | 4.0 | * | * | 0.3 | 0.8 | 2.1 | 5.0 | 15.1 | 46.2 | 134.3 | 442.0 |
| 2000..... | 23.4 | 4.3 | * | * | 0.3 | 0.8 | 1.9 | 4.7 | 14.2 | 44.2 | 132.2 | 434.2 |
| 1999..... | 22.9 | 5.9 | * | * | 0.3 | 0.9 | 1.8 | 4.5 | 13.8 | 43.8 | 127.2 | 437.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 23.1 | 3.1 | * | * | 0.2 | 0.6 | 1.7 | 4.3 | 12.4 | 42.1 | 146.2 | 455.7 |
| 2003..... | 23.1 | 3.9 | * | * | 0.2 | 0.7 | 1.5 | 4.2 | 12.7 | 43.0 | 146.5 | 465.1 |
| 2002..... | 22.7 | 3.6 | * | * | 0.2 | 0.6 | 1.5 | 4.0 | 12.0 | 42.9 | 146.3 | 462.7 |
| 2001..... | 22.2 | 3.0 | * | * | 0.2 | 0.6 | 1.5 | 3.9 | 11.9 | 43.7 | 141.1 | 467.7 |
| 2000..... | 21.4 | 3.4 | * | * | 0.3 | 0.6 | 1.4 | 3.8 | 11.7 | 40.6 | 140.0 | 462.6 |
| 1999..... | 20.8 | 4.1 | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.4 | 11.4 | 40.8 | 136.0 | 456.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 20.9 | 3.8 | * | * | 0.2 | 0.5 | 1.9 | 4.7 | 13.8 | 46.4 | 173.6 | 534.7 |
| 2003..... | 20.9 | 4.9 | * | * | 0.2 | 0.7 | 1.7 | 4.6 | 14.0 | 47.5 | 174.2 | 552.8 |
| 2002..... | 20.3 | 4.3 | * | * | 0.2 | 0.6 | 1.5 | 4.5 | 12.6 | 48.7 | 172.0 | 551.9 |
| 2001..... | 19.8 | 3.2 | * | * | 0.2 | 0.5 | 1.7 | 4.4 | 12.7 | 49.7 | 166.6 | 561.9 |
| 2000..... | 19.1 | 4.2 | * | * | 0.2 | 0.6 | 1.5 | 4.2 | 12.8 | 45.9 | 165.9 | 566.3 |
| 1999..... | 18.3 | 4.3 | * | * | 0.2 | 0.6 | 1.4 | 3.7 | 12.5 | 45.9 | 162.5 | 543.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 25.3 | 2.4 | * | * | 0.2 | 0.8 | 1.6 | 3.8 | 11.1 | 38.4 | 127.6 | 420.3 |
| 2003..... | 25.3 | 2.9 | * | * | 0.3 | 0.7 | 1.4 | 3.8 | 11.6 | 39.2 | 127.8 | 426.5 |
| 2002..... | 25.0 | 2.8 | * | * | 0.2 | 0.6 | 1.4 | 3.5 | 11.5 | 37.9 | 129.0 | 424.2 |
| 2001..... | 24.5 | 2.8 | * | * | 0.3 | 0.7 | 1.3 | 3.4 | 11.1 | 38.6 | 124.2 | 428.5 |
| 2000..... | 23.7 | 2.6 | * | * | 0.3 | 0.6 | 1.3 | 3.4 | 10.6 | 36.1 | 123.0 | 420.8 |
| 1999..... | 23.2 | 3.8 | * | * | 0.2 | 0.6 | 1.2 | 3.1 | 10.3 | 36.6 | 118.6 | 422.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XIV. Diseases of the genitourinary system (N00-N98) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 26.6 | 12.0 | * | * | 0.6 | 2.3 | 6.6 | 19.7 | 51.9 | 115.4 | 269.6 | 643.6 |
| 2003..... | 26.9 | 11.5 | * | * | 0.7 | 2.6 | 7.3 | 19.3 | 51.2 | 124.0 | 274.4 | 652.3 |
| 2002..... | 26.0 | 8.7 | * | * | 0.8 | 2.6 | 6.4 | 18.7 | 47.9 | 119.2 | 270.9 | 667.5 |
| 2001..... | 26.0 | 9.7 | * | * | 0.5 | 2.2 | 6.3 | 18.8 | 48.2 | 123.9 | 274.7 | 670.5 |
| 2000..... | 25.2 | 13.2 | * | * | 0.5 | 2.2 | 6.4 | 18.2 | 47.0 | 120.4 | 270.1 | 645.2 |
| 1999..... | 25.2 | 11.6 | * | * | 0.7 | 2.3 | 6.4 | 18.0 | 44.1 | 119.8 | 266.7 | 685.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 24.3 | 13.3 | * | * | 0.7 | 2.4 | 7.3 | 24.0 | 58.3 | 130.1 | 305.7 | 729.6 |
| 2003..... | 24.9 | 13.0 | * | * | 0.7 | 2.8 | 7.9 | 24.0 | 59.7 | 142.7 | 325.8 | 689.3 |
| 2002..... | 24.0 | 10.9 | * | * | * | 2.3 | 6.7 | 22.6 | 56.6 | 135.6 | 309.8 | 804.1 |
| 2001..... | 23.0 | 9.6 | * | * | * | 2.7 | 5.9 | 22.3 | 52.7 | 137.9 | 304.3 | 764.6 |
| 2000..... | 22.7 | 14.3 | * | * | * | 2.2 | 7.4 | 23.0 | 52.8 | 126.8 | 312.5 | 726.8 |
| 1999..... | 23.2 | 8.7 | * | * | * | 2.0 | 7.4 | 21.8 | 48.5 | 134.1 | 330.2 | 812.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 28.6 | 10.6 | * | * | * | 2.3 | 6.0 | 16.1 | 46.7 | 104.8 | 249.2 | 609.6 |
| 2003..... | 28.8 | 10.0 | * | * | 0.7 | 2.5 | 6.8 | 15.2 | 44.5 | 110.7 | 245.4 | 638.0 |
| 2002..... | 27.9 | 6.3 | * | * | 1.0 | 2.8 | 6.2 | 15.4 | 40.9 | 107.6 | 248.9 | 615.4 |
| 2001..... | 28.8 | 9.7 | * | * | * | 1.7 | 6.6 | 15.8 | 44.6 | 114.0 | 258.1 | 635.6 |
| 2000..... | 27.6 | 12.0 | * | * | * | 2.1 | 5.4 | 14.1 | 42.3 | 116.0 | 246.3 | 614.9 |
| 1999..... | 27.0 | 14.6 | * | * | 0.8 | 2.5 | 5.5 | 14.7 | 40.6 | 109.7 | 231.1 | 637.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of urinary system (N00-N39) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 20.2 | 4.8 | 0.2 | 0.1 | 0.3 | 0.8 | 2.2 | 6.0 | 16.3 | 48.5 | 150.8 | 448.9 |
| 2003..... | 20.3 | 5.0 | 0.1 | 0.1 | 0.3 | 0.9 | 2.1 | 5.9 | 16.5 | 50.0 | 151.5 | 460.5 |
| 2002..... | 19.9 | 4.6 | * | 0.1 | 0.3 | 0.8 | 2.0 | 5.6 | 15.7 | 49.1 | 151.2 | 460.8 |
| 2001..... | 19.7 | 3.7 | 0.2 | * | 0.3 | 0.7 | 2.1 | 5.6 | 15.6 | 50.5 | 147.6 | 467.4 |
| 2000..... | 19.1 | 4.8 | 0.1 | 0.1 | 0.3 | 0.7 | 2.0 | 5.3 | 15.4 | 47.4 | 145.9 | 462.5 |
| 1999..... | 18.7 | 4.7 | 0.1 | 0.1 | 0.2 | 0.8 | 1.9 | 4.9 | 14.8 | 47.5 | 142.3 | 462.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 18.0 | 5.8 | * | * | 0.3 | 0.7 | 2.4 | 6.9 | 17.8 | 52.8 | 174.3 | 513.6 |
| 2003..... | 18.2 | 5.9 | * | * | 0.3 | 0.9 | 2.4 | 6.8 | 18.1 | 54.4 | 177.2 | 528.3 |
| 2002..... | 17.7 | 5.6 | * | * | 0.2 | 0.8 | 2.1 | 6.4 | 16.8 | 55.0 | 174.6 | 533.8 |
| 2001..... | 17.3 | 4.2 | * | * | 0.3 | 0.7 | 2.2 | 6.3 | 16.5 | 56.0 | 169.8 | 542.9 |
| 2000..... | 16.8 | 6.0 | * | * | 0.3 | 0.8 | 2.2 | 6.2 | 16.5 | 51.7 | 169.2 | 546.9 |
| 1999..... | 16.4 | 5.0 | * | 0.1 | 0.2 | 0.8 | 2.1 | 5.5 | 16.2 | 52.4 | 167.7 | 537.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 22.2 | 3.8 | * | * | 0.2 | 0.9 | 2.0 | 5.2 | 14.8 | 44.9 | 135.1 | 419.9 |
| 2003..... | 22.4 | 4.1 | * | * | 0.3 | 0.9 | 1.9 | 5.0 | 15.1 | 46.2 | 134.3 | 430.5 |
| 2002..... | 22.0 | 3.5 | * | * | 0.3 | 0.9 | 1.9 | 4.9 | 14.6 | 44.2 | 135.7 | 429.1 |
| 2001..... | 22.0 | 3.2 | * | * | 0.3 | 0.8 | 1.9 | 4.9 | 14.8 | 46.0 | 133.0 | 435.8 |
| 2000..... | 21.2 | 3.6 | * | * | 0.3 | 0.7 | 1.7 | 4.4 | 14.3 | 43.8 | 130.6 | 428.1 |
| 1999..... | 20.9 | 4.3 | * | 0.1 | 0.3 | 0.8 | 1.7 | 4.4 | 13.5 | 43.4 | 125.7 | 432.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.1 | 4.5 | * | * | * | 0.3 | 1.4 | 5.5 | 16.7 | 51.2 | 125.1 | 299.7 |
| 2003..... | 7.0 | 4.3 | * | * | 0.3 | 0.5 | 1.5 | 5.2 | 17.3 | 50.3 | 121.8 | 331.1 |
| 2002..... | 6.4 | 4.6 | * | * | * | 0.6 | 1.4 | 4.8 | 15.5 | 45.1 | 118.3 | 323.7 |
| 2001..... | 6.7 | 2.6 | * | * | * | 0.4 | 1.7 | 4.8 | 15.3 | 50.9 | 128.8 | 356.4 |
| 2000..... | 6.3 | 3.8 | * | * | * | 0.5 | 1.2 | 4.4 | 16.2 | 45.9 | 124.0 | 364.3 |
| 1999..... | 6.4 | 3.0 | * | * | * | 0.4 | 1.6 | 4.6 | 16.8 | 45.3 | 124.8 | 379.6 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.5 | 5.4 | * | * | * | * | 1.7 | 6.3 | 19.7 | 55.2 | 137.0 | 346.0 |
| 2003..... | 6.3 | 4.5 | * | * | * | * | 1.6 | 6.4 | 17.9 | 52.8 | 136.9 | 378.7 |
| 2002..... | 5.7 | 5.4 | * | * | * | * | 1.6 | 5.3 | 17.0 | 52.2 | 136.2 | 314.6 |
| 2001..... | 5.8 | * | * | * | * | * | 2.0 | 4.8 | 17.4 | 52.7 | 137.7 | 394.2 |
| 2000..... | 5.5 | 5.1 | * | * | * | * | 1.5 | 5.6 | 16.1 | 48.5 | 136.7 | 389.0 |
| 1999..... | 5.9 | * | * | * | * | * | 1.6 | 5.2 | 20.8 | 48.6 | 145.5 | 448.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.7 | * | * | * | * | * | 1.2 | 4.7 | 14.0 | 48.1 | 116.8 | 275.9 |
| 2003..... | 7.7 | * | * | * | * | * | 1.4 | 4.1 | 16.8 | 48.4 | 111.4 | 307.0 |
| 2002..... | 7.2 | * | * | * | * | 0.7 | 1.2 | 4.3 | 14.2 | 39.5 | 106.0 | 328.3 |
| 2001..... | 7.6 | * | * | * | * | * | 1.4 | 4.8 | 13.4 | 49.4 | 122.7 | 337.9 |
| 2000..... | 7.2 | * | * | * | * | * | 0.8 | 3.2 | 16.3 | 43.8 | 115.6 | 352.2 |
| 1999..... | 7.0 | * | * | * | * | * | 1.7 | 4.1 | 13.2 | 42.7 | 111.0 | 346.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of urinary system (N00-N39) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 22.3 | 4.9 | 0.2 | * | 0.3 | 0.9 | 2.3 | 6.0 | 16.2 | 48.1 | 151.9 | 455.0 |
| 2003..... | 22.4 | 5.2 | 0.2 | 0.1 | 0.3 | 1.0 | 2.2 | 5.9 | 16.4 | 49.8 | 152.6 | 464.7 |
| 2002..... | 21.9 | 4.5 | * | 0.1 | 0.3 | 0.9 | 2.1 | 5.7 | 15.6 | 49.2 | 152.4 | 465.2 |
| 2001..... | 21.6 | 3.9 | 0.2 | * | 0.3 | 0.8 | 2.1 | 5.6 | 15.6 | 50.3 | 148.0 | 470.4 |
| 2000..... | 20.8 | 5.0 | * | 0.1 | 0.3 | 0.8 | 2.0 | 5.4 | 15.3 | 47.3 | 146.4 | 465.4 |
| 1999..... | 20.3 | 5.0 | * | 0.1 | 0.3 | 0.8 | 1.9 | 5.0 | 14.6 | 47.4 | 142.6 | 464.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 20.0 | 5.9 | * | * | 0.3 | 0.8 | 2.5 | 6.9 | 17.7 | 52.4 | 176.0 | 521.0 |
| 2003..... | 20.1 | 6.3 | * | * | 0.3 | 1.0 | 2.5 | 6.8 | 18.0 | 54.4 | 178.9 | 533.8 |
| 2002..... | 19.6 | 5.7 | * | * | 0.3 | 0.9 | 2.2 | 6.5 | 16.7 | 54.9 | 175.9 | 542.4 |
| 2001..... | 19.0 | 4.2 | * | * | 0.3 | 0.8 | 2.2 | 6.4 | 16.4 | 56.0 | 170.9 | 547.9 |
| 2000..... | 18.4 | 6.0 | * | * | 0.3 | 0.8 | 2.3 | 6.2 | 16.5 | 51.7 | 170.2 | 552.8 |
| 1999..... | 17.9 | 5.1 | * | * | 0.2 | 0.8 | 2.1 | 5.5 | 15.8 | 52.4 | 168.3 | 538.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 24.4 | 3.8 | * | * | 0.3 | 1.0 | 2.1 | 5.2 | 14.8 | 44.5 | 135.7 | 425.5 |
| 2003..... | 24.5 | 4.0 | * | * | 0.3 | 1.0 | 2.0 | 5.1 | 14.9 | 46.0 | 135.1 | 434.4 |
| 2002..... | 24.1 | 3.3 | * | * | 0.3 | 0.9 | 2.0 | 4.9 | 14.6 | 44.5 | 136.9 | 431.9 |
| 2001..... | 23.9 | 3.6 | * | * | 0.3 | 0.8 | 2.0 | 4.9 | 14.9 | 45.6 | 133.0 | 438.2 |
| 2000..... | 23.1 | 3.9 | * | * | 0.3 | 0.7 | 1.8 | 4.5 | 14.1 | 43.7 | 130.9 | 430.0 |
| 1999..... | 22.7 | 5.0 | * | * | 0.3 | 0.8 | 1.7 | 4.4 | 13.5 | 43.4 | 126.0 | 434.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 22.7 | 3.1 | * | * | 0.2 | 0.6 | 1.7 | 4.2 | 12.2 | 41.5 | 143.6 | 445.9 |
| 2003..... | 22.7 | 3.9 | * | * | 0.2 | 0.7 | 1.4 | 4.1 | 12.5 | 42.4 | 144.1 | 455.6 |
| 2002..... | 22.3 | 3.5 | * | * | 0.2 | 0.6 | 1.4 | 3.9 | 11.9 | 42.4 | 144.1 | 453.9 |
| 2001..... | 21.8 | 2.8 | * | * | 0.2 | 0.6 | 1.5 | 3.8 | 11.7 | 43.0 | 138.8 | 458.5 |
| 2000..... | 21.0 | 3.3 | * | * | 0.3 | 0.6 | 1.4 | 3.7 | 11.6 | 40.0 | 137.6 | 453.7 |
| 1999..... | 20.4 | 3.8 | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.3 | 11.2 | 40.3 | 133.5 | 448.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 20.2 | 3.8 | * | * | 0.2 | 0.5 | 1.8 | 4.6 | 13.5 | 45.4 | 168.9 | 513.6 |
| 2003..... | 20.3 | 4.9 | * | * | 0.2 | 0.7 | 1.6 | 4.5 | 13.8 | 46.6 | 170.4 | 532.1 |
| 2002..... | 19.8 | 4.3 | * | * | 0.2 | 0.6 | 1.5 | 4.4 | 12.5 | 48.0 | 168.3 | 532.0 |
| 2001..... | 19.2 | 3.1 | * | * | 0.2 | 0.5 | 1.7 | 4.3 | 12.5 | 48.8 | 162.8 | 539.4 |
| 2000..... | 18.6 | 4.1 | * | * | 0.2 | 0.6 | 1.5 | 4.2 | 12.6 | 45.2 | 161.9 | 545.9 |
| 1999..... | 17.8 | 4.3 | * | * | 0.2 | 0.6 | 1.4 | 3.6 | 12.3 | 45.1 | 158.0 | 521.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 25.0 | 2.4 | * | * | 0.2 | 0.7 | 1.5 | 3.7 | 10.9 | 38.0 | 126.4 | 415.6 |
| 2003..... | 25.0 | 2.9 | * | * | 0.2 | 0.7 | 1.2 | 3.6 | 11.3 | 38.8 | 126.3 | 421.9 |
| 2002..... | 24.7 | 2.6 | * | * | 0.2 | 0.6 | 1.3 | 3.4 | 11.3 | 37.7 | 127.8 | 420.2 |
| 2001..... | 24.2 | 2.5 | * | * | 0.2 | 0.7 | 1.3 | 3.3 | 10.9 | 38.1 | 122.9 | 424.8 |
| 2000..... | 23.4 | 2.4 | * | * | 0.3 | 0.5 | 1.3 | 3.2 | 10.5 | 35.6 | 121.7 | 416.5 |
| 1999..... | 23.0 | 3.2 | * | * | 0.2 | 0.6 | 1.1 | 3.0 | 10.1 | 36.3 | 117.4 | 418.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Diseases of urinary system (N00-N39) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 26.2 | 11.5 | * | * | 0.6 | 2.3 | 6.5 | 19.3 | 51.5 | 113.7 | 266.3 | 634.4 |
| 2003..... | 26.6 | 11.1 | * | * | 0.7 | 2.5 | 7.1 | 19.0 | 50.7 | 123.0 | 272.2 | 646.4 |
| 2002..... | 25.7 | 8.7 | * | * | 0.7 | 2.5 | 6.2 | 18.5 | 47.5 | 117.7 | 268.0 | 657.3 |
| 2001..... | 25.7 | 9.3 | * | * | 0.5 | 2.2 | 6.0 | 18.4 | 47.5 | 123.1 | 271.1 | 661.1 |
| 2000..... | 24.9 | 12.7 | * | * | 0.5 | 2.1 | 6.1 | 18.1 | 46.5 | 119.2 | 267.0 | 636.0 |
| 1999..... | 24.8 | 10.4 | * | * | 0.6 | 2.2 | 6.3 | 17.6 | 43.4 | 118.1 | 262.7 | 675.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 23.9 | 13.3 | * | * | 0.6 | 2.3 | 7.2 | 23.6 | 57.6 | 128.3 | 298.9 | 712.3 |
| 2003..... | 24.6 | 12.7 | * | * | 0.7 | 2.8 | 7.9 | 23.8 | 59.0 | 141.7 | 320.9 | 675.5 |
| 2002..... | 23.6 | 10.9 | * | * | * | 2.3 | 6.6 | 22.5 | 56.1 | 133.2 | 303.0 | 785.5 |
| 2001..... | 22.6 | 9.6 | * | * | * | 2.7 | 5.7 | 22.0 | 51.8 | 136.9 | 298.6 | 745.0 |
| 2000..... | 22.3 | 14.3 | * | * | * | 2.2 | 7.4 | 22.9 | 51.8 | 125.2 | 306.7 | 705.7 |
| 1999..... | 22.8 | 8.7 | * | * | * | 2.0 | 7.4 | 21.5 | 47.5 | 131.7 | 324.3 | 789.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 28.3 | 9.7 | * | * | * | 2.3 | 5.9 | 15.6 | 46.6 | 103.3 | 248.0 | 603.6 |
| 2003..... | 28.4 | 9.3 | * | * | * | 2.2 | 6.5 | 14.9 | 44.0 | 109.7 | 244.8 | 635.1 |
| 2002..... | 27.5 | 6.3 | * | * | 0.9 | 2.7 | 5.7 | 15.1 | 40.6 | 106.8 | 248.2 | 608.3 |
| 2001..... | 28.4 | 9.1 | * | * | * | 1.7 | 6.3 | 15.2 | 44.0 | 113.3 | 255.7 | 630.1 |
| 2000..... | 27.3 | 10.9 | * | * | * | 2.0 | 5.0 | 14.0 | 42.3 | 115.0 | 244.7 | 610.1 |
| 1999..... | 26.6 | 12.1 | * | * | * | 2.3 | 5.2 | 14.3 | 40.2 | 108.5 | 228.3 | 632.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Glomerular and renal tubulo-interstitial diseases (N00-N15) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.4 | 4.8 | 12.8 |
| 2003..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.5 | 4.8 | 13.4 |
| 2002..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 4.8 | 13.9 |
| 2001..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 1.6 | 4.6 | 13.8 |
| 2000..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 4.8 | 15.7 |
| 1999..... | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.7 | 5.4 | 16.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 5.2 | 17.1 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.4 | 5.2 | 18.0 |
| 2002..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.7 | 5.4 | 18.1 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.8 | 5.4 | 17.2 |
| 2000..... | 0.6 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 5.4 | 20.0 |
| 1999..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 6.7 | 22.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 1.3 | 4.5 | 10.8 |
| 2003..... | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.5 | 4.5 | 11.4 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.3 | 4.5 | 12.1 |
| 2001..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 4.1 | 12.4 |
| 2000..... | 0.8 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.6 | 4.5 | 14.0 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.8 | 4.5 | 13.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.4 | 4.7 | 12.6 |
| 2003..... | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 4.6 | 13.3 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 4.8 | 13.9 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.5 | 4.5 | 14.0 |
| 2000..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.6 | 15.3 |
| 1999..... | 0.8 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.5 | 5.3 | 16.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 5.1 | 16.5 |
| 2003..... | 0.7 | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.4 | 4.8 | 18.1 |
| 2002..... | 0.7 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.7 | 5.2 | 18.4 |
| 2001..... | 0.7 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.7 | 5.2 | 17.5 |
| 2000..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 5.3 | 19.4 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.3 | 0.5 | 1.3 | 6.6 | 22.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.4 | 1.3 | 4.5 | 10.8 |
| 2003..... | 0.8 | * | * | * | * | * | * | 0.3 | 0.5 | 1.5 | 4.4 | 11.1 |
| 2002..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 4.5 | 11.9 |
| 2001..... | 0.8 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 4.1 | 12.5 |
| 2000..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 4.2 | 13.6 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.3 | 0.5 | 1.7 | 4.5 | 14.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Glomerular and renal tubulo-interstitial diseases (N00-N15) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.1 | 2.0 | 5.8 | 15.0 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.6 | 1.0 | 1.9 | 7.0 | 15.2 |
| 2002..... | 0.6 | * | * | * | * | * | * | 0.6 | 0.9 | 1.3 | 5.5 | 15.7 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.6 | 0.9 | 2.5 | 5.9 | 12.5 |
| 2000..... | 0.8 | * | * | * | * | * | 0.4 | 0.9 | 1.0 | 2.7 | 7.2 | 18.4 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.6 | 1.3 | 3.4 | 5.9 | 20.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | * | * | 1.6 | * | 6.3 | 22.4 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | * | 10.4 | * |
| 2002..... | 0.5 | * | * | * | * | * | * | * | * | * | 7.6 | * |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | 3.9 | 6.6 | * |
| 2000..... | 0.7 | * | * | * | * | * | * | 1.1 | * | 3.0 | 7.1 | 24.7 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 4.1 | 7.5 | 35.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | * | * | * | * | 5.5 | 12.0 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 5.1 | 14.0 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | * | 4.3 | 14.6 |
| 2001..... | 0.6 | * | * | * | * | * | * | * | * | * | 5.6 | 12.0 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | 2.4 | 7.3 | 16.1 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 2.9 | 5.1 | 14.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Glomerular diseases (N00-N07) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.6 | 3.3 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.5 | 3.1 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.7 | 3.3 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.7 | 1.5 | 3.3 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.7 | 3.6 |
| 1999..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 2.0 | 3.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.9 | 4.0 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.7 | 4.1 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.9 | 4.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.9 | 2.0 | 4.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.8 | 2.1 | 5.4 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.7 | 5.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.4 | 2.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.6 |
| 2002..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 2.8 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.2 | 2.7 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.9 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 2.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.5 | 3.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 2.8 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.6 | 2.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.7 | 1.4 | 3.3 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 3.2 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 3.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 3.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.6 | 1.5 | 3.8 |
| 2002..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.7 | 4.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.8 | 2.0 | 4.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 4.5 |
| 1999..... | 0.3 | * | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 2.6 | 5.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 2.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.4 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.4 | 2.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.1 | 2.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.4 | 2.7 |
| 1999..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|

Acute and rapidly progressive nephritic syndrome (N00-N01)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, both sexes.

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, male.

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, female.

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, both sexes.

Non-Hispanic white, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, male.

Non-Hispanic white, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified glomerular diseases (N02-N03,N05-N07) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.2 | 2.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.2 | 2.5 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.3 | 2.3 |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.6 | 1.1 | 2.5 |
| 2000..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.3 | 2.4 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.5 | 2.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.3 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.5 | 1.4 | 3.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.6 | 3.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.8 | 1.6 | 3.4 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.5 | 3.6 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.3 | 0.4 | 2.1 | 4.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.0 | 2.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.1 | 1.9 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 0.8 | 2.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.0 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 2.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.2 | 2.4 |
| 2003..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 2.3 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.5 | 1.2 | 2.1 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.1 | 2.5 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.1 |
| 1999..... | 0.2 | * | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.4 | 2.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 3.0 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.2 | 3.5 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.4 | 3.2 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.8 | 1.6 | 3.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.6 | 1.5 | 2.9 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 2.0 | 4.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 2.1 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.9 | 1.8 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 1.7 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.7 | 2.0 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.0 | 1.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | * | 0.5 | 1.0 | 2.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Renal tubulo-interstitial diseases (N10-N15) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 3.2 | 9.5 |
| 2003..... | 0.5 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 3.2 | 10.4 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 3.2 | 10.7 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 3.1 | 10.5 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 3.1 | 12.1 |
| 1999..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.0 | 3.4 | 12.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.4 | 0.9 | 3.2 | 13.0 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 3.5 | 13.9 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 1.0 | 3.5 | 13.8 | |
| 2001..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.9 | 3.4 | 12.5 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.6 | 3.3 | 14.7 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 4.0 | 17.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.8 | 3.1 | 7.9 |
| 2003..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 3.1 | 8.8 |
| 2002..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.0 | 9.3 |
| 2001..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 3.0 | 9.7 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 3.0 | 11.1 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.0 | 11.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.9 | 3.2 | 9.5 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.8 | 3.2 | 10.5 |
| 2002..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 1.0 | 3.2 | 11.0 |
| 2001..... | 0.5 | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.8 | 3.1 | 10.7 |
| 2000..... | 0.5 | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 3.0 | 12.0 |
| 1999..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.4 | 12.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.8 | 3.3 | 12.9 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.2 | 0.8 | 3.3 | 14.4 |
| 2002..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.4 | 14.2 |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | 0.8 | 3.2 | 12.7 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.5 | 3.2 | 14.9 |
| 1999..... | 0.5 | * | * | * | * | * | * | * | 0.3 | 0.7 | 4.0 | 17.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.9 | 3.2 | 7.9 |
| 2003..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.0 | 8.7 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | 0.3 | 1.0 | 3.1 | 9.5 |
| 2001..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 3.0 | 9.9 |
| 2000..... | 0.6 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.8 | 10.9 |
| 1999..... | 0.7 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 3.1 | 11.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Renal failure (N17-N19) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 14.2 | 4.0 | * | * | 0.2 | 0.6 | 1.7 | 4.8 | 13.4 | 37.9 | 106.6 | 282.9 |
| 2003..... | 14.4 | 4.2 | * | 0.1 | 0.2 | 0.7 | 1.7 | 4.8 | 13.4 | 39.3 | 107.8 | 289.7 |
| 2002..... | 13.9 | 4.1 | * | * | 0.2 | 0.6 | 1.6 | 4.6 | 12.7 | 38.6 | 107.2 | 284.8 |
| 2001..... | 13.6 | 3.1 | * | * | 0.2 | 0.5 | 1.6 | 4.5 | 12.7 | 39.4 | 102.6 | 284.0 |
| 2000..... | 13.0 | 4.0 | * | * | 0.2 | 0.6 | 1.6 | 4.2 | 12.5 | 37.2 | 98.9 | 273.7 |
| 1999..... | 12.4 | 4.3 | * | 0.1 | 0.2 | 0.6 | 1.5 | 3.9 | 11.7 | 36.4 | 95.3 | 264.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 13.8 | 5.1 | * | * | 0.2 | 0.5 | 2.0 | 5.7 | 14.8 | 42.9 | 132.1 | 367.4 |
| 2003..... | 14.1 | 4.8 | * | * | 0.2 | 0.7 | 2.0 | 5.7 | 15.2 | 44.5 | 134.8 | 383.1 |
| 2002..... | 13.7 | 5.1 | * | * | 0.2 | 0.6 | 1.8 | 5.4 | 14.0 | 45.0 | 134.0 | 380.7 |
| 2001..... | 13.2 | 3.5 | * | * | 0.2 | 0.5 | 1.8 | 5.3 | 13.8 | 45.2 | 128.0 | 388.7 |
| 2000..... | 12.6 | 5.1 | * | * | 0.2 | 0.6 | 1.8 | 5.1 | 13.8 | 42.1 | 125.1 | 374.2 |
| 1999..... | 12.1 | 4.5 | * | * | 0.1 | 0.5 | 1.7 | 4.5 | 13.1 | 41.5 | 122.3 | 364.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 14.6 | 3.0 | * | * | 0.2 | 0.6 | 1.5 | 3.9 | 12.0 | 33.7 | 89.5 | 244.9 |
| 2003..... | 14.6 | 3.7 | * | * | 0.2 | 0.6 | 1.5 | 3.8 | 11.8 | 34.9 | 89.7 | 248.4 |
| 2002..... | 14.2 | 3.0 | * | * | 0.2 | 0.7 | 1.4 | 3.8 | 11.6 | 33.2 | 89.5 | 243.2 |
| 2001..... | 14.0 | 2.7 | * | * | 0.2 | 0.5 | 1.5 | 3.7 | 11.8 | 34.5 | 85.9 | 240.1 |
| 2000..... | 13.3 | 2.8 | * | * | 0.2 | 0.5 | 1.3 | 3.4 | 11.2 | 33.3 | 81.8 | 232.8 |
| 1999..... | 12.7 | 4.1 | * | * | 0.2 | 0.6 | 1.3 | 3.3 | 10.4 | 32.2 | 77.8 | 224.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 5.2 | 3.9 | * | * | * | 0.3 | 1.2 | 4.4 | 14.0 | 40.5 | 91.0 | 179.3 |
| 2003..... | 5.3 | 3.8 | * | * | * | 0.4 | 1.3 | 4.4 | 14.8 | 39.0 | 88.5 | 230.6 |
| 2002..... | 4.8 | 4.2 | * | * | * | 0.4 | 1.1 | 4.2 | 13.2 | 35.7 | 86.9 | 198.4 |
| 2001..... | 4.9 | * | * | * | * | 0.3 | 1.2 | 3.7 | 12.5 | 40.5 | 93.2 | 217.6 |
| 2000..... | 4.5 | 2.9 | * | * | * | 0.4 | 1.0 | 3.3 | 13.3 | 36.9 | 83.0 | 225.6 |
| 1999..... | 4.5 | * | * | * | * | * | 1.2 | 3.5 | 13.6 | 34.4 | 86.9 | 228.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 5.0 | 5.2 | * | * | * | * | 1.4 | 5.2 | 16.4 | 43.5 | 108.4 | 224.2 |
| 2003..... | 5.1 | * | * | * | * | * | 1.5 | 5.3 | 15.8 | 41.9 | 105.1 | 295.5 |
| 2002..... | 4.6 | 4.9 | * | * | * | * | 1.3 | 4.8 | 14.4 | 40.7 | 109.2 | 232.6 |
| 2001..... | 4.6 | * | * | * | * | * | 1.4 | 3.7 | 14.6 | 42.5 | 110.4 | 278.1 |
| 2000..... | 4.3 | * | * | * | * | * | 1.3 | 4.2 | 13.8 | 40.5 | 99.3 | 280.1 |
| 1999..... | 4.5 | * | * | * | * | * | 1.1 | 4.3 | 17.2 | 38.2 | 109.0 | 330.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | * | * | * | 0.9 | 3.6 | 11.9 | 38.0 | 79.0 | 156.3 |
| 2003..... | 5.6 | * | * | * | * | * | 0.9 | 3.5 | 13.9 | 36.6 | 77.2 | 197.7 |
| 2002..... | 5.0 | * | * | * | * | 0.6 | 0.8 | 3.6 | 12.0 | 31.8 | 71.7 | 181.5 |
| 2001..... | 5.2 | * | * | * | * | * | 1.0 | 3.7 | 10.5 | 38.9 | 81.6 | 188.0 |
| 2000..... | 4.8 | * | * | * | * | * | * | 2.4 | 12.8 | 34.0 | 72.0 | 198.8 |
| 1999..... | 4.5 | * | * | * | * | * | 1.3 | 2.7 | 10.4 | 31.3 | 72.2 | 178.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Renal failure (N17-N19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.7 | 4.1 | * | * | 0.2 | 0.6 | 1.8 | 4.8 | 13.3 | 37.6 | 107.2 | 287.2 |
| 2003..... | 15.7 | 4.3 | * | * | 0.2 | 0.7 | 1.8 | 4.8 | 13.2 | 39.2 | 108.5 | 291.4 |
| 2002..... | 15.3 | 4.1 | * | * | 0.2 | 0.7 | 1.7 | 4.6 | 12.6 | 38.6 | 107.9 | 287.4 |
| 2001..... | 14.9 | 3.3 | * | * | 0.2 | 0.6 | 1.7 | 4.6 | 12.7 | 39.2 | 102.7 | 285.9 |
| 2000..... | 14.1 | 4.2 | * | * | 0.2 | 0.6 | 1.6 | 4.3 | 12.3 | 37.1 | 99.3 | 274.9 |
| 1999..... | 13.5 | 4.7 | * | * | 0.2 | 0.6 | 1.5 | 3.9 | 11.5 | 36.4 | 95.4 | 265.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 15.3 | 5.1 | * | * | 0.2 | 0.6 | 2.1 | 5.8 | 14.6 | 42.7 | 133.1 | 374.0 |
| 2003..... | 15.5 | 5.0 | * | * | 0.2 | 0.8 | 2.1 | 5.8 | 15.0 | 44.6 | 136.0 | 386.2 |
| 2002..... | 15.1 | 5.1 | * | * | 0.2 | 0.6 | 1.8 | 5.4 | 13.9 | 45.1 | 134.7 | 386.2 |
| 2001..... | 14.6 | 3.5 | * | * | 0.2 | 0.6 | 1.8 | 5.4 | 13.7 | 45.3 | 128.5 | 392.7 |
| 2000..... | 13.8 | 5.2 | * | * | 0.2 | 0.6 | 1.9 | 5.1 | 13.7 | 42.0 | 125.9 | 377.7 |
| 1999..... | 13.2 | 4.6 | * | * | 0.1 | 0.6 | 1.8 | 4.5 | 12.7 | 41.5 | 122.6 | 364.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 16.0 | 3.1 | * | * | 0.2 | 0.6 | 1.6 | 4.0 | 12.0 | 33.3 | 89.8 | 248.5 |
| 2003..... | 15.9 | 3.5 | * | * | 0.2 | 0.7 | 1.5 | 3.8 | 11.6 | 34.7 | 90.1 | 249.7 |
| 2002..... | 15.5 | 2.9 | * | * | 0.2 | 0.7 | 1.5 | 3.8 | 11.4 | 33.2 | 90.2 | 244.8 |
| 2001..... | 15.2 | 3.1 | * | * | 0.2 | 0.5 | 1.6 | 3.7 | 11.8 | 34.1 | 85.8 | 241.5 |
| 2000..... | 14.4 | 3.1 | * | * | 0.2 | 0.5 | 1.4 | 3.5 | 11.0 | 33.2 | 82.0 | 233.4 |
| 1999..... | 13.8 | 4.8 | * | * | 0.2 | 0.6 | 1.3 | 3.3 | 10.4 | 32.1 | 77.8 | 225.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 15.5 | 2.5 | * | * | 0.1 | 0.4 | 1.3 | 3.2 | 9.6 | 31.5 | 100.0 | 278.6 |
| 2003..... | 15.5 | 3.2 | * | * | 0.1 | 0.5 | 1.1 | 3.1 | 9.8 | 32.5 | 101.1 | 282.6 |
| 2002..... | 15.2 | 3.1 | * | * | 0.1 | 0.4 | 1.1 | 3.0 | 9.2 | 32.3 | 100.8 | 277.6 |
| 2001..... | 14.6 | 2.2 | * | * | 0.2 | 0.4 | 1.1 | 3.0 | 9.2 | 32.7 | 95.1 | 276.1 |
| 2000..... | 13.9 | 2.5 | * | * | 0.2 | 0.4 | 1.1 | 2.8 | 9.0 | 30.7 | 92.3 | 265.7 |
| 1999..... | 13.1 | 3.5 | * | * | 0.1 | 0.4 | 0.9 | 2.4 | 8.4 | 30.4 | 87.8 | 251.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 15.3 | 3.0 | * | * | 0.2 | 0.3 | 1.4 | 3.7 | 10.9 | 36.5 | 127.2 | 368.8 |
| 2003..... | 15.4 | 4.0 | * | * | 0.2 | 0.5 | 1.3 | 3.7 | 11.2 | 37.8 | 129.3 | 381.9 |
| 2002..... | 15.0 | 3.9 | * | * | * | 0.4 | 1.2 | 3.5 | 10.0 | 38.8 | 128.8 | 377.8 |
| 2001..... | 14.5 | 2.3 | * | * | * | 0.3 | 1.3 | 3.6 | 10.3 | 39.1 | 121.7 | 387.6 |
| 2000..... | 13.7 | 3.3 | * | * | 0.2 | 0.4 | 1.2 | 3.3 | 10.3 | 36.5 | 119.4 | 373.4 |
| 1999..... | 13.0 | 3.8 | * | * | * | 0.4 | 1.1 | 2.8 | 9.6 | 35.9 | 114.4 | 353.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 15.7 | 2.0 | * | * | * | 0.4 | 1.1 | 2.6 | 8.4 | 27.2 | 81.5 | 238.2 |
| 2003..... | 15.6 | 2.3 | * | * | * | 0.5 | 0.9 | 2.5 | 8.5 | 28.0 | 82.0 | 238.9 |
| 2002..... | 15.3 | 2.2 | * | * | * | 0.4 | 1.0 | 2.5 | 8.5 | 26.7 | 82.0 | 234.3 |
| 2001..... | 14.7 | 2.1 | * | * | 0.2 | 0.4 | 0.9 | 2.4 | 8.1 | 27.3 | 77.5 | 229.7 |
| 2000..... | 14.0 | * | * | * | * | 0.3 | 0.9 | 2.3 | 7.7 | 25.8 | 74.4 | 222.2 |
| 1999..... | 13.2 | 3.1 | * | * | * | 0.4 | 0.8 | 2.1 | 7.3 | 25.8 | 70.3 | 210.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Renal failure (N17-N19) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 20.7 | 10.1 | * | * | 0.5 | 1.8 | 5.5 | 16.7 | 45.0 | 95.4 | 202.5 | 432.6 |
| 2003..... | 21.1 | 9.3 | * | * | 0.5 | 2.2 | 6.3 | 16.5 | 43.3 | 104.3 | 207.1 | 442.3 |
| 2002..... | 20.2 | 8.0 | * | * | 0.6 | 2.0 | 5.4 | 16.0 | 41.4 | 100.9 | 202.7 | 436.5 |
| 2001..... | 20.0 | 8.5 | * | * | 0.4 | 1.7 | 5.3 | 15.8 | 42.1 | 102.9 | 201.9 | 428.9 |
| 2000..... | 19.2 | 11.4 | * | * | 0.4 | 1.8 | 5.3 | 15.6 | 40.7 | 99.9 | 194.6 | 404.2 |
| 1999..... | 18.8 | 10.2 | * | * | 0.5 | 1.9 | 5.4 | 15.2 | 37.0 | 95.3 | 193.0 | 428.7 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 19.6 | 12.1 | * | * | * | 2.0 | 6.2 | 21.2 | 50.2 | 108.6 | 233.0 | 514.2 |
| 2003..... | 20.5 | 9.6 | * | * | * | 2.5 | 7.3 | 21.3 | 52.1 | 121.6 | 244.4 | 521.0 |
| 2002..... | 19.4 | 10.0 | * | * | * | 1.8 | 6.1 | 20.1 | 49.8 | 115.4 | 233.3 | 562.3 |
| 2001..... | 18.4 | 8.9 | * | * | * | 2.1 | 5.1 | 19.3 | 45.7 | 114.3 | 233.5 | 514.0 |
| 2000..... | 17.9 | 13.6 | * | * | * | 1.9 | 6.5 | 19.9 | 45.0 | 104.2 | 231.7 | 487.3 |
| 1999..... | 17.7 | 8.7 | * | * | * | 1.8 | 6.4 | 19.0 | 40.5 | 102.5 | 246.7 | 519.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 21.7 | 8.0 | * | * | * | 1.6 | 4.9 | 12.9 | 40.9 | 86.0 | 185.3 | 400.4 |
| 2003..... | 21.7 | 9.0 | * | * | * | 1.9 | 5.5 | 12.4 | 36.3 | 92.0 | 186.1 | 411.9 |
| 2002..... | 20.9 | * | * | * | * | 2.1 | 4.8 | 12.5 | 34.7 | 90.6 | 185.4 | 388.4 |
| 2001..... | 21.4 | 8.1 | * | * | * | 1.4 | 5.5 | 12.8 | 39.2 | 94.9 | 184.1 | 397.4 |
| 2000..... | 20.3 | 9.2 | * | * | * | 1.7 | 4.1 | 11.8 | 37.3 | 96.9 | 173.7 | 373.3 |
| 1999..... | 19.8 | 11.8 | * | * | * | 2.0 | 4.5 | 12.0 | 34.3 | 90.2 | 163.0 | 394.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Urolithiasis (N20-N23) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 2.3 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 2.6 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.6 | 2.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 1.9 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.9 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.9 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 2.0 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.0 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.8 | 2.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 0.8 | 1.9 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.8 | 2.2 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.8 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 1.8 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 1.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Urinary tract infection, site not specified (N39.0) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.7 | 7.7 | 35.6 | 143.5 |
| 2003..... | 4.8 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.9 | 7.9 | 35.5 | 147.7 |
| 2002..... | 4.8 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.9 | 7.7 | 35.6 | 152.2 |
| 2001..... | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 8.2 | 37.0 | 160.4 |
| 2000..... | 4.9 | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.7 | 7.3 | 38.4 | 163.4 |
| 1999..... | 5.0 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 8.0 | 38.1 | 172.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 6.8 | 32.1 | 117.7 |
| 2003..... | 3.0 | * | * | * | * | * | 0.1 | 0.4 | 1.7 | 6.9 | 33.1 | 114.6 |
| 2002..... | 3.0 | * | * | * | * | * | 0.2 | 0.4 | 1.6 | 6.7 | 31.0 | 123.5 |
| 2001..... | 3.0 | * | * | * | * | * | 0.1 | 0.4 | 1.6 | 7.4 | 32.6 | 125.8 |
| 2000..... | 3.1 | * | * | * | * | * | 0.1 | 0.5 | 1.5 | 6.7 | 34.3 | 139.8 |
| 1999..... | 3.1 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 7.5 | 34.7 | 137.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.9 | 8.4 | 38.0 | 155.1 |
| 2003..... | 6.5 | * | * | * | * | * | 0.2 | 0.7 | 2.1 | 8.7 | 37.0 | 162.3 |
| 2002..... | 6.5 | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 8.5 | 38.7 | 164.7 |
| 2001..... | 6.7 | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 8.9 | 39.8 | 174.9 |
| 2000..... | 6.6 | * | * | * | * | * | 0.1 | 0.6 | 1.9 | 7.8 | 41.1 | 173.0 |
| 1999..... | 6.9 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 8.4 | 40.3 | 185.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | * | 0.5 | 1.3 | 7.7 | 27.6 | 102.7 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | 1.4 | 8.3 | 27.8 | 83.4 |
| 2002..... | 1.2 | * | * | * | * | * | * | * | 1.5 | 6.3 | 24.0 | 104.2 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | 2.1 | 7.0 | 28.5 | 115.7 |
| 2000..... | 1.3 | * | * | * | * | * | * | * | 1.8 | 6.5 | 32.4 | 120.1 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | 1.5 | 7.7 | 29.8 | 124.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | * | 8.5 | 20.7 | 98.1 |
| 2003..... | 0.8 | * | * | * | * | * | * | * | * | 8.3 | 25.7 | 61.6 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 8.0 | 17.2 | 51.9 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | 6.2 | 20.1 | 83.8 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | 5.5 | 27.5 | 84.6 |
| 1999..... | 0.9 | * | * | * | * | * | * | * | * | 7.2 | 28.2 | 92.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | * | * | * | * | 7.1 | 32.4 | 105.2 |
| 2003..... | 1.7 | * | * | * | * | * | * | * | * | 8.2 | 29.2 | 94.5 |
| 2002..... | 1.7 | * | * | * | * | * | * | * | * | 4.9 | 28.7 | 130.1 |
| 2001..... | 1.9 | * | * | * | * | * | * | * | 2.1 | 7.6 | 34.2 | 131.2 |
| 2000..... | 1.9 | * | * | * | * | * | * | * | 2.5 | 7.3 | 35.7 | 137.5 |
| 1999..... | 1.8 | * | * | * | * | * | * | * | * | 8.1 | 30.9 | 140.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Urinary tract infection, site not specified (N39.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.8 | 7.6 | 36.0 | 145.1 |
| 2003..... | 5.3 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.9 | 7.8 | 35.8 | 150.0 |
| 2002..... | 5.3 | * | * | * | * | * | 0.2 | 0.5 | 1.9 | 7.8 | 36.1 | 154.0 |
| 2001..... | 5.4 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.8 | 8.3 | 37.2 | 161.6 |
| 2000..... | 5.4 | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.7 | 7.3 | 38.6 | 164.8 |
| 1999..... | 5.5 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.9 | 8.0 | 38.4 | 173.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 6.6 | 32.8 | 118.4 |
| 2003..... | 3.4 | * | * | * | * | * | 0.2 | 0.4 | 1.7 | 6.8 | 33.5 | 116.6 |
| 2002..... | 3.3 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 6.6 | 31.6 | 126.6 |
| 2001..... | 3.3 | * | * | * | * | * | 0.1 | 0.4 | 1.5 | 7.5 | 33.0 | 127.1 |
| 2000..... | 3.4 | * | * | * | * | * | 0.1 | 0.5 | 1.6 | 6.8 | 34.5 | 141.8 |
| 1999..... | 3.4 | * | * | * | * | * | 0.1 | 0.4 | 1.8 | 7.5 | 34.8 | 139.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | * | * | 0.1 | 0.2 | 0.7 | 1.9 | 8.5 | 38.2 | 157.0 |
| 2003..... | 7.2 | * | * | * | * | * | 0.2 | 0.7 | 2.2 | 8.7 | 37.4 | 164.7 |
| 2002..... | 7.2 | * | * | * | * | * | 0.2 | 0.6 | 2.1 | 8.7 | 39.1 | 165.8 |
| 2001..... | 7.4 | * | * | * | * | * | 0.2 | 0.6 | 2.0 | 9.0 | 40.0 | 175.9 |
| 2000..... | 7.2 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 7.8 | 41.2 | 174.1 |
| 1999..... | 7.5 | * | * | * | * | * | 0.2 | 0.6 | 1.9 | 8.4 | 40.7 | 187.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.8 | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.6 | 7.2 | 35.0 | 144.7 |
| 2003..... | 5.8 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 7.2 | 34.8 | 149.6 |
| 2002..... | 5.8 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 7.2 | 35.0 | 152.5 |
| 2001..... | 5.9 | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.6 | 7.6 | 35.8 | 159.2 |
| 2000..... | 5.8 | * | * | * | * | 0.1 | 0.1 | 0.5 | 1.5 | 6.7 | 37.2 | 163.0 |
| 1999..... | 6.0 | * | * | * | * | 0.1 | 0.1 | 0.4 | 1.6 | 7.2 | 37.2 | 171.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.2 | 0.5 | 1.4 | 6.0 | 31.6 | 116.4 |
| 2003..... | 3.6 | * | * | * | * | * | 0.2 | 0.4 | 1.5 | 6.0 | 32.1 | 118.6 |
| 2002..... | 3.6 | * | * | * | * | * | 0.1 | 0.4 | 1.5 | 6.0 | 30.3 | 124.3 |
| 2001..... | 3.5 | * | * | * | * | * | 0.2 | 0.3 | 1.3 | 6.7 | 32.0 | 123.2 |
| 2000..... | 3.6 | * | * | * | * | * | * | 0.4 | 1.3 | 6.1 | 32.8 | 139.7 |
| 1999..... | 3.6 | * | * | * | * | * | * | 0.4 | 1.6 | 6.3 | 33.1 | 133.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.8 | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 8.2 | 37.4 | 157.4 |
| 2003..... | 7.9 | * | * | * | * | * | 0.1 | 0.6 | 1.9 | 8.2 | 36.6 | 163.2 |
| 2002..... | 7.9 | * | * | * | * | * | 0.1 | 0.5 | 1.9 | 8.3 | 38.3 | 164.7 |
| 2001..... | 8.1 | * | * | * | * | * | 0.2 | 0.5 | 1.8 | 8.4 | 38.3 | 174.2 |
| 2000..... | 7.9 | * | * | * | * | * | 0.1 | 0.5 | 1.7 | 7.2 | 40.0 | 172.5 |
| 1999..... | 8.2 | * | * | * | * | * | 0.2 | 0.5 | 1.7 | 7.9 | 40.0 | 186.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Urinary tract infection, site not specified (N39.0) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | * | * | 0.4 | 1.4 | 4.0 | 13.2 | 52.5 | 175.3 |
| 2003..... | 4.3 | * | * | * | * | * | * | 1.2 | 4.4 | 14.7 | 54.7 | 180.0 |
| 2002..... | 4.3 | * | * | * | * | * | 0.4 | 1.0 | 3.8 | 13.5 | 54.5 | 193.9 |
| 2001..... | 4.5 | * | * | * | * | * | * | 1.2 | 3.4 | 14.6 | 58.9 | 210.7 |
| 2000..... | 4.4 | * | * | * | * | * | * | 1.1 | 3.8 | 13.8 | 60.3 | 200.0 |
| 1999..... | 4.6 | * | * | * | * | * | 0.4 | 1.2 | 3.8 | 16.6 | 57.1 | 215.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | * | 1.2 | 4.2 | 13.7 | 53.6 | 165.6 |
| 2003..... | 2.9 | * | * | * | * | * | * | * | 3.9 | 15.5 | 59.4 | 126.8 |
| 2002..... | 3.0 | * | * | * | * | * | * | * | 3.6 | 13.8 | 55.4 | 190.3 |
| 2001..... | 3.1 | * | * | * | * | * | * | 1.0 | 3.7 | 15.0 | 53.7 | 206.7 |
| 2000..... | 3.2 | * | * | * | * | * | * | 1.4 | 4.3 | 14.4 | 62.3 | 178.5 |
| 1999..... | 3.6 | * | * | * | * | * | * | * | 4.1 | 20.7 | 62.5 | 218.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 5.2 | * | * | * | * | * | * | 1.5 | 3.8 | 12.9 | 51.8 | 179.1 |
| 2003..... | 5.6 | * | * | * | * | * | * | 1.5 | 4.8 | 14.1 | 52.0 | 200.6 |
| 2002..... | 5.4 | * | * | * | * | * | * | 1.2 | 3.9 | 13.3 | 54.1 | 195.2 |
| 2001..... | 5.8 | * | * | * | * | * | * | 1.4 | 3.1 | 14.3 | 61.8 | 212.2 |
| 2000..... | 5.5 | * | * | * | * | * | * | * | 3.3 | 13.4 | 59.2 | 208.0 |
| 1999..... | 5.5 | * | * | * | * | * | * | 1.5 | 3.6 | 13.7 | 54.1 | 214.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other diseases of urinary system (N28-N36,N39.1-N39.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.9 | 7.7 |
| 2003..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 2.6 | 7.1 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 7.5 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.7 | 6.9 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.7 | 7.7 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 2.6 | 7.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 1.3 | 3.8 | 9.1 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.6 | 1.2 | 3.3 | 9.9 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 3.3 | 9.0 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.2 | 3.2 | 8.7 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.3 | 3.5 | 10.6 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 3.2 | 10.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.3 | 7.1 |
| 2003..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.2 | 6.0 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.3 | 0.9 | 2.3 | 6.8 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.3 | 6.2 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 2.2 | 6.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.2 | 5.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.4 | 1.0 | 2.8 | 7.7 |
| 2003..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.7 | 7.2 |
| 2002..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.3 | 1.0 | 2.6 | 7.4 |
| 2001..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 7.0 |
| 2000..... | 0.4 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.9 | 2.6 | 7.5 |
| 1999..... | 0.4 | * | * | * | * | * | 0.1 | 0.1 | 0.4 | 0.9 | 2.4 | 7.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 3.9 | 9.3 |
| 2003..... | 0.5 | * | * | * | * | * | * | * | 0.5 | 1.1 | 3.3 | 9.9 |
| 2002..... | 0.4 | * | * | * | * | * | * | 0.3 | 0.4 | 1.1 | 3.2 | 8.8 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.3 | 8.8 |
| 2000..... | 0.5 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 1.1 | 3.4 | 10.5 |
| 1999..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.1 | 10.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.0 | 7.0 |
| 2003..... | 0.5 | * | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 2.3 | 6.0 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | 0.2 | 0.8 | 2.2 | 6.8 |
| 2001..... | 0.4 | * | * | * | * | * | * | 0.1 | 0.4 | 0.7 | 2.2 | 6.2 |
| 2000..... | 0.4 | * | * | * | * | * | * | 0.2 | 0.4 | 0.7 | 2.1 | 6.3 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | 0.4 | 0.7 | 2.0 | 5.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Disorders of female genital organs (N70-N98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 2.6 |
| 2003..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.8 | 2.6 |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | * | 0.1 | 0.6 | 2.4 |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.7 | 2.1 |
| 2000..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | * | 0.2 | 0.7 | 2.6 |
| 1999..... | 0.1 | * | * | * | * | * | 0.0 | * | 0.1 | 0.2 | 0.7 | 2.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 3.7 |
| 2003..... | 0.3 | * | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.3 | 3.7 |
| 2002..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | * | 0.3 | 1.0 | 3.4 |
| 2001..... | 0.2 | * | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 2.9 |
| 2000..... | 0.3 | * | * | * | * | * | 0.1 | 0.2 | * | 0.5 | 1.2 | 3.6 |
| 1999..... | 0.2 | * | * | * | * | * | 0.1 | * | 0.2 | 0.4 | 1.1 | 2.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.5 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.8 | 2.8 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.3 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.0 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.2 | 0.7 | 2.6 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | 0.1 | 0.2 | 0.6 | 2.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 3.6 |
| 2003..... | 0.3 | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.4 | 4.0 |
| 2002..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.1 | 3.3 |
| 2001..... | 0.2 | * | * | * | * | * | * | * | * | 0.4 | 1.1 | 2.8 |
| 2000..... | 0.3 | * | * | * | * | * | * | 0.2 | * | 0.4 | 1.2 | 3.7 |
| 1999..... | 0.2 | * | * | * | * | * | * | * | 0.2 | 0.3 | 1.0 | 2.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Noninflammatory disorders of female genital tract (N80-N98) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.5 | 1.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.6 | 1.9 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.6 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.4 | 1.2 |
| 2000..... | 0.1 | * | * | * | * | * | * | 0.1 | * | 0.1 | 0.5 | 1.9 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.4 | 1.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.8 | 2.7 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | 0.3 | 0.9 | 2.7 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.2 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.7 |
| 2000..... | 0.2 | * | * | * | * | * | * | 0.1 | * | 0.3 | 0.8 | 2.7 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | 0.2 | 0.7 | 2.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.5 | 1.8 |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.1 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.5 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.4 | 1.1 |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | 0.1 | 0.5 | 2.0 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.4 | 1.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | * | * | * | * | * | * | 0.9 | 2.6 |
| 2003..... | 0.2 | * | * | * | * | * | * | * | * | * | 1.0 | 3.0 |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.7 | 2.1 |
| 2001..... | 0.1 | * | * | * | * | * | * | * | * | 0.3 | 0.7 | 1.6 |
| 2000..... | 0.2 | * | * | * | * | * | * | * | * | 0.2 | 0.8 | 2.8 |
| 1999..... | 0.1 | * | * | * | * | * | * | * | * | * | 0.6 | 2.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XV. Pregnancy, childbirth and the puerperium (000-099) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.4 | 0.2 | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | 0.3 | 0.6 | 0.4 | 0.1 | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.3 | 0.5 | 0.2 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.5 | 0.3 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.3 | 0.5 | 0.3 | * | * | * | * | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | ... | ... | * | 0.8 | 1.5 | 0.7 | 0.4 | * | * | * | * |
| 2003..... | 0.4 | ... | ... | * | 0.6 | 1.2 | 0.7 | 0.1 | * | * | * | * |
| 2002..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | * | * | * | * |
| 2001..... | 0.3 | ... | ... | * | 0.6 | 1.0 | 0.5 | * | * | * | * | * |
| 2000..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | * | * | * | * |
| 1999..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | * | * | * | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | ... | ... | * | 0.4 | 0.6 | 0.4 | * | * | * | * | * |
| 2003..... | 0.3 | ... | ... | * | 0.3 | 0.6 | 0.5 | * | * | * | * | * |
| 2002..... | 0.2 | ... | ... | * | * | 0.4 | 0.4 | * | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | 0.4 | 0.6 | * | * | * | * | * | * |
| 2000..... | 0.2 | ... | ... | * | 0.4 | 0.6 | 0.4 | * | * | * | * | * |
| 1999..... | 0.2 | ... | ... | * | * | 0.4 | 0.5 | * | * | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | ... | ... | * | 0.9 | 1.4 | 0.9 | * | * | * | * | * |
| 2003..... | 0.5 | ... | ... | * | 0.7 | 1.4 | 1.0 | * | * | * | * | * |
| 2002..... | 0.4 | ... | ... | * | * | 0.8 | 0.8 | * | * | * | * | * |
| 2001..... | 0.5 | ... | ... | * | 0.8 | 1.2 | * | * | * | * | * | * |
| 2000..... | 0.5 | ... | ... | * | 0.8 | 1.2 | 0.9 | * | * | * | * | * |
| 1999..... | 0.4 | ... | ... | * | * | 0.9 | 1.0 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XV. Pregnancy, childbirth and the puerperium (000-099) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.3 | 0.2 | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | 0.3 | 0.6 | 0.4 | 0.1 | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.3 | 0.5 | 0.2 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.2 | 0.5 | 0.2 | * | * | * | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.8 | 1.5 | 0.7 | 0.4 | * | * | * | * |
| 2003..... | 0.3 | ... | ... | * | 0.5 | 1.1 | 0.7 | 0.1 | * | * | * | * |
| 2002..... | 0.2 | ... | ... | * | 0.4 | 0.9 | 0.4 | * | * | * | * | * |
| 2001..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.4 | * | * | * | * | * |
| 2000..... | 0.3 | ... | ... | * | 0.4 | 0.9 | 0.5 | * | * | * | * | * |
| 1999..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.5 | * | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.2 | 0.5 | 0.2 | 0.1 | * | * | * | * |
| 2003..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.1 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.1 | * | * | * | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | ... | ... | * | 0.5 | 1.0 | 0.5 | 0.3 | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.4 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | 0.2 | 0.6 | 0.3 | * | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | 0.3 | 0.6 | 0.2 | * | * | * | * | * |
| 2000..... | 0.2 | ... | ... | * | 0.3 | 0.6 | 0.3 | * | * | * | * | * |
| 1999..... | 0.2 | ... | ... | * | 0.2 | 0.6 | 0.3 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XV. Pregnancy, childbirth and the puerperium (000-099) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | ... | ... | * | 1.2 | 2.0 | 0.8 | 0.5 | * | * | * | * |
| 2003..... | 0.5 | ... | ... | * | 0.7 | 1.6 | 1.1 | * | * | * | * | * |
| 2002..... | 0.4 | ... | ... | * | 0.8 | 1.2 | 0.7 | * | * | * | * | * |
| 2001..... | 0.4 | ... | ... | * | 0.8 | 1.3 | 0.7 | * | * | * | * | * |
| 2000..... | 0.4 | ... | ... | * | 0.6 | 1.3 | 0.7 | * | * | * | * | * |
| 1999..... | 0.5 | ... | ... | * | 0.8 | 1.3 | 0.7 | * | * | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | ... | ... | * | 2.4 | 3.9 | 1.5 | 0.9 | * | * | * | * |
| 2003..... | 1.0 | ... | ... | * | 1.4 | 3.0 | 2.0 | * | * | * | * | * |
| 2002..... | 0.8 | ... | ... | * | 1.5 | 2.4 | 1.4 | * | * | * | * | * |
| 2001..... | 0.8 | ... | ... | * | 1.6 | 2.4 | 1.4 | * | * | * | * | * |
| 2000..... | 0.8 | ... | ... | * | 1.2 | 2.4 | 1.3 | * | * | * | * | * |
| 1999..... | 0.9 | ... | ... | * | 1.6 | 2.5 | 1.4 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pregnancy with abortive outcome (000-007)

All origins 3/, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for pregnancy with abortive outcome, all origins, both sexes.

All origins 3/, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for pregnancy with abortive outcome, all origins, male.

All origins 3/, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for pregnancy with abortive outcome, all origins, female.

Hispanic, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for pregnancy with abortive outcome, Hispanic, both sexes.

Hispanic, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for pregnancy with abortive outcome, Hispanic, male.

Hispanic, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for pregnancy with abortive outcome, Hispanic, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other direct obstetric deaths (O10-O92) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 2003..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.3 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.4 | 0.8 | 0.4 | * | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.5 | * | * | * | * | * |
| 2002..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.4 | * | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | 0.4 | 0.8 | 0.3 | * | * | * | * | * |
| 2000..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.4 | * | * | * | * | * |
| 1999..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.4 | * | * | * | * | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | * | 0.3 | * | * | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | * | 0.4 | 0.4 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | 0.4 | 0.5 | * | * | * | * | * | * |
| 2000..... | 0.2 | ... | ... | * | * | 0.4 | * | * | * | * | * | * |
| 1999..... | 0.2 | ... | ... | * | * | 0.3 | 0.4 | * | * | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.3 | ... | ... | * | * | 0.7 | * | * | * | * | * | * |
| 2003..... | 0.4 | ... | ... | * | * | 0.9 | 0.9 | * | * | * | * | * |
| 2002..... | 0.3 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.4 | ... | ... | * | 0.8 | 1.1 | * | * | * | * | * | * |
| 2000..... | 0.4 | ... | ... | * | * | 0.9 | * | * | * | * | * | * |
| 1999..... | 0.3 | ... | ... | * | * | 0.8 | 0.8 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other direct obstetric deaths (O10-O92) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 2003..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.2 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.4 | 0.8 | 0.3 | * | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.5 | * | * | * | * | * |
| 2002..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.3 | * | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.3 | * | * | * | * | * |
| 2000..... | 0.2 | ... | ... | * | 0.3 | 0.7 | 0.3 | * | * | * | * | * |
| 1999..... | 0.2 | ... | ... | * | 0.4 | 0.7 | 0.4 | * | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | 0.1 | 0.3 | 0.1 | * | * | * | * | * |
| 2003..... | 0.1 | ... | ... | * | 0.1 | 0.2 | 0.1 | * | * | * | * | * |
| 2002..... | 0.0 | ... | ... | * | * | 0.2 | 0.1 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.1 | 0.2 | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.1 | 0.2 | 0.1 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.1 | 0.2 | 0.1 | * | * | * | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | 0.2 | 0.5 | 0.2 | * | * | * | * | * |
| 2003..... | 0.1 | ... | ... | * | 0.2 | 0.3 | 0.2 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | * | 0.4 | 0.2 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.2 | 0.5 | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | 0.2 | 0.4 | 0.2 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other direct obstetric deaths (O10-O92) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.7 | 1.3 | * | * | * | * | * | * |
| 2003..... | 0.4 | ... | ... | * | 0.4 | 1.1 | 0.8 | * | * | * | * | * |
| 2002..... | 0.3 | ... | ... | * | 0.6 | 0.9 | 0.5 | * | * | * | * | * |
| 2001..... | 0.3 | ... | ... | * | 0.5 | 1.0 | 0.5 | * | * | * | * | * |
| 2000..... | 0.3 | ... | ... | * | 0.5 | 0.9 | 0.4 | * | * | * | * | * |
| 1999..... | 0.3 | ... | ... | * | 0.6 | 1.0 | 0.6 | * | * | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | ... | ... | * | 1.3 | 2.5 | * | * | * | * | * | * |
| 2003..... | 0.7 | ... | ... | * | 0.8 | 2.2 | 1.5 | * | * | * | * | * |
| 2002..... | 0.6 | ... | ... | * | 1.2 | 1.8 | 1.0 | * | * | * | * | * |
| 2001..... | 0.6 | ... | ... | * | 1.1 | 1.8 | 0.9 | * | * | * | * | * |
| 2000..... | 0.5 | ... | ... | * | 1.0 | 1.6 | 0.8 | * | * | * | * | * |
| 1999..... | 0.6 | ... | ... | * | 1.2 | 1.9 | 1.1 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|

Complications predominately related to the puerperium (085-092)

Non-Hispanic 4/, both sexes

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|-----|-----|-----|---|---|---|---|---|
| 2004..... | 0.0 | ... | ... | * | * | 0.1 | 0.1 | * | * | * | * | * |
| 2003..... | 0.0 | ... | ... | * | * | 0.1 | 0.1 | * | * | * | * | * |
| 2002..... | 0.0 | ... | ... | * | * | 0.1 | 0.1 | * | * | * | * | * |
| 2001..... | 0.0 | ... | ... | * | 0.1 | 0.1 | 0.1 | * | * | * | * | * |
| 2000..... | 0.0 | ... | ... | * | 0.1 | 0.1 | * | * | * | * | * | * |
| 1999..... | 0.0 | ... | ... | * | * | 0.1 | * | * | * | * | * | * |

Non-Hispanic 4/, male

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

Non-Hispanic 4/, female

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|-----|-----|-----|---|---|---|---|---|
| 2004..... | 0.1 | ... | ... | * | * | 0.3 | 0.1 | * | * | * | * | * |
| 2003..... | 0.1 | ... | ... | * | * | 0.2 | 0.1 | * | * | * | * | * |
| 2002..... | 0.1 | ... | ... | * | * | 0.2 | 0.1 | * | * | * | * | * |
| 2001..... | 0.1 | ... | ... | * | 0.1 | 0.2 | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.1 | 0.2 | * | * | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | * | 0.2 | * | * | * | * | * | * |

Non-Hispanic white, both sexes

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|---|-----|---|---|---|---|---|---|
| 2004..... | 0.0 | ... | ... | * | * | 0.1 | * | * | * | * | * | * |
| 2003..... | 0.0 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | ... | ... | * | * | 0.1 | * | * | * | * | * | * |
| 2000..... | 0.0 | ... | ... | * | * | 0.1 | * | * | * | * | * | * |
| 1999..... | 0.0 | ... | ... | * | * | 0.1 | * | * | * | * | * | * |

Non-Hispanic white, male

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 2004..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2003..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2002..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2001..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 2000..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| 1999..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |

Non-Hispanic white, female

| | | | | | | | | | | | | |
|-----------|-----|-----|-----|---|---|-----|---|---|---|---|---|---|
| 2004..... | 0.0 | ... | ... | * | * | 0.2 | * | * | * | * | * | * |
| 2003..... | 0.0 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.0 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.0 | ... | ... | * | * | 0.2 | * | * | * | * | * | * |
| 2000..... | 0.0 | ... | ... | * | * | 0.2 | * | * | * | * | * | * |
| 1999..... | 0.0 | ... | ... | * | * | 0.2 | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, both sexes.

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, male.

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, female.

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, both sexes.

Non-Hispanic white, male

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, male.

Non-Hispanic white, female

Table with 13 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | 137.9 | 3.6 | 1.0 | 1.2 | 1.0 | 1.1 | 1.7 | 2.2 | 2.3 | 3.6 | 6.1 |
| 2003..... | 3.6 | 140.4 | 3.4 | 0.9 | 1.1 | 1.1 | 1.3 | 1.6 | 2.4 | 2.3 | 3.6 | 6.8 |
| 2002..... | 3.7 | 139.4 | 3.4 | 1.0 | 1.2 | 1.2 | 1.3 | 1.8 | 2.4 | 2.4 | 3.8 | 6.6 |
| 2001..... | 3.7 | 136.7 | 3.6 | 0.9 | 1.3 | 1.2 | 1.3 | 1.6 | 2.3 | 2.4 | 3.7 | 7.2 |
| 2000..... | 3.8 | 150.9 | 3.2 | 1.0 | 1.1 | 1.2 | 1.3 | 1.6 | 2.3 | 2.5 | 3.9 | 7.5 |
| 1999..... | 3.7 | 144.2 | 3.6 | 1.0 | 1.1 | 1.2 | 1.2 | 1.8 | 2.3 | 2.5 | 4.0 | 7.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.8 | 139.0 | 3.9 | 1.0 | 1.3 | 1.2 | 1.3 | 1.8 | 2.3 | 2.5 | 3.7 | 6.8 |
| 2003..... | 3.8 | 147.6 | 3.4 | 0.9 | 1.3 | 1.2 | 1.4 | 1.7 | 2.5 | 2.4 | 3.5 | 6.7 |
| 2002..... | 3.9 | 142.6 | 3.5 | 1.0 | 1.4 | 1.4 | 1.3 | 1.9 | 2.5 | 2.6 | 4.0 | 7.0 |
| 2001..... | 3.9 | 143.1 | 3.6 | 1.0 | 1.5 | 1.2 | 1.4 | 1.7 | 2.3 | 2.7 | 4.1 | 7.7 |
| 2000..... | 4.0 | 155.4 | 3.3 | 1.0 | 1.3 | 1.4 | 1.4 | 1.9 | 2.3 | 2.7 | 4.0 | 10.6 |
| 1999..... | 3.9 | 147.9 | 4.0 | 1.0 | 1.3 | 1.2 | 1.3 | 1.9 | 2.5 | 2.5 | 4.2 | 8.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.4 | 136.8 | 3.2 | 0.9 | 1.0 | 0.9 | 1.0 | 1.5 | 2.1 | 2.2 | 3.6 | 5.8 |
| 2003..... | 3.4 | 132.8 | 3.5 | 0.9 | 0.9 | 1.0 | 1.2 | 1.5 | 2.3 | 2.2 | 3.7 | 6.8 |
| 2002..... | 3.5 | 136.1 | 3.3 | 1.0 | 1.1 | 1.0 | 1.2 | 1.6 | 2.3 | 2.3 | 3.8 | 6.5 |
| 2001..... | 3.4 | 129.9 | 3.7 | 0.8 | 1.0 | 1.1 | 1.2 | 1.6 | 2.3 | 2.2 | 3.5 | 7.0 |
| 2000..... | 3.5 | 146.2 | 3.1 | 1.0 | 0.9 | 1.0 | 1.1 | 1.4 | 2.4 | 2.4 | 3.9 | 6.3 |
| 1999..... | 3.5 | 140.3 | 3.2 | 1.1 | 0.9 | 1.1 | 1.1 | 1.6 | 2.2 | 2.6 | 3.9 | 7.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | 141.0 | 4.0 | 0.9 | 0.9 | 0.7 | 0.6 | 1.1 | 1.4 | 1.6 | 3.4 | * |
| 2003..... | 4.5 | 153.0 | 3.7 | 0.9 | 0.8 | 0.8 | 0.7 | 1.0 | 1.7 | * | * | * |
| 2002..... | 4.4 | 150.7 | 3.5 | 1.0 | 1.0 | 0.7 | 0.7 | 1.1 | 1.3 | * | 3.8 | * |
| 2001..... | 4.4 | 146.2 | 3.4 | 0.8 | 1.1 | 0.7 | 0.8 | 1.1 | 1.3 | 1.9 | 4.3 | * |
| 2000..... | 4.6 | 153.3 | 3.8 | 0.8 | 0.8 | 0.7 | 0.8 | 1.1 | 1.8 | 2.0 | 4.7 | * |
| 1999..... | 4.4 | 141.8 | 3.5 | 0.9 | 0.9 | 0.9 | 0.7 | 1.4 | 1.6 | 2.9 | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.2 | 134.3 | 4.6 | 1.3 | 1.1 | 0.6 | 0.7 | 1.3 | * | * | * | * |
| 2003..... | 4.5 | 156.9 | 3.7 | 0.7 | 0.8 | 0.9 | 0.8 | 1.2 | * | * | * | * |
| 2002..... | 4.5 | 153.1 | 3.8 | 0.9 | 1.1 | 0.9 | * | 1.3 | * | * | * | * |
| 2001..... | 4.7 | 153.4 | 3.5 | 1.0 | 1.1 | 0.7 | 1.0 | 1.5 | * | * | * | * |
| 2000..... | 4.7 | 153.3 | 4.0 | 0.8 | 1.1 | 0.7 | 1.0 | 1.4 | * | * | * | * |
| 1999..... | 4.3 | 138.1 | 4.3 | 0.9 | 0.9 | 0.7 | * | 1.6 | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 4.3 | 147.9 | 3.3 | 0.6 | 0.6 | 0.8 | * | * | * | * | * | * |
| 2003..... | 4.4 | 148.9 | 3.7 | 1.0 | 0.8 | 0.6 | * | * | * | * | * | * |
| 2002..... | 4.3 | 148.2 | 3.3 | 1.0 | 0.9 | * | 0.8 | * | * | * | * | * |
| 2001..... | 4.1 | 138.8 | 3.3 | 0.6 | 1.1 | * | * | * | * | * | * | * |
| 2000..... | 4.4 | 153.3 | 3.5 | 0.7 | * | 0.7 | * | * | * | * | * | * |
| 1999..... | 4.5 | 145.8 | 2.7 | 0.9 | 0.9 | 1.0 | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | 135.9 | 3.4 | 1.0 | 1.2 | 1.1 | 1.2 | 1.7 | 2.2 | 2.4 | 3.7 | 6.2 |
| 2003..... | 3.5 | 136.0 | 3.4 | 0.9 | 1.1 | 1.1 | 1.3 | 1.7 | 2.4 | 2.3 | 3.7 | 6.7 |
| 2002..... | 3.6 | 134.9 | 3.3 | 1.0 | 1.2 | 1.3 | 1.4 | 1.8 | 2.5 | 2.5 | 3.8 | 6.7 |
| 2001..... | 3.5 | 132.8 | 3.7 | 0.9 | 1.3 | 1.3 | 1.4 | 1.7 | 2.4 | 2.5 | 3.7 | 7.2 |
| 2000..... | 3.6 | 148.8 | 3.1 | 1.0 | 1.2 | 1.3 | 1.3 | 1.7 | 2.4 | 2.6 | 3.9 | 7.5 |
| 1999..... | 3.6 | 143.1 | 3.6 | 1.1 | 1.2 | 1.2 | 1.3 | 1.8 | 2.4 | 2.5 | 4.0 | 7.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.6 | 139.1 | 3.7 | 0.9 | 1.4 | 1.3 | 1.4 | 1.9 | 2.3 | 2.5 | 3.7 | 7.0 |
| 2003..... | 3.7 | 144.2 | 3.3 | 1.0 | 1.4 | 1.2 | 1.5 | 1.8 | 2.5 | 2.4 | 3.5 | 6.7 |
| 2002..... | 3.8 | 138.1 | 3.5 | 1.0 | 1.4 | 1.5 | 1.4 | 2.0 | 2.6 | 2.6 | 3.9 | 7.0 |
| 2001..... | 3.7 | 139.4 | 3.6 | 1.0 | 1.6 | 1.4 | 1.5 | 1.7 | 2.3 | 2.7 | 4.0 | 7.6 |
| 2000..... | 3.9 | 154.4 | 3.1 | 1.0 | 1.4 | 1.5 | 1.5 | 1.9 | 2.2 | 2.7 | 3.9 | 10.4 |
| 1999..... | 3.8 | 148.4 | 3.9 | 1.1 | 1.4 | 1.3 | 1.4 | 1.9 | 2.6 | 2.4 | 4.2 | 8.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.2 | 132.6 | 3.2 | 1.0 | 1.1 | 1.0 | 1.0 | 1.6 | 2.1 | 2.2 | 3.6 | 5.9 |
| 2003..... | 3.3 | 127.6 | 3.4 | 0.9 | 0.9 | 1.0 | 1.2 | 1.6 | 2.3 | 2.2 | 3.8 | 6.8 |
| 2002..... | 3.4 | 131.5 | 3.2 | 1.0 | 1.1 | 1.1 | 1.3 | 1.7 | 2.4 | 2.3 | 3.8 | 6.6 |
| 2001..... | 3.3 | 125.8 | 3.7 | 0.9 | 1.0 | 1.2 | 1.3 | 1.7 | 2.4 | 2.2 | 3.5 | 7.0 |
| 2000..... | 3.4 | 142.8 | 3.0 | 1.0 | 1.0 | 1.1 | 1.2 | 1.4 | 2.5 | 2.4 | 3.9 | 6.4 |
| 1999..... | 3.4 | 137.4 | 3.3 | 1.1 | 0.9 | 1.1 | 1.2 | 1.6 | 2.2 | 2.6 | 3.9 | 7.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.3 | 130.5 | 2.9 | 0.9 | 1.1 | 1.2 | 1.2 | 1.8 | 2.4 | 2.5 | 3.7 | 6.5 |
| 2003..... | 3.3 | 129.8 | 3.0 | 0.9 | 1.1 | 1.2 | 1.4 | 1.7 | 2.6 | 2.3 | 3.8 | 6.9 |
| 2002..... | 3.4 | 131.2 | 3.0 | 1.0 | 1.2 | 1.3 | 1.4 | 1.9 | 2.5 | 2.5 | 3.9 | 6.9 |
| 2001..... | 3.4 | 127.4 | 3.1 | 0.9 | 1.2 | 1.3 | 1.4 | 1.7 | 2.5 | 2.5 | 3.8 | 7.4 |
| 2000..... | 3.4 | 141.2 | 2.6 | 0.9 | 1.1 | 1.3 | 1.3 | 1.7 | 2.4 | 2.6 | 4.0 | 7.8 |
| 1999..... | 3.4 | 136.8 | 3.0 | 0.9 | 1.1 | 1.3 | 1.3 | 1.8 | 2.5 | 2.5 | 4.0 | 8.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.4 | 132.8 | 3.1 | 0.9 | 1.2 | 1.4 | 1.4 | 2.0 | 2.4 | 2.7 | 3.7 | 7.4 |
| 2003..... | 3.4 | 136.1 | 2.9 | 0.9 | 1.3 | 1.2 | 1.5 | 1.8 | 2.6 | 2.3 | 3.4 | 6.8 |
| 2002..... | 3.6 | 137.2 | 3.2 | 1.0 | 1.4 | 1.6 | 1.4 | 2.0 | 2.6 | 2.7 | 3.8 | 7.3 |
| 2001..... | 3.6 | 135.5 | 3.1 | 1.0 | 1.4 | 1.5 | 1.5 | 1.7 | 2.4 | 2.9 | 4.1 | 7.4 |
| 2000..... | 3.6 | 147.9 | 2.6 | 0.9 | 1.3 | 1.6 | 1.4 | 1.9 | 2.3 | 2.6 | 4.1 | 10.8 |
| 1999..... | 3.6 | 144.4 | 3.2 | 0.9 | 1.4 | 1.3 | 1.4 | 2.0 | 2.7 | 2.4 | 4.1 | 8.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.2 | 128.0 | 2.8 | 0.9 | 1.0 | 1.1 | 1.1 | 1.6 | 2.3 | 2.4 | 3.7 | 6.1 |
| 2003..... | 3.2 | 123.2 | 3.0 | 0.9 | 0.9 | 1.1 | 1.3 | 1.7 | 2.5 | 2.3 | 4.0 | 6.9 |
| 2002..... | 3.3 | 125.0 | 2.9 | 1.0 | 1.1 | 1.1 | 1.3 | 1.8 | 2.5 | 2.4 | 3.9 | 6.7 |
| 2001..... | 3.2 | 118.8 | 3.2 | 0.7 | 1.0 | 1.2 | 1.3 | 1.7 | 2.6 | 2.3 | 3.6 | 7.4 |
| 2000..... | 3.2 | 134.2 | 2.6 | 1.0 | 1.0 | 1.1 | 1.2 | 1.6 | 2.5 | 2.6 | 4.0 | 6.6 |
| 1999..... | 3.3 | 128.9 | 2.9 | 1.0 | 0.8 | 1.2 | 1.2 | 1.7 | 2.3 | 2.6 | 4.0 | 7.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 4.6 | 164.3 | 5.1 | 1.3 | 1.8 | 1.1 | 1.3 | 1.7 | 1.7 | 1.6 | 4.0 | * |
| 2003..... | 4.6 | 160.5 | 4.8 | 1.1 | 1.5 | 1.4 | 1.2 | 1.7 | 2.2 | 2.4 | 3.1 | 5.9 |
| 2002..... | 4.7 | 156.9 | 4.7 | 1.3 | 1.7 | 1.5 | 1.6 | 1.8 | 2.7 | 2.4 | 3.4 | * |
| 2001..... | 4.7 | 163.5 | 5.4 | 1.1 | 2.0 | 1.3 | 1.6 | 1.8 | 1.9 | 2.1 | 3.0 | * |
| 2000..... | 5.0 | 184.1 | 5.2 | 1.4 | 1.6 | 1.4 | 1.6 | 1.6 | 2.4 | 2.7 | 2.8 | * |
| 1999..... | 4.8 | 174.8 | 5.8 | 1.6 | 1.5 | 1.1 | 1.4 | 1.9 | 2.0 | 2.9 | 4.3 | 7.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 5.2 | 170.5 | 5.9 | 1.2 | 2.4 | 1.4 | 1.6 | 1.8 | 2.0 | * | * | * |
| 2003..... | 5.2 | 173.1 | 5.0 | 1.2 | 1.8 | 1.5 | 1.2 | 2.0 | 2.8 | 3.6 | * | * |
| 2002..... | 5.2 | 153.3 | 4.8 | 1.4 | 2.0 | 1.8 | 1.7 | 2.3 | 3.5 | 3.1 | * | * |
| 2001..... | 5.1 | 167.5 | 5.5 | 1.0 | 2.5 | 1.2 | 1.8 | 1.7 | 2.1 | * | * | * |
| 2000..... | 5.5 | 186.0 | 5.6 | 1.5 | 2.1 | 1.7 | 1.9 | 2.2 | * | 3.9 | * | * |
| 1999..... | 5.3 | 179.5 | 6.6 | 1.6 | 1.8 | 1.3 | 1.6 | 2.3 | 2.1 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.1 | 157.8 | 4.3 | 1.4 | 1.2 | 0.8 | 0.9 | 1.7 | 1.4 | * | 3.4 | * |
| 2003..... | 4.0 | 147.3 | 4.7 | 1.1 | 1.2 | 1.4 | 1.3 | 1.5 | 1.8 | * | * | * |
| 2002..... | 4.4 | 160.6 | 4.7 | 1.1 | 1.3 | 1.2 | 1.6 | 1.5 | 2.0 | * | * | * |
| 2001..... | 4.4 | 159.2 | 5.4 | 1.3 | 1.4 | 1.4 | 1.5 | 1.8 | 1.8 | 2.4 | * | * |
| 2000..... | 4.5 | 182.2 | 4.8 | 1.2 | 1.2 | 1.1 | 1.4 | 1.2 | 3.1 | * | * | * |
| 1999..... | 4.4 | 170.0 | 5.0 | 1.6 | 1.1 | 1.0 | 1.2 | 1.5 | 1.9 | 3.2 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Congenital malformations of heart (Q20-Q24) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | 35.0 | 1.1 | 0.4 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 1.0 | 1.6 |
| 2003..... | 1.1 | 35.8 | 1.4 | 0.4 | 0.5 | 0.5 | 0.6 | 0.5 | 0.5 | 0.7 | 1.1 | 2.0 |
| 2002..... | 1.1 | 36.1 | 1.2 | 0.4 | 0.5 | 0.6 | 0.6 | 0.5 | 0.5 | 0.6 | 1.2 | 2.3 |
| 2001..... | 1.1 | 37.6 | 1.3 | 0.3 | 0.7 | 0.6 | 0.6 | 0.6 | 0.5 | 0.7 | 1.2 | 2.6 |
| 2000..... | 1.2 | 43.0 | 1.2 | 0.4 | 0.6 | 0.6 | 0.6 | 0.5 | 0.6 | 0.7 | 1.4 | 2.4 |
| 1999..... | 1.2 | 41.7 | 1.3 | 0.4 | 0.6 | 0.6 | 0.6 | 0.6 | 0.7 | 0.9 | 1.4 | 3.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | 38.6 | 1.3 | 0.3 | 0.7 | 0.7 | 0.5 | 0.5 | 0.5 | 0.5 | 0.9 | 1.5 |
| 2003..... | 1.1 | 39.5 | 1.4 | 0.4 | 0.6 | 0.6 | 0.7 | 0.5 | 0.5 | 0.7 | 0.9 | * |
| 2002..... | 1.2 | 39.0 | 1.5 | 0.4 | 0.7 | 0.8 | 0.6 | 0.5 | 0.5 | 0.6 | 1.1 | * |
| 2001..... | 1.3 | 41.1 | 1.4 | 0.3 | 0.9 | 0.7 | 0.6 | 0.6 | 0.5 | 0.8 | 1.2 | 1.6 |
| 2000..... | 1.3 | 45.5 | 1.2 | 0.4 | 0.7 | 0.8 | 0.6 | 0.6 | 0.6 | 0.8 | 1.4 | 2.5 |
| 1999..... | 1.4 | 46.2 | 1.4 | 0.4 | 0.8 | 0.7 | 0.7 | 0.7 | 0.7 | 0.8 | 1.3 | 3.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | 31.3 | 0.9 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 1.1 | 1.6 |
| 2003..... | 1.0 | 31.9 | 1.3 | 0.3 | 0.3 | 0.5 | 0.6 | 0.5 | 0.5 | 0.7 | 1.3 | 2.3 |
| 2002..... | 1.0 | 33.0 | 0.9 | 0.3 | 0.4 | 0.5 | 0.5 | 0.6 | 0.5 | 0.7 | 1.4 | 2.7 |
| 2001..... | 1.0 | 34.0 | 1.2 | 0.3 | 0.4 | 0.5 | 0.5 | 0.6 | 0.5 | 0.6 | 1.3 | 3.0 |
| 2000..... | 1.1 | 40.4 | 1.1 | 0.3 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.7 | 1.4 | 2.4 |
| 1999..... | 1.1 | 37.0 | 1.2 | 0.3 | 0.4 | 0.6 | 0.6 | 0.5 | 0.7 | 0.9 | 1.5 | 3.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | 34.0 | 0.9 | 0.3 | 0.5 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | 1.1 | 1.6 |
| 2003..... | 1.0 | 33.9 | 1.2 | 0.3 | 0.4 | 0.6 | 0.7 | 0.5 | 0.5 | 0.7 | 1.2 | 2.2 |
| 2002..... | 1.0 | 33.9 | 1.2 | 0.3 | 0.5 | 0.7 | 0.6 | 0.6 | 0.5 | 0.7 | 1.2 | 2.4 |
| 2001..... | 1.1 | 35.1 | 1.1 | 0.3 | 0.6 | 0.6 | 0.6 | 0.6 | 0.5 | 0.7 | 1.3 | 2.7 |
| 2000..... | 1.1 | 40.6 | 1.0 | 0.3 | 0.5 | 0.7 | 0.6 | 0.6 | 0.6 | 0.7 | 1.5 | 2.6 |
| 1999..... | 1.2 | 40.7 | 1.0 | 0.3 | 0.6 | 0.7 | 0.6 | 0.6 | 0.7 | 0.9 | 1.5 | 3.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.1 | 37.9 | 1.1 | 0.3 | 0.6 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.9 | 1.6 |
| 2003..... | 1.1 | 37.8 | 1.2 | 0.4 | 0.5 | 0.6 | 0.7 | 0.5 | 0.5 | 0.7 | 0.9 | * |
| 2002..... | 1.1 | 37.7 | 1.6 | 0.4 | 0.7 | 0.8 | 0.6 | 0.5 | 0.5 | 0.6 | 1.0 | * |
| 2001..... | 1.2 | 39.5 | 1.3 | 0.3 | 0.8 | 0.8 | 0.6 | 0.6 | 0.5 | 0.9 | 1.3 | * |
| 2000..... | 1.2 | 44.1 | 1.0 | 0.3 | 0.6 | 0.8 | 0.6 | 0.6 | 0.6 | 0.7 | 1.5 | 2.7 |
| 1999..... | 1.3 | 46.2 | 1.1 | 0.4 | 0.8 | 0.8 | 0.7 | 0.6 | 0.7 | 0.8 | 1.2 | 3.0 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | 29.9 | 0.8 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | 0.5 | 1.2 | 1.6 |
| 2003..... | 0.9 | 29.9 | 1.2 | 0.3 | 0.2 | 0.5 | 0.6 | 0.5 | 0.5 | 0.7 | 1.4 | 2.5 |
| 2002..... | 1.0 | 29.8 | 0.9 | 0.3 | 0.4 | 0.5 | 0.5 | 0.6 | 0.5 | 0.7 | 1.4 | 2.8 |
| 2001..... | 1.0 | 30.5 | 1.0 | 0.2 | 0.4 | 0.5 | 0.6 | 0.6 | 0.4 | 0.6 | 1.3 | 3.2 |
| 2000..... | 1.0 | 36.8 | 1.0 | 0.3 | 0.4 | 0.5 | 0.5 | 0.5 | 0.6 | 0.7 | 1.5 | 2.5 |
| 1999..... | 1.1 | 35.0 | 1.0 | 0.3 | 0.4 | 0.6 | 0.6 | 0.6 | 0.7 | 0.9 | 1.6 | 3.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Congenital malformations of heart (Q20-Q24) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | 42.9 | 1.7 | 0.5 | 0.9 | 0.6 | 0.5 | * | * | * | * | * |
| 2003..... | 1.4 | 42.7 | 2.2 | 0.4 | 0.8 | 0.7 | 0.5 | 0.6 | * | * | * | * |
| 2002..... | 1.5 | 45.3 | 1.4 | 0.5 | 0.7 | 0.7 | 0.6 | * | * | * | * | * |
| 2001..... | 1.6 | 49.7 | 2.0 | 0.4 | 1.2 | 0.6 | 0.7 | 0.6 | * | * | * | * |
| 2000..... | 1.6 | 52.9 | 1.9 | 0.5 | 0.9 | 0.7 | 0.7 | 0.5 | * | * | * | * |
| 1999..... | 1.6 | 50.1 | 2.3 | 0.6 | 0.8 | 0.5 | 0.7 | 0.5 | * | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.6 | 45.8 | 1.9 | * | 1.1 | 0.8 | * | * | * | * | * | * |
| 2003..... | 1.6 | 45.9 | 2.3 | * | 1.0 | 0.8 | * | * | * | * | * | * |
| 2002..... | 1.6 | 43.6 | * | 0.7 | 1.0 | 1.0 | * | * | * | * | * | * |
| 2001..... | 1.8 | 50.7 | 1.9 | * | 1.7 | * | * | * | * | * | * | * |
| 2000..... | 1.8 | 50.8 | 2.2 | * | 1.3 | 0.9 | 0.8 | * | * | * | * | * |
| 1999..... | 1.9 | 52.7 | 2.6 | * | 1.1 | * | 0.8 | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | 39.9 | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 1.3 | 39.3 | 2.0 | * | * | * | 0.7 | * | * | * | * | * |
| 2002..... | 1.3 | 47.0 | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 1.5 | 48.7 | 2.1 | * | 0.7 | * | * | * | * | * | * | * |
| 2000..... | 1.4 | 55.0 | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 1.4 | 47.5 | 2.1 | 0.7 | * | * | 0.7 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other congenital malformations of circulatory system (Q25-Q28) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 6.3 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.9 | 1.7 |
| 2003..... | 0.3 | 5.9 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.5 |
| 2002..... | 0.3 | 5.5 | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 1.0 | 1.6 |
| 2001..... | 0.3 | 4.5 | * | 0.1 | 0.1 | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.7 |
| 2000..... | 0.3 | 5.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.4 | 0.9 | 2.1 |
| 1999..... | 0.3 | 6.2 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.5 | 0.9 | 1.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | 7.2 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 1.1 | 1.9 |
| 2003..... | 0.3 | 6.3 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.5 | 1.0 | 1.7 |
| 2002..... | 0.3 | 6.2 | * | * | * | 0.2 | 0.1 | 0.3 | 0.4 | 0.6 | 1.0 | 1.8 |
| 2001..... | 0.3 | 4.6 | * | 0.1 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 | 2.1 |
| 2000..... | 0.3 | 5.1 | * | * | * | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 1.0 | 3.1 |
| 1999..... | 0.3 | 6.4 | * | * | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 1.1 | 2.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 5.4 | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.8 | 1.5 |
| 2003..... | 0.3 | 5.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.7 | 1.4 |
| 2002..... | 0.3 | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 1.0 | 1.5 |
| 2001..... | 0.3 | 4.5 | * | * | * | * | * | 0.2 | 0.2 | 0.6 | 0.5 | 1.6 |
| 2000..... | 0.3 | 5.3 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.8 | 1.7 |
| 1999..... | 0.3 | 5.9 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 0.8 | 1.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 5.5 | * | 0.1 | * | * | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 1.8 |
| 2003..... | 0.3 | 5.8 | * | 0.1 | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.8 | 1.5 |
| 2002..... | 0.3 | 4.8 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 1.0 | 1.7 |
| 2001..... | 0.3 | 4.2 | * | 0.1 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.7 |
| 2000..... | 0.3 | 4.6 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 | 0.4 | 0.9 | 2.2 |
| 1999..... | 0.3 | 5.4 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.5 | 0.9 | 1.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.3 | 6.3 | * | * | * | * | 0.1 | 0.3 | 0.2 | 0.4 | 1.0 | 2.1 |
| 2003..... | 0.3 | 6.6 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.4 | 1.0 | 1.7 |
| 2002..... | 0.3 | 4.9 | * | * | * | 0.2 | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.9 |
| 2001..... | 0.3 | 4.4 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.0 | 2.0 |
| 2000..... | 0.3 | 4.6 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.5 | 1.1 | 3.3 |
| 1999..... | 0.3 | 5.9 | * | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 1.0 | 2.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 4.6 | * | * | * | * | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.7 |
| 2003..... | 0.3 | 5.0 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.7 | 1.4 |
| 2002..... | 0.3 | 4.7 | * | * | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 1.5 |
| 2001..... | 0.3 | 4.0 | * | * | * | * | * | 0.1 | 0.3 | 0.5 | 0.6 | 1.6 |
| 2000..... | 0.3 | 4.7 | * | * | * | * | 0.2 | 0.1 | 0.3 | 0.4 | 0.8 | 1.7 |
| 1999..... | 0.3 | 4.9 | * | * | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other congenital malformations and deformations (Q10-Q18,Q35-Q89) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | 35.4 | 0.8 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.4 | 0.9 | 1.4 | 2.7 |
| 2003..... | 0.9 | 36.7 | 0.7 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 1.5 | 2.8 |
| 2002..... | 0.9 | 35.3 | 0.7 | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 0.5 | 0.8 | 1.4 | 2.7 |
| 2001..... | 0.9 | 36.3 | 0.8 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.5 | 2.5 |
| 2000..... | 0.9 | 36.9 | 0.7 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.4 | 2.8 |
| 1999..... | 0.9 | 37.6 | 1.0 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.5 | 2.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | 38.0 | 0.9 | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 0.5 | 0.9 | 1.4 | 3.3 |
| 2003..... | 1.0 | 41.8 | 0.7 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.8 | 1.4 | 3.2 |
| 2002..... | 1.0 | 39.1 | 0.6 | 0.2 | 0.3 | 0.2 | 0.3 | 0.5 | 0.6 | 0.9 | 1.6 | 3.7 |
| 2001..... | 1.0 | 41.2 | 0.7 | 0.2 | 0.3 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.6 | 3.7 |
| 2000..... | 1.0 | 41.2 | 0.7 | 0.2 | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 | 0.8 | 1.3 | 4.4 |
| 1999..... | 1.0 | 41.7 | 1.1 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | 0.5 | 0.8 | 1.6 | 3.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | 32.6 | 0.6 | 0.1 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | 0.8 | 1.5 | 2.4 |
| 2003..... | 0.8 | 31.4 | 0.6 | 0.1 | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 0.6 | 1.6 | 2.6 |
| 2002..... | 0.8 | 31.4 | 0.8 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 1.2 | 2.2 |
| 2001..... | 0.8 | 31.0 | 0.8 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 0.7 | 1.5 | 2.0 |
| 2000..... | 0.8 | 32.4 | 0.6 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.6 | 0.9 | 1.4 | 2.1 |
| 1999..... | 0.8 | 33.2 | 0.8 | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.4 | 0.7 | 1.4 | 2.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | 34.0 | 0.7 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.9 | 1.4 | 2.8 |
| 2003..... | 0.8 | 34.5 | 0.6 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 0.7 | 1.6 | 2.8 |
| 2002..... | 0.9 | 35.8 | 0.6 | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 0.6 | 0.8 | 1.4 | 2.7 |
| 2001..... | 0.9 | 35.1 | 0.7 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.5 | 2.6 |
| 2000..... | 0.9 | 35.3 | 0.6 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.6 | 0.9 | 1.4 | 2.8 |
| 1999..... | 0.9 | 35.1 | 0.9 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | 0.5 | 0.8 | 1.5 | 2.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.9 | 36.2 | 0.7 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 1.0 | 1.3 | 3.4 |
| 2003..... | 0.9 | 38.6 | 0.6 | * | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.8 | 1.3 | 3.3 |
| 2002..... | 1.0 | 40.4 | 0.5 | 0.2 | 0.3 | 0.3 | 0.3 | 0.5 | 0.6 | 0.9 | 1.5 | 3.7 |
| 2001..... | 0.9 | 39.8 | 0.6 | 0.2 | 0.3 | 0.2 | 0.3 | 0.3 | 0.5 | 0.9 | 1.5 | 3.8 |
| 2000..... | 0.9 | 39.2 | 0.6 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.8 | 1.4 | 4.4 |
| 1999..... | 1.0 | 40.3 | 1.0 | 0.2 | 0.2 | 0.2 | 0.3 | 0.4 | 0.5 | 0.9 | 1.6 | 3.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | 31.6 | 0.6 | * | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 0.9 | 1.5 | 2.5 |
| 2003..... | 0.8 | 30.2 | 0.6 | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.5 | 0.6 | 1.7 | 2.6 |
| 2002..... | 0.8 | 31.0 | 0.7 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 1.3 | 2.3 |
| 2001..... | 0.8 | 30.2 | 0.7 | * | * | 0.2 | 0.3 | 0.3 | 0.5 | 0.7 | 1.5 | 2.1 |
| 2000..... | 0.8 | 31.1 | 0.6 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.6 | 0.9 | 1.4 | 2.1 |
| 1999..... | 0.8 | 29.5 | 0.8 | 0.2 | * | 0.2 | 0.2 | 0.3 | 0.4 | 0.8 | 1.4 | 2.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Down's syndrome (Q90) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.5 | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 0.9 | 0.5 | * | * |
| 2003..... | 0.3 | 2.6 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 0.3 | * | * |
| 2002..... | 0.3 | 2.8 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 0.4 | * | * |
| 2001..... | 0.3 | 3.1 | * | * | 0.1 | 0.2 | 0.2 | 0.5 | 1.0 | 0.3 | * | * |
| 2000..... | 0.3 | 2.3 | * | * | * | 0.1 | 0.1 | 0.5 | 0.8 | 0.4 | * | * |
| 1999..... | 0.2 | 2.5 | * | * | * | 0.1 | 0.1 | 0.4 | 0.8 | 0.2 | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.6 | * | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 0.5 | * | * |
| 2003..... | 0.3 | 2.8 | * | * | * | * | 0.2 | 0.5 | 1.1 | 0.4 | * | * |
| 2002..... | 0.3 | 2.7 | * | * | * | * | 0.2 | 0.6 | 1.0 | 0.4 | * | * |
| 2001..... | 0.3 | 3.5 | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 0.4 | * | * |
| 2000..... | 0.3 | 2.6 | * | * | * | * | 0.2 | 0.5 | 0.8 | 0.3 | * | * |
| 1999..... | 0.2 | 2.7 | * | * | * | * | 0.1 | 0.4 | 0.9 | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.4 | * | * | * | * | 0.2 | 0.6 | 0.9 | 0.4 | * | * |
| 2003..... | 0.3 | 2.5 | * | * | * | * | 0.2 | 0.6 | 1.0 | 0.3 | * | * |
| 2002..... | 0.3 | 2.8 | * | * | * | * | 0.2 | 0.5 | 0.9 | 0.4 | * | * |
| 2001..... | 0.3 | 2.7 | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 0.3 | * | * |
| 2000..... | 0.3 | 2.0 | * | * | * | 0.1 | 0.1 | 0.5 | 0.8 | 0.4 | * | * |
| 1999..... | 0.2 | 2.4 | * | * | * | * | 0.1 | 0.4 | 0.7 | 0.3 | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.2 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 0.5 | * | * |
| 2003..... | 0.3 | 2.6 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 0.4 | * | * |
| 2002..... | 0.3 | 3.0 | * | * | * | 0.1 | 0.2 | 0.6 | 1.1 | 0.4 | * | * |
| 2001..... | 0.3 | 3.2 | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 0.3 | * | * |
| 2000..... | 0.3 | 2.2 | * | * | * | 0.1 | 0.2 | 0.5 | 0.9 | 0.4 | * | * |
| 1999..... | 0.2 | 2.7 | * | * | * | 0.1 | 0.1 | 0.5 | 0.9 | 0.2 | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | 2.2 | * | * | * | * | 0.2 | 0.7 | 1.1 | 0.6 | * | * |
| 2003..... | 0.3 | 2.6 | * | * | * | * | 0.2 | 0.6 | 1.2 | 0.4 | * | * |
| 2002..... | 0.3 | 2.8 | * | * | * | * | 0.2 | 0.6 | 1.1 | 0.4 | * | * |
| 2001..... | 0.3 | 3.7 | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 0.4 | * | * |
| 2000..... | 0.3 | 2.8 | * | * | * | * | 0.2 | 0.5 | 0.8 | 0.3 | * | * |
| 1999..... | 0.3 | 2.9 | * | * | * | * | 0.1 | 0.5 | 1.0 | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.2 | * | * | * | * | 0.2 | 0.6 | 1.0 | 0.5 | * | * |
| 2003..... | 0.3 | 2.5 | * | * | * | * | 0.2 | 0.6 | 1.1 | 0.4 | * | * |
| 2002..... | 0.3 | 3.2 | * | * | * | * | 0.3 | 0.5 | 1.1 | 0.4 | * | * |
| 2001..... | 0.3 | 2.7 | * | * | * | * | 0.2 | 0.6 | 1.2 | 0.3 | * | * |
| 2000..... | 0.3 | * | * | * | * | 0.2 | 0.1 | 0.6 | 0.9 | 0.5 | * | * |
| 1999..... | 0.2 | 2.5 | * | * | * | * | 0.1 | 0.5 | 0.8 | 0.2 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 10.3 | 82.4 | 1.2 | 0.3 | 1.6 | 2.7 | 4.3 | 5.8 | 7.0 | 12.6 | 39.8 | 221.5 |
| 2003..... | 10.8 | 82.9 | 1.6 | 0.4 | 2.0 | 3.3 | 5.4 | 7.0 | 7.4 | 13.8 | 38.9 | 224.1 |
| 2002..... | 10.4 | 85.7 | 1.6 | 0.3 | 1.8 | 3.3 | 5.3 | 6.7 | 7.5 | 13.1 | 38.8 | 208.0 |
| 2001..... | 11.4 | 86.5 | 1.8 | 0.4 | 2.4 | 3.9 | 6.9 | 8.9 | 9.1 | 15.1 | 39.8 | 206.4 |
| 2000..... | 11.3 | 95.6 | 1.6 | 0.4 | 2.2 | 3.9 | 6.6 | 8.3 | 9.3 | 16.5 | 40.7 | 208.1 |
| 1999..... | 9.6 | 93.2 | 1.5 | 0.3 | 1.6 | 2.7 | 4.6 | 5.9 | 7.8 | 14.6 | 36.1 | 187.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 9.5 | 96.9 | 1.2 | 0.3 | 2.1 | 3.6 | 5.6 | 7.6 | 9.1 | 15.0 | 41.8 | 183.6 |
| 2003..... | 10.2 | 96.5 | 1.8 | 0.4 | 2.7 | 4.5 | 6.9 | 9.2 | 9.6 | 17.1 | 41.9 | 177.9 |
| 2002..... | 10.1 | 98.5 | 1.8 | 0.4 | 2.4 | 4.5 | 6.9 | 8.8 | 10.0 | 15.9 | 43.8 | 175.5 |
| 2001..... | 11.3 | 98.9 | 2.1 | 0.5 | 3.2 | 5.2 | 9.0 | 11.9 | 12.0 | 18.0 | 43.8 | 175.0 |
| 2000..... | 11.5 | 110.6 | 1.7 | 0.5 | 2.9 | 5.3 | 8.9 | 11.5 | 12.3 | 20.6 | 46.0 | 182.7 |
| 1999..... | 9.4 | 106.6 | 1.7 | 0.4 | 2.1 | 3.6 | 6.0 | 8.1 | 10.4 | 18.2 | 40.1 | 166.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 11.0 | 67.3 | 1.1 | 0.3 | 1.0 | 1.8 | 3.0 | 4.0 | 5.1 | 10.6 | 38.4 | 238.5 |
| 2003..... | 11.4 | 68.6 | 1.5 | 0.4 | 1.4 | 2.1 | 3.8 | 5.0 | 5.3 | 10.9 | 36.9 | 244.6 |
| 2002..... | 10.7 | 72.3 | 1.4 | 0.3 | 1.2 | 2.1 | 3.8 | 4.7 | 5.3 | 10.8 | 35.4 | 222.1 |
| 2001..... | 11.4 | 73.6 | 1.5 | 0.4 | 1.5 | 2.5 | 4.9 | 6.1 | 6.3 | 12.7 | 37.2 | 219.5 |
| 2000..... | 11.2 | 79.9 | 1.5 | 0.4 | 1.5 | 2.5 | 4.4 | 5.3 | 6.5 | 13.2 | 37.2 | 218.5 |
| 1999..... | 9.7 | 79.2 | 1.3 | 0.3 | 1.0 | 1.9 | 3.2 | 3.8 | 5.5 | 11.7 | 33.5 | 196.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.9 | 50.4 | 1.1 | * | 1.6 | 2.1 | 3.1 | 3.6 | 3.9 | 6.2 | 16.4 | 99.4 |
| 2003..... | 4.1 | 56.8 | 0.9 | 0.3 | 1.4 | 2.3 | 3.1 | 3.7 | 3.5 | 6.1 | 18.4 | 110.9 |
| 2002..... | 4.0 | 56.6 | 1.3 | * | 1.3 | 2.0 | 3.0 | 3.9 | 4.8 | 7.0 | 18.7 | 91.5 |
| 2001..... | 5.9 | 60.7 | 1.6 | 0.5 | 2.8 | 4.2 | 6.0 | 8.3 | 6.9 | 9.9 | 19.3 | 108.2 |
| 2000..... | 5.9 | 65.6 | 1.7 | 0.5 | 2.1 | 4.0 | 6.7 | 8.0 | 6.7 | 9.9 | 20.5 | 102.2 |
| 1999..... | 3.8 | 54.9 | 1.0 | * | 1.3 | 2.2 | 3.0 | 3.1 | 3.8 | 7.0 | 19.4 | 87.0 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.5 | 60.0 | 1.2 | * | 2.3 | 2.8 | 4.5 | 4.9 | 5.8 | 7.1 | 15.7 | 86.9 |
| 2003..... | 4.8 | 69.9 | * | * | 1.7 | 3.4 | 4.4 | 5.7 | 5.6 | 8.1 | 18.8 | 95.4 |
| 2002..... | 4.6 | 66.4 | 1.6 | * | 1.7 | 2.9 | 4.0 | 4.7 | 6.3 | 9.6 | 23.4 | 88.7 |
| 2001..... | 7.4 | 67.2 | 1.8 | * | 3.9 | 6.0 | 8.5 | 12.6 | 10.6 | 14.2 | 25.0 | 87.6 |
| 2000..... | 7.5 | 78.6 | 2.1 | * | 2.9 | 5.8 | 9.9 | 12.4 | 9.0 | 14.3 | 25.6 | 106.8 |
| 1999..... | 4.4 | 61.3 | * | * | 1.9 | 2.9 | 4.3 | 4.7 | 5.5 | 8.3 | 24.0 | 77.6 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 3.3 | 40.4 | * | * | 0.8 | 1.3 | 1.6 | 2.4 | 2.2 | 5.5 | 16.8 | 105.9 |
| 2003..... | 3.2 | 43.2 | * | * | 1.0 | 1.0 | 1.6 | 1.7 | * | 4.6 | 18.2 | 118.8 |
| 2002..... | 3.3 | 46.4 | * | * | 0.9 | 1.0 | 2.0 | 3.2 | 3.5 | 5.0 | 15.5 | 92.8 |
| 2001..... | 4.3 | 53.9 | * | 0.6 | 1.5 | 2.1 | 3.4 | 4.1 | 3.6 | 6.5 | 15.4 | 118.2 |
| 2000..... | 4.1 | 52.1 | 1.4 | * | 1.3 | 2.0 | 3.2 | 3.7 | 4.6 | 6.5 | 17.2 | 99.9 |
| 1999..... | 3.1 | 48.2 | * | * | * | 1.3 | 1.6 | 1.7 | 2.4 | 6.0 | 16.3 | 91.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 11.2 | 90.9 | 1.2 | 0.3 | 1.6 | 2.8 | 4.5 | 5.9 | 7.2 | 13.0 | 40.9 | 226.7 |
| 2003..... | 11.8 | 89.4 | 1.8 | 0.5 | 2.1 | 3.5 | 5.7 | 7.3 | 7.7 | 14.3 | 39.9 | 228.4 |
| 2002..... | 11.3 | 92.8 | 1.7 | 0.4 | 1.9 | 3.6 | 5.6 | 6.9 | 7.7 | 13.5 | 39.6 | 211.9 |
| 2001..... | 12.1 | 92.6 | 1.8 | 0.4 | 2.3 | 3.7 | 7.0 | 8.9 | 9.2 | 15.4 | 40.7 | 209.6 |
| 2000..... | 12.0 | 102.4 | 1.6 | 0.4 | 2.2 | 3.8 | 6.5 | 8.2 | 9.3 | 16.8 | 41.4 | 211.6 |
| 1999..... | 10.3 | 101.9 | 1.6 | 0.4 | 1.6 | 2.8 | 4.7 | 6.0 | 8.0 | 15.0 | 36.7 | 190.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 10.2 | 106.6 | 1.2 | 0.3 | 2.0 | 3.8 | 5.7 | 7.8 | 9.2 | 15.5 | 43.1 | 188.0 |
| 2003..... | 11.0 | 102.9 | 2.0 | 0.5 | 2.8 | 4.7 | 7.2 | 9.4 | 9.9 | 17.7 | 43.0 | 181.1 |
| 2002..... | 10.9 | 106.2 | 1.9 | 0.4 | 2.5 | 4.8 | 7.3 | 9.1 | 10.1 | 16.3 | 44.6 | 178.6 |
| 2001..... | 11.8 | 106.2 | 2.1 | 0.5 | 3.0 | 4.9 | 9.0 | 11.7 | 12.0 | 18.2 | 44.6 | 178.2 |
| 2000..... | 11.9 | 118.0 | 1.7 | 0.5 | 2.9 | 5.1 | 8.6 | 11.2 | 12.3 | 20.7 | 46.6 | 185.4 |
| 1999..... | 10.0 | 116.7 | 1.9 | 0.4 | 2.2 | 3.7 | 6.1 | 8.2 | 10.6 | 18.7 | 40.6 | 169.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 12.1 | 74.3 | 1.2 | 0.3 | 1.0 | 1.9 | 3.2 | 4.2 | 5.3 | 10.9 | 39.4 | 244.0 |
| 2003..... | 12.6 | 75.3 | 1.6 | 0.5 | 1.4 | 2.4 | 4.1 | 5.3 | 5.6 | 11.4 | 37.8 | 249.2 |
| 2002..... | 11.7 | 78.7 | 1.4 | 0.3 | 1.3 | 2.3 | 4.0 | 4.8 | 5.4 | 11.2 | 36.3 | 226.3 |
| 2001..... | 12.3 | 78.3 | 1.5 | 0.4 | 1.5 | 2.6 | 5.1 | 6.2 | 6.6 | 13.1 | 38.0 | 222.7 |
| 2000..... | 12.1 | 86.1 | 1.5 | 0.4 | 1.5 | 2.6 | 4.5 | 5.4 | 6.6 | 13.6 | 38.0 | 222.3 |
| 1999..... | 10.5 | 86.3 | 1.4 | 0.3 | 1.0 | 2.0 | 3.3 | 3.9 | 5.7 | 11.9 | 34.2 | 199.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 11.7 | 71.6 | 1.0 | 0.2 | 1.5 | 2.8 | 4.3 | 5.6 | 6.7 | 12.5 | 40.7 | 230.0 |
| 2003..... | 12.3 | 72.6 | 1.5 | 0.4 | 2.2 | 3.4 | 5.5 | 6.7 | 7.1 | 13.7 | 40.1 | 232.9 |
| 2002..... | 11.6 | 76.0 | 1.3 | 0.3 | 1.8 | 3.4 | 5.4 | 6.1 | 7.1 | 13.1 | 39.4 | 215.8 |
| 2001..... | 12.4 | 73.1 | 1.5 | 0.3 | 2.2 | 3.6 | 6.9 | 8.5 | 8.6 | 14.7 | 40.4 | 213.3 |
| 2000..... | 12.2 | 79.7 | 1.4 | 0.3 | 2.1 | 3.6 | 6.3 | 7.6 | 8.6 | 15.9 | 40.9 | 215.8 |
| 1999..... | 10.2 | 79.3 | 1.4 | 0.3 | 1.4 | 2.5 | 4.1 | 5.2 | 7.3 | 14.3 | 35.7 | 193.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 10.3 | 83.1 | 1.0 | 0.2 | 2.1 | 3.7 | 5.6 | 7.2 | 8.4 | 14.9 | 42.8 | 191.0 |
| 2003..... | 11.0 | 87.4 | 1.6 | 0.4 | 2.9 | 4.6 | 6.9 | 8.6 | 9.0 | 16.8 | 42.8 | 183.3 |
| 2002..... | 10.7 | 88.2 | 1.4 | 0.3 | 2.5 | 4.7 | 6.9 | 7.9 | 9.3 | 15.7 | 43.6 | 181.5 |
| 2001..... | 11.8 | 85.2 | 1.7 | 0.4 | 2.9 | 4.9 | 8.8 | 11.1 | 10.9 | 17.3 | 44.7 | 180.4 |
| 2000..... | 11.8 | 91.8 | 1.7 | 0.4 | 2.8 | 4.8 | 8.5 | 10.3 | 11.1 | 19.3 | 46.0 | 189.6 |
| 1999..... | 9.6 | 92.3 | 1.7 | 0.3 | 1.9 | 3.3 | 5.5 | 6.9 | 9.5 | 17.7 | 39.2 | 171.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 13.0 | 59.4 | 1.0 | 0.2 | 1.0 | 1.8 | 3.0 | 4.0 | 5.0 | 10.4 | 39.3 | 247.5 |
| 2003..... | 13.5 | 57.1 | 1.4 | 0.4 | 1.4 | 2.2 | 4.0 | 4.8 | 5.4 | 11.0 | 38.3 | 254.7 |
| 2002..... | 12.5 | 63.2 | 1.3 | 0.2 | 1.1 | 2.2 | 3.8 | 4.3 | 5.1 | 10.9 | 36.6 | 230.6 |
| 2001..... | 13.1 | 60.4 | 1.3 | 0.3 | 1.5 | 2.4 | 5.0 | 5.9 | 6.4 | 12.5 | 37.6 | 227.0 |
| 2000..... | 12.6 | 66.9 | 1.2 | 0.3 | 1.3 | 2.3 | 4.1 | 4.9 | 6.2 | 13.0 | 37.5 | 226.4 |
| 1999..... | 10.9 | 65.5 | 1.0 | 0.3 | 0.9 | 1.8 | 2.8 | 3.5 | 5.2 | 11.4 | 33.4 | 202.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 11.9 | 174.5 | 2.0 | 0.6 | 1.8 | 4.1 | 6.6 | 9.8 | 13.4 | 20.8 | 53.1 | 223.3 |
| 2003..... | 12.7 | 161.6 | 3.3 | 0.8 | 2.5 | 5.1 | 8.6 | 13.6 | 14.3 | 24.0 | 45.7 | 214.5 |
| 2002..... | 12.6 | 165.4 | 2.7 | 0.8 | 2.6 | 5.3 | 8.8 | 13.9 | 13.9 | 20.4 | 49.9 | 198.1 |
| 2001..... | 12.9 | 176.9 | 3.0 | 0.6 | 2.7 | 5.2 | 9.3 | 14.0 | 15.1 | 24.6 | 50.3 | 194.2 |
| 2000..... | 13.9 | 205.6 | 2.3 | 0.7 | 3.2 | 5.9 | 9.3 | 14.8 | 17.3 | 29.2 | 56.1 | 189.2 |
| 1999..... | 13.0 | 202.0 | 2.9 | 0.5 | 2.7 | 5.0 | 9.1 | 13.2 | 16.0 | 24.6 | 54.8 | 182.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 12.8 | 208.3 | 2.2 | * | 2.3 | 5.3 | 8.1 | 13.8 | 18.9 | 25.4 | 59.3 | 182.9 |
| 2003..... | 13.8 | 171.9 | 3.6 | 0.8 | 3.0 | 7.0 | 11.2 | 17.9 | 20.7 | 32.2 | 52.8 | 197.1 |
| 2002..... | 14.4 | 186.3 | 3.6 | 0.9 | 3.1 | 7.1 | 11.7 | 19.6 | 20.0 | 26.4 | 68.6 | 182.7 |
| 2001..... | 14.2 | 195.9 | 3.7 | 0.8 | 3.8 | 6.1 | 11.4 | 18.2 | 23.8 | 30.7 | 52.2 | 186.0 |
| 2000..... | 15.7 | 236.8 | 2.2 | 0.9 | 3.8 | 7.8 | 10.9 | 20.6 | 25.1 | 40.0 | 66.0 | 163.2 |
| 1999..... | 14.9 | 228.0 | 2.9 | 0.6 | 3.6 | 6.9 | 11.6 | 19.4 | 23.0 | 32.8 | 66.9 | 180.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 11.0 | 139.4 | 1.7 | * | 1.3 | 3.1 | 5.3 | 6.3 | 8.9 | 17.6 | 49.6 | 239.3 |
| 2003..... | 11.6 | 150.9 | 3.0 | 0.8 | 1.9 | 3.5 | 6.3 | 10.0 | 9.3 | 18.2 | 41.7 | 221.2 |
| 2002..... | 10.9 | 143.6 | 1.9 | 0.7 | 2.0 | 3.7 | 6.2 | 9.0 | 8.9 | 16.1 | 39.3 | 204.0 |
| 2001..... | 11.7 | 156.8 | 2.2 | * | 1.7 | 4.3 | 7.4 | 10.3 | 8.1 | 20.3 | 49.2 | 197.2 |
| 2000..... | 12.4 | 173.4 | 2.4 | * | 2.6 | 4.3 | 7.8 | 9.9 | 11.0 | 21.6 | 50.5 | 198.9 |
| 1999..... | 11.2 | 175.3 | 2.8 | * | 1.9 | 3.2 | 7.0 | 7.9 | 10.3 | 18.8 | 48.0 | 182.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Senility (R54) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | * | * | 0.1 | 0.8 | 7.4 | 78.1 |
| 2003..... | 1.8 | * | * | * | * | * | * | * | 0.1 | 0.9 | 6.9 | 78.3 |
| 2002..... | 1.7 | * | * | * | * | * | * | * | * | 0.8 | 6.4 | 73.8 |
| 2001..... | 1.7 | * | * | * | * | * | * | * | * | 0.9 | 6.5 | 74.3 |
| 2000..... | 1.5 | * | * | * | * | * | * | * | * | 0.8 | 6.5 | 68.4 |
| 1999..... | 1.3 | * | * | * | * | * | * | * | * | 0.7 | 5.4 | 57.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | * | 0.9 | 7.6 | 58.8 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | * | 1.1 | 6.7 | 57.2 |
| 2002..... | 0.9 | * | * | * | * | * | * | * | * | 0.8 | 6.4 | 57.9 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 1.0 | 6.5 | 58.2 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | 0.9 | 6.5 | 52.7 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 1.0 | 5.4 | 47.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | * | * | * | * | * | 0.7 | 7.2 | 86.8 |
| 2003..... | 2.6 | * | * | * | * | * | * | * | * | 0.8 | 7.1 | 87.6 |
| 2002..... | 2.4 | * | * | * | * | * | * | * | * | 0.8 | 6.5 | 80.6 |
| 2001..... | 2.3 | * | * | * | * | * | * | * | * | 0.8 | 6.5 | 81.0 |
| 2000..... | 2.2 | * | * | * | * | * | * | * | * | 0.8 | 6.4 | 74.7 |
| 1999..... | 1.8 | * | * | * | * | * | * | * | * | 0.5 | 5.5 | 61.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | * | * | * | * | * | 0.8 | 7.5 | 81.0 |
| 2003..... | 2.1 | * | * | * | * | * | * | * | 0.1 | 1.0 | 7.3 | 81.1 |
| 2002..... | 2.0 | * | * | * | * | * | * | * | * | 0.8 | 6.7 | 76.1 |
| 2001..... | 1.9 | * | * | * | * | * | * | * | * | 0.9 | 6.6 | 76.6 |
| 2000..... | 1.7 | * | * | * | * | * | * | * | * | 0.8 | 6.4 | 70.1 |
| 1999..... | 1.5 | * | * | * | * | * | * | * | * | 0.8 | 5.3 | 59.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | * | * | * | 0.9 | 7.6 | 61.8 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | * | 1.0 | 6.9 | 59.3 |
| 2002..... | 1.1 | * | * | * | * | * | * | * | * | 0.8 | 6.6 | 59.5 |
| 2001..... | 1.1 | * | * | * | * | * | * | * | * | 1.0 | 6.7 | 59.4 |
| 2000..... | 1.0 | * | * | * | * | * | * | * | * | 0.9 | 6.5 | 54.9 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | 1.0 | 5.2 | 48.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | * | * | * | * | * | 0.7 | 7.4 | 89.7 |
| 2003..... | 3.1 | * | * | * | * | * | * | * | * | 0.9 | 7.5 | 90.6 |
| 2002..... | 2.8 | * | * | * | * | * | * | * | * | 0.8 | 6.7 | 83.3 |
| 2001..... | 2.7 | * | * | * | * | * | * | * | * | 0.7 | 6.6 | 83.7 |
| 2000..... | 2.5 | * | * | * | * | * | * | * | * | 0.7 | 6.4 | 76.3 |
| 1999..... | 2.1 | * | * | * | * | * | * | * | * | 0.5 | 5.4 | 64.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Senility (R54) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | * | 8.0 | 56.1 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 4.6 | 56.8 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | * | 5.3 | 56.0 |
| 2001..... | 0.8 | * | * | * | * | * | * | * | * | * | 6.8 | 57.7 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | 1.5 | 8.7 | 56.6 |
| 1999..... | 0.6 | * | * | * | * | * | * | * | * | * | 8.0 | 39.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | * | * | * | * | * | 10.3 | 30.5 |
| 2003..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 38.4 |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | 50.3 |
| 2001..... | 0.5 | * | * | * | * | * | * | * | * | * | 6.3 | 58.9 |
| 2000..... | 0.4 | * | * | * | * | * | * | * | * | * | 8.9 | 35.2 |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | 9.4 | 36.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | * | * | * | * | 6.8 | 66.3 |
| 2003..... | 1.0 | * | * | * | * | * | * | * | * | * | 4.2 | 64.0 |
| 2002..... | 1.0 | * | * | * | * | * | * | * | * | * | 5.5 | 58.1 |
| 2001..... | 1.0 | * | * | * | * | * | * | * | * | * | 7.1 | 57.3 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | * | * | 8.5 | 64.5 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | * | 7.2 | 40.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Ill-defined and unknown causes of mortality (R95-R99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | 80.7 | 1.0 | 0.2 | 1.2 | 2.1 | 3.5 | 4.6 | 4.9 | 6.8 | 14.3 | 52.4 |
| 2003..... | 6.0 | 81.2 | 1.3 | 0.3 | 1.6 | 2.6 | 4.5 | 5.7 | 5.4 | 7.8 | 15.1 | 59.5 |
| 2002..... | 6.0 | 83.8 | 1.3 | 0.2 | 1.5 | 2.6 | 4.4 | 5.4 | 5.6 | 7.6 | 16.5 | 58.3 |
| 2001..... | 7.1 | 84.8 | 1.5 | 0.3 | 2.0 | 3.3 | 6.0 | 7.8 | 7.2 | 9.6 | 18.1 | 62.1 |
| 2000..... | 7.3 | 93.9 | 1.2 | 0.3 | 1.8 | 3.2 | 5.7 | 7.1 | 7.2 | 11.0 | 20.5 | 73.0 |
| 1999..... | 5.9 | 91.3 | 1.3 | 0.3 | 1.3 | 2.2 | 3.7 | 4.8 | 5.7 | 9.2 | 17.3 | 66.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.0 | 94.9 | 1.1 | 0.2 | 1.7 | 2.9 | 4.7 | 6.1 | 6.6 | 8.7 | 16.7 | 50.7 |
| 2003..... | 6.8 | 94.3 | 1.4 | 0.3 | 2.1 | 3.6 | 5.8 | 7.6 | 7.3 | 10.7 | 18.4 | 53.6 |
| 2002..... | 6.8 | 96.5 | 1.5 | 0.3 | 2.0 | 3.5 | 5.8 | 7.3 | 7.7 | 9.8 | 20.0 | 55.0 |
| 2001..... | 8.2 | 96.5 | 1.8 | 0.3 | 2.7 | 4.5 | 7.8 | 10.4 | 9.6 | 12.2 | 21.8 | 58.1 |
| 2000..... | 8.5 | 108.9 | 1.4 | 0.4 | 2.4 | 4.4 | 7.6 | 10.0 | 9.8 | 14.0 | 24.3 | 72.1 |
| 1999..... | 6.7 | 104.3 | 1.4 | 0.3 | 1.8 | 2.9 | 4.9 | 6.6 | 7.8 | 12.0 | 20.5 | 64.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.6 | 65.8 | 0.9 | 0.2 | 0.8 | 1.4 | 2.4 | 3.1 | 3.3 | 5.2 | 12.8 | 53.2 |
| 2003..... | 5.2 | 67.6 | 1.2 | 0.3 | 1.0 | 1.6 | 3.1 | 4.0 | 3.6 | 5.4 | 12.9 | 62.1 |
| 2002..... | 5.1 | 70.4 | 1.1 | 0.2 | 0.9 | 1.6 | 3.0 | 3.5 | 3.8 | 5.8 | 14.1 | 59.8 |
| 2001..... | 6.1 | 72.4 | 1.2 | 0.3 | 1.2 | 2.0 | 4.3 | 5.3 | 5.0 | 7.4 | 15.7 | 63.8 |
| 2000..... | 6.2 | 78.0 | 1.1 | 0.3 | 1.2 | 2.0 | 3.8 | 4.4 | 4.9 | 8.4 | 17.9 | 73.3 |
| 1999..... | 5.2 | 77.7 | 1.1 | 0.3 | 0.7 | 1.4 | 2.6 | 3.0 | 3.8 | 6.8 | 15.2 | 67.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.0 | 49.4 | 0.9 | * | 1.3 | 1.7 | 2.7 | 3.1 | 3.0 | 3.6 | 7.2 | 26.2 |
| 2003..... | 3.0 | 55.2 | 0.7 | * | 1.0 | 1.8 | 2.4 | 3.1 | 2.8 | 3.7 | 7.3 | 35.8 |
| 2002..... | 3.1 | 55.2 | 0.9 | * | 1.1 | 1.6 | 2.3 | 3.1 | 3.8 | 5.0 | 9.8 | 33.8 |
| 2001..... | 5.1 | 60.3 | 1.4 | 0.4 | 2.6 | 3.6 | 5.4 | 7.8 | 6.2 | 8.1 | 10.8 | 40.0 |
| 2000..... | 5.1 | 63.8 | 1.5 | 0.4 | 1.9 | 3.6 | 6.1 | 7.4 | 6.1 | 7.7 | 13.6 | 39.1 |
| 1999..... | 3.0 | 54.3 | 0.8 | * | 1.1 | 1.9 | 2.5 | 2.7 | 2.9 | 4.2 | 11.9 | 28.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.8 | 59.2 | * | * | 1.8 | 2.4 | 4.0 | 4.1 | 4.4 | 4.2 | 8.2 | 30.8 |
| 2003..... | 4.0 | 67.9 | * | * | 1.3 | 2.7 | 3.7 | 4.8 | 4.4 | 5.7 | 9.6 | 32.3 |
| 2002..... | 3.8 | 65.4 | * | * | 1.4 | 2.4 | 3.2 | 3.8 | 5.0 | 7.1 | 13.5 | 45.2 |
| 2001..... | 6.7 | 66.7 | 1.6 | * | 3.6 | 5.4 | 7.9 | 11.9 | 9.9 | 12.4 | 16.1 | * |
| 2000..... | 6.8 | 77.3 | 1.9 | * | 2.6 | 5.2 | 8.9 | 11.5 | 8.6 | 11.0 | 16.2 | 52.4 |
| 1999..... | 3.8 | 60.2 | * | * | 1.6 | 2.6 | 3.7 | 4.1 | 4.2 | 5.5 | 16.7 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.1 | 39.2 | * | * | 0.6 | 1.0 | 1.3 | 2.0 | * | 3.1 | 6.4 | 23.8 |
| 2003..... | 2.1 | 42.0 | * | * | * | 0.7 | 1.0 | 1.5 | * | * | 5.8 | 37.5 |
| 2002..... | 2.3 | 44.4 | * | * | 0.7 | 0.7 | 1.4 | 2.3 | 2.8 | 3.3 | 7.3 | 28.2 |
| 2001..... | 3.4 | 53.7 | * | * | 1.3 | 1.5 | 2.8 | 3.8 | 2.9 | 4.7 | 7.3 | 44.7 |
| 2000..... | 3.3 | 49.7 | * | * | 1.1 | 1.6 | 3.0 | 3.3 | 3.9 | 5.1 | 11.9 | 32.6 |
| 1999..... | 2.3 | 48.0 | * | * | * | 1.1 | 1.1 | 1.4 | * | * | 8.7 | 28.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Ill-defined and unknown causes of mortality (R95-R99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.6 | 88.9 | 1.0 | 0.2 | 1.2 | 2.2 | 3.6 | 4.7 | 5.0 | 7.0 | 14.6 | 53.5 |
| 2003..... | 6.4 | 87.8 | 1.4 | 0.3 | 1.7 | 2.8 | 4.7 | 6.0 | 5.6 | 8.1 | 15.4 | 60.1 |
| 2002..... | 6.3 | 90.8 | 1.4 | 0.3 | 1.5 | 2.8 | 4.7 | 5.5 | 5.7 | 7.8 | 16.7 | 58.7 |
| 2001..... | 7.4 | 90.4 | 1.5 | 0.3 | 1.8 | 3.2 | 6.1 | 7.7 | 7.2 | 9.6 | 18.4 | 62.8 |
| 2000..... | 7.6 | 100.7 | 1.2 | 0.3 | 1.8 | 3.1 | 5.5 | 7.0 | 7.2 | 11.0 | 20.7 | 74.2 |
| 1999..... | 6.3 | 99.7 | 1.3 | 0.3 | 1.3 | 2.2 | 3.8 | 4.9 | 5.8 | 9.3 | 17.5 | 67.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.3 | 104.4 | 1.0 | 0.1 | 1.6 | 3.0 | 4.7 | 6.2 | 6.6 | 8.9 | 17.0 | 51.4 |
| 2003..... | 7.2 | 100.7 | 1.6 | 0.3 | 2.3 | 3.8 | 6.1 | 7.7 | 7.4 | 11.0 | 18.7 | 54.3 |
| 2002..... | 7.2 | 104.1 | 1.6 | 0.3 | 2.0 | 3.8 | 6.2 | 7.5 | 7.8 | 9.9 | 20.1 | 55.0 |
| 2001..... | 8.4 | 103.4 | 1.8 | 0.3 | 2.4 | 4.2 | 7.7 | 10.2 | 9.5 | 12.1 | 22.0 | 59.0 |
| 2000..... | 8.6 | 116.2 | 1.3 | 0.3 | 2.3 | 4.2 | 7.3 | 9.7 | 9.7 | 14.0 | 24.6 | 72.9 |
| 1999..... | 7.0 | 114.1 | 1.6 | 0.3 | 1.8 | 2.9 | 5.0 | 6.6 | 7.9 | 12.3 | 20.5 | 65.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.9 | 72.8 | 0.9 | 0.2 | 0.8 | 1.4 | 2.6 | 3.1 | 3.4 | 5.3 | 13.0 | 54.4 |
| 2003..... | 5.6 | 74.2 | 1.3 | 0.4 | 1.1 | 1.8 | 3.4 | 4.2 | 3.8 | 5.6 | 13.3 | 62.7 |
| 2002..... | 5.5 | 76.8 | 1.2 | 0.2 | 1.0 | 1.8 | 3.2 | 3.6 | 3.8 | 5.9 | 14.4 | 60.3 |
| 2001..... | 6.4 | 76.8 | 1.2 | 0.3 | 1.2 | 2.1 | 4.5 | 5.3 | 5.1 | 7.6 | 16.1 | 64.4 |
| 2000..... | 6.6 | 84.4 | 1.0 | 0.2 | 1.2 | 2.0 | 3.8 | 4.4 | 4.9 | 8.6 | 18.2 | 74.7 |
| 1999..... | 5.6 | 84.6 | 1.1 | 0.3 | 0.8 | 1.5 | 2.7 | 3.2 | 3.9 | 6.9 | 15.5 | 68.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | 70.0 | 0.8 | 0.1 | 1.3 | 2.3 | 3.6 | 4.5 | 4.7 | 6.7 | 14.8 | 54.1 |
| 2003..... | 6.3 | 70.8 | 1.2 | 0.3 | 1.8 | 2.8 | 4.7 | 5.6 | 5.3 | 7.9 | 15.6 | 61.7 |
| 2002..... | 6.2 | 74.5 | 1.1 | 0.2 | 1.5 | 2.7 | 4.6 | 5.0 | 5.4 | 7.6 | 16.7 | 59.6 |
| 2001..... | 7.3 | 71.1 | 1.2 | 0.2 | 1.8 | 3.1 | 6.1 | 7.5 | 6.9 | 9.4 | 18.3 | 63.9 |
| 2000..... | 7.4 | 78.2 | 1.1 | 0.2 | 1.7 | 3.0 | 5.4 | 6.6 | 6.8 | 10.6 | 20.7 | 75.7 |
| 1999..... | 6.0 | 77.4 | 1.1 | 0.2 | 1.2 | 2.0 | 3.4 | 4.3 | 5.3 | 9.0 | 17.1 | 68.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 6.1 | 81.8 | 0.9 | * | 1.7 | 3.2 | 4.8 | 5.9 | 6.3 | 8.7 | 17.1 | 52.2 |
| 2003..... | 7.0 | 85.3 | 1.2 | 0.3 | 2.4 | 3.8 | 6.0 | 7.3 | 6.8 | 10.5 | 18.8 | 55.4 |
| 2002..... | 6.9 | 86.6 | 1.2 | 0.2 | 2.1 | 3.8 | 5.9 | 6.7 | 7.3 | 9.6 | 19.6 | 56.1 |
| 2001..... | 8.2 | 82.5 | 1.4 | 0.2 | 2.4 | 4.2 | 7.7 | 9.8 | 8.9 | 11.7 | 21.8 | 60.4 |
| 2000..... | 8.3 | 90.1 | 1.3 | 0.3 | 2.3 | 4.1 | 7.3 | 9.1 | 9.1 | 13.1 | 24.4 | 73.8 |
| 1999..... | 6.5 | 90.1 | 1.4 | 0.2 | 1.7 | 2.7 | 4.5 | 5.7 | 7.2 | 11.7 | 19.9 | 64.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.0 | 57.7 | 0.7 | 0.2 | 0.8 | 1.4 | 2.4 | 3.1 | 3.3 | 5.0 | 13.1 | 55.0 |
| 2003..... | 5.7 | 55.8 | 1.2 | 0.3 | 1.1 | 1.8 | 3.4 | 3.9 | 3.8 | 5.7 | 13.5 | 64.4 |
| 2002..... | 5.5 | 61.8 | 1.0 | * | 0.9 | 1.7 | 3.2 | 3.3 | 3.7 | 5.8 | 14.7 | 61.1 |
| 2001..... | 6.5 | 59.0 | 1.1 | 0.2 | 1.2 | 2.0 | 4.4 | 5.1 | 5.0 | 7.4 | 15.9 | 65.4 |
| 2000..... | 6.6 | 65.7 | 0.8 | 0.2 | 1.1 | 1.9 | 3.6 | 4.1 | 4.7 | 8.5 | 18.2 | 76.4 |
| 1999..... | 5.5 | 64.0 | 0.8 | 0.2 | 0.6 | 1.3 | 2.4 | 2.9 | 3.6 | 6.8 | 15.3 | 69.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Ill-defined and unknown causes of mortality (R95-R99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 7.4 | 171.2 | 1.7 | * | 1.1 | 2.6 | 4.8 | 7.1 | 8.0 | 10.7 | 16.7 | 55.8 |
| 2003..... | 8.4 | 160.3 | 2.6 | 0.5 | 1.9 | 3.7 | 6.7 | 9.9 | 9.8 | 12.1 | 16.6 | 55.0 |
| 2002..... | 8.6 | 161.0 | 2.4 | 0.6 | 1.6 | 3.6 | 6.7 | 10.7 | 9.7 | 10.9 | 20.3 | 57.2 |
| 2001..... | 9.0 | 174.6 | 2.4 | 0.5 | 1.9 | 4.0 | 7.0 | 11.1 | 10.5 | 12.9 | 22.7 | 56.2 |
| 2000..... | 10.0 | 202.3 | 1.7 | 0.5 | 2.4 | 4.3 | 7.5 | 11.8 | 12.0 | 17.1 | 25.1 | 67.1 |
| 1999..... | 9.3 | 198.3 | 2.4 | 0.4 | 2.0 | 3.5 | 7.1 | 10.1 | 11.2 | 13.7 | 24.5 | 71.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 8.8 | 202.7 | 1.8 | * | 1.4 | 3.3 | 5.6 | 10.4 | 11.7 | 13.0 | 20.0 | 53.9 |
| 2003..... | 10.1 | 169.7 | 3.0 | * | 2.3 | 5.1 | 8.7 | 13.0 | 15.1 | 19.4 | 21.9 | 59.7 |
| 2002..... | 10.6 | 181.5 | 3.0 | * | 2.0 | 4.7 | 9.2 | 15.5 | 14.5 | 15.4 | 31.1 | 54.7 |
| 2001..... | 10.6 | 193.1 | 3.1 | * | 2.6 | 4.9 | 8.3 | 14.2 | 16.2 | 17.6 | 27.6 | 49.7 |
| 2000..... | 12.1 | 233.7 | 1.8 | 0.7 | 2.9 | 5.8 | 8.7 | 16.4 | 17.3 | 25.3 | 33.0 | 71.6 |
| 1999..... | 11.4 | 223.1 | 2.4 | * | 2.7 | 5.0 | 9.0 | 15.0 | 16.6 | 20.1 | 31.6 | 86.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.1 | 138.4 | * | * | 0.8 | 2.0 | 4.2 | 4.3 | 5.0 | 9.1 | 14.8 | 56.6 |
| 2003..... | 6.9 | 150.5 | 2.3 | * | 1.5 | 2.4 | 5.0 | 7.3 | 5.4 | 6.9 | 13.5 | 53.2 |
| 2002..... | 6.7 | 139.8 | 1.8 | * | 1.3 | 2.7 | 4.4 | 6.6 | 5.8 | 7.8 | 14.3 | 58.1 |
| 2001..... | 7.6 | 155.2 | 1.7 | * | 1.1 | 3.2 | 5.9 | 8.5 | 5.9 | 9.5 | 19.9 | 58.6 |
| 2000..... | 8.1 | 169.8 | 1.7 | * | 2.0 | 2.9 | 6.4 | 7.8 | 7.7 | 11.3 | 20.7 | 65.4 |
| 1999..... | 7.4 | 172.8 | 2.4 | * | 1.4 | 2.1 | 5.4 | 5.9 | 6.8 | 9.1 | 20.5 | 65.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other ill-defined and unknown causes of mortality (R96-R99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.5 | 25.6 | 1.0 | 0.2 | 1.2 | 2.1 | 3.5 | 4.6 | 4.9 | 6.8 | 14.3 | 52.4 |
| 2003..... | 5.2 | 27.2 | 1.3 | 0.3 | 1.6 | 2.6 | 4.5 | 5.7 | 5.4 | 7.8 | 15.1 | 59.5 |
| 2002..... | 5.2 | 26.9 | 1.3 | 0.2 | 1.5 | 2.6 | 4.4 | 5.4 | 5.6 | 7.6 | 16.5 | 58.3 |
| 2001..... | 6.3 | 29.4 | 1.5 | 0.3 | 2.0 | 3.3 | 6.0 | 7.8 | 7.2 | 9.6 | 18.1 | 62.1 |
| 2000..... | 6.4 | 27.6 | 1.2 | 0.3 | 1.8 | 3.2 | 5.7 | 7.1 | 7.2 | 11.0 | 20.5 | 73.0 |
| 1999..... | 5.0 | 21.6 | 1.3 | 0.3 | 1.3 | 2.2 | 3.7 | 4.8 | 5.7 | 9.2 | 17.3 | 66.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 5.1 | 31.2 | 1.1 | 0.2 | 1.7 | 2.9 | 4.7 | 6.1 | 6.6 | 8.7 | 16.7 | 50.7 |
| 2003..... | 5.9 | 31.7 | 1.4 | 0.3 | 2.1 | 3.6 | 5.8 | 7.6 | 7.3 | 10.7 | 18.4 | 53.6 |
| 2002..... | 5.9 | 31.1 | 1.5 | 0.3 | 2.0 | 3.5 | 5.8 | 7.3 | 7.7 | 9.8 | 20.0 | 55.0 |
| 2001..... | 7.3 | 32.7 | 1.8 | 0.3 | 2.7 | 4.5 | 7.8 | 10.4 | 9.6 | 12.2 | 21.8 | 58.1 |
| 2000..... | 7.4 | 31.5 | 1.4 | 0.4 | 2.4 | 4.4 | 7.6 | 10.0 | 9.8 | 14.0 | 24.3 | 72.1 |
| 1999..... | 5.5 | 24.4 | 1.4 | 0.3 | 1.8 | 2.9 | 4.9 | 6.6 | 7.8 | 12.0 | 20.5 | 64.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.0 | 19.7 | 0.9 | 0.2 | 0.8 | 1.4 | 2.4 | 3.1 | 3.3 | 5.2 | 12.8 | 53.2 |
| 2003..... | 4.6 | 22.5 | 1.2 | 0.3 | 1.0 | 1.6 | 3.1 | 4.0 | 3.6 | 5.4 | 12.9 | 62.1 |
| 2002..... | 4.5 | 22.5 | 1.1 | 0.2 | 0.9 | 1.6 | 3.0 | 3.5 | 3.8 | 5.8 | 14.1 | 59.8 |
| 2001..... | 5.5 | 25.8 | 1.2 | 0.3 | 1.2 | 2.0 | 4.3 | 5.3 | 5.0 | 7.4 | 15.7 | 63.8 |
| 2000..... | 5.5 | 23.4 | 1.1 | 0.3 | 1.2 | 2.0 | 3.8 | 4.4 | 4.9 | 8.4 | 17.9 | 73.3 |
| 1999..... | 4.5 | 18.6 | 1.1 | 0.3 | 0.7 | 1.4 | 2.6 | 3.0 | 3.8 | 6.8 | 15.2 | 67.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | 18.1 | 0.9 | * | 1.3 | 1.7 | 2.7 | 3.1 | 3.0 | 3.6 | 7.2 | 26.2 |
| 2003..... | 2.3 | 21.7 | 0.7 | * | 1.0 | 1.8 | 2.4 | 3.1 | 2.8 | 3.7 | 7.3 | 35.8 |
| 2002..... | 2.3 | 19.8 | 0.9 | * | 1.1 | 1.6 | 2.3 | 3.1 | 3.8 | 5.0 | 9.8 | 33.8 |
| 2001..... | 4.4 | 29.7 | 1.4 | 0.4 | 2.6 | 3.6 | 5.4 | 7.8 | 6.2 | 8.1 | 10.8 | 40.0 |
| 2000..... | 4.2 | 24.6 | 1.5 | 0.4 | 1.9 | 3.6 | 6.1 | 7.4 | 6.1 | 7.7 | 13.6 | 39.1 |
| 1999..... | 2.1 | 13.0 | 0.8 | * | 1.1 | 1.9 | 2.5 | 2.7 | 2.9 | 4.2 | 11.9 | 28.8 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.0 | 22.7 | * | * | 1.8 | 2.4 | 4.0 | 4.1 | 4.4 | 4.2 | 8.2 | 30.8 |
| 2003..... | 3.1 | 26.7 | * | * | 1.3 | 2.7 | 3.7 | 4.8 | 4.4 | 5.7 | 9.6 | 32.3 |
| 2002..... | 2.9 | 22.7 | * | * | 1.4 | 2.4 | 3.2 | 3.8 | 5.0 | 7.1 | 13.5 | 45.2 |
| 2001..... | 6.0 | 32.6 | 1.6 | * | 3.6 | 5.4 | 7.9 | 11.9 | 9.9 | 12.4 | 16.1 | * |
| 2000..... | 5.8 | 30.9 | 1.9 | * | 2.6 | 5.2 | 8.9 | 11.5 | 8.6 | 11.0 | 16.2 | 52.4 |
| 1999..... | 2.7 | 14.5 | * | * | 1.6 | 2.6 | 3.7 | 4.1 | 4.2 | 5.5 | 16.7 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.6 | 13.3 | * | * | 0.6 | 1.0 | 1.3 | 2.0 | * | 3.1 | 6.4 | 23.8 |
| 2003..... | 1.5 | 16.5 | * | * | * | 0.7 | 1.0 | 1.5 | * | * | 5.8 | 37.5 |
| 2002..... | 1.7 | 16.7 | * | * | 0.7 | 0.7 | 1.4 | 2.3 | 2.8 | 3.3 | 7.3 | 28.2 |
| 2001..... | 2.8 | 26.7 | * | * | 1.3 | 1.5 | 2.8 | 3.8 | 2.9 | 4.7 | 7.3 | 44.7 |
| 2000..... | 2.6 | 18.1 | * | * | 1.1 | 1.6 | 3.0 | 3.3 | 3.9 | 5.1 | 11.9 | 32.6 |
| 1999..... | 1.5 | 11.4 | * | * | * | 1.1 | 1.1 | 1.4 | * | * | 8.7 | 28.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other ill-defined and unknown causes of mortality (R96-R99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.8 | 27.1 | 1.0 | 0.2 | 1.2 | 2.2 | 3.6 | 4.7 | 5.0 | 7.0 | 14.6 | 53.5 |
| 2003..... | 5.6 | 28.3 | 1.4 | 0.3 | 1.7 | 2.8 | 4.7 | 6.0 | 5.6 | 8.1 | 15.4 | 60.1 |
| 2002..... | 5.5 | 28.5 | 1.4 | 0.3 | 1.5 | 2.8 | 4.7 | 5.5 | 5.7 | 7.8 | 16.7 | 58.7 |
| 2001..... | 6.6 | 28.9 | 1.5 | 0.3 | 1.8 | 3.2 | 6.1 | 7.7 | 7.2 | 9.6 | 18.4 | 62.8 |
| 2000..... | 6.7 | 27.7 | 1.2 | 0.3 | 1.8 | 3.1 | 5.5 | 7.0 | 7.2 | 11.0 | 20.7 | 74.2 |
| 1999..... | 5.3 | 23.3 | 1.3 | 0.3 | 1.3 | 2.2 | 3.8 | 4.9 | 5.8 | 9.3 | 17.5 | 67.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 5.4 | 33.0 | 1.0 | 0.1 | 1.6 | 3.0 | 4.7 | 6.2 | 6.6 | 8.9 | 17.0 | 51.4 |
| 2003..... | 6.3 | 32.5 | 1.6 | 0.3 | 2.3 | 3.8 | 6.1 | 7.7 | 7.4 | 11.0 | 18.7 | 54.3 |
| 2002..... | 6.3 | 33.0 | 1.6 | 0.3 | 2.0 | 3.8 | 6.2 | 7.5 | 7.8 | 9.9 | 20.1 | 55.0 |
| 2001..... | 7.4 | 32.3 | 1.8 | 0.3 | 2.4 | 4.2 | 7.7 | 10.2 | 9.5 | 12.1 | 22.0 | 59.0 |
| 2000..... | 7.5 | 31.1 | 1.3 | 0.3 | 2.3 | 4.2 | 7.3 | 9.7 | 9.7 | 14.0 | 24.6 | 72.9 |
| 1999..... | 5.8 | 26.3 | 1.6 | 0.3 | 1.8 | 2.9 | 5.0 | 6.6 | 7.9 | 12.3 | 20.5 | 65.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.3 | 20.9 | 0.9 | 0.2 | 0.8 | 1.4 | 2.6 | 3.1 | 3.4 | 5.3 | 13.0 | 54.4 |
| 2003..... | 5.0 | 23.9 | 1.3 | 0.4 | 1.1 | 1.8 | 3.4 | 4.2 | 3.8 | 5.6 | 13.3 | 62.7 |
| 2002..... | 4.8 | 23.7 | 1.2 | 0.2 | 1.0 | 1.8 | 3.2 | 3.6 | 3.8 | 5.9 | 14.4 | 60.3 |
| 2001..... | 5.8 | 25.4 | 1.2 | 0.3 | 1.2 | 2.1 | 4.5 | 5.3 | 5.1 | 7.6 | 16.1 | 64.4 |
| 2000..... | 5.9 | 24.3 | 1.0 | 0.2 | 1.2 | 2.0 | 3.8 | 4.4 | 4.9 | 8.6 | 18.2 | 74.7 |
| 1999..... | 4.8 | 20.1 | 1.1 | 0.3 | 0.8 | 1.5 | 2.7 | 3.2 | 3.9 | 6.9 | 15.5 | 68.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.9 | 19.3 | 0.8 | 0.1 | 1.3 | 2.3 | 3.6 | 4.5 | 4.7 | 6.7 | 14.8 | 54.1 |
| 2003..... | 5.8 | 22.7 | 1.2 | 0.3 | 1.8 | 2.8 | 4.7 | 5.6 | 5.3 | 7.9 | 15.6 | 61.7 |
| 2002..... | 5.6 | 22.4 | 1.1 | 0.2 | 1.5 | 2.7 | 4.6 | 5.0 | 5.4 | 7.6 | 16.7 | 59.6 |
| 2001..... | 6.7 | 22.1 | 1.2 | 0.2 | 1.8 | 3.1 | 6.1 | 7.5 | 6.9 | 9.4 | 18.3 | 63.9 |
| 2000..... | 6.8 | 19.6 | 1.1 | 0.2 | 1.7 | 3.0 | 5.4 | 6.6 | 6.8 | 10.6 | 20.7 | 75.7 |
| 1999..... | 5.2 | 16.0 | 1.1 | 0.2 | 1.2 | 2.0 | 3.4 | 4.3 | 5.3 | 9.0 | 17.1 | 68.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 5.4 | 23.6 | 0.9 | * | 1.7 | 3.2 | 4.8 | 5.9 | 6.3 | 8.7 | 17.1 | 52.2 |
| 2003..... | 6.3 | 27.1 | 1.2 | 0.3 | 2.4 | 3.8 | 6.0 | 7.3 | 6.8 | 10.5 | 18.8 | 55.4 |
| 2002..... | 6.2 | 26.4 | 1.2 | 0.2 | 2.1 | 3.8 | 5.9 | 6.7 | 7.3 | 9.6 | 19.6 | 56.1 |
| 2001..... | 7.5 | 24.9 | 1.4 | 0.2 | 2.4 | 4.2 | 7.7 | 9.8 | 8.9 | 11.7 | 21.8 | 60.4 |
| 2000..... | 7.5 | 21.5 | 1.3 | 0.3 | 2.3 | 4.1 | 7.3 | 9.1 | 9.1 | 13.1 | 24.4 | 73.8 |
| 1999..... | 5.6 | 18.8 | 1.4 | 0.2 | 1.7 | 2.7 | 4.5 | 5.7 | 7.2 | 11.7 | 19.9 | 64.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.5 | 14.8 | 0.7 | 0.2 | 0.8 | 1.4 | 2.4 | 3.1 | 3.3 | 5.0 | 13.1 | 55.0 |
| 2003..... | 5.3 | 18.1 | 1.2 | 0.3 | 1.1 | 1.8 | 3.4 | 3.9 | 3.8 | 5.7 | 13.5 | 64.4 |
| 2002..... | 5.0 | 18.2 | 1.0 | * | 0.9 | 1.7 | 3.2 | 3.3 | 3.7 | 5.8 | 14.7 | 61.1 |
| 2001..... | 6.0 | 19.2 | 1.1 | 0.2 | 1.2 | 2.0 | 4.4 | 5.1 | 5.0 | 7.4 | 15.9 | 65.4 |
| 2000..... | 6.1 | 17.7 | 0.8 | 0.2 | 1.1 | 1.9 | 3.6 | 4.1 | 4.7 | 8.5 | 18.2 | 76.4 |
| 1999..... | 4.9 | 13.1 | 0.8 | 0.2 | 0.6 | 1.3 | 2.4 | 2.9 | 3.6 | 6.8 | 15.3 | 69.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other ill-defined and unknown causes of mortality (R96-R99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | 58.4 | 1.7 | * | 1.1 | 2.6 | 4.8 | 7.1 | 8.0 | 10.7 | 16.7 | 55.8 |
| 2003..... | 6.5 | 53.3 | 2.6 | 0.5 | 1.9 | 3.7 | 6.7 | 9.9 | 9.8 | 12.1 | 16.6 | 55.0 |
| 2002..... | 6.6 | 54.6 | 2.4 | 0.6 | 1.6 | 3.6 | 6.7 | 10.7 | 9.7 | 10.9 | 20.3 | 57.2 |
| 2001..... | 7.0 | 55.0 | 2.4 | 0.5 | 1.9 | 4.0 | 7.0 | 11.1 | 10.5 | 12.9 | 22.7 | 56.2 |
| 2000..... | 7.7 | 62.1 | 1.7 | 0.5 | 2.4 | 4.3 | 7.5 | 11.8 | 12.0 | 17.1 | 25.1 | 67.1 |
| 1999..... | 6.9 | 52.9 | 2.4 | 0.4 | 2.0 | 3.5 | 7.1 | 10.1 | 11.2 | 13.7 | 24.5 | 71.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 6.4 | 69.9 | 1.8 | * | 1.4 | 3.3 | 5.6 | 10.4 | 11.7 | 13.0 | 20.0 | 53.9 |
| 2003..... | 8.0 | 59.0 | 3.0 | * | 2.3 | 5.1 | 8.7 | 13.0 | 15.1 | 19.4 | 21.9 | 59.7 |
| 2002..... | 8.3 | 62.1 | 3.0 | * | 2.0 | 4.7 | 9.2 | 15.5 | 14.5 | 15.4 | 31.1 | 54.7 |
| 2001..... | 8.1 | 60.9 | 3.1 | * | 2.6 | 4.9 | 8.3 | 14.2 | 16.2 | 17.6 | 27.6 | 49.7 |
| 2000..... | 9.3 | 72.3 | 1.8 | 0.7 | 2.9 | 5.8 | 8.7 | 16.4 | 17.3 | 25.3 | 33.0 | 71.6 |
| 1999..... | 8.5 | 60.3 | 2.4 | * | 2.7 | 5.0 | 9.0 | 15.0 | 16.6 | 20.1 | 31.6 | 86.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.6 | 46.4 | * | * | 0.8 | 2.0 | 4.2 | 4.3 | 5.0 | 9.1 | 14.8 | 56.6 |
| 2003..... | 5.2 | 47.4 | 2.3 | * | 1.5 | 2.4 | 5.0 | 7.3 | 5.4 | 6.9 | 13.5 | 53.2 |
| 2002..... | 5.1 | 46.7 | 1.8 | * | 1.3 | 2.7 | 4.4 | 6.6 | 5.8 | 7.8 | 14.3 | 58.1 |
| 2001..... | 5.9 | 48.7 | 1.7 | * | 1.1 | 3.2 | 5.9 | 8.5 | 5.9 | 9.5 | 19.9 | 58.6 |
| 2000..... | 6.3 | 51.4 | 1.7 | * | 2.0 | 2.9 | 6.4 | 7.8 | 7.7 | 11.3 | 20.7 | 65.4 |
| 1999..... | 5.4 | 45.3 | 2.4 | * | 1.4 | 2.1 | 5.4 | 5.9 | 6.8 | 9.1 | 20.5 | 65.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | 1.9 | 0.2 | 0.1 | 0.3 | 0.6 | 0.8 | 1.3 | 2.2 | 5.2 | 18.9 | 95.1 |
| 2003..... | 3.6 | 1.6 | 0.4 | 0.1 | 0.4 | 0.8 | 0.9 | 1.4 | 2.0 | 5.3 | 17.5 | 89.9 |
| 2002..... | 3.3 | 2.0 | 0.3 | 0.1 | 0.4 | 0.8 | 1.0 | 1.3 | 1.9 | 4.9 | 16.5 | 79.4 |
| 2001..... | 3.1 | 2.1 | 0.3 | 0.1 | 0.5 | 0.6 | 1.0 | 1.2 | 1.9 | 4.9 | 15.7 | 72.5 |
| 2000..... | 2.9 | 1.7 | 0.4 | 0.1 | 0.4 | 0.7 | 1.0 | 1.2 | 2.1 | 4.9 | 14.2 | 69.1 |
| 1999..... | 2.8 | 2.2 | 0.3 | 0.1 | 0.3 | 0.6 | 0.9 | 1.1 | 2.1 | 4.9 | 13.8 | 65.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.9 | 2.3 | * | 0.1 | 0.5 | 0.8 | 1.0 | 1.5 | 2.5 | 5.6 | 18.5 | 77.8 |
| 2003..... | 2.8 | 2.2 | 0.4 | 0.1 | 0.5 | 1.0 | 1.1 | 1.7 | 2.4 | 5.6 | 17.5 | 69.6 |
| 2002..... | 2.7 | 2.1 | * | 0.1 | 0.5 | 1.1 | 1.1 | 1.5 | 2.3 | 5.5 | 18.1 | 65.8 |
| 2001..... | 2.5 | 2.8 | * | 0.2 | 0.6 | 0.7 | 1.3 | 1.6 | 2.4 | 5.1 | 16.1 | 60.9 |
| 2000..... | 2.5 | 1.8 | 0.4 | 0.1 | 0.5 | 0.9 | 1.3 | 1.6 | 2.6 | 5.8 | 15.5 | 59.9 |
| 1999..... | 2.3 | 2.6 | 0.3 | 0.1 | 0.4 | 0.7 | 1.1 | 1.5 | 2.6 | 5.4 | 14.7 | 56.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.6 | 1.6 | * | * | 0.2 | 0.5 | 0.6 | 1.0 | 1.9 | 4.9 | 19.2 | 102.8 |
| 2003..... | 4.4 | * | * | * | 0.3 | 0.6 | 0.7 | 1.1 | 1.7 | 5.0 | 17.5 | 98.9 |
| 2002..... | 3.9 | 1.9 | * | * | 0.3 | 0.6 | 0.8 | 1.1 | 1.5 | 4.4 | 15.4 | 85.3 |
| 2001..... | 3.6 | 1.5 | 0.3 | * | 0.3 | 0.5 | 0.7 | 0.9 | 1.4 | 4.7 | 15.5 | 77.3 |
| 2000..... | 3.3 | 1.7 | 0.4 | 0.1 | 0.3 | 0.6 | 0.7 | 0.9 | 1.6 | 4.2 | 13.4 | 72.9 |
| 1999..... | 3.2 | 1.8 | * | * | 0.3 | 0.5 | 0.6 | 0.8 | 1.7 | 4.5 | 13.2 | 69.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | 1.5 | * | 0.1 | 0.3 | 0.5 | 0.7 | 1.1 | 1.9 | 5.0 | 18.5 | 94.8 |
| 2003..... | 3.8 | 1.7 | 0.3 | 0.1 | 0.4 | 0.7 | 0.8 | 1.1 | 1.8 | 4.8 | 17.2 | 90.2 |
| 2002..... | 3.5 | 1.5 | 0.2 | 0.1 | 0.3 | 0.7 | 0.8 | 1.1 | 1.6 | 4.7 | 16.1 | 80.1 |
| 2001..... | 3.2 | 2.0 | 0.3 | 0.1 | 0.4 | 0.5 | 0.8 | 1.0 | 1.6 | 4.4 | 15.5 | 72.8 |
| 2000..... | 3.0 | 1.5 | 0.4 | 0.1 | 0.4 | 0.6 | 0.9 | 1.0 | 1.8 | 4.5 | 13.8 | 70.0 |
| 1999..... | 2.8 | 1.9 | 0.3 | 0.1 | 0.3 | 0.5 | 0.7 | 0.9 | 1.9 | 4.5 | 13.3 | 65.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | * | 0.4 | 0.6 | 0.8 | 1.3 | 2.1 | 5.2 | 18.1 | 76.9 |
| 2003..... | 2.8 | 2.1 | * | * | 0.5 | 0.8 | 0.9 | 1.3 | 2.1 | 5.3 | 17.1 | 68.5 |
| 2002..... | 2.7 | * | * | * | 0.4 | 0.9 | 1.0 | 1.2 | 2.0 | 5.2 | 17.4 | 66.0 |
| 2001..... | 2.5 | 2.7 | * | 0.2 | 0.5 | 0.6 | 1.0 | 1.3 | 1.9 | 4.5 | 16.2 | 60.6 |
| 2000..... | 2.5 | * | * | * | 0.5 | 0.7 | 1.2 | 1.2 | 2.0 | 5.3 | 15.1 | 60.8 |
| 1999..... | 2.3 | 2.2 | * | * | 0.3 | 0.6 | 0.9 | 1.2 | 2.3 | 5.0 | 14.1 | 57.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.0 | * | * | * | 0.2 | 0.4 | 0.6 | 0.9 | 1.7 | 4.7 | 18.8 | 102.8 |
| 2003..... | 4.7 | * | * | * | 0.3 | 0.5 | 0.7 | 0.8 | 1.5 | 4.4 | 17.3 | 99.7 |
| 2002..... | 4.2 | * | * | * | 0.2 | 0.5 | 0.6 | 1.0 | 1.4 | 4.3 | 15.3 | 86.1 |
| 2001..... | 3.9 | * | * | * | 0.3 | 0.4 | 0.6 | 0.8 | 1.3 | 4.4 | 15.0 | 77.9 |
| 2000..... | 3.5 | * | * | * | 0.3 | 0.4 | 0.5 | 0.8 | 1.5 | 3.8 | 13.0 | 73.7 |
| 1999..... | 3.3 | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 1.6 | 4.1 | 12.7 | 68.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | 3.3 | * | * | 0.7 | 1.5 | 1.8 | 2.6 | 5.1 | 9.1 | 28.4 | 111.4 |
| 2003..... | 3.6 | * | * | * | 0.6 | 1.4 | 1.9 | 3.7 | 4.4 | 11.0 | 24.6 | 102.6 |
| 2002..... | 3.3 | 4.3 | * | * | 0.9 | 1.7 | 2.1 | 3.2 | 4.2 | 8.3 | 24.2 | 85.0 |
| 2001..... | 3.2 | * | * | * | 0.9 | 1.2 | 2.2 | 2.7 | 4.4 | 10.6 | 20.8 | 80.2 |
| 2000..... | 3.1 | * | * | * | 0.7 | 1.6 | 1.7 | 3.1 | 5.2 | 10.6 | 22.3 | 65.5 |
| 1999..... | 3.1 | 3.7 | * | * | 0.7 | 1.5 | 2.1 | 3.1 | 4.6 | 10.1 | 22.3 | 71.1 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | 0.9 | 2.0 | 2.6 | 3.4 | 6.8 | 11.1 | 29.1 | 98.6 |
| 2003..... | 3.4 | * | * | * | 0.7 | 1.9 | 2.5 | 4.9 | 5.4 | 11.3 | 25.7 | 99.1 |
| 2002..... | 3.4 | * | * | * | 1.1 | 2.4 | 2.5 | 4.0 | 5.5 | 9.7 | 32.5 | 77.7 |
| 2001..... | 3.2 | * | * | * | 1.2 | 1.3 | 3.1 | 3.8 | 7.4 | 12.3 | 18.3 | 77.4 |
| 2000..... | 3.1 | * | * | * | 0.9 | 2.0 | 2.1 | 4.2 | 7.6 | 13.5 | 24.1 | 56.4 |
| 1999..... | 3.1 | * | * | * | 0.9 | 1.9 | 2.6 | 4.3 | 6.1 | 11.6 | 26.0 | 57.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | * | 1.1 | 1.1 | 2.0 | 3.8 | 7.6 | 28.0 | 116.5 |
| 2003..... | 3.8 | * | * | * | * | 1.0 | 1.4 | 2.6 | 3.6 | 10.8 | 24.0 | 104.0 |
| 2002..... | 3.3 | * | * | * | 0.8 | 1.1 | 1.8 | 2.4 | 3.1 | 7.4 | 19.6 | 87.8 |
| 2001..... | 3.1 | * | * | * | * | 1.1 | 1.5 | 1.8 | 2.0 | 9.3 | 22.2 | 81.3 |
| 2000..... | 3.1 | * | * | * | * | 1.4 | 1.4 | 2.1 | 3.3 | 8.5 | 21.4 | 68.9 |
| 1999..... | 3.1 | * | * | * | * | 1.1 | 1.6 | 2.0 | 3.4 | 9.1 | 20.3 | 76.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XX. External causes of mortality (*U01-*U03,V01-Y89) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 57.9 | 36.7 | 13.0 | 8.3 | 61.1 | 58.9 | 62.9 | 66.1 | 52.8 | 62.6 | 129.5 | 307.5 |
| 2003..... | 57.4 | 34.4 | 13.6 | 8.0 | 61.5 | 58.1 | 63.9 | 63.5 | 52.2 | 62.8 | 127.7 | 310.4 |
| 2002..... | 56.9 | 33.8 | 13.7 | 8.4 | 62.2 | 57.9 | 63.7 | 60.8 | 50.7 | 63.9 | 128.4 | 308.3 |
| 2001 5/..... | 56.2 | 35.1 | 14.3 | 8.5 | 60.6 | 58.1 | 63.4 | 58.9 | 49.9 | 62.7 | 128.6 | 310.1 |
| 2000..... | 53.8 | 34.5 | 14.6 | 9.1 | 60.0 | 54.1 | 59.2 | 54.8 | 48.6 | 60.9 | 122.6 | 310.5 |
| 1999..... | 54.2 | 33.9 | 15.3 | 9.4 | 59.7 | 55.2 | 58.7 | 53.2 | 48.2 | 64.4 | 128.6 | 318.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 80.3 | 40.4 | 14.7 | 10.1 | 92.8 | 90.8 | 90.8 | 95.7 | 76.8 | 89.9 | 180.9 | 407.1 |
| 2003..... | 80.2 | 39.2 | 15.9 | 10.1 | 93.6 | 89.9 | 93.4 | 92.7 | 76.9 | 89.8 | 179.3 | 417.7 |
| 2002..... | 79.8 | 37.6 | 16.3 | 10.4 | 95.1 | 89.9 | 93.3 | 89.2 | 75.3 | 92.9 | 180.3 | 414.8 |
| 2001 5/..... | 79.3 | 38.9 | 16.8 | 10.7 | 93.7 | 90.5 | 93.9 | 88.0 | 73.0 | 89.9 | 183.3 | 429.1 |
| 2000..... | 75.8 | 38.1 | 17.0 | 11.4 | 91.9 | 84.5 | 88.8 | 82.5 | 71.1 | 86.9 | 175.5 | 438.2 |
| 1999..... | 76.2 | 38.3 | 17.3 | 11.8 | 90.8 | 85.7 | 87.7 | 80.1 | 71.5 | 92.9 | 183.6 | 454.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 36.2 | 32.9 | 11.3 | 6.3 | 27.6 | 26.0 | 35.1 | 37.5 | 30.5 | 39.7 | 94.8 | 262.7 |
| 2003..... | 35.3 | 29.3 | 11.2 | 5.7 | 27.5 | 25.4 | 34.6 | 35.4 | 29.4 | 40.3 | 93.3 | 262.9 |
| 2002..... | 34.8 | 29.8 | 11.1 | 6.3 | 27.5 | 25.1 | 34.3 | 33.3 | 28.0 | 39.8 | 94.0 | 262.0 |
| 2001 5/..... | 34.0 | 31.2 | 11.8 | 6.2 | 25.8 | 25.1 | 33.1 | 30.8 | 28.5 | 40.1 | 92.7 | 260.3 |
| 2000..... | 32.5 | 30.8 | 12.1 | 6.7 | 26.5 | 23.3 | 29.9 | 28.1 | 27.9 | 39.4 | 88.1 | 258.5 |
| 1999..... | 33.0 | 29.2 | 13.3 | 6.9 | 27.0 | 24.3 | 30.1 | 27.3 | 26.8 | 41.0 | 92.8 | 263.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 39.8 | 27.7 | 11.6 | 6.6 | 60.1 | 48.7 | 45.9 | 46.6 | 43.5 | 49.2 | 90.0 | 182.7 |
| 2003..... | 41.0 | 23.1 | 12.7 | 7.2 | 60.4 | 51.0 | 48.9 | 48.6 | 43.1 | 52.1 | 87.0 | 173.1 |
| 2002..... | 40.5 | 23.4 | 14.1 | 6.9 | 61.0 | 48.0 | 48.7 | 46.9 | 42.0 | 52.7 | 94.9 | 170.7 |
| 2001 5/..... | 41.1 | 26.8 | 13.0 | 6.8 | 58.1 | 50.7 | 52.1 | 48.4 | 42.4 | 54.2 | 91.2 | 193.8 |
| 2000..... | 39.5 | 23.3 | 12.6 | 7.3 | 56.6 | 47.4 | 49.4 | 46.2 | 40.6 | 53.8 | 92.4 | 208.3 |
| 1999..... | 40.1 | 26.2 | 14.3 | 7.5 | 55.8 | 49.3 | 50.5 | 46.5 | 41.4 | 54.5 | 88.3 | 209.1 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 60.0 | 30.4 | 13.5 | 8.0 | 95.1 | 75.3 | 69.8 | 72.1 | 65.4 | 73.2 | 128.8 | 227.0 |
| 2003..... | 62.3 | 25.8 | 14.5 | 9.5 | 94.0 | 80.1 | 74.8 | 75.5 | 66.5 | 77.5 | 130.8 | 234.0 |
| 2002..... | 61.8 | 27.9 | 17.2 | 8.3 | 95.6 | 75.5 | 74.6 | 72.1 | 66.7 | 81.2 | 134.6 | 262.7 |
| 2001 5/..... | 62.2 | 26.9 | 15.6 | 8.2 | 90.6 | 78.9 | 81.7 | 74.5 | 65.2 | 79.5 | 135.9 | 264.7 |
| 2000..... | 60.2 | 24.6 | 13.8 | 8.5 | 89.0 | 75.3 | 77.5 | 72.9 | 63.6 | 81.5 | 132.3 | 298.3 |
| 1999..... | 61.5 | 29.0 | 15.8 | 9.4 | 87.0 | 77.8 | 80.1 | 74.4 | 65.9 | 84.6 | 133.0 | 312.7 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 18.3 | 24.8 | 9.7 | 5.3 | 19.9 | 16.7 | 19.4 | 20.9 | 23.5 | 29.9 | 63.2 | 159.9 |
| 2003..... | 18.2 | 20.3 | 10.7 | 4.8 | 21.5 | 16.3 | 20.4 | 21.8 | 21.9 | 31.9 | 56.9 | 142.2 |
| 2002..... | 17.9 | 18.6 | 10.8 | 5.4 | 20.7 | 15.3 | 20.5 | 21.8 | 19.9 | 30.0 | 67.7 | 125.2 |
| 2001 5/..... | 18.7 | 26.7 | 10.4 | 5.4 | 20.0 | 17.9 | 20.3 | 22.8 | 22.2 | 34.3 | 61.0 | 159.2 |
| 2000..... | 17.5 | 22.0 | 11.4 | 6.1 | 18.4 | 15.1 | 19.3 | 20.2 | 20.3 | 32.0 | 65.7 | 164.2 |
| 1999..... | 17.6 | 23.3 | 12.7 | 5.5 | 19.3 | 16.3 | 19.0 | 19.3 | 19.8 | 31.0 | 58.6 | 159.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XX. External causes of mortality (*U01-*U03,V01-Y89) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 60.6 | 39.0 | 13.4 | 8.6 | 61.1 | 61.0 | 65.4 | 67.8 | 53.4 | 63.4 | 131.4 | 312.7 |
| 2003..... | 59.7 | 37.3 | 13.8 | 8.1 | 61.4 | 59.4 | 65.9 | 64.7 | 52.7 | 63.3 | 129.5 | 315.5 |
| 2002..... | 59.1 | 36.3 | 13.6 | 8.7 | 62.2 | 59.7 | 65.5 | 61.8 | 51.1 | 64.3 | 129.6 | 313.0 |
| 2001 5/..... | 58.1 | 36.9 | 14.6 | 8.9 | 60.8 | 59.3 | 64.5 | 59.5 | 50.1 | 62.9 | 130.0 | 313.6 |
| 2000..... | 55.5 | 37.0 | 15.0 | 9.4 | 60.4 | 55.2 | 60.1 | 55.2 | 48.9 | 61.0 | 123.6 | 313.4 |
| 1999..... | 55.7 | 35.5 | 15.5 | 9.7 | 60.2 | 56.0 | 59.3 | 53.3 | 48.3 | 64.7 | 129.7 | 321.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 83.5 | 42.8 | 15.0 | 10.5 | 92.0 | 94.4 | 94.0 | 97.6 | 77.4 | 90.7 | 183.5 | 415.4 |
| 2003..... | 82.7 | 42.6 | 16.2 | 10.2 | 93.1 | 91.9 | 95.9 | 93.9 | 77.2 | 90.1 | 181.4 | 425.2 |
| 2002..... | 82.2 | 39.8 | 16.0 | 10.7 | 94.5 | 92.9 | 95.7 | 90.3 | 75.3 | 93.0 | 181.8 | 420.1 |
| 2001 5/..... | 81.4 | 41.5 | 17.0 | 11.2 | 93.9 | 92.5 | 94.9 | 88.6 | 73.0 | 90.1 | 185.0 | 434.9 |
| 2000..... | 77.7 | 41.2 | 17.7 | 12.0 | 92.1 | 85.9 | 89.7 | 82.7 | 71.0 | 86.7 | 176.9 | 442.7 |
| 1999..... | 77.8 | 40.5 | 17.5 | 12.2 | 91.2 | 86.7 | 88.1 | 79.8 | 71.2 | 92.9 | 184.9 | 458.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 38.8 | 35.0 | 11.6 | 6.5 | 29.0 | 27.8 | 37.3 | 39.1 | 31.1 | 40.4 | 96.3 | 266.8 |
| 2003..... | 37.7 | 31.7 | 11.3 | 5.9 | 28.5 | 27.1 | 36.4 | 36.6 | 29.9 | 40.8 | 94.9 | 267.3 |
| 2002..... | 37.1 | 32.7 | 11.1 | 6.5 | 28.7 | 26.9 | 36.0 | 34.3 | 28.6 | 40.4 | 95.0 | 266.8 |
| 2001 5/..... | 35.9 | 32.1 | 12.1 | 6.4 | 26.8 | 26.3 | 34.7 | 31.4 | 28.8 | 40.3 | 93.9 | 263.1 |
| 2000..... | 34.4 | 32.7 | 12.2 | 6.8 | 27.9 | 24.6 | 31.1 | 28.7 | 28.4 | 39.8 | 88.8 | 261.1 |
| 1999..... | 34.8 | 30.2 | 13.3 | 7.1 | 28.3 | 25.5 | 31.2 | 27.8 | 27.2 | 41.5 | 93.9 | 265.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 62.1 | 32.6 | 12.0 | 7.8 | 58.3 | 57.9 | 66.9 | 68.6 | 53.3 | 63.8 | 134.8 | 327.1 |
| 2003..... | 60.7 | 30.6 | 12.3 | 7.3 | 57.4 | 55.8 | 66.6 | 64.9 | 52.7 | 63.4 | 132.6 | 329.2 |
| 2002..... | 59.8 | 26.9 | 11.6 | 7.7 | 58.4 | 55.8 | 65.8 | 61.6 | 51.0 | 64.6 | 132.6 | 324.9 |
| 2001 5/..... | 58.4 | 28.9 | 13.1 | 8.2 | 56.3 | 55.7 | 64.1 | 58.6 | 49.6 | 62.7 | 132.6 | 323.5 |
| 2000..... | 55.2 | 28.4 | 13.0 | 8.9 | 55.6 | 50.6 | 58.9 | 53.7 | 48.3 | 60.3 | 125.5 | 323.5 |
| 1999..... | 55.2 | 27.9 | 13.1 | 8.9 | 54.8 | 51.6 | 57.9 | 51.6 | 47.0 | 64.0 | 130.6 | 329.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 83.6 | 37.3 | 13.7 | 9.6 | 85.1 | 86.7 | 94.9 | 97.3 | 75.9 | 90.9 | 187.7 | 432.8 |
| 2003..... | 82.1 | 34.9 | 14.5 | 9.2 | 84.1 | 83.7 | 95.5 | 92.8 | 76.4 | 90.3 | 185.1 | 442.8 |
| 2002..... | 81.2 | 31.3 | 13.5 | 9.5 | 86.0 | 84.1 | 94.9 | 88.9 | 74.2 | 92.5 | 184.9 | 433.3 |
| 2001 5/..... | 80.1 | 32.5 | 15.6 | 10.1 | 84.1 | 84.9 | 93.6 | 86.3 | 71.3 | 89.5 | 188.0 | 447.7 |
| 2000..... | 75.5 | 33.3 | 15.6 | 11.2 | 82.0 | 76.8 | 87.1 | 79.3 | 69.1 | 84.9 | 178.8 | 456.7 |
| 1999..... | 75.3 | 33.3 | 14.9 | 11.1 | 80.1 | 78.2 | 85.1 | 76.2 | 68.3 | 91.3 | 185.8 | 470.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 41.5 | 27.6 | 10.2 | 5.9 | 30.3 | 28.5 | 38.9 | 40.3 | 31.8 | 40.5 | 98.6 | 279.8 |
| 2003..... | 40.1 | 26.2 | 10.0 | 5.4 | 29.5 | 27.5 | 37.6 | 37.3 | 30.2 | 40.4 | 97.2 | 279.2 |
| 2002..... | 39.3 | 22.2 | 9.7 | 5.9 | 29.7 | 27.1 | 36.6 | 34.7 | 29.0 | 40.8 | 97.5 | 278.1 |
| 2001 5/..... | 37.6 | 25.2 | 10.4 | 6.1 | 27.4 | 26.0 | 34.5 | 31.5 | 29.1 | 39.9 | 95.9 | 271.8 |
| 2000..... | 35.8 | 23.2 | 10.2 | 6.4 | 28.2 | 23.9 | 30.7 | 28.5 | 28.7 | 39.6 | 90.4 | 269.7 |
| 1999..... | 36.0 | 22.3 | 11.3 | 6.6 | 28.6 | 24.7 | 30.7 | 27.5 | 26.8 | 41.1 | 94.5 | 273.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| XX. External causes of mortality (*U01-*U03,V01-Y89) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 64.0 | 68.0 | 19.7 | 12.0 | 80.2 | 90.8 | 72.3 | 77.5 | 63.6 | 65.7 | 111.2 | 179.0 |
| 2003..... | 64.5 | 65.6 | 20.8 | 11.3 | 84.2 | 90.7 | 76.2 | 76.5 | 60.8 | 67.8 | 103.9 | 185.9 |
| 2002..... | 65.4 | 74.5 | 21.6 | 12.5 | 85.0 | 92.2 | 77.7 | 75.6 | 59.5 | 66.1 | 107.6 | 201.4 |
| 2001 5/..... | 65.8 | 71.3 | 22.1 | 12.0 | 86.9 | 88.1 | 79.1 | 76.6 | 61.1 | 71.2 | 111.8 | 217.3 |
| 2000..... | 65.6 | 73.7 | 23.7 | 12.5 | 88.7 | 87.9 | 78.0 | 74.7 | 59.4 | 73.1 | 109.6 | 215.0 |
| 1999..... | 67.0 | 70.5 | 25.6 | 13.8 | 90.2 | 86.8 | 77.9 | 74.0 | 62.5 | 76.9 | 128.3 | 240.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 99.5 | 69.0 | 21.6 | 14.7 | 133.7 | 158.1 | 112.1 | 121.3 | 104.4 | 103.0 | 170.6 | 253.0 |
| 2003..... | 101.0 | 74.5 | 24.5 | 14.5 | 142.0 | 157.0 | 120.2 | 120.7 | 98.8 | 100.3 | 155.0 | 258.9 |
| 2002..... | 102.4 | 74.8 | 26.6 | 16.1 | 142.6 | 160.4 | 120.5 | 119.5 | 99.0 | 105.3 | 168.2 | 308.5 |
| 2001 5/..... | 102.7 | 80.4 | 25.1 | 16.4 | 148.4 | 151.1 | 121.4 | 124.3 | 99.4 | 107.8 | 171.0 | 307.2 |
| 2000..... | 102.8 | 75.8 | 26.3 | 16.1 | 149.5 | 151.2 | 123.8 | 121.9 | 98.1 | 115.7 | 169.1 | 320.6 |
| 1999..... | 104.5 | 74.8 | 28.7 | 17.9 | 152.5 | 147.8 | 123.2 | 122.5 | 102.4 | 120.7 | 190.4 | 370.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 31.7 | 67.0 | 17.8 | 9.3 | 25.8 | 29.2 | 37.1 | 39.9 | 30.8 | 39.1 | 77.8 | 149.8 |
| 2003..... | 31.4 | 56.4 | 16.9 | 8.0 | 25.8 | 30.3 | 37.3 | 38.5 | 30.2 | 44.7 | 75.1 | 157.6 |
| 2002..... | 31.8 | 74.1 | 16.3 | 8.7 | 27.2 | 30.6 | 39.9 | 37.9 | 27.9 | 38.3 | 73.3 | 160.5 |
| 2001 5/..... | 32.3 | 61.8 | 19.1 | 7.4 | 25.7 | 31.2 | 41.7 | 35.6 | 30.3 | 45.3 | 78.5 | 183.9 |
| 2000..... | 31.8 | 71.5 | 21.0 | 8.7 | 28.8 | 30.8 | 37.6 | 34.1 | 28.2 | 43.2 | 76.1 | 175.7 |
| 1999..... | 33.0 | 66.1 | 22.5 | 9.5 | 28.7 | 31.9 | 37.8 | 32.5 | 30.5 | 46.0 | 93.5 | 192.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidents (unintentional injuries) (V01-X59,Y85-Y86) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 38.1 | 25.8 | 10.3 | 6.5 | 37.0 | 32.6 | 37.3 | 40.7 | 33.2 | 44.0 | 103.7 | 276.7 |
| 2003..... | 37.6 | 23.6 | 10.9 | 6.4 | 37.1 | 31.5 | 37.8 | 38.8 | 32.9 | 44.1 | 101.9 | 278.9 |
| 2002..... | 37.0 | 23.5 | 10.5 | 6.6 | 38.0 | 31.5 | 37.2 | 36.6 | 31.4 | 44.2 | 101.3 | 275.4 |
| 2001..... | 35.7 | 24.2 | 11.2 | 6.9 | 36.1 | 29.9 | 35.4 | 34.1 | 30.3 | 42.8 | 100.9 | 276.4 |
| 2000..... | 34.8 | 23.1 | 11.9 | 7.3 | 36.0 | 29.5 | 34.1 | 32.6 | 30.9 | 41.9 | 95.1 | 273.5 |
| 1999..... | 35.1 | 22.3 | 12.4 | 7.6 | 35.3 | 29.6 | 33.8 | 31.8 | 30.6 | 44.6 | 100.5 | 282.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 49.8 | 29.0 | 11.8 | 8.0 | 53.3 | 48.5 | 53.0 | 58.6 | 46.9 | 59.2 | 134.7 | 345.5 |
| 2003..... | 49.3 | 26.4 | 13.0 | 7.9 | 53.3 | 47.1 | 54.2 | 55.9 | 47.1 | 58.9 | 132.9 | 352.5 |
| 2002..... | 48.9 | 26.8 | 12.9 | 8.2 | 54.9 | 47.7 | 53.7 | 53.3 | 45.2 | 60.6 | 131.6 | 346.6 |
| 2001..... | 47.2 | 26.2 | 13.3 | 8.6 | 52.9 | 45.4 | 51.8 | 50.6 | 43.0 | 57.3 | 133.4 | 358.5 |
| 2000..... | 46.2 | 25.7 | 14.0 | 8.8 | 52.1 | 45.0 | 51.0 | 49.0 | 44.1 | 55.9 | 124.3 | 359.7 |
| 1999..... | 46.4 | 25.4 | 14.4 | 9.4 | 50.7 | 45.0 | 50.2 | 47.9 | 44.0 | 60.2 | 132.4 | 375.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 26.8 | 22.5 | 8.7 | 5.1 | 19.9 | 16.0 | 21.7 | 23.4 | 20.5 | 31.1 | 82.9 | 245.8 |
| 2003..... | 26.2 | 20.7 | 8.7 | 4.8 | 19.9 | 15.3 | 21.4 | 22.4 | 19.7 | 31.6 | 81.1 | 246.4 |
| 2002..... | 25.6 | 20.0 | 8.1 | 5.0 | 20.1 | 14.9 | 20.8 | 20.5 | 18.6 | 30.6 | 81.2 | 244.6 |
| 2001..... | 24.5 | 22.1 | 8.9 | 5.2 | 18.4 | 14.0 | 19.2 | 18.0 | 18.5 | 30.8 | 79.6 | 242.0 |
| 2000..... | 23.8 | 20.5 | 9.7 | 5.6 | 19.1 | 13.7 | 17.5 | 16.8 | 18.8 | 30.3 | 76.1 | 238.4 |
| 1999..... | 24.1 | 19.0 | 10.3 | 5.6 | 19.2 | 14.0 | 17.6 | 16.3 | 18.3 | 31.8 | 79.7 | 244.8 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 25.2 | 17.5 | 9.0 | 5.1 | 33.4 | 27.4 | 29.6 | 31.0 | 30.5 | 37.2 | 74.2 | 160.8 |
| 2003..... | 26.1 | 15.1 | 9.9 | 5.8 | 33.8 | 29.4 | 31.7 | 32.8 | 30.6 | 38.1 | 70.9 | 151.8 |
| 2002..... | 26.1 | 15.2 | 10.9 | 5.4 | 35.5 | 27.6 | 31.7 | 32.0 | 28.7 | 39.5 | 77.4 | 153.0 |
| 2001..... | 25.8 | 18.8 | 10.3 | 5.4 | 32.6 | 28.2 | 32.3 | 31.7 | 30.5 | 40.0 | 73.8 | 166.3 |
| 2000..... | 25.0 | 15.3 | 10.4 | 5.6 | 31.8 | 27.3 | 31.5 | 30.4 | 28.8 | 38.8 | 72.1 | 184.5 |
| 1999..... | 25.5 | 17.7 | 11.9 | 5.6 | 30.8 | 28.4 | 32.6 | 31.2 | 28.5 | 42.5 | 69.6 | 182.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 36.5 | 18.8 | 10.5 | 6.2 | 50.8 | 41.6 | 44.9 | 47.9 | 44.8 | 52.0 | 99.4 | 193.3 |
| 2003..... | 38.3 | 17.4 | 11.9 | 7.5 | 50.7 | 45.5 | 48.0 | 50.6 | 45.3 | 53.3 | 102.4 | 195.5 |
| 2002..... | 38.5 | 19.5 | 13.5 | 6.6 | 53.5 | 43.0 | 48.5 | 48.7 | 43.5 | 58.7 | 105.9 | 224.2 |
| 2001..... | 37.6 | 18.0 | 12.7 | 6.2 | 49.2 | 43.8 | 50.1 | 48.8 | 45.0 | 54.9 | 105.1 | 198.1 |
| 2000..... | 36.9 | 17.0 | 11.6 | 6.1 | 48.2 | 42.9 | 49.6 | 48.4 | 43.2 | 57.2 | 93.9 | 251.9 |
| 1999..... | 37.7 | 20.7 | 13.5 | 6.8 | 46.2 | 44.1 | 51.3 | 50.2 | 43.5 | 63.3 | 94.9 | 258.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 13.1 | 16.2 | 7.5 | 4.0 | 13.5 | 10.2 | 12.6 | 14.0 | 17.5 | 25.4 | 56.8 | 144.0 |
| 2003..... | 13.1 | 12.7 | 7.8 | 3.9 | 14.3 | 10.1 | 13.8 | 15.1 | 17.3 | 26.0 | 49.2 | 129.7 |
| 2002..... | 12.8 | 10.8 | 8.2 | 4.1 | 14.5 | 9.5 | 13.4 | 15.3 | 15.3 | 24.2 | 57.9 | 117.7 |
| 2001..... | 13.2 | 19.7 | 7.8 | 4.5 | 13.3 | 10.1 | 13.0 | 14.9 | 17.7 | 28.2 | 52.6 | 150.8 |
| 2000..... | 12.4 | 13.5 | 9.2 | 5.2 | 12.5 | 9.2 | 12.2 | 12.8 | 16.0 | 24.4 | 57.4 | 151.3 |
| 1999..... | 12.6 | 14.6 | 10.3 | 4.2 | 12.8 | 10.1 | 12.6 | 12.8 | 15.2 | 26.2 | 52.7 | 145.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidents (unintentional injuries) (V01-X59,Y85-Y86) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 40.1 | 28.1 | 10.6 | 6.8 | 37.7 | 33.6 | 38.5 | 41.5 | 33.3 | 44.3 | 105.2 | 281.5 |
| 2003..... | 39.2 | 25.9 | 11.1 | 6.5 | 37.6 | 31.8 | 38.5 | 39.2 | 32.9 | 44.3 | 103.2 | 283.7 |
| 2002..... | 38.5 | 25.5 | 10.4 | 6.9 | 38.3 | 32.2 | 37.8 | 36.8 | 31.4 | 44.4 | 102.1 | 279.8 |
| 2001..... | 36.9 | 25.4 | 11.3 | 7.2 | 36.6 | 30.1 | 35.6 | 34.0 | 30.0 | 42.8 | 101.9 | 279.8 |
| 2000..... | 36.0 | 25.0 | 12.2 | 7.5 | 36.7 | 29.8 | 34.2 | 32.5 | 30.9 | 41.9 | 95.8 | 276.1 |
| 1999..... | 36.2 | 23.4 | 12.4 | 7.9 | 36.0 | 29.6 | 33.7 | 31.6 | 30.5 | 44.5 | 101.3 | 285.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 51.9 | 31.8 | 12.1 | 8.4 | 53.7 | 50.1 | 54.2 | 59.4 | 46.8 | 59.5 | 136.4 | 352.5 |
| 2003..... | 50.9 | 28.7 | 13.2 | 8.0 | 53.7 | 47.3 | 54.9 | 56.0 | 47.0 | 59.0 | 134.1 | 358.9 |
| 2002..... | 50.3 | 28.6 | 12.7 | 8.4 | 55.0 | 48.6 | 54.2 | 53.4 | 44.9 | 60.3 | 132.3 | 351.2 |
| 2001..... | 48.5 | 28.0 | 13.4 | 9.0 | 53.5 | 45.6 | 51.7 | 50.4 | 42.5 | 57.1 | 134.3 | 364.5 |
| 2000..... | 47.3 | 27.7 | 14.5 | 9.4 | 52.7 | 45.2 | 50.8 | 48.6 | 43.8 | 55.5 | 125.2 | 363.1 |
| 1999..... | 47.4 | 26.6 | 14.5 | 9.9 | 51.4 | 44.8 | 49.7 | 47.3 | 43.5 | 59.7 | 133.2 | 378.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 28.8 | 24.2 | 9.0 | 5.3 | 21.0 | 17.2 | 23.1 | 24.3 | 20.7 | 31.5 | 84.2 | 249.8 |
| 2003..... | 28.1 | 22.9 | 8.9 | 5.0 | 20.9 | 16.3 | 22.4 | 23.0 | 19.8 | 32.0 | 82.6 | 250.6 |
| 2002..... | 27.3 | 22.3 | 8.1 | 5.2 | 21.1 | 15.9 | 21.8 | 20.9 | 18.8 | 31.0 | 82.2 | 249.0 |
| 2001..... | 26.0 | 22.6 | 9.2 | 5.3 | 19.3 | 14.7 | 19.9 | 18.3 | 18.5 | 30.8 | 80.6 | 244.6 |
| 2000..... | 25.2 | 22.2 | 9.7 | 5.6 | 20.3 | 14.5 | 18.1 | 17.0 | 18.9 | 30.6 | 76.7 | 240.9 |
| 1999..... | 25.5 | 20.0 | 10.2 | 5.9 | 20.2 | 14.6 | 18.0 | 16.5 | 18.5 | 32.0 | 80.6 | 247.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 42.8 | 24.1 | 10.2 | 6.4 | 41.5 | 35.7 | 40.3 | 41.5 | 32.5 | 44.3 | 107.9 | 295.2 |
| 2003..... | 41.6 | 21.6 | 10.7 | 6.2 | 41.1 | 33.5 | 39.8 | 38.7 | 32.2 | 43.9 | 105.5 | 296.5 |
| 2002..... | 40.6 | 19.8 | 9.4 | 6.3 | 41.6 | 33.7 | 38.7 | 36.3 | 30.6 | 43.9 | 104.4 | 290.9 |
| 2001..... | 38.5 | 20.9 | 10.9 | 6.6 | 39.0 | 31.1 | 35.9 | 32.8 | 29.0 | 42.4 | 103.7 | 289.3 |
| 2000..... | 37.3 | 20.1 | 11.1 | 7.2 | 39.2 | 30.0 | 34.2 | 31.0 | 30.0 | 41.0 | 97.2 | 285.6 |
| 1999..... | 37.2 | 19.0 | 11.1 | 7.4 | 38.1 | 29.9 | 33.5 | 30.0 | 28.9 | 43.3 | 101.7 | 293.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 54.6 | 28.3 | 11.8 | 7.9 | 58.9 | 52.5 | 56.2 | 58.7 | 44.8 | 59.0 | 138.8 | 366.8 |
| 2003..... | 53.1 | 24.1 | 12.9 | 7.5 | 58.2 | 49.5 | 56.3 | 54.7 | 45.4 | 58.2 | 135.7 | 372.9 |
| 2002..... | 52.0 | 23.3 | 11.1 | 7.7 | 59.2 | 50.3 | 55.0 | 52.1 | 43.0 | 58.8 | 133.5 | 361.6 |
| 2001..... | 49.7 | 22.9 | 13.2 | 8.1 | 56.6 | 46.7 | 51.8 | 48.0 | 40.3 | 56.1 | 135.2 | 374.7 |
| 2000..... | 48.1 | 22.9 | 13.4 | 8.9 | 55.9 | 45.2 | 50.4 | 45.8 | 41.6 | 53.6 | 125.2 | 373.8 |
| 1999..... | 47.8 | 22.0 | 12.9 | 9.1 | 53.6 | 44.9 | 49.0 | 44.2 | 40.7 | 57.3 | 132.7 | 387.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 31.4 | 19.7 | 8.5 | 4.9 | 23.2 | 18.6 | 24.2 | 24.6 | 20.9 | 31.6 | 86.8 | 263.1 |
| 2003..... | 30.5 | 19.0 | 8.4 | 4.8 | 23.2 | 17.2 | 23.2 | 22.9 | 19.6 | 31.7 | 85.1 | 262.8 |
| 2002..... | 29.6 | 16.0 | 7.5 | 4.8 | 23.2 | 16.8 | 22.4 | 20.6 | 18.8 | 31.2 | 84.9 | 260.3 |
| 2001..... | 27.8 | 18.7 | 8.5 | 5.0 | 20.7 | 15.2 | 19.9 | 17.9 | 18.3 | 30.7 | 82.8 | 253.8 |
| 2000..... | 26.9 | 17.1 | 8.7 | 5.3 | 21.9 | 14.6 | 18.0 | 16.3 | 19.1 | 30.4 | 78.8 | 250.0 |
| 1999..... | 27.0 | 15.9 | 9.3 | 5.6 | 22.0 | 14.7 | 17.9 | 15.9 | 17.7 | 31.5 | 81.5 | 255.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidents (unintentional injuries) (V01-X59,Y85-Y86) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 33.7 | 47.0 | 12.8 | 8.8 | 26.3 | 32.5 | 37.8 | 50.7 | 44.5 | 48.5 | 88.2 | 155.1 |
| 2003..... | 33.3 | 44.7 | 13.7 | 8.0 | 26.4 | 30.9 | 39.7 | 49.3 | 42.9 | 51.1 | 83.1 | 161.8 |
| 2002..... | 34.0 | 50.1 | 14.6 | 9.4 | 28.4 | 32.5 | 39.9 | 47.6 | 41.9 | 49.7 | 84.6 | 175.4 |
| 2001..... | 34.4 | 44.5 | 14.2 | 9.7 | 29.6 | 31.1 | 40.9 | 48.5 | 42.4 | 51.3 | 89.6 | 183.9 |
| 2000..... | 34.4 | 46.5 | 16.8 | 9.6 | 29.6 | 33.5 | 40.6 | 47.8 | 41.3 | 53.0 | 85.3 | 183.2 |
| 1999..... | 36.1 | 44.1 | 18.2 | 10.9 | 30.5 | 33.2 | 40.7 | 48.3 | 45.1 | 58.6 | 103.8 | 208.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 47.9 | 49.2 | 14.4 | 10.6 | 38.5 | 52.0 | 55.6 | 77.9 | 71.9 | 74.1 | 133.2 | 218.5 |
| 2003..... | 47.4 | 49.0 | 15.9 | 10.1 | 39.8 | 48.4 | 59.4 | 75.8 | 69.0 | 74.9 | 123.8 | 228.0 |
| 2002..... | 49.0 | 52.1 | 18.9 | 11.9 | 42.6 | 52.2 | 59.7 | 73.3 | 69.2 | 77.2 | 131.3 | 262.5 |
| 2001..... | 49.5 | 50.4 | 15.8 | 13.1 | 44.5 | 48.8 | 61.3 | 78.6 | 68.2 | 74.9 | 138.3 | 256.4 |
| 2000..... | 50.3 | 49.5 | 19.4 | 12.3 | 44.2 | 53.5 | 63.3 | 76.7 | 69.4 | 80.4 | 133.3 | 277.1 |
| 1999..... | 52.3 | 48.5 | 21.8 | 14.3 | 46.4 | 53.0 | 63.0 | 79.4 | 71.8 | 89.6 | 151.0 | 318.5 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 20.8 | 44.8 | 11.1 | 6.8 | 13.9 | 14.7 | 21.9 | 27.5 | 22.4 | 30.3 | 62.8 | 130.1 |
| 2003..... | 20.5 | 40.3 | 11.4 | 5.9 | 12.9 | 14.9 | 22.4 | 26.6 | 22.0 | 34.1 | 60.1 | 136.2 |
| 2002..... | 20.3 | 48.0 | 10.1 | 6.8 | 14.2 | 14.7 | 22.5 | 25.6 | 20.0 | 30.2 | 58.2 | 142.1 |
| 2001..... | 20.6 | 38.3 | 12.4 | 6.2 | 14.8 | 15.1 | 22.7 | 22.7 | 21.6 | 34.6 | 62.2 | 157.0 |
| 2000..... | 20.1 | 43.3 | 14.2 | 6.7 | 15.1 | 15.5 | 20.5 | 23.0 | 18.7 | 33.7 | 58.4 | 148.3 |
| 1999..... | 21.4 | 39.6 | 14.5 | 7.4 | 14.8 | 15.4 | 21.1 | 21.8 | 23.7 | 36.8 | 77.4 | 167.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Transport accidents (V01-V99,Y85) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 16.4 | 3.6 | 4.1 | 4.3 | 27.3 | 18.8 | 16.4 | 16.7 | 15.5 | 17.5 | 25.5 | 27.2 |
| 2003..... | 16.5 | 3.7 | 4.1 | 4.3 | 27.7 | 18.3 | 17.1 | 16.6 | 15.9 | 17.5 | 26.2 | 29.7 |
| 2002..... | 16.8 | 3.1 | 4.1 | 4.2 | 29.1 | 18.7 | 17.1 | 16.4 | 15.7 | 18.3 | 27.0 | 28.8 |
| 2001..... | 16.6 | 3.7 | 4.4 | 4.4 | 27.9 | 18.8 | 17.4 | 16.0 | 15.3 | 17.9 | 27.9 | 28.9 |
| 2000..... | 16.6 | 4.6 | 4.5 | 4.7 | 28.1 | 18.6 | 17.0 | 15.8 | 16.0 | 18.0 | 27.0 | 31.5 |
| 1999..... | 16.6 | 5.0 | 4.6 | 4.8 | 27.6 | 18.5 | 16.8 | 15.4 | 16.0 | 19.4 | 28.4 | 32.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 23.0 | 3.9 | 4.4 | 5.0 | 37.8 | 27.7 | 24.0 | 24.7 | 22.2 | 23.7 | 36.5 | 46.5 |
| 2003..... | 23.3 | 4.0 | 4.5 | 5.2 | 38.6 | 27.3 | 24.8 | 24.3 | 23.0 | 23.7 | 36.6 | 52.0 |
| 2002..... | 23.7 | 3.5 | 4.8 | 4.9 | 40.7 | 28.1 | 24.6 | 24.0 | 22.5 | 25.3 | 37.7 | 53.0 |
| 2001..... | 23.5 | 3.3 | 5.0 | 5.3 | 39.6 | 28.3 | 25.1 | 23.3 | 21.6 | 24.3 | 39.8 | 52.7 |
| 2000..... | 23.3 | 4.8 | 5.0 | 5.4 | 39.3 | 27.7 | 24.8 | 23.0 | 22.5 | 24.3 | 38.1 | 59.8 |
| 1999..... | 23.1 | 5.2 | 5.1 | 5.8 | 38.1 | 27.2 | 24.0 | 22.4 | 22.5 | 26.1 | 40.7 | 63.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 9.9 | 3.2 | 3.8 | 3.6 | 16.1 | 9.5 | 8.8 | 9.0 | 9.3 | 12.3 | 18.1 | 18.5 |
| 2003..... | 10.0 | 3.4 | 3.7 | 3.4 | 16.3 | 9.0 | 9.4 | 9.1 | 9.4 | 12.4 | 19.3 | 19.9 |
| 2002..... | 10.1 | 2.8 | 3.4 | 3.5 | 16.9 | 9.2 | 9.7 | 9.0 | 9.3 | 12.5 | 19.8 | 18.3 |
| 2001..... | 10.0 | 4.2 | 3.8 | 3.5 | 15.6 | 9.2 | 9.8 | 8.9 | 9.4 | 12.7 | 20.1 | 18.9 |
| 2000..... | 10.2 | 4.3 | 4.0 | 3.9 | 16.4 | 9.3 | 9.3 | 8.9 | 10.0 | 12.8 | 19.8 | 19.9 |
| 1999..... | 10.4 | 4.8 | 4.2 | 3.9 | 16.6 | 9.6 | 9.6 | 8.7 | 10.1 | 14.0 | 20.4 | 19.3 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 14.5 | 4.4 | 4.7 | 3.9 | 26.3 | 18.1 | 14.9 | 14.3 | 15.9 | 17.0 | 21.9 | 20.0 |
| 2003..... | 15.2 | 4.5 | 4.2 | 4.1 | 26.2 | 19.1 | 16.4 | 15.1 | 15.8 | 18.6 | 23.6 | 24.9 |
| 2002..... | 15.3 | 3.1 | 5.8 | 3.8 | 27.6 | 18.3 | 16.0 | 15.2 | 15.2 | 18.4 | 26.2 | 25.5 |
| 2001..... | 15.6 | 5.3 | 5.3 | 3.8 | 26.0 | 18.8 | 17.3 | 16.4 | 17.9 | 20.2 | 30.1 | 23.8 |
| 2000..... | 14.8 | 5.4 | 5.2 | 4.1 | 24.6 | 17.3 | 16.2 | 16.2 | 15.7 | 20.4 | 23.9 | 25.2 |
| 1999..... | 14.4 | 5.3 | 5.5 | 3.8 | 23.1 | 17.8 | 15.4 | 15.0 | 15.5 | 22.0 | 25.0 | 25.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 21.3 | * | 5.4 | 4.6 | 39.3 | 27.0 | 22.2 | 21.6 | 21.6 | 23.9 | 34.7 | 28.0 |
| 2003..... | 22.2 | 5.7 | 4.6 | 5.5 | 38.2 | 29.3 | 23.9 | 22.4 | 21.4 | 26.2 | 35.3 | 46.2 |
| 2002..... | 22.4 | * | 7.1 | 4.6 | 40.4 | 27.5 | 23.9 | 21.6 | 22.1 | 28.5 | 37.3 | 45.2 |
| 2001..... | 22.8 | * | 6.0 | 4.3 | 38.4 | 28.8 | 25.9 | 23.2 | 24.6 | 28.2 | 51.0 | 43.8 |
| 2000..... | 21.6 | 6.1 | 5.6 | 4.5 | 36.1 | 26.4 | 24.1 | 24.1 | 23.5 | 29.5 | 36.4 | 44.3 |
| 1999..... | 20.8 | 5.7 | 5.8 | 4.5 | 33.3 | 26.7 | 23.2 | 22.6 | 21.0 | 31.6 | 40.1 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.3 | 5.0 | 4.1 | 3.2 | 11.3 | 7.5 | 6.7 | 6.9 | 10.7 | 11.5 | 13.1 | 15.8 |
| 2003..... | 7.6 | * | 3.8 | 2.7 | 12.3 | 7.0 | 8.3 | 7.9 | 10.7 | 12.5 | 15.5 | * |
| 2002..... | 7.7 | * | 4.5 | 2.9 | 12.6 | 7.3 | 7.5 | 8.7 | 8.9 | 10.3 | 18.5 | * |
| 2001..... | 8.0 | 6.7 | 4.6 | 3.2 | 11.6 | 7.1 | 8.0 | 9.7 | 12.0 | 13.9 | 16.0 | * |
| 2000..... | 7.6 | * | 4.8 | 3.7 | 11.0 | 6.8 | 7.8 | 8.5 | 8.7 | 13.3 | 15.5 | * |
| 1999..... | 7.6 | * | 5.2 | 3.1 | 11.0 | 7.6 | 7.0 | 7.7 | 10.6 | 14.5 | 14.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Transport accidents (V01-V99,Y85) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 16.6 | 3.3 | 3.9 | 4.4 | 27.4 | 18.9 | 16.6 | 16.9 | 15.4 | 17.5 | 25.6 | 27.5 |
| 2003..... | 16.7 | 3.5 | 4.0 | 4.3 | 28.0 | 18.0 | 17.2 | 16.6 | 15.8 | 17.3 | 26.3 | 29.8 |
| 2002..... | 16.9 | 3.2 | 3.7 | 4.3 | 29.3 | 18.8 | 17.2 | 16.4 | 15.6 | 18.3 | 26.9 | 28.9 |
| 2001..... | 16.7 | 3.4 | 4.2 | 4.6 | 28.2 | 18.8 | 17.3 | 15.8 | 15.0 | 17.7 | 27.7 | 29.0 |
| 2000..... | 16.8 | 4.3 | 4.3 | 4.8 | 28.8 | 18.7 | 17.0 | 15.7 | 15.9 | 17.8 | 27.1 | 31.7 |
| 1999..... | 16.9 | 4.9 | 4.4 | 5.0 | 28.4 | 18.5 | 16.9 | 15.4 | 16.0 | 19.2 | 28.3 | 32.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 23.3 | 3.9 | 4.1 | 5.0 | 37.5 | 27.8 | 24.3 | 24.9 | 22.2 | 23.6 | 36.5 | 47.3 |
| 2003..... | 23.4 | 3.5 | 4.5 | 5.1 | 38.5 | 26.7 | 24.9 | 24.4 | 23.0 | 23.3 | 36.6 | 52.0 |
| 2002..... | 23.8 | 3.4 | 4.2 | 5.0 | 40.6 | 28.1 | 24.6 | 24.1 | 22.4 | 24.9 | 37.5 | 53.2 |
| 2001..... | 23.5 | 3.2 | 4.7 | 5.5 | 39.7 | 28.1 | 24.8 | 23.2 | 21.3 | 24.0 | 39.2 | 52.9 |
| 2000..... | 23.4 | 4.5 | 4.8 | 5.6 | 39.8 | 27.8 | 24.7 | 22.8 | 22.2 | 23.9 | 38.0 | 60.4 |
| 1999..... | 23.3 | 5.1 | 4.9 | 6.0 | 39.0 | 27.2 | 24.0 | 22.2 | 22.4 | 25.6 | 40.4 | 64.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 10.3 | 2.6 | 3.8 | 3.7 | 17.0 | 9.9 | 9.1 | 9.2 | 9.1 | 12.3 | 18.4 | 18.6 |
| 2003..... | 10.3 | 3.5 | 3.6 | 3.6 | 17.0 | 9.4 | 9.6 | 9.2 | 9.2 | 12.3 | 19.5 | 20.1 |
| 2002..... | 10.4 | 2.9 | 3.1 | 3.6 | 17.7 | 9.5 | 10.0 | 9.0 | 9.3 | 12.7 | 19.8 | 18.4 |
| 2001..... | 10.2 | 3.6 | 3.6 | 3.6 | 16.3 | 9.5 | 10.0 | 8.8 | 9.1 | 12.5 | 20.2 | 19.1 |
| 2000..... | 10.5 | 4.2 | 3.8 | 3.9 | 17.4 | 9.7 | 9.4 | 8.9 | 10.1 | 12.8 | 20.0 | 20.1 |
| 1999..... | 10.7 | 4.7 | 3.9 | 4.0 | 17.5 | 10.0 | 9.9 | 8.8 | 10.0 | 13.9 | 20.5 | 19.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 17.2 | 2.9 | 3.7 | 4.4 | 29.9 | 19.1 | 17.0 | 16.8 | 15.4 | 17.6 | 26.0 | 28.8 |
| 2003..... | 17.2 | 3.2 | 3.8 | 4.4 | 30.2 | 18.2 | 17.3 | 16.6 | 15.6 | 17.3 | 26.6 | 30.6 |
| 2002..... | 17.5 | 2.5 | 3.3 | 4.3 | 31.8 | 19.0 | 17.3 | 16.4 | 15.5 | 18.3 | 27.4 | 29.7 |
| 2001..... | 17.0 | 3.1 | 3.8 | 4.4 | 29.9 | 19.0 | 17.4 | 15.6 | 14.7 | 17.6 | 28.3 | 29.9 |
| 2000..... | 17.1 | 3.5 | 3.8 | 4.8 | 30.9 | 18.5 | 17.0 | 15.4 | 15.8 | 17.6 | 27.6 | 32.4 |
| 1999..... | 17.1 | 4.3 | 3.7 | 5.0 | 30.3 | 18.6 | 16.7 | 15.0 | 15.6 | 19.1 | 28.6 | 32.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 23.9 | 3.3 | 3.8 | 5.2 | 40.6 | 27.6 | 24.6 | 24.5 | 21.8 | 23.5 | 36.6 | 49.6 |
| 2003..... | 23.9 | 3.3 | 4.4 | 5.1 | 41.1 | 26.8 | 24.9 | 24.1 | 22.6 | 23.3 | 36.6 | 52.8 |
| 2002..... | 24.3 | 2.9 | 3.6 | 5.0 | 43.6 | 28.1 | 24.6 | 24.0 | 22.0 | 24.5 | 37.5 | 54.0 |
| 2001..... | 23.8 | 2.9 | 4.4 | 5.2 | 41.9 | 28.2 | 24.6 | 22.7 | 20.8 | 23.7 | 39.6 | 54.3 |
| 2000..... | 23.7 | 3.8 | 4.3 | 5.7 | 42.5 | 27.1 | 24.5 | 22.2 | 21.6 | 23.2 | 38.1 | 62.3 |
| 1999..... | 23.5 | 4.9 | 4.0 | 5.9 | 41.2 | 26.9 | 23.6 | 21.6 | 21.7 | 25.2 | 40.3 | 64.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 10.7 | 2.4 | 3.6 | 3.6 | 18.6 | 10.4 | 9.3 | 9.2 | 9.3 | 12.5 | 18.8 | 19.5 |
| 2003..... | 10.7 | 3.0 | 3.2 | 3.7 | 18.8 | 9.5 | 9.7 | 9.2 | 8.9 | 12.1 | 19.8 | 20.8 |
| 2002..... | 10.9 | 2.1 | 2.9 | 3.5 | 19.4 | 9.8 | 10.1 | 8.9 | 9.4 | 13.0 | 20.7 | 19.2 |
| 2001..... | 10.6 | 3.3 | 3.2 | 3.5 | 17.4 | 9.7 | 10.2 | 8.6 | 9.1 | 12.5 | 20.8 | 19.8 |
| 2000..... | 10.9 | 3.2 | 3.4 | 4.0 | 18.9 | 9.7 | 9.5 | 8.6 | 10.4 | 12.8 | 20.7 | 20.4 |
| 1999..... | 11.0 | 3.6 | 3.4 | 4.0 | 19.1 | 10.2 | 9.9 | 8.6 | 9.9 | 13.9 | 20.9 | 19.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Transport accidents (V01-V99,Y85) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 15.6 | 5.0 | 5.2 | 4.5 | 20.0 | 21.6 | 17.4 | 19.9 | 17.6 | 16.4 | 21.7 | 15.0 |
| 2003..... | 15.6 | 4.4 | 5.1 | 4.3 | 20.3 | 20.0 | 18.8 | 18.6 | 18.7 | 18.2 | 22.6 | 21.4 |
| 2002..... | 15.7 | 5.6 | 5.0 | 4.7 | 21.6 | 21.1 | 18.5 | 18.0 | 17.3 | 17.3 | 21.8 | 18.8 |
| 2001..... | 16.3 | * | 5.4 | 5.5 | 22.8 | 20.5 | 19.2 | 19.1 | 18.4 | 18.8 | 21.4 | 17.8 |
| 2000..... | 16.4 | 6.9 | 5.9 | 4.8 | 22.4 | 22.2 | 19.1 | 19.7 | 17.5 | 19.9 | 19.9 | 21.9 |
| 1999..... | 16.7 | 7.7 | 7.1 | 5.6 | 22.7 | 20.6 | 19.5 | 19.2 | 18.5 | 19.5 | 26.0 | 27.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 23.4 | 6.2 | 5.4 | 4.9 | 28.6 | 35.1 | 27.3 | 31.5 | 28.2 | 26.4 | 38.5 | 29.5 |
| 2003..... | 23.3 | * | 5.1 | 5.3 | 30.3 | 31.8 | 29.1 | 29.7 | 29.6 | 26.5 | 38.1 | 43.7 |
| 2002..... | 23.6 | * | 6.5 | 5.4 | 31.0 | 34.1 | 28.0 | 28.0 | 28.6 | 28.7 | 40.7 | 47.0 |
| 2001..... | 24.5 | * | 5.8 | 7.1 | 33.4 | 32.6 | 29.7 | 30.7 | 30.2 | 27.4 | 36.9 | 35.8 |
| 2000..... | 24.8 | 7.2 | 6.4 | 5.8 | 32.5 | 35.6 | 30.5 | 30.1 | 29.6 | 32.6 | 36.4 | 45.8 |
| 1999..... | 24.9 | * | 8.0 | 7.0 | 32.9 | 33.0 | 29.8 | 30.7 | 29.3 | 29.5 | 43.5 | 68.9 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 8.5 | * | 5.0 | 4.0 | 11.3 | 9.3 | 8.7 | 9.9 | 9.2 | 9.2 | 12.2 | 9.2 |
| 2003..... | 8.5 | * | 5.2 | 3.3 | 10.1 | 9.3 | 9.7 | 9.0 | 9.9 | 12.3 | 13.9 | 12.8 |
| 2002..... | 8.5 | * | 3.6 | 3.9 | 12.1 | 9.3 | 10.1 | 9.4 | 8.2 | 9.1 | 11.1 | * |
| 2001..... | 8.9 | * | 5.0 | 3.8 | 12.3 | 9.6 | 9.9 | 9.2 | 9.0 | 12.8 | 12.6 | 11.1 |
| 2000..... | 8.8 | * | 5.4 | 3.7 | 12.5 | 10.1 | 8.9 | 10.8 | 7.8 | 11.0 | 10.6 | 13.1 |
| 1999..... | 9.4 | 9.3 | 6.2 | 4.1 | 12.6 | 9.4 | 10.4 | 9.5 | 9.8 | 12.5 | 16.3 | 12.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|---|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.3 | 3.5 | 4.0 | 4.1 | 26.3 | 17.6 | 15.1 | 15.1 | 14.1 | 16.1 | 24.5 | 26.1 |
| 2003..... | 15.4 | 3.6 | 3.9 | 4.0 | 26.6 | 17.1 | 15.7 | 14.9 | 14.2 | 16.2 | 24.9 | 28.8 |
| 2002..... | 15.7 | 3.0 | 3.9 | 3.9 | 28.2 | 17.8 | 15.8 | 14.8 | 14.1 | 17.0 | 25.7 | 28.0 |
| 2001..... | 15.4 | 3.6 | 4.1 | 4.1 | 26.8 | 17.5 | 15.7 | 14.2 | 13.7 | 16.3 | 26.3 | 27.8 |
| 2000..... | 15.4 | 4.4 | 4.2 | 4.3 | 26.9 | 17.3 | 15.3 | 14.2 | 14.4 | 16.5 | 25.7 | 30.4 |
| 1999..... | 15.2 | 4.8 | 4.2 | 4.3 | 26.2 | 16.9 | 14.9 | 13.6 | 14.2 | 17.8 | 26.5 | 30.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 21.3 | 3.8 | 4.2 | 4.7 | 36.4 | 25.8 | 21.9 | 21.9 | 19.8 | 21.3 | 34.7 | 44.2 |
| 2003..... | 21.4 | 3.9 | 4.4 | 4.8 | 36.9 | 25.5 | 22.5 | 21.5 | 20.0 | 21.3 | 34.3 | 50.0 |
| 2002..... | 21.9 | 3.3 | 4.5 | 4.6 | 39.3 | 26.5 | 22.3 | 21.3 | 19.9 | 22.7 | 35.3 | 51.7 |
| 2001..... | 21.5 | 3.2 | 4.5 | 4.9 | 38.0 | 26.2 | 22.4 | 20.5 | 19.0 | 21.5 | 37.3 | 51.1 |
| 2000..... | 21.3 | 4.6 | 4.7 | 5.0 | 37.4 | 25.5 | 22.0 | 20.4 | 19.8 | 21.7 | 35.6 | 57.5 |
| 1999..... | 20.9 | 5.0 | 4.7 | 5.1 | 36.0 | 24.7 | 21.1 | 19.4 | 19.3 | 23.3 | 38.0 | 59.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 9.5 | 3.2 | 3.7 | 3.4 | 15.7 | 9.1 | 8.3 | 8.5 | 8.8 | 11.7 | 17.7 | 17.9 |
| 2003..... | 9.5 | 3.3 | 3.5 | 3.2 | 15.8 | 8.5 | 8.9 | 8.5 | 8.8 | 11.9 | 18.7 | 19.5 |
| 2002..... | 9.8 | 2.8 | 3.3 | 3.3 | 16.6 | 8.8 | 9.3 | 8.6 | 8.9 | 12.3 | 19.3 | 17.7 |
| 2001..... | 9.5 | 4.0 | 3.6 | 3.3 | 15.1 | 8.7 | 9.1 | 8.1 | 8.8 | 12.0 | 19.2 | 18.0 |
| 2000..... | 9.7 | 4.2 | 3.8 | 3.6 | 15.9 | 8.8 | 8.8 | 8.2 | 9.5 | 12.3 | 19.2 | 19.3 |
| 1999..... | 9.7 | 4.7 | 3.8 | 3.6 | 15.9 | 8.9 | 8.8 | 8.0 | 9.5 | 13.3 | 19.1 | 18.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 13.9 | 4.4 | 4.6 | 3.8 | 25.6 | 17.2 | 13.9 | 13.2 | 14.9 | 16.1 | 21.3 | 19.5 |
| 2003..... | 14.4 | 4.5 | 4.1 | 3.9 | 25.5 | 18.1 | 15.4 | 14.2 | 14.6 | 17.4 | 23.1 | 24.4 |
| 2002..... | 14.6 | 3.1 | 5.6 | 3.6 | 26.6 | 17.6 | 15.2 | 13.9 | 14.3 | 17.9 | 24.4 | 24.9 |
| 2001..... | 14.3 | 4.9 | 4.7 | 3.5 | 25.1 | 17.4 | 15.3 | 13.8 | 15.1 | 17.7 | 26.3 | 20.0 |
| 2000..... | 13.9 | 5.2 | 4.9 | 3.8 | 23.7 | 16.4 | 14.7 | 15.1 | 14.4 | 18.4 | 22.3 | 22.6 |
| 1999..... | 13.3 | 5.3 | 5.2 | 3.6 | 21.9 | 16.4 | 13.9 | 13.7 | 14.5 | 19.7 | 23.5 | 23.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 20.2 | * | 5.3 | 4.4 | 38.3 | 25.4 | 20.8 | 19.6 | 19.7 | 22.3 | 33.3 | 28.0 |
| 2003..... | 21.1 | 5.7 | 4.3 | 5.2 | 37.2 | 27.6 | 22.1 | 20.8 | 19.8 | 24.7 | 34.1 | 46.2 |
| 2002..... | 21.3 | * | 6.8 | 4.4 | 38.9 | 26.4 | 22.6 | 19.6 | 20.5 | 27.5 | 34.5 | 43.5 |
| 2001..... | 21.0 | * | 5.3 | 4.0 | 36.9 | 26.8 | 23.1 | 20.2 | 21.2 | 24.6 | 44.3 | * |
| 2000..... | 20.1 | 6.1 | 5.2 | 4.1 | 34.7 | 24.9 | 21.6 | 22.0 | 21.2 | 26.0 | 33.4 | * |
| 1999..... | 19.1 | 5.7 | 5.6 | 4.1 | 31.6 | 24.4 | 20.8 | 20.1 | 19.0 | 27.2 | 37.0 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 7.1 | 5.0 | 3.9 | 3.2 | 11.1 | 7.3 | 6.4 | 6.7 | 10.5 | 11.2 | 13.1 | 15.1 |
| 2003..... | 7.3 | * | 3.8 | 2.7 | 11.9 | 6.6 | 8.0 | 7.5 | 9.8 | 11.6 | 15.5 | * |
| 2002..... | 7.4 | * | 4.4 | 2.7 | 12.4 | 7.2 | 7.3 | 8.2 | 8.7 | 10.2 | 17.4 | * |
| 2001..... | 7.2 | 6.2 | 4.1 | 3.0 | 11.2 | 6.5 | 6.9 | 7.6 | 9.7 | 12.3 | 14.2 | * |
| 2000..... | 7.2 | * | 4.6 | 3.6 | 10.6 | 6.5 | 7.3 | 8.3 | 8.4 | 12.4 | 14.9 | * |
| 1999..... | 7.2 | * | 4.7 | 3.0 | 10.6 | 7.1 | 6.5 | 7.4 | 10.6 | 13.8 | 14.6 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|---|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 15.5 | 3.2 | 3.8 | 4.1 | 26.4 | 17.6 | 15.2 | 15.2 | 14.0 | 16.1 | 24.6 | 26.3 |
| 2003..... | 15.5 | 3.4 | 3.9 | 4.0 | 26.8 | 16.9 | 15.7 | 14.9 | 14.1 | 16.0 | 25.0 | 28.9 |
| 2002..... | 15.9 | 3.0 | 3.5 | 4.0 | 28.5 | 17.7 | 15.8 | 14.8 | 14.1 | 16.9 | 25.7 | 28.1 |
| 2001..... | 15.5 | 3.2 | 3.9 | 4.2 | 27.1 | 17.5 | 15.7 | 14.2 | 13.5 | 16.2 | 26.3 | 28.0 |
| 2000..... | 15.6 | 4.2 | 4.1 | 4.4 | 27.5 | 17.4 | 15.4 | 14.1 | 14.4 | 16.3 | 25.8 | 30.6 |
| 1999..... | 15.4 | 4.7 | 4.0 | 4.5 | 26.9 | 16.9 | 15.0 | 13.5 | 14.0 | 17.6 | 26.5 | 30.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 21.5 | 3.8 | 3.9 | 4.7 | 36.0 | 25.8 | 22.0 | 22.1 | 19.7 | 21.2 | 34.6 | 45.0 |
| 2003..... | 21.4 | 3.4 | 4.3 | 4.7 | 36.7 | 24.9 | 22.5 | 21.4 | 19.9 | 20.9 | 34.2 | 49.9 |
| 2002..... | 21.9 | 3.2 | 4.0 | 4.6 | 39.2 | 26.4 | 22.2 | 21.3 | 19.7 | 22.3 | 35.1 | 51.9 |
| 2001..... | 21.5 | 3.1 | 4.3 | 5.1 | 38.1 | 25.9 | 22.2 | 20.4 | 18.8 | 21.3 | 36.9 | 51.6 |
| 2000..... | 21.4 | 4.2 | 4.5 | 5.1 | 37.9 | 25.5 | 22.0 | 20.2 | 19.6 | 21.3 | 35.7 | 58.3 |
| 1999..... | 21.0 | 4.8 | 4.4 | 5.2 | 36.7 | 24.6 | 21.1 | 19.2 | 19.1 | 22.9 | 37.7 | 60.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 9.8 | 2.6 | 3.7 | 3.4 | 16.5 | 9.4 | 8.6 | 8.6 | 8.6 | 11.7 | 17.9 | 18.0 |
| 2003..... | 9.8 | 3.3 | 3.4 | 3.3 | 16.5 | 8.9 | 9.0 | 8.6 | 8.7 | 11.9 | 18.8 | 19.6 |
| 2002..... | 10.1 | 2.9 | 3.0 | 3.4 | 17.4 | 9.1 | 9.5 | 8.6 | 8.9 | 12.4 | 19.4 | 17.8 |
| 2001..... | 9.8 | 3.4 | 3.5 | 3.3 | 15.8 | 9.1 | 9.4 | 8.2 | 8.7 | 12.0 | 19.3 | 18.2 |
| 2000..... | 10.0 | 4.2 | 3.6 | 3.6 | 16.9 | 9.2 | 8.9 | 8.2 | 9.6 | 12.2 | 19.3 | 19.4 |
| 1999..... | 10.0 | 4.7 | 3.6 | 3.7 | 16.8 | 9.2 | 9.1 | 8.1 | 9.3 | 13.2 | 19.2 | 18.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 16.0 | 2.8 | 3.5 | 4.1 | 28.8 | 17.7 | 15.5 | 15.1 | 13.9 | 16.1 | 25.0 | 27.6 |
| 2003..... | 15.9 | 3.1 | 3.6 | 4.0 | 28.9 | 17.0 | 15.7 | 14.8 | 13.7 | 15.9 | 25.2 | 29.6 |
| 2002..... | 16.3 | 2.4 | 3.0 | 4.0 | 30.8 | 17.8 | 15.8 | 14.8 | 14.0 | 16.9 | 26.2 | 28.9 |
| 2001..... | 15.8 | 3.0 | 3.5 | 4.1 | 28.7 | 17.6 | 15.7 | 13.9 | 13.3 | 16.1 | 26.9 | 29.0 |
| 2000..... | 15.8 | 3.3 | 3.7 | 4.4 | 29.5 | 17.0 | 15.3 | 13.7 | 14.2 | 16.1 | 26.2 | 31.3 |
| 1999..... | 15.6 | 4.1 | 3.4 | 4.4 | 28.7 | 16.8 | 14.8 | 13.1 | 13.7 | 17.5 | 26.8 | 30.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 22.0 | 3.3 | 3.6 | 4.8 | 39.1 | 25.5 | 22.2 | 21.7 | 19.3 | 21.0 | 34.8 | 47.1 |
| 2003..... | 21.8 | 3.2 | 4.2 | 4.7 | 39.1 | 24.9 | 22.4 | 21.1 | 19.4 | 20.9 | 34.3 | 50.7 |
| 2002..... | 22.3 | 2.7 | 3.3 | 4.6 | 42.1 | 26.1 | 22.0 | 21.2 | 19.4 | 21.8 | 35.2 | 52.6 |
| 2001..... | 21.7 | 2.9 | 3.9 | 4.8 | 40.1 | 25.9 | 21.9 | 19.9 | 18.2 | 21.0 | 37.3 | 53.1 |
| 2000..... | 21.5 | 3.4 | 4.1 | 5.1 | 40.3 | 24.7 | 21.6 | 19.6 | 18.9 | 20.7 | 35.6 | 60.0 |
| 1999..... | 21.1 | 4.7 | 3.6 | 5.1 | 38.8 | 24.2 | 20.6 | 18.5 | 18.5 | 22.6 | 37.6 | 61.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 10.2 | 2.3 | 3.5 | 3.3 | 18.0 | 9.8 | 8.7 | 8.6 | 8.7 | 11.9 | 18.4 | 18.9 |
| 2003..... | 10.2 | 2.9 | 3.0 | 3.3 | 18.3 | 9.0 | 9.1 | 8.5 | 8.4 | 11.6 | 19.1 | 20.3 |
| 2002..... | 10.5 | 2.1 | 2.8 | 3.3 | 19.0 | 9.4 | 9.7 | 8.4 | 8.8 | 12.7 | 20.2 | 18.6 |
| 2001..... | 10.1 | 3.1 | 3.1 | 3.3 | 17.0 | 9.3 | 9.6 | 8.1 | 8.7 | 11.9 | 20.0 | 19.0 |
| 2000..... | 10.3 | 3.2 | 3.2 | 3.6 | 18.4 | 9.3 | 9.0 | 7.9 | 9.8 | 12.3 | 20.0 | 19.8 |
| 1999..... | 10.3 | 3.5 | 3.1 | 3.7 | 18.3 | 9.4 | 9.0 | 7.9 | 9.2 | 13.2 | 19.7 | 18.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|---|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 14.6 | 4.9 | 5.0 | 4.3 | 19.1 | 20.5 | 16.3 | 18.1 | 16.1 | 15.4 | 20.4 | 15.0 |
| 2003..... | 14.7 | 4.4 | 5.0 | 4.1 | 19.4 | 19.1 | 17.6 | 16.9 | 17.3 | 17.1 | 21.1 | 21.1 |
| 2002..... | 14.8 | 5.4 | 4.9 | 4.3 | 21.0 | 20.3 | 17.2 | 16.3 | 15.7 | 16.0 | 20.2 | 18.5 |
| 2001..... | 15.2 | * | 5.3 | 5.0 | 22.2 | 19.6 | 17.6 | 17.2 | 16.6 | 17.4 | 19.9 | 16.9 |
| 2000..... | 15.5 | 6.9 | 5.6 | 4.5 | 21.6 | 21.1 | 17.6 | 18.3 | 16.3 | 18.4 | 18.8 | 20.7 |
| 1999..... | 15.5 | 7.6 | 6.5 | 5.1 | 21.7 | 19.2 | 17.9 | 17.4 | 16.7 | 18.1 | 24.0 | 25.5 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 21.7 | * | 5.2 | 4.7 | 27.0 | 33.0 | 25.2 | 28.5 | 25.3 | 24.7 | 36.2 | 29.5 |
| 2003..... | 21.7 | * | 5.0 | 4.9 | 28.8 | 30.2 | 27.1 | 26.5 | 26.7 | 24.1 | 35.5 | 42.6 |
| 2002..... | 22.0 | * | 6.3 | 5.0 | 30.1 | 32.7 | 25.7 | 24.8 | 25.4 | 25.6 | 36.6 | 45.9 |
| 2001..... | 22.7 | * | 5.6 | 6.5 | 32.4 | 31.0 | 26.9 | 27.5 | 26.8 | 24.2 | 34.8 | 35.8 |
| 2000..... | 23.1 | 7.2 | 6.0 | 5.4 | 31.1 | 33.6 | 27.9 | 27.6 | 27.2 | 29.8 | 34.6 | 44.6 |
| 1999..... | 22.8 | * | 7.0 | 6.2 | 31.5 | 30.7 | 26.9 | 27.5 | 25.8 | 26.8 | 41.3 | 61.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 8.1 | * | 4.9 | 3.8 | 11.1 | 9.1 | 8.3 | 9.2 | 8.6 | 8.8 | 11.4 | 9.2 |
| 2003..... | 8.3 | * | 4.9 | 3.2 | 9.9 | 9.0 | 9.3 | 8.6 | 9.7 | 12.1 | 13.0 | 12.8 |
| 2002..... | 8.3 | * | 3.5 | 3.6 | 11.9 | 9.1 | 9.7 | 9.0 | 8.0 | 9.1 | 10.9 | * |
| 2001..... | 8.5 | * | 4.9 | 3.4 | 12.1 | 9.2 | 9.4 | 8.4 | 8.4 | 12.6 | 11.5 | 9.8 |
| 2000..... | 8.5 | * | 5.2 | 3.6 | 12.3 | 9.8 | 8.5 | 10.2 | 7.5 | 10.4 | 9.9 | 11.8 |
| 1999..... | 8.9 | 9.3 | 5.9 | 4.0 | 12.1 | 8.9 | 9.9 | 8.8 | 9.3 | 12.0 | 14.4 | 12.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pedestrian involved in collision with motor vehicle (V02-V04) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | 1.0 | 0.5 | 0.9 | 0.9 | 1.2 | 1.4 | 1.4 | 1.5 | 2.5 | 2.7 |
| 2003..... | 1.2 | * | 1.1 | 0.5 | 1.1 | 1.0 | 1.3 | 1.4 | 1.5 | 1.6 | 2.6 | 2.9 |
| 2002..... | 1.3 | * | 1.0 | 0.6 | 1.1 | 1.0 | 1.4 | 1.4 | 1.4 | 2.0 | 2.6 | 3.3 |
| 2001..... | 1.2 | * | 1.1 | 0.7 | 1.0 | 1.0 | 1.4 | 1.4 | 1.4 | 1.6 | 3.0 | 3.0 |
| 2000..... | 1.2 | * | 1.2 | 0.6 | 0.9 | 1.0 | 1.3 | 1.3 | 1.5 | 1.6 | 2.8 | 3.6 |
| 1999..... | 1.2 | * | 1.1 | 0.7 | 0.9 | 1.0 | 1.3 | 1.3 | 1.5 | 1.7 | 2.5 | 3.7 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | 1.2 | 0.6 | 1.3 | 1.4 | 1.7 | 2.1 | 2.1 | 2.0 | 3.4 | 4.9 |
| 2003..... | 1.7 | * | 1.4 | 0.7 | 1.5 | 1.5 | 1.9 | 2.1 | 2.2 | 2.3 | 3.3 | 5.7 |
| 2002..... | 1.7 | * | 1.3 | 0.7 | 1.6 | 1.6 | 1.9 | 2.1 | 2.0 | 2.7 | 3.4 | 5.8 |
| 2001..... | 1.8 | * | 1.4 | 0.9 | 1.5 | 1.5 | 2.0 | 2.0 | 2.0 | 2.1 | 4.1 | 5.9 |
| 2000..... | 1.7 | * | 1.4 | 0.8 | 1.3 | 1.5 | 1.9 | 2.0 | 2.2 | 2.2 | 3.8 | 7.0 |
| 1999..... | 1.7 | * | 1.2 | 0.8 | 1.4 | 1.5 | 1.9 | 2.0 | 2.2 | 2.3 | 3.7 | 6.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | 0.8 | 0.4 | 0.4 | 0.5 | 0.6 | 0.7 | 0.9 | 1.2 | 1.8 | 1.7 |
| 2003..... | 0.7 | * | 0.8 | 0.4 | 0.6 | 0.5 | 0.7 | 0.8 | 0.8 | 1.0 | 2.2 | 1.7 |
| 2002..... | 0.8 | * | 0.6 | 0.5 | 0.5 | 0.5 | 0.8 | 0.7 | 0.9 | 1.4 | 2.0 | 2.2 |
| 2001..... | 0.8 | * | 0.8 | 0.4 | 0.5 | 0.4 | 0.8 | 0.8 | 0.7 | 1.1 | 2.2 | 1.8 |
| 2000..... | 0.8 | * | 0.9 | 0.5 | 0.5 | 0.4 | 0.7 | 0.7 | 0.8 | 1.1 | 2.1 | 2.3 |
| 1999..... | 0.7 | * | 1.0 | 0.5 | 0.5 | 0.5 | 0.6 | 0.6 | 0.9 | 1.2 | 1.7 | 2.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | 1.6 | 0.6 | 1.2 | 1.5 | 1.6 | 2.1 | 2.9 | 3.1 | 4.5 | * |
| 2003..... | 1.7 | * | 1.3 | 0.4 | 1.4 | 1.6 | 2.5 | 2.1 | 3.4 | 3.8 | 5.9 | * |
| 2002..... | 1.7 | * | 1.8 | 0.7 | 1.3 | 1.5 | 1.9 | 2.1 | 2.8 | 5.0 | 5.5 | * |
| 2001..... | 1.9 | * | 1.8 | 0.6 | 1.3 | 1.8 | 2.5 | 2.4 | 3.1 | 4.0 | 7.9 | * |
| 2000..... | 1.6 | * | 1.5 | 0.6 | 1.1 | 1.4 | 1.9 | 2.1 | 2.7 | 4.3 | 6.7 | * |
| 1999..... | 1.5 | * | 1.9 | 0.5 | 1.2 | 1.5 | 1.5 | 2.2 | 3.0 | 3.7 | 5.6 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.2 | * | 2.0 | 0.7 | 2.0 | 2.3 | 2.5 | 3.3 | 4.2 | 4.6 | 7.5 | * |
| 2003..... | 2.6 | * | 1.6 | 0.6 | 2.2 | 2.5 | 4.0 | 3.3 | 4.9 | 5.7 | * | * |
| 2002..... | 2.6 | * | 2.4 | 0.9 | 2.1 | 2.2 | 3.0 | 3.7 | 4.4 | 8.6 | 9.8 | * |
| 2001..... | 2.8 | * | 2.2 | 0.7 | 2.0 | 3.0 | 4.0 | 3.9 | 4.9 | 6.0 | 10.7 | * |
| 2000..... | 2.3 | * | 1.5 | 0.6 | 1.7 | 2.3 | 2.8 | 3.3 | 4.4 | 6.3 | 10.3 | * |
| 1999..... | 2.2 | * | 2.1 | 0.6 | 2.0 | 2.4 | 2.1 | 3.6 | 4.3 | 5.7 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | 1.2 | * | * | * | 0.7 | * | 1.8 | * | * | * |
| 2003..... | 0.8 | * | * | * | * | * | 0.9 | * | 2.0 | * | * | * |
| 2002..... | 0.8 | * | * | * | * | 0.6 | 0.7 | * | * | * | * | * |
| 2001..... | 0.9 | * | 1.4 | * | * | * | 1.0 | * | * | * | 6.0 | * |
| 2000..... | 0.9 | * | 1.4 | * | * | * | 0.9 | * | * | * | * | * |
| 1999..... | 0.8 | * | 1.6 | * | * | * | 0.8 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pedestrian involved in collision with motor vehicle (V02-V04) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | 0.8 | 0.5 | 0.8 | 0.8 | 1.1 | 1.3 | 1.3 | 1.4 | 2.3 | 2.7 |
| 2003..... | 1.1 | * | 1.1 | 0.6 | 1.0 | 0.9 | 1.1 | 1.3 | 1.3 | 1.4 | 2.5 | 2.8 |
| 2002..... | 1.2 | * | 0.8 | 0.5 | 1.0 | 0.9 | 1.3 | 1.3 | 1.3 | 1.8 | 2.4 | 3.1 |
| 2001..... | 1.1 | * | 0.9 | 0.7 | 0.9 | 0.8 | 1.2 | 1.3 | 1.2 | 1.4 | 2.7 | 3.0 |
| 2000..... | 1.2 | * | 1.1 | 0.6 | 0.8 | 0.9 | 1.2 | 1.3 | 1.4 | 1.4 | 2.6 | 3.4 |
| 1999..... | 1.2 | * | 0.9 | 0.7 | 0.9 | 0.9 | 1.2 | 1.2 | 1.4 | 1.5 | 2.3 | 3.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | 1.0 | 0.6 | 1.1 | 1.2 | 1.6 | 2.0 | 1.9 | 1.8 | 3.2 | 4.9 |
| 2003..... | 1.6 | * | 1.4 | 0.7 | 1.3 | 1.2 | 1.5 | 1.9 | 2.0 | 2.0 | 3.0 | 5.4 |
| 2002..... | 1.6 | * | 1.1 | 0.6 | 1.4 | 1.4 | 1.7 | 1.9 | 1.8 | 2.3 | 3.0 | 5.5 |
| 2001..... | 1.6 | * | 1.2 | 0.9 | 1.4 | 1.2 | 1.7 | 1.8 | 1.8 | 1.8 | 3.8 | 5.7 |
| 2000..... | 1.6 | * | 1.4 | 0.8 | 1.2 | 1.3 | 1.7 | 1.9 | 2.0 | 1.9 | 3.5 | 6.5 |
| 1999..... | 1.6 | * | 1.0 | 0.8 | 1.3 | 1.3 | 1.9 | 1.8 | 2.0 | 2.0 | 3.5 | 6.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | 0.6 | 0.3 | 0.5 | 0.4 | 0.6 | 0.6 | 0.8 | 1.1 | 1.8 | 1.7 |
| 2003..... | 0.7 | * | 0.8 | 0.4 | 0.6 | 0.5 | 0.6 | 0.7 | 0.6 | 0.9 | 2.1 | 1.7 |
| 2002..... | 0.8 | * | 0.5 | 0.5 | 0.5 | 0.5 | 0.8 | 0.7 | 0.8 | 1.3 | 2.0 | 2.0 |
| 2001..... | 0.7 | * | 0.6 | 0.4 | 0.5 | 0.4 | 0.7 | 0.8 | 0.7 | 1.0 | 2.0 | 1.8 |
| 2000..... | 0.7 | * | 0.8 | 0.5 | 0.5 | 0.5 | 0.7 | 0.6 | 0.8 | 1.0 | 2.0 | 2.1 |
| 1999..... | 0.7 | * | 0.8 | 0.6 | 0.5 | 0.5 | 0.6 | 0.6 | 0.8 | 1.1 | 1.6 | 2.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | 0.7 | 0.4 | 0.8 | 0.7 | 1.0 | 1.1 | 1.2 | 1.3 | 2.2 | 2.6 |
| 2003..... | 1.0 | * | 1.0 | 0.5 | 1.0 | 0.8 | 0.9 | 1.1 | 1.2 | 1.2 | 2.3 | 2.8 |
| 2002..... | 1.1 | * | 0.6 | 0.5 | 0.9 | 0.8 | 1.0 | 1.1 | 1.2 | 1.6 | 2.2 | 2.9 |
| 2001..... | 1.0 | * | 0.8 | 0.6 | 0.9 | 0.7 | 1.0 | 1.1 | 1.1 | 1.2 | 2.5 | 2.9 |
| 2000..... | 1.0 | * | 0.9 | 0.6 | 0.9 | 0.8 | 1.0 | 1.1 | 1.2 | 1.3 | 2.5 | 3.1 |
| 1999..... | 1.0 | * | 0.7 | 0.6 | 0.9 | 0.7 | 1.0 | 1.0 | 1.2 | 1.3 | 2.2 | 3.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 0.9 | 0.5 | 1.1 | 1.0 | 1.5 | 1.7 | 1.6 | 1.6 | 2.9 | 4.6 |
| 2003..... | 1.4 | * | 1.3 | 0.6 | 1.3 | 1.1 | 1.2 | 1.6 | 1.9 | 1.7 | 2.7 | 5.1 |
| 2002..... | 1.4 | * | 0.8 | 0.5 | 1.4 | 1.3 | 1.4 | 1.6 | 1.6 | 2.0 | 2.7 | 5.0 |
| 2001..... | 1.4 | * | 1.1 | 0.8 | 1.3 | 1.1 | 1.4 | 1.5 | 1.5 | 1.6 | 3.5 | 5.3 |
| 2000..... | 1.4 | * | 1.1 | 0.7 | 1.1 | 1.1 | 1.3 | 1.6 | 1.7 | 1.6 | 3.1 | 5.7 |
| 1999..... | 1.4 | * | 0.7 | 0.7 | 1.2 | 1.1 | 1.4 | 1.5 | 1.7 | 1.8 | 3.1 | 5.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | 0.6 | 0.3 | 0.4 | 0.4 | 0.6 | 0.5 | 0.8 | 1.1 | 1.8 | 1.7 |
| 2003..... | 0.7 | * | 0.7 | 0.4 | 0.6 | 0.4 | 0.6 | 0.6 | 0.6 | 0.8 | 2.0 | 1.8 |
| 2002..... | 0.7 | * | 0.4 | 0.4 | 0.5 | 0.4 | 0.7 | 0.6 | 0.7 | 1.2 | 1.9 | 1.9 |
| 2001..... | 0.7 | * | 0.4 | 0.3 | 0.5 | 0.4 | 0.6 | 0.7 | 0.7 | 0.9 | 1.8 | 1.9 |
| 2000..... | 0.7 | * | 0.6 | 0.4 | 0.6 | 0.4 | 0.6 | 0.5 | 0.7 | 1.0 | 2.1 | 2.1 |
| 1999..... | 0.7 | * | 0.8 | 0.5 | 0.5 | 0.4 | 0.5 | 0.5 | 0.7 | 1.0 | 1.5 | 2.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Pedestrian involved in collision with motor vehicle (V02-V04) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 1.1 | 0.7 | 0.8 | 1.1 | 1.5 | 2.6 | 2.3 | 1.5 | 2.4 | * |
| 2003..... | 1.6 | * | 1.6 | 0.9 | 1.0 | 1.3 | 2.1 | 2.3 | 1.9 | 2.4 | 2.6 | * |
| 2002..... | 1.6 | * | 1.2 | 0.8 | 1.2 | 1.4 | 2.2 | 2.4 | 2.0 | 2.4 | 3.0 | * |
| 2001..... | 1.7 | * | 1.6 | 1.1 | 1.1 | 1.1 | 2.5 | 2.6 | 2.3 | 2.0 | 2.7 | * |
| 2000..... | 1.7 | * | 1.9 | 1.0 | 0.8 | 1.3 | 2.5 | 2.5 | 2.5 | 2.2 | 2.6 | * |
| 1999..... | 1.9 | * | 1.7 | 1.2 | 1.0 | 1.6 | 2.8 | 2.3 | 2.7 | 2.3 | 2.7 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | 0.9 | 1.2 | 1.8 | 2.2 | 4.3 | 4.2 | * | * | * |
| 2003..... | 2.3 | * | 1.9 | 1.1 | 1.3 | 1.8 | 3.3 | 3.8 | 3.1 | 4.9 | * | * |
| 2002..... | 2.4 | * | 1.6 | 0.9 | 1.7 | 2.2 | 3.2 | 4.0 | 3.4 | 3.8 | 6.2 | * |
| 2001..... | 2.5 | * | 1.8 | 1.7 | 1.5 | 1.6 | 3.7 | 3.8 | 4.3 | 3.1 | * | * |
| 2000..... | 2.6 | * | 2.3 | 1.3 | 1.4 | 2.0 | 3.8 | 3.9 | 4.1 | 4.3 | * | * |
| 1999..... | 2.9 | * | 2.1 | 1.4 | 1.4 | 2.6 | 4.6 | 4.2 | 4.7 | 3.9 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 1.0 | 1.1 | * | * | * | * |
| 2003..... | 0.9 | * | * | 0.7 | 0.7 | 0.8 | 1.0 | 1.0 | * | * | * | * |
| 2002..... | 0.9 | * | * | 0.7 | 0.7 | 0.7 | 1.3 | 1.1 | * | * | * | * |
| 2001..... | 1.0 | * | * | * | * | * | 1.5 | 1.5 | * | * | * | * |
| 2000..... | 0.9 | * | * | 0.7 | * | * | 1.4 | 1.2 | * | * | * | * |
| 1999..... | 0.9 | * | * | 1.0 | * | * | 1.2 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | 0.1 | 2.0 | 2.4 | 2.2 | 2.2 | 1.3 | 0.6 | 0.3 | * |
| 2003..... | 1.4 | * | * | 0.1 | 1.9 | 2.2 | 2.1 | 2.0 | 1.3 | 0.5 | 0.2 | * |
| 2002..... | 1.2 | * | * | 0.1 | 1.6 | 2.1 | 1.9 | 1.8 | 1.1 | 0.4 | 0.2 | * |
| 2001..... | 1.1 | * | * | 0.1 | 1.7 | 2.2 | 1.8 | 1.5 | 0.8 | 0.4 | 0.2 | * |
| 2000..... | 1.1 | * | * | 0.1 | 1.6 | 2.0 | 1.7 | 1.3 | 0.8 | 0.4 | * | * |
| 1999..... | 0.9 | * | * | 0.1 | 1.4 | 1.7 | 1.4 | 1.1 | 0.7 | 0.3 | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | 0.3 | 3.7 | 4.4 | 3.9 | 4.0 | 2.5 | 1.2 | 0.5 | * |
| 2003..... | 2.5 | * | * | 0.2 | 3.3 | 4.1 | 3.8 | 3.5 | 2.4 | 1.1 | 0.4 | * |
| 2002..... | 2.2 | * | * | 0.1 | 3.0 | 3.9 | 3.3 | 3.2 | 2.0 | 0.9 | 0.4 | * |
| 2001..... | 2.1 | * | * | 0.2 | 3.1 | 4.0 | 3.2 | 2.7 | 1.5 | 0.7 | * | * |
| 2000..... | 2.0 | * | * | 0.1 | 2.9 | 3.7 | 3.0 | 2.4 | 1.5 | 0.8 | * | * |
| 1999..... | 1.7 | * | * | 0.1 | 2.5 | 3.1 | 2.5 | 1.9 | 1.4 | 0.6 | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.3 | 0.4 | 0.6 | 0.5 | 0.3 | * | * | * |
| 2003..... | 0.3 | * | * | * | 0.3 | 0.4 | 0.5 | 0.5 | 0.2 | * | * | * |
| 2002..... | 0.2 | * | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.2 | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.2 | 0.4 | 0.5 | 0.3 | * | * | * | * |
| 2000..... | 0.2 | * | * | * | 0.2 | 0.3 | 0.4 | 0.3 | 0.2 | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.3 | 0.3 | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.2 | 2.3 | 2.4 | 2.4 | 2.6 | 1.5 | 0.7 | 0.3 | * |
| 2003..... | 1.5 | * | * | 0.1 | 2.1 | 2.2 | 2.3 | 2.3 | 1.4 | 0.6 | 0.2 | * |
| 2002..... | 1.3 | * | * | 0.1 | 1.8 | 2.1 | 2.1 | 2.0 | 1.2 | 0.4 | 0.2 | * |
| 2001..... | 1.2 | * | * | 0.1 | 1.8 | 2.2 | 2.1 | 1.7 | 0.9 | 0.4 | * | * |
| 2000..... | 1.1 | * | * | 0.1 | 1.7 | 1.9 | 1.9 | 1.5 | 0.9 | 0.4 | * | * |
| 1999..... | 1.0 | * | * | 0.1 | 1.4 | 1.6 | 1.6 | 1.3 | 0.8 | 0.3 | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.9 | * | * | 0.3 | 4.1 | 4.4 | 4.2 | 4.5 | 2.8 | 1.4 | 0.5 | * |
| 2003..... | 2.7 | * | * | 0.2 | 3.7 | 3.8 | 3.9 | 4.1 | 2.6 | 1.3 | 0.5 | * |
| 2002..... | 2.4 | * | * | * | 3.3 | 3.8 | 3.7 | 3.7 | 2.2 | 0.9 | * | * |
| 2001..... | 2.3 | * | * | 0.2 | 3.3 | 3.9 | 3.5 | 3.0 | 1.6 | 0.8 | * | * |
| 2000..... | 2.1 | * | * | 0.2 | 3.1 | 3.5 | 3.2 | 2.7 | 1.6 | 0.9 | * | * |
| 1999..... | 1.7 | * | * | 0.2 | 2.7 | 2.9 | 2.7 | 2.2 | 1.5 | 0.6 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.4 | 0.4 | 0.7 | 0.7 | 0.3 | * | * | * |
| 2003..... | 0.3 | * | * | * | 0.4 | 0.4 | 0.7 | 0.6 | 0.3 | * | * | * |
| 2002..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.5 | 0.4 | 0.2 | * | * | * |
| 2001..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.6 | 0.4 | * | * | * | * |
| 2000..... | 0.2 | * | * | * | 0.2 | 0.3 | 0.5 | 0.4 | 0.2 | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.2 | 0.3 | 0.4 | 0.3 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|

Motorcyclist involved in collision with railway train (V25,V35)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, both sexes. All cells contain an asterisk (*).

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, male. All cells contain an asterisk (*).

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic 4/, female. All cells contain an asterisk (*).

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, both sexes. All cells contain an asterisk (*).

Non-Hispanic white, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, male. All cells contain an asterisk (*).

Non-Hispanic white, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Non-Hispanic white, female. All cells contain an asterisk (*).

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | 0.2 | 0.2 | 4.1 | 2.1 | 1.5 | 1.2 | 1.1 | 1.2 | 1.6 | 1.2 |
| 2003..... | 1.7 | * | 0.2 | 0.3 | 4.2 | 2.3 | 1.6 | 1.4 | 1.1 | 1.3 | 1.6 | 1.9 |
| 2002..... | 1.8 | * | 0.2 | 0.3 | 4.5 | 2.3 | 1.7 | 1.4 | 1.3 | 1.2 | 1.6 | 1.5 |
| 2001..... | 1.5 | * | 0.1 | 0.2 | 3.8 | 2.0 | 1.4 | 1.2 | 0.9 | 1.1 | 1.6 | 1.7 |
| 2000..... | 1.4 | * | 0.2 | 0.2 | 3.5 | 1.7 | 1.4 | 1.0 | 1.0 | 1.2 | 1.4 | 1.7 |
| 1999..... | 1.4 | * | 0.1 | 0.2 | 3.3 | 1.8 | 1.3 | 1.0 | 1.0 | 1.2 | 1.5 | 1.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | 0.3 | 6.0 | 3.1 | 2.3 | 1.8 | 1.7 | 1.8 | 2.5 | 1.9 |
| 2003..... | 2.6 | * | * | 0.4 | 6.3 | 3.6 | 2.4 | 2.1 | 1.7 | 1.9 | 2.3 | 3.7 |
| 2002..... | 2.6 | * | * | 0.3 | 6.7 | 3.6 | 2.5 | 1.9 | 2.0 | 1.7 | 2.2 | 3.2 |
| 2001..... | 2.2 | * | * | 0.2 | 5.6 | 3.0 | 2.2 | 1.8 | 1.5 | 1.5 | 2.5 | 3.1 |
| 2000..... | 2.0 | * | * | 0.2 | 5.2 | 2.7 | 2.0 | 1.5 | 1.4 | 1.6 | 2.1 | 3.0 |
| 1999..... | 2.0 | * | * | 0.3 | 4.9 | 2.8 | 1.9 | 1.6 | 1.4 | 1.8 | 2.3 | 2.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | 0.2 | 2.1 | 1.0 | 0.7 | 0.6 | 0.6 | 0.8 | 1.0 | 0.9 |
| 2003..... | 0.9 | * | 0.3 | 0.2 | 2.0 | 1.1 | 0.8 | 0.8 | 0.5 | 0.9 | 1.1 | 1.1 |
| 2002..... | 0.9 | * | * | 0.3 | 2.3 | 0.9 | 0.9 | 0.8 | 0.6 | 0.8 | 1.2 | 0.8 |
| 2001..... | 0.8 | * | * | 0.2 | 1.9 | 0.9 | 0.7 | 0.6 | 0.4 | 0.7 | 1.1 | 1.2 |
| 2000..... | 0.8 | * | * | 0.2 | 1.6 | 0.8 | 0.8 | 0.6 | 0.6 | 0.8 | 0.9 | 1.2 |
| 1999..... | 0.7 | * | * | 0.2 | 1.7 | 0.7 | 0.7 | 0.5 | 0.6 | 0.8 | 1.0 | 0.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 4.3 | 2.0 | 1.6 | 0.9 | * | * | * | * |
| 2003..... | 1.7 | * | * | 0.4 | 4.4 | 2.2 | 1.4 | 1.1 | * | * | * | * |
| 2002..... | 1.8 | * | * | * | 4.4 | 2.6 | 1.6 | 1.0 | * | * | * | * |
| 2001..... | 1.6 | * | * | * | 3.7 | 2.1 | 1.4 | 1.4 | 1.1 | * | * | * |
| 2000..... | 1.3 | * | * | * | 3.2 | 1.6 | 1.1 | 1.0 | * | * | * | * |
| 1999..... | 1.3 | * | * | * | 3.1 | 1.8 | 1.3 | 0.9 | * | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.5 | * | * | * | 6.8 | 3.1 | 2.4 | 1.6 | * | * | * | * |
| 2003..... | 2.6 | * | * | 0.6 | 6.9 | 3.4 | 2.0 | 1.6 | * | * | * | * |
| 2002..... | 2.8 | * | * | * | 6.9 | 4.2 | 2.6 | 1.2 | * | * | * | * |
| 2001..... | 2.4 | * | * | * | 5.7 | 3.2 | 2.2 | 2.1 | * | * | * | * |
| 2000..... | 1.9 | * | * | * | 4.8 | 2.6 | 1.5 | 1.4 | * | * | * | * |
| 1999..... | 2.1 | * | * | * | 4.8 | 2.9 | 2.1 | 1.4 | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | 1.4 | 0.8 | 0.7 | * | * | * | * | * |
| 2003..... | 0.6 | * | * | * | 1.5 | 0.7 | * | * | * | * | * | * |
| 2002..... | 0.7 | * | * | * | 1.5 | 0.8 | * | * | * | * | * | * |
| 2001..... | 0.7 | * | * | * | 1.4 | 0.8 | * | * | * | * | * | * |
| 2000..... | 0.6 | * | * | * | 1.3 | * | * | * | * | * | * | * |
| 1999..... | 0.5 | * | * | * | 1.1 | * | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.2 | 4.1 | 2.1 | 1.5 | 1.2 | 1.2 | 1.3 | 1.7 | 1.2 |
| 2003..... | 1.7 | * | 0.2 | 0.3 | 4.2 | 2.4 | 1.6 | 1.5 | 1.1 | 1.4 | 1.6 | 1.9 |
| 2002..... | 1.7 | * | 0.2 | 0.3 | 4.5 | 2.2 | 1.7 | 1.4 | 1.3 | 1.2 | 1.6 | 1.6 |
| 2001..... | 1.5 | * | * | 0.2 | 3.8 | 1.9 | 1.4 | 1.2 | 0.9 | 1.1 | 1.6 | 1.7 |
| 2000..... | 1.4 | * | 0.2 | 0.2 | 3.5 | 1.7 | 1.5 | 1.0 | 1.0 | 1.2 | 1.4 | 1.8 |
| 1999..... | 1.4 | * | * | 0.2 | 3.4 | 1.8 | 1.3 | 1.1 | 1.0 | 1.2 | 1.5 | 1.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | 0.2 | 5.8 | 3.1 | 2.3 | 1.8 | 1.8 | 1.8 | 2.7 | 2.0 |
| 2003..... | 2.5 | * | * | 0.3 | 6.1 | 3.6 | 2.4 | 2.1 | 1.8 | 1.9 | 2.4 | 3.7 |
| 2002..... | 2.6 | * | * | 0.3 | 6.6 | 3.4 | 2.5 | 2.0 | 2.0 | 1.8 | 2.3 | 3.3 |
| 2001..... | 2.2 | * | * | 0.2 | 5.6 | 3.0 | 2.1 | 1.8 | 1.5 | 1.5 | 2.5 | 3.0 |
| 2000..... | 2.1 | * | * | 0.2 | 5.3 | 2.7 | 2.1 | 1.5 | 1.4 | 1.7 | 2.1 | 3.1 |
| 1999..... | 2.0 | * | * | 0.3 | 4.8 | 2.8 | 1.9 | 1.6 | 1.5 | 1.8 | 2.4 | 2.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | 0.2 | 2.3 | 1.0 | 0.7 | 0.7 | 0.6 | 0.8 | 1.0 | 0.8 |
| 2003..... | 0.9 | * | * | 0.2 | 2.1 | 1.2 | 0.8 | 0.8 | 0.6 | 0.9 | 1.1 | 1.1 |
| 2002..... | 1.0 | * | * | 0.3 | 2.4 | 1.0 | 1.0 | 0.8 | 0.6 | 0.8 | 1.1 | 0.8 |
| 2001..... | 0.8 | * | * | 0.2 | 2.0 | 0.9 | 0.7 | 0.6 | 0.4 | 0.7 | 1.0 | 1.2 |
| 2000..... | 0.8 | * | * | 0.2 | 1.6 | 0.8 | 0.9 | 0.6 | 0.6 | 0.8 | 0.9 | 1.3 |
| 1999..... | 0.7 | * | * | 0.2 | 1.9 | 0.8 | 0.7 | 0.5 | 0.7 | 0.8 | 0.9 | 0.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.2 | 4.4 | 2.0 | 1.5 | 1.2 | 1.1 | 1.3 | 1.7 | 1.2 |
| 2003..... | 1.8 | * | * | 0.3 | 4.5 | 2.5 | 1.7 | 1.4 | 1.1 | 1.3 | 1.7 | 2.0 |
| 2002..... | 1.8 | * | * | 0.3 | 4.8 | 2.2 | 1.8 | 1.4 | 1.2 | 1.2 | 1.7 | 1.7 |
| 2001..... | 1.5 | * | * | 0.2 | 3.8 | 1.9 | 1.4 | 1.1 | 0.9 | 1.1 | 1.6 | 1.9 |
| 2000..... | 1.4 | * | * | 0.2 | 3.6 | 1.7 | 1.5 | 1.0 | 1.0 | 1.1 | 1.4 | 1.9 |
| 1999..... | 1.3 | * | * | 0.2 | 3.5 | 1.7 | 1.3 | 1.0 | 1.0 | 1.2 | 1.5 | 1.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | 0.3 | 6.3 | 3.0 | 2.3 | 1.9 | 1.6 | 1.7 | 2.7 | 2.2 |
| 2003..... | 2.6 | * | * | 0.4 | 6.6 | 3.6 | 2.6 | 2.0 | 1.7 | 1.8 | 2.4 | 3.9 |
| 2002..... | 2.6 | * | * | 0.3 | 6.9 | 3.4 | 2.5 | 1.9 | 1.9 | 1.6 | 2.3 | 3.3 |
| 2001..... | 2.2 | * | * | 0.2 | 5.6 | 2.9 | 2.1 | 1.7 | 1.5 | 1.5 | 2.5 | 3.2 |
| 2000..... | 2.0 | * | * | 0.2 | 5.4 | 2.5 | 2.1 | 1.4 | 1.3 | 1.5 | 2.0 | 3.3 |
| 1999..... | 2.0 | * | * | 0.3 | 4.9 | 2.7 | 1.9 | 1.5 | 1.4 | 1.7 | 2.4 | 2.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | 0.2 | 2.4 | 1.1 | 0.7 | 0.6 | 0.6 | 0.9 | 1.1 | 0.8 |
| 2003..... | 1.0 | * | * | 0.2 | 2.4 | 1.3 | 0.9 | 0.8 | 0.6 | 0.9 | 1.2 | 1.2 |
| 2002..... | 1.0 | * | * | 0.3 | 2.7 | 1.0 | 1.0 | 0.8 | 0.6 | 0.9 | 1.2 | 0.9 |
| 2001..... | 0.8 | * | * | 0.2 | 2.0 | 0.9 | 0.7 | 0.6 | 0.4 | 0.8 | 1.1 | 1.3 |
| 2000..... | 0.8 | * | * | 0.2 | 1.8 | 0.9 | 0.9 | 0.5 | 0.6 | 0.8 | 0.9 | 1.3 |
| 1999..... | 0.7 | * | * | 0.2 | 1.9 | 0.8 | 0.7 | 0.5 | 0.6 | 0.8 | 0.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 3.0 | 2.8 | 1.8 | 1.5 | 2.0 | 1.4 | * | * |
| 2003..... | 1.7 | * | * | * | 3.2 | 2.5 | 1.3 | 2.0 | 1.6 | 2.1 | * | * |
| 2002..... | 1.9 | * | * | 0.5 | 3.9 | 2.7 | 1.8 | 1.8 | 2.3 | 1.8 | * | * |
| 2001..... | 1.8 | * | * | * | 4.2 | 2.5 | 1.9 | 1.7 | 1.1 | * | 2.2 | * |
| 2000..... | 1.6 | * | * | * | 3.2 | 2.1 | 1.7 | 1.7 | 1.5 | 2.0 | * | * |
| 1999..... | 1.7 | * | * | * | 3.5 | 2.3 | 1.7 | 1.5 | 1.6 | 1.4 | 2.2 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 4.3 | 4.8 | 2.8 | 2.2 | 3.5 | * | * | * |
| 2003..... | 2.7 | * | * | * | 5.1 | 4.3 | 2.2 | 3.1 | 2.8 | * | * | * |
| 2002..... | 3.1 | * | * | * | 6.0 | 4.7 | 2.7 | 3.0 | 4.1 | 3.7 | * | * |
| 2001..... | 2.8 | * | * | * | 6.4 | 4.1 | 3.0 | 2.7 | 2.0 | * | * | * |
| 2000..... | 2.5 | * | * | * | 4.9 | 3.8 | 2.7 | 2.5 | 2.6 | 3.3 | * | * |
| 1999..... | 2.6 | * | * | * | 5.2 | 4.0 | 2.6 | 2.5 | 2.6 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | 1.8 | 1.0 | 0.9 | 1.0 | * | * | * | * |
| 2003..... | 0.7 | * | * | * | 1.3 | 0.9 | * | 1.0 | * | * | * | * |
| 2002..... | 0.9 | * | * | * | 1.7 | 0.9 | 1.1 | 0.9 | * | * | * | * |
| 2001..... | 0.8 | * | * | * | 2.0 | 1.1 | 0.9 | * | * | * | * | * |
| 2000..... | 0.8 | * | * | * | 1.5 | * | 0.8 | 1.0 | * | * | * | * |
| 1999..... | 0.8 | * | * | * | 1.7 | 0.8 | 0.9 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | 1.5 | 0.8 | 0.9 | 6.0 | 3.6 | 3.0 | 3.1 | 3.5 | 5.0 | 8.3 | 9.6 |
| 2003..... | 3.7 | 1.5 | 0.9 | 0.8 | 6.1 | 3.7 | 3.2 | 3.2 | 3.5 | 4.9 | 8.7 | 10.2 |
| 2002..... | 3.9 | 1.1 | 1.0 | 0.9 | 6.8 | 3.8 | 3.4 | 3.3 | 3.8 | 5.2 | 9.2 | 9.9 |
| 2001..... | 3.7 | 1.6 | 1.0 | 0.9 | 6.1 | 3.7 | 3.3 | 3.1 | 3.5 | 5.1 | 8.4 | 10.1 |
| 2000..... | 3.7 | 1.5 | 0.9 | 0.9 | 6.1 | 3.7 | 3.1 | 3.2 | 3.8 | 5.0 | 9.0 | 10.1 |
| 1999..... | 3.7 | 1.6 | 0.9 | 0.9 | 5.8 | 3.7 | 3.1 | 3.2 | 3.8 | 5.3 | 8.8 | 9.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.4 | 1.9 | 0.8 | 1.0 | 7.4 | 4.7 | 3.8 | 3.8 | 4.0 | 5.8 | 10.9 | 15.9 |
| 2003..... | 4.5 | 1.4 | 0.9 | 0.8 | 7.4 | 4.9 | 4.1 | 4.0 | 4.2 | 5.4 | 11.2 | 16.6 |
| 2002..... | 4.8 | 1.5 | 1.0 | 0.9 | 8.7 | 5.1 | 4.2 | 4.1 | 4.7 | 6.0 | 12.2 | 16.9 |
| 2001..... | 4.5 | 1.6 | 0.9 | 0.9 | 7.8 | 5.0 | 4.2 | 3.8 | 4.1 | 6.1 | 10.9 | 18.1 |
| 2000..... | 4.5 | 1.5 | 0.9 | 0.9 | 7.5 | 4.8 | 3.9 | 4.0 | 4.5 | 5.8 | 11.6 | 18.2 |
| 1999..... | 4.4 | 1.7 | 1.0 | 0.9 | 7.3 | 4.9 | 3.8 | 3.9 | 4.3 | 6.2 | 11.6 | 17.8 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.9 | 1.1 | 0.9 | 0.9 | 4.4 | 2.4 | 2.1 | 2.5 | 3.0 | 4.3 | 6.5 | 6.7 |
| 2003..... | 2.9 | 1.5 | 0.9 | 0.8 | 4.7 | 2.5 | 2.3 | 2.4 | 2.9 | 4.4 | 7.0 | 7.4 |
| 2002..... | 3.0 | * | 0.9 | 0.8 | 4.9 | 2.4 | 2.6 | 2.5 | 3.0 | 4.4 | 7.2 | 6.8 |
| 2001..... | 2.9 | 1.6 | 1.1 | 0.9 | 4.3 | 2.3 | 2.5 | 2.4 | 3.0 | 4.3 | 6.7 | 6.8 |
| 2000..... | 2.9 | 1.5 | 0.9 | 0.8 | 4.5 | 2.6 | 2.3 | 2.4 | 3.2 | 4.3 | 7.3 | 6.8 |
| 1999..... | 2.9 | 1.4 | 0.7 | 1.0 | 4.3 | 2.5 | 2.4 | 2.5 | 3.3 | 4.6 | 7.0 | 5.7 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.4 | * | 0.8 | 1.0 | 6.3 | 4.1 | 3.3 | 3.6 | 3.6 | 3.9 | 6.1 | * |
| 2003..... | 3.5 | * | 0.9 | 0.9 | 5.9 | 4.8 | 3.4 | 3.6 | 3.8 | 4.1 | 6.1 | * |
| 2002..... | 3.9 | * | 1.3 | 0.9 | 7.6 | 4.4 | 4.0 | 3.7 | 4.1 | 4.8 | 6.7 | * |
| 2001..... | 3.6 | * | 1.0 | 0.9 | 6.7 | 4.1 | 3.6 | 3.4 | 3.5 | 5.0 | 6.3 | * |
| 2000..... | 3.1 | * | 1.2 | 0.9 | 4.8 | 3.6 | 3.0 | 3.7 | 3.6 | 5.1 | 7.7 | * |
| 1999..... | 2.9 | * | 0.9 | 0.9 | 4.8 | 3.8 | 2.4 | 3.0 | 3.8 | 5.2 | 6.5 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | 1.0 | 8.4 | 5.6 | 4.4 | 4.4 | 4.3 | 5.0 | 8.2 | * |
| 2003..... | 4.8 | * | * | 1.2 | 8.0 | 7.0 | 4.4 | 5.1 | 5.1 | 4.2 | 8.8 | * |
| 2002..... | 5.4 | * | 1.6 | 1.2 | 10.5 | 6.4 | 5.4 | 4.7 | 4.8 | 7.1 | 11.9 | * |
| 2001..... | 4.9 | * | * | 1.0 | 9.5 | 5.9 | 5.1 | 4.2 | 4.3 | 6.4 | 13.0 | * |
| 2000..... | 4.2 | * | * | 0.9 | 6.8 | 4.8 | 4.3 | 5.2 | 5.0 | 6.5 | 12.8 | * |
| 1999..... | 3.9 | * | * | 0.8 | 6.2 | 5.6 | 3.2 | 4.2 | 4.7 | 5.7 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | 1.0 | 4.0 | 2.1 | 2.1 | 2.8 | 3.0 | 3.1 | * | * |
| 2003..... | 2.1 | * | * | 0.6 | 3.5 | 2.0 | 2.3 | 2.1 | 2.6 | 4.0 | * | * |
| 2002..... | 2.3 | * | * | * | 4.1 | 2.1 | 2.4 | 2.7 | 3.4 | 3.0 | * | * |
| 2001..... | 2.2 | * | * | 0.8 | 3.5 | 2.1 | 2.1 | 2.7 | 2.8 | 3.9 | * | * |
| 2000..... | 1.9 | * | * | 0.8 | 2.5 | 2.2 | 1.6 | 2.2 | 2.4 | 4.0 | * | * |
| 1999..... | 1.9 | * | * | 0.9 | 3.1 | 1.8 | 1.5 | 1.8 | 3.0 | 4.8 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.6 | 1.4 | 0.8 | 0.9 | 5.9 | 3.5 | 2.9 | 3.1 | 3.5 | 5.1 | 8.4 | 9.7 |
| 2003..... | 3.7 | 1.5 | 0.9 | 0.8 | 6.1 | 3.5 | 3.2 | 3.2 | 3.5 | 4.9 | 8.8 | 10.2 |
| 2002..... | 3.9 | 1.1 | 0.9 | 0.9 | 6.7 | 3.6 | 3.3 | 3.2 | 3.8 | 5.2 | 9.3 | 10.1 |
| 2001..... | 3.7 | 1.4 | 1.0 | 0.9 | 6.0 | 3.6 | 3.2 | 3.1 | 3.6 | 5.1 | 8.4 | 10.2 |
| 2000..... | 3.8 | 1.4 | 0.8 | 0.9 | 6.3 | 3.7 | 3.1 | 3.1 | 3.8 | 5.0 | 9.0 | 10.3 |
| 1999..... | 3.8 | 1.7 | 0.8 | 1.0 | 6.0 | 3.7 | 3.2 | 3.2 | 3.8 | 5.3 | 8.9 | 9.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.3 | 2.0 | 0.7 | 0.9 | 7.1 | 4.5 | 3.7 | 3.7 | 4.0 | 5.8 | 11.1 | 16.3 |
| 2003..... | 4.4 | 1.4 | 0.8 | 0.7 | 7.3 | 4.4 | 4.1 | 3.9 | 4.1 | 5.5 | 11.3 | 16.7 |
| 2002..... | 4.7 | 1.3 | 0.9 | 0.8 | 8.3 | 4.8 | 4.0 | 4.0 | 4.6 | 5.9 | 12.2 | 17.2 |
| 2001..... | 4.5 | 1.5 | 0.9 | 0.9 | 7.5 | 4.7 | 4.0 | 3.8 | 4.1 | 6.1 | 10.8 | 18.4 |
| 2000..... | 4.5 | 1.5 | 0.9 | 0.9 | 7.7 | 4.7 | 3.9 | 3.9 | 4.4 | 5.8 | 11.5 | 18.8 |
| 1999..... | 4.5 | 1.7 | 1.0 | 0.9 | 7.4 | 4.7 | 3.9 | 3.9 | 4.3 | 6.2 | 11.7 | 18.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.9 | * | 0.9 | 0.8 | 4.5 | 2.4 | 2.1 | 2.5 | 3.0 | 4.4 | 6.6 | 6.8 |
| 2003..... | 3.1 | 1.6 | 1.0 | 0.8 | 4.9 | 2.6 | 2.3 | 2.5 | 2.9 | 4.4 | 7.1 | 7.4 |
| 2002..... | 3.1 | * | 0.9 | 0.9 | 5.0 | 2.5 | 2.6 | 2.5 | 3.0 | 4.5 | 7.4 | 7.0 |
| 2001..... | 3.0 | 1.3 | 1.1 | 1.0 | 4.4 | 2.4 | 2.5 | 2.4 | 3.0 | 4.3 | 6.8 | 6.7 |
| 2000..... | 3.1 | * | 0.8 | 0.8 | 4.9 | 2.6 | 2.4 | 2.4 | 3.2 | 4.4 | 7.4 | 6.9 |
| 1999..... | 3.0 | 1.6 | 0.7 | 1.0 | 4.5 | 2.7 | 2.5 | 2.5 | 3.3 | 4.5 | 7.1 | 5.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | 1.3 | 0.8 | 0.9 | 6.5 | 3.4 | 2.9 | 3.0 | 3.5 | 5.2 | 8.6 | 10.2 |
| 2003..... | 3.9 | 1.5 | 0.8 | 0.7 | 6.7 | 3.6 | 3.2 | 3.2 | 3.3 | 4.9 | 9.1 | 10.6 |
| 2002..... | 4.1 | 0.9 | 0.8 | 0.9 | 7.2 | 3.5 | 3.3 | 3.2 | 3.8 | 5.3 | 9.7 | 10.6 |
| 2001..... | 3.8 | 1.3 | 1.0 | 0.9 | 6.3 | 3.6 | 3.2 | 3.0 | 3.6 | 5.1 | 8.8 | 10.7 |
| 2000..... | 3.9 | 1.3 | 0.8 | 0.9 | 6.8 | 3.7 | 3.1 | 3.0 | 3.8 | 5.1 | 9.3 | 10.9 |
| 1999..... | 3.9 | 1.6 | 0.8 | 1.0 | 6.5 | 3.7 | 3.1 | 3.2 | 3.7 | 5.3 | 9.2 | 9.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.5 | 2.0 | 0.8 | 1.0 | 7.8 | 4.3 | 3.7 | 3.6 | 3.9 | 5.9 | 11.1 | 17.0 |
| 2003..... | 4.6 | * | 0.8 | 0.7 | 7.8 | 4.5 | 4.1 | 3.9 | 3.8 | 5.5 | 11.5 | 17.4 |
| 2002..... | 4.9 | * | 0.7 | 0.9 | 8.8 | 4.6 | 4.0 | 3.9 | 4.6 | 5.9 | 12.5 | 18.2 |
| 2001..... | 4.6 | * | 0.8 | 0.9 | 7.8 | 4.7 | 3.9 | 3.5 | 4.1 | 6.1 | 11.1 | 19.2 |
| 2000..... | 4.6 | * | 0.8 | 0.9 | 8.1 | 4.6 | 3.8 | 3.7 | 4.2 | 5.9 | 11.7 | 20.0 |
| 1999..... | 4.6 | 1.9 | 0.9 | 0.9 | 7.9 | 4.7 | 3.6 | 3.8 | 4.1 | 6.1 | 11.9 | 18.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | 0.8 | 0.9 | 5.1 | 2.5 | 2.1 | 2.5 | 3.1 | 4.6 | 6.9 | 7.2 |
| 2003..... | 3.2 | * | 0.9 | 0.8 | 5.5 | 2.7 | 2.3 | 2.5 | 2.8 | 4.4 | 7.5 | 7.6 |
| 2002..... | 3.3 | * | 0.8 | 0.9 | 5.6 | 2.4 | 2.6 | 2.4 | 3.0 | 4.8 | 7.9 | 7.3 |
| 2001..... | 3.1 | * | 1.1 | 1.0 | 4.8 | 2.5 | 2.6 | 2.4 | 3.0 | 4.3 | 7.3 | 7.2 |
| 2000..... | 3.2 | * | 0.8 | 0.9 | 5.4 | 2.6 | 2.4 | 2.3 | 3.4 | 4.5 | 7.8 | 7.2 |
| 1999..... | 3.2 | * | 0.6 | 1.1 | 5.0 | 2.7 | 2.6 | 2.6 | 3.4 | 4.7 | 7.4 | 5.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 3.1 | * | 0.9 | 0.8 | 4.0 | 4.4 | 3.0 | 3.5 | 3.8 | 4.5 | 6.9 | 5.8 |
| 2003..... | 3.3 | * | 1.0 | 0.8 | 4.3 | 3.7 | 3.6 | 3.5 | 4.8 | 5.1 | 6.7 | 6.5 |
| 2002..... | 3.4 | * | 1.3 | 0.8 | 4.9 | 4.8 | 3.6 | 3.6 | 3.8 | 4.0 | 5.8 | * |
| 2001..... | 3.4 | * | 1.2 | 0.9 | 4.8 | 4.0 | 3.8 | 4.0 | 3.9 | 6.0 | 5.2 | * |
| 2000..... | 3.4 | * | 1.1 | 0.7 | 4.8 | 4.3 | 3.5 | 4.3 | 4.1 | 4.5 | 5.7 | * |
| 1999..... | 3.5 | * | 1.1 | 1.0 | 4.4 | 4.3 | 4.1 | 3.7 | 4.1 | 5.1 | 6.3 | 7.8 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | 0.7 | 5.2 | 6.5 | 3.8 | 4.5 | 5.0 | 5.8 | 12.3 | * |
| 2003..... | 4.2 | * | * | 0.6 | 5.9 | 5.1 | 4.6 | 4.6 | 6.4 | 6.2 | 11.3 | * |
| 2002..... | 4.5 | * | 1.7 | * | 6.4 | 6.8 | 4.8 | 4.9 | 5.3 | 5.4 | 10.0 | * |
| 2001..... | 4.6 | * | * | 0.9 | 6.4 | 5.7 | 5.3 | 6.0 | 4.9 | 7.3 | 8.7 | * |
| 2000..... | 4.5 | * | * | 0.9 | 6.4 | 6.0 | 4.8 | 5.9 | 6.8 | 5.7 | 8.6 | * |
| 1999..... | 4.6 | * | * | 1.1 | 6.0 | 5.8 | 5.8 | 5.1 | 5.5 | 7.8 | 10.9 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | 1.0 | 2.9 | 2.4 | 2.2 | 2.6 | 2.8 | 3.6 | 3.9 | * |
| 2003..... | 2.5 | * | * | 0.9 | 2.8 | 2.5 | 2.8 | 2.5 | 3.5 | 4.3 | 4.1 | * |
| 2002..... | 2.4 | * | * | 0.9 | 3.4 | 3.0 | 2.6 | 2.6 | 2.5 | 3.0 | 3.5 | * |
| 2001..... | 2.4 | * | * | 0.9 | 3.2 | 2.5 | 2.5 | 2.3 | 3.1 | 5.1 | * | * |
| 2000..... | 2.3 | * | * | * | 3.2 | 2.7 | 2.4 | 2.9 | 1.9 | 3.6 | 4.0 | * |
| 1999..... | 2.4 | * | * | 0.9 | 2.9 | 2.9 | 2.6 | 2.5 | 2.9 | 3.2 | 3.7 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 0.3 | 0.4 | 3.1 | 1.7 | 1.5 | 1.2 | 1.1 | 1.0 | 0.9 | 1.0 |
| 2003..... | 1.5 | * | 0.3 | 0.4 | 3.3 | 1.9 | 1.6 | 1.4 | 1.2 | 1.0 | 1.0 | 1.3 |
| 2002..... | 1.5 | * | 0.3 | 0.4 | 3.5 | 2.0 | 1.6 | 1.3 | 1.2 | 1.0 | 1.0 | 0.9 |
| 2001..... | 1.4 | * | 0.2 | 0.4 | 3.0 | 1.8 | 1.5 | 1.3 | 1.1 | 1.0 | 1.0 | 0.7 |
| 2000..... | 1.3 | 0.6 | 0.2 | 0.4 | 3.1 | 1.8 | 1.4 | 1.2 | 0.9 | 0.8 | 0.9 | 1.0 |
| 1999..... | 1.3 | * | 0.3 | 0.3 | 3.1 | 1.7 | 1.5 | 1.1 | 1.1 | 0.9 | 0.8 | 1.0 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | 0.3 | 0.4 | 4.2 | 2.5 | 2.2 | 1.8 | 1.6 | 1.4 | 1.2 | 1.7 |
| 2003..... | 2.1 | * | 0.3 | 0.4 | 4.5 | 2.8 | 2.2 | 1.9 | 1.6 | 1.5 | 1.6 | 2.0 |
| 2002..... | 2.2 | * | 0.4 | 0.4 | 4.8 | 3.0 | 2.3 | 2.0 | 1.6 | 1.4 | 1.5 | 1.8 |
| 2001..... | 2.0 | * | * | 0.4 | 4.2 | 2.7 | 2.1 | 1.9 | 1.6 | 1.4 | 1.5 | * |
| 2000..... | 1.9 | * | * | 0.4 | 4.3 | 2.8 | 2.0 | 1.8 | 1.3 | 1.2 | 1.4 | 2.2 |
| 1999..... | 1.9 | * | 0.3 | 0.3 | 4.2 | 2.6 | 2.1 | 1.6 | 1.6 | 1.3 | 1.2 | 2.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | 0.3 | 0.4 | 1.9 | 0.9 | 0.9 | 0.7 | 0.6 | 0.6 | 0.6 | 0.7 |
| 2003..... | 0.9 | * | 0.4 | 0.4 | 2.1 | 1.0 | 1.0 | 0.8 | 0.8 | 0.6 | 0.6 | 1.0 |
| 2002..... | 0.9 | * | * | 0.4 | 2.2 | 1.0 | 1.0 | 0.7 | 0.7 | 0.6 | 0.7 | * |
| 2001..... | 0.8 | * | * | 0.4 | 1.8 | 0.9 | 0.9 | 0.6 | 0.6 | 0.6 | 0.8 | * |
| 2000..... | 0.7 | * | * | 0.3 | 1.7 | 0.8 | 0.8 | 0.6 | 0.6 | 0.5 | 0.6 | * |
| 1999..... | 0.8 | * | 0.3 | 0.3 | 1.9 | 0.9 | 0.8 | 0.6 | 0.6 | 0.7 | 0.5 | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | 0.5 | 3.2 | 2.1 | 1.5 | 1.1 | 1.8 | * | * | * |
| 2003..... | 1.7 | * | * | 0.5 | 3.7 | 2.3 | 1.7 | 1.2 | 1.6 | * | * | * |
| 2002..... | 1.7 | * | 0.7 | 0.6 | 3.3 | 2.3 | 1.8 | 1.5 | 1.3 | * | * | * |
| 2001..... | 1.6 | * | * | 0.5 | 3.2 | 1.8 | 1.9 | 1.7 | 1.1 | * | * | * |
| 2000..... | 1.5 | * | * | 0.5 | 2.7 | 1.8 | 1.7 | 1.5 | * | * | * | * |
| 1999..... | 1.3 | * | * | 0.3 | 2.6 | 1.6 | 1.4 | 1.2 | * | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | 0.5 | 4.7 | 3.2 | 2.4 | 1.9 | 2.6 | * | * | * |
| 2003..... | 2.3 | * | * | * | 4.9 | 3.4 | 2.2 | 1.6 | * | * | * | * |
| 2002..... | 2.4 | * | * | 0.6 | 4.5 | 3.2 | 2.5 | 2.1 | 2.5 | * | * | * |
| 2001..... | 2.3 | * | * | * | 4.5 | 2.7 | 2.9 | 2.4 | * | * | * | * |
| 2000..... | 2.2 | * | * | * | 4.2 | 2.8 | 2.6 | 1.9 | * | * | * | * |
| 1999..... | 1.9 | * | * | * | 4.0 | 2.6 | 2.1 | 1.6 | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | 0.6 | 1.6 | 0.9 | * | * | * | * | * | * |
| 2003..... | 1.1 | * | * | 0.5 | 2.3 | 1.0 | 1.1 | * | * | * | * | * |
| 2002..... | 1.0 | * | * | * | 1.9 | 1.2 | 1.0 | * | * | * | * | * |
| 2001..... | 0.8 | * | * | * | 1.6 | 0.7 | 0.8 | * | * | * | * | * |
| 2000..... | 0.7 | * | * | * | 0.9 | 0.7 | * | * | * | * | * | * |
| 1999..... | 0.6 | * | * | * | 1.1 | * | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | 0.3 | 0.3 | 3.0 | 1.6 | 1.5 | 1.2 | 1.0 | 1.0 | 0.9 | 1.0 |
| 2003..... | 1.4 | * | 0.3 | 0.4 | 3.3 | 1.8 | 1.6 | 1.4 | 1.1 | 1.0 | 1.0 | 1.3 |
| 2002..... | 1.5 | * | 0.2 | 0.3 | 3.5 | 2.0 | 1.6 | 1.3 | 1.1 | 1.0 | 1.0 | 0.9 |
| 2001..... | 1.3 | * | 0.2 | 0.3 | 3.0 | 1.8 | 1.4 | 1.2 | 1.1 | 0.9 | 1.0 | 0.8 |
| 2000..... | 1.3 | * | 0.2 | 0.3 | 3.1 | 1.8 | 1.4 | 1.1 | 0.9 | 0.8 | 0.9 | 1.0 |
| 1999..... | 1.3 | * | 0.3 | 0.3 | 3.2 | 1.7 | 1.5 | 1.1 | 1.1 | 0.9 | 0.8 | 1.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | 0.4 | 4.0 | 2.3 | 2.2 | 1.8 | 1.5 | 1.4 | 1.3 | 1.8 |
| 2003..... | 2.0 | * | * | 0.4 | 4.5 | 2.6 | 2.2 | 1.9 | 1.6 | 1.5 | 1.6 | 2.0 |
| 2002..... | 2.1 | * | * | 0.4 | 4.9 | 2.9 | 2.2 | 1.9 | 1.6 | 1.4 | 1.5 | 1.8 |
| 2001..... | 1.9 | * | * | 0.3 | 4.1 | 2.7 | 2.0 | 1.9 | 1.5 | 1.4 | 1.4 | * |
| 2000..... | 1.9 | * | * | 0.4 | 4.3 | 2.8 | 2.0 | 1.8 | 1.3 | 1.2 | 1.3 | 2.2 |
| 1999..... | 1.9 | * | * | 0.3 | 4.3 | 2.5 | 2.1 | 1.6 | 1.6 | 1.3 | 1.2 | 2.0 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | 0.3 | 2.0 | 0.9 | 0.9 | 0.7 | 0.5 | 0.6 | 0.6 | 0.7 |
| 2003..... | 0.9 | * | * | 0.3 | 2.0 | 1.0 | 1.0 | 0.8 | 0.7 | 0.6 | 0.7 | 1.0 |
| 2002..... | 0.9 | * | * | 0.3 | 2.2 | 1.0 | 0.9 | 0.7 | 0.8 | 0.6 | 0.6 | * |
| 2001..... | 0.8 | * | * | 0.3 | 1.9 | 0.9 | 0.9 | 0.6 | 0.6 | 0.5 | 0.7 | * |
| 2000..... | 0.8 | * | * | 0.3 | 1.9 | 0.9 | 0.8 | 0.5 | 0.6 | 0.5 | 0.6 | * |
| 1999..... | 0.8 | * | 0.4 | 0.3 | 2.0 | 0.9 | 0.9 | 0.6 | 0.6 | 0.6 | 0.5 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 0.2 | 0.4 | 3.3 | 1.6 | 1.5 | 1.2 | 1.1 | 0.9 | 0.8 | 1.1 |
| 2003..... | 1.5 | * | 0.2 | 0.3 | 3.6 | 1.8 | 1.6 | 1.4 | 1.1 | 1.0 | 1.0 | 1.3 |
| 2002..... | 1.5 | * | * | 0.4 | 3.8 | 2.1 | 1.6 | 1.3 | 1.2 | 1.0 | 1.0 | 0.9 |
| 2001..... | 1.4 | * | * | 0.3 | 3.3 | 1.8 | 1.5 | 1.2 | 1.0 | 1.0 | 1.1 | 0.7 |
| 2000..... | 1.3 | * | * | 0.3 | 3.4 | 1.9 | 1.4 | 1.1 | 0.9 | 0.8 | 0.9 | 1.0 |
| 1999..... | 1.3 | * | 0.2 | 0.3 | 3.4 | 1.8 | 1.4 | 1.0 | 1.0 | 0.9 | 0.8 | 1.0 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | 0.4 | 4.4 | 2.4 | 2.2 | 1.7 | 1.6 | 1.3 | 1.2 | 2.0 |
| 2003..... | 2.1 | * | * | 0.3 | 4.9 | 2.8 | 2.2 | 1.9 | 1.4 | 1.5 | 1.6 | 2.1 |
| 2002..... | 2.2 | * | * | 0.4 | 5.3 | 3.0 | 2.2 | 1.9 | 1.6 | 1.5 | 1.6 | 1.7 |
| 2001..... | 1.9 | * | * | 0.3 | 4.6 | 2.6 | 2.0 | 1.8 | 1.4 | 1.5 | 1.5 | * |
| 2000..... | 1.9 | * | * | 0.3 | 4.7 | 2.9 | 2.0 | 1.7 | 1.2 | 1.2 | 1.3 | 2.2 |
| 1999..... | 1.9 | * | * | 0.3 | 4.6 | 2.6 | 2.1 | 1.6 | 1.5 | 1.2 | 1.2 | 1.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | 0.3 | 2.1 | 0.9 | 0.9 | 0.8 | 0.5 | 0.6 | 0.6 | 0.8 |
| 2003..... | 0.9 | * | * | 0.4 | 2.2 | 0.9 | 1.1 | 0.8 | 0.7 | 0.6 | 0.6 | 1.0 |
| 2002..... | 0.9 | * | * | 0.3 | 2.3 | 1.2 | 0.9 | 0.7 | 0.8 | 0.7 | 0.6 | * |
| 2001..... | 0.8 | * | * | 0.3 | 2.0 | 0.9 | 0.9 | 0.6 | 0.6 | 0.5 | 0.8 | * |
| 2000..... | 0.7 | * | * | 0.2 | 2.1 | 0.9 | 0.8 | 0.5 | 0.5 | 0.4 | 0.6 | * |
| 1999..... | 0.8 | * | * | 0.3 | 2.2 | 1.0 | 0.8 | 0.5 | 0.6 | 0.7 | 0.5 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 1.8 | 1.7 | 1.4 | 1.4 | 1.0 | 1.4 | * | * |
| 2003..... | 1.2 | * | * | 0.4 | 1.7 | 1.8 | 1.4 | 1.2 | 1.5 | * | * | * |
| 2002..... | 1.2 | * | * | 0.4 | 2.0 | 1.5 | 1.6 | 1.2 | 1.1 | * | * | * |
| 2001..... | 1.1 | * | * | 0.4 | 1.7 | 1.8 | 1.1 | 1.2 | 1.8 | * | * | * |
| 2000..... | 1.1 | * | * | 0.4 | 2.0 | 1.4 | 1.4 | 1.1 | 1.6 | * | * | * |
| 1999..... | 1.2 | * | * | * | 2.0 | 1.5 | 1.6 | 1.3 | 1.0 | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 2.4 | 2.3 | 2.2 | 2.5 | * | * | * | * |
| 2003..... | 1.8 | * | * | * | 2.5 | 2.3 | 2.2 | 2.1 | 2.5 | * | * | * |
| 2002..... | 1.8 | * | * | * | 2.7 | 2.7 | 2.3 | 2.3 | 1.8 | * | * | * |
| 2001..... | 1.7 | * | * | * | 2.1 | 3.0 | 1.8 | 2.2 | 3.2 | * | * | * |
| 2000..... | 1.7 | * | * | * | 2.7 | 2.4 | 2.1 | 1.8 | 2.5 | * | * | * |
| 1999..... | 1.7 | * | * | * | 2.7 | 2.5 | 2.3 | 2.2 | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 1.1 | 1.1 | 0.8 | * | * | * | * | * |
| 2003..... | 0.7 | * | * | * | 1.0 | 1.4 | 0.7 | * | * | * | * | * |
| 2002..... | 0.6 | * | * | * | 1.4 | * | 0.9 | * | * | * | * | * |
| 2001..... | 0.6 | * | * | * | 1.3 | 0.8 | * | * | * | * | * | * |
| 2000..... | 0.6 | * | * | * | 1.3 | * | * | * | * | * | * | * |
| 1999..... | 0.7 | * | * | * | 1.4 | 0.7 | 1.0 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | 1.2 | 1.6 | 1.2 | 9.4 | 6.3 | 5.1 | 5.2 | 4.9 | 6.1 | 10.4 | 10.8 |
| 2003..... | 5.3 | 1.3 | 1.3 | 1.2 | 9.3 | 5.4 | 5.2 | 4.8 | 5.0 | 6.1 | 9.9 | 11.9 |
| 2002..... | 5.5 | 1.1 | 1.3 | 1.1 | 9.9 | 6.0 | 5.2 | 5.0 | 4.8 | 6.6 | 10.3 | 11.4 |
| 2001..... | 5.8 | 1.3 | 1.5 | 1.2 | 10.4 | 6.3 | 5.6 | 5.1 | 5.3 | 6.5 | 11.3 | 11.7 |
| 2000..... | 6.0 | 1.8 | 1.6 | 1.4 | 11.1 | 6.5 | 5.8 | 5.5 | 5.7 | 6.7 | 10.7 | 13.1 |
| 1999..... | 6.0 | 2.3 | 1.6 | 1.5 | 10.8 | 6.4 | 5.7 | 5.2 | 5.3 | 7.4 | 11.8 | 13.9 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 7.3 | 1.2 | 1.6 | 1.2 | 12.6 | 8.7 | 7.0 | 7.2 | 6.6 | 7.9 | 14.5 | 17.9 |
| 2003..... | 7.0 | 1.6 | 1.5 | 1.3 | 12.5 | 7.8 | 7.0 | 6.6 | 6.6 | 7.7 | 13.7 | 20.3 |
| 2002..... | 7.3 | 1.1 | 1.4 | 1.3 | 13.3 | 8.5 | 7.0 | 6.8 | 6.5 | 8.7 | 13.9 | 21.7 |
| 2001..... | 7.7 | 1.1 | 1.6 | 1.4 | 14.4 | 8.9 | 7.6 | 7.1 | 6.9 | 8.1 | 15.8 | 21.2 |
| 2000..... | 8.0 | 1.9 | 1.8 | 1.5 | 14.8 | 9.2 | 8.0 | 7.4 | 7.5 | 8.5 | 14.7 | 24.7 |
| 1999..... | 7.9 | 2.1 | 1.8 | 1.6 | 14.4 | 8.8 | 7.6 | 7.0 | 7.1 | 9.3 | 16.7 | 27.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.7 | 1.3 | 1.5 | 1.2 | 6.1 | 3.8 | 3.2 | 3.2 | 3.4 | 4.7 | 7.6 | 7.7 |
| 2003..... | 3.6 | * | 1.1 | 1.1 | 5.9 | 2.9 | 3.4 | 3.0 | 3.5 | 4.7 | 7.4 | 8.1 |
| 2002..... | 3.7 | 1.1 | 1.3 | 1.0 | 6.3 | 3.4 | 3.4 | 3.3 | 3.3 | 4.8 | 8.0 | 6.9 |
| 2001..... | 3.9 | 1.5 | 1.3 | 1.1 | 6.2 | 3.6 | 3.7 | 3.2 | 3.8 | 5.1 | 8.3 | 7.7 |
| 2000..... | 4.1 | 1.7 | 1.4 | 1.4 | 7.1 | 3.8 | 3.5 | 3.6 | 4.0 | 5.3 | 8.1 | 8.4 |
| 1999..... | 4.2 | 2.5 | 1.5 | 1.3 | 7.0 | 3.9 | 3.8 | 3.5 | 3.7 | 5.9 | 8.7 | 8.5 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.5 | * | 1.4 | 1.0 | 8.4 | 5.7 | 4.2 | 4.1 | 4.9 | 6.0 | 8.6 | * |
| 2003..... | 4.7 | * | 1.2 | 1.3 | 8.2 | 5.7 | 5.0 | 4.6 | 4.5 | 6.5 | 8.9 | * |
| 2002..... | 4.4 | * | 1.5 | 0.9 | 8.4 | 5.0 | 4.4 | 4.3 | 4.3 | 5.5 | 8.0 | * |
| 2001..... | 4.6 | * | 1.5 | 1.0 | 8.5 | 5.6 | 4.6 | 3.9 | 5.4 | 5.5 | 8.5 | * |
| 2000..... | 5.4 | * | 1.8 | 1.3 | 10.4 | 6.4 | 5.5 | 5.5 | 5.9 | 6.4 | 5.5 | * |
| 1999..... | 5.4 | 3.8 | 1.8 | 1.3 | 9.1 | 6.2 | 6.0 | 5.3 | 5.6 | 7.7 | 10.0 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.5 | * | 1.7 | 1.3 | 12.6 | 8.0 | 6.2 | 5.9 | 6.0 | 8.0 | 13.9 | * |
| 2003..... | 6.6 | * | 1.4 | 1.7 | 11.7 | 8.4 | 6.9 | 6.1 | 5.6 | 9.4 | 13.8 | * |
| 2002..... | 6.1 | * | 1.4 | 0.9 | 12.1 | 7.3 | 6.4 | 5.5 | 5.6 | 7.8 | 8.6 | * |
| 2001..... | 6.6 | * | 1.6 | 1.0 | 12.6 | 8.3 | 6.9 | 5.7 | 6.4 | 7.6 | 14.3 | * |
| 2000..... | 7.8 | * | 1.9 | 1.4 | 14.7 | 9.7 | 7.8 | 7.9 | 8.0 | 9.3 | * | * |
| 1999..... | 7.4 | * | 1.7 | 1.5 | 12.7 | 8.5 | 8.9 | 7.3 | 6.4 | 10.8 | 17.7 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | 1.2 | 0.7 | 3.5 | 2.8 | 2.1 | 2.2 | 3.9 | 4.4 | 4.9 | * |
| 2003..... | 2.6 | * | * | 1.0 | 4.0 | 2.3 | 2.8 | 3.0 | 3.6 | 4.1 | 5.5 | * |
| 2002..... | 2.5 | * | 1.7 | 1.0 | 4.1 | 2.3 | 2.1 | 3.0 | 3.2 | 3.6 | 7.6 | * |
| 2001..... | 2.4 | * | 1.4 | 1.0 | 3.9 | 2.4 | 2.2 | 2.0 | 4.5 | 3.8 | * | * |
| 2000..... | 3.0 | * | 1.6 | 1.2 | 5.2 | 2.7 | 3.0 | 3.1 | 4.1 | 4.1 | * | * |
| 1999..... | 3.2 | * | 1.9 | 1.1 | 4.9 | 3.5 | 2.9 | 3.3 | 4.8 | 5.3 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.6 | 1.1 | 1.6 | 1.2 | 9.6 | 6.5 | 5.2 | 5.3 | 4.9 | 6.1 | 10.4 | 10.9 |
| 2003..... | 5.4 | 1.1 | 1.3 | 1.2 | 9.5 | 5.3 | 5.2 | 4.8 | 5.0 | 6.0 | 10.0 | 12.0 |
| 2002..... | 5.6 | 1.2 | 1.3 | 1.2 | 10.2 | 6.2 | 5.3 | 5.1 | 4.8 | 6.6 | 10.4 | 11.5 |
| 2001..... | 5.9 | 1.3 | 1.5 | 1.3 | 10.7 | 6.4 | 5.7 | 5.2 | 5.2 | 6.5 | 11.3 | 11.8 |
| 2000..... | 6.1 | 1.8 | 1.6 | 1.4 | 11.2 | 6.5 | 5.8 | 5.4 | 5.6 | 6.7 | 10.9 | 13.3 |
| 1999..... | 6.1 | 1.9 | 1.6 | 1.5 | 11.1 | 6.4 | 5.6 | 5.1 | 5.2 | 7.4 | 11.8 | 14.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 7.4 | * | 1.6 | 1.2 | 12.5 | 8.9 | 7.1 | 7.3 | 6.7 | 7.8 | 14.5 | 18.2 |
| 2003..... | 7.1 | 1.2 | 1.5 | 1.3 | 12.6 | 7.6 | 7.1 | 6.6 | 6.7 | 7.5 | 13.6 | 20.6 |
| 2002..... | 7.5 | 1.3 | 1.4 | 1.4 | 13.6 | 8.7 | 7.0 | 6.9 | 6.5 | 8.7 | 14.0 | 21.9 |
| 2001..... | 7.9 | 1.2 | 1.6 | 1.4 | 14.7 | 9.0 | 7.6 | 7.2 | 6.9 | 8.1 | 15.9 | 21.5 |
| 2000..... | 8.1 | 1.9 | 1.8 | 1.5 | 14.8 | 9.1 | 8.0 | 7.3 | 7.4 | 8.4 | 15.1 | 25.2 |
| 1999..... | 7.9 | 1.7 | 1.8 | 1.6 | 14.7 | 8.8 | 7.4 | 6.9 | 7.0 | 9.2 | 16.5 | 27.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.9 | * | 1.5 | 1.3 | 6.6 | 4.0 | 3.4 | 3.3 | 3.3 | 4.7 | 7.7 | 7.7 |
| 2003..... | 3.7 | * | 1.1 | 1.1 | 6.2 | 3.0 | 3.5 | 3.0 | 3.5 | 4.7 | 7.5 | 8.3 |
| 2002..... | 3.9 | * | 1.2 | 1.0 | 6.8 | 3.7 | 3.6 | 3.4 | 3.3 | 4.9 | 8.0 | 7.0 |
| 2001..... | 4.1 | 1.5 | 1.3 | 1.1 | 6.6 | 3.9 | 3.9 | 3.3 | 3.8 | 5.2 | 8.4 | 7.8 |
| 2000..... | 4.3 | 1.8 | 1.4 | 1.4 | 7.4 | 4.0 | 3.6 | 3.6 | 4.0 | 5.3 | 8.2 | 8.5 |
| 1999..... | 4.3 | 2.2 | 1.4 | 1.4 | 7.4 | 4.0 | 3.9 | 3.5 | 3.6 | 5.9 | 8.8 | 8.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 5.7 | 0.9 | 1.5 | 1.1 | 10.3 | 6.6 | 5.2 | 5.1 | 4.8 | 6.0 | 10.6 | 11.6 |
| 2003..... | 5.4 | 0.9 | 1.2 | 1.2 | 10.0 | 5.3 | 5.1 | 4.5 | 4.8 | 6.0 | 10.1 | 12.2 |
| 2002..... | 5.8 | 0.9 | 1.2 | 1.1 | 11.1 | 6.2 | 5.3 | 4.9 | 4.7 | 6.7 | 10.6 | 11.9 |
| 2001..... | 6.1 | 1.1 | 1.3 | 1.2 | 11.4 | 6.5 | 5.7 | 5.1 | 5.1 | 6.5 | 11.7 | 12.3 |
| 2000..... | 6.2 | 1.3 | 1.5 | 1.4 | 12.0 | 6.3 | 5.8 | 5.2 | 5.6 | 6.5 | 11.2 | 13.7 |
| 1999..... | 6.2 | 1.7 | 1.3 | 1.5 | 11.9 | 6.4 | 5.6 | 4.9 | 5.1 | 7.4 | 12.0 | 14.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 7.4 | * | 1.4 | 1.0 | 13.4 | 8.9 | 6.9 | 6.9 | 6.4 | 7.6 | 14.7 | 19.3 |
| 2003..... | 7.1 | * | 1.5 | 1.3 | 13.0 | 7.6 | 6.9 | 6.1 | 6.4 | 7.6 | 13.8 | 20.6 |
| 2002..... | 7.6 | * | 1.2 | 1.3 | 14.6 | 8.7 | 6.9 | 6.7 | 6.3 | 8.7 | 14.0 | 22.2 |
| 2001..... | 8.0 | * | 1.4 | 1.3 | 15.5 | 9.1 | 7.4 | 7.0 | 6.5 | 8.1 | 16.1 | 22.6 |
| 2000..... | 8.1 | * | 1.7 | 1.5 | 15.9 | 8.7 | 7.9 | 6.9 | 7.2 | 8.0 | 15.3 | 26.5 |
| 1999..... | 8.0 | * | 1.4 | 1.6 | 15.6 | 8.8 | 7.4 | 6.5 | 6.8 | 9.2 | 16.4 | 28.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.1 | * | 1.5 | 1.2 | 7.1 | 4.2 | 3.4 | 3.3 | 3.4 | 4.7 | 7.9 | 8.1 |
| 2003..... | 3.8 | * | 0.9 | 1.1 | 6.8 | 3.0 | 3.4 | 2.9 | 3.3 | 4.7 | 7.6 | 8.5 |
| 2002..... | 4.1 | * | 1.1 | 1.0 | 7.4 | 3.7 | 3.6 | 3.3 | 3.3 | 4.9 | 8.3 | 7.4 |
| 2001..... | 4.2 | * | 1.2 | 1.1 | 7.2 | 3.9 | 4.0 | 3.3 | 3.7 | 5.2 | 8.7 | 8.1 |
| 2000..... | 4.4 | * | 1.2 | 1.4 | 8.0 | 3.9 | 3.7 | 3.4 | 4.1 | 5.3 | 8.5 | 8.6 |
| 1999..... | 4.5 | 1.8 | 1.3 | 1.3 | 8.1 | 4.1 | 3.9 | 3.3 | 3.5 | 5.9 | 9.0 | 8.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | * | 2.2 | 1.8 | 7.6 | 6.9 | 6.0 | 7.4 | 5.8 | 6.1 | 8.3 | * |
| 2003..... | 5.3 | * | 1.6 | 1.2 | 7.5 | 6.0 | 6.4 | 6.5 | 6.5 | 6.2 | 8.1 | 11.0 |
| 2002..... | 5.2 | * | 1.5 | 1.3 | 7.6 | 6.9 | 5.8 | 6.1 | 5.7 | 6.3 | 9.1 | 7.3 |
| 2001..... | 5.7 | * | 1.8 | 1.8 | 8.5 | 6.9 | 6.6 | 6.3 | 6.8 | 6.8 | 9.3 | 7.2 |
| 2000..... | 6.1 | 3.8 | 2.0 | 1.4 | 9.1 | 8.4 | 6.5 | 7.6 | 5.8 | 8.1 | 7.7 | 8.9 |
| 1999..... | 6.0 | * | 2.6 | 1.8 | 8.9 | 6.7 | 6.1 | 7.4 | 6.4 | 7.7 | 11.9 | 10.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 8.0 | * | 2.4 | 1.9 | 10.6 | 10.3 | 9.0 | 11.4 | 8.9 | 10.1 | 13.4 | * |
| 2003..... | 7.6 | * | 1.6 | 1.3 | 11.2 | 9.2 | 9.1 | 10.2 | 9.6 | 8.4 | 12.1 | 21.3 |
| 2002..... | 7.5 | * | 1.9 | 1.6 | 10.5 | 10.3 | 8.3 | 8.4 | 9.0 | 10.0 | 15.8 | 21.9 |
| 2001..... | 8.2 | * | 1.9 | 2.2 | 12.3 | 10.1 | 9.6 | 9.9 | 10.7 | 9.0 | 16.2 | * |
| 2000..... | 8.7 | * | 2.2 | 1.5 | 12.5 | 12.1 | 10.3 | 11.1 | 9.3 | 13.0 | 13.3 | * |
| 1999..... | 8.4 | * | 3.1 | 2.1 | 12.5 | 10.1 | 8.4 | 11.3 | 9.7 | 9.6 | 21.3 | 23.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.3 | * | 2.0 | 1.6 | 4.6 | 3.8 | 3.3 | 3.9 | 3.4 | 3.3 | 5.5 | * |
| 2003..... | 3.2 | * | * | 1.1 | 3.9 | 3.2 | 4.1 | 3.3 | 4.0 | 4.7 | 5.9 | * |
| 2002..... | 3.2 | * | * | 0.9 | 4.6 | 3.8 | 3.6 | 4.0 | 3.0 | 3.7 | 5.3 | * |
| 2001..... | 3.5 | * | * | 1.3 | 4.8 | 4.0 | 3.8 | 3.3 | 3.6 | 5.2 | 5.4 | * |
| 2000..... | 3.8 | * | 1.9 | 1.3 | 5.8 | 5.1 | 3.2 | 4.6 | 2.9 | 4.6 | 4.5 | * |
| 1999..... | 3.8 | * | 2.2 | 1.5 | 5.3 | 3.7 | 4.1 | 4.1 | 3.7 | 6.4 | 6.7 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Streetcar accidents (V82.2-V82.9)

Non-Hispanic 4/, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic 4/, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic 4/, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic white, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic white, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), showing asterisks for all data points.

Non-Hispanic white, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), showing asterisks for all data points.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | 0.1 | 0.3 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | * |
| 2003..... | 0.3 | * | * | 0.1 | 0.5 | 0.3 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 |
| 2002..... | 0.2 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | * |
| 2001..... | 0.3 | * | * | 0.2 | 0.3 | 0.2 | 0.4 | 0.4 | 0.4 | 0.5 | 0.6 | 0.5 |
| 2000..... | 0.3 | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | 0.6 | 0.7 |
| 1999..... | 0.6 | * | 0.3 | 0.3 | 0.7 | 0.5 | 0.6 | 0.6 | 0.7 | 0.6 | 1.2 | 1.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.1 | 0.5 | 0.3 | 0.4 | 0.4 | 0.5 | 0.5 | 0.6 | * |
| 2003..... | 0.5 | * | * | * | 0.7 | 0.4 | 0.4 | 0.6 | 0.6 | 0.7 | 0.9 | * |
| 2002..... | 0.3 | * | * | 0.2 | 0.2 | 0.2 | 0.4 | 0.5 | 0.4 | 0.6 | 0.8 | * |
| 2001..... | 0.5 | * | * | 0.2 | 0.4 | 0.3 | 0.6 | 0.6 | 0.6 | 0.9 | 0.8 | * |
| 2000..... | 0.5 | * | * | 0.2 | 0.6 | 0.5 | 0.6 | 0.5 | 0.6 | 0.7 | 0.9 | * |
| 1999..... | 0.8 | * | 0.4 | 0.3 | 1.0 | 0.7 | 0.8 | 1.0 | 1.0 | 0.9 | 1.6 | 2.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | 0.2 | * | 0.1 | 0.1 | 0.2 | 0.3 | * | * |
| 2003..... | 0.2 | * | * | * | 0.3 | 0.2 | 0.2 | 0.1 | 0.2 | * | 0.3 | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 2001..... | 0.2 | * | * | 0.1 | 0.1 | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | * |
| 2000..... | 0.2 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | * | * | 0.4 | * |
| 1999..... | 0.4 | * | * | 0.2 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.9 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | 0.1 | 0.3 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | * |
| 2003..... | 0.3 | * | * | 0.1 | 0.6 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.5 | 0.5 |
| 2002..... | 0.2 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | * |
| 2001..... | 0.3 | * | * | 0.1 | 0.3 | 0.2 | 0.3 | 0.4 | 0.3 | 0.5 | 0.6 | * |
| 2000..... | 0.4 | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | 0.6 | 0.7 |
| 1999..... | 0.6 | * | 0.3 | 0.3 | 0.7 | 0.5 | 0.6 | 0.6 | 0.7 | 0.6 | 1.1 | 1.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.2 | 0.4 | 0.3 | 0.4 | 0.4 | 0.5 | 0.5 | 0.7 | * |
| 2003..... | 0.5 | * | * | * | 0.8 | 0.4 | 0.4 | 0.5 | 0.6 | 0.6 | 0.9 | * |
| 2002..... | 0.3 | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.5 | 0.4 | 0.5 | 0.8 | * |
| 2001..... | 0.4 | * | * | 0.2 | 0.4 | 0.3 | 0.5 | 0.6 | 0.5 | 0.8 | 0.8 | * |
| 2000..... | 0.5 | * | * | 0.3 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | 0.7 | 1.0 | * |
| 1999..... | 0.8 | * | * | 0.3 | 1.0 | 0.7 | 0.8 | 0.9 | 1.0 | 0.9 | 1.6 | 2.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.2 | * | 0.2 | 0.1 | 0.2 | 0.3 | * | * |
| 2003..... | 0.2 | * | * | * | 0.3 | * | 0.2 | 0.2 | 0.2 | * | 0.3 | * |
| 2002..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.2 | * | * | * | * | * | 0.1 | 0.2 | * | 0.3 | 0.4 | * |
| 2000..... | 0.2 | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | * | * | 0.4 | * |
| 1999..... | 0.4 | * | * | 0.2 | 0.5 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Air and space transport accidents (V95-V97) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.1 | 0.3 | 0.3 | 0.4 | 0.5 | 0.4 | * | * |
| 2003..... | 0.3 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.6 | 0.4 | 0.2 | * |
| 2002..... | 0.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.4 | 0.2 | * |
| 2001..... | 0.3 | * | * | * | 0.1 | 0.4 | 0.4 | 0.5 | 0.4 | 0.4 | 0.2 | * |
| 2000..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.3 | 0.5 | 0.6 | 0.4 | 0.2 | * |
| 1999..... | 0.3 | * | * | 0.1 | 0.1 | 0.4 | 0.4 | 0.4 | 0.5 | 0.4 | 0.2 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.5 | 0.6 | 0.8 | 0.8 | 0.8 | * | * |
| 2003..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.6 | 0.9 | 1.1 | 0.7 | 0.4 | * |
| 2002..... | 0.5 | * | * | * | 0.2 | 0.4 | 0.6 | 0.8 | 1.0 | 0.8 | * | * |
| 2001..... | 0.5 | * | * | * | 0.2 | 0.6 | 0.7 | 0.8 | 0.8 | 0.7 | * | * |
| 2000..... | 0.5 | * | * | * | 0.3 | 0.6 | 0.6 | 0.9 | 1.0 | 0.7 | 0.5 | * |
| 1999..... | 0.5 | * | * | * | 0.2 | 0.6 | 0.6 | 0.7 | 1.0 | 0.7 | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.1 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | 0.1 | * | 0.2 | 0.2 | * | * | * |
| 1999..... | 0.1 | * | * | * | * | 0.1 | 0.2 | 0.2 | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.4 | 0.5 | 0.5 | 0.4 | * | * |
| 2003..... | 0.3 | * | * | 0.1 | 0.1 | 0.3 | 0.4 | 0.6 | 0.7 | 0.4 | 0.2 | * |
| 2002..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.4 | 0.2 | * |
| 2001..... | 0.3 | * | * | * | 0.2 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 | 0.2 | * |
| 2000..... | 0.4 | * | * | * | 0.2 | 0.5 | 0.4 | 0.6 | 0.7 | 0.4 | 0.2 | * |
| 1999..... | 0.3 | * | * | * | 0.2 | 0.4 | 0.5 | 0.5 | 0.6 | 0.5 | 0.2 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.2 | 0.6 | 0.7 | 0.9 | 0.9 | 0.9 | * | * |
| 2003..... | 0.6 | * | * | * | 0.2 | 0.4 | 0.7 | 1.1 | 1.3 | 0.8 | 0.5 | * |
| 2002..... | 0.6 | * | * | * | 0.3 | 0.4 | 0.7 | 0.9 | 1.1 | 0.9 | * | * |
| 2001..... | 0.6 | * | * | * | 0.2 | 0.8 | 0.8 | 0.9 | 0.9 | 0.8 | * | * |
| 2000..... | 0.6 | * | * | * | 0.4 | 0.8 | 0.8 | 1.0 | 1.2 | 0.7 | 0.5 | * |
| 1999..... | 0.6 | * | * | * | 0.3 | 0.7 | 0.7 | 0.8 | 1.2 | 0.8 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | 0.2 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | * | 0.1 | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | 0.1 | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | 0.2 | * | 0.2 | 0.2 | * | * | * |
| 1999..... | 0.1 | * | * | * | * | 0.2 | 0.2 | 0.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other and unspecified transport accidents and their sequelae (V98-V99,Y85) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | * |
| 2003..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | * |
| 2002..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.3 | 0.2 | 0.4 | * |
| 2001..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.2 | 0.4 | * |
| 2000..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | * |
| 1999..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 0.5 | 0.5 | * | * |
| 2003..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.7 | 0.6 | 0.4 | * |
| 2002..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.5 | 0.4 | 0.7 | * |
| 2001..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.4 | 0.4 | 0.6 | * |
| 2000..... | 0.3 | * | * | * | 0.1 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | * |
| 1999..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.4 | 0.4 | 0.5 | 0.4 | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | 0.1 | 0.1 | 0.2 | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | * | * | * |
| 2001..... | 0.1 | * | * | * | * | 0.1 | 0.1 | 0.1 | * | * | 0.3 | * |
| 2000..... | 0.1 | * | * | * | * | * | 0.2 | 0.1 | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | 0.1 | 0.1 | 0.2 | 0.2 | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.2 | 0.2 | 0.4 | 0.4 | 0.3 | 0.3 | 0.2 | * |
| 2003..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 | 0.3 | * |
| 2002..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.3 | 0.2 | 0.4 | * |
| 2001..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | * |
| 2000..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.3 | * |
| 1999..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.2 | 0.4 | 0.5 | 0.6 | 0.5 | 0.5 | * | * |
| 2003..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.7 | 0.6 | * | * |
| 2002..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.6 | 0.5 | 0.4 | 0.6 | * |
| 2001..... | 0.3 | * | * | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.4 | 0.4 | 0.6 | * |
| 2000..... | 0.3 | * | * | * | * | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | * |
| 1999..... | 0.3 | * | * | * | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.2 | 0.2 | 0.2 | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Falls (W00-W19) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.4 | 0.6 | 0.3 | 0.1 | 0.6 | 0.8 | 1.5 | 2.8 | 4.8 | 12.2 | 43.8 | 143.3 |
| 2003..... | 5.9 | * | 0.3 | 0.1 | 0.6 | 0.7 | 1.4 | 2.6 | 4.4 | 11.2 | 40.8 | 135.9 |
| 2002..... | 5.6 | * | 0.2 | 0.1 | 0.6 | 0.8 | 1.5 | 2.5 | 4.1 | 10.8 | 38.3 | 130.4 |
| 2001..... | 5.3 | 0.6 | 0.2 | 0.2 | 0.6 | 0.9 | 1.4 | 2.6 | 4.0 | 10.0 | 35.3 | 121.5 |
| 2000..... | 4.7 | * | 0.2 | 0.1 | 0.6 | 0.8 | 1.3 | 2.3 | 3.9 | 9.0 | 31.1 | 112.6 |
| 1999..... | 4.7 | * | 0.4 | 0.1 | 0.6 | 0.9 | 1.4 | 2.3 | 3.7 | 9.0 | 31.5 | 110.2 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.8 | * | 0.4 | 0.1 | 1.0 | 1.3 | 2.4 | 4.5 | 7.1 | 16.5 | 54.8 | 174.4 |
| 2003..... | 6.2 | * | 0.5 | 0.1 | 0.9 | 1.2 | 2.4 | 4.0 | 6.2 | 14.5 | 51.6 | 165.1 |
| 2002..... | 6.0 | * | 0.4 | 0.1 | 1.0 | 1.3 | 2.3 | 3.9 | 6.0 | 14.5 | 48.1 | 158.9 |
| 2001..... | 5.8 | * | 0.3 | 0.2 | 1.1 | 1.5 | 2.4 | 4.3 | 5.9 | 13.1 | 45.7 | 155.7 |
| 2000..... | 5.2 | * | 0.3 | 0.1 | 1.0 | 1.4 | 2.2 | 3.7 | 5.8 | 11.9 | 39.9 | 145.4 |
| 1999..... | 5.2 | * | 0.5 | 0.2 | 1.1 | 1.5 | 2.4 | 3.7 | 5.7 | 12.4 | 40.5 | 142.5 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 6.0 | * | * | * | 0.2 | 0.2 | 0.6 | 1.3 | 2.7 | 8.6 | 36.4 | 129.2 |
| 2003..... | 5.6 | * | * | * | 0.2 | 0.2 | 0.5 | 1.2 | 2.7 | 8.4 | 33.5 | 122.9 |
| 2002..... | 5.3 | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 2.3 | 7.7 | 31.8 | 118.1 |
| 2001..... | 4.8 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 2.2 | 7.4 | 28.5 | 107.2 |
| 2000..... | 4.3 | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 2.1 | 6.6 | 25.3 | 99.2 |
| 1999..... | 4.3 | * | * | * | 0.2 | 0.2 | 0.5 | 0.9 | 1.9 | 6.3 | 25.7 | 97.2 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | 0.6 | 1.1 | 1.8 | 2.3 | 4.9 | 10.2 | 30.8 | 87.1 |
| 2003..... | 2.1 | * | * | * | 0.5 | 0.8 | 1.4 | 2.6 | 5.0 | 9.2 | 25.4 | 80.8 |
| 2002..... | 2.1 | * | * | * | 0.7 | 0.9 | 1.6 | 2.5 | 4.4 | 10.0 | 29.2 | 71.5 |
| 2001..... | 2.0 | * | * | * | 0.7 | 1.1 | 1.8 | 2.6 | 4.2 | 9.1 | 23.6 | 70.0 |
| 2000..... | 2.0 | * | * | * | 0.7 | 1.1 | 1.3 | 2.5 | 4.3 | 8.0 | 26.5 | 81.0 |
| 1999..... | 1.9 | * | * | * | 0.7 | 1.1 | 1.4 | 2.2 | 3.8 | 8.1 | 24.2 | 86.3 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | 1.1 | 1.8 | 3.0 | 4.2 | 8.1 | 14.0 | 38.3 | 106.5 |
| 2003..... | 2.7 | * | * | * | 0.9 | 1.4 | 2.6 | 4.3 | 7.5 | 11.8 | 36.0 | 90.8 |
| 2002..... | 2.8 | * | * | * | 1.3 | 1.6 | 2.7 | 3.8 | 7.0 | 13.8 | 36.9 | 97.0 |
| 2001..... | 2.8 | * | * | * | 1.2 | 1.9 | 3.3 | 4.5 | 7.1 | 11.4 | 27.7 | 72.4 |
| 2000..... | 2.7 | * | * | * | 1.2 | 2.0 | 2.3 | 4.4 | 6.6 | 12.7 | 30.5 | 104.8 |
| 1999..... | 2.7 | * | * | * | 1.3 | 1.9 | 2.4 | 3.8 | 7.2 | 11.4 | 31.3 | 120.8 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | * | 2.0 | 7.2 | 25.7 | 77.1 |
| 2003..... | 1.5 | * | * | * | * | * | * | * | 2.8 | 7.2 | 18.2 | 75.8 |
| 2002..... | 1.4 | * | * | * | * | * | * | 1.1 | 2.1 | 7.0 | 23.9 | 58.9 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | * | 7.2 | 20.9 | 68.9 |
| 2000..... | 1.2 | * | * | * | * | * | * | * | 2.3 | 4.3 | 23.8 | 69.2 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | * | 5.5 | 19.4 | 69.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Falls (W00-W19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.0 | * | 0.3 | 0.1 | 0.6 | 0.7 | 1.4 | 2.9 | 4.8 | 12.3 | 44.5 | 145.5 |
| 2003..... | 6.5 | * | 0.3 | 0.1 | 0.6 | 0.7 | 1.4 | 2.5 | 4.3 | 11.3 | 41.5 | 137.9 |
| 2002..... | 6.2 | * | 0.2 | 0.1 | 0.6 | 0.7 | 1.5 | 2.5 | 4.0 | 10.8 | 38.6 | 132.6 |
| 2001..... | 5.7 | * | 0.2 | 0.1 | 0.6 | 0.8 | 1.4 | 2.6 | 3.9 | 10.0 | 35.8 | 123.1 |
| 2000..... | 5.1 | * | 0.2 | 0.1 | 0.6 | 0.7 | 1.3 | 2.3 | 3.9 | 9.1 | 31.2 | 113.5 |
| 1999..... | 5.1 | * | 0.4 | 0.1 | 0.6 | 0.8 | 1.4 | 2.2 | 3.7 | 9.0 | 31.7 | 110.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 7.4 | * | 0.4 | 0.1 | 0.9 | 1.2 | 2.3 | 4.4 | 6.9 | 16.7 | 55.6 | 177.2 |
| 2003..... | 6.8 | * | 0.5 | 0.1 | 0.9 | 1.1 | 2.3 | 3.9 | 6.0 | 14.6 | 52.3 | 168.3 |
| 2002..... | 6.5 | * | 0.3 | 0.1 | 0.9 | 1.3 | 2.3 | 3.9 | 5.9 | 14.4 | 48.4 | 161.3 |
| 2001..... | 6.2 | * | 0.3 | 0.2 | 1.1 | 1.4 | 2.3 | 4.2 | 5.8 | 13.1 | 46.4 | 159.0 |
| 2000..... | 5.5 | * | * | 0.2 | 1.0 | 1.2 | 2.2 | 3.6 | 5.7 | 11.9 | 40.2 | 146.5 |
| 1999..... | 5.5 | * | 0.5 | 0.2 | 1.0 | 1.3 | 2.3 | 3.6 | 5.5 | 12.4 | 40.6 | 142.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | 0.2 | 0.2 | 0.6 | 1.4 | 2.8 | 8.7 | 37.0 | 131.3 |
| 2003..... | 6.2 | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.7 | 8.5 | 34.3 | 124.6 |
| 2002..... | 5.9 | * | * | * | 0.2 | 0.2 | 0.7 | 1.2 | 2.3 | 7.7 | 32.1 | 120.2 |
| 2001..... | 5.3 | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.2 | 7.4 | 28.8 | 108.1 |
| 2000..... | 4.7 | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 2.1 | 6.8 | 25.3 | 100.1 |
| 1999..... | 4.6 | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 2.0 | 6.3 | 25.9 | 97.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 8.2 | * | 0.3 | 0.1 | 0.7 | 0.8 | 1.5 | 2.9 | 4.8 | 12.8 | 47.2 | 155.1 |
| 2003..... | 7.5 | * | 0.3 | 0.1 | 0.6 | 0.8 | 1.5 | 2.5 | 4.3 | 11.6 | 43.6 | 146.8 |
| 2002..... | 7.1 | * | * | 0.1 | 0.7 | 0.8 | 1.5 | 2.5 | 4.1 | 11.1 | 40.8 | 140.4 |
| 2001..... | 6.6 | * | 0.2 | 0.2 | 0.8 | 0.9 | 1.4 | 2.6 | 3.9 | 10.3 | 37.6 | 129.9 |
| 2000..... | 5.8 | * | * | 0.1 | 0.7 | 0.7 | 1.4 | 2.3 | 3.8 | 9.2 | 32.7 | 120.2 |
| 1999..... | 5.7 | * | 0.4 | 0.2 | 0.7 | 0.8 | 1.5 | 2.1 | 3.5 | 9.1 | 33.1 | 116.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 8.5 | * | * | * | 1.1 | 1.3 | 2.4 | 4.5 | 6.7 | 17.3 | 58.6 | 186.9 |
| 2003..... | 7.7 | * | 0.5 | 0.2 | 1.0 | 1.2 | 2.4 | 3.8 | 6.0 | 14.8 | 54.6 | 178.7 |
| 2002..... | 7.3 | * | * | * | 1.0 | 1.4 | 2.3 | 3.9 | 5.9 | 14.8 | 50.6 | 169.2 |
| 2001..... | 7.0 | * | * | 0.2 | 1.3 | 1.5 | 2.3 | 4.1 | 5.7 | 13.5 | 48.2 | 166.0 |
| 2000..... | 6.2 | * | * | 0.2 | 1.1 | 1.3 | 2.2 | 3.6 | 5.6 | 12.0 | 41.7 | 154.1 |
| 1999..... | 6.1 | * | 0.5 | 0.2 | 1.1 | 1.4 | 2.3 | 3.4 | 5.1 | 12.2 | 42.0 | 149.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 7.9 | * | * | * | 0.2 | 0.2 | 0.5 | 1.4 | 2.9 | 9.0 | 39.4 | 140.9 |
| 2003..... | 7.3 | * | * | * | 0.2 | 0.3 | 0.6 | 1.2 | 2.7 | 8.9 | 36.2 | 132.8 |
| 2002..... | 7.0 | * | * | * | 0.2 | 0.2 | 0.7 | 1.2 | 2.4 | 8.1 | 34.2 | 128.0 |
| 2001..... | 6.2 | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.2 | 7.7 | 30.6 | 114.8 |
| 2000..... | 5.5 | * | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 2.1 | 6.9 | 26.8 | 106.5 |
| 1999..... | 5.4 | * | * | * | 0.2 | 0.3 | 0.6 | 0.8 | 1.9 | 6.4 | 27.2 | 103.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Falls (W00-W19) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | * | 0.6 | 1.5 | 2.9 | 5.6 | 10.1 | 20.8 | 47.5 |
| 2003..... | 2.5 | * | * | * | * | 0.5 | 1.3 | 3.2 | 4.9 | 9.0 | 19.3 | 44.6 |
| 2002..... | 2.3 | * | * | * | * | 0.4 | 1.3 | 2.8 | 4.3 | 8.2 | 16.3 | 49.3 |
| 2001..... | 2.3 | * | * | * | * | 0.5 | 1.6 | 3.0 | 4.4 | 8.8 | 18.7 | 44.0 |
| 2000..... | 2.1 | * | * | * | * | 0.4 | 1.3 | 2.7 | 4.8 | 8.3 | 15.2 | 39.1 |
| 1999..... | 2.5 | * | * | * | 0.4 | 0.8 | 1.3 | 3.5 | 5.5 | 9.3 | 17.2 | 45.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | * | 1.1 | 2.2 | 4.8 | 10.2 | 14.5 | 29.9 | 73.2 |
| 2003..... | 3.0 | * | * | * | * | * | 2.2 | 5.2 | 7.2 | 12.6 | 24.5 | 56.5 |
| 2002..... | 3.0 | * | * | * | * | 0.8 | 2.2 | 4.7 | 6.8 | 11.9 | 24.9 | 76.6 |
| 2001..... | 3.2 | * | * | * | * | * | 2.7 | 5.4 | 6.9 | 12.6 | 29.1 | 68.1 |
| 2000..... | 2.8 | * | * | * | * | * | 2.1 | 4.6 | 8.2 | 11.4 | 21.6 | 61.1 |
| 1999..... | 3.4 | * | * | * | * | 1.4 | 2.5 | 5.9 | 8.7 | 15.0 | 24.1 | 58.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | * | * | 0.9 | 1.4 | 1.9 | 7.0 | 15.6 | 37.3 |
| 2003..... | 2.0 | * | * | * | * | * | * | 1.5 | 3.0 | 6.5 | 16.3 | 40.0 |
| 2002..... | 1.6 | * | * | * | * | * | * | 1.2 | 2.2 | 5.6 | 11.4 | 38.9 |
| 2001..... | 1.6 | * | * | * | * | * | * | * | 2.3 | 6.1 | 12.8 | 35.1 |
| 2000..... | 1.5 | * | * | * | * | * | * | 1.0 | 2.0 | 6.1 | 11.6 | 31.0 |
| 1999..... | 1.7 | * | * | * | * | * | * | 1.4 | 2.9 | 5.3 | 13.3 | 40.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Fall on same level (W00-W09,W18) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 3.8 | 15.3 | 54.2 |
| 2003..... | 2.1 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.2 | 3.3 | 14.4 | 50.9 |
| 2002..... | 2.0 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.0 | 3.1 | 12.6 | 51.3 |
| 2001..... | 1.5 | * | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.5 | 1.0 | 2.5 | 9.9 | 37.5 |
| 2000..... | 1.3 | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.8 | 2.0 | 8.0 | 33.3 |
| 1999..... | 0.9 | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.4 | 1.5 | 5.7 | 21.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.3 | * | * | * | 0.1 | 0.3 | 0.4 | 1.0 | 1.7 | 4.9 | 18.8 | 64.3 |
| 2003..... | 2.1 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.6 | 4.0 | 18.2 | 61.2 |
| 2002..... | 1.9 | * | * | * | 0.2 | 0.2 | 0.4 | 0.8 | 1.3 | 4.2 | 15.3 | 61.1 |
| 2001..... | 1.5 | * | * | * | 0.2 | 0.2 | 0.3 | 0.8 | 1.3 | 2.9 | 12.8 | 45.9 |
| 2000..... | 1.2 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 2.5 | 9.7 | 39.9 |
| 1999..... | 0.8 | * | * | * | 0.1 | * | 0.2 | 0.5 | 0.5 | 2.0 | 7.2 | 26.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.4 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.0 | 12.9 | 49.6 |
| 2003..... | 2.2 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.7 | 11.8 | 46.4 |
| 2002..... | 2.1 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.2 | 10.9 | 47.1 |
| 2001..... | 1.6 | * | * | * | * | * | 0.1 | 0.3 | 0.6 | 2.1 | 8.0 | 34.0 |
| 2000..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.6 | 1.6 | 6.8 | 30.6 |
| 1999..... | 0.9 | * | * | * | * | * | * | 0.2 | 0.3 | 1.1 | 4.8 | 19.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 4.0 | 16.1 | 58.2 |
| 2003..... | 2.5 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 3.5 | 15.3 | 54.3 |
| 2002..... | 2.4 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.1 | 3.3 | 13.3 | 54.4 |
| 2001..... | 1.8 | * | * | * | 0.1 | 0.1 | 0.2 | 0.6 | 1.0 | 2.6 | 10.4 | 39.6 |
| 2000..... | 1.5 | * | * | * | 0.1 | * | 0.2 | 0.4 | 0.7 | 2.1 | 8.4 | 35.3 |
| 1999..... | 1.0 | * | * | * | 0.1 | 0.1 | 0.1 | 0.3 | 0.3 | 1.5 | 5.9 | 22.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | * | 0.2 | 0.3 | 0.4 | 1.1 | 1.6 | 5.1 | 19.5 | 68.5 |
| 2003..... | 2.4 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.5 | 4.2 | 19.2 | 64.6 |
| 2002..... | 2.2 | * | * | * | 0.2 | 0.2 | 0.4 | 0.9 | 1.4 | 4.3 | 15.7 | 63.6 |
| 2001..... | 1.7 | * | * | * | 0.2 | 0.2 | 0.3 | 0.8 | 1.3 | 3.1 | 13.2 | 48.3 |
| 2000..... | 1.4 | * | * | * | * | * | 0.3 | 0.6 | 0.9 | 2.6 | 10.1 | 42.3 |
| 1999..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.4 | 2.0 | 7.3 | 27.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 2.8 | * | * | * | * | * | 0.2 | 0.3 | 0.9 | 3.0 | 13.7 | 53.6 |
| 2003..... | 2.6 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.8 | 12.6 | 49.7 |
| 2002..... | 2.5 | * | * | * | * | * | 0.1 | 0.3 | 0.8 | 2.4 | 11.7 | 50.4 |
| 2001..... | 1.8 | * | * | * | * | * | * | 0.3 | 0.6 | 2.2 | 8.5 | 36.0 |
| 2000..... | 1.6 | * | * | * | * | * | * | 0.2 | 0.6 | 1.7 | 7.3 | 32.5 |
| 1999..... | 1.0 | * | * | * | * | * | * | 0.2 | 0.2 | 1.1 | 5.0 | 20.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Fall from one level to another (W10-W17) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.3 | 0.3 | 0.7 | 1.1 | 1.6 | 2.8 | 6.0 | 10.8 |
| 2003..... | 1.2 | * | * | * | 0.4 | 0.4 | 0.6 | 1.0 | 1.3 | 2.8 | 6.0 | 11.4 |
| 2002..... | 1.2 | * | * | * | 0.4 | 0.5 | 0.7 | 0.9 | 1.4 | 2.8 | 6.3 | 11.2 |
| 2001..... | 1.2 | * | * | 0.1 | 0.4 | 0.5 | 0.7 | 1.1 | 1.3 | 2.7 | 6.2 | 11.5 |
| 2000..... | 1.1 | * | * | * | 0.4 | 0.5 | 0.7 | 1.0 | 1.3 | 2.4 | 5.2 | 8.8 |
| 1999..... | 1.2 | * | 0.2 | 0.1 | 0.4 | 0.5 | 0.7 | 0.9 | 1.5 | 2.6 | 5.9 | 10.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | 0.5 | 0.6 | 1.2 | 1.9 | 2.6 | 4.6 | 8.7 | 16.6 |
| 2003..... | 1.7 | * | * | * | 0.7 | 0.7 | 1.2 | 1.8 | 2.2 | 4.3 | 9.0 | 18.4 |
| 2002..... | 1.8 | * | * | * | 0.6 | 0.9 | 1.3 | 1.6 | 2.4 | 4.3 | 9.9 | 17.8 |
| 2001..... | 1.8 | * | * | * | 0.8 | 0.9 | 1.3 | 1.9 | 2.3 | 4.2 | 9.6 | 19.0 |
| 2000..... | 1.6 | * | * | * | 0.7 | 0.9 | 1.2 | 1.7 | 2.2 | 3.8 | 8.6 | 16.4 |
| 1999..... | 1.7 | * | * | * | 0.7 | 1.0 | 1.3 | 1.6 | 2.6 | 4.2 | 9.3 | 18.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 4.1 | 8.2 |
| 2003..... | 0.7 | * | * | * | 0.1 | * | 0.1 | 0.3 | 0.6 | 1.5 | 4.0 | 8.4 |
| 2002..... | 0.7 | * | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 1.6 | 3.8 | 8.3 |
| 2001..... | 0.7 | * | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.4 | 4.0 | 8.4 |
| 2000..... | 0.6 | * | * | * | 0.1 | * | 0.2 | 0.3 | 0.5 | 1.2 | 3.0 | 5.8 |
| 1999..... | 0.7 | * | * | * | * | 0.1 | 0.2 | 0.2 | 0.6 | 1.4 | 3.8 | 7.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 0.4 | 0.4 | 0.7 | 1.1 | 1.6 | 3.0 | 6.3 | 11.4 |
| 2003..... | 1.3 | * | * | * | 0.5 | 0.5 | 0.7 | 1.0 | 1.4 | 2.8 | 6.3 | 12.1 |
| 2002..... | 1.4 | * | * | * | 0.4 | 0.6 | 0.8 | 1.0 | 1.5 | 3.0 | 6.6 | 11.5 |
| 2001..... | 1.4 | * | * | * | 0.5 | 0.6 | 0.7 | 1.1 | 1.3 | 2.8 | 6.5 | 12.0 |
| 2000..... | 1.2 | * | * | * | 0.5 | 0.5 | 0.8 | 1.0 | 1.3 | 2.5 | 5.5 | 9.2 |
| 1999..... | 1.3 | * | 0.2 | * | 0.5 | 0.6 | 0.8 | 0.9 | 1.6 | 2.8 | 6.2 | 11.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.0 | * | * | * | 0.6 | 0.7 | 1.2 | 1.9 | 2.6 | 4.9 | 9.3 | 17.7 |
| 2003..... | 1.9 | * | * | * | 0.8 | 0.9 | 1.2 | 1.7 | 2.3 | 4.3 | 9.4 | 19.8 |
| 2002..... | 2.0 | * | * | * | 0.7 | 1.0 | 1.3 | 1.7 | 2.5 | 4.6 | 10.4 | 18.5 |
| 2001..... | 2.0 | * | * | * | 1.0 | 1.0 | 1.3 | 1.8 | 2.3 | 4.3 | 10.0 | 19.8 |
| 2000..... | 1.8 | * | * | * | 0.9 | 1.0 | 1.3 | 1.7 | 2.2 | 3.9 | 9.0 | 17.0 |
| 1999..... | 1.9 | * | * | * | 0.8 | 1.0 | 1.3 | 1.6 | 2.6 | 4.2 | 9.5 | 19.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 4.3 | 8.6 |
| 2003..... | 0.8 | * | * | * | 0.2 | * | 0.1 | 0.3 | 0.6 | 1.6 | 4.1 | 8.8 |
| 2002..... | 0.8 | * | * | * | 0.2 | * | 0.2 | 0.3 | 0.5 | 1.6 | 4.0 | 8.5 |
| 2001..... | 0.8 | * | * | * | * | 0.2 | 0.2 | 0.3 | 0.5 | 1.5 | 4.2 | 8.7 |
| 2000..... | 0.6 | * | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 3.2 | 6.1 |
| 1999..... | 0.8 | * | * | * | * | * | 0.2 | 0.2 | 0.7 | 1.5 | 4.0 | 7.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified fall (W19) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 1.9 | 5.7 | 22.9 | 79.1 |
| 2003..... | 2.9 | * | * | * | 0.1 | 0.1 | 0.5 | 0.9 | 1.8 | 5.1 | 20.8 | 74.2 |
| 2002..... | 2.7 | * | * | * | 0.1 | 0.1 | 0.5 | 1.0 | 1.6 | 4.8 | 19.6 | 68.8 |
| 2001..... | 2.7 | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.6 | 4.8 | 19.3 | 73.0 |
| 2000..... | 2.5 | * | * | * | 0.1 | 0.1 | 0.4 | 0.9 | 1.8 | 4.6 | 17.9 | 70.7 |
| 1999..... | 2.8 | * | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 1.8 | 4.9 | 19.9 | 78.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | 0.2 | 0.3 | 0.8 | 1.6 | 2.6 | 7.3 | 27.7 | 93.9 |
| 2003..... | 2.7 | * | * | * | 0.1 | 0.2 | 0.7 | 1.4 | 2.3 | 6.2 | 24.7 | 86.9 |
| 2002..... | 2.5 | * | * | * | 0.1 | 0.2 | 0.7 | 1.4 | 2.2 | 5.9 | 23.2 | 80.7 |
| 2001..... | 2.6 | * | * | * | 0.1 | 0.3 | 0.8 | 1.5 | 2.2 | 5.9 | 23.4 | 92.1 |
| 2000..... | 2.5 | * | * | * | 0.2 | 0.2 | 0.7 | 1.4 | 2.6 | 5.6 | 21.7 | 89.4 |
| 1999..... | 2.8 | * | * | * | 0.2 | 0.3 | 0.9 | 1.6 | 2.6 | 6.3 | 24.2 | 98.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | * | * | 0.2 | 0.7 | 1.2 | 4.3 | 19.7 | 72.4 |
| 2003..... | 3.0 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.2 | 18.1 | 68.5 |
| 2002..... | 2.8 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 3.9 | 17.2 | 63.6 |
| 2001..... | 2.7 | * | * | * | * | * | 0.2 | 0.4 | 1.1 | 3.9 | 16.6 | 64.9 |
| 2000..... | 2.6 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.8 | 15.5 | 63.1 |
| 1999..... | 2.8 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 3.8 | 17.1 | 70.4 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.7 | 1.0 | 1.8 | 5.6 | 16.4 | 43.3 |
| 2003..... | 0.8 | * | * | * | * | * | 0.5 | 1.0 | 1.8 | 3.9 | 11.9 | 37.3 |
| 2002..... | 0.8 | * | * | * | * | * | 0.5 | 0.9 | 1.7 | 4.6 | 15.5 | 31.6 |
| 2001..... | 0.8 | * | * | * | * | * | 0.6 | 0.9 | 1.5 | 4.0 | 11.7 | 40.0 |
| 2000..... | 0.8 | * | * | * | * | * | 0.4 | 0.9 | 1.7 | 3.9 | 14.6 | 49.1 |
| 1999..... | 1.0 | * | * | * | * | * | 0.5 | 1.1 | 2.0 | 5.1 | 14.6 | 58.9 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.9 | 1.7 | 2.7 | 7.4 | 19.7 | 40.6 |
| 2003..... | 0.9 | * | * | * | * | * | 0.9 | 1.5 | 2.3 | 5.0 | 15.7 | 43.1 |
| 2002..... | 0.9 | * | * | * | * | * | 0.9 | 1.4 | 2.2 | 5.4 | 20.5 | 35.1 |
| 2001..... | 0.9 | * | * | * | * | * | 1.2 | 1.4 | 2.6 | 4.0 | 10.7 | 41.9 |
| 2000..... | 1.0 | * | * | * | * | * | 0.8 | 1.5 | 2.7 | 6.3 | 15.2 | 60.5 |
| 1999..... | 1.2 | * | * | * | * | * | 0.8 | 1.9 | 4.0 | 6.8 | 18.8 | 82.0 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | * | * | * | 4.1 | 14.1 | 44.7 |
| 2003..... | 0.6 | * | * | * | * | * | * | * | * | 3.1 | 9.2 | 34.4 |
| 2002..... | 0.7 | * | * | * | * | * | * | * | * | 4.0 | 12.1 | 29.8 |
| 2001..... | 0.7 | * | * | * | * | * | * | * | * | 3.9 | 12.4 | 39.1 |
| 2000..... | 0.7 | * | * | * | * | * | * | * | * | * | 14.2 | 43.5 |
| 1999..... | 0.7 | * | * | * | * | * | * | * | * | 3.8 | 11.8 | 47.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified fall (W19) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | * | * | * | 0.1 | 0.2 | 0.5 | 1.1 | 1.9 | 5.7 | 23.2 | 80.5 |
| 2003..... | 3.2 | * | * | * | 0.1 | 0.1 | 0.5 | 0.9 | 1.8 | 5.2 | 21.2 | 75.6 |
| 2002..... | 2.9 | * | * | * | 0.1 | 0.1 | 0.4 | 1.0 | 1.6 | 4.8 | 19.7 | 70.0 |
| 2001..... | 3.0 | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 1.6 | 4.8 | 19.7 | 74.1 |
| 2000..... | 2.8 | * | * | * | 0.1 | 0.1 | 0.4 | 0.9 | 1.8 | 4.6 | 18.1 | 71.4 |
| 1999..... | 3.0 | * | * | * | 0.1 | 0.2 | 0.6 | 1.0 | 1.7 | 4.9 | 20.0 | 78.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 3.4 | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 2.6 | 7.3 | 28.1 | 96.3 |
| 2003..... | 3.0 | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.3 | 6.3 | 25.2 | 88.7 |
| 2002..... | 2.8 | * | * | * | * | 0.2 | 0.6 | 1.4 | 2.2 | 6.0 | 23.2 | 82.4 |
| 2001..... | 2.9 | * | * | * | 0.1 | 0.3 | 0.7 | 1.5 | 2.2 | 6.0 | 24.0 | 94.1 |
| 2000..... | 2.7 | * | * | * | 0.1 | 0.2 | 0.6 | 1.4 | 2.6 | 5.6 | 21.9 | 90.2 |
| 1999..... | 3.0 | * | * | * | 0.2 | 0.3 | 0.9 | 1.5 | 2.5 | 6.2 | 24.2 | 98.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | * | * | 0.2 | 0.7 | 1.2 | 4.4 | 20.0 | 73.5 |
| 2003..... | 3.3 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.3 | 18.5 | 69.8 |
| 2002..... | 3.1 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 3.9 | 17.4 | 64.7 |
| 2001..... | 3.0 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 3.9 | 16.8 | 65.7 |
| 2000..... | 2.9 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.9 | 15.5 | 63.7 |
| 1999..... | 3.1 | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 3.8 | 17.3 | 70.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.1 | * | * | * | 0.2 | 0.2 | 0.5 | 1.1 | 1.9 | 5.8 | 24.8 | 85.5 |
| 2003..... | 3.7 | * | * | * | * | 0.2 | 0.5 | 0.9 | 1.8 | 5.3 | 22.1 | 80.4 |
| 2002..... | 3.4 | * | * | * | 0.1 | 0.1 | 0.4 | 0.9 | 1.5 | 4.9 | 20.9 | 74.5 |
| 2001..... | 3.4 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 1.6 | 4.9 | 20.6 | 78.3 |
| 2000..... | 3.2 | * | * | * | 0.1 | 0.1 | 0.4 | 0.9 | 1.7 | 4.7 | 18.8 | 75.6 |
| 1999..... | 3.4 | * | * | * | 0.1 | 0.2 | 0.6 | 0.9 | 1.5 | 4.8 | 21.0 | 82.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.9 | * | * | * | 0.3 | 0.3 | 0.8 | 1.5 | 2.4 | 7.3 | 29.8 | 100.7 |
| 2003..... | 3.4 | * | * | * | * | 0.2 | 0.8 | 1.3 | 2.3 | 6.4 | 26.0 | 94.3 |
| 2002..... | 3.1 | * | * | * | * | * | 0.6 | 1.3 | 2.0 | 5.9 | 24.4 | 87.1 |
| 2001..... | 3.3 | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 2.1 | 6.0 | 24.9 | 97.9 |
| 2000..... | 3.0 | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.4 | 5.6 | 22.6 | 94.9 |
| 1999..... | 3.3 | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 2.1 | 6.1 | 25.2 | 102.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 4.3 | * | * | * | * | * | 0.2 | 0.8 | 1.3 | 4.5 | 21.4 | 78.8 |
| 2003..... | 3.9 | * | * | * | * | * | 0.2 | 0.5 | 1.3 | 4.5 | 19.4 | 74.3 |
| 2002..... | 3.7 | * | * | * | * | * | 0.3 | 0.6 | 1.0 | 4.0 | 18.6 | 69.1 |
| 2001..... | 3.6 | * | * | * | * | * | 0.2 | 0.5 | 1.1 | 4.0 | 17.8 | 70.1 |
| 2000..... | 3.3 | * | * | * | * | * | 0.2 | 0.5 | 1.0 | 3.9 | 16.3 | 67.9 |
| 1999..... | 3.6 | * | * | * | * | * | 0.3 | 0.4 | 0.9 | 3.8 | 18.2 | 75.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Unspecified fall (W19) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | 0.5 | 1.3 | 2.3 | 5.5 | 9.8 | 30.8 |
| 2003..... | 1.3 | * | * | * | * | * | 0.4 | 1.4 | 2.3 | 5.1 | 12.1 | 28.9 |
| 2002..... | 1.2 | * | * | * | * | * | 0.5 | 1.5 | 2.4 | 4.5 | 8.9 | 25.4 |
| 2001..... | 1.2 | * | * | * | * | * | 0.7 | 1.2 | 2.1 | 4.8 | 11.0 | 26.2 |
| 2000..... | 1.2 | * | * | * | * | * | 0.7 | 1.2 | 2.4 | 4.9 | 10.2 | 26.1 |
| 1999..... | 1.5 | * | * | * | * | * | 0.6 | 1.8 | 3.5 | 5.7 | 11.0 | 33.3 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | * | * | * | 1.9 | 4.5 | 8.0 | 14.3 | 49.8 |
| 2003..... | 1.4 | * | * | * | * | * | * | 2.3 | 3.2 | 6.4 | 13.9 | 36.2 |
| 2002..... | 1.5 | * | * | * | * | * | 0.8 | 2.7 | 4.0 | 6.4 | 10.8 | 38.3 |
| 2001..... | 1.4 | * | * | * | * | * | 1.0 | 2.0 | 3.0 | 6.2 | 15.9 | 46.2 |
| 2000..... | 1.5 | * | * | * | * | * | 1.1 | 1.9 | 4.2 | 6.1 | 14.5 | 38.7 |
| 1999..... | 1.9 | * | * | * | * | * | 1.1 | 3.1 | 5.5 | 8.4 | 14.7 | 46.4 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | * | * | 3.8 | 7.2 | 23.3 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | 1.5 | 4.2 | 11.1 | 26.0 |
| 2002..... | 0.8 | * | * | * | * | * | * | * | * | 3.1 | 7.8 | 20.5 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 3.8 | 8.3 | 18.8 |
| 2000..... | 0.9 | * | * | * | * | * | * | * | * | 4.1 | 7.8 | 21.4 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | 1.9 | 3.7 | 8.9 | 28.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental discharge of firearms (W32-W34) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | 0.1 | 0.4 | 0.3 | 0.2 | 0.3 | 0.1 | 0.2 | 0.2 | * |
| 2003..... | 0.3 | * | * | 0.1 | 0.5 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | * |
| 2002..... | 0.3 | * | * | 0.1 | 0.5 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| 2001..... | 0.3 | * | * | 0.2 | 0.5 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | * |
| 2000..... | 0.3 | * | * | 0.2 | 0.5 | 0.4 | 0.4 | 0.2 | 0.2 | 0.2 | 0.2 | * |
| 1999..... | 0.3 | * | * | 0.2 | 0.7 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.2 | 0.8 | 0.5 | 0.4 | 0.5 | 0.2 | 0.4 | 0.5 | * |
| 2003..... | 0.5 | * | * | 0.2 | 0.9 | 0.6 | 0.5 | 0.4 | 0.5 | 0.4 | 0.7 | * |
| 2002..... | 0.5 | * | * | 0.2 | 1.0 | 0.7 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 | * |
| 2001..... | 0.5 | * | * | 0.3 | 0.9 | 0.5 | 0.6 | 0.5 | 0.5 | 0.3 | 0.7 | * |
| 2000..... | 0.5 | * | * | 0.3 | 1.0 | 0.6 | 0.6 | 0.4 | 0.4 | 0.4 | 0.5 | * |
| 1999..... | 0.5 | * | * | 0.3 | 1.2 | 0.6 | 0.5 | 0.5 | 0.4 | 0.4 | 0.4 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | 0.1 | 0.1 | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | 0.1 | 0.1 | 0.1 | * | * | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | 0.1 | 0.4 | 0.2 | 0.3 | 0.3 | 0.1 | 0.2 | 0.2 | * |
| 2003..... | 0.3 | * | * | 0.1 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | * |
| 2002..... | 0.3 | * | * | 0.1 | 0.5 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| 2001..... | 0.3 | * | * | 0.2 | 0.5 | 0.3 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | * |
| 2000..... | 0.3 | * | * | 0.2 | 0.5 | 0.3 | 0.4 | 0.2 | 0.2 | 0.2 | 0.2 | * |
| 1999..... | 0.3 | * | * | 0.2 | 0.6 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | 0.2 | 0.7 | 0.4 | 0.4 | 0.5 | 0.2 | 0.4 | * | * |
| 2003..... | 0.5 | * | * | 0.2 | 0.8 | 0.5 | 0.6 | 0.4 | 0.5 | 0.4 | 0.6 | * |
| 2002..... | 0.5 | * | * | 0.2 | 0.8 | 0.6 | 0.5 | 0.5 | 0.4 | 0.5 | 0.5 | * |
| 2001..... | 0.5 | * | * | 0.3 | 0.8 | 0.6 | 0.6 | 0.5 | 0.5 | 0.3 | 0.7 | * |
| 2000..... | 0.5 | * | * | 0.3 | 0.8 | 0.5 | 0.6 | 0.4 | 0.4 | 0.4 | 0.5 | * |
| 1999..... | 0.5 | * | * | 0.3 | 1.0 | 0.6 | 0.5 | 0.5 | 0.4 | 0.4 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental drowning and submersion (W65-W74) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | 1.4 | 2.8 | 0.7 | 1.4 | 1.0 | 1.0 | 1.1 | 1.0 | 1.0 | 1.2 | 1.6 |
| 2003..... | 1.1 | 1.5 | 3.0 | 0.7 | 1.3 | 0.8 | 1.1 | 1.0 | 1.0 | 1.1 | 1.3 | 1.8 |
| 2002..... | 1.2 | 1.4 | 3.1 | 0.9 | 1.4 | 1.1 | 1.2 | 0.9 | 0.8 | 0.9 | 1.3 | 1.7 |
| 2001..... | 1.1 | 1.6 | 3.1 | 0.9 | 1.5 | 0.9 | 1.0 | 0.9 | 0.8 | 0.9 | 1.3 | 1.7 |
| 2000..... | 1.2 | 2.1 | 3.4 | 1.0 | 1.5 | 1.1 | 1.1 | 1.0 | 0.9 | 1.0 | 1.3 | 1.5 |
| 1999..... | 1.2 | 1.7 | 3.3 | 0.9 | 1.5 | 1.1 | 1.1 | 1.0 | 0.9 | 1.0 | 1.3 | 1.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.8 | 1.8 | 3.4 | 1.0 | 2.4 | 1.6 | 1.7 | 1.8 | 1.5 | 1.7 | 2.2 | 2.7 |
| 2003..... | 1.8 | 1.7 | 3.7 | 1.0 | 2.4 | 1.5 | 1.8 | 1.6 | 1.6 | 1.8 | 2.1 | 3.4 |
| 2002..... | 1.9 | 1.6 | 4.1 | 1.3 | 2.6 | 1.8 | 2.0 | 1.5 | 1.3 | 1.6 | 2.0 | 3.6 |
| 2001..... | 1.8 | 1.3 | 3.8 | 1.2 | 2.6 | 1.6 | 1.7 | 1.6 | 1.3 | 1.5 | 2.1 | 3.4 |
| 2000..... | 2.0 | 2.4 | 4.3 | 1.4 | 2.7 | 1.8 | 1.9 | 1.6 | 1.6 | 1.6 | 1.8 | 2.8 |
| 1999..... | 2.0 | 1.7 | 4.2 | 1.4 | 2.7 | 1.9 | 1.8 | 1.7 | 1.5 | 1.6 | 2.3 | 2.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | 2.2 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.5 | 0.6 | 1.1 |
| 2003..... | 0.5 | * | 2.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.4 | 0.4 | 0.5 | 0.7 | 1.1 |
| 2002..... | 0.5 | * | 2.1 | 0.4 | 0.3 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 | 0.8 | 0.9 |
| 2001..... | 0.5 | 1.9 | 2.4 | 0.5 | 0.3 | 0.2 | 0.4 | 0.3 | 0.4 | 0.4 | 0.7 | 1.0 |
| 2000..... | 0.5 | 1.9 | 2.4 | 0.5 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.5 | 0.9 | 1.0 |
| 1999..... | 0.5 | 1.7 | 2.4 | 0.5 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | 0.6 | 0.6 | 1.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | 1.5 | 3.2 | 0.5 | 1.2 | 0.9 | 1.0 | 1.1 | 0.9 | 1.0 | 1.2 | 1.5 |
| 2003..... | 1.1 | 1.4 | 3.3 | 0.5 | 1.2 | 0.8 | 1.1 | 1.0 | 0.9 | 1.1 | 1.3 | 1.7 |
| 2002..... | 1.1 | 1.4 | 3.3 | 0.5 | 1.2 | 1.0 | 1.2 | 0.9 | 0.8 | 0.9 | 1.2 | 1.6 |
| 2001..... | 1.1 | 1.5 | 3.5 | 0.5 | 1.3 | 0.8 | 1.0 | 0.9 | 0.8 | 0.9 | 1.2 | 1.7 |
| 2000..... | 1.1 | 1.7 | 3.7 | 0.7 | 1.3 | 1.0 | 1.1 | 0.9 | 0.9 | 1.0 | 1.3 | 1.4 |
| 1999..... | 1.1 | 1.3 | 3.6 | 0.5 | 1.3 | 1.0 | 1.1 | 1.0 | 0.9 | 1.0 | 1.2 | 1.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.7 | 1.8 | 3.8 | 0.7 | 2.1 | 1.5 | 1.6 | 1.7 | 1.4 | 1.6 | 2.2 | 2.6 |
| 2003..... | 1.7 | * | 4.0 | 0.7 | 2.1 | 1.3 | 1.8 | 1.5 | 1.5 | 1.8 | 2.1 | 3.4 |
| 2002..... | 1.7 | * | 4.3 | 0.7 | 2.2 | 1.7 | 1.9 | 1.4 | 1.2 | 1.5 | 2.0 | 3.7 |
| 2001..... | 1.6 | * | 4.3 | 0.7 | 2.2 | 1.4 | 1.5 | 1.5 | 1.2 | 1.5 | 2.1 | 3.5 |
| 2000..... | 1.8 | 1.9 | 4.7 | 1.0 | 2.2 | 1.7 | 1.8 | 1.5 | 1.6 | 1.5 | 1.8 | 2.7 |
| 1999..... | 1.8 | * | 4.5 | 0.7 | 2.2 | 1.6 | 1.8 | 1.7 | 1.5 | 1.4 | 2.2 | 2.4 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | 2.5 | 0.2 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 | 0.5 | 0.5 | 1.0 |
| 2003..... | 0.5 | * | 2.7 | 0.3 | 0.2 | 0.2 | 0.3 | 0.4 | 0.4 | 0.5 | 0.7 | 0.9 |
| 2002..... | 0.5 | * | 2.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.4 | 0.3 | 0.7 | 0.7 |
| 2001..... | 0.5 | 1.7 | 2.6 | 0.4 | 0.3 | 0.2 | 0.4 | 0.3 | 0.4 | 0.3 | 0.7 | 1.0 |
| 2000..... | 0.5 | * | 2.6 | 0.4 | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.5 | 0.9 | 0.9 |
| 1999..... | 0.5 | * | 2.6 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.4 | 0.6 | 0.6 | 1.3 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.7 | 1.5 | 0.4 | 0.1 | 0.2 | 0.2 | 0.4 | 0.8 | 1.4 | 3.0 | 9.9 | 30.6 |
| 2003..... | 1.6 | 1.8 | 0.5 | 0.1 | 0.1 | 0.3 | 0.5 | 0.8 | 1.4 | 3.2 | 9.4 | 28.1 |
| 2002..... | 1.6 | 1.6 | 0.4 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.3 | 3.2 | 9.6 | 28.9 |
| 2001..... | 1.6 | 1.7 | 0.4 | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 1.3 | 3.0 | 9.6 | 30.5 |
| 2000..... | 1.7 | 2.2 | 0.6 | * | 0.1 | 0.3 | 0.4 | 0.8 | 1.4 | 3.2 | 10.1 | 32.8 |
| 1999..... | 1.5 | 1.8 | 0.6 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 | 3.1 | 8.9 | 30.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.7 | 2.2 | 0.5 | * | 0.2 | 0.3 | 0.5 | 1.0 | 1.7 | 3.6 | 12.8 | 39.8 |
| 2003..... | 1.7 | 1.8 | 0.7 | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.8 | 3.8 | 12.5 | 38.5 |
| 2002..... | 1.7 | 2.1 | 0.5 | * | 0.2 | 0.3 | 0.5 | 0.9 | 1.7 | 4.0 | 12.2 | 37.6 |
| 2001..... | 1.7 | 2.2 | 0.4 | * | 0.2 | 0.2 | 0.5 | 1.1 | 1.6 | 3.6 | 12.5 | 40.4 |
| 2000..... | 1.7 | 2.5 | 0.7 | * | 0.2 | 0.3 | 0.5 | 1.1 | 1.7 | 3.7 | 13.3 | 41.2 |
| 1999..... | 1.5 | 2.0 | 0.7 | 0.1 | 0.1 | 0.3 | 0.5 | 0.9 | 1.3 | 3.8 | 11.1 | 39.5 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.6 | * | 0.3 | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.1 | 2.4 | 8.1 | 26.5 |
| 2003..... | 1.5 | 1.7 | 0.4 | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.0 | 2.7 | 7.3 | 23.6 |
| 2002..... | 1.5 | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 2.5 | 7.9 | 25.2 |
| 2001..... | 1.6 | * | 0.4 | * | 0.1 | 0.1 | 0.3 | 0.7 | 1.1 | 2.5 | 7.7 | 26.4 |
| 2000..... | 1.7 | 1.9 | 0.5 | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.7 | 8.1 | 29.4 |
| 1999..... | 1.5 | 1.6 | 0.5 | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 2.6 | 7.4 | 26.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | 1.5 | 0.3 | * | 0.2 | 0.2 | 0.5 | 0.9 | 1.3 | 2.7 | 9.8 | 30.9 |
| 2003..... | 1.8 | 1.6 | 0.5 | * | 0.2 | 0.2 | 0.5 | 0.8 | 1.3 | 3.1 | 9.4 | 28.6 |
| 2002..... | 1.7 | 1.4 | 0.3 | * | 0.1 | 0.3 | 0.4 | 0.7 | 1.2 | 2.9 | 9.5 | 28.9 |
| 2001..... | 1.7 | 1.4 | 0.3 | 0.1 | 0.2 | 0.2 | 0.4 | 0.9 | 1.2 | 2.8 | 9.2 | 30.2 |
| 2000..... | 1.8 | 1.8 | 0.5 | * | 0.1 | 0.2 | 0.4 | 0.8 | 1.3 | 3.0 | 10.0 | 32.7 |
| 1999..... | 1.6 | 1.5 | 0.4 | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.0 | 2.9 | 8.5 | 30.5 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.9 | 2.3 | * | * | 0.2 | 0.3 | 0.5 | 1.0 | 1.7 | 3.3 | 12.7 | 40.0 |
| 2003..... | 1.8 | * | 0.6 | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.7 | 3.7 | 12.4 | 39.3 |
| 2002..... | 1.7 | 2.0 | * | * | 0.2 | 0.3 | 0.5 | 0.8 | 1.6 | 3.6 | 11.9 | 37.8 |
| 2001..... | 1.7 | 2.0 | * | * | 0.2 | 0.2 | 0.5 | 1.0 | 1.5 | 3.4 | 11.7 | 40.0 |
| 2000..... | 1.8 | 2.1 | 0.6 | * | * | 0.3 | 0.6 | 1.0 | 1.5 | 3.4 | 13.0 | 40.4 |
| 1999..... | 1.6 | 1.8 | 0.5 | * | * | 0.2 | 0.5 | 0.8 | 1.1 | 3.5 | 10.6 | 39.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | * | 0.2 | 0.5 | 0.7 | 1.0 | 2.3 | 7.9 | 26.7 |
| 2003..... | 1.7 | * | * | * | * | * | 0.4 | 0.7 | 1.0 | 2.5 | 7.3 | 24.0 |
| 2002..... | 1.6 | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.9 | 2.2 | 7.9 | 25.0 |
| 2001..... | 1.7 | * | * | * | * | * | 0.3 | 0.8 | 1.0 | 2.3 | 7.5 | 26.1 |
| 2000..... | 1.8 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.6 | 8.1 | 29.6 |
| 1999..... | 1.6 | * | * | * | * | * | 0.3 | 0.5 | 0.9 | 2.4 | 7.1 | 26.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 0.8 | * | * | * | 0.4 | 0.9 | 1.9 | 5.3 | 13.5 | 31.1 |
| 2003..... | 1.3 | * | * | * | * | 0.4 | 0.6 | 1.0 | 2.1 | 5.0 | 10.6 | 24.4 |
| 2002..... | 1.5 | * | 0.9 | * | * | * | 0.4 | 1.1 | 2.1 | 6.6 | 13.2 | 33.6 |
| 2001..... | 1.6 | * | * | * | * | * | 0.6 | 1.2 | 2.6 | 5.5 | 14.6 | 37.1 |
| 2000..... | 1.5 | 4.0 | 1.0 | * | * | 0.4 | 0.4 | 1.2 | 2.2 | 6.0 | 13.1 | 35.6 |
| 1999..... | 1.5 | * | 1.1 | * | * | 0.4 | 0.4 | 1.1 | 2.5 | 5.5 | 14.7 | 33.6 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | * | * | * | 1.1 | 2.3 | 7.1 | 17.4 | 43.7 |
| 2003..... | 1.5 | * | * | * | * | * | 0.8 | 1.4 | 2.9 | 5.6 | 16.2 | 29.8 |
| 2002..... | 1.6 | * | * | * | * | * | * | 1.4 | 2.7 | 8.6 | 19.6 | 42.7 |
| 2001..... | 1.8 | * | * | * | * | * | * | 1.6 | 3.6 | 7.0 | 23.7 | 50.8 |
| 2000..... | 1.7 | * | * | * | * | * | * | 1.8 | 2.9 | 7.5 | 20.1 | 51.7 |
| 1999..... | 1.7 | * | * | * | * | * | * | 1.4 | 3.3 | 6.9 | 20.6 | 44.0 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | * | * | * | * | * | * | 0.8 | 1.5 | 4.0 | 11.3 | 26.1 |
| 2003..... | 1.2 | * | * | * | * | * | * | * | 1.5 | 4.6 | 7.5 | 22.3 |
| 2002..... | 1.4 | * | * | * | * | * | * | * | 1.6 | 5.1 | 9.6 | 30.1 |
| 2001..... | 1.3 | * | * | * | * | * | * | * | 1.7 | 4.4 | 9.4 | 32.1 |
| 2000..... | 1.4 | * | * | * | * | * | * | * | 1.7 | 5.0 | 9.2 | 29.7 |
| 1999..... | 1.4 | * | * | * | * | * | * | * | 1.9 | 4.6 | 11.4 | 29.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other accidental and unspecified threats to breathing (W75-W77,W81-W84) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | 18.8 | 0.4 | 0.2 | 0.2 | 0.2 | 0.4 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 |
| 2003..... | 0.5 | 16.1 | 0.5 | 0.2 | 0.2 | 0.3 | 0.4 | 0.3 | 0.2 | 0.2 | 0.3 | 0.8 |
| 2002..... | 0.5 | 16.2 | 0.5 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | 0.4 | 1.1 |
| 2001..... | 0.5 | 15.2 | 0.5 | 0.2 | 0.2 | 0.3 | 0.4 | 0.3 | 0.2 | 0.3 | 0.4 | 0.9 |
| 2000..... | 0.5 | 13.3 | 0.4 | 0.2 | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.5 | 1.2 |
| 1999..... | 0.6 | 11.8 | 0.5 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 | 0.3 | 0.6 | 1.6 | 4.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.7 | 20.4 | 0.5 | 0.4 | 0.4 | 0.4 | 0.6 | 0.5 | 0.3 | 0.3 | * | * |
| 2003..... | 0.7 | 19.0 | 0.6 | 0.3 | 0.4 | 0.5 | 0.6 | 0.5 | 0.4 | 0.3 | * | * |
| 2002..... | 0.7 | 17.5 | 0.6 | 0.3 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 | * |
| 2001..... | 0.7 | 17.0 | 0.6 | 0.3 | 0.4 | 0.5 | 0.7 | 0.6 | 0.3 | 0.4 | 0.5 | * |
| 2000..... | 0.7 | 15.1 | 0.5 | 0.4 | 0.5 | 0.5 | 0.7 | 0.5 | 0.5 | 0.4 | 0.6 | * |
| 1999..... | 0.8 | 14.1 | 0.5 | 0.4 | 0.5 | 0.5 | 0.6 | 0.6 | 0.5 | 0.8 | 1.9 | 5.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 17.2 | * | * | * | 0.1 | 0.1 | 0.1 | * | * | * | * |
| 2003..... | 0.3 | 13.1 | 0.4 | * | * | * | 0.1 | * | * | * | 0.3 | 0.8 |
| 2002..... | 0.3 | 14.8 | 0.4 | * | * | * | 0.1 | * | * | * | * | 1.1 |
| 2001..... | 0.3 | 13.3 | 0.4 | * | * | * | 0.1 | * | * | 0.2 | 0.4 | 0.9 |
| 2000..... | 0.3 | 11.6 | * | * | * | * | 0.1 | 0.1 | * | * | 0.4 | 1.1 |
| 1999..... | 0.4 | 9.4 | 0.5 | * | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 1.4 | 4.5 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | 15.4 | 0.4 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 | 0.2 | 0.2 | 0.3 | 0.6 |
| 2003..... | 0.4 | 12.9 | 0.6 | 0.2 | 0.3 | 0.3 | 0.4 | 0.3 | 0.2 | 0.2 | 0.3 | 0.8 |
| 2002..... | 0.5 | 12.1 | 0.5 | 0.2 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 1.0 |
| 2001..... | 0.5 | 11.8 | 0.4 | 0.3 | 0.3 | 0.3 | 0.5 | 0.3 | 0.2 | 0.3 | 0.5 | 0.9 |
| 2000..... | 0.5 | 10.6 | 0.5 | 0.3 | 0.3 | 0.3 | 0.4 | 0.3 | 0.4 | 0.3 | 0.5 | 1.2 |
| 1999..... | 0.6 | 9.3 | 0.5 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.3 | 0.6 | 1.5 | 4.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.7 | 17.7 | 0.6 | 0.4 | 0.5 | 0.4 | 0.7 | 0.6 | 0.3 | 0.3 | * | * |
| 2003..... | 0.7 | 15.4 | 0.7 | 0.3 | 0.5 | 0.6 | 0.6 | 0.5 | 0.4 | 0.3 | * | * |
| 2002..... | 0.7 | 13.7 | 0.6 | 0.4 | 0.6 | 0.6 | 0.6 | 0.6 | 0.4 | 0.4 | 0.5 | * |
| 2001..... | 0.7 | 13.0 | 0.5 | 0.4 | 0.5 | 0.5 | 0.8 | 0.6 | 0.3 | 0.5 | 0.5 | * |
| 2000..... | 0.7 | 12.4 | 0.6 | 0.4 | 0.5 | 0.6 | 0.7 | 0.5 | 0.6 | 0.4 | 0.6 | * |
| 1999..... | 0.8 | 10.8 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.7 | 0.5 | 0.8 | 1.8 | 5.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 13.1 | * | * | * | * | * | 0.2 | * | * | * | * |
| 2003..... | 0.3 | 10.3 | * | * | * | * | 0.2 | * | * | * | 0.3 | 0.8 |
| 2002..... | 0.3 | 10.3 | 0.4 | * | * | * | 0.1 | * | * | * | * | 1.1 |
| 2001..... | 0.3 | 10.5 | * | * | * | * | 0.1 | * | * | * | 0.4 | 0.9 |
| 2000..... | 0.3 | 8.7 | * | * | * | * | * | * | * | * | 0.4 | 1.1 |
| 1999..... | 0.4 | 7.6 | 0.5 | * | * | * | * | 0.1 | * | 0.5 | 1.3 | 4.6 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Accidental exposure to radiation (W88-W91)

Non-Hispanic 4/, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk.

Non-Hispanic 4/, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk.

Non-Hispanic 4/, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk.

Non-Hispanic white, both sexes

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk.

Non-Hispanic white, male

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk.

Non-Hispanic white, female

Table with 13 columns (years 1999-2004) and 13 rows (ages 1-85+), all cells contain an asterisk.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental exposure to smoke, fire and flames (X00-X09) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | 0.7 | 1.6 | 0.7 | 0.5 | 0.7 | 0.8 | 1.3 | 1.5 | 1.9 | 3.5 | 5.6 |
| 2003..... | 1.3 | 0.8 | 1.6 | 0.6 | 0.5 | 0.7 | 1.0 | 1.3 | 1.5 | 2.4 | 3.6 | 5.4 |
| 2002..... | 1.2 | 1.0 | 1.5 | 0.7 | 0.5 | 0.8 | 0.9 | 1.2 | 1.3 | 2.1 | 3.4 | 4.8 |
| 2001..... | 1.3 | 1.3 | 1.7 | 0.7 | 0.6 | 0.7 | 1.0 | 1.1 | 1.5 | 2.3 | 3.4 | 5.7 |
| 2000..... | 1.3 | 1.0 | 2.2 | 0.7 | 0.5 | 0.7 | 0.9 | 1.2 | 1.5 | 2.2 | 3.8 | 6.3 |
| 1999..... | 1.3 | 1.2 | 2.1 | 0.7 | 0.5 | 0.8 | 1.0 | 1.0 | 1.4 | 2.3 | 4.0 | 6.0 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | 1.7 | 0.8 | 0.6 | 0.9 | 1.0 | 1.7 | 1.9 | 2.5 | 4.5 | 7.3 |
| 2003..... | 1.5 | * | 1.9 | 0.6 | 0.6 | 0.8 | 1.3 | 1.7 | 1.9 | 3.0 | 4.5 | 7.3 |
| 2002..... | 1.5 | * | 1.7 | 0.8 | 0.7 | 1.0 | 1.2 | 1.5 | 1.7 | 2.9 | 4.6 | 7.0 |
| 2001..... | 1.6 | 1.5 | 2.0 | 0.8 | 0.8 | 0.9 | 1.5 | 1.6 | 2.0 | 2.8 | 4.2 | 8.3 |
| 2000..... | 1.6 | * | 2.6 | 0.8 | 0.7 | 0.9 | 1.3 | 1.7 | 2.1 | 2.8 | 5.0 | 9.3 |
| 1999..... | 1.6 | * | 2.3 | 0.8 | 0.7 | 1.1 | 1.4 | 1.4 | 1.7 | 3.0 | 5.1 | 9.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | 1.5 | 0.6 | 0.4 | 0.4 | 0.5 | 0.8 | 1.1 | 1.4 | 2.9 | 4.9 |
| 2003..... | 1.0 | 1.3 | 1.3 | 0.5 | 0.4 | 0.6 | 0.7 | 1.0 | 1.0 | 1.8 | 3.1 | 4.5 |
| 2002..... | 0.9 | * | 1.2 | 0.6 | 0.3 | 0.5 | 0.5 | 0.8 | 0.9 | 1.4 | 2.6 | 3.8 |
| 2001..... | 1.0 | * | 1.3 | 0.5 | 0.4 | 0.5 | 0.6 | 0.7 | 1.1 | 1.9 | 2.9 | 4.6 |
| 2000..... | 1.0 | * | 1.7 | 0.7 | 0.4 | 0.5 | 0.6 | 0.7 | 1.0 | 1.7 | 3.1 | 5.0 |
| 1999..... | 1.0 | * | 1.8 | 0.6 | 0.4 | 0.5 | 0.6 | 0.6 | 1.1 | 1.8 | 3.3 | 4.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | 1.3 | 0.5 | 0.5 | 0.6 | 0.7 | 1.1 | 1.3 | 1.6 | 3.0 | 4.9 |
| 2003..... | 1.2 | * | 1.3 | 0.4 | 0.5 | 0.7 | 1.0 | 1.2 | 1.3 | 2.1 | 3.1 | 4.7 |
| 2002..... | 1.0 | * | 1.0 | 0.4 | 0.4 | 0.7 | 0.8 | 1.1 | 1.1 | 1.8 | 3.0 | 3.9 |
| 2001..... | 1.1 | 1.1 | 1.3 | 0.5 | 0.5 | 0.6 | 0.9 | 1.0 | 1.3 | 2.0 | 2.9 | 4.7 |
| 2000..... | 1.1 | * | 1.5 | 0.5 | 0.5 | 0.6 | 0.8 | 1.1 | 1.3 | 1.8 | 3.1 | 5.1 |
| 1999..... | 1.1 | 0.9 | 1.5 | 0.5 | 0.5 | 0.8 | 0.9 | 0.9 | 1.2 | 1.9 | 3.2 | 4.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.3 | * | 1.4 | 0.6 | 0.6 | 0.9 | 0.9 | 1.5 | 1.6 | 2.0 | 3.9 | 6.5 |
| 2003..... | 1.4 | * | 1.6 | 0.4 | 0.7 | 0.8 | 1.3 | 1.5 | 1.7 | 2.6 | 3.6 | 6.5 |
| 2002..... | 1.3 | * | 1.1 | 0.6 | 0.6 | 0.9 | 1.1 | 1.4 | 1.4 | 2.4 | 4.2 | 5.7 |
| 2001..... | 1.4 | * | 1.7 | 0.6 | 0.7 | 0.8 | 1.3 | 1.3 | 1.7 | 2.3 | 3.5 | 6.9 |
| 2000..... | 1.4 | * | 1.8 | 0.6 | 0.6 | 0.8 | 1.1 | 1.6 | 1.7 | 2.3 | 4.1 | 8.1 |
| 1999..... | 1.4 | * | 1.7 | 0.6 | 0.7 | 1.1 | 1.3 | 1.2 | 1.5 | 2.3 | 4.2 | 7.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | 1.2 | 0.4 | 0.4 | 0.4 | 0.5 | 0.8 | 1.0 | 1.2 | 2.4 | 4.2 |
| 2003..... | 1.0 | * | 1.0 | 0.4 | 0.4 | 0.6 | 0.6 | 0.9 | 0.9 | 1.6 | 2.8 | 3.9 |
| 2002..... | 0.8 | * | 0.8 | 0.3 | 0.3 | 0.5 | 0.5 | 0.7 | 0.8 | 1.3 | 2.2 | 3.2 |
| 2001..... | 0.9 | * | 0.9 | 0.4 | 0.3 | 0.4 | 0.6 | 0.6 | 1.0 | 1.7 | 2.5 | 3.8 |
| 2000..... | 0.9 | * | 1.1 | 0.5 | 0.3 | 0.4 | 0.5 | 0.7 | 0.8 | 1.5 | 2.4 | 3.9 |
| 1999..... | 0.8 | * | 1.3 | 0.5 | 0.3 | 0.4 | 0.5 | 0.6 | 0.8 | 1.5 | 2.6 | 3.5 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental exposure to smoke, fire and flames (X00-X09) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | * | 2.9 | 1.7 | 0.6 | 1.0 | 1.2 | 2.5 | 3.5 | 5.0 | 9.7 | 15.3 |
| 2003..... | 2.1 | * | 3.1 | 1.5 | 0.6 | 0.9 | 1.2 | 2.6 | 2.9 | 5.4 | 10.3 | 14.9 |
| 2002..... | 2.1 | * | 3.7 | 1.5 | 0.8 | 1.2 | 1.1 | 2.2 | 3.3 | 5.0 | 8.7 | 16.0 |
| 2001..... | 2.3 | * | 3.4 | 1.3 | 0.7 | 1.2 | 1.8 | 2.4 | 3.6 | 6.1 | 9.9 | 17.8 |
| 2000..... | 2.6 | * | 5.1 | 1.6 | 0.7 | 1.2 | 1.7 | 2.4 | 4.2 | 6.1 | 13.0 | 20.7 |
| 1999..... | 2.6 | * | 4.7 | 1.6 | 0.7 | 1.0 | 1.8 | 2.2 | 3.4 | 7.0 | 13.5 | 22.0 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | 3.0 | 2.0 | 0.7 | 1.3 | 1.7 | 3.9 | 4.5 | 7.7 | 13.4 | * |
| 2003..... | 2.5 | * | 3.4 | 1.5 | * | 1.1 | 1.5 | 3.4 | 4.4 | 7.7 | 16.2 | * |
| 2002..... | 2.6 | * | 4.3 | 1.6 | 1.0 | 1.4 | 1.7 | 3.1 | 4.6 | 7.8 | 11.1 | 26.3 |
| 2001..... | 3.0 | * | 3.7 | 1.6 | 0.8 | 1.4 | 2.8 | 3.9 | 5.1 | 9.0 | 13.2 | 26.6 |
| 2000..... | 3.2 | * | 5.8 | 1.7 | 0.9 | 1.6 | 2.4 | 3.5 | 6.3 | 9.0 | 17.9 | 28.2 |
| 1999..... | 3.1 | * | 5.3 | 1.8 | 0.9 | 1.2 | 2.7 | 3.3 | 4.3 | 10.4 | 18.4 | 29.7 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | 2.9 | 1.3 | * | * | 0.7 | 1.3 | 2.6 | 3.0 | 7.6 | 13.7 |
| 2003..... | 1.8 | * | 2.8 | 1.4 | 0.7 | * | 0.9 | 2.0 | 1.7 | 3.8 | 7.0 | 13.2 |
| 2002..... | 1.6 | * | 3.0 | 1.3 | * | 1.0 | * | 1.4 | 2.2 | 3.1 | 7.3 | 12.1 |
| 2001..... | 1.8 | * | 3.1 | 1.0 | * | 1.1 | 1.0 | 1.1 | 2.4 | 4.0 | 8.1 | 14.5 |
| 2000..... | 2.0 | * | 4.3 | 1.4 | * | 0.9 | 1.2 | 1.4 | 2.5 | 4.0 | 10.2 | 17.9 |
| 1999..... | 2.1 | * | 4.1 | 1.4 | * | 0.8 | 1.1 | 1.3 | 2.8 | 4.7 | 10.7 | 19.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Earthquake, volcanic eruption, avalanched, landslide and other earth movements (X34-X36)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, both sexes.

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, male.

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, female.

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, both sexes.

Non-Hispanic white, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, male.

Non-Hispanic white, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental poisoning and exposure to noxious substances (X40-X49) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.1 | * | * | 0.1 | 5.4 | 9.1 | 14.6 | 14.5 | 5.4 | 2.3 | 2.4 | 3.4 |
| 2003..... | 6.7 | 0.5 | 0.3 | 0.1 | 4.9 | 8.6 | 14.0 | 13.3 | 4.9 | 2.2 | 2.2 | 3.5 |
| 2002..... | 6.1 | 0.6 | 0.2 | 0.1 | 4.1 | 7.8 | 13.4 | 11.7 | 4.1 | 2.1 | 2.3 | 3.9 |
| 2001..... | 4.9 | * | 0.2 | 0.1 | 3.4 | 6.3 | 11.2 | 9.1 | 3.2 | 1.6 | 2.2 | 3.5 |
| 2000..... | 4.5 | * | 0.2 | 0.1 | 3.0 | 6.0 | 10.3 | 8.1 | 2.8 | 1.5 | 2.0 | 4.3 |
| 1999..... | 4.4 | * | 0.2 | 0.1 | 2.5 | 5.9 | 10.1 | 7.8 | 2.8 | 1.6 | 2.2 | 3.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 9.6 | * | * | 0.1 | 8.2 | 13.1 | 19.0 | 19.1 | 6.6 | 2.6 | 2.7 | 4.2 |
| 2003..... | 9.2 | * | 0.3 | 0.1 | 7.4 | 12.4 | 18.8 | 18.0 | 6.3 | 2.7 | 2.8 | 4.0 |
| 2002..... | 8.5 | 1.0 | * | 0.1 | 6.3 | 11.5 | 18.4 | 15.9 | 5.4 | 2.5 | 2.7 | 5.1 |
| 2001..... | 7.1 | * | 0.3 | 0.1 | 5.3 | 9.3 | 15.7 | 13.0 | 4.1 | 2.0 | 2.7 | 3.8 |
| 2000..... | 6.6 | * | 0.3 | 0.1 | 4.6 | 9.0 | 15.1 | 12.0 | 3.4 | 1.8 | 2.1 | 6.2 |
| 1999..... | 6.5 | * | * | 0.1 | 3.8 | 9.0 | 14.9 | 11.7 | 3.9 | 2.1 | 2.4 | 4.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.7 | * | * | 0.1 | 2.4 | 4.9 | 10.3 | 10.0 | 4.3 | 2.0 | 2.2 | 3.0 |
| 2003..... | 4.3 | * | 0.3 | 0.2 | 2.2 | 4.7 | 9.3 | 8.8 | 3.7 | 1.8 | 1.8 | 3.3 |
| 2002..... | 3.7 | * | * | * | 1.9 | 4.0 | 8.4 | 7.6 | 3.0 | 1.7 | 2.1 | 3.4 |
| 2001..... | 2.9 | * | * | 0.1 | 1.4 | 3.3 | 6.7 | 5.2 | 2.3 | 1.2 | 1.9 | 3.3 |
| 2000..... | 2.5 | * | * | * | 1.3 | 2.8 | 5.6 | 4.4 | 2.3 | 1.3 | 1.9 | 3.6 |
| 1999..... | 2.3 | * | * | * | 1.1 | 2.6 | 5.3 | 4.0 | 1.8 | 1.3 | 2.0 | 3.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | 2.8 | 4.9 | 9.2 | 9.7 | 5.4 | 2.4 | * | * |
| 2003..... | 4.5 | * | * | * | 3.1 | 5.7 | 10.3 | 10.5 | 4.4 | 1.8 | * | * |
| 2002..... | 4.3 | * | * | * | 2.9 | 5.0 | 10.1 | 10.1 | 4.0 | 2.2 | * | * |
| 2001..... | 3.6 | * | * | * | 2.2 | 4.9 | 8.7 | 7.7 | 3.5 | * | * | * |
| 2000..... | 3.7 | * | * | * | 2.0 | 5.3 | 9.6 | 8.0 | 3.1 | * | * | * |
| 1999..... | 4.4 | * | * | * | 2.3 | 6.1 | 11.9 | 9.8 | 3.5 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 6.2 | * | * | * | 4.2 | 7.6 | 13.5 | 14.2 | 8.4 | * | * | * |
| 2003..... | 6.9 | * | * | * | 4.8 | 8.6 | 15.6 | 16.4 | 7.4 | * | * | * |
| 2002..... | 6.6 | * | * | * | 4.4 | 7.8 | 15.4 | 15.8 | 6.4 | * | * | * |
| 2001..... | 5.6 | * | * | * | 3.1 | 7.4 | 13.6 | 12.9 | 5.8 | * | * | * |
| 2000..... | 6.0 | * | * | * | 3.2 | 8.4 | 15.6 | 13.6 | 4.4 | * | * | * |
| 1999..... | 7.3 | * | * | * | 3.6 | 9.8 | 19.4 | 16.4 | 6.6 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.9 | * | * | * | 1.3 | 1.6 | 4.4 | 5.1 | 2.6 | * | * | * |
| 2003..... | 1.9 | * | * | * | 1.2 | 2.2 | 4.5 | 4.6 | * | * | * | * |
| 2002..... | 1.7 | * | * | * | 1.1 | 1.6 | 4.3 | 4.4 | * | * | * | * |
| 2001..... | 1.5 | * | * | * | 1.1 | 1.9 | 3.4 | 2.6 | * | * | * | * |
| 2000..... | 1.3 | * | * | * | * | 1.6 | 3.1 | 2.5 | * | * | * | * |
| 1999..... | 1.4 | * | * | * | 0.8 | 1.7 | 3.9 | 3.3 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental poisoning and exposure to noxious substances (X40-X49) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.6 | * | * | 0.1 | 5.9 | 10.1 | 15.4 | 14.9 | 5.4 | 2.3 | 2.4 | 3.4 |
| 2003..... | 7.0 | * | 0.3 | 0.2 | 5.2 | 9.2 | 14.5 | 13.5 | 4.9 | 2.2 | 2.2 | 3.5 |
| 2002..... | 6.3 | 0.6 | 0.2 | 0.1 | 4.4 | 8.4 | 13.8 | 11.8 | 4.1 | 2.1 | 2.3 | 4.1 |
| 2001..... | 5.1 | * | 0.2 | 0.1 | 3.6 | 6.6 | 11.4 | 9.1 | 3.1 | 1.6 | 2.3 | 3.6 |
| 2000..... | 4.6 | * | 0.2 | 0.1 | 3.1 | 6.0 | 10.3 | 8.0 | 2.8 | 1.5 | 2.0 | 4.4 |
| 1999..... | 4.3 | * | 0.2 | 0.1 | 2.5 | 5.8 | 9.7 | 7.5 | 2.7 | 1.6 | 2.2 | 3.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 10.2 | * | * | 0.1 | 9.1 | 14.5 | 19.8 | 19.5 | 6.4 | 2.6 | 2.7 | 4.2 |
| 2003..... | 9.5 | * | 0.3 | 0.2 | 7.9 | 13.3 | 19.2 | 17.9 | 6.1 | 2.6 | 2.8 | 4.0 |
| 2002..... | 8.8 | * | * | 0.1 | 6.6 | 12.3 | 18.7 | 15.9 | 5.2 | 2.4 | 2.6 | 5.3 |
| 2001..... | 7.2 | * | * | 0.1 | 5.7 | 9.6 | 15.9 | 12.9 | 3.9 | 2.0 | 2.8 | 3.9 |
| 2000..... | 6.6 | * | * | 0.1 | 4.9 | 9.1 | 14.8 | 11.7 | 3.3 | 1.7 | 2.1 | 6.2 |
| 1999..... | 6.3 | * | * | 0.1 | 3.8 | 8.8 | 14.2 | 11.1 | 3.6 | 2.0 | 2.5 | 4.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.1 | * | * | 0.1 | 2.7 | 5.6 | 11.1 | 10.5 | 4.4 | 2.0 | 2.2 | 3.1 |
| 2003..... | 4.6 | * | * | 0.2 | 2.4 | 5.2 | 9.9 | 9.2 | 3.8 | 1.9 | 1.8 | 3.3 |
| 2002..... | 4.0 | * | * | * | 2.0 | 4.5 | 8.9 | 7.9 | 3.0 | 1.8 | 2.1 | 3.5 |
| 2001..... | 3.1 | * | * | * | 1.5 | 3.5 | 7.1 | 5.4 | 2.3 | 1.2 | 1.9 | 3.4 |
| 2000..... | 2.7 | * | * | * | 1.4 | 3.0 | 5.9 | 4.4 | 2.3 | 1.3 | 2.0 | 3.6 |
| 1999..... | 2.4 | * | * | * | 1.2 | 2.8 | 5.4 | 4.0 | 1.8 | 1.3 | 2.0 | 3.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 8.2 | * | * | 0.1 | 7.3 | 11.8 | 16.8 | 15.0 | 5.0 | 2.2 | 2.4 | 3.5 |
| 2003..... | 7.4 | * | * | 0.1 | 6.4 | 10.6 | 15.5 | 13.4 | 4.8 | 2.0 | 2.3 | 3.5 |
| 2002..... | 6.6 | * | * | 0.1 | 5.3 | 9.6 | 14.5 | 11.4 | 3.7 | 2.0 | 2.3 | 4.1 |
| 2001..... | 5.2 | * | * | 0.1 | 4.3 | 7.3 | 11.7 | 8.3 | 2.7 | 1.5 | 2.1 | 3.4 |
| 2000..... | 4.6 | * | * | 0.1 | 3.7 | 6.5 | 10.4 | 7.2 | 2.5 | 1.4 | 2.1 | 4.1 |
| 1999..... | 4.2 | * | * | 0.1 | 2.8 | 6.0 | 9.6 | 6.7 | 2.4 | 1.5 | 2.3 | 3.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 10.9 | * | * | 0.2 | 11.2 | 17.1 | 21.4 | 19.3 | 5.4 | 2.3 | 2.5 | 4.3 |
| 2003..... | 10.1 | * | * | * | 9.8 | 15.3 | 20.3 | 17.5 | 5.6 | 2.1 | 2.9 | 3.6 |
| 2002..... | 9.1 | * | * | 0.2 | 8.0 | 13.9 | 19.6 | 15.0 | 4.3 | 2.2 | 2.5 | 5.4 |
| 2001..... | 7.3 | * | * | * | 6.7 | 10.7 | 16.3 | 11.5 | 3.1 | 1.9 | 2.5 | 3.8 |
| 2000..... | 6.6 | * | * | * | 5.8 | 9.8 | 14.9 | 10.3 | 2.7 | 1.5 | 2.2 | 5.8 |
| 1999..... | 6.1 | * | * | * | 4.3 | 9.2 | 14.0 | 9.6 | 2.9 | 1.8 | 2.5 | 4.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.5 | * | * | * | 3.2 | 6.5 | 12.1 | 10.8 | 4.5 | 2.1 | 2.3 | 3.1 |
| 2003..... | 4.9 | * | * | * | 2.9 | 5.9 | 10.6 | 9.4 | 4.0 | 1.9 | 1.9 | 3.5 |
| 2002..... | 4.3 | * | * | * | 2.5 | 5.2 | 9.3 | 7.8 | 3.1 | 1.9 | 2.2 | 3.5 |
| 2001..... | 3.1 | * | * | * | 1.7 | 3.8 | 7.1 | 5.2 | 2.4 | 1.2 | 1.9 | 3.2 |
| 2000..... | 2.7 | * | * | * | 1.5 | 3.2 | 5.8 | 4.2 | 2.3 | 1.4 | 2.0 | 3.4 |
| 1999..... | 2.4 | * | * | * | 1.3 | 2.8 | 5.1 | 3.8 | 1.9 | 1.2 | 2.1 | 2.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental poisoning and exposure to noxious substances (X40-X49) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 6.7 | * | * | * | 1.8 | 5.5 | 13.4 | 19.2 | 10.0 | 3.8 | 3.4 | * |
| 2003..... | 6.4 | * | * | * | 1.4 | 5.8 | 13.9 | 18.3 | 7.6 | 4.1 | * | * |
| 2002..... | 6.4 | * | * | * | 1.5 | 5.8 | 14.1 | 18.3 | 8.6 | 2.5 | 2.8 | * |
| 2001..... | 6.0 | * | * | * | 1.6 | 5.3 | 13.4 | 17.0 | 6.4 | 2.4 | 3.7 | 6.6 |
| 2000..... | 5.7 | * | * | * | 1.4 | 5.7 | 13.2 | 15.7 | 5.5 | 2.5 | * | 8.3 |
| 1999..... | 5.8 | * | * | * | 1.7 | 6.3 | 13.2 | 15.6 | 6.0 | 3.0 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 9.1 | * | * | * | 2.6 | 7.8 | 17.0 | 27.5 | 16.5 | 6.4 | * | * |
| 2003..... | 9.1 | * | * | * | 2.1 | 8.3 | 19.2 | 26.6 | 12.9 | 7.5 | * | * |
| 2002..... | 9.3 | * | * | * | 2.2 | 8.7 | 19.8 | 27.1 | 15.2 | 4.5 | * | * |
| 2001..... | 8.6 | * | * | * | 2.3 | 7.6 | 18.2 | 27.0 | 11.0 | 3.7 | 6.0 | * |
| 2000..... | 8.4 | * | * | * | 1.8 | 8.5 | 19.0 | 25.3 | 9.5 | 4.3 | * | * |
| 1999..... | 8.8 | * | * | * | 2.8 | 9.3 | 19.5 | 25.9 | 11.0 | 4.4 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.4 | * | * | * | 1.1 | 3.4 | 10.1 | 12.0 | 4.7 | * | * | * |
| 2003..... | 4.1 | * | * | * | 0.7 | 3.6 | 9.3 | 11.1 | 3.4 | * | * | * |
| 2002..... | 3.8 | * | * | * | 0.7 | 3.2 | 9.1 | 10.8 | 3.3 | * | * | * |
| 2001..... | 3.6 | * | * | * | 0.9 | 3.2 | 9.2 | 8.4 | 2.6 | * | * | * |
| 2000..... | 3.2 | * | * | * | 1.0 | 3.1 | 8.0 | 7.5 | 2.4 | * | * | * |
| 1999..... | 3.1 | * | * | * | * | 3.5 | 7.7 | 6.7 | 1.9 | 2.1 | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental poisoning by and exposure to drugs and other biological substances (X40-X44) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.8 | * | * | 0.1 | 5.2 | 8.8 | 14.1 | 13.9 | 5.0 | 1.8 | 1.8 | 2.5 |
| 2003..... | 6.3 | * | 0.2 | 0.1 | 4.6 | 8.3 | 13.4 | 12.7 | 4.4 | 1.7 | 1.7 | 2.8 |
| 2002..... | 5.7 | * | * | * | 3.9 | 7.4 | 12.8 | 11.1 | 3.6 | 1.6 | 1.7 | 3.1 |
| 2001..... | 4.6 | * | * | 0.1 | 3.1 | 6.0 | 10.6 | 8.5 | 2.7 | 1.3 | 1.6 | 2.5 |
| 2000..... | 4.2 | * | * | * | 2.7 | 5.6 | 9.8 | 7.6 | 2.3 | 1.1 | 1.4 | 3.3 |
| 1999..... | 4.0 | * | * | * | 2.2 | 5.6 | 9.5 | 7.2 | 2.3 | 1.2 | 1.6 | 2.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 9.1 | * | * | * | 7.8 | 12.6 | 18.1 | 18.1 | 6.0 | 1.9 | 1.8 | 2.6 |
| 2003..... | 8.6 | * | * | * | 6.9 | 11.9 | 17.9 | 16.9 | 5.4 | 2.0 | 2.1 | 2.7 |
| 2002..... | 7.9 | * | * | * | 5.9 | 10.9 | 17.5 | 15.0 | 4.5 | 1.7 | 1.8 | 3.7 |
| 2001..... | 6.5 | * | * | * | 4.8 | 8.9 | 14.8 | 12.2 | 3.5 | 1.5 | 1.7 | 1.8 |
| 2000..... | 6.1 | * | * | * | 4.1 | 8.5 | 14.3 | 11.2 | 2.7 | 1.3 | 1.3 | 4.4 |
| 1999..... | 5.9 | * | * | * | 3.3 | 8.5 | 14.0 | 10.9 | 3.2 | 1.5 | 1.5 | 3.3 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.5 | * | * | * | 2.3 | 4.8 | 10.1 | 9.7 | 4.1 | 1.7 | 1.8 | 2.5 |
| 2003..... | 4.1 | * | * | * | 2.1 | 4.6 | 9.0 | 8.6 | 3.4 | 1.5 | 1.5 | 2.9 |
| 2002..... | 3.5 | * | * | * | 1.8 | 3.9 | 8.1 | 7.3 | 2.7 | 1.4 | 1.6 | 2.8 |
| 2001..... | 2.7 | * | * | * | 1.3 | 3.1 | 6.4 | 5.0 | 2.0 | 1.1 | 1.4 | 2.7 |
| 2000..... | 2.3 | * | * | * | 1.2 | 2.7 | 5.5 | 4.1 | 1.9 | 1.0 | 1.5 | 2.8 |
| 1999..... | 2.1 | * | * | * | 1.0 | 2.5 | 5.1 | 3.7 | 1.5 | 1.0 | 1.6 | 2.6 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | * | * | * | 2.4 | 4.4 | 8.7 | 9.0 | 4.9 | 1.9 | * | * |
| 2003..... | 4.2 | * | * | * | 2.6 | 5.2 | 9.8 | 10.2 | 4.2 | * | * | * |
| 2002..... | 3.9 | * | * | * | 2.5 | 4.4 | 9.5 | 9.6 | 3.6 | 1.8 | * | * |
| 2001..... | 3.3 | * | * | * | 1.7 | 4.5 | 8.2 | 7.4 | 3.1 | * | * | * |
| 2000..... | 3.4 | * | * | * | 1.6 | 4.9 | 9.2 | 7.6 | 2.8 | * | * | * |
| 1999..... | 4.1 | * | * | * | 1.9 | 5.8 | 11.2 | 9.2 | 3.2 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 5.6 | * | * | * | 3.4 | 6.9 | 12.7 | 13.1 | 7.8 | * | * | * |
| 2003..... | 6.4 | * | * | * | 4.0 | 7.9 | 14.7 | 15.9 | 7.1 | * | * | * |
| 2002..... | 6.0 | * | * | * | 3.9 | 6.9 | 14.5 | 15.0 | 5.7 | * | * | * |
| 2001..... | 5.1 | * | * | * | 2.5 | 6.8 | 12.7 | 12.3 | 5.3 | * | * | * |
| 2000..... | 5.6 | * | * | * | 2.6 | 7.9 | 14.9 | 13.0 | 4.0 | * | * | * |
| 1999..... | 6.8 | * | * | * | 3.1 | 9.4 | 18.2 | 15.5 | 6.0 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.8 | * | * | * | 1.2 | 1.5 | 4.2 | 4.8 | 2.1 | * | * | * |
| 2003..... | 1.8 | * | * | * | 1.0 | 2.1 | 4.3 | 4.6 | * | * | * | * |
| 2002..... | 1.6 | * | * | * | 1.0 | 1.5 | 4.0 | 4.3 | * | * | * | * |
| 2001..... | 1.3 | * | * | * | 0.8 | 1.7 | 3.4 | 2.5 | * | * | * | * |
| 2000..... | 1.2 | * | * | * | * | 1.6 | 3.1 | 2.3 | * | * | * | * |
| 1999..... | 1.3 | * | * | * | * | 1.7 | 3.8 | 3.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental poisoning by and exposure to drugs and other biological substances (X40-X44) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 7.2 | * | * | 0.1 | 5.7 | 9.8 | 14.9 | 14.3 | 5.0 | 1.8 | 1.8 | 2.5 |
| 2003..... | 6.6 | * | 0.2 | 0.1 | 5.0 | 8.9 | 13.9 | 12.8 | 4.4 | 1.7 | 1.7 | 2.8 |
| 2002..... | 5.9 | * | * | * | 4.1 | 8.1 | 13.2 | 11.2 | 3.6 | 1.5 | 1.6 | 3.2 |
| 2001..... | 4.7 | * | * | * | 3.3 | 6.3 | 10.8 | 8.5 | 2.7 | 1.2 | 1.6 | 2.5 |
| 2000..... | 4.2 | * | * | * | 2.9 | 5.7 | 9.8 | 7.4 | 2.2 | 1.1 | 1.5 | 3.3 |
| 1999..... | 3.9 | * | * | * | 2.2 | 5.5 | 9.2 | 7.0 | 2.2 | 1.2 | 1.6 | 2.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 9.6 | * | * | * | 8.8 | 14.1 | 19.0 | 18.6 | 5.8 | 1.9 | 1.8 | 2.6 |
| 2003..... | 8.9 | * | * | * | 7.5 | 12.8 | 18.3 | 16.9 | 5.2 | 1.9 | 2.1 | 2.7 |
| 2002..... | 8.1 | * | * | * | 6.3 | 11.8 | 17.8 | 14.9 | 4.4 | 1.6 | 1.7 | 3.8 |
| 2001..... | 6.7 | * | * | * | 5.2 | 9.3 | 15.0 | 12.0 | 3.3 | 1.4 | 1.7 | 4.9 |
| 2000..... | 6.1 | * | * | * | 4.4 | 8.5 | 14.0 | 10.8 | 2.5 | 1.3 | 1.3 | 4.3 |
| 1999..... | 5.7 | * | * | * | 3.4 | 8.3 | 13.3 | 10.3 | 2.9 | 1.4 | 1.5 | 3.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 4.9 | * | * | * | 2.6 | 5.5 | 10.9 | 10.2 | 4.2 | 1.7 | 1.8 | 2.5 |
| 2003..... | 4.4 | * | * | * | 2.3 | 5.1 | 9.7 | 9.0 | 3.6 | 1.6 | 1.5 | 2.9 |
| 2002..... | 3.8 | * | * | * | 1.9 | 4.4 | 8.7 | 7.6 | 2.8 | 1.4 | 1.6 | 2.9 |
| 2001..... | 2.9 | * | * | * | 1.4 | 3.4 | 6.8 | 5.2 | 2.1 | 1.1 | 1.5 | 2.8 |
| 2000..... | 2.5 | * | * | * | 1.3 | 2.9 | 5.7 | 4.2 | 1.9 | 1.0 | 1.5 | 2.9 |
| 1999..... | 2.2 | * | * | * | 1.1 | 2.6 | 5.2 | 3.7 | 1.6 | 1.0 | 1.6 | 2.6 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 7.8 | * | * | * | 7.1 | 11.5 | 16.2 | 14.4 | 4.6 | 1.7 | 1.8 | 2.6 |
| 2003..... | 7.0 | * | * | * | 6.2 | 10.3 | 14.8 | 12.8 | 4.2 | 1.5 | 1.8 | 2.8 |
| 2002..... | 6.2 | * | * | * | 5.0 | 9.3 | 13.9 | 10.8 | 3.2 | 1.5 | 1.6 | 3.2 |
| 2001..... | 4.8 | * | * | * | 4.0 | 7.0 | 11.1 | 7.8 | 2.3 | 1.2 | 1.6 | 2.4 |
| 2000..... | 4.2 | * | * | * | 3.4 | 6.2 | 9.9 | 6.7 | 2.0 | 1.1 | 1.5 | 3.1 |
| 1999..... | 3.9 | * | * | * | 2.5 | 5.7 | 9.0 | 6.2 | 1.9 | 1.1 | 1.6 | 2.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 10.3 | * | * | * | 10.8 | 16.6 | 20.5 | 18.3 | 4.9 | 1.6 | 1.7 | 2.7 |
| 2003..... | 9.5 | * | * | * | 9.3 | 14.8 | 19.3 | 16.4 | 4.7 | 1.5 | 2.1 | 2.4 |
| 2002..... | 8.5 | * | * | * | 7.6 | 13.4 | 18.7 | 14.1 | 3.6 | 1.5 | 1.5 | 3.9 |
| 2001..... | 6.8 | * | * | * | 6.2 | 10.4 | 15.4 | 10.7 | 2.6 | 1.3 | 1.7 | 1.8 |
| 2000..... | 6.1 | * | * | * | 5.3 | 9.2 | 14.1 | 9.5 | 1.9 | 1.1 | 1.3 | 4.1 |
| 1999..... | 5.6 | * | * | * | 3.8 | 8.7 | 13.1 | 8.9 | 2.2 | 1.2 | 1.6 | 3.3 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.3 | * | * | * | 3.1 | 6.4 | 11.9 | 10.5 | 4.3 | 1.8 | 1.9 | 2.6 |
| 2003..... | 4.7 | * | * | * | 2.8 | 5.8 | 10.3 | 9.1 | 3.7 | 1.6 | 1.5 | 3.0 |
| 2002..... | 4.0 | * | * | * | 2.4 | 5.0 | 9.2 | 7.5 | 2.9 | 1.5 | 1.7 | 2.9 |
| 2001..... | 2.9 | * | * | * | 1.6 | 3.7 | 6.8 | 5.0 | 2.1 | 1.1 | 1.5 | 2.7 |
| 2000..... | 2.5 | * | * | * | 1.4 | 3.1 | 5.6 | 3.9 | 2.0 | 1.0 | 1.6 | 2.7 |
| 1999..... | 2.2 | * | * | * | 1.2 | 2.7 | 5.0 | 3.5 | 1.6 | 1.0 | 1.7 | 2.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Accidental poisoning by and exposure to drugs and other biological substances (X40-X44) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 6.3 | * | * | * | 1.7 | 5.3 | 13.0 | 18.6 | 9.4 | 3.0 | 2.1 | * |
| 2003..... | 6.1 | * | * | * | 1.3 | 5.6 | 13.5 | 17.6 | 7.0 | 3.2 | * | * |
| 2002..... | 6.0 | * | * | * | 1.4 | 5.5 | 13.4 | 17.5 | 7.6 | 1.8 | * | * |
| 2001..... | 5.5 | * | * | * | 1.4 | 5.0 | 12.7 | 16.1 | 5.9 | 1.9 | * | * |
| 2000..... | 5.3 | * | * | * | 1.3 | 5.4 | 12.6 | 15.0 | 4.8 | 1.9 | * | 6.4 |
| 1999..... | 5.4 | * | * | * | 1.5 | 6.0 | 12.7 | 14.7 | 5.1 | 2.2 | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 8.6 | * | * | * | 2.4 | 7.5 | 16.4 | 26.5 | 15.6 | 5.1 | * | * |
| 2003..... | 8.5 | * | * | * | 1.8 | 7.7 | 18.4 | 25.3 | 11.7 | 5.6 | * | * |
| 2002..... | 8.6 | * | * | * | 2.1 | 8.2 | 18.6 | 25.7 | 13.5 | 2.9 | * | * |
| 2001..... | 8.0 | * | * | * | 2.1 | 7.3 | 17.2 | 25.7 | 10.2 | 3.0 | * | * |
| 2000..... | 7.8 | * | * | * | 1.6 | 8.1 | 18.0 | 24.3 | 8.2 | 3.4 | * | * |
| 1999..... | 8.2 | * | * | * | 2.4 | 9.0 | 18.6 | 24.5 | 9.5 | 3.3 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 4.2 | * | * | * | 0.9 | 3.3 | 9.9 | 11.9 | 4.4 | * | * | * |
| 2003..... | 3.9 | * | * | * | 0.7 | 3.6 | 9.2 | 11.1 | 3.2 | * | * | * |
| 2002..... | 3.6 | * | * | * | * | 3.1 | 8.7 | 10.4 | 3.0 | * | * | * |
| 2001..... | 3.3 | * | * | * | * | 3.0 | 8.8 | 7.9 | 2.4 | * | * | * |
| 2000..... | 3.0 | * | * | * | 0.9 | 2.9 | 7.9 | 7.1 | 2.0 | * | * | * |
| 1999..... | 2.8 | * | * | * | * | 3.4 | 7.5 | 6.3 | 1.6 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.5 | 1.3 | 0.7 | 0.3 | 0.9 | 1.3 | 1.8 | 2.7 | 3.2 | 5.5 | 16.6 | 64.5 |
| 2003..... | 3.7 | 1.1 | 0.7 | 0.3 | 1.0 | 1.2 | 1.9 | 2.5 | 3.2 | 6.0 | 17.6 | 73.5 |
| 2002..... | 3.8 | 0.9 | 0.6 | 0.3 | 0.9 | 1.2 | 1.9 | 2.4 | 3.5 | 6.2 | 18.5 | 75.8 |
| 2001..... | 4.0 | 1.3 | 0.8 | 0.3 | 1.0 | 1.2 | 1.9 | 2.7 | 3.5 | 6.4 | 20.1 | 83.4 |
| 2000..... | 3.9 | 1.1 | 0.7 | 0.3 | 1.0 | 1.3 | 2.1 | 2.6 | 3.6 | 6.2 | 18.9 | 82.6 |
| 1999..... | 4.3 | 1.2 | 0.8 | 0.4 | 1.0 | 1.4 | 2.0 | 2.7 | 3.7 | 7.0 | 22.1 | 92.8 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 4.2 | 1.6 | 0.9 | 0.4 | 1.5 | 2.2 | 3.0 | 4.4 | 5.0 | 7.6 | 20.2 | 69.6 |
| 2003..... | 4.3 | 1.1 | 0.8 | 0.4 | 1.6 | 2.0 | 3.1 | 4.1 | 5.2 | 8.6 | 21.6 | 80.4 |
| 2002..... | 4.3 | 1.2 | 0.7 | 0.4 | 1.5 | 2.1 | 3.0 | 4.0 | 5.5 | 8.8 | 22.8 | 79.9 |
| 2001..... | 4.5 | 1.6 | 0.9 | 0.4 | 1.6 | 2.1 | 3.1 | 4.4 | 5.3 | 8.7 | 24.8 | 92.4 |
| 2000..... | 4.5 | 1.2 | 0.8 | 0.4 | 1.7 | 2.3 | 3.6 | 4.5 | 5.6 | 8.5 | 22.6 | 91.7 |
| 1999..... | 4.8 | 1.4 | 1.0 | 0.5 | 1.7 | 2.3 | 3.4 | 4.4 | 5.8 | 9.6 | 27.3 | 107.0 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.9 | * | 0.4 | 0.2 | 0.3 | 0.3 | 0.7 | 1.0 | 1.5 | 3.6 | 14.2 | 62.2 |
| 2003..... | 3.1 | 1.1 | 0.5 | 0.1 | 0.3 | 0.4 | 0.6 | 1.0 | 1.4 | 3.8 | 14.9 | 70.5 |
| 2002..... | 3.2 | * | 0.5 | 0.2 | 0.3 | 0.3 | 0.7 | 0.9 | 1.6 | 4.0 | 15.7 | 74.0 |
| 2001..... | 3.5 | 1.1 | 0.7 | 0.2 | 0.3 | 0.3 | 0.7 | 1.1 | 1.8 | 4.4 | 17.1 | 79.6 |
| 2000..... | 3.3 | * | 0.5 | 0.2 | 0.3 | 0.3 | 0.6 | 0.8 | 1.7 | 4.4 | 16.5 | 78.9 |
| 1999..... | 3.7 | * | 0.5 | 0.3 | 0.3 | 0.4 | 0.7 | 1.0 | 1.8 | 4.8 | 18.7 | 87.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 1.0 | 1.2 | 1.8 | 2.3 | 2.1 | 2.9 | 7.6 | 30.0 |
| 2003..... | 1.5 | * | 0.6 | * | 1.2 | 1.4 | 1.4 | 2.0 | 2.6 | 4.3 | 9.2 | 27.5 |
| 2002..... | 1.5 | * | * | * | 0.9 | 1.3 | 1.8 | 2.0 | 2.7 | 3.7 | 8.0 | 35.5 |
| 2001..... | 1.6 | * | * | * | 1.0 | 1.2 | 1.9 | 2.5 | 2.6 | 4.9 | 9.0 | 47.5 |
| 2000..... | 1.6 | * | * | * | 1.0 | 1.6 | 2.1 | 2.1 | 2.6 | 4.3 | 9.3 | 45.1 |
| 1999..... | 1.5 | * | 0.8 | * | 0.9 | 1.0 | 1.7 | 2.0 | 2.9 | 5.1 | 8.7 | 37.2 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | * | 1.8 | 2.0 | 2.9 | 4.0 | 3.1 | 4.2 | 8.6 | * |
| 2003..... | 2.3 | * | * | * | 2.0 | 2.2 | 2.5 | 3.5 | 4.6 | 6.5 | 12.7 | * |
| 2002..... | 2.3 | * | * | * | 1.6 | 2.3 | 3.1 | 3.5 | 4.9 | 5.5 | 11.1 | 48.5 |
| 2001..... | 2.3 | * | * | * | 1.7 | 2.1 | 3.4 | 4.3 | 4.1 | 6.8 | 12.5 | 47.6 |
| 2000..... | 2.4 | * | * | * | 1.8 | 2.7 | 3.8 | 3.4 | 4.4 | 6.3 | * | 54.4 |
| 1999..... | 2.2 | * | * | * | 1.6 | 1.7 | 2.8 | 3.4 | 4.4 | 7.5 | * | 47.4 |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | * | * | 6.9 | 34.6 |
| 2003..... | 0.7 | * | * | * | * | * | * | * | * | * | 6.8 | 27.3 |
| 2002..... | 0.6 | * | * | * | * | * | * | * | * | * | 5.9 | 29.0 |
| 2001..... | 0.9 | * | * | * | * | * | * | * | * | 3.5 | 6.6 | 47.5 |
| 2000..... | 0.8 | * | * | * | * | * | * | * | * | * | 9.6 | 40.6 |
| 1999..... | 0.8 | * | * | * | * | * | * | * | * | * | 8.3 | 32.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.8 | 1.5 | 0.7 | 0.3 | 0.9 | 1.3 | 1.9 | 2.7 | 3.2 | 5.6 | 17.1 | 66.0 |
| 2003..... | 4.0 | 1.1 | 0.7 | 0.3 | 0.9 | 1.1 | 1.9 | 2.6 | 3.3 | 6.1 | 18.0 | 75.3 |
| 2002..... | 4.1 | 1.0 | 0.7 | 0.4 | 0.9 | 1.2 | 1.9 | 2.4 | 3.5 | 6.3 | 19.0 | 77.2 |
| 2001..... | 4.3 | 1.4 | 0.9 | 0.3 | 1.0 | 1.3 | 1.9 | 2.7 | 3.5 | 6.5 | 20.6 | 84.5 |
| 2000..... | 4.2 | 1.2 | 0.7 | 0.3 | 1.0 | 1.2 | 2.1 | 2.6 | 3.6 | 6.3 | 19.2 | 83.7 |
| 1999..... | 4.6 | 1.3 | 0.7 | 0.4 | 1.0 | 1.4 | 2.1 | 2.7 | 3.7 | 7.1 | 22.6 | 94.6 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.5 | 1.9 | 1.0 | 0.4 | 1.5 | 2.2 | 3.0 | 4.5 | 5.1 | 7.8 | 20.9 | 72.0 |
| 2003..... | 4.6 | * | 0.8 | 0.4 | 1.5 | 1.9 | 3.2 | 4.2 | 5.2 | 8.8 | 22.0 | 82.7 |
| 2002..... | 4.6 | 1.3 | 0.8 | 0.5 | 1.5 | 2.0 | 3.0 | 4.0 | 5.5 | 9.0 | 23.3 | 81.0 |
| 2001..... | 4.8 | 1.8 | 1.0 | 0.4 | 1.6 | 2.2 | 3.1 | 4.4 | 5.4 | 8.8 | 25.3 | 94.3 |
| 2000..... | 4.8 | 1.4 | 0.9 | 0.4 | 1.7 | 2.2 | 3.5 | 4.5 | 5.7 | 8.6 | 23.2 | 93.3 |
| 1999..... | 5.2 | 1.6 | 1.0 | 0.5 | 1.7 | 2.4 | 3.4 | 4.5 | 5.8 | 9.7 | 27.9 | 108.9 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 3.2 | * | 0.5 | 0.2 | 0.3 | 0.3 | 0.7 | 1.0 | 1.5 | 3.8 | 14.5 | 63.3 |
| 2003..... | 3.5 | * | 0.5 | 0.2 | 0.4 | 0.4 | 0.7 | 1.0 | 1.5 | 3.9 | 15.3 | 72.1 |
| 2002..... | 3.6 | * | 0.5 | 0.2 | 0.3 | 0.4 | 0.8 | 0.9 | 1.7 | 4.1 | 16.1 | 75.5 |
| 2001..... | 3.8 | * | 0.8 | 0.2 | 0.4 | 0.4 | 0.7 | 1.1 | 1.8 | 4.5 | 17.6 | 80.5 |
| 2000..... | 3.6 | * | 0.5 | 0.2 | 0.3 | 0.3 | 0.6 | 0.8 | 1.7 | 4.5 | 16.7 | 79.9 |
| 1999..... | 4.1 | * | 0.4 | 0.3 | 0.4 | 0.4 | 0.8 | 1.0 | 1.8 | 4.9 | 19.1 | 88.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 4.3 | 1.4 | 0.8 | 0.3 | 0.9 | 1.3 | 1.9 | 2.7 | 3.2 | 5.7 | 17.4 | 69.3 |
| 2003..... | 4.4 | 0.9 | 0.6 | 0.3 | 0.9 | 1.2 | 1.9 | 2.4 | 3.1 | 6.0 | 18.4 | 79.0 |
| 2002..... | 4.5 | 0.9 | 0.6 | 0.3 | 0.9 | 1.3 | 1.8 | 2.4 | 3.4 | 6.2 | 19.2 | 80.6 |
| 2001..... | 4.8 | 1.3 | 0.9 | 0.3 | 1.1 | 1.2 | 1.8 | 2.6 | 3.5 | 6.4 | 21.1 | 87.7 |
| 2000..... | 4.6 | 1.1 | 0.7 | 0.3 | 1.0 | 1.3 | 2.0 | 2.5 | 3.5 | 6.2 | 19.4 | 87.5 |
| 1999..... | 5.0 | 1.1 | 0.7 | 0.4 | 1.1 | 1.4 | 2.1 | 2.5 | 3.5 | 6.8 | 22.6 | 98.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.8 | 1.9 | 1.1 | 0.4 | 1.5 | 2.3 | 3.1 | 4.5 | 5.0 | 7.9 | 21.0 | 75.2 |
| 2003..... | 4.9 | * | 0.8 | 0.4 | 1.5 | 2.0 | 3.1 | 3.9 | 4.9 | 8.5 | 22.2 | 86.3 |
| 2002..... | 4.9 | * | 0.7 | 0.4 | 1.6 | 2.1 | 3.0 | 4.0 | 5.3 | 8.7 | 23.4 | 83.8 |
| 2001..... | 5.2 | 1.8 | 1.1 | 0.4 | 1.7 | 2.1 | 3.0 | 4.2 | 5.3 | 8.7 | 25.8 | 97.8 |
| 2000..... | 5.0 | * | 0.8 | 0.4 | 1.7 | 2.2 | 3.3 | 4.1 | 5.2 | 8.3 | 22.9 | 96.9 |
| 1999..... | 5.4 | * | 1.0 | 0.5 | 1.7 | 2.4 | 3.4 | 4.1 | 5.4 | 9.1 | 27.8 | 110.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 3.7 | * | * | 0.2 | 0.3 | 0.3 | 0.7 | 1.0 | 1.4 | 3.7 | 14.9 | 66.6 |
| 2003..... | 4.0 | * | 0.5 | * | 0.3 | 0.4 | 0.7 | 0.9 | 1.5 | 3.9 | 15.8 | 75.8 |
| 2002..... | 4.1 | * | 0.4 | 0.3 | 0.3 | 0.4 | 0.7 | 0.9 | 1.7 | 4.0 | 16.4 | 79.3 |
| 2001..... | 4.3 | * | 0.7 | 0.2 | 0.4 | 0.4 | 0.6 | 1.1 | 1.8 | 4.4 | 18.1 | 83.5 |
| 2000..... | 4.2 | * | 0.6 | 0.2 | 0.3 | 0.3 | 0.6 | 0.8 | 1.8 | 4.4 | 17.2 | 83.7 |
| 1999..... | 4.6 | * | 0.5 | 0.3 | 0.4 | 0.4 | 0.8 | 0.9 | 1.6 | 4.8 | 19.2 | 93.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | * | * | 0.3 | 0.7 | 1.4 | 2.1 | 3.3 | 4.4 | 6.2 | 17.0 | 38.9 |
| 2003..... | 2.9 | * | 0.9 | * | 1.0 | 1.1 | 2.3 | 4.1 | 4.8 | 7.7 | 16.4 | 46.7 |
| 2002..... | 2.9 | * | 1.0 | 0.5 | 0.7 | 1.2 | 2.5 | 3.2 | 4.7 | 8.2 | 18.8 | 47.5 |
| 2001..... | 3.0 | * | 0.9 | 0.3 | 0.7 | 1.3 | 2.3 | 3.9 | 4.6 | 7.8 | 18.3 | 58.4 |
| 2000..... | 3.1 | * | 0.8 | 0.3 | 1.0 | 1.4 | 2.8 | 4.0 | 5.5 | 8.1 | 19.4 | 51.5 |
| 1999..... | 3.7 | * | 0.9 | 0.3 | 1.0 | 1.6 | 2.5 | 4.7 | 6.4 | 11.3 | 25.3 | 67.9 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.6 | * | * | * | 1.1 | 2.5 | 3.5 | 5.5 | 7.4 | 9.1 | 24.5 | 43.7 |
| 2003..... | 3.9 | * | * | * | 1.4 | 1.9 | 3.7 | 6.7 | 8.3 | 12.6 | 22.2 | 55.4 |
| 2002..... | 3.8 | * | * | 0.7 | 1.1 | 2.1 | 3.7 | 5.1 | 8.3 | 11.9 | 25.8 | 58.0 |
| 2001..... | 3.8 | * | * | * | 1.1 | 2.3 | 3.7 | 6.4 | 7.3 | 11.4 | 24.0 | 63.5 |
| 2000..... | 4.6 | * | * | * | 1.7 | 2.4 | 4.9 | 7.7 | 10.2 | 11.5 | 30.2 | 69.3 |
| 1999..... | 5.1 | * | * | * | 1.8 | 2.7 | 4.4 | 8.2 | 10.5 | 17.7 | 32.8 | 108.2 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.7 | * | * | * | * | * | 0.9 | 1.4 | 1.9 | 4.1 | 12.7 | 36.9 |
| 2003..... | 2.0 | * | * | * | * | * | 1.1 | 1.8 | 1.9 | 4.2 | 13.0 | 43.3 |
| 2002..... | 2.1 | * | * | * | * | * | 1.4 | 1.5 | 1.8 | 5.6 | 14.9 | 43.5 |
| 2001..... | 2.2 | * | * | * | * | * | 1.0 | 1.8 | 2.5 | 5.2 | 15.2 | 56.5 |
| 2000..... | 1.8 | * | * | * | * | * | 0.9 | * | 1.7 | 5.7 | 13.4 | 44.9 |
| 1999..... | 2.4 | * | * | * | * | * | 0.9 | 1.7 | 3.2 | 6.7 | 21.0 | 52.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 11.0 | ... | ... | 0.7 | 10.3 | 12.7 | 15.0 | 16.6 | 13.8 | 12.3 | 16.3 | 16.4 |
| 2003..... | 10.8 | ... | ... | 0.6 | 9.7 | 12.7 | 14.9 | 15.9 | 13.8 | 12.7 | 16.4 | 16.9 |
| 2002..... | 11.0 | ... | ... | 0.6 | 9.9 | 12.6 | 15.3 | 15.7 | 13.6 | 13.5 | 17.7 | 18.0 |
| 2001 5/..... | 10.8 | ... | ... | 0.7 | 9.9 | 12.8 | 14.7 | 15.2 | 13.1 | 13.3 | 17.4 | 17.5 |
| 2000..... | 10.4 | ... | ... | 0.7 | 10.2 | 12.0 | 14.5 | 14.4 | 12.1 | 12.5 | 17.6 | 19.6 |
| 1999..... | 10.5 | ... | ... | 0.6 | 10.1 | 12.7 | 14.3 | 13.9 | 12.2 | 13.4 | 18.1 | 19.3 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 17.7 | ... | ... | 0.9 | 16.8 | 20.4 | 23.0 | 24.8 | 22.1 | 22.6 | 34.8 | 45.0 |
| 2003..... | 17.6 | ... | ... | 0.9 | 16.0 | 20.6 | 23.2 | 24.4 | 22.3 | 23.4 | 35.1 | 47.8 |
| 2002..... | 17.9 | ... | ... | 0.9 | 16.5 | 20.5 | 23.7 | 24.4 | 22.2 | 24.7 | 38.1 | 50.7 |
| 2001 5/..... | 17.6 | ... | ... | 1.0 | 16.6 | 21.0 | 23.1 | 23.4 | 21.1 | 24.6 | 37.8 | 51.1 |
| 2000..... | 17.1 | ... | ... | 1.2 | 17.1 | 19.6 | 22.8 | 22.4 | 19.4 | 22.7 | 38.6 | 57.5 |
| 1999..... | 17.1 | ... | ... | 0.9 | 16.8 | 20.7 | 22.4 | 21.5 | 19.8 | 24.7 | 38.8 | 57.1 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 4.6 | ... | ... | 0.5 | 3.6 | 4.7 | 7.1 | 8.6 | 6.1 | 3.8 | 3.9 | 3.6 |
| 2003..... | 4.3 | ... | ... | 0.3 | 3.0 | 4.6 | 6.6 | 7.7 | 5.9 | 3.8 | 4.0 | 3.3 |
| 2002..... | 4.3 | ... | ... | 0.3 | 2.9 | 4.6 | 6.9 | 7.4 | 5.7 | 4.1 | 4.2 | 3.8 |
| 2001 5/..... | 4.1 | ... | ... | 0.3 | 2.9 | 4.4 | 6.4 | 7.2 | 5.7 | 3.9 | 4.0 | 3.4 |
| 2000..... | 4.0 | ... | ... | 0.3 | 3.0 | 4.3 | 6.4 | 6.7 | 5.4 | 4.0 | 4.0 | 4.2 |
| 1999..... | 4.0 | ... | ... | 0.3 | 3.0 | 4.6 | 6.4 | 6.6 | 5.2 | 4.1 | 4.7 | 4.1 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | ... | ... | 0.5 | 8.0 | 7.2 | 6.8 | 7.5 | 7.0 | 6.8 | 9.1 | * |
| 2003..... | 5.0 | ... | ... | 0.6 | 7.0 | 6.8 | 6.8 | 7.2 | 6.9 | 6.5 | 8.0 | 10.9 |
| 2002..... | 5.0 | ... | ... | 0.5 | 6.7 | 6.6 | 7.0 | 7.1 | 7.2 | 7.5 | 9.2 | 12.7 |
| 2001 5/..... | 5.0 | ... | ... | 0.5 | 6.2 | 7.1 | 7.3 | 7.0 | 6.2 | 7.6 | 8.8 | 13.8 |
| 2000..... | 5.1 | ... | ... | 0.4 | 6.8 | 6.5 | 7.5 | 7.8 | 6.0 | 7.8 | 11.7 | * |
| 1999..... | 5.0 | ... | ... | 0.5 | 6.2 | 6.7 | 7.5 | 7.1 | 6.8 | 7.0 | 13.1 | 15.4 |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 8.6 | ... | ... | 0.6 | 12.8 | 11.2 | 10.8 | 12.0 | 11.3 | 13.6 | 19.7 | * |
| 2003..... | 8.3 | ... | ... | 0.8 | 11.2 | 10.8 | 11.0 | 11.5 | 13.0 | 13.1 | 18.0 | * |
| 2002..... | 8.3 | ... | ... | 0.6 | 10.6 | 10.7 | 11.2 | 11.3 | 13.2 | 14.7 | 19.3 | 35.1 |
| 2001 5/..... | 8.3 | ... | ... | 0.7 | 9.5 | 11.3 | 12.4 | 11.4 | 11.5 | 15.4 | 20.1 | 41.9 |
| 2000..... | 8.4 | ... | ... | 0.6 | 10.9 | 10.6 | 12.1 | 12.5 | 11.1 | 15.6 | 26.1 | * |
| 1999..... | 8.2 | ... | ... | 0.6 | 9.9 | 10.7 | 12.1 | 11.6 | 12.2 | 13.8 | 28.2 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.8 | ... | ... | * | 2.5 | 2.3 | 2.4 | 3.1 | 3.1 | * | * | * |
| 2003..... | 1.5 | ... | ... | * | 2.2 | 1.9 | 2.2 | 2.9 | * | * | * | * |
| 2002..... | 1.6 | ... | ... | * | 2.1 | 1.7 | 2.4 | 2.9 | * | * | * | * |
| 2001 5/..... | 1.5 | ... | ... | * | 2.3 | 2.2 | 1.8 | 2.7 | * | * | * | * |
| 2000..... | 1.5 | ... | ... | * | 2.0 | 1.7 | 2.6 | 3.2 | * | * | * | * |
| 1999..... | 1.6 | ... | ... | * | 1.8 | 2.0 | 2.6 | 2.7 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 11.9 | ... | ... | 0.7 | 10.8 | 13.9 | 16.3 | 17.5 | 14.3 | 12.7 | 16.7 | 16.8 |
| 2003..... | 11.7 | ... | ... | 0.6 | 10.2 | 14.0 | 16.0 | 16.7 | 14.3 | 13.1 | 16.9 | 17.1 |
| 2002..... | 11.8 | ... | ... | 0.7 | 10.5 | 13.9 | 16.4 | 16.5 | 14.0 | 13.8 | 18.1 | 18.1 |
| 2001 5/..... | 11.6 | ... | ... | 0.7 | 10.7 | 13.9 | 15.7 | 15.9 | 13.6 | 13.6 | 17.8 | 17.5 |
| 2000..... | 11.1 | ... | ... | 0.8 | 10.8 | 13.0 | 15.4 | 15.0 | 12.5 | 12.7 | 17.9 | 19.9 |
| 1999..... | 11.2 | ... | ... | 0.6 | 10.8 | 13.8 | 15.1 | 14.4 | 12.5 | 13.7 | 18.3 | 19.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 19.2 | ... | ... | 1.0 | 17.5 | 22.7 | 25.0 | 26.1 | 22.8 | 23.1 | 35.6 | 46.2 |
| 2003..... | 19.1 | ... | ... | 0.9 | 16.9 | 22.9 | 25.1 | 25.6 | 22.9 | 24.1 | 36.0 | 48.6 |
| 2002..... | 19.4 | ... | ... | 1.0 | 17.6 | 22.7 | 25.5 | 25.6 | 22.8 | 25.2 | 39.0 | 50.9 |
| 2001 5/..... | 19.0 | ... | ... | 1.1 | 18.1 | 23.1 | 24.6 | 24.5 | 21.7 | 25.0 | 38.5 | 51.3 |
| 2000..... | 18.3 | ... | ... | 1.3 | 18.3 | 21.4 | 24.1 | 23.2 | 19.9 | 23.1 | 39.1 | 58.4 |
| 1999..... | 18.4 | ... | ... | 1.0 | 18.2 | 22.7 | 23.6 | 22.2 | 20.2 | 25.2 | 39.1 | 57.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 5.0 | ... | ... | 0.5 | 3.7 | 5.2 | 7.8 | 9.2 | 6.4 | 3.9 | 4.0 | 3.6 |
| 2003..... | 4.6 | ... | ... | 0.3 | 3.2 | 5.2 | 7.2 | 8.1 | 6.3 | 4.0 | 4.1 | 3.3 |
| 2002..... | 4.6 | ... | ... | 0.3 | 3.1 | 5.2 | 7.4 | 7.8 | 5.9 | 4.3 | 4.3 | 3.9 |
| 2001 5/..... | 4.5 | ... | ... | 0.3 | 3.0 | 4.9 | 7.0 | 7.6 | 6.1 | 4.1 | 4.2 | 3.5 |
| 2000..... | 4.3 | ... | ... | 0.3 | 3.1 | 4.8 | 6.8 | 7.0 | 5.7 | 4.1 | 4.1 | 4.3 |
| 1999..... | 4.3 | ... | ... | 0.2 | 3.2 | 5.0 | 6.8 | 6.9 | 5.4 | 4.3 | 4.8 | 4.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 13.5 | ... | ... | 0.7 | 11.6 | 15.4 | 18.7 | 19.9 | 15.8 | 13.8 | 17.9 | 17.9 |
| 2003..... | 13.3 | ... | ... | 0.6 | 10.9 | 15.5 | 18.3 | 19.1 | 15.8 | 14.3 | 18.2 | 18.4 |
| 2002..... | 13.4 | ... | ... | 0.7 | 11.5 | 15.4 | 18.7 | 18.7 | 15.6 | 15.1 | 19.4 | 19.2 |
| 2001 5/..... | 13.0 | ... | ... | 0.7 | 11.6 | 15.4 | 17.7 | 17.9 | 15.0 | 14.7 | 19.0 | 18.7 |
| 2000..... | 12.5 | ... | ... | 0.9 | 11.6 | 14.2 | 17.2 | 16.7 | 13.9 | 13.8 | 19.0 | 21.3 |
| 1999..... | 12.4 | ... | ... | 0.7 | 11.5 | 15.0 | 16.9 | 16.0 | 13.7 | 14.8 | 19.5 | 20.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 21.6 | ... | ... | 0.9 | 19.0 | 24.9 | 28.4 | 29.3 | 24.9 | 24.9 | 38.0 | 49.8 |
| 2003..... | 21.6 | ... | ... | 1.0 | 18.2 | 25.0 | 28.2 | 29.0 | 25.1 | 26.0 | 38.6 | 52.5 |
| 2002..... | 21.9 | ... | ... | 1.1 | 19.3 | 24.6 | 28.7 | 28.6 | 25.0 | 27.5 | 41.7 | 54.3 |
| 2001 5/..... | 21.4 | ... | ... | 1.1 | 19.6 | 25.2 | 27.4 | 27.4 | 23.6 | 26.9 | 41.1 | 55.2 |
| 2000..... | 20.4 | ... | ... | 1.4 | 19.5 | 23.1 | 26.6 | 25.5 | 21.8 | 24.7 | 41.7 | 62.7 |
| 1999..... | 20.3 | ... | ... | 1.1 | 19.2 | 24.3 | 26.0 | 24.4 | 21.9 | 27.0 | 41.6 | 62.2 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 5.7 | ... | ... | 0.5 | 4.0 | 5.6 | 9.0 | 10.6 | 7.1 | 4.3 | 4.2 | 3.6 |
| 2003..... | 5.3 | ... | ... | 0.2 | 3.3 | 5.8 | 8.3 | 9.3 | 7.0 | 4.3 | 4.4 | 3.4 |
| 2002..... | 5.3 | ... | ... | 0.3 | 3.4 | 6.1 | 8.6 | 9.0 | 6.6 | 4.6 | 4.5 | 4.1 |
| 2001 5/..... | 5.0 | ... | ... | 0.3 | 3.3 | 5.5 | 8.1 | 8.6 | 6.8 | 4.4 | 4.4 | 3.5 |
| 2000..... | 4.9 | ... | ... | 0.4 | 3.3 | 5.2 | 7.8 | 7.9 | 6.4 | 4.6 | 4.2 | 4.6 |
| 1999..... | 4.9 | ... | ... | 0.3 | 3.4 | 5.5 | 7.8 | 7.8 | 6.0 | 4.6 | 5.0 | 4.2 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 5.3 | ... | ... | 0.8 | 7.3 | 9.7 | 6.7 | 5.9 | 5.8 | 4.5 | 6.2 | * |
| 2003..... | 5.3 | ... | ... | 0.7 | 7.3 | 9.6 | 7.4 | 5.7 | 4.9 | 4.6 | 4.5 | * |
| 2002..... | 5.2 | ... | ... | 0.5 | 6.6 | 9.6 | 7.7 | 5.9 | 4.9 | 4.9 | 5.4 | * |
| 2001 5/..... | 5.3 | ... | ... | 0.6 | 7.3 | 9.2 | 7.5 | 6.2 | 5.1 | 5.5 | 5.6 | * |
| 2000..... | 5.5 | ... | ... | 0.6 | 8.4 | 9.0 | 7.7 | 6.5 | 4.2 | 5.3 | 5.9 | * |
| 1999..... | 5.5 | ... | ... | 0.5 | 8.3 | 9.5 | 7.8 | 5.7 | 5.0 | 5.8 | 5.8 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 9.2 | ... | ... | 0.9 | 12.4 | 16.8 | 11.4 | 9.7 | 11.2 | 10.1 | 15.4 | * |
| 2003..... | 9.0 | ... | ... | 1.0 | 12.5 | 17.1 | 12.5 | 9.3 | 8.5 | 8.5 | 11.3 | * |
| 2002..... | 9.3 | ... | ... | 0.9 | 11.6 | 18.0 | 13.2 | 10.2 | 8.5 | 9.7 | 13.5 | * |
| 2001 5/..... | 9.3 | ... | ... | 0.9 | 13.4 | 16.6 | 12.5 | 9.8 | 9.3 | 10.8 | 13.2 | * |
| 2000..... | 9.6 | ... | ... | 1.0 | 14.6 | 16.1 | 13.3 | 11.1 | 7.5 | 11.4 | 12.3 | * |
| 1999..... | 9.9 | ... | ... | 0.7 | 14.8 | 17.5 | 13.7 | 10.1 | 9.2 | 11.7 | 14.1 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.8 | ... | ... | * | 2.2 | 3.3 | 2.6 | 2.5 | 1.5 | * | * | * |
| 2003..... | 1.8 | ... | ... | * | 2.0 | 2.7 | 2.9 | 2.7 | 2.0 | * | * | * |
| 2002..... | 1.6 | ... | ... | * | 1.7 | 2.1 | 2.9 | 2.2 | 2.0 | * | * | * |
| 2001 5/..... | 1.7 | ... | ... | * | 1.3 | 2.4 | 3.0 | 3.1 | 1.8 | * | * | * |
| 2000..... | 1.7 | ... | ... | * | 2.3 | 2.6 | 2.7 | 2.5 | 1.6 | * | * | * |
| 1999..... | 1.6 | ... | ... | * | 2.0 | 2.4 | 2.6 | 2.0 | 1.6 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.6 | ... | ... | * | 0.7 | 1.6 | 2.9 | 3.3 | 2.1 | 0.9 | 0.9 | 1.1 |
| 2003..... | 1.5 | ... | ... | * | 0.6 | 1.6 | 2.7 | 3.1 | 1.9 | 1.0 | 0.9 | 0.9 |
| 2002..... | 1.5 | ... | ... | * | 0.6 | 1.5 | 2.8 | 3.0 | 1.8 | 1.0 | 1.0 | 0.9 |
| 2001..... | 1.4 | ... | ... | * | 0.6 | 1.4 | 2.7 | 2.8 | 1.5 | 1.0 | 0.8 | 0.9 |
| 2000..... | 1.3 | ... | ... | * | 0.6 | 1.3 | 2.4 | 2.5 | 1.5 | 0.8 | 0.9 | 1.0 |
| 1999..... | 1.2 | ... | ... | * | 0.5 | 1.3 | 2.4 | 2.4 | 1.4 | 1.0 | 1.1 | 0.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | ... | ... | * | 0.8 | 1.7 | 2.7 | 3.0 | 1.9 | 0.8 | 0.9 | 1.7 |
| 2003..... | 1.4 | ... | ... | * | 0.6 | 1.6 | 2.8 | 2.9 | 1.6 | 0.9 | 0.7 | 1.7 |
| 2002..... | 1.4 | ... | ... | * | 0.7 | 1.7 | 2.7 | 2.8 | 1.6 | 0.9 | 1.2 | * |
| 2001..... | 1.4 | ... | ... | * | 0.6 | 1.5 | 2.9 | 2.5 | 1.2 | 1.0 | 0.7 | * |
| 2000..... | 1.2 | ... | ... | * | 0.6 | 1.4 | 2.3 | 2.4 | 1.3 | 0.7 | 1.0 | * |
| 1999..... | 1.2 | ... | ... | * | 0.5 | 1.3 | 2.4 | 2.3 | 1.3 | 1.0 | 1.0 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | ... | ... | * | 0.6 | 1.5 | 3.0 | 3.6 | 2.2 | 1.0 | 0.9 | 0.8 |
| 2003..... | 1.5 | ... | ... | * | 0.6 | 1.6 | 2.6 | 3.2 | 2.1 | 1.1 | 1.1 | * |
| 2002..... | 1.5 | ... | ... | * | 0.6 | 1.4 | 2.8 | 3.2 | 2.1 | 1.1 | 1.0 | 0.7 |
| 2001..... | 1.4 | ... | ... | * | 0.7 | 1.3 | 2.5 | 3.1 | 1.8 | 0.9 | 0.8 | 0.7 |
| 2000..... | 1.3 | ... | ... | * | 0.6 | 1.3 | 2.5 | 2.6 | 1.7 | 0.9 | 0.8 | 0.9 |
| 1999..... | 1.3 | ... | ... | * | 0.6 | 1.3 | 2.3 | 2.5 | 1.6 | 0.9 | 1.2 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | ... | ... | * | 0.8 | 1.9 | 3.4 | 3.8 | 2.3 | 1.0 | 1.0 | 1.1 |
| 2003..... | 1.7 | ... | ... | * | 0.7 | 1.9 | 3.1 | 3.5 | 2.1 | 1.1 | 1.0 | 1.0 |
| 2002..... | 1.7 | ... | ... | * | 0.8 | 1.8 | 3.2 | 3.5 | 2.1 | 1.1 | 1.1 | 1.0 |
| 2001..... | 1.6 | ... | ... | * | 0.7 | 1.7 | 3.1 | 3.2 | 1.7 | 1.1 | 0.9 | 1.0 |
| 2000..... | 1.5 | ... | ... | * | 0.7 | 1.5 | 2.7 | 2.8 | 1.7 | 0.9 | 0.9 | 1.1 |
| 1999..... | 1.4 | ... | ... | * | 0.6 | 1.4 | 2.7 | 2.7 | 1.6 | 1.1 | 1.2 | 0.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.8 | ... | ... | * | 0.9 | 2.1 | 3.2 | 3.5 | 2.1 | 0.8 | 1.0 | 1.8 |
| 2003..... | 1.7 | ... | ... | * | 0.7 | 1.9 | 3.3 | 3.3 | 1.8 | 0.9 | 0.8 | 1.8 |
| 2002..... | 1.7 | ... | ... | * | 0.8 | 1.9 | 3.1 | 3.3 | 1.8 | 1.0 | 1.2 | * |
| 2001..... | 1.6 | ... | ... | * | 0.7 | 1.9 | 3.3 | 2.9 | 1.4 | 1.1 | 0.8 | * |
| 2000..... | 1.4 | ... | ... | * | 0.7 | 1.6 | 2.7 | 2.6 | 1.4 | 0.7 | 1.1 | * |
| 1999..... | 1.4 | ... | ... | * | 0.6 | 1.4 | 2.7 | 2.6 | 1.4 | 1.1 | 1.0 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.9 | ... | ... | * | 0.8 | 1.8 | 3.5 | 4.2 | 2.5 | 1.2 | 1.0 | 0.8 |
| 2003..... | 1.8 | ... | ... | * | 0.6 | 1.8 | 3.0 | 3.7 | 2.4 | 1.3 | 1.2 | * |
| 2002..... | 1.8 | ... | ... | * | 0.8 | 1.6 | 3.3 | 3.7 | 2.3 | 1.2 | 1.0 | 0.8 |
| 2001..... | 1.6 | ... | ... | * | 0.8 | 1.5 | 2.9 | 3.6 | 2.1 | 1.0 | 0.9 | * |
| 2000..... | 1.5 | ... | ... | * | 0.7 | 1.5 | 2.8 | 3.0 | 2.0 | 1.1 | 0.8 | 1.0 |
| 1999..... | 1.5 | ... | ... | * | 0.6 | 1.5 | 2.7 | 2.9 | 1.8 | 1.1 | 1.3 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | * | 0.7 | 0.8 | 0.8 | * | * | * | * |
| 2003..... | 0.4 | ... | ... | * | * | 0.6 | 1.1 | 1.0 | * | * | * | * |
| 2002..... | 0.5 | ... | ... | * | * | 0.8 | 1.0 | 0.8 | * | * | * | * |
| 2001..... | 0.4 | ... | ... | * | * | 0.5 | 1.0 | 0.9 | * | * | * | * |
| 2000..... | 0.4 | ... | ... | * | * | 0.6 | 1.0 | 1.1 | * | * | * | * |
| 1999..... | 0.4 | ... | ... | * | * | 0.7 | 0.9 | 0.8 | * | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | ... | ... | * | * | * | 1.0 | * | * | * | * | * |
| 2002..... | 0.5 | ... | ... | * | * | 1.1 | 1.0 | * | * | * | * | * |
| 2001..... | 0.4 | ... | ... | * | * | * | 1.0 | * | * | * | * | * |
| 2000..... | 0.5 | ... | ... | * | * | * | 0.9 | 1.3 | * | * | * | * |
| 1999..... | 0.4 | ... | ... | * | * | 0.9 | 0.9 | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.5 | ... | ... | * | * | 0.9 | 1.0 | 0.9 | * | * | * | * |
| 2003..... | 0.6 | ... | ... | * | * | * | 1.2 | 1.4 | * | * | * | * |
| 2002..... | 0.4 | ... | ... | * | * | * | 0.9 | 0.9 | * | * | * | * |
| 2001..... | 0.4 | ... | ... | * | * | * | 1.0 | 1.0 | * | * | * | * |
| 2000..... | 0.4 | ... | ... | * | * | * | 1.0 | 0.9 | * | * | * | * |
| 1999..... | 0.4 | ... | ... | * | * | * | 1.0 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.5 | ... | ... | * | 0.2 | 0.6 | 0.9 | 1.0 | 0.6 | 0.3 | 0.5 | 0.6 |
| 2003..... | 0.5 | ... | ... | * | 0.2 | 0.6 | 1.0 | 0.9 | 0.7 | 0.4 | 0.5 | 0.7 |
| 2002..... | 0.6 | ... | ... | * | 0.2 | 0.6 | 0.9 | 1.0 | 0.6 | 0.4 | 0.7 | 0.8 |
| 2001..... | 0.6 | ... | ... | * | 0.3 | 0.6 | 0.9 | 1.0 | 0.8 | 0.5 | 0.7 | 0.8 |
| 2000..... | 0.6 | ... | ... | * | 0.3 | 0.7 | 1.0 | 1.0 | 0.7 | 0.4 | 0.6 | 0.8 |
| 1999..... | 0.6 | ... | ... | * | 0.4 | 0.8 | 1.0 | 1.0 | 0.7 | 0.4 | 0.8 | 1.2 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.8 | ... | ... | * | 0.3 | 1.0 | 1.4 | 1.6 | 1.0 | 0.5 | 0.9 | * |
| 2003..... | 0.8 | ... | ... | * | 0.3 | 1.1 | 1.6 | 1.4 | 1.1 | 0.8 | 0.8 | 1.7 |
| 2002..... | 0.9 | ... | ... | * | 0.4 | 1.0 | 1.5 | 1.6 | 1.0 | 0.7 | 1.3 | 2.3 |
| 2001..... | 0.9 | ... | ... | * | 0.5 | 1.1 | 1.5 | 1.5 | 1.1 | 0.8 | 1.3 | 2.0 |
| 2000..... | 0.9 | ... | ... | * | 0.5 | 1.1 | 1.6 | 1.5 | 1.1 | 0.6 | 1.3 | 2.4 |
| 1999..... | 1.0 | ... | ... | * | 0.7 | 1.3 | 1.6 | 1.5 | 1.0 | 0.6 | 1.5 | 4.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | * | 0.2 | 0.4 | 0.4 | 0.3 | * | 0.3 | * |
| 2003..... | 0.2 | ... | ... | * | * | 0.2 | 0.4 | 0.4 | 0.3 | * | 0.3 | * |
| 2002..... | 0.2 | ... | ... | * | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.2 | 0.3 | * |
| 2001..... | 0.3 | ... | ... | * | * | 0.2 | 0.4 | 0.5 | 0.5 | 0.2 | * | * |
| 2000..... | 0.2 | ... | ... | * | * | 0.3 | 0.4 | 0.5 | 0.3 | 0.2 | * | * |
| 1999..... | 0.3 | ... | ... | * | * | 0.3 | 0.4 | 0.5 | 0.4 | * | 0.4 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | ... | ... | * | 0.2 | 0.7 | 1.1 | 1.2 | 0.7 | 0.4 | 0.6 | 0.7 |
| 2003..... | 0.6 | ... | ... | * | 0.2 | 0.8 | 1.2 | 1.0 | 0.8 | 0.5 | 0.6 | 0.8 |
| 2002..... | 0.7 | ... | ... | * | 0.3 | 0.8 | 1.1 | 1.2 | 0.7 | 0.5 | 0.7 | 0.9 |
| 2001..... | 0.7 | ... | ... | * | 0.3 | 0.8 | 1.1 | 1.1 | 0.9 | 0.5 | 0.7 | 0.9 |
| 2000..... | 0.7 | ... | ... | * | 0.3 | 0.8 | 1.2 | 1.2 | 0.8 | 0.4 | 0.7 | 0.9 |
| 1999..... | 0.7 | ... | ... | * | 0.5 | 1.0 | 1.2 | 1.2 | 0.8 | 0.4 | 0.9 | 1.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | ... | ... | * | 0.4 | 1.2 | 1.7 | 1.9 | 1.1 | 0.6 | 1.0 | * |
| 2003..... | 1.0 | ... | ... | * | 0.4 | 1.3 | 1.9 | 1.6 | 1.2 | 0.9 | 0.9 | 1.8 |
| 2002..... | 1.1 | ... | ... | * | 0.5 | 1.3 | 1.8 | 1.9 | 1.2 | 0.8 | 1.4 | 2.4 |
| 2001..... | 1.1 | ... | ... | * | 0.6 | 1.3 | 1.7 | 1.8 | 1.2 | 0.8 | 1.4 | 2.2 |
| 2000..... | 1.1 | ... | ... | * | 0.6 | 1.3 | 1.8 | 1.8 | 1.3 | 0.6 | 1.3 | 2.6 |
| 1999..... | 1.2 | ... | ... | * | 0.8 | 1.5 | 1.9 | 1.7 | 1.1 | 0.7 | 1.6 | 4.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | ... | ... | * | * | 0.2 | 0.4 | 0.5 | 0.3 | * | 0.3 | * |
| 2003..... | 0.3 | ... | ... | * | * | 0.2 | 0.5 | 0.5 | 0.4 | * | 0.4 | * |
| 2002..... | 0.3 | ... | ... | * | * | 0.3 | 0.5 | 0.5 | 0.3 | 0.3 | 0.3 | * |
| 2001..... | 0.3 | ... | ... | * | * | 0.3 | 0.5 | 0.5 | 0.5 | 0.3 | * | * |
| 2000..... | 0.3 | ... | ... | * | * | 0.4 | 0.5 | 0.6 | 0.4 | 0.3 | * | * |
| 1999..... | 0.3 | ... | ... | * | * | 0.4 | 0.5 | 0.6 | 0.5 | * | 0.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 2.6 | ... | ... | 0.5 | 3.7 | 4.1 | 3.9 | 3.0 | 2.0 | 1.2 | 1.6 | 2.2 |
| 2003..... | 2.3 | ... | ... | 0.4 | 3.3 | 3.9 | 3.6 | 2.7 | 1.7 | 1.2 | 1.6 | 2.1 |
| 2002..... | 2.3 | ... | ... | 0.4 | 3.2 | 3.8 | 3.5 | 2.7 | 1.6 | 1.3 | 1.6 | 2.2 |
| 2001..... | 2.2 | ... | ... | 0.4 | 3.2 | 3.6 | 3.5 | 2.5 | 1.6 | 1.2 | 1.6 | 2.0 |
| 2000..... | 2.0 | ... | ... | 0.4 | 2.9 | 3.3 | 3.1 | 2.2 | 1.4 | 1.1 | 1.7 | 2.9 |
| 1999..... | 2.0 | ... | ... | 0.3 | 2.6 | 3.4 | 3.1 | 2.0 | 1.3 | 1.3 | 1.8 | 2.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 4.2 | ... | ... | 0.7 | 5.8 | 7.1 | 6.8 | 4.9 | 3.2 | 2.0 | 2.8 | 4.5 |
| 2003..... | 4.0 | ... | ... | 0.6 | 5.4 | 6.7 | 6.2 | 4.5 | 2.8 | 2.1 | 3.0 | 4.9 |
| 2002..... | 3.9 | ... | ... | 0.6 | 5.5 | 6.5 | 5.9 | 4.5 | 2.8 | 2.2 | 2.6 | 5.0 |
| 2001..... | 3.8 | ... | ... | 0.7 | 5.4 | 6.3 | 5.9 | 4.2 | 2.5 | 2.2 | 2.9 | 4.6 |
| 2000..... | 3.5 | ... | ... | 0.7 | 4.9 | 5.7 | 5.3 | 3.8 | 2.1 | 1.6 | 3.1 | 7.0 |
| 1999..... | 3.3 | ... | ... | 0.5 | 4.2 | 5.9 | 5.3 | 3.3 | 2.2 | 2.0 | 3.0 | 6.3 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | ... | ... | 0.3 | 1.5 | 1.3 | 1.2 | 1.2 | 0.8 | 0.5 | 0.7 | 1.1 |
| 2003..... | 0.8 | ... | ... | 0.2 | 1.1 | 1.2 | 1.1 | 1.0 | 0.7 | 0.5 | 0.7 | 0.9 |
| 2002..... | 0.8 | ... | ... | 0.2 | 0.9 | 1.1 | 1.1 | 1.0 | 0.6 | 0.6 | 0.9 | 1.0 |
| 2001..... | 0.7 | ... | ... | 0.1 | 0.9 | 1.0 | 1.1 | 0.8 | 0.6 | 0.5 | 0.8 | 0.9 |
| 2000..... | 0.7 | ... | ... | 0.2 | 0.9 | 0.9 | 0.9 | 0.7 | 0.7 | 0.7 | 0.9 | 1.2 |
| 1999..... | 0.7 | ... | ... | 0.1 | 0.9 | 1.0 | 0.8 | 0.7 | 0.5 | 0.7 | 1.1 | 1.3 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | ... | ... | 0.5 | 3.9 | 4.5 | 4.4 | 3.2 | 2.0 | 1.1 | 1.6 | 2.2 |
| 2003..... | 2.5 | ... | ... | 0.3 | 3.5 | 4.3 | 3.9 | 3.0 | 1.7 | 1.1 | 1.6 | 2.1 |
| 2002..... | 2.5 | ... | ... | 0.4 | 3.6 | 4.1 | 3.9 | 3.0 | 1.7 | 1.3 | 1.6 | 2.2 |
| 2001..... | 2.4 | ... | ... | 0.4 | 3.6 | 3.9 | 3.8 | 2.7 | 1.6 | 1.2 | 1.7 | 1.9 |
| 2000..... | 2.2 | ... | ... | 0.4 | 3.1 | 3.5 | 3.3 | 2.4 | 1.5 | 1.1 | 1.7 | 2.9 |
| 1999..... | 2.1 | ... | ... | 0.3 | 2.7 | 3.6 | 3.3 | 2.1 | 1.3 | 1.3 | 1.8 | 2.8 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 4.5 | ... | ... | 0.6 | 6.2 | 7.7 | 7.6 | 5.2 | 3.3 | 1.9 | 2.8 | 4.6 |
| 2003..... | 4.2 | ... | ... | 0.5 | 5.8 | 7.3 | 6.7 | 4.9 | 2.8 | 1.9 | 3.0 | 5.0 |
| 2002..... | 4.2 | ... | ... | 0.6 | 6.2 | 6.9 | 6.5 | 4.8 | 2.9 | 2.2 | 2.7 | 4.9 |
| 2001..... | 4.1 | ... | ... | 0.6 | 6.1 | 6.6 | 6.4 | 4.6 | 2.6 | 2.1 | 3.0 | 4.6 |
| 2000..... | 3.7 | ... | ... | 0.7 | 5.3 | 6.1 | 5.7 | 4.0 | 2.3 | 1.6 | 3.0 | 7.0 |
| 1999..... | 3.5 | ... | ... | 0.5 | 4.4 | 6.2 | 5.7 | 3.4 | 2.2 | 2.1 | 3.0 | 6.5 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.0 | ... | ... | 0.3 | 1.5 | 1.2 | 1.3 | 1.3 | 0.8 | 0.5 | 0.7 | 1.1 |
| 2003..... | 0.8 | ... | ... | * | 1.1 | 1.3 | 1.2 | 1.1 | 0.7 | 0.4 | 0.7 | 0.8 |
| 2002..... | 0.8 | ... | ... | 0.2 | 1.0 | 1.2 | 1.3 | 1.2 | 0.6 | 0.5 | 0.8 | 1.0 |
| 2001..... | 0.7 | ... | ... | * | 0.9 | 1.0 | 1.2 | 0.8 | 0.7 | 0.5 | 0.8 | 0.8 |
| 2000..... | 0.7 | ... | ... | 0.2 | 0.9 | 0.8 | 0.9 | 0.8 | 0.7 | 0.7 | 0.8 | 1.3 |
| 1999..... | 0.7 | ... | ... | * | 0.9 | 1.0 | 0.9 | 0.8 | 0.5 | 0.7 | 1.0 | 1.4 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) by discharge of firearms (X72-X74) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.7 | ... | ... | 0.1 | 5.0 | 5.7 | 6.5 | 8.0 | 8.0 | 8.8 | 12.2 | 11.1 |
| 2003..... | 5.8 | ... | ... | 0.2 | 5.0 | 6.0 | 6.6 | 8.0 | 8.3 | 9.3 | 12.4 | 11.9 |
| 2002..... | 5.9 | ... | ... | 0.2 | 5.1 | 6.0 | 7.0 | 7.8 | 8.4 | 9.7 | 13.2 | 12.0 |
| 2001..... | 5.9 | ... | ... | 0.2 | 5.3 | 6.5 | 6.7 | 7.7 | 8.2 | 9.6 | 13.1 | 12.2 |
| 2000..... | 5.9 | ... | ... | 0.3 | 5.8 | 6.1 | 7.1 | 7.6 | 7.6 | 9.2 | 13.0 | 13.2 |
| 1999..... | 5.9 | ... | ... | 0.3 | 6.0 | 6.5 | 6.9 | 7.3 | 7.8 | 9.7 | 13.1 | 12.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 10.0 | ... | ... | 0.2 | 8.8 | 9.7 | 10.8 | 13.5 | 14.2 | 17.5 | 28.3 | 34.4 |
| 2003..... | 10.4 | ... | ... | 0.3 | 8.9 | 10.4 | 11.2 | 13.8 | 14.9 | 18.4 | 28.7 | 36.6 |
| 2002..... | 10.6 | ... | ... | 0.3 | 9.1 | 10.3 | 11.9 | 13.6 | 15.1 | 19.5 | 31.0 | 37.8 |
| 2001..... | 10.6 | ... | ... | 0.3 | 9.4 | 11.2 | 11.4 | 13.3 | 14.6 | 19.2 | 30.7 | 38.9 |
| 2000..... | 10.5 | ... | ... | 0.4 | 10.2 | 10.6 | 11.8 | 13.0 | 13.5 | 18.4 | 30.8 | 42.9 |
| 1999..... | 10.6 | ... | ... | 0.4 | 10.5 | 11.2 | 11.5 | 12.5 | 14.0 | 19.5 | 31.1 | 41.6 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.5 | ... | ... | * | 1.1 | 1.5 | 2.2 | 2.8 | 2.2 | 1.6 | 1.4 | * |
| 2003..... | 1.4 | ... | ... | * | 1.0 | 1.4 | 2.1 | 2.5 | 2.2 | 1.6 | 1.5 | 0.9 |
| 2002..... | 1.4 | ... | ... | * | 1.0 | 1.6 | 2.2 | 2.2 | 2.2 | 1.6 | 1.4 | 0.8 |
| 2001..... | 1.5 | ... | ... | 0.1 | 1.0 | 1.6 | 2.1 | 2.4 | 2.4 | 1.7 | 1.5 | 1.0 |
| 2000..... | 1.5 | ... | ... | 0.1 | 1.2 | 1.7 | 2.4 | 2.3 | 2.1 | 1.7 | 1.4 | 1.1 |
| 1999..... | 1.5 | ... | ... | 0.1 | 1.2 | 1.8 | 2.3 | 2.3 | 2.1 | 1.7 | 1.4 | 0.9 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 2.1 | ... | ... | * | 3.7 | 2.7 | 2.4 | 3.0 | 2.7 | 3.5 | 5.0 | * |
| 2003..... | 2.1 | ... | ... | * | 3.3 | 2.8 | 2.5 | 3.0 | 2.7 | 3.1 | 4.1 | * |
| 2002..... | 2.2 | ... | ... | * | 3.3 | 2.7 | 2.5 | 2.7 | 3.4 | 4.4 | 4.7 | * |
| 2001..... | 2.2 | ... | ... | * | 2.9 | 3.3 | 2.5 | 3.0 | 3.2 | 4.0 | 4.7 | * |
| 2000..... | 2.3 | ... | ... | * | 3.4 | 2.8 | 2.8 | 4.0 | 2.7 | 4.6 | 5.3 | * |
| 1999..... | 2.3 | ... | ... | * | 3.5 | 2.9 | 3.0 | 3.2 | 2.9 | 3.6 | 8.1 | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 3.7 | ... | ... | * | 6.5 | 4.3 | 4.1 | 5.3 | 5.0 | 7.1 | 11.4 | * |
| 2003..... | 3.8 | ... | ... | * | 5.6 | 4.8 | 4.3 | 5.3 | 5.6 | 7.0 | 9.6 | * |
| 2002..... | 3.8 | ... | ... | * | 5.7 | 4.6 | 4.2 | 4.8 | 6.6 | 9.4 | 11.5 | * |
| 2001..... | 3.8 | ... | ... | * | 4.8 | 5.5 | 4.5 | 5.3 | 6.6 | 8.8 | 11.6 | * |
| 2000..... | 4.1 | ... | ... | * | 5.8 | 4.7 | 4.7 | 6.8 | 5.7 | 9.7 | 12.8 | * |
| 1999..... | 4.0 | ... | ... | * | 5.8 | 4.8 | 5.0 | 5.5 | 5.9 | 7.7 | 19.8 | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.6 | 0.7 | * | * | * | * | * | * |
| 2003..... | 0.3 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.4 | ... | ... | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.4 | ... | ... | * | 0.7 | 0.7 | * | * | * | * | * | * |
| 2000..... | 0.4 | ... | ... | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.5 | ... | ... | * | 0.8 | * | 0.9 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) by discharge of firearms (X72-X74) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 6.3 | ... | ... | 0.2 | 5.3 | 6.4 | 7.1 | 8.6 | 8.4 | 9.2 | 12.6 | 11.4 |
| 2003..... | 6.4 | ... | ... | 0.2 | 5.4 | 6.7 | 7.2 | 8.5 | 8.7 | 9.7 | 12.8 | 12.1 |
| 2002..... | 6.5 | ... | ... | 0.2 | 5.5 | 6.7 | 7.7 | 8.3 | 8.8 | 10.0 | 13.6 | 12.1 |
| 2001..... | 6.5 | ... | ... | 0.2 | 5.8 | 7.1 | 7.3 | 8.1 | 8.6 | 9.9 | 13.5 | 12.4 |
| 2000..... | 6.4 | ... | ... | 0.3 | 6.2 | 6.8 | 7.6 | 7.9 | 7.9 | 9.5 | 13.4 | 13.4 |
| 1999..... | 6.4 | ... | ... | 0.3 | 6.5 | 7.2 | 7.3 | 7.7 | 8.2 | 10.1 | 13.3 | 12.7 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 11.1 | ... | ... | 0.3 | 9.3 | 11.1 | 11.9 | 14.4 | 15.0 | 18.2 | 29.3 | 35.6 |
| 2003..... | 11.5 | ... | ... | 0.3 | 9.6 | 11.7 | 12.2 | 14.7 | 15.6 | 19.2 | 29.7 | 37.5 |
| 2002..... | 11.7 | ... | ... | 0.4 | 9.8 | 11.6 | 13.1 | 14.4 | 15.7 | 20.1 | 31.9 | 38.4 |
| 2001..... | 11.6 | ... | ... | 0.4 | 10.4 | 12.5 | 12.3 | 14.0 | 15.1 | 19.8 | 31.5 | 39.7 |
| 2000..... | 11.4 | ... | ... | 0.5 | 11.1 | 11.7 | 12.7 | 13.5 | 14.0 | 18.9 | 31.6 | 43.8 |
| 1999..... | 11.5 | ... | ... | 0.4 | 11.5 | 12.5 | 12.3 | 13.1 | 14.5 | 20.1 | 31.5 | 42.2 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.7 | ... | ... | * | 1.1 | 1.7 | 2.5 | 3.0 | 2.3 | 1.6 | 1.5 | * |
| 2003..... | 1.6 | ... | ... | * | 1.0 | 1.6 | 2.3 | 2.6 | 2.3 | 1.7 | 1.6 | 0.9 |
| 2002..... | 1.6 | ... | ... | * | 1.1 | 1.9 | 2.4 | 2.4 | 2.3 | 1.6 | 1.4 | 0.8 |
| 2001..... | 1.6 | ... | ... | 0.1 | 1.1 | 1.8 | 2.4 | 2.5 | 2.5 | 1.8 | 1.6 | 1.0 |
| 2000..... | 1.6 | ... | ... | 0.1 | 1.3 | 1.9 | 2.6 | 2.4 | 2.3 | 1.8 | 1.5 | 1.1 |
| 1999..... | 1.6 | ... | ... | * | 1.3 | 1.9 | 2.4 | 2.4 | 2.3 | 1.8 | 1.5 | 0.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 7.3 | ... | ... | 0.2 | 5.8 | 7.1 | 8.3 | 9.9 | 9.5 | 10.2 | 13.7 | 12.5 |
| 2003..... | 7.4 | ... | ... | 0.2 | 5.8 | 7.4 | 8.4 | 9.9 | 9.9 | 10.8 | 14.0 | 13.2 |
| 2002..... | 7.5 | ... | ... | 0.2 | 5.9 | 7.5 | 8.8 | 9.5 | 9.9 | 11.2 | 14.7 | 13.1 |
| 2001..... | 7.4 | ... | ... | 0.3 | 6.2 | 8.0 | 8.3 | 9.3 | 9.6 | 10.9 | 14.5 | 13.4 |
| 2000..... | 7.3 | ... | ... | 0.4 | 6.7 | 7.4 | 8.6 | 8.9 | 8.7 | 10.4 | 14.5 | 14.5 |
| 1999..... | 7.3 | ... | ... | 0.3 | 6.8 | 7.9 | 8.3 | 8.6 | 9.0 | 11.0 | 14.3 | 13.7 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 12.8 | ... | ... | 0.3 | 10.1 | 12.1 | 13.7 | 16.5 | 16.7 | 19.9 | 31.5 | 39.0 |
| 2003..... | 13.2 | ... | ... | 0.4 | 10.3 | 12.7 | 14.1 | 16.8 | 17.4 | 21.2 | 32.1 | 41.0 |
| 2002..... | 13.4 | ... | ... | 0.4 | 10.5 | 12.6 | 14.9 | 16.3 | 17.4 | 22.2 | 34.3 | 41.3 |
| 2001..... | 13.2 | ... | ... | 0.4 | 11.0 | 13.8 | 13.9 | 15.9 | 16.7 | 21.6 | 33.8 | 43.0 |
| 2000..... | 12.9 | ... | ... | 0.6 | 11.8 | 12.7 | 14.3 | 15.1 | 15.3 | 20.4 | 34.2 | 47.6 |
| 1999..... | 12.9 | ... | ... | 0.5 | 12.0 | 13.4 | 13.7 | 14.6 | 15.9 | 21.8 | 33.8 | 45.7 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.9 | ... | ... | * | 1.3 | 2.0 | 3.0 | 3.5 | 2.7 | 1.9 | 1.6 | * |
| 2003..... | 1.8 | ... | ... | * | 1.1 | 2.0 | 2.8 | 3.1 | 2.7 | 1.9 | 1.7 | 1.0 |
| 2002..... | 1.8 | ... | ... | * | 1.2 | 2.3 | 2.8 | 2.8 | 2.7 | 1.8 | 1.6 | 0.9 |
| 2001..... | 1.9 | ... | ... | * | 1.2 | 2.1 | 2.7 | 2.9 | 2.9 | 1.9 | 1.7 | 1.1 |
| 2000..... | 1.9 | ... | ... | * | 1.4 | 2.1 | 3.0 | 2.8 | 2.6 | 2.0 | 1.5 | 1.2 |
| 1999..... | 1.9 | ... | ... | * | 1.4 | 2.3 | 2.9 | 2.7 | 2.6 | 2.0 | 1.6 | 1.0 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) by discharge of firearms (X72-X74) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | ... | ... | * | 4.1 | 5.0 | 2.8 | 2.4 | 3.1 | 3.6 | 4.9 | * |
| 2003..... | 2.7 | ... | ... | * | 4.4 | 5.4 | 2.7 | 2.5 | 2.8 | 3.4 | 4.0 | * |
| 2002..... | 2.9 | ... | ... | * | 4.2 | 5.3 | 3.6 | 2.9 | 2.8 | 3.3 | 4.3 | * |
| 2001..... | 3.0 | ... | ... | * | 5.1 | 5.0 | 3.5 | 2.8 | 2.7 | 3.8 | 5.1 | * |
| 2000..... | 3.1 | ... | ... | * | 5.4 | 5.2 | 3.5 | 3.2 | 3.0 | 3.8 | 3.6 | * |
| 1999..... | 3.2 | ... | ... | * | 5.8 | 5.1 | 3.6 | 3.0 | 3.5 | 4.0 | 4.8 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 5.1 | ... | ... | * | 7.2 | 9.5 | 5.3 | 4.5 | 6.0 | 8.6 | 13.1 | * |
| 2003..... | 5.1 | ... | ... | * | 8.0 | 10.5 | 5.1 | 4.8 | 5.3 | 6.8 | 9.8 | * |
| 2002..... | 5.5 | ... | ... | * | 7.7 | 10.5 | 6.5 | 5.5 | 5.5 | 7.0 | 11.4 | * |
| 2001..... | 5.6 | ... | ... | * | 9.6 | 9.5 | 6.4 | 4.9 | 5.5 | 7.9 | 12.0 | * |
| 2000..... | 5.7 | ... | ... | * | 9.9 | 9.6 | 6.4 | 5.6 | 5.7 | 9.0 | 8.3 | * |
| 1999..... | 6.1 | ... | ... | * | 10.6 | 10.0 | 6.9 | 5.6 | 7.1 | 8.9 | 12.2 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.5 | ... | ... | * | 0.9 | 1.0 | * | * | * | * | * | * |
| 2003..... | 0.5 | ... | ... | * | 0.7 | * | * | * | * | * | * | * |
| 2002..... | 0.5 | ... | ... | * | 0.7 | * | 1.0 | * | * | * | * | * |
| 2001..... | 0.6 | ... | ... | * | * | 0.9 | 1.0 | 1.0 | * | * | * | * |
| 2000..... | 0.7 | ... | ... | * | 1.0 | 1.1 | 0.8 | 1.1 | * | * | * | * |
| 1999..... | 0.6 | ... | ... | * | 1.0 | 0.8 | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) by jumping from a high place (X80) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | 0.5 |
| 2003..... | 0.3 | ... | ... | * | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | 0.6 |
| 2002..... | 0.3 | ... | ... | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | 0.6 |
| 2001..... | 0.2 | ... | ... | * | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | * |
| 2000..... | 0.2 | ... | ... | * | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 |
| 1999..... | 0.3 | ... | ... | * | 0.3 | 0.3 | 0.4 | 0.3 | 0.2 | 0.3 | 0.4 | 0.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.4 | 0.5 | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | * |
| 2003..... | 0.4 | ... | ... | * | 0.4 | 0.5 | 0.5 | 0.4 | 0.4 | 0.3 | 0.5 | * |
| 2002..... | 0.4 | ... | ... | * | 0.5 | 0.6 | 0.5 | 0.4 | 0.4 | 0.3 | 0.5 | * |
| 2001..... | 0.3 | ... | ... | * | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 | 0.6 | * |
| 2000..... | 0.3 | ... | ... | * | 0.4 | 0.4 | 0.4 | 0.5 | 0.2 | * | 0.5 | * |
| 1999..... | 0.4 | ... | ... | * | 0.5 | 0.5 | 0.6 | 0.5 | 0.2 | 0.5 | 0.6 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | 0.1 | 0.2 | 0.2 | 0.2 | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | * | 0.2 | 0.2 | 0.2 | 0.2 | * | * | * |
| 2002..... | 0.2 | ... | ... | * | 0.1 | 0.2 | 0.2 | 0.2 | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | * | 0.2 | 0.2 | 0.3 | * | 0.2 | * | * |
| 2000..... | 0.1 | ... | ... | * | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | * | * | * |
| 1999..... | 0.1 | ... | ... | * | * | 0.1 | 0.2 | 0.2 | 0.2 | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | ... | ... | * | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | * |
| 2003..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | 0.5 |
| 2002..... | 0.3 | ... | ... | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | 0.6 |
| 2001..... | 0.2 | ... | ... | * | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.4 | * |
| 2000..... | 0.2 | ... | ... | * | 0.3 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 |
| 1999..... | 0.3 | ... | ... | * | 0.3 | 0.3 | 0.4 | 0.3 | 0.2 | 0.3 | 0.4 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.4 | 0.5 | 0.5 | 0.4 | 0.4 | 0.4 | * | * |
| 2003..... | 0.4 | ... | ... | * | 0.4 | 0.5 | 0.5 | 0.5 | 0.4 | * | 0.5 | * |
| 2002..... | 0.4 | ... | ... | * | 0.5 | 0.6 | 0.5 | 0.4 | 0.4 | * | 0.5 | * |
| 2001..... | 0.3 | ... | ... | * | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 | 0.6 | * |
| 2000..... | 0.3 | ... | ... | * | 0.5 | 0.3 | 0.4 | 0.5 | 0.2 | * | 0.5 | * |
| 1999..... | 0.4 | ... | ... | * | 0.5 | 0.5 | 0.6 | 0.5 | * | 0.5 | 0.6 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.1 | ... | ... | * | * | * | 0.2 | 0.2 | * | * | * | * |
| 2003..... | 0.2 | ... | ... | * | * | 0.2 | 0.2 | 0.2 | 0.2 | * | * | * |
| 2002..... | 0.2 | ... | ... | * | * | 0.2 | 0.2 | 0.3 | * | * | * | * |
| 2001..... | 0.2 | ... | ... | * | * | 0.2 | 0.2 | 0.2 | * | 0.2 | * | * |
| 2000..... | 0.1 | ... | ... | * | * | * | * | 0.2 | * | * | * | * |
| 1999..... | 0.1 | ... | ... | * | * | * | 0.2 | 0.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | ... | ... | * | 0.5 | 0.7 | 1.0 | 1.2 | 0.8 | 0.7 | 0.7 | 0.9 |
| 2003..... | 0.6 | ... | ... | * | 0.4 | 0.8 | 1.0 | 1.1 | 0.8 | 0.5 | 0.6 | 0.7 |
| 2002..... | 0.7 | ... | ... | * | 0.5 | 0.7 | 1.0 | 1.0 | 0.8 | 0.7 | 0.8 | 1.3 |
| 2001 5/..... | 0.6 | ... | ... | * | 0.5 | 0.7 | 0.9 | 1.0 | 0.8 | 0.6 | 0.9 | 1.0 |
| 2000..... | 0.6 | ... | ... | * | 0.5 | 0.6 | 0.9 | 0.9 | 0.8 | 0.6 | 0.9 | 1.1 |
| 1999..... | 0.6 | ... | ... | * | 0.5 | 0.7 | 0.9 | 0.9 | 0.6 | 0.7 | 0.8 | 1.3 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | ... | ... | * | 0.8 | 1.1 | 1.5 | 1.7 | 1.1 | 1.1 | 1.1 | 2.2 |
| 2003..... | 1.0 | ... | ... | * | 0.6 | 1.2 | 1.6 | 1.6 | 1.2 | 0.8 | 1.2 | 1.7 |
| 2002..... | 1.0 | ... | ... | * | 0.8 | 1.1 | 1.6 | 1.6 | 1.2 | 1.0 | 1.4 | 2.8 |
| 2001 5/..... | 1.0 | ... | ... | * | 0.8 | 1.1 | 1.4 | 1.6 | 1.1 | 0.9 | 1.5 | 2.3 |
| 2000..... | 0.9 | ... | ... | * | 0.7 | 0.9 | 1.5 | 1.3 | 1.2 | 1.0 | 1.5 | 2.4 |
| 1999..... | 0.9 | ... | ... | * | 0.8 | 1.1 | 1.2 | 1.3 | 0.9 | 1.0 | 1.4 | 2.6 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.2 | 0.3 | 0.5 | 0.7 | 0.5 | 0.4 | 0.4 | * |
| 2003..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.5 | 0.3 | * | * |
| 2002..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.5 | 0.4 | 0.5 | 0.4 | 0.5 | 0.7 |
| 2001 5/..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 | * |
| 2000..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.4 | 0.4 | 0.5 | * |
| 1999..... | 0.3 | ... | ... | * | 0.2 | 0.2 | 0.6 | 0.5 | 0.4 | 0.5 | 0.4 | 0.7 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | ... | ... | * | 0.5 | 0.7 | 1.0 | 1.2 | 0.8 | 0.8 | 0.6 | 0.9 |
| 2003..... | 0.7 | ... | ... | * | 0.4 | 0.7 | 1.1 | 1.2 | 0.9 | 0.6 | 0.6 | 0.7 |
| 2002..... | 0.7 | ... | ... | * | 0.5 | 0.7 | 1.1 | 1.1 | 0.9 | 0.7 | 0.9 | 1.4 |
| 2001 5/..... | 0.7 | ... | ... | * | 0.5 | 0.7 | 1.0 | 1.1 | 0.8 | 0.6 | 0.9 | 1.0 |
| 2000..... | 0.6 | ... | ... | * | 0.4 | 0.6 | 0.9 | 0.9 | 0.8 | 0.7 | 0.9 | 1.2 |
| 1999..... | 0.6 | ... | ... | * | 0.5 | 0.6 | 0.9 | 0.9 | 0.6 | 0.7 | 0.8 | 1.3 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.0 | ... | ... | * | 0.8 | 1.1 | 1.5 | 1.7 | 1.1 | 1.2 | 1.0 | 2.2 |
| 2003..... | 1.0 | ... | ... | * | 0.5 | 1.1 | 1.6 | 1.7 | 1.3 | 0.9 | 1.3 | 1.9 |
| 2002..... | 1.1 | ... | ... | * | 0.8 | 1.1 | 1.6 | 1.7 | 1.2 | 1.0 | 1.4 | 2.9 |
| 2001 5/..... | 1.0 | ... | ... | * | 0.8 | 1.2 | 1.4 | 1.6 | 1.2 | 0.9 | 1.5 | 2.6 |
| 2000..... | 0.9 | ... | ... | * | 0.7 | 0.9 | 1.5 | 1.3 | 1.2 | 1.0 | 1.5 | 2.6 |
| 1999..... | 0.9 | ... | ... | * | 0.8 | 1.1 | 1.2 | 1.4 | 0.9 | 0.9 | 1.5 | 2.9 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | ... | ... | * | 0.2 | 0.3 | 0.5 | 0.7 | 0.5 | 0.4 | 0.3 | * |
| 2003..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.6 | 0.6 | 0.5 | 0.3 | * | * |
| 2002..... | 0.4 | ... | ... | * | 0.2 | 0.4 | 0.5 | 0.5 | 0.5 | 0.4 | 0.5 | 0.7 |
| 2001 5/..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 | * |
| 2000..... | 0.3 | ... | ... | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 | 0.4 | 0.5 | * |
| 1999..... | 0.3 | ... | ... | * | 0.2 | 0.2 | 0.6 | 0.5 | 0.3 | 0.4 | 0.4 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.9 | 8.0 | 2.4 | 0.8 | 12.2 | 11.2 | 6.8 | 4.8 | 3.0 | 2.4 | 2.2 | 2.1 |
| 2003..... | 6.1 | 8.5 | 2.4 | 0.8 | 13.0 | 11.3 | 7.0 | 4.9 | 2.8 | 2.4 | 2.5 | 2.2 |
| 2002..... | 6.1 | 7.5 | 2.7 | 0.9 | 12.9 | 11.2 | 7.2 | 4.8 | 3.2 | 2.3 | 2.3 | 2.1 |
| 2001 5/..... | 7.1 | 8.2 | 2.7 | 0.8 | 13.3 | 13.1 | 9.5 | 6.3 | 4.0 | 2.9 | 2.5 | 2.4 |
| 2000..... | 6.0 | 9.2 | 2.3 | 0.9 | 12.6 | 10.4 | 7.1 | 4.7 | 3.0 | 2.4 | 2.4 | 2.4 |
| 1999..... | 6.1 | 8.7 | 2.5 | 1.1 | 12.9 | 10.5 | 7.1 | 4.6 | 3.0 | 2.6 | 2.5 | 2.4 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 9.4 | 8.0 | 2.5 | 1.0 | 20.4 | 18.5 | 10.1 | 7.3 | 4.6 | 3.6 | 2.6 | 2.7 |
| 2003..... | 9.7 | 10.0 | 2.5 | 1.0 | 21.8 | 18.5 | 10.4 | 7.4 | 4.2 | 3.4 | 3.3 | 3.2 |
| 2002..... | 9.6 | 7.9 | 2.9 | 0.9 | 21.5 | 18.2 | 10.7 | 7.1 | 4.8 | 3.3 | 3.1 | 3.0 |
| 2001 5/..... | 11.1 | 9.5 | 3.0 | 0.9 | 22.2 | 20.8 | 13.9 | 9.5 | 5.9 | 4.0 | 3.1 | 3.2 |
| 2000..... | 9.3 | 10.4 | 2.5 | 1.1 | 20.9 | 16.7 | 10.3 | 6.9 | 4.6 | 3.4 | 3.2 | 3.3 |
| 1999..... | 9.3 | 9.6 | 2.5 | 1.2 | 21.1 | 16.5 | 10.3 | 6.8 | 4.7 | 3.7 | 3.4 | 3.4 |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 2.5 | 7.9 | 2.2 | 0.6 | 3.5 | 3.7 | 3.4 | 2.4 | 1.6 | 1.4 | 1.9 | 1.9 |
| 2003..... | 2.6 | 6.9 | 2.3 | 0.5 | 3.7 | 3.9 | 3.6 | 2.5 | 1.5 | 1.5 | 1.9 | 1.8 |
| 2002..... | 2.7 | 7.1 | 2.5 | 0.8 | 3.8 | 4.2 | 3.8 | 2.6 | 1.6 | 1.4 | 1.8 | 1.7 |
| 2001 5/..... | 3.3 | 6.9 | 2.4 | 0.7 | 3.9 | 5.3 | 5.1 | 3.2 | 2.2 | 2.0 | 2.1 | 2.0 |
| 2000..... | 2.8 | 7.9 | 2.1 | 0.7 | 3.9 | 4.1 | 4.0 | 2.5 | 1.6 | 1.6 | 2.0 | 2.0 |
| 1999..... | 2.9 | 7.8 | 2.4 | 0.9 | 4.3 | 4.4 | 4.0 | 2.4 | 1.5 | 1.7 | 1.9 | 2.0 |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 7.9 | 8.1 | 2.3 | 0.9 | 17.4 | 12.6 | 7.7 | 5.9 | 3.8 | 2.3 | * | * |
| 2003..... | 8.4 | 6.9 | 2.5 | 0.7 | 18.4 | 13.2 | 8.5 | 6.4 | 3.6 | 3.8 | 3.6 | * |
| 2002..... | 8.1 | 6.2 | 2.6 | 0.9 | 17.6 | 12.3 | 8.0 | 5.9 | 4.4 | 2.5 | * | * |
| 2001 5/..... | 9.0 | 6.7 | 2.3 | 0.8 | 18.3 | 14.0 | 10.4 | 7.3 | 4.4 | 4.0 | * | * |
| 2000..... | 8.3 | 7.0 | 1.9 | 1.1 | 17.1 | 12.4 | 8.7 | 6.7 | 4.5 | 3.3 | * | * |
| 1999..... | 8.4 | 7.0 | 1.8 | 1.4 | 17.7 | 12.7 | 8.8 | 6.6 | 4.2 | 3.2 | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 13.0 | 9.1 | 2.5 | 1.0 | 29.6 | 20.3 | 11.8 | 9.5 | 6.5 | 4.1 | * | * |
| 2003..... | 13.6 | 7.9 | 2.1 | 0.9 | 30.3 | 21.2 | 12.9 | 10.2 | 6.0 | 6.8 | * | * |
| 2002..... | 13.2 | 6.6 | 3.1 | 1.0 | 29.6 | 19.8 | 12.1 | 9.1 | 7.7 | 4.8 | * | * |
| 2001 5/..... | 14.5 | 7.9 | 2.5 | 1.1 | 30.5 | 21.6 | 15.9 | 11.3 | 6.8 | 6.6 | * | * |
| 2000..... | 13.4 | 6.6 | 2.0 | 1.6 | 28.5 | 19.9 | 13.5 | 10.2 | 7.1 | 4.6 | * | * |
| 1999..... | 13.8 | 6.7 | 1.7 | 1.8 | 29.1 | 20.6 | 13.9 | 10.3 | 7.3 | 5.7 | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 2.4 | 7.0 | 2.1 | 0.7 | 3.3 | 3.4 | 3.2 | 2.2 | * | * | * | * |
| 2003..... | 2.8 | 5.9 | 2.9 | * | 4.5 | 3.5 | 3.6 | 2.7 | * | * | * | * |
| 2002..... | 2.6 | 5.9 | 2.1 | 0.8 | 3.8 | 3.4 | 3.5 | 2.7 | * | * | * | * |
| 2001 5/..... | 3.2 | 5.5 | 2.1 | * | 4.0 | 5.1 | 4.4 | 3.4 | 2.4 | * | * | * |
| 2000..... | 2.8 | 7.4 | 1.9 | 0.6 | 3.7 | 3.7 | 3.6 | 3.3 | 2.2 | * | * | * |
| 1999..... | 2.8 | 7.3 | 1.9 | 0.9 | 4.4 | 3.5 | 3.3 | 3.1 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 5.5 | 7.8 | 2.4 | 0.8 | 11.1 | 10.8 | 6.6 | 4.6 | 3.0 | 2.4 | 2.3 | 2.0 |
| 2003..... | 5.7 | 8.9 | 2.3 | 0.8 | 11.8 | 10.8 | 6.7 | 4.8 | 2.7 | 2.2 | 2.4 | 2.1 |
| 2002..... | 5.7 | 7.7 | 2.7 | 0.9 | 11.8 | 10.9 | 7.0 | 4.6 | 3.0 | 2.3 | 2.2 | 2.1 |
| 2001 5/..... | 6.7 | 8.4 | 2.8 | 0.8 | 12.1 | 12.8 | 9.2 | 6.1 | 3.9 | 2.8 | 2.5 | 2.4 |
| 2000..... | 5.6 | 9.6 | 2.4 | 0.9 | 11.6 | 10.0 | 6.9 | 4.4 | 2.9 | 2.3 | 2.4 | 2.3 |
| 1999..... | 5.7 | 8.9 | 2.6 | 1.0 | 11.9 | 10.0 | 6.8 | 4.3 | 2.9 | 2.5 | 2.5 | 2.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 8.7 | 7.5 | 2.5 | 0.9 | 18.4 | 17.9 | 9.7 | 6.9 | 4.4 | 3.5 | 2.7 | 2.4 |
| 2003..... | 9.0 | 10.5 | 2.6 | 1.0 | 19.8 | 17.7 | 10.0 | 7.1 | 4.0 | 3.1 | 3.2 | 3.0 |
| 2002..... | 8.9 | 8.0 | 2.9 | 0.9 | 19.6 | 17.6 | 10.4 | 6.8 | 4.5 | 3.2 | 3.1 | 2.9 |
| 2001 5/..... | 10.4 | 9.8 | 3.2 | 0.9 | 20.1 | 20.3 | 13.4 | 9.2 | 5.8 | 3.8 | 3.1 | 3.1 |
| 2000..... | 8.6 | 11.2 | 2.6 | 1.0 | 19.1 | 15.9 | 9.8 | 6.6 | 4.4 | 3.2 | 3.1 | 3.1 |
| 1999..... | 8.6 | 10.1 | 2.6 | 1.1 | 19.4 | 15.6 | 9.7 | 6.4 | 4.5 | 3.5 | 3.3 | 3.4 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 2.5 | 8.1 | 2.3 | 0.6 | 3.5 | 3.7 | 3.5 | 2.4 | 1.6 | 1.5 | 2.0 | 1.8 |
| 2003..... | 2.5 | 7.2 | 2.1 | 0.6 | 3.5 | 4.0 | 3.6 | 2.5 | 1.5 | 1.5 | 1.8 | 1.8 |
| 2002..... | 2.7 | 7.4 | 2.6 | 0.8 | 3.7 | 4.3 | 3.8 | 2.5 | 1.7 | 1.5 | 1.7 | 1.7 |
| 2001 5/..... | 3.2 | 6.9 | 2.4 | 0.7 | 3.8 | 5.3 | 5.1 | 3.1 | 2.2 | 2.0 | 2.1 | 2.1 |
| 2000..... | 2.7 | 7.9 | 2.2 | 0.7 | 3.8 | 4.1 | 4.0 | 2.3 | 1.5 | 1.6 | 1.9 | 2.0 |
| 1999..... | 2.9 | 7.6 | 2.5 | 0.9 | 4.2 | 4.6 | 4.0 | 2.3 | 1.4 | 1.7 | 2.0 | 2.0 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 2.7 | 5.9 | 1.4 | 0.5 | 3.5 | 4.0 | 3.8 | 3.0 | 2.2 | 2.0 | 1.9 | 1.8 |
| 2003..... | 2.7 | 6.8 | 1.3 | 0.4 | 3.5 | 3.9 | 3.8 | 3.1 | 2.1 | 1.7 | 2.0 | 2.0 |
| 2002..... | 2.8 | 5.0 | 1.8 | 0.5 | 3.7 | 3.9 | 4.1 | 3.0 | 2.4 | 1.9 | 1.8 | 1.8 |
| 2001 5/..... | 4.0 | 5.8 | 1.7 | 0.6 | 4.2 | 6.5 | 6.6 | 4.6 | 3.3 | 2.3 | 2.1 | 2.2 |
| 2000..... | 2.8 | 6.2 | 1.6 | 0.6 | 3.5 | 4.0 | 4.0 | 2.9 | 2.2 | 1.8 | 2.0 | 2.1 |
| 1999..... | 2.9 | 6.3 | 1.7 | 0.7 | 3.9 | 4.3 | 4.1 | 2.9 | 2.2 | 2.1 | 2.0 | 2.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 3.6 | 6.1 | 1.6 | 0.6 | 4.7 | 5.5 | 5.0 | 4.3 | 3.1 | 2.6 | 2.1 | 1.9 |
| 2003..... | 3.6 | 8.0 | 1.3 | 0.6 | 5.0 | 5.2 | 4.9 | 4.3 | 2.8 | 2.3 | 2.6 | 2.9 |
| 2002..... | 3.8 | 5.8 | 1.9 | 0.5 | 5.2 | 5.3 | 5.6 | 3.9 | 3.3 | 2.4 | 2.3 | 2.7 |
| 2001 5/..... | 5.6 | 6.8 | 2.0 | 0.6 | 5.7 | 9.3 | 9.5 | 6.7 | 4.7 | 2.9 | 2.6 | 2.8 |
| 2000..... | 3.6 | 8.3 | 1.8 | 0.7 | 4.7 | 5.2 | 5.2 | 3.9 | 3.1 | 2.2 | 2.4 | 2.6 |
| 1999..... | 3.9 | 8.0 | 1.7 | 0.7 | 5.1 | 5.7 | 5.4 | 4.2 | 3.2 | 2.7 | 2.6 | 2.6 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.8 | 5.8 | 1.3 | 0.4 | 2.3 | 2.5 | 2.5 | 1.8 | 1.4 | 1.4 | 1.8 | 1.7 |
| 2003..... | 1.8 | 5.5 | 1.4 | 0.3 | 1.9 | 2.5 | 2.6 | 2.0 | 1.4 | 1.3 | 1.6 | 1.6 |
| 2002..... | 1.9 | 4.1 | 1.8 | 0.6 | 2.2 | 2.6 | 2.5 | 2.1 | 1.5 | 1.4 | 1.5 | 1.5 |
| 2001 5/..... | 2.4 | 4.7 | 1.5 | 0.7 | 2.7 | 3.7 | 3.7 | 2.6 | 1.9 | 1.8 | 1.8 | 2.0 |
| 2000..... | 1.9 | 4.1 | 1.3 | 0.6 | 2.3 | 2.7 | 2.7 | 1.9 | 1.3 | 1.4 | 1.7 | 1.9 |
| 1999..... | 2.0 | 4.4 | 1.6 | 0.7 | 2.6 | 2.9 | 2.8 | 1.7 | 1.3 | 1.6 | 1.6 | 1.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 21.7 | 15.8 | 6.2 | 2.0 | 45.1 | 45.9 | 23.3 | 15.4 | 8.8 | 6.2 | 6.4 | * |
| 2003..... | 22.7 | 16.8 | 6.4 | 2.3 | 49.1 | 47.1 | 24.2 | 15.8 | 8.1 | 6.3 | 6.6 | * |
| 2002..... | 22.5 | 17.6 | 6.5 | 2.0 | 48.3 | 47.0 | 24.3 | 15.8 | 8.4 | 5.8 | 7.3 | * |
| 2001 5/..... | 22.7 | 19.5 | 7.1 | 1.4 | 48.7 | 45.3 | 25.5 | 16.2 | 8.9 | 7.2 | 6.8 | * |
| 2000..... | 22.1 | 24.1 | 6.1 | 1.9 | 49.3 | 42.2 | 24.5 | 15.0 | 8.4 | 7.6 | 7.1 | * |
| 1999..... | 21.7 | 20.0 | 6.4 | 2.3 | 49.5 | 40.6 | 23.7 | 14.1 | 8.4 | 6.4 | 7.5 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 38.3 | 13.9 | 6.3 | 2.5 | 80.7 | 85.1 | 39.0 | 26.1 | 15.2 | 11.8 | 10.3 | * |
| 2003..... | 40.1 | 18.6 | 7.8 | 3.0 | 87.3 | 86.1 | 41.6 | 26.9 | 14.9 | 11.3 | 11.3 | * |
| 2002..... | 39.4 | 16.4 | 7.3 | 2.6 | 85.8 | 85.3 | 39.8 | 27.5 | 15.8 | 10.5 | 13.8 | * |
| 2001 5/..... | 39.3 | 21.7 | 8.4 | 2.0 | 88.3 | 81.7 | 39.5 | 27.2 | 15.0 | 12.6 | 10.2 | * |
| 2000..... | 38.3 | 24.9 | 6.1 | 2.1 | 88.1 | 76.5 | 39.4 | 26.3 | 15.0 | 13.9 | 12.0 | * |
| 1999..... | 37.2 | 20.1 | 6.2 | 2.7 | 88.1 | 71.5 | 37.9 | 23.7 | 15.4 | 11.9 | 12.2 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 6.6 | 17.7 | 6.1 | 1.6 | 9.1 | 10.1 | 9.3 | 6.2 | 3.7 | 2.1 | 4.2 | * |
| 2003..... | 6.8 | 14.8 | 4.9 | 1.5 | 10.4 | 11.6 | 8.8 | 6.2 | 2.7 | 2.8 | 3.9 | * |
| 2002..... | 7.2 | 18.9 | 5.7 | 1.5 | 10.6 | 12.3 | 10.6 | 5.7 | 2.5 | 2.4 | 3.6 | * |
| 2001 5/..... | 7.6 | 17.1 | 5.7 | 0.8 | 9.2 | 12.5 | 13.2 | 6.9 | 4.0 | 3.3 | 4.9 | * |
| 2000..... | 7.4 | 23.3 | 6.0 | 1.6 | 11.0 | 11.4 | 11.3 | 5.3 | 3.2 | 3.2 | 4.3 | * |
| 1999..... | 7.7 | 20.0 | 6.7 | 1.8 | 11.4 | 12.8 | 11.1 | 5.9 | 2.7 | 2.5 | 4.9 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by hanging, strangulation and suffocation (X91) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | 1.0 | 0.2 | 0.1 | 0.3 | 0.3 | 0.3 | 0.2 | 0.1 | 0.2 | 0.2 | * |
| 2003..... | 0.2 | 0.9 | * | 0.1 | 0.3 | 0.3 | 0.3 | 0.2 | 0.1 | 0.2 | 0.3 | * |
| 2002..... | 0.2 | 0.9 | 0.2 | 0.1 | 0.3 | 0.3 | 0.3 | 0.3 | 0.1 | 0.1 | * | * |
| 2001..... | 0.2 | 1.0 | * | * | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | * |
| 2000..... | 0.2 | 0.7 | 0.2 | 0.1 | 0.3 | 0.3 | 0.3 | 0.2 | 0.1 | 0.2 | 0.2 | * |
| 1999..... | 0.2 | 1.0 | 0.2 | 0.1 | 0.3 | 0.4 | 0.3 | 0.2 | 0.1 | 0.2 | 0.2 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | 0.1 | 0.2 | 0.2 | 0.2 | * | * | * | * |
| 2003..... | 0.1 | * | * | * | 0.1 | * | 0.2 | 0.1 | * | * | * | * |
| 2002..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.2 | 0.2 | * | * | * | * |
| 2001..... | 0.2 | 1.3 | * | * | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | * | * |
| 2000..... | 0.2 | * | * | * | * | 0.2 | 0.2 | 0.1 | * | * | * | * |
| 1999..... | 0.2 | 1.4 | * | * | * | 0.2 | 0.2 | * | * | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.4 | 0.5 | 0.5 | 0.3 | * | 0.2 | * | * |
| 2003..... | 0.3 | * | * | * | 0.4 | 0.5 | 0.4 | 0.3 | 0.2 | 0.2 | * | * |
| 2002..... | 0.3 | * | * | * | 0.4 | 0.4 | 0.5 | 0.3 | * | * | * | * |
| 2001..... | 0.3 | * | * | * | 0.4 | 0.5 | 0.5 | 0.2 | 0.2 | 0.2 | 0.3 | * |
| 2000..... | 0.3 | * | * | * | 0.5 | 0.5 | 0.4 | 0.2 | * | * | * | * |
| 1999..... | 0.3 | * | * | * | 0.5 | 0.6 | 0.5 | 0.2 | * | 0.2 | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.2 | 0.9 | * | * | 0.2 | 0.3 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | * |
| 2003..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.3 | 0.1 | 0.1 | * | 0.2 | * |
| 2002..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.2 | 0.3 | 0.2 | 0.2 | 0.1 | 0.2 | * | * |
| 2000..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | * | * | * |
| 1999..... | 0.2 | * | * | 0.1 | 0.2 | 0.3 | 0.2 | 0.1 | * | 0.2 | 0.2 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.1 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.1 | * | * | * | * | * | 0.2 | * | * | * | * | * |
| 2002..... | 0.1 | * | * | * | * | * | 0.1 | 0.2 | * | * | * | * |
| 2001..... | 0.1 | * | * | * | * | * | * | 0.2 | * | * | * | * |
| 2000..... | 0.1 | * | * | * | * | 0.2 | 0.2 | * | * | * | * | * |
| 1999..... | 0.1 | * | * | * | * | * | 0.1 | * | * | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.2 | * | * | * | 0.3 | 0.3 | 0.3 | 0.2 | * | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.3 | 0.4 | 0.3 | 0.2 | * | * | * | * |
| 2002..... | 0.2 | * | * | * | 0.3 | 0.3 | 0.3 | 0.3 | * | * | * | * |
| 2001..... | 0.2 | * | * | * | 0.3 | 0.4 | 0.3 | 0.2 | * | * | * | * |
| 2000..... | 0.2 | * | * | * | 0.4 | 0.3 | 0.3 | 0.2 | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.4 | 0.4 | 0.3 | 0.2 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by hanging, strangulation and suffocation (X91) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | * | 0.5 | 0.7 | 1.1 | 0.5 | * | * | * | * |
| 2003..... | 0.5 | * | * | * | 0.7 | 0.7 | 0.7 | 0.7 | * | * | * | * |
| 2002..... | 0.6 | * | * | * | 0.6 | 0.9 | 1.0 | 0.7 | * | * | * | * |
| 2001..... | 0.6 | * | * | * | 0.7 | 0.8 | 1.0 | 0.6 | * | * | * | * |
| 2000..... | 0.6 | * | * | * | 0.7 | 1.0 | 0.9 | * | * | * | * | * |
| 1999..... | 0.6 | * | * | * | 0.7 | 1.2 | 1.0 | * | * | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2003..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 2002..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2001..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| 2000..... | 0.3 | * | * | * | * | * | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | 0.8 | 1.0 | 1.5 | * | * | * | * | * |
| 2003..... | 0.7 | * | * | * | 1.1 | 1.1 | 0.9 | 1.0 | * | * | * | * |
| 2002..... | 0.8 | * | * | * | 0.8 | 1.3 | 1.5 | * | * | * | * | * |
| 2001..... | 0.7 | * | * | * | 0.9 | 1.1 | 1.4 | * | * | * | * | * |
| 2000..... | 0.8 | * | * | * | 1.2 | 1.5 | 1.3 | * | * | * | * | * |
| 1999..... | 0.8 | * | * | * | 1.0 | 1.7 | 1.4 | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by discharge of firearms (*U01.4,X93-X95) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 4.0 | * | 0.2 | 0.5 | 9.9 | 8.8 | 4.3 | 2.6 | 1.6 | 1.0 | 0.8 | 0.6 |
| 2003..... | 4.1 | * | 0.3 | 0.5 | 10.7 | 8.9 | 4.4 | 2.7 | 1.4 | 0.9 | 0.7 | 0.5 |
| 2002..... | 4.1 | * | 0.3 | 0.5 | 10.6 | 8.7 | 4.5 | 2.6 | 1.5 | 0.9 | 0.9 | * |
| 2001..... | 4.0 | * | 0.4 | 0.4 | 10.5 | 8.4 | 4.4 | 2.4 | 1.4 | 1.0 | 0.8 | * |
| 2000..... | 3.8 | * | 0.2 | 0.5 | 10.1 | 7.7 | 4.2 | 2.5 | 1.5 | 1.0 | 0.7 | 0.7 |
| 1999..... | 3.9 | * | 0.3 | 0.5 | 10.4 | 7.6 | 4.1 | 2.4 | 1.6 | 1.0 | 0.8 | 0.6 |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 6.9 | * | * | 0.6 | 17.3 | 15.3 | 7.1 | 4.0 | 2.5 | 1.6 | 0.8 | * |
| 2003..... | 7.1 | * | 0.2 | 0.6 | 18.8 | 15.5 | 6.9 | 4.3 | 2.2 | 1.4 | 0.7 | * |
| 2002..... | 7.0 | * | 0.4 | 0.6 | 18.5 | 14.9 | 7.3 | 3.9 | 2.4 | 1.2 | 1.3 | * |
| 2001..... | 6.8 | * | 0.3 | 0.6 | 18.5 | 14.3 | 6.8 | 3.8 | 2.2 | 1.5 | 1.1 | * |
| 2000..... | 6.5 | * | * | 0.7 | 17.7 | 13.2 | 6.6 | 3.9 | 2.3 | 1.3 | 0.8 | * |
| 1999..... | 6.5 | * | 0.3 | 0.7 | 18.0 | 12.9 | 6.3 | 3.8 | 2.6 | 1.5 | 1.0 | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | 0.3 | 2.0 | 2.0 | 1.5 | 1.1 | 0.8 | 0.5 | 0.8 | * |
| 2003..... | 1.2 | * | 0.3 | 0.3 | 2.2 | 2.1 | 1.8 | 1.2 | 0.7 | 0.5 | 0.7 | * |
| 2002..... | 1.3 | * | * | 0.4 | 2.4 | 2.3 | 1.9 | 1.3 | 0.8 | 0.6 | 0.7 | * |
| 2001..... | 1.3 | * | 0.4 | 0.3 | 2.1 | 2.3 | 2.0 | 1.0 | 0.8 | 0.7 | 0.6 | * |
| 2000..... | 1.3 | * | * | 0.2 | 2.2 | 2.1 | 1.8 | 1.2 | 0.8 | 0.8 | 0.6 | * |
| 1999..... | 1.3 | * | 0.3 | 0.3 | 2.5 | 2.2 | 2.0 | 1.2 | 0.6 | 0.7 | 0.6 | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 5.4 | * | * | 0.5 | 14.0 | 9.4 | 5.0 | 2.7 | 2.0 | * | * | * |
| 2003..... | 5.8 | * | * | 0.4 | 14.9 | 9.9 | 5.3 | 3.6 | 1.2 | * | * | * |
| 2002..... | 5.6 | * | * | 0.6 | 14.3 | 9.3 | 5.1 | 3.2 | 1.6 | * | * | * |
| 2001..... | 5.7 | * | * | 0.4 | 14.6 | 9.5 | 5.6 | 3.0 | 1.1 | * | * | * |
| 2000..... | 5.5 | * | * | 0.6 | 13.7 | 8.9 | 4.9 | 4.0 | 2.0 | * | * | * |
| 1999..... | 5.7 | * | * | 0.8 | 14.4 | 8.7 | 5.2 | 3.8 | 2.1 | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 9.5 | * | * | 0.7 | 24.4 | 15.9 | 8.3 | 4.5 | 3.5 | * | * | * |
| 2003..... | 10.0 | * | * | 0.6 | 25.5 | 16.8 | 8.4 | 5.9 | 2.2 | * | * | * |
| 2002..... | 9.7 | * | * | 0.8 | 24.9 | 15.6 | 8.1 | 5.0 | 2.8 | * | * | * |
| 2001..... | 9.9 | * | * | 0.6 | 24.9 | 16.1 | 9.1 | 4.8 | * | * | * | * |
| 2000..... | 9.6 | * | * | 1.0 | 23.5 | 15.0 | 7.8 | 6.4 | 3.2 | * | * | * |
| 1999..... | 9.8 | * | * | 1.2 | 24.2 | 14.7 | 8.6 | 5.7 | 3.5 | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 2.0 | 1.6 | 1.4 | * | * | * | * | * |
| 2003..... | 1.3 | * | * | * | 2.7 | 1.7 | 1.9 | 1.4 | * | * | * | * |
| 2002..... | 1.2 | * | * | * | 2.1 | 1.8 | 1.8 | 1.4 | * | * | * | * |
| 2001..... | 1.3 | * | * | * | 2.5 | 1.8 | 1.9 | 1.2 | * | * | * | * |
| 2000..... | 1.3 | * | * | * | 2.2 | 1.9 | 1.7 | 1.7 | * | * | * | * |
| 1999..... | 1.4 | * | * | * | 2.9 | 1.7 | 1.7 | 1.9 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by discharge of firearms (*U01.4,X93-X95) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 3.7 | * | 0.2 | 0.4 | 9.0 | 8.5 | 4.1 | 2.5 | 1.6 | 1.1 | 0.8 | 0.5 |
| 2003..... | 3.8 | * | 0.2 | 0.5 | 9.8 | 8.6 | 4.2 | 2.6 | 1.4 | 0.8 | 0.7 | 0.5 |
| 2002..... | 3.8 | * | 0.3 | 0.5 | 9.8 | 8.5 | 4.4 | 2.5 | 1.5 | 0.9 | 0.9 | * |
| 2001..... | 3.7 | * | 0.3 | 0.4 | 9.6 | 8.0 | 4.2 | 2.3 | 1.5 | 1.0 | 0.8 | * |
| 2000..... | 3.6 | * | 0.2 | 0.4 | 9.3 | 7.4 | 4.1 | 2.4 | 1.5 | 1.0 | 0.7 | 0.7 |
| 1999..... | 3.6 | * | 0.4 | 0.5 | 9.6 | 7.3 | 4.0 | 2.3 | 1.5 | 1.1 | 0.8 | 0.5 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 6.4 | * | * | 0.6 | 15.8 | 15.1 | 6.8 | 3.9 | 2.4 | 1.6 | 0.8 | * |
| 2003..... | 6.5 | * | * | 0.7 | 17.2 | 15.0 | 6.7 | 4.1 | 2.1 | 1.3 | 0.7 | * |
| 2002..... | 6.5 | * | 0.3 | 0.6 | 17.0 | 14.7 | 7.1 | 3.7 | 2.3 | 1.2 | 1.2 | * |
| 2001..... | 6.3 | * | 0.3 | 0.6 | 17.0 | 13.8 | 6.5 | 3.7 | 2.2 | 1.4 | 1.1 | * |
| 2000..... | 6.0 | * | * | 0.6 | 16.3 | 12.7 | 6.4 | 3.7 | 2.2 | 1.3 | 0.8 | * |
| 1999..... | 6.0 | * | 0.4 | 0.7 | 16.7 | 12.4 | 6.0 | 3.6 | 2.5 | 1.5 | 0.9 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | 0.2 | 2.0 | 2.1 | 1.5 | 1.2 | 0.8 | 0.6 | 0.8 | * |
| 2003..... | 1.2 | * | * | 0.3 | 2.0 | 2.2 | 1.8 | 1.2 | 0.7 | 0.5 | 0.7 | * |
| 2002..... | 1.3 | * | * | 0.4 | 2.4 | 2.3 | 1.9 | 1.3 | 0.8 | 0.6 | 0.7 | * |
| 2001..... | 1.2 | * | 0.3 | 0.3 | 2.0 | 2.3 | 2.0 | 1.0 | 0.8 | 0.7 | 0.6 | * |
| 2000..... | 1.2 | * | * | 0.2 | 2.2 | 2.2 | 1.9 | 1.1 | 0.8 | 0.7 | 0.6 | * |
| 1999..... | 1.3 | * | 0.4 | 0.3 | 2.4 | 2.3 | 2.0 | 1.1 | 0.6 | 0.7 | 0.6 | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | 0.2 | 2.1 | 2.6 | 2.2 | 1.6 | 1.2 | 0.9 | 0.7 | * |
| 2003..... | 1.4 | * | * | 0.2 | 2.4 | 2.5 | 2.1 | 1.7 | 1.1 | 0.7 | 0.6 | 0.5 |
| 2002..... | 1.5 | * | 0.2 | 0.3 | 2.5 | 2.6 | 2.4 | 1.6 | 1.2 | 0.7 | 0.8 | * |
| 2001..... | 1.6 | * | 0.2 | 0.4 | 2.4 | 2.7 | 2.4 | 1.6 | 1.2 | 0.8 | 0.7 | * |
| 2000..... | 1.4 | * | * | 0.3 | 2.2 | 2.4 | 2.3 | 1.6 | 1.2 | 0.8 | 0.6 | 0.6 |
| 1999..... | 1.5 | * | 0.2 | 0.3 | 2.4 | 2.6 | 2.3 | 1.5 | 1.1 | 0.9 | 0.6 | 0.6 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.1 | * | * | 0.3 | 3.1 | 4.0 | 3.1 | 2.3 | 1.7 | 1.2 | 0.7 | * |
| 2003..... | 2.1 | * | * | 0.3 | 3.7 | 3.7 | 2.9 | 2.4 | 1.6 | 1.0 | 0.5 | * |
| 2002..... | 2.2 | * | * | 0.3 | 3.7 | 3.7 | 3.4 | 2.2 | 1.7 | 0.9 | 1.0 | * |
| 2001..... | 2.2 | * | * | 0.4 | 3.6 | 3.9 | 3.4 | 2.3 | 1.6 | 1.0 | 0.9 | * |
| 2000..... | 2.0 | * | * | 0.4 | 3.2 | 3.4 | 3.2 | 2.2 | 1.6 | 0.9 | 0.6 | * |
| 1999..... | 2.2 | * | * | 0.4 | 3.5 | 3.9 | 3.2 | 2.2 | 1.7 | 1.2 | 0.6 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | 1.1 | 1.2 | 1.2 | 0.9 | 0.7 | 0.7 | 0.8 | * |
| 2003..... | 0.8 | * | * | * | 1.0 | 1.3 | 1.3 | 1.0 | 0.7 | 0.4 | 0.6 | * |
| 2002..... | 0.9 | * | * | 0.3 | 1.2 | 1.4 | 1.3 | 1.1 | 0.7 | 0.6 | 0.7 | * |
| 2001..... | 0.9 | * | * | 0.3 | 1.2 | 1.5 | 1.4 | 0.8 | 0.8 | 0.7 | 0.6 | * |
| 2000..... | 0.9 | * | * | 0.2 | 1.2 | 1.4 | 1.4 | 0.9 | 0.8 | 0.7 | 0.6 | * |
| 1999..... | 0.9 | * | * | 0.2 | 1.2 | 1.4 | 1.5 | 0.8 | 0.6 | 0.7 | 0.6 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by discharge of firearms (*U01.4,X93-X95) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 16.6 | * | * | 1.3 | 40.3 | 39.4 | 16.0 | 8.6 | 4.7 | 2.4 | * | * |
| 2003..... | 17.3 | * | * | 1.5 | 43.3 | 40.5 | 16.5 | 8.9 | 4.0 | 2.5 | * | * |
| 2002..... | 17.1 | * | * | 1.2 | 43.1 | 39.7 | 16.7 | 8.6 | 4.2 | 2.4 | 2.1 | * |
| 2001..... | 16.3 | * | * | 0.9 | 43.0 | 36.4 | 15.5 | 7.7 | 3.7 | 2.6 | * | * |
| 2000..... | 16.0 | * | * | 1.0 | 42.6 | 34.5 | 15.4 | 8.5 | 3.6 | 2.8 | * | * |
| 1999..... | 15.9 | * | * | 1.3 | 43.6 | 32.4 | 14.4 | 8.1 | 4.5 | 2.4 | 2.4 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 31.4 | * | * | 1.9 | 74.1 | 75.4 | 29.8 | 15.5 | 8.5 | 5.6 | * | * |
| 2003..... | 32.6 | * | * | 2.1 | 79.4 | 77.4 | 30.1 | 16.4 | 7.4 | 4.8 | * | * |
| 2002..... | 31.9 | * | * | 1.6 | 78.6 | 75.6 | 30.0 | 15.3 | 7.7 | 4.7 | * | * |
| 2001..... | 30.6 | * | * | 1.3 | 80.6 | 69.4 | 26.7 | 14.0 | 7.1 | 5.5 | * | * |
| 2000..... | 30.1 | * | * | 1.6 | 79.1 | 65.7 | 27.4 | 15.3 | 6.5 | 5.5 | * | * |
| 1999..... | 29.3 | * | * | 1.8 | 80.0 | 61.1 | 24.6 | 14.1 | 9.1 | 4.7 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 3.1 | * | * | * | 6.0 | 6.4 | 3.8 | 2.6 | 1.6 | * | * | * |
| 2003..... | 3.4 | * | * | 0.9 | 6.7 | 6.9 | 4.4 | 2.6 | * | * | * | * |
| 2002..... | 3.7 | * | * | 0.8 | 7.5 | 7.2 | 5.0 | 2.8 | * | * | * | * |
| 2001..... | 3.2 | * | * | * | 5.6 | 6.5 | 5.5 | 2.2 | * | * | * | * |
| 2000..... | 3.3 | * | * | * | 6.7 | 6.4 | 4.7 | 2.6 | * | * | * | * |
| 1999..... | 3.7 | * | * | 0.7 | 7.8 | 6.6 | 5.5 | 3.0 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by sharp object (X99) | | | | | | | | | | | | |
| All origins 3/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | 0.1 | 1.2 | 1.2 | 1.0 | 0.8 | 0.4 | 0.4 | 0.4 | * |
| 2003..... | 0.7 | * | * | 0.1 | 1.1 | 1.2 | 1.1 | 0.8 | 0.4 | 0.4 | 0.4 | * |
| 2002..... | 0.7 | * | * | 0.1 | 1.0 | 1.2 | 1.1 | 0.8 | 0.5 | 0.4 | 0.4 | * |
| 2001..... | 0.7 | * | * | 0.1 | 1.2 | 1.2 | 1.0 | 0.7 | 0.4 | 0.4 | 0.4 | * |
| 2000..... | 0.6 | * | * | 0.1 | 1.0 | 1.1 | 1.0 | 0.6 | 0.4 | 0.3 | 0.5 | * |
| 1999..... | 0.7 | * | * | 0.1 | 1.1 | 1.2 | 1.0 | 0.7 | 0.4 | 0.3 | 0.3 | * |
| All origins 3/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | 1.8 | 1.7 | 1.4 | 1.1 | 0.7 | 0.6 | 0.6 | * |
| 2003..... | 1.0 | * | * | * | 1.7 | 1.7 | 1.5 | 1.2 | 0.6 | 0.5 | 0.6 | * |
| 2002..... | 1.0 | * | * | 0.1 | 1.6 | 1.7 | 1.5 | 1.2 | 0.8 | 0.7 | 0.4 | * |
| 2001..... | 1.0 | * | * | * | 1.8 | 1.7 | 1.3 | 1.0 | 0.6 | 0.5 | 0.4 | * |
| 2000..... | 0.9 | * | * | * | 1.6 | 1.5 | 1.4 | 0.8 | 0.7 | 0.5 | 0.7 | * |
| 1999..... | 1.0 | * | * | * | 1.6 | 1.7 | 1.6 | 1.0 | 0.7 | 0.5 | 0.5 | * |
| All origins 3/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.5 | 0.7 | 0.6 | 0.5 | 0.3 | 0.2 | 0.3 | * |
| 2003..... | 0.4 | * | * | * | 0.5 | 0.7 | 0.6 | 0.4 | 0.3 | 0.3 | 0.3 | * |
| 2002..... | 0.4 | * | * | * | 0.4 | 0.8 | 0.7 | 0.4 | 0.3 | 0.3 | 0.3 | * |
| 2001..... | 0.4 | * | * | * | 0.5 | 0.7 | 0.7 | 0.3 | 0.3 | 0.4 | 0.4 | * |
| 2000..... | 0.4 | * | * | * | 0.4 | 0.6 | 0.7 | 0.4 | 0.2 | * | 0.3 | * |
| 1999..... | 0.4 | * | * | * | 0.5 | 0.6 | 0.5 | 0.4 | 0.3 | 0.2 | * | * |
| Hispanic, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 2.1 | 1.8 | 1.2 | 1.1 | * | * | * | * |
| 2003..... | 1.1 | * | * | * | 1.9 | 1.8 | 1.4 | 1.0 | * | * | * | * |
| 2002..... | 1.1 | * | * | * | 2.0 | 1.7 | 1.5 | 1.0 | * | * | * | * |
| 2001..... | 1.1 | * | * | * | 2.0 | 1.7 | 1.4 | 1.0 | 1.2 | * | * | * |
| 2000..... | 1.2 | * | * | * | 2.0 | 1.9 | 1.8 | 1.0 | * | * | * | * |
| 1999..... | 1.0 | * | * | * | 1.7 | 1.9 | 1.3 | 1.1 | * | * | * | * |
| Hispanic, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 3.5 | 2.5 | 1.8 | 1.7 | * | * | * | * |
| 2003..... | 1.6 | * | * | * | 2.8 | 2.6 | 2.0 | 1.3 | * | * | * | * |
| 2002..... | 1.7 | * | * | * | 3.2 | 2.4 | 2.1 | 1.4 | * | * | * | * |
| 2001..... | 1.7 | * | * | * | 3.3 | 2.4 | 1.9 | 1.5 | * | * | * | * |
| 2000..... | 1.9 | * | * | * | 3.3 | 2.9 | 2.8 | 1.4 | * | * | * | * |
| 1999..... | 1.6 | * | * | * | 2.7 | 2.9 | 2.1 | 1.6 | * | * | * | * |
| Hispanic, female | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | * | 1.1 | * | * | * | * | * | * |
| 2003..... | 0.5 | * | * | * | 0.8 | 0.8 | * | * | * | * | * | * |
| 2002..... | 0.5 | * | * | * | 0.6 | 0.8 | 0.8 | * | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | 0.8 | 0.8 | * | * | * | * | * |
| 2000..... | 0.5 | * | * | * | * | 0.7 | * | * | * | * | * | * |
| 1999..... | 0.4 | * | * | * | * | 0.7 | * | * | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by sharp object (X99) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | * | * | 0.1 | 1.0 | 1.0 | 1.0 | 0.7 | 0.4 | 0.4 | 0.4 | * |
| 2003..... | 0.6 | * | * | 0.1 | 1.0 | 1.0 | 1.0 | 0.8 | 0.4 | 0.4 | 0.4 | * |
| 2002..... | 0.7 | * | * | 0.1 | 0.9 | 1.1 | 1.1 | 0.7 | 0.5 | 0.5 | 0.3 | * |
| 2001..... | 0.6 | * | * | * | 1.0 | 1.1 | 1.0 | 0.6 | 0.4 | 0.4 | 0.4 | * |
| 2000..... | 0.6 | * | * | 0.1 | 0.8 | 0.9 | 0.9 | 0.5 | 0.4 | 0.3 | 0.5 | * |
| 1999..... | 0.6 | * | * | 0.1 | 0.9 | 1.0 | 1.0 | 0.6 | 0.4 | 0.3 | 0.3 | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | 1.4 | 1.5 | 1.3 | 1.1 | 0.6 | 0.6 | 0.6 | * |
| 2003..... | 0.9 | * | * | * | 1.5 | 1.4 | 1.4 | 1.1 | 0.5 | 0.4 | 0.6 | * |
| 2002..... | 0.9 | * | * | 0.1 | 1.3 | 1.5 | 1.5 | 1.1 | 0.7 | 0.6 | * | * |
| 2001..... | 0.9 | * | * | * | 1.5 | 1.5 | 1.2 | 1.0 | 0.5 | 0.5 | * | * |
| 2000..... | 0.8 | * | * | * | 1.2 | 1.2 | 1.2 | 0.7 | 0.6 | 0.5 | 0.7 | * |
| 1999..... | 0.9 | * | * | * | 1.3 | 1.4 | 1.5 | 0.9 | 0.6 | 0.4 | 0.5 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.5 | 0.6 | 0.7 | 0.5 | 0.3 | 0.2 | 0.3 | * |
| 2003..... | 0.4 | * | * | * | 0.5 | 0.6 | 0.6 | 0.4 | 0.3 | 0.3 | 0.3 | * |
| 2002..... | 0.4 | * | * | * | 0.4 | 0.8 | 0.7 | 0.3 | 0.3 | 0.3 | 0.3 | * |
| 2001..... | 0.4 | * | * | * | 0.6 | 0.6 | 0.7 | 0.3 | 0.3 | 0.4 | 0.4 | * |
| 2000..... | 0.3 | * | * | * | 0.4 | 0.5 | 0.7 | 0.3 | 0.2 | * | * | * |
| 1999..... | 0.4 | * | * | * | 0.5 | 0.6 | 0.5 | 0.3 | 0.2 | 0.3 | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | * | * | * | 0.5 | 0.5 | 0.6 | 0.4 | 0.3 | 0.3 | 0.3 | * |
| 2003..... | 0.3 | * | * | * | 0.4 | 0.5 | 0.5 | 0.5 | 0.2 | 0.3 | 0.3 | * |
| 2002..... | 0.4 | * | * | * | 0.4 | 0.5 | 0.6 | 0.4 | 0.3 | 0.4 | 0.2 | * |
| 2001..... | 0.4 | * | * | * | 0.6 | 0.6 | 0.5 | 0.4 | 0.3 | 0.3 | 0.3 | * |
| 2000..... | 0.3 | * | * | * | 0.4 | 0.5 | 0.5 | 0.3 | 0.2 | 0.2 | 0.3 | * |
| 1999..... | 0.4 | * | * | * | 0.5 | 0.6 | 0.5 | 0.4 | 0.3 | 0.2 | 0.2 | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | * | * | * | 0.7 | 0.6 | 0.7 | 0.5 | 0.4 | 0.4 | * | * |
| 2003..... | 0.4 | * | * | * | 0.6 | 0.7 | 0.6 | 0.6 | 0.3 | * | * | * |
| 2002..... | 0.5 | * | * | * | 0.6 | 0.6 | 0.9 | 0.6 | 0.4 | 0.5 | * | * |
| 2001..... | 0.5 | * | * | * | 0.8 | 0.8 | 0.6 | 0.5 | 0.4 | 0.4 | * | * |
| 2000..... | 0.4 | * | * | * | 0.6 | 0.6 | 0.6 | 0.4 | 0.4 | 0.3 | 0.6 | * |
| 1999..... | 0.5 | * | * | * | 0.7 | 0.8 | 0.7 | 0.6 | 0.4 | 0.3 | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | * | * | * | 0.3 | 0.4 | 0.4 | 0.3 | 0.2 | * | * | * |
| 2003..... | 0.2 | * | * | * | 0.2 | 0.4 | 0.4 | 0.3 | 0.2 | 0.3 | * | * |
| 2002..... | 0.2 | * | * | * | 0.2 | 0.4 | 0.4 | 0.2 | 0.3 | 0.3 | * | * |
| 2001..... | 0.3 | * | * | * | 0.3 | 0.4 | 0.4 | 0.2 | * | 0.3 | 0.3 | * |
| 2000..... | 0.2 | * | * | * | 0.2 | 0.4 | 0.4 | 0.2 | * | * | * | * |
| 1999..... | 0.2 | * | * | * | 0.3 | 0.4 | 0.4 | 0.2 | 0.2 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by sharp object (X99) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.2 | * | * | * | 2.8 | 3.5 | 3.5 | 3.0 | 1.6 | 1.6 | 2.4 | * |
| 2003..... | 2.2 | * | * | * | 3.3 | 3.5 | 4.0 | 2.8 | 1.5 | 1.2 | * | * |
| 2002..... | 2.2 | * | * | * | 2.7 | 3.9 | 3.6 | 3.0 | 1.8 | 1.3 | * | * |
| 2001..... | 2.1 | * | * | * | 3.0 | 3.6 | 3.6 | 2.4 | 1.3 | 1.2 | * | * |
| 2000..... | 1.9 | * | * | * | 2.5 | 2.8 | 3.7 | 2.3 | 1.6 | 1.4 | * | * |
| 1999..... | 2.1 | * | * | * | 2.9 | 3.2 | 3.7 | 2.4 | 1.4 | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 3.2 | * | * | * | 4.1 | 5.6 | 4.9 | 4.6 | 2.8 | * | * | * |
| 2003..... | 3.5 | * | * | * | 5.1 | 5.2 | 6.3 | 4.6 | 2.4 | * | * | * |
| 2002..... | 3.3 | * | * | * | 4.3 | 5.8 | 4.9 | 5.1 | 3.3 | * | * | * |
| 2001..... | 3.1 | * | * | * | 4.5 | 5.4 | 5.1 | 4.5 | 2.0 | * | * | * |
| 2000..... | 2.9 | * | * | * | 3.8 | 4.5 | 5.0 | 3.7 | 2.5 | * | * | * |
| 1999..... | 3.2 | * | * | * | 4.3 | 4.7 | 6.4 | 3.6 | 2.7 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 1.4 | 1.5 | 2.2 | 1.6 | * | * | * | * |
| 2003..... | 1.1 | * | * | * | 1.6 | 1.9 | 2.0 | 1.2 | * | * | * | * |
| 2002..... | 1.2 | * | * | * | 1.2 | 2.2 | 2.5 | 1.1 | * | * | * | * |
| 2001..... | 1.1 | * | * | * | 1.6 | 1.8 | 2.2 | * | * | * | * | * |
| 2000..... | 1.1 | * | * | * | 1.3 | 1.3 | 2.6 | 1.1 | * | * | * | * |
| 1999..... | 1.0 | * | * | * | 1.6 | 1.8 | 1.3 | 1.3 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Sexual assault (homicide) by bodily force (Y05)

All origins 3/, both sexes

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), all cells contain an asterisk.

All origins 3/, male

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), all cells contain an asterisk.

All origins 3/, female

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), all cells contain an asterisk.

Hispanic, both sexes

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), all cells contain an asterisk.

Hispanic, male

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), all cells contain an asterisk.

Hispanic, female

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+), all cells contain an asterisk.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.8 | 3.7 | 1.3 | 0.1 | 0.7 | 0.8 | 1.0 | 1.1 | 0.7 | 0.7 | 0.7 | 0.9 |
| 2003..... | 0.8 | 4.7 | 1.3 | 0.2 | 0.7 | 0.8 | 1.1 | 1.0 | 0.7 | 0.7 | 0.9 | 1.0 |
| 2002..... | 0.8 | 3.9 | 1.5 | 0.1 | 0.7 | 0.8 | 1.0 | 1.0 | 0.8 | 0.7 | 0.7 | 0.9 |
| 2001 5/..... | 2.0 | 3.6 | 1.7 | 0.2 | 1.1 | 3.2 | 3.5 | 2.7 | 1.8 | 0.9 | 0.9 | 1.2 |
| 2000..... | 1.0 | 4.5 | 1.4 | 0.2 | 1.0 | 1.1 | 1.3 | 1.1 | 0.7 | 0.7 | 1.0 | 0.9 |
| 1999..... | 0.9 | 3.8 | 1.3 | 0.3 | 0.9 | 1.1 | 1.3 | 1.0 | 0.7 | 0.7 | 1.0 | 1.1 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.1 | 3.5 | 1.3 | 0.1 | 0.9 | 1.1 | 1.3 | 1.6 | 1.1 | 1.1 | 1.0 | * |
| 2003..... | 1.2 | 5.0 | 1.6 | 0.2 | 0.9 | 1.0 | 1.5 | 1.5 | 1.2 | 1.1 | 1.4 | 1.8 |
| 2002..... | 1.1 | 4.3 | 1.6 | 0.1 | 0.9 | 1.1 | 1.4 | 1.4 | 1.3 | 1.0 | 1.2 | * |
| 2001 5/..... | 2.8 | 3.9 | 1.8 | 0.2 | 1.4 | 4.6 | 5.3 | 4.0 | 2.7 | 1.4 | 1.2 | 1.6 |
| 2000..... | 1.3 | 5.0 | 1.6 | 0.2 | 1.3 | 1.5 | 1.7 | 1.6 | 1.2 | 1.1 | 1.4 | * |
| 1999..... | 1.2 | 4.1 | 1.5 | 0.2 | 1.1 | 1.2 | 1.7 | 1.5 | 1.1 | 1.0 | 1.6 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.6 | 3.9 | 1.3 | 0.1 | 0.5 | 0.5 | 0.7 | 0.5 | 0.4 | 0.4 | 0.5 | 0.8 |
| 2003..... | 0.5 | 4.3 | 1.0 | 0.1 | 0.5 | 0.6 | 0.7 | 0.6 | 0.3 | 0.4 | 0.5 | 0.6 |
| 2002..... | 0.5 | 3.6 | 1.3 | 0.2 | 0.5 | 0.6 | 0.6 | 0.5 | 0.3 | 0.4 | 0.4 | 0.8 |
| 2001 5/..... | 1.1 | 3.3 | 1.5 | 0.2 | 0.8 | 1.7 | 1.8 | 1.4 | 0.9 | 0.5 | 0.7 | 1.0 |
| 2000..... | 0.7 | 4.0 | 1.2 | 0.3 | 0.7 | 0.8 | 0.9 | 0.5 | 0.3 | 0.3 | 0.8 | 0.7 |
| 1999..... | 0.7 | 3.6 | 1.2 | 0.3 | 0.7 | 0.9 | 0.9 | 0.5 | 0.4 | 0.4 | 0.6 | 0.9 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.6 | 2.7 | 0.7 | * | 0.5 | 0.6 | 0.7 | 0.8 | 0.6 | 0.6 | 0.6 | 0.9 |
| 2003..... | 0.6 | 3.5 | 0.7 | 0.1 | 0.5 | 0.5 | 0.8 | 0.7 | 0.5 | 0.5 | 0.8 | 1.0 |
| 2002..... | 0.6 | 2.6 | 0.9 | 0.1 | 0.5 | 0.5 | 0.8 | 0.7 | 0.6 | 0.5 | 0.6 | 0.8 |
| 2001 5/..... | 1.7 | 2.6 | 1.0 | 0.2 | 0.9 | 2.9 | 3.3 | 2.4 | 1.6 | 0.8 | 0.8 | 1.0 |
| 2000..... | 0.6 | 2.7 | 0.7 | 0.2 | 0.6 | 0.7 | 0.9 | 0.7 | 0.5 | 0.5 | 0.8 | 0.8 |
| 1999..... | 0.7 | 2.7 | 0.9 | 0.2 | 0.6 | 0.7 | 0.8 | 0.7 | 0.6 | 0.6 | 0.8 | 0.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.8 | 3.0 | 0.8 | * | 0.6 | 0.7 | 0.9 | 1.2 | 0.8 | 0.8 | 0.8 | * |
| 2003..... | 0.8 | 3.8 | 0.8 | * | 0.6 | 0.6 | 1.0 | 1.0 | 0.9 | 0.8 | 1.3 | 1.8 |
| 2002..... | 0.8 | 3.0 | 1.0 | * | 0.6 | 0.7 | 1.0 | 0.9 | 0.9 | 0.8 | 0.8 | * |
| 2001 5/..... | 2.6 | 2.9 | 1.1 | * | 1.0 | 4.3 | 5.2 | 3.5 | 2.4 | 1.2 | 1.0 | * |
| 2000..... | 0.8 | 3.4 | 0.9 | 0.2 | 0.6 | 0.8 | 1.1 | 1.0 | 0.8 | 0.8 | 1.0 | * |
| 1999..... | 0.8 | 3.0 | 1.0 | * | 0.7 | 0.7 | 1.1 | 1.0 | 0.9 | 0.8 | 1.4 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.4 | 2.3 | 0.6 | * | 0.5 | 0.5 | 0.5 | 0.4 | 0.3 | 0.4 | 0.5 | 0.8 |
| 2003..... | 0.4 | 3.2 | 0.6 | * | 0.4 | 0.5 | 0.5 | 0.5 | 0.2 | 0.4 | 0.5 | * |
| 2002..... | 0.4 | 2.1 | 0.7 | * | 0.4 | 0.4 | 0.5 | 0.5 | 0.3 | 0.3 | 0.4 | 0.7 |
| 2001 5/..... | 0.9 | 2.3 | 0.9 | 0.2 | 0.7 | 1.4 | 1.4 | 1.2 | 0.8 | 0.4 | 0.6 | 0.9 |
| 2000..... | 0.5 | 2.0 | 0.5 | 0.2 | 0.5 | 0.5 | 0.6 | 0.5 | 0.3 | 0.3 | 0.7 | * |
| 1999..... | 0.5 | 2.3 | 0.8 | 0.2 | 0.6 | 0.6 | 0.6 | 0.5 | 0.3 | 0.4 | 0.5 | 0.9 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 2.0 | 7.7 | 3.7 | 0.4 | 1.4 | 2.2 | 2.4 | 3.0 | 2.0 | 1.5 | 2.1 | * |
| 2003..... | 2.2 | 9.2 | 3.7 | 0.5 | 1.5 | 2.2 | 2.8 | 2.9 | 2.2 | 1.8 | 2.2 | * |
| 2002..... | 2.2 | 9.3 | 3.9 | 0.4 | 1.6 | 2.2 | 2.6 | 3.1 | 2.0 | 1.6 | 2.6 | * |
| 2001 5/..... | 3.2 | 8.2 | 4.3 | 0.3 | 1.8 | 4.3 | 5.1 | 5.0 | 2.9 | 2.5 | 2.3 | * |
| 2000..... | 3.0 | 12.0 | 4.0 | 0.6 | 3.1 | 3.4 | 3.9 | 3.5 | 2.4 | 2.2 | 3.4 | * |
| 1999..... | 2.5 | 9.0 | 3.4 | 0.6 | 1.9 | 3.2 | 3.9 | 2.9 | 1.6 | 2.1 | 2.4 | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.8 | * | 3.4 | * | 1.9 | 3.5 | 3.4 | 5.1 | 3.4 | 2.9 | * | * |
| 2003..... | 3.1 | 9.0 | 4.7 | 0.6 | 2.1 | 3.0 | 4.2 | 4.8 | 4.2 | 3.6 | * | * |
| 2002..... | 3.2 | 9.1 | 4.3 | * | 2.3 | 2.9 | 3.8 | 5.5 | 4.1 | 2.9 | 6.2 | * |
| 2001 5/..... | 4.4 | 8.3 | 5.1 | * | 2.5 | 6.0 | 6.5 | 7.4 | 4.8 | 4.3 | * | * |
| 2000..... | 4.3 | 11.9 | 4.1 | * | 4.4 | 5.2 | 5.4 | 6.4 | 4.8 | 4.3 | 6.2 | * |
| 1999..... | 3.5 | 9.4 | 3.7 | 0.6 | 3.0 | 4.2 | 5.4 | 5.0 | 2.5 | 4.2 | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.3 | 10.0 | 3.9 | * | 0.9 | 1.0 | 1.5 | 1.2 | * | * | * | * |
| 2003..... | 1.3 | 9.3 | 2.8 | * | 1.0 | 1.5 | 1.5 | 1.3 | * | * | * | * |
| 2002..... | 1.3 | 9.5 | 3.4 | * | 0.9 | 1.5 | 1.6 | 1.1 | * | * | * | * |
| 2001 5/..... | 2.1 | 8.1 | 3.5 | * | 1.0 | 2.7 | 3.8 | 3.0 | * | * | * | * |
| 2000..... | 1.8 | 12.0 | 4.0 | 0.8 | 1.7 | 1.9 | 2.5 | 1.0 | * | * | * | * |
| 1999..... | 1.6 | 8.6 | 3.0 | 0.6 | 0.9 | 2.2 | 2.6 | 1.0 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Event of undetermined intent (Y10-Y34,Y87.2,Y89.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.8 | 2.8 | 0.3 | 0.2 | 1.3 | 2.2 | 3.4 | 3.3 | 1.4 | 0.9 | 0.9 | 1.6 |
| 2003..... | 1.9 | 2.2 | 0.2 | 0.1 | 1.5 | 2.3 | 3.9 | 3.2 | 1.3 | 0.7 | 0.7 | 1.1 |
| 2002..... | 1.8 | 2.7 | 0.4 | 0.2 | 1.3 | 2.3 | 3.6 | 3.0 | 1.2 | 0.7 | 0.8 | 1.4 |
| 2001..... | 1.6 | 2.4 | 0.3 | 0.1 | 1.1 | 2.0 | 3.3 | 2.6 | 0.9 | 0.5 | 0.9 | 0.9 |
| 2000..... | 1.5 | 1.9 | 0.3 | 0.2 | 1.0 | 1.8 | 3.0 | 2.4 | 0.9 | 0.6 | 0.6 | 1.4 |
| 1999..... | 1.5 | 1.6 | 0.3 | 0.1 | 1.2 | 2.0 | 3.1 | 2.3 | 0.9 | 0.6 | 0.8 | 1.4 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 2.4 | 3.3 | 0.3 | 0.2 | 1.9 | 3.1 | 4.3 | 4.1 | 1.7 | 1.1 | 1.0 | 1.7 |
| 2003..... | 2.5 | 3.0 | * | 0.2 | 2.2 | 3.1 | 4.9 | 4.2 | 1.7 | 0.8 | 0.9 | * |
| 2002..... | 2.3 | 2.8 | 0.3 | 0.2 | 1.8 | 3.2 | 4.7 | 3.6 | 1.6 | 1.0 | 0.9 | 2.1 |
| 2001..... | 2.1 | 2.7 | * | 0.2 | 1.7 | 2.8 | 4.4 | 3.5 | 1.2 | 0.7 | 1.2 | * |
| 2000..... | 2.0 | 1.6 | 0.4 | 0.3 | 1.5 | 2.6 | 4.2 | 3.2 | 1.3 | 0.8 | 1.0 | 1.7 |
| 1999..... | 2.1 | 1.9 | * | 0.2 | 1.8 | 2.7 | 4.2 | 3.1 | 1.3 | 0.9 | 1.1 | 2.1 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.3 | 2.3 | 0.3 | * | 0.6 | 1.4 | 2.6 | 2.5 | 1.1 | 0.7 | 0.9 | 1.5 |
| 2003..... | 1.3 | 1.3 | * | * | 0.8 | 1.5 | 2.8 | 2.3 | 1.0 | 0.7 | 0.6 | 1.0 |
| 2002..... | 1.3 | 2.6 | 0.4 | 0.2 | 0.7 | 1.4 | 2.6 | 2.3 | 0.9 | 0.5 | 0.7 | 1.1 |
| 2001..... | 1.0 | 2.0 | 0.4 | * | 0.5 | 1.2 | 2.2 | 1.7 | 0.6 | 0.4 | 0.7 | 0.8 |
| 2000..... | 0.9 | 2.2 | * | * | 0.5 | 1.1 | 1.9 | 1.5 | 0.6 | 0.3 | 0.4 | 1.2 |
| 1999..... | 0.9 | * | * | * | 0.5 | 1.2 | 2.0 | 1.5 | 0.6 | 0.4 | 0.6 | 1.1 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | 2.3 | 0.3 | 0.1 | 1.5 | 2.5 | 3.6 | 3.4 | 1.4 | 0.8 | 0.9 | 1.6 |
| 2003..... | 2.0 | 2.0 | * | 0.1 | 1.7 | 2.6 | 4.1 | 3.3 | 1.3 | 0.7 | 0.7 | 1.2 |
| 2002..... | 1.8 | 1.8 | 0.4 | 0.2 | 1.4 | 2.4 | 3.7 | 2.9 | 1.2 | 0.7 | 0.7 | 1.5 |
| 2001..... | 1.6 | 1.7 | 0.3 | 0.1 | 1.3 | 2.2 | 3.3 | 2.5 | 0.9 | 0.5 | 0.9 | 1.0 |
| 2000..... | 1.4 | 1.5 | 0.2 | 0.1 | 1.1 | 1.9 | 3.0 | 2.3 | 0.8 | 0.5 | 0.7 | 1.3 |
| 1999..... | 1.4 | 1.3 | * | 0.1 | 1.2 | 2.0 | 3.0 | 2.1 | 0.9 | 0.6 | 0.8 | 1.4 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 2.4 | 2.7 | * | 0.2 | 2.2 | 3.4 | 4.5 | 4.1 | 1.6 | 1.0 | 0.9 | 1.9 |
| 2003..... | 2.5 | 2.4 | * | 0.2 | 2.4 | 3.4 | 5.1 | 4.0 | 1.5 | 0.7 | 0.9 | * |
| 2002..... | 2.3 | 1.9 | * | 0.2 | 2.0 | 3.4 | 4.8 | 3.4 | 1.5 | 0.8 | 0.9 | 2.2 |
| 2001..... | 2.1 | 2.0 | * | 0.2 | 1.8 | 3.0 | 4.3 | 3.3 | 1.1 | 0.7 | 1.2 | * |
| 2000..... | 2.0 | * | * | 0.2 | 1.6 | 2.6 | 4.1 | 3.0 | 1.1 | 0.7 | 1.0 | * |
| 1999..... | 2.0 | * | * | 0.2 | 1.8 | 2.7 | 4.0 | 2.6 | 1.1 | 0.8 | 1.1 | 2.1 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.4 | 2.0 | * | * | 0.7 | 1.6 | 2.8 | 2.7 | 1.2 | 0.6 | 0.9 | 1.5 |
| 2003..... | 1.4 | * | * | * | 1.0 | 1.7 | 3.1 | 2.5 | 1.1 | 0.6 | 0.6 | 1.1 |
| 2002..... | 1.3 | 1.8 | * | 0.2 | 0.8 | 1.5 | 2.7 | 2.4 | 0.9 | 0.6 | 0.6 | 1.1 |
| 2001..... | 1.1 | * | * | * | 0.6 | 1.4 | 2.4 | 1.8 | 0.7 | 0.4 | 0.6 | 0.8 |
| 2000..... | 0.9 | * | * | * | 0.6 | 1.2 | 1.9 | 1.5 | 0.6 | 0.3 | 0.4 | 1.2 |
| 1999..... | 1.0 | * | * | * | 0.5 | 1.3 | 1.9 | 1.5 | 0.7 | 0.4 | 0.6 | 1.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Event of undetermined intent (Y10-Y34,Y87.2,Y89.9) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.9 | 4.9 | * | 0.4 | 0.9 | 1.9 | 3.4 | 4.0 | 1.8 | 1.3 | * | * |
| 2003..... | 1.8 | 3.3 | * | * | 0.8 | 1.8 | 3.9 | 4.1 | 2.0 | 1.3 | * | * |
| 2002..... | 2.0 | 5.7 | * | * | 0.9 | 2.3 | 4.3 | 4.3 | 1.5 | * | * | * |
| 2001..... | 1.8 | 5.9 | * | * | 0.7 | 1.7 | 4.0 | 3.8 | 1.2 | * | * | * |
| 2000..... | 1.9 | * | * | 0.3 | 0.9 | 2.0 | 4.1 | 3.7 | 1.8 | * | * | * |
| 1999..... | 2.0 | * | * | * | 1.3 | 2.3 | 4.3 | 4.2 | 1.2 | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 2.7 | * | * | * | 1.4 | 2.8 | 4.8 | 6.0 | 3.0 | * | * | * |
| 2003..... | 2.8 | * | * | * | 1.4 | 3.0 | 5.4 | 6.6 | 3.3 | * | * | * |
| 2002..... | 2.9 | * | * | * | 1.3 | 3.4 | 6.2 | 6.5 | 2.6 | * | * | * |
| 2001..... | 2.8 | 6.4 | * | * | 1.3 | 2.7 | 6.3 | 6.3 | 2.4 | * | * | * |
| 2000..... | 2.9 | * | * | * | 1.6 | 3.2 | 6.1 | 6.0 | 3.0 | * | * | * |
| 1999..... | 3.1 | * | * | * | 2.0 | 3.7 | 6.6 | 7.3 | 2.6 | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 1.0 | 2.2 | 2.3 | * | * | * | * |
| 2003..... | 1.0 | * | * | * | * | * | 2.5 | 2.0 | * | * | * | * |
| 2002..... | 1.2 | * | * | * | * | 1.3 | 2.7 | 2.4 | * | * | * | * |
| 2001..... | 0.9 | * | * | * | * | 0.9 | 1.9 | 1.5 | * | * | * | * |
| 2000..... | 1.0 | * | * | * | * | 0.9 | 2.4 | 1.8 | * | * | * | * |
| 1999..... | 0.9 | * | * | * | * | 1.0 | 2.4 | 1.6 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | 0.8 | 1.7 | 2.7 | 2.5 | 0.9 | 0.3 | * | * |
| 2003..... | 1.3 | * | * | * | 1.0 | 1.8 | 3.2 | 2.5 | 0.8 | 0.2 | * | * |
| 2002..... | 1.2 | * | * | * | 0.8 | 1.7 | 2.8 | 2.3 | 0.7 | 0.2 | * | * |
| 2001..... | 1.1 | * | * | * | 0.7 | 1.5 | 2.6 | 1.9 | 0.5 | 0.2 | 0.2 | * |
| 2000..... | 0.9 | * | * | * | 0.5 | 1.2 | 2.4 | 1.7 | 0.4 | * | * | * |
| 1999..... | 1.0 | * | * | * | 0.6 | 1.4 | 2.4 | 1.6 | 0.5 | 0.2 | * | * |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.5 | * | * | * | 1.2 | 2.3 | 3.3 | 3.0 | 0.8 | 0.3 | * | * |
| 2003..... | 1.7 | * | * | * | 1.5 | 2.4 | 4.0 | 3.1 | 0.9 | * | * | * |
| 2002..... | 1.5 | * | * | * | 1.1 | 2.3 | 3.5 | 2.6 | 0.8 | * | * | * |
| 2001..... | 1.4 | * | * | * | 1.0 | 1.9 | 3.4 | 2.6 | 0.6 | * | * | * |
| 2000..... | 1.3 | * | * | * | 0.8 | 1.7 | 3.4 | 2.3 | 0.5 | * | * | * |
| 1999..... | 1.3 | * | * | * | 0.8 | 1.9 | 3.2 | 2.1 | 0.6 | * | * | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | 0.4 | 1.1 | 2.2 | 2.1 | 0.9 | 0.3 | * | * |
| 2003..... | 1.0 | * | * | * | 0.6 | 1.2 | 2.5 | 1.9 | 0.7 | 0.3 | * | * |
| 2002..... | 0.9 | * | * | * | 0.5 | 1.1 | 2.1 | 2.0 | 0.6 | 0.2 | * | * |
| 2001..... | 0.7 | * | * | * | 0.4 | 1.0 | 1.9 | 1.3 | 0.5 | * | * | * |
| 2000..... | 0.6 | * | * | * | 0.3 | 0.8 | 1.5 | 1.2 | 0.3 | * | * | * |
| 1999..... | 0.7 | * | * | * | 0.3 | 0.9 | 1.6 | 1.2 | 0.4 | * | * | * |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | 1.0 | 2.0 | 2.9 | 2.6 | 0.9 | 0.3 | * | * |
| 2003..... | 1.5 | * | * | * | 1.3 | 2.1 | 3.5 | 2.5 | 0.8 | 0.2 | * | * |
| 2002..... | 1.3 | * | * | * | 1.0 | 1.9 | 3.0 | 2.3 | 0.7 | 0.2 | * | * |
| 2001..... | 1.1 | * | * | * | 0.8 | 1.6 | 2.7 | 1.9 | 0.5 | 0.1 | * | * |
| 2000..... | 1.0 | * | * | * | 0.7 | 1.4 | 2.4 | 1.7 | 0.4 | * | * | * |
| 1999..... | 1.0 | * | * | * | 0.7 | 1.5 | 2.3 | 1.5 | 0.5 | 0.2 | * | * |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.6 | * | * | * | 1.5 | 2.7 | 3.5 | 3.0 | 0.8 | * | * | * |
| 2003..... | 1.8 | * | * | * | 1.8 | 2.8 | 4.3 | 3.0 | 0.8 | * | * | * |
| 2002..... | 1.6 | * | * | * | 1.3 | 2.6 | 3.7 | 2.5 | 0.8 | * | * | * |
| 2001..... | 1.4 | * | * | * | 1.2 | 2.1 | 3.3 | 2.5 | 0.5 | * | * | * |
| 2000..... | 1.3 | * | * | * | 0.9 | 1.8 | 3.4 | 2.2 | 0.5 | * | * | * |
| 1999..... | 1.2 | * | * | * | 1.0 | 2.0 | 3.1 | 1.8 | 0.5 | * | * | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 0.5 | 1.3 | 2.4 | 2.3 | 1.0 | 0.4 | * | * |
| 2003..... | 1.1 | * | * | * | 0.7 | 1.5 | 2.7 | 2.1 | 0.8 | 0.3 | * | * |
| 2002..... | 1.0 | * | * | * | 0.6 | 1.3 | 2.3 | 2.1 | 0.7 | 0.3 | * | * |
| 2001..... | 0.8 | * | * | * | 0.5 | 1.2 | 2.0 | 1.4 | 0.5 | * | * | * |
| 2000..... | 0.6 | * | * | * | 0.4 | 0.9 | 1.5 | 1.2 | 0.4 | * | * | * |
| 1999..... | 0.7 | * | * | * | 0.4 | 1.1 | 1.6 | 1.2 | 0.5 | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 1.0 | 2.6 | 2.7 | 1.0 | * | * | * |
| 2003..... | 1.2 | * | * | * | * | 1.0 | 3.0 | 3.2 | 1.0 | * | * | * |
| 2002..... | 1.3 | * | * | * | 0.3 | 1.4 | 3.2 | 3.3 | 0.9 | * | * | * |
| 2001..... | 1.1 | * | * | * | * | 1.2 | 3.1 | 2.8 | * | * | * | * |
| 2000..... | 1.1 | * | * | * | * | 1.1 | 3.2 | 2.7 | * | * | * | * |
| 1999..... | 1.2 | * | * | * | * | 1.4 | 3.3 | 3.0 | * | * | * | * |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | 1.4 | 3.6 | 3.8 | 1.8 | * | * | * |
| 2003..... | 1.7 | * | * | * | * | 1.6 | 4.1 | 5.0 | 1.7 | * | * | * |
| 2002..... | 1.7 | * | * | * | * | 1.9 | 4.3 | 4.6 | * | * | * | * |
| 2001..... | 1.7 | * | * | * | * | 1.7 | 4.9 | 4.6 | * | * | * | * |
| 2000..... | 1.6 | * | * | * | * | 1.7 | 4.6 | 4.1 | * | * | * | * |
| 1999..... | 1.8 | * | * | * | * | 2.3 | 4.9 | 5.1 | * | * | * | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 0.7 | * | * | * | * | * | 1.8 | 1.8 | * | * | * | * |
| 2003..... | 0.7 | * | * | * | * | * | 2.1 | 1.6 | * | * | * | * |
| 2002..... | 0.9 | * | * | * | * | 1.0 | 2.3 | 2.1 | * | * | * | * |
| 2001..... | 0.5 | * | * | * | * | 0.8 | 1.5 | 1.2 | * | * | * | * |
| 2000..... | 0.6 | * | * | * | * | * | 2.0 | 1.5 | * | * | * | * |
| 1999..... | 0.6 | * | * | * | * | * | 2.0 | 1.3 | * | * | * | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex
All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Non-Hispanic 4/, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, both sexes.

Non-Hispanic 4/, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, male.

Non-Hispanic 4/, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic 4/, female.

Non-Hispanic white, both sexes

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, both sexes.

Non-Hispanic white, male

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, male.

Non-Hispanic white, female

Table with 13 columns (Year, All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over) and 7 rows (2004-1999) for Non-Hispanic white, female.

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | 2.5 | 0.2 | 0.1 | 0.2 | 0.4 | 0.5 | 0.6 | 0.4 | 0.4 | 0.7 | 1.2 |
| 2003..... | 0.4 | 1.9 | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.4 | 0.4 | 0.5 | 0.8 |
| 2002..... | 0.4 | 2.2 | 0.3 | 0.1 | 0.3 | 0.3 | 0.5 | 0.5 | 0.4 | 0.4 | 0.5 | 1.1 |
| 2001..... | 0.3 | 2.0 | 0.3 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.3 | 0.3 | 0.5 | 0.6 |
| 2000..... | 0.4 | 1.7 | 0.2 | 0.1 | 0.3 | 0.4 | 0.4 | 0.5 | 0.4 | 0.3 | 0.4 | 1.0 |
| 1999..... | 0.3 | 1.4 | 0.2 | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.6 | 3.0 | * | 0.2 | 0.4 | 0.5 | 0.7 | 0.9 | 0.6 | 0.5 | 0.7 | * |
| 2003..... | 0.5 | 2.7 | * | 0.1 | 0.3 | 0.5 | 0.6 | 0.8 | 0.6 | 0.4 | 0.6 | * |
| 2002..... | 0.5 | 2.2 | * | 0.1 | 0.3 | 0.5 | 0.8 | 0.7 | 0.6 | 0.6 | 0.7 | 1.7 |
| 2001..... | 0.5 | 2.2 | * | 0.1 | 0.4 | 0.5 | 0.6 | 0.6 | 0.4 | 0.4 | 0.7 | * |
| 2000..... | 0.5 | 1.4 | * | 0.2 | 0.4 | 0.5 | 0.5 | 0.7 | 0.6 | 0.5 | 0.6 | * |
| 1999..... | 0.5 | 1.6 | * | 0.1 | 0.4 | 0.5 | 0.6 | 0.7 | 0.5 | 0.4 | 0.6 | * |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 2.1 | * | * | * | 0.2 | 0.3 | 0.3 | 0.2 | 0.3 | 0.7 | 1.2 |
| 2003..... | 0.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.2 | 0.4 | 0.4 | 0.8 |
| 2002..... | 0.3 | 2.2 | * | 0.1 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.4 | 0.9 |
| 2001..... | 0.2 | 1.7 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | * | * | 0.4 | * |
| 2000..... | 0.2 | 2.0 | * | * | 0.1 | 0.2 | 0.3 | 0.3 | 0.2 | * | 0.3 | 0.9 |
| 1999..... | 0.2 | * | * | * | * | 0.2 | 0.2 | 0.2 | * | * | 0.3 | 0.8 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.4 | 2.2 | 0.2 | * | 0.2 | 0.3 | 0.5 | 0.5 | 0.3 | 0.3 | 0.7 | 1.2 |
| 2003..... | 0.3 | 1.7 | * | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.3 | 0.3 | 0.4 | 0.9 |
| 2002..... | 0.4 | 1.6 | 0.3 | 0.1 | 0.3 | 0.3 | 0.5 | 0.4 | 0.3 | 0.3 | 0.5 | 1.2 |
| 2001..... | 0.3 | 1.3 | 0.2 | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.2 | 0.2 | 0.5 | 0.7 |
| 2000..... | 0.3 | 1.4 | * | 0.1 | 0.3 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.8 |
| 1999..... | 0.3 | 1.2 | * | 0.1 | 0.2 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.9 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.5 | 2.5 | * | * | 0.4 | 0.4 | 0.7 | 0.8 | 0.5 | 0.4 | 0.7 | * |
| 2003..... | 0.4 | 2.1 | * | * | 0.3 | 0.4 | 0.6 | 0.7 | 0.4 | 0.4 | 0.6 | * |
| 2002..... | 0.5 | 1.7 | * | * | 0.3 | 0.4 | 0.7 | 0.5 | 0.5 | 0.4 | 0.6 | 1.8 |
| 2001..... | 0.4 | * | * | * | 0.3 | 0.4 | 0.6 | 0.6 | 0.4 | 0.3 | 0.7 | * |
| 2000..... | 0.4 | * | * | 0.2 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | 0.4 | 0.6 | * |
| 1999..... | 0.4 | * | * | * | 0.4 | 0.4 | 0.5 | 0.6 | 0.5 | 0.4 | 0.5 | * |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.3 | 1.8 | * | * | * | 0.2 | 0.3 | 0.3 | 0.2 | * | 0.7 | 1.2 |
| 2003..... | 0.2 | * | * | * | 0.2 | 0.2 | 0.2 | 0.3 | 0.2 | 0.3 | 0.4 | 0.8 |
| 2002..... | 0.3 | * | * | * | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | * | 0.4 | 0.9 |
| 2001..... | 0.2 | * | * | * | * | * | 0.2 | 0.3 | * | * | 0.4 | * |
| 2000..... | 0.2 | * | * | * | * | 0.2 | 0.3 | 0.3 | * | * | * | 0.8 |
| 1999..... | 0.2 | * | * | * | * | 0.2 | 0.2 | 0.2 | * | * | 0.3 | * |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Complications of medical and surgical care (Y40-Y84,Y88) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 3.1 | 6.1 | 10.8 |
| 2003..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.4 | 2.9 | 6.3 | 11.4 |
| 2002..... | 1.1 | * | * | 0.1 | 0.1 | 0.2 | 0.4 | 0.7 | 1.3 | 3.1 | 6.2 | 11.6 |
| 2001..... | 1.2 | 0.7 | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.6 | 3.1 | 6.9 | 12.9 |
| 2000..... | 1.2 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.8 | 1.6 | 3.5 | 6.7 | 13.7 |
| 1999..... | 1.1 | 1.6 | 0.2 | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.3 | 3.3 | 6.8 | 12.8 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.6 | 3.4 | 7.4 | 12.5 |
| 2003..... | 1.0 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.5 | 3.2 | 7.0 | 13.2 |
| 2002..... | 1.0 | * | * | 0.1 | * | 0.3 | 0.4 | 0.7 | 1.4 | 3.1 | 6.3 | 12.9 |
| 2001..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.7 | 3.3 | 7.7 | 14.5 |
| 2000..... | 1.1 | * | * | * | * | 0.3 | 0.4 | 0.8 | 1.6 | 4.0 | 8.2 | 16.1 |
| 1999..... | 1.0 | 1.8 | * | * | * | 0.2 | 0.4 | 0.6 | 1.5 | 3.5 | 7.9 | 16.8 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 | 2.8 | 5.3 | 10.1 |
| 2003..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.3 | 2.7 | 5.8 | 10.6 |
| 2002..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.3 | 3.1 | 6.2 | 11.1 |
| 2001..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.7 | 1.5 | 3.0 | 6.4 | 12.2 |
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.6 | 3.1 | 5.7 | 12.7 |
| 1999..... | 1.2 | 1.4 | * | * | * | 0.2 | 0.4 | 0.6 | 1.2 | 3.1 | 6.1 | 11.2 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.3 | 0.7 | 1.3 | 2.9 | 5.9 | 10.6 |
| 2003..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.4 | 0.6 | 1.3 | 2.8 | 6.2 | 11.2 |
| 2002..... | 1.1 | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.2 | 3.0 | 6.1 | 11.5 |
| 2001..... | 1.2 | * | * | * | 0.1 | 0.2 | 0.3 | 0.6 | 1.4 | 2.8 | 6.8 | 12.2 |
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.3 | 0.8 | 1.4 | 3.3 | 6.5 | 13.2 |
| 1999..... | 1.1 | 1.3 | * | * | 0.1 | 0.2 | 0.3 | 0.5 | 1.2 | 3.1 | 6.5 | 12.1 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.5 | 3.3 | 7.4 | 12.3 |
| 2003..... | 1.1 | * | * | * | * | 0.2 | 0.4 | 0.6 | 1.4 | 3.1 | 7.1 | 13.1 |
| 2002..... | 1.0 | * | * | * | * | 0.3 | 0.4 | 0.6 | 1.3 | 2.9 | 6.3 | 12.4 |
| 2001..... | 1.0 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.9 | 7.8 | 13.5 |
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.4 | 0.8 | 1.4 | 3.7 | 8.2 | 15.9 |
| 1999..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 3.3 | 7.7 | 15.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.1 | 2.6 | 5.0 | 9.9 |
| 2003..... | 1.1 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.5 | 5.5 | 10.4 |
| 2002..... | 1.2 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 3.0 | 6.0 | 11.1 |
| 2001..... | 1.3 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.7 | 6.2 | 11.6 |
| 2000..... | 1.2 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.4 | 3.0 | 5.4 | 12.0 |
| 1999..... | 1.2 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 3.0 | 5.7 | 10.7 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Complications of medical and surgical care (Y40-Y84,Y88) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.2 | * | * | * | * | * | 0.8 | 1.3 | 2.6 | 5.1 | 9.5 | 14.1 |
| 2003..... | 1.2 | * | * | * | * | * | 0.6 | 1.4 | 2.8 | 4.6 | 9.0 | 15.2 |
| 2002..... | 1.2 | * | * | * | * | * | 0.9 | 1.7 | 2.7 | 4.7 | 8.5 | 14.2 |
| 2001..... | 1.4 | * | * | * | * | 0.4 | 0.9 | 1.6 | 3.4 | 6.4 | 9.3 | 23.7 |
| 2000..... | 1.4 | * | * | * | * | 0.4 | 0.6 | 1.4 | 3.6 | 6.2 | 10.6 | 21.3 |
| 1999..... | 1.3 | * | * | * | * | * | 0.6 | 1.3 | 2.6 | 5.2 | 10.6 | 20.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | * | 1.0 | 3.0 | 4.7 | 10.3 | * |
| 2003..... | 1.0 | * | * | * | * | * | * | 1.9 | 2.9 | 3.9 | 7.8 | * |
| 2002..... | 1.1 | * | * | * | * | * | * | 1.4 | 2.6 | 5.4 | 7.3 | * |
| 2001..... | 1.5 | * | * | * | * | * | 1.0 | 1.9 | 4.3 | 7.9 | 9.0 | 30.0 |
| 2000..... | 1.2 | * | * | * | * | * | * | 1.3 | 3.2 | 7.8 | 10.2 | * |
| 1999..... | 1.2 | * | * | * | * | * | * | 1.4 | 2.9 | 6.0 | 11.9 | 23.8 |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.4 | * | * | * | * | * | 1.0 | 1.4 | 2.4 | 5.4 | 9.0 | 13.7 |
| 2003..... | 1.3 | * | * | * | * | * | 0.8 | 1.0 | 2.6 | 5.0 | 9.6 | 15.3 |
| 2002..... | 1.4 | * | * | * | * | * | 1.1 | 1.9 | 2.8 | 4.2 | 9.1 | 11.7 |
| 2001..... | 1.4 | * | * | * | * | * | 0.7 | 1.4 | 2.7 | 5.3 | 9.4 | 21.4 |
| 2000..... | 1.6 | * | * | * | * | * | * | 1.5 | 3.9 | 5.2 | 10.8 | 22.2 |
| 1999..... | 1.4 | * | * | * | * | * | * | 1.2 | 2.4 | 4.7 | 9.8 | 19.1 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3) | | | | | | | | | | | | |
| Non-Hispanic 4/, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 2.7 | 5.0 | 9.5 |
| 2003..... | 0.9 | * | * | * | 0.1 | 0.1 | 0.3 | 0.5 | 1.1 | 2.5 | 5.5 | 10.0 |
| 2002..... | 0.9 | * | * | 0.1 | * | 0.1 | 0.3 | 0.6 | 1.1 | 2.7 | 5.6 | 10.7 |
| 2001..... | 1.0 | * | * | * | 0.1 | 0.1 | 0.3 | 0.6 | 1.4 | 2.7 | 6.1 | 11.5 |
| 2000..... | 1.0 | * | * | 0.1 | 0.1 | 0.2 | 0.3 | 0.7 | 1.4 | 3.1 | 5.9 | 12.4 |
| 1999..... | 0.9 | 1.3 | * | * | 0.1 | 0.1 | 0.2 | 0.5 | 1.1 | 2.8 | 6.1 | 11.9 |
| Non-Hispanic 4/, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.3 | 0.6 | 1.3 | 3.0 | 6.2 | 11.1 |
| 2003..... | 0.8 | * | * | * | * | * | 0.3 | 0.6 | 1.2 | 2.7 | 6.3 | 11.7 |
| 2002..... | 0.8 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.7 | 5.7 | 12.2 |
| 2001..... | 0.9 | * | * | * | * | 0.1 | 0.3 | 0.6 | 1.4 | 2.9 | 6.8 | 13.2 |
| 2000..... | 1.0 | * | * | * | * | 0.2 | 0.3 | 0.7 | 1.4 | 3.6 | 7.1 | 14.4 |
| 1999..... | 0.9 | 1.5 | * | * | * | 0.2 | 0.2 | 0.4 | 1.3 | 3.1 | 7.1 | 15.7 |
| Non-Hispanic 4/, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.3 | 0.5 | 0.9 | 2.4 | 4.3 | 8.8 |
| 2003..... | 0.9 | * | * | * | * | * | 0.3 | 0.5 | 1.1 | 2.3 | 5.0 | 9.3 |
| 2002..... | 1.0 | * | * | * | * | * | 0.3 | 0.6 | 1.1 | 2.6 | 5.5 | 10.0 |
| 2001..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.4 | 2.5 | 5.6 | 10.7 |
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.6 | 1.4 | 2.7 | 5.1 | 11.6 |
| 1999..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.6 | 5.4 | 10.4 |
| Non-Hispanic white, both sexes | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.0 | 2.5 | 4.9 | 9.3 |
| 2003..... | 0.9 | * | * | * | * | 0.1 | 0.3 | 0.4 | 1.0 | 2.4 | 5.4 | 9.7 |
| 2002..... | 1.0 | * | * | * | * | 0.1 | 0.3 | 0.5 | 1.0 | 2.6 | 5.5 | 10.5 |
| 2001..... | 1.0 | * | * | * | * | 0.1 | 0.2 | 0.5 | 1.2 | 2.4 | 6.0 | 10.7 |
| 2000..... | 1.0 | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 2.9 | 5.6 | 11.9 |
| 1999..... | 1.0 | 1.1 | * | * | * | 0.1 | 0.2 | 0.4 | 1.0 | 2.7 | 5.8 | 11.2 |
| Non-Hispanic white, male | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 2.9 | 6.2 | 10.9 |
| 2003..... | 0.9 | * | * | * | * | * | 0.3 | 0.4 | 1.1 | 2.7 | 6.3 | 11.7 |
| 2002..... | 0.9 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.1 | 2.5 | 5.7 | 11.7 |
| 2001..... | 0.9 | * | * | * | * | * | 0.2 | 0.5 | 1.2 | 2.6 | 6.8 | 12.3 |
| 2000..... | 1.0 | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 3.4 | 7.0 | 14.0 |
| 1999..... | 0.9 | * | * | * | * | 0.2 | 0.2 | 0.4 | 1.1 | 2.9 | 6.9 | 14.8 |
| Non-Hispanic white, female | | | | | | | | | | | | |
| 2004..... | 0.9 | * | * | * | * | * | 0.2 | 0.4 | 0.8 | 2.2 | 4.0 | 8.5 |
| 2003..... | 1.0 | * | * | * | * | * | 0.3 | 0.4 | 1.0 | 2.1 | 4.8 | 8.9 |
| 2002..... | 1.0 | * | * | * | * | * | 0.3 | 0.5 | 1.0 | 2.6 | 5.3 | 9.9 |
| 2001..... | 1.1 | * | * | * | * | 0.2 | 0.3 | 0.5 | 1.3 | 2.3 | 5.5 | 10.0 |
| 2000..... | 1.1 | * | * | * | * | 0.2 | 0.2 | 0.6 | 1.2 | 2.6 | 4.7 | 11.0 |
| 1999..... | 1.0 | * | * | * | * | * | 0.2 | 0.4 | 0.9 | 2.6 | 5.1 | 9.8 |

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-14 years | 15-24 years | 25-34 years | 35-44 years | 45-54 years | 55-64 years | 65-74 years | 75-84 years | 85 years and over |
|--|-------------|-----------------|-----------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3) | | | | | | | | | | | | |
| Non-Hispanic black, both sexes | | | | | | | | | | | | |
| 2004..... | 1.0 | * | * | * | * | * | 0.6 | 0.9 | 2.0 | 4.4 | 8.0 | 13.2 |
| 2003..... | 1.0 | * | * | * | * | * | 0.4 | 1.2 | 2.2 | 3.9 | 8.0 | 14.0 |
| 2002..... | 1.0 | * | * | * | * | * | 0.7 | 1.3 | 2.1 | 4.1 | 7.8 | 13.6 |
| 2001..... | 1.2 | * | * | * | * | * | 0.7 | 1.2 | 3.0 | 5.3 | 8.0 | 22.5 |
| 2000..... | 1.2 | * | * | * | * | * | 0.5 | 1.0 | 3.0 | 5.3 | 9.6 | 19.4 |
| 1999..... | 1.1 | * | * | * | * | * | 0.5 | 1.0 | 2.4 | 4.2 | 9.3 | 19.4 |
| Non-Hispanic black, male | | | | | | | | | | | | |
| 2004..... | 0.8 | * | * | * | * | * | * | * | 2.3 | 4.0 | 8.6 | * |
| 2003..... | 0.8 | * | * | * | * | * | * | 1.6 | 2.2 | 3.2 | 7.2 | * |
| 2002..... | 0.9 | * | * | * | * | * | * | 1.3 | 2.2 | 4.8 | 7.0 | * |
| 2001..... | 1.2 | * | * | * | * | * | * | 1.4 | 4.0 | 6.8 | 7.8 | 28.9 |
| 2000..... | 1.0 | * | * | * | * | * | * | 1.1 | 2.7 | 6.4 | 9.6 | * |
| 1999..... | 1.0 | * | * | * | * | * | * | 1.2 | 2.6 | 5.3 | 10.6 | * |
| Non-Hispanic black, female | | | | | | | | | | | | |
| 2004..... | 1.1 | * | * | * | * | * | 0.7 | 1.1 | 1.8 | 4.8 | 7.7 | 12.9 |
| 2003..... | 1.1 | * | * | * | * | * | * | 0.8 | 2.2 | 4.4 | 8.5 | 14.9 |
| 2002..... | 1.1 | * | * | * | * | * | 0.9 | 1.4 | 2.0 | 3.5 | 8.3 | 11.3 |
| 2001..... | 1.2 | * | * | * | * | * | 0.7 | 0.9 | 2.3 | 4.3 | 8.1 | 20.1 |
| 2000..... | 1.3 | * | * | * | * | * | * | 0.9 | 3.2 | 4.5 | 9.6 | 19.6 |
| 1999..... | 1.1 | * | * | * | * | * | * | * | 2.2 | 3.5 | 8.6 | 18.2 |

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes races other than white and black.

5/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.