

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All causes											
All origins 3/	877.0	731.4	34.7	17.4	21.1	69.8	93.6	97.2	118.4	163.2	235.5
Male	882.0	801.5	38.5	19.2	25.3	96.3	137.6	138.5	161.0	211.1	302.9
Female	872.2	658.1	30.8	15.5	16.7	41.7	48.0	56.4	76.6	115.8	169.3
Hispanic	331.0	611.5	32.1	14.3	20.1	67.3	97.9	98.3	107.9	140.3	192.8
Male	367.9	655.3	34.3	16.1	23.3	99.1	151.6	148.3	154.7	195.8	262.6
Female	293.7	565.9	29.8	12.5	16.7	32.9	40.1	45.6	57.7	81.5	121.4
Mexican	280.3	616.7	32.1	14.3	19.7	67.8	99.2	95.1	102.1	130.6	176.0
Male	313.8	659.2	35.4	16.4	23.0	100.0	152.2	144.0	147.4	181.8	239.8
Female	244.7	572.4	28.8	12.1	16.1	32.1	38.9	41.8	51.3	71.7	109.9
Puerto Rican	472.3	*	30.7	17.8	18.6	64.0	104.3	127.0	172.1	250.6	358.1
Male	550.0	*	*	*	*	94.2	153.4	183.1	255.8	373.8	487.2
Female	399.9	*	*	*	*	37.0	*	*	106.1	154.4	233.9
Cuban	820.4	*	*	*	*	*	*	*	*	134.7	173.8
Male	859.7	*	*	*	*	*	*	*	*	*	*
Female	783.9	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	325.9	574.5	34.0	13.2	22.4	71.7	95.0	102.1	106.1	124.7	175.2
Male	366.2	*	31.0	14.9	25.7	106.8	153.2	153.0	153.2	176.9	241.5
Female	288.8	*	37.5	11.6	19.4	33.5	40.2	51.2	56.3	75.0	110.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All causes											
Non-Hispanic 5/	944.4	746.4	35.1	18.0	21.2	69.9	92.5	96.5	119.4	165.3	238.5
Male	946.9	820.5	39.2	19.8	25.5	95.6	134.5	136.0	161.1	211.9	305.0
Female	942.0	668.9	30.8	16.1	16.7	42.8	49.1	57.8	79.1	119.6	173.5
Non-Hispanic white	996.3	572.7	29.7	15.6	19.3	64.7	78.8	82.6	103.2	144.5	208.0
Male	984.7	636.5	32.9	17.0	23.1	85.9	112.9	116.0	139.5	187.1	268.6
Female	1,007.4	505.6	26.2	14.1	15.4	42.3	43.0	49.2	67.1	101.7	147.4
Non-Hispanic black	852.1	1,623.6	61.8	29.3	30.8	97.3	163.0	177.4	222.1	311.6	466.2
Male	918.0	1,773.3	69.8	33.8	38.6	145.2	249.7	257.1	304.9	396.8	596.0
Female	792.8	1,469.1	53.7	24.8	22.8	47.8	78.4	104.5	149.5	236.7	352.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic 5/	0.4	*	*	*	*	0.0	*	*	*	*	*
Male	0.3	*	*	*	*	0.0	*	*	*	*	*
Female	0.5	*	*	0.0	0.0	0.0	*	*	0.0	*	*
Non-Hispanic white	0.5	*	*	0.0	*	0.0	*	*	*	*	*
Male	0.4	*	*	0.0	*	0.0	*	*	*	*	*
Female	0.6	*	*	0.0	0.0	0.0	*	*	0.0	*	0.0
Non-Hispanic black	0.2	*	0.0	*	0.0	0.0	*	0.0	0.0	0.0	*
Male	0.2	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.3	*	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	---------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Tuberculosis (A16-A19)

Non-Hispanic 5/	0.3	*	*	*	*	0.0	*	*	*	*	0.1
Male	0.4	0.0	*	0.0	*	0.0	0.0	0.0	*	*	*
Female	0.2	*	0.0	*	0.0	0.0	*	*	*	*	*
Non-Hispanic white	0.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	0.2	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Non-Hispanic black	0.7	*	*	*	0.0	0.0	*	*	*	*	*
Male	1.0	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	0.5	*	0.0	*	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Respiratory tuberculosis (A16)											
Non-Hispanic 5/ 6/	0.3	*	*	0.0	*	0.0	*	*	*	*	0.1
Male	0.4	0.0	*	0.0	*	0.0	0.0	0.0	*	*	*
Female	0.2	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Non-Hispanic white	0.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	0.1	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Non-Hispanic black	0.6	*	*	0.0	0.0	0.0	*	*	*	*	*
Male	0.8	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	0.4	*	0.0	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Septicemia (A40-A41)

All origins 3/	11.3	7.4	0.6	0.2	0.2	0.2	0.5	0.6	0.9	1.4	2.2
Male	10.1	9.0	0.6	0.2	*	*	0.5	0.6	1.0	1.5	2.6
Female	12.4	5.8	0.5	0.3	*	*	0.4	0.6	0.8	1.3	1.9
Hispanic	3.9	6.0	*	*	*	*	*	*	0.7	1.2	2.0
Male	3.8	7.3	*	*	*	*	*	*	*	1.5	2.8
Female	3.9	*	*	*	*	*	*	*	*	*	*
Mexican	3.1	6.6	*	*	*	*	*	*	*	*	2.2
Male	3.2	*	*	*	0.0	*	*	*	*	*	3.2
Female	2.9	*	*	*	*	*	0.0	*	*	*	*
Puerto Rican	7.2	*	*	*	0.0	0.0	*	*	*	*	*
Male	6.8	*	*	*	0.0	0.0	0.0	*	*	*	*
Female	7.5	0.0	*	0.0	0.0	0.0	*	*	*	*	*
Cuban	10.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	10.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	0.0
Female	10.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	3.6	*	*	0.0	*	*	*	*	*	*	*
Male	3.2	*	*	0.0	*	*	*	0.0	*	*	*
Female	3.9	*	*	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Septicemia (A40-A41)											
Non-Hispanic 5/	12.2	7.7	0.6	0.2	0.2	0.2	0.6	0.6	0.9	1.4	2.3
Male	10.8	9.3	0.6	0.2	*	*	0.6	0.6	1.0	1.5	2.6
Female	13.4	6.1	0.5	0.3	*	*	0.5	0.6	0.8	1.3	1.9
Non-Hispanic white	11.7	5.1	0.5	0.2	*	0.2	0.4	0.5	0.7	1.0	1.7
Male	10.3	6.5	0.5	*	*	*	0.5	0.5	0.8	1.1	1.9
Female	13.1	3.6	0.5	*	*	*	*	0.4	0.6	0.9	1.5
Non-Hispanic black	17.4	20.0	*	*	*	*	1.1	1.5	1.9	4.0	6.5
Male	16.5	22.3	*	*	*	*	*	*	2.3	4.3	8.1
Female	18.2	17.7	*	*	*	*	*	1.8	1.6	3.7	5.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Viral hepatitis (B15-B19)

All origins 3/	1.8	*	*	*	*	*	*	0.1	0.3	0.9	2.3
Male	2.3	*	*	0.0	*	*	*	*	0.5	1.2	3.5
Female	1.3	0.0	*	*	0.0	0.0	*	*	0.3	0.6	1.2
Hispanic	2.1	0.0	*	0.0	0.0	0.0	*	*	*	1.5	3.0
Male	2.6	0.0	*	0.0	0.0	0.0	*	0.0	*	2.3	4.9
Female	1.6	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Mexican	1.7	0.0	*	0.0	0.0	0.0	*	*	*	1.3	2.9
Male	2.0	0.0	*	0.0	0.0	0.0	*	0.0	*	*	4.8
Female	1.3	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Puerto Rican	5.1	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	7.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	2.7	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Cuban	3.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	3.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	3.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	1.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Viral hepatitis (B15-B19)											
Non-Hispanic 5/	1.7	*	*	*	*	*	*	*	0.3	0.8	2.2
Male	2.2	*	*	0.0	*	*	*	*	0.4	1.1	3.2
Female	1.3	0.0	*	*	0.0	0.0	*	*	0.3	0.6	1.2
Non-Hispanic white	1.6	*	*	*	*	*	*	*	0.3	0.8	2.0
Male	2.1	*	*	0.0	*	*	*	*	0.4	1.1	3.0
Female	1.1	0.0	*	*	0.0	0.0	0.0	*	*	0.5	1.1
Non-Hispanic black	2.2	*	0.0	0.0	*	0.0	*	*	*	0.9	3.6
Male	2.8	*	0.0	0.0	*	0.0	*	*	*	*	5.2
Female	1.6	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/	5.4	*	0.2	0.2	0.1	0.2	0.9	4.2	10.0	13.2	14.6
Male	8.4	*	*	0.2	*	*	0.9	5.6	14.4	19.8	22.5
Female	2.6	*	*	*	*	*	1.0	2.8	5.6	6.7	6.8
Hispanic	6.2	*	*	*	*	*	1.1	4.5	11.0	16.3	20.0
Male	9.7	*	*	*	*	*	*	6.6	16.5	25.4	30.0
Female	2.8	*	*	*	*	*	*	2.2	5.1	6.7	9.8
Mexican	3.0	0.0	*	*	*	*	*	3.3	6.7	9.2	10.1
Male	5.1	0.0	*	*	*	*	*	5.5	11.2	15.8	17.3
Female	0.8	0.0	*	*	0.0	0.0	*	*	*	*	*
Puerto Rican	29.0	*	0.0	*	*	*	*	*	45.5	73.5	98.0
Male	41.2	*	0.0	*	0.0	0.0	*	*	*	*	130.6
Female	17.6	*	0.0	*	*	*	*	*	*	*	*
Cuban	8.4	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	15.6	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Other Hispanic 4/	5.7	0.0	*	*	0.0	*	*	4.5	10.2	11.5	14.7
Male	9.5	0.0	*	0.0	0.0	*	*	*	16.6	19.2	23.9
Female	2.2	0.0	0.0	*	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Human immunodeficiency virus (HIV) disease (B20-B24)											
Non-Hispanic 5/	5.2	*	0.2	0.2	0.1	0.2	0.9	4.1	9.7	12.7	13.8
Male	8.1	*	*	*	*	*	0.8	5.3	14.0	18.9	21.3
Female	2.5	*	*	*	*	*	1.0	2.9	5.6	6.6	6.4
Non-Hispanic white	2.4	*	*	*	*	*	0.2	1.5	4.5	6.5	6.3
Male	4.2	*	*	*	*	*	*	2.3	7.5	10.9	10.8
Female	0.7	0.0	*	*	*	*	*	0.8	1.5	2.0	1.8
Non-Hispanic black	23.4	*	*	0.9	*	0.7	4.3	18.0	41.0	53.1	64.8
Male	34.3	*	*	*	*	*	3.5	22.7	55.2	74.4	98.2
Female	13.6	*	*	*	*	*	5.0	13.8	28.4	34.3	35.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malaria (B50-B54)

Non-Hispanic 5/	*	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Male	*	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Non-Hispanic white	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Non-Hispanic black	*	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Male	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins 3/	2.7	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.0	1.4
Male	2.7	6.1	0.7	0.2	0.2	0.2	0.3	0.5	0.9	1.3	1.7
Female	2.6	4.5	0.4	0.2	*	0.2	0.2	0.5	0.7	0.8	1.1
Hispanic	1.6	5.0	*	*	*	*	*	*	0.8	1.4	1.4
Male	1.6	6.0	*	*	*	*	*	*	*	2.2	1.8
Female	1.5	*	*	*	*	*	*	*	*	*	*
Mexican	1.4	4.9	*	*	*	*	*	*	*	*	*
Male	1.5	*	*	*	*	*	*	*	*	*	*
Female	1.4	*	*	*	*	*	*	*	*	*	*
Puerto Rican	2.1	*	*	0.0	*	0.0	0.0	*	*	*	*
Male	2.3	*	*	0.0	0.0	0.0	0.0	*	*	*	*
Female	1.9	*	0.0	0.0	*	0.0	0.0	*	*	*	*
Cuban	2.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.7	*	*	*	*	*	*	*	*	*	*
Male	1.8	*	*	0.0	*	*	*	*	*	*	*
Female	1.6	*	0.0	*	0.0	0.0	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)											
Non-Hispanic 5/	2.8	5.4	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0	1.4
Male	2.9	6.1	0.6	*	0.2	0.2	0.3	0.5	0.9	1.2	1.6
Female	2.7	4.6	0.4	*	*	*	*	0.5	0.6	0.8	1.1
Non-Hispanic white	2.7	4.3	0.5	0.2	*	0.2	0.2	0.3	0.5	0.7	1.0
Male	2.8	5.3	0.6	*	*	*	*	*	0.6	1.0	1.3
Female	2.6	3.2	*	*	*	*	*	0.4	0.4	0.5	0.7
Non-Hispanic black	3.8	10.8	1.1	*	*	*	*	1.3	2.3	2.7	3.7
Male	3.9	10.8	*	*	*	*	*	*	2.9	2.8	3.7
Female	3.6	10.8	*	*	*	*	*	*	1.8	2.5	3.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

All origins 3/	201.6	1.8	2.8	2.6	2.6	3.8	5.4	7.8	13.1	25.6	49.3
Male	214.5	1.8	2.9	2.7	2.8	4.4	6.3	8.3	12.3	22.0	45.3
Female	189.4	1.7	2.6	2.4	2.4	3.1	4.6	7.4	13.8	29.1	53.1
Hispanic	64.6	*	2.6	2.7	3.2	3.4	6.1	7.4	10.4	18.2	33.9
Male	67.7	*	2.4	2.7	3.4	3.8	7.4	7.4	9.0	14.9	28.0
Female	61.4	*	2.8	2.6	3.1	2.9	4.8	7.3	11.9	21.8	39.9
Mexican	52.6	*	2.4	2.8	2.9	3.1	6.3	6.8	10.3	17.4	30.0
Male	54.5	*	2.3	3.1	2.9	2.8	7.4	6.4	9.4	13.8	24.3
Female	50.6	*	2.4	2.4	2.8	3.4	4.9	7.3	11.2	21.5	36.0
Puerto Rican	88.0	*	*	*	*	*	*	*	*	24.9	43.8
Male	99.6	*	*	*	*	*	*	*	*	*	*
Female	77.2	0.0	*	*	*	*	*	*	*	*	*
Cuban	175.7	0.0	*	*	*	*	*	*	*	*	*
Male	200.4	0.0	*	*	*	*	*	*	*	*	*
Female	152.8	0.0	0.0	0.0	*	*	*	*	*	*	*
Other Hispanic 4/	68.5	*	*	*	4.7	*	5.8	9.0	11.3	16.9	38.3
Male	70.1	*	*	*	*	*	*	11.5	8.9	15.1	30.9
Female	67.1	0.0	*	*	*	*	*	*	13.8	18.5	45.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

Non-Hispanic 5/	218.9	1.7	2.8	2.5	2.5	3.8	5.3	7.9	13.4	26.5	50.7
Male	233.5	1.6	3.0	2.7	2.7	4.5	6.0	8.5	12.8	22.9	47.1
Female	204.9	1.8	2.5	2.3	2.2	3.1	4.5	7.4	14.0	29.9	54.3
Non-Hispanic white	232.8	1.6	3.0	2.4	2.3	3.6	5.0	7.8	12.8	24.9	47.5
Male	246.3	1.6	3.3	2.8	2.5	4.2	5.7	8.4	12.3	22.1	44.1
Female	219.7	*	2.6	2.1	2.1	3.0	4.3	7.3	13.3	27.7	50.8
Non-Hispanic black	185.6	*	2.3	2.8	3.2	4.7	7.0	9.2	17.2	38.2	76.4
Male	207.6	*	2.3	2.7	3.6	5.7	7.7	9.6	15.4	29.7	71.5
Female	165.7	*	2.2	3.0	2.8	3.7	6.3	8.8	18.8	45.6	80.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)											
Non-Hispanic 5/	3.0	0.0	*	*	*	*	*	*	0.2	0.4	0.9
Male	4.0	0.0	*	0.0	*	*	*	*	*	0.5	1.4
Female	2.0	0.0	*	*	*	*	*	*	*	0.3	0.4
Non-Hispanic white	3.0	0.0	*	*	*	*	*	*	0.1	0.3	0.8
Male	3.9	0.0	*	0.0	*	*	*	*	*	0.4	1.2
Female	2.1	0.0	*	*	*	*	*	*	*	0.3	0.4
Non-Hispanic black	3.4	0.0	0.0	0.0	*	*	*	*	*	0.9	1.6
Male	5.2	0.0	0.0	0.0	*	*	*	*	*	*	2.5
Female	1.8	0.0	0.0	0.0	*	*	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of stomach (C16)											
All origins 3/	4.7	0.0	0.0	0.0	*	*	*	0.3	0.5	0.8	1.4
Male	5.6	0.0	0.0	0.0	*	*	*	0.2	0.6	0.9	1.6
Female	3.8	0.0	0.0	0.0	0.0	*	*	0.3	0.4	0.8	1.2
Hispanic	3.4	0.0	0.0	0.0	0.0	*	*	*	0.9	1.3	2.4
Male	3.9	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.5
Female	2.9	0.0	0.0	0.0	0.0	*	*	*	*	1.6	2.2
Mexican	2.8	0.0	0.0	0.0	0.0	*	*	*	*	*	1.9
Male	3.4	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	2.3	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Puerto Rican	4.5	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	4.9	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	4.1	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Cuban	5.4	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*
Male	5.7	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*	0.0
Female	5.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	4.1	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	4.6	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	3.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of stomach (C16)											
Non-Hispanic 5/	4.8	0.0	0.0	0.0	*	*	*	0.2	0.4	0.8	1.3
Male	5.8	0.0	0.0	0.0	*	*	*	*	0.4	0.9	1.6
Female	3.9	0.0	0.0	0.0	0.0	*	*	0.2	0.3	0.7	1.1
Non-Hispanic white	4.4	0.0	0.0	0.0	0.0	*	*	*	0.3	0.6	0.9
Male	5.4	0.0	0.0	0.0	0.0	*	*	*	0.3	0.7	1.0
Female	3.5	0.0	0.0	0.0	0.0	0.0	*	*	*	0.5	0.7
Non-Hispanic black	6.6	0.0	0.0	0.0	*	*	*	*	0.9	1.5	3.3
Male	8.0	0.0	0.0	0.0	*	0.0	*	*	*	1.6	4.3
Female	5.3	0.0	0.0	0.0	0.0	*	*	*	*	1.4	2.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/	21.0	*	0.0	0.0	*	*	0.2	0.5	1.0	1.9	4.0
Male	21.2	0.0	0.0	0.0	*	*	0.2	0.6	1.1	2.0	4.1
Female	20.7	*	0.0	0.0	0.0	*	*	0.3	0.9	1.9	3.9
Hispanic	6.5	0.0	0.0	0.0	0.0	*	*	*	0.9	1.3	2.4
Male	6.7	0.0	0.0	0.0	0.0	*	*	*	*	*	2.2
Female	6.2	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.7
Mexican	4.8	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.0
Male	5.3	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	4.3	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Puerto Rican	10.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Male	10.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Female	10.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Cuban	22.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	22.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	21.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	6.9	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Male	6.7	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Female	7.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of colon, rectum and anus (C18-C21)											
Non-Hispanic 5/	22.8	*	0.0	0.0	*	*	0.2	0.5	1.0	2.0	4.2
Male	23.1	0.0	0.0	0.0	*	*	*	0.7	1.1	2.1	4.3
Female	22.5	*	0.0	0.0	0.0	*	*	0.3	0.9	1.9	4.1
Non-Hispanic white	24.1	0.0	0.0	0.0	*	*	0.2	0.5	0.9	1.9	3.8
Male	24.4	0.0	0.0	0.0	*	*	*	0.6	1.0	2.1	4.1
Female	23.8	0.0	0.0	0.0	0.0	*	*	*	0.9	1.7	3.6
Non-Hispanic black	20.3	*	0.0	0.0	0.0	*	*	0.8	1.4	3.3	6.8
Male	20.2	0.0	0.0	0.0	0.0	*	*	*	*	3.3	6.6
Female	20.4	*	0.0	0.0	0.0	*	*	*	*	3.3	6.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/	4.5	*	0.2	*	*	0.1	0.1	0.2	0.2	0.6	1.4
Male	5.8	*	0.3	*	*	*	*	0.2	0.3	0.8	2.1
Female	3.4	*	*	0.0	*	*	*	*	0.2	0.3	0.8
Hispanic	3.5	*	*	*	*	*	*	*	*	*	1.4
Male	4.4	*	*	*	*	*	*	*	*	*	1.9
Female	2.5	0.0	*	0.0	0.0	*	0.0	0.0	*	*	*
Mexican	3.2	*	*	*	*	*	*	*	*	*	*
Male	3.9	*	*	*	*	*	*	*	*	*	*
Female	2.5	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Puerto Rican	4.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	8.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	1.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Cuban	6.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	6.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	5.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	3.1	0.0	*	0.0	0.0	0.0	0.0	*	*	*	*
Male	3.9	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	2.3	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic 5/	4.7	*	0.2	*	*	*	0.1	0.2	0.2	0.6	1.4
Male	5.9	*	*	*	*	*	*	0.2	0.3	0.8	2.1
Female	3.5	*	*	0.0	*	*	*	*	*	0.3	0.7
Non-Hispanic white	4.5	*	0.2	*	*	*	*	*	*	0.4	1.1
Male	5.7	*	*	*	*	*	*	*	*	0.5	1.6
Female	3.4	*	*	0.0	*	*	*	*	*	0.3	0.5
Non-Hispanic black	4.5	*	*	*	*	*	*	*	*	1.1	2.3
Male	6.0	*	*	*	*	0.0	*	*	*	1.6	3.6
Female	3.2	0.0	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Malignant neoplasm of pancreas (C25)

All origins 3/	10.7	*	0.0	0.0	0.0	0.0	*	*	0.2	0.8	1.9
Male	10.6	0.0	0.0	0.0	0.0	0.0	*	*	0.3	1.0	2.4
Female	10.7	*	0.0	0.0	0.0	0.0	*	*	0.2	0.6	1.4
Hispanic	3.6	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	1.2
Male	3.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	3.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Mexican	3.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	1.5
Male	3.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	3.1	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Puerto Rican	3.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	3.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	3.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Cuban	9.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	8.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	9.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	3.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	3.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	4.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of pancreas (C25)											
Non-Hispanic 5/	11.5	*	0.0	0.0	0.0	0.0	*	*	0.2	0.8	1.9
Male	11.5	0.0	0.0	0.0	0.0	0.0	*	*	0.3	1.0	2.4
Female	11.5	*	0.0	0.0	0.0	0.0	*	*	*	0.6	1.4
Non-Hispanic white	12.2	0.0	0.0	0.0	0.0	0.0	*	*	0.2	0.7	1.8
Male	12.2	0.0	0.0	0.0	0.0	0.0	*	*	*	0.9	2.3
Female	12.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.6	1.3
Non-Hispanic black	10.2	*	0.0	0.0	0.0	0.0	0.0	*	*	1.3	2.9
Male	10.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	1.5	3.6
Female	10.1	*	0.0	0.0	0.0	0.0	0.0	*	*	*	2.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins 3/	55.8	0.0	0.0	*	*	*	*	0.2	0.7	3.0	9.2
Male	67.1	0.0	0.0	*	*	*	*	0.3	0.7	2.9	10.4
Female	45.0	0.0	0.0	*	0.0	*	*	0.2	0.6	3.0	8.1
Hispanic	11.3	0.0	0.0	0.0	0.0	0.0	*	*	*	1.2	3.0
Male	15.1	0.0	0.0	0.0	0.0	0.0	*	*	*	*	3.7
Female	7.5	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	2.3
Mexican	8.6	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.5
Male	11.1	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	5.9	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Puerto Rican	17.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	23.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	11.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	39.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	59.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	20.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	11.5	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	15.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	8.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of trachea, bronchus and lung (C33-C34)											
Non-Hispanic 5/	61.4	0.0	0.0	*	*	*	*	0.3	0.7	3.2	9.9
Male	73.9	0.0	0.0	*	*	*	*	0.3	0.8	3.1	11.1
Female	49.6	0.0	0.0	*	0.0	*	*	0.2	0.6	3.3	8.6
Non-Hispanic white	66.3	0.0	0.0	*	*	*	*	0.2	0.7	3.0	9.4
Male	78.5	0.0	0.0	*	*	*	*	*	0.8	2.9	10.3
Female	54.6	0.0	0.0	*	0.0	0.0	*	*	0.6	3.1	8.5
Non-Hispanic black	47.8	0.0	0.0	*	*	*	*	*	*	4.9	15.2
Male	64.0	0.0	0.0	*	*	*	*	*	*	4.9	19.5
Female	33.3	0.0	0.0	0.0	0.0	*	*	*	*	4.9	11.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Malignant melanoma of skin (C43)

All origins 3/	2.6	*	0.0	*	0.0	*	0.2	0.3	0.7	1.1	1.7
Male	3.4	*	0.0	0.0	0.0	*	*	0.4	0.7	1.3	2.2
Female	1.9	0.0	0.0	*	0.0	*	*	0.3	0.6	0.8	1.2
Hispanic	0.4	*	0.0	*	0.0	*	*	*	*	*	*
Male	0.5	*	0.0	0.0	0.0	*	0.0	*	*	*	*
Female	0.3	0.0	0.0	*	0.0	0.0	*	0.0	*	*	*
Mexican	0.3	0.0	0.0	*	0.0	*	0.0	*	*	*	*
Male	0.4	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Female	0.3	0.0	0.0	*	0.0	0.0	0.0	0.0	*	*	*
Puerto Rican	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.5	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0	0.0
Male	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant melanoma of skin (C43)											
Non-Hispanic 5/	2.9	0.0	0.0	0.0	0.0	*	0.2	0.4	0.8	1.2	1.9
Male	3.8	0.0	0.0	0.0	0.0	*	*	0.5	0.8	1.5	2.4
Female	2.1	0.0	0.0	0.0	0.0	*	*	0.3	0.7	0.9	1.3
Non-Hispanic white	3.5	0.0	0.0	0.0	0.0	*	0.2	0.5	0.9	1.4	2.3
Male	4.6	0.0	0.0	0.0	0.0	*	*	0.6	1.0	1.8	2.9
Female	2.6	0.0	0.0	0.0	0.0	*	*	0.4	0.8	1.1	1.6
Non-Hispanic black	0.3	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Male	0.3	0.0	0.0	0.0	0.0	*	0.0	*	*	*	0.0
Female	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of breast (C50)											
All origins 3/	15.2	0.0	*	0.0	*	*	*	0.5	1.8	4.4	7.9
Male	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	29.5	0.0	*	0.0	*	*	*	0.9	3.5	8.7	15.5
Hispanic	4.9	0.0	0.0	0.0	0.0	*	*	*	1.3	2.9	5.7
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	9.9	0.0	0.0	0.0	0.0	*	*	*	2.6	5.9	11.5
Mexican	3.9	0.0	0.0	0.0	0.0	*	*	*	1.3	2.8	5.4
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	7.9	0.0	0.0	0.0	0.0	*	*	*	2.7	6.1	11.1
Puerto Rican	6.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	12.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Cuban	12.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	24.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	5.9	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	6.5
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	11.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	12.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of breast (C50)											
Non-Hispanic 5/	16.5	0.0	*	0.0	*	*	*	0.5	1.9	4.6	8.1
Male	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	31.9	0.0	*	0.0	*	*	*	0.9	3.6	9.0	15.9
Non-Hispanic white	17.2	0.0	0.0	0.0	*	*	*	0.4	1.6	3.9	7.1
Male	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	33.5	0.0	0.0	0.0	*	*	*	0.8	3.1	7.8	14.2
Non-Hispanic black	16.3	0.0	*	0.0	0.0	0.0	*	1.0	4.0	9.4	15.4
Male	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	30.5	0.0	*	0.0	0.0	0.0	*	1.9	7.4	17.3	28.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Malignant neoplasm of cervix uteri (C53)

All origins 3/	1.5	0.0	0.0	0.0	0.0	*	*	0.3	0.8	1.3	1.8
Female	3.0	0.0	0.0	0.0	0.0	*	*	0.7	1.6	2.6	3.6
Hispanic	1.2	0.0	0.0	0.0	0.0	0.0	*	*	0.9	1.3	2.4
Female	2.4	0.0	0.0	0.0	0.0	0.0	*	*	1.9	2.7	5.0
Mexican	1.0	0.0	0.0	0.0	0.0	0.0	*	*	*	1.3	2.2
Female	2.2	0.0	0.0	0.0	0.0	0.0	*	*	*	*	4.4
Puerto Rican	1.8	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	3.5	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Other Hispanic 4/	1.4	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	2.8	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of cervix uteri (C53)											
Non-Hispanic 5/	1.6	0.0	0.0	0.0	0.0	*	*	0.4	0.8	1.3	1.8
Female	3.1	0.0	0.0	0.0	0.0	*	*	0.7	1.5	2.6	3.5
Non-Hispanic white	1.5	0.0	0.0	0.0	0.0	0.0	*	0.4	0.7	1.2	1.6
Female	2.9	0.0	0.0	0.0	0.0	0.0	*	0.8	1.5	2.3	3.2
Non-Hispanic black	2.5	0.0	0.0	0.0	0.0	*	*	*	1.1	2.2	3.1
Female	4.7	0.0	0.0	0.0	0.0	*	*	*	2.0	4.1	5.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 5/	2.6	0.0	0.0	0.0	0.0	0.0	*	*	*	0.2	0.5
Female	5.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.3	0.9
Non-Hispanic white	2.6	0.0	0.0	0.0	0.0	0.0	*	*	*	0.2	0.4
Female	5.1	0.0	0.0	0.0	0.0	0.0	*	*	*	0.3	0.8
Non-Hispanic black	3.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.8
Female	5.6	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	1.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of ovary (C56)											
All origins 3/	5.0	0.0	*	0.0	*	*	*	0.1	0.3	0.7	1.5
Female	9.8	0.0	*	0.0	*	*	*	0.3	0.5	1.4	3.0
Hispanic	1.5	0.0	*	0.0	0.0	*	*	*	*	*	*
Female	3.1	0.0	*	0.0	0.0	*	*	*	*	*	*
Mexican	1.3	0.0	*	0.0	0.0	*	*	*	0.0	*	*
Female	2.7	0.0	*	0.0	0.0	*	*	*	0.0	*	*
Puerto Rican	1.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	3.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	4.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	8.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	2.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of ovary (C56)											
Non-Hispanic 5/	5.4	0.0	0.0	0.0	*	*	*	0.1	0.3	0.8	1.6
Female	10.6	0.0	0.0	0.0	*	*	*	0.3	0.6	1.5	3.1
Non-Hispanic white	6.0	0.0	0.0	0.0	*	*	*	*	0.3	0.8	1.6
Female	11.8	0.0	0.0	0.0	*	*	*	*	0.5	1.5	3.3
Non-Hispanic black	3.2	0.0	0.0	0.0	0.0	*	*	*	*	*	1.2
Female	6.1	0.0	0.0	0.0	0.0	*	*	*	*	*	2.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of prostate (C61)											
All origins 3/	11.6	*	0.0	0.0	0.0	0.0	*	*	*	*	0.1
Male	23.8	*	0.0	0.0	0.0	0.0	*	*	*	*	0.3
Hispanic	3.4	0.0	0.0	0.0	0.0	0.0	*	*	0.0	0.0	*
Male	6.7	0.0	0.0	0.0	0.0	0.0	*	*	0.0	0.0	*
Mexican	2.5	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Male	4.8	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Puerto Rican	4.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	10.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	11.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	24.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	3.7	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*
Male	7.7	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic 5/	4.4	*	*	0.1	*	*	*	0.1	0.2	0.5	1.1
Male	5.5	*	*	*	*	*	*	*	0.2	0.6	1.5
Female	3.3	*	*	*	*	*	*	*	*	0.3	0.6
Non-Hispanic white	4.8	*	*	*	*	*	*	*	*	0.4	1.0
Male	6.0	*	*	*	*	*	*	*	*	0.5	1.5
Female	3.6	*	*	*	*	0.0	*	*	*	0.3	0.6
Non-Hispanic black	3.1	0.0	*	*	*	*	*	*	*	1.0	1.3
Male	3.9	0.0	*	*	*	0.0	*	*	*	1.5	1.9
Female	2.4	0.0	*	*	*	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of bladder (C67)											
Non-Hispanic 5/	4.8	0.0	0.0	0.0	*	0.0	*	*	*	0.2	0.3
Male	6.7	0.0	0.0	0.0	*	0.0	0.0	*	*	0.2	0.4
Female	3.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*	0.2
Non-Hispanic white	5.4	0.0	0.0	0.0	*	0.0	0.0	*	*	0.1	0.3
Male	7.6	0.0	0.0	0.0	*	0.0	0.0	*	*	*	0.4
Female	3.3	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.3
Non-Hispanic black	2.7	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	3.1	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	2.4	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/	4.7	*	0.7	0.9	0.6	0.6	0.6	1.0	1.4	2.0	2.8
Male	5.3	*	0.7	0.9	0.6	0.7	0.7	1.1	1.8	2.4	3.4
Female	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	1.0	1.5	2.2
Hispanic	1.8	*	1.0	0.9	*	*	*	0.8	*	1.2	1.6
Male	1.9	*	*	*	*	*	*	*	*	1.5	2.3
Female	1.8	0.0	*	*	*	0.0	*	*	*	*	*
Mexican	1.6	*	*	0.9	*	*	*	*	*	1.2	1.4
Male	1.6	*	*	*	*	*	*	*	*	*	*
Female	1.6	0.0	*	*	*	0.0	*	*	*	*	*
Puerto Rican	1.7	0.0	*	*	*	0.0	*	*	*	*	*
Male	1.8	0.0	*	*	0.0	0.0	*	*	0.0	*	*
Female	1.5	0.0	*	*	*	0.0	0.0	0.0	*	0.0	0.0
Cuban	3.6	0.0	0.0	*	0.0	0.0	0.0	*	0.0	0.0	*
Male	4.3	0.0	0.0	*	0.0	0.0	0.0	*	0.0	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Other Hispanic 4/	2.1	0.0	*	*	*	*	*	*	*	*	*
Male	2.1	0.0	*	*	*	*	*	*	*	*	*
Female	2.2	0.0	*	*	*	0.0	*	0.0	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 5/ 6/	5.0	*	0.7	0.9	0.6	0.7	0.6	1.0	1.5	2.1	2.9
Male	5.7	*	0.7	1.0	0.6	0.8	0.7	1.2	2.0	2.5	3.5
Female	4.4	*	0.7	0.9	0.7	0.5	0.4	0.8	1.0	1.6	2.4
Non-Hispanic white	5.7	*	0.7	0.9	0.7	0.6	0.6	1.2	1.7	2.3	3.2
Male	6.5	*	0.8	1.0	0.6	0.7	0.7	1.4	2.2	2.8	3.7
Female	5.0	*	0.7	0.8	0.7	0.5	0.4	1.0	1.2	1.8	2.7
Non-Hispanic black	2.3	*	*	1.2	*	0.8	*	*	1.0	0.9	1.9
Male	2.5	*	*	*	*	*	*	*	*	*	2.6
Female	2.0	*	*	1.3	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/	20.7	0.8	0.9	0.7	1.1	1.5	2.3	2.3	2.7	3.7	4.8
Male	22.3	*	1.0	0.8	1.2	1.8	2.8	2.7	3.2	4.4	5.9
Female	19.1	*	0.8	0.6	0.9	1.3	1.8	2.0	2.2	3.0	3.8
Hispanic	8.1	*	1.0	0.9	1.7	1.6	3.1	2.5	2.7	3.4	4.7
Male	8.4	*	*	1.1	1.6	1.6	3.6	2.6	3.0	4.3	4.7
Female	7.8	*	*	*	1.8	1.6	2.5	2.3	2.4	2.5	4.8
Mexican	6.7	*	1.0	1.0	1.6	1.5	3.3	2.1	2.5	3.0	4.0
Male	7.0	*	*	*	*	*	3.7	2.2	2.9	4.0	4.1
Female	6.5	*	*	*	*	*	2.9	*	*	*	3.8
Puerto Rican	11.0	0.0	*	*	*	*	*	*	*	*	*
Male	12.1	0.0	*	*	*	*	*	*	*	*	*
Female	10.0	0.0	0.0	*	*	*	*	*	*	*	*
Cuban	22.2	0.0	*	0.0	*	*	*	*	*	*	*
Male	23.7	0.0	*	0.0	*	*	0.0	*	*	*	*
Female	20.9	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Other Hispanic 4/	8.1	*	*	*	*	*	*	*	*	*	4.8
Male	8.2	*	*	*	*	*	*	*	*	*	*
Female	8.1	0.0	*	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)											
Non-Hispanic 5/	22.2	0.7	0.9	0.7	1.0	1.5	2.2	2.3	2.7	3.8	4.8
Male	24.1	*	0.9	0.8	1.1	1.8	2.7	2.7	3.3	4.5	6.0
Female	20.4	*	0.8	0.6	0.8	1.2	1.7	1.9	2.2	3.1	3.7
Non-Hispanic white	24.3	*	1.0	0.7	0.9	1.4	2.1	2.2	2.6	3.7	4.6
Male	26.3	*	1.0	0.8	1.0	1.7	2.7	2.5	3.0	4.3	5.6
Female	22.4	*	0.9	0.6	0.7	1.2	1.5	1.9	2.1	3.0	3.6
Non-Hispanic black	15.2	*	*	0.7	1.4	1.8	3.1	2.9	3.4	4.9	6.6
Male	16.4	0.0	*	*	1.8	2.2	3.2	3.6	4.6	5.8	8.7
Female	14.2	*	*	*	*	1.5	3.0	2.2	2.4	4.1	4.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Hodgkin's disease (C81)

All origins 3/	0.5	0.0	*	0.0	*	0.2	0.3	0.4	0.4	0.4	0.3
Male	0.6	0.0	*	0.0	*	0.2	0.4	0.5	0.5	0.5	0.4
Female	0.4	0.0	0.0	0.0	*	*	0.3	0.4	0.3	0.3	0.2
Hispanic	0.4	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	0.4	0.0	0.0	0.0	*	*	*	*	*	*	*
Female	0.3	0.0	0.0	0.0	*	0.0	*	*	*	*	*
Mexican	0.3	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	0.3	0.0	0.0	0.0	*	*	*	*	*	*	*
Female	0.3	0.0	0.0	0.0	*	0.0	*	0.0	0.0	0.0	*
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*
Cuban	*	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0	*
Male	*	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.4	0.0	0.0	0.0	0.0	*	*	0.0	*	*	0.0
Male	*	0.0	0.0	0.0	0.0	*	*	0.0	0.0	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hodgkin's disease (C81)											
Non-Hispanic 5/	0.5	0.0	*	0.0	*	0.2	0.4	0.5	0.4	0.4	0.3
Male	0.6	0.0	*	0.0	*	*	0.4	0.5	0.5	0.5	0.4
Female	0.5	0.0	0.0	0.0	*	*	0.3	0.4	0.3	0.3	0.2
Non-Hispanic white	0.6	0.0	*	0.0	*	0.2	0.4	0.5	0.4	0.5	0.4
Male	0.7	0.0	*	0.0	*	*	0.4	0.5	0.6	0.6	0.5
Female	0.5	0.0	0.0	0.0	*	*	0.3	0.4	0.3	0.4	0.2
Non-Hispanic black	0.4	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.5	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.3	0.0	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Non-Hodgkin's lymphoma (C82-C85)

All origins 3/	8.4	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4	2.1
Male	8.8	0.0	*	*	0.2	0.3	0.7	0.8	1.1	1.8	2.8
Female	7.9	*	*	*	*	0.2	0.4	0.4	0.7	1.1	1.4
Hispanic	3.1	0.0	*	*	*	*	*	*	1.0	1.0	2.0
Male	3.1	0.0	0.0	*	*	*	*	*	*	1.4	2.2
Female	3.0	0.0	*	0.0	*	*	*	*	*	*	1.8
Mexican	2.6	0.0	0.0	*	*	*	*	*	*	*	1.5
Male	2.7	0.0	0.0	*	*	*	*	*	*	*	*
Female	2.4	0.0	0.0	0.0	*	0.0	*	*	*	*	*
Puerto Rican	4.1	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Male	4.4	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Female	3.9	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Cuban	7.9	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Male	7.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	8.2	0.0	0.0	0.0	0.0	*	*	0.0	*	0.0	*
Other Hispanic 4/	3.1	0.0	*	0.0	0.0	*	*	*	*	*	*
Male	2.9	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	3.3	0.0	*	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Non-Hodgkin's lymphoma (C82-C85)											
Non-Hispanic 5/	9.0	*	*	*	0.1	0.3	0.5	0.6	0.9	1.5	2.1
Male	9.6	0.0	*	*	0.2	0.3	0.7	0.8	1.1	1.9	2.8
Female	8.5	*	*	*	*	0.2	0.3	0.4	0.7	1.1	1.4
Non-Hispanic white	10.2	0.0	*	*	*	0.2	0.5	0.6	0.8	1.4	2.0
Male	10.7	0.0	*	*	*	*	0.7	0.7	1.0	1.8	2.6
Female	9.7	0.0	*	*	*	*	*	0.4	0.6	1.0	1.4
Non-Hispanic black	4.3	*	*	*	*	*	0.7	1.0	1.3	2.3	2.8
Male	5.0	0.0	*	*	*	*	*	*	1.9	2.8	4.1
Female	3.7	*	*	*	*	*	*	*	*	1.8	1.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Leukemia (C91-C95)											
All origins 3/	7.7	0.7	0.8	0.7	0.9	1.1	1.4	1.3	1.3	1.7	1.9
Male	8.6	*	0.8	0.7	1.0	1.3	1.8	1.4	1.5	1.8	2.1
Female	6.8	*	0.7	0.6	0.8	0.9	1.1	1.2	1.2	1.5	1.7
Hispanic	3.2	*	0.9	0.8	1.5	1.4	2.3	1.9	1.4	2.0	2.1
Male	3.4	*	*	*	1.4	*	2.9	2.0	1.7	2.1	1.8
Female	3.0	*	*	*	1.6	1.6	1.8	1.8	*	1.9	2.4
Mexican	2.6	*	*	0.9	1.4	1.4	2.6	1.7	1.5	1.6	1.9
Male	2.7	*	*	*	*	*	2.9	*	*	*	*
Female	2.5	*	*	*	*	*	2.2	*	*	*	*
Puerto Rican	4.2	0.0	*	*	*	*	*	*	*	*	*
Male	5.1	0.0	*	*	*	*	*	*	*	*	*
Female	3.5	0.0	0.0	*	*	*	*	*	0.0	*	*
Cuban	9.1	0.0	*	0.0	0.0	*	0.0	*	*	*	*
Male	10.0	0.0	*	0.0	0.0	*	0.0	*	*	*	0.0
Female	8.3	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Other Hispanic 4/	3.1	*	*	*	*	*	*	*	*	*	*
Male	3.4	*	*	*	*	*	*	*	*	*	*
Female	2.8	0.0	*	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Leukemia (C91-C95)											
Non-Hispanic 5/	8.3	0.6	0.7	0.6	0.8	1.1	1.3	1.2	1.3	1.6	1.9
Male	9.3	*	0.8	0.7	0.9	1.3	1.6	1.3	1.5	1.8	2.1
Female	7.3	*	0.7	0.6	0.7	0.8	1.0	1.1	1.2	1.4	1.7
Non-Hispanic white	9.1	*	0.8	0.6	0.7	1.0	1.2	1.2	1.3	1.6	1.9
Male	10.3	*	0.8	0.7	0.8	1.2	1.6	1.3	1.4	1.8	2.1
Female	8.0	*	0.8	0.5	0.6	0.8	0.9	1.1	1.2	1.4	1.7
Non-Hispanic black	5.1	*	*	*	1.2	1.2	1.8	1.2	1.5	1.9	2.0
Male	5.7	0.0	*	*	1.5	1.6	1.7	*	1.9	1.9	2.4
Female	4.6	*	*	*	*	*	1.9	*	*	1.8	1.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 5/	4.4	0.0	0.0	0.0	0.0	0.0	0.0	*	*	0.2	0.5
Male	4.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.3	0.6
Female	4.2	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.4
Non-Hispanic white	4.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.2	0.4
Male	4.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.2	0.4
Female	4.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	0.3
Non-Hispanic black	5.4	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	1.5
Male	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	1.9
Female	5.5	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97
)

Non-Hispanic 5/	26.2	*	0.8	0.8	0.7	1.2	1.5	1.6	2.3	3.8	6.8
Male	27.2	*	1.0	0.8	0.8	1.5	1.7	2.2	3.0	4.3	7.8
Female	25.3	*	0.7	0.7	0.6	0.9	1.2	1.1	1.7	3.3	5.8
Non-Hispanic white	28.0	*	0.9	0.7	0.6	1.2	1.4	1.7	2.3	3.6	6.3
Male	28.9	*	1.1	0.8	0.7	1.4	1.6	2.4	2.9	4.2	7.3
Female	27.2	*	0.8	0.6	0.5	1.0	1.2	1.1	1.6	3.1	5.3
Non-Hispanic black	21.2	*	*	0.7	0.9	1.3	1.7	1.7	2.6	5.2	10.6
Male	22.4	0.0	*	*	*	2.0	2.1	1.7	3.4	5.4	12.2
Female	20.1	*	*	*	*	*	*	1.6	1.9	5.1	9.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/	4.9	1.5	0.4	0.3	0.2	0.3	0.3	0.3	0.6	0.6	0.9
Male	4.9	1.5	0.4	0.3	0.2	0.3	0.3	0.4	0.7	0.7	1.1
Female	4.8	1.5	0.4	0.3	*	0.3	*	0.3	0.5	0.5	0.7
Hispanic	1.6	*	*	*	*	*	*	*	*	*	1.1
Male	1.5	*	*	*	*	*	*	*	*	*	*
Female	1.7	*	*	*	*	*	*	*	*	*	*
Mexican	1.3	*	*	*	*	*	*	*	*	*	*
Male	1.3	*	*	*	*	*	*	*	*	*	*
Female	1.3	*	*	*	*	0.0	*	0.0	*	*	*
Puerto Rican	2.0	*	*	0.0	0.0	*	*	*	0.0	*	*
Male	2.3	0.0	0.0	0.0	0.0	*	0.0	*	0.0	*	*
Female	1.8	*	*	0.0	0.0	0.0	*	*	0.0	0.0	*
Cuban	4.3	0.0	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Male	4.0	0.0	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	4.6	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.7	*	0.0	*	*	*	*	0.0	*	*	*
Male	1.4	*	0.0	0.0	*	0.0	*	0.0	*	0.0	*
Female	2.1	0.0	0.0	*	0.0	*	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Anemias (D50-D64)

All origins 3/	1.7	*	0.2	0.1	0.1	0.2	0.4	0.4	0.5	0.4	0.5
Male	1.3	*	*	*	*	0.2	0.5	0.4	0.5	0.4	0.5
Female	2.0	*	*	*	*	0.2	0.4	0.3	0.5	0.4	0.4
Hispanic	0.5	0.0	*	*	*	*	*	*	*	*	*
Male	0.5	0.0	*	*	0.0	*	*	*	*	*	*
Female	0.5	0.0	*	*	*	*	*	*	*	*	*
Mexican	0.4	0.0	*	0.0	*	0.0	*	0.0	*	*	*
Male	0.4	0.0	*	0.0	0.0	0.0	*	0.0	*	*	*
Female	0.5	0.0	*	0.0	*	0.0	*	0.0	*	*	*
Puerto Rican	0.7	0.0	*	*	0.0	0.0	0.0	*	0.0	*	0.0
Male	*	0.0	*	*	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	*	0.0	0.0	*	0.0	0.0	0.0	*	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	*	0.0	0.0	0.0	0.0	*	*	*	*	*	0.0
Female	0.6	0.0	0.0	0.0	0.0	*	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Anemias (D50-D64)

Non-Hispanic 5/	1.8	*	0.2	0.1	0.2	0.2	0.5	0.4	0.5	0.4	0.5
Male	1.4	*	*	*	*	0.3	0.5	0.4	0.5	0.4	0.5
Female	2.1	*	*	*	*	*	0.4	0.4	0.5	0.4	0.5
Non-Hispanic white	1.6	*	*	*	*	*	*	*	*	*	0.1
Male	1.2	*	*	0.0	*	*	*	0.0	*	*	*
Female	2.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black	3.1	*	1.0	*	*	0.8	2.4	2.2	3.3	2.4	2.9
Male	2.9	*	*	*	*	*	2.9	2.5	3.6	2.7	3.1
Female	3.2	*	*	*	*	*	1.9	2.0	3.1	2.1	2.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Diabetes mellitus (E10-E14)

All origins 3/	25.1	*	*	*	0.1	0.3	0.6	1.0	2.0	3.2	5.4
Male	23.4	0.0	*	*	*	0.2	0.6	1.0	2.3	3.9	6.4
Female	26.7	*	*	*	*	0.3	0.6	1.0	1.7	2.6	4.5
Hispanic	16.5	0.0	0.0	0.0	*	*	*	*	1.1	2.2	4.7
Male	14.8	0.0	0.0	0.0	*	0.0	*	*	*	2.3	4.8
Female	18.3	0.0	0.0	0.0	*	*	*	*	*	2.0	4.5
Mexican	16.0	0.0	0.0	0.0	*	*	*	*	*	2.0	4.7
Male	13.7	0.0	0.0	0.0	0.0	0.0	*	*	*	*	5.1
Female	18.4	0.0	0.0	0.0	*	*	*	*	*	*	4.3
Puerto Rican	22.5	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	21.6	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	23.3	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Cuban	32.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	31.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	33.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	12.3	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	12.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*	*
Female	12.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

Non-Hispanic 5/	26.1	*	*	*	0.1	0.3	0.6	1.1	2.1	3.4	5.5
Male	24.4	0.0	*	*	*	0.3	0.6	1.1	2.5	4.1	6.5
Female	27.7	*	*	*	*	0.3	0.6	1.1	1.7	2.6	4.5
Non-Hispanic white	25.2	*	*	*	*	0.3	0.5	1.0	1.8	2.9	4.7
Male	24.2	0.0	*	*	*	0.3	0.5	1.0	2.1	3.6	5.6
Female	26.1	*	*	*	*	*	0.5	1.0	1.5	2.1	3.8
Non-Hispanic black	35.7	0.0	*	*	*	*	1.2	1.9	4.4	7.0	11.8
Male	30.1	0.0	*	0.0	*	*	*	1.8	5.4	8.3	13.8
Female	40.8	0.0	*	*	*	*	*	2.1	3.6	5.8	10.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Nutritional deficiencies (E40-E64)

Non-Hispanic 5/	1.7	*	*	*	0.0	*	*	*	*	0.1	0.2
Male	1.2	*	*	*	0.0	*	*	*	*	*	0.2
Female	2.2	*	0.0	0.0	0.0	0.0	*	*	*	*	0.2
Non-Hispanic white	1.8	*	*	0.0	0.0	*	*	*	*	*	0.2
Male	1.2	*	*	0.0	0.0	*	*	0.0	*	*	*
Female	2.3	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Non-Hispanic black	1.8	*	*	*	0.0	0.0	*	*	*	*	*
Male	1.6	*	*	*	0.0	0.0	*	*	0.0	*	*
Female	1.9	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malnutrition (E40-E46)											
Non-Hispanic 5/	1.6	*	*	*	0.0	*	*	*	*	0.1	0.2
Male	1.1	*	*	*	0.0	*	*	*	*	*	*
Female	2.0	*	0.0	0.0	0.0	0.0	*	*	*	*	0.2
Non-Hispanic white	1.6	*	*	0.0	0.0	*	*	*	*	*	0.2
Male	1.1	*	*	0.0	0.0	*	*	0.0	*	*	*
Female	2.2	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Non-Hispanic black	1.6	*	*	*	0.0	0.0	*	*	*	*	*
Male	1.5	*	*	*	0.0	0.0	*	*	0.0	*	*
Female	1.8	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Meningitis (G00,G03)

All origins 3/	0.3	3.0	0.3	*	*	0.1	0.1	0.1	0.1	0.1	0.2
Male	0.3	3.1	0.3	*	*	*	*	*	*	*	0.2
Female	0.3	2.9	0.3	*	*	*	*	*	*	*	*
Hispanic	0.2	*	*	*	0.0	*	*	*	*	*	*
Male	0.3	*	*	*	0.0	0.0	*	*	*	*	*
Female	0.2	*	*	0.0	0.0	*	*	*	*	*	0.0
Mexican	0.2	*	*	0.0	0.0	*	*	*	*	0.0	*
Male	0.3	*	*	0.0	0.0	0.0	*	*	*	0.0	*
Female	0.2	*	*	0.0	0.0	*	*	*	*	0.0	0.0
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	*	0.0	*	0.0	*	0.0	0.0	0.0	*	0.0
Male	*	*	0.0	*	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	*	*	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

All origins 3/	348.5	16.9	1.5	0.7	1.1	2.8	4.3	7.0	13.2	26.3	49.8
Male	331.5	17.8	1.6	0.8	1.1	3.4	4.9	8.5	17.1	34.8	68.6
Female	364.7	15.9	1.4	0.6	1.0	2.1	3.5	5.5	9.3	17.8	31.3
Hispanic	107.4	15.9	1.5	0.7	0.9	2.4	3.5	5.8	9.7	16.5	29.1
Male	109.8	13.3	1.5	*	*	2.8	4.4	8.1	13.5	22.3	41.9
Female	105.0	18.7	1.5	*	*	1.9	2.4	3.4	5.8	10.4	16.1
Mexican	86.3	15.8	1.6	*	*	2.2	3.3	6.1	9.9	16.7	28.9
Male	88.8	12.3	1.9	*	*	2.6	4.2	8.8	13.0	22.2	40.8
Female	83.6	19.5	*	*	*	*	2.2	3.0	6.3	10.5	16.4
Puerto Rican	153.2	*	*	*	*	*	*	*	15.9	23.2	43.3
Male	164.6	*	*	*	*	*	*	*	*	*	*
Female	142.7	*	*	0.0	*	*	*	*	*	*	*
Cuban	353.0	*	0.0	0.0	*	*	*	0.0	*	*	*
Male	348.8	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Female	357.0	*	0.0	0.0	*	*	0.0	0.0	*	*	*
Other Hispanic 4/	102.6	*	*	*	*	*	*	4.7	7.7	13.8	24.9
Male	106.3	*	0.0	*	*	*	*	*	12.4	20.1	37.8
Female	99.3	*	*	0.0	*	*	*	*	*	7.8	12.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 5/	378.5	17.0	1.6	0.7	1.1	2.8	4.4	7.1	13.7	27.4	51.8
Male	359.8	18.8	1.7	0.8	1.1	3.5	5.0	8.5	17.7	36.2	71.1
Female	396.2	15.2	1.4	0.6	1.0	2.1	3.7	5.8	9.8	18.7	32.8
Non-Hispanic white	405.0	13.7	1.3	0.7	0.9	2.4	3.5	5.6	11.1	22.8	43.2
Male	381.7	15.6	1.4	0.8	0.9	2.9	3.8	6.7	14.7	30.9	61.9
Female	427.3	11.7	1.1	0.6	0.9	1.9	3.1	4.5	7.5	14.6	24.6
Non-Hispanic black	313.2	32.5	2.9	1.2	2.0	5.1	9.4	15.8	29.4	58.8	114.4
Male	306.1	33.9	3.0	1.3	2.3	6.7	11.2	19.3	36.6	74.2	144.2
Female	319.5	31.1	2.7	*	1.7	3.4	7.7	12.5	23.1	45.2	88.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/	265.9	13.7	1.2	0.6	0.8	2.3	3.4	5.4	10.5	20.9	40.0
Male	263.8	13.8	1.3	0.6	0.9	2.9	4.0	6.8	14.0	29.0	57.9
Female	268.0	13.6	1.1	0.5	0.7	1.7	2.7	4.1	7.1	12.8	22.3
Hispanic	82.5	13.3	1.2	*	*	1.9	2.4	3.6	7.3	12.4	20.4
Male	86.0	10.9	1.4	*	*	2.1	3.0	5.8	10.3	17.7	30.8
Female	79.0	15.8	*	*	*	1.6	1.7	*	4.0	6.8	9.7
Mexican	64.5	13.5	1.3	*	*	1.8	2.2	3.8	7.1	12.4	19.4
Male	68.2	10.3	*	*	*	2.1	2.8	6.4	9.7	17.3	29.1
Female	60.6	16.7	*	*	*	*	*	*	4.2	6.8	9.3
Puerto Rican	124.5	*	*	0.0	*	*	*	*	12.9	18.2	33.5
Male	134.3	*	*	0.0	*	0.0	*	*	*	*	*
Female	115.4	*	*	0.0	*	*	*	*	*	*	*
Cuban	290.8	*	0.0	0.0	*	*	*	0.0	*	*	*
Male	287.5	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Female	293.7	*	0.0	0.0	*	*	0.0	0.0	*	*	*
Other Hispanic 4/	77.4	*	*	*	*	*	*	*	6.1	10.1	17.8
Male	82.4	*	0.0	*	*	*	*	*	10.2	15.7	28.2
Female	72.7	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)											
Non-Hispanic 5/	288.7	13.8	1.2	0.6	0.8	2.4	3.5	5.7	11.0	21.8	41.8
Male	286.4	14.5	1.3	0.6	0.9	3.0	4.2	7.0	14.6	30.3	60.5
Female	290.8	13.0	1.1	0.5	0.8	1.7	2.9	4.5	7.5	13.5	23.6
Non-Hispanic white	310.0	10.9	1.0	0.5	0.7	2.1	2.7	4.5	9.0	18.6	36.0
Male	305.5	12.0	1.1	0.6	0.7	2.5	3.1	5.4	12.2	26.4	54.0
Female	314.2	9.9	0.8	0.5	0.7	1.6	2.2	3.5	5.8	10.7	18.0
Non-Hispanic black	234.8	26.7	2.4	1.1	1.5	4.4	8.1	12.8	23.6	45.3	86.3
Male	236.6	26.4	2.6	*	1.8	6.1	9.7	16.2	30.1	60.1	114.0
Female	233.3	26.9	2.2	*	*	2.7	6.7	9.8	17.9	32.4	62.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All origins 3/	1.3	*	*	*	*	*	*	*	0.2	0.2	0.3
Male	0.8	*	*	*	*	*	*	*	*	0.2	0.3
Female	1.9	*	0.0	*	*	*	*	*	*	0.2	0.4
Hispanic	0.5	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	0.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Female	0.8	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Mexican	0.5	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Male	0.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Female	0.7	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	0.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Female	1.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic 5/	1.4	*	*	*	*	*	*	*	0.2	0.2	0.3
Male	0.9	*	*	*	*	*	*	*	*	*	0.3
Female	2.0	*	0.0	*	*	*	*	*	*	0.2	0.4
Non-Hispanic white	1.6	*	*	*	*	*	*	*	*	0.1	0.3
Male	0.9	*	*	0.0	0.0	*	*	*	*	*	0.2
Female	2.2	*	0.0	*	*	*	*	*	*	*	0.3
Non-Hispanic black	0.8	*	0.0	*	*	*	*	*	*	*	0.7
Male	0.5	*	0.0	*	*	*	*	*	*	*	*
Female	1.1	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hypertensive heart disease (I11)											
Non-Hispanic 5/	9.0	*	0.0	0.0	0.0	*	0.2	0.3	0.9	1.6	2.9
Male	7.5	0.0	0.0	0.0	0.0	*	*	0.5	1.2	2.1	4.0
Female	10.4	*	0.0	0.0	0.0	0.0	*	*	0.6	1.1	1.8
Non-Hispanic white	8.0	*	0.0	0.0	0.0	*	*	0.2	0.5	0.9	1.8
Male	6.3	0.0	0.0	0.0	0.0	*	*	*	0.9	1.2	2.7
Female	9.6	*	0.0	0.0	0.0	0.0	*	*	*	0.5	0.8
Non-Hispanic black	16.8	0.0	0.0	0.0	0.0	*	*	1.1	2.9	6.2	10.6
Male	16.4	0.0	0.0	0.0	0.0	*	*	1.6	3.1	7.9	13.6
Female	17.1	0.0	0.0	0.0	0.0	0.0	*	*	2.7	4.7	8.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hypertensive heart and renal disease (I13)											
Non-Hispanic 5/	1.3	0.0	0.0	*	0.0	*	*	*	*	0.2	0.3
Male	1.1	0.0	0.0	*	0.0	*	*	*	*	0.2	0.4
Female	1.5	0.0	0.0	*	0.0	*	*	*	*	*	0.3
Non-Hispanic white	1.1	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
Male	0.9	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
Female	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Non-Hispanic black	3.0	0.0	0.0	*	0.0	*	*	*	*	1.0	2.1
Male	2.7	0.0	0.0	0.0	0.0	*	*	*	*	*	2.5
Female	3.2	0.0	0.0	*	0.0	*	*	*	*	*	1.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Ischemic heart diseases (I20-I25)											
All origins 3/	194.2	0.7	*	*	*	0.3	0.5	1.3	3.9	10.6	24.1
Male	200.5	*	*	*	*	0.4	0.6	1.9	5.8	16.1	37.8
Female	188.2	*	*	*	*	*	0.3	0.8	2.0	5.2	10.6
Hispanic	63.6	*	0.0	*	0.0	*	*	0.9	2.4	5.0	11.0
Male	67.3	0.0	0.0	*	0.0	*	*	1.7	3.6	8.4	18.1
Female	59.9	*	0.0	0.0	0.0	*	*	*	*	*	3.7
Mexican	49.2	*	0.0	*	0.0	*	*	1.1	2.2	5.1	10.4
Male	53.0	0.0	0.0	*	0.0	*	*	2.0	3.1	8.5	16.5
Female	45.0	*	0.0	0.0	0.0	*	*	*	*	*	4.0
Puerto Rican	95.7	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	103.8	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	88.2	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Cuban	238.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	233.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	243.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	58.4	*	0.0	0.0	0.0	*	*	*	*	3.7	11.1
Male	64.3	0.0	0.0	0.0	0.0	*	*	*	*	*	18.5
Female	53.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Ischemic heart diseases (I20-I25)											
Non-Hispanic 5/	210.4	0.8	*	*	*	0.3	0.5	1.4	4.1	11.2	25.4
Male	217.4	*	*	*	*	0.4	0.7	1.9	6.2	17.0	39.7
Female	203.6	*	*	*	*	*	0.3	0.9	2.1	5.6	11.3
Non-Hispanic white	228.5	*	*	*	*	0.2	0.4	1.3	3.8	10.7	23.6
Male	235.4	*	*	*	*	0.3	0.6	1.6	5.7	16.2	37.7
Female	221.9	*	*	*	0.0	*	*	0.9	1.9	5.1	9.4
Non-Hispanic black	155.2	*	*	*	*	*	1.3	2.1	6.3	17.0	41.5
Male	157.8	*	0.0	*	*	*	*	3.2	9.3	24.6	59.4
Female	152.9	*	*	*	*	*	*	*	3.7	10.2	25.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute myocardial infarction (I21-I22)											
Non-Hispanic 5/ 6/	79.4	*	*	*	*	0.1	0.2	0.6	1.8	4.4	11.0
Male	85.1	*	*	*	*	*	0.3	0.7	2.6	6.7	16.8
Female	74.1	*	*	*	*	*	*	0.4	1.0	2.2	5.2
Non-Hispanic white	86.5	*	*	*	*	*	*	0.5	1.7	4.4	10.5
Male	92.9	*	*	*	*	*	*	0.7	2.4	6.7	16.5
Female	80.3	*	*	*	0.0	*	*	0.4	1.0	2.0	4.6
Non-Hispanic black	57.8	*	0.0	*	*	*	*	0.9	2.6	5.8	16.1
Male	58.1	*	0.0	*	*	*	*	*	3.8	8.1	22.1
Female	57.5	*	0.0	0.0	*	*	*	*	1.6	3.8	10.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other acute ischemic heart diseases (I24)											
Non-Hispanic 5/	1.3	*	*	0.0	0.0	0.0	*	*	*	0.1	0.3
Male	1.4	*	*	0.0	0.0	0.0	*	*	*	0.2	0.4
Female	1.2	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Non-Hispanic white	1.4	*	*	0.0	0.0	0.0	*	*	*	*	0.2
Male	1.4	*	*	0.0	0.0	0.0	*	*	*	*	0.4
Female	1.3	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Non-Hispanic black	1.3	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	1.1	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/	119.9	*	*	*	*	0.1	0.3	0.8	2.2	6.3	13.4
Male	121.1	*	*	*	*	0.2	0.3	1.1	3.4	9.6	21.4
Female	118.7	*	*	*	0.0	*	*	0.5	1.0	3.1	5.6
Hispanic	40.6	0.0	0.0	*	0.0	*	*	*	1.5	3.2	6.5
Male	42.8	0.0	0.0	*	0.0	*	*	*	2.3	5.6	10.9
Female	38.4	0.0	0.0	0.0	0.0	*	*	*	*	*	2.1
Mexican	30.0	0.0	0.0	*	0.0	*	*	*	1.3	3.0	5.8
Male	32.3	0.0	0.0	*	0.0	0.0	0.0	*	*	5.2	9.7
Female	27.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Puerto Rican	64.4	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	69.5	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	59.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Cuban	154.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	151.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	156.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	39.7	0.0	0.0	0.0	0.0	*	*	0.0	*	*	7.3
Male	43.4	0.0	0.0	0.0	0.0	*	*	0.0	*	*	12.3
Female	36.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 5/	129.6	*	*	*	*	0.1	0.3	0.8	2.3	6.7	14.1
Male	130.9	*	*	*	*	*	0.4	1.1	3.5	10.1	22.4
Female	128.3	*	*	*	0.0	*	*	0.5	1.1	3.4	6.0
Non-Hispanic white	140.6	*	*	*	*	*	0.2	0.7	2.1	6.2	12.8
Male	141.1	*	*	*	*	*	*	1.0	3.2	9.3	20.8
Female	140.2	*	0.0	0.0	0.0	*	*	0.5	0.9	3.1	4.8
Non-Hispanic black	96.2	*	*	*	*	*	0.7	1.3	3.7	10.7	24.9
Male	98.1	*	0.0	*	*	*	*	1.9	5.5	15.9	36.4
Female	94.4	*	*	*	0.0	*	*	*	2.1	6.2	14.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerotic cardiovascular disease, so described (I25.0)											
Non-Hispanic 5/	28.3	0.0	*	0.0	*	0.0	*	0.2	0.7	2.1	4.9
Male	29.3	0.0	0.0	0.0	*	0.0	*	*	1.1	3.3	7.9
Female	27.4	0.0	*	0.0	0.0	0.0	*	*	0.3	1.0	1.9
Non-Hispanic white	28.8	0.0	0.0	0.0	*	0.0	*	*	0.6	1.9	4.3
Male	29.1	0.0	0.0	0.0	*	0.0	*	*	0.9	3.0	7.2
Female	28.4	0.0	0.0	0.0	0.0	0.0	0.0	*	*	0.8	1.4
Non-Hispanic black	32.2	0.0	*	0.0	0.0	0.0	*	*	1.3	4.0	9.5
Male	36.3	0.0	0.0	0.0	0.0	0.0	*	*	1.9	6.2	14.2
Female	28.4	0.0	*	0.0	0.0	0.0	*	*	*	2.1	5.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/	93.5	*	*	*	*	0.1	0.2	0.6	1.5	4.3	8.8
Male	93.6	*	*	*	*	0.2	0.2	0.9	2.4	6.4	13.8
Female	93.4	*	0.0	*	0.0	*	*	0.3	0.7	2.2	3.9
Hispanic	30.8	0.0	0.0	*	0.0	*	*	*	1.2	2.3	4.2
Male	31.1	0.0	0.0	*	0.0	*	*	*	1.7	3.9	7.1
Female	30.6	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Mexican	22.3	0.0	0.0	*	0.0	*	*	*	*	2.2	3.6
Male	23.0	0.0	0.0	*	0.0	0.0	0.0	*	*	3.7	5.9
Female	21.5	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Puerto Rican	48.8	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	49.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	47.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Cuban	137.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	132.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	141.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	27.8	0.0	0.0	0.0	0.0	*	*	0.0	*	*	4.8
Male	28.9	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Female	26.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 5/ Male	101.3	*	*	*	*	0.1	0.2	0.6	1.6	4.6	9.3
Female	101.6	*	*	*	*	*	0.2	0.9	2.5	6.7	14.5
Non-Hispanic white Male	100.9	*	0.0	*	0.0	*	*	0.4	0.7	2.4	4.1
Female	111.9	*	*	*	*	*	0.2	0.6	1.5	4.3	8.5
Non-Hispanic black Male	112.0	*	*	*	*	*	*	0.8	2.3	6.3	13.6
Female	111.8	*	0.0	0.0	0.0	*	*	0.4	0.6	2.2	3.4
Non-Hispanic black Male	64.0	*	0.0	*	*	*	*	0.9	2.4	6.7	15.4
Female	61.8	*	0.0	*	*	*	*	*	3.6	9.6	22.1
	65.9	*	0.0	*	0.0	*	*	*	*	4.2	9.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other heart diseases (I26-I51)											
All origins 3/	60.8	12.8	1.2	0.5	0.8	2.0	2.7	3.7	5.6	8.3	12.4
Male	54.5	13.1	1.3	0.5	0.8	2.5	3.1	4.4	6.8	10.5	15.6
Female	66.9	12.6	1.0	0.5	0.7	1.5	2.3	3.1	4.3	6.2	9.4
Hispanic	15.0	12.8	1.2	*	*	1.6	2.0	2.5	4.2	5.7	7.5
Male	15.2	10.9	1.4	*	*	1.7	2.5	3.7	5.7	7.2	10.3
Female	14.7	14.7	*	*	*	1.5	1.5	*	2.6	4.1	4.6
Mexican	12.3	12.9	1.3	*	*	1.6	1.9	2.5	4.3	5.5	7.2
Male	12.5	10.3	*	*	*	*	2.3	3.9	5.7	6.4	10.0
Female	12.0	15.5	*	*	*	*	*	*	2.9	4.5	4.3
Puerto Rican	22.0	*	*	0.0	*	*	*	*	*	*	*
Male	24.0	*	*	0.0	*	0.0	*	*	*	*	*
Female	20.1	*	*	0.0	*	*	*	*	*	*	*
Cuban	40.9	*	0.0	0.0	*	*	*	0.0	*	*	*
Male	42.5	*	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*
Female	39.4	*	0.0	0.0	*	*	0.0	0.0	*	*	*
Other Hispanic 4/	14.9	*	*	*	*	*	*	*	*	5.2	6.1
Male	14.5	*	0.0	*	*	*	*	*	*	*	*
Female	15.2	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other heart diseases (I26-I51)											
Non-Hispanic 5/	66.6	12.8	1.2	0.5	0.8	2.1	2.8	3.9	5.7	8.6	12.9
Male	59.6	13.6	1.3	0.5	0.8	2.6	3.2	4.5	7.0	10.9	16.1
Female	73.3	12.0	1.0	0.5	0.7	1.5	2.4	3.3	4.6	6.4	9.8
Non-Hispanic white	70.8	10.1	0.9	0.4	0.7	1.8	2.2	3.0	4.5	6.8	10.4
Male	62.1	11.2	1.0	0.5	0.6	2.1	2.4	3.4	5.5	8.7	13.4
Female	79.3	9.0	0.8	0.4	0.7	1.4	1.9	2.5	3.6	4.9	7.4
Non-Hispanic black	59.0	25.0	2.3	0.9	1.4	3.8	6.0	9.4	13.6	20.8	31.4
Male	59.1	24.6	2.6	*	1.6	5.2	7.1	11.2	16.9	26.0	37.8
Female	59.0	25.4	2.1	*	*	2.3	4.9	7.7	10.8	16.2	25.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All origins 3/	0.3	0.7	0.2	*	*	*	0.1	0.1	0.2	0.2	0.3
Male	0.3	*	*	*	*	*	*	*	0.2	0.3	0.3
Female	0.3	*	*	*	*	*	*	*	*	0.2	0.2
Hispanic	0.2	*	*	*	*	*	*	*	*	*	*
Male	0.2	*	*	*	*	*	*	*	*	*	*
Female	0.2	*	*	0.0	0.0	0.0	*	0.0	*	*	0.0
Mexican	0.2	*	*	*	*	*	*	*	*	*	*
Male	0.2	0.0	*	*	*	*	*	*	*	*	*
Female	*	*	0.0	0.0	0.0	0.0	*	0.0	*	0.0	0.0
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	*	*	0.0	0.0	*	*	0.0	0.0	*	0.0
Male	*	*	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0
Female	*	0.0	*	0.0	0.0	0.0	*	0.0	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/	39.9	11.8	0.9	0.4	0.7	1.9	2.4	3.4	5.1	7.5	11.1
Male	38.1	12.0	1.0	0.5	0.8	2.3	2.8	4.0	6.3	9.4	13.9
Female	41.7	11.5	0.9	0.4	0.6	1.5	2.1	2.8	4.0	5.5	8.3
Hispanic	10.6	12.1	0.9	*	*	1.4	1.6	2.3	3.8	5.0	6.6
Male	11.4	10.3	*	*	*	1.5	2.2	3.3	5.1	6.2	9.1
Female	9.8	13.9	*	*	*	1.4	*	*	2.3	3.7	4.0
Mexican	8.6	12.1	1.0	*	*	1.5	1.4	2.3	3.9	4.7	6.6
Male	9.3	10.0	*	*	*	*	1.9	3.5	5.2	5.4	9.2
Female	7.8	14.3	*	*	*	*	*	*	*	3.9	4.0
Puerto Rican	15.5	*	*	0.0	*	*	*	*	*	*	*
Male	18.0	*	*	0.0	*	0.0	*	*	*	*	*
Female	13.2	*	*	0.0	*	*	*	*	*	*	*
Cuban	33.2	*	0.0	0.0	*	*	*	0.0	*	*	*
Male	36.3	*	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*
Female	30.2	*	0.0	0.0	*	*	0.0	0.0	*	*	*
Other Hispanic 4/	10.0	*	*	*	*	*	*	*	*	4.4	4.6
Male	10.4	*	0.0	*	*	*	*	*	*	*	*
Female	9.6	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 5/	43.6	11.7	0.9	0.4	0.7	1.9	2.6	3.6	5.3	7.7	11.5
Male	41.5	12.4	1.0	0.5	0.8	2.4	2.9	4.1	6.4	9.8	14.3
Female	45.6	10.9	0.9	0.4	0.6	1.5	2.3	3.1	4.2	5.7	8.7
Non-Hispanic white	45.6	9.5	0.8	0.4	0.6	1.7	2.0	2.7	4.2	6.1	9.3
Male	42.6	10.5	0.9	0.4	0.6	2.0	2.2	3.2	5.1	7.9	12.0
Female	48.5	8.4	0.6	0.4	0.6	1.4	1.8	2.3	3.3	4.3	6.7
Non-Hispanic black	42.9	22.5	1.7	0.7	1.2	3.5	5.5	8.5	12.5	18.6	27.5
Male	45.1	22.3	*	*	1.6	4.9	6.5	10.1	15.4	23.3	33.3
Female	41.0	22.7	1.9	*	*	2.1	4.6	7.1	10.0	14.5	22.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All origins 3/	6.2	*	*	0.0	*	*	*	0.1	0.3	0.5	0.9
Male	4.9	*	*	0.0	*	*	*	*	0.4	0.6	1.1
Female	7.5	*	*	0.0	0.0	*	*	*	0.2	0.3	0.7
Hispanic	2.4	*	*	0.0	0.0	*	*	*	*	*	*
Male	2.2	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	2.7	*	*	0.0	0.0	0.0	*	*	*	0.0	*
Mexican	2.0	*	*	0.0	0.0	*	*	*	*	*	*
Male	1.8	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Female	2.3	*	*	0.0	0.0	0.0	*	0.0	*	0.0	*
Puerto Rican	3.6	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*	*
Male	3.6	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*	*
Female	3.7	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Cuban	7.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	7.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	6.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	2.1	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Male	1.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	2.4	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Essential (primary) hypertension and hypertensive renal disease (I10,I12)											
Non-Hispanic 5/	6.7	*	*	0.0	*	*	*	0.1	0.3	0.5	1.0
Male	5.2	*	*	0.0	*	*	*	*	0.4	0.7	1.1
Female	8.1	0.0	*	0.0	0.0	*	*	*	*	0.4	0.8
Non-Hispanic white	6.1	*	0.0	0.0	0.0	*	*	*	0.1	0.2	0.5
Male	4.6	*	0.0	0.0	0.0	*	*	*	*	0.3	0.6
Female	7.6	0.0	0.0	0.0	0.0	*	*	*	*	*	0.3
Non-Hispanic black	11.2	*	*	0.0	*	0.0	*	*	1.0	2.4	4.2
Male	9.9	*	*	0.0	*	0.0	*	*	*	2.9	4.9
Female	12.4	0.0	0.0	0.0	0.0	0.0	*	*	*	1.9	3.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Cerebrovascular diseases (I60-I69)											
All origins 3/	61.4	2.7	0.3	0.1	0.2	0.3	0.6	1.1	1.9	4.1	7.4
Male	48.4	3.3	0.3	*	*	0.4	0.6	1.1	2.0	4.0	7.8
Female	73.8	2.0	0.3	*	0.2	0.3	0.6	1.1	1.8	4.1	7.1
Hispanic	18.8	*	*	*	*	*	0.8	1.8	2.1	3.4	7.0
Male	17.8	*	*	*	*	*	*	1.9	2.7	3.7	8.9
Female	19.9	*	*	*	*	*	*	1.7	1.5	3.2	5.1
Mexican	16.7	*	*	*	*	*	*	1.9	2.3	3.7	7.9
Male	15.8	*	*	*	*	*	*	2.1	2.9	4.2	9.7
Female	17.6	*	*	*	*	*	*	*	*	3.1	6.1
Puerto Rican	21.6	*	*	*	0.0	*	*	*	*	*	*
Male	22.8	*	0.0	*	0.0	*	0.0	*	*	*	*
Female	20.4	0.0	*	0.0	0.0	*	*	*	*	*	*
Cuban	44.6	*	0.0	0.0	*	0.0	0.0	0.0	*	*	*
Male	41.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	47.7	*	0.0	0.0	*	0.0	0.0	0.0	*	0.0	*
Other Hispanic 4/	19.1	*	0.0	*	0.0	*	*	*	*	*	4.6
Male	17.6	*	0.0	*	0.0	*	*	*	*	*	*
Female	20.5	*	0.0	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Cerebrovascular diseases (I60-I69)

Non-Hispanic 5/ Male	66.7	2.8	0.3	0.1	0.2	0.3	0.6	1.0	1.9	4.1	7.4
Female	52.3	3.6	0.3	*	*	0.3	0.6	1.0	1.9	4.0	7.6
Non-Hispanic white Male	80.4	2.0	*	*	0.2	0.3	0.6	1.0	1.9	4.2	7.3
Female	70.8	2.3	0.3	0.1	0.2	0.3	0.6	0.8	1.5	3.2	5.4
Non-Hispanic black Male	54.2	3.0	*	*	*	0.3	0.5	0.8	1.5	3.1	5.6
Female	86.7	1.6	*	*	*	*	0.7	0.9	1.4	3.2	5.2
Non-Hispanic black Male	56.6	5.1	*	*	*	*	0.9	2.0	4.0	9.7	20.7
Female	49.9	*	*	*	*	*	*	2.0	4.1	9.6	21.6
	62.6	*	*	*	*	*	*	1.9	3.9	9.9	19.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerosis (I70)											
Non-Hispanic 5/	6.0	*	0.0	0.0	0.0	*	*	*	*	*	0.1
Male	4.5	*	0.0	0.0	0.0	0.0	0.0	*	*	*	0.2
Female	7.5	*	0.0	0.0	0.0	*	*	0.0	*	*	*
Non-Hispanic white	6.8	*	0.0	0.0	0.0	*	*	*	*	*	*
Male	4.9	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	8.6	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Non-Hispanic black	3.2	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	2.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	3.6	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other diseases of circulatory system (I71-I78)

All origins 3/	9.5	*	*	*	*	*	0.2	0.3	0.5	0.8	1.3
Male	10.3	*	0.0	*	*	*	0.2	0.4	0.7	1.0	1.6
Female	8.6	*	*	*	*	*	*	*	0.2	0.5	1.1
Hispanic	2.3	*	0.0	0.0	*	*	*	*	*	*	0.9
Male	2.6	*	0.0	0.0	0.0	*	*	*	*	*	*
Female	2.0	*	0.0	0.0	*	*	0.0	*	*	*	*
Mexican	1.8	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	1.9	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	1.7	0.0	0.0	0.0	*	*	0.0	*	*	*	*
Puerto Rican	2.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	3.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Female	2.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	7.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	9.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	5.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	2.8	*	0.0	0.0	0.0	*	*	*	*	*	*
Male	3.6	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	2.1	*	0.0	0.0	0.0	*	0.0	*	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic 5/ Male	10.4	*	*	*	*	*	0.2	0.3	0.5	0.8	1.4
Female	11.4	*	0.0	*	*	*	*	0.4	0.8	1.1	1.7
Non-Hispanic white Male	9.4	*	*	*	*	*	*	*	0.2	0.6	1.1
Female	11.3	*	*	*	*	*	0.2	0.3	0.5	0.8	1.2
Non-Hispanic black Male	12.5	*	0.0	*	*	*	*	0.4	0.7	1.0	1.5
Female	10.2	*	*	*	*	*	*	*	*	0.5	0.9
Non-Hispanic black Male	7.3	*	0.0	0.0	0.0	*	*	*	*	1.3	2.9
Female	7.0	*	0.0	0.0	0.0	*	*	*	*	1.6	3.3
	7.5	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Aortic aneurysm and dissection (I71)

All origins 3/	5.8	*	*	0.0	*	*	0.1	0.2	0.3	0.6	0.9
Male	7.2	*	0.0	0.0	*	*	*	0.3	0.5	0.9	1.2
Female	4.4	0.0	*	0.0	*	*	*	*	*	0.3	0.5
Hispanic	1.2	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	1.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.8	0.0	0.0	0.0	*	*	0.0	*	*	*	*
Mexican	0.8	0.0	0.0	0.0	*	*	0.0	*	*	*	*
Male	1.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0	*	*
Female	0.6	0.0	0.0	0.0	*	0.0	0.0	0.0	*	*	0.0
Puerto Rican	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Male	2.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Cuban	4.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	7.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.7	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	2.5	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	1.1	0.0	0.0	0.0	0.0	*	0.0	*	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Aortic aneurysm and dissection (I71)											
Non-Hispanic 5/	6.4	*	*	0.0	*	*	0.1	0.2	0.3	0.6	0.9
Male	7.9	*	0.0	0.0	*	*	*	0.3	0.6	0.9	1.2
Female	4.9	0.0	*	0.0	0.0	*	*	*	*	0.3	0.6
Non-Hispanic white	7.1	*	*	0.0	*	*	*	0.2	0.3	0.6	0.7
Male	8.9	*	0.0	0.0	*	*	*	*	0.5	0.9	1.0
Female	5.4	0.0	*	0.0	0.0	*	*	*	*	0.3	0.4
Non-Hispanic black	3.5	0.0	0.0	0.0	0.0	0.0	*	*	*	0.9	2.1
Male	3.9	0.0	0.0	0.0	0.0	0.0	*	*	*	*	2.9
Female	3.2	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*	1.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/	3.7	*	*	*	*	*	*	*	0.2	0.2	0.5
Male	3.1	*	0.0	*	*	*	*	*	*	0.2	0.4
Female	4.2	*	*	*	*	*	*	*	*	0.2	0.5
Hispanic	1.1	*	0.0	0.0	*	*	*	*	*	*	*
Male	1.0	*	0.0	0.0	0.0	*	*	*	*	*	*
Female	1.2	*	0.0	0.0	*	*	0.0	*	*	*	*
Mexican	1.0	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	0.9	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	1.1	0.0	0.0	0.0	*	*	0.0	*	*	0.0	*
Puerto Rican	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Cuban	2.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.1	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Male	1.1	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	1.1	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 5/	4.0	*	*	*	*	*	*	*	0.2	0.2	0.5
Male	3.4	*	0.0	*	*	*	*	*	*	*	0.4
Female	4.5	*	*	*	*	*	*	*	*	0.2	0.6
Non-Hispanic white	4.2	*	*	*	*	*	*	*	0.1	0.2	0.5
Male	3.6	*	0.0	*	*	*	*	*	*	*	0.4
Female	4.8	*	*	*	*	*	*	*	*	*	0.5
Non-Hispanic black	3.8	*	0.0	0.0	0.0	*	*	*	*	*	0.8
Male	3.1	*	0.0	0.0	0.0	*	*	*	*	*	*
Female	4.3	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic 5/	1.6	0.7	*	0.0	*	*	*	0.2	0.3	0.5	0.8
Male	1.4	*	*	0.0	0.0	*	*	*	0.3	0.5	1.0
Female	1.8	*	0.0	0.0	*	*	*	*	0.3	0.4	0.6
Non-Hispanic white	1.6	*	0.0	0.0	*	*	*	0.2	0.3	0.4	0.7
Male	1.4	*	0.0	0.0	0.0	*	*	*	*	0.4	0.8
Female	1.8	*	0.0	0.0	*	*	*	*	0.3	0.3	0.5
Non-Hispanic black	1.9	*	*	0.0	0.0	*	*	*	0.8	1.1	2.0
Male	1.8	*	*	0.0	0.0	*	*	*	*	*	2.7
Female	1.9	0.0	0.0	0.0	0.0	*	*	*	*	*	1.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Influenza and pneumonia (J10-J18)

All origins 3/	23.4	8.4	0.9	0.2	0.2	0.4	0.6	0.7	1.0	2.0	2.7
Male	20.8	8.8	0.9	0.3	0.3	0.3	0.6	0.8	1.2	2.2	3.1
Female	25.8	8.0	0.9	0.2	0.2	0.4	0.6	0.6	0.8	1.9	2.3
Hispanic	7.2	8.9	*	*	*	*	*	*	*	1.8	1.3
Male	6.9	10.9	*	*	*	*	*	*	*	2.6	1.8
Female	7.4	6.8	*	*	*	*	*	*	*	*	*
Mexican	5.3	9.0	*	*	*	*	*	*	*	1.7	*
Male	5.3	10.7	*	*	*	*	*	*	*	*	*
Female	5.3	*	*	0.0	*	*	*	*	*	*	*
Puerto Rican	12.0	*	*	*	0.0	*	*	*	*	*	*
Male	11.1	*	*	0.0	0.0	*	*	0.0	0.0	*	*
Female	12.9	*	0.0	*	0.0	*	*	*	*	*	*
Cuban	23.1	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	20.7	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	25.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	7.6	*	*	0.0	*	*	*	*	*	*	*
Male	7.6	*	*	0.0	*	*	0.0	*	*	*	*
Female	7.5	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

Non-Hispanic 5/	25.4	8.2	0.9	0.2	0.2	0.4	0.6	0.8	1.1	2.1	2.8
Male	22.5	8.3	0.9	0.3	0.3	0.3	0.6	0.9	1.3	2.2	3.2
Female	28.0	8.2	0.9	*	*	0.4	0.5	0.7	0.9	2.0	2.5
Non-Hispanic white	27.7	5.1	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.8	2.1
Male	24.1	5.3	0.5	*	0.3	0.3	0.5	0.7	1.1	1.8	2.3
Female	31.2	4.9	0.8	*	*	0.4	0.4	0.5	0.7	1.7	2.0
Non-Hispanic black	17.5	21.6	1.9	*	*	*	1.4	1.9	2.4	4.2	7.2
Male	17.8	21.6	2.4	*	*	*	*	2.1	2.8	4.5	9.4
Female	17.3	21.5	*	*	*	*	*	1.7	2.0	4.0	5.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Influenza (J10-J11)											
All origins 3/	0.6	*	*	*	*	*	*	*	*	*	*
Male	0.5	*	*	*	*	*	*	0.0	*	*	*
Female	0.8	*	*	*	*	*	*	*	*	*	*
Hispanic	0.1	*	*	0.0	0.0	*	*	0.0	0.0	*	*
Male	0.1	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	0.2	*	0.0	0.0	0.0	*	*	0.0	0.0	0.0	0.0
Mexican	0.1	*	*	0.0	0.0	0.0	*	0.0	0.0	0.0	*
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	*	*	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Puerto Rican	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	0.0	*	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Influenza (J10-J11)											
Non-Hispanic 5/	0.7	*	*	*	*	*	*	*	*	*	*
Male	0.5	*	*	*	*	*	*	0.0	*	*	*
Female	0.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white	0.8	*	*	*	*	*	*	*	*	*	*
Male	0.6	*	*	*	*	*	*	0.0	*	*	*
Female	1.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black	0.2	*	*	0.0	0.0	*	0.0	*	0.0	*	*
Male	0.2	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	0.2	*	0.0	0.0	0.0	*	0.0	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Pneumonia (J12-J18)											
All origins 3/	22.8	8.0	0.8	0.2	0.2	0.3	0.6	0.7	1.0	2.0	2.7
Male	20.3	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.2	2.2	3.1
Female	25.1	7.6	0.8	*	*	0.3	0.5	0.6	0.8	1.8	2.3
Hispanic	7.0	8.3	*	*	*	*	*	*	*	1.7	1.2
Male	6.8	10.9	*	*	*	*	*	*	*	2.5	1.7
Female	7.3	5.7	*	*	*	*	*	*	*	*	*
Mexican	5.2	8.4	*	*	*	*	*	*	*	1.7	*
Male	5.2	10.7	*	*	*	*	*	*	*	*	*
Female	5.1	*	*	0.0	*	*	*	*	*	*	*
Puerto Rican	11.9	*	*	*	0.0	*	*	*	*	*	*
Male	10.9	*	*	0.0	0.0	*	*	0.0	0.0	*	*
Female	12.8	0.0	0.0	*	0.0	*	*	*	*	*	*
Cuban	23.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	20.7	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	25.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	7.3	*	*	0.0	*	*	*	*	*	*	*
Male	7.4	*	*	0.0	*	*	0.0	*	*	*	*
Female	7.3	*	*	0.0	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Pneumonia (J12-J18)											
Non-Hispanic 5/	24.7	7.9	0.8	0.2	0.2	0.3	0.6	0.8	1.1	2.0	2.8
Male	22.0	7.9	0.8	0.2	0.3	0.3	0.6	0.9	1.3	2.1	3.2
Female	27.2	8.0	0.9	*	*	0.4	0.5	0.7	0.9	1.9	2.4
Non-Hispanic white	26.9	5.0	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.7	2.1
Male	23.5	5.2	0.4	*	*	0.3	0.5	0.7	1.0	1.8	2.2
Female	30.2	4.8	0.7	*	*	0.3	0.3	0.5	0.7	1.7	2.0
Non-Hispanic black	17.3	20.8	1.9	*	*	*	1.4	1.9	2.4	4.1	7.1
Male	17.6	20.5	2.3	*	*	*	*	2.1	2.8	4.4	9.3
Female	17.0	21.1	*	*	*	*	*	1.6	2.0	3.8	5.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

Non-Hispanic 5/	0.2	1.7	0.2	*	*	*	*	*	*	*	*
Male	0.2	2.0	*	0.0	*	*	0.0	*	*	*	*
Female	0.3	1.5	*	*	*	*	*	0.0	*	*	*
Non-Hispanic white	0.2	1.1	*	*	*	*	*	*	*	*	*
Male	0.2	*	*	0.0	*	*	0.0	*	*	*	*
Female	0.3	*	*	*	*	*	*	0.0	*	*	*
Non-Hispanic black	0.2	4.5	*	0.0	0.0	*	0.0	*	0.0	0.0	*
Male	0.2	*	*	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Female	0.2	*	*	0.0	0.0	*	0.0	0.0	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute bronchitis and bronchiolitis (J20-J21)											
Non-Hispanic 5/	0.2	1.7	0.2	*	*	*	*	*	*	*	*
Male	0.1	2.0	*	0.0	*	*	0.0	*	*	*	*
Female	0.2	1.4	*	*	0.0	*	*	0.0	*	*	*
Non-Hispanic white	0.2	1.1	*	*	*	*	*	*	*	*	*
Male	0.1	*	*	0.0	*	*	0.0	*	*	*	0.0
Female	0.2	*	*	*	0.0	*	*	0.0	*	*	*
Non-Hispanic black	0.2	4.5	*	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Male	0.2	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.2	*	*	0.0	0.0	*	0.0	0.0	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Chronic lower respiratory diseases (J40-J47)

All origins 3/	45.5	0.9	0.4	0.2	0.5	0.5	0.6	0.8	0.9	1.3	2.6
Male	46.8	1.0	0.4	0.3	0.6	0.6	0.7	0.8	0.8	1.3	2.4
Female	44.3	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4	2.8
Hispanic	9.1	*	*	*	*	*	*	*	*	1.3	2.1
Male	9.8	*	*	*	*	*	*	*	*	*	2.0
Female	8.4	*	*	*	*	*	*	*	*	1.6	2.1
Mexican	6.3	*	*	*	*	*	*	*	*	*	*
Male	6.9	*	*	*	*	*	*	*	*	*	*
Female	5.7	*	*	*	*	*	0.0	*	*	*	*
Puerto Rican	14.8	0.0	*	*	*	*	*	*	*	*	*
Male	14.9	0.0	0.0	*	*	0.0	*	*	*	*	*
Female	14.6	0.0	*	*	0.0	*	*	*	*	*	*
Cuban	36.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	43.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	29.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	9.7	*	*	*	*	*	*	*	*	*	*
Male	10.3	*	*	0.0	*	*	0.0	0.0	*	*	*
Female	9.2	0.0	*	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 5/	50.1	1.0	0.4	0.3	0.5	0.6	0.6	0.8	1.0	1.3	2.7
Male	51.6	*	0.4	0.3	0.7	0.6	0.7	0.8	0.9	1.3	2.5
Female	48.7	*	0.3	*	0.3	0.5	0.5	0.8	1.1	1.4	2.9
Non-Hispanic white	56.9	*	0.2	0.2	0.2	0.5	0.4	0.5	0.7	1.0	2.2
Male	57.6	*	*	*	0.3	0.5	0.4	0.4	0.6	1.0	2.0
Female	56.3	*	*	*	*	0.4	0.4	0.5	0.7	1.1	2.5
Non-Hispanic black	23.7	*	*	0.6	1.6	1.2	1.6	2.7	2.9	3.4	6.3
Male	28.4	*	*	*	2.2	1.6	2.3	2.9	2.8	3.4	6.3
Female	19.5	*	*	*	*	*	*	2.6	3.0	3.4	6.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Bronchitis, chronic and unspecified (J40-J42)

All origins 3/	0.4	0.5	0.1	*	*	*	*	*	*	*	*
Male	0.4	*	*	*	*	*	*	0.0	*	*	*
Female	0.5	*	*	*	0.0	*	*	*	*	*	*
Hispanic	0.2	*	*	0.0	*	0.0	0.0	*	0.0	*	*
Male	*	*	*	0.0	*	0.0	0.0	0.0	0.0	*	*
Female	0.2	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Mexican	0.1	*	*	0.0	*	0.0	0.0	*	0.0	*	*
Male	*	*	*	0.0	*	0.0	0.0	0.0	0.0	*	0.0
Female	0.2	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	0.0	*	0.0	0.0	0.0	0.0	*	0.0	*	0.0
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Bronchitis, chronic and unspecified (J40-J42)											
Non-Hispanic 5/	0.5	*	*	*	0.0	*	*	*	*	*	*
Male	0.4	*	*	*	0.0	*	*	0.0	*	*	*
Female	0.5	*	*	*	0.0	*	*	*	*	*	*
Non-Hispanic white	0.5	*	*	*	0.0	*	*	0.0	*	*	*
Male	0.4	*	*	*	0.0	*	*	0.0	*	*	*
Female	0.6	*	*	*	0.0	*	*	0.0	*	*	*
Non-Hispanic black	0.3	*	*	0.0	0.0	0.0	*	*	*	0.0	*
Male	0.3	*	*	0.0	0.0	0.0	*	0.0	*	0.0	0.0
Female	0.3	*	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Emphysema (J43)

All origins 3/	6.5	*	0.0	*	0.0	0.0	*	*	*	0.1	0.4
Male	7.1	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	0.4
Female	6.0	*	0.0	*	0.0	0.0	0.0	*	*	*	0.3
Hispanic	0.9	0.0	0.0	*	0.0	0.0	0.0	0.0	*	*	*
Male	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	0.7	0.0	0.0	*	0.0	0.0	0.0	0.0	*	*	*
Mexican	0.7	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*
Male	0.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.5	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*
Puerto Rican	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	4.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	5.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Male	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Asthma (J45-J46)

All origins 3/	1.7	*	0.2	0.2	0.4	0.5	0.5	0.7	0.7	0.9	1.1
Male	1.2	*	*	0.2	0.6	0.5	0.6	0.7	0.7	0.8	0.8
Female	2.2	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0	1.3
Hispanic	1.0	0.0	*	*	*	*	*	*	*	0.9	1.0
Male	0.8	0.0	0.0	*	*	*	*	*	*	*	*
Female	1.3	0.0	*	*	*	*	*	*	*	*	*
Mexican	0.5	0.0	*	*	*	*	*	*	*	*	*
Male	0.4	0.0	0.0	*	*	*	*	*	*	*	*
Female	0.7	0.0	*	0.0	*	*	0.0	*	*	*	*
Puerto Rican	3.7	0.0	*	*	*	*	*	*	*	*	*
Male	2.9	0.0	0.0	0.0	*	0.0	*	*	*	*	*
Female	4.5	0.0	*	*	0.0	*	*	*	*	*	*
Cuban	2.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	3.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	1.1	0.0	0.0	*	*	*	*	*	*	*	*
Male	0.8	0.0	0.0	0.0	*	*	0.0	0.0	*	*	*
Female	1.3	0.0	0.0	*	*	0.0	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Asthma (J45-J46)											
Non-Hispanic 5/	1.8	*	0.2	0.2	0.5	0.5	0.5	0.7	0.8	0.9	1.1
Male	1.3	*	*	0.3	0.6	0.6	0.7	0.7	0.7	0.9	0.8
Female	2.3	*	*	*	0.3	0.5	0.4	0.6	0.8	1.0	1.4
Non-Hispanic white	1.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.6	0.7
Male	1.0	*	*	*	0.3	0.4	0.4	0.4	0.4	0.5	0.4
Female	2.1	0.0	*	*	*	0.4	*	0.4	0.5	0.7	1.1
Non-Hispanic black	3.4	*	*	0.6	1.6	1.1	1.5	2.5	2.5	2.8	3.4
Male	3.0	*	*	*	2.2	1.4	2.1	2.9	2.7	2.8	2.9
Female	3.8	*	*	*	*	*	*	2.2	2.4	2.8	3.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other chronic lower respiratory diseases (J44,J47)

All origins 3/	36.9	*	*	*	*	*	*	0.1	0.1	0.3	1.2
Male	38.1	*	*	*	*	*	*	*	*	0.3	1.2
Female	35.7	*	*	*	0.0	*	*	*	*	0.3	1.1
Hispanic	7.0	*	*	*	0.0	0.0	*	*	*	*	*
Male	7.8	*	*	*	0.0	0.0	*	0.0	*	*	*
Female	6.2	0.0	*	*	0.0	0.0	0.0	*	*	0.0	*
Mexican	5.0	*	*	*	0.0	0.0	*	0.0	*	*	*
Male	5.6	*	*	0.0	0.0	0.0	*	0.0	*	*	*
Female	4.3	0.0	0.0	*	0.0	0.0	0.0	0.0	*	0.0	*
Puerto Rican	10.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0	0.0	*
Male	10.6	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	9.4	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Cuban	28.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	35.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	22.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	7.5	*	*	0.0	0.0	0.0	0.0	*	*	*	*
Male	8.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	7.0	0.0	*	0.0	0.0	0.0	0.0	*	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other chronic lower respiratory diseases (J44,J47)											
Non-Hispanic 5/	40.6	*	*	*	*	*	*	*	0.1	0.3	1.2
Male	42.1	*	*	*	*	*	*	*	*	0.3	1.3
Female	39.3	*	*	*	0.0	*	*	*	*	0.3	1.1
Non-Hispanic white	46.6	*	*	*	*	*	*	*	*	0.3	1.1
Male	47.3	0.0	0.0	*	*	*	*	*	*	0.3	1.2
Female	45.8	*	*	*	0.0	*	*	*	*	0.3	1.1
Non-Hispanic black	17.2	*	*	0.0	*	*	*	*	*	*	2.1
Male	21.3	*	*	0.0	*	*	*	0.0	0.0	*	2.6
Female	13.5	*	*	0.0	0.0	0.0	*	*	*	*	1.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

Non-Hispanic 5/	0.5	0.0	*	0.0	0.0	*	0.0	0.0	0.0	*	*
Male	0.9	0.0	*	0.0	0.0	*	0.0	0.0	0.0	*	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Non-Hispanic white	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Non-Hispanic black	0.2	0.0	*	0.0	0.0	*	0.0	0.0	0.0	*	*
Male	0.4	0.0	*	0.0	0.0	*	0.0	0.0	0.0	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Pneumonitis due to solids and liquids (J69)

All origins 3/	5.6	*	*	*	*	*	0.1	0.2	0.2	0.3	0.5
Male	5.8	*	*	0.0	*	*	*	0.2	0.3	0.3	0.6
Female	5.4	*	*	*	*	*	*	*	0.2	0.2	0.4
Hispanic	1.3	*	*	*	*	*	*	*	*	*	*
Male	1.2	*	*	0.0	*	0.0	0.0	*	*	*	0.0
Female	1.4	*	*	*	*	*	*	*	0.0	0.0	*
Mexican	1.0	*	0.0	*	*	0.0	0.0	*	*	*	0.0
Male	1.0	*	0.0	0.0	*	0.0	0.0	*	*	*	0.0
Female	1.0	*	0.0	*	0.0	0.0	0.0	*	0.0	0.0	0.0
Puerto Rican	1.9	0.0	*	*	*	*	*	*	0.0	0.0	*
Male	1.6	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Female	2.2	0.0	*	*	*	*	*	0.0	0.0	0.0	*
Cuban	5.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	4.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	7.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.2	0.0	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
Male	1.4	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	1.1	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Pneumonitis due to solids and liquids (J69)											
Non-Hispanic 5/	6.1	*	*	*	*	*	0.1	0.2	0.3	0.3	0.5
Male	6.4	*	*	0.0	*	*	*	0.2	0.3	0.3	0.6
Female	5.9	*	*	*	*	*	*	*	0.2	0.2	0.4
Non-Hispanic white	6.7	*	*	*	*	*	*	0.2	0.2	0.3	0.5
Male	6.9	*	*	0.0	*	*	*	*	0.3	0.3	0.5
Female	6.5	*	*	*	*	*	*	*	*	*	0.4
Non-Hispanic black	4.5	*	*	0.0	*	*	*	*	*	*	1.0
Male	4.9	*	*	0.0	*	*	*	*	*	*	*
Female	4.2	*	*	0.0	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/	9.2	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.6
Male	9.1	7.4	0.7	0.2	0.2	0.4	0.3	0.5	0.8	1.1	1.7
Female	9.3	5.7	0.5	*	0.2	0.2	0.3	0.6	0.8	0.9	1.4
Hispanic	3.7	5.3	*	*	*	*	*	*	*	*	1.5
Male	3.7	5.4	*	0.0	*	*	*	*	*	*	2.1
Female	3.7	*	*	*	0.0	*	*	*	*	*	*
Mexican	3.1	5.3	*	*	*	*	*	*	*	*	1.7
Male	3.2	*	*	0.0	*	*	*	*	*	*	*
Female	3.0	*	*	*	0.0	*	*	*	*	*	*
Puerto Rican	4.9	*	0.0	*	0.0	*	0.0	*	*	*	*
Male	5.6	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	4.2	*	0.0	*	0.0	*	0.0	*	*	*	0.0
Cuban	9.8	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Male	8.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	11.5	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Other Hispanic 4/	3.7	*	*	0.0	0.0	*	*	*	*	*	*
Male	3.7	*	0.0	0.0	0.0	*	*	*	*	*	*
Female	3.8	*	*	0.0	0.0	*	*	0.0	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 5/	9.9	6.8	0.7	0.2	0.2	0.3	0.3	0.6	0.8	1.1	1.6
Male	9.8	7.8	0.8	0.2	*	0.4	0.4	0.5	0.9	1.2	1.7
Female	9.9	5.8	0.5	*	0.2	*	0.3	0.6	0.8	1.0	1.4
Non-Hispanic white	10.7	4.7	0.5	0.2	0.2	0.3	0.3	0.5	0.7	0.9	1.3
Male	10.6	5.4	0.6	*	*	0.4	0.4	0.6	0.8	1.0	1.4
Female	10.8	3.9	0.5	*	*	*	*	0.4	0.7	0.8	1.1
Non-Hispanic black	7.1	17.0	1.3	*	*	*	*	1.2	1.7	2.5	3.5
Male	6.9	19.4	1.8	*	*	*	*	*	1.9	2.6	3.3
Female	7.2	14.6	*	*	*	*	*	1.9	1.6	2.3	3.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Peptic ulcer (K25-K28)

Non-Hispanic 5/	1.8	*	*	0.0	0.0	*	*	*	*	0.2	0.4
Male	1.8	0.0	0.0	0.0	0.0	*	*	*	*	0.3	0.5
Female	1.9	*	*	0.0	0.0	0.0	0.0	*	*	*	0.3
Non-Hispanic white	2.0	*	*	0.0	0.0	*	*	*	*	0.1	0.3
Male	1.9	0.0	0.0	0.0	0.0	*	*	*	*	*	0.3
Female	2.1	*	*	0.0	0.0	0.0	0.0	0.0	*	*	0.3
Non-Hispanic black	1.3	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	1.0
Male	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	1.6
Female	1.2	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Diseases of appendix (K35-K38)

Non-Hispanic 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
Male	0.2	0.0	*	*	*	*	*	*	*	*	*	*
Female	0.1	*	0.0	*	*	0.0	0.0	0.0	*	*	*	*
Non-Hispanic white	0.1	*	0.0	*	*	*	0.0	*	*	*	*	*
Male	0.2	0.0	0.0	*	*	*	0.0	*	0.0	*	*	*
Female	0.1	*	0.0	*	*	0.0	0.0	0.0	*	*	*	*
Non-Hispanic black	0.2	0.0	*	*	*	0.0	*	0.0	*	*	*	*
Male	0.3	0.0	*	*	*	0.0	*	0.0	*	*	*	*
Female	0.1	0.0	0.0	*	*	0.0	0.0	0.0	0.0	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hernia (K40-K46)											
Non-Hispanic 5/	0.6	1.3	*	*	*	*	*	*	*	*	0.1
Male	0.5	1.5	*	*	0.0	*	*	*	*	*	*
Female	0.7	*	*	0.0	*	0.0	0.0	*	*	*	*
Non-Hispanic white	0.6	1.3	*	*	*	*	*	*	*	*	*
Male	0.5	*	*	*	0.0	*	*	*	*	*	*
Female	0.7	*	*	0.0	*	0.0	0.0	*	*	*	*
Non-Hispanic black	0.4	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Male	0.3	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Female	0.5	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/	9.6	*	*	*	*	*	*	0.4	1.7	4.9	9.9
Male	12.8	*	0.0	*	0.0	*	*	0.5	2.2	6.4	14.5
Female	6.6	*	*	0.0	*	*	*	0.3	1.2	3.4	5.4
Hispanic	9.5	0.0	0.0	0.0	0.0	*	*	*	2.7	7.4	15.2
Male	13.6	0.0	0.0	0.0	0.0	*	*	*	4.1	11.6	24.4
Female	5.3	0.0	0.0	0.0	0.0	*	*	*	*	2.9	5.7
Mexican	9.1	0.0	0.0	0.0	0.0	*	*	*	2.8	8.5	16.0
Male	13.0	0.0	0.0	0.0	0.0	*	*	*	4.1	13.1	25.3
Female	5.0	0.0	0.0	0.0	0.0	*	*	*	*	3.3	6.2
Puerto Rican	14.3	0.0	0.0	0.0	0.0	*	*	0.0	*	*	22.7
Male	22.9	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Female	6.4	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Cuban	8.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	11.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Female	7.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	8.5	0.0	0.0	0.0	0.0	*	0.0	*	3.6	5.5	12.4
Male	11.9	0.0	0.0	0.0	0.0	*	0.0	*	*	8.8	19.3
Female	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 5/	9.6	*	*	*	*	*	*	0.4	1.5	4.6	9.2
Male	12.7	*	0.0	*	0.0	0.0	*	0.5	1.9	5.7	13.3
Female	6.7	*	*	0.0	*	*	*	0.4	1.2	3.5	5.3
Non-Hispanic white	10.0	*	0.0	0.0	*	*	*	0.4	1.5	4.4	8.7
Male	13.1	*	0.0	0.0	0.0	0.0	*	0.5	1.8	5.5	12.5
Female	7.0	0.0	0.0	0.0	*	*	*	*	1.1	3.3	4.9
Non-Hispanic black	8.4	0.0	*	*	0.0	*	*	*	1.5	4.7	12.4
Male	11.6	0.0	0.0	*	0.0	0.0	*	*	1.9	5.8	17.8
Female	5.6	0.0	*	0.0	0.0	*	*	*	*	3.7	7.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Alcoholic liver disease (K70)											
All origins 3/	4.4	0.0	0.0	0.0	0.0	0.0	*	0.2	1.1	3.4	6.4
Male	6.7	0.0	0.0	0.0	0.0	0.0	*	0.3	1.4	4.4	9.3
Female	2.2	0.0	0.0	0.0	0.0	0.0	*	*	0.8	2.4	3.5
Hispanic	4.9	0.0	0.0	0.0	0.0	0.0	*	*	1.7	5.0	9.9
Male	8.2	0.0	0.0	0.0	0.0	0.0	*	*	2.7	7.6	16.0
Female	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	2.2	3.6
Mexican	4.8	0.0	0.0	0.0	0.0	0.0	*	*	1.6	5.7	10.4
Male	7.9	0.0	0.0	0.0	0.0	0.0	*	*	2.4	8.4	17.0
Female	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	3.5
Puerto Rican	7.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	13.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	2.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	2.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Male	4.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	4.7	0.0	0.0	0.0	0.0	0.0	0.0	*	*	3.8	9.0
Male	7.8	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	13.5
Female	1.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Alcoholic liver disease (K70)											
Non-Hispanic 5/	4.3	0.0	0.0	0.0	0.0	0.0	*	0.2	1.0	3.2	6.0
Male	6.5	0.0	0.0	0.0	0.0	0.0	*	0.3	1.2	3.9	8.5
Female	2.2	0.0	0.0	0.0	0.0	0.0	*	*	0.8	2.4	3.5
Non-Hispanic white	4.3	0.0	0.0	0.0	0.0	0.0	*	0.2	0.9	3.0	5.6
Male	6.6	0.0	0.0	0.0	0.0	0.0	*	*	1.2	3.8	8.0
Female	2.2	0.0	0.0	0.0	0.0	0.0	*	*	0.7	2.3	3.2
Non-Hispanic black	4.4	0.0	0.0	0.0	0.0	0.0	*	*	1.0	3.1	7.5
Male	6.4	0.0	0.0	0.0	0.0	0.0	*	*	*	3.8	10.6
Female	2.5	0.0	0.0	0.0	0.0	0.0	0.0	*	*	2.4	4.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other chronic liver disease and cirrhosis (K73-K74)

All origins 3/	5.2	*	*	*	*	*	*	0.2	0.6	1.5	3.5
Male	6.1	*	0.0	*	0.0	*	*	*	0.8	2.0	5.2
Female	4.4	*	*	0.0	*	*	*	*	0.4	1.0	1.8
Hispanic	4.5	0.0	0.0	0.0	0.0	*	*	*	1.0	2.4	5.3
Male	5.4	0.0	0.0	0.0	0.0	*	0.0	*	1.4	4.0	8.4
Female	3.7	0.0	0.0	0.0	0.0	*	*	*	*	*	2.1
Mexican	4.4	0.0	0.0	0.0	0.0	*	*	*	1.2	2.8	5.6
Male	5.1	0.0	0.0	0.0	0.0	*	0.0	*	*	4.7	8.3
Female	3.5	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Puerto Rican	6.7	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Male	9.5	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Female	4.1	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Cuban	6.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Male	6.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	6.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	3.8	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Male	4.1	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Female	3.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other chronic liver disease and cirrhosis (K73-K74)											
Non-Hispanic 5/	5.3	*	*	*	*	*	*	0.2	0.5	1.4	3.3
Male	6.2	*	0.0	*	0.0	0.0	*	*	0.7	1.7	4.8
Female	4.5	*	*	0.0	*	*	*	*	0.4	1.1	1.8
Non-Hispanic white	5.7	*	0.0	0.0	*	*	*	*	0.5	1.4	3.1
Male	6.6	*	0.0	0.0	0.0	0.0	*	*	0.7	1.7	4.5
Female	4.8	0.0	0.0	0.0	*	*	0.0	*	0.4	1.0	1.7
Non-Hispanic black	4.1	0.0	*	*	0.0	*	*	*	*	1.6	4.9
Male	5.2	0.0	0.0	*	0.0	0.0	*	*	*	2.0	7.3
Female	3.1	0.0	*	0.0	0.0	*	*	*	*	1.3	2.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic 5/	1.1	*	0.0	0.0	*	*	*	*	*	0.1	0.1
Male	1.0	*	0.0	0.0	*	*	*	0.0	*	*	*
Female	1.2	*	0.0	0.0	0.0	*	*	*	*	*	*
Non-Hispanic white	1.2	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	1.1	0.0	0.0	0.0	*	*	*	0.0	*	*	*
Female	1.3	0.0	0.0	0.0	0.0	*	0.0	*	*	*	*
Non-Hispanic black	0.8	*	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.7	*	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Female	0.9	*	0.0	0.0	0.0	0.0	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/	13.0	4.3	*	*	*	0.1	0.3	0.5	0.8	1.2	2.0
Male	12.8	4.6	*	*	*	*	0.2	0.5	0.8	1.4	2.3
Female	13.3	4.1	*	*	*	*	0.3	0.6	0.7	1.0	1.7
Hispanic	5.1	2.8	*	*	*	*	*	*	*	0.9	1.8
Male	5.2	*	*	*	*	0.0	*	*	*	*	1.7
Female	4.9	*	*	*	*	*	*	*	*	*	1.9
Mexican	4.2	*	*	*	*	*	*	*	*	*	*
Male	4.3	*	*	*	*	0.0	0.0	*	*	*	*
Female	4.1	*	*	*	0.0	*	*	*	*	*	*
Puerto Rican	8.0	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	8.0	*	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Female	7.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Cuban	12.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	13.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	10.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	4.9	*	*	0.0	*	0.0	*	*	*	*	*
Male	4.9	*	*	0.0	*	0.0	0.0	0.0	0.0	*	*
Female	5.0	*	0.0	0.0	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 5/	14.0	4.7	*	*	*	0.1	0.3	0.6	0.8	1.2	2.0
Male	13.7	4.7	*	*	*	*	0.3	0.5	0.9	1.4	2.3
Female	14.3	4.7	*	*	*	*	0.3	0.6	0.8	1.0	1.7
Non-Hispanic white	13.5	3.4	*	*	*	*	0.2	0.3	0.5	0.8	1.2
Male	13.3	3.8	*	*	*	*	*	0.3	0.6	0.9	1.4
Female	13.6	3.0	*	*	*	*	*	0.4	0.5	0.6	1.0
Non-Hispanic black	20.1	11.0	*	*	*	*	0.9	1.6	2.5	4.3	7.3
Male	19.0	9.3	*	*	*	*	*	*	2.5	5.2	8.7
Female	21.1	12.7	0.0	*	*	*	*	1.8	2.5	3.5	6.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07, N26)

Non-Hispanic 5/	0.2	0.0	0.0	*	0.0	*	*	*	*	*	*
Male	0.2	0.0	0.0	*	0.0	*	*	*	*	*	*
Female	0.2	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Non-Hispanic white	0.2	0.0	0.0	*	0.0	*	*	*	*	*	*
Male	0.2	0.0	0.0	*	0.0	*	0.0	*	*	*	*
Female	0.2	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Non-Hispanic black	0.3	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Male	0.3	0.0	0.0	0.0	0.0	*	*	0.0	*	*	*
Female	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Renal failure (N17-N19)

All origins 3/	12.7	4.3	*	*	*	0.1	0.2	0.5	0.7	1.1	1.9
Male	12.5	4.5	*	*	*	*	*	0.4	0.8	1.3	2.2
Female	13.0	4.1	*	*	*	*	0.3	0.5	0.7	0.9	1.6
Hispanic	4.9	2.6	*	*	*	*	*	*	*	0.9	1.7
Male	5.0	*	*	*	*	0.0	*	*	*	*	*
Female	4.8	*	0.0	0.0	*	*	*	*	*	*	1.9
Mexican	4.1	*	*	*	*	*	*	*	*	*	*
Male	4.2	*	*	*	*	0.0	0.0	*	*	*	*
Female	3.9	*	0.0	0.0	0.0	*	*	*	*	*	*
Puerto Rican	7.7	*	0.0	0.0	0.0	0.0	*	*	*	*	*
Male	7.7	*	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Female	7.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Cuban	11.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	13.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	10.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	4.8	*	*	0.0	*	0.0	*	*	*	*	*
Male	4.7	*	*	0.0	*	0.0	0.0	0.0	0.0	*	*
Female	4.8	*	0.0	0.0	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Renal failure (N17-N19)											
Non-Hispanic 5/	13.7	4.6	*	*	*	0.1	0.2	0.5	0.8	1.1	1.9
Male	13.4	4.5	*	*	*	*	*	0.4	0.8	1.3	2.2
Female	14.0	4.7	*	*	*	*	0.3	0.5	0.8	1.0	1.6
Non-Hispanic white	13.2	3.3	*	*	*	*	0.2	0.3	0.5	0.7	1.2
Male	13.0	3.6	*	*	*	*	*	*	0.5	0.8	1.4
Female	13.3	3.0	*	*	*	*	*	*	0.5	0.6	0.9
Non-Hispanic black	19.7	11.0	*	*	*	*	0.7	1.5	2.4	4.1	7.1
Male	18.6	9.3	*	*	*	*	*	*	2.5	4.9	8.4
Female	20.7	12.7	0.0	*	*	*	*	1.8	2.4	3.4	5.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic 5/	0.3	*	*	*	0.0	*	*	*	*	*	*
Male	0.2	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	0.4	*	*	*	0.0	*	*	*	*	*	*
Non-Hispanic white	0.4	*	*	*	0.0	*	*	*	*	*	*
Male	0.2	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	0.5	0.0	*	*	0.0	0.0	*	*	*	*	*
Non-Hispanic black	0.3	*	0.0	0.0	0.0	*	0.0	*	0.0	*	*
Male	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.3	*	0.0	0.0	0.0	*	0.0	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Non-Hispanic 5/	0.0	*	*	0.0	0.0	*	*	*	*	*	*
Female	0.1	*	*	0.0	0.0	*	*	*	*	*	*
Non-Hispanic white	0.0	*	*	0.0	0.0	*	*	*	*	*	*
Female	0.1	*	*	0.0	0.0	*	*	*	*	*	*
Non-Hispanic black	0.1	*	0.0	0.0	0.0	0.0	*	*	0.0	*	*
Female	0.1	*	0.0	0.0	0.0	0.0	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

All origins 3/	0.1	0.0	0.0	0.0	0.0	0.2	0.4	0.5	0.5	0.4	0.1
Female	0.3	0.0	0.0	0.0	0.0	0.3	0.7	1.0	0.9	0.8	0.3
Hispanic	0.2	0.0	0.0	0.0	0.0	*	*	*	*	0.8	*
Female	0.4	0.0	0.0	0.0	0.0	*	*	*	*	1.7	*
Mexican	0.2	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.4	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Puerto Rican	...	0.0	0.0	0.0	0.0	*	*	*	*	0.0	*
Female	...	0.0	0.0	0.0	0.0	*	*	*	*	0.0	*
Cuban	...	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	...	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	0.3	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic 5/	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.4	0.5	0.3	0.1
Female	0.3	0.0	0.0	0.0	0.0	0.3	0.6	0.9	1.0	0.7	0.3
Non-Hispanic white	0.1	0.0	0.0	0.0	0.0	*	*	0.2	0.3	0.2	*
Female	0.2	0.0	0.0	0.0	0.0	*	*	0.5	0.7	0.4	*
Non-Hispanic black	0.5	0.0	0.0	0.0	0.0	*	1.1	1.5	1.3	1.0	*
Female	0.9	0.0	0.0	0.0	0.0	*	2.2	2.9	2.5	1.8	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy with abortive outcome (000-007)

All origins 3/	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Hispanic	...	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0
Female	...	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0
Mexican	...	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0
Female	...	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*	0.0
Puerto Rican	...	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Female	...	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	...	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	...	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic 5/	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Non-Hispanic white	...	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	...	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Non-Hispanic black	...	0.0	0.0	0.0	0.0	*	*	*	*	*	0.0
Female	...	0.0	0.0	0.0	0.0	*	*	*	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All origins 3/	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.5	0.4	0.4	0.1
Female	0.3	0.0	0.0	0.0	0.0	0.3	0.7	0.9	0.8	0.7	0.3
Hispanic	0.2	0.0	0.0	0.0	0.0	*	*	*	*	0.7	*
Female	0.4	0.0	0.0	0.0	0.0	*	*	*	*	1.5	*
Mexican	0.2	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.4	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Puerto Rican	...	0.0	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Female	...	0.0	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Cuban	...	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	...	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	0.3	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Non-Hispanic 5/	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.4	0.4	0.3	0.1
Female	0.2	0.0	0.0	0.0	0.0	0.3	0.6	0.8	0.9	0.6	0.3
Non-Hispanic white	0.1	0.0	0.0	0.0	0.0	*	*	0.2	0.3	0.2	*
Female	0.1	0.0	0.0	0.0	0.0	*	*	0.5	0.6	0.4	*
Non-Hispanic black	0.4	0.0	0.0	0.0	0.0	*	1.0	1.3	1.1	0.9	*
Female	0.8	0.0	0.0	0.0	0.0	*	2.0	2.6	2.2	1.6	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

Non-Hispanic 5/	4.9	376.6	0.6	0.1	*	*	*	*	*	*	*
Male	5.7	415.9	0.8	*	*	*	*	0.0	*	*	0.0
Female	4.1	335.4	0.4	*	*	*	*	*	*	0.0	*
Non-Hispanic white	3.3	265.6	0.4	*	*	*	*	*	*	*	*
Male	3.8	294.4	0.5	*	*	*	*	0.0	*	*	0.0
Female	2.8	235.3	*	*	*	*	0.0	*	*	0.0	*
Non-Hispanic black	15.1	938.2	1.4	*	*	0.0	*	0.0	0.0	0.0	0.0
Male	18.0	1,039.8	1.8	*	*	0.0	0.0	0.0	0.0	0.0	0.0
Female	12.5	833.3	*	0.0	*	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/	3.8	143.3	3.6	1.0	1.1	1.1	1.2	1.2	1.3	1.2	1.2
Male	4.0	147.3	4.0	1.0	1.1	1.4	1.3	1.2	1.4	1.4	1.3
Female	3.6	139.1	3.2	1.1	1.1	0.9	1.0	1.1	1.2	1.1	1.1
Hispanic	4.8	148.6	3.7	1.0	0.9	1.0	1.1	1.1	1.0	*	0.9
Male	4.8	145.1	4.5	*	*	*	*	*	*	*	*
Female	4.8	152.2	2.8	*	*	*	*	*	*	*	*
Mexican	5.3	159.7	3.5	0.8	1.0	1.1	1.0	*	*	*	*
Male	5.2	156.7	4.1	*	*	*	*	*	*	*	*
Female	5.4	162.8	2.8	*	*	*	*	*	*	*	*
Puerto Rican	4.0	*	*	*	*	*	*	*	*	*	*
Male	3.9	*	*	*	0.0	*	*	*	*	0.0	*
Female	4.1	*	*	0.0	*	0.0	*	*	0.0	*	*
Cuban	2.4	*	*	*	*	0.0	0.0	0.0	*	*	0.0
Male	*	*	*	0.0	*	0.0	0.0	0.0	0.0	*	0.0
Female	*	*	0.0	*	0.0	0.0	0.0	0.0	*	0.0	0.0
Other Hispanic 4/	4.0	129.9	4.8	*	*	*	*	*	*	*	*
Male	4.2	*	*	*	*	*	0.0	*	*	*	*
Female	3.8	*	*	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 5/	3.7	140.4	3.6	1.0	1.2	1.1	1.2	1.2	1.3	1.3	1.3
Male	3.9	145.9	3.9	1.0	1.2	1.4	1.4	1.2	1.5	1.5	1.4
Female	3.5	134.6	3.3	1.1	1.2	0.9	1.0	1.1	1.2	1.2	1.2
Non-Hispanic white	3.5	132.1	3.0	0.9	1.0	1.1	1.1	1.2	1.4	1.3	1.3
Male	3.6	139.6	3.2	0.8	1.0	1.4	1.3	1.3	1.5	1.5	1.3
Female	3.3	124.3	2.8	1.0	1.0	0.8	0.9	1.2	1.2	1.2	1.2
Non-Hispanic black	5.1	187.9	6.6	1.7	1.9	1.5	1.6	1.1	1.3	1.5	1.3
Male	5.6	192.8	7.5	1.6	2.0	1.8	1.9	*	*	1.5	1.8
Female	4.6	182.9	5.7	1.7	1.9	*	*	*	*	1.6	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/ 2/	UNDER 1 YEAR 2/ 3/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------------	--------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/	9.8	92.6	1.6	0.3	0.4	1.2	2.1	2.4	3.3	4.2	5.0
Male	9.7	106.1	1.8	0.3	0.4	1.5	2.9	3.2	4.4	5.3	6.7
Female	9.9	78.5	1.4	0.3	0.3	0.8	1.3	1.7	2.2	3.0	3.4
Hispanic	4.1	57.5	1.1	*	*	1.1	2.0	2.3	3.0	3.1	3.1
Male	4.9	64.4	*	*	*	1.5	3.1	3.2	4.1	4.4	4.7
Female	3.3	50.3	*	*	*	*	*	*	1.8	1.7	*
Mexican	3.4	56.4	1.1	*	*	*	1.7	2.3	2.2	2.2	1.9
Male	4.2	63.6	*	*	*	*	2.5	3.5	3.2	3.4	*
Female	2.6	49.0	*	*	*	*	*	*	*	*	*
Puerto Rican	6.5	*	*	*	*	*	*	*	*	*	*
Male	7.8	*	0.0	*	*	*	*	*	*	*	*
Female	5.3	*	*	0.0	0.0	*	*	*	*	*	*
Cuban	7.0	*	0.0	0.0	0.0	*	*	*	*	*	*
Male	7.7	*	0.0	0.0	0.0	*	*	*	*	*	*
Female	6.3	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Other Hispanic 4/	4.5	52.1	*	*	*	*	*	*	3.9	*	*
Male	5.2	*	*	*	*	*	*	*	*	*	*
Female	3.9	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 5/	10.4	100.0	1.6	0.4	0.4	1.2	2.1	2.5	3.3	4.2	5.2
Male	10.2	114.7	1.9	0.4	0.4	1.5	2.9	3.2	4.5	5.4	6.8
Female	10.7	84.5	1.4	0.4	0.3	0.8	1.3	1.8	2.2	3.1	3.5
Non-Hispanic white	10.3	76.6	1.4	0.3	0.3	1.0	1.8	2.1	3.0	3.8	4.4
Male	9.6	89.3	1.7	0.3	0.4	1.3	2.5	2.8	3.9	4.9	5.9
Female	10.9	63.2	1.0	0.4	*	0.7	1.0	1.4	2.1	2.6	3.0
Non-Hispanic black	13.6	217.2	3.2	*	*	1.9	3.8	4.6	5.8	8.2	10.7
Male	15.7	244.9	3.3	*	*	2.6	4.9	5.7	8.9	9.9	14.1
Female	11.8	188.6	3.1	*	*	*	2.8	3.6	3.1	6.7	7.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)											
Non-Hispanic 5/	67.9	30.2	4.3	2.3	2.5	4.1	5.6	6.8	9.2	14.4	21.7
Male	56.0	34.5	4.8	2.3	2.6	4.9	6.3	7.9	10.7	17.0	26.5
Female	79.1	25.7	3.9	2.3	2.4	3.2	4.9	5.8	7.9	11.9	17.0
Non-Hispanic white	71.9	23.4	3.9	2.1	2.4	3.9	4.9	6.2	8.1	12.1	18.7
Male	57.9	26.5	4.4	2.1	2.3	4.9	5.7	7.5	9.7	14.8	23.6
Female	85.4	20.2	3.5	2.1	2.4	2.8	4.1	4.8	6.5	9.4	13.8
Non-Hispanic black	60.9	62.8	6.2	3.4	3.5	5.0	9.4	11.4	16.7	30.5	44.2
Male	57.2	71.5	6.4	3.7	4.2	5.1	10.1	11.4	17.7	33.7	49.6
Female	64.1	53.8	6.1	3.0	2.7	4.9	8.8	11.4	15.9	27.7	39.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/	35.9	22.1	12.6	7.3	8.3	33.9	38.7	32.3	30.5	32.7	35.3
Male	47.7	25.3	14.6	8.8	10.8	45.6	59.0	50.3	46.5	48.2	52.9
Female	24.6	18.8	10.4	5.8	5.8	21.5	17.6	14.5	14.7	17.3	18.1
Hispanic	27.6	18.6	12.4	5.2	7.0	28.8	42.6	36.4	32.1	32.7	35.2
Male	41.7	21.7	14.1	6.5	8.6	42.1	69.3	59.7	51.4	52.1	56.2
Female	13.3	15.3	10.7	4.0	5.4	14.4	13.8	11.9	11.4	12.1	13.8
Mexican	27.8	20.3	12.9	5.3	7.1	30.6	46.2	36.6	33.7	33.5	34.9
Male	41.7	23.7	15.0	6.5	8.5	44.9	73.9	59.0	53.0	52.2	55.9
Female	13.0	16.7	10.7	4.0	5.5	14.6	14.6	12.2	12.2	11.9	13.1
Puerto Rican	27.2	*	*	*	*	23.1	26.8	32.1	33.4	40.2	49.7
Male	42.3	*	*	*	*	*	*	*	*	*	*
Female	13.2	*	*	*	*	*	*	*	*	*	*
Cuban	24.5	0.0	*	*	*	*	*	*	*	*	*
Male	34.8	0.0	*	*	*	*	*	*	*	*	*
Female	15.1	0.0	*	*	*	*	*	*	*	*	0.0
Other Hispanic 4/	27.9	*	11.9	5.3	7.9	26.5	38.3	39.9	30.0	29.9	33.5
Male	43.1	*	12.0	*	*	38.4	66.2	65.9	48.1	48.2	52.1
Female	13.8	*	*	*	*	13.6	12.0	13.9	10.8	12.6	15.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 5/	36.7	22.9	12.5	7.7	8.5	34.6	37.8	31.4	30.0	32.5	35.1
Male	48.1	26.1	14.6	9.2	11.1	46.0	56.8	48.4	45.3	47.4	52.1
Female	25.9	19.5	10.3	6.1	5.9	22.5	18.1	14.8	15.2	17.8	18.3
Non-Hispanic white	37.4	18.4	11.0	6.6	8.2	36.9	38.4	31.7	29.6	32.1	34.3
Male	48.1	21.3	12.7	7.7	10.6	48.1	57.4	48.5	44.3	46.6	50.7
Female	27.2	15.3	9.2	5.4	5.8	25.0	18.4	14.8	15.1	17.6	18.0
Non-Hispanic black	37.9	47.4	20.6	13.2	10.6	26.4	36.4	34.0	35.6	38.7	45.1
Male	55.1	52.1	24.7	16.9	14.3	38.7	56.5	54.8	56.5	58.6	71.6
Female	22.4	42.6	16.4	9.5	6.7	13.8	16.8	15.0	17.2	21.1	22.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Transport accidents (V01-V99,Y85)

All origins 3/	17.0	4.9	4.7	4.5	5.5	27.6	29.1	21.5	17.8	17.2	16.5
Male	23.7	5.2	5.2	5.2	6.7	35.4	43.2	32.6	26.2	24.4	23.9
Female	10.6	4.7	4.2	3.7	4.2	19.4	14.4	10.6	9.6	10.0	9.3
Hispanic	15.6	5.5	5.7	3.3	5.2	22.1	31.3	24.9	18.2	15.2	16.9
Male	23.0	6.0	6.1	3.9	6.1	31.3	49.0	39.6	27.8	23.1	25.9
Female	8.0	*	5.4	2.7	4.2	12.2	12.2	9.4	8.0	6.8	7.8
Mexican	16.8	7.2	6.1	3.4	5.1	23.6	35.0	25.5	20.5	16.4	18.4
Male	24.6	*	6.8	4.1	6.0	33.5	54.4	39.8	30.5	24.3	28.5
Female	8.4	*	5.3	2.7	4.1	12.5	12.9	10.0	9.3	7.2	7.9
Puerto Rican	10.7	*	*	*	*	17.4	17.4	15.6	12.1	11.4	*
Male	16.1	0.0	*	*	*	*	*	*	*	*	*
Female	5.8	*	*	*	*	*	*	*	*	*	*
Cuban	13.5	0.0	*	*	*	*	*	*	*	*	*
Male	20.1	0.0	*	0.0	*	*	*	*	*	*	*
Female	7.5	0.0	*	*	*	*	*	*	*	*	0.0
Other Hispanic 4/	14.5	*	5.2	*	6.1	19.8	25.0	28.0	15.6	13.6	15.3
Male	21.4	*	*	*	*	27.4	40.0	45.0	23.9	20.5	22.0
Female	8.1	*	*	*	*	11.6	10.9	11.1	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Transport accidents (V01-V99,Y85)											
Non-Hispanic 5/	17.1	4.8	4.5	4.7	5.6	28.4	28.5	20.8	17.6	17.3	16.4
Male	23.7	5.0	4.9	5.4	6.8	36.0	41.9	31.2	25.8	24.5	23.6
Female	10.9	4.6	4.0	4.0	4.2	20.4	14.7	10.7	9.8	10.3	9.5
Non-Hispanic white	17.2	4.1	3.7	4.4	5.7	30.8	29.0	20.8	17.5	16.9	16.3
Male	23.6	4.8	3.9	5.0	6.9	38.4	42.5	30.8	25.0	23.7	23.2
Female	11.1	3.5	3.4	3.7	4.4	22.8	14.8	10.7	10.0	10.2	9.4
Non-Hispanic black	17.6	8.3	8.1	6.7	5.5	19.8	27.0	23.0	20.1	21.2	18.8
Male	26.1	*	9.1	8.3	7.1	27.2	40.3	36.9	32.3	31.8	29.5
Female	9.8	10.0	7.0	5.1	3.9	12.0	14.1	10.3	9.3	11.9	9.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)											
All origins 3/	15.5	4.8	4.3	4.0	5.0	26.3	27.3	19.8	16.1	15.3	14.7
Male	21.4	5.0	4.7	4.6	5.9	33.6	40.5	29.7	23.6	21.6	20.9
Female	9.9	4.7	3.8	3.4	3.9	18.6	13.7	10.0	8.7	9.1	8.6
Hispanic	14.4	5.5	5.4	3.1	4.8	21.1	29.6	23.1	16.6	13.7	15.2
Male	21.2	6.0	5.8	3.6	5.5	29.8	46.3	36.4	25.3	20.7	23.3
Female	7.6	*	4.9	2.6	4.0	11.7	11.7	9.0	7.3	6.3	7.0
Mexican	15.6	7.2	5.8	3.2	4.8	22.7	33.0	23.9	18.7	15.1	16.7
Male	22.8	*	6.6	3.7	5.6	32.1	51.1	36.8	27.9	22.3	26.1
Female	8.0	*	5.0	2.6	4.0	12.2	12.3	9.9	8.5	6.8	7.0
Puerto Rican	9.8	*	*	*	*	16.4	16.9	*	*	*	*
Male	14.4	0.0	*	*	*	*	*	*	*	*	*
Female	5.5	*	*	*	*	*	*	*	*	*	*
Cuban	11.8	0.0	*	*	*	*	*	*	*	*	*
Male	17.0	0.0	*	0.0	*	*	*	*	*	*	*
Female	7.0	0.0	*	*	*	*	*	*	*	*	0.0
Other Hispanic 4/	13.3	*	*	*	5.3	18.2	23.9	25.2	14.6	11.8	14.0
Male	19.5	*	*	*	*	25.5	38.5	40.9	22.3	17.6	19.7
Female	7.6	*	*	*	*	*	10.1	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)											
Non-Hispanic 5/	15.6	4.6	4.0	4.2	5.0	27.1	26.8	19.1	15.9	15.5	14.6
Male	21.3	4.7	4.5	4.8	6.0	34.1	39.2	28.4	23.1	21.6	20.6
Female	10.2	4.6	3.6	3.6	3.9	19.6	14.0	10.0	9.0	9.5	8.8
Non-Hispanic white	15.7	4.0	3.3	3.9	5.1	29.4	27.2	18.9	15.7	15.0	14.4
Male	21.2	4.5	3.6	4.3	6.0	36.5	39.7	27.8	22.4	20.7	20.1
Female	10.4	3.4	3.0	3.3	4.0	21.9	14.1	10.0	9.1	9.3	8.6
Non-Hispanic black	16.3	8.1	7.3	6.1	5.1	18.8	26.0	21.8	18.5	19.4	17.2
Male	24.0	*	8.0	7.3	6.3	25.8	38.8	34.9	29.6	28.8	26.6
Female	9.3	10.0	6.6	5.0	3.8	11.6	13.5	9.9	8.7	11.2	9.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins 3/	0.8	*	0.3	0.3	0.3	0.9	1.1	0.8	0.8	0.9	0.9
Male	1.2	*	0.3	0.4	0.5	1.2	1.7	1.4	1.2	1.4	1.5
Female	0.4	0.0	0.3	0.2	*	0.6	0.3	0.3	0.5	0.4	0.4
Hispanic	0.8	0.0	*	*	*	0.7	1.3	1.1	1.2	1.1	1.3
Male	1.4	0.0	*	*	*	*	2.1	2.1	1.8	1.8	2.1
Female	0.3	0.0	*	*	*	*	*	*	*	*	*
Mexican	0.9	0.0	*	*	*	*	1.6	1.1	1.3	*	1.6
Male	1.4	0.0	*	*	*	*	2.5	2.0	*	*	*
Female	0.3	0.0	*	0.0	*	*	*	0.0	*	*	*
Puerto Rican	*	0.0	0.0	*	*	*	*	*	*	*	*
Male	*	0.0	0.0	0.0	*	*	*	*	*	*	*
Female	*	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Other Hispanic 4/	0.8	0.0	*	*	*	*	*	*	*	*	*
Male	1.3	0.0	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	0.0	0.0	*	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)											
Non-Hispanic 5/	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9	0.9
Male	1.1	*	0.4	0.4	0.5	1.2	1.6	1.2	1.1	1.3	1.4
Female	0.4	0.0	*	*	*	0.6	0.3	0.3	0.5	0.4	0.3
Non-Hispanic white	0.8	*	0.3	0.3	0.4	0.9	1.0	0.8	0.7	0.9	0.9
Male	1.1	*	*	0.4	0.5	1.2	1.6	1.2	1.0	1.3	1.4
Female	0.4	0.0	*	*	*	0.6	0.4	0.3	0.5	0.4	0.4
Non-Hispanic black	0.7	*	*	*	*	*	*	*	1.0	1.1	0.9
Male	1.2	*	*	*	*	*	*	*	*	1.7	1.6
Female	0.3	0.0	*	*	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All origins 3/	0.7	*	*	0.2	0.2	0.4	0.6	0.9	0.9	0.9	0.9
Male	1.1	0.0	*	0.2	0.3	0.6	1.0	1.5	1.4	1.4	1.5
Female	0.3	*	*	*	*	0.2	0.3	0.3	0.4	0.4	0.4
Hispanic	0.3	0.0	*	*	*	*	*	*	*	*	*
Male	0.5	0.0	0.0	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*	*	*
Mexican	0.3	0.0	*	*	*	*	*	*	*	*	*
Male	0.4	0.0	0.0	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	0.0	*	0.0	*	*	*	*
Puerto Rican	*	0.0	0.0	0.0	*	*	0.0	*	*	*	*
Male	*	0.0	0.0	0.0	*	*	0.0	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.4	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	*	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	*	0.0	0.0	0.0	*	*	*	*	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Nontransport accidents (W00-X59,Y86)											
Non-Hispanic 5/	19.6	18.1	8.1	3.0	3.0	6.2	9.3	10.6	12.4	15.1	18.6
Male	24.4	21.1	9.7	3.8	4.3	10.0	14.9	17.2	19.6	22.9	28.6
Female	15.1	14.9	6.3	2.2	1.6	2.1	3.4	4.1	5.4	7.5	8.9
Non-Hispanic white	20.2	14.2	7.3	2.2	2.6	6.1	9.4	10.9	12.1	15.2	18.0
Male	24.4	16.5	8.7	2.8	3.7	9.7	15.0	17.7	19.2	22.9	27.5
Female	16.1	11.9	5.8	1.7	1.4	2.2	3.6	4.1	5.1	7.4	8.5
Non-Hispanic black	20.3	39.1	12.5	6.5	5.1	6.7	9.4	11.0	15.5	17.5	26.4
Male	28.9	45.4	15.6	8.7	7.3	11.4	16.2	17.9	24.2	26.8	42.1
Female	12.6	32.7	9.4	4.3	2.8	1.8	2.7	4.7	7.9	9.2	12.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Falls (W00-W19)											
All origins 3/	4.8	*	0.4	0.1	0.1	0.6	0.7	0.9	0.9	1.2	1.7
Male	5.3	*	0.5	0.2	0.2	0.9	1.2	1.5	1.6	2.0	2.7
Female	4.3	*	*	*	*	*	*	0.2	0.3	0.5	0.6
Hispanic	2.1	*	*	*	*	*	1.2	1.4	1.2	1.3	1.7
Male	2.9	*	*	*	*	*	2.2	2.6	2.3	2.3	2.9
Female	1.2	0.0	*	0.0	0.0	*	*	*	*	*	*
Mexican	1.8	*	*	0.0	*	*	1.3	1.4	*	*	1.9
Male	2.6	*	*	0.0	*	*	2.3	2.6	*	*	3.4
Female	0.9	0.0	*	0.0	0.0	0.0	*	*	*	*	*
Puerto Rican	2.1	0.0	0.0	*	0.0	*	0.0	*	*	*	*
Male	2.8	0.0	0.0	*	0.0	*	0.0	*	*	*	*
Female	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	3.9	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Male	4.9	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	2.7	0.0	*	0.0	*	*	*	*	*	*	*
Male	3.9	0.0	*	0.0	*	*	*	*	*	*	*
Female	1.6	0.0	0.0	0.0	0.0	*	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Accidental discharge of firearms (W32-W34)

All origins 3/	0.3	0.0	*	0.1	0.3	0.6	0.7	0.4	0.3	0.3	0.3
Male	0.5	0.0	*	*	0.5	1.1	1.2	0.7	0.6	0.5	0.5
Female	0.1	0.0	*	*	*	*	*	*	*	*	*
Hispanic	0.2	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	0.4	0.0	0.0	0.0	*	*	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	*	*	0.0	*	0.0
Mexican	0.2	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	0.5	0.0	0.0	0.0	*	*	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	0.0	0.0	0.0	0.0	*	*	0.0	0.0	*	*
Male	*	0.0	0.0	0.0	0.0	*	*	0.0	0.0	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental discharge of firearms (W32-W34)											
Non-Hispanic 5/	0.3	0.0	*	0.1	0.3	0.7	0.7	0.4	0.3	0.3	0.3
Male	0.5	0.0	*	*	0.5	1.2	1.3	0.7	0.6	0.5	0.5
Female	0.1	0.0	*	*	*	*	*	*	*	*	*
Non-Hispanic white	0.3	0.0	*	*	0.3	0.6	0.6	0.5	0.3	0.3	0.3
Male	0.5	0.0	*	*	0.6	1.0	1.1	0.7	0.5	0.5	0.5
Female	0.1	0.0	*	*	*	*	*	*	*	*	*
Non-Hispanic black	0.5	0.0	*	*	*	1.2	1.2	*	*	*	*
Male	0.9	0.0	*	*	*	2.2	2.2	*	*	*	*
Female	0.1	0.0	*	*	*	*	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Accidental drowning and submersion (W65-W74)

All origins 3/	1.3	1.8	3.2	1.0	0.9	1.8	1.6	1.2	1.2	1.1	1.0
Male	2.1	2.0	4.0	1.4	1.4	3.3	2.9	2.1	2.1	1.9	1.7
Female	0.5	1.6	2.4	0.5	0.4	0.3	0.3	0.2	0.3	0.3	0.3
Hispanic	1.5	*	2.8	0.8	0.8	2.5	2.8	1.5	1.5	1.0	*
Male	2.6	*	3.1	*	*	4.6	5.2	2.8	2.8	1.8	*
Female	0.5	*	2.6	*	*	*	*	*	*	*	*
Mexican	1.5	*	2.7	*	*	2.8	2.6	1.6	1.4	1.4	*
Male	2.5	*	2.6	*	*	5.1	4.7	3.0	2.6	*	*
Female	0.5	*	2.8	*	*	*	*	*	*	*	*
Puerto Rican	1.1	*	*	*	*	*	*	*	*	*	*
Male	2.1	*	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Cuban	*	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*	0.0
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*	0.0
Female	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	1.9	*	*	*	*	*	4.9	*	*	*	*
Male	3.3	*	*	*	*	*	*	*	*	*	*
Female	*	*	*	0.0	0.0	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/	1.2	1.1	2.0	0.9	0.5	0.4	0.6	0.8	0.7	0.9	1.0
Male	1.5	1.1	2.2	0.9	0.5	0.6	0.8	1.0	1.1	1.2	1.5
Female	0.9	1.1	1.7	0.8	0.4	0.3	0.4	0.5	0.4	0.5	0.5
Hispanic	0.6	*	1.5	*	*	*	*	*	*	*	*
Male	0.8	*	1.6	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*	*	*
Mexican	0.6	*	1.7	*	*	*	*	*	*	*	*
Male	0.9	*	2.0	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*	*	*
Puerto Rican	0.6	0.0	*	*	*	*	*	0.0	*	0.0	*
Male	*	0.0	0.0	*	0.0	0.0	*	0.0	*	0.0	*
Female	*	0.0	*	*	*	*	0.0	0.0	*	0.0	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.6	0.0	*	*	*	0.0	*	*	*	*	*
Male	0.8	0.0	*	0.0	0.0	0.0	*	*	*	*	*
Female	*	0.0	*	*	*	0.0	0.0	*	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental exposure to smoke, fire and flames (X00-X09)											
Non-Hispanic 5/	1.3	1.2	2.1	0.9	0.5	0.4	0.7	0.9	0.8	0.9	1.0
Male	1.6	*	2.4	1.0	0.6	0.6	0.8	1.2	1.1	1.3	1.5
Female	1.0	*	1.8	0.8	0.4	0.3	0.5	0.6	0.4	0.6	0.6
Non-Hispanic white	1.1	0.9	1.4	0.6	0.4	0.4	0.6	0.8	0.7	0.8	1.0
Male	1.4	*	1.6	0.6	0.5	0.5	0.8	1.2	1.1	1.1	1.4
Female	0.8	*	1.2	0.6	0.4	*	0.4	0.5	0.3	0.5	0.5
Non-Hispanic black	2.7	*	5.3	2.5	1.0	*	0.9	1.2	0.9	2.0	1.8
Male	3.2	*	6.0	2.9	*	*	*	*	*	3.0	2.5
Female	2.2	*	4.6	2.1	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/	4.5	*	0.2	*	0.1	1.3	3.9	5.2	7.1	9.2	11.1
Male	6.7	*	*	*	0.2	1.9	6.1	8.4	11.0	13.6	16.5
Female	2.4	*	*	*	*	0.7	1.6	2.1	3.3	4.9	5.7
Hispanic	4.8	*	*	*	*	1.5	3.9	6.0	8.6	12.0	12.7
Male	8.1	*	*	*	*	1.9	6.8	10.1	14.5	19.7	21.1
Female	1.5	0.0	0.0	*	*	*	*	1.8	2.3	3.9	4.0
Mexican	4.2	*	*	*	*	1.4	3.6	5.2	8.0	11.1	10.6
Male	6.9	*	*	0.0	*	1.9	6.0	8.6	13.7	17.9	17.6
Female	1.2	0.0	0.0	*	*	*	*	*	*	3.3	3.3
Puerto Rican	9.4	0.0	0.0	0.0	0.0	*	*	*	17.5	22.8	29.6
Male	16.1	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	3.3	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Cuban	2.1	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*	0.0
Other Hispanic 4/	5.2	*	0.0	*	0.0	*	4.1	6.5	7.7	11.7	12.6
Male	9.3	*	0.0	*	0.0	*	*	11.0	12.7	19.5	21.6
Female	1.4	0.0	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental poisoning and exposure to noxious substances (X40-X49)											
Non-Hispanic 5/	4.4	*	0.3	*	0.1	1.3	3.9	5.1	6.8	8.8	10.7
Male	6.4	*	*	*	*	1.9	5.9	8.0	10.3	12.6	15.8
Female	2.4	*	*	*	*	0.6	1.8	2.2	3.5	5.0	5.8
Non-Hispanic white	4.2	*	*	*	*	1.4	4.2	5.5	6.8	8.8	10.2
Male	6.2	*	*	*	*	2.1	6.4	8.6	10.3	12.6	15.1
Female	2.4	*	*	*	*	0.7	1.9	2.3	3.3	4.9	5.3
Non-Hispanic black	6.1	*	*	*	*	*	2.9	4.5	8.6	10.7	16.6
Male	9.3	*	*	*	*	*	4.8	7.2	12.4	15.5	24.9
Female	3.2	0.0	*	*	0.0	*	*	2.1	5.2	6.6	9.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/	6.7	13.7	1.9	0.7	0.9	1.5	2.1	2.3	2.4	2.8	3.7
Male	7.8	16.1	2.4	0.9	1.3	2.3	3.6	4.0	4.0	4.6	6.1
Female	5.8	11.2	1.5	0.5	0.5	0.6	0.5	0.7	0.8	1.0	1.4
Hispanic	2.8	9.0	1.9	*	*	1.3	2.3	2.1	1.8	2.6	2.4
Male	3.9	9.5	2.6	*	*	1.9	4.3	3.9	2.8	4.5	3.7
Female	1.7	8.5	*	*	*	*	*	*	*	*	*
Mexican	2.7	9.0	1.9	*	*	1.4	2.6	2.3	1.8	3.1	2.8
Male	3.8	*	2.9	*	*	2.1	4.5	4.1	2.7	5.2	4.4
Female	1.5	*	*	*	*	*	*	*	*	*	*
Puerto Rican	3.0	*	*	*	*	*	*	*	*	*	*
Male	4.3	*	*	0.0	*	*	*	*	*	*	*
Female	1.8	*	*	*	0.0	*	0.0	0.0	0.0	*	*
Cuban	3.6	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
Male	4.8	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	2.9	*	*	*	*	*	*	*	*	*	*
Male	4.1	*	*	*	*	*	*	*	*	*	*
Female	1.8	*	*	*	*	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Intentional self-harm (suicide) (X60-X84,Y87.0)

All origins 3/	10.7	0.0	0.0	*	1.2	8.2	12.7	13.2	13.7	14.6	14.3
Male	17.6	0.0	0.0	*	1.9	13.3	21.6	22.1	22.4	23.1	22.0
Female	4.1	0.0	0.0	0.0	0.5	2.8	3.5	4.3	5.2	6.1	6.7
Hispanic	5.4	0.0	0.0	0.0	1.1	6.3	8.0	8.6	7.6	8.8	6.5
Male	9.1	0.0	0.0	0.0	1.5	10.1	13.8	14.2	12.9	14.2	10.9
Female	1.7	0.0	0.0	0.0	*	2.2	1.9	2.6	2.0	3.2	2.1
Mexican	4.5	0.0	0.0	0.0	1.2	5.5	7.1	7.7	7.1	8.1	5.7
Male	7.6	0.0	0.0	0.0	*	9.2	12.1	13.0	11.9	13.0	9.2
Female	1.2	0.0	0.0	0.0	*	*	*	*	*	*	*
Puerto Rican	6.7	0.0	0.0	0.0	*	*	*	*	*	12.3	*
Male	11.5	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	2.2	0.0	0.0	0.0	*	*	*	*	*	*	*
Cuban	10.6	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	18.2	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	3.4	0.0	0.0	0.0	0.0	*	*	*	0.0	*	*
Other Hispanic 4/	6.6	0.0	0.0	0.0	*	10.5	10.8	9.6	8.8	9.0	7.1
Male	11.0	0.0	0.0	0.0	*	16.0	19.2	14.3	14.3	14.2	13.1
Female	2.6	0.0	0.0	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Intentional self-harm (suicide) (X60-X84,Y87.0)											
Non-Hispanic 5/	11.3	0.0	0.0	*	1.3	8.5	13.5	13.9	14.6	15.3	15.0
Male	18.7	0.0	0.0	*	2.0	13.8	22.9	23.4	23.9	24.2	23.1
Female	4.4	0.0	0.0	0.0	0.5	2.9	3.7	4.5	5.7	6.5	7.1
Non-Hispanic white	12.5	0.0	0.0	*	1.3	8.9	14.0	14.6	15.9	16.7	16.8
Male	20.4	0.0	0.0	*	2.1	14.5	23.6	24.3	25.7	25.9	25.6
Female	4.9	0.0	0.0	0.0	0.5	3.0	3.8	4.9	6.3	7.5	8.0
Non-Hispanic black	5.8	0.0	0.0	0.0	1.0	6.1	11.2	10.4	9.5	8.9	7.1
Male	10.4	0.0	0.0	0.0	1.5	10.3	20.3	19.4	17.2	16.7	11.3
Female	1.7	0.0	0.0	0.0	*	1.7	2.4	2.2	2.7	2.0	3.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/	6.1	0.0	0.0	0.0	0.5	4.9	7.4	7.1	6.7	7.2	6.6
Male	10.9	0.0	0.0	0.0	0.8	8.5	13.2	12.6	11.6	12.0	11.2
Female	1.5	0.0	0.0	0.0	0.2	1.1	1.4	1.7	2.0	2.4	2.2
Hispanic	2.5	0.0	0.0	0.0	*	3.4	4.6	4.0	2.9	3.5	2.6
Male	4.5	0.0	0.0	0.0	*	5.6	8.2	7.0	5.2	5.7	4.7
Female	0.6	0.0	0.0	0.0	*	*	*	*	*	*	*
Mexican	2.3	0.0	0.0	0.0	*	3.1	4.0	3.9	3.0	3.4	2.5
Male	4.0	0.0	0.0	0.0	*	5.3	7.1	6.9	5.4	5.5	4.5
Female	0.4	0.0	0.0	0.0	*	*	*	*	*	*	*
Puerto Rican	2.2	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	4.1	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	...	0.0	0.0	0.0	0.0	*	*	*	*	*	0.0
Cuban	4.8	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	8.3	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	...	0.0	0.0	0.0	0.0	*	0.0	*	0.0	*	*
Other Hispanic 4/	3.0	0.0	0.0	0.0	*	5.3	7.1	*	*	3.8	*
Male	5.3	0.0	0.0	0.0	*	*	12.7	*	*	*	*
Female	0.9	0.0	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Intentional self-harm (suicide) by discharge of firearms (X72-X74)											
Non-Hispanic 5/	6.5	0.0	0.0	0.0	0.5	5.2	7.9	7.6	7.3	7.6	7.0
Male	11.7	0.0	0.0	0.0	0.8	9.0	14.1	13.5	12.6	12.8	11.9
Female	1.6	0.0	0.0	0.0	*	1.1	1.5	1.8	2.2	2.5	2.3
Non-Hispanic white	7.3	0.0	0.0	0.0	0.6	5.5	8.0	8.0	8.1	8.4	8.0
Male	13.0	0.0	0.0	0.0	1.0	9.6	14.3	14.0	13.7	13.8	13.3
Female	1.9	0.0	0.0	0.0	*	1.3	1.5	2.0	2.5	2.9	2.7
Non-Hispanic black	3.4	0.0	0.0	0.0	*	4.1	8.0	6.2	4.6	4.5	2.8
Male	6.4	0.0	0.0	0.0	*	7.3	14.8	12.0	8.9	8.7	5.4
Female	0.6	0.0	0.0	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All origins 3/	4.6	0.0	0.0	*	0.7	3.2	5.2	6.1	7.0	7.4	7.6
Male	6.7	0.0	0.0	*	1.1	4.7	8.3	9.6	10.8	11.1	10.8
Female	2.6	0.0	0.0	0.0	0.3	1.7	2.1	2.6	3.2	3.8	4.5
Hispanic	2.9	0.0	0.0	0.0	*	2.9	3.4	4.5	4.7	5.3	3.9
Male	4.6	0.0	0.0	0.0	*	4.5	5.5	7.2	7.6	8.4	6.2
Female	1.1	0.0	0.0	0.0	*	*	*	1.7	1.6	2.0	*
Mexican	2.2	0.0	0.0	0.0	*	2.3	3.0	3.7	4.0	4.7	3.2
Male	3.6	0.0	0.0	0.0	*	4.0	5.0	6.1	6.4	7.4	4.7
Female	0.8	0.0	0.0	0.0	*	*	*	*	*	*	*
Puerto Rican	4.5	0.0	0.0	0.0	*	*	*	*	*	*	*
Male	7.4	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	1.8	0.0	0.0	0.0	*	*	*	*	*	*	*
Cuban	5.8	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	9.9	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	...	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*	0.0
Other Hispanic 4/	3.6	0.0	0.0	0.0	*	5.1	*	5.7	6.4	5.2	*
Male	5.7	0.0	0.0	0.0	*	*	*	*	10.2	8.2	*
Female	1.7	0.0	0.0	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

Non-Hispanic 5/	4.8	0.0	0.0	*	0.7	3.3	5.6	6.3	7.3	7.7	8.0
Male	7.0	0.0	0.0	*	1.1	4.8	8.8	9.9	11.3	11.4	11.2
Female	2.8	0.0	0.0	0.0	0.3	1.7	2.2	2.7	3.5	4.0	4.8
Non-Hispanic white	5.2	0.0	0.0	*	0.7	3.4	5.9	6.6	7.8	8.3	8.8
Male	7.5	0.0	0.0	*	1.1	4.9	9.3	10.3	12.0	12.1	12.4
Female	3.0	0.0	0.0	0.0	*	1.7	2.3	2.9	3.7	4.5	5.2
Non-Hispanic black	2.5	0.0	0.0	0.0	0.8	2.0	3.2	4.2	4.9	4.4	4.3
Male	4.0	0.0	0.0	0.0	*	3.1	5.5	7.4	8.3	8.0	5.9
Female	1.1	0.0	0.0	0.0	*	*	*	*	1.9	*	2.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (X85-Y09,Y87.1)

Non-Hispanic 5/	5.7	8.7	2.6	0.9	1.1	9.4	14.7	12.0	9.0	7.4	6.2
Male	8.7	10.0	2.7	0.9	1.3	15.1	24.0	19.2	13.6	10.5	8.9
Female	2.9	7.4	2.6	0.9	0.9	3.4	5.1	4.9	4.5	4.4	3.6
Non-Hispanic white	2.9	6.0	1.6	0.6	0.8	3.1	4.6	4.4	4.5	4.3	3.8
Male	3.9	7.8	1.7	0.6	0.8	4.2	5.9	5.9	5.8	5.6	5.2
Female	2.0	4.2	1.6	0.6	0.7	2.0	3.3	2.8	3.1	3.0	2.5
Non-Hispanic black	22.8	21.6	7.3	2.2	2.7	38.6	64.0	51.0	34.1	26.6	22.0
Male	39.1	21.6	7.0	2.3	3.7	66.0	115.9	90.5	59.5	42.6	35.4
Female	8.1	21.5	7.5	2.1	1.7	10.4	13.2	14.8	11.8	12.5	10.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (X93-X95)

Non-Hispanic 5/	3.7	*	0.4	0.3	0.7	7.6	11.9	9.2	6.1	4.5	3.5
Male	6.1	*	0.4	0.4	1.0	12.8	20.7	16.0	10.1	6.8	5.2
Female	1.3	*	0.4	0.2	0.5	2.1	2.8	2.6	2.1	2.3	1.8
Non-Hispanic white	1.5	*	0.2	0.2	0.4	2.0	2.8	2.8	2.7	2.5	2.1
Male	2.2	*	*	*	0.5	2.9	4.0	4.2	3.9	3.4	2.9
Female	0.9	0.0	*	*	0.4	0.9	1.5	1.4	1.4	1.6	1.3
Non-Hispanic black	16.6	*	*	0.7	2.0	33.9	56.6	42.5	25.5	16.8	12.7
Male	30.8	*	*	*	3.1	59.5	105.6	80.1	47.9	29.1	21.5
Female	3.9	*	*	*	*	7.4	8.7	8.2	5.7	6.1	5.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All origins 3/	2.2	8.5	2.2	0.6	0.4	1.9	3.2	3.1	3.1	3.1	2.9
Male	2.9	9.3	2.2	0.6	0.4	2.5	4.1	3.9	3.9	4.0	4.0
Female	1.6	7.5	2.1	0.7	0.5	1.4	2.3	2.3	2.3	2.2	1.8
Hispanic	3.0	7.1	1.7	*	0.7	2.6	5.2	5.4	4.2	3.8	3.5
Male	4.4	6.5	1.6	*	*	3.6	8.2	8.3	6.5	5.4	5.8
Female	1.5	7.6	1.9	*	*	1.5	1.9	2.3	1.8	2.2	*
Mexican	2.9	7.8	1.6	*	*	2.6	5.0	5.5	4.5	3.1	2.7
Male	4.3	*	*	*	*	3.7	7.7	8.8	6.9	5.0	4.4
Female	1.4	*	*	*	*	*	*	*	*	*	*
Puerto Rican	3.5	*	*	*	*	*	*	*	*	*	*
Male	4.4	*	*	0.0	*	*	*	*	*	*	*
Female	2.6	*	*	*	*	*	*	*	*	*	*
Cuban	2.0	*	*	*	0.0	0.0	0.0	*	*	*	*
Male	3.6	*	0.0	*	0.0	0.0	0.0	*	*	*	*
Female	*	0.0	*	0.0	0.0	0.0	0.0	*	*	*	0.0
Other Hispanic 4/	3.2	*	*	*	*	*	6.5	4.7	*	4.6	5.2
Male	4.8	*	*	*	*	*	10.4	*	*	*	*
Female	1.6	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

Non-Hispanic 5/	2.1	8.5	2.2	0.6	0.4	1.8	2.8	2.7	2.9	3.0	2.8
Male	2.6	9.8	2.3	0.6	0.3	2.3	3.3	3.2	3.4	3.8	3.8
Female	1.6	7.2	2.2	0.7	0.5	1.4	2.3	2.3	2.4	2.2	1.8
Non-Hispanic white	1.4	6.0	1.4	0.4	0.3	1.2	1.8	1.6	1.8	1.8	1.7
Male	1.7	7.6	1.5	0.4	*	1.3	1.9	1.8	1.9	2.2	2.3
Female	1.1	4.2	1.4	0.5	0.4	1.1	1.8	1.4	1.7	1.4	1.2
Non-Hispanic black	6.1	21.2	6.4	1.5	0.7	4.8	7.4	8.4	8.6	9.7	9.2
Male	8.3	21.2	6.4	1.3	*	6.5	10.3	10.4	11.5	13.5	13.9
Female	4.2	21.1	6.5	1.6	*	3.0	4.5	6.7	6.1	6.4	5.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS	
Legal intervention (Y35,Y89.0)												
Non-Hispanic 5/	0.1	0.0	0.0	0.0	*	*	0.3	0.4	0.3	0.3	0.2	
Male	0.3	0.0	0.0	0.0	*	*	0.5	0.7	0.7	0.6	0.4	
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
Non-Hispanic white	0.1	0.0	0.0	0.0	0.0	*	*	0.2	0.3	0.2	0.2	
Male	0.2	0.0	0.0	0.0	0.0	*	*	0.5	0.5	0.4	0.3	
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	
Non-Hispanic black	0.4	0.0	0.0	0.0	*	*	0.8	1.0	*	0.8	*	
Male	0.8	0.0	0.0	0.0	*	*	1.7	2.0	*	1.6	*	
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/	1.4	1.4	0.3	*	0.2	0.8	1.5	1.7	2.2	2.9	2.9
Male	2.0	1.7	0.3	*	0.3	1.3	2.3	2.5	3.0	4.0	4.0
Female	0.9	1.2	0.3	*	*	0.3	0.7	0.9	1.4	1.7	1.9
Hispanic	0.8	*	*	*	*	0.7	1.2	1.2	1.4	1.4	1.2
Male	1.3	*	*	*	*	*	2.1	2.0	2.0	2.2	1.8
Female	0.3	*	*	0.0	*	*	*	*	*	*	*
Mexican	0.6	*	*	*	*	*	1.3	*	*	*	*
Male	1.0	*	*	*	*	*	2.1	*	*	*	*
Female	0.3	*	*	0.0	*	*	*	*	*	*	*
Puerto Rican	1.7	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	2.7	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Other Hispanic 4/	1.0	*	*	0.0	*	*	*	*	*	*	*
Male	1.7	0.0	*	0.0	*	*	*	*	*	*	*
Female	*	*	*	0.0	0.0	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	1.5	1.6	0.3	*	0.2	0.8	1.5	1.8	2.3	3.0	3.1
Male	2.1	1.9	*	*	0.3	1.3	2.3	2.6	3.1	4.2	4.2
Female	0.9	1.3	0.3	*	*	0.3	0.7	1.0	1.5	1.9	2.0
Non-Hispanic white	1.5	1.3	*	*	0.2	0.8	1.5	1.8	2.3	3.0	2.9
Male	2.0	1.6	*	*	0.4	1.3	2.3	2.5	3.1	3.9	4.0
Female	1.0	*	*	*	*	*	0.7	1.1	1.6	2.0	1.9
Non-Hispanic black	2.1	*	*	*	*	0.9	1.8	2.1	2.7	4.1	4.8
Male	3.3	*	*	*	*	1.4	2.8	3.7	4.0	6.7	6.9
Female	1.0	*	*	*	*	*	*	*	1.6	1.8	3.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Discharge of firearms, undetermined intent (Y22-Y24)											
Non-Hispanic 5/	0.1	0.0	*	0.0	*	0.3	0.3	0.2	*	0.1	0.1
Male	0.2	0.0	*	0.0	*	0.6	0.5	0.3	*	0.2	0.2
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Non-Hispanic white	0.1	0.0	*	0.0	*	0.3	0.3	0.2	*	0.1	0.1
Male	0.2	0.0	*	0.0	*	0.5	0.5	*	*	0.2	0.2
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Non-Hispanic black	0.2	0.0	*	0.0	*	0.7	*	*	*	*	*
Male	0.4	0.0	*	0.0	*	*	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	*	*	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/	1.3	1.4	0.3	*	0.1	0.5	1.2	1.5	2.1	2.7	2.8
Male	1.8	1.7	*	*	0.2	0.7	1.8	2.2	2.8	3.8	3.8
Female	0.8	1.2	0.3	*	*	0.2	0.6	0.9	1.3	1.7	1.8
Hispanic	0.7	*	*	*	*	*	0.9	1.2	1.4	1.3	1.1
Male	1.1	*	*	*	*	*	1.5	1.9	2.0	2.0	1.8
Female	0.3	*	*	0.0	*	*	*	*	*	*	*
Mexican	0.5	*	*	*	*	*	*	*	*	*	*
Male	0.8	*	*	*	*	*	*	*	*	*	*
Female	0.2	*	*	0.0	*	0.0	*	*	*	*	*
Puerto Rican	1.7	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Male	2.7	0.0	0.0	0.0	0.0	*	*	*	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Other Hispanic 4/	0.8	*	*	0.0	0.0	*	*	*	*	*	*
Male	1.4	0.0	*	0.0	0.0	*	*	*	*	*	*
Female	*	*	*	0.0	0.0	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	1.4	1.6	0.3	*	0.1	0.5	1.2	1.6	2.2	2.9	3.0
Male	1.9	1.9	*	*	0.2	0.7	1.8	2.2	2.9	4.0	4.0
Female	0.9	1.3	0.3	*	*	*	0.6	0.9	1.4	1.9	1.9
Non-Hispanic white	1.3	1.3	*	*	0.1	0.6	1.3	1.6	2.2	2.8	2.8
Male	1.8	1.6	*	*	*	0.9	1.9	2.2	2.9	3.7	3.7
Female	0.9	*	*	*	*	*	0.6	1.1	1.5	1.9	1.8
Non-Hispanic black	1.9	*	*	*	*	*	1.2	1.7	2.5	4.0	4.7
Male	2.9	*	*	*	*	*	1.8	3.0	3.7	6.5	6.8
Female	0.9	*	*	*	*	*	*	*	1.5	1.8	2.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Complications of medical and surgical care (Y40-Y84,Y88)

All origins 3/	1.0	1.4	0.2	*	*	*	0.1	0.2	0.2	0.3	0.4
Male	1.0	1.6	*	*	*	*	*	*	0.3	0.3	0.4
Female	1.1	1.2	*	*	*	*	*	0.2	*	0.3	0.4
Hispanic	0.3	*	*	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	0.0	*	*	*
Female	0.3	*	*	*	0.0	*	*	*	*	*	0.0
Mexican	0.2	*	*	*	0.0	*	*	*	*	*	*
Male	0.2	*	*	*	0.0	*	0.0	0.0	*	*	*
Female	0.2	*	*	*	0.0	*	*	*	*	*	0.0
Puerto Rican	*	0.0	*	0.0	*	0.0	0.0	0.0	*	*	0.0
Male	*	0.0	*	0.0	*	0.0	0.0	0.0	*	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Male	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.5	0.0	*	0.0	0.0	0.0	0.0	0.0	*	*	*
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Female	0.6	0.0	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 5/	1.1	1.6	0.2	*	*	*	*	0.2	0.3	0.3	0.4
Male	1.1	1.8	*	*	*	*	*	*	0.4	0.3	0.4
Female	1.2	1.4	*	*	*	*	*	*	*	0.3	0.4
Non-Hispanic white	1.1	1.3	*	*	*	*	*	*	0.2	0.3	0.4
Male	1.1	1.6	*	*	*	*	*	*	0.4	0.3	0.3
Female	1.2	*	*	*	*	*	*	*	*	0.2	0.4
Non-Hispanic black	1.3	*	*	*	*	*	*	*	*	*	0.8
Male	1.2	*	*	*	*	*	*	*	*	*	*
Female	1.4	*	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

All causes

All origins 3/	351.9	516.0	814.6	1,275.4	1,976.2	3,031.6	4,616.3	7,478.2	15,476.1
Male	455.8	654.6	1,026.1	1,595.9	2,479.5	3,816.4	5,719.7	9,156.8	16,931.3
Female	251.8	384.7	619.2	988.3	1,549.1	2,414.2	3,826.7	6,464.2	14,861.1
Hispanic	287.6	406.4	589.4	913.8	1,444.8	2,115.3	3,154.8	4,838.4	9,175.2
Male	392.5	542.6	780.8	1,195.0	1,841.1	2,704.7	3,913.0	5,696.9	9,842.3
Female	185.2	280.0	420.9	676.9	1,125.8	1,662.1	2,591.2	4,300.8	8,838.0
Mexican	263.6	402.7	560.7	939.6	1,492.6	2,135.1	3,225.4	4,556.5	*
Male	347.1	507.0	697.8	1,145.8	1,801.6	2,638.4	*	*	*
Female	174.8	293.8	428.9	746.4	1,225.4	1,732.9	2,783.7	*	*
Puerto Rican	497.2	637.2	943.8	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	672.5	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	257.7	332.6	516.4	827.7	1,342.1	2,048.1	*	*	*
Male	367.0	466.1	*	*	*	*	*	*	*
Female	168.6	233.3	335.1	595.2	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All causes

Non-Hispanic 5/	355.9	522.2	828.4	1,297.5	2,006.1	3,075.7	4,675.7	7,573.7	15,731.3
Male	458.6	659.4	1,039.3	1,617.3	2,513.0	3,866.4	5,794.2	9,288.0	17,262.3
Female	256.9	392.1	632.8	1,009.7	1,574.4	2,452.8	3,876.8	6,539.8	15,090.0
Non-Hispanic white	312.7	471.6	764.4	1,225.8	1,942.0	2,997.8	4,602.0	7,546.5	15,945.8
Male	403.1	593.3	955.2	1,521.3	2,433.4	3,780.8	5,712.0	9,286.8	17,539.1
Female	223.4	353.5	583.3	953.0	1,515.0	2,372.3	3,802.6	6,492.2	15,284.6
Non-Hispanic black	706.9	1,004.9	1,470.9	2,060.8	2,802.3	4,227.5	6,031.7	8,702.4	15,015.7
Male	948.9	1,349.2	1,970.1	2,748.4	3,567.2	5,236.8	7,455.2	10,546.3	16,320.6
Female	503.9	726.1	1,084.2	1,550.5	2,231.3	3,516.5	5,123.2	7,714.3	14,474.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Salmonella infections (A01-A02)

Non-Hispanic 5/	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	0.0	0.0	*	0.0	0.0	*	*	*	*
Non-Hispanic white	0.0	*	*	0.0	0.0	*	*	*	*
Male	0.0	*	0.0	0.0	0.0	*	*	*	*
Female	0.0	0.0	*	0.0	0.0	*	*	*	*
Non-Hispanic black	0.0	0.0	0.0	0.0	*	0.0	*	*	0.0
Male	0.0	0.0	0.0	0.0	*	0.0	0.0	*	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Certain other intestinal infections (A04,A07-A09)

All origins 3/	*	*	*	0.3	0.7	1.2	2.3	4.6	10.3
Male	*	*	*	*	0.6	1.2	2.6	4.9	9.1
Female	*	*	*	*	0.8	1.3	2.1	4.4	10.8
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	0.0	*	*	*	*	0.0	*
Female	*	*	*	*	*	*	*	*	*
Mexican	0.0	*	*	*	*	*	*	0.0	*
Male	0.0	*	0.0	*	*	0.0	*	0.0	*
Female	0.0	0.0	*	0.0	*	*	*	0.0	*
Puerto Rican	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Cuban	*	0.0	0.0	*	0.0	0.0	0.0	*	*
Male	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	*	0.0	0.0	0.0	*	*
Other Hispanic 4/	*	*	*	*	*	0.0	*	0.0	*
Male	0.0	0.0	0.0	*	*	0.0	*	0.0	*
Female	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic 5/	*	*	*	0.3	0.7	1.3	2.4	4.8	10.5
Male	*	*	*	*	0.6	1.2	2.7	5.1	9.3
Female	*	*	*	*	0.8	1.3	2.1	4.6	11.0
Non-Hispanic white	*	*	*	0.3	0.8	1.3	2.4	5.0	11.2
Male	*	*	*	*	0.7	1.4	2.8	5.4	9.9
Female	*	*	*	*	0.8	1.3	2.2	4.7	11.7
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Tuberculosis (A16-A19)

All origins 3/	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.5	4.1
Male	0.4	0.6	0.8	1.0	1.1	1.7	2.3	3.6	6.9
Female	*	*	0.3	*	0.4	0.7	1.5	1.8	3.0
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	0.0	0.0	*	0.0	*	*	*	*	*
Male	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*
Female	0.0	0.0	*	0.0	0.0	*	*	*	*
Cuban	*	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	0.0	*	0.0	*	*	*	*
Female	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Respiratory tuberculosis (A16)

All origins 3/	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.2	3.7
Male	0.3	0.4	0.6	0.9	0.9	1.5	2.0	3.3	6.1
Female	*	*	*	*	*	0.5	1.2	1.5	2.6
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*
Puerto Rican	0.0	0.0	*	0.0	0.0	*	*	*	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	*	0.0	0.0	*	*	*	*
Cuban	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	*	0.0	*	*	*	*	*	*	*
Male	*	0.0	0.0	*	0.0	*	*	*	*
Female	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Respiratory tuberculosis (A16)

Non-Hispanic 5/	0.2	0.2	0.4	0.5	0.6	0.8	1.4	2.0	3.5
Male	0.3	0.4	0.6	0.8	0.9	1.3	1.8	3.2	5.9
Female	*	*	*	*	*	0.5	1.1	1.3	2.5
Non-Hispanic white	*	*	0.2	0.3	0.4	0.5	0.9	1.6	2.8
Male	*	*	*	*	0.6	0.8	1.1	2.5	4.7
Female	*	*	*	*	*	*	0.7	1.0	2.0
Non-Hispanic black	*	*	1.6	*	*	2.6	4.8	*	7.2
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other tuberculosis (A17-A19)

All origins 3/	*	*	*	*	*	0.3	0.3	*	0.5
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	0.0	*	*	*
Female	*	*	*	0.0	*	*	0.0	*	0.0
Mexican	*	*	*	*	*	*	*	*	0.0
Male	*	*	*	*	*	0.0	*	*	0.0
Female	0.0	*	*	0.0	*	*	0.0	*	0.0
Puerto Rican	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.0	*	*	0.0	0.0	0.0	0.0	0.0	*
Male	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Other tuberculosis (A17-A19)

Non-Hispanic 5/	*	*	*	*	*	0.2	0.3	*	0.5
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Non-Hispanic white	*	*	*	*	*	*	*	*	*
Male	*	*	*	0.0	*	*	*	*	*
Female	0.0	0.0	*	*	*	*	*	*	*
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	0.0	*
Female	*	*	0.0	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Scarlet fever and erysipelas (A38,A46)

Non-Hispanic 5/	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Non-Hispanic white	0.0	0.0	0.0	0.0	*	0.0	*	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	*	0.0	*	0.0	*
Non-Hispanic black	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Septicemia (A40-A41)

Non-Hispanic 5/	3.9	5.9	9.3	15.0	25.2	39.1	64.6	105.6	222.7
Male	4.4	6.4	10.0	16.4	28.2	44.7	73.9	121.7	228.6
Female	3.4	5.3	8.6	13.7	22.6	34.8	57.9	95.9	220.2
Non-Hispanic white	2.9	4.6	7.3	12.5	21.6	34.1	58.0	97.6	208.0
Male	3.2	5.0	7.7	13.0	23.9	38.9	67.2	112.3	214.0
Female	2.5	4.2	7.0	11.9	19.6	30.2	51.3	88.6	205.5
Non-Hispanic black	11.7	16.5	26.6	38.2	57.8	96.4	146.5	224.0	419.1
Male	13.9	19.8	32.1	50.7	71.2	112.9	162.8	271.5	424.0
Female	9.9	13.9	22.4	28.9	47.7	84.8	136.0	198.5	417.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Syphilis (A50-A53)

Non-Hispanic 5/	*	*	*	*	*	*	*	*	*
Male	*	0.0	*	*	*	*	0.0	0.0	*
Female	0.0	*	0.0	*	0.0	*	*	*	*
Non-Hispanic white	*	*	0.0	*	*	*	*	*	*
Male	*	0.0	0.0	*	*	0.0	0.0	0.0	0.0
Female	0.0	*	0.0	*	0.0	*	*	*	*
Non-Hispanic black	0.0	0.0	*	0.0	0.0	*	*	*	*
Male	0.0	0.0	*	0.0	0.0	*	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

Non-Hispanic 5/	4.3	3.5	3.1	3.0	4.1	4.7	5.0	5.2	4.7
Male	6.6	5.5	4.4	3.5	4.5	5.0	5.8	7.4	6.6
Female	2.2	1.6	1.8	2.5	3.6	4.6	4.3	3.9	4.0
Non-Hispanic white	3.9	2.9	2.5	2.5	3.4	4.2	4.6	5.1	4.5
Male	5.8	4.6	3.6	3.2	4.1	4.6	5.6	7.7	6.4
Female	2.1	1.3	1.4	1.9	2.9	3.8	3.8	3.4	3.8
Non-Hispanic black	7.5	7.2	5.9	4.4	6.6	7.0	6.7	*	6.2
Male	12.8	12.4	8.2	4.9	5.5	*	*	*	*
Female	3.0	3.0	4.2	4.1	7.4	7.8	7.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/	12.6	8.9	6.0	3.5	2.8	1.6	0.7	*	*
Male	20.4	15.0	10.4	6.1	4.9	2.7	1.3	*	*
Female	5.1	3.2	2.0	1.3	1.0	0.6	*	*	*
Hispanic	17.3	13.4	9.5	6.4	5.7	*	*	*	*
Male	27.4	21.5	17.4	12.3	10.3	*	*	*	0.0
Female	7.5	5.8	*	*	*	*	0.0	0.0	*
Mexican	6.9	4.5	*	*	*	*	0.0	*	*
Male	11.8	7.5	*	*	*	*	0.0	*	0.0
Female	*	*	*	*	*	0.0	0.0	0.0	*
Puerto Rican	86.4	65.2	*	*	*	*	*	*	0.0
Male	*	*	*	*	*	*	*	*	0.0
Female	*	*	*	*	*	*	0.0	0.0	0.0
Cuban	*	*	*	*	*	*	0.0	0.0	0.0
Male	*	*	*	*	*	*	0.0	0.0	0.0
Female	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	14.5	8.1	*	*	*	0.0	*	0.0	0.0
Male	25.6	*	*	*	*	0.0	*	0.0	0.0
Female	*	*	*	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 5/	11.9	8.4	5.7	3.2	2.5	1.5	0.7	*	*
Male	19.3	14.1	9.7	5.4	4.4	2.5	1.2	*	*
Female	4.8	2.9	1.9	1.2	0.9	0.6	*	*	*
Non-Hispanic white	5.2	4.0	2.5	1.4	1.1	0.5	0.4	*	*
Male	9.1	7.4	4.5	2.7	2.1	0.9	0.7	*	*
Female	1.4	0.8	0.5	*	*	*	*	*	*
Non-Hispanic black	61.8	45.5	34.1	18.9	14.7	11.3	5.0	*	*
Male	101.6	77.4	61.4	32.4	25.7	20.6	*	*	*
Female	28.4	19.6	12.9	8.9	6.4	4.7	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malaria (B50-B54)

Non-Hispanic 5/	0.0	0.0	*	0.0	*	*	0.0	0.0	0.0
Male	0.0	0.0	*	0.0	0.0	*	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Non-Hispanic white	0.0	0.0	*	0.0	0.0	*	0.0	0.0	0.0
Male	0.0	0.0	*	0.0	0.0	*	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Non-Hispanic black	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 5/	2.0	2.7	3.3	4.4	6.8	9.1	12.6	17.0	27.2
Male	2.5	3.0	3.6	5.2	8.0	10.9	15.1	20.7	31.4
Female	1.6	2.4	3.0	3.8	5.8	7.8	10.9	14.8	25.4
Non-Hispanic white	1.5	2.2	2.9	3.9	6.4	8.7	12.0	16.4	26.7
Male	2.0	2.5	3.3	4.5	7.5	10.5	14.6	18.9	30.3
Female	1.1	1.9	2.6	3.4	5.4	7.3	10.1	14.9	25.2
Non-Hispanic black	5.4	6.4	6.7	8.5	10.7	13.4	19.4	25.2	34.7
Male	6.3	7.2	6.3	11.1	12.2	15.8	21.8	42.7	43.5
Female	4.6	5.7	7.0	6.5	9.5	11.6	17.9	15.7	31.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Malignant neoplasms (C00-C97)

All origins 3/	95.1	171.9	300.1	479.7	707.2	975.1	1,226.2	1,513.2	1,796.7
Male	96.3	181.1	337.9	562.5	860.4	1,222.9	1,576.2	2,032.7	2,619.2
Female	94.0	163.2	265.2	405.5	577.2	780.1	975.8	1,199.4	1,449.1
Hispanic	65.9	113.2	175.4	274.6	424.5	571.0	736.2	916.9	1,054.5
Male	65.8	115.3	206.4	339.4	538.1	740.4	958.9	1,236.0	1,419.8
Female	66.0	111.2	148.1	220.1	333.0	440.7	570.8	717.1	869.9
Mexican	58.4	108.6	159.8	276.0	428.8	562.3	749.4	842.9	*
Male	57.7	101.6	178.4	316.2	531.4	713.9	*	*	*
Female	59.2	115.9	142.0	238.4	340.1	441.2	598.5	*	*
Puerto Rican	88.8	168.7	248.6	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	240.6	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	69.1	101.8	174.4	273.1	415.8	563.9	*	*	*
Male	61.8	102.0	*	*	*	*	*	*	*
Female	75.0	101.6	136.2	215.9	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

Non-Hispanic 5/	97.6	176.3	309.0	493.9	725.0	996.4	1,247.5	1,535.6	1,827.7
Male	98.8	185.9	347.0	577.3	879.8	1,247.4	1,603.4	2,063.9	2,678.0
Female	96.3	167.1	273.8	418.9	593.2	798.7	993.3	1,216.9	1,471.6
Non-Hispanic white	91.2	167.0	297.3	482.3	718.9	987.9	1,234.9	1,527.0	1,825.5
Male	90.7	172.6	328.7	554.7	863.6	1,227.6	1,573.8	2,042.3	2,670.7
Female	91.7	161.6	267.5	415.5	593.1	796.3	990.8	1,214.8	1,474.8
Non-Hispanic black	153.2	276.6	452.1	664.5	876.6	1,217.0	1,543.5	1,815.6	2,048.1
Male	172.8	333.3	577.9	883.9	1,150.0	1,619.3	2,184.9	2,670.8	3,155.2
Female	136.7	230.6	354.7	501.7	672.5	933.5	1,134.2	1,357.3	1,588.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins 3/	2.3	3.7	5.8	8.2	9.4	11.5	13.5	15.6	19.2
Male	3.6	5.9	9.2	13.3	14.5	17.6	19.8	22.2	26.0
Female	1.0	1.6	2.8	3.6	5.1	6.7	9.1	11.6	16.3
Hispanic	1.1	1.8	3.1	5.0	6.0	7.1	10.2	13.3	10.6
Male	*	*	6.2	8.2	12.1	13.6	18.0	*	*
Female	*	*	*	*	*	*	*	15.2	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	*	0.0	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	0.0	*
Female	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic 5/	2.4	3.9	6.0	8.4	9.6	11.7	13.7	15.7	19.5
Male	3.8	6.2	9.4	13.7	14.6	17.8	19.8	22.7	26.6
Female	1.1	1.7	2.9	3.7	5.4	6.9	9.3	11.5	16.5
Non-Hispanic white	1.8	3.2	5.2	7.8	9.2	11.3	13.4	15.7	20.1
Male	2.8	5.0	7.8	12.5	13.7	17.1	18.9	22.4	27.0
Female	0.9	1.3	2.7	3.4	5.2	6.7	9.4	11.7	17.2
Non-Hispanic black	6.2	9.5	13.1	15.2	14.2	16.7	17.8	16.8	14.7
Male	11.0	16.2	23.2	27.3	23.7	26.3	32.8	29.4	23.4
Female	2.2	4.1	5.3	6.2	7.2	10.0	8.2	10.0	11.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasm of esophagus (C15)

All origins 3/	2.4	4.8	8.8	13.2	18.4	22.3	24.3	25.0	26.7
Male	4.3	8.6	15.7	22.4	32.2	39.6	43.4	44.5	46.2
Female	0.6	1.3	2.5	5.0	6.7	8.7	10.7	13.1	18.4
Hispanic	1.3	1.8	3.9	6.6	9.1	8.0	15.4	16.2	20.1
Male	2.4	3.1	7.8	12.9	15.4	15.0	28.0	26.8	36.2
Female	*	*	*	*	*	*	*	*	*
Mexican	*	*	*	6.1	9.5	*	19.9	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	0.0	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of esophagus (C15)

Non-Hispanic 5/	2.5	5.1	9.2	13.7	19.0	23.1	24.7	25.3	26.9
Male	4.5	9.0	16.2	23.0	33.2	40.9	44.1	45.3	46.7
Female	0.6	1.3	2.7	5.3	6.9	9.0	10.9	13.2	18.7
Non-Hispanic white	2.3	4.6	8.4	12.7	18.2	22.5	24.1	25.5	27.5
Male	4.1	8.2	14.8	21.4	32.4	40.1	43.0	46.2	47.1
Female	0.5	1.1	2.3	4.7	5.9	8.5	10.5	12.9	19.3
Non-Hispanic black	4.6	10.0	18.0	24.7	28.6	31.9	36.6	25.5	23.9
Male	8.1	18.0	32.8	43.5	45.8	55.3	66.9	37.4	46.9
Female	1.7	3.6	6.5	10.7	15.7	15.4	17.3	19.1	14.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of stomach (C16)									
Non-Hispanic 5/	2.2	3.8	6.1	9.6	13.8	19.9	27.6	36.6	49.5
Male	3.0	5.4	8.7	13.4	19.8	28.2	40.0	53.8	69.2
Female	1.5	2.3	3.6	6.2	8.7	13.3	18.7	26.3	41.3
Non-Hispanic white	1.8	2.9	4.9	7.8	12.2	17.1	24.0	32.4	43.9
Male	2.5	4.2	7.3	10.9	17.7	24.6	35.0	47.7	61.5
Female	1.1	1.7	2.7	5.0	7.4	11.1	16.0	23.2	36.6
Non-Hispanic black	4.2	9.2	13.8	20.0	23.6	41.1	60.3	78.9	104.4
Male	6.1	13.5	20.2	30.6	34.8	59.0	89.2	122.0	140.6
Female	2.5	5.6	8.8	12.2	15.1	28.5	41.8	55.8	89.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 5/	8.5	15.1	26.5	45.4	65.8	94.5	131.6	177.9	276.0
Male	9.4	17.5	32.3	55.6	80.5	116.9	161.6	217.1	319.2
Female	7.7	12.8	21.2	36.3	53.2	76.9	110.1	154.3	257.9
Non-Hispanic white	7.8	14.0	24.8	43.5	63.9	92.5	129.2	175.2	276.0
Male	8.7	16.4	30.1	53.2	78.7	116.0	158.4	215.4	322.1
Female	7.0	11.6	19.8	34.6	51.0	73.8	108.2	150.8	256.9
Non-Hispanic black	14.1	25.4	44.3	69.2	89.8	127.6	177.2	240.7	312.5
Male	16.2	29.9	58.4	87.3	109.2	144.9	225.5	276.9	330.3
Female	12.3	21.7	33.3	55.7	75.3	115.4	146.3	221.4	305.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic 5/	3.3	4.8	7.0	10.6	15.7	21.4	25.5	30.4	29.9
Male	5.5	7.4	10.4	15.2	23.2	31.2	35.4	41.7	37.7
Female	1.3	2.4	3.8	6.5	9.2	13.7	18.4	23.6	26.6
Non-Hispanic white	2.5	3.9	5.6	9.2	14.3	19.9	24.4	29.4	29.1
Male	4.0	5.8	8.3	13.0	21.2	29.2	33.9	40.9	38.3
Female	1.1	2.2	3.0	5.6	8.2	12.5	17.5	22.4	25.3
Non-Hispanic black	7.0	8.8	12.6	15.4	21.0	28.7	27.4	36.3	35.3
Male	12.7	15.8	20.2	23.4	31.7	41.5	39.9	43.6	29.0
Female	2.1	3.1	6.8	9.5	13.0	19.7	19.4	32.4	38.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of pancreas (C25)

Non-Hispanic 5/	4.6	9.1	16.3	26.0	39.2	53.9	68.0	84.3	100.7
Male	6.1	11.7	19.9	31.4	46.4	62.0	75.9	93.1	105.5
Female	3.2	6.7	12.9	21.1	33.1	47.5	62.4	79.0	98.7
Non-Hispanic white	4.5	8.6	15.3	25.2	38.4	52.7	66.5	82.7	99.8
Male	5.9	10.8	18.5	30.5	45.5	61.3	74.7	91.9	106.1
Female	3.1	6.5	12.3	20.4	32.2	45.9	60.5	77.0	97.2
Non-Hispanic black	6.6	15.3	26.5	33.7	51.0	71.5	89.9	111.8	120.4
Male	9.0	21.8	37.1	43.1	59.8	76.5	99.7	119.3	108.2
Female	4.6	10.0	18.2	26.8	44.4	67.9	83.6	107.8	125.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Malignant neoplasm of larynx (C32)

All origins 3/	0.8	1.5	3.2	4.7	6.2	7.5	7.2	7.5	7.0
Male	1.4	2.7	5.5	8.0	10.6	12.7	13.0	15.7	17.2
Female	0.2	0.4	1.1	1.8	2.5	3.4	3.1	2.6	2.7
Hispanic	*	*	*	*	4.9	5.1	6.8	*	*
Male	*	*	*	*	9.5	10.3	13.4	*	*
Female	0.0	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	0.0	*	0.0	0.0	*	*	*	0.0
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	0.0
Female	0.0	0.0	0.0	*	*	*	0.0	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	*	0.0	*	0.0	0.0
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	0.0	0.0	0.0	*	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of larynx (C32)

Non-Hispanic 5/	0.8	1.6	3.3	4.9	6.2	7.6	7.2	7.5	6.8
Male	1.3	2.8	5.6	8.3	10.6	12.8	12.9	15.6	16.8
Female	0.2	0.5	1.1	1.8	2.6	3.5	3.1	2.5	2.7
Non-Hispanic white	0.6	1.3	2.8	4.4	5.8	7.3	7.1	7.3	6.5
Male	1.0	2.3	4.7	7.4	9.7	12.1	12.6	15.1	15.7
Female	*	0.4	1.0	1.6	2.4	3.4	3.1	2.7	2.7
Non-Hispanic black	2.2	4.3	7.9	10.4	11.3	12.0	10.5	9.9	8.5
Male	4.2	7.7	15.8	19.2	20.8	22.6	19.9	26.7	23.4
Female	*	*	*	3.8	4.3	4.5	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
Non-Hispanic 5/	21.4	48.0	99.0	171.2	256.0	337.1	371.6	375.3	300.8
Male	26.1	58.8	125.7	220.8	338.4	456.2	517.2	558.0	521.1
Female	17.0	37.7	74.3	126.6	185.9	243.3	267.5	265.0	208.5
Non-Hispanic white	19.8	45.9	97.8	171.1	258.2	340.1	372.9	377.4	303.4
Male	23.4	54.2	120.9	214.8	334.8	453.1	513.0	555.1	526.9
Female	16.2	37.8	75.9	130.8	191.7	249.7	271.9	269.7	210.6
Non-Hispanic black	37.1	76.3	131.8	208.7	281.4	359.2	413.7	387.0	296.7
Male	52.4	114.2	201.7	327.0	425.6	549.3	650.8	673.1	544.5
Female	24.3	45.7	77.6	120.8	173.7	225.3	262.3	233.8	194.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Malignant melanoma of skin (C43)

All origins 3/	2.7	3.7	4.8	6.2	8.4	9.8	12.9	14.4	16.3
Male	3.5	4.8	6.9	8.6	12.1	14.8	19.5	22.6	26.4
Female	1.8	2.6	2.8	4.0	5.2	5.9	8.2	9.5	12.0
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	0.0	*
Puerto Rican	0.0	*	*	*	*	0.0	0.0	0.0	0.0
Male	0.0	*	*	*	*	0.0	0.0	0.0	0.0
Female	0.0	*	0.0	*	*	0.0	0.0	0.0	0.0
Cuban	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	0.0	*	0.0
Female	0.0	0.0	*	*	*	*	*	0.0	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Malignant neoplasm of breast (C50)

All origins 3/	13.5	21.6	28.6	35.4	43.9	56.6	70.1	93.6	143.7
Male	*	0.4	0.6	0.7	1.1	1.5	1.8	2.3	4.4
Female	26.5	41.7	54.5	66.4	80.2	99.9	119.0	148.8	202.6
Hispanic	9.7	16.1	17.6	21.0	23.9	29.4	32.7	46.3	65.0
Male	*	*	*	0.0	*	*	*	*	0.0
Female	19.0	30.7	32.5	38.7	42.5	50.9	56.6	74.4	97.9
Mexican	8.0	15.6	14.8	21.2	22.1	28.4	29.6	*	*
Male	*	*	*	0.0	0.0	*	*	*	0.0
Female	16.2	31.6	28.7	41.1	41.2	50.3	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	0.0	*	0.0	0.0	*	*	0.0	0.0	0.0
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	12.0	16.0	19.6	26.2	*	*	*	*	*
Male	0.0	0.0	*	0.0	*	*	0.0	0.0	0.0
Female	21.8	27.9	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of breast (C50)									
Non-Hispanic 5/	13.9	22.0	29.4	36.3	45.2	58.1	71.8	95.5	147.1
Male	*	0.4	0.6	0.7	1.1	1.4	1.9	2.4	4.6
Female	27.1	42.6	56.1	68.4	82.7	102.8	121.7	151.7	206.8
Non-Hispanic white	12.4	20.5	27.8	35.1	44.5	57.8	71.5	95.8	149.2
Male	*	0.3	0.5	0.6	1.1	1.3	1.8	2.3	4.5
Female	24.5	40.2	53.6	67.0	82.3	102.9	121.7	152.4	209.2
Non-Hispanic black	25.5	37.4	46.7	51.1	58.0	71.7	90.6	112.4	150.8
Male	*	*	*	*	*	*	*	*	*
Female	46.7	66.8	81.9	87.5	100.1	119.9	146.6	170.3	210.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic 5/	2.3	2.7	3.0	3.5	3.6	3.9	4.6	4.6	7.9
Female	4.5	5.2	5.8	6.7	6.7	7.0	7.9	7.4	11.2
Non-Hispanic white	2.1	2.4	2.6	2.9	3.1	3.0	3.9	4.1	6.7
Female	4.3	4.8	5.0	5.6	5.8	5.4	6.7	6.5	9.5
Non-Hispanic black	3.2	4.5	6.5	8.6	8.6	13.4	12.6	10.3	21.9
Female	5.8	8.1	11.5	15.0	14.9	22.8	20.6	15.7	31.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 5/	1.0	1.7	3.3	5.7	8.9	11.2	14.7	20.0	24.5
Female	1.9	3.4	6.4	10.9	16.6	20.1	25.3	32.1	34.7
Non-Hispanic white	1.0	1.7	3.1	5.3	7.7	10.4	14.0	19.4	23.9
Female	1.9	3.3	6.0	10.1	14.4	18.8	24.0	31.1	33.8
Non-Hispanic black	*	2.4	6.2	11.1	20.8	22.1	26.3	32.0	36.3
Female	*	4.4	10.9	19.4	36.3	37.8	43.0	49.1	51.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasm of ovary (C56)

Non-Hispanic 5/	3.3	5.8	8.7	13.1	17.8	23.0	29.7	36.5	38.3
Female	6.5	11.3	16.7	24.9	32.9	41.1	50.9	58.5	54.4
Non-Hispanic white	3.4	5.9	9.0	13.6	18.4	23.5	30.2	37.4	39.6
Female	6.8	11.6	17.5	26.2	34.3	42.3	51.9	60.0	56.0
Non-Hispanic black	2.5	5.5	7.6	11.5	15.0	23.1	29.4	30.1	29.4
Female	4.6	10.0	13.5	20.1	26.2	39.4	48.2	46.3	41.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasm of prostate (C61)

Non-Hispanic 5/	0.5	1.7	5.0	12.8	29.1	53.5	90.4	145.6	209.5
Male	1.1	3.4	10.4	27.1	63.3	121.4	217.0	386.9	709.8
Non-Hispanic white	0.4	1.4	4.2	10.5	25.3	47.6	82.0	136.4	197.3
Male	0.9	2.8	8.6	21.9	54.4	107.2	195.9	361.5	672.7
Non-Hispanic black	1.4	4.6	12.7	35.0	68.8	126.1	201.9	290.4	373.0
Male	3.2	10.2	29.1	82.2	160.9	305.0	518.3	832.4	1,271.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic 5/	2.2	4.4	7.5	11.1	14.7	19.7	23.3	27.5	31.6
Male	3.2	6.4	11.0	16.2	20.7	27.8	33.5	39.3	48.5
Female	1.2	2.5	4.3	6.5	9.5	13.3	15.9	20.4	24.6
Non-Hispanic white	2.1	4.4	7.6	11.4	15.1	20.0	23.5	27.8	31.8
Male	3.1	6.2	11.0	16.7	21.1	28.1	33.9	40.5	49.2
Female	1.2	2.6	4.4	6.6	9.8	13.5	16.0	20.1	24.6
Non-Hispanic black	3.1	5.4	8.0	10.2	13.6	19.6	23.9	26.1	32.1
Male	4.5	9.0	13.3	15.1	20.0	27.6	34.2	28.5	46.9
Female	2.0	2.5	3.9	6.5	8.7	14.0	17.3	24.8	25.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasm of bladder (C67)

All origins 3/	0.7	2.0	3.8	7.2	11.9	20.5	29.7	44.4	68.1
Male	1.1	3.0	6.0	11.3	19.6	33.2	51.5	79.0	129.7
Female	0.3	0.9	1.8	3.5	5.4	10.5	14.1	23.5	42.1
Hispanic	*	*	2.3	4.0	5.7	10.8	11.7	18.7	29.1
Male	*	*	*	7.3	11.0	18.3	18.7	30.7	56.7
Female	*	*	*	*	*	*	*	*	15.1
Mexican	0.0	*	*	*	*	11.0	*	*	*
Male	0.0	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*
Puerto Rican	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	0.0	*	*	*	*	*	*	*
Female	0.0	*	0.0	*	*	*	*	0.0	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	0.0	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of bladder (C67)									
Non-Hispanic 5/	0.8	2.0	3.9	7.4	12.3	21.1	30.5	45.5	69.7
Male	1.2	3.1	6.2	11.5	20.1	34.1	53.0	81.1	133.2
Female	0.4	1.0	1.9	3.6	5.7	10.8	14.4	24.0	43.2
Non-Hispanic white	0.7	2.1	4.0	7.6	12.7	21.7	31.4	46.7	71.4
Male	1.1	3.1	6.4	11.9	21.0	35.7	55.1	83.9	140.4
Female	0.4	1.0	1.6	3.6	5.5	10.5	14.3	24.2	42.8
Non-Hispanic black	1.1	2.3	4.8	7.0	11.7	19.4	25.9	37.9	61.5
Male	*	4.1	5.9	10.2	15.3	23.3	38.4	60.5	81.4
Female	*	*	3.9	4.6	8.9	16.6	17.9	25.8	53.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 5/	4.5	6.5	9.7	12.2	16.4	18.8	21.5	19.9	16.2
Male	5.6	8.0	11.2	15.3	20.5	23.2	26.3	24.6	20.7
Female	3.5	5.1	8.3	9.4	12.9	15.4	18.1	17.0	14.2
Non-Hispanic white	5.0	7.0	10.5	13.1	17.7	20.3	22.7	20.8	16.9
Male	6.1	8.4	12.0	16.3	22.0	24.9	27.7	26.0	22.0
Female	3.9	5.5	9.1	10.2	14.0	16.7	19.2	17.6	14.7
Non-Hispanic black	2.8	4.3	5.7	7.1	8.6	7.7	12.2	11.8	10.1
Male	4.0	5.8	6.5	9.5	10.9	8.8	14.2	*	*
Female	1.7	3.0	5.0	5.3	6.8	6.9	10.9	13.4	9.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 5/	8.5	14.1	26.4	40.8	64.5	96.7	133.3	179.1	202.7
Male	10.4	17.4	32.2	50.7	80.8	122.1	170.4	224.7	277.1
Female	6.7	10.9	21.1	31.9	50.6	76.6	106.8	151.6	171.5
Non-Hispanic white	8.3	13.6	26.0	40.5	65.6	98.5	135.6	183.3	208.4
Male	10.0	17.0	31.5	50.3	82.3	124.6	173.2	231.7	286.4
Female	6.6	10.4	20.7	31.4	51.1	77.6	108.5	154.0	176.0
Non-Hispanic black	11.0	19.7	34.9	47.9	64.7	92.6	123.0	148.2	164.2
Male	14.1	24.9	44.9	59.8	80.3	113.2	158.1	167.4	207.5
Female	8.4	15.4	27.1	39.2	53.0	78.1	100.6	137.9	146.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Hodgkin's disease (C81)

All origins 3/	0.4	0.5	0.7	0.9	1.3	1.9	2.0	2.3	2.3
Male	0.5	0.6	0.9	1.2	1.7	2.3	2.3	2.8	3.6
Female	0.3	0.4	0.5	0.5	0.9	1.5	1.9	2.1	1.8
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	0.0	*	*	*	*	*	*	0.0	0.0
Male	0.0	*	*	*	*	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	*	0.0	*	*	0.0	0.0
Cuban	*	*	0.0	0.0	*	*	*	*	*
Male	*	*	0.0	0.0	*	*	*	*	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	0.0
Female	*	*	*	*	0.0	*	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Non-Hodgkin's lymphoma (C82-C85)

Non-Hispanic 5/	3.6	6.4	11.3	16.9	26.2	39.3	55.6	74.7	78.8
Male	4.8	8.2	14.1	20.4	31.9	47.9	68.3	86.6	103.3
Female	2.4	4.7	8.7	13.7	21.4	32.6	46.5	67.5	68.6
Non-Hispanic white	3.5	6.6	11.6	17.7	27.9	41.2	58.0	78.8	82.7
Male	4.7	8.4	14.3	20.9	33.6	50.2	70.5	92.0	107.9
Female	2.3	4.9	9.1	14.7	23.0	34.0	49.1	70.8	72.3
Non-Hispanic black	4.3	6.0	11.2	11.9	17.0	24.7	33.1	33.2	40.9
Male	5.9	8.6	15.0	16.7	23.4	28.3	47.0	28.5	55.8
Female	2.9	3.9	8.2	8.4	12.2	22.1	24.2	35.8	34.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Leukemia (C91-C95)

Non-Hispanic 5/	3.2	4.5	8.8	13.8	22.5	34.3	47.4	64.8	84.2
Male	3.6	5.5	10.5	17.7	30.2	46.9	65.3	89.2	119.0
Female	2.9	3.6	7.2	10.3	16.0	24.3	34.6	50.1	69.6
Non-Hispanic white	3.2	4.4	8.9	13.9	23.0	35.5	49.1	66.2	87.7
Male	3.5	5.3	10.6	17.8	30.8	48.6	67.4	91.9	124.8
Female	2.9	3.5	7.3	10.3	16.1	25.0	35.9	50.6	72.3
Non-Hispanic black	3.7	5.9	8.6	14.4	22.2	28.6	36.2	54.4	56.6
Male	4.8	7.7	11.3	18.8	28.9	37.4	51.3	62.3	73.6
Female	2.7	4.4	6.5	11.2	17.3	22.3	26.7	50.1	49.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 5/	1.2	2.6	5.7	9.3	14.4	21.2	28.1	37.1	37.0
Male	1.4	3.1	6.6	11.4	17.0	24.9	34.3	46.0	50.9
Female	1.1	2.1	4.7	7.4	12.2	18.2	23.6	31.7	31.2
Non-Hispanic white	1.1	2.1	4.7	7.9	13.4	19.7	26.1	35.7	35.3
Male	1.3	2.7	5.6	10.2	16.2	23.3	32.6	44.8	49.7
Female	0.9	1.5	3.9	5.9	11.0	16.9	21.5	30.2	29.4
Non-Hispanic black	2.6	7.2	14.3	21.1	24.2	38.4	52.5	60.3	64.1
Male	2.8	7.3	17.4	23.6	26.0	46.2	58.4	75.7	75.9
Female	2.4	7.1	11.9	19.3	22.9	33.0	48.8	52.0	59.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 5/	11.8	20.1	33.7	54.7	78.4	111.0	144.6	193.5	253.2
Male	13.8	23.5	39.9	65.5	93.9	136.0	174.2	234.3	313.8
Female	10.0	16.9	27.9	44.9	65.2	91.4	123.5	168.9	227.9
Non-Hispanic white	11.0	19.0	32.0	53.1	78.7	110.3	144.3	193.9	256.0
Male	12.5	21.9	37.9	63.2	93.6	135.2	174.8	236.6	320.8
Female	9.6	16.2	26.5	43.9	65.8	90.4	122.4	167.9	229.1
Non-Hispanic black	19.4	31.7	50.9	76.8	85.1	130.7	162.0	205.9	247.7
Male	25.0	40.8	66.9	101.9	108.7	160.8	190.3	236.8	275.6
Female	14.8	24.3	38.5	58.1	67.5	109.5	143.9	189.4	236.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 5/	1.5	2.4	3.6	6.6	11.2	19.1	29.9	49.0	84.3
Male	1.5	2.7	4.2	8.3	14.2	24.6	37.6	61.9	108.6
Female	1.4	2.0	3.0	5.0	8.6	14.8	24.4	41.2	74.2
Non-Hispanic white	1.4	2.3	3.5	6.5	11.1	19.5	30.3	50.1	87.1
Male	1.5	2.7	4.0	8.4	14.2	25.3	38.2	63.9	114.7
Female	1.2	1.9	3.0	4.7	8.4	14.9	24.7	41.8	75.7
Non-Hispanic black	2.3	3.3	5.2	8.3	13.8	18.4	30.0	41.3	63.5
Male	1.9	3.5	6.7	9.0	16.9	21.2	35.6	43.6	60.2
Female	2.7	3.2	4.0	7.7	11.5	16.4	26.4	40.1	64.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Anemias (D50-D64)

All origins 3/	0.7	0.6	0.8	1.2	1.9	3.5	6.8	13.8	43.6
Male	0.7	0.6	0.7	1.2	2.1	3.5	7.1	14.7	41.8
Female	0.7	0.6	0.9	1.2	1.6	3.6	6.6	13.3	44.4
Hispanic	*	*	*	*	*	4.1	*	9.4	26.4
Male	*	0.0	0.0	*	*	*	*	*	29.9
Female	*	*	*	*	*	*	*	*	24.7
Mexican	*	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*
Puerto Rican	*	*	*	0.0	*	*	*	*	*
Male	*	0.0	0.0	0.0	*	*	*	*	*
Female	0.0	*	*	0.0	0.0	0.0	*	*	*
Cuban	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Male	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Other Hispanic 4/	*	*	0.0	*	*	*	0.0	*	*
Male	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	*	*	0.0	*	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Anemias (D50-D64)

Non-Hispanic 5/	0.7	0.6	0.8	1.2	1.9	3.5	6.8	14.0	44.3
Male	0.7	0.6	0.7	1.3	2.1	3.3	7.2	14.7	42.2
Female	0.7	0.6	0.9	1.2	1.6	3.7	6.6	13.6	45.1
Non-Hispanic white	0.3	0.3	0.4	0.8	1.7	3.2	6.6	14.0	43.6
Male	0.4	0.3	*	0.7	2.0	2.9	7.0	14.7	42.6
Female	0.3	*	0.5	1.0	1.4	3.5	6.2	13.5	44.0
Non-Hispanic black	3.4	3.2	4.5	4.4	3.7	6.7	11.6	17.4	53.7
Male	3.6	3.6	4.6	6.5	*	7.8	10.4	16.9	41.3
Female	3.3	2.9	4.3	*	3.7	5.9	12.4	17.7	58.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

Non-Hispanic 5/	10.0	16.7	29.2	47.7	73.3	106.8	149.8	213.9	312.0
Male	12.5	19.5	33.3	52.9	80.9	116.7	166.5	236.9	330.9
Female	7.6	14.0	25.4	42.9	66.8	98.9	137.9	200.0	304.1
Non-Hispanic white	8.1	13.5	23.3	39.3	63.2	93.5	136.2	198.0	293.0
Male	10.4	16.4	27.7	45.3	72.4	107.6	157.1	228.0	320.0
Female	5.9	10.7	19.1	33.8	55.2	82.3	121.1	179.7	281.7
Non-Hispanic black	24.0	43.7	76.7	117.2	159.8	234.3	302.3	422.4	537.2
Male	29.7	48.5	84.7	122.5	163.8	213.0	279.1	381.0	475.3
Female	19.2	39.7	70.5	113.3	156.8	249.3	317.2	444.6	562.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Nutritional deficiencies (E40-E64)

All origins 3/	0.2	0.4	0.6	0.9	1.6	3.2	6.4	16.1	53.7
Male	0.2	0.4	0.7	1.1	1.7	3.9	6.7	15.9	46.5
Female	0.2	0.4	0.5	0.7	1.5	2.7	6.2	16.2	56.7
Hispanic	*	*	*	*	*	*	*	*	38.6
Male	*	*	*	*	*	*	*	*	37.8
Female	0.0	*	*	*	*	*	*	*	39.0
Mexican	*	*	*	*	0.0	*	*	*	*
Male	*	*	*	0.0	0.0	*	*	*	*
Female	0.0	0.0	*	*	0.0	*	*	*	*
Puerto Rican	0.0	*	0.0	0.0	*	*	*	0.0	*
Male	0.0	*	0.0	0.0	*	*	0.0	0.0	*
Female	0.0	*	0.0	0.0	*	*	*	0.0	*
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	0.0	*	*	0.0	*
Female	0.0	0.0	0.0	0.0	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Nutritional deficiencies (E40-E64)

Non-Hispanic 5/	0.2	0.4	0.6	0.9	1.6	3.2	6.5	16.5	54.2
Male	0.2	0.4	0.7	1.2	1.7	3.9	6.7	16.4	46.9
Female	0.3	0.4	0.5	0.7	1.6	2.7	6.4	16.5	57.3
Non-Hispanic white	0.2	0.3	0.4	0.8	1.5	2.7	6.0	15.3	53.3
Male	*	0.3	0.4	1.0	1.4	3.0	5.9	15.1	45.3
Female	*	0.4	0.4	0.6	1.5	2.4	6.1	15.5	56.6
Non-Hispanic black	*	*	2.1	2.1	3.2	8.5	12.8	32.9	71.7
Male	*	*	*	*	*	12.8	16.1	35.6	71.4
Female	*	*	*	*	*	5.5	10.6	31.5	71.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Malnutrition (E40-E46)

All origins 3/	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.9	49.7
Male	0.2	0.3	0.6	1.1	1.6	3.6	6.1	14.7	42.7
Female	0.2	0.4	0.4	0.6	1.4	2.6	5.9	15.1	52.7
Hispanic	*	*	*	*	*	*	*	*	35.9
Male	*	*	*	*	*	*	*	*	33.1
Female	0.0	*	*	*	*	*	*	*	37.4
Mexican	*	*	*	*	0.0	*	*	*	*
Male	*	*	*	0.0	0.0	*	*	*	*
Female	0.0	0.0	*	*	0.0	*	*	*	*
Puerto Rican	0.0	*	0.0	0.0	*	*	*	0.0	*
Male	0.0	*	0.0	0.0	*	*	0.0	0.0	*
Female	0.0	*	0.0	0.0	*	*	*	0.0	*
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	0.0	*	*	0.0	*
Female	0.0	0.0	0.0	0.0	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Malnutrition (E40-E46)

Non-Hispanic 5/	0.2	0.3	0.5	0.9	1.6	3.0	6.1	15.3	50.3
Male	0.2	0.3	0.6	1.1	1.6	3.6	6.1	15.1	43.2
Female	0.2	0.4	0.4	0.6	1.5	2.6	6.1	15.4	53.3
Non-Hispanic white	0.2	0.3	0.3	0.8	1.4	2.5	5.6	14.2	49.3
Male	*	*	*	1.0	1.3	2.8	5.4	13.7	41.3
Female	*	0.4	*	0.6	1.4	2.3	5.7	14.5	52.7
Non-Hispanic black	*	*	1.7	2.1	3.2	8.2	11.5	30.8	66.7
Male	*	*	*	*	*	12.1	13.3	33.8	70.3
Female	*	*	*	*	*	5.5	10.3	29.1	65.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other nutritional deficiencies (E50-E64)

Non-Hispanic 5/	*	*	*	*	*	*	0.5	1.2	4.0
Male	*	*	*	*	*	*	*	1.3	3.7
Female	*	*	*	*	*	*	*	1.1	4.1
Non-Hispanic white	*	*	*	*	*	*	0.4	1.1	4.0
Male	*	*	*	*	*	*	*	1.3	4.1
Female	0.0	0.0	*	*	*	*	*	1.0	3.9
Non-Hispanic black	*	*	*	0.0	0.0	*	*	*	*
Male	0.0	*	*	0.0	0.0	*	*	*	*
Female	*	0.0	*	0.0	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Meningitis (G00,G03)

All origins 3/	0.3	0.3	0.3	0.5	0.5	0.8	0.9	1.2	1.5
Male	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.6
Female	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4
Hispanic	*	*	*	*	*	*	*	0.0	*
Male	*	*	*	0.0	*	*	0.0	0.0	*
Female	*	*	0.0	*	*	0.0	*	0.0	*
Mexican	*	*	*	*	*	*	*	0.0	*
Male	*	*	*	0.0	0.0	*	0.0	0.0	*
Female	*	*	0.0	*	*	0.0	*	0.0	*
Puerto Rican	*	*	0.0	*	*	*	0.0	0.0	0.0
Male	0.0	*	0.0	0.0	*	*	0.0	0.0	0.0
Female	*	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Cuban	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	*	0.0	*	*	*	0.0	*	0.0	0.0
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	*	*	0.0	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Parkinson's disease (G20-G21)

All origins 3/	*	0.2	0.6	1.6	5.7	16.9	41.8	83.9	123.8
Male	*	0.3	0.7	2.4	7.8	25.0	63.5	130.5	197.9
Female	*	*	0.4	0.9	3.9	10.5	26.4	55.8	92.5
Hispanic	0.0	*	*	*	3.9	12.3	27.6	42.8	58.1
Male	0.0	*	*	*	7.0	15.5	40.1	67.7	77.1
Female	0.0	0.0	*	*	*	9.8	18.4	27.2	48.5
Mexican	0.0	*	*	*	*	12.5	24.2	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	0.0	0.0	*	*	*	*	*	*	*
Puerto Rican	0.0	0.0	*	0.0	*	*	*	*	*
Male	0.0	0.0	*	0.0	*	0.0	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	*	0.0	*
Cuban	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	*	*	*	*	*
Other Hispanic 4/	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Parkinson's disease (G20-G21)

Non-Hispanic 5/	*	0.2	0.6	1.7	5.8	17.1	42.5	85.6	126.6
Male	*	0.3	0.7	2.4	7.8	25.5	64.6	133.1	204.1
Female	*	*	0.5	1.0	4.0	10.6	26.8	56.9	94.1
Non-Hispanic white	*	0.2	0.6	1.7	6.2	18.3	44.9	90.2	134.1
Male	*	0.3	0.8	2.5	8.3	27.3	67.9	140.7	216.9
Female	*	*	0.4	1.0	4.3	11.0	28.3	59.7	99.8
Non-Hispanic black	0.0	*	*	*	3.6	8.8	20.2	34.8	52.3
Male	0.0	*	*	*	5.7	12.1	31.8	49.9	82.6
Female	0.0	*	*	*	*	6.4	12.7	26.7	39.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Alzheimer's disease (G30)

All origins 3/	0.1	0.2	1.0	3.1	8.8	27.1	78.7	209.0	598.3
Male	*	*	1.1	3.1	9.0	27.1	76.3	196.9	467.3
Female	*	0.3	0.9	3.0	8.6	27.0	80.4	216.3	653.7
Hispanic	0.0	*	*	*	4.1	15.3	47.3	96.5	264.8
Male	0.0	0.0	0.0	*	*	16.0	44.7	78.0	203.0
Female	0.0	*	*	*	*	14.8	49.1	108.0	296.1
Mexican	0.0	*	0.0	*	*	14.4	43.1	*	*
Male	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.0	*	0.0	*	*	*	*	*	*
Puerto Rican	0.0	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.0	0.0	*	*	*	*	*	*	*
Cuban	0.0	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	0.0	0.0	*	*	*	*	*	*	*
Other Hispanic 4/	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	*	*	*
Female	0.0	*	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Alzheimer's disease (G30)

Non-Hispanic 5/	0.1	0.2	1.0	3.2	9.1	27.7	80.2	213.5	612.8
Male	*	*	1.2	3.2	9.2	27.7	77.8	202.0	480.3
Female	*	0.2	0.9	3.2	9.1	27.7	81.8	220.5	668.3
Non-Hispanic white	*	0.3	1.1	3.4	9.6	28.6	83.0	220.9	642.6
Male	*	*	1.2	3.5	9.5	28.5	80.6	209.9	505.5
Female	*	*	1.0	3.4	9.6	28.6	84.7	227.5	699.6
Non-Hispanic black	*	*	*	1.9	6.9	26.0	66.2	163.1	379.2
Male	0.0	*	*	*	8.1	27.3	61.7	142.4	301.2
Female	*	0.0	*	*	6.0	25.2	69.1	174.1	411.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Major cardiovascular diseases (I00-I78)

All origins 3/	89.7	153.0	259.0	423.1	685.7	1,124.8	1,892.1	3,401.8	8,155.7
Male	127.9	220.0	366.4	581.2	921.9	1,474.0	2,356.9	4,052.4	8,277.6
Female	52.8	89.5	159.7	281.6	485.2	850.0	1,559.5	3,008.8	8,104.2
Hispanic	60.0	110.7	181.9	304.9	514.2	817.5	1,340.4	2,255.5	4,773.7
Male	85.0	165.9	257.4	420.5	692.3	1,090.9	1,684.7	2,551.3	4,813.3
Female	35.5	59.5	115.4	207.5	370.7	607.2	1,084.4	2,070.4	4,753.6
Mexican	57.0	114.6	171.6	308.8	534.5	814.9	1,366.2	2,087.9	*
Male	75.3	159.7	224.7	399.7	661.6	1,050.6	*	*	*
Female	37.6	67.5	120.6	223.5	424.7	626.7	1,145.8	*	*
Puerto Rican	88.2	153.2	302.1	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	215.0	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	50.3	85.2	156.5	260.3	450.7	791.2	*	*	*
Male	82.4	138.7	*	*	*	*	*	*	*
Female	24.1	45.3	82.7	166.5	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 5/	92.0	155.7	263.7	429.9	694.8	1,138.2	1,913.5	3,442.2	8,292.6
Male	131.2	223.4	372.7	589.3	933.3	1,489.3	2,382.9	4,108.0	8,438.6
Female	54.2	91.6	162.6	286.4	491.6	861.7	1,578.3	3,040.6	8,231.5
Non-Hispanic white	78.5	136.4	235.4	394.4	658.4	1,091.8	1,868.9	3,421.7	8,422.4
Male	116.3	199.9	338.8	547.1	897.1	1,447.5	2,351.1	4,120.9	8,613.0
Female	41.1	74.7	137.3	253.6	451.0	807.6	1,521.6	2,998.2	8,343.3
Non-Hispanic black	199.6	337.0	535.1	777.0	1,087.8	1,734.6	2,616.6	4,031.6	7,715.1
Male	264.7	469.9	741.7	1,067.3	1,385.6	2,102.3	3,005.6	4,448.7	7,465.2
Female	145.0	229.4	375.0	561.5	865.6	1,475.6	2,368.3	3,808.0	7,818.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/	72.8	127.1	214.4	347.8	552.6	878.4	1,430.4	2,518.3	6,032.4
Male	108.9	189.3	313.3	489.0	763.7	1,183.8	1,837.7	3,103.0	6,313.3
Female	38.0	68.1	123.0	221.2	373.5	638.2	1,139.0	2,165.1	5,913.8
Hispanic	43.2	86.7	142.8	244.3	402.6	631.6	1,030.1	1,687.1	3,683.2
Male	64.9	135.0	211.6	341.7	557.5	861.2	1,306.8	1,950.5	3,658.0
Female	22.1	41.9	82.1	162.1	277.8	455.2	824.4	1,522.2	3,696.0
Mexican	40.5	87.9	133.0	244.8	405.2	612.8	1,019.8	1,521.7	*
Male	58.3	127.4	182.2	322.0	518.8	815.1	*	*	*
Female	21.6	46.6	85.7	172.5	306.9	451.1	834.8	*	*
Puerto Rican	66.3	131.1	246.6	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	178.1	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	35.3	63.6	120.9	209.1	346.7	605.9	*	*	*
Male	60.1	110.1	*	*	*	*	*	*	*
Female	15.0	29.0	54.9	132.6	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic 5/	75.2	129.8	218.9	353.7	560.7	889.0	1,445.3	2,546.7	6,125.3
Male	112.4	192.8	319.3	496.6	774.2	1,196.4	1,857.6	3,145.3	6,435.0
Female	39.4	70.0	125.8	225.0	378.9	646.9	1,150.8	2,185.7	5,995.7
Non-Hispanic white	66.4	116.8	199.0	329.7	536.4	858.7	1,417.7	2,539.8	6,235.4
Male	102.7	177.0	294.7	467.7	750.8	1,171.6	1,843.6	3,169.1	6,589.0
Female	30.6	58.3	108.1	202.4	350.0	608.8	1,110.9	2,158.6	6,088.7
Non-Hispanic black	151.3	260.3	421.0	606.2	847.0	1,308.9	1,922.8	2,923.0	5,580.3
Male	207.3	372.1	601.6	848.2	1,098.8	1,614.3	2,242.8	3,286.0	5,517.2
Female	104.4	169.8	281.1	426.6	659.1	1,093.8	1,718.5	2,728.4	5,606.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic 5/	0.5	0.9	1.5	2.1	3.5	6.0	8.5	13.0	18.4
Male	0.4	0.6	1.1	1.3	2.5	4.9	5.5	9.2	10.5
Female	0.6	1.1	1.9	2.8	4.4	6.8	10.5	15.3	21.7
Non-Hispanic white	0.5	0.7	1.4	2.0	3.6	6.2	8.9	13.6	19.4
Male	0.4	0.6	1.0	1.3	2.7	5.1	5.6	9.7	11.4
Female	0.6	0.9	1.8	2.7	4.5	7.1	11.2	16.0	22.7
Non-Hispanic black	0.9	1.8	2.3	2.2	2.7	4.2	4.3	*	8.5
Male	*	*	*	*	*	*	*	*	*
Female	*	2.1	2.9	*	*	5.0	*	*	11.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

Non-Hispanic 5/	4.7	6.4	9.8	12.2	16.4	22.6	36.7	65.7	186.5
Male	6.3	8.6	12.9	15.2	19.6	24.8	39.0	58.9	130.8
Female	3.2	4.4	6.9	9.5	13.6	20.8	35.0	69.8	209.8
Non-Hispanic white	2.9	4.1	6.3	8.6	11.9	17.0	29.8	57.9	176.9
Male	4.1	5.9	8.8	11.1	14.2	19.0	31.9	51.3	121.5
Female	1.6	2.4	4.0	6.4	9.8	15.4	28.4	61.8	199.8
Non-Hispanic black	18.8	26.9	39.7	43.5	56.5	80.2	116.9	170.8	312.1
Male	24.1	34.8	53.2	55.8	71.2	88.3	128.2	167.4	249.9
Female	14.4	20.5	29.3	34.4	45.4	74.6	109.7	172.7	337.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Hypertensive heart and renal disease (I13)

All origins 3/	0.4	0.5	0.9	1.4	2.1	3.1	6.6	12.5	29.0
Male	0.5	0.6	1.1	1.6	2.3	3.2	7.2	13.2	30.2
Female	0.3	0.3	0.7	1.2	2.0	3.0	6.2	12.0	28.5
Hispanic	*	*	*	*	*	*	5.7	9.4	17.4
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	15.9
Mexican	*	*	*	*	*	*	*	*	*
Male	*	0.0	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	0.0	*	*
Female	0.0	*	0.0	0.0	*	*	*	*	*
Cuban	*	0.0	*	*	0.0	*	0.0	*	*
Male	*	0.0	*	*	0.0	0.0	0.0	0.0	*
Female	*	0.0	0.0	0.0	0.0	*	0.0	*	*
Other Hispanic 4/	*	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Hypertensive heart and renal disease (I13)

Non-Hispanic 5/	0.4	0.5	0.9	1.4	2.1	3.1	6.7	12.6	29.5
Male	0.5	0.7	1.1	1.6	2.3	3.2	7.3	13.4	30.6
Female	0.3	0.3	0.7	1.2	1.9	3.0	6.2	12.1	29.0
Non-Hispanic white	*	0.2	0.4	0.6	1.1	1.8	5.2	10.6	26.4
Male	*	*	0.4	0.7	1.3	1.9	5.8	12.2	28.2
Female	*	*	*	0.6	0.9	1.7	4.7	9.6	25.6
Non-Hispanic black	2.4	2.5	5.3	7.6	10.1	15.9	22.4	37.9	66.7
Male	3.3	3.9	6.9	10.2	10.7	17.2	23.3	30.3	56.9
Female	1.7	*	4.2	5.7	9.7	15.0	21.8	42.0	70.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Ischemic heart diseases (I20-I25)

All origins 3/	49.2	92.7	159.8	264.4	424.3	669.9	1,083.0	1,866.6	4,296.3
Male	78.4	145.2	243.0	384.5	603.8	932.2	1,440.7	2,384.8	4,654.4
Female	21.1	43.1	83.0	156.9	271.9	463.6	827.1	1,553.5	4,145.0
Hispanic	30.3	65.7	109.2	193.5	322.0	514.8	836.2	1,337.7	2,920.5
Male	47.8	107.6	169.5	274.4	458.6	713.2	1,087.8	1,584.9	2,910.3
Female	13.2	26.8	56.1	125.4	212.1	362.3	649.2	1,182.8	2,925.6
Mexican	29.1	65.1	101.5	194.2	330.7	503.8	832.3	1,195.4	*
Male	44.5	99.4	146.2	261.2	437.5	677.4	*	*	*
Female	12.7	29.3	58.5	131.3	238.3	365.0	657.8	*	*
Puerto Rican	46.2	96.8	184.9	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	141.3	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	25.5	51.4	92.3	160.3	279.2	481.7	*	*	*
Male	45.1	93.2	*	*	*	*	*	*	*
Female	*	20.2	36.4	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

Non-Hispanic 5/	50.7	94.5	162.9	268.3	429.5	675.9	1,091.1	1,883.3	4,347.3
Male	80.9	147.5	247.2	390.1	610.8	939.9	1,452.7	2,413.3	4,731.7
Female	21.7	44.2	84.8	158.8	275.1	467.9	832.9	1,563.6	4,186.2
Non-Hispanic white	47.7	88.4	153.1	255.7	417.4	659.9	1,078.7	1,886.7	4,432.0
Male	77.8	140.1	234.9	374.7	601.7	929.7	1,454.0	2,445.1	4,862.0
Female	18.1	38.3	75.5	145.8	257.3	444.4	808.4	1,548.4	4,253.5
Non-Hispanic black	81.7	161.7	271.8	412.7	592.4	919.0	1,351.0	2,045.1	3,863.3
Male	118.8	241.1	402.9	593.9	779.0	1,159.6	1,604.4	2,324.5	3,832.5
Female	50.5	97.4	170.3	278.3	453.2	749.5	1,189.3	1,895.4	3,876.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

Non-Hispanic 5/	22.6	41.3	73.1	119.6	187.1	278.7	427.6	707.1	1,389.9
Male	35.6	64.9	110.9	172.7	262.3	382.0	560.3	892.7	1,590.2
Female	10.0	18.9	38.0	71.8	123.2	197.3	332.9	595.2	1,306.0
Non-Hispanic white	22.0	40.1	71.2	117.5	186.2	275.2	424.9	708.4	1,407.5
Male	35.5	64.2	108.8	171.7	263.4	383.1	564.5	905.1	1,623.1
Female	8.6	16.7	35.4	67.5	119.1	189.0	324.4	589.3	1,318.0
Non-Hispanic black	32.0	59.9	102.1	157.9	223.5	352.5	508.7	778.1	1,344.4
Male	44.4	85.8	150.5	215.6	289.4	425.3	586.7	868.0	1,417.0
Female	21.6	38.8	64.6	115.0	174.3	301.3	458.9	729.9	1,314.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Other acute ischemic heart diseases (I24)

All origins 3/	0.5	1.0	1.6	2.3	3.1	4.3	5.5	9.3	21.7
Male	0.7	1.4	2.6	3.1	4.4	5.7	7.5	10.7	21.4
Female	0.2	0.6	0.7	1.5	2.0	3.3	4.1	8.5	21.9
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	0.0	*	*	*
Puerto Rican	0.0	*	*	*	0.0	*	*	0.0	*
Male	0.0	*	0.0	0.0	0.0	0.0	*	0.0	0.0
Female	0.0	*	*	*	0.0	*	0.0	0.0	*
Cuban	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	0.0	0.0	*	*	*	*	*	0.0	*
Male	0.0	0.0	*	*	*	0.0	*	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other acute ischemic heart diseases (I24)

Non-Hispanic 5/	0.5	1.0	1.7	2.3	3.3	4.5	5.7	9.7	22.4
Male	0.8	1.4	2.7	3.3	4.6	6.0	7.6	11.1	22.4
Female	0.3	0.6	0.8	1.5	2.1	3.4	4.2	8.8	22.4
Non-Hispanic white	0.4	0.9	1.5	2.2	3.2	4.3	5.5	9.4	22.9
Male	0.6	1.3	2.4	3.0	4.4	5.6	7.6	10.6	23.1
Female	*	0.5	0.7	1.5	2.1	3.2	4.0	8.7	22.9
Non-Hispanic black	1.6	2.4	3.4	4.0	4.3	7.5	9.1	14.9	17.7
Male	2.7	3.2	6.1	6.9	6.8	10.1	10.0	17.8	*
Female	*	*	*	*	*	5.7	8.5	13.4	17.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 5/	27.6	52.2	88.1	146.4	239.1	392.6	657.8	1,166.5	2,935.0
Male	44.5	81.2	133.5	214.1	343.9	551.9	884.8	1,509.5	3,119.1
Female	11.4	24.7	46.0	85.5	149.9	267.2	495.8	959.6	2,857.9
Non-Hispanic white	25.4	47.4	80.4	136.0	228.1	380.5	648.3	1,168.8	3,001.5
Male	41.7	74.6	123.7	200.0	333.9	541.0	881.9	1,529.5	3,215.9
Female	9.3	21.0	39.3	76.9	136.1	252.2	480.1	950.4	2,912.6
Non-Hispanic black	48.1	99.4	166.3	250.8	364.6	559.0	833.2	1,252.1	2,501.3
Male	71.7	152.1	246.2	371.3	482.8	724.2	1,007.7	1,438.7	2,396.5
Female	28.3	56.8	104.4	161.4	276.4	442.6	721.9	1,152.1	2,544.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 5/	9.4	17.9	29.1	43.8	59.8	87.5	131.8	222.5	568.5
Male	15.4	28.6	45.9	65.5	87.1	120.1	169.4	263.8	565.5
Female	3.7	7.7	13.5	24.4	36.5	61.7	105.0	197.6	569.8
Non-Hispanic white	8.1	15.2	24.5	37.2	51.8	78.0	121.1	213.2	564.1
Male	13.5	24.8	39.4	56.0	76.6	108.1	156.2	253.2	562.2
Female	2.8	6.0	10.4	20.0	30.4	54.1	95.8	189.0	564.9
Non-Hispanic black	20.2	41.6	70.9	101.4	132.7	189.3	265.4	359.7	678.2
Male	32.1	66.2	112.0	156.1	194.0	260.2	340.3	438.9	670.5
Female	10.3	21.6	39.1	60.8	86.9	139.4	217.5	317.3	681.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 5/	18.2	34.3	59.0	102.6	179.3	305.2	526.0	944.0	2,366.5
Male	29.0	52.6	87.7	148.7	256.8	431.7	715.5	1,245.7	2,553.6
Female	7.8	17.0	32.5	61.1	113.3	205.5	390.8	762.0	2,288.1
Non-Hispanic white	17.3	32.2	55.9	98.7	176.2	302.4	527.2	955.6	2,437.4
Male	28.3	49.9	84.3	144.1	257.4	433.0	725.7	1,276.3	2,653.7
Female	6.5	15.1	28.9	56.9	105.8	198.1	384.3	761.4	2,347.7
Non-Hispanic black	27.8	57.8	95.4	149.4	231.9	369.7	567.9	892.4	1,823.0
Male	39.6	85.8	134.2	215.2	288.9	464.1	667.4	999.8	1,726.0
Female	18.0	35.1	65.3	100.5	189.4	303.2	504.4	834.9	1,863.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

Non-Hispanic 5/	18.9	27.5	43.8	69.6	109.2	181.5	302.4	572.1	1,543.7
Male	24.4	35.4	57.1	88.4	139.0	223.7	353.0	650.5	1,531.5
Female	13.5	20.1	31.5	52.7	83.9	148.3	266.2	524.8	1,548.9
Non-Hispanic white	15.2	23.3	37.7	62.7	102.4	173.8	295.1	571.1	1,580.8
Male	20.3	30.1	49.6	79.9	130.9	215.9	346.3	650.8	1,565.9
Female	10.2	16.6	26.5	46.8	77.6	140.2	258.2	522.8	1,587.0
Non-Hispanic black	47.5	67.5	101.7	140.2	185.4	289.6	428.3	664.1	1,329.6
Male	60.3	91.0	137.0	186.7	236.3	346.1	482.3	760.3	1,376.8
Female	36.7	48.5	74.4	105.7	147.3	249.8	393.8	612.6	1,310.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Acute and subacute endocarditis (I33)

All origins 3/	0.4	0.5	0.7	0.7	1.1	1.6	2.1	2.8	2.9
Male	0.5	0.7	1.0	0.8	1.2	1.7	2.7	3.9	4.5
Female	0.2	0.2	0.5	0.7	1.0	1.4	1.7	2.1	2.1
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	0.0
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	0.0	*	*	*	*	*
Female	*	0.0	*	*	*	0.0	0.0	0.0	0.0
Puerto Rican	0.0	*	0.0	*	*	*	*	0.0	0.0
Male	0.0	*	0.0	*	*	*	*	0.0	0.0
Female	0.0	0.0	0.0	*	0.0	*	*	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Male	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Other Hispanic 4/	0.0	0.0	*	0.0	*	*	0.0	*	0.0
Male	0.0	0.0	*	0.0	0.0	*	0.0	0.0	0.0
Female	0.0	0.0	*	0.0	*	0.0	0.0	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Heart failure (I50)									
Non-Hispanic 5/	1.6	3.1	6.8	12.2	24.5	50.0	96.6	208.9	695.4
Male	2.1	3.9	8.4	14.9	29.3	60.8	109.0	228.8	660.0
Female	1.2	2.2	5.4	9.7	20.4	41.6	87.7	196.8	710.2
Non-Hispanic white	1.2	2.4	5.8	11.3	23.0	48.5	94.7	209.4	715.8
Male	1.5	3.1	7.4	13.8	27.3	60.0	107.9	230.4	678.0
Female	0.9	1.7	4.3	9.0	19.2	39.3	85.1	196.7	731.5
Non-Hispanic black	5.1	9.2	17.2	23.1	42.2	75.5	133.9	231.1	564.0
Male	6.9	12.3	20.6	29.4	51.5	82.2	144.3	243.0	572.4
Female	3.6	6.6	14.5	18.4	35.3	70.8	127.2	224.7	560.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 5/	16.6	23.7	35.7	56.1	82.9	129.0	202.5	358.9	843.3
Male	21.5	30.5	47.0	72.0	107.8	160.3	240.1	416.2	864.4
Female	11.9	17.3	25.3	41.7	61.7	104.4	175.6	324.4	834.5
Non-Hispanic white	13.5	20.3	30.9	50.2	77.7	123.0	197.2	357.3	859.8
Male	18.1	26.3	40.8	64.7	101.6	153.4	234.2	414.8	880.3
Female	9.0	14.4	21.5	36.7	57.0	98.8	170.5	322.6	851.3
Non-Hispanic black	40.8	56.2	81.4	114.6	139.7	209.8	290.7	429.0	761.7
Male	51.4	75.1	112.0	154.7	181.7	259.2	336.5	511.9	803.3
Female	31.9	41.0	57.7	84.9	108.3	175.0	261.4	384.5	744.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic 5/	1.7	2.9	4.2	7.4	12.1	18.9	32.8	60.4	153.3
Male	2.1	3.6	5.3	8.3	13.4	20.3	32.3	58.6	128.6
Female	1.3	2.2	3.3	6.6	11.0	17.7	33.1	61.4	163.6
Non-Hispanic white	1.0	1.6	2.7	4.9	9.0	15.1	27.3	53.2	143.1
Male	1.3	2.0	3.6	5.7	10.2	16.5	27.2	51.3	117.6
Female	0.7	1.1	1.8	4.2	7.9	14.0	27.4	54.3	153.7
Non-Hispanic black	6.7	14.0	17.9	28.1	38.0	58.8	97.6	148.5	275.5
Male	8.5	18.7	21.1	32.9	42.9	61.7	96.8	157.6	252.2
Female	5.2	10.2	15.4	24.6	34.4	56.7	98.2	143.6	285.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Cerebrovascular diseases (I60-I69)

Non-Hispanic 5/	12.5	18.8	32.2	52.7	92.2	177.1	345.7	683.1	1,640.9
Male	13.1	21.1	36.3	61.3	104.2	200.3	374.6	713.4	1,486.3
Female	11.9	16.7	28.4	44.9	82.0	158.9	325.1	664.8	1,705.6
Non-Hispanic white	8.9	14.3	25.9	43.7	83.1	163.9	333.4	675.4	1,663.7
Male	9.1	15.7	29.2	50.5	93.6	185.2	360.1	707.1	1,506.6
Female	8.6	12.9	22.8	37.5	74.0	146.8	314.2	656.2	1,728.9
Non-Hispanic black	36.4	54.6	82.3	123.5	169.9	315.2	502.1	807.3	1,519.4
Male	41.9	67.9	100.3	160.7	203.3	362.3	554.9	828.9	1,366.7
Female	31.9	43.8	68.4	95.9	145.0	282.1	468.3	795.8	1,582.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Atherosclerosis (I70)

All origins 3/	0.3	0.8	1.3	3.0	5.1	11.3	24.1	54.3	201.0
Male	0.4	1.1	1.8	3.9	6.4	13.4	27.6	59.6	181.8
Female	0.2	0.5	0.9	2.2	4.1	9.6	21.7	51.0	209.0
Hispanic	*	*	*	*	*	8.0	17.1	34.5	97.3
Male	*	*	*	*	*	10.3	20.0	38.3	110.2
Female	*	*	*	*	*	*	14.9	32.0	90.7
Mexican	*	*	*	*	*	9.9	21.5	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	0.0	*	*	*	*
Female	0.0	0.0	*	0.0	*	*	*	*	*
Cuban	*	0.0	*	0.0	*	*	*	*	*
Male	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	*	0.0	*	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	0.0	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Atherosclerosis (I70)

Non-Hispanic 5/	0.3	0.8	1.3	3.0	5.3	11.5	24.4	55.0	205.2
Male	0.4	1.1	1.8	3.9	6.5	13.6	28.0	60.3	185.1
Female	*	0.5	0.9	2.3	4.2	9.8	21.9	51.7	213.7
Non-Hispanic white	0.3	0.7	1.3	2.9	5.2	11.6	24.5	56.0	211.8
Male	0.4	1.0	1.7	3.8	6.4	13.5	27.6	61.1	191.0
Female	*	0.4	0.9	2.1	4.1	10.0	22.3	52.9	220.5
Non-Hispanic black	*	1.4	2.1	4.9	6.9	12.3	28.1	49.7	157.7
Male	*	*	3.5	6.3	8.8	15.5	37.0	57.0	156.2
Female	*	*	*	4.0	5.4	10.0	22.4	45.8	158.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic 5/	2.3	3.5	7.0	13.1	24.5	41.7	65.3	97.0	167.9
Male	3.2	4.8	9.9	19.1	35.0	58.8	90.3	130.4	203.7
Female	1.4	2.2	4.2	7.6	15.5	28.3	47.5	76.9	152.9
Non-Hispanic white	2.0	3.1	6.6	13.2	24.8	42.5	66.0	97.4	168.3
Male	2.8	4.2	9.6	19.5	36.1	60.7	92.5	132.2	208.8
Female	1.1	2.0	3.7	7.3	14.9	28.0	46.9	76.2	151.5
Non-Hispanic black	4.9	6.7	11.8	14.2	26.0	39.4	66.0	103.1	182.2
Male	6.7	9.7	15.2	19.2	31.7	48.5	74.0	119.3	172.9
Female	3.4	4.3	9.2	10.5	21.7	33.0	60.9	94.5	186.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Aortic aneurysm and dissection (I71)

Non-Hispanic 5/	1.5	2.2	4.5	8.5	17.3	28.6	43.5	60.1	82.2
Male	2.4	3.5	7.2	13.7	26.6	42.6	64.4	89.8	123.8
Female	0.7	1.0	2.1	3.9	9.4	17.5	28.6	42.2	64.8
Non-Hispanic white	1.3	2.0	4.4	8.9	18.0	29.7	45.0	61.9	84.5
Male	2.0	3.1	7.1	14.3	28.0	44.9	67.7	92.9	129.4
Female	0.5	0.9	1.8	3.9	9.3	17.6	28.6	43.1	65.8
Non-Hispanic black	3.8	4.4	6.6	6.9	14.0	20.2	30.3	43.2	61.8
Male	5.8	7.2	9.6	10.2	19.0	24.9	29.4	55.2	72.5
Female	2.1	2.1	4.3	4.5	10.3	16.9	30.9	36.7	57.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 5/	0.7	1.3	2.4	4.5	7.2	13.2	21.8	36.9	85.7
Male	0.8	1.3	2.8	5.4	8.4	16.1	25.9	40.6	79.9
Female	0.7	1.2	2.1	3.7	6.1	10.9	18.9	34.7	88.1
Non-Hispanic white	0.7	1.1	2.2	4.3	6.8	12.8	21.0	35.5	83.8
Male	0.9	1.2	2.5	5.2	8.1	15.8	24.8	39.3	79.5
Female	0.6	1.1	1.8	3.4	5.6	10.3	18.3	33.2	85.6
Non-Hispanic black	1.1	2.3	5.2	7.3	12.0	19.2	35.7	59.9	120.4
Male	*	*	5.6	9.0	12.7	23.6	44.6	64.1	100.4
Female	*	2.1	4.9	6.0	11.5	16.1	30.0	57.7	128.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other disorders of circulatory system (I80-I99)

All origins 3/	1.0	1.1	1.6	2.3	3.3	4.3	6.8	12.8	23.3
Male	1.1	1.2	1.7	2.4	3.8	4.8	7.4	13.1	20.0
Female	1.0	0.9	1.5	2.1	2.9	3.9	6.4	12.7	24.7
Hispanic	1.2	*	*	*	*	5.5	*	*	18.5
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	18.3
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	0.0	*	*
Male	*	*	0.0	*	0.0	*	0.0	*	*
Female	0.0	*	*	*	*	*	0.0	*	*
Cuban	*	*	0.0	*	0.0	*	*	*	*
Male	*	*	0.0	*	0.0	*	0.0	*	0.0
Female	0.0	*	0.0	0.0	0.0	*	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	0.0	*	*
Male	*	*	*	*	*	*	0.0	*	*
Female	*	*	*	*	*	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other disorders of circulatory system (I80-I99)

Non-Hispanic 5/	1.0	1.1	1.7	2.3	3.4	4.3	7.0	13.0	23.4
Male	1.1	1.2	1.8	2.4	3.9	4.8	7.5	13.3	20.0
Female	1.0	0.9	1.6	2.2	2.9	3.8	6.6	12.8	24.9
Non-Hispanic white	0.9	0.9	1.6	2.1	3.0	4.1	6.7	12.6	23.2
Male	0.9	1.0	1.7	2.3	3.3	4.6	7.1	12.8	19.9
Female	0.8	0.8	1.5	1.9	2.7	3.6	6.4	12.5	24.6
Non-Hispanic black	2.5	2.5	2.2	4.1	6.9	6.5	11.8	18.9	28.8
Male	2.9	3.3	*	*	9.4	6.7	14.7	21.4	21.2
Female	2.2	*	*	4.3	5.0	6.4	10.0	17.7	31.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Influenza and pneumonia (J10-J18)

Non-Hispanic 5/	4.2	5.5	8.6	14.5	25.2	51.6	107.4	239.7	758.8
Male	5.3	6.9	10.0	17.2	32.1	65.3	133.3	303.2	875.2
Female	3.1	4.1	7.4	12.1	19.3	40.8	88.9	201.4	710.1
Non-Hispanic white	3.4	4.7	7.8	13.1	24.2	49.7	106.8	240.9	774.2
Male	4.2	5.7	8.8	15.3	30.3	61.9	131.8	303.6	892.4
Female	2.6	3.7	6.9	11.1	18.8	40.0	88.8	203.0	725.2
Non-Hispanic black	10.3	12.7	16.8	27.4	36.3	77.0	121.9	233.3	642.2
Male	14.3	18.9	23.2	36.4	51.5	107.8	159.5	313.4	754.2
Female	6.9	7.6	11.9	20.8	25.0	55.3	97.8	190.4	595.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Influenza (J10-J11)

All origins 3/	*	*	0.2	0.4	0.6	1.2	2.1	6.2	21.1
Male	*	*	*	0.5	0.6	1.2	2.3	7.2	19.2
Female	*	*	*	*	0.6	1.2	1.9	5.5	21.9
Hispanic	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	*	0.0	*	*	*	*
Mexican	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	*	0.0	*	*	0.0	0.0	*
Female	0.0	0.0	0.0	*	0.0	0.0	*	*	*
Puerto Rican	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Other Hispanic 4/	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	*	*	*	*	*	0.0
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Influenza (J10-J11)

Non-Hispanic 5/	*	*	0.2	0.4	0.6	1.2	2.1	6.3	21.8
Male	*	*	*	0.5	0.6	1.2	2.4	7.5	20.1
Female	*	*	*	*	0.7	1.2	1.9	5.6	22.5
Non-Hispanic white	*	*	0.2	0.4	0.7	1.2	2.3	6.6	23.5
Male	*	*	*	0.5	0.6	1.2	2.6	7.8	22.1
Female	*	*	*	*	0.8	1.2	2.0	5.9	24.0
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	*	*	0.0	*	*	*	*	*	*
Female	0.0	*	*	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Pneumonia (J12-J18)

Non-Hispanic 5/	4.1	5.4	8.4	14.1	24.5	50.4	105.3	233.4	737.1
Male	5.2	6.9	9.8	16.7	31.5	64.2	130.9	295.7	855.0
Female	3.0	4.1	7.1	11.8	18.6	39.6	87.0	195.8	687.7
Non-Hispanic white	3.3	4.6	7.6	12.7	23.5	48.5	104.6	234.3	750.8
Male	4.1	5.6	8.5	14.7	29.7	60.7	129.2	295.8	870.3
Female	2.6	3.7	6.6	10.8	18.1	38.8	86.9	197.1	701.2
Non-Hispanic black	10.2	12.6	16.6	27.2	36.1	75.6	120.9	230.2	636.4
Male	14.2	18.8	23.2	36.1	51.0	106.5	159.0	308.9	750.9
Female	6.9	7.5	11.5	20.6	25.0	53.9	96.6	188.0	588.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute lower respiratory infections (J20-J22)

Non-Hispanic 5/	*	*	*	*	*	0.2	0.6	1.4	6.1
Male	*	*	*	*	*	*	0.8	1.7	4.0
Female	*	*	*	*	*	*	0.5	1.2	7.0
Non-Hispanic white	*	*	*	*	*	*	0.7	1.4	6.5
Male	*	*	*	*	*	*	0.8	1.7	4.0
Female	*	*	*	*	*	*	0.5	1.2	7.5
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	0.0	0.0	*	0.0	*	*	0.0	*	*
Female	*	*	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Acute bronchitis and bronchiolitis (J20-J21)

Non-Hispanic 5/	*	*	*	*	*	*	0.4	0.9	3.7
Male	*	*	*	*	*	*	*	1.1	2.5
Female	*	*	*	*	*	*	*	0.8	4.1
Non-Hispanic white	*	*	*	*	*	*	0.4	0.9	3.9
Male	*	*	*	*	*	*	*	*	2.6
Female	*	*	*	*	*	*	*	0.8	4.4
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	0.0	0.0	*	0.0	*	*	0.0	0.0	*
Female	*	*	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 5/	5.8	12.9	34.1	71.1	137.5	238.6	350.0	501.8	652.2
Male	6.0	13.0	36.8	79.6	159.5	289.3	437.1	657.6	974.9
Female	5.5	12.7	31.7	63.4	118.7	198.6	287.8	407.8	517.0
Non-Hispanic white	5.1	12.6	34.8	74.4	145.5	250.1	363.8	522.3	682.0
Male	5.3	12.4	36.7	81.2	165.4	298.3	446.7	674.9	1,014.3
Female	4.9	12.8	32.9	68.2	128.2	211.6	304.1	429.8	544.1
Non-Hispanic black	12.0	18.6	38.2	60.5	99.9	165.9	249.8	314.7	381.5
Male	12.9	21.8	47.1	85.7	138.6	237.6	384.5	527.0	663.8
Female	11.2	15.9	31.3	41.8	71.0	115.4	163.9	200.8	264.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Bronchitis, chronic and unspecified (J40-J42)

All origins 3/	*	0.1	0.3	0.6	0.7	1.3	1.9	4.2	10.7
Male	*	*	0.4	0.5	0.8	1.5	2.4	4.6	11.4
Female	*	*	*	0.6	0.5	1.1	1.5	3.9	10.4
Hispanic	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	0.0	0.0	0.0	*	0.0	*	*	*	*
Mexican	0.0	0.0	0.0	*	0.0	*	*	*	*
Male	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Female	0.0	0.0	0.0	*	0.0	*	*	*	*
Puerto Rican	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Cuban	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*
Male	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
Other Hispanic 4/	0.0	0.0	0.0	*	*	0.0	*	0.0	*
Male	0.0	0.0	0.0	0.0	*	0.0	*	0.0	*
Female	0.0	0.0	0.0	*	0.0	0.0	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Emphysema (J43)

All origins 3/	0.7	2.0	5.6	12.3	23.8	38.9	51.4	66.2	61.8
Male	0.9	2.3	6.4	13.9	28.4	49.4	66.3	91.0	101.4
Female	0.5	1.8	4.9	10.9	19.9	30.6	40.7	51.2	45.0
Hispanic	*	*	*	*	4.6	9.4	15.9	28.1	33.3
Male	*	*	*	*	*	15.5	22.7	44.7	56.7
Female	*	*	*	*	*	*	10.9	17.6	21.5
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	0.0	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*
Puerto Rican	*	0.0	*	*	*	*	*	*	*
Male	*	0.0	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	*	*	*	0.0	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	0.0	0.0	*	*	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	*	*	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Emphysema (J43)									
Non-Hispanic 5/	0.7	2.1	6.0	13.1	25.0	40.5	53.0	67.7	62.9
Male	0.9	2.4	6.8	14.5	29.7	51.2	68.3	92.9	103.4
Female	0.6	1.9	5.3	11.7	21.1	32.2	42.1	52.5	45.9
Non-Hispanic white	0.7	2.1	6.3	14.1	27.2	43.3	55.6	71.3	65.8
Male	0.8	2.2	6.9	15.2	31.4	53.2	70.8	95.9	107.9
Female	0.5	2.0	5.8	13.0	23.5	35.3	44.7	56.4	48.3
Non-Hispanic black	1.5	2.7	4.9	8.5	14.0	21.3	32.4	33.5	38.3
Male	2.2	4.6	7.0	13.0	22.6	36.4	48.9	66.8	69.2
Female	*	*	3.2	5.2	7.6	10.7	21.8	15.7	25.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

Non-Hispanic 5/	1.6	1.8	2.7	2.3	3.6	4.7	6.1	9.7	17.4
Male	1.0	1.2	1.9	1.8	2.6	3.9	4.7	7.1	12.3
Female	2.2	2.3	3.4	2.7	4.5	5.4	7.1	11.3	19.5
Non-Hispanic white	1.1	1.2	2.0	1.7	2.9	3.9	5.7	9.4	16.6
Male	0.6	0.8	1.4	1.3	1.9	3.0	4.1	7.0	10.4
Female	1.6	1.7	2.5	2.2	3.8	4.7	6.8	10.8	19.2
Non-Hispanic black	5.1	6.2	7.8	6.5	8.4	11.3	10.2	12.1	20.9
Male	3.9	4.7	5.6	6.3	6.8	11.8	9.5	*	*
Female	6.2	7.5	9.5	6.7	9.7	10.9	10.6	15.3	22.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic 5/	3.4	8.8	25.2	55.1	108.1	192.0	288.9	420.3	561.2
Male	4.0	9.3	27.7	62.7	126.4	232.7	361.6	552.9	847.6
Female	2.7	8.4	22.8	48.3	92.5	159.9	237.0	340.3	441.2
Non-Hispanic white	3.2	9.1	26.1	58.0	114.7	201.6	300.6	437.5	588.3
Male	3.8	9.3	28.0	64.1	131.2	240.6	369.4	567.6	883.8
Female	2.6	9.0	24.4	52.3	100.4	170.5	251.1	358.7	465.7
Non-Hispanic black	5.1	9.2	25.3	45.1	77.0	131.8	205.1	264.6	317.7
Male	6.6	12.0	34.1	66.0	108.9	187.4	323.3	445.1	573.5
Female	4.0	6.8	18.5	29.6	53.2	92.6	129.7	167.9	211.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Pneumoconioses and chemical effects (J60-J66,J68)

Non-Hispanic 5/	*	*	0.2	0.4	1.2	2.0	3.6	5.5	6.3
Male	*	*	0.3	0.8	2.4	4.6	8.3	14.0	20.6
Female	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Non-Hispanic white	*	*	0.2	0.3	1.2	2.1	3.8	5.8	6.5
Male	*	*	0.4	0.7	2.5	4.8	8.7	14.9	21.4
Female	0.0	0.0	0.0	0.0	*	0.0	*	*	*
Non-Hispanic black	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Pneumonitis due to solids and liquids (J69)

All origins 3/	0.7	0.9	2.0	3.1	6.2	13.4	28.2	61.5	174.7
Male	0.9	1.0	2.4	4.3	8.5	18.5	41.0	89.5	248.6
Female	0.5	0.9	1.6	2.0	4.2	9.3	19.1	44.6	143.5
Hispanic	*	*	*	*	3.4	7.4	16.2	30.0	98.3
Male	*	*	*	*	*	8.9	22.7	39.6	116.5
Female	*	*	*	*	*	*	11.4	24.0	89.1
Mexican	*	*	*	*	*	*	18.3	*	*
Male	*	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*	*
Puerto Rican	*	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	*	*	0.0	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*
Cuban	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	*	0.0	0.0	*	*	*	*	*
Female	0.0	0.0	0.0	*	*	*	*	*	*
Other Hispanic 4/	0.0	0.0	0.0	*	*	*	*	*	*
Male	0.0	0.0	0.0	0.0	*	*	*	*	*
Female	0.0	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Pneumonitis due to solids and liquids (J69)

Non-Hispanic 5/	0.7	1.0	2.1	3.2	6.3	13.7	28.7	62.7	177.9
Male	0.9	1.1	2.5	4.4	8.7	19.0	41.8	91.5	254.9
Female	0.5	0.9	1.7	2.0	4.3	9.5	19.4	45.2	145.7
Non-Hispanic white	0.6	0.9	1.9	3.0	6.0	12.9	28.1	62.1	181.0
Male	0.8	0.9	2.3	4.1	8.1	17.7	40.6	90.4	262.4
Female	0.4	0.9	1.6	2.0	4.2	9.0	19.1	45.0	147.2
Non-Hispanic black	1.3	1.9	4.1	5.3	10.0	24.2	41.6	75.8	170.5
Male	*	*	5.6	8.8	15.3	36.7	64.1	112.2	215.3
Female	*	*	3.0	*	6.0	15.4	27.3	56.3	151.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 5/	2.7	4.2	7.1	12.8	22.7	37.7	59.9	89.6	148.7
Male	2.8	4.6	7.5	14.8	27.8	47.4	74.2	115.0	178.1
Female	2.6	3.8	6.6	11.0	18.4	30.0	49.8	74.2	136.3
Non-Hispanic white	2.3	3.8	6.5	12.6	22.9	37.8	61.0	91.3	152.2
Male	2.3	4.3	6.6	14.7	28.2	48.1	75.9	117.6	182.7
Female	2.2	3.3	6.3	10.7	18.2	29.6	50.3	75.3	139.5
Non-Hispanic black	6.2	7.7	12.5	16.1	24.8	39.0	54.6	70.8	114.2
Male	7.2	8.6	15.0	19.2	28.1	43.5	66.0	85.5	137.2
Female	5.4	7.1	10.5	13.8	22.3	35.9	47.3	63.0	104.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Peptic ulcer (K25-K28)

All origins 3/	0.6	0.9	1.4	2.2	3.3	6.0	9.1	17.3	34.5
Male	0.7	1.3	1.8	3.1	4.5	8.0	10.8	21.0	36.1
Female	0.5	0.5	1.1	1.5	2.3	4.4	7.8	15.1	33.8
Hispanic	*	*	*	*	*	4.3	5.7	11.8	20.1
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	17.5
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	*	*	0.0	0.0	*
Cuban	0.0	0.0	0.0	*	0.0	*	*	*	*
Male	0.0	0.0	0.0	*	0.0	*	*	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*	*
Female	*	0.0	*	0.0	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Peptic ulcer (K25-K28)									
Non-Hispanic 5/	0.6	0.9	1.4	2.3	3.4	6.0	9.2	17.5	35.1
Male	0.7	1.3	1.8	3.1	4.6	8.1	10.8	21.1	36.7
Female	0.5	0.5	1.1	1.6	2.3	4.4	8.0	15.4	34.4
Non-Hispanic white	0.5	0.8	1.3	2.2	3.2	5.9	9.2	17.5	35.9
Male	0.6	1.1	1.6	2.9	4.2	8.0	11.0	21.2	37.8
Female	0.4	0.4	1.1	1.6	2.3	4.3	8.0	15.3	35.1
Non-Hispanic black	1.3	2.0	2.6	3.3	5.0	6.5	7.8	17.4	25.8
Male	2.0	3.2	4.1	5.6	7.5	8.8	*	18.7	26.8
Female	*	*	*	*	*	5.0	7.3	16.7	25.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Diseases of appendix (K35-K38)

All origins 3/	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5
Male	*	0.2	*	*	*	0.5	0.9	1.0	2.8
Female	*	*	*	*	*	0.4	*	1.1	0.9
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	0.0	*	0.0	*
Female	*	0.0	0.0	*	0.0	*	*	*	0.0
Mexican	0.0	*	*	*	*	0.0	*	*	0.0
Male	0.0	*	*	0.0	*	0.0	*	0.0	0.0
Female	0.0	0.0	0.0	*	0.0	0.0	*	*	0.0
Puerto Rican	*	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	0.0	0.0	0.0	*	*	0.0	0.0	0.0
Male	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of appendix (K35-K38)

Non-Hispanic 5/	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5
Male	*	*	*	*	*	0.5	0.9	1.1	2.9
Female	*	*	*	*	*	*	*	1.1	1.0
Non-Hispanic white	*	0.2	*	*	0.3	0.4	0.6	1.1	1.6
Male	*	*	*	*	*	*	0.9	*	3.0
Female	*	*	*	*	*	*	*	1.1	1.0
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Hernia (K40-K46)

All origins 3/	*	0.2	0.3	0.7	0.8	1.6	2.5	5.5	12.7
Male	*	0.3	0.3	0.8	0.8	1.5	2.5	6.4	12.7
Female	*	0.2	*	0.6	0.8	1.6	2.6	4.9	12.7
Hispanic	*	*	0.0	*	*	*	*	*	*
Male	*	*	0.0	*	*	*	*	*	*
Female	0.0	*	0.0	*	*	*	*	*	*
Mexican	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	0.0	*	0.0	*	*	*	*	*	*
Puerto Rican	0.0	0.0	0.0	0.0	0.0	*	0.0	*	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Female	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	*
Cuban	0.0	0.0	0.0	*	0.0	*	*	*	*
Male	0.0	0.0	0.0	*	0.0	*	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	*	*	0.0	0.0	*	*	0.0	*	*
Male	*	*	0.0	0.0	0.0	*	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 5/	15.8	18.1	20.7	25.7	28.5	30.6	31.5	30.5	22.3
Male	24.0	28.1	31.0	35.9	40.1	39.7	38.2	37.7	30.8
Female	7.9	8.7	11.0	16.6	18.5	23.3	26.8	26.1	18.7
Non-Hispanic white	14.8	17.5	19.7	25.1	28.7	31.3	32.4	31.5	22.8
Male	22.5	27.0	29.5	35.3	40.3	40.6	39.0	39.2	32.1
Female	7.1	8.3	10.4	15.8	18.7	23.9	27.7	26.8	19.0
Non-Hispanic black	23.7	24.6	29.4	32.4	27.8	25.4	23.7	20.2	17.0
Male	36.6	40.4	47.1	46.1	43.2	36.1	31.8	25.8	*
Female	12.9	11.9	15.7	22.2	16.3	17.8	18.5	17.2	16.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Alcoholic liver disease (K70)

Non-Hispanic 5/	9.5	10.5	11.0	12.1	11.0	8.9	6.9	5.2	2.6
Male	14.5	16.6	17.4	19.0	18.7	15.0	11.1	9.2	5.4
Female	4.7	4.8	5.0	5.9	4.5	4.1	3.8	2.7	1.4
Non-Hispanic white	9.1	10.2	10.4	11.7	10.9	8.9	6.7	5.1	2.4
Male	14.0	16.1	16.4	18.6	18.2	15.0	10.8	9.3	5.0
Female	4.2	4.5	4.7	5.4	4.5	4.1	3.8	2.6	1.3
Non-Hispanic black	13.3	13.7	15.5	16.4	13.0	10.3	8.3	5.9	*
Male	20.2	22.3	26.9	25.2	24.2	17.9	14.2	*	*
Female	7.6	6.8	6.8	9.8	4.7	5.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 5/	6.2	7.6	9.7	13.6	17.4	21.6	24.7	25.3	19.7
Male	9.4	11.5	13.6	16.9	21.5	24.7	27.1	28.5	25.5
Female	3.2	4.0	6.0	10.6	14.0	19.2	22.9	23.4	17.3
Non-Hispanic white	5.7	7.3	9.3	13.4	17.9	22.4	25.7	26.4	20.4
Male	8.6	11.0	13.0	16.7	22.1	25.6	28.1	29.9	27.0
Female	2.8	3.8	5.7	10.4	14.2	19.9	23.9	24.2	17.7
Non-Hispanic black	10.4	10.9	13.8	16.0	14.8	15.0	15.3	14.3	12.4
Male	16.4	18.1	20.2	20.8	19.0	18.2	17.6	*	*
Female	5.3	5.0	8.9	12.4	11.6	12.8	13.9	13.4	13.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Cholelithiasis and other disorders of gallbladder (K80-K82)

All origins 3/	0.3	0.4	0.7	1.0	1.8	3.5	5.7	11.0	23.8
Male	0.3	0.4	0.9	1.3	2.1	4.2	6.8	13.2	26.6
Female	0.3	0.5	0.6	0.8	1.6	2.9	4.9	9.7	22.6
Hispanic	*	*	*	*	*	4.1	8.5	13.3	24.3
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	24.7
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	*	0.0	*	0.0	0.0	*	*	*	0.0
Male	0.0	0.0	*	0.0	0.0	0.0	*	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	*	*	*	0.0
Cuban	*	0.0	0.0	0.0	*	*	*	*	*
Male	*	0.0	0.0	0.0	*	0.0	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Other Hispanic 4/	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	0.0	*	*	*	*	*	*
Female	0.0	*	*	*	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic 5/	0.3	0.4	0.7	1.0	1.8	3.4	5.5	10.9	23.7
Male	0.3	0.4	0.9	1.2	2.0	4.2	6.5	13.1	26.8
Female	0.2	0.5	0.6	0.8	1.6	2.7	4.8	9.5	22.4
Non-Hispanic white	0.2	0.4	0.6	0.8	1.6	3.3	5.6	10.9	24.2
Male	0.3	0.3	0.7	1.1	1.8	4.2	6.6	13.7	27.3
Female	*	0.4	0.6	0.6	1.3	2.6	4.8	9.2	22.9
Non-Hispanic black	*	*	*	2.1	3.2	4.2	4.8	12.1	15.7
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	13.4	16.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 5/	3.1	5.3	9.0	15.8	28.5	47.5	78.2	130.4	270.9
Male	3.8	6.0	9.8	17.7	31.4	55.8	98.0	170.5	361.5
Female	2.5	4.7	8.2	14.2	25.9	41.0	64.1	106.2	232.9
Non-Hispanic white	1.9	3.5	6.3	11.8	22.8	40.7	70.9	120.8	257.8
Male	2.2	3.8	7.3	13.4	26.0	49.5	90.0	161.1	352.9
Female	1.5	3.1	5.3	10.4	20.0	33.7	57.2	96.4	218.3
Non-Hispanic black	12.6	21.4	30.6	49.4	77.6	119.2	168.5	260.6	442.0
Male	16.7	26.3	31.7	59.1	80.6	128.4	206.5	327.6	499.8
Female	9.1	17.5	29.7	42.3	75.3	112.8	144.2	224.7	418.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All origins 3/	*	*	*	*	*	0.3	0.5	0.5	0.8
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	0.5	*	0.7
Hispanic	*	0.0	0.0	*	*	*	*	0.0	*
Male	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	*	0.0	0.0	*	*	*	*	0.0	0.0
Mexican	0.0	0.0	0.0	*	*	*	0.0	0.0	*
Male	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	*	*	0.0	0.0	0.0
Puerto Rican	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Non-Hispanic 5/	*	*	*	*	*	0.3	0.5	0.5	0.8
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	0.5	*	0.7
Non-Hispanic white	*	*	*	*	*	0.3	0.5	0.5	0.8
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	0.7
Non-Hispanic black	*	*	*	*	0.0	*	*	*	*
Male	*	*	0.0	*	0.0	*	*	*	*
Female	*	*	*	*	0.0	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

Non-Hispanic 5/	3.0	5.1	8.7	15.4	28.0	46.5	76.2	127.7	266.7
Male	3.7	5.7	9.6	17.3	30.9	54.8	95.5	166.5	355.2
Female	2.4	4.5	7.9	13.7	25.5	40.0	62.4	104.2	229.6
Non-Hispanic white	1.7	3.3	6.1	11.5	22.4	39.7	69.0	118.4	254.0
Male	2.1	3.6	7.2	13.1	25.5	48.6	87.6	157.6	347.2
Female	1.4	2.9	5.1	10.1	19.7	32.6	55.6	94.6	215.3
Non-Hispanic black	12.4	20.7	29.9	48.3	76.3	117.8	165.7	256.0	433.8
Male	16.5	25.4	31.1	57.9	79.6	126.7	203.2	321.4	487.6
Female	9.0	16.8	29.0	41.3	73.9	111.6	141.8	220.9	411.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other disorders of kidney (N25,N27)

Non-Hispanic 5/	0.0	*	*	*	*	*	*	*	*
Male	0.0	*	0.0	0.0	*	*	0.0	*	*
Female	0.0	*	*	*	*	*	*	*	*
Non-Hispanic white	0.0	*	0.0	*	*	*	*	*	*
Male	0.0	*	0.0	0.0	0.0	*	0.0	*	*
Female	0.0	*	0.0	*	*	*	*	*	*
Non-Hispanic black	0.0	*	*	*	*	0.0	*	*	*
Male	0.0	0.0	0.0	0.0	*	0.0	0.0	*	0.0
Female	0.0	*	*	*	0.0	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Infections of kidney (N10-N12,N13.6,N15.1)

All origins 3/	0.1	0.2	0.2	0.3	0.6	0.9	1.5	2.8	7.4
Male	*	*	*	*	0.4	0.6	1.2	2.5	6.5
Female	*	0.3	*	*	0.6	1.1	1.7	2.9	7.8
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	0.0	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Mexican	*	0.0	*	*	*	*	*	*	*
Male	*	0.0	*	*	*	*	*	*	*
Female	*	0.0	0.0	*	0.0	*	*	*	*
Puerto Rican	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Cuban	0.0	0.0	*	0.0	0.0	*	0.0	*	0.0
Male	0.0	0.0	0.0	0.0	0.0	*	0.0	*	0.0
Female	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	*	*	*	0.0	*	0.0	*	*
Male	*	0.0	*	*	0.0	0.0	0.0	0.0	0.0
Female	0.0	*	*	0.0	0.0	*	0.0	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Hyperplasia of prostate (N40)

All origins 3/	0.0	*	*	*	*	0.4	0.8	2.1	5.0
Male	0.0	*	*	*	*	0.8	1.9	5.6	16.9
Hispanic	0.0	0.0	*	0.0	*	*	0.0	*	*
Male	0.0	0.0	*	0.0	*	*	0.0	*	*
Mexican	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*
Male	0.0	0.0	0.0	0.0	*	0.0	0.0	*	*
Puerto Rican	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Male	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0
Other Hispanic 4/	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Hyperplasia of prostate (N40)

Non-Hispanic 5/	0.0	0.0	*	*	*	0.4	0.8	2.1	5.2
Male	0.0	0.0	*	*	*	0.8	2.0	5.6	17.5
Non-Hispanic white	0.0	0.0	*	*	*	0.4	0.9	2.1	5.3
Male	0.0	0.0	*	*	*	0.8	2.1	5.5	18.0
Non-Hispanic black	0.0	0.0	*	*	*	*	*	*	*
Male	0.0	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Inflammatory diseases of female pelvic organs (N70-N76)

All origins 3/	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Hispanic	*	*	*	0.0	0.0	0.0	*	0.0	*
Female	*	*	*	0.0	0.0	0.0	*	0.0	*
Mexican	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Female	*	0.0	0.0	0.0	0.0	0.0	*	0.0	*
Puerto Rican	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	*
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Inflammatory diseases of female pelvic organs (N70-N76)

Non-Hispanic 5/	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Non-Hispanic white	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Non-Hispanic black	*	0.0	0.0	*	*	*	*	*	*
Female	*	0.0	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/	1.8	1.9	2.4	2.4	2.7	2.5	3.4	5.0	7.8
Male	1.9	2.0	2.5	2.5	2.4	2.7	3.3	5.6	8.2
Female	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.7	7.6
Hispanic	1.6	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Mexican	1.9	*	*	*	*	*	*	*	*
Male	*	*	*	0.0	*	*	*	*	*
Female	*	*	*	*	*	*	0.0	*	*
Puerto Rican	*	*	*	*	*	*	0.0	*	0.0
Male	*	0.0	*	0.0	0.0	*	0.0	0.0	0.0
Female	*	*	*	*	*	*	0.0	*	0.0
Cuban	*	*	*	*	*	*	0.0	*	*
Male	0.0	*	0.0	0.0	*	*	0.0	*	0.0
Female	*	0.0	*	*	*	*	0.0	0.0	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	0.0	*	0.0	*
Female	0.0	0.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 5/	6.1	6.2	7.3	9.3	12.3	18.2	27.2	52.3	192.1
Male	8.4	8.4	9.6	12.4	15.7	22.6	31.5	55.3	165.2
Female	4.0	4.1	5.1	6.6	9.5	14.8	24.1	50.5	203.4
Non-Hispanic white	5.3	5.3	6.4	8.6	11.5	17.6	26.1	51.1	195.6
Male	7.0	7.1	8.3	11.4	14.5	21.8	30.3	53.4	168.0
Female	3.6	3.5	4.7	5.9	8.9	14.3	23.1	49.6	207.1
Non-Hispanic black	13.7	14.0	15.5	17.7	21.6	28.0	44.0	77.7	184.2
Male	21.0	20.3	23.7	25.5	29.4	35.4	53.6	89.9	169.6
Female	7.6	8.8	9.2	11.9	15.7	22.8	37.9	71.1	190.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All origins 3/	30.4	39.6	51.9	73.4	108.7	169.8	281.8	517.7	1,403.5
Male	36.9	47.0	60.8	82.7	118.8	185.9	293.6	524.7	1,179.0
Female	24.0	32.5	43.5	65.0	100.1	157.2	273.4	513.5	1,498.4
Hispanic	26.4	31.8	42.4	59.9	83.8	126.6	178.5	314.0	783.4
Male	36.2	40.5	52.7	73.5	91.6	147.5	188.3	317.0	673.7
Female	16.8	23.8	33.3	48.4	77.6	110.5	171.2	312.1	838.8
Mexican	24.8	33.5	41.1	62.4	83.1	133.7	177.8	293.5	*
Male	32.5	39.4	48.6	74.6	88.6	148.4	*	*	*
Female	16.7	27.2	33.8	51.0	78.3	121.9	185.2	*	*
Puerto Rican	50.3	47.0	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	22.8	25.7	40.6	55.5	87.3	131.2	*	*	*
Male	32.9	36.0	*	*	*	*	*	*	*
Female	*	18.0	30.7	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

Non-Hispanic 5/	30.5	40.0	52.3	74.0	110.0	171.9	286.3	525.7	1,429.2
Male	36.7	47.2	61.1	82.8	120.1	187.5	298.3	533.0	1,202.8
Female	24.5	33.1	44.2	66.1	101.4	159.6	277.7	521.3	1,524.1
Non-Hispanic white	26.2	35.5	48.0	69.7	105.2	167.3	281.6	525.8	1,457.6
Male	32.1	42.3	55.9	77.7	114.3	182.3	290.8	530.7	1,216.8
Female	20.5	28.9	40.5	62.4	97.3	155.3	274.9	522.8	1,557.5
Non-Hispanic black	65.4	82.2	96.7	122.5	166.5	244.2	379.6	593.3	1,310.7
Male	78.3	99.8	119.8	144.5	188.5	267.2	431.5	641.9	1,223.9
Female	54.5	67.9	78.8	106.2	150.0	227.9	346.6	567.2	1,346.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 5/	34.5	29.8	30.6	31.7	38.8	52.0	79.8	137.2	286.8
Male	51.9	44.4	44.2	45.2	53.2	69.1	102.6	182.8	369.1
Female	17.6	15.9	18.0	19.7	26.6	38.6	63.4	109.6	252.3
Non-Hispanic white	32.6	27.9	29.0	29.7	37.7	50.5	79.0	138.6	296.1
Male	48.3	41.1	41.0	42.1	50.9	66.4	100.4	184.6	379.9
Female	17.0	15.2	17.7	18.3	26.2	37.8	63.6	110.8	261.3
Non-Hispanic black	53.2	47.3	45.5	47.9	50.5	68.1	92.3	132.3	211.4
Male	88.5	79.8	77.5	75.0	77.2	100.8	133.9	178.9	299.0
Female	23.5	21.0	20.7	27.7	30.5	45.1	65.7	107.3	175.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Transport accidents (V01-V99,Y85)

Non-Hispanic 5/	16.2	15.2	16.5	16.1	18.0	21.1	26.1	32.7	32.4
Male	23.5	22.0	23.4	22.3	24.7	27.6	35.8	47.8	62.7
Female	9.1	8.8	10.0	10.4	12.3	16.0	19.2	23.6	19.7
Non-Hispanic white	15.7	14.8	16.1	15.6	17.8	21.0	26.1	33.0	32.8
Male	22.5	21.2	22.6	21.4	24.3	27.1	35.5	47.9	63.4
Female	9.0	8.5	9.8	10.2	12.2	16.1	19.4	24.0	20.1
Non-Hispanic black	20.7	19.5	19.7	18.4	17.9	21.3	25.1	29.8	28.1
Male	33.5	31.8	31.7	30.6	26.8	31.3	39.9	49.0	64.7
Female	10.0	9.5	10.5	9.3	11.3	14.2	15.8	19.6	13.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic 5/	14.2	13.4	14.5	14.1	16.3	19.5	24.3	30.7	30.7
Male	20.3	19.0	20.0	19.1	21.9	24.9	33.3	44.8	59.1
Female	8.3	8.1	9.3	9.7	11.5	15.3	17.9	22.3	18.8
Non-Hispanic white	13.7	12.9	14.0	13.7	16.2	19.4	24.4	31.1	31.2
Male	19.3	18.2	19.2	18.3	21.5	24.5	32.9	45.0	60.0
Female	8.1	7.8	9.1	9.4	11.5	15.4	18.3	22.7	19.2
Non-Hispanic black	18.8	17.4	18.0	16.3	16.4	19.9	23.3	27.3	25.8
Male	30.3	28.1	28.4	26.4	24.4	28.3	38.4	45.4	58.0
Female	9.2	8.8	9.9	8.8	10.5	14.0	13.6	17.7	12.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic 5/	1.1	1.1	1.2	1.2	0.9	0.9	0.5	0.6	*
Male	1.8	1.9	2.1	2.1	1.7	1.7	0.9	*	*
Female	0.4	0.4	*	*	*	*	*	*	*
Non-Hispanic white	1.1	1.1	1.2	1.1	1.0	0.9	0.5	0.5	*
Male	1.9	2.0	2.1	2.0	1.8	1.7	1.0	*	*
Female	0.4	0.4	*	*	*	*	*	*	*
Non-Hispanic black	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	0.0	0.0	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Nontransport accidents (W00-X59,Y86)									
Non-Hispanic 5/	18.3	14.5	14.1	15.7	20.8	30.9	53.7	104.5	254.4
Male	28.5	22.3	20.7	22.8	28.5	41.5	66.9	135.1	306.4
Female	8.5	7.1	8.0	9.3	14.3	22.6	44.3	86.0	232.7
Non-Hispanic white	16.9	13.2	13.0	14.1	19.8	29.5	52.9	105.6	263.3
Male	25.8	19.9	18.3	20.7	26.6	39.2	65.0	136.7	316.6
Female	8.0	6.7	7.9	8.1	14.0	21.7	44.2	86.8	241.2
Non-Hispanic black	32.5	27.8	25.7	29.5	32.6	46.8	67.1	102.5	183.2
Male	55.0	48.0	45.8	44.5	50.4	69.4	94.0	130.0	234.3
Female	13.6	11.5	10.2	18.4	19.2	30.9	50.0	87.8	162.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Falls (W00-W19)

All origins 3/	2.3	2.3	3.1	4.7	7.0	11.4	22.3	46.1	109.7
Male	3.9	3.7	4.8	7.1	10.0	15.4	27.5	61.2	137.2
Female	0.8	1.0	1.5	2.4	4.5	8.3	18.6	37.0	98.0
Hispanic	2.1	2.5	3.2	4.4	7.7	7.6	15.7	30.0	65.0
Male	3.4	4.6	6.4	7.9	10.3	11.3	19.4	39.6	88.1
Female	*	*	*	*	5.6	*	12.9	24.0	53.3
Mexican	2.0	*	*	*	9.2	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	0.0	*	0.0	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	0.0	0.0	0.0	0.0	*	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	0.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Falls (W00-W19)									
Non-Hispanic 5/	2.3	2.3	3.0	4.6	6.9	11.6	22.5	46.6	111.4
Male	3.8	3.6	4.6	7.0	9.8	15.6	27.7	61.8	139.3
Female	0.8	1.0	1.6	2.5	4.4	8.5	18.8	37.4	99.7
Non-Hispanic white	2.2	2.1	2.8	4.3	6.9	11.7	23.3	48.8	117.7
Male	3.6	3.3	4.2	6.5	9.6	15.6	28.2	64.5	146.9
Female	0.8	0.9	1.6	2.3	4.5	8.6	19.7	39.2	105.5
Non-Hispanic black	3.4	3.9	4.6	6.9	8.9	9.8	13.5	24.8	45.8
Male	6.0	6.6	8.0	10.9	14.8	14.5	19.5	32.1	54.7
Female	*	*	*	4.0	4.5	6.4	9.7	21.0	42.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Accidental discharge of firearms (W32-W34)

Non-Hispanic 5/	0.3	0.3	0.2	0.2	*	*	*	*	*
Male	0.5	0.4	0.4	0.4	*	*	*	*	*
Female	*	*	*	*	*	0.0	*	*	*
Non-Hispanic white	0.3	0.3	0.2	0.2	*	*	*	*	*
Male	0.5	0.4	*	*	*	*	*	*	*
Female	*	*	*	*	*	0.0	*	*	*
Non-Hispanic black	*	*	*	*	*	*	*	*	0.0
Male	*	*	*	*	*	*	*	*	0.0
Female	*	0.0	*	*	0.0	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Accidental drowning and submersion (W65-W74)

All origins 3/	1.0	1.0	0.9	0.9	0.9	1.2	1.2	1.5	1.6
Male	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.6	2.3
Female	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4
Hispanic	*	*	*	*	*	*	*	*	0.0
Male	*	*	*	*	*	*	*	0.0	0.0
Female	*	0.0	*	0.0	0.0	0.0	*	*	0.0
Mexican	*	*	*	*	*	*	*	0.0	0.0
Male	*	*	*	*	*	*	*	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Puerto Rican	*	*	0.0	*	*	0.0	0.0	0.0	0.0
Male	*	*	0.0	*	*	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	*	0.0	*	*	0.0	0.0	*	0.0
Male	*	*	0.0	*	*	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	*	*	*	0.0	0.0	*	*	0.0	0.0
Male	*	*	*	0.0	0.0	*	*	0.0	0.0
Female	*	0.0	*	0.0	0.0	0.0	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.8	5.9
Male	1.5	1.4	1.6	1.9	2.6	3.4	4.2	6.6	8.9
Female	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.7	4.6
Hispanic	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	0.0
Mexican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	0.0	*	0.0	*	*	0.0
Puerto Rican	0.0	0.0	0.0	*	*	0.0	*	0.0	*
Male	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	*	*	0.0	*	0.0	0.0
Cuban	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	*	0.0	*	*	*	*	*	*
Male	*	*	0.0	*	*	0.0	*	*	*
Female	*	0.0	0.0	0.0	0.0	*	*	*	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic 5/	1.1	1.1	1.3	1.5	2.1	2.6	3.6	4.8	6.0
Male	1.5	1.4	1.7	1.8	2.6	3.6	4.2	6.6	9.0
Female	0.6	0.7	0.9	1.3	1.7	1.9	3.1	3.7	4.8
Non-Hispanic white	0.9	0.9	1.2	1.2	1.7	2.1	2.8	3.9	4.8
Male	1.3	1.2	1.4	1.6	2.0	2.7	3.2	5.8	7.7
Female	0.5	0.6	0.9	0.8	1.4	1.6	2.6	2.8	3.5
Non-Hispanic black	2.1	2.5	2.8	4.5	5.8	8.5	12.4	16.5	22.2
Male	3.3	3.7	4.4	4.6	7.8	13.1	17.1	20.5	27.9
Female	*	*	*	4.5	4.3	5.2	9.4	14.3	19.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/	10.0	5.6	3.5	2.1	1.7	1.6	1.9	2.6	3.4
Male	15.3	8.1	4.8	3.0	2.3	1.9	2.1	2.9	4.3
Female	4.8	3.2	2.3	1.2	1.2	1.4	1.8	2.5	3.1
Hispanic	11.5	7.8	4.0	2.8	*	*	*	*	*
Male	20.2	12.4	7.6	*	*	*	*	0.0	*
Female	3.1	3.5	*	*	*	*	*	*	*
Mexican	10.5	6.4	4.1	*	*	*	*	*	*
Male	17.4	9.7	*	*	*	*	0.0	0.0	*
Female	*	*	*	*	*	0.0	*	*	0.0
Puerto Rican	*	*	*	*	*	0.0	0.0	0.0	0.0
Male	*	*	*	*	*	0.0	0.0	0.0	0.0
Female	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	*	*	0.0	0.0	*	0.0	0.0	*
Male	*	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	*	0.0	0.0	0.0	*	0.0	0.0	*
Other Hispanic 4/	10.8	8.1	*	*	*	*	*	0.0	0.0
Male	23.4	*	*	*	*	*	*	0.0	0.0
Female	*	*	0.0	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 5/	9.7	5.3	3.4	2.0	1.6	1.6	1.9	2.7	3.5
Male	14.7	7.5	4.5	2.8	2.1	1.9	2.1	3.1	4.3
Female	4.9	3.1	2.3	1.3	1.2	1.4	1.8	2.5	3.1
Non-Hispanic white	8.8	4.5	3.0	1.6	1.5	1.5	2.0	2.7	3.3
Male	13.0	6.1	3.7	2.0	1.9	1.8	2.1	3.1	4.6
Female	4.6	2.9	2.4	1.2	1.2	1.3	1.9	2.5	2.8
Non-Hispanic black	18.9	12.9	6.7	5.5	2.8	3.3	*	*	*
Male	31.5	22.5	13.0	10.2	*	*	*	*	*
Female	8.2	5.1	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 5/	4.0	4.6	5.2	6.4	9.1	13.6	24.3	48.7	131.6
Male	6.2	7.6	8.0	9.3	12.2	18.2	30.4	60.4	150.8
Female	1.8	1.8	2.7	3.7	6.5	10.0	19.9	41.7	123.5
Non-Hispanic white	3.6	4.4	4.8	5.8	8.7	12.9	23.5	48.5	135.6
Male	5.5	7.2	7.2	8.6	11.4	17.1	29.2	60.2	154.4
Female	1.7	1.7	2.5	3.3	6.4	9.6	19.4	41.5	127.8
Non-Hispanic black	7.0	7.1	9.8	11.3	13.9	23.4	37.0	57.2	108.0
Male	11.9	12.0	16.9	17.1	20.8	34.4	50.3	70.3	146.2
Female	2.8	3.0	4.3	7.0	8.7	15.7	28.5	50.1	92.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Intentional self-harm (suicide) (X60-X84,Y87.0)

All origins 3/	14.6	13.8	13.0	11.6	12.6	14.6	17.5	19.5	19.2
Male	22.7	21.3	20.8	19.5	22.4	27.9	35.1	43.8	54.9
Female	6.7	6.7	5.9	4.5	4.2	4.2	4.8	4.7	4.1
Hispanic	7.3	7.1	7.0	6.3	7.5	5.5	10.0	13.8	11.6
Male	12.7	10.7	12.6	11.4	15.0	10.3	20.0	30.7	*
Female	*	3.7	*	*	*	*	*	*	*
Mexican	5.8	5.4	5.1	*	*	*	*	*	*
Male	9.9	7.8	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	0.0	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	0.0	0.0	*	*	0.0
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	9.8	8.5	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/	7.6	7.4	8.2	7.6	8.8	11.0	12.9	13.7	12.6
Male	12.9	12.9	14.3	14.3	17.1	22.6	28.6	34.4	40.0
Female	2.4	2.2	2.6	1.6	1.7	1.9	1.7	1.2	1.0
Hispanic	3.9	2.5	3.1	2.5	3.9	*	6.8	*	*
Male	6.7	4.3	6.0	5.6	8.1	*	16.0	*	*
Female	*	*	*	0.0	*	0.0	0.0	*	0.0
Mexican	3.2	*	*	*	*	*	*	*	*
Male	5.6	*	*	*	*	*	*	*	*
Female	*	*	0.0	0.0	*	0.0	0.0	0.0	0.0
Puerto Rican	*	*	*	*	*	*	*	*	0.0
Male	*	*	*	*	*	*	*	*	0.0
Female	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	0.0	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic 5/	7.9	7.8	8.6	8.0	9.1	11.5	13.2	14.0	12.8
Male	13.4	13.5	14.9	14.9	17.6	23.5	29.1	35.1	41.1
Female	2.6	2.3	2.8	1.8	1.7	2.0	1.7	1.2	1.0
Non-Hispanic white	8.8	8.7	9.5	8.8	9.9	12.5	14.2	14.9	13.9
Male	14.8	14.9	16.1	16.2	19.2	25.3	31.3	37.4	44.9
Female	2.9	2.6	3.2	1.9	1.9	2.2	1.9	1.3	1.0
Non-Hispanic black	3.6	2.6	3.7	3.5	4.0	4.0	5.2	*	*
Male	6.5	5.4	7.8	7.2	7.8	9.8	11.9	*	*
Female	*	*	*	*	*	0.0	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All origins 3/	7.0	6.4	4.8	4.0	3.8	3.6	4.6	5.8	6.6
Male	9.8	8.4	6.5	5.2	5.3	5.3	6.5	9.4	14.9
Female	4.3	4.5	3.3	2.8	2.5	2.3	3.2	3.5	3.1
Hispanic	3.4	4.5	3.9	3.7	3.6	*	*	*	*
Male	6.0	6.4	6.6	5.9	7.0	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Mexican	2.6	3.7	*	*	*	*	*	*	*
Male	4.3	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	0.0	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	*	0.0	0.0	*	*	0.0
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.0	*	*	0.0	0.0	0.0	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*	0.0
Male	*	*	*	*	*	0.0	0.0	*	0.0
Female	*	*	*	*	*	*	*	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Assault (homicide) (X85-Y09,Y87.1)

All origins 3/	5.2	4.1	3.3	2.9	2.7	2.4	2.5	2.5	2.4
Male	7.9	6.0	5.2	4.4	4.2	3.1	3.3	3.4	3.3
Female	2.6	2.2	1.4	1.5	1.5	1.9	2.0	1.9	2.0
Hispanic	8.0	5.1	5.0	2.9	3.3	*	*	*	*
Male	12.2	8.3	8.5	5.6	*	*	*	*	*
Female	3.9	*	*	*	*	*	*	*	*
Mexican	8.4	5.7	4.5	*	*	*	*	*	0.0
Male	12.4	8.9	*	*	*	*	*	*	0.0
Female	*	*	*	*	*	*	*	*	0.0
Puerto Rican	*	*	*	*	*	*	*	0.0	*
Male	*	*	*	*	*	*	*	0.0	0.0
Female	*	*	0.0	0.0	*	*	0.0	0.0	*
Cuban	*	*	*	*	*	*	0.0	*	0.0
Male	*	*	*	*	*	*	0.0	0.0	0.0
Female	*	0.0	*	*	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	7.5	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	0.0	*	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Assault (homicide) (X85-Y09,Y87.1)

Non-Hispanic 5/	4.8	3.9	3.1	2.8	2.7	2.4	2.6	2.5	2.4
Male	7.4	5.7	4.9	4.2	4.0	3.1	3.3	3.3	3.3
Female	2.4	2.2	1.4	1.6	1.5	1.9	2.0	1.9	2.0
Non-Hispanic white	3.1	2.8	2.3	2.2	2.3	2.0	2.0	2.1	2.1
Male	4.5	3.9	3.4	3.0	3.1	2.4	2.4	2.8	2.6
Female	1.7	1.7	1.2	1.4	1.5	1.7	1.7	1.6	1.9
Non-Hispanic black	17.0	11.9	9.3	7.8	6.6	6.1	8.5	6.5	*
Male	29.2	20.2	18.4	13.9	12.7	10.1	13.8	*	*
Female	6.8	5.3	*	3.3	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Assault (homicide) by discharge of firearms (X93-X95)

All origins 3/	2.8	2.1	1.6	1.6	1.1	1.0	0.8	0.7	0.6
Male	4.5	3.1	2.7	2.6	1.6	1.5	1.1	*	*
Female	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.7	*
Hispanic	4.8	2.5	2.5	*	*	*	*	*	*
Male	7.1	4.3	*	*	*	*	*	*	*
Female	2.6	*	*	*	*	0.0	0.0	*	0.0
Mexican	5.0	*	*	*	*	*	*	0.0	0.0
Male	7.0	*	*	*	*	*	*	0.0	0.0
Female	*	*	*	*	*	0.0	0.0	0.0	0.0
Puerto Rican	*	*	0.0	*	*	0.0	0.0	0.0	0.0
Male	*	*	0.0	*	*	0.0	0.0	0.0	0.0
Female	*	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
Cuban	*	*	*	*	*	*	0.0	*	0.0
Male	*	*	*	*	*	*	0.0	0.0	0.0
Female	*	0.0	*	*	0.0	0.0	0.0	*	0.0
Other Hispanic 4/	*	*	*	*	*	0.0	*	*	*
Male	*	*	*	*	0.0	0.0	*	*	*
Female	*	*	*	0.0	*	0.0	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Assault (homicide) by discharge of firearms (X93-X95)

Non-Hispanic 5/	2.6	2.0	1.5	1.6	1.1	1.1	0.8	0.7	0.6
Male	4.2	3.0	2.6	2.5	1.6	1.5	1.0	*	*
Female	1.1	1.1	0.5	0.7	0.6	0.8	0.7	0.7	*
Non-Hispanic white	1.6	1.4	1.1	1.2	1.0	0.9	0.6	0.7	0.6
Male	2.4	2.1	1.7	1.7	1.3	1.2	*	*	*
Female	0.9	0.7	0.5	0.7	0.7	0.7	0.6	*	*
Non-Hispanic black	10.1	6.4	4.8	4.4	2.2	2.6	*	*	*
Male	18.3	10.6	10.2	9.0	*	*	*	*	*
Female	3.2	3.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All origins 3/	2.3	2.0	1.7	1.3	1.7	1.4	1.7	1.8	1.8
Male	3.4	2.9	2.5	1.8	2.7	1.7	2.2	2.6	2.5
Female	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.3	1.5
Hispanic	3.2	2.5	2.6	*	*	*	*	*	*
Male	5.1	4.0	4.4	*	*	*	*	*	*
Female	*	*	*	0.0	0.0	*	*	*	*
Mexican	3.4	*	*	*	*	*	*	*	0.0
Male	5.4	*	*	*	*	*	*	*	0.0
Female	*	*	*	0.0	0.0	*	*	*	0.0
Puerto Rican	*	*	*	0.0	*	*	*	0.0	*
Male	*	*	*	0.0	*	*	*	0.0	0.0
Female	*	*	0.0	0.0	0.0	*	0.0	0.0	*
Cuban	*	0.0	*	*	*	*	0.0	0.0	0.0
Male	*	0.0	*	*	*	*	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Hispanic 4/	*	*	*	*	*	*	0.0	*	*
Male	*	*	*	*	*	*	0.0	*	*
Female	*	*	*	0.0	0.0	*	0.0	0.0	0.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

Non-Hispanic 5/	2.2	1.9	1.6	1.2	1.6	1.4	1.7	1.8	1.8
Male	3.2	2.7	2.3	1.7	2.4	1.6	2.2	2.6	2.5
Female	1.3	1.1	0.8	0.9	0.9	1.2	1.4	1.3	1.5
Non-Hispanic white	1.5	1.4	1.2	1.0	1.3	1.1	1.4	1.4	1.6
Male	2.1	1.8	1.7	1.3	1.8	1.2	1.8	2.2	1.8
Female	0.9	1.0	0.7	0.7	0.8	1.0	1.1	1.0	1.5
Non-Hispanic black	6.9	5.5	4.5	3.4	4.3	3.5	5.4	*	*
Male	10.9	9.5	8.2	4.9	8.3	*	*	*	*
Female	3.6	2.2	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	2.9	1.6	1.1	0.8	0.8	0.5	0.7	1.0	1.4
Male	3.9	2.2	1.5	1.1	1.1	0.6	1.0	1.4	2.0
Female	1.9	1.1	0.7	0.6	0.5	*	0.5	0.7	1.1
Non-Hispanic white	2.7	1.5	1.0	0.8	0.8	0.4	0.7	1.0	1.4
Male	3.4	1.8	1.2	1.0	1.1	*	1.0	1.3	2.1
Female	1.9	1.1	0.8	0.7	0.5	*	0.5	0.8	1.1
Non-Hispanic black	5.0	3.6	1.6	*	*	*	*	*	*
Male	8.8	6.4	3.5	*	*	*	*	*	*
Female	1.8	*	*	0.0	*	*	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85 YEARS AND OVER

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	2.8	1.6	1.0	0.7	0.7	0.4	0.6	0.8	1.3
Male	3.8	2.1	1.4	1.0	0.9	*	0.8	*	1.7
Female	1.8	1.1	0.6	0.6	0.5	*	0.5	0.7	1.1
Non-Hispanic white	2.6	1.4	1.0	0.8	0.7	0.4	0.6	0.8	1.3
Male	3.3	1.7	1.2	0.8	0.9	*	0.8	*	*
Female	1.9	1.0	0.7	0.7	0.5	*	*	0.8	1.1
Non-Hispanic black	4.8	3.6	1.5	*	*	*	*	0.0	*
Male	8.4	6.4	*	*	*	*	*	0.0	*
Female	1.8	*	*	0.0	*	*	*	0.0	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY
 RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY
 RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 5/	0.5	0.8	1.1	1.7	2.5	4.2	5.6	8.8	12.9
Male	0.4	0.8	1.3	1.8	2.4	4.8	6.4	10.3	16.4
Female	0.6	0.8	0.9	1.6	2.6	3.8	5.1	7.8	11.5
Non-Hispanic white	0.4	0.7	0.9	1.6	2.4	4.0	5.4	8.4	12.3
Male	0.3	0.6	1.0	1.6	2.3	4.5	6.3	9.9	15.6
Female	0.5	0.7	0.8	1.5	2.5	3.6	4.8	7.4	10.9
Non-Hispanic black	0.9	1.9	2.8	2.7	3.4	7.4	9.6	13.0	20.6
Male	*	*	*	*	*	9.1	10.0	*	22.3
Female	*	*	*	*	*	6.2	9.4	11.9	19.9

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ ALL ORIGINS INCLUDES ORIGIN NOT STATED; SPECIFIED ORIGINS EXCLUDE ORIGIN NOT STATED.

4/ INCLUDES CENTRAL AND SOUTH AMERICAN AND OTHER AND UNKNOWN HISPANIC.

5/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY