

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other bacterial food-borne intoxications (A05)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Amebiasis (A06)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Intestinal infections due to other specified organisms (A04,A07-A08)									
2002.....	0.8	0.8	0.8	0.8	0.9	0.8	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
2000.....	0.5	0.5	0.4	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Diarrhea and gastroenteritis of infectious origin (A09)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Tuberculosis (A16-A19)									
2002.....	0.3	0.4	0.2	0.2	0.3	0.1	0.7	1.1	0.5
2001.....	0.3	0.4	0.2	0.2	0.2	0.1	0.8	1.2	0.5
2000.....	0.3	0.4	0.2	0.2	0.2	0.1	0.9	1.5	0.6
1999.....	0.3	0.5	0.2	0.2	0.3	0.1	0.9	1.5	0.6
Respiratory tuberculosis (A16)									
2002.....	0.2	0.3	0.1	0.2	0.2	0.1	0.6	0.9	0.4
2001.....	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3
2000.....	0.2	0.3	0.1	0.1	0.2	0.1	0.7	1.3	0.4
1999.....	0.3	0.4	0.2	0.2	0.3	0.1	0.8	1.2	0.5
Other tuberculosis (A17-A19)									
2002.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.1
2000.....	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.2
1999.....	0.1	0.1	0.0	0.0	0.1	0.0	0.2	0.3	*
Zoonotic and other bacterial diseases (A20-A49)									
2002.....	12.3	13.5	11.4	11.2	12.4	10.4	25.1	28.8	22.8
2001.....	12.0	13.4	11.0	10.9	12.2	10.0	24.5	28.8	22.0
2000.....	11.9	13.1	11.0	10.7	11.8	9.9	25.5	29.2	23.2
1999.....	12.0	13.4	11.0	10.7	11.9	9.9	26.0	30.6	23.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Whooping cough (A37)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)									
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2000.....	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Septicemia (A40-A41)									
2002.....	11.7	12.9	10.9	10.7	11.7	9.9	24.2	27.8	22.1
2001.....	11.4	12.7	10.5	10.3	11.5	9.5	23.6	27.7	21.2
2000.....	11.3	12.4	10.5	10.2	11.2	9.4	24.6	28.0	22.5
1999.....	11.3	12.5	10.4	10.0	11.1	9.3	24.6	28.9	21.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Yellow fever (A95)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other and unspecified arthropod-borne viral and hemorrhagic fevers (A90-A94,A96-A99)									
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Herpesviral (herpes simplex) infections (B00)									
2002.....	0.0	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Zoster (herpes zoster)(B02)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Smallpox (B03)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Measles (B05)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Rubella (German measles) (B06)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)									
2002.....	2.0	2.7	1.3	1.9	2.6	1.2	2.8	3.8	1.9
2001.....	2.0	2.6	1.3	1.8	2.5	1.2	2.7	3.4	2.1
2000.....	1.9	2.6	1.3	1.8	2.4	1.2	2.7	3.8	1.8
1999.....	1.8	2.4	1.2	1.6	2.2	1.1	2.6	3.4	1.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease (B20-B24)									
2002.....	4.9	7.4	2.5	2.6	4.3	0.9	22.5	33.3	13.4
2001.....	5.0	7.5	2.5	2.6	4.4	0.9	22.8	33.8	13.4
2000.....	5.2	7.9	2.5	2.8	4.6	1.0	23.3	35.1	13.2
1999.....	5.3	8.2	2.5	2.9	4.9	1.0	23.6	36.1	13.1
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)									
2002.....	1.9	2.9	1.0	1.1	1.7	0.4	8.6	12.8	5.1
2001.....	1.9	2.9	1.0	1.1	1.8	0.4	8.6	12.6	5.1
2000.....	2.0	3.1	1.0	1.1	1.8	0.4	9.0	13.4	5.2
1999.....	2.1	3.3	1.1	1.2	1.9	0.4	9.3	14.1	5.2
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)									
2002.....	0.3	0.5	0.1	0.2	0.4	0.0	1.2	1.8	0.6
2001.....	0.3	0.6	0.1	0.2	0.4	0.1	1.2	2.0	0.6
2000.....	0.4	0.6	0.1	0.3	0.5	0.0	1.3	2.1	0.5
1999.....	0.4	0.7	0.1	0.3	0.5	0.1	1.3	2.2	0.5
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)									
2002.....	0.5	0.7	0.2	0.3	0.5	0.1	1.8	2.8	1.0
2001.....	0.5	0.8	0.2	0.3	0.5	0.1	2.1	3.1	1.2
2000.....	0.6	0.9	0.2	0.3	0.6	0.1	2.3	3.6	1.2
1999.....	0.6	0.9	0.2	0.4	0.6	0.1	2.3	3.6	1.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)									
2002.....	1.2	1.7	0.7	0.5	0.8	0.2	6.2	9.1	3.8
2001.....	1.2	1.8	0.7	0.6	0.9	0.2	6.2	9.1	3.8
2000.....	1.2	1.8	0.7	0.6	0.9	0.2	6.0	8.9	3.7
1999.....	1.2	1.8	0.7	0.6	0.9	0.2	6.4	9.5	3.7
Unspecified human immunodeficiency virus (HIV) disease (B24)									
2002.....	1.0	1.4	0.5	0.5	0.8	0.2	4.6	6.8	2.9
2001.....	1.0	1.5	0.5	0.5	0.8	0.1	4.7	7.0	2.8
2000.....	1.0	1.5	0.5	0.5	0.8	0.2	4.7	7.1	2.6
1999.....	1.0	1.5	0.5	0.5	0.9	0.2	4.4	6.6	2.5
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)									
2002.....	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.4
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)									
2002.....	1.3	1.7	0.9	1.2	1.6	0.9	1.8	2.5	1.3
2001.....	1.1	1.3	0.9	1.0	1.2	0.8	1.7	2.1	1.4
2000.....	1.1	1.3	0.8	1.0	1.2	0.8	1.7	2.2	1.3
1999.....	1.5	1.8	1.2	1.3	1.6	1.1	2.9	3.3	2.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Schistosomiasis (bilharziasis) (B65)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other and unspecified helminthiasis (B66-B83)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	0.0	*	0.0	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Sequelae of tuberculosis (B90)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Sequelae of poliomyelitis (B91)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)									
2002.....	0.6	0.8	0.4	0.6	0.8	0.4	0.7	1.0	0.6
2001.....	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.6	0.5
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.7	0.4
1999.....	0.5	0.6	0.5	0.4	0.5	0.4	1.2	1.4	1.1
II. Neoplasms (C00-D48)									
2002.....	198.1	244.6	167.0	196.4	241.0	166.4	243.2	324.7	194.4
2001.....	200.9	249.8	168.8	198.8	245.4	168.1	247.8	336.5	195.5
2000.....	204.5	255.0	171.8	202.2	250.1	171.2	253.1	345.7	197.9
1999.....	205.6	258.0	171.7	202.9	252.7	170.5	257.3	351.4	201.3
Malignant neoplasms (C00-C97)									
2002.....	193.5	238.9	163.1	191.7	235.2	162.4	238.8	319.6	190.3
2001.....	196.0	243.7	164.7	193.9	239.2	163.9	243.1	330.9	191.3
2000.....	199.6	248.9	167.6	197.2	243.9	166.9	248.5	340.3	193.8
1999.....	200.8	251.9	167.6	197.9	246.5	166.4	252.5	346.1	196.9
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
2002.....	2.7	4.1	1.5	2.5	3.8	1.5	3.6	6.3	1.7
2001.....	2.7	4.1	1.6	2.6	3.9	1.5	3.9	6.7	1.8
2000.....	2.7	4.1	1.6	2.5	3.7	1.6	4.1	7.3	1.8
1999.....	2.7	4.1	1.6	2.5	3.8	1.5	4.3	7.5	2.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of lip (C00)									
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Of tongue (C01-C02)									
2002.....	0.7	0.9	0.4	0.6	0.9	0.4	0.7	1.3	0.3
2001.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2000.....	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.3
1999.....	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.4
Of pharynx (C10-C13,C14.0)									
2002.....	1.1	1.7	0.6	1.0	1.5	0.5	1.8	3.3	0.8
2001.....	1.1	1.8	0.6	1.0	1.6	0.5	2.0	3.6	0.9
2000.....	1.1	1.8	0.6	1.0	1.5	0.5	2.2	4.1	0.9
1999.....	1.1	1.8	0.6	1.0	1.6	0.5	2.3	4.0	1.0
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)									
2002.....	0.9	1.3	0.6	0.9	1.3	0.6	1.0	1.7	0.5
2001.....	0.9	1.3	0.6	0.9	1.3	0.6	1.2	1.9	0.6
2000.....	0.9	1.3	0.6	0.9	1.3	0.6	1.1	1.9	0.5
1999.....	0.9	1.3	0.6	0.9	1.3	0.6	1.2	2.0	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of digestive organs (C15-C26)									
2002.....	45.7	58.8	35.6	44.0	56.6	34.1	63.4	82.9	50.1
2001.....	46.4	59.4	36.5	44.6	57.2	34.9	64.8	83.3	52.1
2000.....	47.2	60.5	37.2	45.3	58.1	35.6	66.2	86.2	52.5
1999.....	47.3	60.7	37.3	45.4	58.4	35.7	66.7	87.1	52.7
Of esophagus (C15)									
2002.....	4.4	7.8	1.7	4.3	7.7	1.6	5.9	10.1	2.9
2001.....	4.4	7.8	1.8	4.3	7.6	1.7	6.4	11.3	3.0
2000.....	4.4	7.7	1.8	4.3	7.6	1.7	6.5	11.0	3.3
1999.....	4.4	7.7	1.8	4.2	7.4	1.6	6.5	11.5	3.1
Of stomach (C16)									
2002.....	4.2	5.9	3.0	3.7	5.2	2.5	8.4	11.9	6.1
2001.....	4.4	6.1	3.1	3.8	5.4	2.7	8.3	11.9	5.8
2000.....	4.6	6.4	3.3	4.0	5.6	2.8	9.0	12.7	6.5
1999.....	4.6	6.6	3.3	4.1	5.8	2.8	9.2	13.5	6.4
Of small intestine (C17)									
2002.....	0.4	0.4	0.3	0.3	0.4	0.3	0.6	0.7	0.5
2001.....	0.4	0.5	0.3	0.4	0.4	0.3	0.6	0.8	0.5
2000.....	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of colon, rectum and anus (C18-C21)									
2002.....	19.7	23.7	16.7	19.2	23.1	16.2	26.8	33.1	22.8
2001.....	20.1	24.2	17.2	19.6	23.7	16.6	27.6	33.1	24.1
2000.....	20.8	25.1	17.7	20.3	24.5	17.2	28.2	34.9	24.0
1999.....	20.9	25.3	17.8	20.4	24.8	17.3	28.4	34.2	24.7
Colon (C18)									
2002.....	16.5	19.7	14.2	16.0	19.1	13.7	23.2	28.3	19.9
2001.....	16.9	20.1	14.7	16.4	19.6	14.1	24.0	28.4	21.3
2000.....	17.6	21.0	15.1	17.1	20.4	14.7	24.6	29.9	21.3
1999.....	17.7	21.2	15.3	17.3	20.8	14.7	24.8	29.2	22.0
Rectosigmoid junction and rectum (C19-C20)									
2002.....	3.0	3.9	2.3	3.0	3.8	2.3	3.4	4.6	2.7
2001.....	3.0	4.0	2.3	3.0	4.0	2.3	3.4	4.5	2.7
2000.....	3.0	4.0	2.3	3.0	3.9	2.3	3.4	4.7	2.5
1999.....	3.0	3.9	2.3	3.0	3.9	2.4	3.4	4.8	2.5
Anus and anus canal (C21)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of liver and intrahepatic bile ducts (C22)									
2002.....	4.9	7.1	3.0	4.5	6.5	2.8	6.3	9.9	3.6
2001.....	4.7	6.9	3.0	4.3	6.3	2.7	6.2	9.1	4.1
2000.....	4.7	6.8	3.0	4.2	6.2	2.7	6.1	9.5	3.6
1999.....	4.5	6.5	3.0	4.2	5.9	2.7	5.9	8.7	3.8
Liver (C22.0,C22.2-C22.9)									
2002.....	3.8	5.9	2.1	3.5	5.3	1.9	5.4	8.8	2.8
2001.....	3.7	5.7	2.1	3.3	5.2	1.9	5.3	8.0	3.3
2000.....	3.7	5.7	2.1	3.3	5.0	1.9	5.3	8.6	2.8
1999.....	3.6	5.4	2.1	3.2	4.9	1.9	5.1	7.9	3.1
Intrahepatic bile duct carcinoma (C22.1)									
2002.....	1.0	1.2	0.9	1.0	1.2	0.9	0.9	1.1	0.8
2001.....	1.0	1.2	0.9	1.0	1.2	0.9	0.9	1.1	0.8
2000.....	1.0	1.1	0.9	1.0	1.2	0.9	0.8	0.9	0.8
1999.....	0.9	1.0	0.9	0.9	1.1	0.9	0.8	0.8	0.7
Of gallbladder and extrahepatic bile ducts (C23-C24)									
2002.....	1.2	1.1	1.3	1.1	1.1	1.2	1.4	1.1	1.5
2001.....	1.3	1.2	1.4	1.3	1.1	1.4	1.3	1.2	1.5
2000.....	1.3	1.2	1.4	1.3	1.2	1.4	1.2	1.0	1.4
1999.....	1.3	1.2	1.4	1.3	1.2	1.4	1.3	1.0	1.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of pancreas (C25)									
2002.....	10.5	12.1	9.2	10.3	12.0	9.0	13.6	15.3	12.3
2001.....	10.5	12.1	9.3	10.4	12.0	9.1	13.6	15.0	12.5
2000.....	10.6	12.2	9.3	10.4	12.0	9.0	14.1	15.8	12.7
1999.....	10.6	12.3	9.3	10.4	12.0	9.1	14.1	16.4	12.3
Of other and ill-defined digestive organs (C26)									
2002.....	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.8	0.5
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.8	0.6
2000.....	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.7	0.6
1999.....	0.6	0.7	0.5	0.6	0.7	0.5	0.7	0.9	0.5
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)									
2002.....	56.5	76.0	42.3	56.8	75.1	43.3	64.7	100.3	41.2
2001.....	57.0	78.1	41.8	57.1	76.9	42.8	65.4	104.7	39.9
2000.....	57.8	79.7	42.1	57.8	78.4	43.0	67.2	107.1	41.1
1999.....	57.2	79.8	41.1	57.0	78.4	41.8	67.7	108.5	41.1
Of nasal cavity, middle ear and accessory sinuses (C30-C31)									
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of larynx (C32)									
2002.....	1.3	2.3	0.5	1.2	2.1	0.5	2.5	4.8	0.8
2001.....	1.3	2.5	0.5	1.2	2.3	0.5	2.5	5.0	0.9
2000.....	1.4	2.5	0.5	1.3	2.3	0.5	2.8	5.5	0.9
1999.....	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.1	0.8
Of trachea, bronchus and lung (C33-C34)									
2002.....	54.9	73.2	41.6	55.3	72.5	42.6	61.9	95.0	40.1
2001.....	55.3	75.2	41.0	55.6	74.2	42.1	62.5	99.1	38.7
2000.....	56.1	76.7	41.3	56.2	75.7	42.3	64.0	101.1	39.8
1999.....	55.5	76.9	40.2	55.4	75.6	41.0	64.8	103.0	39.9
Of thymus, heart, mediastinum and pleura (C37-C38)									
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2001.....	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)									
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of bone and articular cartilage (C40-C41)									
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.4
2001.....	0.5	0.6	0.4	0.5	0.6	0.3	0.5	0.6	0.4
2000.....	0.4	0.6	0.3	0.4	0.5	0.3	0.4	0.6	0.3
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
Melanoma and other malignant neoplasms of skin (C43-C44)									
2002.....	3.4	5.1	2.1	3.8	5.7	2.4	0.9	1.5	0.6
2001.....	3.5	5.3	2.2	3.9	5.9	2.5	1.0	1.4	0.7
2000.....	3.5	5.2	2.2	3.8	5.8	2.5	1.1	1.5	0.8
1999.....	3.5	5.2	2.2	3.9	5.8	2.5	0.9	1.6	0.6
Malignant melanoma of skin (C43)									
2002.....	2.6	3.8	1.7	2.9	4.2	2.0	0.4	0.5	0.3
2001.....	2.7	3.9	1.8	3.0	4.3	2.0	0.4	0.3	0.4
2000.....	2.7	3.8	1.8	3.0	4.3	2.0	0.6	0.6	0.5
1999.....	2.6	3.8	1.7	3.0	4.2	2.0	0.4	0.5	0.3
Other malignant neoplasms of skin (C44)									
2002.....	0.8	1.4	0.4	0.9	1.5	0.5	0.6	1.0	0.3
2001.....	0.9	1.4	0.5	0.9	1.5	0.5	0.6	1.0	0.2
2000.....	0.8	1.4	0.4	0.9	1.5	0.4	0.6	0.9	0.3
1999.....	0.8	1.4	0.4	0.9	1.5	0.4	0.5	1.0	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of mesothelial and soft tissue (C45-C49)									
2002.....	2.4	3.2	1.8	2.4	3.4	1.8	2.0	2.4	1.7
2001.....	2.4	3.2	1.8	2.5	3.4	1.8	1.9	2.2	1.7
2000.....	2.5	3.3	1.8	2.5	3.5	1.9	2.1	2.6	1.9
1999.....	2.4	3.3	1.8	2.5	3.4	1.8	2.2	2.6	1.9
Mesothelioma (C45)									
2002.....	0.9	1.7	0.2	0.9	1.8	0.3	0.4	0.8	*
2001.....	0.8	1.6	0.3	0.9	1.8	0.3	0.2	0.5	*
2000.....	0.9	1.7	0.3	0.9	1.8	0.3	0.4	0.8	0.1
1999.....	0.9	1.6	0.3	0.9	1.8	0.3	0.4	0.8	0.2
Kaposi's sarcoma (C46)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Of other mesothelial and soft tissue (C47-C49)									
2002.....	1.5	1.5	1.5	1.5	1.5	1.5	1.6	1.6	1.6
2001.....	1.5	1.5	1.5	1.5	1.6	1.5	1.6	1.7	1.6
2000.....	1.6	1.6	1.5	1.6	1.7	1.6	1.6	1.6	1.7
1999.....	1.6	1.6	1.5	1.5	1.6	1.5	1.7	1.8	1.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of breast (C50)									
2002.....	14.5	0.3	25.6	14.1	0.3	25.0	20.2	0.6	34.0
2001.....	14.8	0.3	26.0	14.4	0.3	25.5	20.4	0.6	34.4
2000.....	15.3	0.4	26.8	14.9	0.4	26.3	20.5	0.6	34.5
1999.....	15.2	0.3	26.6	14.8	0.3	26.0	20.8	0.7	35.1
Malignant neoplasms of female genital organs (C51-C58)									
2002.....	9.4	...	16.7	9.2	...	16.5	12.2	...	20.5
2001.....	9.5	...	16.8	9.4	...	16.6	12.1	...	20.4
2000.....	9.5	...	16.8	9.3	...	16.6	12.5	...	20.8
1999.....	9.4	...	16.7	9.3	...	16.5	12.3	...	20.6
Of vulva and vagina (C51-C52)									
2002.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.5
2001.....	0.4	...	0.7	0.4	...	0.7	0.5	...	0.8
2000.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
1999.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
Of cervix uteri (C53)									
2002.....	1.3	...	2.6	1.2	...	2.3	2.9	...	4.9
2001.....	1.4	...	2.7	1.3	...	2.5	2.8	...	4.8
2000.....	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4
1999.....	1.5	...	2.8	1.4	...	2.6	3.1	...	5.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of corpus uteri and uterus, part unspecified (C54-C55)									
2002.....	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3
2001.....	2.4	...	4.2	2.3	...	4.0	4.2	...	7.0
2000.....	2.4	...	4.1	2.2	...	3.9	4.4	...	7.2
1999.....	2.4	...	4.1	2.2	...	3.9	4.2	...	6.8
Of ovary (C56)									
2002.....	5.1	...	9.0	5.3	...	9.4	4.4	...	7.4
2001.....	5.1	...	9.0	5.2	...	9.3	4.5	...	7.5
2000.....	5.1	...	8.9	5.3	...	9.3	4.3	...	7.2
1999.....	5.0	...	8.8	5.1	...	9.1	4.4	...	7.3
Of other and unspecified female genital organs (C57)									
2002.....	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3
2001.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
1999.....	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3
Of placenta (C58)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of male genital organs (C60-C63)									
2002.....	10.8	28.4	...	10.0	26.2	...	22.1	62.4	...
2001.....	11.1	29.6	...	10.2	27.1	...	23.2	66.5	...
2000.....	11.5	30.8	...	10.5	28.3	...	24.1	68.6	...
1999.....	11.9	31.8	...	10.9	29.3	...	24.4	69.5	...
Of prostate (C61)									
2002.....	10.6	27.9	...	9.8	25.7	...	22.0	62.0	...
2001.....	10.9	29.1	...	10.0	26.6	...	23.0	66.1	...
2000.....	11.3	30.4	...	10.3	27.8	...	23.9	68.1	...
1999.....	11.6	31.3	...	10.7	28.7	...	24.2	69.0	...
Of testis (C62)									
2002.....	0.1	0.3	...	0.2	0.3	...	0.1	0.2	...
2001.....	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2000.....	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
1999.....	0.1	0.3	...	0.2	0.3	...	0.1	0.1	...
Of penis and other and unspecified male genital organs (C60,C63)									
2002.....	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2001.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2000.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
1999.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of urinary tract (C64-C68)									
2002.....	8.9	13.9	5.3	9.1	14.4	5.4	8.1	11.4	6.1
2001.....	8.8	14.1	5.2	9.0	14.5	5.2	8.2	11.7	5.9
2000.....	8.8	14.1	5.3	9.0	14.5	5.3	8.0	12.1	5.6
1999.....	8.7	13.8	5.2	8.8	14.2	5.1	8.3	12.0	6.0
Of kidney and renal pelvis (C64-C65)									
2002.....	4.2	6.1	2.8	4.3	6.3	2.8	4.0	5.8	2.8
2001.....	4.3	6.2	2.8	4.3	6.3	2.8	4.2	6.4	2.8
2000.....	4.2	6.2	2.8	4.3	6.3	2.8	4.1	6.3	2.8
1999.....	4.1	5.9	2.7	4.1	5.9	2.7	4.1	6.1	2.8
Of bladder (C67)									
2002.....	4.4	7.5	2.4	4.5	7.8	2.4	3.9	5.5	3.0
2001.....	4.3	7.5	2.3	4.5	7.9	2.2	3.7	5.1	2.9
2000.....	4.3	7.6	2.3	4.5	7.9	2.3	3.7	5.6	2.7
1999.....	4.4	7.6	2.3	4.5	7.9	2.3	4.0	5.8	3.0
Of other and unspecified sites within the urinary tract (C66,C68)									
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
2001.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
1999.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)									
2002.....	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.7
2001.....	0.8	0.8	0.7	0.8	0.8	0.7	0.7	0.6	0.8
2000.....	0.8	0.8	0.7	0.7	0.8	0.7	0.8	0.6	0.9
1999.....	0.7	0.8	0.7	0.7	0.8	0.7	0.7	0.7	0.7
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)									
2002.....	14.1	17.2	11.9	13.9	16.9	11.7	18.2	23.1	15.0
2001.....	14.2	17.2	12.0	13.9	16.8	11.9	18.4	24.3	14.5
2000.....	14.7	17.8	12.5	14.5	17.4	12.4	18.7	24.8	14.7
1999.....	16.0	19.5	13.6	15.7	19.1	13.4	20.6	26.5	16.8
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
2002.....	19.5	25.1	15.5	19.9	25.6	15.7	19.4	24.4	16.1
2001.....	20.0	25.5	16.0	20.2	25.9	16.2	20.0	25.3	16.5
2000.....	20.4	26.0	16.5	20.7	26.4	16.7	19.9	25.0	16.6
1999.....	20.6	26.1	16.7	20.8	26.5	16.8	20.3	25.6	16.8
Hodgkin's disease (C81)									
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000.....	0.5	0.6	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999.....	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.5	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)									
2002.....	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001.....	7.9	10.0	6.4	8.2	10.3	6.7	5.6	7.3	4.5
2000.....	8.3	10.3	6.8	8.6	10.7	7.1	5.2	6.7	4.2
1999.....	8.4	10.4	6.9	8.7	10.7	7.2	5.6	7.5	4.4
Non-Hodgkin's lymphoma (C82-C85)									
2002.....	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001.....	7.9	9.9	6.4	8.2	10.2	6.7	5.6	7.2	4.5
2000.....	8.2	10.2	6.7	8.6	10.7	7.1	5.2	6.6	4.2
1999.....	8.3	10.3	6.9	8.7	10.7	7.2	5.6	7.4	4.4
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Malignant immunoproliferative diseases (C88)									
2002.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Multiple myeloma and malignant plasma cell neoplasms (C90)									
2002.....	3.8	4.8	3.2	3.6	4.5	2.9	6.9	8.2	6.0
2001.....	3.8	4.7	3.2	3.6	4.4	2.9	7.3	8.7	6.3
2000.....	3.9	4.7	3.3	3.6	4.4	3.0	7.6	8.8	6.8
1999.....	3.9	4.7	3.3	3.5	4.4	2.9	7.5	8.8	6.6
Leukemia (C91-C95)									
2002.....	7.5	10.0	5.7	7.7	10.3	5.8	6.6	8.6	5.3
2001.....	7.6	10.1	5.9	7.8	10.4	6.0	6.7	8.7	5.4
2000.....	7.7	10.3	5.9	7.9	10.5	6.1	6.7	9.0	5.2
1999.....	7.6	10.2	6.0	7.8	10.4	6.1	6.7	8.7	5.4
Lymphoid leukemia (C91)									
2002.....	2.2	3.1	1.6	2.3	3.2	1.6	1.9	2.7	1.4
2001.....	2.2	3.1	1.6	2.3	3.1	1.6	2.1	2.9	1.5
2000.....	2.2	3.1	1.6	2.3	3.2	1.6	1.9	2.8	1.3
1999.....	2.3	3.2	1.6	2.4	3.3	1.7	2.0	2.9	1.4
Myeloid leukemia (C92)									
2002.....	3.4	4.5	2.7	3.5	4.6	2.8	2.8	3.7	2.3
2001.....	3.5	4.5	2.8	3.6	4.7	2.9	2.9	3.6	2.4
2000.....	3.5	4.6	2.8	3.6	4.7	2.8	3.0	3.7	2.6
1999.....	3.4	4.3	2.7	3.4	4.4	2.8	2.8	3.6	2.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Carcinoma in situ of other and unspecified sites (D00-D04,D09)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Benign neoplasms (D10-D36)									
2002.....	0.5	0.5	0.6	0.5	0.5	0.5	0.8	0.9	0.9
2001.....	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.1	0.9
2000.....	0.6	0.5	0.6	0.5	0.5	0.6	0.9	0.8	1.0
1999.....	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.0	0.9
Of female genital organs (D25-D28)									
2002.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.2
1999.....	0.0	...	0.0	*	...	*	*	...	*
Of eye, brain and other parts of central nervous system (D31-D33)									
2002.....	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.6	0.6
2000.....	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.5	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
2001.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.5	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)									
2002.....	3.8	5.0	3.1	4.0	5.2	3.2	3.3	4.1	2.9
2001.....	4.1	5.3	3.3	4.2	5.4	3.4	3.4	4.3	2.9
2000.....	4.1	5.3	3.3	4.2	5.4	3.4	3.4	4.4	2.9
1999.....	4.1	5.3	3.3	4.2	5.5	3.4	3.6	4.1	3.2
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)									
2002.....	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
1999.....	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.3	0.3
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)									
2002.....	3.4	3.4	3.3	2.8	2.9	2.8	7.5	7.7	7.4
2001.....	3.4	3.4	3.3	2.9	3.0	2.8	7.4	7.2	7.5
2000.....	3.4	3.4	3.3	2.9	3.0	2.8	7.4	7.2	7.5
1999.....	3.3	3.4	3.2	2.8	3.0	2.7	7.3	7.3	7.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Anemias (D50-D64)									
2002.....	1.6	1.6	1.6	1.3	1.3	1.4	3.7	3.9	3.5
2001.....	1.6	1.6	1.6	1.4	1.4	1.4	3.5	3.6	3.4
2000.....	1.6	1.6	1.6	1.4	1.4	1.4	3.4	3.4	3.5
1999.....	1.6	1.7	1.6	1.4	1.4	1.4	3.6	3.7	3.5
Nutritional anemias (D50-D53)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Hemolytic anemias (D55-D59)									
2002.....	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.5	1.5
2001.....	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.3
2000.....	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.3	1.5
1999.....	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.7	1.4
Aplastic anemias (D60-D61)									
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
2001.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3
2000.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4
1999.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute posthemorrhagic and other anemias (D62,D64)									
2002.....	0.8	0.8	0.9	0.8	0.7	0.8	1.7	1.8	1.5
2001.....	0.9	0.8	0.9	0.8	0.7	0.8	1.5	1.5	1.6
2000.....	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.4	1.4
1999.....	0.8	0.8	0.9	0.8	0.7	0.8	1.5	1.3	1.5
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)									
2002.....	0.9	1.0	0.9	0.9	1.0	0.8	1.4	1.4	1.4
2001.....	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.4	1.5
2000.....	1.0	1.0	0.9	0.9	1.0	0.9	1.5	1.5	1.5
1999.....	1.0	1.0	0.9	0.9	0.9	0.8	1.6	1.8	1.5
Other diseases of blood and blood-forming organs (D70-D76)									
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000.....	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
Certain disorders involving the immune mechanism (D80-D89)									
2002.....	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.3
2001.....	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.4
2000.....	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.2
1999.....	0.4	0.4	0.4	0.3	0.3	0.2	1.7	1.5	1.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
IV. Endocrine, nutritional and metabolic diseases (E00-E88)									
2002.....	34.4	37.7	31.7	31.9	35.7	28.8	62.3	62.3	61.2
2001.....	34.3	37.1	31.8	31.8	35.0	29.0	61.7	63.0	60.1
2000.....	34.1	36.9	31.8	31.6	34.9	28.9	62.1	61.4	61.8
1999.....	34.0	36.6	31.8	31.3	34.4	28.8	62.2	63.0	61.3
Endocrine diseases (E00-E34)									
2002.....	26.4	29.3	24.1	24.1	27.4	21.5	50.8	50.4	50.0
2001.....	26.3	28.9	24.2	24.0	27.0	21.7	50.6	51.2	49.5
2000.....	26.0	28.4	24.1	23.7	26.6	21.5	51.0	49.9	51.0
1999.....	26.0	28.5	24.1	23.5	26.5	21.3	51.2	50.8	51.0
Disorders of thyroid gland (E00-E07)									
2002.....	0.7	0.4	0.8	0.7	0.4	0.8	0.7	0.4	0.9
2001.....	0.7	0.4	0.8	0.7	0.4	0.8	0.8	0.6	0.9
2000.....	0.6	0.4	0.8	0.6	0.3	0.8	0.8	0.6	1.0
1999.....	0.7	0.4	0.8	0.6	0.4	0.8	0.9	0.6	1.0
Diabetes mellitus (E10-E14)									
2002.....	25.4	28.6	23.0	23.1	26.8	20.3	49.5	49.4	48.6
2001.....	25.3	28.1	23.1	23.0	26.2	20.5	49.2	49.9	48.1
2000.....	25.0	27.8	23.0	22.8	26.0	20.4	49.5	48.7	49.4
1999.....	25.0	27.8	23.0	22.6	25.8	20.2	49.7	49.4	49.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Disorders of pituitary gland (E22-E23)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
Disorders of adrenal glands (E24-E27)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Diseases of thymus (E32)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other endocrine diseases (E15-E21,E28-E31,E34)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.3	0.4
2001.....	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.4	0.3
2000.....	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.3	0.4
1999.....	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.5	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Nutritional diseases (E40-E68)									
2002.....	2.6	2.5	2.6	2.4	2.4	2.4	4.5	4.0	4.7
2001.....	2.4	2.2	2.5	2.3	2.1	2.3	4.1	4.0	4.3
2000.....	2.6	2.5	2.6	2.5	2.4	2.5	4.3	4.2	4.4
1999.....	2.5	2.5	2.5	2.4	2.3	2.4	4.4	4.5	4.3
Malnutrition (E40-E46)									
2002.....	1.2	1.1	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001.....	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000.....	1.4	1.3	1.5	1.4	1.2	1.4	2.2	2.5	2.1
1999.....	1.5	1.4	1.5	1.4	1.3	1.4	2.5	2.9	2.2
Nutritional marasmus (E41)									
2002.....	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Other and unspecified malnutrition (E40,E42-E46)									
2002.....	1.1	1.1	1.1	1.1	1.0	1.1	1.9	1.9	1.9
2001.....	1.1	1.0	1.2	1.1	1.0	1.2	1.8	2.1	1.6
2000.....	1.3	1.2	1.4	1.3	1.2	1.3	2.1	2.3	2.0
1999.....	1.4	1.3	1.4	1.3	1.2	1.3	2.3	2.8	2.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other nutritional deficiencies (E50-E64)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Obesity and other hyperalimentionation (E65-E68)									
2002.....	1.3	1.2	1.3	1.2	1.2	1.1	2.3	1.8	2.7
2001.....	1.1	1.0	1.2	1.0	1.0	1.0	2.1	1.6	2.5
2000.....	1.1	1.1	1.1	1.0	1.0	0.9	1.9	1.6	2.2
1999.....	0.9	0.9	1.0	0.9	0.9	0.8	1.7	1.3	2.0
Metabolic disorders (E70-E88)									
2002.....	5.4	5.9	5.0	5.4	5.9	4.9	7.1	8.0	6.5
2001.....	5.5	6.0	5.0	5.5	6.0	5.0	7.0	7.9	6.3
2000.....	5.5	5.9	5.0	5.5	5.9	5.0	6.8	7.3	6.3
1999.....	5.5	5.7	5.1	5.4	5.6	5.1	6.6	7.7	6.0
Cystic fibrosis (E84)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)									
2002.....	2.5	2.3	2.6	2.4	2.1	2.6	4.0	4.2	3.8
2001.....	2.7	2.5	2.8	2.6	2.3	2.8	4.4	4.5	4.2
2000.....	2.8	2.7	2.8	2.7	2.6	2.7	4.2	4.5	4.0
1999.....	3.0	2.8	3.1	2.9	2.6	3.1	4.3	5.1	3.7
Other metabolic disorders (E70-E83,E85,E88)									
2002.....	2.8	3.5	2.2	2.8	3.5	2.2	3.1	3.7	2.6
2001.....	2.6	3.4	2.0	2.7	3.4	2.1	2.6	3.3	2.1
2000.....	2.5	3.1	2.0	2.5	3.2	2.0	2.5	2.7	2.3
1999.....	2.3	2.8	1.9	2.3	2.8	1.9	2.4	2.5	2.2
V. Mental and behavioral disorders (F01-F99)									
2002.....	19.2	18.8	18.4	19.3	18.6	18.8	20.1	23.5	17.0
2001.....	18.0	17.8	17.1	18.1	17.6	17.4	18.9	22.7	15.6
2000.....	16.7	16.8	15.7	16.8	16.5	16.0	17.9	22.0	14.4
1999.....	15.5	16.0	14.3	15.6	15.8	14.5	15.9	20.4	12.3
Organic dementia (F01,F03)									
2002.....	14.3	12.2	15.3	14.6	12.3	15.7	13.5	13.2	13.4
2001.....	13.2	11.4	14.0	13.5	11.5	14.3	12.0	11.7	12.0
2000.....	12.0	10.3	12.6	12.2	10.3	12.9	11.1	11.3	10.8
1999.....	10.6	9.3	11.1	10.9	9.5	11.5	8.7	8.9	8.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified organic mental disorders (F04-F09)									
2002.....	0.7	0.6	0.7	0.7	0.6	0.7	0.6	0.5	0.6
2001.....	0.7	0.7	0.8	0.8	0.7	0.8	0.8	0.8	0.7
2000.....	0.8	0.7	0.9	0.8	0.7	0.9	0.8	0.8	0.7
1999.....	0.9	0.8	0.9	0.9	0.8	1.0	0.9	1.1	0.8
Mental and behavioral disorders due to psychoactive substance use (F10-F19)									
2002.....	3.3	5.2	1.5	3.1	4.9	1.5	5.2	8.9	2.3
2001.....	3.2	5.0	1.5	3.0	4.6	1.4	5.4	9.4	2.3
2000.....	3.1	5.0	1.4	2.9	4.6	1.3	5.4	9.3	2.3
1999.....	3.1	5.1	1.4	2.9	4.6	1.2	5.7	9.7	2.6
Mental and behavioral disorders due to use of alcohol (F10)									
2002.....	2.3	3.9	1.0	2.3	3.7	0.9	3.3	6.1	1.2
2001.....	2.3	3.8	1.0	2.2	3.5	0.9	3.7	6.7	1.3
2000.....	2.2	3.7	0.9	2.1	3.5	0.8	3.6	6.6	1.3
1999.....	2.2	3.7	0.9	2.1	3.4	0.8	3.8	6.8	1.5
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)									
2002.....	0.9	1.3	0.6	0.9	1.2	0.5	1.9	2.8	1.1
2001.....	0.9	1.2	0.5	0.8	1.1	0.5	1.8	2.6	1.1
2000.....	0.9	1.2	0.6	0.8	1.1	0.5	1.8	2.7	1.1
1999.....	0.9	1.3	0.5	0.8	1.1	0.4	1.9	2.9	1.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Schizophrenia, schizotypal and delusional disorders (F20-F29)									
2002.....	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.3	0.2
2001.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Mood (affective) disorders (F30-F39)									
2002.....	0.3	0.2	0.3	0.3	0.2	0.4	0.2	*	0.1
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.2	0.1
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	0.1
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	*
Neurotic, stress-related and somatoform disorders (F40-F48)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Mental retardation (F70-F79)									
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)									
2002.....	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	0.1
2001.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*
2000.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*
1999.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*
VI. Diseases of the nervous system (G00-G98)									
2002.....	35.6	36.5	34.4	36.7	37.6	35.6	29.3	30.5	27.9
2001.....	34.2	35.8	32.7	35.2	36.8	33.7	28.6	30.2	27.3
2000.....	33.0	34.8	31.5	34.0	35.8	32.4	27.0	28.8	25.7
1999.....	30.9	33.2	29.1	31.8	34.2	29.9	25.0	26.6	23.7
Meningitis (G00,G03)									
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.5	0.7	0.4
2001.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.8	0.4
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6
Other inflammatory diseases of central nervous system (G04-G09)									
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.4
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
1999.....	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Parkinson's disease (G20-G21)									
2002.....	5.9	9.0	4.1	6.3	9.5	4.3	2.5	4.0	1.7
2001.....	5.9	9.1	4.0	6.2	9.7	4.3	2.5	3.9	1.8
2000.....	5.7	8.8	3.9	6.0	9.3	4.2	2.5	4.2	1.6
1999.....	5.4	8.4	3.7	5.7	8.9	3.9	2.4	3.7	1.7
Alzheimer's disease (G30)									
2002.....	20.2	16.6	22.0	21.1	17.3	22.9	15.2	13.2	15.9
2001.....	19.1	16.1	20.5	19.8	16.7	21.3	14.6	13.1	15.1
2000.....	18.1	15.2	19.3	18.8	15.8	20.1	13.0	11.4	13.7
1999.....	16.5	14.4	17.4	17.2	15.0	18.1	11.3	10.2	11.8
Multiple sclerosis (G35)									
2002.....	1.1	0.8	1.3	1.1	0.8	1.4	1.1	0.6	1.5
2001.....	1.1	0.8	1.3	1.1	0.8	1.4	1.0	0.6	1.3
2000.....	1.0	0.8	1.2	1.1	0.8	1.3	1.0	0.7	1.3
1999.....	1.0	0.7	1.2	1.0	0.7	1.2	0.8	0.6	1.0
Epilepsy (G40-G41)									
2002.....	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8
2001.....	0.4	0.5	0.4	0.4	0.5	0.4	0.9	1.2	0.8
2000.....	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.3	0.9
1999.....	0.5	0.6	0.4	0.4	0.5	0.4	1.1	1.3	0.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Transient cerebral ischemic attacks and related syndromes (G45)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Infantile cerebral palsy (G80)									
2002.....	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.7	0.5
2001.....	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4
2000.....	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.6	0.5
1999.....	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.4
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)									
2002.....	6.9	8.3	5.8	6.9	8.3	5.8	7.9	9.4	6.8
2001.....	6.7	8.1	5.7	6.7	8.1	5.6	8.1	9.6	7.2
2000.....	6.8	8.3	5.7	6.8	8.3	5.6	7.9	9.5	6.8
1999.....	6.6	8.1	5.6	6.6	8.1	5.5	7.7	8.8	7.0
VII. Diseases of the eye and adnexa (H00-H57)									
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
VIII. Diseases of the ear and mastoid process (H60-H93)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
IX. Diseases of the circulatory system (I00-I99)									
2002.....	319.0	378.8	272.1	312.0	372.2	264.3	418.9	490.6	366.4
2001.....	328.2	389.8	280.4	320.8	382.6	272.3	430.2	508.3	374.9
2000.....	341.3	409.0	290.0	334.0	402.3	281.7	441.6	520.7	386.1
1999.....	350.8	420.5	297.9	343.3	413.2	289.8	450.0	534.9	390.8
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)									
2002.....	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001.....	1.2	0.9	1.5	1.2	0.9	1.5	1.1	0.7	1.3
2000.....	1.3	1.0	1.5	1.3	1.0	1.6	1.0	0.7	1.2
1999.....	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.8	1.2
Acute rheumatic fever (I00-I02)									
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	*	0.0	0.0	*	*	*	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)									
2002.....	0.4	0.2	0.4	0.4	0.3	0.4	0.3	0.2	0.3
2001.....	0.4	0.3	0.5	0.4	0.3	0.5	0.3	0.2	0.4
2000.....	0.4	0.3	0.5	0.4	0.3	0.5	0.4	0.2	0.5
1999.....	0.4	0.3	0.5	0.5	0.3	0.5	0.3	0.2	0.3
Hypertensive diseases (I10-I13)									
2002.....	17.1	17.4	16.3	14.4	14.4	13.8	44.8	49.6	40.6
2001.....	16.5	16.6	15.8	13.9	13.7	13.4	43.0	47.8	38.9
2000.....	16.2	16.3	15.5	13.5	13.4	12.9	43.3	47.2	39.9
1999.....	15.8	15.9	15.2	13.1	12.9	12.6	43.1	47.5	39.4
Essential (primary) hypertension (I10)									
2002.....	4.0	3.5	4.2	3.7	3.1	3.9	7.7	7.5	7.5
2001.....	3.9	3.5	4.0	3.6	3.1	3.7	7.6	8.2	7.1
2000.....	3.7	3.4	3.8	3.4	3.0	3.5	7.3	7.6	6.9
1999.....	3.6	3.3	3.7	3.3	2.9	3.4	7.2	7.4	7.0
Hypertensive heart disease (I11)									
2002.....	9.1	9.5	8.3	7.6	7.8	7.0	24.4	28.0	21.2
2001.....	8.7	8.9	8.1	7.3	7.3	6.8	23.2	26.3	20.5
2000.....	8.6	8.7	8.1	7.1	7.1	6.7	23.7	26.3	21.2
1999.....	8.3	8.3	7.9	6.9	6.7	6.6	22.9	25.7	20.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive renal disease (I12)									
2002.....	3.0	3.3	2.8	2.4	2.7	2.2	9.3	10.2	8.6
2001.....	2.9	3.2	2.7	2.3	2.5	2.1	8.9	9.7	8.3
2000.....	2.8	3.1	2.6	2.2	2.5	2.0	9.0	9.5	8.7
1999.....	2.6	2.9	2.4	2.0	2.3	1.8	8.8	9.8	8.0
Hypertensive heart and renal disease (I13)									
2002.....	1.0	1.0	0.9	0.7	0.8	0.7	3.5	3.9	3.2
2001.....	1.0	1.1	1.0	0.8	0.8	0.7	3.3	3.6	3.0
2000.....	1.0	1.1	0.9	0.8	0.9	0.7	3.3	3.8	3.1
1999.....	1.2	1.4	1.1	0.9	1.0	0.9	4.2	4.5	3.9
Ischemic heart diseases (I20-I25)									
2002.....	170.8	220.4	133.6	169.8	220.5	131.2	203.0	250.6	169.7
2001.....	177.8	228.5	139.9	176.5	228.4	137.4	211.6	262.0	176.7
2000.....	186.8	241.4	146.5	185.6	241.6	143.9	218.3	269.5	183.2
1999.....	194.6	251.2	152.9	193.4	251.2	150.5	224.9	278.7	188.0
Acute myocardial infarction (I21-I22)									
2002.....	62.1	80.6	48.1	61.9	81.0	47.1	73.9	89.5	63.2
2001.....	65.4	84.9	50.9	65.1	85.4	49.9	76.8	93.1	66.0
2000.....	69.9	91.3	54.1	69.6	91.8	53.1	81.0	98.9	69.2
1999.....	73.2	96.1	56.3	73.1	96.8	55.4	83.1	102.1	70.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other acute ischemic heart diseases (I24)									
2002.....	1.2	1.5	0.9	1.1	1.4	0.9	1.7	2.3	1.3
2001.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.3	1.3
2000.....	1.2	1.5	0.9	1.2	1.5	0.9	1.8	2.5	1.3
1999.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.4	1.3
Other forms of chronic ischemic heart disease (I20,I25)									
2002.....	107.6	138.3	84.5	106.8	138.0	83.2	127.5	158.9	105.2
2001.....	111.2	142.1	88.1	110.2	141.6	86.7	133.0	166.7	109.3
2000.....	115.7	148.6	91.5	114.8	148.3	90.0	135.4	168.1	112.6
1999.....	120.2	153.6	95.6	119.2	153.0	94.2	140.0	174.2	116.2
Atherosclerotic cardiovascular disease, so described (I25.0)									
2002.....	23.5	30.4	17.8	22.1	28.5	16.7	40.3	54.8	30.0
2001.....	24.1	30.8	18.5	22.5	28.8	17.3	41.8	56.0	31.7
2000.....	25.2	32.3	19.4	23.6	30.2	18.2	43.5	58.7	32.9
1999.....	26.4	33.5	20.6	24.7	31.2	19.3	45.8	61.5	34.9
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
2002.....	84.1	108.0	66.7	84.7	109.5	66.4	87.2	104.1	75.2
2001.....	87.1	111.3	69.6	87.6	112.8	69.4	91.1	110.6	77.6
2000.....	90.5	116.3	72.0	91.2	118.1	71.8	91.9	109.5	79.7
1999.....	93.8	120.1	75.0	94.5	121.8	74.9	94.1	112.7	81.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other heart diseases (I26-I51)									
2002.....	58.5	65.5	52.8	57.3	64.1	51.7	76.6	87.8	68.1
2001.....	59.1	66.1	53.4	57.7	64.4	52.3	77.8	91.8	68.2
2000.....	60.0	67.9	53.8	58.6	66.3	52.6	78.4	92.2	69.0
1999.....	60.9	69.2	54.5	59.3	67.2	53.1	81.2	97.5	70.1
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)									
2002.....	4.8	4.4	5.0	4.5	4.2	4.7	8.1	7.2	8.8
2001.....	4.7	4.5	4.9	4.5	4.2	4.6	8.0	7.9	8.3
2000.....	4.7	4.5	4.9	4.5	4.3	4.6	8.0	7.6	8.3
1999.....	4.8	4.7	5.0	4.5	4.5	4.6	8.7	8.4	8.9
Pulmonary embolism (I26)									
2002.....	3.0	2.9	3.1	2.8	2.7	2.9	5.2	5.0	5.4
2001.....	3.1	3.0	3.1	2.9	2.8	2.9	5.3	5.7	5.2
2000.....	3.0	3.0	3.0	2.9	2.9	2.8	5.4	5.7	5.4
1999.....	3.3	3.3	3.3	3.1	3.1	3.1	6.1	6.5	5.9
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)									
2002.....	1.8	1.5	1.9	1.7	1.5	1.8	2.9	2.2	3.3
2001.....	1.7	1.5	1.8	1.6	1.4	1.7	2.7	2.3	3.0
2000.....	1.7	1.5	1.8	1.6	1.4	1.7	2.5	2.0	2.9
1999.....	1.5	1.4	1.6	1.5	1.4	1.5	2.6	1.9	3.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other non-pulmonary forms of heart disease (I30-I51)									
2002.....	53.8	61.1	47.8	52.8	59.9	47.0	68.5	80.6	59.4
2001.....	54.4	61.6	48.5	53.3	60.2	47.7	69.8	83.9	60.0
2000.....	55.2	63.4	48.9	54.2	62.0	48.0	70.5	84.5	60.7
1999.....	56.1	64.5	49.6	54.9	62.7	48.6	72.5	89.1	61.2
Acute and subacute endocarditis (I33)									
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2001.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.1	0.6
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.7
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
Diseases of pericardium and acute myocarditis (I30-I31,I40)									
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4
2000.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.5	0.5
Other diseases of endocardium (I34-I38)									
2002.....	6.9	7.1	6.7	7.2	7.4	7.0	4.8	4.6	4.8
2001.....	6.9	7.2	6.6	7.2	7.5	6.9	4.9	5.3	4.6
2000.....	7.2	7.5	6.9	7.4	7.7	7.1	5.3	5.4	5.1
1999.....	7.2	7.6	6.9	7.4	7.8	7.1	5.3	5.4	5.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Nonrheumatic mitral valve disorders (I34)									
2002.....	1.0	0.8	1.0	1.0	0.9	1.1	0.6	0.6	0.7
2001.....	1.0	0.9	1.1	1.0	0.9	1.1	0.7	0.5	0.8
2000.....	1.0	1.0	1.1	1.1	1.0	1.1	0.8	0.6	0.9
1999.....	1.1	1.0	1.1	1.1	1.0	1.2	0.8	0.6	0.9
Nonrheumatic aortic valve disorders (I35)									
2002.....	4.3	4.6	4.0	4.5	4.9	4.2	2.4	2.4	2.4
2001.....	4.3	4.7	4.0	4.5	4.9	4.2	2.5	3.0	2.1
2000.....	4.5	4.9	4.2	4.7	5.1	4.4	2.7	2.9	2.5
1999.....	4.5	5.0	4.2	4.7	5.2	4.3	2.7	3.1	2.5
All other diseases of endocardium (I36-I38)									
2002.....	1.6	1.6	1.7	1.6	1.6	1.7	1.7	1.6	1.7
2001.....	1.6	1.7	1.6	1.6	1.7	1.6	1.7	1.8	1.6
2000.....	1.6	1.6	1.6	1.6	1.6	1.6	1.8	2.0	1.7
1999.....	1.6	1.6	1.6	1.6	1.6	1.6	1.8	1.8	1.8
Cardiomyopathy (I42)									
2002.....	9.6	13.4	6.8	8.9	12.5	6.2	16.6	23.0	12.0
2001.....	9.5	13.1	6.8	8.7	12.2	6.2	16.9	23.3	12.5
2000.....	9.8	13.8	7.0	9.0	12.7	6.4	17.8	24.9	12.8
1999.....	10.0	13.9	7.1	9.1	12.7	6.5	18.2	25.9	12.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Conduction disorders and cardiac dysrhythmias (I44-I49)									
2002.....	13.6	15.1	12.2	13.2	14.6	11.9	18.8	21.7	16.7
2001.....	13.6	15.1	12.3	13.2	14.5	12.0	19.1	22.8	16.6
2000.....	13.9	15.6	12.4	13.6	15.2	12.1	18.8	21.8	16.6
1999.....	14.7	16.7	13.0	14.3	16.2	12.8	20.0	24.6	17.1
Heart failure (I50)									
2002.....	19.5	20.4	18.6	19.5	20.5	18.6	21.8	23.3	20.5
2001.....	20.2	21.2	19.2	20.2	21.2	19.3	22.1	24.4	20.6
2000.....	20.3	21.5	19.2	20.3	21.6	19.3	22.0	24.2	20.7
1999.....	20.3	21.4	19.3	20.3	21.4	19.3	22.5	25.3	20.9
Congestive heart failure (I50.0)									
2002.....	18.0	18.8	17.3	18.1	19.0	17.4	19.6	20.7	18.5
2001.....	18.7	19.5	18.0	18.8	19.6	18.1	19.9	21.7	18.8
2000.....	18.8	19.8	17.9	18.9	20.0	18.0	19.7	21.3	18.8
1999.....	18.8	19.8	17.9	18.8	19.7	18.0	20.3	22.7	18.9
Other and unspecified heart failure (I50.1-I50.9)									
2002.....	1.4	1.6	1.3	1.3	1.5	1.2	2.2	2.6	2.0
2001.....	1.5	1.7	1.3	1.4	1.6	1.2	2.2	2.7	1.8
2000.....	1.5	1.7	1.4	1.4	1.6	1.3	2.3	2.8	1.9
1999.....	1.5	1.7	1.3	1.4	1.6	1.3	2.2	2.5	2.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)									
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.6	0.4
2001.....	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.4
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3
1999.....	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)									
2002.....	3.2	4.0	2.6	3.1	3.8	2.5	4.9	6.1	3.9
2001.....	3.2	3.8	2.6	3.0	3.7	2.5	5.0	6.0	4.2
2000.....	3.2	3.9	2.6	3.0	3.7	2.4	5.0	6.5	4.0
1999.....	3.0	3.7	2.4	2.8	3.5	2.2	4.7	6.0	3.8
Cerebrovascular diseases (I60-I69)									
2002.....	56.2	56.5	55.2	54.2	54.2	53.4	76.3	81.7	71.8
2001.....	57.9	59.0	56.4	55.8	56.5	54.5	78.8	85.4	73.7
2000.....	60.9	62.4	59.1	58.8	59.8	57.3	81.9	89.6	76.2
1999.....	61.6	63.2	59.8	59.6	60.8	58.0	81.8	89.6	76.2
Subarachnoid hemorrhage (I60)									
2002.....	2.1	1.6	2.6	2.0	1.5	2.4	2.9	2.1	3.5
2001.....	2.2	1.7	2.5	2.0	1.6	2.4	3.1	2.3	3.7
2000.....	2.4	1.8	2.8	2.2	1.7	2.6	3.4	2.3	4.2
1999.....	2.4	1.8	2.8	2.2	1.7	2.6	3.3	2.4	4.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intracerebral and other intracranial hemorrhage (I61-I62)									
2002.....	9.4	10.4	8.5	8.8	9.7	8.0	13.2	15.3	11.6
2001.....	9.4	10.5	8.6	8.8	9.8	8.0	13.7	15.8	12.1
2000.....	9.9	11.1	9.0	9.2	10.3	8.5	14.3	17.0	12.3
1999.....	9.3	10.3	8.5	8.7	9.6	8.0	13.3	15.5	11.7
Cerebral infarction (I63)									
2002.....	4.3	4.4	4.1	4.1	4.3	4.0	5.5	6.1	5.0
2001.....	4.5	4.6	4.3	4.4	4.5	4.2	5.8	6.8	5.2
2000.....	5.0	5.3	4.7	4.9	5.1	4.6	6.6	7.5	5.9
1999.....	5.4	5.7	5.2	5.3	5.5	5.1	7.1	8.3	6.3
Stroke, not specified as hemorrhage or infarction (I64)									
2002.....	30.0	29.6	29.8	29.1	28.4	29.0	41.2	44.1	38.8
2001.....	31.4	31.4	30.9	30.5	30.2	30.2	42.7	46.4	39.8
2000.....	32.8	33.0	32.2	31.9	31.8	31.4	44.2	48.5	41.1
1999.....	33.6	34.0	32.8	32.7	32.8	32.0	44.6	48.4	41.8
Other cerebrovascular diseases and their sequelae (I67,I69)									
2002.....	10.4	10.6	10.2	10.2	10.3	10.0	13.4	14.0	12.9
2001.....	10.4	10.7	10.1	10.2	10.5	9.8	13.5	14.1	13.0
2000.....	10.8	11.3	10.3	10.5	11.0	10.1	13.4	14.4	12.7
1999.....	10.9	11.4	10.5	10.7	11.1	10.3	13.6	15.0	12.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of arteries, arterioles and capillaries (I70-I78)									
2002.....	13.4	16.4	11.2	13.5	16.6	11.2	14.4	17.0	12.6
2001.....	14.0	17.1	11.7	14.0	17.2	11.7	15.1	17.4	13.4
2000.....	14.7	18.4	12.0	14.7	18.6	11.9	15.7	18.4	13.9
1999.....	15.0	18.5	12.5	15.1	18.7	12.5	15.5	18.2	13.7
Atherosclerosis (I70)									
2002.....	4.7	4.9	4.6	4.8	5.0	4.7	4.4	4.9	4.0
2001.....	5.0	5.1	4.8	5.1	5.2	5.0	4.3	4.6	4.1
2000.....	5.2	5.5	5.0	5.3	5.6	5.1	4.8	5.4	4.4
1999.....	5.5	5.7	5.3	5.6	5.8	5.4	5.0	5.8	4.5
Aortic aneurysm and dissection (I71)									
2002.....	5.2	7.7	3.3	5.2	7.9	3.4	4.6	6.1	3.5
2001.....	5.4	8.0	3.6	5.5	8.2	3.6	5.0	6.4	3.9
2000.....	5.7	8.7	3.7	5.8	9.0	3.6	5.2	6.8	4.1
1999.....	5.8	8.8	3.7	5.9	9.1	3.7	4.9	6.3	3.9
Other diseases of arteries, arterioles and capillaries (I72-I78)									
2002.....	3.5	3.8	3.3	3.4	3.7	3.1	5.4	6.0	5.0
2001.....	3.6	3.9	3.3	3.4	3.8	3.2	5.9	6.4	5.5
2000.....	3.7	4.2	3.4	3.6	4.0	3.2	5.7	6.2	5.3
1999.....	3.7	4.0	3.4	3.5	3.9	3.3	5.6	6.0	5.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other disorders of circulatory system (I80-I99)									
2002.....	1.6	1.6	1.6	1.5	1.5	1.6	2.9	3.2	2.6
2001.....	1.6	1.7	1.6	1.5	1.6	1.5	2.7	3.1	2.5
2000.....	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.0	2.8
1999.....	1.5	1.5	1.4	1.4	1.4	1.3	2.4	2.6	2.2
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)									
2002.....	1.2	1.1	1.3	1.2	1.0	1.2	2.3	2.4	2.2
2001.....	1.2	1.2	1.3	1.2	1.1	1.2	2.2	2.3	2.1
2000.....	1.2	1.2	1.2	1.1	1.1	1.2	2.2	2.1	2.3
1999.....	1.0	0.9	1.0	0.9	0.9	0.9	1.7	1.7	1.6
All other and unspecified disorders of circulatory system (I83-I99)									
2002.....	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.8	0.4
2001.....	0.4	0.5	0.3	0.4	0.5	0.3	0.5	0.8	0.4
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.9	0.5
1999.....	0.5	0.6	0.4	0.5	0.6	0.4	0.7	1.0	0.6
X. Diseases of the respiratory system (J00-J98)									
2002.....	81.4	100.4	69.7	83.4	101.7	72.1	70.5	97.1	55.3
2001.....	81.6	101.2	69.8	83.3	102.3	71.9	71.8	99.5	56.3
2000.....	83.7	105.7	70.9	85.4	106.8	73.0	73.5	102.9	56.9
1999.....	84.3	107.7	70.6	85.9	108.7	72.5	75.5	106.9	57.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Acute upper respiratory infections (J00-J06)									
2002.....	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2001.....	0.0	0.0	0.1	0.0	0.0	0.1	0.1	*	*
2000.....	0.0	0.0	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Acute pharyngitis and tonsillitis (J02-J03)									
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	*	*	0.0	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Acute laryngitis and tracheitis (J04)									
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
1999.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other diseases of the respiratory system (J10-J98)									
2002.....	81.3	100.4	69.7	83.3	101.7	72.1	70.5	97.0	55.3
2001.....	81.5	101.2	69.7	83.2	102.2	71.8	71.8	99.4	56.2
2000.....	83.7	105.7	70.8	85.3	106.7	72.9	73.5	102.8	56.8
1999.....	84.3	107.6	70.5	85.8	108.6	72.4	75.4	106.8	57.8
Influenza (J10-J11)									
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.6	0.7	0.6	0.6	0.7	0.6	0.3	0.4	0.3
1999.....	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3
Pneumonia (J12-J18)									
2002.....	22.4	26.7	19.6	22.3	26.4	19.7	23.9	30.7	19.9
2001.....	21.9	26.5	19.1	21.6	25.9	19.0	24.1	32.2	19.4
2000.....	23.1	28.2	20.1	22.9	27.7	20.0	25.3	33.5	20.5
1999.....	22.9	27.9	20.0	22.7	27.5	19.9	25.1	33.2	20.5
Viral pneumonia, not elsewhere classified (J12)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Bacterial pneumonia (J13-J15)									
2002.....	1.6	2.0	1.3	1.6	2.0	1.3	1.7	2.3	1.4
2001.....	1.6	2.0	1.2	1.5	2.0	1.3	1.9	2.8	1.3
2000.....	1.6	2.2	1.3	1.6	2.1	1.2	2.0	2.9	1.5
1999.....	1.4	1.8	1.2	1.4	1.7	1.2	1.7	2.3	1.3
Pneumonia due to other or unspecified organisms (J16,J18)									
2002.....	20.7	24.6	18.2	20.6	24.3	18.3	22.1	28.3	18.5
2001.....	20.2	24.3	17.7	20.0	23.8	17.7	22.2	29.3	18.0
2000.....	21.3	25.9	18.7	21.1	25.4	18.6	23.2	30.5	18.9
1999.....	21.3	26.0	18.6	21.2	25.6	18.6	23.3	30.7	19.0
Other acute lower respiratory infections (J20-J22)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.1
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Acute bronchitis and bronchiolitis (J20-J21)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Unspecified acute lower respiratory infection (J22)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Other diseases of upper respiratory tract (J30-J39)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2001.....	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
2000.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.2
1999.....	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2
Chronic lower respiratory diseases (J40-J47)									
2002.....	43.5	53.5	37.4	45.4	54.9	39.7	31.2	46.3	22.6
2001.....	43.7	54.0	37.6	45.6	55.5	39.8	30.9	46.3	22.4
2000.....	44.2	55.8	37.4	46.0	57.2	39.5	31.6	47.5	22.7
1999.....	45.4	58.7	37.7	47.1	60.0	39.7	33.5	51.5	23.4
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)									
2002.....	43.1	53.2	37.0	45.1	54.7	39.3	31.0	46.2	22.4
2001.....	43.3	53.8	37.2	45.3	55.2	39.4	30.8	46.1	22.3
2000.....	43.8	55.5	37.0	45.6	56.9	39.1	31.4	47.3	22.6
1999.....	45.1	58.4	37.3	46.8	59.7	39.3	33.3	51.3	23.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Bronchitis, not specified as acute or chronic (J40)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	0.1
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	*
2000.....	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Chronic bronchitis (J41-J42)									
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001.....	0.1	0.2	0.1	0.2	0.2	0.1	0.1	*	*
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
Emphysema (J43)									
2002.....	5.4	7.0	4.4	5.7	7.2	4.8	3.4	5.8	2.0
2001.....	5.8	7.4	4.7	6.1	7.8	5.1	3.3	5.3	2.2
2000.....	6.0	7.8	4.9	6.4	8.1	5.3	3.8	6.2	2.4
1999.....	6.5	8.6	5.2	6.9	8.9	5.6	4.0	6.7	2.3
Other chronic obstructive pulmonary disease (J44)									
2002.....	35.9	44.7	30.7	37.8	46.1	32.8	24.0	36.8	16.8
2001.....	35.7	44.8	30.3	37.5	46.2	32.4	23.6	37.3	16.1
2000.....	35.8	45.9	29.9	37.5	47.3	31.9	23.5	37.3	15.8
1999.....	36.5	47.9	29.8	38.1	49.3	31.6	25.1	40.6	16.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Asthma (J45-J46)									
2002.....	1.5	1.2	1.7	1.2	0.9	1.4	3.4	3.3	3.4
2001.....	1.5	1.2	1.7	1.2	0.9	1.5	3.6	3.2	3.8
2000.....	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.1
1999.....	1.7	1.3	2.0	1.4	1.0	1.7	3.9	3.5	4.1
Bronchiectasis (J47)									
2002.....	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2001.....	0.3	0.2	0.4	0.4	0.2	0.4	0.1	*	*
2000.....	0.3	0.2	0.4	0.4	0.3	0.4	0.1	*	*
1999.....	0.3	0.3	0.4	0.4	0.3	0.4	0.2	*	0.2
Lung diseases due to external agents (J60-J70)									
2002.....	6.5	9.4	4.8	6.5	9.5	4.8	6.8	9.9	5.2
2001.....	6.6	9.5	4.9	6.6	9.6	4.8	7.3	10.1	5.8
2000.....	6.5	9.7	4.7	6.5	9.8	4.7	7.0	10.2	5.2
1999.....	6.1	9.2	4.4	6.0	9.1	4.3	7.1	10.7	5.1
Pneumoconioses and chemical effects (J60-J66,J68)									
2002.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*
2001.....	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*
2000.....	0.4	1.1	0.0	0.4	1.1	0.0	0.2	0.7	*
1999.....	0.4	1.0	0.0	0.4	1.1	0.0	0.3	0.8	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pneumonitis due to solids and liquids (J69)									
2002.....	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1
2001.....	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7
2000.....	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2
1999.....	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1
Pneumonitis due to food and vomit (J69.0)									
2002.....	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1
2001.....	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7
2000.....	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2
1999.....	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1
Other pneumonitis due to solids and liquids (J69.1-J69.8)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other lung diseases due to external agents (J67,J70)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)									
2002.....	0.2	0.4	0.2	0.2	0.3	0.1	0.4	0.7	0.3
2001.....	0.3	0.4	0.2	0.2	0.3	0.2	0.4	0.6	0.2
2000.....	0.2	0.4	0.2	0.2	0.4	0.2	0.4	0.7	0.3
1999.....	0.3	0.4	0.2	0.3	0.4	0.2	0.5	0.8	0.3
Pleural effusion and plaque (J90,J92)									
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.3
2001.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.5	0.4
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.3
1999.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
All other diseases of respiratory system (J80-J84,J93-J98)									
2002.....	7.9	9.4	6.9	8.0	9.5	6.9	7.3	8.5	6.6
2001.....	8.4	10.0	7.3	8.5	10.1	7.3	8.2	9.2	7.7
2000.....	8.4	10.2	7.2	8.5	10.3	7.3	8.1	9.6	7.2
1999.....	8.3	10.0	7.2	8.3	10.1	7.1	8.2	9.5	7.5
XI. Diseases of the digestive system (K00-K92)									
2002.....	29.9	35.0	25.3	30.0	34.9	25.4	32.4	40.1	26.5
2001.....	30.3	35.5	25.7	30.1	34.9	25.6	34.6	44.0	27.5
2000.....	30.4	35.8	25.5	30.4	35.3	25.3	35.3	43.8	28.8
1999.....	30.3	36.0	25.3	30.0	35.5	25.0	35.3	44.1	28.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of oral cavity, salivary glands and jaws (K00-K14)									
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Diseases of other parts of digestive system (K20-K92)									
2002.....	29.9	35.0	25.3	29.9	34.8	25.3	32.3	40.1	26.4
2001.....	30.2	35.4	25.6	30.1	34.9	25.6	34.5	44.0	27.4
2000.....	30.3	35.7	25.5	30.0	35.2	25.2	35.2	43.8	28.7
1999.....	30.3	35.9	25.3	30.0	35.4	25.0	35.3	44.0	28.6
Diseases of esophagus (K20-K22)									
2002.....	0.7	0.9	0.6	0.8	0.9	0.6	0.7	0.8	0.5
2001.....	0.7	0.9	0.7	0.8	0.9	0.7	0.6	0.9	0.5
2000.....	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5
1999.....	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5
Peptic ulcer (K25-K28)									
2002.....	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1
2001.....	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1
2000.....	1.6	2.0	1.4	1.6	2.0	1.4	1.7	2.3	1.3
1999.....	1.7	2.0	1.4	1.7	2.0	1.4	1.9	2.4	1.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Gastric ulcer (K25)									
2002.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.4
2001.....	0.5	0.4	0.4	0.4	0.4	0.4	0.6	0.8	0.4
2000.....	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.4
1999.....	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.5
Duodenal ulcer (K26)									
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.7	0.4	0.6	0.7	0.4	0.4	0.6	0.3
2000.....	0.5	0.7	0.4	0.6	0.7	0.4	0.5	0.6	0.3
1999.....	0.5	0.7	0.4	0.5	0.7	0.4	0.5	0.6	0.4
Peptic ulcer, site unspecified (K27)									
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.3
2001.....	0.6	0.7	0.5	0.6	0.7	0.5	0.6	0.9	0.4
2000.....	0.6	0.8	0.5	0.6	0.8	0.5	0.6	0.7	0.5
1999.....	0.6	0.8	0.6	0.6	0.8	0.6	0.6	0.8	0.5
Gastrojejunal ulcer (K28)									
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Gastritis and duodenitis (K29)									
2002.....	0.1	0.1	0.1	0.1	0.2	0.1	0.1	*	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	*	0.1
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
Dyspepsia and other diseases of stomach and duodenum (K30-K31)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Diseases of appendix (K35-K38)									
2002.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.2
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
Hernia (K40-K46)									
2002.....	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.5
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.8	0.5
2000.....	0.5	0.5	0.5	0.6	0.6	0.6	0.6	0.6	0.6
1999.....	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Crohn's disease and ulcerative colitis (K50-K51)									
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.1
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*	0.2
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
Vascular disorders and obstruction of intestine without hernia (K55-K56)									
2002.....	5.0	4.5	5.3	5.0	4.4	5.3	6.0	6.3	5.7
2001.....	5.1	4.6	5.3	5.0	4.5	5.3	6.2	7.0	5.8
2000.....	5.2	4.7	5.4	5.1	4.5	5.4	6.6	7.3	6.1
1999.....	5.1	4.6	5.4	5.1	4.4	5.4	6.4	6.6	6.2
Diverticular disease of intestine (K57)									
2002.....	1.2	0.9	1.3	1.2	1.0	1.3	1.1	1.0	1.1
2001.....	1.2	1.0	1.3	1.2	1.0	1.4	1.1	0.8	1.2
2000.....	1.2	1.0	1.3	1.2	1.0	1.3	1.0	0.8	1.2
1999.....	1.3	1.0	1.4	1.3	1.1	1.4	1.1	0.8	1.2
Other diseases of intestines and peritoneum (K52,K58-K63)									
2002.....	1.7	1.6	1.7	1.7	1.6	1.7	1.7	1.9	1.5
2001.....	1.7	1.6	1.7	1.7	1.5	1.8	1.9	2.1	1.8
2000.....	1.6	1.5	1.7	1.6	1.5	1.7	1.8	1.9	1.7
1999.....	1.6	1.6	1.7	1.6	1.5	1.7	1.7	2.0	1.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of peritoneum (K65-K66)									
2002.....	0.7	0.6	0.7	0.6	0.6	0.7	1.0	0.9	1.0
2001.....	0.7	0.7	0.6	0.6	0.7	0.6	1.1	1.2	1.0
2000.....	0.7	0.7	0.7	0.7	0.7	0.6	1.1	1.3	1.0
1999.....	0.7	0.8	0.6	0.7	0.7	0.6	1.1	1.2	1.0
Diseases of liver (K70-K76)									
2002.....	12.3	16.7	8.4	12.4	16.9	8.4	12.3	16.9	8.6
2001.....	12.3	16.7	8.3	12.3	16.7	8.3	13.0	18.9	8.5
2000.....	12.2	16.7	8.2	12.1	16.7	8.0	13.2	18.4	9.1
1999.....	12.3	17.0	8.1	12.1	16.9	7.9	13.9	19.6	9.5
Alcoholic liver disease (K70)									
2002.....	4.2	6.3	2.2	4.2	6.5	2.2	3.8	5.8	2.3
2001.....	4.3	6.6	2.2	4.3	6.6	2.2	4.4	6.9	2.4
2000.....	4.3	6.8	2.2	4.3	6.9	2.0	4.6	7.0	2.8
1999.....	4.3	6.9	2.1	4.3	6.8	2.0	5.1	8.1	2.8
Chronic hepatitis, not elsewhere classified (K73)									
2002.....	0.2	0.1	0.2	0.2	0.1	0.3	0.2	*	0.2
2001.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2
2000.....	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.2
1999.....	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Fibrosis and cirrhosis of liver (K74)									
2002.....	5.0	6.4	3.8	5.2	6.6	3.9	4.5	6.1	3.2
2001.....	5.1	6.5	3.8	5.2	6.5	3.9	4.7	6.8	3.0
2000.....	5.1	6.5	3.8	5.1	6.6	3.8	4.7	6.4	3.3
1999.....	5.0	6.6	3.7	5.1	6.6	3.8	4.8	6.7	3.2
Other diseases of liver (K71-K72,K75-K76)									
2002.....	2.9	3.8	2.1	2.8	3.7	2.1	3.8	5.0	2.9
2001.....	2.7	3.5	2.1	2.7	3.4	2.0	3.8	5.0	2.9
2000.....	2.6	3.3	2.0	2.5	3.2	1.9	3.8	4.9	2.9
1999.....	2.7	3.4	2.1	2.5	3.3	1.9	3.9	4.7	3.2
Cholelithiasis and other disorders of gallbladder (K80-K82)									
2002.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
2001.....	1.0	1.2	0.9	1.0	1.2	0.9	1.3	1.4	1.2
2000.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.1	1.2
1999.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.0
Disorders of biliary tract and pancreas (K83-K86)									
2002.....	1.7	2.1	1.3	1.6	2.0	1.3	2.2	3.1	1.7
2001.....	1.8	2.1	1.5	1.7	2.0	1.4	2.5	3.1	2.0
2000.....	1.7	2.1	1.4	1.6	1.9	1.3	2.5	3.3	1.9
1999.....	1.6	2.0	1.3	1.5	1.9	1.3	2.4	3.2	1.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pancreatitis (K85,K86.0-K86.1)									
2002.....	1.2	1.5	0.9	1.2	1.5	0.9	1.6	2.3	1.1
2001.....	1.3	1.5	1.1	1.2	1.5	1.0	1.9	2.4	1.5
2000.....	1.2	1.5	1.0	1.2	1.4	0.9	1.9	2.6	1.3
1999.....	1.2	1.5	1.0	1.1	1.4	0.9	1.9	2.6	1.3
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)									
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.7	0.6
2001.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.7	0.5
2000.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.7	0.5
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.6	0.5
All other diseases of digestive system (K90,K92)									
2002.....	2.9	3.3	2.5	2.8	3.2	2.5	3.6	4.4	3.0
2001.....	2.9	3.4	2.5	2.9	3.3	2.5	3.8	4.7	3.1
2000.....	2.9	3.4	2.6	2.9	3.3	2.5	4.0	4.8	3.4
1999.....	2.9	3.3	2.5	2.8	3.3	2.5	3.6	4.4	3.0
XII. Diseases of the skin and subcutaneous tissue (L00-L98)									
2002.....	1.4	1.3	1.4	1.1	1.1	1.1	3.9	3.3	4.2
2001.....	1.3	1.2	1.4	1.1	1.0	1.2	3.6	3.4	3.7
2000.....	1.4	1.2	1.4	1.1	1.0	1.2	3.9	3.7	4.0
1999.....	1.3	1.3	1.4	1.1	1.0	1.1	4.2	4.0	4.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Infections of skin and subcutaneous tissue (L00-L08)									
2002.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.6
2001.....	0.4	0.4	0.4	0.4	0.4	0.4	0.6	0.5	0.6
2000.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5
1999.....	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.6	0.5
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)									
2002.....	0.9	0.8	1.0	0.7	0.6	0.8	3.3	2.7	3.6
2001.....	0.9	0.7	1.0	0.7	0.6	0.8	3.1	2.9	3.1
2000.....	0.9	0.8	1.1	0.7	0.6	0.8	3.4	3.2	3.5
1999.....	1.0	0.9	1.1	0.8	0.7	0.8	3.8	3.4	3.9
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)									
2002.....	5.0	3.6	5.9	4.8	3.5	5.7	6.4	5.1	7.4
2001.....	5.0	3.6	5.9	4.8	3.5	5.7	6.4	5.1	7.5
2000.....	5.0	3.7	5.9	4.8	3.5	5.7	6.6	5.7	7.4
1999.....	4.9	3.5	5.9	4.7	3.4	5.6	6.3	5.0	7.4
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)									
2002.....	1.0	0.6	1.2	1.0	0.6	1.3	0.7	0.4	0.8
2001.....	0.9	0.5	1.2	1.0	0.5	1.3	0.7	0.4	0.8
2000.....	1.0	0.6	1.2	1.0	0.6	1.3	0.6	0.4	0.8
1999.....	1.0	0.6	1.3	1.0	0.6	1.3	0.6	0.4	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Systemic lupus erythematosus (M32)									
2002.....	0.4	0.1	0.7	0.3	0.1	0.5	1.4	0.2	2.3
2001.....	0.5	0.2	0.8	0.3	0.1	0.6	1.3	0.3	2.1
2000.....	0.5	0.2	0.8	0.3	0.1	0.6	1.3	0.4	2.2
1999.....	0.5	0.2	0.8	0.4	0.1	0.6	1.3	0.3	2.2
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)									
2002.....	1.7	1.2	2.0	1.6	1.1	1.9	2.2	1.7	2.6
2001.....	1.7	1.3	2.0	1.6	1.2	1.9	2.3	1.8	2.6
2000.....	1.7	1.3	2.0	1.7	1.2	2.0	2.4	2.1	2.6
1999.....	1.7	1.3	2.0	1.6	1.2	1.9	2.4	1.9	2.9
Dorsopathies (M40-M54)									
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.2
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2
1999.....	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.4	0.2
Soft tissue disorders (M60-M79)									
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.5
2001.....	0.4	0.5	0.3	0.3	0.5	0.3	0.7	0.9	0.6
2000.....	0.4	0.5	0.3	0.3	0.4	0.2	0.8	0.9	0.7
1999.....	0.4	0.5	0.3	0.3	0.4	0.2	0.7	1.0	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)									
2002.....	1.2	0.8	1.3	1.2	0.8	1.3	1.1	1.3	1.0
2001.....	1.1	0.8	1.3	1.2	0.8	1.3	1.1	1.2	1.0
2000.....	1.1	0.8	1.2	1.1	0.8	1.3	1.1	1.4	0.9
1999.....	1.1	0.7	1.2	1.1	0.7	1.3	0.9	1.0	0.9
Osteoporosis (M80-M81)									
2002.....	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	0.2
2001.....	0.6	0.2	0.8	0.6	0.2	0.8	0.1	*	0.1
2000.....	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*
1999.....	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)									
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.3	0.8
2001.....	0.5	0.6	0.5	0.5	0.6	0.4	1.0	1.2	0.9
2000.....	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.8	0.9	0.7
XIV. Diseases of the genitourinary system (N00-N98)									
2002.....	20.2	23.2	18.3	18.6	21.6	16.7	38.2	44.4	34.6
2001.....	20.2	23.2	18.5	18.5	21.6	16.7	38.6	43.1	36.0
2000.....	19.8	23.0	18.0	18.1	21.4	16.3	37.7	42.6	34.8
1999.....	19.5	22.7	17.7	17.8	20.9	16.1	37.9	44.6	34.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of urinary system (N00-N39)									
2002.....	19.9	22.6	18.1	18.3	21.0	16.6	37.7	43.5	34.2
2001.....	19.9	22.6	18.2	18.2	21.0	16.5	38.1	42.4	35.6
2000.....	19.5	22.4	17.7	17.8	20.8	16.1	37.2	41.8	34.4
1999.....	19.2	22.0	17.5	17.5	20.2	15.9	37.3	43.7	33.7
Glomerular and renal tubulo-interstitial diseases (N00-N15)									
2002.....	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7
2001.....	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7
2000.....	0.7	0.8	0.7	0.6	0.7	0.6	1.1	1.2	1.0
1999.....	0.7	0.9	0.7	0.7	0.9	0.7	1.1	1.5	0.9
Glomerular diseases (N00-N07)									
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.4
2001.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.4
1999.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4
Acute and rapidly progressive nephritic syndrome (N00-N01)									
2002.....	0.0	*	*	0.0	*	*	*	*	*
2001.....	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	*	*	0.0	*	*	*	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Nephrotic syndrome (N04)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
1999.....	0.1	0.1	0.0	0.0	0.0	0.0	0.1	*	*
Other and unspecified glomerular diseases (N02-N03,N05-N07)									
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.3	0.3	0.2
2001.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.4	0.4	0.3
Renal tubulo-interstitial diseases (N10-N15)									
2002.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.4
2001.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.5
2000.....	0.5	0.5	0.5	0.4	0.4	0.4	0.6	0.7	0.6
1999.....	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.9	0.5
Renal failure (N17-N19)									
2002.....	13.9	17.3	11.9	12.5	15.8	10.5	29.3	34.8	25.9
2001.....	13.7	17.1	11.8	12.3	15.8	10.2	29.2	33.4	26.7
2000.....	13.2	16.6	11.3	11.8	15.2	9.8	28.2	32.6	25.5
1999.....	12.8	16.1	10.8	11.3	14.6	9.4	27.9	33.0	24.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Urolithiasis (N20-N23)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Other and unspecified disorders of kidney (N25-N27)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Urinary tract infection, site not specified (N39.0)									
2002.....	4.7	4.0	5.2	4.6	3.8	5.0	6.8	6.7	6.9
2001.....	5.0	4.1	5.4	4.8	4.0	5.2	7.3	7.0	7.4
2000.....	5.0	4.4	5.3	4.9	4.2	5.2	7.2	7.0	7.1
1999.....	5.2	4.4	5.5	5.0	4.2	5.4	7.5	8.0	7.2
Other diseases of urinary system (N28-N36,N39.1-N39.9)									
2002.....	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.8	0.6
2001.....	0.4	0.4	0.3	0.4	0.4	0.3	0.7	0.8	0.6
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
1999.....	0.4	0.5	0.3	0.3	0.5	0.3	0.7	0.9	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of male genital organs (N40-N50)									
2002.....	0.2	0.6	...	0.2	0.5	...	0.3	0.8	...
2001.....	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
2000.....	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
1999.....	0.2	0.6	...	0.2	0.6	...	0.3	0.9	...
Hyperplasia of prostate (N40)									
2002.....	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001.....	0.1	0.5	...	0.2	0.5	...	0.1	0.4	...
2000.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
Other diseases of male genital organs (N41-N50)									
2002.....	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2001.....	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2000.....	0.0	0.1	...	0.0	0.1	...	0.1	0.3	...
1999.....	0.1	0.2	...	0.1	0.1	...	0.1	0.4	...
Disorders of breast (N60-N64)									
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	*	0.0	0.0	*	0.0	*	*	*
1999.....	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Disorders of female genital organs (N70-N98)									
2002.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2001.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.4
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
1999.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.4
Inflammatory diseases of female pelvic organs (N70-N76)									
2002.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Noninflammatory disorders of female genital tract (N80-N98)									
2002.....	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2
2001.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2000.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
1999.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.3
XV. Pregnancy, childbirth and the puerperium (000-099)									
2002.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2001.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2000.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999.....	0.2	...	0.3	0.1	...	0.2	0.4	...	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pregnancy with abortive outcome (000-007)									
2002.....	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	*	...	*
Ectopic pregnancy (000)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Spontaneous abortion (003)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Medical abortion (004)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other abortion (005)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Other and unspecified pregnancy with abortive outcome (001-002,006-007)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Other direct obstetric deaths (010-092)									
2002.....	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
2001.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.6
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.5
1999.....	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
Eclampsia and pre-eclampsia (011,013-016)									
2002.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)									
2002.....	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*
Complications predominately related to the puerperium (085-092)									
2002.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2001.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2000.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Obstetric embolism (088)									
2002.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*
Other complications predominately related to the puerperium (085-087,089-092)									
2002.....	0.0	...	0.0	*	...	*	0.1	...	0.1
2001.....	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)									
2002.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2001.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Obstetric death of unspecified cause (095)									
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Other deaths related to pregnancy, childbirth and the puerperium (096-097)									
2002.....	0.0	...	0.0	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Indirect obstetric deaths (098-099)									
2002.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intrauterine hypoxia and birth asphyxia (P20-P21)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2000.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
Respiratory distress of newborn (P22)									
2002.....	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.7	0.7
2001.....	0.3	0.4	0.3	0.3	0.3	0.3	0.7	0.9	0.6
2000.....	0.4	0.5	0.3	0.3	0.4	0.2	0.8	0.9	0.6
1999.....	0.4	0.5	0.3	0.3	0.4	0.3	0.9	1.0	0.7
Other respiratory conditions originating in the perinatal period (P23-P28)									
2002.....	0.4	0.5	0.4	0.4	0.4	0.3	0.9	1.0	0.8
2001.....	0.5	0.6	0.5	0.4	0.5	0.4	1.1	1.2	1.0
2000.....	0.6	0.6	0.5	0.5	0.5	0.4	1.2	1.4	1.0
1999.....	0.6	0.7	0.5	0.5	0.6	0.4	1.2	1.4	1.1
Infections specific to the perinatal period (P35-P39)									
2002.....	0.3	0.4	0.3	0.3	0.3	0.2	0.6	0.6	0.6
2001.....	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.6	0.6
2000.....	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
1999.....	0.3	0.4	0.3	0.2	0.3	0.2	0.7	0.8	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)									
2002.....	0.8	0.9	0.7	0.7	0.8	0.6	1.4	1.6	1.3
2001.....	0.8	1.0	0.7	0.7	0.8	0.6	1.5	1.8	1.3
2000.....	1.0	1.1	0.8	0.8	0.9	0.7	1.9	2.3	1.6
1999.....	1.0	1.1	0.8	0.8	0.9	0.7	2.0	2.2	1.9
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)									
2002.....	3.7	3.8	3.5	3.7	3.8	3.5	4.1	4.4	3.9
2001.....	3.6	3.8	3.4	3.6	3.8	3.4	4.1	4.3	3.9
2000.....	3.8	4.1	3.6	3.7	4.0	3.5	4.4	4.8	4.1
1999.....	3.8	3.9	3.6	3.7	3.9	3.6	4.3	4.7	4.1
Anencephaly and similar malformations (Q00)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
Spina bifida (Q05)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.1	0.0	0.1	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)									
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.4
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.4
Congenital malformations of heart (Q20-Q24)									
2002.....	1.1	1.2	1.0	1.1	1.2	1.0	1.2	1.3	1.2
2001.....	1.1	1.2	1.0	1.1	1.2	1.0	1.4	1.4	1.3
2000.....	1.2	1.3	1.2	1.2	1.3	1.1	1.4	1.6	1.3
1999.....	1.3	1.4	1.2	1.3	1.4	1.2	1.5	1.7	1.3
Other congenital malformations of circulatory system (Q25-Q28)									
2002.....	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.6	0.4
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.4
Congenital malformations of respiratory system (Q30-Q34)									
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)									
2002.....	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9
2001.....	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9
2000.....	0.9	1.1	0.8	0.9	1.0	0.8	1.1	1.2	0.9
1999.....	0.9	1.1	0.9	0.9	1.0	0.8	1.1	1.2	0.9
Down's syndrome (Q90)									
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)									
2002.....	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5
2001.....	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4
2000.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5
1999.....	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)									
2002.....	10.3	11.4	9.0	9.9	10.8	8.7	14.5	18.0	11.8
2001.....	11.3	12.7	9.8	11.0	12.2	9.5	15.1	17.7	12.8
2000.....	11.4	13.1	9.7	11.0	12.6	9.3	16.5	19.7	13.6
1999.....	9.7	11.0	8.4	9.1	10.1	7.9	15.4	19.1	12.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Senility (R54)									
2002.....	1.5	1.2	1.6	1.5	1.3	1.6	1.2	1.1	1.2
2001.....	1.5	1.2	1.6	1.5	1.3	1.6	1.3	1.3	1.3
2000.....	1.4	1.1	1.5	1.4	1.2	1.5	1.4	1.0	1.5
1999.....	1.2	1.0	1.2	1.2	1.0	1.3	1.0	1.1	1.0
Ill-defined and unknown causes of mortality (R95-R99)									
2002.....	5.9	7.2	4.6	5.6	6.8	4.4	8.9	11.7	6.7
2001.....	7.1	8.7	5.6	6.8	8.4	5.3	9.5	11.6	7.7
2000.....	7.4	9.1	5.8	7.0	8.7	5.4	10.9	13.9	8.4
1999.....	6.0	7.3	4.8	5.5	6.6	4.4	10.1	13.3	7.6
Sudden infant death syndrome (R95)									
2002.....	0.8	0.9	0.7	0.7	0.8	0.5	1.4	1.6	1.3
2001.....	0.8	0.9	0.6	0.6	0.7	0.5	1.6	1.7	1.4
2000.....	0.9	1.1	0.8	0.7	0.9	0.6	1.9	2.1	1.6
1999.....	1.0	1.1	0.8	0.8	0.9	0.7	1.9	2.1	1.7
Other ill-defined and unknown causes of mortality (R96-R99)									
2002.....	5.1	6.3	4.0	4.9	6.0	3.8	7.5	10.0	5.4
2001.....	6.3	7.8	4.9	6.2	7.6	4.8	7.9	9.8	6.3
2000.....	6.5	8.1	5.0	6.3	7.8	4.8	9.0	11.8	6.8
1999.....	5.1	6.2	4.0	4.7	5.7	3.7	8.2	11.1	6.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)									
2002.....	2.9	3.0	2.8	2.8	2.8	2.7	4.5	5.3	3.9
2001.....	2.8	2.8	2.7	2.6	2.6	2.6	4.3	4.8	3.8
2000.....	2.7	2.8	2.5	2.6	2.7	2.4	4.2	4.8	3.7
1999.....	2.6	2.7	2.4	2.4	2.5	2.3	4.2	4.8	3.8
XX. External causes of mortality (*U01-*U03,V01-Y89)									
2002.....	56.7	82.7	32.6	55.9	80.7	32.9	67.2	107.7	32.6
2001 2/.....	56.1	82.6	31.8	55.2	80.3	31.8	68.2	108.4	33.7
2000.....	53.9	79.5	30.5	52.5	76.6	30.4	67.9	109.1	32.9
1999.....	54.3	80.1	31.0	52.7	76.9	30.7	69.8	111.7	34.6
Accidents (unintentional injuries) (V01-X59,Y85-Y86)									
2002.....	36.9	51.5	23.5	37.5	52.0	24.0	36.9	56.2	21.3
2001.....	35.7	50.2	22.5	36.0	50.3	22.8	37.6	56.9	22.0
2000.....	34.9	49.3	22.0	35.1	49.2	22.2	37.7	58.1	21.3
1999.....	35.3	49.8	22.3	35.2	49.3	22.4	40.1	61.3	23.1
Transport accidents (V01-V99,Y85)									
2002.....	16.7	23.9	9.9	17.1	24.2	10.2	16.0	25.0	8.5
2001.....	16.6	23.7	9.8	16.8	24.0	10.0	16.5	25.5	8.9
2000.....	16.6	23.7	10.0	16.8	23.8	10.2	16.7	26.1	8.7
1999.....	16.6	23.6	10.2	16.8	23.6	10.4	17.1	26.5	9.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Railway accidents (V05,V15,V80.6,V81.2-V81.9)									
2002.....	0.2	0.3	0.0	0.2	0.3	0.0	0.3	0.5	*
2001.....	0.2	0.3	0.1	0.2	0.3	0.0	0.2	0.4	*
2000.....	0.2	0.3	0.1	0.2	0.3	0.0	0.3	0.5	*
1999.....	0.2	0.3	0.0	0.2	0.3	0.1	0.2	0.4	*
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)									
2002.....	15.7	22.1	9.6	16.0	22.4	9.8	15.0	23.2	8.2
2001.....	15.3	21.8	9.3	15.6	22.0	9.5	15.4	23.5	8.5
2000.....	15.4	21.7	9.5	15.6	21.8	9.8	15.7	24.4	8.4
1999.....	15.2	21.3	9.6	15.3	21.3	9.7	15.9	24.3	9.0
Pedestrian involved in collision with motor vehicle (V02-V04)									
2002.....	1.3	1.8	0.8	1.2	1.7	0.7	1.7	2.8	0.9
2001.....	1.3	1.8	0.7	1.2	1.7	0.6	1.8	2.7	1.0
2000.....	1.2	1.8	0.7	1.1	1.6	0.7	1.8	3.0	0.9
1999.....	1.2	1.8	0.7	1.1	1.6	0.7	1.9	3.2	0.9
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)									
2002.....	0.2	0.3	0.0	0.2	0.4	0.0	0.2	0.4	*
2001.....	0.2	0.4	0.0	0.2	0.4	0.0	0.3	0.6	*
2000.....	0.2	0.4	0.0	0.2	0.4	0.0	0.3	0.6	*
1999.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51, V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)									
2002.....	1.7	2.6	0.9	1.8	2.6	1.0	1.9	3.1	0.8
2001.....	1.5	2.2	0.8	1.5	2.2	0.8	1.7	2.8	0.8
2000.....	1.4	2.0	0.7	1.4	2.0	0.8	1.6	2.6	0.7
1999.....	1.3	2.0	0.7	1.4	2.0	0.7	1.7	2.7	0.8
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)									
2002.....	3.9	5.0	2.9	4.0	5.1	3.0	3.5	4.8	2.4
2001.....	3.7	4.7	2.8	3.8	4.8	2.9	3.6	4.9	2.4
2000.....	3.7	4.6	2.9	3.8	4.7	3.0	3.5	4.9	2.3
1999.....	3.7	4.6	2.9	3.7	4.6	3.0	3.6	5.1	2.4
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)									
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)									
2002.....	1.5	2.1	0.9	1.6	2.2	1.0	1.2	1.9	0.6
2001.....	1.4	1.9	0.8	1.4	2.0	0.8	1.1	1.7	0.5
2000.....	1.3	1.9	0.7	1.4	2.0	0.8	1.1	1.7	0.6
1999.....	1.3	1.9	0.8	1.3	1.9	0.8	1.2	1.7	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Occupant of special-use motor vehicle involved in any accident (V83-V86)									
2002.....	0.3	0.6	0.1	0.4	0.7	0.1	0.1	0.1	*
2001.....	0.4	0.7	0.1	0.4	0.7	0.1	0.1	0.2	*
2000.....	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.3	*
1999.....	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.3	*
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)									
2002.....	5.4	7.4	3.6	5.5	7.5	3.7	5.4	8.1	3.2
2001.....	5.7	7.9	3.8	5.8	7.9	3.9	5.8	8.7	3.5
2000.....	6.0	8.2	4.1	6.1	8.2	4.1	6.2	9.2	3.7
1999.....	6.0	8.1	4.1	6.1	8.2	4.2	6.2	9.1	3.9
Streetcar accidents (V82.2-V82.9)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)									
2002.....	0.2	0.4	0.1	0.2	0.3	0.1	0.2	0.4	*
2001.....	0.3	0.5	0.2	0.3	0.5	0.1	0.4	0.6	0.2
2000.....	0.4	0.5	0.2	0.4	0.5	0.2	0.3	0.5	0.2
1999.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Water transport accidents (V90-V94)									
2002.....	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*
2001.....	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.4	*
2000.....	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*
1999.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*
Air and space transport accidents (V95-V97)									
2002.....	0.2	0.4	0.0	0.3	0.5	0.1	*	*	*
2001.....	0.3	0.5	0.2	0.4	0.6	0.2	0.1	*	*
2000.....	0.3	0.5	0.1	0.3	0.5	0.1	*	*	*
1999.....	0.2	0.4	0.1	0.3	0.5	0.1	*	*	*
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)									
2002.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.5	*
2001.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	*
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*
1999.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1
Falls (W00-W19)									
2002.....	5.6	7.4	4.3	5.9	7.6	4.6	3.1	4.9	2.0
2001.....	5.3	7.2	3.9	5.6	7.5	4.1	3.3	5.1	2.0
2000.....	4.8	6.6	3.6	5.0	6.8	3.7	2.9	4.6	1.8
1999.....	4.8	6.6	3.5	5.0	6.8	3.7	3.4	5.3	2.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Fall on same level (W00-W09,W18)									
2002.....	1.8	2.3	1.5	1.9	2.4	1.6	0.8	1.3	0.5
2001.....	1.4	1.8	1.1	1.5	1.9	1.2	0.7	1.0	0.5
2000.....	1.2	1.5	1.0	1.2	1.5	1.0	0.5	0.7	0.4
1999.....	0.8	1.1	0.7	0.8	1.1	0.7	0.6	0.8	0.4
Fall from one level to another (W10-W17)									
2002.....	1.1	1.9	0.5	1.2	2.0	0.6	0.7	1.2	0.4
2001.....	1.2	2.0	0.5	1.2	2.0	0.6	0.9	1.5	0.3
2000.....	1.0	1.8	0.4	1.1	1.9	0.5	0.7	1.3	0.3
1999.....	1.1	1.9	0.5	1.2	1.9	0.6	0.8	1.3	0.3
Unspecified fall (W19)									
2002.....	2.6	3.2	2.2	2.8	3.3	2.4	1.6	2.4	1.1
2001.....	2.7	3.5	2.2	2.8	3.6	2.3	1.7	2.6	1.2
2000.....	2.6	3.3	2.1	2.7	3.4	2.2	1.7	2.5	1.2
1999.....	2.9	3.7	2.3	3.0	3.8	2.4	2.1	3.1	1.4
Accidental discharge of firearms (W32-W34)									
2002.....	0.3	0.4	0.1	0.2	0.4	0.1	0.4	0.7	*
2001.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.5	*
2000.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1
1999.....	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.7	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental explosion (W39-W40)									
2002.....	0.0	0.1	*	0.1	0.1	*	*	*	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Accidental drowning and submersion (W65-W74)									
2002.....	1.2	2.0	0.5	1.1	1.8	0.4	1.6	2.7	0.5
2001.....	1.1	1.8	0.5	1.1	1.7	0.5	1.5	2.6	0.5
2000.....	1.2	2.0	0.5	1.2	1.8	0.5	1.5	2.6	0.6
1999.....	1.3	2.0	0.5	1.2	1.9	0.5	1.7	3.0	0.5
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)									
2002.....	1.4	1.8	1.2	1.4	1.8	1.1	2.1	2.7	1.6
2001.....	1.5	1.9	1.2	1.4	1.8	1.2	2.2	3.1	1.6
2000.....	1.6	2.0	1.3	1.5	1.9	1.3	2.1	2.9	1.7
1999.....	1.4	1.8	1.2	1.4	1.7	1.2	2.2	2.9	1.7
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)									
2002.....	0.5	0.7	0.3	0.5	0.6	0.3	0.7	0.8	0.5
2001.....	0.5	0.6	0.3	0.5	0.6	0.3	0.6	0.8	0.5
2000.....	0.5	0.7	0.3	0.5	0.7	0.3	0.6	0.8	0.4
1999.....	0.6	0.8	0.4	0.6	0.8	0.4	0.7	1.0	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental exposure to electric current (W85-W87)									
2002.....	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*
2001.....	0.1	0.3	*	0.2	0.3	*	0.1	0.2	*
2000.....	0.1	0.3	0.0	0.2	0.3	*	0.1	0.2	*
1999.....	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*
Accidental exposure to radiation (W88-W91)									
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)									
2002.....	1.1	1.4	0.8	0.9	1.3	0.7	2.4	3.3	1.7
2001.....	1.2	1.5	0.9	1.0	1.3	0.7	2.7	3.8	1.9
2000.....	1.2	1.6	0.9	1.0	1.3	0.7	3.0	4.2	2.2
1999.....	1.2	1.6	0.9	1.0	1.4	0.7	3.0	4.0	2.3
Accidental contact with heat and hot substances (X10-X19)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Lightning (X33)									
2002.....	0.0	0.0	*	0.0	0.1	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.1	*	0.0	0.1	*	*	*	*
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)									
2002.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Cataclysmic storm and flood (X37-X38)									
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)									
2002.....	6.1	8.4	3.7	6.3	8.7	3.9	6.8	10.1	3.9
2001.....	4.9	7.0	2.9	5.0	7.1	2.9	6.3	9.5	3.7
2000.....	4.5	6.6	2.5	4.5	6.6	2.5	6.0	9.3	3.3
1999.....	4.4	6.4	2.3	4.3	6.3	2.3	6.2	9.8	3.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)									
2002.....	5.7	7.8	3.5	5.9	8.1	3.7	6.3	9.3	3.7
2001.....	4.6	6.5	2.7	4.7	6.5	2.7	5.9	8.7	3.3
2000.....	4.1	6.0	2.3	4.2	6.0	2.3	5.6	8.6	3.1
1999.....	4.0	5.9	2.1	3.9	5.8	2.1	5.8	9.0	2.9
Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)									
2002.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2000.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
Accidental poisoning by and exposure to other gases and vapors (X47)									
2002.....	0.2	0.4	0.1	0.3	0.3	0.1	0.2	0.4	0.1
2001.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1
1999.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)									
2002.....	3.7	5.0	2.6	3.8	5.0	2.7	3.7	5.4	2.4
2001.....	4.0	5.3	2.9	4.1	5.4	2.9	3.9	5.5	2.7
2000.....	4.0	5.3	2.8	4.0	5.3	2.8	4.0	6.5	2.3
1999.....	4.4	5.9	3.1	4.4	5.8	3.1	4.9	7.7	3.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)									
2002.....	10.9	18.4	4.2	12.0	20.0	4.7	5.3	9.8	1.6
2001 2/.....	10.7	18.2	4.0	11.7	19.6	4.5	5.5	9.8	1.8
2000.....	10.4	17.7	4.0	11.3	19.1	4.3	5.5	10.0	1.8
1999.....	10.5	17.8	4.0	11.3	19.0	4.4	5.6	10.3	1.6
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)									
2002.....	1.3	1.3	1.4	1.5	1.4	1.6	0.4	0.5	0.4
2001.....	1.3	1.2	1.3	1.4	1.4	1.4	0.5	0.4	0.5
2000.....	1.2	1.1	1.2	1.3	1.3	1.4	0.5	0.5	0.4
1999.....	1.1	1.1	1.2	1.3	1.2	1.3	0.4	0.5	0.4
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)									
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)									
2002.....	0.5	0.8	0.2	0.6	0.9	0.2	0.1	0.2	*
2001.....	0.5	0.8	0.2	0.6	0.9	0.3	0.1	0.2	*
2000.....	0.5	0.8	0.2	0.6	0.9	0.3	0.1	0.2	*
1999.....	0.5	0.9	0.3	0.6	1.0	0.3	0.1	0.2	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)									
2002.....	2.2	3.7	0.8	2.4	4.0	0.8	1.1	2.1	0.3
2001.....	2.2	3.7	0.7	2.3	3.9	0.7	1.1	2.0	0.3
2000.....	2.0	3.4	0.7	2.1	3.6	0.7	1.2	2.1	0.3
1999.....	1.9	3.3	0.6	2.0	3.4	0.7	1.0	2.0	0.2
Intentional self-harm (suicide) by discharge of firearms (X72-X74)									
2002.....	5.9	11.1	1.4	6.5	12.1	1.6	3.0	5.9	0.5
2001.....	5.9	11.0	1.4	6.5	12.0	1.6	3.1	6.0	0.6
2000.....	5.9	11.0	1.5	6.5	11.9	1.6	3.1	6.0	0.7
1999.....	6.0	11.1	1.5	6.5	12.0	1.7	3.3	6.5	0.6
Intentional self-harm (suicide) by jumping from a high place (X80)									
2002.....	0.2	0.4	0.1	0.3	0.4	0.1	0.2	0.3	*
2001.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
1999.....	0.3	0.4	0.1	0.3	0.4	0.1	0.2	0.3	*
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)									
2002.....	0.6	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.2
2001 2/.....	0.6	0.9	0.3	0.6	1.0	0.3	0.5	0.7	0.3
2000.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2
1999.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)									
2002.....	6.1	9.4	2.8	3.7	5.3	2.0	21.0	36.4	6.9
2001 2/.....	7.1	10.8	3.3	4.9	7.1	2.6	21.2	36.2	7.4
2000.....	5.9	9.0	2.8	3.6	5.2	2.1	20.5	35.4	7.1
1999.....	6.0	9.1	2.9	3.8	5.4	2.1	20.1	34.2	7.3
Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)									
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Assault (homicide) by hanging, strangulation and suffocation (X91)									
2002.....	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2001.....	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2000.....	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.3	0.8
1999.....	0.2	0.2	0.3	0.2	0.2	0.2	0.6	0.4	0.8
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)									
2002.....	4.1	6.8	1.3	2.2	3.4	1.0	15.6	28.7	3.5
2001.....	3.9	6.6	1.3	2.2	3.5	1.0	14.8	27.4	3.0
2000.....	3.8	6.3	1.2	2.1	3.2	0.9	14.6	26.9	3.1
1999.....	3.8	6.3	1.3	2.1	3.3	1.0	14.3	26.1	3.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2001.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
2000.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
1999.....	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.2
Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09, Y87.1)									
2002.....	0.8	1.1	0.5	0.6	0.9	0.4	2.2	3.4	1.2
2001 2/.....	2.0	2.8	1.1	1.8	2.6	1.0	3.2	4.6	2.1
2000.....	1.0	1.4	0.7	0.7	0.9	0.5	2.9	4.4	1.7
1999.....	1.0	1.3	0.7	0.8	1.0	0.5	2.5	3.6	1.5
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)									
2002.....	1.7	2.2	1.2	1.7	2.2	1.2	2.1	3.1	1.2
2001.....	1.5	2.0	0.9	1.5	2.0	1.0	1.8	2.9	0.9
2000.....	1.3	1.9	0.8	1.3	1.8	0.8	1.9	3.0	1.0
1999.....	1.4	2.0	0.9	1.4	1.8	0.9	2.0	3.2	0.9
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)									
2002.....	1.1	1.3	0.8	1.1	1.4	0.9	1.3	1.7	0.9
2001.....	1.0	1.3	0.7	1.0	1.3	0.7	1.1	1.8	0.5
2000.....	0.8	1.2	0.6	0.9	1.2	0.6	1.1	1.6	0.6
1999.....	0.9	1.2	0.6	0.9	1.1	0.6	1.2	1.9	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)									
2002.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)									
2002.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.1	*	0.0	0.1	*	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)									
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999.....	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*
Falling, jumping or pushed from a high place, undetermined intent (Y30)									
2002.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)									
2002.....	0.4	0.5	0.2	0.4	0.5	0.2	0.6	1.1	0.3
2001.....	0.3	0.5	0.2	0.3	0.4	0.2	0.5	0.8	0.3
2000.....	0.3	0.5	0.2	0.3	0.4	0.2	0.6	1.0	0.3
1999.....	0.3	0.5	0.2	0.3	0.4	0.2	0.5	0.9	0.2
Legal intervention (Y35,Y89.0)									
2002.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2001.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2000.....	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*
1999.....	0.1	0.3	*	0.1	0.2	*	0.3	0.7	*
Legal execution (Y35.5)									
2002.....	0.0	0.1	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.1	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	0.1	0.2	*
1999.....	0.0	0.1	*	0.0	0.1	*	0.1	0.2	*
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)									
2002.....	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*
2001.....	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*
2000.....	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*
1999.....	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Operations of war and their sequelae (Y36,Y89.1)									
2002.....	0.0	0.0	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)									
2002.....	1.0	1.0	1.0	0.9	1.0	0.9	1.6	1.6	1.6
2001.....	1.1	1.1	1.0	1.0	1.1	0.9	2.0	2.3	1.7
2000.....	1.1	1.2	1.0	1.0	1.2	0.9	1.9	1.9	1.9
1999.....	1.0	1.1	0.9	1.0	1.1	0.9	1.8	2.0	1.7
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)									
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)									
2002.....	0.0	0.0	0.1	0.0	0.0	0.0	0.1	*	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999.....	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2002

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992) and year	All races 1/			White			Black		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)									
2002.....	0.8	0.9	0.8	0.8	0.9	0.8	1.4	1.4	1.3
2001.....	0.9	1.0	0.9	0.8	0.9	0.8	1.6	2.0	1.4
2000.....	0.9	1.1	0.9	0.9	1.0	0.8	1.6	1.7	1.6
1999.....	0.9	1.0	0.8	0.8	0.9	0.8	1.5	1.7	1.4

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Includes races other than white and black.

2/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.