

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
All races, both sexes												
2003.....	2,448,288	28,025	4,965	2,898	4,056	13,595	19,973	18,250	23,050	34,055	55,406	78,173
2002.....	2,443,387	28,034	4,858	3,018	4,132	13,812	19,234	17,959	23,396	35,347	55,793	76,065
2001.....	2,416,425	27,568	5,107	3,093	4,002	13,555	18,697	18,211	23,472	36,895	54,779	73,684
2000.....	2,403,351	28,035	4,979	3,253	4,160	13,563	17,744	17,681	22,770	36,140	53,658	70,832
1999.....	2,391,399	27,937	5,249	3,474	4,121	13,778	16,878	17,703	23,363	36,804	52,452	68,120
All races, male												
2003.....	1,201,964	15,902	2,826	1,640	2,510	9,706	14,964	13,043	15,559	21,741	34,694	49,139
2002.....	1,199,264	15,717	2,806	1,702	2,496	9,844	14,572	12,954	15,782	22,626	34,967	47,701
2001.....	1,183,421	15,477	2,899	1,727	2,441	9,766	14,197	13,006	15,751	23,727	34,437	46,567
2000.....	1,177,578	15,718	2,824	1,850	2,551	9,697	13,374	12,619	15,271	23,252	34,045	45,121
1999.....	1,175,460	15,646	2,975	1,964	2,528	9,780	12,634	12,542	15,734	23,681	33,437	43,303
All races, female												
2003.....	1,246,324	12,123	2,139	1,258	1,546	3,889	5,009	5,207	7,491	12,314	20,712	29,034
2002.....	1,244,123	12,317	2,052	1,316	1,636	3,968	4,662	5,005	7,614	12,721	20,826	28,364
2001.....	1,233,004	12,091	2,208	1,366	1,561	3,789	4,500	5,205	7,721	13,168	20,342	27,117
2000.....	1,225,773	12,317	2,155	1,403	1,609	3,866	4,370	5,062	7,499	12,888	19,613	25,711
1999.....	1,215,939	12,291	2,274	1,510	1,593	3,998	4,244	5,161	7,629	13,123	19,015	24,817

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

White, both sexes

2003.....	2,103,714	18,440	3,509	2,111	2,920	10,392	14,584	12,991	16,698	25,012	41,736	59,135
2002.....	2,102,589	18,369	3,406	2,168	2,970	10,613	14,028	12,703	16,868	25,912	42,181	57,690
2001.....	2,079,691	17,955	3,670	2,250	2,920	10,235	13,513	12,988	17,006	27,238	40,899	55,634
2000.....	2,071,287	18,144	3,494	2,359	3,091	10,273	12,745	12,427	16,292	26,633	39,863	53,131
1999.....	2,061,348	18,067	3,690	2,450	3,043	10,355	11,997	12,569	16,831	27,024	38,583	50,895

White, male

2003.....	1,025,650	10,500	1,981	1,205	1,803	7,308	10,911	9,342	11,472	16,344	26,739	37,898
2002.....	1,025,196	10,433	1,958	1,219	1,786	7,474	10,608	9,219	11,579	16,840	27,115	36,894
2001.....	1,011,218	10,099	2,094	1,232	1,782	7,264	10,265	9,381	11,597	17,880	26,363	35,774
2000.....	1,007,191	10,177	2,004	1,348	1,907	7,242	9,626	8,943	11,197	17,529	25,849	34,599
1999.....	1,005,335	10,197	2,084	1,376	1,852	7,222	8,981	9,015	11,534	17,792	25,122	32,778

White, female

2003.....	1,078,064	7,940	1,528	906	1,117	3,084	3,673	3,649	5,226	8,668	14,997	21,237
2002.....	1,077,393	7,936	1,448	949	1,184	3,139	3,420	3,484	5,289	9,072	15,066	20,796
2001.....	1,068,473	7,856	1,576	1,018	1,138	2,971	3,248	3,607	5,409	9,358	14,536	19,860
2000.....	1,064,096	7,967	1,490	1,011	1,184	3,031	3,119	3,484	5,095	9,104	14,014	18,532
1999.....	1,056,013	7,870	1,606	1,074	1,191	3,133	3,016	3,554	5,297	9,232	13,461	18,117

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Black, both sexes												
2003.....	291,300	8,402	1,199	637	914	2,572	4,573	4,524	5,418	7,823	11,961	16,680
2002.....	290,051	8,524	1,196	701	965	2,602	4,443	4,550	5,621	8,227	12,008	16,184
2001.....	287,709	8,498	1,181	700	887	2,710	4,437	4,467	5,561	8,385	12,302	15,867
2000.....	285,826	8,771	1,248	756	887	2,717	4,332	4,541	5,698	8,352	12,205	15,735
1999.....	285,064	8,822	1,311	880	909	2,849	4,217	4,444	5,671	8,663	12,378	15,304
Black, male												
2003.....	148,022	4,740	699	355	567	1,974	3,474	3,219	3,474	4,620	6,924	9,824
2002.....	146,835	4,652	702	402	596	1,963	3,401	3,229	3,619	4,996	6,853	9,501
2001.....	145,908	4,764	672	409	551	2,080	3,399	3,117	3,556	5,022	7,113	9,495
2000.....	145,184	4,901	692	431	543	2,045	3,273	3,163	3,587	5,015	7,236	9,350
1999.....	145,703	4,897	746	519	577	2,157	3,191	3,073	3,650	5,171	7,396	9,392
Black, female												
2003.....	143,278	3,662	500	282	347	598	1,099	1,305	1,944	3,203	5,037	6,856
2002.....	143,216	3,872	494	299	369	639	1,042	1,321	2,002	3,231	5,155	6,683
2001.....	141,801	3,734	509	291	336	630	1,038	1,350	2,005	3,363	5,189	6,372
2000.....	140,642	3,870	556	325	344	672	1,059	1,378	2,111	3,337	4,969	6,385
1999.....	139,361	3,925	565	361	332	692	1,026	1,371	2,021	3,492	4,982	5,912

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

American Indian 1/, both sexes

2003.....	13,147	335	95	55	82	286	335	288	355	539	705	842
2002.....	12,415	343	90	48	78	267	316	290	362	512	657	751
2001.....	11,977	373	103	44	85	273	284	265	331	531	601	747
2000.....	11,363	323	91	50	63	251	268	272	324	472	609	611
1999.....	11,312	343	82	48	57	245	295	281	364	453	574	628

American Indian 1/, male

2003.....	7,106	195	55	30	56	195	255	193	251	351	433	516
2002.....	6,750	191	49	21	44	178	238	218	239	335	414	455
2001.....	6,466	211	52	29	46	186	214	194	223	355	367	437
2000.....	6,185	193	49	28	33	178	191	192	218	284	391	389
1999.....	6,092	180	48	21	34	179	217	180	238	293	352	415

American Indian 1/, female

2003.....	6,041	140	40	25	26	91	80	95	104	188	272	326
2002.....	5,665	152	41	27	34	89	78	72	123	177	243	296
2001.....	5,511	162	51	15	39	87	70	71	108	176	234	310
2000.....	5,178	130	42	22	30	73	77	80	106	188	218	222
1999.....	5,220	163	34	27	23	66	78	101	126	160	222	213

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

Asian or Pacific Islander 2/, both sexes

2003.....	40,127	848	162	95	140	345	481	447	579	681	1,004	1,516
2002.....	38,332	798	166	101	119	330	447	416	545	696	947	1,440
2001.....	37,048	742	153	99	110	337	463	491	574	741	977	1,436
2000.....	34,875	797	146	88	119	322	399	441	456	683	981	1,355
1999.....	33,675	705	166	96	112	329	369	409	497	664	917	1,293

Asian or Pacific Islander 2/, male

2003.....	21,186	467	91	50	84	229	324	289	362	426	598	901
2002.....	20,483	441	97	60	70	229	325	288	345	455	585	851
2001.....	19,829	403	81	57	62	236	319	314	375	470	594	861
2000.....	19,018	447	79	43	68	232	284	321	269	424	569	783
1999.....	18,330	372	97	48	65	222	245	274	312	425	567	718

Asian or Pacific Islander 2/, female

2003.....	18,941	381	71	45	56	116	157	158	217	255	406	615
2002.....	17,849	357	69	41	49	101	122	128	200	241	362	589
2001.....	17,219	339	72	42	48	101	144	177	199	271	383	575
2000.....	15,857	350	67	45	51	90	115	120	187	259	412	572
1999.....	15,345	333	69	48	47	107	124	135	185	239	350	575

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Certain other intestinal infections (A04,A07-A09)												
All races, both sexes												
2003.....	3,090	10	4	4	3	1	3	-	8	4	10	21
2002.....	2,465	13	8	1	2	4	2	3	1	5	13	25
2001.....	1,537	10	18	1	3	-	-	1	3	4	8	9
2000.....	1,328	13	10	-	-	-	1	2	1	7	12	11
1999.....	1,108	12	8	2	1	1	4	3	3	2	5	11
All races, male												
2003.....	1,109	6	4	1	1	1	1	-	4	3	2	11
2002.....	947	3	5	1	-	3	-	1	-	2	6	12
2001.....	572	7	11	1	1	-	-	-	2	4	4	4
2000.....	524	10	6	-	-	-	-	2	1	4	6	7
1999.....	420	8	4	2	1	1	2	2	3	1	3	5
All races, female												
2003.....	1,981	4	-	3	2	-	2	-	4	1	8	10
2002.....	1,518	10	3	-	2	1	2	2	1	3	7	13
2001.....	965	3	7	-	2	-	-	1	1	-	4	5
2000.....	804	3	4	-	-	-	1	-	-	3	6	4
1999.....	688	4	4	-	-	-	2	1	-	1	2	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2003.....	2,877	5	3	4	2	1	2	-	6	4	7	18
2002.....	2,287	7	6	1	2	4	2	2	-	4	11	21
2001.....	1,417	6	15	1	2	-	-	-	2	3	7	6
2000.....	1,224	8	7	-	-	-	1	1	1	3	8	8
1999.....	1,026	8	7	-	1	1	3	3	3	2	2	8

White, male

2003.....	1,036	2	3	1	1	1	1	-	3	3	2	9
2002.....	872	1	4	1	-	3	-	1	-	1	5	11
2001.....	518	4	9	1	-	-	-	-	2	3	3	2
2000.....	478	6	5	-	-	-	-	1	1	2	5	5
1999.....	389	6	3	-	1	1	2	2	3	1	2	3

White, female

2003.....	1,841	3	-	3	1	-	1	-	3	1	5	9
2002.....	1,415	6	2	-	2	1	2	1	-	3	6	10
2001.....	899	2	6	-	2	-	-	-	-	-	4	4
2000.....	746	2	2	-	-	-	1	-	-	1	3	3
1999.....	637	2	4	-	-	-	1	1	-	1	-	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain other intestinal infections (A04,A07-A09)

Black, both sexes

2003.....	186	5	1	-	1	-	1	-	1	-	2	3
2002.....	156	5	1	-	-	-	-	1	1	1	2	3
2001.....	101	3	3	-	1	-	-	1	1	1	1	3
2000.....	84	4	2	-	-	-	-	1	-	3	3	3
1999.....	74	4	-	2	-	-	1	-	-	-	3	2

Black, male

2003.....	66	4	1	-	-	-	-	-	1	-	-	2
2002.....	66	1	1	-	-	-	-	-	-	1	1	1
2001.....	43	2	2	-	1	-	-	-	-	1	1	2
2000.....	37	3	1	-	-	-	-	1	-	1	1	2
1999.....	25	2	-	2	-	-	-	-	-	-	1	1

Black, female

2003.....	120	1	-	-	1	-	1	-	-	-	2	1
2002.....	90	4	-	-	-	-	-	1	1	-	1	2
2001.....	58	1	1	-	-	-	-	1	1	-	-	1
2000.....	47	1	1	-	-	-	-	-	-	2	2	1
1999.....	49	2	-	-	-	-	1	-	-	-	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
All races, both sexes												
2003.....	711	-	1	-	2	3	7	8	10	11	24	54
2002.....	784	1	-	-	-	-	9	7	11	18	34	35
2001.....	764	2	1	-	-	6	6	11	14	18	22	37
2000.....	776	1	6	-	1	2	9	7	13	17	25	38
1999.....	930	2	2	1	2	1	10	3	16	25	32	51
All races, male												
2003.....	427	-	1	-	2	1	2	6	7	8	17	39
2002.....	462	-	-	-	-	-	5	4	7	11	27	20
2001.....	453	-	-	-	-	5	3	7	7	12	15	27
2000.....	468	-	1	-	-	2	5	3	11	11	17	35
1999.....	584	-	2	-	2	1	3	1	10	20	23	42
All races, female												
2003.....	284	-	-	-	-	2	5	2	3	3	7	15
2002.....	322	1	-	-	-	-	4	3	4	7	7	15
2001.....	311	2	1	-	-	1	3	4	7	6	7	10
2000.....	308	1	5	-	1	-	4	4	2	6	8	3
1999.....	346	2	-	1	-	-	7	2	6	5	9	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Tuberculosis (A16-A19)

White, both sexes

2003.....	437	-	1	-	2	2	4	4	4	4	7	22
2002.....	461	-	-	-	-	-	3	3	3	8	14	13
2001.....	469	1	1	-	-	4	4	7	3	7	15	20
2000.....	439	-	4	-	1	1	5	5	6	8	5	13
1999.....	564	1	-	-	1	1	4	2	3	12	13	25

White, male

2003.....	255	-	1	-	2	1	2	3	2	4	5	18
2002.....	272	-	-	-	-	-	2	2	1	5	12	7
2001.....	271	-	-	-	-	4	2	6	2	5	11	14
2000.....	253	-	-	-	-	1	3	2	5	8	3	12
1999.....	348	-	-	-	1	1	3	1	3	9	10	21

White, female

2003.....	182	-	-	-	-	1	2	1	2	-	2	4
2002.....	189	-	-	-	-	-	1	1	2	3	2	6
2001.....	198	1	1	-	-	-	2	1	1	2	4	6
2000.....	186	-	4	-	1	-	2	3	1	-	2	1
1999.....	216	1	-	-	-	-	1	1	-	3	3	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Tuberculosis (A16-A19)

Black, both sexes

2003.....	172	-	-	-	-	1	3	1	5	5	11	24
2002.....	204	-	-	-	-	-	4	4	4	9	15	17
2001.....	203	-	-	-	-	-	1	3	7	9	7	14
2000.....	244	-	2	-	-	-	2	2	5	8	17	21
1999.....	245	1	2	1	-	-	4	1	8	10	15	20

Black, male

2003.....	114	-	-	-	-	-	-	1	4	3	8	17
2002.....	127	-	-	-	-	-	2	2	3	5	12	11
2001.....	127	-	-	-	-	-	-	1	3	6	4	11
2000.....	161	-	1	-	-	-	1	1	4	2	12	19
1999.....	159	-	2	-	-	-	-	-	4	8	11	17

Black, female

2003.....	58	-	-	-	-	1	3	-	1	2	3	7
2002.....	77	-	-	-	-	-	2	2	1	4	3	6
2001.....	76	-	-	-	-	-	1	2	4	3	3	3
2000.....	83	-	1	-	-	-	1	1	1	6	5	2
1999.....	86	1	-	1	-	-	4	1	4	2	4	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Tuberculosis (A16-A19)

American Indian 1/, both sexes

2003.....	15	-	-	-	-	-	-	-	-	1	1	4
2002.....	27	-	-	-	-	-	-	-	1	-	2	2
2001.....	13	-	-	-	-	-	-	1	-	-	-	2
2000.....	20	1	-	-	-	-	-	-	-	-	-	2
1999.....	20	-	-	-	-	-	-	-	1	1	-	3

American Indian 1/, male

2003.....	6	-	-	-	-	-	-	-	-	-	1	2
2002.....	14	-	-	-	-	-	-	-	1	-	-	1
2001.....	5	-	-	-	-	-	-	-	-	-	-	1
2000.....	12	-	-	-	-	-	-	-	-	-	-	2
1999.....	11	-	-	-	-	-	-	-	-	1	-	3

American Indian 1/, female

2003.....	9	-	-	-	-	-	-	-	-	1	-	2
2002.....	13	-	-	-	-	-	-	-	-	-	2	1
2001.....	8	-	-	-	-	-	-	1	-	-	-	1
2000.....	8	1	-	-	-	-	-	-	-	-	-	-
1999.....	9	-	-	-	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Tuberculosis (A16-A19)

Asian or Pacific Islander 2/, both sexes

2003.....	87	-	-	-	-	-	-	3	1	1	5	4
2002.....	92	1	-	-	-	-	2	-	3	1	3	3
2001.....	79	1	-	-	-	2	1	-	4	2	-	1
2000.....	73	-	-	-	-	1	2	-	2	1	3	2
1999.....	101	-	-	-	1	-	2	-	4	2	4	3

Asian or Pacific Islander 2/, male

2003.....	52	-	-	-	-	-	-	2	1	1	3	2
2002.....	49	-	-	-	-	-	1	-	2	1	3	1
2001.....	50	-	-	-	-	1	1	-	2	1	-	1
2000.....	42	-	-	-	-	1	1	-	2	1	2	2
1999.....	66	-	-	-	1	-	-	-	3	2	2	1

Asian or Pacific Islander 2/, female

2003.....	35	-	-	-	-	-	-	1	-	-	2	2
2002.....	43	1	-	-	-	-	1	-	1	-	-	2
2001.....	29	1	-	-	-	1	-	-	2	1	-	-
2000.....	31	-	-	-	-	-	1	-	-	-	1	-
1999.....	35	-	-	-	-	-	2	-	1	-	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
All races, both sexes												
2003.....	562	-	1	-	2	2	7	2	4	5	21	44
2002.....	626	-	-	-	-	-	6	6	8	13	27	28
2001.....	600	2	1	-	-	3	4	8	9	14	14	27
2000.....	615	-	3	-	-	2	5	5	6	15	20	30
1999.....	764	1	1	-	1	-	8	2	12	19	26	38
All races, male												
2003.....	344	-	1	-	2	1	2	2	4	3	14	32
2002.....	372	-	-	-	-	-	4	4	5	9	22	16
2001.....	371	-	-	-	-	3	2	5	5	9	9	20
2000.....	393	-	1	-	-	2	2	3	5	11	15	29
1999.....	486	-	1	-	1	-	2	-	8	15	18	32
All races, female												
2003.....	218	-	-	-	-	1	5	-	-	2	7	12
2002.....	254	-	-	-	-	-	2	2	3	4	5	12
2001.....	229	2	1	-	-	-	2	3	4	5	5	7
2000.....	222	-	2	-	-	-	3	2	1	4	5	1
1999.....	278	1	-	-	-	-	6	2	4	4	8	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
White, both sexes												
2003.....	347	-	1	-	2	2	4	2	2	3	6	18
2002.....	374	-	-	-	-	-	2	2	3	8	12	12
2001.....	368	1	1	-	-	2	2	5	2	4	10	14
2000.....	351	-	1	-	-	1	3	3	2	8	3	9
1999.....	472	-	-	-	-	-	3	1	2	8	12	20
White, male												
2003.....	208	-	1	-	2	1	2	2	2	3	4	16
2002.....	223	-	-	-	-	-	2	2	1	5	11	6
2001.....	221	-	-	-	-	2	1	4	1	3	7	9
2000.....	216	-	-	-	-	1	2	2	2	8	3	9
1999.....	296	-	-	-	-	-	2	-	2	6	9	17
White, female												
2003.....	139	-	-	-	-	1	2	-	-	-	2	2
2002.....	151	-	-	-	-	-	-	-	2	3	1	6
2001.....	147	1	1	-	-	-	1	1	1	1	3	5
2000.....	135	-	1	-	-	-	1	1	-	-	-	-
1999.....	176	-	-	-	-	-	1	1	-	2	3	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Black, both sexes												
2003.....	138	-	-	-	-	-	3	-	2	1	10	20
2002.....	164	-	-	-	-	-	2	4	4	4	11	13
2001.....	160	-	-	-	-	-	1	3	4	8	4	12
2000.....	192	-	2	-	-	-	1	2	4	6	14	17
1999.....	199	1	1	-	-	-	4	1	7	9	11	15
Black, male												
2003.....	93	-	-	-	-	-	-	-	2	-	7	13
2002.....	104	-	-	-	-	-	1	2	3	3	8	8
2001.....	106	-	-	-	-	-	-	1	2	5	2	10
2000.....	134	-	1	-	-	-	-	1	3	2	10	16
1999.....	126	-	1	-	-	-	-	-	3	7	7	13
Black, female												
2003.....	45	-	-	-	-	-	3	-	-	1	3	7
2002.....	60	-	-	-	-	-	1	2	1	1	3	5
2001.....	54	-	-	-	-	-	1	2	2	3	2	2
2000.....	58	-	1	-	-	-	1	1	1	4	4	1
1999.....	73	1	-	-	-	-	4	1	4	2	4	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Asian or Pacific Islander 2/, both sexes												
2003.....	67	-	-	-	-	-	-	-	-	-	5	3
2002.....	70	-	-	-	-	-	2	-	1	1	3	2
2001.....	64	1	-	-	-	1	1	-	3	2	-	1
2000.....	60	-	-	-	-	1	1	-	-	1	3	2
1999.....	83	-	-	-	1	-	1	-	3	2	3	2
Asian or Pacific Islander 2/, male												
2003.....	39	-	-	-	-	-	-	-	-	-	3	1
2002.....	36	-	-	-	-	-	1	-	1	1	3	1
2001.....	41	-	-	-	-	1	1	-	2	1	-	1
2000.....	35	-	-	-	-	1	-	-	-	1	2	2
1999.....	59	-	-	-	1	-	-	-	3	2	2	1
Asian or Pacific Islander 2/, female												
2003.....	28	-	-	-	-	-	-	-	-	-	2	2
2002.....	34	-	-	-	-	-	1	-	-	-	-	1
2001.....	23	1	-	-	-	-	-	-	1	1	-	-
2000.....	25	-	-	-	-	-	1	-	-	-	1	-
1999.....	24	-	-	-	-	-	1	-	-	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
All races, both sexes												
2003.....	149	-	-	-	-	1	-	6	6	6	3	10
2002.....	158	1	-	-	-	-	3	1	3	5	7	7
2001.....	164	-	-	-	-	3	2	3	5	4	8	10
2000.....	161	1	3	-	1	-	4	2	7	2	5	8
1999.....	166	1	1	1	1	1	2	1	4	6	6	13
All races, male												
2003.....	83	-	-	-	-	-	-	4	3	5	3	7
2002.....	90	-	-	-	-	-	1	-	2	2	5	4
2001.....	82	-	-	-	-	2	1	2	2	3	6	7
2000.....	75	-	-	-	-	-	3	-	6	-	2	6
1999.....	98	-	1	-	1	1	1	1	2	5	5	10
All races, female												
2003.....	66	-	-	-	-	1	-	2	3	1	-	3
2002.....	68	1	-	-	-	-	2	1	1	3	2	3
2001.....	82	-	-	-	-	1	1	1	3	1	2	3
2000.....	86	1	3	-	1	-	1	2	1	2	3	2
1999.....	68	1	-	1	-	-	1	-	2	1	1	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
White, both sexes												
2003.....	90	-	-	-	-	-	-	2	2	1	1	4
2002.....	87	-	-	-	-	-	1	1	-	-	2	1
2001.....	101	-	-	-	-	2	2	2	1	3	5	6
2000.....	88	-	3	-	1	-	2	2	4	-	2	4
1999.....	92	1	-	-	1	1	1	1	1	4	1	5
White, male												
2003.....	47	-	-	-	-	-	-	1	-	1	1	2
2002.....	49	-	-	-	-	-	-	-	-	-	1	1
2001.....	50	-	-	-	-	2	1	2	1	2	4	5
2000.....	37	-	-	-	-	-	1	-	3	-	-	3
1999.....	52	-	-	-	1	1	1	1	1	3	1	4
White, female												
2003.....	43	-	-	-	-	-	-	1	2	-	-	2
2002.....	38	-	-	-	-	-	1	1	-	-	1	-
2001.....	51	-	-	-	-	-	1	-	-	1	1	1
2000.....	51	-	3	-	1	-	1	2	1	-	2	1
1999.....	40	1	-	-	-	-	-	-	-	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
American Indian 1/, both sexes												
2003.....	5	-	-	-	-	-	-	-	-	-	1	1
2002.....	9	-	-	-	-	-	-	-	1	-	1	1
2001.....	5	-	-	-	-	-	-	1	-	-	-	2
2000.....	8	1	-	-	-	-	-	-	-	-	-	-
1999.....	10	-	-	-	-	-	-	-	1	1	-	2
American Indian 1/, male												
2003.....	2	-	-	-	-	-	-	-	-	-	1	-
2002.....	5	-	-	-	-	-	-	-	1	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	1
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, female												
2003.....	3	-	-	-	-	-	-	-	-	-	-	1
2002.....	4	-	-	-	-	-	-	-	-	-	1	1
2001.....	3	-	-	-	-	-	-	1	-	-	-	1
2000.....	4	1	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other tuberculosis (A17-A19)

Asian or Pacific Islander 2/, both sexes

2003.....	20	-	-	-	-	-	-	3	1	1	-	1
2002.....	22	1	-	-	-	-	-	-	2	-	-	1
2001.....	15	-	-	-	-	1	-	-	1	-	-	-
2000.....	13	-	-	-	-	-	1	-	2	-	-	-
1999.....	18	-	-	-	-	-	1	-	1	-	1	1

Asian or Pacific Islander 2/, male

2003.....	13	-	-	-	-	-	-	2	1	1	-	1
2002.....	13	-	-	-	-	-	-	-	1	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	-
2000.....	7	-	-	-	-	-	1	-	2	-	-	-
1999.....	7	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	7	-	-	-	-	-	-	1	-	-	-	-
2002.....	9	1	-	-	-	-	-	-	1	-	-	1
2001.....	6	-	-	-	-	1	-	-	1	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	-
1999.....	11	-	-	-	-	-	1	-	1	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
All races, both sexes												
2003.....	161	12	18	2	9	18	21	8	4	10	9	8
2002.....	161	17	12	6	14	21	12	12	4	4	5	10
2001.....	199	16	19	12	14	43	12	6	4	8	5	9
2000.....	211	22	19	12	6	34	22	9	6	7	6	13
1999.....	227	32	30	13	12	28	11	11	12	12	15	11
All races, male												
2003.....	75	9	10	1	2	8	10	3	4	4	5	5
2002.....	83	8	9	5	8	11	6	8	2	2	3	5
2001.....	94	6	9	6	9	20	7	4	2	3	3	6
2000.....	106	14	12	5	2	24	11	3	3	6	3	5
1999.....	101	14	19	7	6	13	6	8	4	3	6	1
All races, female												
2003.....	86	3	8	1	7	10	11	5	-	6	4	3
2002.....	78	9	3	1	6	10	6	4	2	2	2	5
2001.....	105	10	10	6	5	23	5	2	2	5	2	3
2000.....	105	8	7	7	4	10	11	6	3	1	3	8
1999.....	126	18	11	6	6	15	5	3	8	9	9	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Meningococcal infection (A39)

White, both sexes

2003.....	121	12	11	1	6	14	16	6	2	9	5	6
2002.....	125	14	8	5	5	17	10	9	4	1	4	9
2001.....	158	14	14	8	12	31	9	5	2	6	5	8
2000.....	161	19	14	5	5	29	16	7	3	3	4	7
1999.....	184	30	26	10	8	20	9	6	8	8	11	10

White, male

2003.....	56	9	6	-	2	6	8	3	2	4	3	4
2002.....	67	8	6	5	4	8	6	7	2	1	2	5
2001.....	72	5	7	4	8	15	5	3	1	1	3	5
2000.....	75	11	8	2	2	20	8	2	-	2	3	2
1999.....	79	14	17	5	4	8	5	4	2	2	4	1

White, female

2003.....	65	3	5	1	4	8	8	3	-	5	2	2
2002.....	58	6	2	-	1	9	4	2	2	-	2	4
2001.....	86	9	7	4	4	16	4	2	1	5	2	3
2000.....	86	8	6	3	3	9	8	5	3	1	1	5
1999.....	105	16	9	5	4	12	4	2	6	6	7	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Black, both sexes												
2003.....	36	-	7	1	3	3	4	2	1	1	4	2
2002.....	29	3	3	-	8	3	2	2	-	3	1	1
2001.....	36	2	4	4	2	10	3	-	2	1	-	1
2000.....	44	3	5	7	1	3	4	1	3	4	2	5
1999.....	39	2	4	2	4	7	2	4	4	4	4	1
Black, male												
2003.....	17	-	4	1	-	1	2	-	1	-	2	1
2002.....	12	-	2	-	4	2	-	-	-	1	1	-
2001.....	18	1	1	2	1	4	2	-	1	1	-	1
2000.....	27	3	4	3	-	3	2	-	3	4	-	2
1999.....	20	-	2	1	2	5	1	3	2	1	2	-
Black, female												
2003.....	19	-	3	-	3	2	2	2	-	1	2	1
2002.....	17	3	1	-	4	1	2	2	-	2	-	1
2001.....	18	1	3	2	1	6	1	-	1	-	-	-
2000.....	17	-	1	4	1	-	2	1	-	-	2	3
1999.....	19	2	2	1	2	2	1	1	2	3	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Meningococcal infection (A39)

Asian or Pacific Islander 2/, both sexes

2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	5	-	-	1	-	1	-	1	-	-	-	-
2001.....	4	-	-	-	-	2	-	1	-	1	-	-
2000.....	5	-	-	-	-	1	2	1	-	-	-	1
1999.....	2	-	-	-	-	1	-	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	3	-	-	-	-	1	-	1	-	-	-	-
2001.....	3	-	-	-	-	1	-	1	-	1	-	-
2000.....	4	-	-	-	-	1	1	1	-	-	-	1
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	1	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	1	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
All races, both sexes												
2003.....	34,069	278	85	39	38	59	95	123	186	315	595	916
2002.....	33,865	296	79	42	53	49	69	119	195	318	538	904
2001.....	32,238	312	108	29	40	57	76	96	185	309	520	837
2000.....	31,224	274	99	38	25	40	60	104	176	324	553	781
1999.....	30,680	283	87	47	30	32	89	106	169	314	498	731
All races, male												
2003.....	14,987	151	54	15	19	35	47	56	105	167	288	498
2002.....	14,947	158	41	26	26	22	37	63	103	159	282	495
2001.....	14,307	174	60	17	22	31	44	50	89	170	281	439
2000.....	13,537	139	53	24	14	23	37	62	96	162	289	439
1999.....	13,395	175	48	22	14	17	50	53	94	170	290	406
All races, female												
2003.....	19,082	127	31	24	19	24	48	67	81	148	307	418
2002.....	18,918	138	38	16	27	27	32	56	92	159	256	409
2001.....	17,931	138	48	12	18	26	32	46	96	139	239	398
2000.....	17,687	135	46	14	11	17	23	42	80	162	264	342
1999.....	17,285	108	39	25	16	15	39	53	75	144	208	325

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
White, both sexes												
2003.....	27,180	142	53	26	28	37	66	74	116	202	396	613
2002.....	27,110	191	58	31	38	38	48	75	127	210	364	628
2001.....	25,806	171	80	25	29	37	45	61	108	191	340	547
2000.....	24,804	169	66	28	21	24	36	61	104	204	374	495
1999.....	24,287	164	60	35	20	26	53	66	116	194	316	457
White, male												
2003.....	11,909	74	35	11	15	25	31	32	63	112	191	343
2002.....	11,919	100	34	21	21	16	29	45	64	104	190	354
2001.....	11,371	100	46	14	18	22	24	33	52	115	194	286
2000.....	10,716	88	35	19	11	15	24	34	61	110	201	289
1999.....	10,489	106	32	18	10	14	35	37	65	109	185	256
White, female												
2003.....	15,271	68	18	15	13	12	35	42	53	90	205	270
2002.....	15,191	91	24	10	17	22	19	30	63	106	174	274
2001.....	14,435	71	34	11	11	15	21	28	56	76	146	261
2000.....	14,088	81	31	9	10	9	12	27	43	94	173	206
1999.....	13,798	58	28	17	10	12	18	29	51	85	131	201

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Black, both sexes												
2003.....	6,206	123	24	10	5	18	23	42	60	101	184	278
2002.....	6,137	91	17	7	13	9	15	41	62	97	161	243
2001.....	5,880	126	22	2	8	19	22	28	70	109	160	261
2000.....	5,928	101	26	8	3	12	21	40	68	109	163	259
1999.....	5,826	107	17	7	7	5	28	37	48	110	172	251
Black, male												
2003.....	2,775	71	14	4	2	8	11	20	36	48	87	145
2002.....	2,703	51	5	3	4	5	5	15	33	50	83	122
2001.....	2,635	67	12	1	2	9	16	13	33	51	77	133
2000.....	2,587	50	13	4	2	6	12	27	32	48	77	136
1999.....	2,623	61	11	4	3	3	11	14	27	56	99	136
Black, female												
2003.....	3,431	52	10	6	3	10	12	22	24	53	97	133
2002.....	3,434	40	12	4	9	4	10	26	29	47	78	121
2001.....	3,245	59	10	1	6	10	6	15	37	58	83	128
2000.....	3,341	51	13	4	1	6	9	13	36	61	86	123
1999.....	3,203	46	6	3	4	2	17	23	21	54	73	115

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
American Indian 1/, both sexes												
2003.....	227	4	4	2	2	2	2	3	5	9	13	9
2002.....	195	4	3	1	1	-	2	1	3	7	6	16
2001.....	155	4	3	1	1	-	3	2	2	6	9	18
2000.....	162	1	3	1	-	1	1	1	2	6	7	12
1999.....	169	3	1	2	1	-	4	1	2	7	5	10
American Indian 1/, male												
2003.....	89	2	2	-	1	2	2	1	4	5	9	2
2002.....	95	2	2	1	1	-	-	1	3	3	3	8
2001.....	83	2	1	1	1	-	1	1	1	2	5	12
2000.....	74	1	2	1	-	1	-	-	2	2	5	7
1999.....	81	3	-	-	-	-	2	1	1	3	4	6
American Indian 1/, female												
2003.....	138	2	2	2	1	-	-	2	1	4	4	7
2002.....	100	2	1	-	-	-	2	-	-	4	3	8
2001.....	72	2	2	-	-	-	2	1	1	4	4	6
2000.....	88	-	1	-	-	-	1	1	-	4	2	5
1999.....	88	-	1	2	1	-	2	-	1	4	1	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Asian or Pacific Islander 2/, both sexes												
2003.....	456	9	4	1	3	2	4	4	5	3	2	16
2002.....	423	10	1	3	1	2	4	2	3	4	7	17
2001.....	397	11	3	1	2	1	6	5	5	3	11	11
2000.....	330	3	4	1	1	3	2	2	2	5	9	15
1999.....	398	9	9	3	2	1	4	2	3	3	5	13
Asian or Pacific Islander 2/, male												
2003.....	214	4	3	-	1	-	3	3	2	2	1	8
2002.....	230	5	-	1	-	1	3	2	3	2	6	11
2001.....	218	5	1	1	1	-	3	3	3	2	5	8
2000.....	160	-	3	-	1	1	1	1	1	2	6	7
1999.....	202	5	5	-	1	-	2	1	1	2	2	8
Asian or Pacific Islander 2/, female												
2003.....	242	5	1	1	2	2	1	1	3	1	1	8
2002.....	193	5	1	2	1	1	1	-	-	2	1	6
2001.....	179	6	2	-	1	1	3	2	2	1	6	3
2000.....	170	3	1	1	-	2	1	1	1	3	3	8
1999.....	196	4	4	3	1	1	2	1	2	1	3	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

All races, both sexes

2003.....	5,431	1	-	-	-	5	9	21	43	174	478	1,053
2002.....	5,793	2	3	-	2	12	11	37	57	216	524	1,192
2001.....	5,585	1	-	3	-	9	11	19	55	202	595	1,101
2000.....	5,357	6	1	2	-	4	14	25	54	219	560	1,044
1999.....	4,853	2	3	1	2	3	11	19	69	204	518	871

All races, male

2003.....	3,546	-	-	-	-	2	5	12	30	126	335	763
2002.....	3,770	1	3	-	1	4	9	20	43	154	365	887
2001.....	3,539	1	-	2	-	6	5	7	35	133	420	833
2000.....	3,381	2	1	1	-	2	9	13	33	151	389	787
1999.....	3,043	2	2	-	2	3	7	9	44	139	381	651

All races, female

2003.....	1,885	1	-	-	-	3	4	9	13	48	143	290
2002.....	2,023	1	-	-	1	8	2	17	14	62	159	305
2001.....	2,046	-	-	1	-	3	6	12	20	69	175	268
2000.....	1,976	4	-	1	-	2	5	12	21	68	171	257
1999.....	1,810	-	1	1	-	-	4	10	25	65	137	220

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

White, both sexes

2003.....	4,295	-	-	-	-	5	4	17	32	146	385	878
2002.....	4,620	1	-	-	2	9	11	21	44	178	422	961
2001.....	4,451	1	-	3	-	6	7	12	41	166	480	898
2000.....	4,216	3	-	2	-	2	6	17	34	180	445	827
1999.....	3,835	1	3	1	1	3	6	11	57	169	402	683

White, male

2003.....	2,848	-	-	-	-	2	1	10	20	104	275	648
2002.....	3,049	1	-	-	1	2	9	12	34	133	296	727
2001.....	2,900	1	-	2	-	4	2	5	26	111	345	691
2000.....	2,686	1	-	1	-	2	4	9	21	123	316	636
1999.....	2,436	1	2	-	1	3	4	4	37	118	301	501

White, female

2003.....	1,447	-	-	-	-	3	3	7	12	42	110	230
2002.....	1,571	-	-	-	1	7	2	9	10	45	126	234
2001.....	1,551	-	-	1	-	2	5	7	15	55	135	207
2000.....	1,530	2	-	1	-	-	2	8	13	57	129	191
1999.....	1,399	-	1	1	-	-	2	7	20	51	101	182

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

Black, both sexes

2003.....	854	-	-	-	-	-	4	2	9	20	78	138
2002.....	865	1	1	-	-	2	-	15	7	31	81	185
2001.....	831	-	-	-	-	2	4	7	8	27	94	167
2000.....	793	3	-	-	-	2	8	7	14	30	88	172
1999.....	731	1	-	-	1	-	4	6	11	25	96	161

Black, male

2003.....	544	-	-	-	-	-	3	1	8	15	49	93
2002.....	541	-	1	-	-	1	-	7	4	16	52	125
2001.....	484	-	-	-	-	1	3	2	5	15	64	115
2000.....	501	1	-	-	-	-	5	3	8	22	56	121
1999.....	443	1	-	-	1	-	2	3	6	15	66	126

Black, female

2003.....	310	-	-	-	-	-	1	1	1	5	29	45
2002.....	324	1	-	-	-	1	-	8	3	15	29	60
2001.....	347	-	-	-	-	1	1	5	3	12	30	52
2000.....	292	2	-	-	-	2	3	4	6	8	32	51
1999.....	288	-	-	-	-	-	2	3	5	10	30	35

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

American Indian 1/, both sexes

2003.....	62	-	-	-	-	-	1	1	2	2	7	17
2002.....	56	-	-	-	-	-	-	-	2	3	9	17
2001.....	55	-	-	-	-	1	-	-	4	4	9	10
2000.....	63	-	-	-	-	-	-	-	4	3	12	12
1999.....	37	-	-	-	-	-	1	-	1	2	6	8

American Indian 1/, male

2003.....	32	-	-	-	-	-	1	-	2	1	5	4
2002.....	36	-	-	-	-	-	-	-	2	1	7	11
2001.....	30	-	-	-	-	1	-	-	2	3	3	6
2000.....	33	-	-	-	-	-	-	-	2	2	6	7
1999.....	22	-	-	-	-	-	1	-	1	-	3	8

American Indian 1/, female

2003.....	30	-	-	-	-	-	-	1	-	1	2	13
2002.....	20	-	-	-	-	-	-	-	-	2	2	6
2001.....	25	-	-	-	-	-	-	-	2	1	6	4
2000.....	30	-	-	-	-	-	-	-	2	1	6	5
1999.....	15	-	-	-	-	-	-	-	-	2	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

Asian or Pacific Islander 2/, both sexes

2003.....	220	1	-	-	-	-	-	1	-	6	8	20
2002.....	252	-	2	-	-	1	-	1	4	4	12	29
2001.....	248	-	-	-	-	-	-	-	2	5	12	26
2000.....	285	-	1	-	-	-	-	1	2	6	15	33
1999.....	250	-	-	-	-	-	-	2	-	8	14	19

Asian or Pacific Islander 2/, male

2003.....	122	-	-	-	-	-	-	1	-	6	6	18
2002.....	144	-	2	-	-	1	-	1	3	4	10	24
2001.....	125	-	-	-	-	-	-	-	2	4	8	21
2000.....	161	-	1	-	-	-	-	1	2	4	11	23
1999.....	142	-	-	-	-	-	-	2	-	6	11	16

Asian or Pacific Islander 2/, female

2003.....	98	1	-	-	-	-	-	-	-	-	2	2
2002.....	108	-	-	-	-	-	-	-	1	-	2	5
2001.....	123	-	-	-	-	-	-	-	-	1	4	5
2000.....	124	-	-	-	-	-	-	-	-	2	4	10
1999.....	108	-	-	-	-	-	-	-	-	2	3	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races, both sexes

2003.....	13,658	5	2	15	21	32	146	444	1,144	2,401	2,939	2,636
2002.....	14,095	7	5	4	21	38	140	500	1,339	2,618	3,089	2,680
2001.....	14,175	10	13	17	21	42	183	584	1,517	2,741	3,126	2,511
2000.....	14,478	10	14	17	29	35	144	647	1,790	2,896	3,023	2,551
1999.....	14,802	14	24	38	28	31	167	763	1,966	2,981	3,251	2,440

All races, male

2003.....	10,119	2	2	7	12	15	80	272	766	1,710	2,174	2,013
2002.....	10,468	4	1	4	10	17	76	288	899	1,943	2,255	2,039
2001.....	10,590	7	8	11	12	21	84	354	1,067	2,042	2,318	1,942
2000.....	10,868	8	6	9	13	19	80	371	1,231	2,172	2,269	2,008
1999.....	11,234	9	16	20	16	15	80	504	1,410	2,224	2,482	1,936

All races, female

2003.....	3,539	3	-	8	9	17	66	172	378	691	765	623
2002.....	3,627	3	4	-	11	21	64	212	440	675	834	641
2001.....	3,585	3	5	6	9	21	99	230	450	699	808	569
2000.....	3,610	2	8	8	16	16	64	276	559	724	754	543
1999.....	3,568	5	8	18	12	16	87	259	556	757	769	504

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2003.....	6,019	1	1	2	3	12	45	139	480	1,049	1,383	1,199
2002.....	6,089	4	-	-	6	12	40	184	504	1,192	1,378	1,193
2001.....	6,171	3	3	7	4	6	54	191	623	1,275	1,406	1,097
2000.....	6,498	1	1	8	13	13	40	235	748	1,402	1,383	1,150
1999.....	6,752	4	7	12	13	10	53	304	898	1,472	1,479	1,065

White, male

2003.....	4,964	1	1	-	2	6	29	106	373	846	1,140	1,014
2002.....	5,028	3	-	-	4	5	26	135	397	990	1,133	997
2001.....	5,135	2	-	5	2	2	33	136	492	1,059	1,179	924
2000.....	5,360	1	-	4	5	8	30	158	591	1,165	1,135	977
1999.....	5,613	3	5	6	7	7	33	226	730	1,222	1,227	897

White, female

2003.....	1,055	-	-	2	1	6	16	33	107	203	243	185
2002.....	1,061	1	-	-	2	7	14	49	107	202	245	196
2001.....	1,036	1	3	2	2	4	21	55	131	216	227	173
2000.....	1,138	-	1	4	8	5	10	77	157	237	248	173
1999.....	1,139	1	2	6	6	3	20	78	168	250	252	168

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2003.....	7,479	3	1	13	18	20	99	301	647	1,309	1,523	1,407
2002.....	7,835	3	5	4	15	25	99	309	814	1,397	1,666	1,466
2001.....	7,844	7	10	9	17	34	129	384	874	1,432	1,687	1,393
2000.....	7,848	9	12	9	14	22	103	409	1,023	1,470	1,610	1,385
1999.....	7,893	10	17	26	15	21	112	452	1,045	1,473	1,733	1,350

Black, male

2003.....	5,033	1	1	7	10	9	50	164	379	830	1,010	975
2002.....	5,301	1	1	4	6	11	49	148	486	932	1,085	1,023
2001.....	5,328	5	8	5	10	18	51	212	555	955	1,116	1,001
2000.....	5,400	7	6	5	7	11	49	210	625	988	1,108	1,016
1999.....	5,493	6	11	14	9	8	46	273	662	969	1,225	1,019

Black, female

2003.....	2,446	2	-	6	8	11	49	137	268	479	513	432
2002.....	2,534	2	4	-	9	14	50	161	328	465	581	443
2001.....	2,516	2	2	4	7	16	78	172	319	477	571	392
2000.....	2,448	2	6	4	7	11	54	199	398	482	502	369
1999.....	2,400	4	6	12	6	13	66	179	383	504	508	331

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

American Indian 1/, both sexes

2003.....	70	-	-	-	-	-	-	-	8	20	17	12
2002.....	64	-	-	-	-	-	1	3	10	16	20	8
2001.....	74	-	-	1	-	-	-	2	9	21	14	14
2000.....	57	-	1	-	1	-	-	1	7	12	12	12
1999.....	68	-	-	-	-	-	-	4	13	19	15	8

American Indian 1/, male

2003.....	49	-	-	-	-	-	-	-	7	14	9	8
2002.....	50	-	-	-	-	-	1	2	8	12	17	7
2001.....	57	-	-	1	-	-	-	1	9	15	9	11
2000.....	43	-	-	-	-	-	-	1	4	9	11	11
1999.....	54	-	-	-	-	-	-	3	10	17	10	6

American Indian 1/, female

2003.....	21	-	-	-	-	-	-	-	1	6	8	4
2002.....	14	-	-	-	-	-	-	1	2	4	3	1
2001.....	17	-	-	-	-	-	-	1	-	6	5	3
2000.....	14	-	1	-	1	-	-	-	3	3	1	1
1999.....	14	-	-	-	-	-	-	1	3	2	5	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Asian or Pacific Islander 2/, both sexes

2003.....	90	1	-	-	-	-	2	4	9	23	16	18
2002.....	107	-	-	-	-	1	-	4	11	13	25	13
2001.....	86	-	-	-	-	2	-	7	11	13	19	7
2000.....	75	-	-	-	1	-	1	2	12	12	18	4
1999.....	89	-	-	-	-	-	2	3	10	17	24	17

Asian or Pacific Islander 2/, male

2003.....	73	-	-	-	-	-	1	2	7	20	15	16
2002.....	89	-	-	-	-	1	-	3	8	9	20	12
2001.....	70	-	-	-	-	1	-	5	11	13	14	6
2000.....	65	-	-	-	1	-	1	2	11	10	15	4
1999.....	74	-	-	-	-	-	1	2	8	16	20	14

Asian or Pacific Islander 2/, female

2003.....	17	1	-	-	-	-	1	2	2	3	1	2
2002.....	18	-	-	-	-	-	-	1	3	4	5	1
2001.....	16	-	-	-	-	1	-	2	-	-	5	1
2000.....	10	-	-	-	-	-	-	-	1	2	3	-
1999.....	15	-	-	-	-	-	1	1	2	1	4	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malaria (B50-B54)

All races, both sexes

2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	12	-	-	-	-	-	1	-	1	-	1	1
2001.....	9	-	-	-	1	-	-	-	-	2	-	2
2000.....	3	-	-	-	-	-	1	-	-	-	-	-
1999.....	7	1	-	-	-	-	-	2	-	-	1	-

All races, male

2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	8	-	-	-	-	-	-	-	-	-	1	1
2001.....	5	-	-	-	1	-	-	-	-	-	-	1
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	1	-	-	-	-	-	1	-	-	-	-

All races, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	-	-	1	-	1	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	2	-	1
2000.....	1	-	-	-	-	-	1	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	1	-	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malaria (B50-B54)

Black, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	-	-	1	-	1	-	-	1
2001.....	3	-	-	-	1	-	-	-	-	-	-	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	1	-	-	-	-	-	1	-	-	-	-

Black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	1
2001.....	2	-	-	-	1	-	-	-	-	-	-	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	1	-	-	-	-	-	-	-	-	-	-

Black, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	1	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races, both sexes

2003.....	7,427	174	85	29	22	49	39	54	86	158	289	526
2002.....	6,707	224	61	41	43	42	55	67	102	163	290	403
2001.....	5,715	191	64	34	30	40	46	58	89	162	207	307
2000.....	5,532	167	71	30	37	44	52	56	85	162	188	277
1999.....	7,279	203	81	40	39	43	47	92	150	234	311	392

All races, male

2003.....	3,964	86	43	14	11	30	22	30	53	96	188	351
2002.....	3,622	125	39	16	24	20	32	38	58	97	194	278
2001.....	2,931	101	37	14	15	23	28	31	52	93	134	211
2000.....	2,861	94	41	16	19	25	29	29	46	105	115	164
1999.....	3,655	119	51	20	24	23	26	46	85	149	184	231

All races, female

2003.....	3,463	88	42	15	11	19	17	24	33	62	101	175
2002.....	3,085	99	22	25	19	22	23	29	44	66	96	125
2001.....	2,784	90	27	20	15	17	18	27	37	69	73	96
2000.....	2,671	73	30	14	18	19	23	27	39	57	73	113
1999.....	3,624	84	30	20	15	20	21	46	65	85	127	161

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes

2003.....	6,182	110	53	22	17	36	27	37	60	110	224	418
2002.....	5,528	155	45	32	34	30	34	47	70	103	204	314
2001.....	4,688	120	44	28	21	27	31	39	57	111	147	237
2000.....	4,550	101	50	24	28	32	39	34	50	104	138	212
1999.....	5,794	135	56	32	24	32	32	56	88	156	197	262

White, male

2003.....	3,289	51	26	12	7	24	16	22	35	71	149	287
2002.....	2,967	85	28	14	21	16	17	24	39	64	143	220
2001.....	2,405	62	25	11	11	14	19	24	37	65	95	166
2000.....	2,334	59	27	12	13	16	25	19	29	68	85	127
1999.....	2,910	85	37	18	13	16	18	27	51	110	129	163

White, female

2003.....	2,893	59	27	10	10	12	11	15	25	39	75	131
2002.....	2,561	70	17	18	13	14	17	23	31	39	61	94
2001.....	2,283	58	19	17	10	13	12	15	20	46	52	71
2000.....	2,216	42	23	12	15	16	14	15	21	36	53	85
1999.....	2,884	50	19	14	11	16	14	29	37	46	68	99

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2003.....	960	53	25	5	4	11	7	13	20	40	61	94
2002.....	943	59	14	5	5	9	18	18	29	53	72	81
2001.....	825	62	16	6	7	9	14	17	28	42	52	62
2000.....	826	61	14	5	5	9	13	16	34	48	47	62
1999.....	1,260	60	23	7	10	7	13	31	58	74	98	117

Black, male

2003.....	523	30	13	2	3	4	5	5	15	21	36	57
2002.....	526	33	10	1	2	3	12	13	16	28	43	53
2001.....	427	36	10	3	3	7	8	7	12	23	36	41
2000.....	432	31	10	3	2	6	4	6	16	30	28	35
1999.....	624	31	14	2	8	5	7	14	34	37	46	62

Black, female

2003.....	437	23	12	3	1	7	2	8	5	19	25	37
2002.....	417	26	4	4	3	6	6	5	13	25	29	28
2001.....	398	26	6	3	4	2	6	10	16	19	16	21
2000.....	394	30	4	2	3	3	9	10	18	18	19	27
1999.....	636	29	9	5	2	2	6	17	24	37	52	55

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

American Indian 1/, both sexes

2003.....	72	4	4	1	1	-	2	1	1	4	1	7
2002.....	54	4	1	-	1	-	-	-	-	1	8	3
2001.....	51	5	3	-	2	1	1	-	-	4	4	3
2000.....	34	2	2	-	-	-	-	1	-	5	-	1
1999.....	53	4	1	-	1	1	-	-	-	1	6	3

American Indian 1/, male

2003.....	33	1	2	-	1	-	-	1	1	2	-	3
2002.....	28	3	1	-	1	-	-	-	-	-	5	3
2001.....	21	2	2	-	1	1	1	-	-	2	1	1
2000.....	13	1	-	-	-	-	-	-	-	2	-	1
1999.....	20	1	-	-	-	-	-	-	-	1	2	-

American Indian 1/, female

2003.....	39	3	2	1	-	-	2	-	-	2	1	4
2002.....	26	1	-	-	-	-	-	-	-	1	3	-
2001.....	30	3	1	-	1	-	-	-	-	2	3	2
2000.....	21	1	2	-	-	-	-	1	-	3	-	-
1999.....	33	3	1	-	1	1	-	-	-	-	4	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Asian or Pacific Islander 2/, both sexes

2003.....	213	7	3	1	-	2	3	3	5	4	3	7
2002.....	182	6	1	4	3	3	3	2	3	6	6	5
2001.....	151	4	1	-	-	3	-	2	4	5	4	5
2000.....	122	3	5	1	4	3	-	5	1	5	3	2
1999.....	172	4	1	1	4	3	2	5	4	3	10	10

Asian or Pacific Islander 2/, male

2003.....	119	4	2	-	-	2	1	2	2	2	3	4
2002.....	101	4	-	1	-	1	3	1	3	5	3	2
2001.....	78	1	-	-	-	1	-	-	3	3	2	3
2000.....	82	3	4	1	4	3	-	4	1	5	2	1
1999.....	101	2	-	-	3	2	1	5	-	1	7	6

Asian or Pacific Islander 2/, female

2003.....	94	3	1	1	-	-	2	1	3	2	-	3
2002.....	81	2	1	3	3	2	-	1	-	1	3	3
2001.....	73	3	1	-	-	2	-	2	1	2	2	2
2000.....	40	-	1	-	-	-	-	1	-	-	1	1
1999.....	71	2	1	1	1	1	1	-	4	2	3	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

All races, both sexes

2003.....	556,902	75	392	516	560	690	961	1,307	2,434	4,919	10,590	19,552
2002.....	557,271	74	402	537	535	723	1,007	1,258	2,614	5,150	10,935	19,295
2001.....	553,768	66	420	493	515	732	972	1,389	2,605	5,466	11,103	19,199
2000.....	553,091	92	420	489	525	745	968	1,359	2,557	5,565	10,955	18,803
1999.....	549,838	68	418	509	503	745	979	1,429	2,576	5,764	10,968	18,412

All races, male

2003.....	287,990	34	227	293	298	438	541	664	1,140	2,070	4,755	9,727
2002.....	288,768	42	216	297	303	424	592	678	1,190	2,166	4,890	9,418
2001.....	287,075	31	229	253	280	447	577	694	1,166	2,331	4,966	9,377
2000.....	286,082	50	233	269	300	446	576	692	1,162	2,402	4,942	9,243
1999.....	285,832	36	227	279	276	448	575	755	1,204	2,467	5,004	9,151

All races, female

2003.....	268,912	41	165	223	262	252	420	643	1,294	2,849	5,835	9,825
2002.....	268,503	32	186	240	232	299	415	580	1,424	2,984	6,045	9,877
2001.....	266,693	35	191	240	235	285	395	695	1,439	3,135	6,137	9,822
2000.....	267,009	42	187	220	225	299	392	667	1,395	3,163	6,013	9,560
1999.....	264,006	32	191	230	227	297	404	674	1,372	3,297	5,964	9,261

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

White, both sexes

2003.....	481,556	64	317	427	426	563	761	1,015	1,853	3,818	8,248	15,226
2002.....	482,481	56	306	428	418	574	825	975	2,022	4,009	8,573	15,180
2001.....	479,651	54	353	387	393	587	772	1,074	2,020	4,278	8,681	15,127
2000.....	480,011	75	330	397	421	576	757	1,081	1,991	4,316	8,448	14,743
1999.....	477,238	53	353	397	383	568	759	1,137	2,010	4,495	8,565	14,513

White, male

2003.....	249,053	29	182	248	224	358	432	543	889	1,679	3,743	7,612
2002.....	249,867	35	168	229	230	344	486	531	941	1,710	3,894	7,492
2001.....	248,146	24	189	195	203	360	464	549	944	1,861	3,922	7,421
2000.....	247,403	42	190	215	241	344	454	559	941	1,907	3,832	7,292
1999.....	247,396	31	194	224	210	335	449	606	955	1,984	3,950	7,171

White, female

2003.....	232,503	35	135	179	202	205	329	472	964	2,139	4,505	7,614
2002.....	232,614	21	138	199	188	230	339	444	1,081	2,299	4,679	7,688
2001.....	231,505	30	164	192	190	227	308	525	1,076	2,417	4,759	7,706
2000.....	232,608	33	140	182	180	232	303	522	1,050	2,409	4,616	7,451
1999.....	229,842	22	159	173	173	233	310	531	1,055	2,511	4,615	7,342

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

Black, both sexes

2003.....	62,660	7	57	73	100	94	153	229	438	880	1,937	3,646
2002.....	62,617	12	74	85	87	115	135	212	457	918	1,985	3,449
2001.....	62,170	6	48	80	96	113	149	242	455	959	2,022	3,395
2000.....	61,945	11	70	72	78	136	165	215	434	1,017	2,091	3,437
1999.....	61,951	11	49	85	98	139	179	233	438	1,058	2,022	3,308

Black, male

2003.....	32,442	4	31	40	56	57	86	90	179	291	836	1,786
2002.....	32,627	4	34	55	51	66	74	103	187	349	823	1,625
2001.....	32,679	3	30	43	59	68	85	107	163	370	847	1,642
2000.....	32,817	4	34	42	45	82	97	94	168	383	926	1,664
1999.....	32,850	3	26	41	55	87	97	114	183	385	882	1,702

Black, female

2003.....	30,218	3	26	33	44	37	67	139	259	589	1,101	1,860
2002.....	29,990	8	40	30	36	49	61	109	270	569	1,162	1,824
2001.....	29,491	3	18	37	37	45	64	135	292	589	1,175	1,753
2000.....	29,128	7	36	30	33	54	68	121	266	634	1,165	1,773
1999.....	29,101	8	23	44	43	52	82	119	255	673	1,140	1,606

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

American Indian 1/, both sexes

2003.....	2,154	-	7	4	4	9	10	10	14	36	65	114
2002.....	2,175	1	3	6	6	9	11	11	24	39	61	109
2001.....	2,155	3	4	-	5	10	12	16	18	45	69	105
2000.....	1,914	2	3	5	3	6	7	9	17	34	71	88
1999.....	1,836	1	4	2	4	4	9	13	13	28	57	86

American Indian 1/, male

2003.....	1,092	-	7	1	1	6	7	4	7	15	29	61
2002.....	1,081	-	2	3	4	4	6	6	8	14	30	44
2001.....	1,103	3	3	-	3	7	8	12	7	16	37	47
2000.....	997	2	2	3	1	4	3	5	8	16	33	44
1999.....	949	-	1	1	4	1	6	7	7	13	22	51

American Indian 1/, female

2003.....	1,062	-	-	3	3	3	3	6	7	21	36	53
2002.....	1,094	1	1	3	2	5	5	5	16	25	31	65
2001.....	1,052	-	1	-	2	3	4	4	11	29	32	58
2000.....	917	-	1	2	2	2	4	4	9	18	38	44
1999.....	887	1	3	1	-	3	3	6	6	15	35	35

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

Asian or Pacific Islander 2/, both sexes

2003.....	10,532	4	11	12	30	24	37	53	129	185	340	566
2002.....	9,998	5	19	18	24	25	36	60	111	184	316	557
2001.....	9,792	3	15	26	21	22	39	57	112	184	331	572
2000.....	9,221	4	17	15	23	27	39	54	115	198	345	535
1999.....	8,813	3	12	25	18	34	32	46	115	183	324	505

Asian or Pacific Islander 2/, male

2003.....	5,403	1	7	4	17	17	16	27	65	85	147	268
2002.....	5,193	3	12	10	18	10	26	38	54	93	143	257
2001.....	5,147	1	7	15	15	12	20	26	52	84	160	267
2000.....	4,865	2	7	9	13	16	22	34	45	96	151	243
1999.....	4,637	2	6	13	7	25	23	28	59	85	150	227

Asian or Pacific Islander 2/, female

2003.....	5,129	3	4	8	13	7	21	26	64	100	193	298
2002.....	4,805	2	7	8	6	15	10	22	57	91	173	300
2001.....	4,645	2	8	11	6	10	19	31	60	100	171	305
2000.....	4,356	2	10	6	10	11	17	20	70	102	194	292
1999.....	4,176	1	6	12	11	9	9	18	56	98	174	278

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All races, both sexes												
2003.....	7,778	-	4	4	4	11	19	19	40	64	210	445
2002.....	7,737	-	3	1	7	12	11	22	35	77	181	431
2001.....	7,702	-	-	2	2	10	19	19	37	93	190	427
2000.....	7,492	-	-	-	4	9	17	18	32	88	205	408
1999.....	7,486	-	2	1	5	9	18	14	29	96	195	449
All races, male												
2003.....	5,298	-	2	1	3	8	13	11	28	40	155	346
2002.....	5,184	-	2	1	2	7	4	14	25	47	132	336
2001.....	5,156	-	-	1	-	3	9	12	23	61	141	325
2000.....	4,952	-	-	-	4	6	7	12	16	60	155	309
1999.....	4,944	-	1	-	3	6	14	10	17	58	146	346
All races, female												
2003.....	2,480	-	2	3	1	3	6	8	12	24	55	99
2002.....	2,553	-	1	-	5	5	7	8	10	30	49	95
2001.....	2,546	-	-	1	2	7	10	7	14	32	49	102
2000.....	2,540	-	-	-	-	3	10	6	16	28	50	99
1999.....	2,542	-	1	1	2	3	4	4	12	38	49	103

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
White, both sexes												
2003.....	6,409	-	3	4	1	5	14	14	27	47	149	322
2002.....	6,432	-	2	1	4	8	7	14	25	52	128	304
2001.....	6,369	-	-	1	1	7	14	14	23	55	123	296
2000.....	6,135	-	-	-	2	6	12	12	23	60	138	291
1999.....	6,116	-	2	1	3	5	9	10	20	60	134	287
White, male												
2003.....	4,282	-	1	1	1	4	9	8	17	32	121	243
2002.....	4,256	-	2	1	1	5	3	9	19	35	92	240
2001.....	4,183	-	-	1	-	1	6	10	13	37	93	227
2000.....	3,942	-	-	-	2	3	4	8	12	38	100	222
1999.....	3,955	-	1	-	2	3	6	6	11	34	100	218
White, female												
2003.....	2,127	-	2	3	-	1	5	6	10	15	28	79
2002.....	2,176	-	-	-	3	3	4	5	6	17	36	64
2001.....	2,186	-	-	-	1	6	8	4	10	18	30	69
2000.....	2,193	-	-	-	-	3	8	4	11	22	38	69
1999.....	2,161	-	1	1	1	2	3	4	9	26	34	69

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Black, both sexes												
2003.....	1,107	-	1	-	3	5	4	4	7	14	49	104
2002.....	1,044	-	-	-	2	4	2	3	8	22	43	106
2001.....	1,065	-	-	1	-	1	3	2	4	24	48	106
2000.....	1,102	-	-	-	2	3	4	5	2	15	46	96
1999.....	1,145	-	-	-	2	4	4	2	6	25	42	135
Black, male												
2003.....	837	-	1	-	2	3	3	3	7	6	26	86
2002.....	766	-	-	-	1	2	1	2	5	10	33	80
2001.....	782	-	-	-	-	1	1	1	4	14	32	80
2000.....	833	-	-	-	2	3	2	3	1	11	36	69
1999.....	828	-	-	-	1	3	3	2	4	17	30	109
Black, female												
2003.....	270	-	-	-	1	2	1	1	-	8	23	18
2002.....	278	-	-	-	1	2	1	1	3	12	10	26
2001.....	283	-	-	1	-	-	2	1	-	10	16	26
2000.....	269	-	-	-	-	-	2	2	1	4	10	27
1999.....	317	-	-	-	1	1	1	-	2	8	12	26

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Asian or Pacific Islander 2/, both sexes

2003.....	229	-	-	-	-	1	1	1	6	3	12	18
2002.....	218	-	1	-	1	-	2	4	2	3	8	20
2001.....	228	-	-	-	1	2	2	3	8	12	16	24
2000.....	214	-	-	-	-	-	1	1	7	11	19	21
1999.....	192	-	-	-	-	-	5	1	3	11	18	24

Asian or Pacific Islander 2/, male

2003.....	162	-	-	-	-	1	1	-	4	2	8	16
2002.....	135	-	-	-	-	-	-	2	1	2	6	15
2001.....	158	-	-	-	-	1	2	1	5	8	14	17
2000.....	150	-	-	-	-	-	1	1	3	9	17	18
1999.....	139	-	-	-	-	-	5	1	2	7	15	16

Asian or Pacific Islander 2/, female

2003.....	67	-	-	-	-	-	-	1	2	1	4	2
2002.....	83	-	1	-	1	-	2	2	1	1	2	5
2001.....	70	-	-	-	1	1	-	2	3	4	2	7
2000.....	64	-	-	-	-	-	-	-	4	2	2	3
1999.....	53	-	-	-	-	-	-	-	1	4	3	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of esophagus (C15)												
All races, both sexes												
2003.....	12,860	-	-	-	-	2	3	11	18	76	212	526
2002.....	12,701	-	-	-	1	-	6	4	15	84	227	503
2001.....	12,530	-	-	-	-	2	4	9	20	68	241	492
2000.....	12,232	1	-	-	-	1	2	9	30	78	219	478
1999.....	11,918	-	-	-	-	-	-	10	27	75	218	469
All races, male												
2003.....	9,921	-	-	-	-	2	2	8	15	66	174	445
2002.....	9,789	-	-	-	-	-	3	4	10	71	186	426
2001.....	9,607	-	-	-	-	1	4	8	18	52	203	417
2000.....	9,279	1	-	-	-	-	2	7	24	69	196	415
1999.....	9,094	-	-	-	-	-	-	7	22	65	190	410
All races, female												
2003.....	2,939	-	-	-	-	-	1	3	3	10	38	81
2002.....	2,912	-	-	-	1	-	3	-	5	13	41	77
2001.....	2,923	-	-	-	-	1	-	1	2	16	38	75
2000.....	2,953	-	-	-	-	1	-	2	6	9	23	63
1999.....	2,824	-	-	-	-	-	-	3	5	10	28	59

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of esophagus (C15)												
White, both sexes												
2003.....	11,058	-	-	-	-	2	3	9	16	65	174	406
2002.....	10,910	-	-	-	1	-	6	3	14	73	178	403
2001.....	10,624	-	-	-	-	2	4	7	17	59	191	389
2000.....	10,406	-	-	-	-	1	2	9	29	74	175	377
1999.....	10,079	-	-	-	-	-	-	9	23	67	162	362
White, male												
2003.....	8,617	-	-	-	-	2	2	7	14	58	146	349
2002.....	8,510	-	-	-	-	-	3	3	10	65	151	353
2001.....	8,205	-	-	-	-	1	4	6	15	45	164	338
2000.....	7,986	-	-	-	-	-	2	7	24	67	162	341
1999.....	7,740	-	-	-	-	-	-	6	18	60	143	323
White, female												
2003.....	2,441	-	-	-	-	-	1	2	2	7	28	57
2002.....	2,400	-	-	-	1	-	3	-	4	8	27	50
2001.....	2,419	-	-	-	-	1	-	1	2	14	27	51
2000.....	2,420	-	-	-	-	1	-	2	5	7	13	36
1999.....	2,339	-	-	-	-	-	-	3	5	7	19	39

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of esophagus (C15)												
Black, both sexes												
2003.....	1,591	-	-	-	-	-	-	1	2	10	37	109
2002.....	1,590	-	-	-	-	-	-	1	1	9	47	93
2001.....	1,688	-	-	-	-	-	-	2	1	9	46	92
2000.....	1,655	1	-	-	-	-	-	-	1	3	41	96
1999.....	1,648	-	-	-	-	-	-	-	3	6	48	99
Black, male												
2003.....	1,133	-	-	-	-	-	-	1	1	8	28	87
2002.....	1,133	-	-	-	-	-	-	1	-	4	34	68
2001.....	1,231	-	-	-	-	-	-	2	1	7	35	69
2000.....	1,170	1	-	-	-	-	-	-	-	1	31	69
1999.....	1,203	-	-	-	-	-	-	-	3	4	41	79
Black, female												
2003.....	458	-	-	-	-	-	-	-	1	2	9	22
2002.....	457	-	-	-	-	-	-	-	1	5	13	25
2001.....	457	-	-	-	-	-	-	-	-	2	11	23
2000.....	485	-	-	-	-	-	-	-	1	2	10	27
1999.....	445	-	-	-	-	-	-	-	-	2	7	20

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of esophagus (C15)

Asian or Pacific Islander 2/, both sexes

2003.....	162	-	-	-	-	-	-	1	-	1	1	5
2002.....	154	-	-	-	-	-	-	-	-	2	-	6
2001.....	166	-	-	-	-	-	-	-	2	-	1	9
2000.....	128	-	-	-	-	-	-	-	-	1	1	2
1999.....	153	-	-	-	-	-	-	1	1	2	7	5

Asian or Pacific Islander 2/, male

2003.....	133	-	-	-	-	-	-	-	-	-	-	4
2002.....	109	-	-	-	-	-	-	-	-	2	-	4
2001.....	133	-	-	-	-	-	-	-	2	-	1	8
2000.....	87	-	-	-	-	-	-	-	-	1	1	2
1999.....	120	-	-	-	-	-	-	1	1	1	5	5

Asian or Pacific Islander 2/, female

2003.....	29	-	-	-	-	-	-	1	-	1	1	1
2002.....	45	-	-	-	-	-	-	-	-	-	-	2
2001.....	33	-	-	-	-	-	-	-	-	-	-	1
2000.....	41	-	-	-	-	-	-	-	-	-	-	-
1999.....	33	-	-	-	-	-	-	-	-	1	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of stomach (C16)												
All races, both sexes												
2003.....	12,110	-	-	-	1	5	14	34	93	178	302	493
2002.....	12,198	-	1	-	-	7	22	48	90	160	276	443
2001.....	12,319	-	-	-	-	5	15	33	89	178	286	452
2000.....	12,645	-	-	-	1	3	11	36	88	167	317	468
1999.....	12,711	-	-	-	1	5	8	47	89	189	316	450
All races, male												
2003.....	7,055	-	-	-	1	3	4	16	47	91	171	305
2002.....	7,133	-	1	-	-	-	8	31	49	82	157	285
2001.....	7,191	-	-	-	-	4	5	20	51	98	155	288
2000.....	7,328	-	-	-	-	-	6	22	47	100	184	290
1999.....	7,459	-	-	-	1	2	4	22	54	101	182	303
All races, female												
2003.....	5,055	-	-	-	-	2	10	18	46	87	131	188
2002.....	5,065	-	-	-	-	7	14	17	41	78	119	158
2001.....	5,128	-	-	-	-	1	10	13	38	80	131	164
2000.....	5,317	-	-	-	1	3	5	14	41	67	133	178
1999.....	5,252	-	-	-	-	3	4	25	35	88	134	147

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of stomach (C16)												
White, both sexes												
2003.....	9,282	-	-	-	1	5	10	24	64	123	200	330
2002.....	9,319	-	1	-	-	5	16	33	61	119	182	317
2001.....	9,501	-	-	-	-	2	8	25	59	112	198	321
2000.....	9,723	-	-	-	1	2	7	21	57	113	205	314
1999.....	9,779	-	-	-	-	2	6	29	59	128	193	320
White, male												
2003.....	5,428	-	-	-	1	3	4	12	34	71	112	206
2002.....	5,542	-	1	-	-	-	5	21	36	60	108	209
2001.....	5,568	-	-	-	-	2	2	14	34	67	109	207
2000.....	5,688	-	-	-	-	-	3	13	32	66	115	192
1999.....	5,786	-	-	-	-	1	3	15	40	70	112	222
White, female												
2003.....	3,854	-	-	-	-	2	6	12	30	52	88	124
2002.....	3,777	-	-	-	-	5	11	12	25	59	74	108
2001.....	3,933	-	-	-	-	-	6	11	25	45	89	114
2000.....	4,035	-	-	-	1	2	4	8	25	47	90	122
1999.....	3,993	-	-	-	-	1	3	14	19	58	81	98

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of stomach (C16)												
Black, both sexes												
2003.....	2,041	-	-	-	-	-	1	6	16	34	69	114
2002.....	2,131	-	-	-	-	2	6	13	20	27	65	92
2001.....	2,067	-	-	-	-	1	4	4	21	49	72	90
2000.....	2,176	-	-	-	-	1	3	11	25	37	81	110
1999.....	2,203	-	-	-	1	1	2	14	22	40	87	89
Black, male												
2003.....	1,187	-	-	-	-	-	-	3	7	10	45	76
2002.....	1,194	-	-	-	-	-	3	10	10	16	33	60
2001.....	1,186	-	-	-	-	1	1	3	11	25	38	57
2000.....	1,218	-	-	-	-	-	2	5	15	26	52	72
1999.....	1,263	-	-	-	1	-	1	7	12	20	52	60
Black, female												
2003.....	854	-	-	-	-	-	1	3	9	24	24	38
2002.....	937	-	-	-	-	2	3	3	10	11	32	32
2001.....	881	-	-	-	-	-	3	1	10	24	34	33
2000.....	958	-	-	-	-	1	1	6	10	11	29	38
1999.....	940	-	-	-	-	1	1	7	10	20	35	29

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of stomach (C16)

American Indian 1/, both sexes

2003.....	84	-	-	-	-	-	1	1	3	1	6	8
2002.....	91	-	-	-	-	-	-	-	1	1	5	6
2001.....	79	-	-	-	-	-	1	1	1	1	-	4
2000.....	73	-	-	-	-	-	-	-	-	1	3	-
1999.....	67	-	-	-	-	1	-	-	1	2	3	3

American Indian 1/, male

2003.....	50	-	-	-	-	-	-	1	1	-	2	4
2002.....	45	-	-	-	-	-	-	-	1	1	3	3
2001.....	51	-	-	-	-	-	1	1	1	1	-	2
2000.....	41	-	-	-	-	-	-	-	-	-	3	-
1999.....	44	-	-	-	-	-	-	-	1	1	2	2

American Indian 1/, female

2003.....	34	-	-	-	-	-	1	-	2	1	4	4
2002.....	46	-	-	-	-	-	-	-	-	-	2	3
2001.....	28	-	-	-	-	-	-	-	-	-	-	2
2000.....	32	-	-	-	-	-	-	-	-	1	-	-
1999.....	23	-	-	-	-	1	-	-	-	1	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of stomach (C16)												
Asian or Pacific Islander 2/, both sexes												
2003.....	703	-	-	-	-	-	2	3	10	20	27	41
2002.....	657	-	-	-	-	-	-	2	8	13	24	28
2001.....	672	-	-	-	-	2	2	3	8	16	16	37
2000.....	673	-	-	-	-	-	1	4	6	16	28	44
1999.....	662	-	-	-	-	1	-	4	7	19	33	38
Asian or Pacific Islander 2/, male												
2003.....	390	-	-	-	-	-	-	-	5	10	12	19
2002.....	352	-	-	-	-	-	-	-	2	5	13	13
2001.....	386	-	-	-	-	1	1	2	5	5	8	22
2000.....	381	-	-	-	-	-	1	4	-	8	14	26
1999.....	366	-	-	-	-	1	-	-	1	10	16	19
Asian or Pacific Islander 2/, female												
2003.....	313	-	-	-	-	-	2	3	5	10	15	22
2002.....	305	-	-	-	-	-	-	2	6	8	11	15
2001.....	286	-	-	-	-	1	1	1	3	11	8	15
2000.....	292	-	-	-	-	-	-	-	6	8	14	18
1999.....	296	-	-	-	-	-	-	4	6	9	17	19

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races, both sexes

2003.....	55,958	-	-	1	3	8	32	96	195	445	870	1,724
2002.....	56,741	1	-	-	1	9	41	88	202	439	986	1,755
2001.....	56,887	-	-	-	-	11	35	81	218	505	928	1,744
2000.....	57,477	-	-	-	3	13	27	79	193	480	889	1,656
1999.....	57,155	1	-	-	1	11	38	87	196	439	894	1,623

All races, male

2003.....	28,007	-	-	-	3	6	19	59	99	235	479	965
2002.....	28,501	1	-	-	1	6	28	48	113	233	564	958
2001.....	28,225	-	-	-	-	6	18	38	119	290	491	967
2000.....	28,462	-	-	-	2	8	16	42	111	271	488	893
1999.....	28,240	-	-	-	1	5	22	57	103	226	452	876

All races, female

2003.....	27,951	-	-	1	-	2	13	37	96	210	391	759
2002.....	28,240	-	-	-	-	3	13	40	89	206	422	797
2001.....	28,662	-	-	-	-	5	17	43	99	215	437	777
2000.....	29,015	-	-	-	1	5	11	37	82	209	401	763
1999.....	28,915	1	-	-	-	6	16	30	93	213	442	747

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of colon, rectum and anus (C18-C21)												
White, both sexes												
2003.....	47,711	-	-	1	3	7	22	72	145	329	636	1,310
2002.....	48,562	1	-	-	1	7	31	70	164	337	734	1,313
2001.....	48,748	-	-	-	-	7	27	60	163	384	707	1,324
2000.....	49,488	-	-	-	3	12	19	57	144	349	666	1,250
1999.....	49,374	-	-	-	1	6	26	62	150	336	684	1,264
White, male												
2003.....	24,013	-	-	-	3	5	15	45	73	177	362	746
2002.....	24,439	1	-	-	1	4	20	39	87	175	425	736
2001.....	24,335	-	-	-	-	5	11	28	91	226	373	746
2000.....	24,517	-	-	-	2	7	9	32	87	204	363	692
1999.....	24,539	-	-	-	1	2	14	40	78	180	360	692
White, female												
2003.....	23,698	-	-	1	-	2	7	27	72	152	274	564
2002.....	24,123	-	-	-	-	3	11	31	77	162	309	577
2001.....	24,413	-	-	-	-	2	16	32	72	158	334	578
2000.....	24,971	-	-	-	1	5	10	25	57	145	303	558
1999.....	24,835	-	-	-	-	4	12	22	72	156	324	572

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Black, both sexes												
2003.....	6,943	-	-	-	-	1	7	17	33	96	192	348
2002.....	6,874	-	-	-	-	2	8	11	29	75	209	370
2001.....	6,905	-	-	-	-	2	6	13	43	99	187	356
2000.....	6,867	-	-	-	-	-	6	12	42	109	184	359
1999.....	6,777	1	-	-	-	5	11	21	35	92	181	305
Black, male												
2003.....	3,335	-	-	-	-	1	3	9	16	44	97	187
2002.....	3,351	-	-	-	-	2	6	5	20	43	114	186
2001.....	3,251	-	-	-	-	1	6	6	24	53	102	187
2000.....	3,326	-	-	-	-	-	5	4	22	56	105	180
1999.....	3,196	-	-	-	-	3	7	14	17	43	82	159
Black, female												
2003.....	3,608	-	-	-	-	-	4	8	17	52	95	161
2002.....	3,523	-	-	-	-	-	2	6	9	32	95	184
2001.....	3,654	-	-	-	-	1	-	7	19	46	85	169
2000.....	3,541	-	-	-	-	-	1	8	20	53	79	179
1999.....	3,581	1	-	-	-	2	4	7	18	49	99	146

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

American Indian 1/, both sexes

2003.....	205	-	-	-	-	-	-	-	-	3	7	18
2002.....	246	-	-	-	-	-	-	2	1	7	4	11
2001.....	192	-	-	-	-	-	-	-	1	5	6	3
2000.....	194	-	-	-	-	-	-	2	1	2	11	7
1999.....	170	-	-	-	-	-	1	-	2	1	6	7

American Indian 1/, male

2003.....	111	-	-	-	-	-	-	-	-	2	3	11
2002.....	131	-	-	-	-	-	-	1	-	5	1	6
2001.....	95	-	-	-	-	-	-	-	-	3	2	2
2000.....	110	-	-	-	-	-	-	1	-	1	7	4
1999.....	87	-	-	-	-	-	1	-	2	-	2	6

American Indian 1/, female

2003.....	94	-	-	-	-	-	-	-	-	1	4	7
2002.....	115	-	-	-	-	-	-	1	1	2	3	5
2001.....	97	-	-	-	-	-	-	-	1	2	4	1
2000.....	84	-	-	-	-	-	-	1	1	1	4	3
1999.....	83	-	-	-	-	-	-	-	-	1	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Asian or Pacific Islander 2/, both sexes

2003.....	1,099	-	-	-	-	-	3	7	17	17	35	48
2002.....	1,059	-	-	-	-	-	2	5	8	20	39	61
2001.....	1,042	-	-	-	-	2	2	8	11	17	28	61
2000.....	928	-	-	-	-	1	2	8	6	20	28	40
1999.....	834	-	-	-	-	-	-	4	9	10	23	47

Asian or Pacific Islander 2/, male

2003.....	548	-	-	-	-	-	1	5	10	12	17	21
2002.....	580	-	-	-	-	-	2	3	6	10	24	30
2001.....	544	-	-	-	-	-	1	4	4	8	14	32
2000.....	509	-	-	-	-	1	2	5	2	10	13	17
1999.....	418	-	-	-	-	-	-	3	6	3	8	19

Asian or Pacific Islander 2/, female

2003.....	551	-	-	-	-	-	2	2	7	5	18	27
2002.....	479	-	-	-	-	-	-	2	2	10	15	31
2001.....	498	-	-	-	-	2	1	4	7	9	14	29
2000.....	419	-	-	-	-	-	-	3	4	10	15	23
1999.....	416	-	-	-	-	-	-	1	3	7	15	28

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All races, both sexes												
2003.....	14,706	6	23	6	8	10	26	25	64	113	282	815
2002.....	14,047	3	28	7	11	10	17	33	72	123	291	731
2001.....	13,351	5	18	7	8	14	15	29	46	137	264	795
2000.....	12,916	2	23	9	11	15	15	28	58	113	316	742
1999.....	12,382	8	29	5	6	19	24	33	45	127	317	666
All races, male												
2003.....	9,482	2	18	5	4	6	14	12	41	83	205	643
2002.....	9,011	2	19	4	8	9	9	20	42	85	223	596
2001.....	8,483	3	7	4	3	6	9	16	27	100	202	618
2000.....	8,176	1	14	6	2	11	9	18	41	74	223	595
1999.....	7,689	5	20	5	4	12	12	22	26	88	230	533
All races, female												
2003.....	5,224	4	5	1	4	4	12	13	23	30	77	172
2002.....	5,036	1	9	3	3	1	8	13	30	38	68	135
2001.....	4,868	2	11	3	5	8	6	13	19	37	62	177
2000.....	4,740	1	9	3	9	4	6	10	17	39	93	147
1999.....	4,693	3	9	-	2	7	12	11	19	39	87	133

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
White, both sexes												
2003.....	11,660	4	19	4	4	7	16	12	33	73	191	555
2002.....	11,250	2	22	7	6	6	13	18	39	74	201	521
2001.....	10,697	5	14	2	6	10	12	19	29	82	155	549
2000.....	10,304	1	17	8	5	10	10	12	32	65	196	510
1999.....	9,986	6	26	3	3	14	15	17	19	73	210	448
White, male												
2003.....	7,489	2	15	4	3	4	9	3	20	50	133	441
2002.....	7,114	2	14	4	3	5	5	8	19	48	153	426
2001.....	6,774	3	5	1	3	5	7	9	15	56	119	427
2000.....	6,447	1	11	5	2	7	6	8	19	37	142	401
1999.....	6,151	3	18	3	2	10	7	11	10	45	155	351
White, female												
2003.....	4,171	2	4	-	1	3	7	9	13	23	58	114
2002.....	4,136	-	8	3	3	1	8	10	20	26	48	95
2001.....	3,923	2	9	1	3	5	5	10	14	26	36	122
2000.....	3,857	-	6	3	3	3	4	4	13	28	54	109
1999.....	3,835	3	8	-	1	4	8	6	9	28	55	97

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, both sexes

2003.....	1,877	2	1	2	1	2	5	6	19	23	55	164
2002.....	1,734	1	4	-	3	3	1	9	18	23	61	147
2001.....	1,670	-	3	3	2	2	2	6	9	30	66	175
2000.....	1,598	1	2	-	2	4	2	10	11	25	77	166
1999.....	1,507	1	3	1	3	2	5	9	12	31	60	150

Black, male

2003.....	1,234	-	-	1	1	1	3	3	12	18	41	126
2002.....	1,165	-	3	-	3	3	1	6	11	16	45	116
2001.....	1,048	-	2	2	-	-	1	4	6	23	48	133
2000.....	1,061	-	1	-	-	3	2	5	10	16	50	138
1999.....	947	1	2	1	2	-	2	7	7	21	44	125

Black, female

2003.....	643	2	1	1	-	1	2	3	7	5	14	38
2002.....	569	1	1	-	-	-	-	3	7	7	16	31
2001.....	622	-	1	1	2	2	1	2	3	7	18	42
2000.....	537	1	1	-	2	1	-	5	1	9	27	28
1999.....	560	-	1	-	1	2	3	2	5	10	16	25

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

American Indian 1/, both sexes

2003.....	105	-	2	-	1	-	-	-	1	1	3	9
2002.....	94	-	1	-	-	-	-	-	1	-	2	6
2001.....	109	-	-	-	-	1	-	-	1	2	1	9
2000.....	98	-	-	1	-	-	-	-	2	1	3	7
1999.....	76	-	-	-	-	-	1	1	-	3	-	7

American Indian 1/, male

2003.....	71	-	2	-	-	-	-	-	1	1	3	7
2002.....	60	-	1	-	-	-	-	-	1	-	2	4
2001.....	71	-	-	-	-	-	-	-	-	1	-	7
2000.....	65	-	-	1	-	-	-	-	1	1	3	7
1999.....	42	-	-	-	-	-	1	-	-	2	-	5

American Indian 1/, female

2003.....	34	-	-	-	1	-	-	-	-	-	-	2
2002.....	34	-	-	-	-	-	-	-	-	-	-	2
2001.....	38	-	-	-	-	1	-	-	1	1	1	2
2000.....	33	-	-	-	-	-	-	-	1	-	-	-
1999.....	34	-	-	-	-	-	-	1	-	1	-	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Asian or Pacific Islander 2/, both sexes

2003.....	1,064	-	1	-	2	1	5	7	11	16	33	87
2002.....	969	-	1	-	2	1	3	6	14	26	27	57
2001.....	875	-	1	2	-	1	1	4	7	23	42	62
2000.....	916	-	4	-	4	1	3	6	13	22	40	59
1999.....	813	1	-	1	-	3	3	6	14	20	47	61

Asian or Pacific Islander 2/, male

2003.....	688	-	1	-	-	1	2	6	8	14	28	69
2002.....	672	-	1	-	2	1	3	6	11	21	23	50
2001.....	590	-	-	1	-	1	1	3	6	20	35	51
2000.....	603	-	2	-	-	1	1	5	11	20	28	49
1999.....	549	1	-	1	-	2	2	4	9	20	31	52

Asian or Pacific Islander 2/, female

2003.....	376	-	-	-	2	-	3	1	3	2	5	18
2002.....	297	-	-	-	-	-	-	-	3	5	4	7
2001.....	285	-	1	1	-	-	-	1	1	3	7	11
2000.....	313	-	2	-	4	-	2	1	2	2	12	10
1999.....	264	-	-	-	-	1	1	2	5	-	16	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of pancreas (C25)												
All races, both sexes												
2003.....	30,777	-	-	-	1	-	10	20	40	153	395	891
2002.....	30,264	1	-	-	-	2	5	7	32	156	440	889
2001.....	29,803	-	1	1	1	1	1	11	44	167	407	833
2000.....	29,332	2	1	-	1	1	4	17	38	140	408	834
1999.....	29,082	1	-	-	-	-	3	9	46	178	416	887
All races, male												
2003.....	15,060	-	-	-	1	-	6	11	24	97	232	556
2002.....	14,877	-	-	-	-	2	3	3	23	89	275	555
2001.....	14,467	-	1	-	-	1	1	5	25	99	222	503
2000.....	14,238	1	1	-	-	1	2	11	22	87	237	531
1999.....	14,176	-	-	-	-	-	1	7	26	107	262	575
All races, female												
2003.....	15,717	-	-	-	-	-	4	9	16	56	163	335
2002.....	15,387	1	-	-	-	-	2	4	9	67	165	334
2001.....	15,336	-	-	1	1	-	-	6	19	68	185	330
2000.....	15,094	1	-	-	1	-	2	6	16	53	171	303
1999.....	14,906	1	-	-	-	-	2	2	20	71	154	312

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of pancreas (C25)												
White, both sexes												
2003.....	26,418	-	-	-	-	-	7	18	30	121	300	699
2002.....	26,031	-	-	-	-	2	5	4	24	119	354	715
2001.....	25,769	-	1	1	1	1	1	7	35	134	327	652
2000.....	25,301	2	1	-	1	1	3	15	30	110	319	679
1999.....	25,086	-	-	-	-	-	2	7	33	137	331	729
White, male												
2003.....	13,033	-	-	-	-	-	3	10	16	78	173	434
2002.....	12,921	-	-	-	-	2	3	1	18	65	226	451
2001.....	12,641	-	1	-	-	1	1	4	21	77	182	403
2000.....	12,392	1	1	-	-	1	2	9	18	69	189	437
1999.....	12,273	-	-	-	-	-	1	6	19	85	212	476
White, female												
2003.....	13,385	-	-	-	-	-	4	8	14	43	127	265
2002.....	13,110	-	-	-	-	-	2	3	6	54	128	264
2001.....	13,128	-	-	1	1	-	-	3	14	57	145	249
2000.....	12,909	1	-	-	1	-	1	6	12	41	130	242
1999.....	12,813	-	-	-	-	-	1	1	14	52	119	253

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of pancreas (C25)												
Black, both sexes												
2003.....	3,616	-	-	-	1	-	3	1	9	23	80	168
2002.....	3,494	1	-	-	-	-	-	2	4	30	72	144
2001.....	3,394	-	-	-	-	-	-	4	6	26	64	157
2000.....	3,419	-	-	-	-	-	1	1	8	24	73	136
1999.....	3,391	1	-	-	-	-	-	2	10	36	76	141
Black, male												
2003.....	1,675	-	-	-	1	-	3	1	7	16	48	110
2002.....	1,617	-	-	-	-	-	-	1	4	19	39	85
2001.....	1,523	-	-	-	-	-	-	1	3	16	31	90
2000.....	1,561	-	-	-	-	-	-	1	4	15	39	80
1999.....	1,611	-	-	-	-	-	-	1	7	19	44	88
Black, female												
2003.....	1,941	-	-	-	-	-	-	-	2	7	32	58
2002.....	1,877	1	-	-	-	-	-	1	-	11	33	59
2001.....	1,871	-	-	-	-	-	-	3	3	10	33	67
2000.....	1,858	-	-	-	-	-	1	-	4	9	34	56
1999.....	1,780	1	-	-	-	-	-	1	3	17	32	53

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of pancreas (C25)												
American Indian 1/, both sexes												
2003.....	103	-	-	-	-	-	-	1	-	4	3	6
2002.....	112	-	-	-	-	-	-	-	1	1	3	6
2001.....	89	-	-	-	-	-	-	-	1	-	3	5
2000.....	91	-	-	-	-	-	-	-	-	2	6	1
1999.....	84	-	-	-	-	-	-	-	1	-	2	3
American Indian 1/, male												
2003.....	51	-	-	-	-	-	-	-	-	1	2	3
2002.....	50	-	-	-	-	-	-	-	-	1	3	5
2001.....	38	-	-	-	-	-	-	-	-	-	1	1
2000.....	46	-	-	-	-	-	-	-	-	-	5	1
1999.....	40	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, female												
2003.....	52	-	-	-	-	-	-	1	-	3	1	3
2002.....	62	-	-	-	-	-	-	-	1	-	-	1
2001.....	51	-	-	-	-	-	-	-	1	-	2	4
2000.....	45	-	-	-	-	-	-	-	-	2	1	-
1999.....	44	-	-	-	-	-	-	-	1	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of pancreas (C25)												
Asian or Pacific Islander 2/, both sexes												
2003.....	640	-	-	-	-	-	-	-	1	5	12	18
2002.....	627	-	-	-	-	-	-	1	3	6	11	24
2001.....	551	-	-	-	-	-	-	-	2	7	13	19
2000.....	521	-	-	-	-	-	-	1	-	4	10	18
1999.....	521	-	-	-	-	-	1	-	2	5	7	14
Asian or Pacific Islander 2/, male												
2003.....	301	-	-	-	-	-	-	-	1	2	9	9
2002.....	289	-	-	-	-	-	-	1	1	4	7	14
2001.....	265	-	-	-	-	-	-	-	1	6	8	9
2000.....	239	-	-	-	-	-	-	1	-	3	4	13
1999.....	252	-	-	-	-	-	-	-	-	3	5	9
Asian or Pacific Islander 2/, female												
2003.....	339	-	-	-	-	-	-	-	-	3	3	9
2002.....	338	-	-	-	-	-	-	-	2	2	4	10
2001.....	286	-	-	-	-	-	-	-	1	1	5	10
2000.....	282	-	-	-	-	-	-	-	-	1	6	5
1999.....	269	-	-	-	-	-	1	-	2	2	2	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of larynx (C32)												
All races, both sexes												
2003.....	3,792	-	-	-	-	-	-	-	1	10	69	157
2002.....	3,723	-	-	-	-	-	-	3	8	19	57	156
2001.....	3,797	-	-	-	-	-	-	1	2	18	68	149
2000.....	3,861	-	-	-	2	1	-	-	4	9	55	156
1999.....	3,815	-	-	-	-	-	-	-	6	20	58	151
All races, male												
2003.....	3,020	-	-	-	-	-	-	-	1	7	53	129
2002.....	2,922	-	-	-	-	-	-	3	3	16	42	130
2001.....	3,018	-	-	-	-	-	-	1	1	12	54	123
2000.....	3,041	-	-	-	-	-	-	-	2	6	43	122
1999.....	2,992	-	-	-	-	-	-	-	5	18	46	130
All races, female												
2003.....	772	-	-	-	-	-	-	-	-	3	16	28
2002.....	801	-	-	-	-	-	-	-	5	3	15	26
2001.....	779	-	-	-	-	-	-	-	1	6	14	26
2000.....	820	-	-	-	2	1	-	-	2	3	12	34
1999.....	823	-	-	-	-	-	-	-	1	2	12	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of larynx (C32)												
White, both sexes												
2003.....	3,052	-	-	-	-	-	-	-	1	9	53	110
2002.....	3,012	-	-	-	-	-	-	2	5	10	38	104
2001.....	3,049	-	-	-	-	-	-	1	2	11	50	98
2000.....	3,096	-	-	-	2	1	-	-	4	8	41	108
1999.....	3,106	-	-	-	-	-	-	-	4	14	31	101
White, male												
2003.....	2,410	-	-	-	-	-	-	-	1	6	43	94
2002.....	2,351	-	-	-	-	-	-	2	1	8	29	88
2001.....	2,417	-	-	-	-	-	-	1	1	10	40	79
2000.....	2,426	-	-	-	-	-	-	-	2	5	30	86
1999.....	2,419	-	-	-	-	-	-	-	3	14	24	87
White, female												
2003.....	642	-	-	-	-	-	-	-	-	3	10	16
2002.....	661	-	-	-	-	-	-	-	4	2	9	16
2001.....	632	-	-	-	-	-	-	-	1	1	10	19
2000.....	670	-	-	-	2	1	-	-	2	3	11	22
1999.....	687	-	-	-	-	-	-	-	1	-	7	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of larynx (C32)												
Black, both sexes												
2003.....	693	-	-	-	-	-	-	-	-	1	15	44
2002.....	675	-	-	-	-	-	-	1	3	8	19	48
2001.....	678	-	-	-	-	-	-	-	-	7	17	48
2000.....	719	-	-	-	-	-	-	-	-	1	14	47
1999.....	663	-	-	-	-	-	-	-	2	6	26	48
Black, male												
2003.....	571	-	-	-	-	-	-	-	-	1	9	33
2002.....	542	-	-	-	-	-	-	1	2	7	13	38
2001.....	541	-	-	-	-	-	-	-	-	2	13	41
2000.....	578	-	-	-	-	-	-	-	-	1	13	35
1999.....	536	-	-	-	-	-	-	-	2	4	21	41
Black, female												
2003.....	122	-	-	-	-	-	-	-	-	-	6	11
2002.....	133	-	-	-	-	-	-	-	1	1	6	10
2001.....	137	-	-	-	-	-	-	-	-	5	4	7
2000.....	141	-	-	-	-	-	-	-	-	-	1	12
1999.....	127	-	-	-	-	-	-	-	-	2	5	7

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All races, both sexes												
2003.....	158,086	-	2	-	2	7	18	29	125	520	1,958	4,617
2002.....	157,713	-	1	2	2	4	17	35	130	583	2,115	4,341
2001.....	156,058	-	5	1	1	6	14	36	121	640	2,156	4,255
2000.....	155,521	2	2	-	1	3	19	37	160	664	2,092	4,220
1999.....	152,156	-	-	3	2	6	14	43	136	668	2,056	3,916
All races, male												
2003.....	89,964	-	1	-	1	4	10	9	69	258	1,086	2,659
2002.....	90,171	-	1	1	2	2	14	19	71	290	1,096	2,471
2001.....	90,426	-	4	1	-	2	10	17	60	315	1,189	2,487
2000.....	90,469	2	1	-	-	3	11	28	74	370	1,172	2,540
1999.....	89,453	-	-	2	2	5	8	24	73	323	1,151	2,346
All races, female												
2003.....	68,122	-	1	-	1	3	8	20	56	262	872	1,958
2002.....	67,542	-	-	1	-	2	3	16	59	293	1,019	1,870
2001.....	65,632	-	1	-	1	4	4	19	61	325	967	1,768
2000.....	65,052	-	1	-	1	-	8	9	86	294	920	1,680
1999.....	62,703	-	-	1	-	1	6	19	63	345	905	1,570

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
White, both sexes												
2003.....	138,743	-	2	-	1	7	15	24	103	409	1,541	3,571
2002.....	138,661	-	1	2	1	3	14	25	102	465	1,655	3,426
2001.....	137,274	-	4	1	1	5	11	29	101	522	1,692	3,328
2000.....	136,912	1	2	-	1	1	13	27	128	499	1,594	3,265
1999.....	133,752	-	-	2	1	3	8	33	116	511	1,617	3,022
White, male												
2003.....	78,215	-	1	-	-	4	8	8	54	203	838	2,042
2002.....	78,465	-	1	1	1	2	11	14	58	222	841	1,913
2001.....	78,665	-	3	1	-	2	8	15	50	253	912	1,907
2000.....	78,747	1	1	-	-	1	9	20	62	277	859	1,953
1999.....	77,846	-	-	1	1	3	4	17	64	247	891	1,782
White, female												
2003.....	60,528	-	1	-	1	3	7	16	49	206	703	1,529
2002.....	60,196	-	-	1	-	1	3	11	44	243	814	1,513
2001.....	58,609	-	1	-	1	3	3	14	51	269	780	1,421
2000.....	58,165	-	1	-	1	-	4	7	66	222	735	1,312
1999.....	55,906	-	-	1	-	-	4	16	52	264	726	1,240

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Black, both sexes												
2003.....	16,372	-	-	-	1	-	3	2	17	81	361	932
2002.....	16,316	-	-	-	1	1	3	4	19	96	409	827
2001.....	16,019	-	1	-	-	1	2	3	13	103	416	804
2000.....	16,066	1	-	-	-	2	3	8	24	142	442	837
1999.....	15,963	-	-	1	1	3	5	5	15	134	399	804
Black, male												
2003.....	9,988	-	-	-	1	-	2	1	12	39	213	553
2002.....	10,075	-	-	-	1	-	3	2	12	57	232	505
2001.....	10,131	-	1	-	-	-	1	1	6	54	247	515
2000.....	10,153	1	-	-	-	2	-	6	9	80	282	520
1999.....	10,135	-	-	1	1	2	3	4	7	63	240	518
Black, female												
2003.....	6,384	-	-	-	-	-	1	1	5	42	148	379
2002.....	6,241	-	-	-	-	1	-	2	7	39	177	322
2001.....	5,888	-	-	-	-	1	1	2	7	49	169	289
2000.....	5,913	-	-	-	-	-	3	2	15	62	160	317
1999.....	5,828	-	-	-	-	1	2	1	8	71	159	286

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

American Indian 1/, both sexes

2003.....	556	-	-	-	-	-	-	-	-	3	8	21
2002.....	556	-	-	-	-	-	-	-	1	-	5	14
2001.....	547	-	-	-	-	-	-	-	1	1	7	24
2000.....	480	-	-	-	-	-	-	-	1	3	8	17
1999.....	497	-	-	-	-	-	-	-	-	3	5	13

American Indian 1/, male

2003.....	301	-	-	-	-	-	-	-	-	2	5	11
2002.....	304	-	-	-	-	-	-	-	-	-	2	7
2001.....	299	-	-	-	-	-	-	-	-	1	7	13
2000.....	270	-	-	-	-	-	-	-	1	1	4	9
1999.....	285	-	-	-	-	-	-	-	-	2	1	8

American Indian 1/, female

2003.....	255	-	-	-	-	-	-	-	-	1	3	10
2002.....	252	-	-	-	-	-	-	-	1	-	3	7
2001.....	248	-	-	-	-	-	-	-	1	-	-	11
2000.....	210	-	-	-	-	-	-	-	-	2	4	8
1999.....	212	-	-	-	-	-	-	-	-	1	4	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Asian or Pacific Islander 2/, both sexes

2003.....	2,415	-	-	-	-	-	-	3	5	27	48	93
2002.....	2,180	-	-	-	-	-	-	6	8	22	46	74
2001.....	2,218	-	-	-	-	-	1	4	6	14	41	99
2000.....	2,063	-	-	-	-	-	3	2	7	20	48	101
1999.....	1,944	-	-	-	-	-	1	5	5	20	35	77

Asian or Pacific Islander 2/, male

2003.....	1,460	-	-	-	-	-	-	-	3	14	30	53
2002.....	1,327	-	-	-	-	-	-	3	1	11	21	46
2001.....	1,331	-	-	-	-	-	1	1	4	7	23	52
2000.....	1,299	-	-	-	-	-	2	2	2	12	27	58
1999.....	1,187	-	-	-	-	-	1	3	2	11	19	38

Asian or Pacific Islander 2/, female

2003.....	955	-	-	-	-	-	-	3	2	13	18	40
2002.....	853	-	-	-	-	-	-	3	7	11	25	28
2001.....	887	-	-	-	-	-	-	3	2	7	18	47
2000.....	764	-	-	-	-	-	1	-	5	8	21	43
1999.....	757	-	-	-	-	-	-	2	3	9	16	39

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant melanoma of skin (C43)												
All races, both sexes												
2003.....	7,818	-	1	1	1	9	29	63	128	189	372	541
2002.....	7,514	-	3	-	1	7	34	70	124	219	385	495
2001.....	7,542	-	-	-	-	11	30	72	114	213	409	536
2000.....	7,420	-	-	-	1	7	34	58	128	234	392	534
1999.....	7,215	1	-	1	-	9	30	62	132	239	386	518
All races, male												
2003.....	4,963	-	-	-	-	6	18	40	79	114	222	328
2002.....	4,725	-	2	-	1	3	17	38	81	120	243	343
2001.....	4,763	-	-	-	-	9	15	40	63	130	250	331
2000.....	4,592	-	-	-	1	4	17	33	77	118	237	321
1999.....	4,529	1	-	-	-	7	16	38	72	151	246	337
All races, female												
2003.....	2,855	-	1	1	1	3	11	23	49	75	150	213
2002.....	2,789	-	1	-	-	4	17	32	43	99	142	152
2001.....	2,779	-	-	-	-	2	15	32	51	83	159	205
2000.....	2,828	-	-	-	-	3	17	25	51	116	155	213
1999.....	2,686	-	-	1	-	2	14	24	60	88	140	181

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant melanoma of skin (C43)												
White, both sexes												
2003.....	7,648	-	1	1	1	9	27	61	125	186	363	528
2002.....	7,353	-	3	-	1	6	34	68	124	210	380	485
2001.....	7,403	-	-	-	-	10	30	70	114	211	404	529
2000.....	7,245	-	-	-	1	6	34	55	125	229	377	525
1999.....	7,086	1	-	1	-	8	28	61	127	236	382	509
White, male												
2003.....	4,887	-	-	-	-	6	18	40	77	113	219	322
2002.....	4,639	-	2	-	1	2	17	36	81	118	239	337
2001.....	4,702	-	-	-	-	8	15	38	63	129	247	328
2000.....	4,514	-	-	-	1	4	17	32	76	116	229	319
1999.....	4,459	1	-	-	-	6	14	37	69	149	245	331
White, female												
2003.....	2,761	-	1	1	1	3	9	21	48	73	144	206
2002.....	2,714	-	1	-	-	4	17	32	43	92	141	148
2001.....	2,701	-	-	-	-	2	15	32	51	82	157	201
2000.....	2,731	-	-	-	-	2	17	23	49	113	148	206
1999.....	2,627	-	-	1	-	2	14	24	58	87	137	178

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant melanoma of skin (C43)												
Black, both sexes												
2003.....	121	-	-	-	-	-	1	1	2	2	9	8
2002.....	102	-	-	-	-	1	-	1	-	7	2	3
2001.....	104	-	-	-	-	1	-	1	-	2	5	3
2000.....	135	-	-	-	-	1	-	2	1	4	11	6
1999.....	96	-	-	-	-	1	-	1	3	1	3	6
Black, male												
2003.....	50	-	-	-	-	-	-	-	2	1	3	4
2002.....	52	-	-	-	-	1	-	1	-	2	2	2
2001.....	37	-	-	-	-	1	-	1	-	1	3	2
2000.....	60	-	-	-	-	-	-	1	1	2	7	1
1999.....	48	-	-	-	-	1	-	1	1	1	-	4
Black, female												
2003.....	71	-	-	-	-	-	1	1	-	1	6	4
2002.....	50	-	-	-	-	-	-	-	-	5	-	1
2001.....	67	-	-	-	-	-	-	-	-	1	2	1
2000.....	75	-	-	-	-	1	-	1	-	2	4	5
1999.....	48	-	-	-	-	-	-	-	2	-	3	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant melanoma of skin (C43)

Asian or Pacific Islander 2/, both sexes

2003.....	35	-	-	-	-	-	-	1	1	1	-	5
2002.....	42	-	-	-	-	-	-	1	-	2	3	6
2001.....	25	-	-	-	-	-	-	-	-	-	-	4
2000.....	32	-	-	-	-	-	-	1	2	1	4	2
1999.....	23	-	-	-	-	-	1	-	2	2	1	1

Asian or Pacific Islander 2/, male

2003.....	21	-	-	-	-	-	-	-	-	-	-	2
2002.....	25	-	-	-	-	-	-	1	-	-	2	3
2001.....	15	-	-	-	-	-	-	-	-	-	-	1
2000.....	13	-	-	-	-	-	-	-	-	-	1	1
1999.....	15	-	-	-	-	-	1	-	2	1	1	-

Asian or Pacific Islander 2/, female

2003.....	14	-	-	-	-	-	-	1	1	1	-	3
2002.....	17	-	-	-	-	-	-	-	-	2	1	3
2001.....	10	-	-	-	-	-	-	-	-	-	-	3
2000.....	19	-	-	-	-	-	-	1	2	1	3	1
1999.....	8	-	-	-	-	-	-	-	-	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
All races, both sexes												
2003.....	42,000	-	-	-	-	1	15	55	352	912	1,804	2,734
2002.....	41,883	-	-	-	-	1	9	63	362	904	1,818	2,747
2001.....	41,809	-	-	-	-	1	8	88	375	960	1,863	2,790
2000.....	42,300	-	-	-	-	1	10	83	384	982	1,859	2,730
1999.....	41,528	-	1	-	1	3	10	86	353	986	1,755	2,619
All races, male												
2003.....	380	-	-	-	-	-	-	1	1	2	7	14
2002.....	369	-	-	-	-	-	-	2	1	4	8	16
2001.....	415	-	-	-	-	-	-	-	2	5	9	13
2000.....	428	-	-	-	-	-	-	2	1	12	7	18
1999.....	384	-	-	-	-	-	-	-	3	6	9	11
All races, female												
2003.....	41,620	-	-	-	-	1	15	54	351	910	1,797	2,720
2002.....	41,514	-	-	-	-	1	9	61	361	900	1,810	2,731
2001.....	41,394	-	-	-	-	1	8	88	373	955	1,854	2,777
2000.....	41,872	-	-	-	-	1	10	81	383	970	1,852	2,712
1999.....	41,144	-	1	-	1	3	10	86	350	980	1,746	2,608

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
White, both sexes												
2003.....	35,358	-	-	-	-	1	10	27	242	652	1,329	2,052
2002.....	35,382	-	-	-	-	-	6	40	245	634	1,316	2,059
2001.....	35,422	-	-	-	-	-	4	58	248	697	1,372	2,135
2000.....	36,127	-	-	-	-	1	4	60	265	717	1,380	2,066
1999.....	35,385	-	-	-	1	3	5	59	245	700	1,301	1,977
White, male												
2003.....	326	-	-	-	-	-	-	-	1	1	3	9
2002.....	300	-	-	-	-	-	-	2	-	2	7	11
2001.....	344	-	-	-	-	-	-	-	1	4	4	10
2000.....	360	-	-	-	-	-	-	-	1	8	3	13
1999.....	312	-	-	-	-	-	-	-	2	-	8	8
White, female												
2003.....	35,032	-	-	-	-	1	10	27	241	651	1,326	2,043
2002.....	35,082	-	-	-	-	-	6	38	245	632	1,309	2,048
2001.....	35,078	-	-	-	-	-	4	58	247	693	1,368	2,125
2000.....	35,767	-	-	-	-	1	4	60	264	709	1,377	2,053
1999.....	35,073	-	-	-	1	3	5	59	243	700	1,293	1,969

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

Black, both sexes

2003.....	5,746	-	-	-	-	-	5	24	95	228	412	595
2002.....	5,636	-	-	-	-	1	2	22	103	234	445	585
2001.....	5,571	-	-	-	-	1	4	30	112	221	424	556
2000.....	5,424	-	-	-	-	-	5	20	103	228	417	564
1999.....	5,425	-	1	-	-	-	5	25	101	261	404	553

Black, male

2003.....	51	-	-	-	-	-	-	1	-	1	4	4
2002.....	64	-	-	-	-	-	-	-	-	2	1	5
2001.....	65	-	-	-	-	-	-	-	1	-	5	3
2000.....	63	-	-	-	-	-	-	2	-	4	4	4
1999.....	67	-	-	-	-	-	-	-	1	6	1	1

Black, female

2003.....	5,695	-	-	-	-	-	5	23	95	227	408	591
2002.....	5,572	-	-	-	-	1	2	22	102	232	444	580
2001.....	5,506	-	-	-	-	1	4	30	111	221	419	553
2000.....	5,361	-	-	-	-	-	5	18	103	224	413	560
1999.....	5,358	-	1	-	-	-	5	25	100	255	403	552

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
American Indian 1/, both sexes												
2003.....	148	-	-	-	-	-	-	2	2	4	5	8
2002.....	148	-	-	-	-	-	-	1	4	10	7	14
2001.....	127	-	-	-	-	-	-	-	2	9	6	15
2000.....	131	-	-	-	-	-	-	1	1	3	13	11
1999.....	131	-	-	-	-	-	-	1	1	3	8	9
American Indian 1/, male												
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2003.....	147	-	-	-	-	-	-	2	2	4	5	8
2002.....	148	-	-	-	-	-	-	1	4	10	7	14
2001.....	126	-	-	-	-	-	-	-	2	9	6	15
2000.....	130	-	-	-	-	-	-	1	1	3	13	11
1999.....	130	-	-	-	-	-	-	1	1	3	8	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
Asian or Pacific Islander 2/, both sexes												
2003.....	748	-	-	-	-	-	-	2	13	28	58	79
2002.....	717	-	-	-	-	-	1	-	10	26	50	89
2001.....	689	-	-	-	-	-	-	-	13	33	61	84
2000.....	618	-	-	-	-	-	1	2	15	34	49	89
1999.....	587	-	-	-	-	-	-	1	6	22	42	80
Asian or Pacific Islander 2/, male												
2003.....	2	-	-	-	-	-	-	-	-	-	-	1
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	1	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	1
1999.....	4	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, female												
2003.....	746	-	-	-	-	-	-	2	13	28	58	78
2002.....	712	-	-	-	-	-	1	-	10	26	50	89
2001.....	684	-	-	-	-	-	-	-	13	32	61	84
2000.....	614	-	-	-	-	-	1	2	15	34	49	88
1999.....	583	-	-	-	-	-	-	1	6	22	42	78

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of cervix uteri (C53)												
All races, both sexes												
2003.....	3,919	-	-	-	-	1	15	59	142	250	383	440
2002.....	3,952	-	-	-	-	1	16	47	161	275	391	439
2001.....	4,092	-	-	-	1	-	12	51	158	291	420	475
2000.....	4,200	-	-	-	1	3	14	77	153	294	400	448
1999.....	4,205	-	-	-	-	2	10	62	155	292	407	462
All races, male												
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2003.....	3,919	-	-	-	-	1	15	59	142	250	383	440
2002.....	3,952	-	-	-	-	1	16	47	161	275	391	439
2001.....	4,092	-	-	-	1	-	12	51	158	291	420	475
2000.....	4,200	-	-	-	1	3	14	77	153	294	400	448
1999.....	4,205	-	-	-	-	2	10	62	155	292	407	462

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2003.....	2,953	-	-	-	-	1	12	40	99	182	296	334
2002.....	2,967	-	-	-	-	-	12	34	126	211	292	324
2001.....	3,136	-	-	-	-	-	11	40	122	226	325	371
2000.....	3,158	-	-	-	-	3	10	63	111	223	288	334
1999.....	3,229	-	-	-	-	-	9	52	124	221	314	371

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	2,953	-	-	-	-	1	12	40	99	182	296	334
2002.....	2,967	-	-	-	-	-	12	34	126	211	292	324
2001.....	3,136	-	-	-	-	-	11	40	122	226	325	371
2000.....	3,158	-	-	-	-	3	10	63	111	223	288	334
1999.....	3,229	-	-	-	-	-	9	52	124	221	314	371

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2003.....	794	-	-	-	-	-	2	17	33	57	70	92
2002.....	829	-	-	-	-	1	2	13	30	60	86	95
2001.....	787	-	-	-	1	-	1	11	31	54	85	80
2000.....	869	-	-	-	1	-	4	12	39	62	88	88
1999.....	838	-	-	-	-	2	1	9	27	61	83	68

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	794	-	-	-	-	-	2	17	33	57	70	92
2002.....	829	-	-	-	-	1	2	13	30	60	86	95
2001.....	787	-	-	-	1	-	1	11	31	54	85	80
2000.....	869	-	-	-	1	-	4	12	39	62	88	88
1999.....	838	-	-	-	-	2	1	9	27	61	83	68

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

American Indian 1/, both sexes

2003.....	31	-	-	-	-	-	-	1	1	3	4	2
2002.....	34	-	-	-	-	-	2	-	2	-	3	9
2001.....	32	-	-	-	-	-	-	-	1	5	2	9
2000.....	30	-	-	-	-	-	-	-	2	2	5	3
1999.....	19	-	-	-	-	-	-	1	1	3	-	4

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	31	-	-	-	-	-	-	1	1	3	4	2
2002.....	34	-	-	-	-	-	2	-	2	-	3	9
2001.....	32	-	-	-	-	-	-	-	1	5	2	9
2000.....	30	-	-	-	-	-	-	-	2	2	5	3
1999.....	19	-	-	-	-	-	-	1	1	3	-	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

Asian or Pacific Islander 2/, both sexes

2003.....	141	-	-	-	-	-	1	1	9	8	13	12
2002.....	122	-	-	-	-	-	-	-	3	4	10	11
2001.....	137	-	-	-	-	-	-	-	4	6	8	15
2000.....	143	-	-	-	-	-	-	2	1	7	19	23
1999.....	119	-	-	-	-	-	-	-	3	7	10	19

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	141	-	-	-	-	-	1	1	9	8	13	12
2002.....	122	-	-	-	-	-	-	-	3	4	10	11
2001.....	137	-	-	-	-	-	-	-	4	6	8	15
2000.....	143	-	-	-	-	-	-	2	1	7	19	23
1999.....	119	-	-	-	-	-	-	-	3	7	10	19

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races, both sexes

2003.....	6,899	-	-	-	-	1	3	8	19	37	117	171
2002.....	6,853	-	-	-	-	-	-	2	21	46	100	189
2001.....	6,784	-	-	-	-	-	-	10	28	47	92	192
2000.....	6,586	-	-	-	-	-	3	2	25	41	97	190
1999.....	6,468	-	-	-	-	-	2	11	12	38	104	182

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	6,899	-	-	-	-	1	3	8	19	37	117	171
2002.....	6,853	-	-	-	-	-	-	2	21	46	100	189
2001.....	6,784	-	-	-	-	-	-	10	28	47	92	192
2000.....	6,586	-	-	-	-	-	3	2	25	41	97	190
1999.....	6,468	-	-	-	-	-	2	11	12	38	104	182

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2003.....	5,641	-	-	-	-	1	2	5	11	29	79	133
2002.....	5,567	-	-	-	-	-	-	1	16	34	82	147
2001.....	5,606	-	-	-	-	-	-	3	23	33	74	150
2000.....	5,397	-	-	-	-	-	1	2	18	33	73	158
1999.....	5,386	-	-	-	-	-	2	8	10	31	78	150

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	5,641	-	-	-	-	1	2	5	11	29	79	133
2002.....	5,567	-	-	-	-	-	-	1	16	34	82	147
2001.....	5,606	-	-	-	-	-	-	3	23	33	74	150
2000.....	5,397	-	-	-	-	-	1	2	18	33	73	158
1999.....	5,386	-	-	-	-	-	2	8	10	31	78	150

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2003.....	1,107	-	-	-	-	-	1	2	6	6	25	29
2002.....	1,119	-	-	-	-	-	-	1	4	8	14	28
2001.....	1,047	-	-	-	-	-	-	6	3	8	14	34
2000.....	1,061	-	-	-	-	-	1	-	6	6	16	24
1999.....	987	-	-	-	-	-	-	3	2	5	23	18

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	1,107	-	-	-	-	-	1	2	6	6	25	29
2002.....	1,119	-	-	-	-	-	-	1	4	8	14	28
2001.....	1,047	-	-	-	-	-	-	6	3	8	14	34
2000.....	1,061	-	-	-	-	-	1	-	6	6	16	24
1999.....	987	-	-	-	-	-	-	3	2	5	23	18

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

American Indian 1/, both sexes

2003.....	11	-	-	-	-	-	-	1	-	-	1	1
2002.....	32	-	-	-	-	-	-	-	-	3	1	4
2001.....	25	-	-	-	-	-	-	-	-	2	-	-
2000.....	19	-	-	-	-	-	-	-	-	-	1	1
1999.....	13	-	-	-	-	-	-	-	-	1	1	2

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	11	-	-	-	-	-	-	1	-	-	1	1
2002.....	32	-	-	-	-	-	-	-	-	3	1	4
2001.....	25	-	-	-	-	-	-	-	-	2	-	-
2000.....	19	-	-	-	-	-	-	-	-	-	1	1
1999.....	13	-	-	-	-	-	-	-	-	1	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Asian or Pacific Islander 2/, both sexes

2003.....	140	-	-	-	-	-	-	-	2	2	12	8
2002.....	135	-	-	-	-	-	-	-	1	1	3	10
2001.....	106	-	-	-	-	-	-	1	2	4	4	8
2000.....	109	-	-	-	-	-	1	-	1	2	7	7
1999.....	82	-	-	-	-	-	-	-	-	1	2	12

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	140	-	-	-	-	-	-	-	2	2	12	8
2002.....	135	-	-	-	-	-	-	-	1	1	3	10
2001.....	106	-	-	-	-	-	-	1	2	4	4	8
2000.....	109	-	-	-	-	-	1	-	1	2	7	7
1999.....	82	-	-	-	-	-	-	-	-	1	2	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of ovary (C56)												
All races, both sexes												
2003.....	14,657	-	-	-	-	10	18	31	53	134	318	632
2002.....	14,682	1	1	1	1	12	25	33	64	150	319	688
2001.....	14,414	-	-	-	2	5	16	37	65	152	320	610
2000.....	14,060	-	-	2	-	7	16	36	65	145	327	617
1999.....	13,627	-	1	-	2	9	18	26	51	162	334	616
All races, male												
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2003.....	14,657	-	-	-	-	10	18	31	53	134	318	632
2002.....	14,682	1	1	1	1	12	25	33	64	150	319	688
2001.....	14,414	-	-	-	2	5	16	37	65	152	320	610
2000.....	14,060	-	-	2	-	7	16	36	65	145	327	617
1999.....	13,627	-	1	-	2	9	18	26	51	162	334	616

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of ovary (C56)												
White, both sexes												
2003.....	13,123	-	-	-	-	8	13	23	46	103	270	536
2002.....	13,202	1	1	1	1	9	21	29	48	122	269	582
2001.....	12,950	-	-	-	1	3	11	32	49	120	270	536
2000.....	12,731	-	-	1	-	5	12	32	52	120	277	541
1999.....	12,302	-	1	-	2	8	14	20	37	136	288	535
White, male												
2003.....
2002.....
2001.....
2000.....
1999.....
White, female												
2003.....	13,123	-	-	-	-	8	13	23	46	103	270	536
2002.....	13,202	1	1	1	1	9	21	29	48	122	269	582
2001.....	12,950	-	-	-	1	3	11	32	49	120	270	536
2000.....	12,731	-	-	1	-	5	12	32	52	120	277	541
1999.....	12,302	-	1	-	2	8	14	20	37	136	288	535

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of ovary (C56)

Black, both sexes

2003.....	1,196	-	-	-	-	2	4	7	5	23	35	74
2002.....	1,163	-	-	-	-	3	4	3	10	24	35	73
2001.....	1,153	-	-	-	1	1	3	5	10	25	41	48
2000.....	1,073	-	-	1	-	-	2	4	8	17	43	51
1999.....	1,079	-	-	-	-	1	3	6	7	16	31	55

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	1,196	-	-	-	-	2	4	7	5	23	35	74
2002.....	1,163	-	-	-	-	3	4	3	10	24	35	73
2001.....	1,153	-	-	-	1	1	3	5	10	25	41	48
2000.....	1,073	-	-	1	-	-	2	4	8	17	43	51
1999.....	1,079	-	-	-	-	1	3	6	7	16	31	55

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of ovary (C56)

Asian or Pacific Islander 2/, both sexes

2003.....	276	-	-	-	-	-	1	1	2	7	12	18
2002.....	267	-	-	-	-	-	-	-	5	3	12	27
2001.....	243	-	-	-	-	1	2	-	6	4	8	21
2000.....	228	-	-	-	-	2	1	-	5	7	7	23
1999.....	217	-	-	-	-	-	1	-	7	10	14	25

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	276	-	-	-	-	-	1	1	2	7	12	18
2002.....	267	-	-	-	-	-	-	-	5	3	12	27
2001.....	243	-	-	-	-	1	2	-	6	4	8	21
2000.....	228	-	-	-	-	2	1	-	5	7	7	23
1999.....	217	-	-	-	-	-	1	-	7	10	14	25

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All races, both sexes												
2003.....	12,286	1	14	28	11	11	13	19	37	81	240	438
2002.....	12,165	4	8	23	7	7	13	14	34	91	215	514
2001.....	12,078	1	22	30	5	8	12	19	33	74	284	498
2000.....	11,736	2	15	21	8	8	12	14	28	83	251	450
1999.....	11,116	2	19	23	10	8	9	25	29	105	237	427
All races, male												
2003.....	7,667	1	4	15	4	7	9	9	27	56	175	324
2002.....	7,587	1	-	14	5	2	5	8	23	63	154	355
2001.....	7,524	1	10	7	2	7	9	11	16	58	182	355
2000.....	7,249	1	8	12	3	4	6	7	19	53	175	309
1999.....	6,867	1	11	12	4	4	7	15	22	70	166	310
All races, female												
2003.....	4,619	-	10	13	7	4	4	10	10	25	65	114
2002.....	4,578	3	8	9	2	5	8	6	11	28	61	159
2001.....	4,554	-	12	23	3	1	3	8	17	16	102	143
2000.....	4,487	1	7	9	5	4	6	7	9	30	76	141
1999.....	4,249	1	8	11	6	4	2	10	7	35	71	117

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
White, both sexes												
2003.....	10,930	1	14	24	6	8	8	12	23	56	191	358
2002.....	10,856	4	8	18	6	5	11	10	20	66	182	428
2001.....	10,720	-	16	21	2	2	6	10	21	51	223	424
2000.....	10,480	-	10	12	7	6	7	8	23	59	211	381
1999.....	9,878	2	12	19	6	5	5	14	18	76	195	346
White, male												
2003.....	6,840	1	4	12	2	5	6	4	16	41	139	274
2002.....	6,803	1	-	11	4	1	4	5	14	45	131	304
2001.....	6,695	-	6	4	1	1	4	7	9	41	151	312
2000.....	6,475	-	5	6	3	3	4	2	15	42	150	266
1999.....	6,112	1	8	10	1	4	3	7	13	51	138	255
White, female												
2003.....	4,090	-	10	12	4	3	2	8	7	15	52	84
2002.....	4,053	3	8	7	2	4	7	5	6	21	51	124
2001.....	4,025	-	10	17	1	1	2	3	12	10	72	112
2000.....	4,005	-	5	6	4	3	3	6	8	17	61	115
1999.....	3,766	1	4	9	5	1	2	7	5	25	57	91

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, both sexes

2003.....	1,116	-	-	4	5	3	5	7	12	20	44	65
2002.....	1,069	-	-	5	1	2	1	3	14	21	25	72
2001.....	1,113	-	4	9	3	5	5	9	10	21	51	64
2000.....	1,044	1	5	7	1	2	4	6	4	21	31	58
1999.....	1,035	-	4	3	3	3	4	11	10	27	34	67

Black, male

2003.....	677	-	-	3	2	2	3	5	9	10	34	44
2002.....	630	-	-	3	1	1	1	2	9	15	17	41
2001.....	678	-	2	3	1	5	5	4	6	15	25	40
2000.....	632	-	3	4	-	1	1	5	3	9	18	35
1999.....	622	-	2	1	2	-	4	8	8	19	23	44

Black, female

2003.....	439	-	-	1	3	1	2	2	3	10	10	21
2002.....	439	-	-	2	-	1	-	1	5	6	8	31
2001.....	435	-	2	6	2	-	-	5	4	6	26	24
2000.....	412	1	2	3	1	1	3	1	1	12	13	23
1999.....	413	-	2	2	1	3	-	3	2	8	11	23

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Asian or Pacific Islander 2/, both sexes

2003.....	161	-	-	-	-	-	-	-	2	2	3	11
2002.....	142	-	-	-	-	-	1	1	-	3	2	7
2001.....	158	-	1	-	-	-	1	-	2	1	7	6
2000.....	148	-	-	1	-	-	-	-	-	2	5	5
1999.....	125	-	1	1	1	-	-	-	1	2	5	10

Asian or Pacific Islander 2/, male

2003.....	102	-	-	-	-	-	-	-	2	2	2	4
2002.....	91	-	-	-	-	-	-	1	-	3	2	6
2001.....	97	-	1	-	-	-	-	-	1	1	5	1
2000.....	99	-	-	1	-	-	-	-	-	1	4	5
1999.....	83	-	1	1	1	-	-	-	1	-	4	8

Asian or Pacific Islander 2/, female

2003.....	59	-	-	-	-	-	-	-	-	-	1	7
2002.....	51	-	-	-	-	-	1	-	-	-	-	1
2001.....	61	-	-	-	-	-	1	-	1	-	2	5
2000.....	49	-	-	-	-	-	-	-	-	1	1	-
1999.....	42	-	-	-	-	-	-	-	-	2	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of bladder (C67)												
All races, both sexes												
2003.....	12,483	-	-	-	-	-	1	1	3	24	74	181
2002.....	12,628	-	-	2	1	-	1	3	11	31	92	182
2001.....	12,225	-	-	1	-	1	2	6	9	35	93	161
2000.....	12,002	-	-	-	1	2	1	1	9	40	71	160
1999.....	11,911	-	-	-	1	-	1	6	10	32	68	140
All races, male												
2003.....	8,576	-	-	-	-	-	1	1	1	15	49	149
2002.....	8,504	-	-	1	-	-	-	2	7	21	57	129
2001.....	8,343	-	-	-	-	1	2	1	7	14	56	118
2000.....	8,163	-	-	-	-	2	1	-	2	21	41	110
1999.....	8,105	-	-	-	1	-	-	3	6	20	43	107
All races, female												
2003.....	3,907	-	-	-	-	-	-	-	2	9	25	32
2002.....	4,124	-	-	1	1	-	1	1	4	10	35	53
2001.....	3,882	-	-	1	-	-	-	5	2	21	37	43
2000.....	3,839	-	-	-	1	-	-	1	7	19	30	50
1999.....	3,806	-	-	-	-	-	1	3	4	12	25	33

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of bladder (C67)												
White, both sexes												
2003.....	11,409	-	-	-	-	-	1	1	3	21	54	145
2002.....	11,512	-	-	2	1	-	-	2	8	29	76	163
2001.....	11,180	-	-	1	-	1	2	6	8	30	74	133
2000.....	11,036	-	-	-	1	1	1	1	8	36	63	137
1999.....	10,863	-	-	-	1	-	-	3	7	23	57	114
White, male												
2003.....	7,965	-	-	-	-	-	1	1	1	13	36	123
2002.....	7,908	-	-	1	-	-	-	1	5	19	49	116
2001.....	7,780	-	-	-	-	1	2	1	7	13	44	103
2000.....	7,620	-	-	-	-	1	1	-	2	20	37	96
1999.....	7,520	-	-	-	1	-	-	1	4	13	36	87
White, female												
2003.....	3,444	-	-	-	-	-	-	-	2	8	18	22
2002.....	3,604	-	-	1	1	-	-	1	3	10	27	47
2001.....	3,400	-	-	1	-	-	-	5	1	17	30	30
2000.....	3,416	-	-	-	1	-	-	1	6	16	26	41
1999.....	3,343	-	-	-	-	-	-	2	3	10	21	27

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of bladder (C67)												
Black, both sexes												
2003.....	914	-	-	-	-	-	-	-	-	3	18	31
2002.....	940	-	-	-	-	-	-	1	3	1	12	16
2001.....	877	-	-	-	-	-	-	-	1	4	19	21
2000.....	852	-	-	-	-	1	-	-	-	3	7	20
1999.....	918	-	-	-	-	-	1	2	2	9	9	23
Black, male												
2003.....	502	-	-	-	-	-	-	-	-	2	11	21
2002.....	481	-	-	-	-	-	-	1	2	1	6	10
2001.....	447	-	-	-	-	-	-	-	-	1	12	10
2000.....	466	-	-	-	-	1	-	-	-	-	4	11
1999.....	496	-	-	-	-	-	-	1	1	7	6	17
Black, female												
2003.....	412	-	-	-	-	-	-	-	-	1	7	10
2002.....	459	-	-	-	-	-	-	-	1	-	6	6
2001.....	430	-	-	-	-	-	-	-	1	3	7	11
2000.....	386	-	-	-	-	-	-	-	-	3	3	9
1999.....	422	-	-	-	-	-	1	1	1	2	3	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races, both sexes

2003.....	12,901	18	104	175	161	119	117	165	240	411	668	857
2002.....	12,830	12	114	202	135	108	102	129	259	375	634	900
2001.....	12,609	16	100	170	130	104	120	134	282	399	629	830
2000.....	12,655	20	104	166	150	105	120	159	247	454	642	815
1999.....	12,765	10	113	184	124	115	103	177	267	443	622	842

All races, male

2003.....	7,122	9	56	89	86	68	68	85	147	246	393	544
2002.....	7,125	8	63	112	79	65	71	67	156	222	396	560
2001.....	7,015	8	52	81	64	63	66	88	174	245	393	500
2000.....	7,011	10	58	89	87	63	69	94	170	293	411	481
1999.....	7,001	8	56	96	58	69	62	104	172	268	370	507

All races, female

2003.....	5,779	9	48	86	75	51	49	80	93	165	275	313
2002.....	5,705	4	51	90	56	43	31	62	103	153	238	340
2001.....	5,594	8	48	89	66	41	54	46	108	154	236	330
2000.....	5,644	10	46	77	63	42	51	65	77	161	231	334
1999.....	5,764	2	57	88	66	46	41	73	95	175	252	335

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
White, both sexes												
2003.....	11,849	15	85	140	131	99	98	144	211	366	610	767
2002.....	11,794	7	80	159	106	88	86	109	229	341	581	826
2001.....	11,638	12	84	144	100	88	95	120	254	374	568	753
2000.....	11,672	16	82	135	120	84	99	140	220	405	564	745
1999.....	11,819	7	96	139	103	86	85	165	238	408	558	775
White, male												
2003.....	6,543	8	48	70	69	55	57	75	131	217	355	495
2002.....	6,554	5	46	84	59	55	62	56	135	203	369	514
2001.....	6,498	6	46	66	47	54	53	81	157	232	357	461
2000.....	6,486	9	46	72	69	52	57	82	145	264	366	449
1999.....	6,492	6	49	75	50	50	51	97	153	249	328	463
White, female												
2003.....	5,306	7	37	70	62	44	41	69	80	149	255	272
2002.....	5,240	2	34	75	47	33	24	53	94	138	212	312
2001.....	5,140	6	38	78	53	34	42	39	97	142	211	292
2000.....	5,186	7	36	63	51	32	42	58	75	141	198	296
1999.....	5,327	1	47	64	53	36	34	68	85	159	230	312

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2003.....	811	2	17	27	21	17	17	19	17	35	48	70
2002.....	781	3	27	34	20	16	12	11	22	30	38	49
2001.....	741	1	10	19	25	12	17	8	20	18	45	64
2000.....	772	2	18	26	26	16	19	15	21	34	57	52
1999.....	764	3	12	37	18	23	14	10	26	26	51	59

Black, male

2003.....	447	1	6	16	11	11	10	9	10	23	33	38
2002.....	409	2	12	23	12	9	5	5	17	16	19	25
2001.....	382	-	3	12	13	6	10	4	10	8	25	34
2000.....	414	1	10	15	16	9	12	9	19	19	35	24
1999.....	404	2	5	18	7	15	9	6	17	15	33	39

Black, female

2003.....	364	1	11	11	10	6	7	10	7	12	15	32
2002.....	372	1	15	11	8	7	7	6	5	14	19	24
2001.....	359	1	7	7	12	6	7	4	10	10	20	30
2000.....	358	1	8	11	10	7	7	6	2	15	22	28
1999.....	360	1	7	19	11	8	5	4	9	11	18	20

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

American Indian 1/, both sexes

2003.....	45	-	1	2	2	1	1	-	1	2	1	5
2002.....	41	1	1	4	3	2	-	1	3	-	3	4
2001.....	43	2	1	-	-	1	-	1	1	5	2	2
2000.....	41	-	2	2	2	2	-	1	-	7	4	4
1999.....	34	-	1	1	-	1	1	-	-	2	7	-

American Indian 1/, male

2003.....	26	-	1	1	1	1	-	-	-	1	1	4
2002.....	23	-	-	2	2	1	-	1	-	-	2	4
2001.....	25	2	1	-	-	-	-	1	1	4	2	-
2000.....	20	-	1	1	1	1	-	-	-	4	1	1
1999.....	19	-	-	-	-	1	-	-	-	2	5	-

American Indian 1/, female

2003.....	19	-	-	1	1	-	1	-	1	1	-	1
2002.....	18	1	1	2	1	1	-	-	3	-	1	-
2001.....	18	-	-	-	-	1	-	-	-	1	-	2
2000.....	21	-	1	1	1	1	-	1	-	3	3	3
1999.....	15	-	1	1	-	-	1	-	-	-	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Asian or Pacific Islander 2/, both sexes

2003.....	196	1	1	6	7	2	1	2	11	8	9	15
2002.....	214	1	6	5	6	2	4	8	5	4	12	21
2001.....	187	1	5	7	5	3	8	5	7	2	14	11
2000.....	170	2	2	3	2	3	2	3	6	8	17	14
1999.....	148	-	4	7	3	5	3	2	3	7	6	8

Asian or Pacific Islander 2/, male

2003.....	106	-	1	2	5	1	1	1	6	5	4	7
2002.....	139	1	5	3	6	-	4	5	4	3	6	17
2001.....	110	-	2	3	4	3	3	2	6	1	9	5
2000.....	91	-	1	1	1	1	-	3	6	6	9	7
1999.....	86	-	2	3	1	3	2	1	2	2	4	5

Asian or Pacific Islander 2/, female

2003.....	90	1	-	4	2	1	-	1	5	3	5	8
2002.....	75	-	1	2	-	2	-	3	1	1	6	4
2001.....	77	1	3	4	1	-	5	3	1	1	5	6
2000.....	79	2	1	2	1	2	2	-	-	2	8	7
1999.....	62	-	2	4	2	2	1	1	1	5	2	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All races, both sexes												
2003.....	55,679	31	138	173	221	252	354	373	492	655	1,073	1,557
2002.....	56,225	28	119	166	213	310	393	368	548	695	1,056	1,599
2001.....	56,341	27	157	165	202	311	394	436	552	768	1,089	1,674
2000.....	56,518	39	146	168	204	323	395	412	506	776	1,077	1,675
1999.....	56,318	30	137	142	209	303	417	423	537	837	1,076	1,648
All races, male												
2003.....	30,076	10	80	102	123	166	209	220	315	380	663	947
2002.....	30,302	15	63	87	130	188	229	225	320	409	629	953
2001.....	30,060	12	82	95	122	192	234	246	328	464	669	1,002
2000.....	29,888	21	76	98	125	202	253	226	313	438	637	1,015
1999.....	29,721	12	75	84	122	180	257	241	316	497	647	986
All races, female												
2003.....	25,603	21	58	71	98	86	145	153	177	275	410	610
2002.....	25,923	13	56	79	83	122	164	143	228	286	427	646
2001.....	26,281	15	75	70	80	119	160	190	224	304	420	672
2000.....	26,630	18	70	70	79	121	142	186	193	338	440	660
1999.....	26,597	18	62	58	87	123	160	182	221	340	429	662

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
White, both sexes												
2003.....	49,420	29	115	146	168	201	282	292	378	509	857	1,258
2002.....	49,943	19	98	137	166	244	331	287	423	542	863	1,276
2001.....	49,989	22	136	125	156	253	309	318	435	612	858	1,364
2000.....	50,394	36	121	136	170	249	300	315	392	617	868	1,340
1999.....	50,189	24	118	114	157	232	332	332	419	677	862	1,362
White, male												
2003.....	26,943	8	67	92	92	136	165	175	244	300	536	775
2002.....	27,087	11	53	69	101	155	197	180	246	316	516	774
2001.....	26,767	9	69	72	85	159	187	182	269	360	532	833
2000.....	26,766	19	64	77	103	155	196	180	240	354	536	827
1999.....	26,558	12	63	70	90	136	213	186	245	408	518	816
White, female												
2003.....	22,477	21	48	54	76	65	117	117	134	209	321	483
2002.....	22,856	8	45	68	65	89	134	107	177	226	347	502
2001.....	23,222	13	67	53	71	94	122	136	166	252	326	531
2000.....	23,628	17	57	59	67	94	104	135	152	263	332	513
1999.....	23,631	12	55	44	67	96	119	146	174	269	344	546

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2003.....	5,103	1	14	20	37	34	54	65	91	117	184	250
2002.....	5,151	5	16	19	34	45	45	62	95	111	165	268
2001.....	5,227	4	14	29	36	48	63	85	98	133	190	257
2000.....	5,071	1	20	21	22	58	74	75	85	130	176	295
1999.....	5,088	4	15	21	42	54	80	72	88	135	179	239

Black, male

2003.....	2,537	1	8	9	22	19	33	32	56	63	109	146
2002.....	2,622	2	7	13	20	25	20	35	58	70	98	155
2001.....	2,676	3	9	14	30	29	32	48	52	90	110	136
2000.....	2,558	-	9	14	12	36	44	33	58	66	87	163
1999.....	2,590	-	9	11	28	33	41	43	55	74	108	141

Black, female

2003.....	2,566	-	6	11	15	15	21	33	35	54	75	104
2002.....	2,529	3	9	6	14	20	25	27	37	41	67	113
2001.....	2,551	1	5	15	6	19	31	37	46	43	80	121
2000.....	2,513	1	11	7	10	22	30	42	27	64	89	132
1999.....	2,498	4	6	10	14	21	39	29	33	61	71	98

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

American Indian 1/, both sexes

2003.....	210	-	3	2	1	7	5	2	3	5	7	9
2002.....	171	-	1	-	2	4	7	3	4	10	4	7
2001.....	225	-	1	-	4	3	7	11	3	5	13	7
2000.....	199	1	1	1	1	3	2	2	7	4	5	9
1999.....	161	1	1	-	2	1	2	4	3	3	6	7

American Indian 1/, male

2003.....	99	-	3	-	-	4	5	2	3	2	4	5
2002.....	89	-	1	-	1	2	6	1	2	6	4	3
2001.....	113	-	1	-	2	2	5	8	1	-	9	5
2000.....	101	1	1	-	-	2	1	2	4	3	1	6
1999.....	92	-	1	-	2	-	1	3	1	1	3	5

American Indian 1/, female

2003.....	111	-	-	2	1	3	-	-	-	3	3	4
2002.....	82	-	-	-	1	2	1	2	2	4	-	4
2001.....	112	-	-	-	2	1	2	3	2	5	4	2
2000.....	98	-	-	1	1	1	1	-	3	1	4	3
1999.....	69	1	-	-	-	1	1	1	2	2	3	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Asian or Pacific Islander 2/, both sexes

2003.....	946	1	6	5	15	10	13	14	20	24	25	40
2002.....	960	4	4	10	11	17	10	16	26	32	24	48
2001.....	900	1	6	11	6	7	15	22	16	18	28	46
2000.....	854	1	4	10	11	13	19	20	22	25	28	31
1999.....	880	1	3	7	8	16	3	15	27	22	29	40

Asian or Pacific Islander 2/, male

2003.....	497	1	2	1	9	7	6	11	12	15	14	21
2002.....	504	2	2	5	8	6	6	9	14	17	11	21
2001.....	504	-	3	9	5	2	10	8	6	14	18	28
2000.....	463	1	2	7	10	9	12	11	11	15	13	19
1999.....	481	-	2	3	2	11	2	9	15	14	18	24

Asian or Pacific Islander 2/, female

2003.....	449	-	4	4	6	3	7	3	8	9	11	19
2002.....	456	2	2	5	3	11	4	7	12	15	13	27
2001.....	396	1	3	2	1	5	5	14	10	4	10	18
2000.....	391	-	2	3	1	4	7	9	11	10	15	12
1999.....	399	1	1	4	6	5	1	6	12	8	11	16

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
All races, both sexes												
2003.....	1,347	-	-	-	5	16	48	71	96	76	78	71
2002.....	1,352	-	-	3	3	27	59	64	88	81	78	83
2001.....	1,323	-	-	1	5	25	53	76	91	81	80	91
2000.....	1,287	-	-	3	5	21	52	60	85	78	73	73
1999.....	1,403	-	1	-	9	31	63	77	71	90	75	82
All races, male												
2003.....	725	-	-	-	4	7	23	35	53	43	52	44
2002.....	751	-	-	2	3	15	26	37	44	46	46	53
2001.....	725	-	-	1	2	15	28	39	51	50	51	54
2000.....	686	-	-	1	3	6	29	31	48	42	40	46
1999.....	785	-	1	-	3	20	35	42	44	56	47	48
All races, female												
2003.....	622	-	-	-	1	9	25	36	43	33	26	27
2002.....	601	-	-	1	-	12	33	27	44	35	32	30
2001.....	598	-	-	-	3	10	25	37	40	31	29	37
2000.....	601	-	-	2	2	15	23	29	37	36	33	27
1999.....	618	-	-	-	6	11	28	35	27	34	28	34

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
White, both sexes												
2003.....	1,192	-	-	-	5	15	36	56	78	65	67	64
2002.....	1,193	-	-	3	2	21	51	52	75	64	66	72
2001.....	1,170	-	-	1	4	21	41	56	69	69	69	77
2000.....	1,156	-	-	2	3	17	40	54	76	66	61	66
1999.....	1,263	-	1	-	7	26	49	60	62	80	66	72
White, male												
2003.....	638	-	-	-	4	7	17	30	44	39	45	40
2002.....	655	-	-	2	2	13	24	28	37	36	37	44
2001.....	644	-	-	1	2	13	25	27	42	41	44	46
2000.....	616	-	-	1	1	4	26	28	42	36	34	42
1999.....	708	-	1	-	3	17	28	32	40	49	43	41
White, female												
2003.....	554	-	-	-	1	8	19	26	34	26	22	24
2002.....	538	-	-	1	-	8	27	24	38	28	29	28
2001.....	526	-	-	-	2	8	16	29	27	28	25	31
2000.....	540	-	-	1	2	13	14	26	34	30	27	24
1999.....	555	-	-	-	4	9	21	28	22	31	23	31

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
Black, both sexes												
2003.....	120	-	-	-	-	-	10	15	16	9	9	6
2002.....	130	-	-	-	1	4	7	11	13	15	11	10
2001.....	137	-	-	-	1	4	11	19	21	11	10	12
2000.....	110	-	-	1	1	4	10	6	7	12	12	7
1999.....	128	-	-	-	1	5	14	17	9	10	8	10
Black, male												
2003.....	68	-	-	-	-	-	5	5	7	4	6	3
2002.....	82	-	-	-	1	2	2	8	7	9	9	8
2001.....	73	-	-	-	-	2	2	12	9	8	7	7
2000.....	57	-	-	-	1	2	3	3	5	6	6	4
1999.....	75	-	-	-	-	3	7	10	4	7	4	7
Black, female												
2003.....	52	-	-	-	-	-	5	10	9	5	3	3
2002.....	48	-	-	-	-	2	5	3	6	6	2	2
2001.....	64	-	-	-	1	2	9	7	12	3	3	5
2000.....	53	-	-	1	-	2	7	3	2	6	6	3
1999.....	53	-	-	-	1	2	7	7	5	3	4	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hodgkin's disease (C81)

Asian or Pacific Islander 2/, both sexes

2003.....	26	-	-	-	-	1	2	-	2	2	1	1
2002.....	28	-	-	-	-	2	1	1	-	2	1	1
2001.....	13	-	-	-	-	-	1	1	-	1	-	2
2000.....	16	-	-	-	1	-	2	-	2	-	-	-
1999.....	7	-	-	-	1	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	15	-	-	-	-	-	1	-	2	-	-	1
2002.....	14	-	-	-	-	-	-	1	-	1	-	1
2001.....	8	-	-	-	-	-	1	-	-	1	-	1
2000.....	9	-	-	-	1	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	11	-	-	-	-	1	1	-	-	2	1	-
2002.....	14	-	-	-	-	2	1	-	-	1	1	-
2001.....	5	-	-	-	-	-	-	1	-	-	-	1
2000.....	7	-	-	-	-	-	2	-	1	-	-	-
1999.....	6	-	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
All races, both sexes												
2003.....	21,475	-	5	17	28	53	87	96	156	233	397	649
2002.....	21,910	2	15	15	42	44	98	111	170	253	434	638
2001.....	22,305	1	1	11	27	61	87	129	169	301	455	711
2000.....	22,729	1	7	23	24	60	120	124	169	290	461	730
1999.....	22,802	2	16	12	25	51	93	109	180	324	465	696
All races, male												
2003.....	11,399	-	5	12	14	34	54	59	106	153	253	414
2002.....	11,538	1	6	10	30	31	62	71	107	161	278	404
2001.....	11,710	1	1	8	21	43	51	72	107	195	305	450
2000.....	11,812	-	5	16	12	46	80	64	108	192	290	466
1999.....	11,794	-	9	9	21	30	61	69	109	204	306	459
All races, female												
2003.....	10,076	-	-	5	14	19	33	37	50	80	144	235
2002.....	10,372	1	9	5	12	13	36	40	63	92	156	234
2001.....	10,595	-	-	3	6	18	36	57	62	106	150	261
2000.....	10,917	1	2	7	12	14	40	60	61	98	171	264
1999.....	11,008	2	7	3	4	21	32	40	71	120	159	237

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
White, both sexes												
2003.....	19,598	-	4	14	17	39	73	66	111	183	316	524
2002.....	19,996	1	14	15	30	28	78	84	119	194	346	532
2001.....	20,335	1	1	10	22	44	65	90	131	234	367	584
2000.....	20,892	-	4	18	22	46	85	89	129	226	384	603
1999.....	20,941	-	13	8	18	36	74	79	135	253	374	582
White, male												
2003.....	10,447	-	4	10	7	27	45	44	76	125	202	332
2002.....	10,535	-	6	10	22	23	52	58	73	123	220	349
2001.....	10,642	1	1	7	16	31	37	52	88	149	247	377
2000.....	10,838	-	4	12	12	38	58	50	81	150	251	388
1999.....	10,768	-	8	7	16	22	49	48	82	163	244	386
White, female												
2003.....	9,151	-	-	4	10	12	28	22	35	58	114	192
2002.....	9,461	1	8	5	8	5	26	26	46	71	126	183
2001.....	9,693	-	-	3	6	13	28	38	43	85	120	207
2000.....	10,054	-	-	6	10	8	27	39	48	76	133	215
1999.....	10,173	-	5	1	2	14	25	31	53	90	130	196

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
Black, both sexes												
2003.....	1,371	-	-	3	7	11	12	24	34	42	66	103
2002.....	1,474	1	1	-	10	11	14	22	41	40	78	90
2001.....	1,496	-	-	1	4	16	16	29	34	59	70	105
2000.....	1,369	1	2	4	1	8	27	30	26	56	65	107
1999.....	1,443	2	2	3	6	13	19	24	34	62	76	93
Black, male												
2003.....	687	-	-	2	6	4	8	10	24	22	41	70
2002.....	775	1	-	-	7	4	6	11	29	27	52	48
2001.....	804	-	-	1	4	11	11	15	18	42	46	58
2000.....	728	-	-	3	-	3	16	12	19	37	34	67
1999.....	786	-	1	2	4	6	12	17	22	36	51	59
Black, female												
2003.....	684	-	-	1	1	7	4	14	10	20	25	33
2002.....	699	-	1	-	3	7	8	11	12	13	26	42
2001.....	692	-	-	-	-	5	5	14	16	17	24	47
2000.....	641	1	2	1	1	5	11	18	7	19	31	40
1999.....	657	2	1	1	2	7	7	7	12	26	25	34

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
American Indian 1/, both sexes												
2003.....	69	-	1	-	-	-	-	1	1	1	2	7
2002.....	53	-	-	-	1	-	1	-	1	6	2	3
2001.....	93	-	-	-	1	1	-	3	1	3	7	3
2000.....	83	-	-	-	-	1	-	-	4	1	3	5
1999.....	57	-	-	-	1	-	-	1	2	1	5	2
American Indian 1/, male												
2003.....	34	-	1	-	-	-	-	1	1	-	2	3
2002.....	29	-	-	-	1	-	1	-	-	5	2	1
2001.....	48	-	-	-	1	1	-	2	-	-	5	3
2000.....	41	-	-	-	-	1	-	-	3	1	-	3
1999.....	31	-	-	-	1	-	-	1	-	1	3	1
American Indian 1/, female												
2003.....	35	-	-	-	-	-	-	-	-	1	-	4
2002.....	24	-	-	-	-	-	-	-	1	1	-	2
2001.....	45	-	-	-	-	-	-	1	1	3	2	-
2000.....	42	-	-	-	-	-	-	-	1	-	3	2
1999.....	26	-	-	-	-	-	-	-	2	-	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
Asian or Pacific Islander 2/, both sexes												
2003.....	437	-	-	-	4	3	2	5	10	7	13	15
2002.....	387	-	-	-	1	5	5	5	9	13	8	13
2001.....	381	-	-	-	-	-	6	7	3	5	11	19
2000.....	385	-	1	1	1	5	8	5	10	7	9	15
1999.....	361	-	1	1	-	2	-	5	9	8	10	19
Asian or Pacific Islander 2/, male												
2003.....	231	-	-	-	1	3	1	4	5	6	8	9
2002.....	199	-	-	-	-	4	3	2	5	6	4	6
2001.....	216	-	-	-	-	-	3	3	1	4	7	12
2000.....	205	-	1	1	-	4	6	2	5	4	5	8
1999.....	209	-	-	-	-	2	-	3	5	4	8	13
Asian or Pacific Islander 2/, female												
2003.....	206	-	-	-	3	-	1	1	5	1	5	6
2002.....	188	-	-	-	1	1	2	3	4	7	4	7
2001.....	165	-	-	-	-	-	3	4	2	1	4	7
2000.....	180	-	-	-	1	1	2	3	5	3	4	7
1999.....	152	-	1	1	-	-	-	2	4	4	2	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
All races, both sexes												
2003.....	21,535	30	133	156	186	183	219	205	228	321	475	595
2002.....	21,498	21	104	148	168	237	235	190	281	325	442	606
2001.....	21,451	25	155	153	170	225	254	230	283	346	433	630
2000.....	21,339	38	137	142	173	242	222	228	245	368	433	622
1999.....	21,014	27	119	130	175	221	261	234	266	374	425	626
All races, male												
2003.....	12,073	9	75	90	105	125	132	125	149	167	277	347
2002.....	12,016	11	57	75	97	142	140	117	163	178	251	335
2001.....	11,850	10	81	86	99	134	155	135	165	196	236	346
2000.....	11,777	21	69	81	109	150	143	131	152	185	238	356
1999.....	11,519	11	65	75	98	130	161	128	148	204	231	346
All races, female												
2003.....	9,462	21	58	66	81	58	87	80	79	154	198	248
2002.....	9,482	10	47	73	71	95	95	73	118	147	191	271
2001.....	9,601	15	74	67	71	91	99	95	118	150	197	284
2000.....	9,562	17	68	61	64	92	79	97	93	183	195	266
1999.....	9,495	16	54	55	77	91	100	106	118	170	194	280

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
White, both sexes												
2003.....	19,353	28	111	132	144	147	173	169	180	244	387	510
2002.....	19,273	16	84	119	134	193	201	149	221	262	382	481
2001.....	19,239	20	134	114	130	188	203	171	231	279	349	531
2000.....	19,191	36	115	116	143	186	175	172	183	300	348	500
1999.....	18,899	23	103	106	132	170	209	191	210	307	353	524
White, male												
2003.....	10,938	7	63	82	81	102	103	100	118	123	228	306
2002.....	10,835	10	47	57	77	119	120	94	131	142	220	263
2001.....	10,672	7	68	64	67	115	125	103	136	155	187	299
2000.....	10,617	19	58	64	89	113	112	102	114	154	199	292
1999.....	10,395	11	54	63	71	97	136	104	115	170	192	286
White, female												
2003.....	8,415	21	48	50	63	45	70	69	62	121	159	204
2002.....	8,438	6	37	62	57	74	81	55	90	120	162	218
2001.....	8,567	13	66	50	63	73	78	68	95	124	162	232
2000.....	8,574	17	57	52	54	73	63	70	69	146	149	208
1999.....	8,504	12	49	43	61	73	73	87	95	137	161	238

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
Black, both sexes												
2003.....	1,762	1	14	17	30	23	32	26	39	58	74	67
2002.....	1,770	4	15	19	23	30	24	28	40	43	48	93
2001.....	1,765	4	14	28	31	28	36	37	38	53	66	75
2000.....	1,742	-	18	16	20	46	36	39	49	49	68	107
1999.....	1,708	2	13	18	35	36	47	30	38	53	54	79
Black, male												
2003.....	918	1	8	7	16	15	20	17	24	33	42	33
2002.....	923	1	7	13	12	19	12	16	21	26	23	57
2001.....	928	3	9	13	26	16	19	21	23	32	36	32
2000.....	929	-	9	11	11	31	24	18	32	20	33	54
1999.....	892	-	8	9	24	24	22	16	22	25	29	47
Black, female												
2003.....	844	-	6	10	14	8	12	9	15	25	32	34
2002.....	847	3	8	6	11	11	12	12	19	17	25	36
2001.....	837	1	5	15	5	12	17	16	15	21	30	43
2000.....	813	-	9	5	9	15	12	21	17	29	35	53
1999.....	816	2	5	9	11	12	25	14	16	28	25	32

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
American Indian 1/, both sexes												
2003.....	77	-	2	2	1	7	5	1	2	4	3	2
2002.....	73	-	1	-	1	4	6	3	3	3	1	3
2001.....	77	-	1	-	3	2	7	8	1	2	3	2
2000.....	72	1	1	1	1	2	2	2	3	3	1	2
1999.....	56	1	1	-	1	1	2	3	1	1	-	3
American Indian 1/, male												
2003.....	38	-	2	-	-	4	5	1	2	2	1	2
2002.....	40	-	1	-	-	2	5	1	2	-	1	2
2001.....	42	-	1	-	1	1	5	6	1	-	3	2
2000.....	38	1	1	-	-	1	1	2	1	2	-	1
1999.....	38	-	1	-	1	-	1	2	1	-	-	3
American Indian 1/, female												
2003.....	39	-	-	2	1	3	-	-	-	2	2	-
2002.....	33	-	-	-	1	2	1	2	1	3	-	1
2001.....	35	-	-	-	2	1	2	2	-	2	-	-
2000.....	34	-	-	1	1	1	1	-	2	1	1	1
1999.....	18	1	-	-	-	1	1	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
Asian or Pacific Islander 2/, both sexes												
2003.....	343	1	6	5	11	6	9	9	7	15	11	16
2002.....	382	1	4	10	10	10	4	10	17	17	11	29
2001.....	370	1	6	11	6	7	8	14	13	12	15	22
2000.....	334	1	3	9	9	8	9	15	10	16	16	13
1999.....	351	1	2	6	7	14	3	10	17	13	18	20
Asian or Pacific Islander 2/, male												
2003.....	179	1	2	1	8	4	4	7	5	9	6	6
2002.....	218	-	2	5	8	2	3	6	9	10	7	13
2001.....	208	-	3	9	5	2	6	5	5	9	10	13
2000.....	193	1	1	6	9	5	6	9	5	9	6	9
1999.....	194	-	2	3	2	9	2	6	10	9	10	10
Asian or Pacific Islander 2/, female												
2003.....	164	-	4	4	3	2	5	2	2	6	5	10
2002.....	164	1	2	5	2	8	1	4	8	7	4	16
2001.....	162	1	3	2	1	5	2	9	8	3	5	9
2000.....	141	-	2	3	-	3	3	6	5	7	10	4
1999.....	157	1	-	3	5	5	1	4	7	4	8	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All races, both sexes												
2003.....	11,267	-	-	-	1	-	-	1	11	25	122	238
2002.....	11,392	2	-	-	-	2	-	3	9	36	102	270
2001.....	11,172	-	-	-	-	-	-	1	8	39	118	241
2000.....	11,063	-	-	-	1	-	-	-	7	38	108	245
1999.....	11,016	-	-	-	-	-	-	3	19	47	109	241
All races, male												
2003.....	5,848	-	-	-	-	-	-	1	6	17	80	139
2002.....	5,953	-	-	-	-	-	-	-	6	24	54	159
2001.....	5,724	-	-	-	-	-	-	-	4	22	75	151
2000.....	5,553	-	-	-	-	-	-	-	5	18	67	143
1999.....	5,584	-	-	-	-	-	-	2	14	33	62	132
All races, female												
2003.....	5,419	-	-	-	1	-	-	-	5	8	42	99
2002.....	5,439	2	-	-	-	2	-	3	3	12	48	111
2001.....	5,448	-	-	-	-	-	-	1	4	17	43	90
2000.....	5,510	-	-	-	1	-	-	-	2	20	41	102
1999.....	5,432	-	-	-	-	-	-	1	5	14	47	109

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
White, both sexes												
2003.....	9,227	-	-	-	1	-	-	1	8	17	86	158
2002.....	9,420	1	-	-	-	2	-	2	8	22	69	189
2001.....	9,168	-	-	-	-	-	-	1	4	30	70	171
2000.....	9,066	-	-	-	1	-	-	-	4	24	73	168
1999.....	9,010	-	-	-	-	-	-	2	11	35	68	182
White, male												
2003.....	4,892	-	-	-	-	-	-	1	5	13	60	96
2002.....	5,024	-	-	-	-	-	-	-	5	15	39	116
2001.....	4,765	-	-	-	-	-	-	-	3	15	52	110
2000.....	4,644	-	-	-	-	-	-	-	3	14	50	103
1999.....	4,651	-	-	-	-	-	-	2	7	26	38	102
White, female												
2003.....	4,335	-	-	-	1	-	-	-	3	4	26	62
2002.....	4,396	1	-	-	-	2	-	2	3	7	30	73
2001.....	4,403	-	-	-	-	-	-	1	1	15	18	61
2000.....	4,422	-	-	-	1	-	-	-	1	10	23	65
1999.....	4,359	-	-	-	-	-	-	-	4	9	30	80

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Black, both sexes

2003.....	1,847	-	-	-	-	-	-	-	2	8	35	73
2002.....	1,771	-	-	-	-	-	-	1	1	13	28	75
2001.....	1,817	-	-	-	-	-	-	-	4	9	44	65
2000.....	1,841	-	-	-	-	-	-	-	3	13	31	72
1999.....	1,803	-	-	-	-	-	-	1	7	10	40	56

Black, male

2003.....	863	-	-	-	-	-	-	-	1	4	20	39
2002.....	840	-	-	-	-	-	-	-	1	8	14	42
2001.....	864	-	-	-	-	-	-	-	1	7	21	39
2000.....	837	-	-	-	-	-	-	-	2	3	14	36
1999.....	834	-	-	-	-	-	-	-	7	6	24	28

Black, female

2003.....	984	-	-	-	-	-	-	-	1	4	15	34
2002.....	931	-	-	-	-	-	-	1	-	5	14	33
2001.....	953	-	-	-	-	-	-	-	3	2	23	26
2000.....	1,004	-	-	-	-	-	-	-	1	10	17	36
1999.....	969	-	-	-	-	-	-	1	-	4	16	28

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

American Indian 1/, both sexes

2003.....	55	-	-	-	-	-	-	-	-	-	1	-
2002.....	43	-	-	-	-	-	-	-	-	1	1	1
2001.....	51	-	-	-	-	-	-	-	-	-	2	2
2000.....	38	-	-	-	-	-	-	-	-	-	1	2
1999.....	43	-	-	-	-	-	-	-	-	1	-	2

American Indian 1/, male

2003.....	23	-	-	-	-	-	-	-	-	-	-	-
2002.....	20	-	-	-	-	-	-	-	-	1	1	-
2001.....	23	-	-	-	-	-	-	-	-	-	1	-
2000.....	17	-	-	-	-	-	-	-	-	-	1	2
1999.....	22	-	-	-	-	-	-	-	-	-	-	1

American Indian 1/, female

2003.....	32	-	-	-	-	-	-	-	-	-	1	-
2002.....	23	-	-	-	-	-	-	-	-	-	-	1
2001.....	28	-	-	-	-	-	-	-	-	-	1	2
2000.....	21	-	-	-	-	-	-	-	-	-	-	-
1999.....	21	-	-	-	-	-	-	-	-	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Asian or Pacific Islander 2/, both sexes

2003.....	138	-	-	-	-	-	-	-	1	-	-	7
2002.....	158	1	-	-	-	-	-	-	-	-	4	5
2001.....	136	-	-	-	-	-	-	-	-	-	2	3
2000.....	118	-	-	-	-	-	-	-	-	1	3	3
1999.....	160	-	-	-	-	-	-	-	1	1	1	1

Asian or Pacific Islander 2/, male

2003.....	70	-	-	-	-	-	-	-	-	-	-	4
2002.....	69	-	-	-	-	-	-	-	-	-	-	1
2001.....	72	-	-	-	-	-	-	-	-	-	1	2
2000.....	55	-	-	-	-	-	-	-	-	1	2	2
1999.....	77	-	-	-	-	-	-	-	-	1	-	1

Asian or Pacific Islander 2/, female

2003.....	68	-	-	-	-	-	-	-	1	-	-	3
2002.....	89	1	-	-	-	-	-	-	-	-	4	4
2001.....	64	-	-	-	-	-	-	-	-	-	1	1
2000.....	63	-	-	-	-	-	-	-	-	-	1	1
1999.....	83	-	-	-	-	-	-	-	1	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
All races, both sexes												
2003.....	55	1	-	-	1	-	-	-	1	-	1	4
2002.....	73	3	-	-	-	-	1	-	-	-	-	2
2001.....	90	1	1	-	-	-	-	-	1	1	3	1
2000.....	100	-	2	-	1	-	1	-	-	2	2	5
1999.....	83	1	1	-	-	-	-	-	1	2	2	3
All races, male												
2003.....	31	1	-	-	-	-	-	-	1	-	1	3
2002.....	44	3	-	-	-	-	1	-	-	-	-	2
2001.....	51	1	-	-	-	-	-	-	1	1	2	1
2000.....	60	-	2	-	1	-	1	-	-	1	2	4
1999.....	39	1	-	-	-	-	-	-	1	-	1	1
All races, female												
2003.....	24	-	-	-	1	-	-	-	-	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	-	-	-
2001.....	39	-	1	-	-	-	-	-	-	-	1	-
2000.....	40	-	-	-	-	-	-	-	-	1	-	1
1999.....	44	-	1	-	-	-	-	-	-	2	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

White, both sexes

2003.....	50	1	-	-	1	-	-	-	1	-	1	2
2002.....	61	1	-	-	-	-	1	-	-	-	-	2
2001.....	77	1	1	-	-	-	-	-	-	-	3	1
2000.....	89	-	2	-	1	-	-	-	-	1	2	3
1999.....	76	1	1	-	-	-	-	-	1	2	1	2

White, male

2003.....	28	1	-	-	-	-	-	-	1	-	1	1
2002.....	38	1	-	-	-	-	1	-	-	-	-	2
2001.....	44	1	-	-	-	-	-	-	-	-	2	1
2000.....	51	-	2	-	1	-	-	-	-	-	2	2
1999.....	36	1	-	-	-	-	-	-	1	-	1	1

White, female

2003.....	22	-	-	-	1	-	-	-	-	-	-	1
2002.....	23	-	-	-	-	-	-	-	-	-	-	-
2001.....	33	-	1	-	-	-	-	-	-	-	1	-
2000.....	38	-	-	-	-	-	-	-	-	1	-	1
1999.....	40	-	1	-	-	-	-	-	-	2	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
All races, both sexes												
2003.....	62,639	19	106	127	147	242	274	298	392	664	1,221	2,234
2002.....	62,969	24	124	133	155	233	294	289	445	716	1,327	2,213
2001.....	62,708	17	117	116	163	242	274	316	412	718	1,337	2,187
2000.....	63,060	24	129	123	137	243	268	292	409	771	1,319	2,114
1999.....	66,251	14	116	150	141	246	273	307	455	833	1,481	2,247
All races, male												
2003.....	31,845	12	66	80	72	161	168	181	246	377	669	1,274
2002.....	32,122	15	65	77	75	140	200	194	265	407	703	1,225
2001.....	31,663	7	73	64	89	152	194	190	252	385	723	1,231
2000.....	31,728	13	75	64	76	142	177	189	243	424	717	1,186
1999.....	33,449	8	64	80	80	158	171	204	286	464	836	1,274
All races, female												
2003.....	30,794	7	40	47	75	81	106	117	146	287	552	960
2002.....	30,847	9	59	56	80	93	94	95	180	309	624	988
2001.....	31,045	10	44	52	74	90	80	126	160	333	614	956
2000.....	31,332	11	54	59	61	101	91	103	166	347	602	928
1999.....	32,802	6	52	70	61	88	102	103	169	369	645	973

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2003.....	54,660	15	78	106	110	201	221	236	296	536	940	1,741
2002.....	54,810	22	90	101	124	191	232	226	348	565	1,043	1,739
2001.....	54,510	15	98	91	125	196	226	255	317	562	1,054	1,705
2000.....	55,066	19	97	105	107	187	223	251	330	595	1,001	1,652
1999.....	57,818	12	98	118	105	196	212	255	360	657	1,147	1,773

White, male

2003.....	27,830	10	46	68	53	133	135	154	190	317	512	988
2002.....	28,060	15	49	58	59	113	156	154	211	323	539	972
2001.....	27,506	6	59	50	67	120	163	153	198	308	579	970
2000.....	27,697	11	62	55	59	110	144	165	206	336	539	928
1999.....	29,229	7	55	65	62	120	132	176	225	375	659	992

White, female

2003.....	26,830	5	32	38	57	68	86	82	106	219	428	753
2002.....	26,750	7	41	43	65	78	76	72	137	242	504	767
2001.....	27,004	9	39	41	58	76	63	102	119	254	475	735
2000.....	27,369	8	35	50	48	77	79	86	124	259	462	724
1999.....	28,589	5	43	53	43	76	80	79	135	282	488	781

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2003.....	6,618	2	24	20	31	30	41	50	74	106	227	421
2002.....	6,824	2	27	27	26	34	48	51	74	131	232	402
2001.....	6,799	1	16	19	28	38	39	52	73	126	221	412
2000.....	6,696	4	25	17	24	48	37	34	54	154	280	397
1999.....	7,067	1	14	22	28	40	44	41	67	146	280	418

Black, male

2003.....	3,324	2	16	11	16	20	26	22	40	48	128	243
2002.....	3,381	-	12	16	13	23	33	31	36	70	131	218
2001.....	3,436	-	13	12	15	24	28	31	39	61	110	217
2000.....	3,378	1	11	9	15	27	29	20	26	75	156	228
1999.....	3,547	-	8	9	13	30	27	20	41	71	151	246

Black, female

2003.....	3,294	-	8	9	15	10	15	28	34	58	99	178
2002.....	3,443	2	15	11	13	11	15	20	38	61	101	184
2001.....	3,363	1	3	7	13	14	11	21	34	65	111	195
2000.....	3,318	3	14	8	9	21	8	14	28	79	124	169
1999.....	3,520	1	6	13	15	10	17	21	26	75	129	172

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

American Indian 1/, both sexes

2003.....	275	-	1	-	-	1	2	2	3	6	17	10
2002.....	265	-	-	2	1	3	1	2	5	5	11	11
2001.....	289	-	1	-	1	4	4	2	4	4	18	12
2000.....	250	-	-	-	-	1	3	3	2	4	4	14
1999.....	300	-	-	1	2	1	3	4	4	7	13	18

American Indian 1/, male

2003.....	147	-	1	-	-	1	2	1	2	3	9	7
2002.....	130	-	-	1	1	1	-	2	4	1	7	4
2001.....	150	-	-	-	1	4	2	1	3	3	9	9
2000.....	118	-	-	-	-	1	1	2	1	2	2	8
1999.....	140	-	-	1	2	-	2	2	3	5	5	12

American Indian 1/, female

2003.....	128	-	-	-	-	-	-	1	1	3	8	3
2002.....	135	-	-	1	-	2	1	-	1	4	4	7
2001.....	139	-	1	-	-	-	2	1	1	1	9	3
2000.....	132	-	-	-	-	-	2	1	1	2	2	6
1999.....	160	-	-	-	-	1	1	2	1	2	8	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Asian or Pacific Islander 2/, both sexes

2003.....	1,086	2	3	1	6	10	10	10	19	16	37	62
2002.....	1,070	-	7	3	4	5	13	10	18	15	41	61
2001.....	1,110	1	2	6	9	4	5	7	18	26	44	58
2000.....	1,048	1	7	1	6	7	5	4	23	18	34	51
1999.....	1,066	1	4	9	6	9	14	7	24	23	41	38

Asian or Pacific Islander 2/, male

2003.....	544	-	3	1	3	7	5	4	14	9	20	36
2002.....	551	-	4	2	2	3	11	7	14	13	26	31
2001.....	571	1	1	2	6	4	1	5	12	13	25	35
2000.....	535	1	2	-	2	4	3	2	10	11	20	22
1999.....	533	1	1	5	3	8	10	6	17	13	21	24

Asian or Pacific Islander 2/, female

2003.....	542	2	-	-	3	3	5	6	5	7	17	26
2002.....	519	-	3	1	2	2	2	3	4	2	15	30
2001.....	539	-	1	4	3	-	4	2	6	13	19	23
2000.....	513	-	5	1	4	3	2	2	13	7	14	29
1999.....	533	-	3	4	3	1	4	1	7	10	20	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races, both sexes

2003.....	13,563	62	51	38	41	41	55	69	109	142	197	295
2002.....	13,299	69	60	44	45	41	46	62	101	137	229	327
2001.....	13,676	51	58	52	53	47	55	60	91	156	210	304
2000.....	13,546	60	53	62	37	40	55	68	97	155	213	293
1999.....	13,227	58	63	64	37	52	46	55	116	140	202	276

All races, male

2003.....	6,687	32	31	21	22	24	30	40	58	79	116	148
2002.....	6,536	30	30	22	27	21	29	33	51	77	119	169
2001.....	6,735	26	30	26	26	20	33	33	44	79	114	173
2000.....	6,559	32	33	32	17	27	35	30	52	86	102	160
1999.....	6,520	30	33	30	19	26	28	32	64	79	118	142

All races, female

2003.....	6,876	30	20	17	19	17	25	29	51	63	81	147
2002.....	6,763	39	30	22	18	20	17	29	50	60	110	158
2001.....	6,941	25	28	26	27	27	22	27	47	77	96	131
2000.....	6,987	28	20	30	20	13	20	38	45	69	111	133
1999.....	6,707	28	30	34	18	26	18	23	52	61	84	134

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2003.....	12,181	48	39	26	29	31	47	54	90	101	161	231
2002.....	11,913	59	43	36	31	34	34	42	79	112	174	274
2001.....	12,261	37	48	38	39	33	41	53	73	124	166	257
2000.....	12,171	48	44	46	30	29	44	47	82	124	168	225
1999.....	11,849	41	48	52	32	37	33	39	85	107	162	214

White, male

2003.....	6,036	25	22	13	17	19	27	31	46	62	96	120
2002.....	5,902	27	21	16	21	18	20	24	46	65	100	146
2001.....	6,089	20	24	19	18	12	25	29	35	64	95	149
2000.....	5,900	25	25	24	15	18	30	24	46	68	79	127
1999.....	5,891	21	22	24	16	17	21	21	46	63	101	116

White, female

2003.....	6,145	23	17	13	12	12	20	23	44	39	65	111
2002.....	6,011	32	22	20	10	16	14	18	33	47	74	128
2001.....	6,172	17	24	19	21	21	16	24	38	60	71	108
2000.....	6,271	23	19	22	15	11	14	23	36	56	89	98
1999.....	5,958	20	26	28	16	20	12	18	39	44	61	98

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes

2003.....	1,123	11	9	10	8	8	8	14	15	37	30	59
2002.....	1,143	9	14	7	11	6	9	13	18	20	50	50
2001.....	1,157	11	5	13	13	12	9	6	12	26	35	35
2000.....	1,136	9	9	13	6	8	8	18	15	24	39	56
1999.....	1,173	15	13	12	5	13	13	9	21	30	36	50

Black, male

2003.....	525	5	8	6	3	4	3	9	9	15	16	26
2002.....	500	2	8	6	3	3	6	5	3	8	18	20
2001.....	525	5	3	6	7	7	6	3	8	13	17	16
2000.....	520	4	8	6	2	7	2	5	6	13	19	24
1999.....	521	8	10	6	3	8	7	8	12	15	15	19

Black, female

2003.....	598	6	1	4	5	4	5	5	6	22	14	33
2002.....	643	7	6	1	8	3	3	8	15	12	32	30
2001.....	632	6	2	7	6	5	3	3	4	13	18	19
2000.....	616	5	1	7	4	1	6	13	9	11	20	32
1999.....	652	7	3	6	2	5	6	1	9	15	21	31

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Asian or Pacific Islander 2/, both sexes

2003.....	220	3	3	2	3	2	-	1	2	3	5	4
2002.....	209	1	1	1	2	1	3	6	4	5	2	3
2001.....	200	2	5	-	1	-	3	1	5	4	5	10
2000.....	194	3	-	2	1	1	1	1	-	4	5	11
1999.....	163	2	2	-	-	1	-	7	9	3	4	11

Asian or Pacific Islander 2/, male

2003.....	106	2	1	2	1	1	-	-	2	2	4	1
2002.....	119	1	1	-	2	-	3	4	2	4	1	3
2001.....	92	-	3	-	1	-	1	1	1	1	1	6
2000.....	119	3	-	1	-	1	1	-	-	4	4	9
1999.....	86	1	1	-	-	-	-	3	5	1	2	6

Asian or Pacific Islander 2/, female

2003.....	114	1	2	-	2	1	-	1	-	1	1	3
2002.....	90	-	-	1	-	1	-	2	2	1	1	-
2001.....	108	2	2	-	-	-	2	-	4	3	4	4
2000.....	75	-	-	1	1	-	-	1	-	-	1	2
1999.....	77	1	1	-	-	1	-	4	4	2	2	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
All races, both sexes												
2003.....	4,594	22	24	16	31	37	68	69	88	81	102	137
2002.....	4,614	18	21	29	24	42	69	83	97	99	97	131
2001.....	4,627	16	22	25	30	48	61	65	95	83	132	112
2000.....	4,494	10	35	24	28	27	58	62	91	85	105	110
1999.....	4,503	12	30	24	29	43	78	64	96	90	104	134
All races, male												
2003.....	1,832	11	14	8	15	19	33	37	46	37	54	77
2002.....	1,781	12	9	13	13	23	44	53	48	53	43	57
2001.....	1,774	11	12	12	15	26	33	40	55	42	49	57
2000.....	1,669	6	16	17	12	9	30	29	41	34	44	52
1999.....	1,746	8	17	8	15	24	43	33	48	47	54	68
All races, female												
2003.....	2,762	11	10	8	16	18	35	32	42	44	48	60
2002.....	2,833	6	12	16	11	19	25	30	49	46	54	74
2001.....	2,853	5	10	13	15	22	28	25	40	41	83	55
2000.....	2,825	4	19	7	16	18	28	33	50	51	61	58
1999.....	2,757	4	13	16	14	19	35	31	48	43	50	66

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
White, both sexes												
2003.....	3,476	9	10	4	17	15	8	16	23	17	28	46
2002.....	3,460	8	12	15	8	13	17	15	28	22	23	49
2001.....	3,543	6	11	11	16	17	16	15	24	16	37	31
2000.....	3,450	4	15	9	12	10	11	15	14	17	26	38
1999.....	3,393	6	9	7	13	18	16	6	11	22	26	55
White, male												
2003.....	1,319	4	5	3	8	6	5	8	14	8	14	29
2002.....	1,267	4	5	6	5	7	10	10	14	14	12	25
2001.....	1,303	5	5	6	10	8	10	9	11	9	12	12
2000.....	1,235	2	8	7	5	2	6	9	6	7	14	16
1999.....	1,244	4	6	1	6	9	5	1	5	10	14	31
White, female												
2003.....	2,157	5	5	1	9	9	3	8	9	9	14	17
2002.....	2,193	4	7	9	3	6	7	5	14	8	11	24
2001.....	2,240	1	6	5	6	9	6	6	13	7	25	19
2000.....	2,215	2	7	2	7	8	5	6	8	10	12	22
1999.....	2,149	2	3	6	7	9	11	5	6	12	12	24

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
Black, both sexes												
2003.....	1,037	6	12	12	13	19	60	52	62	62	72	87
2002.....	1,064	6	9	12	14	27	52	65	65	74	71	80
2001.....	1,001	6	11	13	13	28	45	50	69	62	92	79
2000.....	967	5	19	15	16	16	46	47	77	65	76	69
1999.....	1,019	3	21	17	16	23	61	56	84	66	76	74
Black, male												
2003.....	487	4	7	5	7	12	28	28	30	29	40	48
2002.....	479	5	4	7	7	16	34	40	33	37	31	31
2001.....	446	4	7	5	5	18	23	31	43	30	36	45
2000.....	395	4	8	10	7	7	23	20	35	26	29	35
1999.....	457	2	11	7	9	13	37	30	43	35	38	35
Black, female												
2003.....	550	2	5	7	6	7	32	24	32	33	32	39
2002.....	585	1	5	5	7	11	18	25	32	37	40	49
2001.....	555	2	4	8	8	10	22	19	26	32	56	34
2000.....	572	1	11	5	9	9	23	27	42	39	47	34
1999.....	562	1	10	10	7	10	24	26	41	31	38	39

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Anemias (D50-D64)

American Indian 1/, both sexes

2003.....	15	-	-	-	-	-	-	-	1	1	1	1
2002.....	15	-	-	-	1	1	-	1	1	-	1	-
2001.....	27	-	-	-	1	2	-	-	-	1	3	1
2000.....	20	-	-	-	-	-	-	-	-	-	2	1
1999.....	21	-	-	-	-	1	1	-	1	2	-	2

American Indian 1/, male

2003.....	4	-	-	-	-	-	-	-	1	-	-	-
2002.....	4	-	-	-	-	-	-	1	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	-	1	1	-
2000.....	9	-	-	-	-	-	-	-	-	-	1	-
1999.....	9	-	-	-	-	1	1	-	-	2	-	1

American Indian 1/, female

2003.....	11	-	-	-	-	-	-	-	-	1	1	1
2002.....	11	-	-	-	1	1	-	-	1	-	1	-
2001.....	20	-	-	-	1	2	-	-	-	-	2	1
2000.....	11	-	-	-	-	-	-	-	-	-	1	1
1999.....	12	-	-	-	-	-	-	-	1	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

All races, both sexes

2003.....	74,219	3	8	10	21	46	114	227	430	715	1,334	2,245
2002.....	73,249	1	3	5	29	55	116	229	413	776	1,388	2,106
2001.....	71,372	2	6	7	32	48	103	215	380	735	1,223	2,063
2000.....	69,301	4	6	8	18	48	114	206	417	684	1,242	1,917
1999.....	68,399	2	8	9	23	51	104	189	393	730	1,212	1,942

All races, male

2003.....	35,438	-	6	6	10	28	68	119	260	420	844	1,354
2002.....	34,301	-	1	4	18	30	68	128	257	452	878	1,248
2001.....	32,841	1	2	2	21	27	58	114	227	436	745	1,234
2000.....	31,602	4	2	5	8	31	63	118	258	396	755	1,124
1999.....	31,150	-	4	4	12	25	55	94	225	437	702	1,189

All races, female

2003.....	38,781	3	2	4	11	18	46	108	170	295	490	891
2002.....	38,948	1	2	1	11	25	48	101	156	324	510	858
2001.....	38,531	1	4	5	11	21	45	101	153	299	478	829
2000.....	37,699	-	4	3	10	17	51	88	159	288	487	793
1999.....	37,249	2	4	5	11	26	49	95	168	293	510	753

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

White, both sexes

2003.....	59,099	2	6	9	13	29	82	135	296	487	952	1,583
2002.....	58,459	1	3	4	16	36	73	152	287	549	1,005	1,489
2001.....	57,180	1	6	3	18	27	55	124	259	511	888	1,474
2000.....	55,561	3	6	5	12	33	72	128	292	493	926	1,344
1999.....	54,599	2	6	8	14	38	69	137	272	515	869	1,367

White, male

2003.....	28,939	-	5	5	8	18	49	63	183	295	622	994
2002.....	28,110	-	1	4	13	18	42	88	178	327	651	906
2001.....	26,917	-	2	1	14	18	32	63	149	302	574	898
2000.....	26,009	3	2	3	6	22	42	76	191	285	581	808
1999.....	25,545	-	3	4	6	21	38	72	154	318	511	864

White, female

2003.....	30,160	2	1	4	5	11	33	72	113	192	330	589
2002.....	30,349	1	2	-	3	18	31	64	109	222	354	583
2001.....	30,263	1	4	2	4	9	23	61	110	209	314	576
2000.....	29,552	-	4	2	6	11	30	52	101	208	345	536
1999.....	29,054	2	3	4	8	17	31	65	118	197	358	503

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diabetes mellitus (E10-E14)												
Black, both sexes												
2003.....	12,892	1	2	1	6	16	30	85	123	208	341	583
2002.....	12,687	-	-	1	13	17	42	74	121	212	348	545
2001.....	12,305	1	-	4	11	18	42	86	116	203	308	517
2000.....	12,021	1	-	3	6	14	41	75	114	177	285	517
1999.....	11,927	-	2	1	9	13	31	49	113	195	312	515
Black, male												
2003.....	5,425	-	1	1	2	10	17	53	69	113	197	312
2002.....	5,207	-	-	-	5	10	26	38	75	116	203	301
2001.....	5,049	1	-	1	5	8	22	49	74	120	158	292
2000.....	4,771	1	-	2	2	8	20	40	59	99	155	280
1999.....	4,759	-	1	-	6	4	15	21	65	107	170	292
Black, female												
2003.....	7,467	1	1	-	4	6	13	32	54	95	144	271
2002.....	7,480	-	-	1	8	7	16	36	46	96	145	244
2001.....	7,256	-	-	3	6	10	20	37	42	83	150	225
2000.....	7,250	-	-	1	4	6	21	35	55	78	130	237
1999.....	7,168	-	1	1	3	9	16	28	48	88	142	223

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

American Indian 1/, both sexes

2003.....	783	-	-	-	-	-	-	5	7	18	24	46
2002.....	744	-	-	-	-	1	1	2	1	10	23	43
2001.....	644	-	-	-	1	2	4	2	2	7	11	40
2000.....	616	-	-	-	-	1	1	-	7	7	16	25
1999.....	725	-	-	-	-	-	2	2	6	13	16	35

American Indian 1/, male

2003.....	367	-	-	-	-	-	-	3	5	10	16	25
2002.....	336	-	-	-	-	1	-	2	1	6	15	26
2001.....	276	-	-	-	1	1	2	1	1	5	2	21
2000.....	275	-	-	-	-	1	1	-	6	6	10	18
1999.....	323	-	-	-	-	-	2	-	5	8	12	21

American Indian 1/, female

2003.....	416	-	-	-	-	-	-	2	2	8	8	21
2002.....	408	-	-	-	-	-	1	-	-	4	8	17
2001.....	368	-	-	-	-	1	2	1	1	2	9	19
2000.....	341	-	-	-	-	-	-	-	1	1	6	7
1999.....	402	-	-	-	-	-	-	2	1	5	4	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

Asian or Pacific Islander 2/, both sexes

2003.....	1,445	-	-	-	2	1	2	2	4	2	17	33
2002.....	1,359	-	-	-	-	1	-	1	4	5	12	29
2001.....	1,243	-	-	-	2	1	2	3	3	14	16	32
2000.....	1,103	-	-	-	-	-	-	3	4	7	15	31
1999.....	1,148	-	-	-	-	-	2	1	2	7	15	25

Asian or Pacific Islander 2/, male

2003.....	707	-	-	-	-	-	2	-	3	2	9	23
2002.....	648	-	-	-	-	1	-	-	3	3	9	15
2001.....	599	-	-	-	1	-	2	1	3	9	11	23
2000.....	547	-	-	-	-	-	-	2	2	6	9	18
1999.....	523	-	-	-	-	-	-	1	1	4	9	12

Asian or Pacific Islander 2/, female

2003.....	738	-	-	-	2	1	-	2	1	-	8	10
2002.....	711	-	-	-	-	-	-	1	1	2	3	14
2001.....	644	-	-	-	1	1	-	2	-	5	5	9
2000.....	556	-	-	-	-	-	-	1	2	1	6	13
1999.....	625	-	-	-	-	-	2	-	1	3	6	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nutritional deficiencies (E40-E64)												
All races, both sexes												
2003.....	3,338	10	2	2	-	3	4	5	5	14	21	39
2002.....	3,779	4	1	-	1	1	6	5	11	27	33	39
2001.....	3,704	4	4	-	2	1	9	5	9	16	28	48
2000.....	4,242	14	2	1	4	4	6	9	9	30	26	46
1999.....	4,289	9	2	1	-	2	6	4	8	20	44	45
All races, male												
2003.....	1,176	7	2	2	-	1	3	4	3	7	12	19
2002.....	1,321	3	1	-	-	-	4	3	4	12	19	20
2001.....	1,233	4	1	-	2	-	7	2	2	8	13	27
2000.....	1,412	10	2	-	2	1	1	5	4	9	7	25
1999.....	1,486	4	2	1	-	2	3	1	3	8	20	22
All races, female												
2003.....	2,162	3	-	-	-	2	1	1	2	7	9	20
2002.....	2,458	1	-	-	1	1	2	2	7	15	14	19
2001.....	2,471	-	3	-	-	1	2	3	7	8	15	21
2000.....	2,830	4	-	1	2	3	5	4	5	21	19	21
1999.....	2,803	5	-	-	-	-	3	3	5	12	24	23

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nutritional deficiencies (E40-E64)												
White, both sexes												
2003.....	2,826	6	2	2	-	1	4	4	4	11	18	27
2002.....	3,213	3	1	-	-	1	5	5	8	15	26	32
2001.....	3,152	3	4	-	1	1	7	5	5	11	18	34
2000.....	3,627	8	1	-	2	4	4	5	4	21	19	37
1999.....	3,628	5	1	-	-	2	3	3	6	15	34	28
White, male												
2003.....	969	5	2	2	-	-	3	4	2	6	10	12
2002.....	1,102	2	1	-	-	-	3	3	3	7	15	14
2001.....	1,000	3	1	-	1	-	7	2	1	5	7	14
2000.....	1,175	6	1	-	2	1	1	3	1	7	4	20
1999.....	1,199	2	1	-	-	2	1	-	3	5	15	16
White, female												
2003.....	1,857	1	-	-	-	1	1	-	2	5	8	15
2002.....	2,111	1	-	-	-	1	2	2	5	8	11	18
2001.....	2,152	-	3	-	-	1	-	3	4	6	11	20
2000.....	2,452	2	-	-	-	3	3	2	3	14	15	17
1999.....	2,429	3	-	-	-	-	2	3	3	10	19	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nutritional deficiencies (E40-E64)

Black, both sexes

2003.....	455	3	-	-	-	1	-	1	1	1	3	12
2002.....	502	-	-	-	1	-	1	-	3	9	7	7
2001.....	475	1	-	-	1	-	2	-	4	5	10	12
2000.....	540	5	1	1	2	-	2	4	4	9	7	8
1999.....	589	4	1	1	-	-	3	1	2	5	10	15

Black, male

2003.....	185	1	-	-	-	-	-	-	1	-	2	7
2002.....	188	-	-	-	-	-	1	-	1	3	4	6
2001.....	204	1	-	-	1	-	-	-	1	3	6	11
2000.....	215	3	1	-	-	-	-	2	3	2	3	5
1999.....	250	2	1	1	-	-	2	1	-	3	5	5

Black, female

2003.....	270	2	-	-	-	1	-	1	-	1	1	5
2002.....	314	-	-	-	1	-	-	-	2	6	3	1
2001.....	271	-	-	-	-	-	2	-	3	2	4	1
2000.....	325	2	-	1	2	-	2	2	1	7	4	3
1999.....	339	2	-	-	-	-	1	-	2	2	5	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malnutrition (E40-E46)												
All races, both sexes												
2003.....	3,153	5	1	1	-	2	4	4	4	12	21	36
2002.....	3,510	2	1	-	1	1	5	4	9	25	30	36
2001.....	3,454	2	-	-	2	1	8	4	9	16	23	42
2000.....	3,926	11	1	1	2	3	5	9	7	29	24	44
1999.....	3,985	7	2	1	-	2	5	4	7	20	41	42
All races, male												
2003.....	1,098	4	1	1	-	-	3	3	3	7	12	19
2002.....	1,233	1	1	-	-	-	4	2	3	11	17	18
2001.....	1,148	2	-	-	2	-	6	2	2	8	11	22
2000.....	1,302	8	1	-	1	1	1	5	3	9	6	23
1999.....	1,371	4	2	1	-	2	2	1	3	8	19	21
All races, female												
2003.....	2,055	1	-	-	-	2	1	1	1	5	9	17
2002.....	2,277	1	-	-	1	1	1	2	6	14	13	18
2001.....	2,306	-	-	-	-	1	2	2	7	8	12	20
2000.....	2,624	3	-	1	1	2	4	4	4	20	18	21
1999.....	2,614	3	-	-	-	-	3	3	4	12	22	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malnutrition (E40-E46)												
White, both sexes												
2003.....	2,657	2	1	1	-	1	4	3	3	10	18	25
2002.....	2,979	2	1	-	-	1	4	4	6	15	23	29
2001.....	2,939	1	-	-	1	1	6	4	5	11	14	30
2000.....	3,352	7	-	-	1	3	4	5	3	20	17	35
1999.....	3,368	4	1	-	-	2	3	3	5	15	32	27
White, male												
2003.....	899	2	1	1	-	-	3	3	2	6	10	12
2002.....	1,029	1	1	-	-	-	3	2	2	7	13	12
2001.....	932	1	-	-	1	-	6	2	1	5	5	11
2000.....	1,079	5	-	-	1	1	1	3	1	7	3	18
1999.....	1,100	2	1	-	-	2	1	-	3	5	14	15
White, female												
2003.....	1,758	-	-	-	-	1	1	-	1	4	8	13
2002.....	1,950	1	-	-	-	1	1	2	4	8	10	17
2001.....	2,007	-	-	-	-	1	-	2	4	6	9	19
2000.....	2,273	2	-	-	-	2	3	2	2	13	14	17
1999.....	2,268	2	-	-	-	-	2	3	2	10	18	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malnutrition (E40-E46)												
Black, both sexes												
2003.....	442	2	-	-	-	1	-	1	1	1	3	11
2002.....	472	-	-	-	1	-	1	-	3	7	7	7
2001.....	441	1	-	-	1	-	2	-	4	5	9	10
2000.....	508	3	1	1	1	-	1	4	3	9	7	8
1999.....	547	3	1	1	-	-	2	1	2	5	9	13
Black, male												
2003.....	179	1	-	-	-	-	-	-	1	-	2	7
2002.....	175	-	-	-	-	-	1	-	1	2	4	6
2001.....	188	1	-	-	1	-	-	-	1	3	6	9
2000.....	203	2	1	-	-	-	-	2	2	2	3	5
1999.....	234	2	1	1	-	-	1	1	-	3	5	5
Black, female												
2003.....	263	1	-	-	-	1	-	1	-	1	1	4
2002.....	297	-	-	-	1	-	-	-	2	5	3	1
2001.....	253	-	-	-	-	-	2	-	3	2	3	1
2000.....	305	1	-	1	1	-	1	2	1	7	4	3
1999.....	313	1	-	-	-	-	1	-	2	2	4	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other nutritional deficiencies (E50-E64)

All races, both sexes

2003.....	185	5	1	1	-	1	-	1	1	2	-	3
2002.....	269	2	-	-	-	-	1	1	2	2	3	3
2001.....	250	2	4	-	-	-	1	1	-	-	5	6
2000.....	316	3	1	-	2	1	1	-	2	1	2	2
1999.....	304	2	-	-	-	-	1	-	1	-	3	3

All races, male

2003.....	78	3	1	1	-	1	-	1	-	-	-	-
2002.....	88	2	-	-	-	-	-	1	1	1	2	2
2001.....	85	2	1	-	-	-	1	-	-	-	2	5
2000.....	110	2	1	-	1	-	-	-	1	-	1	2
1999.....	115	-	-	-	-	-	1	-	-	-	1	1

All races, female

2003.....	107	2	-	-	-	-	-	-	1	2	-	3
2002.....	181	-	-	-	-	-	1	-	1	1	1	1
2001.....	165	-	3	-	-	-	-	1	-	-	3	1
2000.....	206	1	-	-	1	1	1	-	1	1	1	-
1999.....	189	2	-	-	-	-	-	-	1	-	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other nutritional deficiencies (E50-E64)

White, both sexes

2003.....	169	4	1	1	-	-	-	1	1	1	-	2
2002.....	234	1	-	-	-	-	1	1	2	-	3	3
2001.....	213	2	4	-	-	-	1	1	-	-	4	4
2000.....	275	1	1	-	1	1	-	-	1	1	2	2
1999.....	260	1	-	-	-	-	-	-	1	-	2	1

White, male

2003.....	70	3	1	1	-	-	-	1	-	-	-	-
2002.....	73	1	-	-	-	-	-	1	1	-	2	2
2001.....	68	2	1	-	-	-	1	-	-	-	2	3
2000.....	96	1	1	-	1	-	-	-	-	-	1	2
1999.....	99	-	-	-	-	-	-	-	-	-	1	1

White, female

2003.....	99	1	-	-	-	-	-	-	1	1	-	2
2002.....	161	-	-	-	-	-	1	-	1	-	1	1
2001.....	145	-	3	-	-	-	-	1	-	-	2	1
2000.....	179	-	-	-	-	1	-	-	1	1	1	-
1999.....	161	1	-	-	-	-	-	-	1	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningitis (G00,G03)												
All races, both sexes												
2003.....	730	77	24	10	7	14	23	18	26	31	36	59
2002.....	700	74	19	10	21	12	14	16	20	23	38	63
2001.....	727	78	26	15	9	19	21	13	28	31	38	55
2000.....	758	91	36	12	18	7	8	16	27	35	37	45
1999.....	850	116	45	14	13	19	22	21	27	33	43	62
All races, male												
2003.....	387	38	12	7	3	9	11	10	18	21	19	35
2002.....	389	41	13	5	11	8	7	9	10	17	26	41
2001.....	394	39	16	9	6	12	11	11	14	23	19	30
2000.....	405	48	20	6	10	2	2	7	13	22	24	25
1999.....	442	61	26	9	9	10	16	8	16	18	27	39
All races, female												
2003.....	343	39	12	3	4	5	12	8	8	10	17	24
2002.....	311	33	6	5	10	4	7	7	10	6	12	22
2001.....	333	39	10	6	3	7	10	2	14	8	19	25
2000.....	353	43	16	6	8	5	6	9	14	13	13	20
1999.....	408	55	19	5	4	9	6	13	11	15	16	23

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningitis (G00,G03)												
White, both sexes												
2003.....	524	52	18	8	5	10	15	10	16	17	26	41
2002.....	498	52	13	9	11	10	8	9	11	18	21	39
2001.....	526	47	19	10	8	14	12	8	21	15	25	37
2000.....	536	54	23	11	10	4	6	8	17	24	20	35
1999.....	601	68	33	9	10	10	11	16	18	22	27	39
White, male												
2003.....	293	32	9	6	3	6	10	6	11	12	14	27
2002.....	270	29	8	5	6	7	3	5	4	14	12	24
2001.....	275	22	13	5	5	7	6	7	11	10	14	20
2000.....	294	31	14	6	4	1	2	3	7	15	12	21
1999.....	308	46	20	5	7	5	7	4	9	12	15	22
White, female												
2003.....	231	20	9	2	2	4	5	4	5	5	12	14
2002.....	228	23	5	4	5	3	5	4	7	4	9	15
2001.....	251	25	6	5	3	7	6	1	10	5	11	17
2000.....	242	23	9	5	6	3	4	5	10	9	8	14
1999.....	293	22	13	4	3	5	4	12	9	10	12	17

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningitis (G00,G03)												
Black, both sexes												
2003.....	187	23	5	2	1	4	8	8	10	13	9	16
2002.....	178	20	6	1	9	1	5	7	8	5	15	21
2001.....	175	29	5	4	-	4	7	5	6	13	13	18
2000.....	204	35	13	1	8	3	2	8	10	10	17	9
1999.....	222	42	11	5	3	9	8	5	6	10	12	22
Black, male												
2003.....	84	6	2	1	-	3	1	4	7	8	4	7
2002.....	107	11	5	-	5	1	4	4	5	3	12	15
2001.....	105	16	2	3	-	4	4	4	2	12	5	10
2000.....	102	16	6	-	6	1	-	4	6	6	12	3
1999.....	121	13	6	4	2	5	6	4	5	6	9	17
Black, female												
2003.....	103	17	3	1	1	1	7	4	3	5	5	9
2002.....	71	9	1	1	4	-	1	3	3	2	3	6
2001.....	70	13	3	1	-	-	3	1	4	1	8	8
2000.....	102	19	7	1	2	2	2	4	4	4	5	6
1999.....	101	29	5	1	1	4	2	1	1	4	3	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Meningitis (G00,G03)

Asian or Pacific Islander 2/, both sexes

2003.....	11	1	-	-	1	-	-	-	-	-	1	1
2002.....	16	1	-	-	-	1	-	-	1	-	1	3
2001.....	16	1	-	1	1	-	2	-	1	3	-	-
2000.....	13	1	-	-	-	-	-	-	-	-	-	1
1999.....	17	4	1	-	-	-	1	-	1	1	3	-

Asian or Pacific Islander 2/, male

2003.....	5	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	1	-	1	2
2001.....	9	1	-	1	1	-	1	-	1	1	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	1
1999.....	6	1	-	-	-	-	1	-	-	-	3	-

Asian or Pacific Islander 2/, female

2003.....	6	1	-	-	1	-	-	-	-	-	-	1
2002.....	9	1	-	-	-	1	-	-	-	-	-	1
2001.....	7	-	-	-	-	-	1	-	-	2	-	-
2000.....	7	1	-	-	-	-	-	-	-	-	-	-
1999.....	11	3	1	-	-	-	-	-	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alzheimer's disease (G30)												
All races, both sexes												
2003.....	63,457	-	-	-	-	-	-	-	-	2	9	21
2002.....	58,866	-	-	-	1	-	-	-	2	3	9	12
2001.....	53,852	-	-	-	-	-	-	-	2	1	2	17
2000.....	49,558	-	-	-	-	-	-	1	1	2	4	17
1999.....	44,536	-	-	-	-	-	1	-	-	1	6	20
All races, male												
2003.....	18,335	-	-	-	-	-	-	-	-	1	3	9
2002.....	16,989	-	-	-	-	-	-	-	2	3	5	5
2001.....	15,762	-	-	-	-	-	-	-	2	1	2	11
2000.....	14,438	-	-	-	-	-	-	1	-	1	3	10
1999.....	13,391	-	-	-	-	-	1	-	-	-	3	10
All races, female												
2003.....	45,122	-	-	-	-	-	-	-	-	1	6	12
2002.....	41,877	-	-	-	1	-	-	-	-	-	4	7
2001.....	38,090	-	-	-	-	-	-	-	-	-	-	6
2000.....	35,120	-	-	-	-	-	-	-	1	1	1	7
1999.....	31,145	-	-	-	-	-	-	-	-	1	3	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alzheimer's disease (G30)												
White, both sexes												
2003.....	59,184	-	-	-	-	-	-	-	-	2	9	19
2002.....	55,058	-	-	-	1	-	-	-	1	1	6	12
2001.....	50,348	-	-	-	-	-	-	-	2	1	2	13
2000.....	46,460	-	-	-	-	-	-	1	1	2	4	14
1999.....	41,877	-	-	-	-	-	1	-	-	1	4	18
White, male												
2003.....	17,086	-	-	-	-	-	-	-	-	1	3	8
2002.....	15,874	-	-	-	-	-	-	-	1	1	4	5
2001.....	14,714	-	-	-	-	-	-	-	2	1	2	9
2000.....	13,524	-	-	-	-	-	-	1	-	1	3	8
1999.....	12,585	-	-	-	-	-	1	-	-	-	2	10
White, female												
2003.....	42,098	-	-	-	-	-	-	-	-	1	6	11
2002.....	39,184	-	-	-	1	-	-	-	-	-	2	7
2001.....	35,634	-	-	-	-	-	-	-	-	-	-	4
2000.....	32,936	-	-	-	-	-	-	-	1	1	1	6
1999.....	29,292	-	-	-	-	-	-	-	-	1	2	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

All races, both sexes

2003.....	902,443	555	234	134	213	519	928	1,352	2,737	5,375	11,555	19,091
2002.....	918,628	640	223	127	231	493	790	1,348	2,638	5,541	11,421	18,716
2001.....	922,334	600	285	139	226	443	837	1,351	2,687	5,568	11,118	17,768
2000.....	936,923	636	234	136	226	493	816	1,295	2,497	5,491	11,133	17,785
1999.....	950,314	644	234	144	210	548	767	1,270	2,603	5,918	11,092	17,354

All races, male

2003.....	423,810	310	116	75	129	323	570	830	1,779	3,585	7,717	13,215
2002.....	429,682	343	122	57	137	328	517	869	1,699	3,602	7,754	12,998
2001.....	428,065	315	149	65	119	278	529	840	1,715	3,672	7,508	12,418
2000.....	435,912	341	128	78	128	295	488	800	1,578	3,538	7,572	12,544
1999.....	441,827	347	127	84	115	345	454	769	1,673	3,898	7,573	12,150

All races, female

2003.....	478,633	245	118	59	84	196	358	522	958	1,790	3,838	5,876
2002.....	488,946	297	101	70	94	165	273	479	939	1,939	3,667	5,718
2001.....	494,269	285	136	74	107	165	308	511	972	1,896	3,610	5,350
2000.....	501,011	295	106	58	98	198	328	495	919	1,953	3,561	5,241
1999.....	508,487	297	107	60	95	203	313	501	930	2,020	3,519	5,204

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

White, both sexes

2003.....	780,020	343	148	88	146	355	596	861	1,802	3,676	8,143	13,669
2002.....	796,748	421	160	92	152	341	522	858	1,707	3,813	8,216	13,501
2001.....	801,052	383	185	100	158	307	531	876	1,802	3,858	7,770	12,789
2000.....	816,219	434	161	95	147	331	527	845	1,660	3,812	7,867	12,794
1999.....	829,464	434	161	108	142	384	502	834	1,764	4,115	7,769	12,559

White, male

2003.....	365,390	184	67	48	92	220	373	544	1,210	2,557	5,701	9,879
2002.....	372,088	230	92	42	89	224	353	577	1,131	2,547	5,852	9,822
2001.....	370,918	212	95	49	87	189	346	563	1,185	2,622	5,535	9,346
2000.....	379,095	233	96	56	90	201	327	540	1,105	2,573	5,616	9,522
1999.....	384,572	234	90	64	76	237	294	517	1,180	2,807	5,576	9,214

White, female

2003.....	414,630	159	81	40	54	135	223	317	592	1,119	2,442	3,790
2002.....	424,660	191	68	50	63	117	169	281	576	1,266	2,364	3,679
2001.....	430,134	171	90	51	71	118	185	313	617	1,236	2,235	3,443
2000.....	437,124	201	65	39	57	130	200	305	555	1,239	2,251	3,272
1999.....	444,892	200	71	44	66	147	208	317	584	1,308	2,193	3,345

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2003.....	104,181	173	73	39	56	141	297	433	818	1,497	3,038	4,845
2002.....	104,215	186	51	26	69	126	237	439	828	1,533	2,881	4,698
2001.....	104,316	186	87	34	54	124	275	434	789	1,509	3,034	4,482
2000.....	104,280	171	63	34	62	144	255	407	764	1,510	2,946	4,531
1999.....	104,693	177	61	35	58	149	245	392	742	1,630	3,030	4,317

Black, male

2003.....	48,853	105	43	23	30	92	179	252	480	881	1,772	2,941
2002.....	48,264	97	25	10	41	83	141	260	489	910	1,672	2,812
2001.....	48,237	94	46	13	27	81	163	256	462	905	1,750	2,713
2000.....	47,980	91	27	18	30	80	139	231	426	854	1,736	2,710
1999.....	48,578	94	32	19	34	99	145	228	432	965	1,786	2,614

Black, female

2003.....	55,328	68	30	16	26	49	118	181	338	616	1,266	1,904
2002.....	55,951	89	26	16	28	43	96	179	339	623	1,209	1,886
2001.....	56,079	92	41	21	27	43	112	178	327	604	1,284	1,769
2000.....	56,300	80	36	16	32	64	116	176	338	656	1,210	1,821
1999.....	56,115	83	29	16	24	50	100	164	310	665	1,244	1,703

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

American Indian 1/, both sexes

2003.....	3,488	9	5	1	2	5	9	13	39	64	112	168
2002.....	3,259	6	6	2	5	10	10	15	26	47	100	128
2001.....	3,162	11	3	3	1	2	8	12	24	63	99	151
2000.....	3,198	7	2	2	5	2	10	12	18	53	97	130
1999.....	3,121	8	5	-	6	2	8	11	32	50	89	119

American Indian 1/, male

2003.....	1,882	5	2	-	-	3	7	6	32	48	67	113
2002.....	1,752	4	2	1	3	7	9	8	21	35	67	93
2001.....	1,663	3	2	2	-	1	7	4	16	39	64	103
2000.....	1,670	3	1	1	2	2	6	5	11	30	66	90
1999.....	1,613	5	2	-	2	1	6	6	13	34	59	81

American Indian 1/, female

2003.....	1,606	4	3	1	2	2	2	7	7	16	45	55
2002.....	1,507	2	4	1	2	3	1	7	5	12	33	35
2001.....	1,499	8	1	1	1	1	1	8	8	24	35	48
2000.....	1,528	4	1	1	3	-	4	7	7	23	31	40
1999.....	1,508	3	3	-	4	1	2	5	19	16	30	38

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

Asian or Pacific Islander 2/, both sexes

2003.....	14,754	30	8	6	9	18	26	45	78	138	262	409
2002.....	14,406	27	6	7	5	16	21	36	77	148	224	389
2001.....	13,804	20	10	2	13	10	23	29	72	138	215	346
2000.....	13,226	24	8	5	12	16	24	31	55	116	223	330
1999.....	13,036	25	7	1	4	13	12	33	65	123	204	359

Asian or Pacific Islander 2/, male

2003.....	7,685	16	4	4	7	8	11	28	57	99	177	282
2002.....	7,578	12	3	4	4	14	14	24	58	110	163	271
2001.....	7,247	6	6	1	5	7	13	17	52	106	159	256
2000.....	7,167	14	4	3	6	12	16	24	36	81	154	222
1999.....	7,064	14	3	1	3	8	9	18	48	92	152	241

Asian or Pacific Islander 2/, female

2003.....	7,069	14	4	2	2	10	15	17	21	39	85	127
2002.....	6,828	15	3	3	1	2	7	12	19	38	61	118
2001.....	6,557	14	4	1	8	3	10	12	20	32	56	90
2000.....	6,059	10	4	2	6	4	8	7	19	35	69	108
1999.....	5,972	11	4	-	1	5	3	15	17	31	52	118

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races, both sexes

2003.....	685,089	439	186	104	160	393	740	1,075	2,175	4,310	9,290	15,520
2002.....	696,947	500	165	92	163	405	617	1,076	2,089	4,448	9,240	15,179
2001.....	700,142	479	225	98	174	347	652	1,052	2,108	4,419	8,907	14,390
2000.....	710,760	493	181	106	165	403	628	1,013	1,945	4,333	8,848	14,313
1999.....	725,192	524	183	116	161	463	606	992	2,074	4,703	8,897	14,096

All races, male

2003.....	336,095	247	89	56	99	249	469	677	1,452	2,977	6,486	11,164
2002.....	340,933	267	87	40	101	275	411	716	1,415	3,048	6,590	10,983
2001.....	339,095	243	121	43	95	223	424	672	1,396	3,038	6,308	10,491
2000.....	344,807	260	107	64	99	248	390	640	1,282	2,945	6,341	10,596
1999.....	351,617	270	104	66	91	297	367	618	1,370	3,252	6,388	10,346

All races, female

2003.....	348,994	192	97	48	61	144	271	398	723	1,333	2,804	4,356
2002.....	356,014	233	78	52	62	130	206	360	674	1,400	2,650	4,196
2001.....	361,047	236	104	55	79	124	228	380	712	1,381	2,599	3,899
2000.....	365,953	233	74	42	66	155	238	373	663	1,388	2,507	3,717
1999.....	373,575	254	79	50	70	166	239	374	704	1,451	2,509	3,750

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of heart (I00-I09,I11,I13,I20-I51)												
White, both sexes												
2003.....	594,842	273	117	66	108	264	472	679	1,441	3,050	6,730	11,428
2002.....	606,876	332	115	65	111	274	393	682	1,358	3,131	6,836	11,271
2001.....	610,638	312	141	68	119	232	407	678	1,408	3,112	6,398	10,646
2000.....	621,719	343	126	71	106	264	392	637	1,300	3,083	6,449	10,645
1999.....	635,118	352	124	83	110	321	379	638	1,410	3,324	6,394	10,500
White, male												
2003.....	291,560	150	52	35	70	166	298	444	1,005	2,189	4,904	8,539
2002.....	296,904	181	65	30	69	185	272	468	952	2,199	5,089	8,497
2001.....	295,556	170	74	31	67	145	269	451	957	2,216	4,765	8,100
2000.....	301,551	182	80	42	69	161	252	418	898	2,185	4,835	8,267
1999.....	307,585	183	74	47	61	198	231	410	966	2,376	4,809	8,052
White, female												
2003.....	303,282	123	65	31	38	98	174	235	436	861	1,826	2,889
2002.....	309,972	151	50	35	42	89	121	214	406	932	1,747	2,774
2001.....	315,082	142	67	37	52	87	138	227	451	896	1,633	2,546
2000.....	320,168	161	46	29	37	103	140	219	402	898	1,614	2,378
1999.....	327,533	169	50	36	49	123	148	228	444	948	1,585	2,448

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Black, both sexes

2003.....	77,372	133	57	33	43	111	243	348	643	1,125	2,303	3,680
2002.....	77,621	144	40	19	46	109	197	355	649	1,181	2,175	3,541
2001.....	77,674	141	74	28	45	105	221	342	628	1,168	2,283	3,380
2000.....	77,523	122	49	30	46	125	208	346	591	1,127	2,178	3,356
1999.....	78,574	144	50	32	45	130	210	319	595	1,259	2,287	3,275

Black, male

2003.....	37,466	81	32	18	24	74	158	202	378	685	1,402	2,308
2002.....	37,094	76	18	6	27	72	120	221	398	742	1,318	2,208
2001.....	37,016	65	41	11	24	70	139	202	384	719	1,373	2,109
2000.....	36,740	62	24	18	24	75	118	201	346	674	1,335	2,091
1999.....	37,576	72	27	18	27	91	124	191	355	781	1,412	2,050

Black, female

2003.....	39,906	52	25	15	19	37	85	146	265	440	901	1,372
2002.....	40,527	68	22	13	19	37	77	134	251	439	857	1,333
2001.....	40,658	76	33	17	21	35	82	140	244	449	910	1,271
2000.....	40,783	60	25	12	22	50	90	145	245	453	843	1,265
1999.....	40,998	72	23	14	18	39	86	128	240	478	875	1,225

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

American Indian 1/, both sexes

2003.....	2,712	9	4	-	2	4	7	10	34	48	87	135
2002.....	2,467	5	6	2	3	8	9	12	22	36	71	95
2001.....	2,402	9	3	2	1	2	8	10	19	49	76	120
2000.....	2,417	7	1	1	4	2	9	8	16	44	78	103
1999.....	2,404	6	5	-	3	2	7	10	20	40	66	96

American Indian 1/, male

2003.....	1,529	5	1	-	-	2	5	5	27	37	58	96
2002.....	1,412	3	2	1	3	6	8	7	18	29	53	74
2001.....	1,358	3	2	1	-	1	7	4	14	33	52	92
2000.....	1,341	3	-	1	2	2	6	5	11	26	57	77
1999.....	1,302	3	2	-	1	1	5	5	9	29	46	70

American Indian 1/, female

2003.....	1,183	4	3	-	2	2	2	5	7	11	29	39
2002.....	1,055	2	4	1	-	2	1	5	4	7	18	21
2001.....	1,044	6	1	1	1	1	1	6	5	16	24	28
2000.....	1,076	4	1	-	2	-	3	3	5	18	21	26
1999.....	1,102	3	3	-	2	1	2	5	11	11	20	26

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Asian or Pacific Islander 2/, both sexes

2003.....	10,163	24	8	5	7	14	18	38	57	87	170	277
2002.....	9,983	19	4	6	3	14	18	27	60	100	158	272
2001.....	9,428	17	7	-	9	8	16	22	53	90	150	244
2000.....	9,101	21	5	4	9	12	19	22	38	79	143	209
1999.....	9,096	22	4	1	3	10	10	25	49	80	150	225

Asian or Pacific Islander 2/, male

2003.....	5,540	11	4	3	5	7	8	26	42	66	122	221
2002.....	5,523	7	2	3	2	12	11	20	47	78	130	204
2001.....	5,165	5	4	-	4	7	9	15	41	70	118	190
2000.....	5,175	13	3	3	4	10	14	16	27	60	114	161
1999.....	5,154	12	1	1	2	7	7	12	40	66	121	174

Asian or Pacific Islander 2/, female

2003.....	4,623	13	4	2	2	7	10	12	15	21	48	56
2002.....	4,460	12	2	3	1	2	7	7	13	22	28	68
2001.....	4,263	12	3	-	5	1	7	7	12	20	32	54
2000.....	3,926	8	2	1	5	2	5	6	11	19	29	48
1999.....	3,942	10	3	-	1	3	3	13	9	14	29	51

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races, both sexes

2003.....	3,624	1	1	2	4	4	7	11	14	41	58	107
2002.....	3,579	4	1	4	2	8	10	10	26	38	57	97
2001.....	3,489	3	1	1	4	3	8	11	23	45	85	107
2000.....	3,582	6	2	2	2	9	8	10	23	49	92	119
1999.....	3,676	6	1	2	3	5	8	12	30	47	77	101

All races, male

2003.....	1,153	1	-	2	2	3	4	8	6	19	25	39
2002.....	1,078	3	1	1	1	3	5	6	12	19	26	40
2001.....	1,031	3	1	-	2	2	5	4	11	18	36	46
2000.....	1,072	5	2	1	2	5	6	2	10	25	36	49
1999.....	1,042	2	1	1	1	4	6	5	12	19	30	40

All races, female

2003.....	2,471	-	1	-	2	1	3	3	8	22	33	68
2002.....	2,501	1	-	3	1	5	5	4	14	19	31	57
2001.....	2,458	-	-	1	2	1	3	7	12	27	49	61
2000.....	2,510	1	-	1	-	4	2	8	13	24	56	70
1999.....	2,634	4	-	1	2	1	2	7	18	28	47	61

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
White, both sexes												
2003.....	3,226	1	1	2	2	-	5	9	9	25	36	71
2002.....	3,204	4	1	1	2	3	7	6	18	32	36	70
2001.....	3,086	1	1	1	2	3	7	5	10	28	57	70
2000.....	3,229	5	2	2	2	6	4	7	18	34	62	90
1999.....	3,302	4	1	1	2	2	6	8	17	33	51	79
White, male												
2003.....	1,023	1	-	2	1	-	4	6	3	12	14	29
2002.....	950	3	1	1	1	-	4	3	8	15	15	30
2001.....	910	1	1	-	1	2	5	2	7	12	28	32
2000.....	965	4	2	1	2	3	3	1	9	19	27	39
1999.....	933	1	1	-	-	1	4	4	8	14	21	32
White, female												
2003.....	2,203	-	1	-	1	-	1	3	6	13	22	42
2002.....	2,254	1	-	-	1	3	3	3	10	17	21	40
2001.....	2,176	-	-	1	1	1	2	3	3	16	29	38
2000.....	2,264	1	-	1	-	3	1	6	9	15	35	51
1999.....	2,369	3	-	1	2	1	2	4	9	19	30	47

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2003.....	308	-	-	-	2	2	2	2	4	13	16	32
2002.....	252	-	-	2	-	3	2	3	6	4	16	21
2001.....	302	2	-	-	1	-	1	3	10	15	26	27
2000.....	265	1	-	-	-	3	4	3	5	11	24	23
1999.....	275	1	-	1	1	2	2	4	9	12	20	20

Black, male

2003.....	110	-	-	-	1	2	-	2	2	6	9	9
2002.....	93	-	-	-	-	2	1	2	2	2	10	7
2001.....	86	2	-	-	-	-	-	-	3	6	7	11
2000.....	79	1	-	-	-	2	3	1	1	5	7	8
1999.....	88	1	-	1	1	2	2	1	2	4	8	8

Black, female

2003.....	198	-	-	-	1	-	2	-	2	7	7	23
2002.....	159	-	-	2	-	1	1	1	4	2	6	14
2001.....	216	-	-	-	1	-	1	3	7	9	19	16
2000.....	186	-	-	-	-	1	1	2	4	6	17	15
1999.....	187	-	-	-	-	-	-	3	7	8	12	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

American Indian 1/, both sexes

2003.....	11	-	-	-	-	-	-	-	-	1	-	-
2002.....	23	-	-	-	-	1	-	1	-	-	1	-
2001.....	21	-	-	-	-	-	-	-	-	-	1	3
2000.....	18	-	-	-	-	-	-	-	-	1	1	1
1999.....	19	-	-	-	-	-	-	-	2	1	-	-

American Indian 1/, male

2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	9	-	-	-	-	-	-	1	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	-	-	1	2
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-

American Indian 1/, female

2003.....	8	-	-	-	-	-	-	-	-	1	-	-
2002.....	14	-	-	-	-	1	-	-	-	-	1	-
2001.....	10	-	-	-	-	-	-	-	-	-	-	1
2000.....	15	-	-	-	-	-	-	-	-	1	1	1
1999.....	17	-	-	-	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Asian or Pacific Islander 2/, both sexes

2003.....	79	-	-	-	-	2	-	-	1	2	6	4
2002.....	100	-	-	1	-	1	1	-	2	2	4	6
2001.....	80	-	-	-	1	-	-	3	3	2	1	7
2000.....	70	-	-	-	-	-	-	-	-	3	5	5
1999.....	80	1	-	-	-	1	-	-	2	1	6	2

Asian or Pacific Islander 2/, male

2003.....	17	-	-	-	-	1	-	-	1	1	2	1
2002.....	26	-	-	-	-	1	-	-	2	2	1	3
2001.....	24	-	-	-	1	-	-	2	1	-	-	1
2000.....	25	-	-	-	-	-	-	-	-	1	2	2
1999.....	19	-	-	-	-	1	-	-	1	-	1	-

Asian or Pacific Islander 2/, female

2003.....	62	-	-	-	-	1	-	-	-	1	4	3
2002.....	74	-	-	1	-	-	1	-	-	-	3	3
2001.....	56	-	-	-	-	-	-	1	2	2	1	6
2000.....	45	-	-	-	-	-	-	-	-	2	3	3
1999.....	61	1	-	-	-	-	-	-	1	1	5	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
All races, both sexes												
2003.....	28,345	-	1	-	-	9	44	90	215	409	897	1,446
2002.....	26,551	-	-	-	2	4	26	81	216	402	811	1,266
2001.....	24,689	-	-	1	-	6	18	65	181	352	739	1,050
2000.....	23,761	-	1	-	2	5	27	69	144	323	658	952
1999.....	22,702	1	-	-	-	6	27	50	160	347	617	889
All races, male												
2003.....	12,536	-	1	-	-	6	34	64	144	285	592	962
2002.....	11,660	-	-	-	1	4	22	64	151	289	560	844
2001.....	10,451	-	-	-	-	4	13	40	129	234	506	715
2000.....	9,876	-	-	-	1	3	21	43	96	210	433	625
1999.....	9,316	-	-	-	-	6	19	41	108	232	428	586
All races, female												
2003.....	15,809	-	-	-	-	3	10	26	71	124	305	484
2002.....	14,891	-	-	-	1	-	4	17	65	113	251	422
2001.....	14,238	-	-	1	-	2	5	25	52	118	233	335
2000.....	13,885	-	1	-	1	2	6	26	48	113	225	327
1999.....	13,386	1	-	-	-	-	8	9	52	115	189	303

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
White, both sexes												
2003.....	20,904	-	-	-	-	4	23	49	108	231	503	785
2002.....	19,530	-	-	-	1	1	12	39	102	222	458	753
2001.....	18,232	-	-	-	-	4	6	36	89	197	389	572
2000.....	17,425	-	-	-	2	1	14	31	64	174	368	521
1999.....	16,646	1	-	-	-	2	10	24	85	170	321	469
White, male												
2003.....	8,948	-	-	-	-	2	19	41	80	179	377	572
2002.....	8,234	-	-	-	1	1	11	32	76	164	347	539
2001.....	7,395	-	-	-	-	3	4	24	72	149	292	412
2000.....	6,927	-	-	-	1	1	11	22	47	126	259	383
1999.....	6,499	-	-	-	-	2	7	22	69	125	245	340
White, female												
2003.....	11,956	-	-	-	-	2	4	8	28	52	126	213
2002.....	11,296	-	-	-	-	-	1	7	26	58	111	214
2001.....	10,837	-	-	-	-	1	2	12	17	48	97	160
2000.....	10,498	-	-	-	1	-	3	9	17	48	109	138
1999.....	10,147	1	-	-	-	-	3	2	16	45	76	129

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Black, both sexes												
2003.....	6,843	-	1	-	-	5	20	40	97	170	374	629
2002.....	6,480	-	-	-	1	3	12	40	110	168	332	493
2001.....	5,970	-	-	1	-	2	12	28	87	149	330	459
2000.....	5,886	-	1	-	-	4	13	37	77	145	279	420
1999.....	5,619	-	-	-	-	3	17	26	72	169	281	409
Black, male												
2003.....	3,298	-	1	-	-	4	15	22	57	98	204	365
2002.....	3,146	-	-	-	-	3	9	30	71	116	199	289
2001.....	2,813	-	-	-	-	1	9	15	53	81	199	287
2000.....	2,722	-	-	-	-	2	10	20	46	81	167	235
1999.....	2,606	-	-	-	-	3	12	19	36	101	169	239
Black, female												
2003.....	3,545	-	-	-	-	1	5	18	40	72	170	264
2002.....	3,334	-	-	-	1	-	3	10	39	52	133	204
2001.....	3,157	-	-	1	-	1	3	13	34	68	131	172
2000.....	3,164	-	1	-	-	2	3	17	31	64	112	185
1999.....	3,013	-	-	-	-	-	5	7	36	68	112	170

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive heart disease (I11)

American Indian 1/, both sexes

2003.....	112	-	-	-	-	-	-	1	5	3	5	9
2002.....	68	-	-	-	-	-	2	-	-	6	5	4
2001.....	64	-	-	-	-	-	-	-	2	1	5	2
2000.....	59	-	-	-	-	-	-	-	-	-	-	1
1999.....	56	-	-	-	-	-	-	-	-	3	3	-

American Indian 1/, male

2003.....	54	-	-	-	-	-	-	1	3	3	2	6
2002.....	32	-	-	-	-	-	2	-	-	4	2	4
2001.....	29	-	-	-	-	-	-	-	1	1	2	2
2000.....	32	-	-	-	-	-	-	-	-	-	-	1
1999.....	26	-	-	-	-	-	-	-	-	2	3	-

American Indian 1/, female

2003.....	58	-	-	-	-	-	-	-	2	-	3	3
2002.....	36	-	-	-	-	-	-	-	-	2	3	-
2001.....	35	-	-	-	-	-	-	-	1	-	3	-
2000.....	27	-	-	-	-	-	-	-	-	-	-	-
1999.....	30	-	-	-	-	-	-	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Asian or Pacific Islander 2/, both sexes												
2003.....	486	-	-	-	-	-	1	-	5	5	15	23
2002.....	473	-	-	-	-	-	-	2	4	6	16	16
2001.....	423	-	-	-	-	-	-	1	3	5	15	17
2000.....	391	-	-	-	-	-	-	1	3	4	11	10
1999.....	381	-	-	-	-	1	-	-	3	5	12	11
Asian or Pacific Islander 2/, male												
2003.....	236	-	-	-	-	-	-	-	4	5	9	19
2002.....	248	-	-	-	-	-	-	2	4	5	12	12
2001.....	214	-	-	-	-	-	-	1	3	3	13	14
2000.....	195	-	-	-	-	-	-	1	3	3	7	6
1999.....	185	-	-	-	-	1	-	-	3	4	11	7
Asian or Pacific Islander 2/, female												
2003.....	250	-	-	-	-	-	1	-	1	-	6	4
2002.....	225	-	-	-	-	-	-	-	-	1	4	4
2001.....	209	-	-	-	-	-	-	-	-	2	2	3
2000.....	196	-	-	-	-	-	-	-	-	1	4	4
1999.....	196	-	-	-	-	-	-	-	-	1	1	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
All races, both sexes												
2003.....	3,108	-	1	-	-	-	5	8	20	35	58	87
2002.....	2,895	-	-	-	-	1	4	11	17	28	49	75
2001.....	2,826	1	-	-	-	2	5	10	18	40	59	64
2000.....	2,785	-	-	-	1	1	5	11	17	22	61	68
1999.....	3,327	-	-	2	-	2	4	5	16	43	71	80
All races, male												
2003.....	1,326	-	1	-	-	-	4	4	9	26	36	63
2002.....	1,205	-	-	-	-	-	3	9	10	19	25	49
2001.....	1,169	1	-	-	-	-	4	4	13	20	43	46
2000.....	1,163	-	-	-	-	1	3	7	11	12	37	43
1999.....	1,390	-	-	1	-	1	2	3	13	24	43	50
All races, female												
2003.....	1,782	-	-	-	-	-	1	4	11	9	22	24
2002.....	1,690	-	-	-	-	1	1	2	7	9	24	26
2001.....	1,657	-	-	-	-	2	1	6	5	20	16	18
2000.....	1,622	-	-	-	1	-	2	4	6	10	24	25
1999.....	1,937	-	-	1	-	1	2	2	3	19	28	30

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
White, both sexes												
2003.....	2,103	-	1	-	-	-	2	4	3	8	13	25
2002.....	1,925	-	-	-	-	1	1	6	5	5	16	19
2001.....	1,922	-	-	-	-	-	3	2	5	7	18	17
2000.....	1,904	-	-	-	-	-	1	6	2	9	28	21
1999.....	2,235	-	-	1	-	-	-	1	6	14	14	21
White, male												
2003.....	865	-	1	-	-	-	1	2	-	8	6	19
2002.....	760	-	-	-	-	-	1	4	2	5	10	13
2001.....	762	-	-	-	-	-	3	1	2	3	14	11
2000.....	775	-	-	-	-	-	1	4	2	5	17	12
1999.....	906	-	-	1	-	-	-	1	5	7	10	11
White, female												
2003.....	1,238	-	-	-	-	-	1	2	3	-	7	6
2002.....	1,165	-	-	-	-	1	-	2	3	-	6	6
2001.....	1,160	-	-	-	-	-	-	1	3	4	4	6
2000.....	1,129	-	-	-	-	-	-	2	-	4	11	9
1999.....	1,329	-	-	-	-	-	-	-	1	7	4	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Black, both sexes												
2003.....	913	-	-	-	-	-	3	4	16	27	44	60
2002.....	903	-	-	-	-	-	2	5	12	22	32	56
2001.....	832	1	-	-	-	2	2	8	13	29	39	46
2000.....	809	-	-	-	1	1	4	4	15	13	33	45
1999.....	1,000	-	-	1	-	2	4	3	10	28	57	57
Black, male												
2003.....	413	-	-	-	-	-	3	2	8	18	29	42
2002.....	405	-	-	-	-	-	1	5	8	13	14	36
2001.....	376	1	-	-	-	-	1	3	11	15	28	34
2000.....	356	-	-	-	-	1	2	3	9	7	20	30
1999.....	434	-	-	-	-	1	2	1	8	16	33	37
Black, female												
2003.....	500	-	-	-	-	-	-	2	8	9	15	18
2002.....	498	-	-	-	-	-	1	-	4	9	18	20
2001.....	456	-	-	-	-	2	1	5	2	14	11	12
2000.....	453	-	-	-	1	-	2	1	6	6	13	15
1999.....	566	-	-	1	-	1	2	2	2	12	24	20

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
All races, both sexes												
2003.....	480,028	22	12	12	12	32	128	274	808	2,152	5,395	9,989
2002.....	494,382	23	7	4	16	30	103	251	745	2,211	5,472	9,892
2001.....	502,189	26	9	4	16	31	132	279	837	2,212	5,322	9,461
2000.....	515,204	30	10	8	8	33	90	236	699	2,233	5,286	9,638
1999.....	529,659	27	5	13	9	52	85	245	769	2,387	5,364	9,524
All races, male												
2003.....	246,342	13	6	4	6	21	86	197	616	1,630	4,049	7,661
2002.....	252,760	15	4	1	12	18	74	191	550	1,670	4,170	7,582
2001.....	254,005	17	3	2	11	24	95	211	619	1,667	4,045	7,323
2000.....	260,574	17	7	5	4	21	60	181	523	1,665	4,122	7,601
1999.....	267,268	13	3	10	7	36	57	169	570	1,801	4,169	7,447
All races, female												
2003.....	233,686	9	6	8	6	11	42	77	192	522	1,346	2,328
2002.....	241,622	8	3	3	4	12	29	60	195	541	1,302	2,310
2001.....	248,184	9	6	2	5	7	37	68	218	545	1,277	2,138
2000.....	254,630	13	3	3	4	12	30	55	176	568	1,164	2,037
1999.....	262,391	14	2	3	2	16	28	76	199	586	1,195	2,077

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
White, both sexes												
2003.....	421,482	16	6	7	8	25	87	192	604	1,680	4,206	7,811
2002.....	435,170	14	5	3	12	24	70	185	555	1,700	4,310	7,782
2001.....	442,536	19	6	3	10	22	82	197	623	1,714	4,128	7,426
2000.....	455,400	23	10	6	7	28	64	160	516	1,745	4,106	7,570
1999.....	469,149	18	4	9	5	36	49	180	573	1,855	4,133	7,541
White, male												
2003.....	216,751	9	3	2	5	17	59	143	465	1,304	3,238	6,156
2002.....	223,262	10	2	1	9	14	52	149	424	1,306	3,390	6,172
2001.....	224,464	14	2	2	7	16	62	156	468	1,316	3,244	5,958
2000.....	230,951	14	7	3	4	18	43	124	405	1,336	3,326	6,194
1999.....	237,193	8	3	7	5	25	38	124	431	1,425	3,317	6,089
White, female												
2003.....	204,731	7	3	5	3	8	28	49	139	376	968	1,655
2002.....	211,908	4	3	2	3	10	18	36	131	394	920	1,610
2001.....	218,072	5	4	1	3	6	20	41	155	398	884	1,468
2000.....	224,449	9	3	3	3	10	21	36	111	409	780	1,376
1999.....	231,956	10	1	2	-	11	11	56	142	430	816	1,452

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Black, both sexes												
2003.....	49,274	4	5	4	4	7	36	64	170	399	1,026	1,884
2002.....	50,174	8	2	1	4	4	27	55	158	438	1,022	1,858
2001.....	50,933	7	3	1	5	9	44	73	177	437	1,049	1,787
2000.....	51,265	4	-	2	-	4	20	66	166	432	1,041	1,845
1999.....	51,995	8	1	4	4	14	33	53	160	474	1,096	1,761
Black, male												
2003.....	24,273	3	3	2	1	4	24	40	122	262	686	1,266
2002.....	24,322	4	2	-	3	2	16	34	99	300	659	1,204
2001.....	24,593	3	1	-	3	8	28	49	122	298	685	1,165
2000.....	24,625	2	-	2	-	2	12	48	104	281	682	1,223
1999.....	25,104	4	-	3	2	9	17	38	110	323	734	1,171
Black, female												
2003.....	25,001	1	2	2	3	3	12	24	48	137	340	618
2002.....	25,852	4	-	1	1	2	11	21	59	138	363	654
2001.....	26,340	4	2	1	2	1	16	24	55	139	364	622
2000.....	26,640	2	-	-	-	2	8	18	62	151	359	622
1999.....	26,891	4	1	1	2	5	16	15	50	151	362	590

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
American Indian 1/, both sexes												
2003.....	1,905	-	-	-	-	-	2	2	14	25	57	95
2002.....	1,777	1	-	-	-	1	3	6	10	15	44	61
2001.....	1,710	-	-	-	-	-	3	1	10	18	46	77
2000.....	1,740	1	-	-	-	-	4	4	7	18	55	75
1999.....	1,720	1	-	-	-	-	1	5	11	17	39	65
American Indian 1/, male												
2003.....	1,133	-	-	-	-	-	1	1	10	22	41	70
2002.....	1,041	1	-	-	-	1	3	4	9	13	35	49
2001.....	995	-	-	-	-	-	3	-	8	12	34	58
2000.....	1,008	-	-	-	-	-	3	3	6	14	42	61
1999.....	977	1	-	-	-	-	1	2	6	13	32	52
American Indian 1/, female												
2003.....	772	-	-	-	-	-	1	1	4	3	16	25
2002.....	736	-	-	-	-	-	-	2	1	2	9	12
2001.....	715	-	-	-	-	-	-	1	2	6	12	19
2000.....	732	1	-	-	-	-	1	1	1	4	13	14
1999.....	743	-	-	-	-	-	-	3	5	4	7	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Asian or Pacific Islander 2/, both sexes												
2003.....	7,367	2	1	1	-	-	3	16	20	48	106	199
2002.....	7,261	-	-	-	-	1	3	5	22	58	96	191
2001.....	7,010	-	-	-	1	-	3	8	27	43	99	171
2000.....	6,799	2	-	-	1	1	2	6	10	38	84	148
1999.....	6,795	-	-	-	-	2	2	7	25	41	96	157
Asian or Pacific Islander 2/, male												
2003.....	4,185	1	-	-	-	-	2	13	19	42	84	169
2002.....	4,135	-	-	-	-	1	3	4	18	51	86	157
2001.....	3,953	-	-	-	1	-	2	6	21	41	82	142
2000.....	3,990	1	-	-	-	1	2	6	8	34	72	123
1999.....	3,994	-	-	-	-	2	1	5	23	40	86	135
Asian or Pacific Islander 2/, female												
2003.....	3,182	1	1	1	-	-	1	3	1	6	22	30
2002.....	3,126	-	-	-	-	-	-	1	4	7	10	34
2001.....	3,057	-	-	-	-	-	1	2	6	2	17	29
2000.....	2,809	1	-	-	1	-	-	-	2	4	12	25
1999.....	2,801	-	-	-	-	-	1	2	2	1	10	22

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
All races, both sexes												
2003.....	170,564	9	6	6	6	17	56	105	308	881	2,213	4,092
2002.....	179,514	11	5	2	5	15	47	91	309	882	2,279	4,055
2001.....	184,757	15	2	1	6	15	49	105	370	924	2,252	4,097
2000.....	192,898	20	1	5	1	14	38	108	267	904	2,228	4,230
1999.....	199,454	13	2	5	6	25	35	101	333	935	2,314	4,225
All races, male												
2003.....	89,515	4	2	2	4	11	35	75	225	664	1,637	3,130
2002.....	93,830	7	2	-	2	10	31	67	225	656	1,691	3,066
2001.....	95,875	10	2	-	4	14	34	69	264	691	1,693	3,146
2000.....	100,306	10	1	3	1	8	22	75	194	657	1,730	3,319
1999.....	104,169	6	1	3	4	16	24	68	238	707	1,763	3,261
All races, female												
2003.....	81,049	5	4	4	2	6	21	30	83	217	576	962
2002.....	85,684	4	3	2	3	5	16	24	84	226	588	989
2001.....	88,882	5	-	1	2	1	15	36	106	233	559	951
2000.....	92,592	10	-	2	-	6	16	33	73	247	498	911
1999.....	95,285	7	1	2	2	9	11	33	95	228	551	964

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
White, both sexes												
2003.....	149,915	7	3	5	4	14	39	75	235	705	1,747	3,263
2002.....	157,922	9	5	1	4	13	30	63	238	698	1,826	3,247
2001.....	162,943	12	2	-	2	12	30	70	285	730	1,783	3,321
2000.....	170,584	16	1	3	1	11	28	71	201	744	1,752	3,442
1999.....	177,075	9	2	3	3	19	20	76	249	757	1,839	3,453
White, male												
2003.....	79,157	3	1	1	3	10	25	58	171	548	1,328	2,573
2002.....	83,288	6	2	-	2	8	22	50	179	531	1,390	2,543
2001.....	85,280	9	2	-	1	11	23	46	208	556	1,395	2,654
2000.....	89,383	9	1	1	1	6	18	50	155	556	1,413	2,797
1999.....	93,189	3	1	1	3	13	16	51	178	585	1,447	2,757
White, female												
2003.....	70,758	4	2	4	1	4	14	17	64	157	419	690
2002.....	74,634	3	3	1	2	5	8	13	59	167	436	704
2001.....	77,663	3	-	-	1	1	7	24	77	174	388	667
2000.....	81,201	7	-	2	-	5	10	21	46	188	339	645
1999.....	83,886	6	1	2	-	6	4	25	71	172	392	696

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Black, both sexes												
2003.....	17,441	1	2	1	2	3	16	24	64	160	420	726
2002.....	18,322	2	-	1	1	-	15	25	64	158	403	707
2001.....	18,581	3	-	1	3	3	18	29	70	169	410	684
2000.....	19,112	2	-	2	-	2	9	32	62	146	423	699
1999.....	19,276	4	-	2	3	6	13	22	66	163	424	688
Black, male												
2003.....	8,472	-	1	1	1	1	10	13	46	101	273	473
2002.....	8,680	1	-	-	-	-	7	15	40	103	259	440
2001.....	8,717	1	-	-	2	3	10	19	45	110	250	416
2000.....	9,045	1	-	2	-	1	4	21	35	90	272	449
1999.....	9,186	3	-	2	1	3	7	16	45	107	272	436
Black, female												
2003.....	8,969	1	1	-	1	2	6	11	18	59	147	253
2002.....	9,642	1	-	1	1	-	8	10	24	55	144	267
2001.....	9,864	2	-	1	1	-	8	10	25	59	160	268
2000.....	10,067	1	-	-	-	1	5	11	27	56	151	250
1999.....	10,090	1	-	-	2	3	6	6	21	56	152	252

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
American Indian 1/, both sexes												
2003.....	706	-	-	-	-	-	-	2	3	7	20	40
2002.....	657	-	-	-	-	1	1	-	3	6	15	32
2001.....	660	-	-	-	-	-	-	1	6	9	21	30
2000.....	656	1	-	-	-	-	1	2	3	7	23	30
1999.....	674	-	-	-	-	-	-	2	6	3	19	26
American Indian 1/, male												
2003.....	430	-	-	-	-	-	-	1	2	6	16	32
2002.....	384	-	-	-	-	1	1	-	3	5	12	26
2001.....	404	-	-	-	-	-	-	-	5	9	17	24
2000.....	384	-	-	-	-	-	-	1	3	5	18	25
1999.....	382	-	-	-	-	-	-	-	4	3	17	20
American Indian 1/, female												
2003.....	276	-	-	-	-	-	-	1	1	1	4	8
2002.....	273	-	-	-	-	-	-	-	-	1	3	6
2001.....	256	-	-	-	-	-	-	1	1	-	4	6
2000.....	272	1	-	-	-	-	1	1	-	2	5	5
1999.....	292	-	-	-	-	-	-	2	2	-	2	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Asian or Pacific Islander 2/, both sexes												
2003.....	2,502	1	1	-	-	-	1	4	6	9	26	63
2002.....	2,613	-	-	-	-	1	1	3	4	20	35	69
2001.....	2,573	-	-	-	1	-	1	5	9	16	38	62
2000.....	2,546	1	-	-	-	1	-	3	1	7	30	59
1999.....	2,429	-	-	-	-	-	2	1	12	12	32	58
Asian or Pacific Islander 2/, male												
2003.....	1,456	1	-	-	-	-	-	3	6	9	20	52
2002.....	1,478	-	-	-	-	1	1	2	3	17	30	57
2001.....	1,474	-	-	-	1	-	1	4	6	16	31	52
2000.....	1,494	-	-	-	-	1	-	3	1	6	27	48
1999.....	1,412	-	-	-	-	-	1	1	11	12	27	48
Asian or Pacific Islander 2/, female												
2003.....	1,046	-	1	-	-	-	1	1	-	-	6	11
2002.....	1,135	-	-	-	-	-	-	1	1	3	5	12
2001.....	1,099	-	-	-	-	-	-	1	3	-	7	10
2000.....	1,052	1	-	-	-	-	-	-	-	1	3	11
1999.....	1,017	-	-	-	-	-	1	-	1	-	5	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
All races, both sexes												
2003.....	3,158	4	1	1	-	-	5	3	8	27	71	106
2002.....	3,407	2	-	1	-	2	2	4	14	17	60	119
2001.....	3,351	3	-	-	2	-	7	2	12	32	62	120
2000.....	3,363	1	1	-	-	4	4	4	12	30	62	93
1999.....	3,232	3	1	1	-	-	3	2	4	25	56	92
All races, male												
2003.....	1,621	2	1	-	-	-	5	2	5	20	53	75
2002.....	1,767	1	-	1	-	-	1	2	10	9	41	88
2001.....	1,720	2	-	-	1	-	6	2	8	18	47	89
2000.....	1,749	1	1	-	-	3	3	3	10	24	47	76
1999.....	1,665	1	1	1	-	-	3	1	3	19	44	69
All races, female												
2003.....	1,537	2	-	1	-	-	-	1	3	7	18	31
2002.....	1,640	1	-	-	-	2	1	2	4	8	19	31
2001.....	1,631	1	-	-	1	-	1	-	4	14	15	31
2000.....	1,614	-	-	-	-	1	1	1	2	6	15	17
1999.....	1,567	2	-	-	-	-	-	1	1	6	12	23

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
White, both sexes												
2003.....	2,663	2	-	1	-	-	4	2	4	20	49	73
2002.....	2,907	1	-	1	-	2	1	3	9	15	40	90
2001.....	2,821	3	-	-	2	-	4	2	9	23	40	89
2000.....	2,839	1	1	-	-	4	2	4	8	23	45	69
1999.....	2,752	1	1	1	-	-	3	2	3	13	40	56
White, male												
2003.....	1,338	1	-	-	-	-	4	2	3	16	40	51
2002.....	1,494	1	-	1	-	-	1	2	6	8	34	64
2001.....	1,431	2	-	-	1	-	3	2	7	14	32	69
2000.....	1,453	1	1	-	-	3	1	3	6	19	35	61
1999.....	1,389	1	1	1	-	-	3	1	3	10	31	42
White, female												
2003.....	1,325	1	-	1	-	-	-	-	1	4	9	22
2002.....	1,413	-	-	-	-	2	-	1	3	7	6	26
2001.....	1,390	1	-	-	1	-	1	-	2	9	8	20
2000.....	1,386	-	-	-	-	1	1	1	2	4	10	8
1999.....	1,363	-	-	-	-	-	-	1	-	3	9	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Black, both sexes												
2003.....	436	1	1	-	-	-	1	1	2	7	20	30
2002.....	446	1	-	-	-	-	1	1	5	2	18	27
2001.....	454	-	-	-	-	-	3	-	2	8	21	27
2000.....	456	-	-	-	-	-	1	-	4	7	15	23
1999.....	431	2	-	-	-	-	-	-	1	12	14	34
Black, male												
2003.....	256	1	1	-	-	-	1	-	1	4	12	21
2002.....	243	-	-	-	-	-	-	-	4	1	5	23
2001.....	249	-	-	-	-	-	3	-	-	4	15	17
2000.....	258	-	-	-	-	-	1	-	4	5	11	15
1999.....	248	-	-	-	-	-	-	-	-	9	11	26
Black, female												
2003.....	180	-	-	-	-	-	-	1	1	3	8	9
2002.....	203	1	-	-	-	-	1	1	1	1	13	4
2001.....	205	-	-	-	-	-	-	-	2	4	6	10
2000.....	198	-	-	-	-	-	-	-	-	2	4	8
1999.....	183	2	-	-	-	-	-	-	1	3	3	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
All races, both sexes												
2003.....	306,306	9	5	5	6	15	67	166	492	1,244	3,111	5,791
2002.....	311,461	10	2	1	11	13	54	156	422	1,312	3,133	5,718
2001.....	314,081	8	7	3	8	16	76	172	455	1,256	3,008	5,244
2000.....	318,943	9	8	3	7	15	48	124	420	1,299	2,996	5,315
1999.....	326,973	11	2	7	3	27	47	142	432	1,427	2,994	5,207
All races, male												
2003.....	155,206	7	3	2	2	10	46	120	386	946	2,359	4,456
2002.....	157,163	7	2	-	10	8	42	122	315	1,005	2,438	4,428
2001.....	156,410	5	1	2	6	10	55	140	347	958	2,305	4,088
2000.....	158,519	6	5	2	3	10	35	103	319	984	2,345	4,206
1999.....	161,434	6	1	6	3	20	30	100	329	1,075	2,362	4,117
All races, female												
2003.....	151,100	2	2	3	4	5	21	46	106	298	752	1,335
2002.....	154,298	3	-	1	1	5	12	34	107	307	695	1,290
2001.....	157,671	3	6	1	2	6	21	32	108	298	703	1,156
2000.....	160,424	3	3	1	4	5	13	21	101	315	651	1,109
1999.....	165,539	5	1	1	-	7	17	42	103	352	632	1,090

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
White, both sexes												
2003.....	268,904	7	3	1	4	11	44	115	365	955	2,410	4,475
2002.....	274,341	4	-	1	8	9	39	119	308	987	2,444	4,445
2001.....	276,772	4	4	3	6	10	48	125	329	961	2,305	4,016
2000.....	281,977	6	8	3	6	13	34	85	307	978	2,309	4,059
1999.....	289,322	8	1	5	2	17	26	102	321	1,085	2,254	4,032
White, male												
2003.....	136,256	5	2	1	2	7	30	83	291	740	1,870	3,532
2002.....	138,480	3	-	-	7	6	29	97	239	767	1,966	3,565
2001.....	137,753	3	-	2	5	5	36	108	253	746	1,817	3,235
2000.....	140,115	4	5	2	3	9	24	71	244	761	1,878	3,336
1999.....	142,615	4	1	5	2	12	19	72	250	830	1,839	3,290
White, female												
2003.....	132,648	2	1	-	2	4	14	32	74	215	540	943
2002.....	135,861	1	-	1	1	3	10	22	69	220	478	880
2001.....	139,019	1	4	1	1	5	12	17	76	215	488	781
2000.....	141,862	2	3	1	3	4	10	14	63	217	431	723
1999.....	146,707	4	-	-	-	5	7	30	71	255	415	742

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Black, both sexes												
2003.....	31,397	2	2	3	2	4	19	39	104	232	586	1,128
2002.....	31,406	5	2	-	3	4	11	29	89	278	601	1,124
2001.....	31,898	4	3	-	2	6	23	44	105	260	618	1,076
2000.....	31,697	2	-	-	-	2	10	34	100	279	603	1,123
1999.....	32,288	2	1	2	1	8	20	31	93	299	658	1,039
Black, male												
2003.....	15,545	2	1	1	-	3	13	27	75	157	401	772
2002.....	15,399	3	2	-	3	2	9	19	55	196	395	741
2001.....	15,627	2	1	-	1	5	15	30	77	184	420	732
2000.....	15,322	1	-	-	-	1	7	27	65	186	399	759
1999.....	15,670	1	-	1	1	6	10	22	65	207	451	709
Black, female												
2003.....	15,852	-	1	2	2	1	6	12	29	75	185	356
2002.....	16,007	2	-	-	-	2	2	10	34	82	206	383
2001.....	16,271	2	2	-	1	1	8	14	28	76	198	344
2000.....	16,375	1	-	-	-	1	3	7	35	93	204	364
1999.....	16,618	1	1	1	-	2	10	9	28	92	207	330

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
American Indian 1/, both sexes												
2003.....	1,162	-	-	-	-	-	2	-	10	18	35	53
2002.....	1,084	1	-	-	-	-	2	6	7	9	27	27
2001.....	1,003	-	-	-	-	-	3	-	3	8	25	44
2000.....	1,041	-	-	-	-	-	2	2	4	11	30	44
1999.....	1,009	1	-	-	-	-	1	3	5	14	19	37
American Indian 1/, male												
2003.....	688	-	-	-	-	-	1	-	8	16	24	36
2002.....	636	1	-	-	-	-	2	4	6	8	21	22
2001.....	563	-	-	-	-	-	3	-	2	3	17	32
2000.....	598	-	-	-	-	-	2	2	3	9	23	36
1999.....	575	1	-	-	-	-	1	2	2	10	14	31
American Indian 1/, female												
2003.....	474	-	-	-	-	-	1	-	2	2	11	17
2002.....	448	-	-	-	-	-	-	2	1	1	6	5
2001.....	440	-	-	-	-	-	-	-	1	5	8	12
2000.....	443	-	-	-	-	-	-	-	1	2	7	8
1999.....	434	-	-	-	-	-	-	1	3	4	5	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Asian or Pacific Islander 2/, both sexes												
2003.....	4,843	-	-	1	-	-	2	12	13	39	80	135
2002.....	4,630	-	-	-	-	-	2	2	18	38	61	122
2001.....	4,408	-	-	-	-	-	2	3	18	27	60	108
2000.....	4,228	1	-	-	1	-	2	3	9	31	54	89
1999.....	4,354	-	-	-	-	2	-	6	13	29	63	99
Asian or Pacific Islander 2/, male												
2003.....	2,717	-	-	-	-	-	2	10	12	33	64	116
2002.....	2,648	-	-	-	-	-	2	2	15	34	56	100
2001.....	2,467	-	-	-	-	-	1	2	15	25	51	89
2000.....	2,484	1	-	-	-	-	2	3	7	28	45	75
1999.....	2,574	-	-	-	-	2	-	4	12	28	58	87
Asian or Pacific Islander 2/, female												
2003.....	2,126	-	-	1	-	-	-	2	1	6	16	19
2002.....	1,982	-	-	-	-	-	-	-	3	4	5	22
2001.....	1,941	-	-	-	-	-	1	1	3	2	9	19
2000.....	1,744	-	-	-	1	-	-	-	2	3	9	14
1999.....	1,780	-	-	-	-	-	-	2	1	1	5	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races, both sexes												
2003.....	67,232	2	-	-	-	2	14	50	161	414	1,074	2,151
2002.....	68,129	2	-	-	1	4	13	39	142	410	1,105	2,081
2001.....	68,103	-	-	-	-	3	17	39	140	402	1,055	1,838
2000.....	69,554	1	-	-	1	-	15	33	111	391	985	1,894
1999.....	71,926	-	1	-	1	-	13	32	128	453	1,035	1,792
All races, male												
2003.....	36,042	2	-	-	-	1	11	40	141	315	842	1,715
2002.....	36,047	1	-	-	-	3	10	31	113	325	891	1,634
2001.....	35,440	-	-	-	-	2	14	30	103	315	815	1,488
2000.....	35,896	-	-	-	1	-	12	26	89	312	787	1,533
1999.....	36,633	-	-	-	1	-	10	21	98	355	837	1,438
All races, female												
2003.....	31,190	-	-	-	-	1	3	10	20	99	232	436
2002.....	32,082	1	-	-	1	1	3	8	29	85	214	447
2001.....	32,663	-	-	-	-	1	3	9	37	87	240	350
2000.....	33,658	1	-	-	-	-	3	7	22	79	198	361
1999.....	35,293	-	1	-	-	-	3	11	30	98	198	354

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
White, both sexes												
2003.....	55,760	1	-	-	-	2	10	35	121	324	820	1,609
2002.....	56,569	1	-	-	1	4	11	32	100	296	834	1,602
2001.....	56,414	-	-	-	-	2	10	28	105	304	785	1,370
2000.....	57,794	-	-	-	1	-	13	19	83	291	743	1,393
1999.....	59,711	-	-	-	1	-	5	20	90	336	761	1,305
White, male												
2003.....	29,560	1	-	-	-	1	8	27	106	246	653	1,323
2002.....	29,604	1	-	-	-	3	8	26	83	239	696	1,293
2001.....	29,039	-	-	-	-	1	7	23	80	244	620	1,145
2000.....	29,467	-	-	-	1	-	10	14	69	237	615	1,164
1999.....	29,977	-	-	-	1	-	5	14	71	269	640	1,076
White, female												
2003.....	26,200	-	-	-	-	1	2	8	15	78	167	286
2002.....	26,965	-	-	-	1	1	3	6	17	57	138	309
2001.....	27,375	-	-	-	-	1	3	5	25	60	165	225
2000.....	28,327	-	-	-	-	-	3	5	14	54	128	229
1999.....	29,734	-	-	-	-	-	-	6	19	67	121	229

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Black, both sexes												
2003.....	9,923	1	-	-	-	-	3	12	35	73	212	459
2002.....	10,195	1	-	-	-	-	2	5	36	100	235	434
2001.....	10,321	-	-	-	-	1	6	11	27	89	236	413
2000.....	10,418	1	-	-	-	-	1	12	24	87	208	459
1999.....	10,823	-	1	-	-	-	8	8	33	110	251	437
Black, male												
2003.....	5,543	1	-	-	-	-	2	10	30	52	155	324
2002.....	5,598	-	-	-	-	-	2	3	25	73	161	302
2001.....	5,578	-	-	-	-	1	6	7	17	64	167	294
2000.....	5,615	-	-	-	-	-	1	10	16	65	145	332
1999.....	5,815	-	-	-	-	-	5	5	22	80	177	317
Black, female												
2003.....	4,380	-	-	-	-	-	1	2	5	21	57	135
2002.....	4,597	1	-	-	-	-	-	2	11	27	74	132
2001.....	4,743	-	-	-	-	-	-	4	10	25	69	119
2000.....	4,803	1	-	-	-	-	-	2	8	22	63	127
1999.....	5,008	-	1	-	-	-	3	3	11	30	74	120

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
American Indian 1/, both sexes												
2003.....	358	-	-	-	-	-	-	-	2	6	12	22
2002.....	318	-	-	-	-	-	-	2	1	2	16	8
2001.....	335	-	-	-	-	-	1	-	2	3	11	15
2000.....	266	-	-	-	-	-	1	1	3	3	9	13
1999.....	295	-	-	-	-	-	-	1	2	3	3	13
American Indian 1/, male												
2003.....	227	-	-	-	-	-	-	-	2	6	9	16
2002.....	196	-	-	-	-	-	-	2	-	2	14	7
2001.....	196	-	-	-	-	-	1	-	1	1	8	10
2000.....	159	-	-	-	-	-	1	1	3	2	6	11
1999.....	179	-	-	-	-	-	-	-	2	2	2	12
American Indian 1/, female												
2003.....	131	-	-	-	-	-	-	-	-	-	3	6
2002.....	122	-	-	-	-	-	-	-	1	-	2	1
2001.....	139	-	-	-	-	-	-	-	1	2	3	5
2000.....	107	-	-	-	-	-	-	-	-	1	3	2
1999.....	116	-	-	-	-	-	-	1	-	1	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Asian or Pacific Islander 2/, both sexes												
2003.....	1,191	-	-	-	-	-	1	3	3	11	30	61
2002.....	1,047	-	-	-	-	-	-	-	5	12	20	37
2001.....	1,033	-	-	-	-	-	-	-	6	6	23	40
2000.....	1,076	-	-	-	-	-	-	1	1	10	25	29
1999.....	1,097	-	-	-	-	-	-	3	3	4	20	37
Asian or Pacific Islander 2/, male												
2003.....	712	-	-	-	-	-	1	3	3	11	25	52
2002.....	649	-	-	-	-	-	-	-	5	11	20	32
2001.....	627	-	-	-	-	-	-	-	5	6	20	39
2000.....	655	-	-	-	-	-	-	1	1	8	21	26
1999.....	662	-	-	-	-	-	-	2	3	4	18	33
Asian or Pacific Islander 2/, female												
2003.....	479	-	-	-	-	-	-	-	-	-	5	9
2002.....	398	-	-	-	-	-	-	-	-	1	-	5
2001.....	406	-	-	-	-	-	-	-	1	-	3	1
2000.....	421	-	-	-	-	-	-	-	-	2	4	3
1999.....	435	-	-	-	-	-	-	1	-	-	2	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All races, both sexes												
2003.....	239,074	7	5	5	6	13	53	116	331	830	2,037	3,640
2002.....	243,332	8	2	1	10	9	41	117	280	902	2,028	3,637
2001.....	245,978	8	7	3	8	13	59	133	315	854	1,953	3,406
2000.....	249,389	8	8	3	6	15	33	91	309	908	2,011	3,421
1999.....	255,047	11	1	7	2	27	34	110	304	974	1,959	3,415
All races, male												
2003.....	119,164	5	3	2	2	9	35	80	245	631	1,517	2,741
2002.....	121,116	6	2	-	10	5	32	91	202	680	1,547	2,794
2001.....	120,970	5	1	2	6	8	41	110	244	643	1,490	2,600
2000.....	122,623	6	5	2	2	10	23	77	230	672	1,558	2,673
1999.....	124,801	6	1	6	2	20	20	79	231	720	1,525	2,679
All races, female												
2003.....	119,910	2	2	3	4	4	18	36	86	199	520	899
2002.....	122,216	2	-	1	-	4	9	26	78	222	481	843
2001.....	125,008	3	6	1	2	5	18	23	71	211	463	806
2000.....	126,766	2	3	1	4	5	10	14	79	236	453	748
1999.....	130,246	5	-	1	-	7	14	31	73	254	434	736

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, both sexes

2003.....	213,144	6	3	1	4	9	34	80	244	631	1,590	2,866
2002.....	217,772	3	-	1	7	5	28	87	208	691	1,610	2,843
2001.....	220,358	4	4	3	6	8	38	97	224	657	1,520	2,646
2000.....	224,183	6	8	3	5	13	21	66	224	687	1,566	2,666
1999.....	229,611	8	1	5	1	17	21	82	231	749	1,493	2,727

White, male

2003.....	106,696	4	2	1	2	6	22	56	185	494	1,217	2,209
2002.....	108,876	2	-	-	7	3	21	71	156	528	1,270	2,272
2001.....	108,714	3	-	2	5	4	29	85	173	502	1,197	2,090
2000.....	110,648	4	5	2	2	9	14	57	175	524	1,263	2,172
1999.....	112,638	4	1	5	1	12	14	58	179	561	1,199	2,214

White, female

2003.....	106,448	2	1	-	2	3	12	24	59	137	373	657
2002.....	108,896	1	-	1	-	2	7	16	52	163	340	571
2001.....	111,644	1	4	1	1	4	9	12	51	155	323	556
2000.....	113,535	2	3	1	3	4	7	9	49	163	303	494
1999.....	116,973	4	-	-	-	5	7	24	52	188	294	513

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, both sexes

2003.....	21,474	1	2	3	2	4	16	27	69	159	374	669
2002.....	21,211	4	2	-	3	4	9	24	53	178	366	690
2001.....	21,577	4	3	-	2	5	17	33	78	171	382	663
2000.....	21,279	1	-	-	-	2	9	22	76	192	395	664
1999.....	21,465	2	-	2	1	8	12	23	60	189	407	602

Black, male

2003.....	10,002	1	1	1	-	3	11	17	45	105	246	448
2002.....	9,801	3	2	-	3	2	7	16	30	123	234	439
2001.....	10,049	2	1	-	1	4	9	23	60	120	253	438
2000.....	9,707	1	-	-	-	1	6	17	49	121	254	427
1999.....	9,855	1	-	1	1	6	5	17	43	127	274	392

Black, female

2003.....	11,472	-	1	2	2	1	5	10	24	54	128	221
2002.....	11,410	1	-	-	-	2	2	8	23	55	132	251
2001.....	11,528	2	2	-	1	1	8	10	18	51	129	225
2000.....	11,572	-	-	-	-	1	3	5	27	71	141	237
1999.....	11,610	1	-	1	-	2	7	6	17	62	133	210

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
American Indian 1/, both sexes												
2003.....	804	-	-	-	-	-	2	-	8	12	23	31
2002.....	766	1	-	-	-	-	2	4	6	7	11	19
2001.....	668	-	-	-	-	-	2	-	1	5	14	29
2000.....	775	-	-	-	-	-	1	1	1	8	21	31
1999.....	714	1	-	-	-	-	1	2	3	11	16	24
American Indian 1/, male												
2003.....	461	-	-	-	-	-	1	-	6	10	15	20
2002.....	440	1	-	-	-	-	2	2	6	6	7	15
2001.....	367	-	-	-	-	-	2	-	1	2	9	22
2000.....	439	-	-	-	-	-	1	1	-	7	17	25
1999.....	396	1	-	-	-	-	1	2	-	8	12	19
American Indian 1/, female												
2003.....	343	-	-	-	-	-	1	-	2	2	8	11
2002.....	326	-	-	-	-	-	-	2	-	1	4	4
2001.....	301	-	-	-	-	-	-	-	-	3	5	7
2000.....	336	-	-	-	-	-	-	-	1	1	4	6
1999.....	318	-	-	-	-	-	-	-	3	3	4	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Asian or Pacific Islander 2/, both sexes												
2003.....	3,652	-	-	1	-	-	1	9	10	28	50	74
2002.....	3,583	-	-	-	-	-	2	2	13	26	41	85
2001.....	3,375	-	-	-	-	-	2	3	12	21	37	68
2000.....	3,152	1	-	-	1	-	2	2	8	21	29	60
1999.....	3,257	-	-	-	-	2	-	3	10	25	43	62
Asian or Pacific Islander 2/, male												
2003.....	2,005	-	-	-	-	-	1	7	9	22	39	64
2002.....	1,999	-	-	-	-	-	2	2	10	23	36	68
2001.....	1,840	-	-	-	-	-	1	2	10	19	31	50
2000.....	1,829	1	-	-	-	-	2	2	6	20	24	49
1999.....	1,912	-	-	-	-	2	-	2	9	24	40	54
Asian or Pacific Islander 2/, female												
2003.....	1,647	-	-	1	-	-	-	2	1	6	11	10
2002.....	1,584	-	-	-	-	-	-	-	3	3	5	17
2001.....	1,535	-	-	-	-	-	1	1	2	2	6	18
2000.....	1,323	-	-	-	1	-	-	-	2	1	5	11
1999.....	1,345	-	-	-	-	-	-	1	1	1	3	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
All races, both sexes												
2003.....	169,984	416	171	90	144	348	556	692	1,118	1,673	2,882	3,891
2002.....	169,540	473	157	84	143	362	474	723	1,085	1,769	2,851	3,849
2001.....	166,949	449	215	92	154	305	489	687	1,049	1,770	2,702	3,708
2000.....	165,428	457	168	96	152	355	498	687	1,062	1,706	2,751	3,536
1999.....	165,828	490	177	99	149	398	482	680	1,099	1,879	2,768	3,502
All races, male												
2003.....	74,738	233	81	50	91	219	341	404	677	1,017	1,784	2,439
2002.....	74,230	249	82	38	87	250	307	446	692	1,051	1,809	2,468
2001.....	72,439	222	117	41	82	193	307	413	624	1,099	1,678	2,361
2000.....	72,122	238	98	58	92	218	300	407	642	1,033	1,713	2,278
1999.....	72,601	255	100	54	83	250	283	400	667	1,176	1,718	2,223
All races, female												
2003.....	95,246	183	90	40	53	129	215	288	441	656	1,098	1,452
2002.....	95,310	224	75	46	56	112	167	277	393	718	1,042	1,381
2001.....	94,510	227	98	51	72	112	182	274	425	671	1,024	1,347
2000.....	93,306	219	70	38	60	137	198	280	420	673	1,038	1,258
1999.....	93,227	235	77	45	66	148	199	280	432	703	1,050	1,279

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
White, both sexes												
2003.....	147,127	256	109	57	98	235	355	425	717	1,106	1,972	2,736
2002.....	147,047	314	109	61	96	245	303	446	678	1,172	2,016	2,647
2001.....	144,862	292	134	64	107	203	309	438	681	1,166	1,806	2,561
2000.....	143,761	315	114	63	95	229	309	433	700	1,121	1,885	2,443
1999.....	143,786	329	119	72	103	281	314	425	729	1,252	1,875	2,390
White, male												
2003.....	63,973	140	48	31	64	147	215	252	457	686	1,269	1,763
2002.....	63,698	168	62	28	58	170	204	280	442	709	1,327	1,743
2001.....	62,025	155	71	29	59	124	195	268	408	736	1,187	1,687
2000.....	61,933	164	71	38	62	139	194	267	435	699	1,206	1,639
1999.....	62,054	174	70	39	56	170	182	259	453	805	1,216	1,580
White, female												
2003.....	83,154	116	61	26	34	88	140	173	260	420	703	973
2002.....	83,349	146	47	33	38	75	99	166	236	463	689	904
2001.....	82,837	137	63	35	48	79	114	170	273	430	619	874
2000.....	81,828	151	43	25	33	90	115	166	265	422	679	804
1999.....	81,732	155	49	33	47	111	132	166	276	447	659	810

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Black, both sexes												
2003.....	20,034	129	51	29	37	97	182	238	356	516	843	1,075
2002.....	19,812	136	38	16	41	99	154	252	363	549	773	1,113
2001.....	19,637	131	71	26	39	92	162	230	341	538	839	1,061
2000.....	19,298	117	48	28	45	113	167	236	328	526	801	1,023
1999.....	19,685	135	49	26	40	109	154	233	344	576	833	1,028
Black, male												
2003.....	9,372	78	28	16	22	64	116	136	189	301	474	626
2002.....	9,128	72	16	6	24	65	93	150	218	311	436	672
2001.....	9,148	59	40	11	21	61	101	135	195	319	454	612
2000.....	8,958	59	24	16	24	68	91	129	186	300	459	595
1999.....	9,344	67	27	14	24	76	91	132	199	337	468	595
Black, female												
2003.....	10,662	51	23	13	15	33	66	102	167	215	369	449
2002.....	10,684	64	22	10	17	34	61	102	145	238	337	441
2001.....	10,489	72	31	15	18	31	61	95	146	219	385	449
2000.....	10,340	58	24	12	21	45	76	107	142	226	342	428
1999.....	10,341	68	22	12	16	33	63	101	145	239	365	433

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
American Indian 1/, both sexes												
2003.....	676	9	4	-	2	4	5	7	14	19	25	31
2002.....	594	4	6	2	3	6	4	5	12	15	21	30
2001.....	599	9	3	2	1	2	5	9	7	28	24	38
2000.....	591	6	1	1	4	2	5	4	9	25	22	25
1999.....	595	5	5	-	3	2	6	5	7	19	24	30
American Indian 1/, male												
2003.....	333	5	1	-	-	2	4	3	13	12	15	20
2002.....	328	2	2	1	3	5	3	2	9	12	16	21
2001.....	318	3	2	1	-	1	4	4	5	18	15	30
2000.....	295	3	-	1	2	2	3	2	5	12	15	15
1999.....	289	2	2	-	1	1	4	3	2	13	11	17
American Indian 1/, female												
2003.....	343	4	3	-	2	2	1	4	1	7	10	11
2002.....	266	2	4	1	-	1	1	3	3	3	5	9
2001.....	281	6	1	1	1	1	1	5	2	10	9	8
2000.....	296	3	1	-	2	-	2	2	4	13	7	10
1999.....	306	3	3	-	2	1	2	2	5	6	13	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Asian or Pacific Islander 2/, both sexes												
2003.....	2,147	22	7	4	7	12	14	22	31	32	42	49
2002.....	2,087	19	4	5	3	12	13	20	32	33	41	59
2001.....	1,851	17	7	-	7	8	13	10	20	38	33	48
2000.....	1,778	19	5	4	8	11	17	14	25	34	43	45
1999.....	1,762	21	4	1	3	6	8	17	19	32	36	54
Asian or Pacific Islander 2/, male												
2003.....	1,060	10	4	3	5	6	6	13	18	18	26	30
2002.....	1,076	7	2	3	2	10	7	14	23	19	30	32
2001.....	948	5	4	-	2	7	7	6	16	26	22	32
2000.....	936	12	3	3	4	9	12	9	16	22	33	29
1999.....	914	12	1	1	2	3	6	6	13	21	23	31
Asian or Pacific Islander 2/, female												
2003.....	1,087	12	3	1	2	6	8	9	13	14	16	19
2002.....	1,011	12	2	2	1	2	6	6	9	14	11	27
2001.....	903	12	3	-	5	1	6	4	4	12	11	16
2000.....	842	7	2	1	4	2	5	5	9	12	10	16
1999.....	848	9	3	-	1	3	2	11	6	11	13	23

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
All races, both sexes												
2003.....	1,245	1	-	-	1	3	13	18	15	43	69	89
2002.....	1,154	2	2	1	1	2	5	16	28	26	64	84
2001.....	1,178	1	-	1	2	3	5	9	20	46	71	93
2000.....	1,172	3	1	2	1	5	8	13	27	41	75	89
1999.....	1,116	2	1	-	-	2	5	12	18	54	60	68
All races, male												
2003.....	682	1	-	-	1	-	6	11	8	22	41	60
2002.....	643	1	1	-	1	2	4	9	22	19	41	55
2001.....	668	1	-	-	1	2	3	5	12	25	37	68
2000.....	665	2	-	1	-	1	6	7	17	24	49	59
1999.....	639	2	1	-	-	-	3	9	15	34	41	49
All races, female												
2003.....	563	-	-	-	-	3	7	7	7	21	28	29
2002.....	511	1	1	1	-	-	1	7	6	7	23	29
2001.....	510	-	-	1	1	1	2	4	8	21	34	25
2000.....	507	1	1	1	1	4	2	6	10	17	26	30
1999.....	477	-	-	-	-	2	2	3	3	20	19	19

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
White, both sexes												
2003.....	990	-	-	-	1	2	12	14	11	27	51	57
2002.....	930	1	2	-	-	2	4	11	23	20	46	49
2001.....	917	1	-	1	2	1	5	8	13	30	44	64
2000.....	927	1	1	1	1	1	5	10	17	27	53	57
1999.....	879	1	1	-	-	2	4	10	14	32	37	41
White, male												
2003.....	541	-	-	-	1	-	6	9	5	14	32	36
2002.....	524	-	1	-	-	2	3	7	19	14	31	34
2001.....	517	1	-	-	1	1	3	4	8	16	25	45
2000.....	536	1	-	-	-	-	3	6	13	14	36	39
1999.....	509	1	1	-	-	-	2	7	11	21	24	32
White, female												
2003.....	449	-	-	-	-	2	6	5	6	13	19	21
2002.....	406	1	1	-	-	-	1	4	4	6	15	15
2001.....	400	-	-	1	1	-	2	4	5	14	19	19
2000.....	391	-	1	1	1	1	2	4	4	13	17	18
1999.....	370	-	-	-	-	2	2	3	3	11	13	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Black, both sexes												
2003.....	225	1	-	-	-	1	1	2	3	15	17	29
2002.....	199	-	-	1	-	-	1	5	5	6	17	29
2001.....	235	-	-	-	-	1	-	1	7	15	26	27
2000.....	214	2	-	-	-	3	3	3	8	11	20	30
1999.....	210	-	-	-	-	-	1	2	4	20	21	26
Black, male												
2003.....	122	1	-	-	-	-	-	2	2	7	8	21
2002.....	106	-	-	-	-	-	1	2	3	5	10	18
2001.....	135	-	-	-	-	-	-	1	4	8	12	21
2000.....	111	1	-	-	-	1	3	1	2	9	12	20
1999.....	117	-	-	-	-	-	1	2	4	12	15	16
Black, female												
2003.....	103	-	-	-	-	1	1	-	1	8	9	8
2002.....	93	-	-	1	-	-	-	3	2	1	7	11
2001.....	100	-	-	-	-	1	-	-	3	7	14	6
2000.....	103	1	-	-	-	2	-	2	6	2	8	10
1999.....	93	-	-	-	-	-	-	-	-	8	6	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute and subacute endocarditis (I33)

American Indian 1/, both sexes

2003.....	12	-	-	-	-	-	-	1	1	1	1	2
2002.....	6	-	-	-	-	-	-	-	-	-	-	3
2001.....	6	-	-	-	-	1	-	-	-	-	-	2
2000.....	13	-	-	-	-	-	-	-	-	1	2	2
1999.....	7	-	-	-	-	-	-	-	-	2	1	-

American Indian 1/, male

2003.....	7	-	-	-	-	-	-	-	1	1	1	2
2002.....	2	-	-	-	-	-	-	-	-	-	-	1
2001.....	5	-	-	-	-	1	-	-	-	-	-	2
2000.....	7	-	-	-	-	-	-	-	-	1	1	-
1999.....	4	-	-	-	-	-	-	-	-	1	1	-

American Indian 1/, female

2003.....	5	-	-	-	-	-	-	1	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	2
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	1	2
1999.....	3	-	-	-	-	-	-	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All races, both sexes												
2003.....	868	33	22	14	14	21	34	23	25	32	54	63
2002.....	848	26	22	12	10	17	19	16	30	35	40	56
2001.....	839	32	26	12	9	14	26	20	30	43	50	57
2000.....	767	20	24	9	11	17	21	29	24	45	41	45
1999.....	890	27	25	11	10	12	20	19	31	54	60	59
All races, male												
2003.....	449	15	8	4	8	14	20	13	16	21	31	41
2002.....	434	9	11	4	3	12	12	9	23	17	32	40
2001.....	401	18	15	2	5	7	15	16	17	24	17	30
2000.....	404	10	13	4	5	10	11	17	12	30	28	28
1999.....	442	13	14	5	2	8	12	14	20	32	32	31
All races, female												
2003.....	419	18	14	10	6	7	14	10	9	11	23	22
2002.....	414	17	11	8	7	5	7	7	7	18	8	16
2001.....	438	14	11	10	4	7	11	4	13	19	33	27
2000.....	363	10	11	5	6	7	10	12	12	15	13	17
1999.....	448	14	11	6	8	4	8	5	11	22	28	28

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
White, both sexes												
2003.....	677	15	13	9	8	17	21	10	14	17	38	46
2002.....	666	13	13	8	4	10	16	14	19	22	33	43
2001.....	658	14	14	9	7	10	17	15	23	28	32	43
2000.....	590	11	15	8	2	15	10	21	13	27	31	26
1999.....	703	12	12	7	6	9	15	13	22	41	38	47
White, male												
2003.....	359	9	4	2	5	13	12	5	8	13	23	31
2002.....	344	6	7	2	1	7	10	8	16	9	27	29
2001.....	310	10	8	2	4	3	9	12	13	15	14	21
2000.....	309	6	11	4	1	8	5	11	7	18	21	17
1999.....	351	7	5	4	2	7	8	10	14	24	23	24
White, female												
2003.....	318	6	9	7	3	4	9	5	6	4	15	15
2002.....	322	7	6	6	3	3	6	6	3	13	6	14
2001.....	348	4	6	7	3	7	8	3	10	13	18	22
2000.....	281	5	4	4	1	7	5	10	6	9	10	9
1999.....	352	5	7	3	4	2	7	3	8	17	15	23

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Black, both sexes												
2003.....	168	14	9	4	5	3	12	12	10	14	16	16
2002.....	154	12	7	2	6	5	3	1	9	10	7	12
2001.....	158	14	12	3	1	4	9	5	5	10	15	14
2000.....	142	7	8	1	7	1	7	6	10	14	10	14
1999.....	160	10	12	4	4	3	3	6	8	12	19	9
Black, male												
2003.....	81	4	4	2	3	1	8	8	7	8	8	9
2002.....	75	3	3	1	2	3	2	-	6	5	5	10
2001.....	79	5	7	-	1	4	6	4	3	5	3	9
2000.....	72	4	2	-	3	1	2	5	5	9	7	7
1999.....	77	4	9	1	-	1	2	4	6	7	7	5
Black, female												
2003.....	87	10	5	2	2	2	4	4	3	6	8	7
2002.....	79	9	4	1	4	2	1	1	3	5	2	2
2001.....	79	9	5	3	-	-	3	1	2	5	12	5
2000.....	70	3	6	1	4	-	5	1	5	5	3	7
1999.....	83	6	3	3	4	2	1	2	2	5	12	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

American Indian 1/, both sexes

2003.....	6	3	-	-	-	-	-	-	-	-	-	-
2002.....	8	-	1	1	-	1	-	-	-	1	-	1
2001.....	8	3	-	-	1	-	-	-	-	2	1	-
2000.....	6	1	-	-	-	-	1	-	-	-	-	1
1999.....	9	1	1	-	-	-	1	-	1	-	1	1

American Indian 1/, male

2003.....	5	2	-	-	-	-	-	-	-	-	-	-
2002.....	5	-	-	1	-	1	-	-	-	1	-	1
2001.....	5	2	-	-	-	-	-	-	-	2	-	-
2000.....	4	-	-	-	-	-	1	-	-	-	-	1
1999.....	2	-	-	-	-	-	1	-	-	-	-	1

American Indian 1/, female

2003.....	1	1	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	1	-	-	-	-	-	-	-	-	-
2001.....	3	1	-	-	1	-	-	-	-	-	1	-
2000.....	2	1	-	-	-	-	-	-	-	-	-	-
1999.....	7	1	1	-	-	-	-	-	1	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
All races, both sexes												
2003.....	57,448	38	5	5	7	18	20	28	69	91	224	358
2002.....	56,494	23	11	3	7	10	17	33	68	102	180	355
2001.....	56,934	24	17	8	7	19	15	41	51	115	192	353
2000.....	55,704	31	9	10	11	21	13	27	36	77	185	283
1999.....	54,913	11	10	5	7	14	17	27	38	88	184	298
All races, male												
2003.....	22,427	28	3	4	3	13	11	16	44	50	143	223
2002.....	21,698	14	6	1	4	6	9	22	38	56	117	210
2001.....	21,632	16	5	5	3	11	5	25	31	76	106	227
2000.....	21,175	22	7	4	5	11	9	15	21	40	119	191
1999.....	20,755	5	7	3	3	12	11	13	19	52	112	186
All races, female												
2003.....	35,021	10	2	1	4	5	9	12	25	41	81	135
2002.....	34,796	9	5	2	3	4	8	11	30	46	63	145
2001.....	35,302	8	12	3	4	8	10	16	20	39	86	126
2000.....	34,529	9	2	6	6	10	4	12	15	37	66	92
1999.....	34,158	6	3	2	4	2	6	14	19	36	72	112

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
White, both sexes												
2003.....	51,454	20	4	3	4	9	11	19	39	56	122	229
2002.....	50,672	15	8	3	4	5	11	16	37	61	113	211
2001.....	51,252	19	15	5	6	13	5	25	32	76	112	237
2000.....	50,142	19	5	7	9	12	5	15	23	44	120	180
1999.....	49,317	8	8	3	6	9	9	13	21	58	116	183
White, male												
2003.....	19,987	13	2	2	2	5	7	13	25	30	86	151
2002.....	19,335	10	5	1	3	2	5	11	19	34	78	130
2001.....	19,332	12	5	3	2	9	1	12	19	49	69	166
2000.....	18,949	14	5	3	4	5	5	9	13	25	84	127
1999.....	18,469	3	5	2	3	8	6	6	11	34	77	116
White, female												
2003.....	31,467	7	2	1	2	4	4	6	14	26	36	78
2002.....	31,337	5	3	2	1	3	6	5	18	27	35	81
2001.....	31,920	7	10	2	4	4	4	13	13	27	43	71
2000.....	31,193	5	-	4	5	7	-	6	10	19	36	53
1999.....	30,848	5	3	1	3	1	3	7	10	24	39	67

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Black, both sexes												
2003.....	5,330	15	-	2	3	9	9	8	29	34	97	124
2002.....	5,186	8	3	-	3	3	6	16	29	40	64	140
2001.....	5,122	5	2	3	-	6	10	16	18	35	77	111
2000.....	4,965	12	4	2	2	8	5	12	12	32	62	98
1999.....	4,998	3	2	2	1	4	8	13	17	28	66	111
Black, male												
2003.....	2,150	13	-	2	1	8	4	2	18	20	53	68
2002.....	2,069	4	1	-	1	2	4	10	17	22	37	78
2001.....	2,052	4	-	2	-	2	4	13	11	26	35	57
2000.....	1,951	8	2	1	1	5	3	6	7	14	33	60
1999.....	2,021	2	2	1	-	3	5	7	8	16	35	68
Black, female												
2003.....	3,180	2	-	-	2	1	5	6	11	14	44	56
2002.....	3,117	4	2	-	2	1	2	6	12	18	27	62
2001.....	3,070	1	2	1	-	4	6	3	7	9	42	54
2000.....	3,014	4	2	1	1	3	2	6	5	18	29	38
1999.....	2,977	1	-	1	1	1	3	6	9	12	31	43

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Asian or Pacific Islander 2/, both sexes												
2003.....	484	2	1	-	-	-	-	1	-	1	3	2
2002.....	452	-	-	-	-	1	-	-	1	-	1	2
2001.....	396	-	-	-	1	-	-	-	1	2	2	2
2000.....	422	-	-	1	-	1	2	-	1	-	3	3
1999.....	428	-	-	-	-	-	-	1	-	-	-	2
Asian or Pacific Islander 2/, male												
2003.....	215	1	1	-	-	-	-	1	-	-	3	2
2002.....	202	-	-	-	-	1	-	-	1	-	-	-
2001.....	177	-	-	-	1	-	-	-	1	1	2	1
2000.....	196	-	-	-	-	1	-	-	1	-	2	3
1999.....	200	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, female												
2003.....	269	1	-	-	-	-	-	-	-	1	-	-
2002.....	250	-	-	-	-	-	-	-	-	-	1	2
2001.....	219	-	-	-	-	-	-	-	-	1	-	1
2000.....	226	-	-	1	-	-	2	-	-	-	1	-
1999.....	228	-	-	-	-	-	-	1	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All races, both sexes												
2003.....	110,423	344	144	71	122	306	489	623	1,009	1,507	2,535	3,381
2002.....	111,044	422	122	68	125	333	433	658	959	1,606	2,567	3,354
2001.....	107,998	392	172	71	136	269	443	617	948	1,566	2,389	3,205
2000.....	107,785	403	134	75	129	312	456	618	975	1,543	2,450	3,119
1999.....	108,909	450	141	83	132	370	440	622	1,012	1,683	2,464	3,077
All races, male												
2003.....	51,180	189	70	42	79	192	304	364	609	924	1,569	2,115
2002.....	51,455	225	64	33	79	230	282	406	609	959	1,619	2,163
2001.....	49,738	187	97	34	73	173	284	367	564	974	1,518	2,036
2000.....	49,878	204	78	49	82	196	274	368	592	939	1,517	2,000
1999.....	50,765	235	78	46	78	230	257	364	613	1,058	1,533	1,957
All races, female												
2003.....	59,243	155	74	29	43	114	185	259	400	583	966	1,266
2002.....	59,589	197	58	35	46	103	151	252	350	647	948	1,191
2001.....	58,260	205	75	37	63	96	159	250	384	592	871	1,169
2000.....	57,907	199	56	26	47	116	182	250	383	604	933	1,119
1999.....	58,144	215	63	37	54	140	183	258	399	625	931	1,120

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, both sexes

2003.....	94,006	221	92	45	85	207	311	382	653	1,006	1,761	2,404
2002.....	94,779	285	86	50	88	228	272	405	599	1,069	1,824	2,344
2001.....	92,035	258	105	49	92	179	282	390	613	1,032	1,618	2,217
2000.....	92,102	284	93	47	83	201	289	387	647	1,023	1,681	2,180
1999.....	92,887	308	98	62	91	261	286	389	672	1,121	1,684	2,119

White, male

2003.....	43,086	118	42	27	56	129	190	225	419	629	1,128	1,545
2002.....	43,495	152	49	25	54	159	186	254	388	652	1,191	1,550
2001.....	41,866	132	58	24	52	111	182	240	368	656	1,079	1,455
2000.....	42,139	143	55	31	57	126	181	241	402	642	1,065	1,456
1999.....	42,725	163	59	33	51	155	166	236	417	726	1,092	1,408

White, female

2003.....	50,920	103	50	18	29	78	121	157	234	377	633	859
2002.....	51,284	133	37	25	34	69	86	151	211	417	633	794
2001.....	50,169	126	47	25	40	68	100	150	245	376	539	762
2000.....	49,963	141	38	16	26	75	108	146	245	381	616	724
1999.....	50,162	145	39	29	40	106	120	153	255	395	592	711

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes

2003.....	14,311	99	42	23	29	84	160	216	314	453	713	906
2002.....	14,273	116	28	13	32	91	144	230	320	493	685	932
2001.....	14,122	112	57	20	38	81	143	208	311	478	721	909
2000.....	13,977	96	36	25	36	101	152	215	298	469	709	881
1999.....	14,317	122	35	20	35	102	142	212	315	516	727	882

Black, male

2003.....	7,019	60	24	12	18	55	104	124	162	266	405	528
2002.....	6,878	65	12	5	21	60	86	138	192	279	384	566
2001.....	6,882	50	33	9	20	55	91	117	177	280	404	525
2000.....	6,824	46	20	15	20	61	83	117	172	268	407	508
1999.....	7,129	61	16	12	24	72	83	119	181	302	411	506

Black, female

2003.....	7,292	39	18	11	11	29	56	92	152	187	308	378
2002.....	7,395	51	16	8	11	31	58	92	128	214	301	366
2001.....	7,240	62	24	11	18	26	52	91	134	198	317	384
2000.....	7,153	50	16	10	16	40	69	98	126	201	302	373
1999.....	7,188	61	19	8	11	30	59	93	134	214	316	376

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
American Indian 1/, both sexes												
2003.....	478	5	4	-	2	4	5	6	12	18	22	26
2002.....	396	4	5	1	3	4	4	4	11	13	19	24
2001.....	421	6	3	2	-	1	5	9	7	24	22	33
2000.....	397	5	1	1	4	2	3	4	9	23	20	20
1999.....	409	4	4	-	3	1	5	5	6	15	20	27
American Indian 1/, male												
2003.....	246	2	1	-	-	2	4	3	11	11	13	16
2002.....	229	2	2	-	3	3	3	1	8	11	14	17
2001.....	237	1	2	1	-	-	4	4	5	16	15	25
2000.....	205	3	-	1	2	2	1	2	5	10	14	13
1999.....	218	2	2	-	1	-	3	3	2	10	10	15
American Indian 1/, female												
2003.....	232	3	3	-	2	2	1	3	1	7	9	10
2002.....	167	2	3	1	-	1	1	3	3	2	5	7
2001.....	184	5	1	1	-	1	1	5	2	8	7	8
2000.....	192	2	1	-	2	-	2	2	4	13	6	7
1999.....	191	2	2	-	2	1	2	2	4	5	10	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Asian or Pacific Islander 2/, both sexes

2003.....	1,628	19	6	3	6	11	13	19	30	30	39	45
2002.....	1,596	17	3	4	2	10	13	19	29	31	39	54
2001.....	1,420	16	7	-	6	8	13	10	17	32	28	46
2000.....	1,309	18	4	2	6	8	12	12	21	28	40	38
1999.....	1,296	16	4	1	3	6	7	16	19	31	33	49

Asian or Pacific Islander 2/, male

2003.....	829	9	3	3	5	6	6	12	17	18	23	26
2002.....	853	6	1	3	1	8	7	13	21	17	30	30
2001.....	753	4	4	-	1	7	7	6	14	22	20	31
2000.....	710	12	3	2	3	7	9	8	13	19	31	23
1999.....	693	9	1	1	2	3	5	6	13	20	20	28

Asian or Pacific Islander 2/, female

2003.....	799	10	3	-	1	5	7	7	13	12	16	19
2002.....	743	11	2	1	1	2	6	6	8	14	9	24
2001.....	667	12	3	-	5	1	6	4	3	10	8	15
2000.....	599	6	1	-	3	1	3	4	8	9	9	15
1999.....	603	7	3	-	1	3	2	10	6	11	13	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races, both sexes

2003.....	21,940	1	2	-	2	7	15	27	71	126	246	400
2002.....	20,261	5	1	1	-	4	17	36	49	126	228	382
2001.....	19,250	-	1	-	1	5	18	32	72	101	230	385
2000.....	18,073	1	1	2	2	6	11	21	49	105	243	352
1999.....	16,968	3	3	-	1	3	10	24	54	105	208	329

All races, male

2003.....	8,213	1	2	-	1	5	7	9	49	72	143	265
2002.....	7,647	2	1	1	-	-	11	21	26	75	144	238
2001.....	7,254	-	-	-	-	2	14	20	39	59	147	239
2000.....	6,895	1	-	1	2	-	4	12	27	61	155	224
1999.....	6,488	2	1	-	1	2	6	13	35	69	126	196

All races, female

2003.....	13,727	-	-	-	1	2	8	18	22	54	103	135
2002.....	12,614	3	-	-	-	4	6	15	23	51	84	144
2001.....	11,996	-	1	-	1	3	4	12	33	42	83	146
2000.....	11,178	-	1	1	-	6	7	9	22	44	88	128
1999.....	10,480	1	2	-	-	1	4	11	19	36	82	133

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

White, both sexes

2003.....	16,927	-	1	-	1	5	7	12	31	58	128	208
2002.....	15,587	4	1	1	-	3	11	15	20	44	112	185
2001.....	14,752	-	1	-	1	4	10	13	32	45	95	183
2000.....	13,775	1	1	1	2	2	5	12	20	41	117	169
1999.....	12,830	2	1	-	-	3	4	12	24	37	94	172

White, male

2003.....	6,108	-	1	-	-	3	4	4	17	39	79	144
2002.....	5,719	2	1	1	-	-	8	10	11	24	76	123
2001.....	5,351	-	-	-	-	2	7	8	20	27	66	119
2000.....	5,120	1	-	1	2	-	3	8	14	30	79	117
1999.....	4,744	1	-	-	-	2	1	7	20	29	63	109

White, female

2003.....	10,819	-	-	-	1	2	3	8	14	19	49	64
2002.....	9,868	2	-	-	-	3	3	5	9	20	36	62
2001.....	9,401	-	1	-	1	2	3	5	12	18	29	64
2000.....	8,655	-	1	-	-	2	2	4	6	11	38	52
1999.....	8,086	1	1	-	-	1	3	5	4	8	31	63

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Black, both sexes

2003.....	4,489	1	1	-	-	2	8	15	38	63	113	186
2002.....	4,196	1	-	-	-	1	6	18	27	80	111	189
2001.....	4,038	-	-	-	-	1	8	18	37	53	126	191
2000.....	3,887	-	-	1	-	4	6	7	28	62	117	171
1999.....	3,732	1	1	-	1	-	6	11	27	66	111	145

Black, male

2003.....	1,895	1	1	-	-	2	3	5	30	31	62	120
2002.....	1,717	-	-	-	-	-	3	10	14	49	64	109
2001.....	1,715	-	-	-	-	-	7	12	17	30	74	115
2000.....	1,592	-	-	-	-	-	1	3	13	30	70	99
1999.....	1,565	1	1	-	1	-	5	5	13	38	60	83

Black, female

2003.....	2,594	-	-	-	-	-	5	10	8	32	51	66
2002.....	2,479	1	-	-	-	1	3	8	13	31	47	80
2001.....	2,323	-	-	-	-	1	1	6	20	23	52	76
2000.....	2,295	-	-	1	-	4	5	4	15	32	47	72
1999.....	2,167	-	-	-	-	-	1	6	14	28	51	62

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

American Indian 1/, both sexes

2003.....	88	-	-	-	-	-	-	-	-	1	2	2
2002.....	76	-	-	-	-	-	-	1	-	1	3	2
2001.....	63	-	-	-	-	-	-	1	1	2	3	1
2000.....	73	-	-	-	-	-	-	1	-	1	2	2
1999.....	55	-	-	-	-	-	-	1	1	1	2	1

American Indian 1/, male

2003.....	39	-	-	-	-	-	-	-	-	-	-	-
2002.....	30	-	-	-	-	-	-	-	-	1	3	1
2001.....	25	-	-	-	-	-	-	-	1	1	2	-
2000.....	24	-	-	-	-	-	-	-	-	-	2	1
1999.....	25	-	-	-	-	-	-	1	-	1	2	1

American Indian 1/, female

2003.....	49	-	-	-	-	-	-	-	-	1	2	2
2002.....	46	-	-	-	-	-	-	1	-	-	-	1
2001.....	38	-	-	-	-	-	-	1	-	1	1	1
2000.....	49	-	-	-	-	-	-	1	-	1	-	1
1999.....	30	-	-	-	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
All races, both sexes												
2003.....	157,689	101	46	29	40	88	133	188	395	761	1,699	2,577
2002.....	162,672	117	53	33	58	53	118	183	384	797	1,628	2,615
2001.....	163,538	108	54	38	42	68	128	199	402	840	1,651	2,507
2000.....	167,661	124	47	25	51	67	132	209	393	880	1,719	2,625
1999.....	167,366	103	46	26	39	67	115	202	378	916	1,658	2,455
All races, male												
2003.....	61,426	57	25	19	22	51	65	101	213	411	872	1,389
2002.....	62,622	67	30	16	31	34	69	100	184	389	789	1,402
2001.....	63,177	63	25	21	21	36	67	105	206	435	832	1,354
2000.....	64,769	74	19	12	21	38	68	111	197	430	867	1,401
1999.....	64,485	65	22	17	17	36	59	102	197	449	859	1,282
All races, female												
2003.....	96,263	44	21	10	18	37	68	87	182	350	827	1,188
2002.....	100,050	50	23	17	27	19	49	83	200	408	839	1,213
2001.....	100,361	45	29	17	21	32	61	94	196	405	819	1,153
2000.....	102,892	50	28	13	30	29	64	98	196	450	852	1,224
1999.....	102,881	38	24	9	22	31	56	100	181	467	799	1,173

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
White, both sexes												
2003.....	134,705	63	30	21	30	63	89	126	262	457	1,051	1,611
2002.....	139,719	71	40	25	38	38	92	124	252	509	1,038	1,671
2001.....	140,465	60	39	29	30	53	85	133	283	553	1,052	1,597
2000.....	144,580	75	30	21	34	51	97	160	260	558	1,087	1,627
1999.....	144,827	70	34	23	26	46	89	145	257	602	1,044	1,535
White, male												
2003.....	51,646	32	14	13	18	36	51	66	144	245	557	903
2002.....	52,959	41	22	11	18	23	56	75	120	254	519	930
2001.....	53,428	35	18	17	17	28	53	68	152	281	553	876
2000.....	54,938	46	14	12	14	33	51	89	136	282	560	894
1999.....	54,867	43	16	16	11	28	44	73	140	300	556	802
White, female												
2003.....	83,059	31	16	8	12	27	38	60	118	212	494	708
2002.....	86,760	30	18	14	20	15	36	49	132	255	519	741
2001.....	87,037	25	21	12	13	25	32	65	131	272	499	721
2000.....	89,642	29	16	9	20	18	46	71	124	276	527	733
1999.....	89,960	27	18	7	15	18	45	72	117	302	488	733

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

Black, both sexes

2003.....	18,806	32	15	6	10	21	38	54	114	250	543	828
2002.....	18,856	37	11	7	17	13	25	51	117	238	506	818
2001.....	19,002	43	12	6	8	13	38	59	99	236	529	802
2000.....	19,221	46	14	2	13	13	30	41	119	283	553	878
1999.....	18,884	28	10	3	12	18	23	50	100	268	548	787

Black, male

2003.....	7,866	20	10	5	4	13	13	33	54	130	257	419
2002.....	7,828	20	7	4	12	10	12	21	53	102	230	407
2001.....	7,907	27	5	2	3	8	12	35	44	121	240	418
2000.....	8,026	27	3	-	5	4	15	18	53	127	270	449
1999.....	7,894	18	4	1	6	7	12	24	48	124	269	412

Black, female

2003.....	10,940	12	5	1	6	8	25	21	60	120	286	409
2002.....	11,028	17	4	3	5	3	13	30	64	136	276	411
2001.....	11,095	16	7	4	5	5	26	24	55	115	289	384
2000.....	11,195	19	11	2	8	9	15	23	66	156	283	429
1999.....	10,990	10	6	2	6	11	11	26	52	144	279	375

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

American Indian 1/, both sexes

2003.....	552	-	1	1	-	1	-	2	3	13	21	26
2002.....	567	1	-	-	2	2	-	2	3	8	25	27
2001.....	574	2	-	1	-	-	-	1	4	11	19	27
2000.....	572	-	1	1	1	-	1	3	2	7	15	22
1999.....	546	2	-	-	1	-	1	-	11	9	21	21

American Indian 1/, male

2003.....	242	-	1	-	-	1	-	1	3	10	8	15
2002.....	236	1	-	-	-	1	-	1	3	3	11	14
2001.....	217	-	-	1	-	-	-	-	1	4	9	9
2000.....	250	-	1	-	-	-	-	-	-	4	7	10
1999.....	236	2	-	-	-	-	1	-	4	4	11	10

American Indian 1/, female

2003.....	310	-	-	1	-	-	-	1	-	3	13	11
2002.....	331	-	-	-	2	1	-	1	-	5	14	13
2001.....	357	2	-	-	-	-	-	1	3	7	10	18
2000.....	322	-	-	1	1	-	1	3	2	3	8	12
1999.....	310	-	-	-	1	-	-	-	7	5	10	11

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

Asian or Pacific Islander 2/, both sexes

2003.....	3,626	6	-	1	-	3	6	6	16	41	84	112
2002.....	3,530	8	2	1	1	-	1	6	12	42	59	99
2001.....	3,497	3	3	2	4	2	5	6	16	40	51	81
2000.....	3,288	3	2	1	3	3	4	5	12	32	64	98
1999.....	3,109	3	2	-	-	3	2	7	10	37	45	112

Asian or Pacific Islander 2/, male

2003.....	1,672	5	-	1	-	1	1	1	12	26	50	52
2002.....	1,599	5	1	1	1	-	1	3	8	30	29	51
2001.....	1,625	1	2	1	1	-	2	2	9	29	30	51
2000.....	1,555	1	1	-	2	1	2	4	8	17	30	48
1999.....	1,488	2	2	-	-	1	2	5	5	21	23	58

Asian or Pacific Islander 2/, female

2003.....	1,954	1	-	-	-	2	5	5	4	15	34	60
2002.....	1,931	3	1	-	-	-	-	3	4	12	30	48
2001.....	1,872	2	1	1	3	2	3	4	7	11	21	30
2000.....	1,733	2	1	1	1	2	2	1	4	15	34	50
1999.....	1,621	1	-	-	-	2	-	2	5	16	22	54

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
All races, both sexes												
2003.....	13,053	3	-	-	-	1	2	-	7	11	36	70
2002.....	13,821	4	-	-	-	-	3	2	7	16	34	61
2001.....	14,086	1	1	1	-	-	1	1	2	14	26	65
2000.....	14,393	2	-	-	-	-	3	3	5	16	32	71
1999.....	14,979	2	-	-	-	1	1	1	7	16	29	56
All races, male												
2003.....	4,859	2	-	-	-	1	1	-	5	8	27	54
2002.....	5,171	1	-	-	-	-	2	2	3	9	21	44
2001.....	5,143	1	1	1	-	-	1	-	2	8	18	43
2000.....	5,357	-	-	-	-	-	2	1	5	10	25	48
1999.....	5,443	1	-	-	-	-	-	1	4	11	21	37
All races, female												
2003.....	8,194	1	-	-	-	-	1	-	2	3	9	16
2002.....	8,650	3	-	-	-	-	1	-	4	7	13	17
2001.....	8,943	-	-	-	-	-	-	1	-	6	8	22
2000.....	9,036	2	-	-	-	-	1	2	-	6	7	23
1999.....	9,536	1	-	-	-	1	1	-	3	5	8	19

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
White, both sexes												
2003.....	11,885	1	-	-	-	-	2	-	5	7	32	58
2002.....	12,631	3	-	-	-	-	3	2	4	14	26	44
2001.....	12,961	1	1	1	-	-	-	-	2	11	22	54
2000.....	13,182	2	-	-	-	-	3	1	4	12	24	50
1999.....	13,737	1	-	-	-	1	1	1	5	14	19	48
White, male												
2003.....	4,368	-	-	-	-	-	1	-	4	5	23	43
2002.....	4,688	1	-	-	-	-	2	2	2	7	17	34
2001.....	4,696	1	1	1	-	-	-	-	2	5	16	35
2000.....	4,875	-	-	-	-	-	2	1	4	9	19	36
1999.....	4,932	1	-	-	-	-	-	1	3	10	14	32
White, female												
2003.....	7,517	1	-	-	-	-	1	-	1	2	9	15
2002.....	7,943	2	-	-	-	-	1	-	2	7	9	10
2001.....	8,265	-	-	-	-	-	-	-	-	6	6	19
2000.....	8,307	2	-	-	-	-	1	-	-	3	5	14
1999.....	8,805	-	-	-	-	1	1	-	2	4	5	16

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Atherosclerosis (I70)

Black, both sexes

2003.....	995	2	-	-	-	1	-	-	2	2	3	11
2002.....	1,017	1	-	-	-	-	-	-	3	2	7	15
2001.....	970	-	-	-	-	-	1	1	-	2	4	11
2000.....	1,054	-	-	-	-	-	-	2	1	4	8	19
1999.....	1,069	1	-	-	-	-	-	-	2	2	7	6

Black, male

2003.....	410	2	-	-	-	1	-	-	1	1	3	10
2002.....	414	-	-	-	-	-	-	-	1	2	4	8
2001.....	370	-	-	-	-	-	1	-	-	2	2	8
2000.....	415	-	-	-	-	-	-	-	1	1	6	11
1999.....	432	-	-	-	-	-	-	-	1	1	5	4

Black, female

2003.....	585	-	-	-	-	-	-	-	1	1	-	1
2002.....	603	1	-	-	-	-	-	-	2	-	3	7
2001.....	600	-	-	-	-	-	-	1	-	-	2	3
2000.....	639	-	-	-	-	-	-	2	-	3	2	8
1999.....	637	1	-	-	-	-	-	-	1	1	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of circulatory system (I71-I78)

All races, both sexes

2003.....	24,672	11	-	1	11	30	38	62	89	167	284	524
2002.....	24,927	14	4	1	10	31	35	51	109	154	291	479
2001.....	25,318	12	4	2	9	23	38	67	103	194	304	421
2000.....	26,036	16	5	3	8	17	42	49	105	157	291	424
1999.....	25,809	12	2	2	9	14	35	51	90	178	300	418

All races, male

2003.....	13,217	3	-	-	7	17	28	43	60	117	189	343
2002.....	13,309	6	4	-	5	19	24	30	71	81	210	331
2001.....	13,396	8	2	-	3	17	23	43	72	132	203	291
2000.....	14,084	6	2	1	6	9	24	36	67	92	184	275
1999.....	13,794	9	-	1	6	10	22	35	67	117	179	289

All races, female

2003.....	11,455	8	-	1	4	13	10	19	29	50	95	181
2002.....	11,618	8	-	1	5	12	11	21	38	73	81	148
2001.....	11,922	4	2	2	6	6	15	24	31	62	101	130
2000.....	11,952	10	3	2	2	8	18	13	38	65	107	149
1999.....	12,015	3	2	1	3	4	13	16	23	61	121	129

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of circulatory system (I71-I78)

White, both sexes

2003.....	21,661	6	-	1	7	23	26	44	63	104	202	364
2002.....	21,935	11	4	1	3	26	23	35	73	115	204	330
2001.....	22,236	10	3	2	8	18	29	52	77	137	203	309
2000.....	22,963	13	4	2	5	14	30	35	76	118	190	303
1999.....	22,952	9	2	2	6	13	29	38	68	138	218	304

White, male

2003.....	11,708	2	-	-	4	15	19	30	40	79	138	250
2002.....	11,818	5	4	-	2	16	15	22	46	63	151	238
2001.....	11,887	6	2	-	3	14	17	36	54	93	135	216
2000.....	12,611	4	2	1	5	7	19	24	53	67	123	208
1999.....	12,444	6	-	1	4	9	18	26	51	92	134	219

White, female

2003.....	9,953	4	-	1	3	8	7	14	23	25	64	114
2002.....	10,117	6	-	1	1	10	8	13	27	52	53	92
2001.....	10,349	4	1	2	5	4	12	16	23	44	68	93
2000.....	10,352	9	2	1	-	7	11	11	23	51	67	95
1999.....	10,508	3	2	1	2	4	11	12	17	46	84	85

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2003.....	2,519	5	-	-	3	6	8	16	21	57	76	140
2002.....	2,525	3	-	-	6	3	9	15	32	32	82	135
2001.....	2,632	2	1	-	1	5	7	14	25	50	92	98
2000.....	2,595	3	-	1	3	2	11	11	25	34	90	107
1999.....	2,434	3	-	-	-	1	6	12	18	35	77	104

Black, male

2003.....	1,216	1	-	-	2	2	5	12	17	34	48	84
2002.....	1,211	1	-	-	2	1	6	8	23	15	56	80
2001.....	1,229	2	-	-	-	3	4	7	17	33	61	63
2000.....	1,207	2	-	-	1	1	5	9	13	22	55	60
1999.....	1,111	3	-	-	-	1	4	8	15	21	40	65

Black, female

2003.....	1,303	4	-	-	1	4	3	4	4	23	28	56
2002.....	1,314	2	-	-	4	2	3	7	9	17	26	55
2001.....	1,403	-	1	-	1	2	3	7	8	17	31	35
2000.....	1,388	1	-	1	2	1	6	2	12	12	35	47
1999.....	1,323	-	-	-	-	-	2	4	3	14	37	39

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of circulatory system (I71-I78)

American Indian 1/, both sexes

2003.....	102	-	-	-	-	-	2	1	2	1	1	5
2002.....	95	-	-	-	-	-	1	-	1	2	-	3
2001.....	79	-	-	-	-	-	-	-	-	1	1	3
2000.....	87	-	-	-	-	-	-	-	-	1	2	3
1999.....	72	-	-	-	2	-	-	-	-	-	-	-

American Indian 1/, male

2003.....	55	-	-	-	-	-	2	-	2	-	-	2
2002.....	53	-	-	-	-	-	1	-	-	2	-	3
2001.....	42	-	-	-	-	-	-	-	-	1	1	2
2000.....	37	-	-	-	-	-	-	-	-	-	-	2
1999.....	32	-	-	-	1	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	47	-	-	-	-	-	-	1	-	1	1	3
2002.....	42	-	-	-	-	-	-	-	1	-	-	-
2001.....	37	-	-	-	-	-	-	-	-	-	-	1
2000.....	50	-	-	-	-	-	-	-	-	1	2	1
1999.....	40	-	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of circulatory system (I71-I78)

Asian or Pacific Islander 2/, both sexes

2003.....	390	-	-	-	1	1	2	1	3	5	5	15
2002.....	372	-	-	-	1	2	2	1	3	5	5	11
2001.....	371	-	-	-	-	-	2	1	1	6	8	11
2000.....	391	-	1	-	-	1	1	3	4	4	9	11
1999.....	351	-	-	-	1	-	-	1	4	5	5	10

Asian or Pacific Islander 2/, male

2003.....	238	-	-	-	1	-	2	1	1	4	3	7
2002.....	227	-	-	-	1	2	2	-	2	1	3	10
2001.....	238	-	-	-	-	-	2	-	1	5	6	10
2000.....	229	-	-	-	-	1	-	3	1	3	6	5
1999.....	207	-	-	-	1	-	-	1	1	4	5	5

Asian or Pacific Islander 2/, female

2003.....	152	-	-	-	-	1	-	-	2	1	2	8
2002.....	145	-	-	-	-	-	-	1	1	4	2	1
2001.....	133	-	-	-	-	-	-	1	-	1	2	1
2000.....	162	-	1	-	-	-	1	-	3	1	3	6
1999.....	144	-	-	-	-	-	-	-	3	1	-	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
All races, both sexes												
2003.....	14,810	1	-	-	3	16	25	43	69	124	192	368
2002.....	14,818	2	-	1	4	20	19	30	80	104	201	327
2001.....	15,234	-	2	-	4	15	28	46	71	134	198	287
2000.....	15,810	-	-	-	5	10	26	38	69	103	200	268
1999.....	15,807	1	1	-	3	8	23	33	60	134	195	283
All races, male												
2003.....	8,976	-	-	-	2	10	21	32	51	96	150	269
2002.....	9,013	-	-	-	3	17	15	25	57	67	162	249
2001.....	9,114	-	2	-	3	12	19	34	56	105	159	219
2000.....	9,661	-	-	-	3	7	16	30	48	72	147	191
1999.....	9,615	1	-	-	2	6	14	25	53	98	134	216
All races, female												
2003.....	5,834	1	-	-	1	6	4	11	18	28	42	99
2002.....	5,805	2	-	1	1	3	4	5	23	37	39	78
2001.....	6,120	-	-	-	1	3	9	12	15	29	39	68
2000.....	6,149	-	-	-	2	3	10	8	21	31	53	77
1999.....	6,192	-	1	-	1	2	9	8	7	36	61	67

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
White, both sexes												
2003.....	13,246	-	-	-	1	14	19	28	49	77	137	252
2002.....	13,285	2	-	1	1	18	13	21	52	82	137	220
2001.....	13,652	-	2	-	4	11	21	37	55	96	131	208
2000.....	14,210	-	-	-	2	9	17	27	50	75	129	193
1999.....	14,336	1	1	-	2	8	19	25	45	105	135	194
White, male												
2003.....	8,074	-	-	-	-	10	15	20	34	64	110	189
2002.....	8,132	-	-	-	1	15	9	18	37	52	112	172
2001.....	8,224	-	2	-	3	9	14	29	44	77	106	163
2000.....	8,797	-	-	-	2	6	12	19	37	52	97	147
1999.....	8,817	1	-	-	1	6	11	19	39	77	95	154
White, female												
2003.....	5,172	-	-	-	1	4	4	8	15	13	27	63
2002.....	5,153	2	-	1	-	3	4	3	15	30	25	48
2001.....	5,428	-	-	-	1	2	7	8	11	19	25	45
2000.....	5,413	-	-	-	-	3	5	8	13	23	32	46
1999.....	5,519	-	1	-	1	2	8	6	6	28	40	40

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Black, both sexes												
2003.....	1,230	1	-	-	1	1	3	13	16	43	52	101
2002.....	1,218	-	-	-	3	1	4	9	25	19	61	96
2001.....	1,251	-	-	-	-	4	6	8	15	33	59	67
2000.....	1,277	-	-	-	3	1	8	8	17	25	65	68
1999.....	1,181	-	-	-	-	-	4	8	13	24	56	81
Black, male												
2003.....	686	-	-	-	1	-	3	11	15	28	38	71
2002.....	677	-	-	-	2	1	4	7	18	13	48	67
2001.....	671	-	-	-	-	3	4	5	11	24	46	45
2000.....	671	-	-	-	1	1	4	8	11	18	46	39
1999.....	619	-	-	-	-	-	3	6	13	17	35	57
Black, female												
2003.....	544	1	-	-	-	1	-	2	1	15	14	30
2002.....	541	-	-	-	1	-	-	2	7	6	13	29
2001.....	580	-	-	-	-	1	2	3	4	9	13	22
2000.....	606	-	-	-	2	-	4	-	6	7	19	29
1999.....	562	-	-	-	-	-	1	2	-	7	21	24

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Aortic aneurysm and dissection (I71)

Asian or Pacific Islander 2/, both sexes

2003.....	288	-	-	-	1	1	2	1	3	4	2	12
2002.....	266	-	-	-	-	1	2	-	2	2	3	9
2001.....	284	-	-	-	-	-	1	1	1	4	7	10
2000.....	282	-	-	-	-	-	1	3	2	3	6	6
1999.....	254	-	-	-	-	-	-	-	2	5	4	8

Asian or Pacific Islander 2/, male

2003.....	190	-	-	-	1	-	2	1	1	4	2	7
2002.....	174	-	-	-	-	1	2	-	2	1	2	8
2001.....	189	-	-	-	-	-	1	-	1	3	6	9
2000.....	172	-	-	-	-	-	-	3	-	2	4	4
1999.....	160	-	-	-	-	-	-	-	1	4	4	5

Asian or Pacific Islander 2/, female

2003.....	98	-	-	-	-	1	-	-	2	-	-	5
2002.....	92	-	-	-	-	-	-	-	-	1	1	1
2001.....	95	-	-	-	-	-	-	1	-	1	1	1
2000.....	110	-	-	-	-	-	1	-	2	1	2	2
1999.....	94	-	-	-	-	-	-	-	1	1	-	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races, both sexes

2003.....	9,862	10	-	1	8	14	13	19	20	43	92	156
2002.....	10,109	12	4	-	6	11	16	21	29	50	90	152
2001.....	10,084	12	2	2	5	8	10	21	32	60	106	134
2000.....	10,226	16	5	3	3	7	16	11	36	54	91	156
1999.....	10,002	11	1	2	6	6	12	18	30	44	105	135

All races, male

2003.....	4,241	3	-	-	5	7	7	11	9	21	39	74
2002.....	4,296	6	4	-	2	2	9	5	14	14	48	82
2001.....	4,282	8	-	-	-	5	4	9	16	27	44	72
2000.....	4,423	6	2	1	3	2	8	6	19	20	37	84
1999.....	4,179	8	-	1	4	4	8	10	14	19	45	73

All races, female

2003.....	5,621	7	-	1	3	7	6	8	11	22	53	82
2002.....	5,813	6	-	-	4	9	7	16	15	36	42	70
2001.....	5,802	4	2	2	5	3	6	12	16	33	62	62
2000.....	5,803	10	3	2	-	5	8	5	17	34	54	72
1999.....	5,823	3	1	1	2	2	4	8	16	25	60	62

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, both sexes

2003.....	8,415	6	-	1	6	9	7	16	14	27	65	112
2002.....	8,650	9	4	-	2	8	10	14	21	33	67	110
2001.....	8,584	10	1	2	4	7	8	15	22	41	72	101
2000.....	8,753	13	4	2	3	5	13	8	26	43	61	110
1999.....	8,616	8	1	2	4	5	10	13	23	33	83	110

White, male

2003.....	3,634	2	-	-	4	5	4	10	6	15	28	61
2002.....	3,686	5	4	-	1	1	6	4	9	11	39	66
2001.....	3,663	6	-	-	-	5	3	7	10	16	29	53
2000.....	3,814	4	2	1	3	1	7	5	16	15	26	61
1999.....	3,627	5	-	1	3	3	7	7	12	15	39	65

White, female

2003.....	4,781	4	-	1	2	4	3	6	8	12	37	51
2002.....	4,964	4	-	-	1	7	4	10	12	22	28	44
2001.....	4,921	4	1	2	4	2	5	8	12	25	43	48
2000.....	4,939	9	2	1	-	4	6	3	10	28	35	49
1999.....	4,989	3	1	1	1	2	3	6	11	18	44	45

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2003.....	1,289	4	-	-	2	5	5	3	5	14	24	39
2002.....	1,307	3	-	-	3	2	5	6	7	13	21	39
2001.....	1,381	2	1	-	1	1	1	6	10	17	33	31
2000.....	1,318	3	-	1	-	1	3	3	8	9	25	39
1999.....	1,253	3	-	-	-	1	2	4	5	11	21	23

Black, male

2003.....	530	1	-	-	1	2	2	1	2	6	10	13
2002.....	534	1	-	-	-	-	2	1	5	2	8	13
2001.....	558	2	-	-	-	-	-	2	6	9	15	18
2000.....	536	2	-	-	-	-	1	1	2	4	9	21
1999.....	492	3	-	-	-	1	1	2	2	4	5	8

Black, female

2003.....	759	3	-	-	1	3	3	2	3	8	14	26
2002.....	773	2	-	-	3	2	3	5	2	11	13	26
2001.....	823	-	1	-	1	1	1	4	4	8	18	13
2000.....	782	1	-	1	-	1	2	2	6	5	16	18
1999.....	761	-	-	-	-	-	1	2	3	7	16	15

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

American Indian 1/, both sexes

2003.....	56	-	-	-	-	-	1	-	1	1	-	2
2002.....	46	-	-	-	-	-	1	-	-	1	-	1
2001.....	32	-	-	-	-	-	-	-	-	-	-	1
2000.....	46	-	-	-	-	-	-	-	-	1	2	2
1999.....	36	-	-	-	1	-	-	-	-	-	-	-

American Indian 1/, male

2003.....	29	-	-	-	-	-	1	-	1	-	-	-
2002.....	23	-	-	-	-	-	1	-	-	1	-	1
2001.....	12	-	-	-	-	-	-	-	-	-	-	-
2000.....	16	-	-	-	-	-	-	-	-	-	-	1
1999.....	13	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	27	-	-	-	-	-	-	-	-	1	-	2
2002.....	23	-	-	-	-	-	-	-	-	-	-	-
2001.....	20	-	-	-	-	-	-	-	-	-	-	1
2000.....	30	-	-	-	-	-	-	-	-	1	2	1
1999.....	23	-	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Asian or Pacific Islander 2/, both sexes

2003.....	102	-	-	-	-	-	-	-	-	1	3	3
2002.....	106	-	-	-	1	1	-	1	1	3	2	2
2001.....	87	-	-	-	-	-	1	-	-	2	1	1
2000.....	109	-	1	-	-	1	-	-	2	1	3	5
1999.....	97	-	-	-	1	-	-	1	2	-	1	2

Asian or Pacific Islander 2/, male

2003.....	48	-	-	-	-	-	-	-	-	-	1	-
2002.....	53	-	-	-	1	1	-	-	-	-	1	2
2001.....	49	-	-	-	-	-	1	-	-	2	-	1
2000.....	57	-	-	-	-	1	-	-	1	1	2	1
1999.....	47	-	-	-	1	-	-	1	-	-	1	-

Asian or Pacific Islander 2/, female

2003.....	54	-	-	-	-	-	-	-	-	1	2	3
2002.....	53	-	-	-	-	-	-	1	1	3	1	-
2001.....	38	-	-	-	-	-	-	-	-	-	1	-
2000.....	52	-	1	-	-	-	-	-	1	-	1	4
1999.....	50	-	-	-	-	-	-	-	2	-	-	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

All races, both sexes

2003.....	4,737	36	4	1	3	14	36	53	86	126	216	273
2002.....	4,711	27	4	1	1	12	23	38	101	128	177	251
2001.....	4,665	22	5	-	-	10	28	45	79	100	185	222
2000.....	4,603	27	4	5	7	10	27	58	75	125	194	249
1999.....	4,025	23	1	1	1	11	19	30	63	98	173	202

All races, male

2003.....	1,966	21	3	1	1	5	15	23	37	57	110	147
2002.....	1,944	15	1	-	-	9	12	12	57	67	87	137
2001.....	1,981	11	5	-	-	5	12	18	35	51	98	127
2000.....	1,981	16	2	3	3	6	10	18	39	70	120	145
1999.....	1,714	13	1	1	-	6	7	14	30	55	104	102

All races, female

2003.....	2,771	15	1	-	2	9	21	30	49	69	106	126
2002.....	2,767	12	3	1	1	3	11	26	44	61	90	114
2001.....	2,684	11	-	-	-	5	16	27	44	49	87	95
2000.....	2,622	11	2	2	4	4	17	40	36	55	74	104
1999.....	2,311	10	-	-	1	5	12	16	33	43	69	100

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2003.....	3,843	28	4	1	2	9	21	36	51	78	146	202
2002.....	3,853	20	2	-	1	8	13	23	61	82	121	185
2001.....	3,843	17	3	-	-	5	16	30	50	61	128	163
2000.....	3,787	20	3	4	5	4	18	33	49	87	143	189
1999.....	3,345	18	-	1	1	4	11	21	43	66	117	146

White, male

2003.....	1,569	16	3	1	-	3	8	11	25	45	84	103
2002.....	1,545	13	-	-	-	6	6	6	35	44	58	97
2001.....	1,603	8	3	-	-	3	7	11	23	32	67	102
2000.....	1,618	13	1	3	2	3	6	9	25	47	94	111
1999.....	1,399	9	-	1	-	3	5	10	20	37	69	73

White, female

2003.....	2,274	12	1	-	2	6	13	25	26	33	62	99
2002.....	2,308	7	2	-	1	2	7	17	26	38	63	88
2001.....	2,240	9	-	-	-	2	9	19	27	29	61	61
2000.....	2,169	7	2	1	3	1	12	24	24	40	49	78
1999.....	1,946	9	-	-	1	1	6	11	23	29	48	73

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

Black, both sexes

2003.....	824	6	-	-	-	5	14	17	32	45	67	69
2002.....	797	7	1	1	-	4	9	15	39	43	52	60
2001.....	742	5	2	-	-	3	11	15	28	36	53	52
2000.....	761	5	1	1	2	6	8	23	26	36	51	58
1999.....	616	4	1	-	-	6	8	9	19	29	52	55

Black, male

2003.....	368	5	-	-	-	2	7	12	11	11	25	42
2002.....	366	2	1	-	-	3	6	6	21	21	28	36
2001.....	339	3	2	-	-	1	4	7	11	17	30	20
2000.....	335	2	1	-	1	3	4	8	14	22	26	34
1999.....	282	3	1	-	-	2	2	4	10	16	33	29

Black, female

2003.....	456	1	-	-	-	3	7	5	21	34	42	27
2002.....	431	5	-	1	-	1	3	9	18	22	24	24
2001.....	403	2	-	-	-	2	7	8	17	19	23	32
2000.....	426	3	-	1	1	3	4	15	12	14	25	24
1999.....	334	1	-	-	-	4	6	5	9	13	19	26

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

American Indian 1/, both sexes

2003.....	15	-	-	-	-	-	-	-	-	-	2	1
2002.....	23	-	-	-	-	-	-	-	-	1	1	4
2001.....	25	-	-	-	-	1	-	-	1	2	-	4
2000.....	18	-	-	-	-	-	1	1	-	-	-	1
1999.....	13	-	-	-	-	-	-	-	1	-	1	-

American Indian 1/, male

2003.....	7	-	-	-	-	-	-	-	-	-	1	1
2002.....	13	-	-	-	-	-	-	-	-	1	-	2
2001.....	14	-	-	-	-	-	-	-	1	1	-	3
2000.....	8	-	-	-	-	-	-	1	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	8	-	-	-	-	-	-	-	-	-	1	-
2002.....	10	-	-	-	-	-	-	-	-	-	1	2
2001.....	11	-	-	-	-	1	-	-	-	1	-	1
2000.....	10	-	-	-	-	-	1	-	-	-	-	1
1999.....	9	-	-	-	-	-	-	-	1	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

Asian or Pacific Islander 2/, both sexes

2003.....	55	2	-	-	1	-	1	-	3	3	1	1
2002.....	38	-	1	-	-	-	1	-	1	2	3	2
2001.....	55	-	-	-	-	1	1	-	-	1	4	3
2000.....	37	2	-	-	-	-	-	1	-	2	-	1
1999.....	51	1	-	-	-	1	-	-	-	3	3	1

Asian or Pacific Islander 2/, male

2003.....	22	-	-	-	1	-	-	-	1	1	-	1
2002.....	20	-	-	-	-	-	-	-	1	1	1	2
2001.....	25	-	-	-	-	1	1	-	-	1	1	2
2000.....	20	1	-	-	-	-	-	-	-	1	-	-
1999.....	29	1	-	-	-	1	-	-	-	2	2	-

Asian or Pacific Islander 2/, female

2003.....	33	2	-	-	-	-	1	-	2	2	1	-
2002.....	18	-	1	-	-	-	1	-	-	1	2	-
2001.....	30	-	-	-	-	-	-	-	-	-	3	1
2000.....	17	1	-	-	-	-	-	1	-	1	-	1
1999.....	22	-	-	-	-	-	-	-	-	1	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza and pneumonia (J10-J18)												
All races, both sexes												
2003.....	65,163	322	163	75	72	80	144	149	224	376	616	918
2002.....	65,681	263	110	38	53	75	92	135	210	367	604	887
2001.....	62,034	299	112	46	46	66	115	124	215	393	590	810
2000.....	65,313	289	103	47	40	65	124	142	222	405	663	783
1999.....	63,730	320	130	46	47	73	106	135	204	462	601	798
All races, male												
2003.....	28,778	192	91	37	33	50	74	90	139	208	352	528
2002.....	28,918	162	63	15	23	41	56	68	122	197	349	519
2001.....	27,342	167	57	23	24	36	70	64	127	219	337	497
2000.....	28,658	163	54	27	21	35	67	82	122	211	419	464
1999.....	27,718	171	66	27	26	34	56	76	121	251	341	500
All races, female												
2003.....	36,385	130	72	38	39	30	70	59	85	168	264	390
2002.....	36,763	101	47	23	30	34	36	67	88	170	255	368
2001.....	34,692	132	55	23	22	30	45	60	88	174	253	313
2000.....	36,655	126	49	20	19	30	57	60	100	194	244	319
1999.....	36,012	149	64	19	21	39	50	59	83	211	260	298

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza and pneumonia (J10-J18)												
White, both sexes												
2003.....	57,645	209	115	50	56	57	116	100	158	272	437	667
2002.....	58,346	162	63	29	40	52	71	102	146	258	439	673
2001.....	54,774	168	78	33	34	46	80	83	149	275	403	571
2000.....	57,914	172	69	36	34	48	90	98	163	305	462	561
1999.....	56,694	184	75	36	36	54	67	85	137	327	387	557
White, male												
2003.....	25,009	127	67	23	27	35	60	67	98	153	243	385
2002.....	25,381	105	35	10	16	31	43	51	86	136	257	387
2001.....	23,744	100	41	17	17	24	55	40	87	151	231	357
2000.....	25,002	98	30	22	17	27	52	51	87	161	294	325
1999.....	24,281	103	33	21	21	26	36	50	84	181	209	346
White, female												
2003.....	32,636	82	48	27	29	22	56	33	60	119	194	282
2002.....	32,965	57	28	19	24	21	28	51	60	122	182	286
2001.....	31,030	68	37	16	17	22	25	43	62	124	172	214
2000.....	32,912	74	39	14	17	21	38	47	76	144	168	236
1999.....	32,413	81	42	15	15	28	31	35	53	146	178	211

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

Black, both sexes

2003.....	5,872	93	28	20	10	17	25	44	58	93	159	225
2002.....	5,871	87	41	7	12	18	20	27	56	101	151	192
2001.....	5,771	106	26	9	8	16	29	32	57	109	167	221
2000.....	5,990	103	30	7	4	14	30	38	55	85	172	197
1999.....	5,876	115	43	8	7	14	35	47	60	121	191	223

Black, male

2003.....	2,906	54	14	12	3	10	14	22	37	47	96	127
2002.....	2,768	48	22	4	6	8	12	13	32	54	84	117
2001.....	2,813	53	14	4	5	11	12	17	36	62	94	130
2000.....	2,915	56	21	4	3	6	15	27	32	42	110	124
1999.....	2,825	58	25	5	3	6	17	25	33	62	116	142

Black, female

2003.....	2,966	39	14	8	7	7	11	22	21	46	63	98
2002.....	3,103	39	19	3	6	10	8	14	24	47	67	75
2001.....	2,958	53	12	5	3	5	17	15	21	47	73	91
2000.....	3,075	47	9	3	1	8	15	11	23	43	62	73
1999.....	3,051	57	18	3	4	8	18	22	27	59	75	81

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

American Indian 1/, both sexes

2003.....	390	12	5	1	2	3	1	3	3	7	14	18
2002.....	293	6	1	-	-	2	1	2	5	5	8	10
2001.....	318	17	3	1	3	2	1	5	4	6	13	6
2000.....	289	6	-	2	1	1	1	4	1	11	17	8
1999.....	315	9	3	1	2	1	4	1	3	4	16	14

American Indian 1/, male

2003.....	184	7	2	1	1	2	-	-	1	4	10	12
2002.....	133	3	1	-	-	-	1	2	2	5	4	7
2001.....	160	11	-	-	1	-	1	5	1	5	8	3
2000.....	149	2	-	-	-	1	-	2	1	6	12	5
1999.....	147	4	3	1	1	1	3	-	1	-	12	8

American Indian 1/, female

2003.....	206	5	3	-	1	1	1	3	2	3	4	6
2002.....	160	3	-	-	-	2	-	-	3	-	4	3
2001.....	158	6	3	1	2	2	-	-	3	1	5	3
2000.....	140	4	-	2	1	-	1	2	-	5	5	3
1999.....	168	5	-	-	1	-	1	1	2	4	4	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

Asian or Pacific Islander 2/, both sexes

2003.....	1,256	8	15	4	4	3	2	2	5	4	6	8
2002.....	1,171	8	5	2	1	3	-	4	3	3	6	12
2001.....	1,171	8	5	3	1	2	5	4	5	3	7	12
2000.....	1,120	8	4	2	1	2	3	2	3	4	12	17
1999.....	845	12	9	1	2	4	-	2	4	10	7	4

Asian or Pacific Islander 2/, male

2003.....	679	4	8	1	2	3	-	1	3	4	3	4
2002.....	636	6	5	1	1	2	-	2	2	2	4	8
2001.....	625	3	2	2	1	1	2	2	3	1	4	7
2000.....	592	7	3	1	1	1	-	2	2	2	3	10
1999.....	465	6	5	-	1	1	-	1	3	8	4	4

Asian or Pacific Islander 2/, female

2003.....	577	4	7	3	2	-	2	1	2	-	3	4
2002.....	535	2	-	1	-	1	-	2	1	1	2	4
2001.....	546	5	3	1	-	1	3	2	2	2	3	5
2000.....	528	1	1	1	-	1	3	-	1	2	9	7
1999.....	380	6	4	1	1	3	-	1	1	2	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
All races, both sexes												
2003.....	1,792	31	59	28	18	18	12	5	8	10	14	29
2002.....	727	7	5	5	7	5	2	3	1	3	5	8
2001.....	257	7	6	9	3	4	3	4	6	1	5	8
2000.....	1,765	9	10	3	8	8	5	8	5	8	14	24
1999.....	1,665	13	12	6	5	7	4	2	7	8	8	13
All races, male												
2003.....	699	18	33	13	6	9	3	4	5	5	5	15
2002.....	289	5	3	1	1	2	2	2	-	2	2	7
2001.....	101	6	2	3	1	3	3	3	3	1	1	5
2000.....	664	5	4	1	4	3	3	4	1	5	9	11
1999.....	602	6	9	4	1	1	1	-	5	5	4	8
All races, female												
2003.....	1,093	13	26	15	12	9	9	1	3	5	9	14
2002.....	438	2	2	4	6	3	-	1	1	1	3	1
2001.....	156	1	4	6	2	1	-	1	3	-	4	3
2000.....	1,101	4	6	2	4	5	2	4	4	3	5	13
1999.....	1,063	7	3	2	4	6	3	2	2	3	4	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
White, both sexes												
2003.....	1,653	23	42	20	10	11	9	3	7	8	12	22
2002.....	686	5	2	4	6	3	2	3	1	3	5	8
2001.....	229	7	5	5	2	3	2	2	5	1	3	8
2000.....	1,662	7	9	2	7	8	4	7	4	7	11	18
1999.....	1,574	7	8	6	4	6	4	1	7	4	6	12
White, male												
2003.....	640	14	25	9	3	5	2	3	5	3	3	11
2002.....	270	3	1	1	1	-	2	2	-	2	2	7
2001.....	88	6	2	2	1	2	2	1	2	1	1	5
2000.....	617	5	4	1	3	3	2	3	-	4	6	7
1999.....	569	1	6	4	1	1	1	-	5	3	3	7
White, female												
2003.....	1,013	9	17	11	7	6	7	-	2	5	9	11
2002.....	416	2	1	3	5	3	-	1	1	1	3	1
2001.....	141	1	3	3	1	1	-	1	3	-	2	3
2000.....	1,045	2	5	1	4	5	2	4	4	3	5	11
1999.....	1,005	6	2	2	3	5	3	1	2	1	3	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
Black, both sexes												
2003.....	97	7	6	6	5	4	3	2	1	-	1	7
2002.....	30	1	3	1	1	2	-	-	-	-	-	-
2001.....	20	-	-	2	-	1	-	1	1	-	1	-
2000.....	81	2	1	-	1	-	1	1	-	-	3	5
1999.....	66	4	1	-	-	1	-	1	-	3	2	1
Black, male												
2003.....	37	3	3	3	1	2	1	1	-	-	1	4
2002.....	15	1	2	-	-	2	-	-	-	-	-	-
2001.....	10	-	-	-	-	1	-	1	1	-	-	-
2000.....	35	-	-	-	1	-	1	1	-	-	3	3
1999.....	24	3	1	-	-	-	-	-	-	1	1	1
Black, female												
2003.....	60	4	3	3	4	2	2	1	1	-	-	3
2002.....	15	-	1	1	1	-	-	-	-	-	-	-
2001.....	10	-	-	2	-	-	-	-	-	-	1	-
2000.....	46	2	1	-	-	-	-	-	-	-	-	2
1999.....	42	1	-	-	-	1	-	1	-	2	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza (J10-J11)

American Indian 1/, both sexes

2003.....	17	1	3	-	1	2	-	-	-	1	-	-
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	1	1	-	-	-	-	-	-	-
2000.....	4	-	-	1	-	-	-	-	-	-	-	-
1999.....	5	1	-	-	1	-	-	-	-	-	-	-

American Indian 1/, male

2003.....	7	1	1	-	-	1	-	-	-	1	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	10	-	2	-	1	1	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	1	1	-	-	-	-	-	-	-
2000.....	3	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
All races, both sexes												
2003.....	63,371	291	104	47	54	62	132	144	216	366	602	889
2002.....	64,954	256	105	33	46	70	90	132	209	364	599	879
2001.....	61,777	292	106	37	43	62	112	120	209	392	585	802
2000.....	63,548	280	93	44	32	57	119	134	217	397	649	759
1999.....	62,065	307	118	40	42	66	102	133	197	454	593	785
All races, male												
2003.....	28,079	174	58	24	27	41	71	86	134	203	347	513
2002.....	28,629	157	60	14	22	39	54	66	122	195	347	512
2001.....	27,241	161	55	20	23	33	67	61	124	218	336	492
2000.....	27,994	158	50	26	17	32	64	78	121	206	410	453
1999.....	27,116	165	57	23	25	33	55	76	116	246	337	492
All races, female												
2003.....	35,292	117	46	23	27	21	61	58	82	163	255	376
2002.....	36,325	99	45	19	24	31	36	66	87	169	252	367
2001.....	34,536	131	51	17	20	29	45	59	85	174	249	310
2000.....	35,554	122	43	18	15	25	55	56	96	191	239	306
1999.....	34,949	142	61	17	17	33	47	57	81	208	256	293

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
White, both sexes												
2003.....	55,992	186	73	30	46	46	107	97	151	264	425	645
2002.....	57,660	157	61	25	34	49	69	99	145	255	434	665
2001.....	54,545	161	73	28	32	43	78	81	144	274	400	563
2000.....	56,252	165	60	34	27	40	86	91	159	298	451	543
1999.....	55,120	177	67	30	32	48	63	84	130	323	381	545
White, male												
2003.....	24,369	113	42	14	24	30	58	64	93	150	240	374
2002.....	25,111	102	34	9	15	31	41	49	86	134	255	380
2001.....	23,656	94	39	15	16	22	53	39	85	150	230	352
2000.....	24,385	93	26	21	14	24	50	48	87	157	288	318
1999.....	23,712	102	27	17	20	25	35	50	79	178	206	339
White, female												
2003.....	31,623	73	31	16	22	16	49	33	58	114	185	271
2002.....	32,549	55	27	16	19	18	28	50	59	121	179	285
2001.....	30,889	67	34	13	16	21	25	42	59	124	170	211
2000.....	31,867	72	34	13	13	16	36	43	72	141	163	225
1999.....	31,408	75	40	13	12	23	28	34	51	145	175	206

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Black, both sexes												
2003.....	5,775	86	22	14	5	13	22	42	57	93	158	218
2002.....	5,841	86	38	6	11	16	20	27	56	101	151	192
2001.....	5,751	106	26	7	8	15	29	31	56	109	166	221
2000.....	5,909	101	29	7	3	14	29	37	55	85	169	192
1999.....	5,810	111	42	8	7	13	35	46	60	118	189	222
Black, male												
2003.....	2,869	51	11	9	2	8	13	21	37	47	95	123
2002.....	2,753	47	20	4	6	6	12	13	32	54	84	117
2001.....	2,803	53	14	4	5	10	12	16	35	62	94	130
2000.....	2,880	56	21	4	2	6	14	26	32	42	107	121
1999.....	2,801	55	24	5	3	6	17	25	33	61	115	141
Black, female												
2003.....	2,906	35	11	5	3	5	9	21	20	46	63	95
2002.....	3,088	39	18	2	5	10	8	14	24	47	67	75
2001.....	2,948	53	12	3	3	5	17	15	21	47	72	91
2000.....	3,029	45	8	3	1	8	15	11	23	43	62	71
1999.....	3,009	56	18	3	4	7	18	21	27	57	74	81

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
American Indian 1/, both sexes												
2003.....	373	11	2	1	1	1	1	3	3	6	14	18
2002.....	288	6	1	-	-	2	1	2	5	5	8	10
2001.....	315	17	2	-	2	2	1	5	4	6	13	6
2000.....	285	6	-	1	1	1	1	4	1	11	17	8
1999.....	310	8	3	1	1	1	4	1	3	4	16	14
American Indian 1/, male												
2003.....	177	6	1	1	1	1	-	-	1	3	10	12
2002.....	130	3	1	-	-	-	1	2	2	5	4	7
2001.....	160	11	-	-	1	-	1	5	1	5	8	3
2000.....	148	2	-	-	-	1	-	2	1	6	12	5
1999.....	145	3	3	1	1	1	3	-	1	-	12	8
American Indian 1/, female												
2003.....	196	5	1	-	-	-	1	3	2	3	4	6
2002.....	158	3	-	-	-	2	-	-	3	-	4	3
2001.....	155	6	2	-	1	2	-	-	3	1	5	3
2000.....	137	4	-	1	1	-	1	2	-	5	5	3
1999.....	165	5	-	-	-	-	1	1	2	4	4	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Asian or Pacific Islander 2/, both sexes												
2003.....	1,231	8	7	2	2	2	2	2	5	3	5	8
2002.....	1,165	7	5	2	1	3	-	4	3	3	6	12
2001.....	1,166	8	5	2	1	2	4	3	5	3	6	12
2000.....	1,102	8	4	2	1	2	3	2	2	3	12	16
1999.....	825	11	6	1	2	4	-	2	4	9	7	4
Asian or Pacific Islander 2/, male												
2003.....	664	4	4	-	-	2	-	1	3	3	2	4
2002.....	635	5	5	1	1	2	-	2	2	2	4	8
2001.....	622	3	2	1	1	1	1	1	3	1	4	7
2000.....	581	7	3	1	1	1	-	2	1	1	3	9
1999.....	458	5	3	-	1	1	-	1	3	7	4	4
Asian or Pacific Islander 2/, female												
2003.....	567	4	3	2	2	-	2	1	2	-	3	4
2002.....	530	2	-	1	-	1	-	2	1	1	2	4
2001.....	544	5	3	1	-	1	3	2	2	2	2	5
2000.....	521	1	1	1	-	1	3	-	1	2	9	7
1999.....	367	6	3	1	1	3	-	1	1	2	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

All races, both sexes

2003.....	421	54	28	3	3	1	5	-	2	4	7	7
2002.....	386	52	23	7	1	1	2	3	2	2	6	6
2001.....	347	50	18	3	-	-	1	1	1	7	5	4
2000.....	425	43	11	3	4	4	1	5	3	3	9	8
1999.....	540	63	25	2	3	3	2	2	2	6	5	7

All races, male

2003.....	170	35	17	2	-	-	2	-	1	3	3	2
2002.....	154	39	9	7	-	-	-	1	-	1	2	1
2001.....	117	26	8	2	-	-	1	-	-	4	3	1
2000.....	160	25	4	2	1	4	-	5	1	1	6	2
1999.....	196	36	15	-	2	1	-	2	1	3	2	5

All races, female

2003.....	251	19	11	1	3	1	3	-	1	1	4	5
2002.....	232	13	14	-	1	1	2	2	2	1	4	5
2001.....	230	24	10	1	-	-	-	1	1	3	2	3
2000.....	265	18	7	1	3	-	1	-	2	2	3	6
1999.....	344	27	10	2	1	2	2	-	1	3	3	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

White, both sexes

2003.....	375	42	17	3	2	1	2	-	1	4	7	6
2002.....	334	34	20	7	1	1	1	2	2	2	5	4
2001.....	299	31	13	3	-	-	1	-	1	7	5	4
2000.....	372	25	8	2	1	4	1	4	1	2	6	5
1999.....	462	36	15	1	3	2	2	1	2	6	3	6

White, male

2003.....	147	27	10	2	-	-	-	-	1	3	3	2
2002.....	130	26	7	7	-	-	-	-	-	1	2	1
2001.....	94	16	5	2	-	-	1	-	-	4	3	1
2000.....	134	12	4	2	-	4	-	4	-	1	5	2
1999.....	156	19	8	-	2	1	-	1	1	3	1	5

White, female

2003.....	228	15	7	1	2	1	2	-	-	1	4	4
2002.....	204	8	13	-	1	1	1	2	2	1	3	3
2001.....	205	15	8	1	-	-	-	-	1	3	2	3
2000.....	238	13	4	-	1	-	1	-	1	1	1	3
1999.....	306	17	7	1	1	1	2	-	1	3	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

Black, both sexes

2003.....	40	12	7	-	1	-	3	-	1	-	-	1
2002.....	45	16	2	-	-	-	1	1	-	-	1	2
2001.....	34	15	2	-	-	-	-	1	-	-	-	-
2000.....	46	14	3	-	3	-	-	1	1	1	3	3
1999.....	64	24	6	-	-	1	-	1	-	-	2	1

Black, male

2003.....	21	8	6	-	-	-	2	-	-	-	-	-
2002.....	23	13	1	-	-	-	-	1	-	-	-	-
2001.....	16	9	1	-	-	-	-	-	-	-	-	-
2000.....	22	10	-	-	1	-	-	1	1	-	1	-
1999.....	30	15	3	-	-	-	-	1	-	-	1	-

Black, female

2003.....	19	4	1	-	1	-	1	-	1	-	-	1
2002.....	22	3	1	-	-	-	1	-	-	-	1	2
2001.....	18	6	1	-	-	-	-	1	-	-	-	-
2000.....	24	4	3	-	2	-	-	-	-	1	2	3
1999.....	34	9	3	-	-	1	-	-	-	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
All races, both sexes												
2003.....	304	53	27	3	3	1	5	-	2	4	5	7
2002.....	279	49	23	7	1	-	2	3	2	2	4	6
2001.....	263	49	18	3	-	-	1	1	1	6	5	4
2000.....	290	42	11	2	4	3	1	3	3	2	8	8
1999.....	388	62	24	2	2	3	2	1	2	6	4	7
All races, male												
2003.....	126	34	16	2	-	-	2	-	1	3	2	2
2002.....	122	37	9	7	-	-	-	1	-	1	2	1
2001.....	89	25	8	2	-	-	1	-	-	3	3	1
2000.....	119	24	4	1	1	3	-	3	1	1	5	2
1999.....	154	36	14	-	2	1	-	1	1	3	1	5
All races, female												
2003.....	178	19	11	1	3	1	3	-	1	1	3	5
2002.....	157	12	14	-	1	-	2	2	2	1	2	5
2001.....	174	24	10	1	-	-	-	1	1	3	2	3
2000.....	171	18	7	1	3	-	1	-	2	1	3	6
1999.....	234	26	10	2	-	2	2	-	1	3	3	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2003.....	263	41	16	3	2	1	2	-	1	4	5	6
2002.....	237	31	20	7	1	-	1	2	2	2	4	4
2001.....	218	30	13	3	-	-	1	-	1	6	5	4
2000.....	244	25	8	1	1	3	1	2	1	1	5	5
1999.....	320	35	15	1	2	2	2	1	2	6	2	6

White, male

2003.....	104	26	9	2	-	-	-	-	1	3	2	2
2002.....	100	24	7	7	-	-	-	-	-	1	2	1
2001.....	67	15	5	2	-	-	1	-	-	3	3	1
2000.....	98	12	4	1	-	3	-	2	-	1	4	2
1999.....	119	19	8	-	2	1	-	1	1	3	-	5

White, female

2003.....	159	15	7	1	2	1	2	-	-	1	3	4
2002.....	137	7	13	-	1	-	1	2	2	1	2	3
2001.....	151	15	8	1	-	-	-	-	1	3	2	3
2000.....	146	13	4	-	1	-	1	-	1	-	1	3
1999.....	201	16	7	1	-	1	2	-	1	3	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Black, both sexes												
2003.....	35	12	7	-	1	-	3	-	1	-	-	1
2002.....	36	16	2	-	-	-	1	1	-	-	-	2
2001.....	32	15	2	-	-	-	-	1	-	-	-	-
2000.....	40	13	3	-	3	-	-	1	1	1	3	3
1999.....	55	24	5	-	-	1	-	-	-	-	2	1
Black, male												
2003.....	20	8	6	-	-	-	2	-	-	-	-	-
2002.....	21	13	1	-	-	-	-	1	-	-	-	-
2001.....	15	9	1	-	-	-	-	-	-	-	-	-
2000.....	18	9	-	-	1	-	-	1	1	-	1	-
1999.....	26	15	2	-	-	-	-	-	-	-	1	-
Black, female												
2003.....	15	4	1	-	1	-	1	-	1	-	-	1
2002.....	15	3	1	-	-	-	1	-	-	-	-	2
2001.....	17	6	1	-	-	-	-	1	-	-	-	-
2000.....	22	4	3	-	2	-	-	-	-	1	2	3
1999.....	29	9	3	-	-	1	-	-	-	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified acute lower respiratory infection (J22)

All races, both sexes

2003.....	117	1	1	-	-	-	-	-	-	-	2	-
2002.....	107	3	-	-	-	1	-	-	-	-	2	-
2001.....	84	1	-	-	-	-	-	-	-	1	-	-
2000.....	135	1	-	1	-	1	-	2	-	1	1	-
1999.....	152	1	1	-	1	-	-	1	-	-	1	-

All races, male

2003.....	44	1	1	-	-	-	-	-	-	-	1	-
2002.....	32	2	-	-	-	-	-	-	-	-	-	-
2001.....	28	1	-	-	-	-	-	-	-	1	-	-
2000.....	41	1	-	1	-	1	-	2	-	-	1	-
1999.....	42	-	1	-	-	-	-	1	-	-	1	-

All races, female

2003.....	73	-	-	-	-	-	-	-	-	-	1	-
2002.....	75	1	-	-	-	1	-	-	-	-	2	-
2001.....	56	-	-	-	-	-	-	-	-	-	-	-
2000.....	94	-	-	-	-	-	-	-	-	1	-	-
1999.....	110	1	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified acute lower respiratory infection (J22)

White, both sexes

2003.....	112	1	1	-	-	-	-	-	-	-	2	-
2002.....	97	3	-	-	-	1	-	-	-	-	1	-
2001.....	81	1	-	-	-	-	-	-	-	1	-	-
2000.....	128	-	-	1	-	1	-	2	-	1	1	-
1999.....	142	1	-	-	1	-	-	-	-	-	1	-

White, male

2003.....	43	1	1	-	-	-	-	-	-	-	1	-
2002.....	30	2	-	-	-	-	-	-	-	-	-	-
2001.....	27	1	-	-	-	-	-	-	-	1	-	-
2000.....	36	-	-	1	-	1	-	2	-	-	1	-
1999.....	37	-	-	-	-	-	-	-	-	-	1	-

White, female

2003.....	69	-	-	-	-	-	-	-	-	-	1	-
2002.....	67	1	-	-	-	1	-	-	-	-	1	-
2001.....	54	-	-	-	-	-	-	-	-	-	-	-
2000.....	92	-	-	-	-	-	-	-	-	1	-	-
1999.....	105	1	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

All races, both sexes

2003.....	126,382	31	55	37	81	84	107	118	164	301	649	1,276
2002.....	124,816	39	65	41	95	93	99	116	185	340	668	1,226
2001.....	123,013	41	43	42	62	74	97	102	188	355	614	1,127
2000.....	122,009	36	51	48	91	86	104	125	171	310	620	1,174
1999.....	124,181	36	54	49	90	107	102	145	185	302	588	1,079

All races, male

2003.....	60,714	23	38	23	52	50	64	67	89	142	287	681
2002.....	60,713	23	44	24	50	68	52	70	95	159	338	617
2001.....	59,697	22	29	24	37	43	56	55	91	171	284	591
2000.....	60,004	23	28	26	54	58	62	72	93	145	292	617
1999.....	62,415	19	30	29	64	60	63	69	83	142	268	547

All races, female

2003.....	65,668	8	17	14	29	34	43	51	75	159	362	595
2002.....	64,103	16	21	17	45	25	47	46	90	181	330	609
2001.....	63,316	19	14	18	25	31	41	47	97	184	330	536
2000.....	62,005	13	23	22	37	28	42	53	78	165	328	557
1999.....	61,766	17	24	20	26	47	39	76	102	160	320	532

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2003.....	116,917	23	24	14	40	38	63	70	99	211	464	1,012
2002.....	115,395	25	30	17	38	47	62	68	121	226	470	970
2001.....	113,819	22	20	18	30	40	45	53	103	212	453	882
2000.....	112,840	23	28	23	40	40	63	67	99	194	430	882
1999.....	114,735	16	31	29	42	68	57	73	103	200	408	801

White, male

2003.....	55,397	20	14	7	24	20	39	34	45	100	205	549
2002.....	55,409	16	21	12	19	31	29	37	64	96	240	489
2001.....	54,561	13	12	10	18	19	26	29	45	109	207	480
2000.....	54,816	16	13	12	21	26	39	36	48	85	207	472
1999.....	57,000	10	14	15	30	36	32	32	46	90	181	410

White, female

2003.....	61,520	3	10	7	16	18	24	36	54	111	259	463
2002.....	59,986	9	9	5	19	16	33	31	57	130	230	481
2001.....	59,258	9	8	8	12	21	19	24	58	103	246	402
2000.....	58,024	7	15	11	19	14	24	31	51	109	223	410
1999.....	57,735	6	17	14	12	32	25	41	57	110	227	391

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2003.....	7,709	8	26	23	40	41	41	45	59	81	166	241
2002.....	7,831	14	30	21	55	43	36	43	61	110	188	233
2001.....	7,589	17	23	23	31	30	47	47	82	132	147	224
2000.....	7,607	11	21	24	46	44	37	51	65	104	171	259
1999.....	7,915	18	18	20	48	36	42	68	76	95	169	257

Black, male

2003.....	4,247	3	21	16	27	26	23	31	41	38	70	120
2002.....	4,341	7	20	11	30	36	23	28	30	61	94	117
2001.....	4,187	7	17	14	18	22	26	26	44	59	69	102
2000.....	4,238	6	14	14	29	31	20	31	41	52	78	125
1999.....	4,500	9	13	14	34	23	29	34	35	46	80	126

Black, female

2003.....	3,462	5	5	7	13	15	18	14	18	43	96	121
2002.....	3,490	7	10	10	25	7	13	15	31	49	94	116
2001.....	3,402	10	6	9	13	8	21	21	38	73	78	122
2000.....	3,369	5	7	10	17	13	17	20	24	52	93	134
1999.....	3,415	9	5	6	14	13	13	34	41	49	89	131

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

American Indian 1/, both sexes

2003.....	512	-	1	-	-	1	1	-	3	5	9	12
2002.....	452	-	1	1	-	1	1	1	-	2	4	10
2001.....	427	1	-	1	-	2	1	1	1	5	5	8
2000.....	429	1	2	-	-	1	2	4	-	4	6	10
1999.....	406	1	-	-	-	-	1	-	2	2	7	8

American Indian 1/, male

2003.....	269	-	-	-	-	1	1	-	1	2	6	6
2002.....	220	-	1	-	-	-	-	1	-	2	-	5
2001.....	200	1	-	-	-	1	1	-	1	-	3	3
2000.....	228	1	1	-	-	1	2	3	-	1	3	7
1999.....	197	-	-	-	-	-	-	-	1	2	3	5

American Indian 1/, female

2003.....	243	-	1	-	-	-	-	-	2	3	3	6
2002.....	232	-	-	1	-	1	1	-	-	-	4	5
2001.....	227	-	-	1	-	1	-	1	-	5	2	5
2000.....	201	-	1	-	-	-	-	1	-	3	3	3
1999.....	209	1	-	-	-	-	1	-	1	-	4	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Asian or Pacific Islander 2/, both sexes

2003.....	1,244	-	4	-	1	4	2	3	3	4	10	11
2002.....	1,138	-	4	2	2	2	-	4	3	2	6	13
2001.....	1,178	1	-	-	1	2	4	1	2	6	9	13
2000.....	1,133	1	-	1	5	1	2	3	7	8	13	23
1999.....	1,125	1	5	-	-	3	2	4	4	5	4	13

Asian or Pacific Islander 2/, male

2003.....	801	-	3	-	1	3	1	2	2	2	6	6
2002.....	743	-	2	1	1	1	-	4	1	-	4	6
2001.....	749	1	-	-	1	1	3	-	1	3	5	6
2000.....	722	-	-	-	4	-	1	2	4	7	4	13
1999.....	718	-	3	-	-	1	2	3	1	4	4	6

Asian or Pacific Islander 2/, female

2003.....	443	-	1	-	-	1	1	1	1	2	4	5
2002.....	395	-	2	1	1	1	-	-	2	2	2	7
2001.....	429	-	-	-	-	1	1	1	1	3	4	7
2000.....	411	1	-	1	1	1	1	1	3	1	9	10
1999.....	407	1	2	-	-	2	-	1	3	1	-	7

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified (J40-J42)												
All races, both sexes												
2003.....	850	20	13	1	5	2	5	5	5	8	9	17
2002.....	955	24	13	3	2	3	3	2	4	12	6	16
2001.....	959	21	8	-	-	3	2	4	2	9	12	21
2000.....	1,167	18	14	2	3	3	2	3	5	7	13	19
1999.....	1,172	20	19	3	1	3	4	3	8	10	8	14
All races, male												
2003.....	351	15	6	-	3	2	4	1	2	5	4	13
2002.....	420	15	8	2	-	2	1	-	2	9	3	13
2001.....	396	11	5	-	-	2	1	2	-	3	5	12
2000.....	503	9	7	2	1	2	2	3	4	4	10	8
1999.....	498	9	11	2	1	2	2	-	3	3	2	9
All races, female												
2003.....	499	5	7	1	2	-	1	4	3	3	5	4
2002.....	535	9	5	1	2	1	2	2	2	3	3	3
2001.....	563	10	3	-	-	1	1	2	2	6	7	9
2000.....	664	9	7	-	2	1	-	-	1	3	3	11
1999.....	674	11	8	1	-	1	2	3	5	7	6	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified (J40-J42)												
White, both sexes												
2003.....	751	15	8	1	2	1	5	1	5	6	7	11
2002.....	876	16	7	2	1	1	3	1	4	11	5	12
2001.....	878	11	6	-	-	2	2	3	2	9	9	16
2000.....	1,073	9	7	2	2	2	2	3	3	7	7	17
1999.....	1,056	11	15	3	1	3	3	2	6	10	6	9
White, male												
2003.....	309	12	5	-	1	1	4	-	2	4	3	9
2002.....	376	11	3	2	-	1	1	-	2	8	2	9
2001.....	354	7	5	-	-	1	1	1	-	3	4	9
2000.....	458	5	3	2	-	1	2	3	2	4	4	8
1999.....	440	6	8	2	1	2	1	-	2	3	2	5
White, female												
2003.....	442	3	3	1	1	-	1	1	3	2	4	2
2002.....	500	5	4	-	1	-	2	1	2	3	3	3
2001.....	524	4	1	-	-	1	1	2	2	6	5	7
2000.....	615	4	4	-	2	1	-	-	1	3	3	9
1999.....	616	5	7	1	-	1	2	2	4	7	4	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified (J40-J42)												
Black, both sexes												
2003.....	75	5	3	-	3	1	-	3	-	2	2	6
2002.....	66	8	6	1	1	2	-	1	-	1	1	4
2001.....	63	10	2	-	-	1	-	1	-	-	3	4
2000.....	80	7	6	-	1	1	-	-	1	-	6	2
1999.....	89	7	1	-	-	-	1	1	2	-	2	4
Black, male												
2003.....	30	3	-	-	2	1	-	1	-	1	1	4
2002.....	35	4	5	-	-	1	-	-	-	1	1	4
2001.....	32	4	-	-	-	1	-	1	-	-	1	3
2000.....	35	3	3	-	1	1	-	-	1	-	6	-
1999.....	44	3	1	-	-	-	1	-	1	-	-	3
Black, female												
2003.....	45	2	3	-	1	-	-	2	-	1	1	2
2002.....	31	4	1	1	1	1	-	1	-	-	-	-
2001.....	31	6	2	-	-	-	-	-	-	-	2	1
2000.....	45	4	3	-	-	-	-	-	-	-	-	2
1999.....	45	4	-	-	-	-	-	1	1	-	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
All races, both sexes												
2003.....	14,861	-	1	-	-	1	3	2	3	21	66	157
2002.....	15,489	1	3	-	-	1	1	2	8	21	73	156
2001.....	16,242	3	-	-	1	1	1	1	6	24	51	141
2000.....	16,700	3	2	-	-	1	4	6	5	23	59	162
1999.....	17,787	1	-	1	-	-	1	1	8	19	78	136
All races, male												
2003.....	7,549	-	1	-	-	-	2	2	1	14	35	105
2002.....	8,084	-	3	-	-	-	-	1	7	14	52	95
2001.....	8,415	1	-	-	1	1	-	-	4	17	35	89
2000.....	8,671	3	-	-	-	1	4	1	4	15	31	101
1999.....	9,480	-	-	-	-	-	1	-	6	9	40	85
All races, female												
2003.....	7,312	-	-	-	-	1	1	-	2	7	31	52
2002.....	7,405	1	-	-	-	1	1	1	1	7	21	61
2001.....	7,827	2	-	-	-	-	1	1	2	7	16	52
2000.....	8,029	-	2	-	-	-	-	5	1	8	28	61
1999.....	8,307	1	-	1	-	-	-	1	2	10	38	51

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
White, both sexes												
2003.....	13,856	-	1	-	-	1	3	1	2	17	53	129
2002.....	14,476	1	1	-	-	1	-	2	6	17	50	129
2001.....	15,243	2	-	-	-	1	-	1	3	22	42	127
2000.....	15,607	3	-	-	-	-	2	6	4	18	45	121
1999.....	16,674	-	-	1	-	-	-	1	6	15	55	104
White, male												
2003.....	6,915	-	1	-	-	-	2	1	-	11	28	86
2002.....	7,426	-	1	-	-	-	-	1	5	11	37	78
2001.....	7,798	-	-	-	-	1	-	-	2	16	28	82
2000.....	7,985	3	-	-	-	-	2	1	3	11	23	72
1999.....	8,754	-	-	-	-	-	-	-	4	5	30	64
White, female												
2003.....	6,941	-	-	-	-	1	1	-	2	6	25	43
2002.....	7,050	1	-	-	-	1	-	1	1	6	13	51
2001.....	7,445	2	-	-	-	-	-	1	1	6	14	45
2000.....	7,622	-	-	-	-	-	-	5	1	7	22	49
1999.....	7,920	-	-	1	-	-	-	1	2	10	25	40

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
All races, both sexes												
2003.....	4,099	7	37	35	75	70	88	98	129	152	259	317
2002.....	4,261	4	43	35	88	84	85	102	133	207	265	303
2001.....	4,269	10	31	41	58	60	80	88	145	207	248	279
2000.....	4,487	8	32	43	86	78	89	105	144	183	275	318
1999.....	4,657	5	27	40	86	95	87	121	141	206	241	312
All races, male												
2003.....	1,493	4	29	23	48	40	51	58	69	60	100	127
2002.....	1,580	3	30	20	49	63	47	63	66	92	106	113
2001.....	1,479	6	22	24	33	35	45	49	67	93	102	103
2000.....	1,632	6	20	22	52	52	50	64	78	81	115	128
1999.....	1,620	3	16	25	60	53	55	64	64	94	90	96
All races, female												
2003.....	2,606	3	8	12	27	30	37	40	60	92	159	190
2002.....	2,681	1	13	15	39	21	38	39	67	115	159	190
2001.....	2,790	4	9	17	25	25	35	39	78	114	146	176
2000.....	2,855	2	12	21	34	26	39	41	66	102	160	190
1999.....	3,037	2	11	15	26	42	32	57	77	112	151	216

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
White, both sexes												
2003.....	2,888	4	13	12	37	31	47	59	72	90	152	197
2002.....	3,014	2	18	15	32	41	49	57	88	121	145	187
2001.....	2,990	5	11	17	28	31	36	43	71	99	148	164
2000.....	3,144	5	18	18	36	36	53	49	83	95	161	178
1999.....	3,328	1	11	20	39	59	47	55	72	123	143	191
White, male												
2003.....	961	4	8	7	22	16	28	29	31	32	59	78
2002.....	1,017	1	15	10	18	28	24	31	43	43	54	68
2001.....	937	4	6	10	16	15	20	25	26	48	56	58
2000.....	1,057	3	9	8	20	24	31	29	41	37	68	79
1999.....	1,046	1	5	11	27	32	28	28	30	53	46	55
White, female												
2003.....	1,927	-	5	5	15	15	19	30	41	58	93	119
2002.....	1,997	1	3	5	14	13	25	26	45	78	91	119
2001.....	2,053	1	5	7	12	16	16	18	45	51	92	106
2000.....	2,087	2	9	10	16	12	22	20	42	58	93	99
1999.....	2,282	-	6	9	12	27	19	27	42	70	97	136

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Black, both sexes												
2003.....	1,030	3	22	23	37	36	39	37	53	57	100	109
2002.....	1,096	2	22	17	54	40	35	41	44	83	116	104
2001.....	1,108	4	20	23	29	27	40	44	72	100	90	107
2000.....	1,158	3	14	24	45	40	34	49	55	79	100	128
1999.....	1,145	4	14	20	47	33	38	63	67	77	91	111
Black, male												
2003.....	445	-	20	16	25	22	21	27	36	26	37	43
2002.....	497	2	14	9	30	34	23	28	23	47	51	40
2001.....	459	1	16	14	16	19	22	24	39	42	39	43
2000.....	481	3	11	14	28	27	17	30	34	38	41	44
1999.....	481	2	10	14	33	20	26	34	34	36	38	38
Black, female												
2003.....	585	3	2	7	12	14	18	10	17	31	63	66
2002.....	599	-	8	8	24	6	12	13	21	36	65	64
2001.....	649	3	4	9	13	8	18	20	33	58	51	64
2000.....	677	-	3	10	17	13	17	19	21	41	59	84
1999.....	664	2	4	6	14	13	12	29	33	41	53	73

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
American Indian 1/, both sexes												
2003.....	38	-	-	-	-	-	1	-	3	3	2	5
2002.....	25	-	-	1	-	1	1	-	-	2	2	3
2001.....	33	1	-	1	-	-	1	1	1	4	3	2
2000.....	31	-	-	-	-	1	1	4	-	2	5	1
1999.....	29	-	-	-	-	-	1	-	1	1	3	2
American Indian 1/, male												
2003.....	18	-	-	-	-	-	1	-	1	2	2	2
2002.....	9	-	-	-	-	-	-	-	-	2	-	1
2001.....	11	1	-	-	-	-	1	-	1	-	2	-
2000.....	14	-	-	-	-	1	1	3	-	-	2	-
1999.....	12	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, female												
2003.....	20	-	-	-	-	-	-	-	2	1	-	3
2002.....	16	-	-	1	-	1	1	-	-	-	2	2
2001.....	22	-	-	1	-	-	-	1	-	4	1	2
2000.....	17	-	-	-	-	-	-	1	-	2	3	1
1999.....	17	-	-	-	-	-	1	-	1	-	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Asian or Pacific Islander 2/, both sexes												
2003.....	143	-	2	-	1	3	1	2	1	2	5	6
2002.....	126	-	3	2	2	2	-	4	1	1	2	9
2001.....	138	-	-	-	1	2	3	-	1	4	7	6
2000.....	154	-	-	1	5	1	1	3	6	7	9	11
1999.....	155	-	2	-	-	3	1	3	1	5	4	8
Asian or Pacific Islander 2/, male												
2003.....	69	-	1	-	1	2	1	2	1	-	2	4
2002.....	57	-	1	1	1	1	-	4	-	-	1	4
2001.....	72	-	-	-	1	1	2	-	1	3	5	2
2000.....	80	-	-	-	4	-	1	2	3	6	4	5
1999.....	81	-	1	-	-	1	1	2	-	4	4	3
Asian or Pacific Islander 2/, female												
2003.....	74	-	1	-	-	1	-	-	-	2	3	2
2002.....	69	-	2	1	1	1	-	-	1	1	1	5
2001.....	66	-	-	-	-	1	1	-	-	1	2	4
2000.....	74	-	-	1	1	1	-	1	3	1	5	6
1999.....	74	-	1	-	-	2	-	1	1	1	-	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic lower respiratory diseases (J44,J47)												
All races, both sexes												
2003.....	106,572	4	4	1	1	11	11	13	27	120	315	785
2002.....	104,111	10	6	3	5	5	10	10	40	100	324	751
2001.....	101,543	7	4	1	3	10	14	9	35	115	303	686
2000.....	99,655	7	3	3	2	4	9	11	17	97	273	675
1999.....	100,565	10	8	5	3	9	10	20	28	67	261	617
All races, male												
2003.....	51,321	4	2	-	1	8	7	6	17	63	148	436
2002.....	50,629	5	3	2	1	3	4	6	20	44	177	396
2001.....	49,407	4	2	-	3	5	10	4	20	58	142	387
2000.....	49,198	5	1	2	1	3	6	4	7	45	136	380
1999.....	50,817	7	3	2	3	5	5	5	10	36	136	357
All races, female												
2003.....	55,251	-	2	1	-	3	4	7	10	57	167	349
2002.....	53,482	5	3	1	4	2	6	4	20	56	147	355
2001.....	52,136	3	2	1	-	5	4	5	15	57	161	299
2000.....	50,457	2	2	1	1	1	3	7	10	52	137	295
1999.....	49,748	3	5	3	-	4	5	15	18	31	125	260

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic lower respiratory diseases (J44,J47)

White, both sexes

2003.....	99,422	4	2	1	1	5	8	9	20	98	252	675
2002.....	97,029	6	4	-	5	4	10	8	23	77	270	642
2001.....	94,708	4	3	1	2	6	7	6	27	82	254	575
2000.....	93,016	6	3	3	2	2	6	9	9	74	217	566
1999.....	93,677	4	5	5	2	6	7	15	19	52	204	497

White, male

2003.....	47,212	4	-	-	1	3	5	4	12	53	115	376
2002.....	46,590	4	2	-	1	2	4	5	14	34	147	334
2001.....	45,472	2	1	-	2	2	5	3	17	42	119	331
2000.....	45,316	5	1	2	1	1	4	3	2	33	112	313
1999.....	46,760	3	1	2	2	2	3	4	10	29	103	286

White, female

2003.....	52,210	-	2	1	-	2	3	5	8	45	137	299
2002.....	50,439	2	2	-	4	2	6	3	9	43	123	308
2001.....	49,236	2	2	1	-	4	2	3	10	40	135	244
2000.....	47,700	1	2	1	1	1	2	6	7	41	105	253
1999.....	46,917	1	4	3	-	4	4	11	9	23	101	211

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic lower respiratory diseases (J44,J47)												
Black, both sexes												
2003.....	5,794	-	1	-	-	4	2	4	5	19	52	99
2002.....	5,819	4	1	3	-	1	-	1	15	22	49	99
2001.....	5,607	2	1	-	1	2	6	2	7	31	45	100
2000.....	5,468	1	-	-	-	2	1	2	8	21	52	91
1999.....	5,751	6	3	-	1	3	3	4	7	15	54	110
Black, male												
2003.....	3,274	-	1	-	-	3	2	2	4	9	26	55
2002.....	3,268	1	-	2	-	1	-	-	5	10	27	57
2001.....	3,209	1	1	-	1	2	4	1	3	16	22	50
2000.....	3,163	-	-	-	-	2	1	1	5	11	23	55
1999.....	3,377	4	2	-	1	3	2	-	-	7	32	64
Black, female												
2003.....	2,520	-	-	-	-	1	-	2	1	10	26	44
2002.....	2,551	3	1	1	-	-	-	1	10	12	22	42
2001.....	2,398	1	-	-	-	-	2	1	4	15	23	50
2000.....	2,305	1	-	-	-	-	-	1	3	10	29	36
1999.....	2,374	2	1	-	-	-	1	4	7	8	22	46

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic lower respiratory diseases (J44,J47)

Asian or Pacific Islander 2/, both sexes

2003.....	933	-	1	-	-	1	1	-	2	1	4	5
2002.....	884	-	-	-	-	-	-	-	2	1	4	3
2001.....	888	1	-	-	-	-	1	1	1	1	2	6
2000.....	834	-	-	-	-	-	1	-	-	1	3	10
1999.....	812	-	-	-	-	-	-	1	2	-	-	5

Asian or Pacific Islander 2/, male

2003.....	613	-	1	-	-	1	-	-	1	1	3	2
2002.....	591	-	-	-	-	-	-	-	1	-	3	1
2001.....	567	1	-	-	-	-	1	-	-	-	-	3
2000.....	536	-	-	-	-	-	-	-	-	1	-	6
1999.....	520	-	-	-	-	-	-	1	-	-	-	3

Asian or Pacific Islander 2/, female

2003.....	320	-	-	-	-	-	1	-	1	-	1	3
2002.....	293	-	-	-	-	-	-	-	1	1	1	2
2001.....	321	-	-	-	-	-	-	1	1	1	2	3
2000.....	298	-	-	-	-	-	1	-	-	-	3	4
1999.....	292	-	-	-	-	-	-	-	2	-	-	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumoconioses and chemical effects (J60-J66,J68)

All races, both sexes

2003.....	1,114	-	-	-	-	-	1	-	1	2	5	4
2002.....	1,114	2	-	-	-	-	-	-	-	-	5	5
2001.....	1,136	-	1	-	-	-	-	-	1	3	2	3
2000.....	1,161	-	-	-	-	-	-	1	2	2	5	5
1999.....	1,113	-	3	-	-	1	-	-	-	3	4	3

All races, male

2003.....	1,065	-	-	-	-	-	1	-	1	2	4	4
2002.....	1,057	-	-	-	-	-	-	-	-	-	4	4
2001.....	1,095	-	1	-	-	-	-	-	1	3	2	2
2000.....	1,113	-	-	-	-	-	-	1	2	1	5	5
1999.....	1,079	-	3	-	-	1	-	-	-	2	3	3

All races, female

2003.....	49	-	-	-	-	-	-	-	-	-	1	-
2002.....	57	2	-	-	-	-	-	-	-	-	1	1
2001.....	41	-	-	-	-	-	-	-	-	-	-	1
2000.....	48	-	-	-	-	-	-	-	-	1	-	-
1999.....	34	-	-	-	-	-	-	-	-	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumoconioses and chemical effects (J60-J66,J68)

Black, both sexes

2003.....	48	-	-	-	-	-	-	-	1	-	1	1
2002.....	58	1	-	-	-	-	-	-	-	-	1	2
2001.....	57	-	-	-	-	-	-	-	-	-	-	2
2000.....	55	-	-	-	-	-	-	-	-	1	2	-
1999.....	67	-	3	-	-	1	-	-	-	2	1	-

Black, male

2003.....	45	-	-	-	-	-	-	-	1	-	-	1
2002.....	52	-	-	-	-	-	-	-	-	-	1	2
2001.....	55	-	-	-	-	-	-	-	-	-	-	1
2000.....	53	-	-	-	-	-	-	-	-	1	2	-
1999.....	64	-	3	-	-	1	-	-	-	1	1	-

Black, female

2003.....	3	-	-	-	-	-	-	-	-	-	1	-
2002.....	6	1	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	1
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
All races, both sexes												
2003.....	17,335	13	10	4	12	13	33	26	43	66	100	188
2002.....	17,593	2	7	7	9	16	28	31	43	72	120	143
2001.....	17,301	12	14	5	10	10	24	31	48	61	118	142
2000.....	16,636	10	11	8	5	12	25	22	40	65	116	157
1999.....	15,268	16	12	4	12	9	23	33	45	59	107	129
All races, male												
2003.....	8,702	8	5	4	5	6	25	16	26	39	50	110
2002.....	8,876	-	3	3	6	10	17	17	25	41	78	90
2001.....	8,576	6	6	3	4	5	17	15	25	42	75	81
2000.....	8,390	4	5	5	4	8	15	11	24	45	70	99
1999.....	7,755	9	3	-	6	7	15	21	26	35	62	83
All races, female												
2003.....	8,633	5	5	-	7	7	8	10	17	27	50	78
2002.....	8,717	2	4	4	3	6	11	14	18	31	42	53
2001.....	8,725	6	8	2	6	5	7	16	23	19	43	61
2000.....	8,246	6	6	3	1	4	10	11	16	20	46	58
1999.....	7,513	7	9	4	6	2	8	12	19	24	45	46

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to solids and liquids (J69)

White, both sexes

2003.....	15,395	7	8	3	10	8	21	19	36	50	74	152
2002.....	15,773	2	5	6	8	13	21	27	32	55	92	118
2001.....	15,425	9	12	4	7	8	17	26	36	45	97	114
2000.....	14,869	8	9	8	4	10	17	16	30	43	99	115
1999.....	13,582	13	8	4	8	8	14	23	34	43	78	98

White, male

2003.....	7,731	4	5	3	4	4	17	11	22	31	40	89
2002.....	7,954	-	3	3	5	8	12	16	19	31	59	74
2001.....	7,705	6	4	3	2	4	14	14	21	31	63	71
2000.....	7,488	4	4	5	3	7	11	7	15	32	61	72
1999.....	6,853	7	2	-	5	6	8	13	20	26	45	66

White, female

2003.....	7,664	3	3	-	6	4	4	8	14	19	34	63
2002.....	7,819	2	2	3	3	5	9	11	13	24	33	44
2001.....	7,720	3	8	1	5	4	3	12	15	14	34	43
2000.....	7,381	4	5	3	1	3	6	9	15	11	38	43
1999.....	6,729	6	6	4	3	2	6	10	14	17	33	32

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to solids and liquids (J69)

Black, both sexes

2003.....	1,607	4	1	1	2	4	10	6	6	15	26	34
2002.....	1,537	-	2	1	1	2	6	4	10	11	27	21
2001.....	1,599	3	2	1	3	1	6	5	11	13	21	25
2000.....	1,521	1	2	-	1	2	7	6	9	21	17	39
1999.....	1,505	2	4	-	3	1	9	9	9	14	27	28

Black, male

2003.....	781	3	-	1	1	2	6	5	4	7	10	21
2002.....	760	-	-	-	1	1	4	1	5	5	18	12
2001.....	741	-	2	-	2	-	2	1	3	8	12	8
2000.....	765	-	1	-	1	1	3	4	8	12	9	26
1999.....	775	1	1	-	1	1	7	7	5	8	15	15

Black, female

2003.....	826	1	1	-	1	2	4	1	2	8	16	13
2002.....	777	-	2	1	-	1	2	3	5	6	9	9
2001.....	858	3	-	1	1	1	4	4	8	5	9	17
2000.....	756	1	1	-	-	1	4	2	1	9	8	13
1999.....	730	1	3	-	2	-	2	2	4	6	12	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to solids and liquids (J69)

American Indian 1/, both sexes

2003.....	70	2	1	-	-	-	-	-	1	-	-	1
2002.....	73	-	-	-	-	1	-	-	1	3	1	2
2001.....	59	-	-	-	-	-	1	-	1	2	-	2
2000.....	49	-	-	-	-	-	-	-	1	-	-	2
1999.....	46	-	-	-	-	-	-	1	2	1	1	1

American Indian 1/, male

2003.....	33	1	-	-	-	-	-	-	-	-	-	-
2002.....	44	-	-	-	-	1	-	-	1	2	1	2
2001.....	23	-	-	-	-	-	1	-	1	2	-	1
2000.....	27	-	-	-	-	-	-	-	1	-	-	-
1999.....	28	-	-	-	-	-	-	1	1	-	1	1

American Indian 1/, female

2003.....	37	1	1	-	-	-	-	-	1	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	1	-	-
2001.....	36	-	-	-	-	-	-	-	-	-	-	1
2000.....	22	-	-	-	-	-	-	-	-	-	-	2
1999.....	18	-	-	-	-	-	-	-	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to solids and liquids (J69)

Asian or Pacific Islander 2/, both sexes

2003.....	263	-	-	-	-	1	2	1	-	1	-	1
2002.....	210	-	-	-	-	-	1	-	-	3	-	2
2001.....	218	-	-	-	-	1	-	-	-	1	-	1
2000.....	197	1	-	-	-	-	1	-	-	1	-	1
1999.....	135	1	-	-	1	-	-	-	-	1	1	2

Asian or Pacific Islander 2/, male

2003.....	157	-	-	-	-	-	2	-	-	1	-	-
2002.....	118	-	-	-	-	-	1	-	-	3	-	2
2001.....	107	-	-	-	-	1	-	-	-	1	-	1
2000.....	110	-	-	-	-	-	1	-	-	1	-	1
1999.....	99	1	-	-	-	-	-	-	-	1	1	1

Asian or Pacific Islander 2/, female

2003.....	106	-	-	-	-	1	-	1	-	-	-	1
2002.....	92	-	-	-	-	-	-	-	-	-	-	-
2001.....	111	-	-	-	-	-	-	-	-	-	-	-
2000.....	87	1	-	-	-	-	-	-	-	-	-	-
1999.....	36	-	-	-	1	-	-	-	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races, both sexes

2003.....	25,520	279	86	25	35	59	77	80	130	188	307	573
2002.....	25,039	278	85	41	43	51	75	70	115	191	337	522
2001.....	26,178	294	93	30	48	55	86	97	119	218	352	542
2000.....	25,535	273	87	36	38	60	71	99	154	231	384	527
1999.....	25,031	252	91	37	39	59	56	97	160	228	348	525

All races, male

2003.....	12,391	163	49	16	22	31	45	49	66	97	154	301
2002.....	12,057	164	56	21	18	29	39	31	57	101	180	263
2001.....	12,495	186	47	18	25	30	46	45	59	105	199	295
2000.....	12,391	154	45	21	17	30	43	48	79	132	203	281
1999.....	12,110	145	55	20	19	37	32	45	82	125	192	269

All races, female

2003.....	13,129	116	37	9	13	28	32	31	64	91	153	272
2002.....	12,982	114	29	20	25	22	36	39	58	90	157	259
2001.....	13,683	108	46	12	23	25	40	52	60	113	153	247
2000.....	13,144	119	42	15	21	30	28	51	75	99	181	246
1999.....	12,921	107	36	17	20	22	24	52	78	103	156	256

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2003.....	22,706	169	53	21	25	40	48	55	92	128	206	402
2002.....	22,184	151	56	29	28	36	45	50	82	132	247	366
2001.....	23,133	156	61	22	37	38	62	61	78	136	252	388
2000.....	22,664	166	63	24	29	39	47	72	110	152	291	374
1999.....	22,161	147	62	25	30	47	43	65	113	155	240	374

White, male

2003.....	11,014	92	28	15	14	17	29	34	49	69	112	225
2002.....	10,693	87	39	17	10	19	29	22	41	70	131	186
2001.....	11,085	95	28	14	20	21	34	26	38	72	155	216
2000.....	10,979	100	30	14	14	17	29	35	58	87	154	199
1999.....	10,764	85	34	13	16	31	27	39	59	88	140	193

White, female

2003.....	11,692	77	25	6	11	23	19	21	43	59	94	177
2002.....	11,491	64	17	12	18	17	16	28	41	62	116	180
2001.....	12,048	61	33	8	17	17	28	35	40	64	97	172
2000.....	11,685	66	33	10	15	22	18	37	52	65	137	175
1999.....	11,397	62	28	12	14	16	16	26	54	67	100	181

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2003.....	2,233	99	27	2	6	15	26	22	34	52	89	149
2002.....	2,279	112	29	12	13	15	26	18	29	51	83	140
2001.....	2,491	117	26	6	11	15	23	29	39	72	94	141
2000.....	2,361	103	21	10	8	18	22	22	37	70	83	142
1999.....	2,356	93	26	10	9	10	11	29	45	68	95	133

Black, male

2003.....	1,097	62	18	1	5	12	15	13	15	26	34	67
2002.....	1,084	67	17	4	6	10	9	7	16	26	47	67
2001.....	1,131	77	14	3	5	8	11	14	20	27	43	71
2000.....	1,135	51	14	7	3	11	12	12	18	41	44	79
1999.....	1,092	54	19	6	3	5	4	5	22	34	43	70

Black, female

2003.....	1,136	37	9	1	1	3	11	9	19	26	55	82
2002.....	1,195	45	12	8	7	5	17	11	13	25	36	73
2001.....	1,360	40	12	3	6	7	12	15	19	45	51	70
2000.....	1,226	52	7	3	5	7	10	10	19	29	39	63
1999.....	1,264	39	7	4	6	5	7	24	23	34	52	63

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

American Indian 1/, both sexes

2003.....	166	2	-	1	1	2	-	-	-	4	3	10
2002.....	146	5	-	-	-	-	2	-	3	3	3	7
2001.....	177	7	3	-	-	1	1	3	1	5	4	6
2000.....	157	1	2	-	1	1	-	3	4	4	4	6
1999.....	148	3	1	-	-	1	-	3	-	-	5	7

American Indian 1/, male

2003.....	74	2	-	-	1	1	-	-	-	1	2	3
2002.....	73	4	-	-	-	-	1	-	-	1	1	4
2001.....	89	5	3	-	-	-	1	3	1	4	1	3
2000.....	73	-	1	-	-	1	-	-	2	3	2	1
1999.....	71	3	1	-	-	1	-	1	-	-	2	3

American Indian 1/, female

2003.....	92	-	-	1	-	1	-	-	-	3	1	7
2002.....	73	1	-	-	-	-	1	-	3	2	2	3
2001.....	88	2	-	-	-	1	-	-	-	1	3	3
2000.....	84	1	1	-	1	-	-	3	2	1	2	5
1999.....	77	-	-	-	-	-	-	2	-	-	3	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Asian or Pacific Islander 2/, both sexes

2003.....	415	9	6	1	3	2	3	3	4	4	9	12
2002.....	430	10	-	-	2	-	2	2	1	5	4	9
2001.....	377	14	3	2	-	1	-	4	1	5	2	7
2000.....	353	3	1	2	-	2	2	2	3	5	6	5
1999.....	366	9	2	2	-	1	2	-	2	5	8	11

Asian or Pacific Islander 2/, male

2003.....	206	7	3	-	2	1	1	2	2	1	6	6
2002.....	207	6	-	-	2	-	-	2	-	4	1	6
2001.....	190	9	2	1	-	1	-	2	-	2	-	5
2000.....	204	3	-	-	-	1	2	1	1	1	3	2
1999.....	183	3	1	1	-	-	1	-	1	3	7	3

Asian or Pacific Islander 2/, female

2003.....	209	2	3	1	1	1	2	1	2	3	3	6
2002.....	223	4	-	-	-	-	2	-	1	1	3	3
2001.....	187	5	1	1	-	-	-	2	1	3	2	2
2000.....	149	-	1	2	-	1	-	1	2	4	3	3
1999.....	183	6	1	1	-	1	1	-	1	2	1	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
All races, both sexes												
2003.....	3,913	2	1	-	3	3	4	8	17	33	68	122
2002.....	4,079	1	1	-	1	-	-	5	17	40	73	118
2001.....	4,491	-	1	3	2	3	3	13	19	36	82	137
2000.....	4,558	-	-	-	-	4	5	3	11	42	71	130
1999.....	4,604	2	2	-	-	1	1	7	14	38	85	113
All races, male												
2003.....	1,880	-	1	-	1	3	3	4	9	20	40	82
2002.....	1,951	-	1	-	1	-	-	2	12	28	44	71
2001.....	2,074	-	1	2	-	2	1	10	11	25	62	91
2000.....	2,201	-	-	-	-	3	4	1	5	31	48	86
1999.....	2,189	-	-	-	-	1	1	5	7	27	49	67
All races, female												
2003.....	2,033	2	-	-	2	-	1	4	8	13	28	40
2002.....	2,128	1	-	-	-	-	-	3	5	12	29	47
2001.....	2,417	-	-	1	2	1	2	3	8	11	20	46
2000.....	2,357	-	-	-	-	1	1	2	6	11	23	44
1999.....	2,415	2	2	-	-	-	-	2	7	11	36	46

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Peptic ulcer (K25-K28)

White, both sexes

2003.....	3,408	1	1	-	2	2	2	6	14	22	50	90
2002.....	3,587	1	1	-	-	-	-	2	10	30	47	94
2001.....	3,942	-	1	3	1	2	1	5	11	19	59	102
2000.....	4,021	-	-	-	-	4	5	3	6	31	50	95
1999.....	4,040	1	2	-	-	1	1	6	9	21	56	80

White, male

2003.....	1,592	-	1	-	1	2	1	4	6	13	29	62
2002.....	1,685	-	1	-	-	-	-	2	8	20	30	56
2001.....	1,751	-	1	2	-	1	1	4	7	12	44	65
2000.....	1,900	-	-	-	-	3	4	1	3	22	34	68
1999.....	1,887	-	-	-	-	1	1	5	3	16	28	45

White, female

2003.....	1,816	1	-	-	1	-	1	2	8	9	21	28
2002.....	1,902	1	-	-	-	-	-	-	2	10	17	38
2001.....	2,191	-	-	1	1	1	-	1	4	7	15	37
2000.....	2,121	-	-	-	-	1	1	2	3	9	16	27
1999.....	2,153	1	2	-	-	-	-	1	6	5	28	35

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Peptic ulcer (K25-K28)

Black, both sexes

2003.....	383	1	-	-	1	1	-	2	3	7	16	32
2002.....	380	-	-	-	1	-	-	3	7	9	26	23
2001.....	409	-	-	-	1	1	1	8	7	16	19	27
2000.....	425	-	-	-	-	-	-	-	4	10	19	32
1999.....	448	1	-	-	-	-	-	-	4	16	28	28

Black, male

2003.....	223	-	-	-	-	1	-	-	3	3	10	20
2002.....	212	-	-	-	1	-	-	-	4	7	14	14
2001.....	243	-	-	-	-	1	-	6	4	13	14	21
2000.....	229	-	-	-	-	-	-	-	2	8	13	17
1999.....	243	-	-	-	-	-	-	-	3	11	20	20

Black, female

2003.....	160	1	-	-	1	-	-	2	-	4	6	12
2002.....	168	-	-	-	-	-	-	3	3	2	12	9
2001.....	166	-	-	-	1	-	1	2	3	3	5	6
2000.....	196	-	-	-	-	-	-	-	2	2	6	15
1999.....	205	1	-	-	-	-	-	-	1	5	8	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Peptic ulcer (K25-K28)

American Indian 1/, both sexes

2003.....	16	-	-	-	-	-	1	-	-	1	2	-
2002.....	17	-	-	-	-	-	-	-	-	-	-	-
2001.....	28	-	-	-	-	-	-	-	1	1	1	4
2000.....	19	-	-	-	-	-	-	-	-	1	-	2
1999.....	10	-	-	-	-	-	-	-	-	1	1	1

American Indian 1/, male

2003.....	7	-	-	-	-	-	1	-	-	1	1	-
2002.....	10	-	-	-	-	-	-	-	-	-	-	-
2001.....	17	-	-	-	-	-	-	-	-	-	1	3
2000.....	11	-	-	-	-	-	-	-	-	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	-	1	-

American Indian 1/, female

2003.....	9	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	1	1	-	1
2000.....	8	-	-	-	-	-	-	-	-	-	-	2
1999.....	5	-	-	-	-	-	-	-	-	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Peptic ulcer (K25-K28)

Asian or Pacific Islander 2/, both sexes

2003.....	106	-	-	-	-	-	1	-	-	3	-	-
2002.....	95	-	-	-	-	-	-	-	-	1	-	1
2001.....	112	-	-	-	-	-	1	-	-	-	3	4
2000.....	93	-	-	-	-	-	-	-	1	-	2	1
1999.....	106	-	-	-	-	-	-	1	1	-	-	4

Asian or Pacific Islander 2/, male

2003.....	58	-	-	-	-	-	1	-	-	3	-	-
2002.....	44	-	-	-	-	-	-	-	-	1	-	1
2001.....	63	-	-	-	-	-	-	-	-	-	3	2
2000.....	61	-	-	-	-	-	-	-	-	-	1	1
1999.....	54	-	-	-	-	-	-	-	1	-	-	2

Asian or Pacific Islander 2/, female

2003.....	48	-	-	-	-	-	-	-	-	-	-	-
2002.....	51	-	-	-	-	-	-	-	-	-	-	-
2001.....	49	-	-	-	-	-	1	-	-	-	-	2
2000.....	32	-	-	-	-	-	-	-	1	-	1	-
1999.....	52	-	-	-	-	-	-	1	-	-	-	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of appendix (K35-K38)

All races, both sexes

2003.....	439	4	5	6	5	6	5	5	11	10	15	25
2002.....	480	3	-	3	4	7	5	4	6	8	19	24
2001.....	428	1	8	3	3	5	10	7	8	9	17	25
2000.....	435	3	5	4	11	5	6	3	8	11	16	18
1999.....	390	3	2	12	14	2	3	3	3	13	11	25

All races, male

2003.....	258	2	2	2	3	4	5	2	7	5	11	14
2002.....	275	1	-	1	2	5	2	4	5	7	10	19
2001.....	252	-	7	3	2	3	5	5	6	7	10	17
2000.....	237	2	2	1	3	2	5	3	7	5	11	12
1999.....	236	1	2	6	9	2	3	3	2	10	7	18

All races, female

2003.....	181	2	3	4	2	2	-	3	4	5	4	11
2002.....	205	2	-	2	2	2	3	-	1	1	9	5
2001.....	176	1	1	-	1	2	5	2	2	2	7	8
2000.....	198	1	3	3	8	3	1	-	1	6	5	6
1999.....	154	2	-	6	5	-	-	-	1	3	4	7

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of appendix (K35-K38)

White, both sexes

2003.....	355	3	3	4	2	4	3	3	7	6	8	19
2002.....	404	2	-	2	2	4	3	3	5	5	13	17
2001.....	342	1	5	2	2	5	5	3	5	7	10	14
2000.....	348	3	4	3	10	5	3	1	3	7	11	10
1999.....	314	3	-	8	10	1	2	3	1	10	9	16

White, male

2003.....	213	2	2	1	-	3	3	1	4	5	7	11
2002.....	225	-	-	1	1	2	1	3	4	4	9	13
2001.....	198	-	4	2	1	3	2	2	3	6	6	10
2000.....	183	2	2	1	3	2	3	1	3	3	6	7
1999.....	184	1	-	4	6	1	2	3	-	8	5	11

White, female

2003.....	142	1	1	3	2	1	-	2	3	1	1	8
2002.....	179	2	-	1	1	2	2	-	1	1	4	4
2001.....	144	1	1	-	1	2	3	1	2	1	4	4
2000.....	165	1	2	2	7	3	-	-	-	4	5	3
1999.....	130	2	-	4	4	-	-	-	1	2	4	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of appendix (K35-K38)

Black, both sexes

2003.....	70	-	2	2	3	2	2	2	3	4	7	5
2002.....	63	1	-	1	2	3	2	1	1	-	5	7
2001.....	73	-	2	1	1	-	5	3	3	2	6	9
2000.....	77	-	-	1	1	-	2	2	5	4	5	8
1999.....	68	-	2	4	3	-	1	-	2	3	2	8

Black, male

2003.....	37	-	-	1	3	1	2	1	2	-	4	3
2002.....	42	1	-	-	1	3	1	1	1	-	1	6
2001.....	47	-	2	1	1	-	3	3	3	1	4	6
2000.....	48	-	-	-	-	-	1	2	4	2	5	5
1999.....	48	-	2	2	2	-	1	-	2	2	2	7

Black, female

2003.....	33	-	2	1	-	1	-	1	1	4	3	2
2002.....	21	-	-	1	1	-	1	-	-	-	4	1
2001.....	26	-	-	-	-	-	2	-	-	1	2	3
2000.....	29	-	-	1	1	-	1	-	1	2	-	3
1999.....	20	-	-	2	1	-	-	-	-	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hernia (K40-K46)

All races, both sexes

2003.....	1,613	37	6	4	2	-	3	1	4	7	21	45
2002.....	1,595	29	3	4	3	2	3	4	7	14	21	33
2001.....	1,522	43	2	1	2	2	1	6	9	13	21	24
2000.....	1,522	46	4	1	2	2	3	2	3	13	18	26
1999.....	1,464	44	3	1	1	1	2	6	10	15	24	16

All races, male

2003.....	668	20	4	2	2	-	1	1	3	6	10	24
2002.....	599	18	3	3	2	1	3	2	5	8	9	16
2001.....	587	28	1	-	-	1	-	4	6	5	11	13
2000.....	586	28	3	1	2	1	2	-	1	8	5	13
1999.....	597	26	2	1	-	1	2	5	3	10	13	12

All races, female

2003.....	945	17	2	2	-	-	2	-	1	1	11	21
2002.....	996	11	-	1	1	1	-	2	2	6	12	17
2001.....	935	15	1	1	2	1	1	2	3	8	10	11
2000.....	936	18	1	-	-	1	1	2	2	5	13	13
1999.....	867	18	1	-	1	-	-	1	7	5	11	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
White, both sexes												
2003.....	1,421	29	3	2	1	-	2	1	2	4	16	37
2002.....	1,433	24	2	4	3	2	3	2	4	11	15	24
2001.....	1,343	31	2	-	1	1	1	6	7	10	15	20
2000.....	1,354	36	2	1	2	2	1	2	2	9	13	20
1999.....	1,298	34	3	1	1	1	1	4	6	14	16	11
White, male												
2003.....	585	16	3	1	1	-	1	1	1	3	8	19
2002.....	533	14	2	3	2	1	3	-	3	6	8	13
2001.....	502	18	1	-	-	1	-	4	4	5	8	9
2000.....	520	23	2	1	2	1	-	-	1	6	4	9
1999.....	535	20	2	1	-	1	1	3	2	9	11	9
White, female												
2003.....	836	13	-	1	-	-	1	-	1	1	8	18
2002.....	900	10	-	1	1	1	-	2	1	5	7	11
2001.....	841	13	1	-	1	-	1	2	3	5	7	11
2000.....	834	13	-	-	-	1	1	2	1	3	9	11
1999.....	763	14	1	-	1	-	-	1	4	5	5	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hernia (K40-K46)

Black, both sexes

2003.....	164	8	3	2	1	-	-	-	2	3	5	7
2002.....	137	4	1	-	-	-	-	2	1	3	5	9
2001.....	157	11	-	1	-	1	-	-	2	3	6	4
2000.....	145	7	2	-	-	-	2	-	1	3	5	5
1999.....	140	9	-	-	-	-	1	-	4	1	8	5

Black, male

2003.....	72	4	1	1	1	-	-	-	2	3	2	4
2002.....	56	3	1	-	-	-	-	2	-	2	1	3
2001.....	76	9	-	-	-	-	-	-	2	-	3	4
2000.....	54	3	1	-	-	-	2	-	-	1	1	4
1999.....	53	6	-	-	-	-	1	-	1	1	2	3

Black, female

2003.....	92	4	2	1	-	-	-	-	-	-	3	3
2002.....	81	1	-	-	-	-	-	-	1	1	4	6
2001.....	81	2	-	1	-	1	-	-	-	3	3	-
2000.....	91	4	1	-	-	-	-	-	1	2	4	1
1999.....	87	3	-	-	-	-	-	-	3	-	6	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All races, both sexes

2003.....	27,503	2	-	-	-	3	13	78	280	882	2,138	3,457
2002.....	27,257	6	3	1	5	7	24	77	297	929	2,225	3,404
2001.....	27,035	3	-	1	1	6	20	77	310	1,108	2,228	3,557
2000.....	26,552	3	3	-	3	7	29	88	327	1,083	2,288	3,374
1999.....	26,259	5	2	1	1	9	17	70	337	1,103	2,199	3,228

All races, male

2003.....	17,912	2	-	-	-	1	7	54	198	574	1,459	2,510
2002.....	17,401	6	1	1	2	3	11	46	190	605	1,474	2,442
2001.....	17,393	1	-	1	1	2	10	53	206	713	1,493	2,581
2000.....	17,214	3	1	-	2	3	16	63	217	716	1,598	2,515
1999.....	17,115	4	-	1	-	3	8	41	216	716	1,598	2,446

All races, female

2003.....	9,591	-	-	-	-	2	6	24	82	308	679	947
2002.....	9,856	-	2	-	3	4	13	31	107	324	751	962
2001.....	9,642	2	-	-	-	4	10	24	104	395	735	976
2000.....	9,338	-	2	-	1	4	13	25	110	367	690	859
1999.....	9,144	1	2	-	1	6	9	29	121	387	601	782

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

2003.....	24,005	1	-	-	-	2	7	53	233	729	1,811	2,910
2002.....	23,809	5	3	1	4	7	18	54	231	766	1,890	2,842
2001.....	23,408	3	-	-	1	2	8	51	256	919	1,826	2,958
2000.....	22,982	-	3	-	2	2	19	57	255	872	1,873	2,817
1999.....	22,620	4	-	-	1	7	8	52	271	905	1,771	2,596

White, male

2003.....	15,662	1	-	-	-	1	4	38	167	486	1,262	2,151
2002.....	15,278	5	1	1	2	3	9	34	159	504	1,279	2,089
2001.....	15,048	1	-	-	1	-	6	38	173	590	1,236	2,184
2000.....	15,002	-	1	-	1	1	10	46	178	588	1,333	2,160
1999.....	14,790	4	-	-	-	3	5	35	178	600	1,306	2,002

White, female

2003.....	8,343	-	-	-	-	1	3	15	66	243	549	759
2002.....	8,531	-	2	-	2	4	9	20	72	262	611	753
2001.....	8,360	2	-	-	-	2	2	13	83	329	590	774
2000.....	7,980	-	2	-	1	1	9	11	77	284	540	657
1999.....	7,830	-	-	-	1	4	3	17	93	305	465	594

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Black, both sexes

2003.....	2,621	1	-	-	-	-	3	18	23	101	225	394
2002.....	2,588	1	-	-	1	-	5	14	37	87	231	444
2001.....	2,775	-	-	1	-	2	8	18	29	118	304	472
2000.....	2,737	2	-	-	1	4	8	21	36	144	307	453
1999.....	2,833	-	2	1	-	2	8	16	39	129	328	516

Black, male

2003.....	1,727	1	-	-	-	-	2	11	14	58	133	263
2002.....	1,622	1	-	-	-	-	1	8	19	50	133	277
2001.....	1,843	-	-	1	-	1	1	10	16	78	196	323
2000.....	1,730	2	-	-	1	1	5	12	15	92	192	289
1999.....	1,842	-	-	1	-	-	3	5	23	75	221	366

Black, female

2003.....	894	-	-	-	-	-	1	7	9	43	92	131
2002.....	966	-	-	-	1	-	4	6	18	37	98	167
2001.....	932	-	-	-	-	1	7	8	13	40	108	149
2000.....	1,007	-	-	-	-	3	3	9	21	52	115	164
1999.....	991	-	2	-	-	2	5	11	16	54	107	150

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

American Indian 1/, both sexes

2003.....	570	-	-	-	-	1	3	6	19	45	82	108
2002.....	547	-	-	-	-	-	1	8	27	62	80	89
2001.....	533	-	-	-	-	2	4	7	18	57	78	95
2000.....	534	-	-	-	-	1	1	10	29	56	81	72
1999.....	513	-	-	-	-	-	1	2	19	57	78	88

American Indian 1/, male

2003.....	317	-	-	-	-	-	1	4	13	24	48	61
2002.....	319	-	-	-	-	-	1	4	11	39	43	54
2001.....	309	-	-	-	-	1	3	5	12	34	48	48
2000.....	298	-	-	-	-	1	-	5	18	27	51	42
1999.....	299	-	-	-	-	-	-	1	9	32	52	54

American Indian 1/, female

2003.....	253	-	-	-	-	1	2	2	6	21	34	47
2002.....	228	-	-	-	-	-	-	4	16	23	37	35
2001.....	224	-	-	-	-	1	1	2	6	23	30	47
2000.....	236	-	-	-	-	-	1	5	11	29	30	30
1999.....	214	-	-	-	-	-	1	1	10	25	26	34

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Asian or Pacific Islander 2/, both sexes

2003.....	307	-	-	-	-	-	-	1	5	7	20	45
2002.....	313	-	-	-	-	-	-	1	2	14	24	29
2001.....	319	-	-	-	-	-	-	1	7	14	20	32
2000.....	299	1	-	-	-	-	1	-	7	11	27	32
1999.....	293	1	-	-	-	-	-	-	8	12	22	28

Asian or Pacific Islander 2/, male

2003.....	206	-	-	-	-	-	-	1	4	6	16	35
2002.....	182	-	-	-	-	-	-	-	1	12	19	22
2001.....	193	-	-	-	-	-	-	-	5	11	13	26
2000.....	184	1	-	-	-	-	1	-	6	9	22	24
1999.....	184	-	-	-	-	-	-	-	6	9	19	24

Asian or Pacific Islander 2/, female

2003.....	101	-	-	-	-	-	-	-	1	1	4	10
2002.....	131	-	-	-	-	-	-	1	1	2	5	7
2001.....	126	-	-	-	-	-	-	1	2	3	7	6
2000.....	115	-	-	-	-	-	-	-	1	2	5	8
1999.....	109	1	-	-	-	-	-	-	2	3	3	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
All races, both sexes												
2003.....	12,360	-	-	-	-	1	5	49	194	612	1,359	2,050
2002.....	12,121	-	1	-	-	-	8	43	199	607	1,364	1,998
2001.....	12,207	-	-	-	-	1	7	45	219	745	1,400	2,121
2000.....	12,109	-	-	-	-	-	8	56	223	741	1,485	2,000
1999.....	11,958	-	-	-	-	-	8	37	222	763	1,423	1,965
All races, male												
2003.....	9,104	-	-	-	-	-	3	39	143	402	915	1,524
2002.....	8,814	-	1	-	-	-	6	32	130	392	913	1,450
2001.....	8,932	-	-	-	-	-	5	31	148	492	954	1,563
2000.....	8,985	-	-	-	-	-	5	45	157	486	1,047	1,510
1999.....	8,922	-	-	-	-	-	6	24	141	491	1,029	1,501
All races, female												
2003.....	3,256	-	-	-	-	1	2	10	51	210	444	526
2002.....	3,307	-	-	-	-	-	2	11	69	215	451	548
2001.....	3,275	-	-	-	-	1	2	14	71	253	446	558
2000.....	3,124	-	-	-	-	-	3	11	66	255	438	490
1999.....	3,036	-	-	-	-	-	2	13	81	272	394	464

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
White, both sexes												
2003.....	10,625	-	-	-	-	-	4	37	161	499	1,159	1,743
2002.....	10,423	-	1	-	-	-	4	31	156	501	1,159	1,700
2001.....	10,387	-	-	-	-	-	2	32	179	613	1,158	1,812
2000.....	10,272	-	-	-	-	-	6	42	176	596	1,214	1,670
1999.....	10,041	-	-	-	-	-	5	31	174	623	1,141	1,605
White, male												
2003.....	7,905	-	-	-	-	-	2	31	118	339	792	1,306
2002.....	7,688	-	1	-	-	-	4	25	110	326	786	1,257
2001.....	7,689	-	-	-	-	-	2	26	125	407	795	1,363
2000.....	7,770	-	-	-	-	-	4	36	132	397	867	1,301
1999.....	7,602	-	-	-	-	-	4	21	115	409	840	1,252
White, female												
2003.....	2,720	-	-	-	-	-	2	6	43	160	367	437
2002.....	2,735	-	-	-	-	-	-	6	46	175	373	443
2001.....	2,698	-	-	-	-	-	-	6	54	206	363	449
2000.....	2,502	-	-	-	-	-	2	6	44	199	347	369
1999.....	2,439	-	-	-	-	-	1	10	59	214	301	353

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Black, both sexes												
2003.....	1,222	-	-	-	-	-	-	6	13	67	121	199
2002.....	1,191	-	-	-	-	-	3	6	20	47	123	209
2001.....	1,348	-	-	-	-	-	1	7	20	75	170	221
2000.....	1,380	-	-	-	-	-	-	7	27	94	193	259
1999.....	1,466	-	-	-	-	-	2	4	25	84	199	292
Black, male												
2003.....	865	-	-	-	-	-	-	4	10	37	74	147
2002.....	797	-	-	-	-	-	1	4	10	27	76	136
2001.....	935	-	-	-	-	-	-	2	10	51	110	148
2000.....	918	-	-	-	-	-	-	5	12	61	126	164
1999.....	1,018	-	-	-	-	-	2	2	13	49	130	204
Black, female												
2003.....	357	-	-	-	-	-	-	2	3	30	47	52
2002.....	394	-	-	-	-	-	2	2	10	20	47	73
2001.....	413	-	-	-	-	-	1	5	10	24	60	73
2000.....	462	-	-	-	-	-	-	2	15	33	67	95
1999.....	448	-	-	-	-	-	-	2	12	35	69	88

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
American Indian 1/, both sexes												
2003.....	403	-	-	-	-	1	1	5	16	42	66	81
2002.....	381	-	-	-	-	-	1	6	22	49	63	71
2001.....	359	-	-	-	-	1	4	5	15	47	63	68
2000.....	339	-	-	-	-	-	1	7	18	42	57	52
1999.....	356	-	-	-	-	-	1	2	17	48	70	54
American Indian 1/, male												
2003.....	241	-	-	-	-	-	1	3	11	23	38	47
2002.....	230	-	-	-	-	-	1	3	10	31	35	41
2001.....	218	-	-	-	-	-	3	3	10	27	41	34
2000.....	205	-	-	-	-	-	-	4	11	20	36	30
1999.....	224	-	-	-	-	-	-	1	8	26	47	33
American Indian 1/, female												
2003.....	162	-	-	-	-	1	-	2	5	19	28	34
2002.....	151	-	-	-	-	-	-	3	12	18	28	30
2001.....	141	-	-	-	-	1	1	2	5	20	22	34
2000.....	134	-	-	-	-	-	1	3	7	22	21	22
1999.....	132	-	-	-	-	-	1	1	9	22	23	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Asian or Pacific Islander 2/, both sexes												
2003.....	110	-	-	-	-	-	-	1	4	4	13	27
2002.....	126	-	-	-	-	-	-	-	1	10	19	18
2001.....	113	-	-	-	-	-	-	1	5	10	9	20
2000.....	118	-	-	-	-	-	1	-	2	9	21	19
1999.....	95	-	-	-	-	-	-	-	6	8	13	14
Asian or Pacific Islander 2/, male												
2003.....	93	-	-	-	-	-	-	1	4	3	11	24
2002.....	99	-	-	-	-	-	-	-	-	8	16	16
2001.....	90	-	-	-	-	-	-	-	3	7	8	18
2000.....	92	-	-	-	-	-	1	-	2	8	18	15
1999.....	78	-	-	-	-	-	-	-	5	7	12	12
Asian or Pacific Islander 2/, female												
2003.....	17	-	-	-	-	-	-	-	-	1	2	3
2002.....	27	-	-	-	-	-	-	-	1	2	3	2
2001.....	23	-	-	-	-	-	-	1	2	3	1	2
2000.....	26	-	-	-	-	-	-	-	-	1	3	4
1999.....	17	-	-	-	-	-	-	-	1	1	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic liver disease and cirrhosis (K73-K74)

All races, both sexes

2003.....	15,143	2	-	-	-	2	8	29	86	270	779	1,407
2002.....	15,136	6	2	1	5	7	16	34	98	322	861	1,406
2001.....	14,828	3	-	1	1	5	13	32	91	363	828	1,436
2000.....	14,443	3	3	-	3	7	21	32	104	342	803	1,374
1999.....	14,301	5	2	1	1	9	9	33	115	340	776	1,263

All races, male

2003.....	8,808	2	-	-	-	1	4	15	55	172	544	986
2002.....	8,587	6	-	1	2	3	5	14	60	213	561	992
2001.....	8,461	1	-	1	1	2	5	22	58	221	539	1,018
2000.....	8,229	3	1	-	2	3	11	18	60	230	551	1,005
1999.....	8,193	4	-	1	-	3	2	17	75	225	569	945

All races, female

2003.....	6,335	-	-	-	-	1	4	14	31	98	235	421
2002.....	6,549	-	2	-	3	4	11	20	38	109	300	414
2001.....	6,367	2	-	-	-	3	8	10	33	142	289	418
2000.....	6,214	-	2	-	1	4	10	14	44	112	252	369
1999.....	6,108	1	2	-	1	6	7	16	40	115	207	318

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2003.....	13,380	1	-	-	-	2	3	16	72	230	652	1,167
2002.....	13,386	5	2	1	4	7	14	23	75	265	731	1,142
2001.....	13,021	3	-	-	1	2	6	19	77	306	668	1,146
2000.....	12,710	-	3	-	2	2	13	15	79	276	659	1,147
1999.....	12,579	4	-	-	1	7	3	21	97	282	630	991

White, male

2003.....	7,757	1	-	-	-	1	2	7	49	147	470	845
2002.....	7,590	5	-	1	2	3	5	9	49	178	493	832
2001.....	7,359	1	-	-	1	-	4	12	48	183	441	821
2000.....	7,232	-	1	-	1	1	6	10	46	191	466	859
1999.....	7,188	4	-	-	-	3	1	14	63	191	466	750

White, female

2003.....	5,623	-	-	-	-	1	1	9	23	83	182	322
2002.....	5,796	-	2	-	2	4	9	14	26	87	238	310
2001.....	5,662	2	-	-	-	2	2	7	29	123	227	325
2000.....	5,478	-	2	-	1	1	7	5	33	85	193	288
1999.....	5,391	-	-	-	1	4	2	7	34	91	164	241

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic liver disease and cirrhosis (K73-K74)												
Black, both sexes												
2003.....	1,399	1	-	-	-	-	3	12	10	34	104	195
2002.....	1,397	1	-	-	1	-	2	8	17	40	108	235
2001.....	1,427	-	-	1	-	2	7	11	9	43	134	251
2000.....	1,357	2	-	-	1	4	8	14	9	50	114	194
1999.....	1,367	-	2	1	-	2	6	12	14	45	129	224
Black, male												
2003.....	862	1	-	-	-	-	2	7	4	21	59	116
2002.....	825	1	-	-	-	-	-	4	9	23	57	141
2001.....	908	-	-	1	-	1	1	8	6	27	86	175
2000.....	812	2	-	-	1	1	5	7	3	31	66	125
1999.....	824	-	-	1	-	-	1	3	10	26	91	162
Black, female												
2003.....	537	-	-	-	-	-	1	5	6	13	45	79
2002.....	572	-	-	-	1	-	2	4	8	17	51	94
2001.....	519	-	-	-	-	1	6	3	3	16	48	76
2000.....	545	-	-	-	-	3	3	7	6	19	48	69
1999.....	543	-	2	-	-	2	5	9	4	19	38	62

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic liver disease and cirrhosis (K73-K74)

American Indian 1/, both sexes

2003.....	167	-	-	-	-	-	2	1	3	3	16	27
2002.....	166	-	-	-	-	-	-	2	5	13	17	18
2001.....	174	-	-	-	-	1	-	2	3	10	15	27
2000.....	195	-	-	-	-	1	-	3	11	14	24	20
1999.....	157	-	-	-	-	-	-	-	2	9	8	34

American Indian 1/, male

2003.....	76	-	-	-	-	-	-	1	2	1	10	14
2002.....	89	-	-	-	-	-	-	1	1	8	8	13
2001.....	91	-	-	-	-	1	-	2	2	7	7	14
2000.....	93	-	-	-	-	1	-	1	7	7	15	12
1999.....	75	-	-	-	-	-	-	-	1	6	5	21

American Indian 1/, female

2003.....	91	-	-	-	-	-	2	-	1	2	6	13
2002.....	77	-	-	-	-	-	-	1	4	5	9	5
2001.....	83	-	-	-	-	-	-	-	1	3	8	13
2000.....	102	-	-	-	-	-	-	2	4	7	9	8
1999.....	82	-	-	-	-	-	-	-	1	3	3	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic liver disease and cirrhosis (K73-K74)

Asian or Pacific Islander 2/, both sexes

2003.....	197	-	-	-	-	-	-	-	1	3	7	18
2002.....	187	-	-	-	-	-	-	1	1	4	5	11
2001.....	206	-	-	-	-	-	-	-	2	4	11	12
2000.....	181	1	-	-	-	-	-	-	5	2	6	13
1999.....	198	1	-	-	-	-	-	-	2	4	9	14

Asian or Pacific Islander 2/, male

2003.....	113	-	-	-	-	-	-	-	-	3	5	11
2002.....	83	-	-	-	-	-	-	-	1	4	3	6
2001.....	103	-	-	-	-	-	-	-	2	4	5	8
2000.....	92	1	-	-	-	-	-	-	4	1	4	9
1999.....	106	-	-	-	-	-	-	-	1	2	7	12

Asian or Pacific Islander 2/, female

2003.....	84	-	-	-	-	-	-	-	1	-	2	7
2002.....	104	-	-	-	-	-	-	1	-	-	2	5
2001.....	103	-	-	-	-	-	-	-	-	-	6	4
2000.....	89	-	-	-	-	-	-	-	1	1	2	4
1999.....	92	1	-	-	-	-	-	-	1	2	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races, both sexes

2003.....	2,948	-	-	-	1	1	3	9	16	14	26	46
2002.....	2,979	-	-	-	-	2	7	9	12	15	39	52
2001.....	2,965	1	-	-	1	2	5	11	8	21	23	69
2000.....	2,810	-	-	1	1	2	7	6	13	22	31	43
1999.....	2,830	2	-	-	1	4	6	5	11	25	34	55

All races, male

2003.....	1,318	-	-	-	-	1	3	3	8	7	14	20
2002.....	1,341	-	-	-	-	-	3	4	7	7	22	35
2001.....	1,341	1	-	-	-	1	2	5	4	9	16	39
2000.....	1,244	-	-	-	1	1	3	2	5	12	15	23
1999.....	1,248	1	-	-	1	2	3	-	4	14	16	28

All races, female

2003.....	1,630	-	-	-	1	-	-	6	8	7	12	26
2002.....	1,638	-	-	-	-	2	4	5	5	8	17	17
2001.....	1,624	-	-	-	1	1	3	6	4	12	7	30
2000.....	1,566	-	-	1	-	1	4	4	8	10	16	20
1999.....	1,582	1	-	-	-	2	3	5	7	11	18	27

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2003.....	2,573	-	-	-	1	1	2	8	10	11	17	33
2002.....	2,614	-	-	-	-	2	5	6	10	9	34	38
2001.....	2,567	1	-	-	1	2	3	9	6	14	19	49
2000.....	2,457	-	-	-	1	1	6	3	7	17	21	26
1999.....	2,487	-	-	-	1	3	3	4	9	17	20	43

White, male

2003.....	1,150	-	-	-	-	1	2	2	4	6	7	16
2002.....	1,179	-	-	-	-	-	3	2	6	5	19	24
2001.....	1,157	1	-	-	-	1	1	3	3	7	13	26
2000.....	1,111	-	-	-	1	1	2	2	3	10	12	16
1999.....	1,101	-	-	-	1	1	3	-	2	10	10	21

White, female

2003.....	1,423	-	-	-	1	-	-	6	6	5	10	17
2002.....	1,435	-	-	-	-	2	2	4	4	4	15	14
2001.....	1,410	-	-	-	1	1	2	6	3	7	6	23
2000.....	1,346	-	-	-	-	-	4	1	4	7	9	10
1999.....	1,386	-	-	-	-	2	-	4	7	7	10	22

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2003.....	286	-	-	-	-	-	1	1	6	3	8	13
2002.....	273	-	-	-	-	-	1	3	2	6	3	10
2001.....	310	-	-	-	-	-	2	2	2	7	4	14
2000.....	280	-	-	1	-	1	1	3	5	4	10	14
1999.....	257	2	-	-	-	1	3	1	2	7	13	8

Black, male

2003.....	127	-	-	-	-	-	1	1	4	1	7	4
2002.....	110	-	-	-	-	-	-	2	1	2	2	8
2001.....	130	-	-	-	-	-	1	2	1	2	3	9
2000.....	96	-	-	-	-	-	1	-	2	1	3	5
1999.....	102	1	-	-	-	1	-	-	2	3	6	5

Black, female

2003.....	159	-	-	-	-	-	-	-	2	2	1	9
2002.....	163	-	-	-	-	-	1	1	1	4	1	2
2001.....	180	-	-	-	-	-	1	-	1	5	1	5
2000.....	184	-	-	1	-	1	-	3	3	3	7	9
1999.....	155	1	-	-	-	-	3	1	-	4	7	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
All races, both sexes												
2003.....	42,453	181	14	7	15	29	66	103	179	270	526	835
2002.....	40,974	173	14	9	12	21	63	107	162	274	475	790
2001.....	39,480	133	16	9	9	21	71	82	143	299	469	749
2000.....	37,251	162	14	11	14	32	56	80	164	278	462	687
1999.....	35,525	166	15	13	18	24	48	97	155	268	439	608
All races, male												
2003.....	20,481	106	7	3	6	19	40	53	94	148	311	482
2002.....	19,695	109	7	5	7	10	31	44	85	148	259	454
2001.....	18,852	76	7	3	3	14	31	45	67	158	256	445
2000.....	17,811	103	7	7	10	18	25	44	85	152	277	433
1999.....	17,016	90	9	8	10	11	22	43	82	152	250	365
All races, female												
2003.....	21,972	75	7	4	9	10	26	50	85	122	215	353
2002.....	21,279	64	7	4	5	11	32	63	77	126	216	336
2001.....	20,628	57	9	6	6	7	40	37	76	141	213	304
2000.....	19,440	59	7	4	4	14	31	36	79	126	185	254
1999.....	18,509	76	6	5	8	13	26	54	73	116	189	243

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
White, both sexes												
2003.....	33,707	117	10	6	9	18	38	63	94	138	279	472
2002.....	32,615	112	10	7	8	10	32	56	96	149	268	463
2001.....	31,345	77	13	5	5	15	46	45	80	166	266	406
2000.....	29,598	87	8	8	12	20	43	48	88	144	268	376
1999.....	28,041	102	13	11	11	16	26	52	85	146	235	320
White, male												
2003.....	16,408	71	5	2	5	14	22	33	47	73	180	280
2002.....	15,850	70	4	4	5	5	17	23	53	82	147	264
2001.....	15,241	45	4	3	2	11	19	22	33	103	153	238
2000.....	14,385	58	5	6	8	11	22	28	45	80	163	237
1999.....	13,632	62	7	7	8	9	10	24	46	83	135	192
White, female												
2003.....	17,299	46	5	4	4	4	16	30	47	65	99	192
2002.....	16,765	42	6	3	3	5	15	33	43	67	121	199
2001.....	16,104	32	9	2	3	4	27	23	47	63	113	168
2000.....	15,213	29	3	2	4	9	21	20	43	64	105	139
1999.....	14,409	40	6	4	3	7	16	28	39	63	100	128

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Black, both sexes												
2003.....	7,855	60	3	1	6	11	23	37	82	125	231	332
2002.....	7,488	55	4	2	3	10	26	49	60	115	194	298
2001.....	7,274	53	3	4	3	5	22	35	60	117	192	309
2000.....	6,911	71	5	3	1	12	12	29	71	130	176	300
1999.....	6,711	58	1	2	6	8	22	40	63	118	192	270
Black, male												
2003.....	3,642	32	1	1	1	5	15	18	46	72	120	181
2002.....	3,427	35	3	1	1	4	11	20	29	60	103	174
2001.....	3,186	29	3	-	-	2	10	22	34	47	94	189
2000.....	3,074	42	2	1	1	7	3	13	37	71	104	191
1999.....	3,008	25	1	1	2	2	12	17	30	67	106	164
Black, female												
2003.....	4,213	28	2	-	5	6	8	19	36	53	111	151
2002.....	4,061	20	1	1	2	6	15	29	31	55	91	124
2001.....	4,088	24	-	4	3	3	12	13	26	70	98	120
2000.....	3,837	29	3	2	-	5	9	16	34	59	72	109
1999.....	3,703	33	-	1	4	6	10	23	33	51	86	106

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Asian or Pacific Islander 2/, both sexes

2003.....	636	3	1	-	-	-	2	3	1	4	8	13
2002.....	649	4	-	-	-	1	3	-	2	6	7	15
2001.....	625	1	-	-	-	1	1	2	2	9	5	17
2000.....	527	3	1	-	1	-	1	1	4	3	7	6
1999.....	579	6	1	-	1	-	-	5	3	4	5	7

Asian or Pacific Islander 2/, male

2003.....	304	2	1	-	-	-	2	2	-	3	5	11
2002.....	320	2	-	-	-	1	1	-	1	3	4	9
2001.....	320	-	-	-	-	1	-	1	-	5	4	9
2000.....	254	3	-	-	1	-	-	1	3	1	3	3
1999.....	297	3	1	-	-	-	-	2	2	2	3	4

Asian or Pacific Islander 2/, female

2003.....	332	1	-	-	-	-	-	1	1	1	3	2
2002.....	329	2	-	-	-	-	2	-	1	3	3	6
2001.....	305	1	-	-	-	-	1	1	2	4	1	8
2000.....	273	-	1	-	-	-	1	-	1	2	4	3
1999.....	282	3	-	-	1	-	-	3	1	2	2	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
All races, both sexes												
2003.....	148	8	2	1	-	1	3	1	1	1	3	3
2002.....	166	6	5	1	1	2	1	-	1	2	2	5
2001.....	156	8	4	1	-	-	-	2	1	4	4	1
2000.....	177	8	3	1	1	-	1	-	4	1	2	5
1999.....	182	3	2	3	2	-	2	3	2	2	3	6
All races, male												
2003.....	65	7	1	-	-	1	3	1	-	-	2	2
2002.....	70	3	2	1	1	1	-	-	1	-	-	4
2001.....	73	4	3	-	-	-	-	2	-	4	2	1
2000.....	91	3	2	1	1	-	1	-	3	-	1	2
1999.....	90	3	2	1	1	-	2	2	2	-	2	3
All races, female												
2003.....	83	1	1	1	-	-	-	-	1	1	1	1
2002.....	96	3	3	-	-	1	1	-	-	2	2	1
2001.....	83	4	1	1	-	-	-	-	1	-	2	-
2000.....	86	5	1	-	-	-	-	-	1	1	1	3
1999.....	92	-	-	2	1	-	-	1	-	2	1	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
White, both sexes												
2003.....	120	6	1	1	-	1	2	-	1	1	2	3
2002.....	130	5	4	1	1	1	-	-	-	1	2	3
2001.....	130	7	3	-	-	-	-	-	1	4	4	1
2000.....	141	6	1	1	-	-	1	-	3	-	1	2
1999.....	144	3	2	3	1	-	1	1	1	2	1	4
White, male												
2003.....	47	5	-	-	-	1	2	-	-	-	2	2
2002.....	56	2	1	1	1	1	-	-	-	-	-	3
2001.....	60	3	2	-	-	-	-	-	-	4	2	1
2000.....	72	3	1	1	-	-	1	-	2	-	1	2
1999.....	67	3	2	1	1	-	1	-	1	-	1	2
White, female												
2003.....	73	1	1	1	-	-	-	-	1	1	-	1
2002.....	74	3	3	-	-	-	-	-	-	1	2	-
2001.....	70	4	1	-	-	-	-	-	1	-	2	-
2000.....	69	3	-	-	-	-	-	-	1	-	-	-
1999.....	77	-	-	2	-	-	-	1	-	2	-	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Black, both sexes

2003.....	18	1	-	-	-	-	-	1	-	-	1	-
2002.....	29	1	1	-	-	1	1	-	1	1	-	2
2001.....	21	-	1	1	-	-	-	2	-	-	-	-
2000.....	30	2	2	-	-	-	-	-	1	1	1	3
1999.....	28	-	-	-	1	-	1	2	-	-	2	2

Black, male

2003.....	12	1	-	-	-	-	-	1	-	-	-	-
2002.....	11	1	1	-	-	-	-	-	1	-	-	1
2001.....	9	-	1	-	-	-	-	2	-	-	-	-
2000.....	13	-	1	-	-	-	-	-	1	-	-	-
1999.....	16	-	-	-	-	-	1	2	-	-	1	1

Black, female

2003.....	6	-	-	-	-	-	-	-	-	-	1	-
2002.....	18	-	-	-	-	1	1	-	-	1	-	1
2001.....	12	-	-	1	-	-	-	-	-	-	-	-
2000.....	17	2	1	-	-	-	-	-	-	1	1	3
1999.....	12	-	-	-	1	-	-	-	-	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races, both sexes

2003.....	537	3	-	-	-	3	5	7	7	8	12	15
2002.....	553	2	-	1	2	2	2	10	13	13	13	28
2001.....	511	-	2	-	1	3	5	4	8	16	11	15
2000.....	556	1	-	1	3	2	7	8	9	14	20	18
1999.....	589	-	1	1	-	3	4	10	5	15	13	16

All races, male

2003.....	269	1	-	-	-	1	4	3	2	6	8	12
2002.....	278	1	-	-	1	1	2	4	9	7	7	10
2001.....	264	-	1	-	-	3	2	-	1	7	6	9
2000.....	284	-	-	-	1	-	2	5	6	8	12	13
1999.....	304	-	-	1	-	3	3	7	3	9	9	10

All races, female

2003.....	268	2	-	-	-	2	1	4	5	2	4	3
2002.....	275	1	-	1	1	1	-	6	4	6	6	18
2001.....	247	-	1	-	1	-	3	4	7	9	5	6
2000.....	272	1	-	1	2	2	5	3	3	6	8	5
1999.....	285	-	1	-	-	-	1	3	2	6	4	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2003.....	430	3	-	-	-	1	2	5	3	8	8	12
2002.....	450	-	-	1	1	2	1	6	9	11	11	19
2001.....	417	-	2	-	-	2	5	3	6	8	7	8
2000.....	439	-	-	-	3	1	6	4	7	7	13	15
1999.....	467	-	1	1	-	2	2	8	3	8	7	14

White, male

2003.....	214	1	-	-	-	-	1	2	1	6	6	9
2002.....	233	-	-	-	-	1	1	3	7	6	5	8
2001.....	216	-	1	-	-	2	2	-	1	5	3	3
2000.....	226	-	-	-	1	-	1	3	4	5	7	11
1999.....	244	-	-	1	-	2	1	6	2	4	6	9

White, female

2003.....	216	2	-	-	-	1	1	3	2	2	2	3
2002.....	217	-	-	1	1	1	-	3	2	5	6	11
2001.....	201	-	1	-	-	-	3	3	5	3	4	5
2000.....	213	-	-	-	2	1	5	1	3	2	6	4
1999.....	223	-	1	-	-	-	1	2	1	4	1	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Black, both sexes

2003.....	76	-	-	-	-	2	2	1	3	-	4	3
2002.....	82	2	-	-	-	-	1	4	3	1	2	9
2001.....	69	-	-	-	1	-	-	1	1	4	4	6
2000.....	90	1	-	1	-	1	1	3	1	7	3	3
1999.....	94	-	-	-	-	1	2	-	2	6	5	1

Black, male

2003.....	39	-	-	-	-	1	2	-	1	-	2	3
2002.....	37	1	-	-	-	-	1	1	2	-	2	2
2001.....	37	-	-	-	-	-	-	-	-	2	3	5
2000.....	44	-	-	-	-	-	1	1	1	3	2	2
1999.....	47	-	-	-	-	1	2	-	1	4	2	1

Black, female

2003.....	37	-	-	-	-	1	-	1	2	-	2	-
2002.....	45	1	-	-	-	-	-	3	1	1	-	7
2001.....	32	-	-	-	1	-	-	1	1	2	1	1
2000.....	46	1	-	1	-	1	-	2	-	4	1	1
1999.....	47	-	-	-	-	-	-	-	1	2	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Asian or Pacific Islander 2/, both sexes

2003.....	25	-	-	-	-	-	1	1	-	-	-	-
2002.....	15	-	-	-	-	-	-	-	-	-	-	-
2001.....	18	-	-	-	-	1	-	-	1	3	-	-
2000.....	19	-	-	-	-	-	-	-	1	-	2	-
1999.....	24	-	-	-	-	-	-	2	-	1	-	1

Asian or Pacific Islander 2/, male

2003.....	11	-	-	-	-	-	1	1	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	-
2001.....	7	-	-	-	-	1	-	-	-	-	-	-
2000.....	8	-	-	-	-	-	-	-	1	-	1	-
1999.....	10	-	-	-	-	-	-	1	-	1	-	-

Asian or Pacific Islander 2/, female

2003.....	14	-	-	-	-	-	-	-	-	-	-	-
2002.....	11	-	-	-	-	-	-	-	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	1	3	-	-
2000.....	11	-	-	-	-	-	-	-	-	-	1	-
1999.....	14	-	-	-	-	-	-	1	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
All races, both sexes												
2003.....	41,737	170	11	6	15	25	57	95	171	261	510	813
2002.....	40,222	165	9	6	9	17	60	95	148	258	460	757
2001.....	38,784	125	9	8	8	18	64	76	133	278	454	729
2000.....	36,471	152	11	9	10	30	48	72	149	262	439	658
1999.....	34,719	163	12	9	16	21	42	83	147	250	423	586
All races, male												
2003.....	20,135	98	5	3	6	17	32	49	92	142	301	465
2002.....	19,341	105	5	3	5	8	29	39	75	141	252	440
2001.....	18,509	72	3	3	3	11	29	43	66	146	248	434
2000.....	17,416	100	5	6	8	18	22	39	76	144	263	415
1999.....	16,611	87	7	6	9	8	17	34	76	143	239	352
All races, female												
2003.....	21,602	72	6	3	9	8	25	46	79	119	209	348
2002.....	20,881	60	4	3	4	9	31	56	73	117	208	317
2001.....	20,275	53	6	5	5	7	35	33	67	132	206	295
2000.....	19,055	52	6	3	2	12	26	33	73	118	176	243
1999.....	18,108	76	5	3	7	13	25	49	71	107	184	234

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
White, both sexes												
2003.....	33,133	108	8	5	9	16	34	58	90	129	268	454
2002.....	32,008	107	6	5	6	7	31	48	87	136	255	441
2001.....	30,777	70	7	5	5	13	40	42	72	153	255	394
2000.....	28,982	80	7	7	9	19	36	44	77	137	253	356
1999.....	27,407	99	10	7	10	14	23	42	80	135	227	302
White, male												
2003.....	16,139	65	4	2	5	13	19	31	46	67	172	267
2002.....	15,558	68	3	3	4	3	16	19	46	76	142	253
2001.....	14,961	42	1	3	2	9	17	22	32	93	148	233
2000.....	14,070	55	4	5	7	11	20	25	39	75	154	223
1999.....	13,312	59	5	5	7	7	8	18	42	79	128	181
White, female												
2003.....	16,994	43	4	3	4	3	15	27	44	62	96	187
2002.....	16,450	39	3	2	2	4	15	29	41	60	113	188
2001.....	15,816	28	6	2	3	4	23	20	40	60	107	161
2000.....	14,912	25	3	2	2	8	16	19	38	62	99	133
1999.....	14,095	40	5	2	3	7	15	24	38	56	99	121

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
Black, both sexes												
2003.....	7,755	59	3	1	6	9	20	35	79	125	226	328
2002.....	7,373	52	3	1	3	9	24	45	56	113	192	287
2001.....	7,178	53	2	3	2	5	21	32	59	113	188	303
2000.....	6,782	68	3	2	1	11	11	26	68	121	172	291
1999.....	6,577	58	1	2	5	7	19	38	61	112	185	267
Black, male												
2003.....	3,587	31	1	1	1	4	12	17	45	72	118	177
2002.....	3,377	33	2	-	1	4	10	19	26	60	101	171
2001.....	3,138	29	2	-	-	2	10	20	34	45	91	184
2000.....	3,015	42	1	1	1	7	2	12	35	68	102	187
1999.....	2,943	25	1	1	2	1	9	15	29	63	103	162
Black, female												
2003.....	4,168	28	2	-	5	5	8	18	34	53	108	151
2002.....	3,996	19	1	1	2	5	14	26	30	53	91	116
2001.....	4,040	24	-	3	2	3	11	12	25	68	97	119
2000.....	3,767	26	2	1	-	4	9	14	33	53	70	104
1999.....	3,634	33	-	1	3	6	10	23	32	49	82	105

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Renal failure (N17-N19)

Asian or Pacific Islander 2/, both sexes

2003.....	601	2	-	-	-	-	-	2	1	4	8	13
2002.....	627	4	-	-	-	1	3	-	2	6	7	15
2001.....	603	1	-	-	-	-	1	2	1	6	5	17
2000.....	502	3	1	-	-	-	1	1	3	3	5	6
1999.....	548	6	1	-	1	-	-	3	3	3	5	6

Asian or Pacific Islander 2/, male

2003.....	287	1	-	-	-	-	-	1	-	3	5	11
2002.....	313	2	-	-	-	1	1	-	1	3	4	9
2001.....	311	-	-	-	-	-	-	1	-	5	4	9
2000.....	241	3	-	-	-	-	-	1	2	1	2	3
1999.....	282	3	1	-	-	-	-	1	2	1	3	4

Asian or Pacific Islander 2/, female

2003.....	314	1	-	-	-	-	-	1	1	1	3	2
2002.....	314	2	-	-	-	-	2	-	1	3	3	6
2001.....	292	1	-	-	-	-	1	1	1	1	1	8
2000.....	261	-	1	-	-	-	1	-	1	2	3	3
1999.....	266	3	-	-	1	-	-	2	1	2	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of kidney (N25,N27)

All races, both sexes

2003.....	31	-	1	-	-	-	1	-	-	-	1	4
2002.....	33	-	-	1	-	-	-	2	-	1	-	-
2001.....	29	-	1	-	-	-	2	-	1	1	-	4
2000.....	47	1	-	-	-	-	-	-	2	1	1	6
1999.....	35	-	-	-	-	-	-	1	1	1	-	-

All races, male

2003.....	12	-	1	-	-	-	1	-	-	-	-	3
2002.....	6	-	-	1	-	-	-	1	-	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	1	-	1
2000.....	20	-	-	-	-	-	-	-	-	-	1	3
1999.....	11	-	-	-	-	-	-	-	1	-	-	-

All races, female

2003.....	19	-	-	-	-	-	-	-	-	-	1	1
2002.....	27	-	-	-	-	-	-	1	-	1	-	-
2001.....	23	-	1	-	-	-	2	-	1	-	-	3
2000.....	27	1	-	-	-	-	-	-	2	1	-	3
1999.....	24	-	-	-	-	-	-	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of kidney (N25,N27)

White, both sexes

2003.....	24	-	1	-	-	-	-	-	-	-	1	3
2002.....	27	-	-	-	-	-	-	2	-	1	-	-
2001.....	21	-	1	-	-	-	1	-	1	1	-	3
2000.....	36	1	-	-	-	-	-	-	1	-	1	3
1999.....	23	-	-	-	-	-	-	1	1	1	-	-

White, male

2003.....	8	-	1	-	-	-	-	-	-	-	-	2
2002.....	3	-	-	-	-	-	-	1	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	1	-	1
2000.....	17	-	-	-	-	-	-	-	-	-	1	1
1999.....	9	-	-	-	-	-	-	-	1	-	-	-

White, female

2003.....	16	-	-	-	-	-	-	-	-	-	1	1
2002.....	24	-	-	-	-	-	-	1	-	1	-	-
2001.....	17	-	1	-	-	-	1	-	1	-	-	2
2000.....	19	1	-	-	-	-	-	-	1	-	-	2
1999.....	14	-	-	-	-	-	-	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races, both sexes

2003.....	823	4	2	-	1	2	8	7	9	10	26	32
2002.....	788	2	-	-	3	2	3	5	8	11	17	19
2001.....	748	2	4	-	-	1	4	5	6	16	17	21
2000.....	813	5	3	-	-	1	1	4	5	9	12	21
1999.....	846	3	1	1	-	3	1	5	4	13	15	22

All races, male

2003.....	246	3	-	-	1	-	2	-	1	3	7	14
2002.....	251	2	-	-	1	-	1	3	3	3	9	5
2001.....	213	1	2	-	-	-	2	-	3	4	6	6
2000.....	214	3	-	-	-	1	-	2	1	2	6	6
1999.....	263	1	-	-	-	1	-	-	-	2	5	8

All races, female

2003.....	577	1	2	-	-	2	6	7	8	7	19	18
2002.....	537	-	-	-	2	2	2	2	5	8	8	14
2001.....	535	1	2	-	-	1	2	5	3	12	11	15
2000.....	599	2	3	-	-	-	1	2	4	7	6	15
1999.....	583	2	1	1	-	2	1	5	4	11	10	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2003.....	705	1	1	-	1	2	6	3	8	6	22	23
2002.....	692	2	-	-	1	-	1	4	6	7	15	10
2001.....	647	1	2	-	-	-	2	4	4	11	15	15
2000.....	689	5	3	-	-	-	1	4	3	5	7	17
1999.....	749	1	1	1	-	1	1	4	4	8	12	19

White, male

2003.....	215	-	-	-	1	-	2	-	1	2	7	12
2002.....	219	2	-	-	-	-	1	3	3	3	7	3
2001.....	176	-	1	-	-	-	1	-	2	3	6	5
2000.....	182	3	-	-	-	-	-	2	1	-	4	6
1999.....	232	1	-	-	-	1	-	-	-	2	3	6

White, female

2003.....	490	1	1	-	-	2	4	3	7	4	15	11
2002.....	473	-	-	-	1	-	-	1	3	4	8	7
2001.....	471	1	1	-	-	-	1	4	2	8	9	10
2000.....	507	2	3	-	-	-	1	2	2	5	3	11
1999.....	517	-	1	1	-	-	1	4	4	6	9	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Infections of kidney (N10-N12,N13.6,N15.1)

Black, both sexes

2003.....	95	3	1	-	-	-	2	2	-	3	4	7
2002.....	73	-	-	-	-	1	2	-	2	4	2	6
2001.....	80	1	-	-	-	-	1	1	2	5	2	4
2000.....	101	-	-	-	-	1	-	-	2	3	4	3
1999.....	85	1	-	-	-	2	-	1	-	3	3	3

Black, male

2003.....	29	3	-	-	-	-	-	-	-	1	-	2
2002.....	22	-	-	-	-	-	-	-	-	-	2	1
2001.....	28	1	-	-	-	-	1	-	1	1	-	1
2000.....	27	-	-	-	-	1	-	-	-	1	1	-
1999.....	29	-	-	-	-	-	-	-	-	-	2	2

Black, female

2003.....	66	-	1	-	-	-	2	2	-	2	4	5
2002.....	51	-	-	-	-	1	2	-	2	4	-	5
2001.....	52	-	-	-	-	-	-	1	1	4	2	3
2000.....	74	-	-	-	-	-	-	-	2	2	3	3
1999.....	56	1	-	-	-	2	-	1	-	3	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Infections of kidney (N10-N12,N13.6,N15.1)

American Indian 1/, both sexes

2003.....	6	-	-	-	-	-	-	1	-	1	-	1
2002.....	8	-	-	-	1	-	-	-	-	-	-	2
2001.....	6	-	1	-	-	-	1	-	-	-	-	1
2000.....	5	-	-	-	-	-	-	-	-	-	-	1
1999.....	3	-	-	-	-	-	-	-	-	1	-	-

American Indian 1/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	1	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	6	-	-	-	-	-	-	1	-	1	-	1
2002.....	5	-	-	-	-	-	-	-	-	-	-	2
2001.....	4	-	1	-	-	-	1	-	-	-	-	1
2000.....	5	-	-	-	-	-	-	-	-	-	-	1
1999.....	3	-	-	-	-	-	-	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Infections of kidney (N10-N12,N13.6,N15.1)

Asian or Pacific Islander 2/, both sexes

2003.....	17	-	-	-	-	-	-	1	1	-	-	1
2002.....	15	-	-	-	1	1	-	1	-	-	-	1
2001.....	15	-	1	-	-	1	-	-	-	-	-	1
2000.....	18	-	-	-	-	-	-	-	-	1	1	-
1999.....	9	1	-	-	-	-	-	-	-	1	-	-

Asian or Pacific Islander 2/, male

2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	7	-	-	-	-	-	-	-	-	-	-	1
2001.....	7	-	1	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	1	1	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	15	-	-	-	-	-	-	1	1	-	-	1
2002.....	8	-	-	-	1	1	-	1	-	-	-	-
2001.....	8	-	-	-	-	1	-	-	-	-	-	1
2000.....	13	-	-	-	-	-	-	-	-	-	-	-
1999.....	7	1	-	-	-	-	-	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Inflammatory diseases of female pelvic organs (N70-N76)

All races, both sexes

2003.....	125	1	-	-	1	1	5	1	2	7	7	7
2002.....	114	1	-	-	-	1	1	-	4	12	5	4
2001.....	127	1	-	-	-	-	2	2	1	2	8	9
2000.....	120	2	-	-	1	-	2	2	4	9	6	6
1999.....	108	2	1	-	-	1	2	3	3	4	8	4

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	125	1	-	-	1	1	5	1	2	7	7	7
2002.....	114	1	-	-	-	1	1	-	4	12	5	4
2001.....	127	1	-	-	-	-	2	2	1	2	8	9
2000.....	120	2	-	-	1	-	2	2	4	9	6	6
1999.....	108	2	1	-	-	1	2	3	3	4	8	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Inflammatory diseases of female pelvic organs (N70-N76)

White, both sexes

2003.....	101	-	-	-	1	1	2	1	1	6	4	5
2002.....	92	1	-	-	-	-	-	-	3	4	5	2
2001.....	103	1	-	-	-	-	2	2	1	-	6	6
2000.....	94	1	-	-	-	-	2	2	2	5	2	4
1999.....	80	1	1	-	-	1	1	1	2	1	3	2

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	101	-	-	-	1	1	2	1	1	6	4	5
2002.....	92	1	-	-	-	-	-	-	3	4	5	2
2001.....	103	1	-	-	-	-	2	2	1	-	6	6
2000.....	94	1	-	-	-	-	2	2	2	5	2	4
1999.....	80	1	1	-	-	1	1	1	2	1	3	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Inflammatory diseases of female pelvic organs (N70-N76)

Black, both sexes

2003.....	20	1	-	-	-	-	2	-	1	1	3	2
2002.....	20	-	-	-	-	1	1	-	1	8	-	1
2001.....	23	-	-	-	-	-	-	-	-	2	2	3
2000.....	23	1	-	-	1	-	-	-	1	3	4	2
1999.....	26	1	-	-	-	-	1	2	-	2	5	2

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	20	1	-	-	-	-	2	-	1	1	3	2
2002.....	20	-	-	-	-	1	1	-	1	8	-	1
2001.....	23	-	-	-	-	-	-	-	-	2	2	3
2000.....	23	1	-	-	1	-	-	-	1	3	4	2
1999.....	26	1	-	-	-	-	1	2	-	2	5	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Inflammatory diseases of female pelvic organs (N70-N76)

Asian or Pacific Islander 2/, both sexes

2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

All races, both sexes

2003.....	545	1	27	89	110	119	106	60	19
2002.....	379	1	28	61	82	93	79	26	8
2001.....	416	3	37	73	93	97	73	33	5
2000.....	404	-	18	77	87	95	86	35	5
1999.....	406	-	33	64	89	92	88	33	5

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	545	1	27	89	110	119	106	60	19
2002.....	379	1	28	61	82	93	79	26	8
2001.....	416	3	37	73	93	97	73	33	5
2000.....	404	-	18	77	87	95	86	35	5
1999.....	406	-	33	64	89	92	88	33	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

White, both sexes

2003.....	317	-	15	51	73	59	56	38	14
2002.....	203	-	12	28	41	56	41	19	5
2001.....	235	2	22	42	50	64	33	18	2
2000.....	243	-	11	44	53	55	56	21	2
1999.....	220	-	14	31	47	53	55	16	4

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	317	-	15	51	73	59	56	38	14
2002.....	203	-	12	28	41	56	41	19	5
2001.....	235	2	22	42	50	64	33	18	2
2000.....	243	-	11	44	53	55	56	21	2
1999.....	220	-	14	31	47	53	55	16	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

Black, both sexes

2003.....	193	1	12	32	32	51	40	18	5
2002.....	157	1	16	30	38	28	35	7	2
2001.....	158	1	14	31	36	30	30	13	3
2000.....	142	-	4	32	31	34	26	12	3
1999.....	162	-	18	29	38	33	28	14	-

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	193	1	12	32	32	51	40	18	5
2002.....	157	1	16	30	38	28	35	7	2
2001.....	158	1	14	31	36	30	30	13	3
2000.....	142	-	4	32	31	34	26	12	3
1999.....	162	-	18	29	38	33	28	14	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

American Indian 1/, both sexes

2003.....	8	-	-	1	1	1	4	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	3	2	-	-	-
2000.....	6	-	1	1	1	3	-	-	-
1999.....	5	-	1	2	-	1	1	-	-

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	8	-	-	1	1	1	4	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	3	2	-	-	-
2000.....	6	-	1	1	1	3	-	-	-
1999.....	5	-	1	2	-	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

Asian or Pacific Islander 2/, both sexes

2003.....	27	-	-	5	4	8	6	3	-
2002.....	19	-	-	3	3	9	3	-	1
2001.....	18	-	1	-	4	1	10	2	-
2000.....	13	-	2	-	2	3	4	2	-
1999.....	19	-	-	2	4	5	4	3	1

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	27	-	-	5	4	8	6	3	-
2002.....	19	-	-	3	3	9	3	-	1
2001.....	18	-	1	-	4	1	10	2	-
2000.....	13	-	2	-	2	3	4	2	-
1999.....	19	-	-	2	4	5	4	3	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

All races, both sexes

2003.....	40	-	6	5	8	11	6	3	1
2002.....	22	1	-	4	4	5	6	1	1
2001.....	38	2	6	7	6	8	8	1	-
2000.....	37	-	-	6	12	5	10	4	-
1999.....	31	-	3	5	5	10	6	2	-

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	40	-	6	5	8	11	6	3	1
2002.....	22	1	-	4	4	5	6	1	1
2001.....	38	2	6	7	6	8	8	1	-
2000.....	37	-	-	6	12	5	10	4	-
1999.....	31	-	3	5	5	10	6	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

White, both sexes

2003.....	18	-	1	2	7	4	3	1	-
2002.....	7	-	-	1	2	3	-	1	-
2001.....	20	1	3	4	4	4	3	1	-
2000.....	18	-	-	1	7	1	6	3	-
1999.....	15	-	2	2	1	5	3	2	-

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	18	-	1	2	7	4	3	1	-
2002.....	7	-	-	1	2	3	-	1	-
2001.....	20	1	3	4	4	4	3	1	-
2000.....	18	-	-	1	7	1	6	3	-
1999.....	15	-	2	2	1	5	3	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Black, both sexes

2003.....	18	-	5	3	-	5	2	2	1
2002.....	12	1	-	2	2	2	5	-	-
2001.....	16	1	3	3	2	3	4	-	-
2000.....	19	-	-	5	5	4	4	1	-
1999.....	15	-	1	3	4	4	3	-	-

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	18	-	5	3	-	5	2	2	1
2002.....	12	1	-	2	2	2	5	-	-
2001.....	16	1	3	3	2	3	4	-	-
2000.....	19	-	-	5	5	4	4	1	-
1999.....	15	-	1	3	4	4	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

American Indian 1/, both sexes

2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races, both sexes

2003.....	505	1	21	84	102	108	100	57	18
2002.....	357	-	28	57	78	88	73	25	7
2001.....	378	1	31	66	87	89	65	32	5
2000.....	367	-	18	71	75	90	76	31	5
1999.....	375	-	30	59	84	82	82	31	5

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	505	1	21	84	102	108	100	57	18
2002.....	357	-	28	57	78	88	73	25	7
2001.....	378	1	31	66	87	89	65	32	5
2000.....	367	-	18	71	75	90	76	31	5
1999.....	375	-	30	59	84	82	82	31	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

White, both sexes

2003.....	299	-	14	49	66	55	53	37	14
2002.....	196	-	12	27	39	53	41	18	5
2001.....	215	1	19	38	46	60	30	17	2
2000.....	225	-	11	43	46	54	50	18	2
1999.....	205	-	12	29	46	48	52	14	4

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	299	-	14	49	66	55	53	37	14
2002.....	196	-	12	27	39	53	41	18	5
2001.....	215	1	19	38	46	60	30	17	2
2000.....	225	-	11	43	46	54	50	18	2
1999.....	205	-	12	29	46	48	52	14	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Black, both sexes

2003.....	175	1	7	29	32	46	38	16	4
2002.....	145	-	16	28	36	26	30	7	2
2001.....	142	-	11	28	34	27	26	13	3
2000.....	123	-	4	27	26	30	22	11	3
1999.....	147	-	17	26	34	29	25	14	-

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	175	1	7	29	32	46	38	16	4
2002.....	145	-	16	28	36	26	30	7	2
2001.....	142	-	11	28	34	27	26	13	3
2000.....	123	-	4	27	26	30	22	11	3
1999.....	147	-	17	26	34	29	25	14	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

American Indian 1/, both sexes

2003.....	7	-	-	1	1	1	3	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	3	1	-	-	-
2000.....	6	-	1	1	1	3	-	-	-
1999.....	4	-	1	2	-	-	1	-	-

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	7	-	-	1	1	1	3	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	3	1	-	-	-
2000.....	6	-	1	1	1	3	-	-	-
1999.....	4	-	1	2	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Asian or Pacific Islander 2/, both sexes

2003.....	24	-	-	5	3	6	6	3	-
2002.....	16	-	-	2	3	9	2	-	-
2001.....	17	-	1	-	4	1	9	2	-
2000.....	13	-	2	-	2	3	4	2	-
1999.....	19	-	-	2	4	5	4	3	1

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	24	-	-	5	3	6	6	3	-
2002.....	16	-	-	2	3	9	2	-	-
2001.....	17	-	1	-	4	1	9	2	-
2000.....	13	-	2	-	2	3	4	2	-
1999.....	19	-	-	2	4	5	4	3	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

All races, both sexes

2003.....	14,378	14,254	79	10	9	12	4	2	1	1	2	1
2002.....	14,254	14,106	65	15	16	14	7	6	4	7	3	2
2001.....	13,887	13,734	72	22	14	8	9	5	1	6	3	3
2000.....	14,069	13,918	79	25	16	10	5	5	2	2	2	1
1999.....	14,259	14,084	92	25	18	11	5	2	5	2	1	2

All races, male

2003.....	8,197	8,129	43	5	6	6	2	1	1	1	-	-
2002.....	8,047	7,960	40	10	9	9	3	3	1	4	1	2
2001.....	7,868	7,775	44	16	8	3	4	2	-	4	2	3
2000.....	7,968	7,872	52	13	10	9	2	4	2	-	2	1
1999.....	8,044	7,941	56	14	9	6	4	-	2	2	-	2

All races, female

2003.....	6,181	6,125	36	5	3	6	2	1	-	-	2	1
2002.....	6,207	6,146	25	5	7	5	4	3	3	3	2	-
2001.....	6,019	5,959	28	6	6	5	5	3	1	2	1	-
2000.....	6,101	6,046	27	12	6	1	3	1	-	2	-	-
1999.....	6,215	6,143	36	11	9	5	1	2	3	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

White, both sexes

2003.....	8,975	8,891	45	9	7	12	3	2	1	1	2	1
2002.....	8,777	8,679	36	9	12	13	4	5	2	5	3	2
2001.....	8,551	8,444	47	14	7	5	8	5	1	6	2	3
2000.....	8,532	8,431	46	18	12	8	5	2	1	2	2	1
1999.....	8,648	8,522	56	19	14	11	4	2	5	2	1	2

White, male

2003.....	5,134	5,087	25	5	5	6	2	1	1	1	-	-
2002.....	5,011	4,959	19	7	5	8	1	3	1	2	1	2
2001.....	4,846	4,780	28	11	5	2	4	2	-	4	1	3
2000.....	4,828	4,762	30	10	10	8	2	1	1	-	2	1
1999.....	4,884	4,815	30	10	7	6	4	-	2	2	-	2

White, female

2003.....	3,841	3,804	20	4	2	6	1	1	-	-	2	1
2002.....	3,766	3,720	17	2	7	5	3	2	1	3	2	-
2001.....	3,705	3,664	19	3	2	3	4	3	1	2	1	-
2000.....	3,704	3,669	16	8	2	-	3	1	-	2	-	-
1999.....	3,764	3,707	26	9	7	5	-	2	3	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

Black, both sexes

2003.....	4,879	4,844	30	-	2	-	1	-	-	-	-	-
2002.....	4,931	4,889	26	4	4	1	2	1	1	1	-	-
2001.....	4,847	4,808	23	6	5	3	1	-	-	-	1	-
2000.....	5,021	4,979	27	6	4	1	-	3	1	-	-	-
1999.....	5,150	5,113	29	2	3	-	1	-	-	-	-	-

Black, male

2003.....	2,775	2,756	16	-	1	-	-	-	-	-	-	-
2002.....	2,725	2,696	18	2	4	1	1	-	-	1	-	-
2001.....	2,746	2,725	14	3	2	1	-	-	-	-	1	-
2000.....	2,844	2,818	18	3	-	1	-	3	1	-	-	-
1999.....	2,907	2,883	19	2	1	-	-	-	-	-	-	-

Black, female

2003.....	2,104	2,088	14	-	1	-	1	-	-	-	-	-
2002.....	2,206	2,193	8	2	-	-	1	1	1	-	-	-
2001.....	2,101	2,083	9	3	3	2	1	-	-	-	-	-
2000.....	2,177	2,161	9	3	4	-	-	-	-	-	-	-
1999.....	2,243	2,230	10	-	2	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, both sexes

2003.....	10,518	5,621	541	180	206	223	228	198	228	242	322	324
2002.....	10,687	5,623	530	199	218	248	244	213	262	264	308	346
2001.....	10,442	5,513	557	182	194	255	250	225	233	259	327	303
2000.....	10,578	5,743	495	198	201	225	216	208	269	277	292	287
1999.....	10,393	5,473	549	207	221	222	212	212	253	279	273	342

All races, male

2003.....	5,460	3,020	275	94	104	144	128	119	114	129	175	177
2002.....	5,515	2,942	282	99	116	146	136	132	152	132	164	195
2001.....	5,448	2,954	282	101	109	161	145	121	128	137	180	156
2000.....	5,517	3,028	259	105	102	137	131	120	159	154	162	157
1999.....	5,346	2,875	310	102	113	139	123	110	132	152	144	185

All races, female

2003.....	5,058	2,601	266	86	102	79	100	79	114	113	147	147
2002.....	5,172	2,681	248	100	102	102	108	81	110	132	144	151
2001.....	4,994	2,559	275	81	85	94	105	104	105	122	147	147
2000.....	5,061	2,715	236	93	99	88	85	88	110	123	130	130
1999.....	5,047	2,598	239	105	108	83	89	102	121	127	129	157

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2003.....	8,378	4,292	392	131	157	171	171	159	184	199	278	264
2002.....	8,556	4,330	395	155	166	189	194	169	215	213	250	291
2001.....	8,338	4,247	387	137	145	189	185	178	197	206	273	249
2000.....	8,396	4,371	347	147	145	159	173	163	228	223	236	245
1999.....	8,297	4,211	385	144	157	170	162	178	208	232	229	292

White, male

2003.....	4,288	2,282	198	67	79	110	96	95	94	108	152	138
2002.....	4,416	2,297	211	75	84	111	107	104	127	107	135	162
2001.....	4,376	2,301	195	79	90	117	105	104	108	113	148	131
2000.....	4,368	2,316	180	72	75	93	108	93	136	127	132	132
1999.....	4,269	2,240	218	69	79	109	95	93	109	128	117	156

White, female

2003.....	4,090	2,010	194	64	78	61	75	64	90	91	126	126
2002.....	4,140	2,033	184	80	82	78	87	65	88	106	115	129
2001.....	3,962	1,946	192	58	55	72	80	74	89	93	125	118
2000.....	4,028	2,055	167	75	70	66	65	70	92	96	104	113
1999.....	4,028	1,971	167	75	78	61	67	85	99	104	112	136

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2003.....	1,700	1,039	120	36	38	40	52	33	43	34	36	51
2002.....	1,758	1,047	114	40	43	54	46	35	40	40	53	47
2001.....	1,720	1,032	131	34	41	61	53	36	32	45	46	44
2000.....	1,790	1,102	126	44	47	56	35	38	39	48	45	39
1999.....	1,720	1,025	138	50	56	43	40	28	33	42	35	43

Black, male

2003.....	923	574	62	20	19	27	28	19	19	15	18	32
2002.....	909	522	59	23	25	34	26	22	21	18	28	27
2001.....	882	537	69	17	18	41	31	12	19	19	28	18
2000.....	944	567	70	29	23	39	19	24	22	25	26	22
1999.....	899	531	79	25	29	27	24	14	18	19	23	27

Black, female

2003.....	777	465	58	16	19	13	24	14	24	19	18	19
2002.....	849	525	55	17	18	20	20	13	19	22	25	20
2001.....	838	495	62	17	23	20	22	24	13	26	18	26
2000.....	846	535	56	15	24	17	16	14	17	23	19	17
1999.....	821	494	59	25	27	16	16	14	15	23	12	16

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

American Indian 1/, both sexes

2003.....	118	72	10	5	6	2	4	1	-	3	3	2
2002.....	93	61	5	3	3	1	-	4	2	1	2	2
2001.....	98	60	15	1	2	-	5	5	-	-	2	3
2000.....	95	68	4	3	-	1	2	-	-	2	2	-
1999.....	106	67	5	3	3	3	4	2	5	-	1	3

American Indian 1/, male

2003.....	58	35	4	2	3	1	3	1	-	2	2	2
2002.....	42	29	2	1	2	-	-	2	1	-	1	1
2001.....	46	30	5	-	1	-	5	1	-	-	1	2
2000.....	56	42	1	2	-	-	2	-	-	-	1	-
1999.....	50	28	3	2	2	1	2	-	3	-	-	2

American Indian 1/, female

2003.....	60	37	6	3	3	1	1	-	-	1	1	-
2002.....	51	32	3	2	1	1	-	2	1	1	1	1
2001.....	52	30	10	1	1	-	-	4	-	-	1	1
2000.....	39	26	3	1	-	1	-	-	-	2	1	-
1999.....	56	39	2	1	1	2	2	2	2	-	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Asian or Pacific Islander 2/, both sexes

2003.....	322	218	19	8	5	10	1	5	1	6	5	7
2002.....	280	185	16	1	6	4	4	5	5	10	3	6
2001.....	286	174	24	10	6	5	7	6	4	8	6	7
2000.....	297	202	18	4	9	9	6	7	2	4	9	3
1999.....	270	170	21	10	5	6	6	4	7	5	8	4

Asian or Pacific Islander 2/, male

2003.....	191	129	11	5	3	6	1	4	1	4	3	5
2002.....	148	94	10	-	5	1	3	4	3	7	-	5
2001.....	144	86	13	5	-	3	4	4	1	5	3	5
2000.....	149	103	8	2	4	5	2	3	1	2	3	3
1999.....	128	76	10	6	3	2	2	3	2	5	4	-

Asian or Pacific Islander 2/, female

2003.....	131	89	8	3	2	4	-	1	-	2	2	2
2002.....	132	91	6	1	1	3	1	1	2	3	3	1
2001.....	142	88	11	5	6	2	3	2	3	3	3	2
2000.....	148	99	10	2	5	4	4	4	1	2	6	-
1999.....	142	94	11	4	2	4	4	1	5	-	4	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All races, both sexes												
2003.....	31,444	3,318	256	75	103	300	537	617	710	1,037	1,345	1,578
2002.....	29,975	3,456	248	61	78	256	484	562	757	1,013	1,389	1,368
2001.....	32,326	3,491	272	79	102	361	588	663	867	1,379	1,747	1,882
2000.....	31,876	3,639	248	85	92	317	551	663	891	1,284	1,703	1,727
1999.....	26,672	3,538	236	65	72	228	381	446	649	940	1,122	1,168
All races, male												
2003.....	14,583	1,974	143	30	62	197	366	434	473	685	841	1,011
2002.....	14,269	2,032	145	37	45	172	329	391	514	670	870	895
2001.....	15,779	2,041	162	44	55	249	405	461	579	888	1,119	1,220
2000.....	15,863	2,156	137	48	51	196	385	456	604	857	1,137	1,157
1999.....	12,877	2,072	136	35	42	153	269	287	434	599	740	796
All races, female												
2003.....	16,861	1,344	113	45	41	103	171	183	237	352	504	567
2002.....	15,706	1,424	103	24	33	84	155	171	243	343	519	473
2001.....	16,547	1,450	110	35	47	112	183	202	288	491	628	662
2000.....	16,013	1,483	111	37	41	121	166	207	287	427	566	570
1999.....	13,795	1,466	100	30	30	75	112	159	215	341	382	372

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2003.....	26,147	2,125	166	57	61	244	415	464	550	806	1,043	1,188
2002.....	24,749	2,224	161	35	49	192	369	422	580	775	1,079	1,024
2001.....	26,877	2,231	185	53	71	288	459	519	683	1,089	1,423	1,517
2000.....	26,285	2,294	182	57	63	240	416	495	690	1,035	1,360	1,364
1999.....	21,591	2,227	159	49	47	163	270	312	486	699	808	849

White, male

2003.....	11,802	1,313	91	26	35	163	290	328	378	534	655	772
2002.....	11,421	1,323	89	22	30	130	261	303	397	517	681	669
2001.....	12,868	1,320	106	25	41	198	318	379	461	702	933	994
2000.....	12,827	1,363	106	35	33	157	301	349	481	714	935	929
1999.....	10,094	1,321	96	26	28	110	199	211	318	463	542	572

White, female

2003.....	14,345	812	75	31	26	81	125	136	172	272	388	416
2002.....	13,328	901	72	13	19	62	108	119	183	258	398	355
2001.....	14,009	911	79	28	30	90	141	140	222	387	490	523
2000.....	13,458	931	76	22	30	83	115	146	209	321	425	435
1999.....	11,497	906	63	23	19	53	71	101	168	236	266	277

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2003.....	4,708	1,051	80	14	37	47	107	132	139	203	284	354
2002.....	4,638	1,095	67	24	26	58	96	121	158	218	277	308
2001.....	4,691	1,104	69	20	22	55	111	123	147	241	282	326
2000.....	5,004	1,212	57	21	25	67	119	142	174	214	314	325
1999.....	4,591	1,167	68	15	18	54	100	116	150	231	283	299

Black, male

2003.....	2,459	574	44	3	24	29	65	94	82	129	171	215
2002.....	2,536	630	45	14	14	39	54	73	104	140	169	203
2001.....	2,465	626	44	15	11	38	75	69	85	145	158	198
2000.....	2,693	710	29	11	17	32	76	91	107	115	178	200
1999.....	2,517	669	35	9	11	38	63	67	109	131	176	209

Black, female

2003.....	2,249	477	36	11	13	18	42	38	57	74	113	139
2002.....	2,102	465	22	10	12	19	42	48	54	78	108	105
2001.....	2,226	478	25	5	11	17	36	54	62	96	124	128
2000.....	2,311	502	28	10	8	35	43	51	67	99	136	125
1999.....	2,074	498	33	6	7	16	37	49	41	100	107	90

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

American Indian 1/, both sexes

2003.....	231	76	4	1	3	5	5	6	6	19	9	17
2002.....	248	72	7	1	-	3	6	8	12	9	22	17
2001.....	263	81	6	1	3	4	5	6	14	14	14	10
2000.....	209	61	3	1	-	4	5	5	12	12	9	13
1999.....	215	72	3	-	2	3	3	8	4	5	10	9

American Indian 1/, male

2003.....	135	44	3	-	3	3	5	4	5	15	6	14
2002.....	130	41	3	-	-	1	5	6	7	4	14	9
2001.....	160	52	5	1	-	3	4	3	13	12	10	7
2000.....	127	42	1	-	-	3	3	2	5	7	7	12
1999.....	117	39	1	-	1	2	3	4	1	3	6	7

American Indian 1/, female

2003.....	96	32	1	1	-	2	-	2	1	4	3	3
2002.....	118	31	4	1	-	2	1	2	5	5	8	8
2001.....	103	29	1	-	3	1	1	3	1	2	4	3
2000.....	82	19	2	1	-	1	2	3	7	5	2	1
1999.....	98	33	2	-	1	1	-	4	3	2	4	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Asian or Pacific Islander 2/, both sexes

2003.....	358	66	6	3	2	4	10	15	15	9	9	19
2002.....	340	65	13	1	3	3	13	11	7	11	11	19
2001.....	495	75	12	5	6	14	13	15	23	35	28	29
2000.....	378	72	6	6	4	6	11	21	15	23	20	25
1999.....	275	72	6	1	5	8	8	10	9	5	21	11

Asian or Pacific Islander 2/, male

2003.....	187	43	5	1	-	2	6	8	8	7	9	10
2002.....	182	38	8	1	1	2	9	9	6	9	6	14
2001.....	286	43	7	3	3	10	8	10	20	29	18	21
2000.....	216	41	1	2	1	4	5	14	11	21	17	16
1999.....	149	43	4	-	2	3	4	5	6	2	16	8

Asian or Pacific Islander 2/, female

2003.....	171	23	1	2	2	2	4	7	7	2	-	9
2002.....	158	27	5	-	2	1	4	2	1	2	5	5
2001.....	209	32	5	2	3	4	5	5	3	6	10	8
2000.....	162	31	5	4	3	2	6	7	4	2	3	9
1999.....	126	29	2	1	3	5	4	5	3	3	5	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, both sexes

2003.....	201,676	1,190	635	408	519	830	1,193	1,413	1,917	3,099	5,308	7,248
2002.....	194,591	1,144	661	386	482	717	1,136	1,256	1,880	3,099	5,407	7,159
2001.....	186,406	1,131	646	389	471	772	1,027	1,225	1,847	3,145	4,939	6,667
2000.....	178,378	1,105	609	376	503	732	1,032	1,257	1,840	3,161	4,952	6,227
1999.....	171,683	1,148	647	436	485	785	986	1,235	1,827	3,203	4,786	5,877

All races, male

2003.....	81,907	711	342	234	307	489	700	802	1,083	1,765	3,167	4,320
2002.....	79,007	690	363	220	266	423	649	714	1,123	1,800	3,219	4,305
2001.....	75,162	649	358	187	267	463	601	699	1,030	1,869	2,905	3,970
2000.....	72,227	629	338	198	276	432	625	737	1,074	1,873	2,928	3,811
1999.....	69,784	680	354	229	254	483	562	717	1,073	1,912	2,923	3,510

All races, female

2003.....	119,769	479	293	174	212	341	493	611	834	1,334	2,141	2,928
2002.....	115,584	454	298	166	216	294	487	542	757	1,299	2,188	2,854
2001.....	111,244	482	288	202	204	309	426	526	817	1,276	2,034	2,697
2000.....	106,151	476	271	178	227	300	407	520	766	1,288	2,024	2,416
1999.....	101,899	468	293	207	231	302	424	518	754	1,291	1,863	2,367

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2003.....	174,660	796	461	307	380	635	895	1,003	1,353	2,228	4,010	5,532
2002.....	168,631	788	470	295	359	551	797	890	1,343	2,252	4,063	5,444
2001.....	161,293	745	471	308	355	575	760	860	1,303	2,274	3,615	4,951
2000.....	154,181	740	422	277	375	558	767	880	1,317	2,269	3,647	4,572
1999.....	148,614	750	482	314	361	609	710	912	1,342	2,264	3,488	4,315

White, male

2003.....	69,977	502	241	187	219	392	561	609	805	1,323	2,485	3,404
2002.....	67,452	493	257	171	195	337	483	526	842	1,384	2,529	3,355
2001.....	64,020	416	272	152	200	358	466	509	755	1,406	2,190	3,025
2000.....	61,390	409	238	152	217	340	482	551	803	1,398	2,220	2,913
1999.....	59,336	446	264	163	181	391	418	557	827	1,423	2,225	2,652

White, female

2003.....	104,683	294	220	120	161	243	334	394	548	905	1,525	2,128
2002.....	101,179	295	213	124	164	214	314	364	501	868	1,534	2,089
2001.....	97,273	329	199	156	155	217	294	351	548	868	1,425	1,926
2000.....	92,791	331	184	125	158	218	285	329	514	871	1,427	1,659
1999.....	89,278	304	218	151	180	218	292	355	515	841	1,263	1,663

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2003.....	23,223	342	137	72	114	159	251	356	471	773	1,127	1,534
2002.....	22,519	305	150	73	103	135	285	323	466	744	1,210	1,538
2001.....	21,970	334	133	67	100	167	225	304	459	781	1,184	1,553
2000.....	21,272	305	148	79	110	150	233	344	472	793	1,176	1,517
1999.....	20,325	343	131	101	101	145	243	283	425	846	1,172	1,416

Black, male

2003.....	10,126	186	80	29	71	77	111	167	226	384	574	809
2002.....	9,847	168	82	38	60	68	140	160	238	360	602	841
2001.....	9,671	195	64	26	57	89	111	155	229	409	633	855
2000.....	9,455	187	78	35	52	80	124	165	239	415	632	819
1999.....	9,071	199	68	57	62	76	129	136	212	437	615	777

Black, female

2003.....	13,097	156	57	43	43	82	140	189	245	389	553	725
2002.....	12,672	137	68	35	43	67	145	163	228	384	608	697
2001.....	12,299	139	69	41	43	78	114	149	230	372	551	698
2000.....	11,817	118	70	44	58	70	109	179	233	378	544	698
1999.....	11,254	144	63	44	39	69	114	147	213	409	557	639

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

American Indian 1/, both sexes

2003.....	1,243	10	12	7	9	18	18	17	42	52	107	98
2002.....	1,102	11	11	6	5	14	20	17	35	56	70	95
2001.....	1,109	15	11	2	6	11	13	30	41	55	83	92
2000.....	945	17	17	2	5	7	9	18	28	58	74	69
1999.....	951	16	12	6	6	12	13	21	29	54	86	72

American Indian 1/, male

2003.....	606	3	8	5	6	11	13	11	21	31	60	62
2002.....	562	4	8	3	3	10	10	14	20	36	45	56
2001.....	547	11	5	1	3	7	11	19	23	36	49	52
2000.....	471	8	7	2	3	4	7	13	24	35	49	38
1999.....	469	11	7	2	3	5	7	15	17	28	55	45

American Indian 1/, female

2003.....	637	7	4	2	3	7	5	6	21	21	47	36
2002.....	540	7	3	3	2	4	10	3	15	20	25	39
2001.....	562	4	6	1	3	4	2	11	18	19	34	40
2000.....	474	9	10	-	2	3	2	5	4	23	25	31
1999.....	482	5	5	4	3	7	6	6	12	26	31	27

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Asian or Pacific Islander 2/, both sexes

2003.....	2,550	42	25	22	16	18	29	37	51	46	64	84
2002.....	2,339	40	30	12	15	17	34	26	36	47	64	82
2001.....	2,034	37	31	12	10	19	29	31	44	35	57	71
2000.....	1,980	43	22	18	13	17	23	15	23	41	55	69
1999.....	1,793	39	22	15	17	19	20	19	31	39	40	74

Asian or Pacific Islander 2/, male

2003.....	1,198	20	13	13	11	9	15	15	31	27	48	45
2002.....	1,146	25	16	8	8	8	16	14	23	20	43	53
2001.....	924	27	17	8	7	9	13	16	23	18	33	38
2000.....	911	25	15	9	4	8	12	8	8	25	27	41
1999.....	908	24	15	7	8	11	8	9	17	24	28	36

Asian or Pacific Islander 2/, female

2003.....	1,352	22	12	9	5	9	14	22	20	19	16	39
2002.....	1,193	15	14	4	7	9	18	12	13	27	21	29
2001.....	1,110	10	14	4	3	10	16	15	21	17	24	33
2000.....	1,069	18	7	9	9	9	11	7	15	16	28	28
1999.....	885	15	7	8	9	8	12	10	14	15	12	38

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races, both sexes

2003.....	109,277	945	1,717	1,096	1,522	6,755	8,517	6,284	6,257	7,495	9,271	8,865
2002.....	106,742	946	1,641	1,176	1,542	7,137	8,275	6,122	6,447	7,642	9,068	8,241
2001.....	101,537	976	1,714	1,283	1,553	6,646	7,765	5,935	5,904	7,589	8,356	7,469
2000.....	97,900	881	1,826	1,391	1,588	6,755	7,358	5,898	5,871	7,471	7,942	6,870
1999.....	97,860	845	1,898	1,459	1,632	6,688	6,968	5,881	6,009	7,366	7,865	6,696

All races, male

2003.....	70,532	539	1,048	650	1,012	4,706	6,585	4,826	4,708	5,390	6,613	6,325
2002.....	69,257	553	1,024	720	993	4,981	6,457	4,768	4,867	5,535	6,477	5,940
2001.....	66,060	541	1,045	786	1,017	4,722	6,110	4,605	4,487	5,502	6,110	5,519
2000.....	63,817	501	1,098	824	1,037	4,757	5,703	4,606	4,453	5,537	5,907	5,103
1999.....	63,535	494	1,127	897	1,077	4,624	5,416	4,555	4,542	5,411	5,837	4,996

All races, female

2003.....	38,745	406	669	446	510	2,049	1,932	1,458	1,549	2,105	2,658	2,540
2002.....	37,485	393	617	456	549	2,156	1,818	1,354	1,580	2,107	2,591	2,301
2001.....	35,477	435	669	497	536	1,924	1,655	1,330	1,417	2,087	2,246	1,950
2000.....	34,083	380	728	567	551	1,998	1,655	1,292	1,418	1,934	2,035	1,767
1999.....	34,325	351	771	562	555	2,064	1,552	1,326	1,467	1,955	2,028	1,700

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2003.....	93,381	624	1,304	805	1,149	5,771	7,219	5,217	5,183	6,226	7,790	7,353
2002.....	90,866	583	1,199	810	1,155	6,072	6,976	5,004	5,374	6,270	7,631	6,784
2001.....	85,964	647	1,309	906	1,154	5,563	6,475	4,877	4,850	6,269	6,874	6,011
2000.....	82,592	569	1,335	996	1,223	5,731	6,092	4,736	4,783	6,171	6,492	5,554
1999.....	82,245	562	1,374	1,011	1,259	5,647	5,717	4,790	4,871	6,051	6,449	5,369

White, male

2003.....	59,912	359	807	472	761	4,005	5,591	4,041	3,923	4,480	5,608	5,247
2002.....	58,467	358	736	497	743	4,214	5,465	3,907	4,079	4,546	5,494	4,949
2001.....	55,493	356	815	536	749	3,930	5,137	3,803	3,715	4,571	5,076	4,443
2000.....	53,329	331	808	587	791	4,007	4,763	3,704	3,665	4,608	4,854	4,143
1999.....	52,898	333	811	611	823	3,862	4,465	3,741	3,701	4,477	4,811	4,002

White, female

2003.....	33,469	265	497	333	388	1,766	1,628	1,176	1,260	1,746	2,182	2,106
2002.....	32,399	225	463	313	412	1,858	1,511	1,097	1,295	1,724	2,137	1,835
2001.....	30,471	291	494	370	405	1,633	1,338	1,074	1,135	1,698	1,798	1,568
2000.....	29,263	238	527	409	432	1,724	1,329	1,032	1,118	1,563	1,638	1,411
1999.....	29,347	229	563	400	436	1,785	1,252	1,049	1,170	1,574	1,638	1,367

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2003.....	12,351	287	343	235	288	667	951	800	835	1,010	1,214	1,263
2002.....	12,513	330	359	306	315	758	968	867	844	1,105	1,163	1,232
2001.....	12,462	281	339	319	325	775	967	806	832	1,049	1,272	1,233
2000.....	12,277	274	405	338	293	737	958	892	878	1,058	1,229	1,109
1999.....	12,728	252	441	396	311	774	945	855	910	1,081	1,204	1,153

Black, male

2003.....	8,385	160	204	142	192	490	737	601	620	735	830	907
2002.....	8,612	176	239	185	215	558	740	677	632	803	789	853
2001.....	8,537	162	194	216	226	576	733	609	611	735	898	934
2000.....	8,531	149	239	209	205	542	719	693	647	772	906	817
1999.....	8,773	141	266	258	214	574	727	658	677	769	891	877

Black, female

2003.....	3,966	127	139	93	96	177	214	199	215	275	384	356
2002.....	3,901	154	120	121	100	200	228	190	212	302	374	379
2001.....	3,925	119	145	103	99	199	234	197	221	314	374	299
2000.....	3,746	125	166	129	88	195	239	199	231	286	323	292
1999.....	3,955	111	175	138	97	200	218	197	233	312	313	276

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

American Indian 1/, both sexes

2003.....	1,573	20	33	28	40	153	162	137	127	157	156	127
2002.....	1,488	15	33	23	37	146	154	132	122	166	167	117
2001.....	1,361	32	30	29	44	145	148	102	114	152	108	113
2000.....	1,353	20	39	31	39	149	150	123	116	129	127	108
1999.....	1,327	13	37	25	26	130	159	119	131	129	123	96

American Indian 1/, male

2003.....	1,043	14	19	19	29	102	126	88	89	111	104	89
2002.....	1,003	8	19	11	20	91	121	101	85	114	114	78
2001.....	908	14	15	20	26	95	108	80	83	117	71	70
2000.....	900	11	25	16	21	102	100	94	78	81	85	80
1999.....	890	8	24	12	16	95	119	72	91	90	77	74

American Indian 1/, female

2003.....	530	6	14	9	11	51	36	49	38	46	52	38
2002.....	485	7	14	12	17	55	33	31	37	52	53	39
2001.....	453	18	15	9	18	50	40	22	31	35	37	43
2000.....	453	9	14	15	18	47	50	29	38	48	42	28
1999.....	437	5	13	13	10	35	40	47	40	39	46	22

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Asian or Pacific Islander 2/, both sexes

2003.....	1,972	14	37	28	45	164	185	130	112	102	111	122
2002.....	1,875	18	50	37	35	161	177	119	107	101	107	108
2001.....	1,750	16	36	29	30	163	175	150	108	119	102	112
2000.....	1,678	18	47	26	33	138	158	147	94	113	94	99
1999.....	1,560	18	46	27	36	137	147	117	97	105	89	78

Asian or Pacific Islander 2/, male

2003.....	1,192	6	18	17	30	109	131	96	76	64	71	82
2002.....	1,175	11	30	27	15	118	131	83	71	72	80	60
2001.....	1,122	9	21	14	16	121	132	113	78	79	65	72
2000.....	1,057	10	26	12	20	106	121	115	63	76	62	63
1999.....	974	12	26	16	24	93	105	84	73	75	58	43

Asian or Pacific Islander 2/, female

2003.....	780	8	19	11	15	55	54	34	36	38	40	40
2002.....	700	7	20	10	20	43	46	36	36	29	27	48
2001.....	628	7	15	15	14	42	43	37	30	40	37	40
2000.....	621	8	21	14	13	32	37	32	31	37	32	36
1999.....	586	6	20	11	12	44	42	33	24	30	31	35

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
All races, both sexes												
2003.....	48,071	148	649	694	1,071	5,471	5,961	3,855	3,425	3,579	4,015	3,656
2002.....	48,366	127	640	705	1,033	5,782	6,034	3,891	3,593	3,750	3,938	3,593
2001.....	47,288	151	678	769	1,054	5,393	5,746	3,953	3,501	3,881	3,946	3,379
2000.....	46,749	174	691	842	1,080	5,475	5,553	3,957	3,443	3,889	3,772	3,248
1999.....	46,423	189	712	897	1,078	5,452	5,237	3,922	3,512	3,869	3,684	3,121
All races, male												
2003.....	33,291	81	366	401	683	3,653	4,518	2,915	2,596	2,585	2,912	2,667
2002.....	33,533	72	381	405	634	3,875	4,596	2,990	2,686	2,703	2,796	2,638
2001.....	32,794	68	392	447	668	3,694	4,417	3,015	2,646	2,744	2,869	2,452
2000.....	32,166	94	391	464	674	3,693	4,197	3,016	2,554	2,802	2,758	2,338
1999.....	31,642	101	400	532	674	3,593	3,965	2,952	2,559	2,741	2,635	2,234
All races, female												
2003.....	14,780	67	283	293	388	1,818	1,443	940	829	994	1,103	989
2002.....	14,833	55	259	300	399	1,907	1,438	901	907	1,047	1,142	955
2001.....	14,494	83	286	322	386	1,699	1,329	938	855	1,137	1,077	927
2000.....	14,583	80	300	378	406	1,782	1,356	941	889	1,087	1,014	910
1999.....	14,781	88	312	365	404	1,859	1,272	970	953	1,128	1,049	887

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
White, both sexes												
2003.....	40,282	111	491	538	859	4,686	4,957	3,125	2,760	2,925	3,328	3,043
2002.....	40,614	84	478	525	822	4,968	5,023	3,101	2,936	3,032	3,290	2,982
2001.....	39,544	115	511	558	820	4,563	4,722	3,227	2,826	3,179	3,272	2,743
2000.....	39,088	121	513	646	873	4,713	4,566	3,121	2,769	3,208	3,110	2,679
1999.....	38,767	137	497	664	878	4,661	4,292	3,171	2,855	3,140	3,059	2,563
White, male												
2003.....	27,925	63	285	305	546	3,114	3,753	2,383	2,106	2,110	2,430	2,225
2002.....	28,118	50	278	305	509	3,316	3,845	2,388	2,202	2,189	2,351	2,220
2001.....	27,407	52	299	320	515	3,107	3,667	2,473	2,147	2,249	2,383	1,992
2000.....	26,815	68	290	358	538	3,158	3,474	2,377	2,064	2,312	2,277	1,945
1999.....	26,385	80	273	390	546	3,047	3,269	2,401	2,076	2,232	2,196	1,837
White, female												
2003.....	12,357	48	206	233	313	1,572	1,204	742	654	815	898	818
2002.....	12,496	34	200	220	313	1,652	1,178	713	734	843	939	762
2001.....	12,137	63	212	238	305	1,456	1,055	754	679	930	889	751
2000.....	12,273	53	223	288	335	1,555	1,092	744	705	896	833	734
1999.....	12,382	57	224	274	332	1,614	1,023	770	779	908	863	726

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Black, both sexes												
2003.....	5,771	28	129	117	161	521	721	546	510	508	546	476
2002.....	5,796	36	126	151	161	569	745	608	504	557	497	482
2001.....	5,906	19	130	179	184	590	757	556	524	531	556	504
2000.....	5,839	40	141	157	160	539	744	634	535	527	542	457
1999.....	5,901	44	177	202	162	580	704	578	511	587	498	449
Black, male												
2003.....	4,119	12	66	73	102	371	562	410	389	381	387	357
2002.....	4,156	18	83	83	100	395	554	473	383	406	343	345
2001.....	4,214	7	72	113	127	423	563	419	394	375	413	387
2000.....	4,199	21	79	87	110	380	548	489	401	396	408	321
1999.....	4,162	18	101	127	105	405	521	443	385	412	366	331
Black, female												
2003.....	1,652	16	63	44	59	150	159	136	121	127	159	119
2002.....	1,640	18	43	68	61	174	191	135	121	151	154	137
2001.....	1,692	12	58	66	57	167	194	137	130	156	143	117
2000.....	1,640	19	62	70	50	159	196	145	134	131	134	136
1999.....	1,739	26	76	75	57	175	183	135	126	175	132	118

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
American Indian 1/, both sexes												
2003.....	915	7	18	20	26	126	137	98	80	82	72	67
2002.....	903	4	16	12	27	123	127	89	82	94	89	56
2001.....	832	11	17	16	32	121	126	73	74	91	55	61
2000.....	829	7	16	19	27	120	121	94	73	80	66	53
1999.....	816	4	15	16	19	107	126	87	76	76	72	52
American Indian 1/, male												
2003.....	595	5	8	11	20	82	103	64	57	55	52	41
2002.....	603	1	9	6	14	74	95	66	56	63	60	36
2001.....	558	5	8	8	18	78	86	56	55	73	39	33
2000.....	537	2	10	9	14	80	82	69	48	50	42	38
1999.....	547	2	12	8	13	77	91	50	50	51	45	39
American Indian 1/, female												
2003.....	320	2	10	9	6	44	34	34	23	27	20	26
2002.....	300	3	7	6	13	49	32	23	26	31	29	20
2001.....	274	6	9	8	14	43	40	17	19	18	16	28
2000.....	292	5	6	10	13	40	39	25	25	30	24	15
1999.....	269	2	3	8	6	30	35	37	26	25	27	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Asian or Pacific Islander 2/, both sexes												
2003.....	1,103	2	11	19	25	138	146	86	75	64	69	70
2002.....	1,053	3	20	17	23	122	139	93	71	67	62	73
2001.....	1,006	6	20	16	18	119	141	97	77	80	63	71
2000.....	993	6	21	20	20	103	122	108	66	74	54	59
1999.....	939	4	23	15	19	104	115	86	70	66	55	57
Asian or Pacific Islander 2/, male												
2003.....	652	1	7	12	15	86	100	58	44	39	43	44
2002.....	656	3	11	11	11	90	102	63	45	45	42	37
2001.....	615	4	13	6	8	86	101	67	50	47	34	40
2000.....	615	3	12	10	12	75	93	81	41	44	31	34
1999.....	548	1	14	7	10	64	84	58	48	46	28	27
Asian or Pacific Islander 2/, female												
2003.....	451	1	4	7	10	52	46	28	31	25	26	26
2002.....	397	-	9	6	12	32	37	30	26	22	20	36
2001.....	391	2	7	10	10	33	40	30	27	33	29	31
2000.....	378	3	9	10	8	28	29	27	25	30	23	25
1999.....	391	3	9	8	9	40	31	28	22	20	27	30

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All races, both sexes												
2003.....	44,757	145	621	646	996	5,269	5,703	3,644	3,189	3,289	3,672	3,286
2002.....	45,380	123	610	664	950	5,634	5,825	3,731	3,361	3,462	3,615	3,240
2001.....	43,788	144	626	718	968	5,208	5,517	3,710	3,227	3,545	3,538	3,011
2000.....	43,354	168	651	780	992	5,251	5,309	3,715	3,169	3,545	3,382	2,936
1999.....	42,401	184	650	802	969	5,198	4,930	3,604	3,174	3,460	3,278	2,747
All races, male												
2003.....	30,655	80	353	373	636	3,500	4,306	2,753	2,403	2,350	2,634	2,366
2002.....	31,064	68	361	385	578	3,765	4,413	2,855	2,494	2,459	2,525	2,328
2001.....	30,064	66	356	418	611	3,546	4,232	2,826	2,410	2,473	2,543	2,164
2000.....	29,451	90	368	429	614	3,527	3,989	2,804	2,331	2,507	2,433	2,091
1999.....	28,552	97	366	469	593	3,412	3,715	2,690	2,303	2,424	2,311	1,936
All races, female												
2003.....	14,102	65	268	273	360	1,769	1,397	891	786	939	1,038	920
2002.....	14,316	55	249	279	372	1,869	1,412	876	867	1,003	1,090	912
2001.....	13,724	78	270	300	357	1,662	1,285	884	817	1,072	995	847
2000.....	13,903	78	283	351	378	1,724	1,320	911	838	1,038	949	845
1999.....	13,849	87	284	333	376	1,786	1,215	914	871	1,036	967	811

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
White, both sexes												
2003.....	37,424	108	467	501	793	4,517	4,740	2,942	2,565	2,675	3,030	2,735
2002.....	38,054	81	452	494	760	4,840	4,840	2,963	2,734	2,787	3,020	2,683
2001.....	36,576	109	466	522	753	4,409	4,520	3,014	2,599	2,904	2,924	2,447
2000.....	36,160	115	484	593	796	4,530	4,351	2,908	2,540	2,918	2,772	2,421
1999.....	35,343	133	455	591	786	4,450	4,035	2,897	2,572	2,791	2,706	2,244
White, male												
2003.....	25,660	62	273	286	508	2,986	3,576	2,242	1,947	1,909	2,187	1,975
2002.....	26,000	47	260	288	467	3,221	3,683	2,273	2,033	1,979	2,125	1,955
2001.....	25,084	50	268	298	472	2,982	3,506	2,306	1,949	2,026	2,107	1,756
2000.....	24,470	64	275	330	485	3,026	3,291	2,191	1,876	2,061	1,996	1,739
1999.....	23,763	77	253	344	479	2,899	3,060	2,175	1,863	1,961	1,918	1,583
White, female												
2003.....	11,764	46	194	215	285	1,531	1,164	700	618	766	843	760
2002.....	12,054	34	192	206	293	1,619	1,157	690	701	808	895	728
2001.....	11,492	59	198	224	281	1,427	1,014	708	650	878	817	691
2000.....	11,690	51	209	263	311	1,504	1,060	717	664	857	776	682
1999.....	11,580	56	202	247	307	1,551	975	722	709	830	788	661

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2003.....	5,431	28	125	110	153	497	692	524	483	476	511	428
2002.....	5,462	35	123	141	146	553	726	591	481	519	461	432
2001.....	5,506	19	123	165	166	568	740	533	492	491	506	447
2000.....	5,499	40	134	149	153	513	721	614	497	485	502	415
1999.....	5,470	43	160	184	150	553	677	547	471	538	457	408

Black, male

2003.....	3,834	12	65	67	94	352	535	393	366	353	361	317
2002.....	3,874	17	81	80	89	382	539	457	364	376	313	303
2001.....	3,900	7	67	106	113	406	547	402	369	343	370	342
2000.....	3,922	21	74	80	105	358	528	470	372	361	374	288
1999.....	3,821	17	89	111	94	384	501	418	353	374	330	299

Black, female

2003.....	1,597	16	60	43	59	145	157	131	117	123	150	111
2002.....	1,588	18	42	61	57	171	187	134	117	143	148	129
2001.....	1,606	12	56	59	53	162	193	131	123	148	136	105
2000.....	1,577	19	60	69	48	155	193	144	125	124	128	127
1999.....	1,649	26	71	73	56	169	176	129	118	164	127	109

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

American Indian 1/, both sexes

2003.....	865	7	18	19	25	121	133	95	69	77	68	63
2002.....	866	4	16	12	26	122	121	87	78	91	77	54
2001.....	764	10	17	16	31	116	121	69	65	79	50	53
2000.....	779	7	14	19	25	113	118	92	71	74	61	45
1999.....	730	4	14	14	16	98	112	82	68	70	63	42

American Indian 1/, male

2003.....	557	5	8	11	19	78	99	61	48	52	49	38
2002.....	572	1	9	6	13	74	90	65	53	61	49	34
2001.....	506	5	8	8	18	74	82	53	47	63	35	30
2000.....	500	2	9	9	14	75	79	67	47	45	38	32
1999.....	481	2	11	7	12	70	79	45	45	46	38	30

American Indian 1/, female

2003.....	308	2	10	8	6	43	34	34	21	25	19	25
2002.....	294	3	7	6	13	48	31	22	25	30	28	20
2001.....	258	5	9	8	13	42	39	16	18	16	15	23
2000.....	279	5	5	10	11	38	39	25	24	29	23	13
1999.....	249	2	3	7	4	28	33	37	23	24	25	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Asian or Pacific Islander 2/, both sexes

2003.....	1,037	2	11	16	25	134	138	83	72	61	63	60
2002.....	998	3	19	17	18	119	138	90	68	65	57	71
2001.....	942	6	20	15	18	115	136	94	71	71	58	64
2000.....	916	6	19	19	18	95	119	101	61	68	47	55
1999.....	858	4	21	13	17	97	106	78	63	61	52	53

Asian or Pacific Islander 2/, male

2003.....	604	1	7	9	15	84	96	57	42	36	37	36
2002.....	618	3	11	11	9	88	101	60	44	43	38	36
2001.....	574	4	13	6	8	84	97	65	45	41	31	36
2000.....	559	3	10	10	10	68	91	76	36	40	25	32
1999.....	487	1	13	7	8	59	75	52	42	43	25	24

Asian or Pacific Islander 2/, female

2003.....	433	1	4	7	10	50	42	26	30	25	26	24
2002.....	380	-	8	6	9	31	37	30	24	22	19	35
2001.....	368	2	7	9	10	31	39	29	26	30	27	28
2000.....	357	3	9	9	8	27	28	25	25	28	22	23
1999.....	371	3	8	6	9	38	31	26	21	18	27	29

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
All races, both sexes												
2003.....	1,357	2	17	23	37	129	135	100	104	107	132	137
2002.....	1,086	2	17	21	49	63	96	59	79	101	122	121
2001.....	1,405	3	28	25	55	95	103	85	92	141	144	142
2000.....	1,492	2	31	39	59	128	115	94	118	149	168	125
1999.....	2,118	4	48	62	66	172	190	152	165	205	203	179
All races, male												
2003.....	1,044	-	9	12	26	95	110	80	86	85	101	114
2002.....	882	2	10	11	36	49	85	47	61	85	102	107
2001.....	1,067	-	20	13	36	72	86	71	81	117	118	105
2000.....	1,125	2	17	21	40	99	95	81	90	121	137	92
1999.....	1,559	4	26	43	48	119	160	126	118	157	162	135
All races, female												
2003.....	313	2	8	11	11	34	25	20	18	22	31	23
2002.....	204	-	7	10	13	14	11	12	18	16	20	14
2001.....	338	3	8	12	19	23	17	14	11	24	26	37
2000.....	367	-	14	18	19	29	20	13	28	28	31	33
1999.....	559	-	22	19	18	53	30	26	47	48	41	44

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2003.....	1,121	2	14	16	33	110	117	85	80	88	106	103
2002.....	870	1	13	17	36	54	82	49	66	80	90	91
2001.....	1,105	2	23	16	40	72	89	74	67	99	109	106
2000.....	1,221	2	21	34	49	109	94	77	94	118	132	94
1999.....	1,755	3	32	44	55	141	154	126	133	169	171	146

White, male

2003.....	856	-	8	7	22	81	96	67	67	69	85	82
2002.....	708	1	8	11	27	43	75	38	51	69	74	79
2001.....	850	-	17	9	25	57	75	63	61	84	89	80
2000.....	924	2	10	18	34	83	78	67	72	98	108	70
1999.....	1,281	3	15	29	39	96	127	104	95	130	136	108

White, female

2003.....	265	2	6	9	11	29	21	18	13	19	21	21
2002.....	162	-	5	6	9	11	7	11	15	11	16	12
2001.....	255	2	6	7	15	15	14	11	6	15	20	26
2000.....	297	-	11	16	15	26	16	10	22	20	24	24
1999.....	474	-	17	15	16	45	27	22	38	39	35	38

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Black, both sexes

2003.....	190	-	3	5	3	14	13	13	17	16	23	29
2002.....	172	1	3	4	10	9	9	10	9	18	22	29
2001.....	226	-	5	9	14	16	7	8	17	28	32	29
2000.....	206	-	7	4	7	14	18	11	21	26	28	25
1999.....	253	1	13	15	6	18	16	18	24	29	23	24

Black, male

2003.....	155	-	1	4	3	10	11	11	14	14	14	27
2002.....	145	1	2	-	7	6	6	9	8	14	20	28
2001.....	168	-	3	4	11	11	6	6	13	22	27	22
2000.....	158	-	5	3	5	11	15	10	15	21	22	20
1999.....	196	1	9	14	6	14	14	16	18	22	19	19

Black, female

2003.....	35	-	2	1	-	4	2	2	3	2	9	2
2002.....	27	-	1	4	3	3	3	1	1	4	2	1
2001.....	58	-	2	5	3	5	1	2	4	6	5	7
2000.....	48	-	2	1	2	3	3	1	6	5	6	5
1999.....	57	-	4	1	-	4	2	2	6	7	4	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

American Indian 1/, both sexes

2003.....	28	-	-	1	1	5	2	2	7	3	1	2
2002.....	21	-	-	-	1	-	5	-	3	3	6	-
2001.....	39	1	-	-	1	4	4	3	6	8	1	4
2000.....	28	-	1	-	2	4	1	2	1	3	5	3
1999.....	60	-	1	1	3	8	12	2	6	5	7	9

American Indian 1/, male

2003.....	21	-	-	-	1	4	2	2	5	2	-	2
2002.....	17	-	-	-	1	-	4	-	2	2	5	-
2001.....	30	-	-	-	-	3	3	2	6	7	1	2
2000.....	19	-	-	-	-	4	1	2	1	2	4	1
1999.....	45	-	1	-	1	6	11	2	3	4	5	8

American Indian 1/, female

2003.....	7	-	-	1	-	1	-	-	2	1	1	-
2002.....	4	-	-	-	-	-	1	-	1	1	1	-
2001.....	9	1	-	-	1	1	1	1	-	1	-	2
2000.....	9	-	1	-	2	-	-	-	-	1	1	2
1999.....	15	-	-	1	2	2	1	-	3	1	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Asian or Pacific Islander 2/, both sexes

2003.....	18	-	-	1	-	-	3	-	-	-	2	3
2002.....	23	-	1	-	2	-	-	-	1	-	4	1
2001.....	35	-	-	-	-	3	3	-	2	6	2	3
2000.....	37	-	2	1	1	1	2	4	2	2	3	3
1999.....	50	-	2	2	2	5	8	6	2	2	2	-

Asian or Pacific Islander 2/, male

2003.....	12	-	-	1	-	-	1	-	-	-	2	3
2002.....	12	-	-	-	1	-	-	-	-	-	3	-
2001.....	19	-	-	-	-	1	2	-	1	4	1	1
2000.....	24	-	2	-	1	1	1	2	2	-	3	1
1999.....	37	-	1	-	2	3	8	4	2	1	2	-

Asian or Pacific Islander 2/, female

2003.....	6	-	-	-	-	-	2	-	-	-	-	-
2002.....	11	-	1	-	1	-	-	-	1	-	1	1
2001.....	16	-	-	-	-	2	1	-	1	2	1	2
2000.....	13	-	-	1	-	-	1	2	-	2	-	2
1999.....	13	-	1	2	-	2	-	2	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races, both sexes

2003.....	1,957	1	11	25	38	73	123	111	132	183	211	233
2002.....	1,900	2	13	20	34	85	113	101	153	187	201	232
2001.....	2,095	4	24	26	31	90	126	158	182	195	264	226
2000.....	1,903	4	9	23	29	96	129	148	156	195	222	187
1999.....	1,904	1	14	33	43	82	117	166	173	204	203	195

All races, male

2003.....	1,592	1	4	16	21	58	102	82	107	150	177	187
2002.....	1,587	2	10	9	20	61	98	88	131	159	169	203
2001.....	1,663	2	16	16	21	76	99	118	155	154	208	183
2000.....	1,590	2	6	14	20	67	113	131	133	174	188	155
1999.....	1,531	-	8	20	33	62	90	136	138	160	162	163

All races, female

2003.....	365	-	7	9	17	15	21	29	25	33	34	46
2002.....	313	-	3	11	14	24	15	13	22	28	32	29
2001.....	432	2	8	10	10	14	27	40	27	41	56	43
2000.....	313	2	3	9	9	29	16	17	23	21	34	32
1999.....	373	1	6	13	10	20	27	30	35	44	41	32

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
White, both sexes												
2003.....	1,737	1	10	21	33	59	100	98	115	162	192	205
2002.....	1,690	2	13	14	26	74	101	89	136	165	180	208
2001.....	1,863	4	22	20	27	82	113	139	160	176	239	190
2000.....	1,707	4	8	19	28	74	121	136	135	172	206	164
1999.....	1,669	1	10	29	37	70	103	148	150	180	182	173
White, male												
2003.....	1,409	1	4	12	16	47	81	74	92	132	158	168
2002.....	1,410	2	10	6	15	52	87	77	118	141	152	186
2001.....	1,473	2	14	13	18	68	86	104	137	139	187	156
2000.....	1,421	2	5	10	19	49	105	119	116	153	173	136
1999.....	1,341	-	5	17	28	52	82	122	118	141	142	146
White, female												
2003.....	328	-	6	9	17	12	19	24	23	30	34	37
2002.....	280	-	3	8	11	22	14	12	18	24	28	22
2001.....	390	2	8	7	9	14	27	35	23	37	52	34
2000.....	286	2	3	9	9	25	16	17	19	19	33	28
1999.....	328	1	5	12	9	18	21	26	32	39	40	27

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Black, both sexes

2003.....	150	-	1	2	5	10	16	9	10	16	12	19
2002.....	162	-	-	6	5	7	10	7	14	20	14	21
2001.....	174	-	2	5	4	6	10	15	15	12	18	28
2000.....	134	-	-	4	-	12	5	9	17	16	12	17
1999.....	178	-	4	3	6	9	11	13	16	20	18	17

Black, male

2003.....	130	-	-	2	5	9	16	6	9	14	12	13
2002.....	137	-	-	3	4	7	9	7	11	16	10	14
2001.....	146	-	2	3	3	6	10	11	12	10	16	23
2000.....	119	-	-	4	-	11	5	9	14	14	12	13
1999.....	145	-	3	2	5	7	6	9	14	16	17	13

Black, female

2003.....	20	-	1	-	-	1	-	3	1	2	-	6
2002.....	25	-	-	3	1	-	1	-	3	4	4	7
2001.....	28	-	-	2	1	-	-	4	3	2	2	5
2000.....	15	-	-	-	-	1	-	-	3	2	-	4
1999.....	33	-	1	1	1	2	5	4	2	4	1	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

American Indian 1/, both sexes

2003.....	22	-	-	-	-	-	2	1	4	2	3	2
2002.....	16	-	-	-	-	1	1	2	1	-	6	2
2001.....	29	-	-	-	-	1	1	1	3	4	4	4
2000.....	22	-	1	-	-	3	2	-	1	3	-	5
1999.....	26	-	-	1	-	1	2	3	2	1	2	1

American Indian 1/, male

2003.....	17	-	-	-	-	-	2	1	4	1	3	1
2002.....	14	-	-	-	-	-	1	1	1	-	6	2
2001.....	22	-	-	-	-	1	1	1	2	3	3	1
2000.....	18	-	1	-	-	1	2	-	-	3	-	5
1999.....	21	-	-	1	-	1	1	3	2	1	2	1

American Indian 1/, female

2003.....	5	-	-	-	-	-	-	-	-	1	-	1
2002.....	2	-	-	-	-	1	-	1	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	1	1	1	3
2000.....	4	-	-	-	-	2	-	-	1	-	-	-
1999.....	5	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Asian or Pacific Islander 2/, both sexes

2003.....	48	-	-	2	-	4	5	3	3	3	4	7
2002.....	32	-	-	-	3	3	1	3	2	2	1	1
2001.....	29	-	-	1	-	1	2	3	4	3	3	4
2000.....	40	-	-	-	1	7	1	3	3	4	4	1
1999.....	31	-	-	-	-	2	1	2	5	3	1	4

Asian or Pacific Islander 2/, male

2003.....	36	-	-	2	-	2	3	1	2	3	4	5
2002.....	26	-	-	-	1	2	1	3	1	2	1	1
2001.....	22	-	-	-	-	1	2	2	4	2	2	3
2000.....	32	-	-	-	1	6	1	3	3	4	3	1
1999.....	24	-	-	-	-	2	1	2	4	2	1	3

Asian or Pacific Islander 2/, female

2003.....	12	-	-	-	-	2	2	2	1	-	-	2
2002.....	6	-	-	-	2	1	-	-	1	-	-	-
2001.....	7	-	-	1	-	-	-	1	-	1	1	1
2000.....	8	-	-	-	-	1	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	-	1	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
All races, both sexes												
2003.....	61,206	797	1,068	402	451	1,284	2,556	2,429	2,832	3,916	5,256	5,209
2002.....	58,376	819	1,001	471	509	1,355	2,241	2,231	2,854	3,892	5,130	4,648
2001.....	54,249	825	1,036	514	499	1,253	2,019	1,982	2,403	3,708	4,410	4,090
2000.....	51,151	707	1,135	549	508	1,280	1,805	1,941	2,428	3,582	4,170	3,622
1999.....	51,437	656	1,186	562	554	1,236	1,731	1,959	2,497	3,497	4,181	3,575
All races, male												
2003.....	37,241	458	682	249	329	1,053	2,067	1,911	2,112	2,805	3,701	3,658
2002.....	35,724	481	643	315	359	1,106	1,861	1,778	2,181	2,832	3,681	3,302
2001.....	33,266	473	653	339	349	1,028	1,693	1,590	1,841	2,758	3,241	3,067
2000.....	31,651	407	707	360	363	1,064	1,506	1,590	1,899	2,735	3,149	2,765
1999.....	31,893	393	727	365	403	1,031	1,451	1,603	1,983	2,670	3,202	2,762
All races, female												
2003.....	23,965	339	386	153	122	231	489	518	720	1,111	1,555	1,551
2002.....	22,652	338	358	156	150	249	380	453	673	1,060	1,449	1,346
2001.....	20,983	352	383	175	150	225	326	392	562	950	1,169	1,023
2000.....	19,500	300	428	189	145	216	299	351	529	847	1,021	857
1999.....	19,544	263	459	197	151	205	280	356	514	827	979	813

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
White, both sexes												
2003.....	53,099	513	813	267	290	1,085	2,262	2,092	2,423	3,301	4,462	4,310
2002.....	50,252	499	721	285	333	1,104	1,953	1,903	2,438	3,238	4,341	3,802
2001.....	46,420	532	798	348	334	1,000	1,753	1,650	2,024	3,090	3,602	3,268
2000.....	43,504	448	822	350	350	1,018	1,526	1,615	2,014	2,963	3,382	2,875
1999.....	43,478	425	877	347	381	986	1,425	1,619	2,016	2,911	3,390	2,806
White, male												
2003.....	31,987	296	522	167	215	891	1,838	1,658	1,817	2,370	3,178	3,022
2002.....	30,349	308	458	192	234	898	1,620	1,519	1,877	2,357	3,143	2,729
2001.....	28,086	304	516	216	234	823	1,470	1,330	1,568	2,322	2,693	2,451
2000.....	26,514	263	518	229	253	849	1,289	1,327	1,601	2,296	2,577	2,198
1999.....	26,513	253	538	221	277	815	1,196	1,340	1,625	2,245	2,615	2,165
White, female												
2003.....	21,112	217	291	100	75	194	424	434	606	931	1,284	1,288
2002.....	19,903	191	263	93	99	206	333	384	561	881	1,198	1,073
2001.....	18,334	228	282	132	100	177	283	320	456	768	909	817
2000.....	16,990	185	304	121	97	169	237	288	413	667	805	677
1999.....	16,965	172	339	126	104	171	229	279	391	666	775	641

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
Black, both sexes												
2003.....	6,580	259	214	118	127	146	230	254	325	502	668	787
2002.....	6,717	294	233	155	154	189	223	259	340	548	666	750
2001.....	6,556	262	209	140	141	185	210	250	308	518	716	729
2000.....	6,438	234	264	181	133	198	214	258	343	531	687	652
1999.....	6,827	208	264	194	149	194	241	277	399	494	706	704
Black, male												
2003.....	4,266	148	138	69	90	119	175	191	231	354	443	550
2002.....	4,456	158	156	102	115	163	186	204	249	397	446	508
2001.....	4,323	155	122	103	99	153	170	190	217	360	485	547
2000.....	4,332	128	160	122	95	162	171	204	246	376	498	496
1999.....	4,611	123	165	131	109	169	206	215	292	357	525	546
Black, female												
2003.....	2,314	111	76	49	37	27	55	63	94	148	225	237
2002.....	2,261	136	77	53	39	26	37	55	91	151	220	242
2001.....	2,233	107	87	37	42	32	40	60	91	158	231	182
2000.....	2,106	106	104	59	38	36	43	54	97	155	189	156
1999.....	2,216	85	99	63	40	25	35	62	107	137	181	158

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
American Indian 1/, both sexes												
2003.....	658	13	15	8	14	27	25	39	47	75	84	60
2002.....	585	11	17	11	10	23	27	43	40	72	78	61
2001.....	529	21	13	13	12	24	22	29	40	61	53	52
2000.....	524	13	23	12	12	29	29	29	43	49	61	55
1999.....	511	9	22	9	7	23	33	32	55	53	51	44
American Indian 1/, male												
2003.....	448	9	11	8	9	20	23	24	32	56	52	48
2002.....	400	7	10	5	6	17	26	35	29	51	54	42
2001.....	350	9	7	12	8	17	22	24	28	44	32	37
2000.....	363	9	15	7	7	22	18	25	30	31	43	42
1999.....	343	6	12	4	3	18	28	22	41	39	32	35
American Indian 1/, female												
2003.....	210	4	4	-	5	7	2	15	15	19	32	12
2002.....	185	4	7	6	4	6	1	8	11	21	24	19
2001.....	179	12	6	1	4	7	-	5	12	17	21	15
2000.....	161	4	8	5	5	7	11	4	13	18	18	13
1999.....	168	3	10	5	4	5	5	10	14	14	19	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nontransport accidents (W00-X59,Y86)

Asian or Pacific Islander 2/, both sexes

2003.....	869	12	26	9	20	26	39	44	37	38	42	52
2002.....	822	15	30	20	12	39	38	26	36	34	45	35
2001.....	744	10	16	13	12	44	34	53	31	39	39	41
2000.....	685	12	26	6	13	35	36	39	28	39	40	40
1999.....	621	14	23	12	17	33	32	31	27	39	34	21

Asian or Pacific Islander 2/, male

2003.....	540	5	11	5	15	23	31	38	32	25	28	38
2002.....	519	8	19	16	4	28	29	20	26	27	38	23
2001.....	507	5	8	8	8	35	31	46	28	32	31	32
2000.....	442	7	14	2	8	31	28	34	22	32	31	29
1999.....	426	11	12	9	14	29	21	26	25	29	30	16

Asian or Pacific Islander 2/, female

2003.....	329	7	15	4	5	3	8	6	5	13	14	14
2002.....	303	7	11	4	8	11	9	6	10	7	7	12
2001.....	237	5	8	5	4	9	3	7	3	7	8	9
2000.....	243	5	12	4	5	4	8	5	6	7	9	11
1999.....	195	3	11	3	3	4	11	5	2	10	4	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
All races, both sexes												
2003.....	17,229	14	54	14	31	90	140	139	146	255	381	508
2002.....	16,257	16	37	18	24	83	164	130	177	256	408	487
2001.....	15,019	23	32	33	33	88	168	160	180	272	375	520
2000.....	13,322	8	36	16	21	99	138	138	165	238	370	409
1999.....	13,162	12	55	25	28	112	130	156	187	276	369	447
All races, male												
2003.....	8,910	9	39	9	21	71	115	123	114	218	307	391
2002.....	8,463	9	28	9	18	65	146	118	151	191	331	378
2001.....	8,089	15	25	22	24	79	153	142	153	233	311	426
2000.....	7,122	6	23	14	16	88	115	122	150	199	297	332
1999.....	7,109	8	37	19	21	95	114	137	157	225	303	366
All races, female												
2003.....	8,319	5	15	5	10	19	25	16	32	37	74	117
2002.....	7,794	7	9	9	6	18	18	12	26	65	77	109
2001.....	6,930	8	7	11	9	9	15	18	27	39	64	94
2000.....	6,200	2	13	2	5	11	23	16	15	39	73	77
1999.....	6,053	4	18	6	7	17	16	19	30	51	66	81

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
White, both sexes												
2003.....	15,855	10	38	13	26	79	126	122	127	211	320	413
2002.....	15,027	12	23	14	19	72	147	119	152	217	343	410
2001.....	13,827	15	28	27	29	80	157	140	157	227	310	429
2000.....	12,252	5	23	12	18	90	127	125	140	194	310	352
1999.....	11,992	8	43	18	24	100	112	135	149	239	308	362
White, male												
2003.....	8,099	6	29	8	17	63	106	108	101	183	261	322
2002.....	7,699	6	17	7	15	56	130	108	131	160	280	321
2001.....	7,331	10	21	20	20	71	146	126	134	195	261	348
2000.....	6,449	3	13	10	15	81	106	112	127	163	253	290
1999.....	6,352	7	29	14	18	86	100	120	126	193	249	298
White, female												
2003.....	7,756	4	9	5	9	16	20	14	26	28	59	91
2002.....	7,328	6	6	7	4	16	17	11	21	57	63	89
2001.....	6,496	5	7	7	9	9	11	14	23	32	49	81
2000.....	5,803	2	10	2	3	9	21	13	13	31	57	62
1999.....	5,640	1	14	4	6	14	12	15	23	46	59	64

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
Black, both sexes												
2003.....	920	2	13	1	2	7	10	13	14	31	45	79
2002.....	838	3	8	3	3	8	9	8	16	31	46	65
2001.....	854	6	3	5	2	4	9	10	18	33	58	64
2000.....	760	2	13	4	3	6	7	5	18	29	45	47
1999.....	878	2	11	6	4	8	13	13	30	27	48	73
Black, male												
2003.....	537	1	9	1	2	5	5	11	9	23	36	59
2002.....	532	2	7	2	3	6	9	7	14	25	35	49
2001.....	547	4	3	2	2	4	5	7	14	27	45	56
2000.....	485	2	10	4	1	4	7	3	16	23	34	37
1999.....	564	1	8	4	3	6	10	11	24	24	43	59
Black, female												
2003.....	383	1	4	-	-	2	5	2	5	8	9	20
2002.....	306	1	1	1	-	2	-	1	2	6	11	16
2001.....	307	2	-	3	-	-	4	3	4	6	13	8
2000.....	275	-	3	-	2	2	-	2	2	6	11	10
1999.....	314	1	3	2	1	2	3	2	6	3	5	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Falls (W00-W19)

American Indian 1/, both sexes

2003.....	120	-	-	-	-	2	2	2	3	8	8	6
2002.....	97	1	-	1	1	1	3	2	4	3	10	9
2001.....	86	-	-	-	1	2	1	4	-	5	4	14
2000.....	81	1	-	-	-	1	-	6	3	7	6	5
1999.....	75	-	1	-	-	2	4	4	2	3	4	5

American Indian 1/, male

2003.....	83	-	-	-	-	2	2	2	3	8	5	3
2002.....	54	1	-	-	-	1	3	2	3	2	8	6
2001.....	51	-	-	-	1	2	1	3	-	5	2	11
2000.....	57	1	-	-	-	1	-	5	3	5	5	1
1999.....	42	-	-	-	-	2	3	3	1	3	4	4

American Indian 1/, female

2003.....	37	-	-	-	-	-	-	-	-	-	3	3
2002.....	43	-	-	1	1	-	-	-	1	1	2	3
2001.....	35	-	-	-	-	-	-	1	-	-	2	3
2000.....	24	-	-	-	-	-	-	1	-	2	1	4
1999.....	33	-	1	-	-	-	1	1	1	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
Asian or Pacific Islander 2/, both sexes												
2003.....	334	2	3	-	3	2	2	2	2	5	8	10
2002.....	295	-	6	-	1	2	5	1	5	5	9	3
2001.....	252	2	1	1	1	2	1	6	5	7	3	13
2000.....	229	-	-	-	-	2	4	2	4	8	9	5
1999.....	217	2	-	1	-	2	1	4	6	7	9	7
Asian or Pacific Islander 2/, male												
2003.....	191	2	1	-	2	1	2	2	1	4	5	7
2002.....	178	-	4	-	-	2	4	1	3	4	8	2
2001.....	160	1	1	-	1	2	1	6	5	6	3	11
2000.....	131	-	-	-	-	2	2	2	4	8	5	4
1999.....	151	-	-	1	-	1	1	3	6	5	7	5
Asian or Pacific Islander 2/, female												
2003.....	143	-	2	-	1	1	-	-	1	1	3	3
2002.....	117	-	2	-	1	-	1	-	2	1	1	1
2001.....	92	1	-	1	-	-	-	-	-	1	-	2
2000.....	98	-	-	-	-	-	2	-	-	-	4	1
1999.....	66	2	-	-	-	1	-	1	-	2	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
All races, both sexes												
2003.....	730	-	7	13	36	95	105	74	46	57	61	48
2002.....	762	1	11	14	34	107	103	75	68	52	71	52
2001.....	802	-	15	18	39	110	96	71	51	70	76	61
2000.....	776	1	18	18	49	107	95	74	57	75	78	38
1999.....	824	-	12	19	57	126	125	76	67	67	69	51
All races, male												
2003.....	656	-	7	11	35	88	94	69	41	50	55	43
2002.....	667	1	7	10	29	101	94	70	59	42	56	45
2001.....	690	-	12	14	35	102	91	63	40	57	64	51
2000.....	671	-	11	16	46	97	86	65	44	60	65	31
1999.....	707	-	7	13	47	115	114	62	60	56	56	46
All races, female												
2003.....	74	-	-	2	1	7	11	5	5	7	6	5
2002.....	95	-	4	4	5	6	9	5	9	10	15	7
2001.....	112	-	3	4	4	8	5	8	11	13	12	10
2000.....	105	1	7	2	3	10	9	9	13	15	13	7
1999.....	117	-	5	6	10	11	11	14	7	11	13	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental discharge of firearms (W32-W34)

White, both sexes

2003.....	586	-	4	10	28	72	75	47	36	49	55	39
2002.....	591	1	9	10	26	69	73	51	49	39	63	44
2001.....	675	-	10	14	31	79	73	55	48	64	63	57
2000.....	611	1	11	11	41	76	65	52	42	60	68	30
1999.....	649	-	5	9	48	88	86	62	53	56	60	45

White, male

2003.....	525	-	4	8	27	66	66	43	32	44	51	34
2002.....	510	1	5	6	22	64	67	46	42	32	48	38
2001.....	580	-	8	11	30	72	69	48	39	52	55	47
2000.....	530	-	6	10	38	70	59	44	31	49	58	26
1999.....	554	-	3	4	42	80	78	49	47	46	48	41

White, female

2003.....	61	-	-	2	1	6	9	4	4	5	4	5
2002.....	81	-	4	4	4	5	6	5	7	7	15	6
2001.....	95	-	2	3	1	7	4	7	9	12	8	10
2000.....	81	1	5	1	3	6	6	8	11	11	10	4
1999.....	95	-	2	5	6	8	8	13	6	10	12	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
Black, both sexes												
2003.....	128	-	3	3	6	22	28	26	8	6	6	4
2002.....	155	-	2	3	6	36	25	23	17	13	7	8
2001.....	112	-	5	2	6	26	23	14	3	4	12	4
2000.....	137	-	5	5	7	28	24	18	14	13	6	7
1999.....	156	-	7	9	8	36	31	12	11	9	9	6
Black, male												
2003.....	116	-	3	3	6	21	26	25	8	4	4	4
2002.....	142	-	2	3	5	35	23	23	15	10	7	7
2001.....	99	-	4	1	5	25	22	13	1	4	9	4
2000.....	117	-	3	5	7	25	21	17	12	9	5	4
1999.....	135	-	4	8	4	33	28	12	10	8	8	5
Black, female												
2003.....	12	-	-	-	-	1	2	1	-	2	2	-
2002.....	13	-	-	-	1	1	2	-	2	3	-	1
2001.....	13	-	1	1	1	1	1	1	2	-	3	-
2000.....	20	-	2	-	-	3	3	1	2	4	1	3
1999.....	21	-	3	1	4	3	3	-	1	1	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental discharge of firearms (W32-W34)

American Indian 1/, both sexes

2003.....	13	-	-	-	2	1	2	1	1	2	-	3
2002.....	9	-	-	1	1	2	1	1	1	-	-	-
2001.....	9	-	-	2	1	3	-	1	-	1	1	-
2000.....	14	-	2	1	1	3	3	1	-	-	1	1
1999.....	14	-	-	-	1	1	8	1	2	1	-	-

American Indian 1/, male

2003.....	12	-	-	-	2	1	2	1	-	2	-	3
2002.....	9	-	-	1	1	2	1	1	1	-	-	-
2001.....	6	-	-	2	-	3	-	1	-	-	-	-
2000.....	12	-	2	1	1	2	3	1	-	-	-	1
1999.....	13	-	-	-	1	1	8	-	2	1	-	-

American Indian 1/, female

2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	1	-	-	-	-	1	1	-
2000.....	2	-	-	-	-	1	-	-	-	-	1	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
All races, both sexes												
2003.....	3,306	58	456	126	142	280	287	191	165	217	245	211
2002.....	3,447	63	454	159	162	320	309	227	206	263	263	205
2001.....	3,281	68	458	168	165	322	274	212	162	221	241	214
2000.....	3,482	75	493	201	174	371	275	213	206	250	230	182
1999.....	3,529	68	490	192	177	359	288	212	234	250	225	198
All races, male												
2003.....	2,632	34	301	90	109	262	263	177	146	186	206	173
2002.....	2,761	40	312	126	118	288	290	201	179	223	216	164
2001.....	2,560	30	291	120	117	295	249	186	142	181	195	182
2000.....	2,735	43	316	137	130	339	244	184	180	212	194	156
1999.....	2,794	39	310	139	136	330	262	193	204	215	187	166
All races, female												
2003.....	674	24	155	36	33	18	24	14	19	31	39	38
2002.....	686	23	142	33	44	32	19	26	27	40	47	41
2001.....	721	38	167	48	48	27	25	26	20	40	46	32
2000.....	747	32	177	64	44	32	31	29	26	38	36	26
1999.....	735	29	180	53	41	29	26	19	30	35	38	32

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
White, both sexes												
2003.....	2,622	44	386	85	75	201	218	144	125	177	195	173
2002.....	2,624	49	368	88	76	220	235	167	163	201	217	158
2001.....	2,515	52	397	97	78	207	217	150	122	173	188	170
2000.....	2,683	47	415	125	93	253	207	162	160	205	180	147
1999.....	2,715	46	416	106	80	247	215	151	179	205	188	175
White, male												
2003.....	2,067	24	252	61	56	188	201	134	108	148	166	138
2002.....	2,093	30	251	68	52	198	219	147	144	167	178	125
2001.....	1,936	25	257	61	54	191	193	128	104	138	154	145
2000.....	2,105	28	268	84	67	231	182	141	143	172	151	125
1999.....	2,120	26	262	71	61	226	193	135	158	172	155	143
White, female												
2003.....	555	20	134	24	19	13	17	10	17	29	29	35
2002.....	531	19	117	20	24	22	16	20	19	34	39	33
2001.....	579	27	140	36	24	16	24	22	18	35	34	25
2000.....	578	19	147	41	26	22	25	21	17	33	29	22
1999.....	595	20	154	35	19	21	22	16	21	33	33	32

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Black, both sexes												
2003.....	480	13	54	33	56	58	43	30	31	24	27	23
2002.....	613	11	63	58	78	72	57	42	33	43	31	33
2001.....	579	12	48	56	78	90	44	33	30	35	37	34
2000.....	593	26	56	67	69	86	48	34	24	36	43	23
1999.....	634	21	60	76	83	85	53	48	40	32	28	17
Black, male												
2003.....	407	9	39	23	44	55	40	27	29	23	25	22
2002.....	509	7	44	46	62	70	55	39	26	38	28	28
2001.....	478	3	28	47	56	83	44	33	29	31	29	27
2000.....	478	14	35	48	54	81	45	29	23	32	37	21
1999.....	530	12	40	60	65	78	51	45	35	30	25	17
Black, female												
2003.....	73	4	15	10	12	3	3	3	2	1	2	1
2002.....	104	4	19	12	16	2	2	3	7	5	3	5
2001.....	101	9	20	9	22	7	-	-	1	4	8	7
2000.....	115	12	21	19	15	5	3	5	1	4	6	2
1999.....	104	9	20	16	18	7	2	3	5	2	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
American Indian 1/, both sexes												
2003.....	62	1	6	4	4	4	6	1	1	10	11	8
2002.....	66	-	8	2	1	8	4	10	2	12	8	5
2001.....	52	2	6	6	3	3	2	8	5	4	5	3
2000.....	79	2	10	7	4	12	6	2	12	4	3	5
1999.....	68	-	7	2	3	6	5	5	10	7	6	4
American Indian 1/, male												
2003.....	51	1	4	4	3	3	6	1	1	9	7	8
2002.....	53	-	6	1	1	7	4	9	2	11	5	4
2001.....	46	1	3	6	3	3	2	8	5	3	4	3
2000.....	61	1	6	4	4	11	4	2	7	4	3	4
1999.....	55	-	4	2	1	6	5	5	6	7	5	4
American Indian 1/, female												
2003.....	11	-	2	-	1	1	-	-	-	1	4	-
2002.....	13	-	2	1	-	1	-	1	-	1	3	1
2001.....	6	1	3	-	-	-	-	-	-	1	1	-
2000.....	18	1	4	3	-	1	2	-	5	-	-	1
1999.....	13	-	3	-	2	-	-	-	4	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental drowning and submersion (W65-W74)

Asian or Pacific Islander 2/, both sexes

2003.....	142	-	10	4	7	17	20	16	8	6	12	7
2002.....	144	3	15	11	7	20	13	8	8	7	7	9
2001.....	135	2	7	9	6	22	11	21	5	9	11	7
2000.....	127	-	12	2	8	20	14	15	10	5	4	7
1999.....	112	1	7	8	11	21	15	8	5	6	3	2

Asian or Pacific Islander 2/, male

2003.....	107	-	6	2	6	16	16	15	8	6	8	5
2002.....	106	3	11	11	3	13	12	6	7	7	5	7
2001.....	100	1	3	6	4	18	10	17	4	9	8	7
2000.....	91	-	7	1	5	16	13	12	7	4	3	6
1999.....	89	1	4	6	9	20	13	8	5	6	2	2

Asian or Pacific Islander 2/, female

2003.....	35	-	4	2	1	1	4	1	-	-	4	2
2002.....	38	-	4	-	4	7	1	2	1	-	2	2
2001.....	35	1	4	3	2	4	1	4	1	-	3	-
2000.....	36	-	5	1	3	4	1	3	3	1	1	1
1999.....	23	-	3	2	2	1	2	-	-	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
All races, both sexes												
2003.....	3,369	30	224	137	78	75	124	118	134	167	228	270
2002.....	3,159	36	221	152	101	86	107	112	154	147	214	213
2001.....	3,309	50	225	164	88	76	133	114	133	190	251	213
2000.....	3,377	37	290	182	84	79	113	120	121	188	214	214
1999.....	3,348	41	302	170	92	88	113	142	140	200	221	198
All races, male												
2003.....	1,948	8	131	72	40	43	78	70	84	115	142	181
2002.....	1,935	21	126	89	62	61	71	75	95	101	150	146
2001.....	1,998	30	142	104	48	45	92	73	88	132	176	151
2000.....	2,025	20	174	109	36	44	77	78	79	133	135	145
1999.....	2,025	21	173	95	52	58	75	93	104	138	162	144
All races, female												
2003.....	1,421	22	93	65	38	32	46	48	50	52	86	89
2002.....	1,224	15	95	63	39	25	36	37	59	46	64	67
2001.....	1,311	20	83	60	40	31	41	41	45	58	75	62
2000.....	1,352	17	116	73	48	35	36	42	42	55	79	69
1999.....	1,323	20	129	75	40	30	38	49	36	62	59	54

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
White, both sexes												
2003.....	2,526	22	144	74	41	53	103	88	107	137	180	207
2002.....	2,287	19	127	78	61	61	77	72	118	110	175	167
2001.....	2,387	34	142	102	57	55	102	71	103	139	187	154
2000.....	2,376	23	160	104	54	60	81	83	86	134	157	155
1999.....	2,380	26	178	96	63	70	83	107	108	141	169	145
White, male												
2003.....	1,480	7	85	39	19	33	67	52	68	96	118	138
2002.....	1,419	14	70	48	42	44	53	52	74	75	119	117
2001.....	1,436	19	95	62	30	33	73	52	68	94	134	107
2000.....	1,435	11	98	61	24	32	58	54	59	97	98	106
1999.....	1,466	14	103	52	36	47	56	74	82	97	126	106
White, female												
2003.....	1,046	15	59	35	22	20	36	36	39	41	62	69
2002.....	868	5	57	30	19	17	24	20	44	35	56	50
2001.....	951	15	47	40	27	22	29	19	35	45	53	47
2000.....	941	12	62	43	30	28	23	29	27	37	59	49
1999.....	914	12	75	44	27	23	27	33	26	44	43	39

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Black, both sexes												
2003.....	775	7	76	60	34	20	19	25	22	28	40	61
2002.....	774	15	88	64	32	21	27	31	30	29	31	40
2001.....	846	15	82	60	28	14	28	38	28	45	59	56
2000.....	930	14	121	74	27	16	28	31	34	44	55	52
1999.....	892	15	111	74	28	16	24	29	24	55	47	46
Black, male												
2003.....	433	1	42	31	19	9	10	16	12	18	21	41
2002.....	461	7	53	37	17	14	17	17	18	21	24	25
2001.....	511	10	46	40	16	8	16	17	18	34	39	42
2000.....	551	9	71	46	11	10	17	19	20	30	35	35
1999.....	510	7	63	43	15	9	15	15	15	39	32	32
Black, female												
2003.....	342	6	34	29	15	11	9	9	10	10	19	20
2002.....	313	8	35	27	15	7	10	14	12	8	7	15
2001.....	335	5	36	20	12	6	12	21	10	11	20	14
2000.....	379	5	50	28	16	6	11	12	14	14	20	17
1999.....	382	8	48	31	13	7	9	14	9	16	15	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
American Indian 1/, both sexes												
2003.....	33	-	4	1	2	1	1	3	2	1	6	-
2002.....	56	1	3	3	5	1	2	8	3	5	6	5
2001.....	45	1	1	1	3	4	1	3	1	6	3	1
2000.....	37	-	5	2	2	3	1	6	1	5	2	1
1999.....	40	-	5	-	-	1	3	1	6	3	1	4
American Indian 1/, male												
2003.....	16	-	4	1	1	-	1	1	1	1	3	-
2002.....	35	-	1	-	3	-	1	5	3	3	5	3
2001.....	30	1	1	1	2	2	1	2	1	4	1	1
2000.....	23	-	3	1	1	2	1	5	-	2	2	1
1999.....	26	-	3	-	-	1	3	-	5	2	-	3
American Indian 1/, female												
2003.....	17	-	-	-	1	1	-	2	1	-	3	-
2002.....	21	1	2	3	2	1	1	3	-	2	1	2
2001.....	15	-	-	-	1	2	-	1	-	2	2	-
2000.....	14	-	2	1	1	1	-	1	1	3	-	-
1999.....	14	-	2	-	-	-	-	1	1	1	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Asian or Pacific Islander 2/, both sexes												
2003.....	35	1	-	2	1	1	1	2	3	1	2	2
2002.....	42	1	3	7	3	3	1	1	3	3	2	1
2001.....	31	-	-	1	-	3	2	2	1	-	2	2
2000.....	34	-	4	2	1	-	3	-	-	5	-	6
1999.....	36	-	8	-	1	1	3	5	2	1	4	3
Asian or Pacific Islander 2/, male												
2003.....	19	-	-	1	1	1	-	1	3	-	-	2
2002.....	20	-	2	4	-	3	-	1	-	2	2	1
2001.....	21	-	-	1	-	2	2	2	1	-	2	1
2000.....	16	-	2	1	-	-	1	-	-	4	-	3
1999.....	23	-	4	-	1	1	1	4	2	-	4	3
Asian or Pacific Islander 2/, female												
2003.....	16	1	-	1	-	-	1	1	-	1	2	-
2002.....	22	1	1	3	3	-	1	-	3	1	-	-
2001.....	10	-	-	-	-	1	-	-	-	-	-	1
2000.....	18	-	2	1	1	-	2	-	-	1	-	3
1999.....	13	-	4	-	-	-	2	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
All races, both sexes												
2003.....	19,457	20	49	18	43	520	1,479	1,517	1,918	2,674	3,556	3,351
2002.....	17,550	26	31	15	28	486	1,193	1,310	1,806	2,597	3,410	2,946
2001.....	14,078	15	31	18	32	406	956	1,078	1,429	2,307	2,729	2,266
2000.....	12,757	14	32	17	28	351	809	980	1,400	2,159	2,504	1,992
1999.....	12,186	12	34	12	28	260	704	952	1,403	2,082	2,467	1,926
All races, male												
2003.....	13,176	9	27	4	27	398	1,164	1,145	1,369	1,804	2,362	2,230
2002.....	12,059	21	14	10	18	366	940	1,003	1,316	1,807	2,315	1,960
2001.....	9,885	8	21	7	16	311	769	832	1,030	1,626	1,896	1,610
2000.....	9,138	11	21	12	15	273	647	779	1,041	1,554	1,830	1,455
1999.....	8,887	11	17	6	19	195	559	757	1,071	1,523	1,826	1,456
All races, female												
2003.....	6,281	11	22	14	16	122	315	372	549	870	1,194	1,121
2002.....	5,491	5	17	5	10	120	253	307	490	790	1,095	986
2001.....	4,193	7	10	11	16	95	187	246	399	681	833	656
2000.....	3,619	3	11	5	13	78	162	201	359	605	674	537
1999.....	3,299	1	17	6	9	65	145	195	332	559	641	470

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
White, both sexes												
2003.....	16,690	12	31	12	23	490	1,390	1,367	1,671	2,278	3,058	2,814
2002.....	14,868	13	19	12	25	452	1,103	1,165	1,580	2,199	2,901	2,407
2001.....	11,662	9	21	14	26	362	866	949	1,227	1,939	2,258	1,801
2000.....	10,508	10	19	10	20	318	732	847	1,184	1,811	2,031	1,568
1999.....	9,883	10	20	4	23	231	612	816	1,149	1,734	1,979	1,493
White, male												
2003.....	11,333	8	16	3	17	379	1,093	1,042	1,206	1,544	2,050	1,870
2002.....	10,213	12	6	8	17	342	870	891	1,161	1,542	1,985	1,609
2001.....	8,232	5	15	6	13	286	694	731	907	1,384	1,600	1,274
2000.....	7,558	8	11	7	11	251	601	674	894	1,330	1,498	1,143
1999.....	7,240	9	9	1	16	173	489	656	895	1,289	1,488	1,125
White, female												
2003.....	5,357	4	15	9	6	111	297	325	465	734	1,008	944
2002.....	4,655	1	13	4	8	110	233	274	419	657	916	798
2001.....	3,430	4	6	8	13	76	172	218	320	555	658	527
2000.....	2,950	2	8	3	9	67	131	173	290	481	533	425
1999.....	2,643	1	11	3	7	58	123	160	254	445	491	368

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Black, both sexes												
2003.....	2,396	7	16	4	14	17	70	107	205	335	442	495
2002.....	2,365	12	11	3	3	19	69	116	190	345	456	494
2001.....	2,169	6	8	4	2	27	68	107	175	326	437	437
2000.....	2,028	4	12	6	3	21	58	115	188	317	429	389
1999.....	2,062	1	12	6	2	18	76	115	221	305	445	411
Black, male												
2003.....	1,609	1	9	-	7	10	54	73	135	224	280	327
2002.....	1,631	8	8	2	1	14	53	89	131	232	294	323
2001.....	1,490	3	5	1	1	15	54	85	108	213	274	320
2000.....	1,433	3	9	5	2	13	37	91	125	206	301	288
1999.....	1,494	1	7	4	2	15	62	87	150	207	312	315
Black, female												
2003.....	787	6	7	4	7	7	16	34	70	111	162	168
2002.....	734	4	3	1	2	5	16	27	59	113	162	171
2001.....	679	3	3	3	1	12	14	22	67	113	163	117
2000.....	595	1	3	1	1	8	21	24	63	111	128	101
1999.....	568	-	5	2	-	3	14	28	71	98	133	96

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
American Indian 1/, both sexes												
2003.....	231	-	-	-	3	10	10	23	26	42	43	26
2002.....	188	1	-	-	-	7	9	16	22	38	33	31
2001.....	149	-	1	-	3	9	9	11	20	28	21	21
2000.....	137	-	-	1	3	6	12	11	19	21	28	23
1999.....	149	1	2	2	2	7	8	13	24	28	28	17
American Indian 1/, male												
2003.....	141	-	-	-	2	7	9	14	15	26	22	20
2002.....	124	1	-	-	-	4	9	12	14	22	19	20
2001.....	90	-	-	-	1	4	9	8	8	18	13	12
2000.....	84	-	-	-	-	3	4	9	15	10	17	16
1999.....	91	1	1	1	-	4	6	8	18	17	12	12
American Indian 1/, female												
2003.....	90	-	-	-	1	3	1	9	11	16	21	6
2002.....	64	-	-	-	-	3	-	4	8	16	14	11
2001.....	59	-	1	-	2	5	-	3	12	10	8	9
2000.....	53	-	-	1	3	3	8	2	4	11	11	7
1999.....	58	-	1	1	2	3	2	5	6	11	16	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Asian or Pacific Islander 2/, both sexes

2003.....	140	1	2	2	3	3	9	20	16	19	13	16
2002.....	129	-	1	-	-	8	12	13	14	15	20	14
2001.....	98	-	1	-	1	8	13	11	7	14	13	7
2000.....	84	-	1	-	2	6	7	7	9	10	16	12
1999.....	92	-	-	-	1	4	8	8	9	15	15	5

Asian or Pacific Islander 2/, male

2003.....	93	-	2	1	1	2	8	16	13	10	10	13
2002.....	91	-	-	-	-	6	8	11	10	11	17	8
2001.....	73	-	1	-	1	6	12	8	7	11	9	4
2000.....	63	-	1	-	2	6	5	5	7	8	14	8
1999.....	62	-	-	-	1	3	2	6	8	10	14	4

Asian or Pacific Islander 2/, female

2003.....	47	1	-	1	2	1	1	4	3	9	3	3
2002.....	38	-	1	-	-	2	4	2	4	4	3	6
2001.....	25	-	-	-	-	2	1	3	-	3	4	3
2000.....	21	-	-	-	-	-	2	2	2	2	2	4
1999.....	30	-	-	-	-	1	6	2	1	5	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
All races, both sexes												
2003.....	17,115	675	278	94	121	224	421	390	423	546	785	821
2002.....	17,201	677	247	113	160	273	365	377	443	577	764	745
2001.....	17,760	669	275	113	142	251	392	347	448	648	738	816
2000.....	17,437	572	266	115	152	273	375	416	479	672	774	787
1999.....	18,388	523	293	144	172	291	371	421	466	622	830	755
All races, male												
2003.....	9,919	398	177	63	97	191	353	327	358	432	629	640
2002.....	9,839	389	156	71	114	225	320	311	381	468	613	609
2001.....	10,044	390	162	72	109	196	339	294	388	529	599	647
2000.....	9,960	327	162	72	120	223	337	362	405	577	628	646
1999.....	10,371	314	183	93	128	238	327	361	387	513	668	584
All races, female												
2003.....	7,196	277	101	31	24	33	68	63	65	114	156	181
2002.....	7,362	288	91	42	46	48	45	66	62	109	151	136
2001.....	7,716	279	113	41	33	55	53	53	60	119	139	169
2000.....	7,477	245	104	43	32	50	38	54	74	95	146	141
1999.....	8,017	209	110	51	44	53	44	60	79	109	162	171

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
White, both sexes												
2003.....	14,820	425	210	73	97	190	350	324	357	449	654	664
2002.....	14,855	405	175	83	126	230	318	329	376	472	642	616
2001.....	15,354	422	200	94	113	217	338	285	367	548	596	657
2000.....	15,074	362	194	88	124	221	314	346	402	559	636	623
1999.....	15,859	335	215	114	143	250	317	348	378	536	686	586
White, male												
2003.....	8,483	251	136	48	79	162	305	279	302	355	532	520
2002.....	8,415	245	109	55	86	194	281	275	325	381	533	519
2001.....	8,571	245	120	56	87	170	295	245	316	459	489	530
2000.....	8,437	213	122	57	98	184	283	302	347	485	519	508
1999.....	8,781	197	132	79	104	203	280	306	317	448	549	452
White, female												
2003.....	6,337	174	74	25	18	28	45	45	55	94	122	144
2002.....	6,440	160	66	28	40	36	37	54	51	91	109	97
2001.....	6,783	177	80	38	26	47	43	40	51	89	107	127
2000.....	6,637	149	72	31	26	37	31	44	55	74	117	115
1999.....	7,078	138	83	35	39	47	37	42	61	88	137	134

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Black, both sexes												
2003.....	1,881	230	52	17	15	22	60	53	45	78	108	125
2002.....	1,972	253	61	24	32	33	36	39	54	87	95	110
2001.....	1,996	223	63	13	25	24	38	48	54	75	113	134
2000.....	1,990	188	57	25	24	41	49	55	65	92	109	134
1999.....	2,205	169	63	23	24	31	44	60	73	66	129	151
Black, male												
2003.....	1,164	136	36	11	12	19	40	39	38	62	77	97
2002.....	1,181	134	42	12	27	24	29	29	45	71	58	76
2001.....	1,198	135	36	12	19	18	29	35	47	51	89	98
2000.....	1,268	100	32	14	20	29	44	45	50	76	86	111
1999.....	1,378	102	43	12	20	28	40	45	58	49	105	118
Black, female												
2003.....	717	94	16	6	3	3	20	14	7	16	31	28
2002.....	791	119	19	12	5	9	7	10	9	16	37	34
2001.....	798	88	27	1	6	6	9	13	7	24	24	36
2000.....	722	88	25	11	4	12	5	10	15	16	23	23
1999.....	827	67	20	11	4	3	4	15	15	17	24	33

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

American Indian 1/, both sexes

2003.....	199	12	5	3	3	9	4	9	14	12	16	17
2002.....	169	8	6	4	2	4	8	6	8	14	21	11
2001.....	188	18	5	4	1	3	9	2	14	17	19	13
2000.....	176	10	6	1	2	4	7	3	8	12	21	20
1999.....	165	8	7	5	1	6	5	8	11	11	12	14

American Indian 1/, male

2003.....	145	8	3	3	1	7	3	5	12	10	15	14
2002.....	125	5	3	3	1	3	8	6	6	13	17	9
2001.....	127	7	3	3	1	3	9	2	14	14	12	10
2000.....	126	7	4	1	1	3	6	3	5	10	16	19
1999.....	116	5	4	1	1	4	3	6	9	9	11	12

American Indian 1/, female

2003.....	54	4	2	-	2	2	1	4	2	2	1	3
2002.....	44	3	3	1	1	1	-	-	2	1	4	2
2001.....	61	11	2	1	-	-	-	-	-	3	7	3
2000.....	50	3	2	-	1	1	1	-	3	2	5	1
1999.....	49	3	3	4	-	2	2	2	2	2	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Asian or Pacific Islander 2/, both sexes

2003.....	215	8	11	1	6	3	7	4	7	7	7	15
2002.....	205	11	5	2	-	6	3	3	5	4	6	8
2001.....	222	6	7	2	3	7	7	12	13	8	10	12
2000.....	197	12	9	1	2	7	5	12	4	9	8	10
1999.....	159	11	8	2	4	4	5	5	4	9	3	4

Asian or Pacific Islander 2/, male

2003.....	127	3	2	1	5	3	5	4	6	5	5	9
2002.....	118	5	2	1	-	4	2	1	5	3	5	5
2001.....	148	3	3	1	2	5	6	12	11	5	9	9
2000.....	129	7	4	-	1	7	4	12	3	6	7	8
1999.....	96	10	4	1	3	3	4	4	3	7	3	2

Asian or Pacific Islander 2/, female

2003.....	88	5	9	-	1	-	2	-	1	2	2	6
2002.....	87	6	3	1	-	2	1	2	-	1	1	3
2001.....	74	3	4	1	1	2	1	-	2	3	1	3
2000.....	68	5	5	1	1	-	1	-	1	3	1	2
1999.....	63	1	4	1	1	1	1	1	1	2	-	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races, both sexes

2003.....	31,484	6	244	1,487	2,501	2,276	2,789	2,958	3,644	3,485
2002.....	31,655	4	260	1,513	2,497	2,423	2,623	3,141	3,710	3,473
2001 3/.....	30,622	7	272	1,611	2,360	2,389	2,681	3,176	3,459	3,260
2000.....	29,350	7	300	1,621	2,373	2,341	2,451	3,112	3,450	2,987
1999.....	29,199	2	242	1,615	2,286	2,398	2,708	3,286	3,180	2,817

All races, male

2003.....	25,203	6	188	1,222	2,159	1,901	2,255	2,347	2,791	2,616
2002.....	25,409	3	196	1,280	2,152	2,003	2,132	2,484	2,821	2,608
2001 3/.....	24,672	7	207	1,345	2,064	2,015	2,184	2,518	2,662	2,484
2000.....	23,618	6	238	1,351	2,073	1,956	1,982	2,457	2,657	2,307
1999.....	23,458	2	192	1,347	1,979	2,005	2,189	2,590	2,429	2,152

All races, female

2003.....	6,281	-	56	265	342	375	534	611	853	869
2002.....	6,246	1	64	233	345	420	491	657	889	865
2001 3/.....	5,950	-	65	266	296	374	497	658	797	776
2000.....	5,732	1	62	270	300	385	469	655	793	680
1999.....	5,741	-	50	268	307	393	519	696	751	665

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, both sexes

2003.....	28,485	5	188	1,274	2,044	1,895	2,410	2,629	3,354	3,254
2002.....	28,731	3	211	1,308	2,098	2,021	2,277	2,799	3,420	3,245
2001 3/.....	27,710	5	213	1,379	1,955	2,012	2,344	2,849	3,168	3,030
2000.....	26,475	6	248	1,354	1,961	1,941	2,143	2,796	3,146	2,762
1999.....	26,300	2	204	1,344	1,875	1,996	2,350	2,933	2,927	2,599

White, male

2003.....	22,830	5	147	1,047	1,781	1,580	1,950	2,080	2,571	2,447
2002.....	23,049	2	157	1,109	1,810	1,651	1,844	2,205	2,608	2,423
2001 3/.....	22,328	5	163	1,156	1,704	1,699	1,905	2,248	2,443	2,325
2000.....	21,293	5	195	1,130	1,723	1,631	1,732	2,197	2,423	2,128
1999.....	21,107	2	162	1,122	1,629	1,667	1,900	2,291	2,244	1,987

White, female

2003.....	5,655	-	41	227	263	315	460	549	783	807
2002.....	5,682	1	54	199	288	370	433	594	812	822
2001 3/.....	5,382	-	50	223	251	313	439	601	725	705
2000.....	5,182	1	53	224	238	310	411	599	723	634
1999.....	5,193	-	42	222	246	329	450	642	683	612

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Black, both sexes												
2003.....	1,955	1	43	121	326	255	254	212	204	141
2002.....	1,939	1	34	128	275	273	231	235	207	150
2001 3/.....	1,957	2	39	136	297	254	238	230	195	159
2000.....	1,962	-	40	172	308	266	210	214	224	156
1999.....	1,950	-	29	179	291	261	242	248	187	146
Black, male												
2003.....	1,597	1	34	107	278	224	208	173	156	109
2002.....	1,633	1	28	111	240	246	201	197	157	130
2001 3/.....	1,627	2	30	116	279	223	202	191	145	115
2000.....	1,636	-	35	150	265	224	178	180	176	129
1999.....	1,651	-	22	155	259	231	205	217	139	116
Black, female												
2003.....	358	-	9	14	48	31	46	39	48	32
2002.....	306	-	6	17	35	27	30	38	50	20
2001 3/.....	330	-	9	20	18	31	36	39	50	44
2000.....	326	-	5	22	43	42	32	34	48	27
1999.....	299	-	7	24	32	30	37	31	48	30

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

American Indian 1/, both sexes

2003.....	322	-	7	50	53	44	42	42	35	15
2002.....	324	-	6	42	58	50	34	44	24	25
2001 3/.....	321	-	12	52	34	40	36	44	36	19
2000.....	297	1	6	44	45	43	35	37	30	16
1999.....	291	-	2	52	43	53	38	35	14	20

American Indian 1/, male

2003.....	265	-	6	37	43	38	38	36	30	11
2002.....	258	-	4	34	46	45	29	35	17	15
2001 3/.....	259	-	7	41	28	35	28	37	28	14
2000.....	237	1	5	34	37	33	26	31	22	11
1999.....	233	-	2	43	36	42	30	29	9	17

American Indian 1/, female

2003.....	57	-	1	13	10	6	4	6	5	4
2002.....	66	-	2	8	12	5	5	9	7	10
2001 3/.....	62	-	5	11	6	5	8	7	8	5
2000.....	60	-	1	10	8	10	9	6	8	5
1999.....	58	-	-	9	7	11	8	6	5	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Asian or Pacific Islander 2/, both sexes

2003.....	722	-	6	42	78	82	83	75	51	75
2002.....	661	-	9	35	66	79	81	63	59	53
2001 3/.....	634	-	8	44	74	83	63	53	60	52
2000.....	616	-	6	51	59	91	63	65	50	53
1999.....	658	-	7	40	77	88	78	70	52	52

Asian or Pacific Islander 2/, male

2003.....	511	-	1	31	57	59	59	58	34	49
2002.....	469	-	7	26	56	61	58	47	39	40
2001 3/.....	458	-	7	32	53	58	49	42	46	30
2000.....	452	-	3	37	48	68	46	49	36	39
1999.....	467	-	6	27	55	65	54	53	37	32

Asian or Pacific Islander 2/, female

2003.....	211	-	5	11	21	23	24	17	17	26
2002.....	192	-	2	9	10	18	23	16	20	13
2001 3/.....	176	-	1	12	21	25	14	11	14	22
2000.....	164	-	3	14	11	23	17	16	14	14
1999.....	191	-	1	13	22	23	24	17	15	20

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes

2003.....	16,907	1	73	736	1,339	1,114	1,267	1,280	1,647	1,705
2002.....	17,108	-	86	742	1,346	1,217	1,182	1,450	1,707	1,652
2001.....	16,869	-	90	838	1,292	1,263	1,301	1,429	1,601	1,552
2000.....	16,586	-	110	897	1,370	1,223	1,230	1,521	1,670	1,491
1999.....	16,599	-	103	975	1,340	1,292	1,329	1,617	1,480	1,462

All races, male

2003.....	14,827	1	62	659	1,220	990	1,109	1,092	1,378	1,419
2002.....	15,045	-	68	668	1,229	1,068	1,010	1,234	1,438	1,394
2001.....	14,758	-	69	743	1,192	1,122	1,124	1,200	1,351	1,309
2000.....	14,454	-	90	793	1,250	1,084	1,039	1,255	1,402	1,263
1999.....	14,479	-	80	867	1,216	1,137	1,133	1,349	1,237	1,221

All races, female

2003.....	2,080	-	11	77	119	124	158	188	269	286
2002.....	2,063	-	18	74	117	149	172	216	269	258
2001.....	2,111	-	21	95	100	141	177	229	250	243
2000.....	2,132	-	20	104	120	139	191	266	268	228
1999.....	2,120	-	23	108	124	155	196	268	243	241

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2003.....	15,587	1	64	638	1,090	927	1,091	1,166	1,550	1,627
2002.....	15,733	-	73	632	1,115	1,011	1,040	1,303	1,597	1,560
2001.....	15,467	-	73	714	1,039	1,068	1,152	1,296	1,474	1,467
2000.....	15,174	-	103	741	1,137	1,012	1,086	1,384	1,549	1,390
1999.....	15,135	-	94	817	1,082	1,081	1,172	1,452	1,381	1,360

White, male

2003.....	13,649	1	55	569	997	815	950	985	1,299	1,351
2002.....	13,809	-	58	570	1,016	869	888	1,104	1,347	1,313
2001.....	13,521	-	58	628	953	946	990	1,082	1,249	1,243
2000.....	13,214	-	86	653	1,040	893	922	1,133	1,298	1,178
1999.....	13,167	-	72	724	983	945	993	1,200	1,149	1,139

White, female

2003.....	1,938	-	9	69	93	112	141	181	251	276
2002.....	1,924	-	15	62	99	142	152	199	250	247
2001.....	1,946	-	15	86	86	122	162	214	225	224
2000.....	1,960	-	17	88	97	119	164	251	251	212
1999.....	1,968	-	22	93	99	136	179	252	232	221

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Black, both sexes												
2003.....	1,004	-	7	67	202	148	133	80	71	50
2002.....	1,059	-	9	78	177	162	112	115	86	73
2001.....	1,091	-	13	88	208	153	114	109	91	64
2000.....	1,090	-	4	117	191	161	111	101	93	77
1999.....	1,125	-	5	120	207	156	116	125	74	77
Black, male												
2003.....	902	-	5	64	184	142	120	74	57	46
2002.....	962	-	8	69	165	157	102	101	71	67
2001.....	971	-	10	80	200	137	105	98	73	50
2000.....	965	-	4	109	172	145	95	89	80	66
1999.....	1,018	-	4	109	189	144	106	113	66	64
Black, female												
2003.....	102	-	2	3	18	6	13	6	14	4
2002.....	97	-	1	9	12	5	10	14	15	6
2001.....	120	-	3	8	8	16	9	11	18	14
2000.....	125	-	-	8	19	16	16	12	13	11
1999.....	107	-	1	11	18	12	10	12	8	13

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

American Indian 1/, both sexes

2003.....	127	-	1	21	18	19	16	13	13	8
2002.....	146	-	1	17	36	20	13	17	10	9
2001.....	132	-	2	22	18	12	17	14	18	9
2000.....	133	-	1	20	23	14	13	12	13	8
1999.....	136	-	1	25	24	24	13	16	6	9

American Indian 1/, male

2003.....	117	-	1	18	17	16	15	13	12	8
2002.....	128	-	-	16	30	20	10	16	9	6
2001.....	115	-	-	21	16	11	15	10	16	9
2000.....	109	-	-	17	20	12	8	10	10	4
1999.....	120	-	1	22	21	21	11	15	5	8

American Indian 1/, female

2003.....	10	-	-	3	1	3	1	-	1	-
2002.....	18	-	1	1	6	-	3	1	1	3
2001.....	17	-	2	1	2	1	2	4	2	-
2000.....	24	-	1	3	3	2	5	2	3	4
1999.....	16	-	-	3	3	3	2	1	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Asian or Pacific Islander 2/, both sexes

2003.....	189	-	1	10	29	20	27	21	13	20
2002.....	170	-	3	15	18	24	17	15	14	10
2001.....	179	-	2	14	27	30	18	10	18	12
2000.....	189	-	2	19	19	36	20	24	15	16
1999.....	203	-	3	13	27	31	28	24	19	16

Asian or Pacific Islander 2/, male

2003.....	159	-	1	8	22	17	24	20	10	14
2002.....	146	-	2	13	18	22	10	13	11	8
2001.....	151	-	1	14	23	28	14	10	13	7
2000.....	166	-	-	14	18	34	14	23	14	15
1999.....	174	-	3	12	23	27	23	21	17	10

Asian or Pacific Islander 2/, female

2003.....	30	-	-	2	7	3	3	1	3	6
2002.....	24	-	1	2	-	2	7	2	3	2
2001.....	28	-	1	-	4	2	4	-	5	5
2000.....	23	-	2	5	1	2	6	1	1	1
1999.....	29	-	-	1	4	4	5	3	2	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races, both sexes

2003.....	14,577	5	171	751	1,162	1,162	1,522	1,678	1,997	1,780
2002.....	14,547	4	174	771	1,151	1,206	1,441	1,691	2,003	1,821
2001 3/.....	13,753	7	182	773	1,068	1,126	1,380	1,747	1,858	1,708
2000.....	12,764	7	190	724	1,003	1,118	1,221	1,591	1,780	1,496
1999.....	12,600	2	139	640	946	1,106	1,379	1,669	1,700	1,355

All races, male

2003.....	10,376	5	126	563	939	911	1,146	1,255	1,413	1,197
2002.....	10,364	3	128	612	923	935	1,122	1,250	1,383	1,214
2001 3/.....	9,914	7	138	602	872	893	1,060	1,318	1,311	1,175
2000.....	9,164	6	148	558	823	872	943	1,202	1,255	1,044
1999.....	8,979	2	112	480	763	868	1,056	1,241	1,192	931

All races, female

2003.....	4,201	-	45	188	223	251	376	423	584	583
2002.....	4,183	1	46	159	228	271	319	441	620	607
2001 3/.....	3,839	-	44	171	196	233	320	429	547	533
2000.....	3,600	1	42	166	180	246	278	389	525	452
1999.....	3,621	-	27	160	183	238	323	428	508	424

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

White, both sexes

2003.....	12,898	4	124	636	954	968	1,319	1,463	1,804	1,627
2002.....	12,998	3	138	676	983	1,010	1,237	1,496	1,823	1,685
2001 3/.....	12,243	5	140	665	916	944	1,192	1,553	1,694	1,563
2000.....	11,301	6	145	613	824	929	1,057	1,412	1,597	1,372
1999.....	11,165	2	110	527	793	915	1,178	1,481	1,546	1,239

White, male

2003.....	9,181	4	92	478	784	765	1,000	1,095	1,272	1,096
2002.....	9,240	2	99	539	794	782	956	1,101	1,261	1,110
2001 3/.....	8,807	5	105	528	751	753	915	1,166	1,194	1,082
2000.....	8,079	5	109	477	683	738	810	1,064	1,125	950
1999.....	7,940	2	90	398	646	722	907	1,091	1,095	848

White, female

2003.....	3,717	-	32	158	170	203	319	368	532	531
2002.....	3,758	1	39	137	189	228	281	395	562	575
2001 3/.....	3,436	-	35	137	165	191	277	387	500	481
2000.....	3,222	1	36	136	141	191	247	348	472	422
1999.....	3,225	-	20	129	147	193	271	390	451	391

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Black, both sexes

2003.....	951	1	36	54	124	107	121	132	133	91
2002.....	880	1	25	50	98	111	119	120	121	77
2001 3/.....	866	2	26	48	89	101	124	121	104	95
2000.....	872	-	36	55	117	105	99	113	131	79
1999.....	825	-	24	59	84	105	126	123	113	69

Black, male

2003.....	695	1	29	43	94	82	88	99	99	63
2002.....	671	1	20	42	75	89	99	96	86	63
2001 3/.....	656	2	20	36	79	86	97	93	72	65
2000.....	671	-	31	41	93	79	83	91	96	63
1999.....	633	-	18	46	70	87	99	104	73	52

Black, female

2003.....	256	-	7	11	30	25	33	33	34	28
2002.....	209	-	5	8	23	22	20	24	35	14
2001 3/.....	210	-	6	12	10	15	27	28	32	30
2000.....	201	-	5	14	24	26	16	22	35	16
1999.....	192	-	6	13	14	18	27	19	40	17

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

American Indian 1/, both sexes

2003.....	195	-	6	29	35	25	26	29	22	7
2002.....	178	-	5	25	22	30	21	27	14	16
2001 3/.....	189	-	10	30	16	28	19	30	18	10
2000.....	164	1	5	24	22	29	22	25	17	8
1999.....	155	-	1	27	19	29	25	19	8	11

American Indian 1/, male

2003.....	148	-	5	19	26	22	23	23	18	3
2002.....	130	-	4	18	16	25	19	19	8	9
2001 3/.....	144	-	7	20	12	24	13	27	12	5
2000.....	128	1	5	17	17	21	18	21	12	7
1999.....	113	-	1	21	15	21	19	14	4	9

American Indian 1/, female

2003.....	47	-	1	10	9	3	3	6	4	4
2002.....	48	-	1	7	6	5	2	8	6	7
2001 3/.....	45	-	3	10	4	4	6	3	6	5
2000.....	36	-	-	7	5	8	4	4	5	1
1999.....	42	-	-	6	4	8	6	5	4	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Asian or Pacific Islander 2/, both sexes

2003.....	533	-	5	32	49	62	56	54	38	55
2002.....	491	-	6	20	48	55	64	48	45	43
2001 3/.....	455	-	6	30	47	53	45	43	42	40
2000.....	427	-	4	32	40	55	43	41	35	37
1999.....	455	-	4	27	50	57	50	46	33	36

Asian or Pacific Islander 2/, male

2003.....	352	-	-	23	35	42	35	38	24	35
2002.....	323	-	5	13	38	39	48	34	28	32
2001 3/.....	307	-	6	18	30	30	35	32	33	23
2000.....	286	-	3	23	30	34	32	26	22	24
1999.....	293	-	3	15	32	38	31	32	20	22

Asian or Pacific Islander 2/, female

2003.....	181	-	5	9	14	20	21	16	14	20
2002.....	168	-	1	7	10	16	16	14	17	11
2001 3/.....	148	-	-	12	17	23	10	11	9	17
2000.....	141	-	1	9	10	21	11	15	13	13
1999.....	162	-	1	12	18	19	19	14	13	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All races, both sexes												
2003.....	17,732	341	376	122	202	1,938	3,430	2,566	1,950	1,636	1,474	1,219
2002.....	17,638	303	423	140	216	1,892	3,327	2,514	1,975	1,734	1,505	1,155
2001 3/.....	20,308	332	415	137	189	1,899	3,398	2,817	2,387	2,306	1,962	1,463
2000.....	16,765	349	356	140	231	1,914	3,025	2,312	1,852	1,732	1,487	1,059
1999.....	16,889	331	376	186	246	2,093	2,905	2,362	1,869	1,751	1,455	998
All races, male												
2003.....	13,882	205	202	70	145	1,677	2,947	2,191	1,550	1,224	1,084	893
2002.....	13,640	164	234	64	133	1,604	2,870	2,121	1,548	1,288	1,098	841
2001 3/.....	15,555	197	238	75	118	1,632	2,909	2,308	1,849	1,698	1,424	1,085
2000.....	12,820	203	198	77	150	1,613	2,590	1,918	1,443	1,235	1,084	773
1999.....	12,785	187	194	95	155	1,748	2,443	1,922	1,424	1,257	1,050	746
All races, female												
2003.....	3,850	136	174	52	57	261	483	375	400	412	390	326
2002.....	3,998	139	189	76	83	288	457	393	427	446	407	314
2001 3/.....	4,753	135	177	62	71	267	489	509	538	608	538	378
2000.....	3,945	146	158	63	81	301	435	394	409	497	403	286
1999.....	4,104	144	182	91	91	345	462	440	445	494	405	252

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2003.....	8,709	216	204	61	98	798	1,373	1,042	869	857	796	702
2002.....	8,685	170	239	83	112	830	1,307	1,015	850	891	848	664
2001 3/.....	11,328	196	228	102	116	809	1,455	1,329	1,294	1,406	1,266	958
2000.....	8,339	197	199	98	142	823	1,158	945	869	909	816	616
1999.....	8,628	200	208	106	155	884	1,180	1,006	920	946	820	587

White, male

2003.....	6,337	129	96	34	72	660	1,121	830	657	601	546	487
2002.....	6,282	100	133	35	63	681	1,065	812	623	624	616	446
2001 3/.....	8,254	118	132	52	67	652	1,159	1,032	956	1,058	914	700
2000.....	5,925	125	114	59	89	659	924	721	635	614	588	423
1999.....	6,162	121	108	55	93	704	918	784	661	659	581	429

White, female

2003.....	2,372	87	108	27	26	138	252	212	212	256	250	215
2002.....	2,403	70	106	48	49	149	242	203	227	267	232	218
2001 3/.....	3,074	78	96	50	49	157	296	297	338	348	352	258
2000.....	2,414	72	85	39	53	164	234	224	234	295	228	193
1999.....	2,466	79	100	51	62	180	262	222	259	287	239	158

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2003.....	8,392	106	156	54	95	1,068	1,938	1,462	1,013	721	624	466
2002.....	8,287	117	161	45	90	983	1,917	1,432	1,037	779	598	449
2001 3/.....	8,226	123	169	29	65	1,027	1,843	1,392	979	827	623	445
2000.....	7,867	140	145	40	84	1,028	1,791	1,301	923	764	612	397
1999.....	7,648	118	152	67	80	1,131	1,650	1,278	879	741	584	371

Black, male

2003.....	7,083	60	96	33	66	962	1,727	1,310	846	585	502	368
2002.....	6,896	56	92	24	62	859	1,723	1,262	867	616	442	365
2001 3/.....	6,780	70	101	20	48	928	1,672	1,211	817	590	461	343
2000.....	6,482	73	75	16	56	899	1,602	1,146	765	590	450	315
1999.....	6,214	60	75	36	55	978	1,475	1,084	716	556	440	289

Black, female

2003.....	1,309	46	60	21	29	106	211	152	167	136	122	98
2002.....	1,391	61	69	21	28	124	194	170	170	163	156	84
2001 3/.....	1,446	53	68	9	17	99	171	181	162	237	162	102
2000.....	1,385	67	70	24	28	129	189	155	158	174	162	82
1999.....	1,434	58	77	31	25	153	175	194	163	185	144	82

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

American Indian 1/, both sexes

2003.....	237	10	7	2	2	31	47	26	25	25	19	23
2002.....	267	10	12	4	7	32	38	29	34	27	25	15
2001 3/.....	211	6	12	1	3	27	32	25	29	17	19	13
2000.....	203	5	8	1	-	25	26	28	31	24	22	9
1999.....	253	5	6	5	4	27	33	34	39	25	19	15

American Indian 1/, male

2003.....	175	8	5	1	1	23	35	23	18	17	12	18
2002.....	185	4	5	-	3	25	29	21	25	20	20	13
2001 3/.....	146	3	5	-	1	21	24	22	17	11	13	9
2000.....	159	3	6	1	-	21	25	22	24	16	16	7
1999.....	179	3	4	1	3	22	23	22	28	18	12	11

American Indian 1/, female

2003.....	62	2	2	1	1	8	12	3	7	8	7	5
2002.....	82	6	7	4	4	7	9	8	9	7	5	2
2001 3/.....	65	3	7	1	2	6	8	3	12	6	6	4
2000.....	44	2	2	-	-	4	1	6	7	8	6	2
1999.....	74	2	2	4	1	5	10	12	11	7	7	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Asian or Pacific Islander 2/, both sexes

2003.....	394	9	9	5	7	41	72	36	43	33	35	28
2002.....	399	6	11	8	7	47	65	38	54	37	34	27
2001 3/.....	543	7	6	5	5	36	68	71	85	56	54	47
2000.....	356	7	4	1	5	38	50	38	29	35	37	37
1999.....	360	8	10	8	7	51	42	44	31	39	32	25

Asian or Pacific Islander 2/, male

2003.....	287	8	5	2	6	32	64	28	29	21	24	20
2002.....	277	4	4	5	5	39	53	26	33	28	20	17
2001 3/.....	375	6	-	3	2	31	54	43	59	39	36	33
2000.....	254	2	3	1	5	34	39	29	19	15	30	28
1999.....	230	3	7	3	4	44	27	32	19	24	17	17

Asian or Pacific Islander 2/, female

2003.....	107	1	4	3	1	9	8	8	14	12	11	8
2002.....	122	2	7	3	2	8	12	12	21	9	14	10
2001 3/.....	168	1	6	2	3	5	14	28	26	17	18	14
2000.....	102	5	1	-	-	4	11	9	10	20	7	9
1999.....	130	5	3	5	3	7	15	12	12	15	15	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All races, both sexes												
2003.....	11,920	8	40	48	139	1,587	2,823	2,070	1,470	1,066	875	683
2002.....	11,829	9	49	55	150	1,567	2,750	2,018	1,447	1,135	907	604
2001.....	11,348	11	55	59	121	1,525	2,675	1,940	1,368	1,142	836	552
2000.....	10,801	12	28	50	137	1,549	2,414	1,789	1,285	1,064	842	582
1999.....	10,828	8	50	61	163	1,708	2,330	1,795	1,253	1,054	814	551
All races, male												
2003.....	10,126	7	20	27	107	1,436	2,541	1,867	1,258	860	675	534
2002.....	9,899	3	30	26	101	1,384	2,467	1,790	1,227	918	705	445
2001.....	9,532	7	27	35	90	1,377	2,419	1,708	1,156	892	636	434
2000.....	9,006	9	17	28	110	1,381	2,167	1,574	1,076	842	645	445
1999.....	8,944	5	26	37	118	1,498	2,071	1,568	1,041	807	607	426
All races, female												
2003.....	1,794	1	20	21	32	151	282	203	212	206	200	149
2002.....	1,930	6	19	29	49	183	283	228	220	217	202	159
2001.....	1,816	4	28	24	31	148	256	232	212	250	200	118
2000.....	1,795	3	11	22	27	168	247	215	209	222	197	137
1999.....	1,884	3	24	24	45	210	259	227	212	247	207	125

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
White, both sexes												
2003.....	5,173	2	20	22	59	604	1,030	742	589	520	446	388
2002.....	5,185	3	33	35	71	649	951	738	563	533	494	343
2001.....	5,188	8	33	47	72	593	985	741	586	583	482	324
2000.....	4,806	5	17	39	75	617	817	645	547	525	460	338
1999.....	4,918	4	26	32	95	662	822	675	562	555	456	313
White, male												
2003.....	4,108	1	8	10	48	528	898	641	476	384	321	281
2002.....	4,050	1	20	18	46	556	824	621	449	399	374	228
2001.....	4,042	5	17	28	50	508	843	627	456	443	348	242
2000.....	3,686	2	9	22	59	531	693	528	427	384	340	246
1999.....	3,820	4	14	20	66	567	688	568	440	408	328	230
White, female												
2003.....	1,065	1	12	12	11	76	132	101	113	136	125	107
2002.....	1,135	2	13	17	25	93	127	117	114	134	120	115
2001.....	1,146	3	16	19	22	85	142	114	130	140	134	82
2000.....	1,120	3	8	17	16	86	124	117	120	141	120	92
1999.....	1,098	-	12	12	29	95	134	107	122	147	128	83

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Black, both sexes												
2003.....	6,397	6	19	23	72	932	1,715	1,286	839	515	400	269
2002.....	6,285	6	14	13	68	865	1,724	1,240	842	564	381	244
2001.....	5,885	2	18	12	45	902	1,635	1,155	746	540	333	207
2000.....	5,699	7	8	11	58	888	1,546	1,097	715	509	354	223
1999.....	5,577	2	18	23	59	991	1,459	1,068	656	468	336	222
Black, male												
2003.....	5,740	6	11	15	52	869	1,574	1,191	746	454	332	234
2002.....	5,575	2	9	6	48	781	1,583	1,139	746	487	310	206
2001.....	5,279	1	9	7	37	842	1,531	1,050	673	434	274	175
2000.....	5,084	7	5	6	47	812	1,431	1,009	628	439	284	183
1999.....	4,883	1	7	15	46	883	1,344	961	576	378	265	183
Black, female												
2003.....	657	-	8	8	20	63	141	95	93	61	68	35
2002.....	710	4	5	7	20	84	141	101	96	77	71	38
2001.....	606	1	9	5	8	60	104	105	73	106	59	32
2000.....	615	-	3	5	11	76	115	88	87	70	70	40
1999.....	694	1	11	8	13	108	115	107	80	90	71	39

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

American Indian 1/, both sexes

2003.....	109	-	-	2	2	20	29	13	12	7	10	9
2002.....	117	-	1	2	5	20	24	15	14	12	10	5
2001.....	87	1	4	-	2	12	17	13	14	4	8	4
2000.....	86	-	2	-	-	14	18	17	10	9	8	2
1999.....	104	2	2	2	4	16	17	16	14	7	4	4

American Indian 1/, male

2003.....	81	-	-	1	1	14	24	12	10	4	6	7
2002.....	85	-	-	-	2	15	17	11	11	10	8	3
2001.....	64	1	1	-	1	10	14	12	8	2	6	2
2000.....	70	-	2	-	-	12	17	13	9	7	4	2
1999.....	82	-	2	1	3	15	14	12	12	5	2	3

American Indian 1/, female

2003.....	28	-	-	1	1	6	5	1	2	3	4	2
2002.....	32	-	1	2	3	5	7	4	3	2	2	2
2001.....	23	-	3	-	1	2	3	1	6	2	2	2
2000.....	16	-	-	-	-	2	1	4	1	2	4	-
1999.....	22	2	-	1	1	1	3	4	2	2	2	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Asian or Pacific Islander 2/, both sexes

2003.....	241	-	1	1	6	31	49	29	30	24	19	17
2002.....	242	-	1	5	6	33	51	25	28	26	22	12
2001.....	188	-	-	-	2	18	38	31	22	15	13	17
2000.....	210	-	1	-	4	30	33	30	13	21	20	19
1999.....	229	-	4	4	5	39	32	36	21	24	18	12

Asian or Pacific Islander 2/, male

2003.....	197	-	1	1	6	25	45	23	26	18	16	12
2002.....	189	-	1	2	5	32	43	19	21	22	13	8
2001.....	147	-	-	-	2	17	31	19	19	13	8	15
2000.....	166	-	1	-	4	26	26	24	12	12	17	14
1999.....	159	-	3	1	3	33	25	27	13	16	12	10

Asian or Pacific Islander 2/, female

2003.....	44	-	-	-	-	6	4	6	4	6	3	5
2002.....	53	-	-	3	1	1	8	6	7	4	9	4
2001.....	41	-	-	-	-	1	7	12	3	2	5	2
2000.....	44	-	-	-	-	4	7	6	1	9	3	5
1999.....	70	-	1	3	2	6	7	9	8	8	6	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races, both sexes

2003.....	5,812	333	336	74	63	351	607	496	480	570	599	536
2002.....	5,809	294	374	85	66	325	577	496	528	599	598	551
2001 3/.....	8,960	321	360	78	68	374	723	877	1,019	1,164	1,126	911
2000.....	5,964	337	328	90	94	365	611	523	567	668	645	477
1999.....	6,061	323	326	125	83	385	575	567	616	697	641	447

All races, male

2003.....	3,756	198	182	43	38	241	406	324	292	364	409	359
2002.....	3,741	161	204	38	32	220	403	331	321	370	393	396
2001 3/.....	6,023	190	211	40	28	255	490	600	693	806	788	651
2000.....	3,814	194	181	49	40	232	423	344	367	393	439	328
1999.....	3,841	182	168	58	37	250	372	354	383	450	443	320

All races, female

2003.....	2,056	135	154	31	25	110	201	172	188	206	190	177
2002.....	2,068	133	170	47	34	105	174	165	207	229	205	155
2001 3/.....	2,937	131	149	38	40	119	233	277	326	358	338	260
2000.....	2,150	143	147	41	54	133	188	179	200	275	206	149
1999.....	2,220	141	158	67	46	135	203	213	233	247	198	127

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
White, both sexes												
2003.....	3,536	214	184	39	39	194	343	300	280	337	350	314
2002.....	3,500	167	206	48	41	181	356	277	287	358	354	321
2001 3/.....	6,140	188	195	55	44	216	470	588	708	823	784	634
2000.....	3,533	192	182	59	67	206	341	300	322	384	356	278
1999.....	3,710	196	182	74	60	222	358	331	358	391	364	274
White, male												
2003.....	2,229	128	88	24	24	132	223	189	181	217	225	206
2002.....	2,232	99	113	17	17	125	241	191	174	225	242	218
2001 3/.....	4,212	113	115	24	17	144	316	405	500	615	566	458
2000.....	2,239	123	105	37	30	128	231	193	208	230	248	177
1999.....	2,342	117	94	35	27	137	230	216	221	251	253	199
White, female												
2003.....	1,307	86	96	15	15	62	120	111	99	120	125	108
2002.....	1,268	68	93	31	24	56	115	86	113	133	112	103
2001 3/.....	1,928	75	80	31	27	72	154	183	208	208	218	176
2000.....	1,294	69	77	22	37	78	110	107	114	154	108	101
1999.....	1,368	79	88	39	33	85	128	115	137	140	111	75

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2003.....	1,995	100	137	31	23	136	223	176	174	206	224	197
2002.....	2,002	111	147	32	22	118	193	192	195	215	217	205
2001 3/.....	2,341	121	151	17	20	125	208	237	233	287	290	238
2000.....	2,168	133	137	29	26	140	245	204	208	255	258	174
1999.....	2,071	116	134	44	21	140	191	210	223	273	248	149

Black, male

2003.....	1,343	54	85	18	14	93	153	119	100	131	170	134
2002.....	1,321	54	83	18	14	78	140	123	121	129	132	159
2001 3/.....	1,501	69	92	13	11	86	141	161	144	156	187	168
2000.....	1,398	66	70	10	9	87	171	137	137	151	166	132
1999.....	1,331	59	68	21	9	95	131	123	140	178	175	106

Black, female

2003.....	652	46	52	13	9	43	70	57	74	75	54	63
2002.....	681	57	64	14	8	40	53	69	74	86	85	46
2001 3/.....	840	52	59	4	9	39	67	76	89	131	103	70
2000.....	770	67	67	19	17	53	74	67	71	104	92	42
1999.....	740	57	66	23	12	45	60	87	83	95	73	43

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

American Indian 1/, both sexes

2003.....	128	10	7	-	-	11	18	13	13	18	9	14
2002.....	150	10	11	2	2	12	14	14	20	15	15	10
2001 3/.....	124	5	8	1	1	15	15	12	15	13	11	9
2000.....	117	5	6	1	-	11	8	11	21	15	14	7
1999.....	149	3	4	3	-	11	16	18	25	18	15	11

American Indian 1/, male

2003.....	94	8	5	-	-	9	11	11	8	13	6	11
2002.....	100	4	5	-	1	10	12	10	14	10	12	10
2001 3/.....	82	2	4	-	-	11	10	10	9	9	7	7
2000.....	89	3	4	1	-	9	8	9	15	9	12	5
1999.....	97	3	2	-	-	7	9	10	16	13	10	8

American Indian 1/, female

2003.....	34	2	2	-	-	2	7	2	5	5	3	3
2002.....	50	6	6	2	1	2	2	4	6	5	3	-
2001 3/.....	42	3	4	1	1	4	5	2	6	4	4	2
2000.....	28	2	2	-	-	2	-	2	6	6	2	2
1999.....	52	-	2	3	-	4	7	8	9	5	5	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Asian or Pacific Islander 2/, both sexes

2003.....	153	9	8	4	1	10	23	7	13	9	16	11
2002.....	157	6	10	3	1	14	14	13	26	11	12	15
2001 3/.....	355	7	6	5	3	18	30	40	63	41	41	30
2000.....	146	7	3	1	1	8	17	8	16	14	17	18
1999.....	131	8	6	4	2	12	10	8	10	15	14	13

Asian or Pacific Islander 2/, male

2003.....	90	8	4	1	-	7	19	5	3	3	8	8
2002.....	88	4	3	3	-	7	10	7	12	6	7	9
2001 3/.....	228	6	-	3	-	14	23	24	40	26	28	18
2000.....	88	2	2	1	1	8	13	5	7	3	13	14
1999.....	71	3	4	2	1	11	2	5	6	8	5	7

Asian or Pacific Islander 2/, female

2003.....	63	1	4	3	1	3	4	2	10	6	8	3
2002.....	69	2	7	-	1	7	4	6	14	5	5	6
2001 3/.....	127	1	6	2	3	4	7	16	23	15	13	12
2000.....	58	5	1	-	-	-	4	3	9	11	4	4
1999.....	60	5	2	2	1	1	8	3	4	7	9	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

All races, both sexes

2003.....	423	-	-	-	2	20	73	63	72	66	51	29
2002.....	384	-	-	-	3	24	67	51	51	64	45	43
2001.....	396	-	-	-	-	26	62	47	60	65	50	33
2000.....	359	-	-	-	3	27	46	59	51	60	41	38
1999.....	398	-	-	-	1	20	47	66	70	72	49	19

All races, male

2003.....	405	-	-	-	1	19	70	61	70	65	49	27
2002.....	369	-	-	-	3	23	64	51	48	60	45	40
2001.....	380	-	-	-	-	25	59	46	58	63	45	33
2000.....	348	-	-	-	3	27	44	58	50	58	38	37
1999.....	392	-	-	-	1	20	47	66	69	69	48	19

All races, female

2003.....	18	-	-	-	1	1	3	2	2	1	2	2
2002.....	15	-	-	-	-	1	3	-	3	4	-	3
2001.....	16	-	-	-	-	1	3	1	2	2	5	-
2000.....	11	-	-	-	-	-	2	1	1	2	3	1
1999.....	6	-	-	-	-	-	-	-	1	3	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

White, both sexes

2003.....	284	-	-	-	1	11	43	30	46	52	38	22
2002.....	257	-	-	-	1	14	35	28	41	45	33	28
2001.....	292	-	-	-	-	15	45	35	43	46	40	23
2000.....	237	-	-	-	1	11	33	36	36	38	25	32
1999.....	255	-	-	-	-	9	22	39	47	47	33	15

White, male

2003.....	271	-	-	-	-	10	42	30	44	51	36	20
2002.....	246	-	-	-	1	13	33	28	39	43	33	25
2001.....	283	-	-	-	-	14	44	34	43	44	37	23
2000.....	231	-	-	-	1	11	31	35	35	37	25	32
1999.....	250	-	-	-	-	9	22	39	47	44	32	15

White, female

2003.....	13	-	-	-	1	1	1	-	2	1	2	2
2002.....	11	-	-	-	-	1	2	-	2	2	-	3
2001.....	9	-	-	-	-	1	1	1	-	2	3	-
2000.....	6	-	-	-	-	-	2	1	1	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	3	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

Black, both sexes

2003.....	120	-	-	-	1	8	27	29	24	12	10	6
2002.....	110	-	-	-	2	10	27	20	9	17	11	11
2001.....	84	-	-	-	-	10	16	9	12	15	8	9
2000.....	111	-	-	-	2	15	13	19	14	20	15	6
1999.....	126	-	-	-	1	10	22	24	18	21	15	4

Black, male

2003.....	117	-	-	-	1	8	25	28	24	12	10	6
2002.....	106	-	-	-	2	10	26	20	8	15	11	11
2001.....	79	-	-	-	-	10	14	9	11	15	6	9
2000.....	108	-	-	-	2	15	13	19	14	20	13	5
1999.....	125	-	-	-	1	10	22	24	17	21	15	4

Black, female

2003.....	3	-	-	-	-	-	2	1	-	-	-	-
2002.....	4	-	-	-	-	-	1	-	1	2	-	-
2001.....	5	-	-	-	-	-	2	-	1	-	2	-
2000.....	3	-	-	-	-	-	-	-	-	-	2	1
1999.....	1	-	-	-	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races, both sexes

2003.....	5,072	75	31	16	39	169	394	351	484	686	895	777
2002.....	4,830	101	57	26	42	143	344	352	476	727	789	709
2001.....	4,198	83	44	17	34	132	280	311	417	674	722	619
2000.....	3,819	65	39	18	42	120	250	306	373	599	687	543
1999.....	3,917	55	43	12	37	159	271	313	430	644	651	528

All races, male

2003.....	3,295	50	17	9	32	124	288	249	331	450	563	493
2002.....	3,114	52	28	11	27	108	250	256	329	494	487	432
2001.....	2,833	49	19	10	28	102	214	231	284	438	493	433
2000.....	2,635	27	26	13	37	93	192	223	266	415	470	374
1999.....	2,692	33	21	6	32	132	213	230	293	446	441	357

All races, female

2003.....	1,777	25	14	7	7	45	106	102	153	236	332	284
2002.....	1,716	49	29	15	15	35	94	96	147	233	302	277
2001.....	1,365	34	25	7	6	30	66	80	133	236	229	186
2000.....	1,184	38	13	5	5	27	58	83	107	184	217	169
1999.....	1,225	22	22	6	5	27	58	83	137	198	210	171

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	4,255	52	21	7	30	149	339	300	419	578	755	654
2002.....	3,962	56	45	16	33	120	298	291	388	599	653	581
2001.....	3,453	45	31	9	30	116	235	259	353	548	598	508
2000.....	3,053	41	27	9	29	103	199	252	301	492	539	432
1999.....	3,122	36	23	10	32	125	217	255	351	516	507	409

White, male

2003.....	2,709	30	11	3	26	108	243	209	280	380	468	405
2002.....	2,524	29	23	5	21	89	217	210	267	399	406	353
2001.....	2,283	28	13	4	25	89	174	190	241	347	398	345
2000.....	2,078	18	17	6	26	79	149	181	211	342	368	292
1999.....	2,105	22	13	5	28	106	172	180	236	349	349	262

White, female

2003.....	1,546	22	10	4	4	41	96	91	139	198	287	249
2002.....	1,438	27	22	11	12	31	81	81	121	200	247	228
2001.....	1,170	17	18	5	5	27	61	69	112	201	200	163
2000.....	975	23	10	3	3	24	50	71	90	150	171	140
1999.....	1,017	14	10	5	4	19	45	75	115	167	158	147

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2003.....	687	21	9	7	7	16	35	38	58	92	124	115
2002.....	759	37	8	9	6	17	40	54	71	118	126	116
2001.....	646	36	13	7	4	12	32	39	54	112	115	101
2000.....	668	15	11	9	12	13	39	45	60	98	136	103
1999.....	696	18	16	2	5	25	46	51	70	114	126	109

Black, male

2003.....	498	19	5	4	4	13	30	32	45	59	85	83
2002.....	521	18	3	5	6	14	29	41	49	89	75	74
2001.....	480	20	6	5	3	10	28	31	36	81	89	80
2000.....	488	3	8	7	10	13	34	35	45	67	95	77
1999.....	523	10	8	1	4	20	36	44	49	87	84	88

Black, female

2003.....	189	2	4	3	3	3	5	6	13	33	39	32
2002.....	238	19	5	4	-	3	11	13	22	29	51	42
2001.....	166	16	7	2	1	2	4	8	18	31	26	21
2000.....	180	12	3	2	2	-	5	10	15	31	41	26
1999.....	173	8	8	1	1	5	10	7	21	27	42	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

American Indian 1/, both sexes

2003.....	64	2	-	1	1	1	7	8	3	11	10	5
2002.....	48	3	2	1	1	4	1	2	10	3	5	7
2001.....	43	1	-	-	-	3	4	2	5	8	6	5
2000.....	46	4	1	-	1	1	4	3	6	6	6	4
1999.....	54	1	2	-	-	4	5	2	9	9	9	4

American Indian 1/, male

2003.....	47	1	-	1	1	-	7	4	3	9	7	3
2002.....	26	1	1	1	-	4	1	-	7	1	2	3
2001.....	32	-	-	-	-	3	4	2	5	6	3	3
2000.....	36	3	1	-	1	-	3	2	6	5	4	3
1999.....	37	1	-	-	-	3	4	2	8	6	4	3

American Indian 1/, female

2003.....	17	1	-	-	-	1	-	4	-	2	3	2
2002.....	22	2	1	-	1	-	-	2	3	2	3	4
2001.....	11	1	-	-	-	-	-	-	-	2	3	2
2000.....	10	1	-	-	-	1	1	1	-	1	2	1
1999.....	17	-	2	-	-	1	1	-	1	3	5	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 2/, both sexes

2003.....	66	-	1	1	1	3	13	5	4	5	6	3
2002.....	61	5	2	-	2	2	5	5	7	7	5	5
2001.....	56	1	-	1	-	1	9	11	5	6	3	5
2000.....	52	5	-	-	-	3	8	6	6	3	6	4
1999.....	45	-	2	-	-	5	3	5	-	5	9	6

Asian or Pacific Islander 2/, male

2003.....	41	-	1	1	1	3	8	4	3	2	3	2
2002.....	43	4	1	-	-	1	3	5	6	5	4	2
2001.....	38	1	-	1	-	-	8	8	2	4	3	5
2000.....	33	3	-	-	-	1	6	5	4	1	3	2
1999.....	27	-	-	-	-	3	1	4	-	4	4	4

Asian or Pacific Islander 2/, female

2003.....	25	-	-	-	-	-	5	1	1	3	3	1
2002.....	18	1	1	-	2	1	2	-	1	2	1	3
2001.....	18	-	-	-	-	1	1	3	3	2	-	-
2000.....	19	2	-	-	-	2	2	1	2	2	3	2
1999.....	18	-	2	-	-	2	2	1	-	1	5	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races, both sexes

2003.....	232	-	1	1	11	31	38	21	17	21	22	19
2002.....	243	-	1	2	4	35	44	27	19	31	19	19
2001.....	231	-	-	2	4	24	40	32	16	33	17	16
2000.....	230	-	-	2	8	26	45	32	20	18	21	9
1999.....	324	-	3	-	12	68	59	32	19	28	29	18

All races, male

2003.....	186	-	1	1	11	29	31	18	15	12	14	11
2002.....	199	-	1	2	3	34	37	23	16	22	17	14
2001.....	190	-	-	2	4	23	36	27	15	20	13	12
2000.....	190	-	-	2	7	25	37	26	17	12	18	6
1999.....	277	-	3	-	12	61	50	28	16	24	21	15

All races, female

2003.....	46	-	-	-	-	2	7	3	2	9	8	8
2002.....	44	-	-	-	1	1	7	4	3	9	2	5
2001.....	41	-	-	-	-	1	4	5	1	13	4	4
2000.....	40	-	-	-	1	1	8	6	3	6	3	3
1999.....	47	-	-	-	-	7	9	4	3	4	8	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Discharge of firearms, undetermined intent (Y22-Y24)												
White, both sexes												
2003.....	183	-	-	-	7	23	24	14	13	18	20	18
2002.....	192	-	-	-	4	24	35	20	13	25	16	18
2001.....	188	-	-	-	4	17	27	26	14	31	14	14
2000.....	167	-	-	-	5	20	28	23	11	13	18	8
1999.....	247	-	1	-	11	42	43	22	14	25	25	13
White, male												
2003.....	144	-	-	-	7	21	19	12	11	10	12	11
2002.....	153	-	-	-	3	23	28	16	11	18	14	13
2001.....	150	-	-	-	4	16	24	22	13	18	11	10
2000.....	137	-	-	-	5	19	23	18	10	8	17	5
1999.....	212	-	1	-	11	39	37	19	12	21	20	10
White, female												
2003.....	39	-	-	-	-	2	5	2	2	8	8	7
2002.....	39	-	-	-	1	1	7	4	2	7	2	5
2001.....	38	-	-	-	-	1	3	4	1	13	3	4
2000.....	30	-	-	-	-	1	5	5	1	5	1	3
1999.....	35	-	-	-	-	3	6	3	2	4	5	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Discharge of firearms, undetermined intent (Y22-Y24)

Black, both sexes

2003.....	35	-	1	1	3	7	10	4	3	2	1	1
2002.....	41	-	1	2	-	9	8	7	4	4	2	-
2001.....	32	-	-	2	-	7	8	4	1	-	2	2
2000.....	54	-	-	2	3	6	14	8	7	4	3	1
1999.....	66	-	2	-	1	20	15	10	4	3	3	4

Black, male

2003.....	32	-	1	1	3	7	9	4	3	1	1	-
2002.....	39	-	1	2	-	9	8	7	4	3	2	-
2001.....	29	-	-	2	-	7	7	3	1	-	1	2
2000.....	47	-	-	2	2	6	13	7	5	4	1	1
1999.....	56	-	2	-	1	17	12	9	3	3	1	4

Black, female

2003.....	3	-	-	-	-	-	1	-	-	1	-	1
2002.....	2	-	-	-	-	-	-	-	-	1	-	-
2001.....	3	-	-	-	-	-	1	1	-	-	1	-
2000.....	7	-	-	-	1	-	1	1	2	-	2	-
1999.....	10	-	-	-	-	3	3	1	1	-	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Discharge of firearms, undetermined intent (Y22-Y24)

Asian or Pacific Islander 2/, both sexes

2003.....	5	-	-	-	-	1	2	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	2	-	1
2001.....	7	-	-	-	-	-	3	2	-	1	1	-
2000.....	5	-	-	-	-	-	1	1	1	1	-	-
1999.....	5	-	-	-	-	3	-	-	-	-	-	1

Asian or Pacific Islander 2/, male

2003.....	2	-	-	-	-	1	1	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	-	1
2001.....	7	-	-	-	-	-	3	2	-	1	1	-
2000.....	3	-	-	-	-	-	-	1	1	-	-	-
1999.....	4	-	-	-	-	2	-	-	-	-	-	1

Asian or Pacific Islander 2/, female

2003.....	3	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	-	1	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races, both sexes

2003.....	4,840	75	30	15	28	138	356	330	467	665	873	758
2002.....	4,587	101	56	24	38	108	300	325	457	696	770	690
2001.....	3,967	83	44	15	30	108	240	279	401	641	705	603
2000.....	3,589	65	39	16	34	94	205	274	353	581	666	534
1999.....	3,593	55	40	12	25	91	212	281	411	616	622	510

All races, male

2003.....	3,109	50	16	8	21	95	257	231	316	438	549	482
2002.....	2,915	52	27	9	24	74	213	233	313	472	470	418
2001.....	2,643	49	19	8	24	79	178	204	269	418	480	421
2000.....	2,445	27	26	11	30	68	155	197	249	403	452	368
1999.....	2,415	33	18	6	20	71	163	202	277	422	420	342

All races, female

2003.....	1,731	25	14	7	7	43	99	99	151	227	324	276
2002.....	1,672	49	29	15	14	34	87	92	144	224	300	272
2001.....	1,324	34	25	7	6	29	62	75	132	223	225	182
2000.....	1,144	38	13	5	4	26	50	77	104	178	214	166
1999.....	1,178	22	22	6	5	20	49	79	134	194	202	168

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	4,072	52	21	7	23	126	315	286	406	560	735	636
2002.....	3,770	56	45	16	29	96	263	271	375	574	637	563
2001.....	3,265	45	31	9	26	99	208	233	339	517	584	494
2000.....	2,886	41	27	9	24	83	171	229	290	479	521	424
1999.....	2,875	36	22	10	21	83	174	233	337	491	482	396

White, male

2003.....	2,565	30	11	3	19	87	224	197	269	370	456	394
2002.....	2,371	29	23	5	18	66	189	194	256	381	392	340
2001.....	2,133	28	13	4	21	73	150	168	228	329	387	335
2000.....	1,941	18	17	6	21	60	126	163	201	334	351	287
1999.....	1,893	22	12	5	17	67	135	161	224	328	329	252

White, female

2003.....	1,507	22	10	4	4	39	91	89	137	190	279	242
2002.....	1,399	27	22	11	11	30	74	77	119	193	245	223
2001.....	1,132	17	18	5	5	26	58	65	111	188	197	159
2000.....	945	23	10	3	3	23	45	66	89	145	170	137
1999.....	982	14	10	5	4	16	39	72	113	163	153	144

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2003.....	652	21	8	6	4	9	25	34	55	90	123	114
2002.....	718	37	7	7	6	8	32	47	67	114	124	116
2001.....	614	36	13	5	4	5	24	35	53	112	113	99
2000.....	614	15	11	7	9	7	25	37	53	94	133	102
1999.....	630	18	14	2	4	5	31	41	66	111	123	105

Black, male

2003.....	466	19	4	3	1	6	21	28	42	58	84	83
2002.....	482	18	2	3	6	5	21	34	45	86	73	74
2001.....	451	20	6	3	3	3	21	28	35	81	88	78
2000.....	441	3	8	5	8	7	21	28	40	63	94	76
1999.....	467	10	6	1	3	3	24	35	46	84	83	84

Black, female

2003.....	186	2	4	3	3	3	4	6	13	32	39	31
2002.....	236	19	5	4	-	3	11	13	22	28	51	42
2001.....	163	16	7	2	1	2	3	7	18	31	25	21
2000.....	173	12	3	2	1	-	4	9	13	31	39	26
1999.....	163	8	8	1	1	2	7	6	20	27	40	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

American Indian 1/, both sexes

2003.....	55	2	-	1	-	1	5	5	2	10	9	5
2002.....	41	3	2	1	1	2	-	2	8	3	4	7
2001.....	39	1	-	-	-	3	2	2	4	7	6	5
2000.....	42	4	1	-	1	1	2	3	5	6	6	4
1999.....	48	1	2	-	-	1	4	2	8	9	8	4

American Indian 1/, male

2003.....	39	1	-	1	-	-	5	2	2	8	6	3
2002.....	21	1	1	1	-	2	-	-	6	1	1	3
2001.....	28	-	-	-	-	3	2	2	4	5	3	3
2000.....	33	3	1	-	1	-	2	2	5	5	4	3
1999.....	32	1	-	-	-	-	3	2	7	6	4	3

American Indian 1/, female

2003.....	16	1	-	-	-	1	-	3	-	2	3	2
2002.....	20	2	1	-	1	-	-	2	2	2	3	4
2001.....	11	1	-	-	-	-	-	-	-	2	3	2
2000.....	9	1	-	-	-	1	-	1	-	1	2	1
1999.....	16	-	2	-	-	1	1	-	1	3	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 2/, both sexes

2003.....	61	-	1	1	1	2	11	5	4	5	6	3
2002.....	58	5	2	-	2	2	5	5	7	5	5	4
2001.....	49	1	-	1	-	1	6	9	5	5	2	5
2000.....	47	5	-	-	-	3	7	5	5	2	6	4
1999.....	40	-	2	-	-	2	3	5	-	5	9	5

Asian or Pacific Islander 2/, male

2003.....	39	-	1	1	1	2	7	4	3	2	3	2
2002.....	41	4	1	-	-	1	3	5	6	4	4	1
2001.....	31	1	-	1	-	-	5	6	2	3	2	5
2000.....	30	3	-	-	-	1	6	4	3	1	3	2
1999.....	23	-	-	-	-	1	1	4	-	4	4	3

Asian or Pacific Islander 2/, female

2003.....	22	-	-	-	-	-	4	1	1	3	3	1
2002.....	17	1	1	-	2	1	2	-	1	1	1	3
2001.....	18	-	-	-	-	1	1	3	3	2	-	-
2000.....	17	2	-	-	-	2	1	1	2	1	3	2
1999.....	17	-	2	-	-	1	2	1	-	1	5	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
All races, both sexes												
2003.....	2,855	15	24	6	6	21	23	26	43	75	104	145
2002.....	2,843	15	19	13	22	16	19	27	52	73	98	114
2001.....	3,021	26	24	5	9	18	25	24	55	71	101	142
2000.....	3,059	19	22	18	13	10	22	37	48	65	84	128
1999.....	2,823	54	30	7	11	17	19	29	49	66	86	91
All races, male												
2003.....	1,333	8	16	2	2	18	8	14	22	36	60	80
2002.....	1,236	8	11	8	18	10	9	16	31	35	47	60
2001.....	1,357	15	14	2	6	6	13	11	31	39	44	75
2000.....	1,430	12	11	11	4	5	11	23	24	33	43	59
1999.....	1,308	31	12	4	5	8	13	9	34	37	41	37
All races, female												
2003.....	1,522	7	8	4	4	3	15	12	21	39	44	65
2002.....	1,607	7	8	5	4	6	10	11	21	38	51	54
2001.....	1,664	11	10	3	3	12	12	13	24	32	57	67
2000.....	1,629	7	11	7	9	5	11	14	24	32	41	69
1999.....	1,515	23	18	3	6	9	6	20	15	29	45	54

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
White, both sexes												
2003.....	2,356	9	13	5	3	18	19	16	32	57	84	103
2002.....	2,331	8	14	7	14	10	14	20	37	51	68	80
2001.....	2,446	17	16	3	7	13	20	16	41	47	70	100
2000.....	2,520	14	12	12	11	8	16	22	38	52	63	97
1999.....	2,314	35	21	6	8	15	15	16	39	52	62	68
White, male												
2003.....	1,121	5	9	2	1	15	7	8	17	29	52	56
2002.....	1,025	5	9	4	12	7	7	12	23	28	37	45
2001.....	1,084	9	10	1	4	5	11	7	22	24	30	53
2000.....	1,210	11	8	7	4	4	9	13	20	27	37	45
1999.....	1,078	22	11	3	4	7	10	4	28	30	30	26
White, female												
2003.....	1,235	4	4	3	2	3	12	8	15	28	32	47
2002.....	1,306	3	5	3	2	3	7	8	14	23	31	35
2001.....	1,362	8	6	2	3	8	9	9	19	23	40	47
2000.....	1,310	3	4	5	7	4	7	9	18	25	26	52
1999.....	1,236	13	10	3	4	8	5	12	11	22	32	42

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2003.....	432	5	10	-	3	2	4	7	9	17	17	34
2002.....	448	7	5	6	8	5	2	6	11	20	29	31
2001.....	522	9	7	1	2	3	4	8	14	22	28	39
2000.....	487	5	8	4	1	2	5	13	10	13	18	27
1999.....	449	18	9	1	3	2	4	10	8	13	22	21

Black, male

2003.....	177	3	6	-	1	2	1	3	5	6	6	19
2002.....	184	3	2	4	6	3	1	4	5	6	10	13
2001.....	250	6	4	-	2	-	2	4	9	14	13	20
2000.....	193	1	2	2	-	1	1	9	4	6	6	12
1999.....	197	8	1	1	1	1	3	4	6	6	10	11

Black, female

2003.....	255	2	4	-	2	-	3	4	4	11	11	15
2002.....	264	4	3	2	2	2	1	2	6	14	19	18
2001.....	272	3	3	1	-	3	2	4	5	8	15	19
2000.....	294	4	6	2	1	1	4	4	6	7	12	15
1999.....	252	10	8	-	2	1	1	6	2	7	12	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

American Indian 1/, both sexes

2003.....	24	1	-	-	-	1	-	2	1	1	-	4
2002.....	26	-	-	-	-	-	1	-	2	1	1	2
2001.....	17	-	1	1	-	1	1	-	-	-	2	-
2000.....	14	-	-	-	1	-	-	-	-	-	1	-
1999.....	20	-	-	-	-	-	-	2	1	1	-	-

American Indian 1/, male

2003.....	10	-	-	-	-	1	-	2	-	1	-	3
2002.....	12	-	-	-	-	-	1	-	2	-	-	1
2001.....	6	-	-	1	-	-	-	-	-	-	1	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	-
1999.....	13	-	-	-	-	-	-	1	-	1	-	-

American Indian 1/, female

2003.....	14	1	-	-	-	-	-	-	1	-	-	1
2002.....	14	-	-	-	-	-	-	-	-	1	1	1
2001.....	11	-	1	-	-	1	1	-	-	-	1	-
2000.....	9	-	-	-	1	-	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

Asian or Pacific Islander 2/, both sexes

2003.....	43	-	1	1	-	-	-	1	1	-	3	4
2002.....	38	-	-	-	-	1	2	1	2	1	-	1
2001.....	36	-	-	-	-	1	-	-	-	2	1	3
2000.....	38	-	2	2	-	-	1	2	-	-	2	4
1999.....	40	1	-	-	-	-	-	1	1	-	2	2

Asian or Pacific Islander 2/, male

2003.....	25	-	1	-	-	-	-	1	-	-	2	2
2002.....	15	-	-	-	-	-	-	-	1	1	-	1
2001.....	17	-	-	-	-	1	-	-	-	1	-	2
2000.....	22	-	1	2	-	-	1	1	-	-	-	2
1999.....	20	1	-	-	-	-	-	-	-	-	1	-

Asian or Pacific Islander 2/, female

2003.....	18	-	-	1	-	-	-	-	1	-	1	2
2002.....	23	-	-	-	-	1	2	1	1	-	-	-
2001.....	19	-	-	-	-	-	-	-	-	1	1	1
2000.....	16	-	1	-	-	-	-	1	-	-	2	2
1999.....	20	-	-	-	-	-	-	1	1	-	1	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

All races, both sexes

2003.....	20,687	-	-	-	5	49	75	167	429	1,144	2,428	3,472
2002 5/.....	20,218	1	2	-	3	33	72	160	441	1,191	2,445	3,321
2001 5/.....	20,114	2	-	-	2	41	65	168	477	1,329	2,417	3,422
2000 5/.....	19,643	3	1	-	4	33	62	175	475	1,355	2,482	3,203
1999 5/.....	19,469	1	-	-	2	25	75	147	494	1,416	2,413	3,149

All races, male

2003.....	15,630	-	-	-	3	37	64	139	333	799	1,735	2,627
2002 5/.....	15,272	-	1	-	2	28	58	133	325	829	1,742	2,485
2001 5/.....	15,149	2	-	-	-	31	57	124	342	940	1,725	2,593
2000 5/.....	14,993	2	-	-	4	27	49	146	358	986	1,822	2,477
1999 5/.....	14,894	1	-	-	1	20	60	121	338	994	1,822	2,436

All races, female

2003.....	5,057	-	-	-	2	12	11	28	96	345	693	845
2002 5/.....	4,946	1	1	-	1	5	14	27	116	362	703	836
2001 5/.....	4,965	-	-	-	2	10	8	44	135	389	692	829
2000 5/.....	4,650	1	1	-	-	6	13	29	117	369	660	726
1999 5/.....	4,575	-	-	-	1	5	15	26	156	422	591	713

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

White, both sexes

2003.....	17,437	-	-	-	4	38	58	133	350	931	2,033	2,913
2002 5/.....	16,988	-	1	-	2	31	55	126	350	965	2,061	2,787
2001 5/.....	16,640	1	-	-	2	32	46	128	379	1,082	1,951	2,840
2000 5/.....	16,223	3	-	-	3	28	47	138	366	1,099	1,996	2,592
1999 5/.....	15,903	1	-	-	2	17	60	122	392	1,119	1,916	2,508

White, male

2003.....	13,218	-	-	-	3	32	50	113	270	662	1,471	2,212
2002 5/.....	12,926	-	1	-	1	27	46	107	265	680	1,492	2,107
2001 5/.....	12,588	1	-	-	-	25	41	99	278	764	1,400	2,179
2000 5/.....	12,509	2	-	-	3	23	39	119	281	812	1,481	2,047
1999 5/.....	12,277	1	-	-	1	13	50	104	274	800	1,476	1,970

White, female

2003.....	4,219	-	-	-	1	6	8	20	80	269	562	701
2002 5/.....	4,062	-	-	-	1	4	9	19	85	285	569	680
2001 5/.....	4,052	-	-	-	2	7	5	29	101	318	551	661
2000 5/.....	3,714	1	-	-	-	5	8	19	85	287	515	545
1999 5/.....	3,626	-	-	-	1	4	10	18	118	319	440	538

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Black, both sexes

2003.....	2,406	-	-	-	1	2	6	24	36	135	254	397
2002 5/.....	2,434	1	1	-	-	-	10	17	54	128	261	404
2001 5/.....	2,723	1	-	-	-	5	12	20	52	153	350	453
2000 5/.....	2,712	-	1	-	1	2	7	24	70	163	364	509
1999 5/.....	2,832	-	-	-	-	4	6	14	60	209	356	523

Black, male

2003.....	1,824	-	-	-	-	2	4	19	29	87	172	303
2002 5/.....	1,798	-	-	-	-	-	8	14	39	84	171	289
2001 5/.....	2,048	1	-	-	-	3	10	15	33	116	245	331
2000 5/.....	1,993	-	-	-	1	2	6	20	45	113	254	361
1999 5/.....	2,100	-	-	-	-	4	4	9	38	141	244	381

Black, female

2003.....	582	-	-	-	1	-	2	5	7	48	82	94
2002 5/.....	636	1	1	-	-	-	2	3	15	44	90	115
2001 5/.....	675	-	-	-	-	2	2	5	19	37	105	122
2000 5/.....	719	-	1	-	-	-	1	4	25	50	110	148
1999 5/.....	732	-	-	-	-	-	2	5	22	68	112	142

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

American Indian 1/, both sexes

2003.....	664	-	-	-	-	8	9	7	29	68	117	126
2002 5/.....	597	-	-	-	1	2	6	17	34	79	94	102
2001 5/.....	591	-	-	-	-	4	6	17	39	81	99	104
2000 5/.....	537	-	-	-	-	2	5	13	35	77	95	75
1999 5/.....	589	-	-	-	-	4	7	9	33	76	119	94

American Indian 1/, male

2003.....	431	-	-	-	-	2	9	5	21	41	70	80
2002 5/.....	387	-	-	-	1	1	3	12	19	50	55	63
2001 5/.....	387	-	-	-	-	3	5	10	26	50	66	61
2000 5/.....	353	-	-	-	-	1	1	7	28	46	65	46
1999 5/.....	397	-	-	-	-	3	4	6	20	42	81	65

American Indian 1/, female

2003.....	233	-	-	-	-	6	-	2	8	27	47	46
2002 5/.....	210	-	-	-	-	1	3	5	15	29	39	39
2001 5/.....	204	-	-	-	-	1	1	7	13	31	33	43
2000 5/.....	184	-	-	-	-	1	4	6	7	31	30	29
1999 5/.....	192	-	-	-	-	1	3	3	13	34	38	29

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Asian or Pacific Islander 2/, both sexes

2003.....	180	-	-	-	-	1	2	3	14	10	24	36
2002 5/.....	199	-	-	-	-	-	1	-	3	19	29	28
2001 5/.....	160	-	-	-	-	-	1	3	7	13	17	25
2000 5/.....	171	-	-	-	-	1	3	-	4	16	27	27
1999 5/.....	145	-	-	-	-	-	2	2	9	12	22	24

Asian or Pacific Islander 2/, male

2003.....	157	-	-	-	-	1	1	2	13	9	22	32
2002 5/.....	161	-	-	-	-	-	1	-	2	15	24	26
2001 5/.....	126	-	-	-	-	-	1	-	5	10	14	22
2000 5/.....	138	-	-	-	-	1	3	-	4	15	22	23
1999 5/.....	120	-	-	-	-	-	2	2	6	11	21	20

Asian or Pacific Islander 2/, female

2003.....	23	-	-	-	-	-	1	1	1	1	2	4
2002 5/.....	38	-	-	-	-	-	-	-	1	4	5	2
2001 5/.....	34	-	-	-	-	-	-	3	2	3	3	3
2000 5/.....	33	-	-	-	-	-	-	-	-	1	5	4
1999 5/.....	25	-	-	-	-	-	-	-	3	1	1	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
All races, both sexes												
2003.....	28,723	25	35	9	32	669	1,950	2,086	2,759	3,918	5,240	4,946
2002 5/.....	26,040	29	30	7	35	625	1,590	1,850	2,599	3,837	5,021	4,451
2001 5/.....	21,705	22	26	8	37	539	1,274	1,552	2,187	3,448	4,221	3,660
2000 5/.....	19,720	18	15	9	35	452	1,115	1,450	2,077	3,270	3,936	3,171
1999 5/.....	19,128	21	25	7	23	367	984	1,452	2,127	3,242	3,853	3,042
All races, male												
2003.....	18,426	17	21	2	18	477	1,478	1,508	1,853	2,549	3,324	3,131
2002 5/.....	16,734	20	9	6	18	447	1,175	1,346	1,793	2,534	3,194	2,765
2001 5/.....	14,253	14	21	3	19	386	964	1,142	1,498	2,302	2,777	2,440
2000 5/.....	13,137	12	13	6	19	318	848	1,057	1,458	2,242	2,648	2,170
1999 5/.....	12,885	14	12	1	15	251	730	1,067	1,512	2,238	2,669	2,121
All races, female												
2003.....	10,297	8	14	7	14	192	472	578	906	1,369	1,916	1,815
2002 5/.....	9,306	9	21	1	17	178	415	504	806	1,303	1,827	1,686
2001 5/.....	7,452	8	5	5	18	153	310	410	689	1,146	1,444	1,220
2000 5/.....	6,583	6	2	3	16	134	267	393	619	1,028	1,288	1,001
1999 5/.....	6,243	7	13	6	8	116	254	385	615	1,004	1,184	921

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
White, both sexes												
2003.....	24,683	16	22	7	24	631	1,820	1,868	2,408	3,337	4,472	4,146
2002 5/.....	22,146	14	18	4	32	572	1,464	1,633	2,257	3,259	4,263	3,671
2001 5/.....	18,195	10	17	6	32	490	1,161	1,364	1,881	2,909	3,499	2,919
2000 5/.....	16,388	11	7	6	31	415	1,002	1,253	1,757	2,736	3,200	2,511
1999 5/.....	15,714	13	14	4	21	333	861	1,229	1,779	2,699	3,085	2,379
White, male												
2003.....	15,824	14	14	2	15	457	1,387	1,372	1,633	2,186	2,853	2,611
2002 5/.....	14,170	11	3	3	17	411	1,088	1,190	1,576	2,160	2,735	2,275
2001 5/.....	11,882	4	15	3	16	363	872	1,003	1,314	1,960	2,319	1,915
2000 5/.....	10,857	7	5	5	16	297	771	911	1,247	1,900	2,161	1,696
1999 5/.....	10,506	9	8	-	14	226	645	901	1,283	1,875	2,158	1,633
White, female												
2003.....	8,859	2	8	5	9	174	433	496	775	1,151	1,619	1,535
2002 5/.....	7,976	3	15	1	15	161	376	443	681	1,099	1,528	1,396
2001 5/.....	6,313	6	2	3	16	127	289	361	567	949	1,180	1,004
2000 5/.....	5,531	4	2	1	15	118	231	342	510	836	1,039	815
1999 5/.....	5,208	4	6	4	7	107	216	328	496	824	927	746

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
Black, both sexes												
2003.....	3,527	9	13	2	7	23	91	155	291	508	688	742
2002 5/.....	3,463	13	10	2	3	32	98	175	286	502	700	720
2001 5/.....	3,165	11	8	2	3	29	87	153	264	489	670	690
2000 5/.....	3,034	6	6	3	2	29	89	166	285	488	671	614
1999 5/.....	3,100	8	11	2	2	23	101	186	311	487	709	636
Black, male												
2003.....	2,303	3	7	-	3	10	68	99	180	325	424	482
2002 5/.....	2,307	8	6	2	1	22	69	125	185	330	424	459
2001 5/.....	2,163	9	5	-	2	15	73	117	162	309	427	495
2000 5/.....	2,094	4	6	1	1	16	63	125	186	316	446	446
1999 5/.....	2,191	5	4	1	1	19	75	142	205	331	475	470
Black, female												
2003.....	1,224	6	6	2	4	13	23	56	111	183	264	260
2002 5/.....	1,156	5	4	-	2	10	29	50	101	172	276	261
2001 5/.....	1,002	2	3	2	1	14	14	36	102	180	243	195
2000 5/.....	940	2	-	2	1	13	26	41	99	172	225	168
1999 5/.....	909	3	7	1	1	4	26	44	106	156	234	166

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

American Indian 1/, both sexes

2003.....	295	-	-	-	1	9	18	31	33	53	56	31
2002 5/.....	230	2	1	1	-	13	11	19	30	46	36	38
2001 5/.....	184	-	-	-	1	10	8	15	26	28	31	34
2000 5/.....	160	1	1	-	2	1	12	16	21	24	38	25
1999 5/.....	164	-	-	1	-	5	7	22	22	33	36	17

American Indian 1/, male

2003.....	168	-	-	-	-	6	12	17	21	28	31	19
2002 5/.....	136	1	-	1	-	9	9	13	18	25	18	20
2001 5/.....	100	-	-	-	-	2	8	9	11	17	14	20
2000 5/.....	99	1	1	-	2	-	6	13	16	13	20	15
1999 5/.....	96	-	-	-	-	2	6	14	14	18	19	11

American Indian 1/, female

2003.....	127	-	-	-	1	3	6	14	12	25	25	12
2002 5/.....	94	1	1	-	-	4	2	6	12	21	18	18
2001 5/.....	84	-	-	-	1	8	-	6	15	11	17	14
2000 5/.....	61	-	-	-	-	1	6	3	5	11	18	10
1999 5/.....	68	-	-	1	-	3	1	8	8	15	17	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
Asian or Pacific Islander 2/, both sexes												
2003.....	218	-	-	-	-	6	21	32	27	20	24	27
2002 5/.....	201	-	1	-	-	8	17	23	26	30	22	22
2001 5/.....	161	1	1	-	1	10	18	20	16	22	21	17
2000 5/.....	138	-	1	-	-	7	12	15	14	22	27	21
1999 5/.....	150	-	-	-	-	6	15	15	15	23	23	10
Asian or Pacific Islander 2/, male												
2003.....	131	-	-	-	-	4	11	20	19	10	16	19
2002 5/.....	121	-	-	-	-	5	9	18	14	19	17	11
2001 5/.....	108	1	1	-	1	6	11	13	11	16	17	10
2000 5/.....	87	-	1	-	-	5	8	8	9	13	21	13
1999 5/.....	92	-	-	-	-	4	4	10	10	14	17	7
Asian or Pacific Islander 2/, female												
2003.....	87	-	-	-	-	2	10	12	8	10	8	8
2002 5/.....	80	-	1	-	-	3	8	5	12	11	5	11
2001 5/.....	53	-	-	-	-	4	7	7	5	6	4	7
2000 5/.....	51	-	-	-	-	2	4	7	5	9	6	8
1999 5/.....	58	-	-	-	-	2	11	5	5	9	6	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

All races, both sexes

2003.....	30,136	8	48	63	261	2,469	4,377	3,336	2,852	2,476	2,642	2,475
2002.....	30,242	10	61	71	277	2,474	4,306	3,380	2,760	2,712	2,734	2,358
2001.....	29,573	11	70	79	254	2,523	4,164	3,350	2,787	2,723	2,565	2,203
2000.....	28,663	13	46	70	307	2,606	3,969	3,165	2,624	2,718	2,640	2,146
1999.....	28,874	8	65	80	336	2,896	3,899	3,248	2,727	2,812	2,422	2,096

All races, male

2003.....	26,124	7	28	40	216	2,231	3,955	2,999	2,473	2,065	2,157	2,025
2002.....	26,098	4	38	38	204	2,209	3,887	2,994	2,353	2,257	2,246	1,927
2001.....	25,480	7	39	51	198	2,270	3,796	2,963	2,384	2,216	2,096	1,828
2000.....	24,582	9	28	46	256	2,323	3,583	2,796	2,207	2,207	2,156	1,770
1999.....	24,700	5	36	50	258	2,560	3,496	2,848	2,308	2,279	1,950	1,722

All races, female

2003.....	4,012	1	20	23	45	238	422	337	379	411	485	450
2002.....	4,144	6	23	33	73	265	419	386	407	455	488	431
2001.....	4,093	4	31	28	56	253	368	387	403	507	469	375
2000.....	4,081	4	18	24	51	283	386	369	417	511	484	376
1999.....	4,174	3	29	30	78	336	403	400	419	533	472	374

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

White, both sexes

2003.....	21,763	2	24	33	159	1,348	2,262	1,757	1,764	1,796	2,098	2,088
2002.....	21,902	4	42	45	175	1,388	2,209	1,844	1,700	1,932	2,193	1,986
2001.....	21,760	8	43	61	180	1,418	2,168	1,922	1,837	2,011	2,062	1,881
2000.....	20,945	6	28	50	225	1,465	2,079	1,764	1,713	2,008	2,115	1,788
1999.....	21,143	4	32	41	248	1,617	2,053	1,874	1,839	2,120	1,944	1,742

White, male

2003.....	18,647	1	12	19	137	1,194	2,022	1,538	1,502	1,465	1,708	1,691
2002.....	18,714	2	25	24	130	1,226	1,968	1,576	1,423	1,583	1,806	1,611
2001.....	18,527	5	25	39	142	1,238	1,932	1,674	1,535	1,630	1,690	1,561
2000.....	17,750	2	15	32	189	1,284	1,845	1,515	1,416	1,599	1,733	1,477
1999.....	17,942	4	18	24	191	1,418	1,806	1,615	1,530	1,704	1,566	1,431

White, female

2003.....	3,116	1	12	14	22	154	240	219	262	331	390	397
2002.....	3,188	2	17	21	45	162	241	268	277	349	387	375
2001.....	3,233	3	18	22	38	180	236	248	302	381	372	320
2000.....	3,195	4	13	18	36	181	234	249	297	409	382	311
1999.....	3,201	-	14	17	57	199	247	259	309	416	378	311

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/												
Black, both sexes												
2003.....	7,659	6	23	27	89	1,036	1,981	1,490	998	610	485	327
2002.....	7,623	6	17	18	85	997	1,958	1,448	983	706	482	331
2001.....	7,184	2	23	16	64	1,033	1,890	1,335	873	662	443	279
2000.....	7,054	7	13	18	74	1,054	1,788	1,296	851	639	464	312
1999.....	7,017	2	27	32	74	1,177	1,734	1,264	803	617	430	312
Black, male												
2003.....	6,882	6	15	19	67	969	1,817	1,387	892	540	401	287
2002.....	6,798	2	12	11	63	903	1,802	1,342	874	610	396	286
2001.....	6,438	1	13	10	52	964	1,774	1,212	788	545	361	233
2000.....	6,284	7	8	13	62	967	1,650	1,190	744	553	376	257
1999.....	6,184	1	13	23	56	1,052	1,595	1,144	710	514	348	259
Black, female												
2003.....	777	-	8	8	22	67	164	103	106	70	84	40
2002.....	825	4	5	7	22	94	156	106	109	96	86	45
2001.....	746	1	10	6	12	69	116	123	85	117	82	46
2000.....	770	-	5	5	12	87	138	106	107	86	88	55
1999.....	833	1	14	9	18	125	139	120	93	103	82	53

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

American Indian 1/, both sexes

2003.....	269	-	-	2	6	43	53	39	31	24	26	20
2002.....	287	-	1	3	7	41	64	37	31	29	22	17
2001.....	240	1	4	2	5	38	38	26	34	22	28	13
2000.....	240	-	4	1	2	38	46	33	24	22	22	11
1999.....	268	2	2	2	6	46	50	42	34	26	11	13

American Indian 1/, male

2003.....	229	-	-	1	5	34	47	34	27	21	21	18
2002.....	235	-	-	1	3	35	51	33	24	26	19	12
2001.....	196	1	1	2	1	35	33	24	25	15	23	11
2000.....	196	-	4	1	1	32	41	27	18	17	14	7
1999.....	228	-	2	1	5	42	44	34	30	23	7	11

American Indian 1/, female

2003.....	40	-	-	1	1	9	6	5	4	3	5	2
2002.....	52	-	1	2	4	6	13	4	7	3	3	5
2001.....	44	-	3	-	4	3	5	2	9	7	5	2
2000.....	44	-	-	-	1	6	5	6	6	5	8	4
1999.....	40	2	-	1	1	4	6	8	4	3	4	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Asian or Pacific Islander 2/, both sexes

2003.....	445	-	1	1	7	42	81	50	59	46	33	40
2002.....	430	-	1	5	10	48	75	51	46	45	37	24
2001.....	389	-	-	-	5	34	68	67	43	28	32	30
2000.....	424	-	1	1	6	49	56	72	36	49	39	35
1999.....	446	-	4	5	8	56	62	68	51	49	37	29

Asian or Pacific Islander 2/, male

2003.....	366	-	1	1	7	34	69	40	52	39	27	29
2002.....	351	-	1	2	8	45	66	43	32	38	25	18
2001.....	319	-	-	-	3	33	57	53	36	26	22	23
2000.....	352	-	1	-	4	40	47	64	29	38	33	29
1999.....	346	-	3	2	6	48	51	55	38	38	29	21

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Asian or Pacific Islander 2/, female

2003.....	79	-	-	-	-	8	12	10	7	7	6	11
2002.....	79	-	-	3	2	3	9	8	14	7	12	6
2001.....	70	-	-	-	2	1	11	14	7	2	10	7
2000.....	72	-	-	1	2	9	9	8	7	11	6	6
1999.....	100	-	1	3	2	8	11	13	13	11	8	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes

All races, both sexes

2003.....	98,608	119,339	143,180	174,817	238,680	325,185	377,839	346,847	231,403	90,271	19,331	342
2002.....	96,320	115,441	137,901	175,591	247,399	330,140	377,514	345,385	228,587	88,282	18,822	357
2001.....	94,381	109,523	134,616	178,232	252,728	333,677	368,252	339,775	221,314	85,954	18,488	422
2000.....	89,509	106,751	134,095	181,739	259,470	337,700	362,745	335,813	219,283	84,641	18,434	356
1999.....	84,854	104,884	134,095	186,695	265,905	338,349	360,241	330,903	213,991	83,334	17,913	356

All races, male

2003.....	61,543	72,188	84,273	100,315	131,106	167,354	174,978	135,177	70,162	19,872	3,001	271
2002.....	60,021	69,830	81,533	100,590	136,431	169,625	173,879	134,637	68,817	19,522	2,930	282
2001.....	58,281	65,801	79,157	102,110	139,471	171,311	169,431	130,405	65,527	18,732	2,869	331
2000.....	55,277	64,425	78,896	103,935	143,473	173,327	166,892	128,877	64,439	18,552	2,874	289
1999.....	52,356	63,441	79,283	107,529	147,391	174,853	166,117	126,531	62,511	18,111	2,836	277

All races, female

2003.....	37,065	47,151	58,907	74,502	107,574	157,831	202,861	211,670	161,241	70,399	16,330	71
2002.....	36,299	45,611	56,368	75,001	110,968	160,515	203,635	210,748	159,770	68,760	15,892	75
2001.....	36,100	43,722	55,459	76,122	113,257	162,366	198,821	209,370	155,787	67,222	15,619	91
2000.....	34,232	42,326	55,199	77,804	115,997	164,373	195,853	206,936	154,844	66,089	15,560	67
1999.....	32,498	41,443	54,812	79,166	118,514	163,496	194,124	204,372	151,480	65,223	15,077	79

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes

White, both sexes

2003.....	75,377	95,720	117,101	145,645	204,904	286,819	340,633	317,904	212,499	82,525	16,767	292
2002.....	74,126	92,712	112,702	146,491	213,556	291,430	340,923	316,444	210,030	80,543	16,438	286
2001.....	73,065	87,807	110,046	148,608	218,520	295,048	332,704	311,165	203,570	78,462	16,068	320
2000.....	69,543	85,840	109,701	152,597	224,389	299,257	328,472	307,495	201,942	77,167	16,157	275
1999.....	66,116	83,876	109,706	157,260	230,177	299,701	326,788	302,550	197,437	76,142	15,820	271

White, male

2003.....	47,689	58,348	69,404	84,313	113,540	148,708	158,342	124,593	64,481	17,945	2,556	228
2002.....	46,757	56,535	67,016	84,741	118,994	150,862	157,792	124,020	63,038	17,634	2,452	230
2001.....	45,629	53,120	65,233	86,088	121,705	152,517	153,761	119,811	60,027	16,916	2,425	255
2000.....	43,267	52,048	65,066	88,182	125,197	154,451	151,919	118,321	59,009	16,624	2,455	231
1999.....	41,130	51,133	65,404	91,520	128,922	155,859	151,591	115,599	57,211	16,329	2,471	213

White, female

2003.....	27,688	37,372	47,697	61,332	91,364	138,111	182,291	193,311	148,018	64,580	14,211	64
2002.....	27,369	36,177	45,686	61,750	94,562	140,568	183,131	192,424	146,992	62,909	13,986	56
2001.....	27,436	34,687	44,813	62,520	96,815	142,531	178,943	191,354	143,543	61,546	13,643	65
2000.....	26,276	33,792	44,635	64,415	99,192	144,806	176,553	189,174	142,933	60,543	13,702	44
1999.....	24,986	32,743	44,302	65,740	101,255	143,842	175,197	186,951	140,226	59,813	13,349	58

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes

Black, both sexes

2003.....	20,326	20,458	22,365	24,747	28,381	31,859	30,513	23,582	15,606	6,517	2,203	40
2002.....	19,439	19,693	21,643	24,783	28,531	32,431	30,431	23,920	15,458	6,578	2,066	57
2001.....	18,651	18,852	21,124	25,367	29,012	32,590	29,643	23,803	14,999	6,466	2,119	88
2000.....	17,554	18,161	21,120	25,064	30,131	32,541	28,903	23,919	14,680	6,407	2,032	72
1999.....	16,455	18,359	21,117	25,461	30,661	32,927	28,274	24,046	14,101	6,281	1,857	77

Black, male

2003.....	12,162	12,040	12,730	13,544	14,747	15,288	13,312	8,197	4,311	1,445	341	35
2002.....	11,621	11,550	12,450	13,465	14,733	15,434	13,044	8,292	4,455	1,454	383	40
2001.....	11,099	10,984	11,921	13,763	15,012	15,616	12,779	8,389	4,292	1,428	380	67
2000.....	10,563	10,787	11,891	13,505	15,566	15,682	12,344	8,428	4,249	1,532	351	50
1999.....	9,906	10,751	12,018	13,862	15,732	15,873	11,966	8,828	4,167	1,462	311	58

Black, female

2003.....	8,164	8,418	9,635	11,203	13,634	16,571	17,201	15,385	11,295	5,072	1,862	5
2002.....	7,818	8,143	9,193	11,318	13,798	16,997	17,387	15,628	11,003	5,124	1,683	17
2001.....	7,552	7,868	9,203	11,604	14,000	16,974	16,864	15,414	10,707	5,038	1,739	21
2000.....	6,991	7,374	9,229	11,559	14,565	16,859	16,559	15,491	10,431	4,875	1,681	22
1999.....	6,549	7,608	9,099	11,599	14,929	17,054	16,308	15,218	9,934	4,819	1,546	19

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes

American Indian 1/, both sexes

2003.....	970	951	1,091	1,111	1,145	1,233	1,130	837	503	175	79	5
2002.....	851	894	992	1,095	1,196	1,160	1,055	746	470	182	53	7
2001.....	812	821	947	1,060	1,146	1,148	998	782	407	167	46	6
2000.....	743	813	885	1,049	1,092	1,130	932	728	442	169	43	3
1999.....	700	783	891	1,037	1,127	1,151	913	733	390	157	59	1

American Indian 1/, male

2003.....	574	543	593	583	608	595	497	326	177	52	23	5
2002.....	523	514	560	590	611	586	462	317	135	47	17	6
2001.....	503	475	529	546	619	547	436	308	131	46	7	5
2000.....	469	459	505	556	557	567	422	286	155	50	10	3
1999.....	430	452	468	558	590	587	400	261	128	46	14	1

American Indian 1/, female

2003.....	396	408	498	528	537	638	633	511	326	123	56	-
2002.....	328	380	432	505	585	574	593	429	335	135	36	1
2001.....	309	346	418	514	527	601	562	474	276	121	39	1
2000.....	274	354	380	493	535	563	510	442	287	119	33	-
1999.....	270	331	423	479	537	564	513	472	262	111	45	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes

Asian or Pacific Islander 2/, both sexes

2003.....	1,935	2,210	2,623	3,314	4,250	5,274	5,563	4,524	2,795	1,054	282	5
2002.....	1,904	2,142	2,564	3,222	4,116	5,119	5,105	4,275	2,629	979	265	7
2001.....	1,853	2,043	2,499	3,197	4,050	4,891	4,907	4,025	2,338	859	255	8
2000.....	1,669	1,937	2,389	3,029	3,858	4,772	4,438	3,671	2,219	898	202	6
1999.....	1,583	1,866	2,381	2,937	3,940	4,570	4,266	3,574	2,063	754	177	7

Asian or Pacific Islander 2/, male

2003.....	1,118	1,257	1,546	1,875	2,211	2,763	2,827	2,061	1,193	430	81	3
2002.....	1,120	1,231	1,507	1,794	2,093	2,743	2,581	2,008	1,189	387	78	6
2001.....	1,050	1,222	1,474	1,713	2,135	2,631	2,455	1,897	1,077	342	57	4
2000.....	978	1,131	1,434	1,692	2,153	2,627	2,207	1,842	1,026	346	58	5
1999.....	890	1,105	1,393	1,589	2,147	2,534	2,160	1,843	1,005	274	40	5

Asian or Pacific Islander 2/, female

2003.....	817	953	1,077	1,439	2,039	2,511	2,736	2,463	1,602	624	201	2
2002.....	784	911	1,057	1,428	2,023	2,376	2,524	2,267	1,440	592	187	1
2001.....	803	821	1,025	1,484	1,915	2,260	2,452	2,128	1,261	517	198	4
2000.....	691	806	955	1,337	1,705	2,145	2,231	1,829	1,193	552	144	1
1999.....	693	761	988	1,348	1,793	2,036	2,106	1,731	1,058	480	137	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Salmonella infections (A01-A02)

All races, both sexes

2003.....	1	4	2	2	6	8	2	5	3	1	-	-
2002.....	2	-	-	3	1	3	5	1	-	-	-	-
2001.....	2	2	1	4	3	5	9	6	3	-	-	-
2000.....	3	2	-	1	6	3	2	1	2	-	-	-
1999.....	2	1	2	2	5	8	4	4	2	1	-	-

All races, male

2003.....	1	3	1	2	2	4	-	4	1	1	-	-
2002.....	1	-	-	1	1	1	4	1	-	-	-	-
2001.....	2	2	-	3	3	3	4	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	-	-	-
1999.....	2	-	2	2	3	2	3	1	1	-	-	-

All races, female

2003.....	-	1	1	-	4	4	2	1	2	-	-	-
2002.....	1	-	-	2	-	2	1	-	-	-	-	-
2001.....	-	-	1	1	-	2	5	5	3	-	-	-
2000.....	2	-	-	-	2	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	6	1	3	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Salmonella infections (A01-A02)

White, both sexes

2003.....	1	3	2	2	4	5	2	4	3	1	-	-
2002.....	2	-	-	3	-	2	5	1	-	-	-	-
2001.....	2	2	1	2	2	5	7	6	2	-	-	-
2000.....	3	2	-	1	5	3	2	1	2	-	-	-
1999.....	2	1	1	1	4	7	3	4	2	1	-	-

White, male

2003.....	1	2	1	2	2	2	-	3	1	1	-	-
2002.....	1	-	-	1	-	1	4	1	-	-	-	-
2001.....	2	2	-	2	2	3	4	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	-	-	-
1999.....	2	-	1	1	2	2	2	1	1	-	-	-

White, female

2003.....	-	1	1	-	2	3	2	1	2	-	-	-
2002.....	1	-	-	2	-	1	1	-	-	-	-	-
2001.....	-	-	1	-	-	2	3	5	2	-	-	-
2000.....	2	-	-	-	1	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	5	1	3	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Salmonella infections (A01-A02)

Black, both sexes

2003.....	-	-	-	-	1	3	-	-	-	-	-	-
2002.....	-	-	-	-	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	2	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	1	-	-	-	-	-

Black, male

2003.....	-	-	-	-	-	2	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-

Black, female

2003.....	-	-	-	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	2	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Shigellosis and amebiasis (A03,A06)

All races, both sexes

2003.....	-	1	1	-	1	1	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	2	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	1	-	2	5	-	-	-	1	-
1999.....	-	-	1	-	1	1	-	1	1	2	-	-

All races, male

2003.....	-	-	1	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	1	-	-	-

All races, female

2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	1	-	1	3	-	-	-	1	-
1999.....	-	-	-	-	-	1	-	1	-	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Shigellosis and amebiasis (A03,A06)

White, both sexes

2003.....	-	1	1	-	1	1	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	2	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	-	-	2	5	-	-	-	1	-
1999.....	-	-	1	-	1	1	-	1	1	2	-	-

White, male

2003.....	-	-	1	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	1	-	-	-

White, female

2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	3	-	-	-	1	-
1999.....	-	-	-	-	-	1	-	1	-	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

All races, both sexes

2003.....	43	70	106	161	283	524	668	647	383	120	17	-
2002.....	31	63	79	133	229	374	515	509	318	114	23	-
2001.....	33	28	53	73	162	237	329	293	187	75	10	-
2000.....	18	33	37	85	137	202	260	267	168	53	11	-
1999.....	12	15	34	67	109	168	222	211	146	58	14	-

All races, male

2003.....	21	35	47	68	117	213	265	198	86	23	2	-
2002.....	17	29	42	67	118	156	202	160	91	27	5	-
2001.....	11	14	21	32	76	107	120	93	50	14	-	-
2000.....	9	20	19	44	61	88	107	81	48	9	2	-
1999.....	6	10	16	28	46	80	89	54	44	13	2	-

All races, female

2003.....	22	35	59	93	166	311	403	449	297	97	15	-
2002.....	14	34	37	66	111	218	313	349	227	87	18	-
2001.....	22	14	32	41	86	130	209	200	137	61	10	-
2000.....	9	13	18	41	76	114	153	186	120	44	9	-
1999.....	6	5	18	39	63	88	133	157	102	45	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2003.....	32	61	89	147	266	478	642	619	363	112	16	-
2002.....	28	47	64	124	205	351	494	482	303	109	20	-
2001.....	31	24	42	60	151	222	305	284	178	68	10	-
2000.....	14	30	32	76	126	190	251	247	162	49	10	-
1999.....	10	13	28	62	100	155	209	204	139	55	13	-

White, male

2003.....	17	28	40	64	109	200	255	189	85	21	2	-
2002.....	16	22	34	61	103	145	195	152	85	27	5	-
2001.....	10	12	19	27	69	99	109	88	49	12	-	-
2000.....	7	17	17	40	56	83	101	76	46	8	2	-
1999.....	4	9	13	27	45	75	85	53	41	12	1	-

White, female

2003.....	15	33	49	83	157	278	387	430	278	91	14	-
2002.....	12	25	30	63	102	206	299	330	218	82	15	-
2001.....	21	12	23	33	82	123	196	196	129	56	10	-
2000.....	7	13	15	36	70	107	150	171	116	41	8	-
1999.....	6	4	15	35	55	80	124	151	98	43	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Black, both sexes

2003.....	8	8	15	13	12	40	24	26	18	7	1	-
2002.....	2	15	14	9	22	19	17	25	12	4	3	-
2001.....	2	4	9	12	11	11	19	7	7	5	-	-
2000.....	4	3	3	8	7	10	7	17	5	4	-	-
1999.....	1	2	4	5	9	13	12	5	7	3	1	-

Black, male

2003.....	3	6	6	4	5	13	10	8	1	2	-	-
2002.....	1	7	8	6	14	8	5	7	5	-	-	-
2001.....	1	2	2	4	7	4	8	3	1	2	-	-
2000.....	2	3	1	3	3	4	5	4	2	1	-	-
1999.....	1	1	1	1	1	5	4	-	3	1	1	-

Black, female

2003.....	5	2	9	9	7	27	14	18	17	5	1	-
2002.....	1	8	6	3	8	11	12	18	7	4	3	-
2001.....	1	2	7	8	4	7	11	4	6	3	-	-
2000.....	2	-	2	5	4	6	2	13	3	3	-	-
1999.....	-	1	3	4	8	8	8	5	4	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

American Indian 1/, both sexes

2003.....	1	-	2	1	2	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	1	1	-	2	-	-	-
2001.....	-	-	1	-	-	2	2	-	-	1	-	-
2000.....	-	-	2	1	-	-	-	1	-	-	-	-
1999.....	1	-	1	-	-	-	1	-	-	-	-	-

American Indian 1/, male

2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	-	1	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	1	-	1	1	1	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	-	2	-	-	-
2001.....	-	-	1	-	-	-	1	-	-	1	-	-
2000.....	-	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Asian or Pacific Islander 2/, both sexes

2003.....	2	1	-	-	3	4	2	2	2	1	-	-
2002.....	1	1	-	-	1	3	3	2	1	1	-	-
2001.....	-	-	1	1	-	2	3	2	2	1	-	-
2000.....	-	-	-	-	4	2	2	2	1	-	1	-
1999.....	-	-	1	-	-	-	-	2	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	1	1	-	-	2	-	-	1	-	-	-	-
2002.....	-	-	-	-	1	2	2	1	1	-	-	-
2001.....	-	-	-	1	-	2	2	2	-	-	-	-
2000.....	-	-	-	-	2	1	1	1	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	-	-	-	1	4	2	1	2	1	-	-
2002.....	1	1	-	-	-	1	1	1	-	1	-	-
2001.....	-	-	1	-	-	-	1	-	2	1	-	-
2000.....	-	-	-	-	2	1	1	1	1	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

All races, both sexes

2003.....	51	63	39	48	66	106	111	70	33	4	-	-
2002.....	46	44	63	82	79	114	120	68	42	9	2	-
2001.....	47	63	53	60	92	102	104	85	30	10	-	1
2000.....	53	31	50	70	85	132	120	74	31	10	-	1
1999.....	55	67	63	70	103	134	119	115	41	16	1	1

All races, male

2003.....	40	46	25	28	44	61	55	32	11	2	-	-
2002.....	35	36	39	54	48	56	65	36	16	2	1	-
2001.....	36	47	38	40	58	53	47	34	18	5	-	1
2000.....	41	23	36	46	41	69	70	38	13	5	-	1
1999.....	45	48	48	49	67	71	66	58	16	11	-	1

All races, female

2003.....	11	17	14	20	22	45	56	38	22	2	-	-
2002.....	11	8	24	28	31	58	55	32	26	7	1	-
2001.....	11	16	15	20	34	49	57	51	12	5	-	-
2000.....	12	8	14	24	44	63	50	36	18	5	-	-
1999.....	10	19	15	21	36	63	53	57	25	5	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

White, both sexes

2003.....	23	39	21	23	53	69	81	50	25	3	-	-
2002.....	23	20	37	41	47	78	85	48	30	7	1	-
2001.....	22	35	32	38	52	73	67	59	21	7	-	1
2000.....	31	8	25	38	49	79	76	53	26	6	-	-
1999.....	27	37	36	48	60	85	87	87	24	11	-	-

White, male

2003.....	19	29	14	12	34	36	41	23	7	2	-	-
2002.....	19	15	22	31	33	37	46	27	11	1	1	-
2001.....	16	29	22	24	29	38	31	22	11	4	-	1
2000.....	22	6	18	26	27	40	38	27	12	3	-	-
1999.....	25	27	25	33	40	42	49	43	9	6	-	-

White, female

2003.....	4	10	7	11	19	33	40	27	18	1	-	-
2002.....	4	5	15	10	14	41	39	21	19	6	-	-
2001.....	6	6	10	14	23	35	36	37	10	3	-	-
2000.....	9	2	7	12	22	39	38	26	14	3	-	-
1999.....	2	10	11	15	20	43	38	44	15	5	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

Black, both sexes

2003.....	23	21	10	14	6	16	17	11	4	-	-	-
2002.....	18	18	16	25	18	17	20	10	9	-	-	-
2001.....	23	23	15	19	25	16	19	11	8	3	-	-
2000.....	21	18	16	25	27	29	31	13	2	4	-	1
1999.....	22	22	18	15	28	31	19	14	8	4	1	1

Black, male

2003.....	18	16	5	10	5	12	8	5	2	-	-	-
2002.....	13	15	10	16	9	11	11	3	4	-	-	-
2001.....	19	16	11	13	17	7	7	5	6	1	-	-
2000.....	18	13	13	17	11	17	23	6	-	2	-	1
1999.....	17	15	17	11	18	17	8	6	3	4	-	1

Black, female

2003.....	5	5	5	4	1	4	9	6	2	-	-	-
2002.....	5	3	6	9	9	6	9	7	5	-	-	-
2001.....	4	7	4	6	8	9	12	6	2	2	-	-
2000.....	3	5	3	8	16	12	8	7	2	2	-	-
1999.....	5	7	1	4	10	14	11	8	5	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

American Indian 1/, both sexes

2003.....	2	-	-	-	1	2	2	1	-	1	-	-
2002.....	1	2	3	4	3	4	3	2	-	-	-	-
2001.....	-	1	1	-	2	1	3	2	-	-	-	-
2000.....	-	3	3	4	1	4	1	-	1	-	-	-
1999.....	1	1	1	1	5	1	-	1	3	1	-	-

American Indian 1/, male

2003.....	1	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	2	1	2	1	2	2	2	-	-	-	-
2001.....	-	-	1	-	1	1	1	-	-	-	-	-
2000.....	-	2	2	2	1	2	1	-	-	-	-	-
1999.....	-	1	-	1	2	-	-	-	2	1	-	-

American Indian 1/, female

2003.....	1	-	-	-	1	2	1	-	-	1	-	-
2002.....	1	-	2	2	2	2	1	-	-	-	-	-
2001.....	-	1	-	-	1	-	2	2	-	-	-	-
2000.....	-	1	1	2	-	2	-	-	1	-	-	-
1999.....	1	-	1	-	3	1	-	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

Asian or Pacific Islander 2/, both sexes

2003.....	3	3	8	11	6	19	11	8	4	-	-	-
2002.....	4	4	7	12	11	15	12	8	3	2	1	-
2001.....	2	4	5	3	13	12	15	13	1	-	-	-
2000.....	1	2	6	3	8	20	12	8	2	-	-	-
1999.....	5	7	8	6	10	17	13	13	6	-	-	-

Asian or Pacific Islander 2/, male

2003.....	2	1	6	6	5	13	5	3	2	-	-	-
2002.....	3	4	6	5	5	6	6	4	1	1	-	-
2001.....	1	2	4	3	11	7	8	7	1	-	-	-
2000.....	1	2	3	1	2	10	8	5	1	-	-	-
1999.....	3	5	6	4	7	12	9	9	2	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	2	2	5	1	6	6	5	2	-	-	-
2002.....	1	-	1	7	6	9	6	4	2	1	1	-
2001.....	1	2	1	-	2	5	7	6	-	-	-	-
2000.....	-	-	3	2	6	10	4	3	1	-	-	-
1999.....	2	2	2	2	3	5	4	4	4	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

All races, both sexes

2003.....	42	52	29	36	52	89	88	55	28	3	-	-
2002.....	36	37	47	61	59	87	105	60	37	7	2	-
2001.....	34	51	45	47	68	82	84	71	28	7	-	1
2000.....	42	24	36	52	67	107	102	64	26	8	-	1
1999.....	38	53	56	58	81	112	104	100	36	16	1	1

All races, male

2003.....	32	38	19	23	38	52	45	26	9	1	-	-
2002.....	27	30	30	44	36	40	56	32	15	1	1	-
2001.....	29	41	37	31	45	46	38	29	17	4	-	1
2000.....	35	21	29	38	35	58	58	33	13	4	-	1
1999.....	32	39	44	41	56	60	60	52	13	11	-	1

All races, female

2003.....	10	14	10	13	14	37	43	29	19	2	-	-
2002.....	9	7	17	17	23	47	49	28	22	6	1	-
2001.....	5	10	8	16	23	36	46	42	11	3	-	-
2000.....	7	3	7	14	32	49	44	31	13	4	-	-
1999.....	6	14	12	17	25	52	44	48	23	5	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

White, both sexes

2003.....	17	31	17	16	41	59	66	38	20	2	-	-
2002.....	17	17	26	34	36	59	70	42	27	6	1	-
2001.....	16	30	27	29	40	59	55	46	19	5	-	1
2000.....	26	7	19	27	39	65	66	45	22	5	-	-
1999.....	20	27	34	38	49	70	75	79	23	11	-	-

White, male

2003.....	13	23	12	10	29	30	34	18	5	1	-	-
2002.....	14	12	17	28	26	25	37	24	11	1	1	-
2001.....	14	26	22	18	24	34	25	17	10	3	-	1
2000.....	21	5	15	20	23	36	31	24	12	2	-	-
1999.....	19	21	24	27	34	36	43	41	9	6	-	-

White, female

2003.....	4	8	5	6	12	29	32	20	15	1	-	-
2002.....	3	5	9	6	10	34	33	18	16	5	-	-
2001.....	2	4	5	11	16	25	30	29	9	2	-	-
2000.....	5	2	4	7	16	29	35	21	10	3	-	-
1999.....	1	6	10	11	15	34	32	38	14	5	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Black, both sexes

2003.....	21	18	7	11	6	12	14	9	4	-	-	-
2002.....	16	15	14	19	13	14	20	8	7	-	-	-
2001.....	17	18	13	15	17	13	15	10	8	2	-	-
2000.....	15	12	10	21	22	24	24	12	2	3	-	1
1999.....	14	20	16	13	20	26	17	11	7	4	1	1

Black, male

2003.....	16	14	4	8	5	10	7	5	2	-	-	-
2002.....	12	13	9	12	7	10	11	2	3	-	-	-
2001.....	15	14	10	10	13	6	6	5	6	1	-	-
2000.....	13	12	10	16	10	13	19	5	-	2	-	1
1999.....	11	13	16	9	13	14	8	4	2	4	-	1

Black, female

2003.....	5	4	3	3	1	2	7	4	2	-	-	-
2002.....	4	2	5	7	6	4	9	6	4	-	-	-
2001.....	2	4	3	5	4	7	9	5	2	1	-	-
2000.....	2	-	-	5	12	11	5	7	2	1	-	-
1999.....	3	7	-	4	7	12	9	7	5	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

American Indian 1/, both sexes

2003.....	2	-	-	-	-	1	2	-	-	1	-	-
2002.....	1	2	2	1	2	3	3	2	-	-	-	-
2001.....	-	-	1	-	1	1	3	2	-	-	-	-
2000.....	-	3	2	2	-	3	-	-	-	-	-	-
1999.....	-	-	1	1	4	1	-	-	1	1	-	-

American Indian 1/, male

2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	2	-	1	-	1	2	2	-	-	-	-
2001.....	-	-	1	-	-	1	1	-	-	-	-	-
2000.....	-	2	2	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	2	-	-	-	-	1	-	-

American Indian 1/, female

2003.....	1	-	-	-	-	1	1	-	-	1	-	-
2002.....	1	-	2	-	2	2	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	2	2	-	-	-	-
2000.....	-	1	-	1	-	2	-	-	-	-	-	-
1999.....	-	-	1	-	2	1	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Asian or Pacific Islander 2/, both sexes

2003.....	2	3	5	9	5	17	6	8	4	-	-	-
2002.....	2	3	5	7	8	11	12	8	3	1	1	-
2001.....	1	3	4	3	10	9	11	13	1	-	-	-
2000.....	1	2	5	2	6	15	12	7	2	-	-	-
1999.....	4	6	5	6	8	15	12	10	5	-	-	-

Asian or Pacific Islander 2/, male

2003.....	2	1	3	5	4	12	3	3	2	-	-	-
2002.....	1	3	4	3	3	4	6	4	1	-	-	-
2001.....	-	1	4	3	8	5	6	7	1	-	-	-
2000.....	1	2	2	1	2	8	8	4	1	-	-	-
1999.....	2	5	4	4	7	10	9	7	2	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	2	2	4	1	5	3	5	2	-	-	-
2002.....	1	-	1	4	5	7	6	4	2	1	1	-
2001.....	1	2	-	-	2	4	5	6	-	-	-	-
2000.....	-	-	3	1	4	7	4	3	1	-	-	-
1999.....	2	1	1	2	1	5	3	3	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

All races, both sexes

2003.....	9	11	10	12	14	17	23	15	5	1	-	-
2002.....	10	7	16	21	20	27	15	8	5	2	-	-
2001.....	13	12	8	13	24	20	20	14	2	3	-	-
2000.....	11	7	14	18	18	25	18	10	5	2	-	-
1999.....	17	14	7	12	22	22	15	15	5	-	-	-

All races, male

2003.....	8	8	6	5	6	9	10	6	2	1	-	-
2002.....	8	6	9	10	12	16	9	4	1	1	-	-
2001.....	7	6	1	9	13	7	9	5	1	1	-	-
2000.....	6	2	7	8	6	11	12	5	-	1	-	-
1999.....	13	9	4	8	11	11	6	6	3	-	-	-

All races, female

2003.....	1	3	4	7	8	8	13	9	3	-	-	-
2002.....	2	1	7	11	8	11	6	4	4	1	-	-
2001.....	6	6	7	4	11	13	11	9	1	2	-	-
2000.....	5	5	7	10	12	14	6	5	5	1	-	-
1999.....	4	5	3	4	11	11	9	9	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

White, both sexes

2003.....	6	8	4	7	12	10	15	12	5	1	-	-
2002.....	6	3	11	7	11	19	15	6	3	1	-	-
2001.....	6	5	5	9	12	14	12	13	2	2	-	-
2000.....	5	1	6	11	10	14	10	8	4	1	-	-
1999.....	7	10	2	10	11	15	12	8	1	-	-	-

White, male

2003.....	6	6	2	2	5	6	7	5	2	1	-	-
2002.....	5	3	5	3	7	12	9	3	-	-	-	-
2001.....	2	3	-	6	5	4	6	5	1	1	-	-
2000.....	1	1	3	6	4	4	7	3	-	1	-	-
1999.....	6	6	1	6	6	6	6	2	-	-	-	-

White, female

2003.....	-	2	2	5	7	4	8	7	3	-	-	-
2002.....	1	-	6	4	4	7	6	3	3	1	-	-
2001.....	4	2	5	3	7	10	6	8	1	1	-	-
2000.....	4	-	3	5	6	10	3	5	4	-	-	-
1999.....	1	4	1	4	5	9	6	6	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Black, both sexes

2003.....	2	3	3	3	-	4	3	2	-	-	-	-
2002.....	2	3	2	6	5	3	-	2	2	-	-	-
2001.....	6	5	2	4	8	3	4	1	-	1	-	-
2000.....	6	6	6	4	5	5	7	1	-	1	-	-
1999.....	8	2	2	2	8	5	2	3	1	-	-	-

Black, male

2003.....	2	2	1	2	-	2	1	-	-	-	-	-
2002.....	1	2	1	4	2	1	-	1	1	-	-	-
2001.....	4	2	1	3	4	1	1	-	-	-	-	-
2000.....	5	1	3	1	1	4	4	1	-	-	-	-
1999.....	6	2	1	2	5	3	-	2	1	-	-	-

Black, female

2003.....	-	1	2	1	-	2	2	2	-	-	-	-
2002.....	1	1	1	2	3	2	-	1	1	-	-	-
2001.....	2	3	1	1	4	2	3	1	-	1	-	-
2000.....	1	5	3	3	4	1	3	-	-	1	-	-
1999.....	2	-	1	-	3	2	2	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

American Indian 1/, both sexes

2003.....	-	-	-	-	1	1	-	1	-	-	-	-
2002.....	-	-	1	3	1	1	-	-	-	-	-	-
2001.....	-	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	1	2	1	1	1	-	1	-	-	-
1999.....	1	1	-	-	1	-	-	1	2	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	1	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	2	-	-	-

American Indian 1/, female

2003.....	-	-	-	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	2	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	1	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Asian or Pacific Islander 2/, both sexes

2003.....	1	-	3	2	1	2	5	-	-	-	-	-
2002.....	2	1	2	5	3	4	-	-	-	1	-	-
2001.....	1	1	1	-	3	3	4	-	-	-	-	-
2000.....	-	-	1	1	2	5	-	1	-	-	-	-
1999.....	1	1	3	-	2	2	1	3	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	3	1	1	1	2	-	-	-	-	-
2002.....	2	1	2	2	2	2	-	-	-	1	-	-
2001.....	1	1	-	-	3	2	2	-	-	-	-	-
2000.....	-	-	1	-	-	2	-	1	-	-	-	-
1999.....	1	-	2	-	-	2	-	2	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	-	-	1	-	1	3	-	-	-	-	-
2002.....	-	-	-	3	1	2	-	-	-	-	-	-
2001.....	-	-	1	-	-	1	2	-	-	-	-	-
2000.....	-	-	-	1	2	3	-	-	-	-	-	-
1999.....	-	1	1	-	2	-	1	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Scarlet fever and erysipelas (A38,A46)

All races, both sexes

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	1	-	-	1	-	1	-

All races, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

All races, female

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	1	1	-	-	1	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Scarlet fever and erysipelas (A38,A46)

White, both sexes

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	1	-	1	-

White, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

White, female

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	1	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Scarlet fever and erysipelas (A38,A46)

Black, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Black, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningococcal infection (A39)

All races, both sexes

2003.....	10	9	2	3	6	4	5	3	-	-	-	-
2002.....	9	9	4	3	4	3	7	2	-	3	-	-
2001.....	4	11	4	7	8	7	1	6	1	2	-	-
2000.....	16	6	3	3	5	7	8	2	4	1	-	-
1999.....	6	3	1	9	-	7	6	3	4	1	-	-

All races, male

2003.....	5	4	1	1	1	-	1	1	-	-	-	-
2002.....	5	2	2	1	1	1	2	1	-	1	-	-
2001.....	2	5	2	2	3	2	1	1	-	1	-	-
2000.....	6	1	1	-	2	4	3	-	-	1	-	-
1999.....	3	-	-	5	-	2	2	-	1	1	-	-

All races, female

2003.....	5	5	1	2	5	4	4	2	-	-	-	-
2002.....	4	7	2	2	3	2	5	1	-	2	-	-
2001.....	2	6	2	5	5	5	-	5	1	1	-	-
2000.....	10	5	2	3	3	3	5	2	4	-	-	-
1999.....	3	3	1	4	-	5	4	3	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningococcal infection (A39)

White, both sexes

2003.....	5	6	1	3	6	4	5	3	-	-	-	-
2002.....	7	7	4	2	4	3	7	2	-	3	-	-
2001.....	4	10	1	6	8	6	1	5	1	2	-	-
2000.....	12	5	3	2	5	7	8	2	4	1	-	-
1999.....	5	3	1	9	-	7	5	3	4	1	-	-

White, male

2003.....	3	2	-	1	1	-	1	1	-	-	-	-
2002.....	3	2	2	-	1	1	2	1	-	1	-	-
2001.....	2	4	1	2	3	1	1	-	-	1	-	-
2000.....	3	1	1	-	2	4	3	-	-	1	-	-
1999.....	2	-	-	5	-	2	2	-	1	1	-	-

White, female

2003.....	2	4	1	2	5	4	4	2	-	-	-	-
2002.....	4	5	2	2	3	2	5	1	-	2	-	-
2001.....	2	6	-	4	5	5	-	5	1	1	-	-
2000.....	9	4	2	2	3	3	5	2	4	-	-	-
1999.....	3	3	1	4	-	5	3	3	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningococcal infection (A39)

American Indian 1/, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Septicemia (A40-A41)

All races, both sexes

2003.....	1,241	1,631	2,020	2,459	3,511	5,096	5,836	5,178	3,064	1,112	189	3
2002.....	1,170	1,499	1,861	2,589	3,747	5,195	5,815	5,026	2,988	1,111	199	3
2001.....	1,140	1,392	1,719	2,409	3,589	5,000	5,351	4,856	3,001	1,020	192	-
2000.....	1,064	1,260	1,639	2,303	3,401	4,745	5,193	4,900	2,949	1,079	216	1
1999.....	954	1,178	1,536	2,352	3,398	4,686	5,024	4,854	2,984	1,133	195	-

All races, male

2003.....	657	864	1,023	1,299	1,724	2,377	2,517	1,920	894	247	28	2
2002.....	629	767	981	1,292	1,862	2,476	2,552	1,857	834	247	35	3
2001.....	604	728	878	1,227	1,817	2,402	2,360	1,788	869	225	32	-
2000.....	600	657	856	1,182	1,685	2,249	2,233	1,690	812	199	35	1
1999.....	513	613	799	1,218	1,701	2,252	2,174	1,712	817	236	21	-

All races, female

2003.....	584	767	997	1,160	1,787	2,719	3,319	3,258	2,170	865	161	1
2002.....	541	732	880	1,297	1,885	2,719	3,263	3,169	2,154	864	164	-
2001.....	536	664	841	1,182	1,772	2,598	2,991	3,068	2,132	795	160	-
2000.....	464	603	783	1,121	1,716	2,496	2,960	3,210	2,137	880	181	-
1999.....	441	565	737	1,134	1,697	2,434	2,850	3,142	2,167	897	174	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Septicemia (A40-A41)

White, both sexes

2003.....	826	1,180	1,486	1,817	2,770	4,218	4,992	4,448	2,625	927	136	2
2002.....	782	1,070	1,340	1,948	2,990	4,259	4,943	4,323	2,569	925	151	2
2001.....	765	980	1,260	1,830	2,847	4,123	4,585	4,179	2,596	861	146	-
2000.....	724	897	1,169	1,716	2,702	3,859	4,364	4,206	2,546	875	163	1
1999.....	650	813	1,115	1,769	2,642	3,799	4,236	4,110	2,546	957	143	-

White, male

2003.....	448	616	762	965	1,379	1,981	2,169	1,661	767	207	21	1
2002.....	422	560	730	972	1,510	2,025	2,168	1,614	709	205	24	2
2001.....	409	502	658	937	1,441	1,995	2,034	1,541	737	193	20	-
2000.....	404	474	623	892	1,372	1,833	1,904	1,448	692	161	25	1
1999.....	351	418	565	917	1,330	1,858	1,837	1,443	690	195	18	-

White, female

2003.....	378	564	724	852	1,391	2,237	2,823	2,787	1,858	720	115	1
2002.....	360	510	610	976	1,480	2,234	2,775	2,709	1,860	720	127	-
2001.....	356	478	602	893	1,406	2,128	2,551	2,638	1,859	668	126	-
2000.....	320	423	546	824	1,330	2,026	2,460	2,758	1,854	714	138	-
1999.....	299	395	550	852	1,312	1,941	2,399	2,667	1,856	762	125	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Septicemia (A40-A41)

Black, both sexes

2003.....	378	400	491	582	665	784	762	662	392	172	49	1
2002.....	357	405	471	584	676	853	798	638	382	169	47	1
2001.....	339	385	429	532	666	801	713	630	366	149	43	-
2000.....	316	328	440	540	649	824	766	635	380	187	53	-
1999.....	274	333	389	524	699	800	725	682	396	163	52	-

Black, male

2003.....	195	227	239	304	305	351	319	232	113	36	7	1
2002.....	193	195	228	287	316	400	340	213	108	36	10	1
2001.....	176	213	204	261	330	368	295	225	111	27	11	-
2000.....	183	169	218	265	290	387	297	218	109	34	10	-
1999.....	147	175	219	275	339	350	306	240	107	37	3	-

Black, female

2003.....	183	173	252	278	360	433	443	430	279	136	42	-
2002.....	164	210	243	297	360	453	458	425	274	133	37	-
2001.....	163	172	225	271	336	433	418	405	255	122	32	-
2000.....	133	159	222	275	359	437	469	417	271	153	43	-
1999.....	127	158	170	249	360	450	419	442	289	126	49	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Septicemia (A40-A41)

American Indian 1/, both sexes

2003.....	14	28	17	23	23	18	24	13	6	3	3	-
2002.....	13	11	16	20	26	20	20	12	8	4	1	-
2001.....	11	12	9	18	21	10	11	9	4	1	-	-
2000.....	9	18	11	22	11	19	15	12	4	6	-	-
1999.....	16	12	17	18	15	18	14	11	8	4	-	-

American Indian 1/, male

2003.....	5	12	8	10	7	4	7	4	1	1	-	-
2002.....	8	4	7	13	10	11	12	2	1	2	1	-
2001.....	5	7	4	10	10	7	6	4	2	1	-	-
2000.....	5	8	6	10	5	6	8	2	2	1	-	-
1999.....	9	8	5	10	8	7	7	2	3	2	-	-

American Indian 1/, female

2003.....	9	16	9	13	16	14	17	9	5	2	3	-
2002.....	5	7	9	7	16	9	8	10	7	2	-	-
2001.....	6	5	5	8	11	3	5	5	2	-	-	-
2000.....	4	10	5	12	6	13	7	10	2	5	-	-
1999.....	7	4	12	8	7	11	7	9	5	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Septicemia (A40-A41)

Asian or Pacific Islander 2/, both sexes

2003.....	23	23	26	37	53	76	58	55	41	10	1	-
2002.....	18	13	34	37	55	63	54	53	29	13	-	-
2001.....	25	15	21	29	55	66	42	38	35	9	3	-
2000.....	15	17	19	25	39	43	48	47	19	11	-	-
1999.....	14	20	15	41	42	69	49	51	34	9	-	-

Asian or Pacific Islander 2/, male

2003.....	9	9	14	20	33	41	22	23	13	3	-	-
2002.....	6	8	16	20	26	40	32	28	16	4	-	-
2001.....	14	6	12	19	36	32	25	18	19	4	1	-
2000.....	8	6	9	15	18	23	24	22	9	3	-	-
1999.....	6	12	10	16	24	37	24	27	17	2	-	-

Asian or Pacific Islander 2/, female

2003.....	14	14	12	17	20	35	36	32	28	7	1	-
2002.....	12	5	18	17	29	23	22	25	13	9	-	-
2001.....	11	9	9	10	19	34	17	20	16	5	2	-
2000.....	7	11	10	10	21	20	24	25	10	8	-	-
1999.....	8	8	5	25	18	32	25	24	17	7	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Syphilis (A50-A53)

All races, both sexes

2003.....	3	2	1	3	3	6	2	5	3	1	-	-
2002.....	4	2	3	4	2	5	7	6	3	-	-	-
2001.....	1	3	1	3	2	4	9	2	2	1	-	-
2000.....	4	2	1	1	6	7	6	6	3	1	-	-
1999.....	2	1	3	2	4	6	4	2	1	1	-	-

All races, male

2003.....	3	1	1	2	3	2	2	2	1	-	-	-
2002.....	4	1	1	4	1	3	1	2	1	-	-	-
2001.....	1	2	1	1	2	2	3	1	-	-	-	-
2000.....	4	2	1	1	1	3	4	3	2	-	-	-
1999.....	1	1	2	2	3	-	-	1	-	-	-	-

All races, female

2003.....	-	1	-	1	-	4	-	3	2	1	-	-
2002.....	-	1	2	-	1	2	6	4	2	-	-	-
2001.....	-	1	-	2	-	2	6	1	2	1	-	-
2000.....	-	-	-	-	5	4	2	3	1	1	-	-
1999.....	1	-	1	-	1	6	4	1	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Syphilis (A50-A53)

White, both sexes

2003.....	1	2	-	2	-	4	1	2	3	1	-	-
2002.....	2	-	1	2	1	1	4	4	2	-	-	-
2001.....	1	2	1	1	1	4	6	2	-	1	-	-
2000.....	1	-	-	-	3	3	4	4	3	1	-	-
1999.....	2	-	3	2	1	2	2	-	-	1	-	-

White, male

2003.....	1	1	-	2	-	2	1	1	1	-	-	-
2002.....	2	-	-	2	1	1	-	2	-	-	-	-
2001.....	1	1	1	-	1	2	1	1	-	-	-	-
2000.....	1	-	-	-	-	-	2	1	2	-	-	-
1999.....	1	-	2	2	-	-	-	-	-	-	-	-

White, female

2003.....	-	1	-	-	-	2	-	1	2	1	-	-
2002.....	-	-	1	-	-	-	4	2	2	-	-	-
2001.....	-	1	-	1	-	2	5	1	-	1	-	-
2000.....	-	-	-	-	3	3	2	3	1	1	-	-
1999.....	1	-	1	-	1	2	2	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Syphilis (A50-A53)

Black, both sexes

2003.....	2	-	1	1	3	2	1	2	-	-	-	-
2002.....	2	2	2	2	1	4	3	2	1	-	-	-
2001.....	-	-	-	2	1	-	3	-	2	-	-	-
2000.....	3	2	1	1	3	4	2	2	-	-	-	-
1999.....	-	1	-	-	3	3	2	2	1	-	-	-

Black, male

2003.....	2	-	1	-	3	-	1	-	-	-	-	-
2002.....	2	1	1	2	-	2	1	-	1	-	-	-
2001.....	-	-	-	1	1	-	2	-	-	-	-	-
2000.....	3	2	1	1	1	3	2	2	-	-	-	-
1999.....	-	1	-	-	3	-	-	1	-	-	-	-

Black, female

2003.....	-	-	-	1	-	2	-	2	-	-	-	-
2002.....	-	1	1	-	1	2	2	2	-	-	-	-
2001.....	-	-	-	1	-	-	1	-	2	-	-	-
2000.....	-	-	-	-	2	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	3	2	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Syphilis (A50-A53)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

All races, both sexes

2003.....	1,206	709	376	330	368	323	220	87	24	4	-	-
2002.....	1,139	644	428	369	393	352	235	136	34	7	-	-
2001.....	959	586	428	407	429	375	238	120	41	5	-	1
2000.....	850	492	412	410	435	401	259	123	35	10	-	1
1999.....	635	438	348	427	454	389	262	145	45	5	2	-

All races, male

2003.....	903	506	223	169	183	147	102	30	8	2	-	-
2002.....	868	448	254	193	180	174	96	56	11	3	-	-
2001.....	737	406	237	207	189	156	107	42	13	2	-	1
2000.....	638	338	239	202	194	198	113	55	11	4	-	1
1999.....	487	298	193	217	209	185	132	61	20	1	-	-

All races, female

2003.....	303	203	153	161	185	176	118	57	16	2	-	-
2002.....	271	196	174	176	213	178	139	80	23	4	-	-
2001.....	222	180	191	200	240	219	131	78	28	3	-	-
2000.....	212	154	173	208	241	203	146	68	24	6	-	-
1999.....	148	140	155	210	245	204	130	84	25	4	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

White, both sexes

2003.....	955	517	278	244	289	262	185	74	21	3	-	-
2002.....	920	487	317	278	311	298	205	121	30	4	-	-
2001.....	734	447	331	314	341	322	205	104	34	5	-	-
2000.....	635	350	315	320	358	350	222	110	30	9	-	1
1999.....	481	323	269	330	364	326	235	124	40	4	2	-

White, male

2003.....	723	378	171	130	145	116	89	27	7	2	-	-
2002.....	705	342	185	154	148	152	87	51	10	-	-	-
2001.....	580	319	189	170	161	143	98	41	10	2	-	-
2000.....	487	246	186	156	161	175	98	49	11	3	-	1
1999.....	371	224	160	177	177	159	126	52	17	1	-	-

White, female

2003.....	232	139	107	114	144	146	96	47	14	1	-	-
2002.....	215	145	132	124	163	146	118	70	20	4	-	-
2001.....	154	128	142	144	180	179	107	63	24	3	-	-
2000.....	148	104	129	164	197	175	124	61	19	6	-	-
1999.....	110	99	109	153	187	167	109	72	23	3	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

Black, both sexes

2003.....	219	148	74	49	43	38	21	8	2	1	-	-
2002.....	189	115	72	57	46	35	15	7	4	2	-	-
2001.....	188	103	67	47	46	35	19	12	4	-	-	1
2000.....	164	93	50	47	47	32	25	9	2	-	-	-
1999.....	118	75	45	60	53	38	17	17	2	1	-	-

Black, male

2003.....	161	106	39	22	21	17	5	3	1	-	-	-
2002.....	141	83	41	32	18	14	3	-	1	2	-	-
2001.....	128	65	36	18	16	9	4	-	2	-	-	1
2000.....	120	63	25	26	22	13	12	4	-	-	-	-
1999.....	91	45	21	21	19	15	2	8	1	-	-	-

Black, female

2003.....	58	42	35	27	22	21	16	5	1	1	-	-
2002.....	48	32	31	25	28	21	12	7	3	-	-	-
2001.....	60	38	31	29	30	26	15	12	2	-	-	-
2000.....	44	30	25	21	25	19	13	5	2	-	-	-
1999.....	27	30	24	39	34	23	15	9	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

American Indian 1/, both sexes

2003.....	10	9	7	3	2	1	-	-	-	-	-	-
2002.....	8	10	3	1	3	-	-	-	-	-	-	-
2001.....	12	1	4	5	3	2	-	-	-	-	-	-
2000.....	13	4	6	4	1	1	2	-	1	-	-	-
1999.....	6	2	2	2	1	2	3	1	-	-	-	-

American Indian 1/, male

2003.....	5	7	3	2	1	1	-	-	-	-	-	-
2002.....	6	5	2	1	1	-	-	-	-	-	-	-
2001.....	10	-	3	2	-	-	-	-	-	-	-	-
2000.....	9	1	4	1	-	-	1	-	-	-	-	-
1999.....	4	1	1	1	-	-	2	-	-	-	-	-

American Indian 1/, female

2003.....	5	2	4	1	1	-	-	-	-	-	-	-
2002.....	2	5	1	-	2	-	-	-	-	-	-	-
2001.....	2	1	1	3	3	2	-	-	-	-	-	-
2000.....	4	3	2	3	1	1	1	-	1	-	-	-
1999.....	2	1	1	1	1	2	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

Asian or Pacific Islander 2/, both sexes

2003.....	22	35	17	34	34	22	14	5	1	-	-	-
2002.....	22	32	36	33	33	19	15	8	-	1	-	-
2001.....	25	35	26	41	39	16	14	4	3	-	-	-
2000.....	38	45	41	39	29	18	10	4	2	1	-	-
1999.....	30	38	32	35	36	23	7	3	3	-	-	-

Asian or Pacific Islander 2/, male

2003.....	14	15	10	15	16	13	8	-	-	-	-	-
2002.....	16	18	26	6	13	8	6	5	-	1	-	-
2001.....	19	22	9	17	12	4	5	1	1	-	-	-
2000.....	22	28	24	19	11	10	2	2	-	1	-	-
1999.....	21	28	11	18	13	11	2	1	2	-	-	-

Asian or Pacific Islander 2/, female

2003.....	8	20	7	19	18	9	6	5	1	-	-	-
2002.....	6	14	10	27	20	11	9	3	-	-	-	-
2001.....	6	13	17	24	27	12	9	3	2	-	-	-
2000.....	16	17	17	20	18	8	8	2	2	-	-	-
1999.....	9	10	21	17	23	12	5	2	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races, both sexes

2003.....	1,806	1,017	500	277	161	71	24	11	3	-	-	3
2002.....	1,794	885	462	275	126	71	25	8	3	1	-	4
2001.....	1,609	821	496	258	126	68	20	5	3	2	-	2
2000.....	1,591	797	442	250	147	69	16	4	3	2	-	1
1999.....	1,467	777	372	261	137	52	18	7	5	1	1	1

All races, male

2003.....	1,425	816	399	209	127	55	20	10	2	-	-	3
2002.....	1,446	710	373	223	100	53	18	4	-	1	-	4
2001.....	1,299	673	378	187	105	55	18	4	1	2	-	2
2000.....	1,288	655	358	199	114	54	11	1	1	-	-	1
1999.....	1,198	646	302	211	106	39	11	6	1	1	-	1

All races, female

2003.....	381	201	101	68	34	16	4	1	1	-	-	-
2002.....	348	175	89	52	26	18	7	4	3	-	-	-
2001.....	310	148	118	71	21	13	2	1	2	-	-	-
2000.....	303	142	84	51	33	15	5	3	2	2	-	-
1999.....	269	131	70	50	31	13	7	1	4	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2003.....	771	474	213	111	78	41	7	5	3	-	-	2
2002.....	728	425	218	106	45	30	12	5	3	1	-	3
2001.....	695	406	194	102	61	31	6	3	2	2	-	-
2000.....	706	391	197	110	56	33	7	2	1	-	-	1
1999.....	700	346	173	119	52	25	12	3	4	-	1	-

White, male

2003.....	663	399	181	87	66	35	7	4	2	-	-	2
2002.....	622	365	181	93	41	21	8	3	-	1	-	3
2001.....	616	350	157	82	58	27	6	2	1	2	-	-
2000.....	613	336	163	96	44	28	5	-	-	-	-	1
1999.....	613	305	156	103	42	22	7	2	-	-	-	-

White, female

2003.....	108	75	32	24	12	6	-	1	1	-	-	-
2002.....	106	60	37	13	4	9	4	2	3	-	-	-
2001.....	79	56	37	20	3	4	-	1	1	-	-	-
2000.....	93	55	34	14	12	5	2	2	1	-	-	-
1999.....	87	41	17	16	10	3	5	1	4	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2003.....	1,021	535	282	164	83	29	17	6	-	-	-	1
2002.....	1,043	450	238	165	78	41	13	3	-	-	-	1
2001.....	897	410	293	151	63	37	13	1	1	-	-	2
2000.....	872	395	241	137	87	35	9	2	2	2	-	-
1999.....	753	429	196	139	83	27	6	3	1	1	-	1

Black, male

2003.....	750	411	214	122	61	19	13	6	-	-	-	1
2002.....	804	336	187	126	58	32	10	1	-	-	-	1
2001.....	667	319	214	104	46	28	11	1	-	-	-	2
2000.....	666	310	191	101	67	25	6	1	1	-	-	-
1999.....	573	339	144	105	63	17	4	3	1	1	-	1

Black, female

2003.....	271	124	68	42	22	10	4	-	-	-	-	-
2002.....	239	114	51	39	20	9	3	2	-	-	-	-
2001.....	230	91	79	47	17	9	2	-	1	-	-	-
2000.....	206	85	50	36	20	10	3	1	1	2	-	-
1999.....	180	90	52	34	20	10	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Asian or Pacific Islander 2/, both sexes

2003.....	8	4	4	1	-	-	-	-	-	-	-	-
2002.....	20	7	6	4	3	-	-	-	-	-	-	-
2001.....	9	4	8	3	2	-	-	1	-	-	-	-
2000.....	7	10	4	1	3	-	-	-	-	-	-	-
1999.....	9	2	1	2	1	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	6	3	3	-	-	-	-	-	-	-	-	-
2002.....	19	7	5	4	1	-	-	-	-	-	-	-
2001.....	8	3	6	1	1	-	-	1	-	-	-	-
2000.....	6	9	4	-	2	-	-	-	-	-	-	-
1999.....	7	2	1	2	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	1	1	1	-	-	-	-	-	-	-	-
2002.....	1	-	1	-	2	-	-	-	-	-	-	-
2001.....	1	1	2	2	1	-	-	-	-	-	-	-
2000.....	1	1	-	1	1	-	-	-	-	-	-	-
1999.....	2	-	-	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races, both sexes

2003.....	659	590	546	577	767	913	802	594	335	113	20	-
2002.....	513	445	444	520	703	859	735	580	289	112	16	-
2001.....	356	338	433	537	606	698	632	483	288	98	18	-
2000.....	308	323	384	451	623	739	636	513	265	93	28	-
1999.....	444	427	471	652	791	927	812	633	330	138	22	-

All races, male

2003.....	441	376	313	316	395	430	386	232	123	24	4	-
2002.....	348	273	259	290	373	429	356	230	110	30	3	-
2001.....	215	209	240	275	290	350	306	185	101	21	-	-
2000.....	194	179	204	253	344	367	298	216	97	22	4	-
1999.....	247	220	262	351	409	460	367	253	99	28	1	-

All races, female

2003.....	218	214	233	261	372	483	416	362	212	89	16	-
2002.....	165	172	185	230	330	430	379	350	179	82	13	-
2001.....	141	129	193	262	316	348	326	298	187	77	18	-
2000.....	114	144	180	198	279	372	338	297	168	71	24	-
1999.....	197	207	209	301	382	467	445	380	231	110	21	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

White, both sexes

2003.....	521	469	440	491	670	781	716	543	310	108	19	-
2002.....	394	337	355	444	604	749	663	536	260	104	14	-
2001.....	271	262	347	447	518	615	565	438	261	88	14	-
2000.....	232	263	297	380	533	655	571	470	235	77	25	-
1999.....	318	329	362	535	673	798	712	565	288	126	18	-

White, male

2003.....	353	302	251	269	339	375	345	212	115	24	4	-
2002.....	272	203	207	254	323	375	320	214	99	27	2	-
2001.....	166	165	195	229	251	312	278	169	92	19	-	-
2000.....	147	150	161	211	300	318	269	197	81	17	3	-
1999.....	182	178	199	289	355	404	306	221	83	26	-	-

White, female

2003.....	168	167	189	222	331	406	371	331	195	84	15	-
2002.....	122	134	148	190	281	374	343	322	161	77	12	-
2001.....	105	97	152	218	267	303	287	269	169	69	14	-
2000.....	85	113	136	169	233	337	302	273	154	60	22	-
1999.....	136	151	163	246	318	394	406	344	205	100	18	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2003.....	114	95	73	56	63	95	64	38	24	4	1	-
2002.....	100	90	62	50	72	89	44	36	27	8	2	-
2001.....	69	63	67	66	73	56	49	35	20	8	4	-
2000.....	65	54	65	55	74	65	57	33	26	15	3	-
1999.....	104	83	88	97	97	105	81	54	37	12	4	-

Black, male

2003.....	70	55	43	36	40	36	30	14	8	-	-	-
2002.....	65	62	37	26	36	40	19	13	10	3	1	-
2001.....	40	35	36	38	31	22	20	10	7	2	-	-
2000.....	41	27	35	33	37	36	24	10	12	5	1	-
1999.....	53	34	49	48	47	46	48	24	12	2	1	-

Black, female

2003.....	44	40	30	20	23	59	34	24	16	4	1	-
2002.....	35	28	25	24	36	49	25	23	17	5	1	-
2001.....	29	28	31	28	42	34	29	25	13	6	4	-
2000.....	24	27	30	22	37	29	33	23	14	10	2	-
1999.....	51	49	39	49	50	59	33	30	25	10	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

American Indian 1/, both sexes

2003.....	12	8	6	6	2	5	3	2	1	1	-	-
2002.....	5	6	6	9	2	3	5	-	-	-	-	-
2001.....	2	2	5	5	3	4	3	3	1	-	-	-
2000.....	4	3	9	1	2	2	1	1	-	-	-	-
1999.....	4	3	6	7	6	3	4	2	1	-	-	-

American Indian 1/, male

2003.....	7	5	3	3	-	3	-	1	-	-	-	-
2002.....	3	3	3	2	1	1	2	-	-	-	-	-
2001.....	1	1	1	1	1	2	1	2	-	-	-	-
2000.....	4	1	2	-	-	1	-	1	-	-	-	-
1999.....	1	-	3	5	2	-	3	1	1	-	-	-

American Indian 1/, female

2003.....	5	3	3	3	2	2	3	1	1	1	-	-
2002.....	2	3	3	7	1	2	3	-	-	-	-	-
2001.....	1	1	4	4	2	2	2	1	1	-	-	-
2000.....	-	2	7	1	2	1	1	-	-	-	-	-
1999.....	3	3	3	2	4	3	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Asian or Pacific Islander 2/, both sexes

2003.....	12	18	27	24	32	32	19	11	-	-	-	-
2002.....	14	12	21	17	25	18	23	8	2	-	-	-
2001.....	14	11	14	19	12	23	15	7	6	2	-	-
2000.....	7	3	13	15	14	17	7	9	4	1	-	-
1999.....	18	12	15	13	15	21	15	12	4	-	-	-

Asian or Pacific Islander 2/, male

2003.....	11	14	16	8	16	16	11	5	-	-	-	-
2002.....	8	5	12	8	13	13	15	3	1	-	-	-
2001.....	8	8	8	7	7	14	7	4	2	-	-	-
2000.....	2	1	6	9	7	12	5	8	4	-	-	-
1999.....	11	8	11	9	5	10	10	7	3	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	4	11	16	16	16	8	6	-	-	-	-
2002.....	6	7	9	9	12	5	8	5	1	-	-	-
2001.....	6	3	6	12	5	9	8	3	4	2	-	-
2000.....	5	2	7	6	7	5	2	1	-	1	-	-
1999.....	7	4	4	4	10	11	5	5	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

All races, both sexes

2003.....	30,291	42,315	53,377	63,141	78,107	89,086	78,531	50,789	22,660	5,827	770	12
2002.....	30,342	41,487	51,904	63,598	81,159	89,565	77,497	50,298	22,323	5,772	789	7
2001.....	30,363	39,827	50,396	64,172	82,846	89,967	75,478	49,680	21,660	5,684	727	8
2000.....	29,231	38,930	50,075	65,843	84,288	90,655	74,444	49,155	21,564	5,653	764	11
1999.....	28,269	38,637	50,430	66,814	85,524	89,874	72,896	48,041	20,702	5,492	779	9

All races, male

2003.....	15,815	22,671	29,256	35,048	42,745	47,516	39,866	23,531	9,124	2,006	217	8
2002.....	15,774	22,407	28,708	35,374	44,711	47,734	39,218	23,345	9,106	1,952	218	5
2001.....	15,694	21,390	27,854	35,900	45,799	47,951	38,175	23,140	8,628	1,951	235	7
2000.....	14,972	20,912	27,515	36,730	46,461	48,284	37,623	22,714	8,506	1,850	190	10
1999.....	14,483	20,891	27,942	37,315	47,229	48,184	36,876	22,147	8,287	1,826	224	6

All races, female

2003.....	14,476	19,644	24,121	28,093	35,362	41,570	38,665	27,258	13,536	3,821	553	4
2002.....	14,568	19,080	23,196	28,224	36,448	41,831	38,279	26,953	13,217	3,820	571	2
2001.....	14,669	18,437	22,542	28,272	37,047	42,016	37,303	26,540	13,032	3,733	492	1
2000.....	14,259	18,018	22,560	29,113	37,827	42,371	36,821	26,441	13,058	3,803	574	1
1999.....	13,786	17,746	22,488	29,499	38,295	41,690	36,020	25,894	12,415	3,666	555	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

White, both sexes

2003.....	24,033	35,179	45,119	54,000	68,223	79,174	70,674	46,109	20,524	5,156	636	11
2002.....	24,336	34,425	43,715	54,326	71,132	79,678	69,847	45,714	20,162	5,114	662	4
2001.....	24,559	33,051	42,552	54,773	72,631	80,120	67,885	45,136	19,572	5,040	602	4
2000.....	23,735	32,440	42,319	56,673	73,897	80,805	67,321	44,566	19,488	4,995	630	7
1999.....	22,976	32,083	42,485	57,500	75,203	80,023	65,986	43,518	18,705	4,869	650	7

White, male

2003.....	12,546	18,836	24,690	29,981	37,449	42,224	35,860	21,387	8,219	1,735	179	8
2002.....	12,665	18,583	24,049	30,313	39,275	42,403	35,246	21,201	8,171	1,723	175	3
2001.....	12,648	17,679	23,476	30,684	40,191	42,585	34,197	20,922	7,722	1,710	196	4
2000.....	12,061	17,275	23,182	31,634	40,650	42,849	33,865	20,503	7,611	1,600	149	7
1999.....	11,668	17,289	23,479	32,130	41,549	42,755	33,311	19,908	7,428	1,579	187	4

White, female

2003.....	11,487	16,343	20,429	24,019	30,774	36,950	34,814	24,722	12,305	3,421	457	3
2002.....	11,671	15,842	19,666	24,013	31,857	37,275	34,601	24,513	11,991	3,391	487	1
2001.....	11,911	15,372	19,076	24,089	32,440	37,535	33,688	24,214	11,850	3,330	406	-
2000.....	11,674	15,165	19,137	25,039	33,247	37,956	33,456	24,063	11,877	3,395	481	-
1999.....	11,308	14,794	19,006	25,370	33,654	37,268	32,675	23,610	11,277	3,290	463	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

Black, both sexes

2003.....	5,257	6,043	6,972	7,652	8,164	8,142	6,464	3,863	1,807	566	116	-
2002.....	5,001	5,954	6,871	7,845	8,400	8,300	6,364	3,856	1,832	552	110	3
2001.....	4,850	5,741	6,582	7,951	8,596	8,251	6,339	3,803	1,801	571	116	4
2000.....	4,710	5,499	6,620	7,797	8,839	8,338	6,023	3,897	1,784	591	118	3
1999.....	4,522	5,635	6,798	7,955	8,809	8,414	5,898	3,930	1,717	531	121	1

Black, male

2003.....	2,804	3,314	3,876	4,248	4,432	4,328	3,274	1,726	740	217	27	-
2002.....	2,615	3,254	3,929	4,329	4,602	4,459	3,290	1,775	776	187	38	2
2001.....	2,593	3,164	3,669	4,468	4,754	4,486	3,296	1,843	745	204	37	3
2000.....	2,528	3,114	3,715	4,371	4,950	4,584	3,183	1,832	742	219	38	2
1999.....	2,442	3,148	3,845	4,467	4,847	4,644	3,030	1,905	715	198	33	1

Black, female

2003.....	2,453	2,729	3,096	3,404	3,732	3,814	3,190	2,137	1,067	349	89	-
2002.....	2,386	2,700	2,942	3,516	3,798	3,841	3,074	2,081	1,056	365	72	1
2001.....	2,257	2,577	2,913	3,483	3,842	3,765	3,043	1,960	1,056	367	79	1
2000.....	2,182	2,385	2,905	3,426	3,889	3,754	2,840	2,065	1,042	372	80	1
1999.....	2,080	2,487	2,953	3,488	3,962	3,770	2,868	2,025	1,002	333	88	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

American Indian 1/, both sexes

2003.....	189	220	245	257	300	283	211	108	48	16	4	-
2002.....	184	219	266	274	315	252	208	104	53	16	4	-
2001.....	177	209	261	298	298	263	190	109	50	13	-	-
2000.....	132	190	227	251	288	248	164	103	46	16	4	-
1999.....	131	187	219	271	264	228	177	91	36	10	1	-

American Indian 1/, male

2003.....	92	109	130	146	155	135	99	55	21	10	2	-
2002.....	94	123	142	141	166	129	107	41	13	3	1	-
2001.....	88	108	136	159	169	127	98	46	23	6	-	-
2000.....	72	106	120	136	155	133	80	44	20	9	1	-
1999.....	62	90	115	146	146	120	88	44	17	7	1	-

American Indian 1/, female

2003.....	97	111	115	111	145	148	112	53	27	6	2	-
2002.....	90	96	124	133	149	123	101	63	40	13	3	-
2001.....	89	101	125	139	129	136	92	63	27	7	-	-
2000.....	60	84	107	115	133	115	84	59	26	7	3	-
1999.....	69	97	104	125	118	108	89	47	19	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

Asian or Pacific Islander 2/, both sexes

2003.....	812	873	1,041	1,232	1,420	1,487	1,182	709	281	89	14	1
2002.....	821	889	1,052	1,153	1,312	1,335	1,078	624	276	90	13	-
2001.....	777	826	1,001	1,150	1,321	1,333	1,064	632	237	60	9	-
2000.....	654	801	909	1,122	1,264	1,264	936	589	246	51	12	1
1999.....	640	732	928	1,088	1,248	1,209	835	502	244	82	7	1

Asian or Pacific Islander 2/, male

2003.....	373	412	560	673	709	829	633	363	144	44	9	-
2002.....	400	447	588	591	668	743	575	328	146	39	4	-
2001.....	365	439	573	589	685	753	584	329	138	31	2	-
2000.....	311	417	498	589	706	718	495	335	133	22	2	1
1999.....	311	364	503	572	687	665	447	290	127	42	3	1

Asian or Pacific Islander 2/, female

2003.....	439	461	481	559	711	658	549	346	137	45	5	1
2002.....	421	442	464	562	644	592	503	296	130	51	9	-
2001.....	412	387	428	561	636	580	480	303	99	29	7	-
2000.....	343	384	411	533	558	546	441	254	113	29	10	-
1999.....	329	368	425	516	561	544	388	212	117	40	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races, both sexes

2003.....	686	867	959	950	924	955	737	543	242	78	16	1
2002.....	670	875	936	846	1,052	973	748	505	262	77	13	-
2001.....	649	861	879	893	1,019	1,004	783	502	240	67	5	1
2000.....	660	770	872	942	1,017	933	693	488	249	74	13	-
1999.....	612	753	863	889	1,007	993	751	501	220	64	15	-

All races, male

2003.....	575	694	729	678	626	589	410	255	108	20	6	1
2002.....	534	681	711	619	698	590	407	245	101	24	4	-
2001.....	521	687	685	631	696	598	427	232	83	18	2	1
2000.....	516	618	652	667	659	569	370	217	96	19	-	-
1999.....	474	568	663	629	679	605	402	229	79	14	1	-

All races, female

2003.....	111	173	230	272	298	366	327	288	134	58	10	-
2002.....	136	194	225	227	354	383	341	260	161	53	9	-
2001.....	128	174	194	262	323	406	356	270	157	49	3	-
2000.....	144	152	220	275	358	364	323	271	153	55	13	-
1999.....	138	185	200	260	328	388	349	272	141	50	14	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, both sexes

2003.....	502	706	761	781	780	832	651	504	216	73	16	1
2002.....	491	683	745	689	938	877	681	458	247	67	11	-
2001.....	497	664	697	720	881	900	720	463	222	66	5	-
2000.....	475	581	674	776	868	832	629	449	229	67	11	-
1999.....	430	560	688	737	866	873	687	457	213	60	14	-

White, male

2003.....	432	562	564	553	515	506	356	234	97	19	6	1
2002.....	390	537	560	499	619	536	368	221	97	19	3	-
2001.....	400	525	537	509	594	529	391	213	77	18	2	-
2000.....	358	467	499	541	555	495	332	202	86	18	-	-
1999.....	337	418	530	521	587	524	362	205	76	13	1	-

White, female

2003.....	70	144	197	228	265	326	295	270	119	54	10	-
2002.....	101	146	185	190	319	341	313	237	150	48	8	-
2001.....	97	139	160	211	287	371	329	250	145	48	3	-
2000.....	117	114	175	235	313	337	297	247	143	49	11	-
1999.....	93	142	158	216	279	349	325	252	137	47	13	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, both sexes

2003.....	140	139	170	148	116	87	63	30	20	3	-	-
2002.....	149	150	169	125	93	66	48	33	12	7	2	-
2001.....	124	173	146	142	117	82	49	25	16	1	-	1
2000.....	156	164	166	135	121	82	56	26	15	6	2	-
1999.....	155	165	157	130	120	99	54	37	5	2	1	-

Black, male

2003.....	113	116	144	112	92	58	41	15	9	-	-	-
2002.....	124	122	136	97	64	36	28	18	3	3	1	-
2001.....	100	142	122	100	87	54	26	12	5	-	-	1
2000.....	137	133	132	106	87	61	34	8	8	-	-	-
1999.....	118	127	119	93	78	70	33	19	1	1	-	-

Black, female

2003.....	27	23	26	36	24	29	22	15	11	3	-	-
2002.....	25	28	33	28	29	30	20	15	9	4	1	-
2001.....	24	31	24	42	30	28	23	13	11	1	-	-
2000.....	19	31	34	29	34	21	22	18	7	6	2	-
1999.....	37	38	38	37	42	29	21	18	4	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

American Indian 1/, both sexes

2003.....	5	4	5	1	5	7	3	1	1	-	-	-
2002.....	4	9	6	8	6	3	3	-	-	-	-	-
2001.....	4	6	7	4	4	4	2	1	-	-	-	-
2000.....	4	6	4	7	10	3	-	3	-	-	-	-
1999.....	7	3	4	6	5	2	-	1	-	-	-	-

American Indian 1/, male

2003.....	4	2	2	1	3	2	1	-	1	-	-	-
2002.....	3	5	4	3	4	3	2	-	-	-	-	-
2001.....	3	6	6	4	4	1	2	1	-	-	-	-
2000.....	3	4	1	5	7	2	-	1	-	-	-	-
1999.....	4	3	2	4	3	-	-	1	-	-	-	-

American Indian 1/, female

2003.....	1	2	3	-	2	5	2	1	-	-	-	-
2002.....	1	4	2	5	2	-	1	-	-	-	-	-
2001.....	1	-	1	-	-	3	-	-	-	-	-	-
2000.....	1	2	3	2	3	1	-	2	-	-	-	-
1999.....	3	-	2	2	2	2	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Asian or Pacific Islander 2/, both sexes

2003.....	39	18	23	20	23	29	20	8	5	2	-	-
2002.....	26	33	16	24	15	27	16	14	3	3	-	-
2001.....	24	18	29	27	17	18	12	13	2	-	-	-
2000.....	25	19	28	24	18	16	8	10	5	1	-	-
1999.....	20	25	14	16	16	19	10	6	2	2	-	-

Asian or Pacific Islander 2/, male

2003.....	26	14	19	12	16	23	12	6	1	1	-	-
2002.....	17	17	11	20	11	15	9	6	1	2	-	-
2001.....	18	14	20	18	11	14	8	6	1	-	-	-
2000.....	18	14	20	15	10	11	4	6	2	1	-	-
1999.....	15	20	12	11	11	11	7	4	2	-	-	-

Asian or Pacific Islander 2/, female

2003.....	13	4	4	8	7	6	8	2	4	1	-	-
2002.....	9	16	5	4	4	12	7	8	2	1	-	-
2001.....	6	4	9	9	6	4	4	7	1	-	-	-
2000.....	7	5	8	9	8	5	4	4	3	-	-	-
1999.....	5	5	2	5	5	8	3	2	-	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

All races, both sexes

2003.....	881	1,389	1,551	1,714	1,905	1,895	1,497	806	306	63	5	-
2002.....	887	1,286	1,477	1,736	1,955	1,909	1,480	766	295	65	4	1
2001.....	916	1,114	1,440	1,765	2,014	1,941	1,366	757	317	56	7	1
2000.....	870	1,180	1,430	1,711	1,960	1,817	1,346	764	274	55	7	-
1999.....	793	1,139	1,392	1,739	1,955	1,784	1,202	755	279	72	8	1

All races, male

2003.....	757	1,175	1,308	1,385	1,466	1,415	1,049	476	156	22	-	-
2002.....	759	1,099	1,222	1,394	1,558	1,437	1,017	452	127	23	1	-
2001.....	787	940	1,197	1,436	1,553	1,423	942	452	153	18	2	1
2000.....	746	988	1,181	1,349	1,482	1,337	896	437	122	23	4	-
1999.....	684	969	1,112	1,396	1,529	1,328	808	421	127	22	3	1

All races, female

2003.....	124	214	243	329	439	480	448	330	150	41	5	-
2002.....	128	187	255	342	397	472	463	314	168	42	3	1
2001.....	129	174	243	329	461	518	424	305	164	38	5	-
2000.....	124	192	249	362	478	480	450	327	152	32	3	-
1999.....	109	170	280	343	426	456	394	334	152	50	5	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

White, both sexes

2003.....	682	1,145	1,279	1,426	1,689	1,710	1,355	751	286	55	5	-
2002.....	725	1,044	1,218	1,442	1,708	1,698	1,353	704	276	61	3	-
2001.....	718	860	1,168	1,455	1,714	1,741	1,251	700	287	53	7	1
2000.....	674	932	1,150	1,441	1,705	1,599	1,218	709	253	51	7	-
1999.....	617	895	1,114	1,448	1,691	1,564	1,102	694	259	67	5	-

White, male

2003.....	601	990	1,098	1,159	1,317	1,292	972	446	147	17	-	-
2002.....	633	906	1,026	1,185	1,378	1,301	935	418	121	22	-	-
2001.....	624	741	985	1,200	1,350	1,290	867	420	135	17	2	1
2000.....	588	791	975	1,158	1,308	1,186	830	412	112	19	4	-
1999.....	542	773	905	1,191	1,339	1,171	751	383	115	18	2	-

White, female

2003.....	81	155	181	267	372	418	383	305	139	38	5	-
2002.....	92	138	192	257	330	397	418	286	155	39	3	-
2001.....	94	119	183	255	364	451	384	280	152	36	5	-
2000.....	86	141	175	283	397	413	388	297	141	32	3	-
1999.....	75	122	209	257	352	393	351	311	144	49	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

Black, both sexes

2003.....	181	222	245	249	188	158	116	50	16	7	-	-
2002.....	145	222	235	259	222	185	100	53	13	3	1	1
2001.....	172	225	245	274	266	177	102	51	23	3	-	-
2000.....	184	228	255	244	222	203	106	48	19	4	-	-
1999.....	166	225	252	260	231	200	82	51	17	5	2	1

Black, male

2003.....	139	166	188	189	126	104	59	26	7	4	-	-
2002.....	111	180	174	182	163	118	63	30	3	1	1	-
2001.....	141	176	191	204	178	119	67	27	13	1	-	-
2000.....	150	185	186	172	149	139	54	21	8	4	-	-
1999.....	133	179	190	179	165	143	42	30	10	4	-	1

Black, female

2003.....	42	56	57	60	62	54	57	24	9	3	-	-
2002.....	34	42	61	77	59	67	37	23	10	2	-	1
2001.....	31	49	54	70	88	58	35	24	10	2	-	-
2000.....	34	43	69	72	73	64	52	27	11	-	-	-
1999.....	33	46	62	81	66	57	40	21	7	1	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of esophagus (C15)												
American Indian 1/, both sexes												
2003.....	5	6	6	9	6	5	6	-	-	-	-	-
2002.....	4	2	7	5	7	7	9	-	2	1	-	-
2001.....	4	7	7	8	8	7	3	2	1	-	-	-
2000.....	3	3	8	6	7	3	6	1	1	-	-	-
1999.....	1	7	7	5	6	5	3	-	-	-	-	-
American Indian 1/, male												
2003.....	5	5	2	8	5	5	3	-	-	-	-	-
2002.....	4	1	7	4	5	4	8	-	2	-	-	-
2001.....	3	6	4	6	7	4	1	1	1	-	-	-
2000.....	3	1	8	5	7	2	4	-	1	-	-	-
1999.....	1	6	4	5	6	4	1	-	-	-	-	-
American Indian 1/, female												
2003.....	-	1	4	1	1	-	3	-	-	-	-	-
2002.....	-	1	-	1	2	3	1	-	-	1	-	-
2001.....	1	1	3	2	1	3	2	1	-	-	-	-
2000.....	-	2	-	1	-	1	2	1	-	-	-	-
1999.....	-	1	3	-	-	1	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

Asian or Pacific Islander 2/, both sexes

2003.....	13	16	21	30	22	22	20	5	4	1	-	-
2002.....	13	18	17	30	18	19	18	9	4	-	-	-
2001.....	22	22	20	28	26	16	10	4	6	-	-	-
2000.....	9	17	17	20	26	12	16	6	1	-	-	-
1999.....	9	12	19	26	27	15	15	10	3	-	1	-

Asian or Pacific Islander 2/, male

2003.....	12	14	20	29	18	14	15	4	2	1	-	-
2002.....	11	12	15	23	12	14	11	4	1	-	-	-
2001.....	19	17	17	26	18	10	7	4	4	-	-	-
2000.....	5	11	12	14	18	10	8	4	1	-	-	-
1999.....	8	11	13	21	19	10	14	8	2	-	1	-

Asian or Pacific Islander 2/, female

2003.....	1	2	1	1	4	8	5	1	2	-	-	-
2002.....	2	6	2	7	6	5	7	5	3	-	-	-
2001.....	3	5	3	2	8	6	3	-	2	-	-	-
2000.....	4	6	5	6	8	2	8	2	-	-	-	-
1999.....	1	1	6	5	8	5	1	2	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

All races, both sexes

2003.....	717	830	1,044	1,211	1,582	1,805	1,744	1,263	619	161	14	-
2002.....	663	841	990	1,179	1,637	1,903	1,753	1,344	664	153	24	-
2001.....	675	751	1,020	1,308	1,713	2,004	1,789	1,216	620	149	16	-
2000.....	629	753	1,051	1,317	1,737	2,009	1,824	1,379	641	188	26	-
1999.....	651	808	1,052	1,359	1,793	2,064	1,796	1,277	607	170	29	-

All races, male

2003.....	462	574	708	810	1,016	1,061	919	574	238	51	4	-
2002.....	440	556	659	804	1,063	1,118	946	628	258	38	10	-
2001.....	443	514	724	858	1,070	1,194	954	550	222	36	5	-
2000.....	413	506	722	860	1,106	1,201	943	617	247	56	8	-
1999.....	443	558	694	893	1,112	1,250	989	568	223	52	8	-

All races, female

2003.....	255	256	336	401	566	744	825	689	381	110	10	-
2002.....	223	285	331	375	574	785	807	716	406	115	14	-
2001.....	232	237	296	450	643	810	835	666	398	113	11	-
2000.....	216	247	329	457	631	808	881	762	394	132	18	-
1999.....	208	250	358	466	681	814	807	709	384	118	21	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

White, both sexes

2003.....	497	622	788	922	1,227	1,420	1,403	1,026	498	114	8	-
2002.....	453	624	730	864	1,260	1,496	1,406	1,086	537	115	14	-
2001.....	473	565	750	964	1,350	1,595	1,441	1,008	503	116	11	-
2000.....	463	561	791	988	1,348	1,571	1,465	1,129	525	149	13	-
1999.....	451	583	765	1,066	1,393	1,636	1,462	1,039	497	131	19	-

White, male

2003.....	323	436	531	610	794	843	749	459	200	36	4	-
2002.....	307	427	491	605	834	904	773	521	207	29	4	-
2001.....	312	385	536	627	850	973	771	463	183	30	3	-
2000.....	311	379	557	660	874	952	770	510	205	45	4	-
1999.....	312	415	508	713	880	1,004	807	452	184	42	6	-

White, female

2003.....	174	186	257	312	433	577	654	567	298	78	4	-
2002.....	146	197	239	259	426	592	633	565	330	86	10	-
2001.....	161	180	214	337	500	622	670	545	320	86	8	-
2000.....	152	182	234	328	474	619	695	619	320	104	9	-
1999.....	139	168	257	353	513	632	655	587	313	89	13	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

Black, both sexes

2003.....	161	152	198	213	250	287	246	157	96	35	6	-
2002.....	158	155	185	232	290	307	241	200	98	31	9	-
2001.....	150	130	198	251	258	294	258	161	94	27	5	-
2000.....	130	149	189	243	286	320	266	197	86	30	12	-
1999.....	151	173	205	214	298	328	257	194	86	32	9	-

Black, male

2003.....	108	106	135	148	165	164	118	71	22	9	-	-
2002.....	106	102	121	148	179	160	118	75	41	7	5	-
2001.....	106	88	135	172	161	163	133	59	25	6	2	-
2000.....	83	100	125	148	171	176	125	80	27	8	3	-
1999.....	99	112	133	135	178	188	139	89	28	7	2	-

Black, female

2003.....	53	46	63	65	85	123	128	86	74	26	6	-
2002.....	52	53	64	84	111	147	123	125	57	24	4	-
2001.....	44	42	63	79	97	131	125	102	69	21	3	-
2000.....	47	49	64	95	115	144	141	117	59	22	9	-
1999.....	52	61	72	79	120	140	118	105	58	25	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

American Indian 1/, both sexes

2003.....	8	3	7	11	11	12	4	5	2	1	-	-
2002.....	10	5	11	11	10	13	12	4	2	-	-	-
2001.....	3	7	8	6	11	18	10	4	4	-	-	-
2000.....	5	7	9	15	10	9	6	6	2	-	-	-
1999.....	2	6	5	12	11	9	7	1	4	-	-	-

American Indian 1/, male

2003.....	6	2	4	9	8	5	3	3	2	-	-	-
2002.....	5	1	4	7	5	5	7	3	-	-	-	-
2001.....	2	7	6	4	6	7	7	2	4	-	-	-
2000.....	4	5	4	8	5	6	2	3	1	-	-	-
1999.....	1	3	4	8	7	5	6	1	3	-	-	-

American Indian 1/, female

2003.....	2	1	3	2	3	7	1	2	-	1	-	-
2002.....	5	4	7	4	5	8	5	1	2	-	-	-
2001.....	1	-	2	2	5	11	3	2	-	-	-	-
2000.....	1	2	5	7	5	3	4	3	1	-	-	-
1999.....	1	3	1	4	4	4	1	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

Asian or Pacific Islander 2/, both sexes

2003.....	51	53	51	65	94	86	91	75	23	11	-	-
2002.....	42	57	64	72	77	87	94	54	27	7	1	-
2001.....	49	49	64	87	94	97	80	43	19	6	-	-
2000.....	31	36	62	71	93	109	87	47	28	9	1	-
1999.....	47	46	77	67	91	91	70	43	20	7	1	-

Asian or Pacific Islander 2/, male

2003.....	25	30	38	43	49	49	49	41	14	6	-	-
2002.....	22	26	43	44	45	49	48	29	10	2	1	-
2001.....	23	34	47	55	53	51	43	26	10	-	-	-
2000.....	15	22	36	44	56	67	46	24	14	3	1	-
1999.....	31	28	49	37	47	53	37	26	8	3	-	-

Asian or Pacific Islander 2/, female

2003.....	26	23	13	22	45	37	42	34	9	5	-	-
2002.....	20	31	21	28	32	38	46	25	17	5	-	-
2001.....	26	15	17	32	41	46	37	17	9	6	-	-
2000.....	16	14	26	27	37	42	41	23	14	6	-	-
1999.....	16	18	28	30	44	38	33	17	12	4	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races, both sexes

2003.....	2,718	3,631	4,673	5,775	7,159	8,720	8,611	6,590	3,547	1,024	135	1
2002.....	2,818	3,694	4,727	5,901	7,491	8,826	8,672	6,453	3,433	1,075	128	1
2001.....	2,804	3,610	4,374	5,997	7,794	8,988	8,539	6,618	3,482	1,028	131	-
2000.....	2,581	3,397	4,543	5,913	8,188	9,410	8,643	6,735	3,518	1,061	148	-
1999.....	2,434	3,309	4,658	6,095	8,154	9,473	8,456	6,672	3,503	973	138	-

All races, male

2003.....	1,544	2,117	2,714	3,393	3,953	4,433	4,038	2,565	1,112	245	27	1
2002.....	1,606	2,185	2,850	3,455	4,125	4,536	3,968	2,508	1,024	267	24	1
2001.....	1,545	2,116	2,536	3,479	4,294	4,568	3,883	2,520	1,068	251	36	-
2000.....	1,466	1,989	2,619	3,407	4,517	4,852	3,908	2,571	1,012	260	30	-
1999.....	1,374	1,932	2,689	3,431	4,452	4,857	3,897	2,525	1,081	236	24	-

All races, female

2003.....	1,174	1,514	1,959	2,382	3,206	4,287	4,573	4,025	2,435	779	108	-
2002.....	1,212	1,509	1,877	2,446	3,366	4,290	4,704	3,945	2,409	808	104	-
2001.....	1,259	1,494	1,838	2,518	3,500	4,420	4,656	4,098	2,414	777	95	-
2000.....	1,115	1,408	1,924	2,506	3,671	4,558	4,735	4,164	2,506	801	118	-
1999.....	1,060	1,377	1,969	2,664	3,702	4,616	4,559	4,147	2,422	737	114	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, both sexes

2003.....	2,060	2,898	3,836	4,853	6,057	7,625	7,705	5,911	3,212	914	114	1
2002.....	2,240	2,985	3,873	4,933	6,460	7,731	7,686	5,857	3,073	955	110	1
2001.....	2,211	2,906	3,609	5,004	6,722	7,831	7,618	6,015	3,146	908	106	-
2000.....	2,050	2,737	3,752	4,991	7,094	8,322	7,710	6,090	3,180	933	129	-
1999.....	1,935	2,676	3,861	5,145	7,083	8,370	7,574	6,030	3,198	864	109	-

White, male

2003.....	1,195	1,730	2,245	2,891	3,407	3,911	3,641	2,326	1,002	215	23	1
2002.....	1,287	1,796	2,360	2,932	3,615	3,998	3,521	2,270	916	237	18	1
2001.....	1,254	1,735	2,132	2,949	3,766	4,036	3,486	2,288	962	217	30	-
2000.....	1,182	1,607	2,181	2,906	3,955	4,332	3,485	2,327	905	218	23	-
1999.....	1,119	1,574	2,254	2,943	3,948	4,303	3,529	2,279	990	211	22	-

White, female

2003.....	865	1,168	1,591	1,962	2,650	3,714	4,064	3,585	2,210	699	91	-
2002.....	953	1,189	1,513	2,001	2,845	3,733	4,165	3,587	2,157	718	92	-
2001.....	957	1,171	1,477	2,055	2,956	3,795	4,132	3,727	2,184	691	76	-
2000.....	868	1,130	1,571	2,085	3,139	3,990	4,225	3,763	2,275	715	106	-
1999.....	816	1,102	1,607	2,202	3,135	4,067	4,045	3,751	2,208	653	87	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Black, both sexes

2003.....	551	622	707	774	942	909	779	563	291	93	18	-
2002.....	493	596	712	817	857	934	836	509	309	94	13	-
2001.....	495	611	651	842	894	994	779	509	296	105	23	-
2000.....	464	538	687	787	945	942	801	561	295	117	18	-
1999.....	414	551	705	819	927	965	785	576	268	92	24	-

Black, male

2003.....	300	329	396	417	477	427	328	184	90	28	2	-
2002.....	269	319	401	438	414	455	373	197	86	19	4	-
2001.....	244	328	333	449	436	448	328	188	86	27	5	-
2000.....	251	312	369	420	479	442	354	199	82	39	7	-
1999.....	218	319	379	426	435	480	315	207	77	15	-	-

Black, female

2003.....	251	293	311	357	465	482	451	379	201	65	16	-
2002.....	224	277	311	379	443	479	463	312	223	75	9	-
2001.....	251	283	318	393	458	546	451	321	210	78	18	-
2000.....	213	226	318	367	466	500	447	362	213	78	11	-
1999.....	196	232	326	393	492	485	470	369	191	77	24	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

American Indian 1/, both sexes

2003.....	15	13	25	24	25	33	18	13	9	2	-	-
2002.....	19	31	33	34	34	25	21	14	7	2	1	-
2001.....	16	17	25	22	39	20	22	12	4	-	-	-
2000.....	11	22	20	22	29	27	16	13	8	3	-	-
1999.....	18	14	16	18	19	24	22	14	7	1	-	-

American Indian 1/, male

2003.....	6	7	13	17	13	17	7	10	5	-	-	-
2002.....	12	23	24	17	16	11	10	3	1	-	1	-
2001.....	11	10	10	14	19	6	10	6	2	-	-	-
2000.....	5	17	13	14	16	15	7	6	3	1	-	-
1999.....	6	7	10	13	7	13	11	6	2	1	-	-

American Indian 1/, female

2003.....	9	6	12	7	12	16	11	3	4	2	-	-
2002.....	7	8	9	17	18	14	11	11	6	2	-	-
2001.....	5	7	15	8	20	14	12	6	2	-	-	-
2000.....	6	5	7	8	13	12	9	7	5	2	-	-
1999.....	12	7	6	5	12	11	11	8	5	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Asian or Pacific Islander 2/, both sexes

2003.....	92	98	105	124	135	153	109	103	35	15	3	-
2002.....	66	82	109	117	140	136	129	73	44	24	4	-
2001.....	82	76	89	129	139	143	120	82	36	15	2	-
2000.....	56	100	84	113	120	119	116	71	35	8	1	-
1999.....	67	68	76	113	125	114	75	52	30	16	5	-

Asian or Pacific Islander 2/, male

2003.....	43	51	60	68	56	78	62	45	15	2	2	-
2002.....	38	47	65	68	80	72	64	38	21	11	1	-
2001.....	36	43	61	67	73	78	59	38	18	7	1	-
2000.....	28	53	56	67	67	63	62	39	22	2	-	-
1999.....	31	32	46	49	62	61	42	33	12	9	2	-

Asian or Pacific Islander 2/, female

2003.....	49	47	45	56	79	75	47	58	20	13	1	-
2002.....	28	35	44	49	60	64	65	35	23	13	3	-
2001.....	46	33	28	62	66	65	61	44	18	8	1	-
2000.....	28	47	28	46	53	56	54	32	13	6	1	-
1999.....	36	36	30	64	63	53	33	19	18	7	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races, both sexes

2003.....	1,408	1,348	1,417	1,612	2,018	2,208	1,741	1,056	396	113	11	-
2002.....	1,245	1,208	1,350	1,618	1,963	2,049	1,679	1,094	419	80	16	-
2001.....	1,073	1,063	1,288	1,556	1,884	2,003	1,646	1,009	386	97	8	-
2000.....	944	1,040	1,216	1,526	1,913	2,011	1,547	943	359	81	4	-
1999.....	826	928	1,163	1,558	1,933	1,914	1,511	863	325	75	7	-

All races, male

2003.....	1,136	999	996	1,054	1,276	1,354	947	495	154	36	2	-
2002.....	1,015	899	959	1,100	1,267	1,212	900	463	155	22	2	-
2001.....	872	780	911	1,036	1,188	1,200	847	481	140	31	2	-
2000.....	729	761	846	1,039	1,207	1,196	856	407	114	26	1	-
1999.....	628	671	794	1,041	1,243	1,103	774	352	109	17	-	-

All races, female

2003.....	272	349	421	558	742	854	794	561	242	77	9	-
2002.....	230	309	391	518	696	837	779	631	264	58	14	-
2001.....	201	283	377	520	696	803	799	528	246	66	6	-
2000.....	215	279	370	487	706	815	691	536	245	55	3	-
1999.....	198	257	369	517	690	811	737	511	216	58	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

White, both sexes

2003.....	1,001	1,011	1,069	1,270	1,633	1,878	1,512	923	349	89	7	-
2002.....	862	875	1,053	1,298	1,611	1,766	1,471	973	354	67	11	-
2001.....	764	772	1,000	1,239	1,566	1,729	1,415	896	344	82	7	-
2000.....	654	756	924	1,204	1,580	1,717	1,376	830	325	70	2	-
1999.....	601	662	892	1,258	1,614	1,660	1,338	766	291	63	7	-

White, male

2003.....	801	752	761	840	1,060	1,162	833	435	132	28	1	-
2002.....	703	651	746	889	1,041	1,049	781	417	131	18	1	-
2001.....	617	561	719	851	1,009	1,037	746	426	129	27	2	-
2000.....	500	542	641	827	1,006	1,048	766	352	105	21	-	-
1999.....	448	481	613	846	1,052	960	698	321	103	14	-	-

White, female

2003.....	200	259	308	430	573	716	679	488	217	61	6	-
2002.....	159	224	307	409	570	717	690	556	223	49	10	-
2001.....	147	211	281	388	557	692	669	470	215	55	5	-
2000.....	154	214	283	377	574	669	610	478	220	49	2	-
1999.....	153	181	279	412	562	700	640	445	188	49	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, both sexes

2003.....	306	232	218	180	211	178	129	91	30	19	3	-
2002.....	258	231	161	184	204	176	122	77	43	6	2	-
2001.....	214	215	160	185	178	155	151	71	32	10	1	-
2000.....	199	171	173	189	202	162	101	71	21	7	2	-
1999.....	143	157	158	191	207	149	117	69	29	10	-	-

Black, male

2003.....	253	171	148	109	129	96	61	39	14	7	1	-
2002.....	209	170	115	129	135	97	67	24	12	3	-	-
2001.....	172	162	105	110	97	85	61	31	6	-	-	-
2000.....	152	134	122	131	124	77	54	34	4	3	1	-
1999.....	115	110	102	124	124	85	49	18	6	2	-	-

Black, female

2003.....	53	61	70	71	82	82	68	52	16	12	2	-
2002.....	49	61	46	55	69	79	55	53	31	3	2	-
2001.....	42	53	55	75	81	70	90	40	26	10	1	-
2000.....	47	37	51	58	78	85	47	37	17	4	1	-
1999.....	28	47	56	67	83	64	68	51	23	8	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

American Indian 1/, both sexes

2003.....	19	15	12	10	12	12	4	3	-	-	1	-
2002.....	11	9	11	13	14	6	11	5	2	2	-	-
2001.....	13	9	11	15	21	9	10	6	1	-	-	-
2000.....	13	14	9	11	10	6	11	8	1	1	-	-
1999.....	4	9	9	12	8	12	5	4	1	-	-	-

American Indian 1/, male

2003.....	15	7	8	7	7	10	2	1	-	-	-	-
2002.....	10	8	7	7	10	2	6	2	-	-	-	-
2001.....	12	6	7	10	17	5	5	1	-	-	-	-
2000.....	10	10	5	8	8	3	5	2	-	1	-	-
1999.....	3	6	7	5	3	7	2	1	-	-	-	-

American Indian 1/, female

2003.....	4	8	4	3	5	2	2	2	-	-	1	-
2002.....	1	1	4	6	4	4	5	3	2	2	-	-
2001.....	1	3	4	5	4	4	5	5	1	-	-	-
2000.....	3	4	4	3	2	3	6	6	1	-	-	-
1999.....	1	3	2	7	5	5	3	3	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Asian or Pacific Islander 2/, both sexes

2003.....	82	90	118	152	162	140	96	39	17	5	-	-
2002.....	114	93	125	123	134	101	75	39	20	5	3	-
2001.....	82	67	117	117	119	110	70	36	9	5	-	-
2000.....	78	99	110	122	121	126	59	34	12	3	-	-
1999.....	78	100	104	97	104	93	51	24	4	2	-	-

Asian or Pacific Islander 2/, male

2003.....	67	69	79	98	80	86	51	20	8	1	-	-
2002.....	93	70	91	75	81	64	46	20	12	1	1	-
2001.....	71	51	80	65	65	73	35	23	5	4	-	-
2000.....	67	75	78	73	69	68	31	19	5	1	-	-
1999.....	62	74	72	66	64	51	25	12	-	1	-	-

Asian or Pacific Islander 2/, female

2003.....	15	21	39	54	82	54	45	19	9	4	-	-
2002.....	21	23	34	48	53	37	29	19	8	4	2	-
2001.....	11	16	37	52	54	37	35	13	4	1	-	-
2000.....	11	24	32	49	52	58	28	15	7	2	-	-
1999.....	16	26	32	31	40	42	26	12	4	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

All races, both sexes

2003.....	1,649	2,327	2,993	3,550	4,554	5,038	4,670	2,897	1,264	293	32	-
2002.....	1,544	2,321	2,784	3,564	4,623	5,210	4,345	2,824	1,217	263	36	1
2001.....	1,577	2,111	2,757	3,525	4,633	4,987	4,335	2,861	1,214	293	42	1
2000.....	1,528	2,030	2,678	3,546	4,557	5,223	4,236	2,571	1,196	286	34	1
1999.....	1,466	2,039	2,668	3,632	4,661	4,931	4,002	2,673	1,120	309	40	1

All races, male

2003.....	992	1,391	1,715	1,924	2,327	2,358	2,044	992	327	55	8	-
2002.....	971	1,394	1,618	1,924	2,346	2,424	1,846	985	351	60	7	1
2001.....	964	1,269	1,605	1,921	2,294	2,360	1,831	984	310	64	7	1
2000.....	955	1,206	1,543	1,965	2,224	2,447	1,742	891	310	59	2	1
1999.....	907	1,204	1,517	1,976	2,371	2,281	1,659	910	311	59	3	-

All races, female

2003.....	657	936	1,278	1,626	2,227	2,680	2,626	1,905	937	238	24	-
2002.....	573	927	1,166	1,640	2,277	2,786	2,499	1,839	866	203	29	-
2001.....	613	842	1,152	1,604	2,339	2,627	2,504	1,877	904	229	35	-
2000.....	573	824	1,135	1,581	2,333	2,776	2,494	1,680	886	227	32	-
1999.....	559	835	1,151	1,656	2,290	2,650	2,343	1,763	809	250	37	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

White, both sexes

2003.....	1,287	1,924	2,475	3,013	3,923	4,449	4,152	2,586	1,150	259	25	-
2002.....	1,223	1,927	2,321	2,975	3,969	4,574	3,904	2,552	1,104	229	30	-
2001.....	1,293	1,753	2,354	2,989	3,998	4,398	3,876	2,581	1,072	261	33	1
2000.....	1,248	1,675	2,250	2,999	3,945	4,603	3,786	2,297	1,059	250	27	1
1999.....	1,184	1,660	2,237	3,087	4,053	4,339	3,574	2,397	1,003	282	30	1

White, male

2003.....	784	1,168	1,443	1,678	2,033	2,115	1,844	898	301	47	8	-
2002.....	785	1,166	1,378	1,645	2,048	2,177	1,679	900	320	52	5	-
2001.....	801	1,081	1,385	1,647	2,008	2,118	1,657	907	281	60	5	1
2000.....	787	1,003	1,314	1,709	1,949	2,197	1,570	812	272	49	2	1
1999.....	728	978	1,287	1,699	2,096	2,031	1,496	820	280	56	3	-

White, female

2003.....	503	756	1,032	1,335	1,890	2,334	2,308	1,688	849	212	17	-
2002.....	438	761	943	1,330	1,921	2,397	2,225	1,652	784	177	25	-
2001.....	492	672	969	1,342	1,990	2,280	2,219	1,674	791	201	28	-
2000.....	461	672	936	1,290	1,996	2,406	2,216	1,485	787	201	25	-
1999.....	456	682	950	1,388	1,957	2,308	2,078	1,577	723	226	27	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

Black, both sexes

2003.....	313	327	427	449	514	491	423	252	100	29	6	-
2002.....	259	334	396	475	550	516	351	226	95	32	6	1
2001.....	244	294	353	449	526	489	387	233	124	29	9	-
2000.....	246	307	365	455	523	528	369	228	118	32	5	-
1999.....	249	328	345	461	518	490	362	236	101	25	10	-

Black, male

2003.....	184	187	222	203	246	195	152	73	22	5	-	-
2002.....	159	195	206	227	252	197	138	62	22	8	2	1
2001.....	140	155	188	236	234	205	136	60	22	4	2	-
2000.....	151	178	195	214	236	215	138	60	27	8	-	-
1999.....	158	201	186	231	231	212	135	73	22	3	-	-

Black, female

2003.....	129	140	205	246	268	296	271	179	78	24	6	-
2002.....	100	139	190	248	298	319	213	164	73	24	4	-
2001.....	104	139	165	213	292	284	251	173	102	25	7	-
2000.....	95	129	170	241	287	313	231	168	91	24	5	-
1999.....	91	127	159	230	287	278	227	163	79	22	10	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

American Indian 1/, both sexes

2003.....	5	11	18	14	16	9	10	2	3	1	-	-
2002.....	14	12	10	11	23	12	13	5	1	-	-	-
2001.....	6	11	4	18	14	12	6	7	1	1	-	-
2000.....	8	16	9	10	14	8	10	3	2	1	1	-
1999.....	6	7	12	14	7	10	12	8	1	1	-	-

American Indian 1/, male

2003.....	3	4	13	8	7	3	5	1	-	1	-	-
2002.....	6	6	2	7	9	6	2	3	-	-	-	-
2001.....	3	5	3	10	7	3	3	2	-	-	-	-
2000.....	4	10	6	5	7	2	4	-	1	1	-	-
1999.....	3	3	6	8	6	4	5	2	-	-	-	-

American Indian 1/, female

2003.....	2	7	5	6	9	6	5	1	3	-	-	-
2002.....	8	6	8	4	14	6	11	2	1	-	-	-
2001.....	3	6	1	8	7	9	3	5	1	1	-	-
2000.....	4	6	3	5	7	6	6	3	1	-	1	-
1999.....	3	4	6	6	1	6	7	6	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

Asian or Pacific Islander 2/, both sexes

2003.....	44	65	73	74	101	89	85	57	11	4	1	-
2002.....	48	48	57	103	81	108	77	41	17	2	-	-
2001.....	34	53	46	69	95	88	66	40	17	2	-	-
2000.....	26	32	54	82	75	84	71	43	17	3	1	-
1999.....	27	44	74	70	83	92	54	32	15	1	-	-

Asian or Pacific Islander 2/, male

2003.....	21	32	37	35	41	45	43	20	4	2	-	-
2002.....	21	27	32	45	37	44	27	20	9	-	-	-
2001.....	20	28	29	28	45	34	35	15	7	-	-	-
2000.....	13	15	28	37	32	33	30	19	10	1	-	-
1999.....	18	22	38	38	38	34	23	15	9	-	-	-

Asian or Pacific Islander 2/, female

2003.....	23	33	36	39	60	44	42	37	7	2	1	-
2002.....	27	21	25	58	44	64	50	21	8	2	-	-
2001.....	14	25	17	41	50	54	31	25	10	2	-	-
2000.....	13	17	26	45	43	51	41	24	7	2	1	-
1999.....	9	22	36	32	45	58	31	17	6	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

All races, both sexes

2003.....	263	404	552	568	568	490	401	225	70	13	-	1
2002.....	269	412	492	508	576	588	381	172	76	5	-	1
2001.....	270	414	499	531	639	553	357	208	74	13	1	-
2000.....	270	384	489	574	645	595	401	191	73	11	1	-
1999.....	254	410	494	584	655	529	363	200	72	16	3	-

All races, male

2003.....	225	324	453	464	435	391	317	164	47	10	-	-
2002.....	219	339	399	395	437	455	300	126	56	1	-	1
2001.....	223	339	398	417	502	442	268	171	58	8	1	-
2000.....	218	319	407	441	504	451	314	155	51	8	-	-
1999.....	216	339	396	458	490	397	284	154	46	10	3	-

All races, female

2003.....	38	80	99	104	133	99	84	61	23	3	-	1
2002.....	50	73	93	113	139	133	81	46	20	4	-	-
2001.....	47	75	101	114	137	111	89	37	16	5	-	-
2000.....	52	65	82	133	141	144	87	36	22	3	1	-
1999.....	38	71	98	126	165	132	79	46	26	6	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

White, both sexes

2003.....	182	312	432	459	464	420	344	199	57	9	-	1
2002.....	192	317	383	416	485	494	333	158	71	4	-	-
2001.....	190	308	392	432	522	465	315	183	68	11	1	-
2000.....	191	279	364	466	521	501	358	171	70	10	1	-
1999.....	183	308	383	475	560	469	325	176	59	15	3	-

White, male

2003.....	154	247	358	365	353	331	268	144	39	7	-	-
2002.....	159	258	303	322	367	381	266	115	52	-	-	-
2001.....	155	251	316	339	412	366	232	152	55	7	1	-
2000.....	151	228	302	356	407	383	281	140	48	7	-	-
1999.....	158	252	310	372	417	352	249	133	36	9	3	-

White, female

2003.....	28	65	74	94	111	89	76	55	18	2	-	1
2002.....	33	59	80	94	118	113	67	43	19	4	-	-
2001.....	35	57	76	93	110	99	83	31	13	4	-	-
2000.....	40	51	62	110	114	118	77	31	22	3	1	-
1999.....	25	56	73	103	143	117	76	43	23	6	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

Black, both sexes

2003.....	77	91	115	101	96	65	50	24	11	3	-	-
2002.....	75	91	107	86	88	89	43	12	3	1	-	1
2001.....	77	101	99	88	103	77	34	20	6	1	-	-
2000.....	76	101	118	96	115	88	41	19	2	1	-	-
1999.....	69	98	107	103	87	57	34	15	10	1	-	-

Black, male

2003.....	67	76	91	92	77	56	43	18	6	2	-	-
2002.....	58	77	95	68	67	72	30	10	2	1	-	1
2001.....	65	83	78	69	78	66	28	15	3	-	-	-
2000.....	65	87	100	76	90	62	32	14	2	1	-	-
1999.....	56	85	84	81	67	42	32	13	7	1	-	-

Black, female

2003.....	10	15	24	9	19	9	7	6	5	1	-	-
2002.....	17	14	12	18	21	17	13	2	1	-	-	-
2001.....	12	18	21	19	25	11	6	5	3	1	-	-
2000.....	11	14	18	20	25	26	9	5	-	-	-	-
1999.....	13	13	23	22	20	15	2	2	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

American Indian 1/, both sexes

2003.....	3	-	2	1	2	2	1	1	-	-	-	-
2002.....	2	3	-	3	1	3	1	-	-	-	-	-
2001.....	1	1	2	6	4	2	3	1	-	1	-	-
2000.....	1	1	5	7	1	1	-	-	-	-	-	-
1999.....	2	1	2	2	2	-	-	3	-	-	-	-

American Indian 1/, male

2003.....	3	-	2	1	-	2	1	1	-	-	-	-
2002.....	2	3	-	3	1	-	1	-	-	-	-	-
2001.....	1	1	1	4	3	1	3	1	-	1	-	-
2000.....	1	1	3	5	-	1	-	-	-	-	-	-
1999.....	2	1	1	2	1	-	-	3	-	-	-	-

American Indian 1/, female

2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	3	-	-	-	-	-	-
2001.....	-	-	1	2	1	1	-	-	-	-	-	-
2000.....	-	-	2	2	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

Asian or Pacific Islander 2/, both sexes

2003.....	1	1	3	7	6	3	6	1	2	1	-	-
2002.....	-	1	2	3	2	2	4	2	2	-	-	-
2001.....	2	4	6	5	10	9	5	4	-	-	-	-
2000.....	2	3	2	5	8	5	2	1	1	-	-	-
1999.....	-	3	2	4	6	3	4	6	3	-	-	-

Asian or Pacific Islander 2/, male

2003.....	1	1	2	6	5	2	5	1	2	1	-	-
2002.....	-	1	1	2	2	2	3	1	2	-	-	-
2001.....	2	4	3	5	9	9	5	3	-	-	-	-
2000.....	1	3	2	4	7	5	1	1	1	-	-	-
1999.....	-	1	1	3	5	3	3	5	3	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	1	1	1	1	1	-	-	-	-	-
2002.....	-	-	1	1	-	-	1	1	-	-	-	-
2001.....	-	-	3	-	1	-	-	1	-	-	-	-
2000.....	1	-	-	1	1	-	1	-	-	-	-	-
1999.....	-	2	1	1	1	-	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races, both sexes

2003.....	7,757	12,920	18,036	22,414	26,972	28,031	20,588	10,076	3,319	617	76	2
2002.....	7,809	12,930	17,754	22,358	27,898	27,969	20,113	9,781	3,214	576	81	-
2001.....	7,786	12,595	17,182	22,850	28,375	27,723	18,977	9,623	3,064	587	57	4
2000.....	7,695	12,453	17,268	23,438	28,820	27,305	18,531	9,302	2,902	542	60	5
1999.....	7,518	12,165	17,179	23,239	28,609	26,556	17,705	8,945	2,766	549	79	2

All races, male

2003.....	4,654	7,758	10,589	13,218	15,717	15,934	11,130	5,099	1,485	255	26	2
2002.....	4,723	7,769	10,646	13,209	16,334	15,833	10,988	5,022	1,439	216	25	-
2001.....	4,703	7,523	10,350	13,798	16,901	15,968	10,438	5,026	1,370	237	23	4
2000.....	4,583	7,546	10,391	14,189	17,193	15,726	10,240	4,840	1,335	204	16	5
1999.....	4,505	7,465	10,548	14,186	17,101	15,461	9,936	4,739	1,314	234	28	2

All races, female

2003.....	3,103	5,162	7,447	9,196	11,255	12,097	9,458	4,977	1,834	362	50	-
2002.....	3,086	5,161	7,108	9,149	11,564	12,136	9,125	4,759	1,775	360	56	-
2001.....	3,083	5,072	6,832	9,052	11,474	11,755	8,539	4,597	1,694	350	34	-
2000.....	3,112	4,907	6,877	9,249	11,627	11,579	8,291	4,462	1,567	338	44	-
1999.....	3,013	4,700	6,631	9,053	11,508	11,095	7,769	4,206	1,452	315	51	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

White, both sexes

2003.....	6,215	10,870	15,565	19,553	24,005	25,257	18,731	9,234	3,023	549	66	2
2002.....	6,288	10,873	15,182	19,480	24,973	25,262	18,416	8,973	2,937	519	64	-
2001.....	6,346	10,669	14,837	19,915	25,344	25,025	17,272	8,812	2,799	514	45	2
2000.....	6,272	10,562	14,925	20,532	25,676	24,672	17,054	8,508	2,649	479	49	3
1999.....	6,125	10,324	14,766	20,284	25,556	23,939	16,192	8,166	2,523	498	65	1

White, male

2003.....	3,682	6,420	9,006	11,435	13,929	14,262	10,068	4,660	1,344	224	25	2
2002.....	3,743	6,441	8,950	11,416	14,558	14,193	9,981	4,589	1,307	200	23	-
2001.....	3,753	6,261	8,834	11,917	14,992	14,290	9,421	4,562	1,247	215	20	2
2000.....	3,641	6,274	8,905	12,313	15,221	14,059	9,356	4,398	1,206	177	11	3
1999.....	3,583	6,252	8,949	12,286	15,164	13,846	9,018	4,306	1,196	215	20	1

White, female

2003.....	2,533	4,450	6,559	8,118	10,076	10,995	8,663	4,574	1,679	325	41	-
2002.....	2,545	4,432	6,232	8,064	10,415	11,069	8,435	4,384	1,630	319	41	-
2001.....	2,593	4,408	6,003	7,998	10,352	10,735	7,851	4,250	1,552	299	25	-
2000.....	2,631	4,288	6,020	8,219	10,455	10,613	7,698	4,110	1,443	302	38	-
1999.....	2,542	4,072	5,817	7,998	10,392	10,093	7,174	3,860	1,327	283	45	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Black, both sexes

2003.....	1,340	1,814	2,165	2,446	2,486	2,275	1,486	662	235	57	9	-
2002.....	1,353	1,797	2,229	2,516	2,467	2,277	1,400	635	223	44	15	-
2001.....	1,275	1,692	2,034	2,534	2,583	2,275	1,367	643	199	61	11	2
2000.....	1,295	1,660	2,066	2,534	2,692	2,222	1,216	651	208	53	9	1
1999.....	1,252	1,643	2,127	2,553	2,595	2,257	1,258	662	198	38	13	-

Black, male

2003.....	858	1,197	1,389	1,528	1,507	1,355	851	343	114	24	1	-
2002.....	882	1,156	1,475	1,598	1,517	1,373	810	336	104	10	2	-
2001.....	854	1,113	1,321	1,648	1,642	1,420	822	377	89	15	3	2
2000.....	864	1,125	1,312	1,661	1,680	1,390	732	355	105	23	5	1
1999.....	841	1,099	1,422	1,653	1,641	1,387	763	377	93	13	7	-

Black, female

2003.....	482	617	776	918	979	920	635	319	121	33	8	-
2002.....	471	641	754	918	950	904	590	299	119	34	13	-
2001.....	421	579	713	886	941	855	545	266	110	46	8	-
2000.....	431	535	754	873	1,012	832	484	296	103	30	4	-
1999.....	411	544	705	900	954	870	495	285	105	25	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

American Indian 1/, both sexes

2003.....	41	67	67	94	97	74	56	19	8	1	-	-
2002.....	35	61	90	93	98	77	47	26	8	-	1	-
2001.....	39	51	82	90	91	82	52	17	8	2	-	-
2000.....	28	56	70	78	87	68	38	21	3	2	-	-
1999.....	27	59	83	89	94	57	46	12	6	2	1	-

American Indian 1/, male

2003.....	21	42	43	47	52	39	28	8	2	1	-	-
2002.....	20	41	53	44	50	45	27	13	2	-	-	-
2001.....	22	28	47	48	52	46	27	7	-	1	-	-
2000.....	15	31	45	40	43	44	24	9	3	1	-	-
1999.....	17	30	49	49	60	30	25	7	4	2	1	-

American Indian 1/, female

2003.....	20	25	24	47	45	35	28	11	6	-	-	-
2002.....	15	20	37	49	48	32	20	13	6	-	1	-
2001.....	17	23	35	42	39	36	25	10	8	1	-	-
2000.....	13	25	25	38	44	24	14	12	-	1	-	-
1999.....	10	29	34	40	34	27	21	5	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Asian or Pacific Islander 2/, both sexes

2003.....	161	169	239	321	384	425	315	161	53	10	1	-
2002.....	133	199	253	269	360	353	250	147	46	13	1	-
2001.....	126	183	229	311	357	341	286	151	58	10	1	-
2000.....	100	175	207	294	365	343	223	122	42	8	2	1
1999.....	114	139	203	313	364	303	209	105	39	11	-	1

Asian or Pacific Islander 2/, male

2003.....	93	99	151	208	229	278	183	88	25	6	-	-
2002.....	78	131	168	151	209	222	170	84	26	6	-	-
2001.....	74	121	148	185	215	212	168	80	34	6	-	-
2000.....	63	116	129	175	249	233	128	78	21	3	-	1
1999.....	64	84	128	198	236	198	130	49	21	4	-	1

Asian or Pacific Islander 2/, female

2003.....	68	70	88	113	155	147	132	73	28	4	1	-
2002.....	55	68	85	118	151	131	80	63	20	7	1	-
2001.....	52	62	81	126	142	129	118	71	24	4	1	-
2000.....	37	59	78	119	116	110	95	44	21	5	2	-
1999.....	50	55	75	115	128	105	79	56	18	7	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

All races, both sexes

2003.....	638	778	769	710	883	1,033	842	500	243	78	10	-
2002.....	640	659	692	748	868	954	794	512	233	58	17	1
2001.....	672	653	680	746	943	919	759	489	213	72	11	-
2000.....	665	627	676	753	884	887	737	495	252	52	4	-
1999.....	602	615	648	791	861	945	694	435	181	53	12	-

All races, male

2003.....	425	528	514	471	579	692	511	291	110	32	3	-
2002.....	403	445	470	500	575	609	469	264	115	21	6	-
2001.....	425	430	463	524	632	600	463	256	99	31	2	-
2000.....	448	420	446	495	572	559	454	277	100	13	-	-
1999.....	380	425	426	526	570	596	410	223	90	11	4	-

All races, female

2003.....	213	250	255	239	304	341	331	209	133	46	7	-
2002.....	237	214	222	248	293	345	325	248	118	37	11	1
2001.....	247	223	217	222	311	319	296	233	114	41	9	-
2000.....	217	207	230	258	312	328	283	218	152	39	4	-
1999.....	222	190	222	265	291	349	284	212	91	42	8	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

White, both sexes

2003.....	623	760	753	694	870	1,005	829	490	236	77	9	-
2002.....	628	642	677	731	843	938	783	501	227	55	16	1
2001.....	663	647	665	724	920	903	742	481	209	71	10	-
2000.....	654	611	662	736	862	871	717	483	243	50	4	-
1999.....	598	608	636	777	846	924	680	424	176	52	12	-

White, male

2003.....	417	519	505	461	577	679	504	288	107	32	3	-
2002.....	395	433	462	490	561	600	466	259	113	21	6	-
2001.....	421	427	456	513	622	592	458	253	99	31	2	-
2000.....	441	411	440	484	563	552	444	276	96	13	-	-
1999.....	377	420	418	519	564	584	404	219	88	10	4	-

White, female

2003.....	206	241	248	233	293	326	325	202	129	45	6	-
2002.....	233	209	215	241	282	338	317	242	114	34	10	1
2001.....	242	220	209	211	298	311	284	228	110	40	8	-
2000.....	213	200	222	252	299	319	273	207	147	37	4	-
1999.....	221	188	218	258	282	340	276	205	88	42	8	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

Black, both sexes

2003.....	8	12	11	16	10	20	8	7	5	-	1	-
2002.....	9	13	8	11	14	10	8	10	5	-	-	-
2001.....	6	5	8	18	19	9	16	5	4	1	1	-
2000.....	7	11	9	11	19	14	18	12	7	2	-	-
1999.....	3	5	9	11	14	14	11	8	5	1	-	-

Black, male

2003.....	4	6	6	10	2	7	3	-	2	-	-	-
2002.....	5	10	3	5	7	6	2	4	2	-	-	-
2001.....	1	3	1	9	8	2	4	1	-	-	-	-
2000.....	5	6	5	7	7	6	9	1	2	-	-	-
1999.....	2	3	7	4	6	8	4	3	2	1	-	-

Black, female

2003.....	4	6	5	6	8	13	5	7	3	-	1	-
2002.....	4	3	5	6	7	4	6	6	3	-	-	-
2001.....	5	2	7	9	11	7	12	4	4	1	1	-
2000.....	2	5	4	4	12	8	9	11	5	2	-	-
1999.....	1	2	2	7	8	6	7	5	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)												
American Indian 1/, both sexes												
2003.....	2	2	2	-	2	2	3	-	-	-	-	-
2002.....	1	2	2	1	5	2	-	-	-	2	1	-
2001.....	1	-	3	-	-	3	-	2	-	-	-	-
2000.....	1	2	1	1	-	1	1	-	-	-	-	-
1999.....	1	-	3	2	-	-	1	-	-	-	-	-
American Indian 1/, male												
2003.....	1	1	1	-	-	-	2	-	-	-	-	-
2002.....	1	2	1	1	3	-	-	-	-	-	-	-
2001.....	1	-	3	-	-	2	-	2	-	-	-	-
2000.....	1	1	1	1	-	-	1	-	-	-	-	-
1999.....	1	-	1	2	-	-	-	-	-	-	-	-
American Indian 1/, female												
2003.....	1	1	1	-	2	2	1	-	-	-	-	-
2002.....	-	-	1	-	2	2	-	-	-	2	1	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	1	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	2	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

Asian or Pacific Islander 2/, both sexes

2003.....	5	4	3	-	1	6	2	3	2	1	-	-
2002.....	2	2	5	5	6	4	3	1	1	1	-	-
2001.....	2	1	4	4	4	4	1	1	-	-	-	-
2000.....	3	3	4	5	3	1	1	-	2	-	-	-
1999.....	-	2	-	1	1	7	2	3	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	3	2	2	-	-	6	2	3	1	-	-	-
2002.....	2	-	4	4	4	3	1	1	-	-	-	-
2001.....	2	-	3	2	2	4	1	-	-	-	-	-
2000.....	1	2	-	3	2	1	-	-	2	-	-	-
1999.....	-	2	-	1	-	4	2	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	2	1	-	1	-	-	-	1	1	-	-
2002.....	-	2	1	1	2	1	2	-	1	1	-	-
2001.....	-	1	1	2	2	-	-	1	-	-	-	-
2000.....	2	1	4	2	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	3	-	2	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

All races, both sexes

2003.....	3,631	4,206	4,061	3,887	4,451	4,916	4,728	3,530	1,965	630	120	2
2002.....	3,707	3,924	3,914	3,954	4,551	4,978	4,770	3,528	1,869	658	126	-
2001.....	3,803	3,843	3,789	3,972	4,743	5,130	4,533	3,330	1,848	644	89	-
2000.....	3,655	3,790	3,783	4,135	4,889	5,256	4,494	3,596	1,880	658	115	-
1999.....	3,555	3,682	3,717	4,147	4,962	5,140	4,511	3,428	1,837	611	124	-

All races, male

2003.....	34	30	41	42	47	46	58	35	17	4	-	1
2002.....	28	29	38	35	49	57	55	31	13	3	-	-
2001.....	33	32	40	54	66	69	46	31	9	6	-	-
2000.....	28	44	41	47	68	65	44	36	13	1	1	-
1999.....	30	36	34	47	56	56	42	33	13	7	1	-

All races, female

2003.....	3,597	4,176	4,020	3,845	4,404	4,870	4,670	3,495	1,948	626	120	1
2002.....	3,679	3,895	3,876	3,919	4,502	4,921	4,715	3,497	1,856	655	126	-
2001.....	3,770	3,811	3,749	3,918	4,677	5,061	4,487	3,299	1,839	638	89	-
2000.....	3,627	3,746	3,742	4,088	4,821	5,191	4,450	3,560	1,867	657	114	-
1999.....	3,525	3,646	3,683	4,100	4,906	5,084	4,469	3,395	1,824	604	123	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

White, both sexes

2003.....	2,823	3,406	3,380	3,279	3,879	4,374	4,264	3,196	1,779	568	96	1
2002.....	2,904	3,206	3,288	3,365	3,968	4,465	4,268	3,218	1,702	590	108	-
2001.....	3,008	3,137	3,170	3,362	4,150	4,604	4,103	3,021	1,699	585	69	-
2000.....	2,934	3,100	3,182	3,584	4,341	4,735	4,107	3,267	1,696	586	102	-
1999.....	2,849	3,006	3,105	3,550	4,372	4,598	4,116	3,154	1,681	555	108	-

White, male

2003.....	30	24	35	36	40	42	53	31	16	4	-	1
2002.....	21	23	28	24	44	47	49	29	10	3	-	-
2001.....	24	27	34	46	55	59	41	26	9	4	-	-
2000.....	24	34	33	43	60	60	38	29	12	1	1	-
1999.....	22	29	25	40	45	49	37	30	11	5	1	-

White, female

2003.....	2,793	3,382	3,345	3,243	3,839	4,332	4,211	3,165	1,763	564	96	-
2002.....	2,883	3,183	3,260	3,341	3,924	4,418	4,219	3,189	1,692	587	108	-
2001.....	2,984	3,110	3,136	3,316	4,095	4,545	4,062	2,995	1,690	581	69	-
2000.....	2,910	3,066	3,149	3,541	4,281	4,675	4,069	3,238	1,684	585	101	-
1999.....	2,827	2,977	3,080	3,510	4,327	4,549	4,079	3,124	1,670	550	107	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

Black, both sexes

2003.....	675	685	569	521	492	472	419	303	169	59	23	-
2002.....	673	604	533	513	501	457	449	286	148	64	16	-
2001.....	665	602	529	537	523	481	391	275	142	59	19	-
2000.....	614	592	518	477	492	474	356	307	173	71	13	-
1999.....	614	582	521	523	517	493	363	252	140	54	16	-

Black, male

2003.....	4	6	6	6	7	3	4	4	1	-	-	-
2002.....	6	6	9	11	5	8	5	2	3	-	-	-
2001.....	9	4	6	7	10	9	5	4	-	2	-	-
2000.....	4	8	8	3	8	5	5	7	1	-	-	-
1999.....	8	7	9	6	10	6	5	3	2	2	-	-

Black, female

2003.....	671	679	563	515	485	469	415	299	168	59	23	-
2002.....	667	598	524	502	496	449	444	284	145	64	16	-
2001.....	656	598	523	530	513	472	386	271	142	57	19	-
2000.....	610	584	510	474	484	469	351	300	172	71	13	-
1999.....	606	575	512	517	507	487	358	249	138	52	16	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

American Indian 1/, both sexes

2003.....	20	17	19	9	15	18	15	10	4	-	-	-
2002.....	22	17	14	18	11	9	9	4	5	2	1	-
2001.....	20	14	18	12	13	3	5	9	1	-	-	-
2000.....	14	20	19	14	9	11	9	2	4	-	-	-
1999.....	17	14	14	18	18	11	6	9	2	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	20	17	19	9	15	18	14	10	4	-	-	-
2002.....	22	17	14	18	11	9	9	4	5	2	1	-
2001.....	20	14	18	12	12	3	5	9	1	-	-	-
2000.....	14	19	19	14	9	11	9	2	4	-	-	-
1999.....	17	14	14	17	18	11	6	9	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

Asian or Pacific Islander 2/, both sexes

2003.....	113	98	93	78	65	52	30	21	13	3	1	1
2002.....	108	97	79	58	71	47	44	20	14	2	1	-
2001.....	110	90	72	61	57	42	34	25	6	-	1	-
2000.....	93	78	64	60	47	36	22	20	7	1	-	-
1999.....	75	80	77	56	55	38	26	13	14	2	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	-	1	-	-	2	1	-	-	-	-	-
2001.....	-	1	-	1	-	1	-	1	-	-	-	-
2000.....	-	1	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	113	98	93	78	65	51	30	21	13	3	1	1
2002.....	107	97	78	58	71	45	43	20	14	2	1	-
2001.....	110	89	72	60	57	41	34	24	6	-	1	-
2000.....	93	77	64	59	47	36	21	20	7	1	-	-
1999.....	75	80	77	56	54	37	26	13	14	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

All races, both sexes

2003.....	450	445	348	308	256	274	278	166	78	22	4	-
2002.....	455	401	324	315	315	305	223	155	99	24	6	-
2001.....	478	371	352	279	306	298	285	193	90	30	2	-
2000.....	467	416	332	342	344	320	279	180	92	35	3	-
1999.....	452	393	376	350	346	345	223	199	98	26	6	1

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	450	445	348	308	256	274	278	166	78	22	4	-
2002.....	455	401	324	315	315	305	223	155	99	24	6	-
2001.....	478	371	352	279	306	298	285	193	90	30	2	-
2000.....	467	416	332	342	344	320	279	180	92	35	3	-
1999.....	452	393	376	350	346	345	223	199	98	26	6	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2003.....	341	348	265	231	183	188	223	132	54	20	4	-
2002.....	343	319	234	226	229	237	170	116	74	15	5	-
2001.....	363	283	272	195	235	227	219	150	69	26	2	-
2000.....	352	314	243	243	255	253	225	147	67	25	2	-
1999.....	360	298	275	261	242	265	181	161	76	14	4	1

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	341	348	265	231	183	188	223	132	54	20	4	-
2002.....	343	319	234	226	229	237	170	116	74	15	5	-
2001.....	363	283	272	195	235	227	219	150	69	26	2	-
2000.....	352	314	243	243	255	253	225	147	67	25	2	-
1999.....	360	298	275	261	242	265	181	161	76	14	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2003.....	83	81	72	68	58	66	45	29	19	2	-	-
2002.....	96	62	69	77	75	56	48	30	19	9	1	-
2001.....	89	65	66	73	57	56	56	39	19	4	-	-
2000.....	92	83	70	85	73	61	48	30	22	10	1	-
1999.....	72	80	89	79	97	69	33	35	22	9	2	-

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	83	81	72	68	58	66	45	29	19	2	-	-
2002.....	96	62	69	77	75	56	48	30	19	9	1	-
2001.....	89	65	66	73	57	56	56	39	19	4	-	-
2000.....	92	83	70	85	73	61	48	30	22	10	1	-
1999.....	72	80	89	79	97	69	33	35	22	9	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

American Indian 1/, both sexes

2003.....	3	6	2	1	2	2	2	-	2	-	-	-
2002.....	3	2	2	2	3	1	2	1	2	-	-	-
2001.....	3	5	1	1	1	1	1	1	1	-	-	-
2000.....	6	2	3	1	2	2	-	1	1	-	-	-
1999.....	2	3	2	1	-	-	2	-	-	-	-	-

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	3	6	2	1	2	2	2	-	2	-	-	-
2002.....	3	2	2	2	3	1	2	1	2	-	-	-
2001.....	3	5	1	1	1	1	1	1	1	-	-	-
2000.....	6	2	3	1	2	2	-	1	1	-	-	-
1999.....	2	3	2	1	-	-	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Asian or Pacific Islander 2/, both sexes

2003.....	23	10	9	8	13	18	8	5	3	-	-	-
2002.....	13	18	19	10	8	11	3	8	4	-	-	-
2001.....	23	18	13	10	13	14	9	3	1	-	-	-
2000.....	17	17	16	13	14	4	6	2	2	-	-	-
1999.....	18	12	10	9	7	11	7	3	-	3	-	-

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	23	10	9	8	13	18	8	5	3	-	-	-
2002.....	13	18	19	10	8	11	3	8	4	-	-	-
2001.....	23	18	13	10	13	14	9	3	1	-	-	-
2000.....	17	17	16	13	14	4	6	2	2	-	-	-
1999.....	18	12	10	9	7	11	7	3	-	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races, both sexes

2003.....	363	557	744	878	973	1,025	926	668	316	76	17	-
2002.....	376	531	668	809	1,026	1,051	906	694	314	109	11	-
2001.....	374	457	648	785	999	1,075	988	662	312	104	10	1
2000.....	317	478	644	800	981	1,039	919	617	333	84	15	1
1999.....	286	427	592	835	972	1,061	947	593	290	100	16	-

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	363	557	744	878	973	1,025	926	668	316	76	17	-
2002.....	376	531	668	809	1,026	1,051	906	694	314	109	11	-
2001.....	374	457	648	785	999	1,075	988	662	312	104	10	1
2000.....	317	478	644	800	981	1,039	919	617	333	84	15	1
1999.....	286	427	592	835	972	1,061	947	593	290	100	16	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2003.....	295	455	561	663	789	860	790	597	287	72	12	-
2002.....	296	412	522	595	824	865	783	609	281	91	9	-
2001.....	296	380	492	595	809	911	865	601	275	92	7	-
2000.....	262	394	496	602	771	866	785	557	294	74	11	-
1999.....	236	342	474	630	805	904	835	523	253	91	14	-

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	295	455	561	663	789	860	790	597	287	72	12	-
2002.....	296	412	522	595	824	865	783	609	281	91	9	-
2001.....	296	380	492	595	809	911	865	601	275	92	7	-
2000.....	262	394	496	602	771	866	785	557	294	74	11	-
1999.....	236	342	474	630	805	904	835	523	253	91	14	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2003.....	55	83	165	190	169	150	123	66	28	4	5	-
2002.....	61	99	124	188	188	162	111	81	31	17	2	-
2001.....	59	65	130	174	180	153	115	55	36	11	3	1
2000.....	44	71	131	177	198	156	124	56	36	10	4	1
1999.....	39	76	113	189	161	144	103	65	35	9	2	-

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	55	83	165	190	169	150	123	66	28	4	5	-
2002.....	61	99	124	188	188	162	111	81	31	17	2	-
2001.....	59	65	130	174	180	153	115	55	36	11	3	1
2000.....	44	71	131	177	198	156	124	56	36	10	4	1
1999.....	39	76	113	189	161	144	103	65	35	9	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

American Indian 1/, both sexes

2003.....	1	-	1	1	3	-	2	-	-	-	-	-
2002.....	3	3	3	3	1	7	2	2	-	-	-	-
2001.....	3	-	6	2	4	3	2	2	-	1	-	-
2000.....	-	-	2	5	3	4	2	-	1	-	-	-
1999.....	-	2	-	2	2	1	-	2	-	-	-	-

American Indian 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2003.....	1	-	1	1	3	-	2	-	-	-	-	-
2002.....	3	3	3	3	1	7	2	2	-	-	-	-
2001.....	3	-	6	2	4	3	2	2	-	1	-	-
2000.....	-	-	2	5	3	4	2	-	1	-	-	-
1999.....	-	2	-	2	2	1	-	2	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Asian or Pacific Islander 2/, both sexes												
2003.....	12	19	17	24	12	15	11	5	1	-	-	-
2002.....	16	17	19	23	13	17	10	2	2	1	-	-
2001.....	16	12	20	14	6	8	6	4	1	-	-	-
2000.....	11	13	15	16	9	13	8	4	2	-	-	-
1999.....	11	7	5	14	4	12	9	3	2	-	-	-
Asian or Pacific Islander 2/, male												
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2003.....	12	19	17	24	12	15	11	5	1	-	-	-
2002.....	16	17	19	23	13	17	10	2	2	1	-	-
2001.....	16	12	20	14	6	8	6	4	1	-	-	-
2000.....	11	13	15	16	9	13	8	4	2	-	-	-
1999.....	11	7	5	14	4	12	9	3	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of ovary (C56)

All races, both sexes

2003.....	997	1,315	1,576	1,687	1,882	2,193	2,001	1,207	490	102	11	-
2002.....	1,029	1,249	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001.....	1,016	1,171	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000.....	966	1,123	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999.....	933	1,081	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	997	1,315	1,576	1,687	1,882	2,193	2,001	1,207	490	102	11	-
2002.....	1,029	1,249	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001.....	1,016	1,171	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000.....	966	1,123	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999.....	933	1,081	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of ovary (C56)

White, both sexes

2003.....	851	1,173	1,398	1,499	1,682	2,004	1,854	1,105	457	93	8	-
2002.....	906	1,102	1,339	1,484	1,751	2,058	1,859	1,087	416	105	11	-
2001.....	890	1,030	1,254	1,486	1,803	2,062	1,794	1,088	426	85	10	-
2000.....	838	1,008	1,179	1,489	1,752	2,130	1,684	1,099	418	86	8	-
1999.....	814	965	1,199	1,455	1,799	1,956	1,614	987	381	84	7	-

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	851	1,173	1,398	1,499	1,682	2,004	1,854	1,105	457	93	8	-
2002.....	906	1,102	1,339	1,484	1,751	2,058	1,859	1,087	416	105	11	-
2001.....	890	1,030	1,254	1,486	1,803	2,062	1,794	1,088	426	85	10	-
2000.....	838	1,008	1,179	1,489	1,752	2,130	1,684	1,099	418	86	8	-
1999.....	814	965	1,199	1,455	1,799	1,956	1,614	987	381	84	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of ovary (C56)												
Black, both sexes												
2003.....	110	108	130	146	156	154	116	83	31	9	3	-
2002.....	87	114	126	140	149	154	124	78	27	9	3	-
2001.....	88	112	141	151	154	151	103	80	24	10	5	-
2000.....	89	86	119	152	147	147	97	66	31	12	1	-
1999.....	89	95	122	136	168	162	98	54	30	6	-	-
Black, male												
2003.....
2002.....
2001.....
2000.....
1999.....
Black, female												
2003.....	110	108	130	146	156	154	116	83	31	9	3	-
2002.....	87	114	126	140	149	154	124	78	27	9	3	-
2001.....	88	112	141	151	154	151	103	80	24	10	5	-
2000.....	89	86	119	152	147	147	97	66	31	12	1	-
1999.....	89	95	122	136	168	162	98	54	30	6	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of ovary (C56)												
American Indian 1/, both sexes												
2003.....	8	4	11	6	6	11	5	5	-	-	-	-
2002.....	7	4	4	3	9	1	8	1	1	-	-	-
2001.....	6	8	8	14	6	8	6	2	-	1	-	-
2000.....	1	3	4	2	3	6	3	2	-	-	-	-
1999.....	2	2	1	8	3	5	4	1	-	1	-	-
American Indian 1/, male												
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2003.....	8	4	11	6	6	11	5	5	-	-	-	-
2002.....	7	4	4	3	9	1	8	1	1	-	-	-
2001.....	6	8	8	14	6	8	6	2	-	1	-	-
2000.....	1	3	4	2	3	6	3	2	-	-	-	-
1999.....	2	2	1	8	3	5	4	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of ovary (C56)

Asian or Pacific Islander 2/, both sexes

2003.....	28	30	37	36	38	24	26	14	2	-	-	-
2002.....	29	29	30	35	30	36	19	8	3	1	-	-
2001.....	32	21	38	31	21	25	21	11	1	-	-	-
2000.....	38	26	19	26	30	12	19	9	4	-	-	-
1999.....	28	19	23	27	16	20	16	10	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2003.....	28	30	37	36	38	24	26	14	2	-	-	-
2002.....	29	29	30	35	30	36	19	8	3	1	-	-
2001.....	32	21	38	31	21	25	21	11	1	-	-	-
2000.....	38	26	19	26	30	12	19	9	4	-	-	-
1999.....	28	19	23	27	16	20	16	10	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races, both sexes

2003.....	763	1,121	1,377	1,409	1,681	1,849	1,626	1,033	420	107	7	-
2002.....	808	1,075	1,265	1,533	1,706	1,829	1,526	967	429	90	7	-
2001.....	813	1,006	1,322	1,395	1,826	1,810	1,496	947	390	82	5	-
2000.....	789	1,026	1,169	1,432	1,714	1,819	1,463	945	397	83	7	-
1999.....	720	959	1,152	1,374	1,702	1,696	1,314	867	341	91	6	-

All races, male

2003.....	535	798	989	916	1,059	1,114	922	486	175	40	2	-
2002.....	592	771	887	1,020	1,067	1,123	801	457	201	34	4	-
2001.....	586	722	912	928	1,165	1,052	814	500	164	23	-	-
2000.....	571	727	803	942	1,068	1,054	811	477	165	31	3	-
1999.....	506	671	803	894	1,064	1,016	701	421	140	29	-	-

All races, female

2003.....	228	323	388	493	622	735	704	547	245	67	5	-
2002.....	216	304	378	513	639	706	725	510	228	56	3	-
2001.....	227	284	410	467	661	758	682	447	226	59	5	-
2000.....	218	299	366	490	646	765	652	468	232	52	4	-
1999.....	214	288	349	480	638	680	613	446	201	62	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

White, both sexes

2003.....	660	967	1,208	1,252	1,520	1,684	1,499	956	380	98	5	-
2002.....	688	951	1,114	1,367	1,543	1,654	1,402	888	403	81	7	-
2001.....	687	878	1,152	1,259	1,649	1,639	1,371	873	354	77	5	-
2000.....	685	903	1,031	1,271	1,542	1,657	1,344	880	361	76	6	-
1999.....	616	841	1,027	1,226	1,531	1,541	1,211	802	303	77	5	-

White, male

2003.....	462	703	871	819	954	1,015	858	456	161	36	1	-
2002.....	511	682	785	917	969	1,033	733	423	195	31	4	-
2001.....	495	640	786	841	1,060	947	750	467	151	22	-	-
2000.....	495	637	716	829	966	965	744	444	152	29	2	-
1999.....	427	586	724	798	961	927	660	394	124	20	-	-

White, female

2003.....	198	264	337	433	566	669	641	500	219	62	4	-
2002.....	177	269	329	450	574	621	669	465	208	50	3	-
2001.....	192	238	366	418	589	692	621	406	203	55	5	-
2000.....	190	266	315	442	576	692	600	436	209	47	4	-
1999.....	189	255	303	428	570	614	551	408	179	57	5	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, both sexes

2003.....	89	130	132	128	122	131	103	71	36	7	2	-
2002.....	92	104	122	135	125	150	102	67	21	7	-	-
2001.....	101	105	136	108	143	138	104	63	29	5	-	-
2000.....	87	100	114	134	139	132	103	57	30	7	1	-
1999.....	87	102	104	122	143	129	84	55	31	11	1	-

Black, male

2003.....	64	82	88	79	79	78	53	26	13	2	1	-
2002.....	62	79	85	80	71	75	51	29	6	2	-	-
2001.....	72	67	101	67	87	84	53	30	10	1	-	-
2000.....	61	69	74	96	84	68	58	30	10	2	1	-
1999.....	65	74	66	77	83	72	32	21	13	8	-	-

Black, female

2003.....	25	48	44	49	43	53	50	45	23	5	1	-
2002.....	30	25	37	55	54	75	51	38	15	5	-	-
2001.....	29	38	35	41	56	54	51	33	19	4	-	-
2000.....	26	31	40	38	55	64	45	27	20	5	-	-
1999.....	22	28	38	45	60	57	52	34	18	3	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

American Indian 1/, both sexes

2003.....	8	10	10	8	15	7	10	1	1	-	-	-
2002.....	11	7	12	16	18	7	8	4	-	1	-	-
2001.....	11	9	16	12	10	9	4	1	4	-	-	-
2000.....	6	5	5	7	12	9	1	3	1	-	-	-
1999.....	8	9	7	10	11	14	6	2	1	1	-	-

American Indian 1/, male

2003.....	4	6	10	5	11	3	4	-	-	-	-	-
2002.....	7	5	6	12	14	4	6	1	-	-	-	-
2001.....	8	5	9	9	8	5	1	-	2	-	-	-
2000.....	5	5	3	5	6	7	-	1	-	-	-	-
1999.....	6	7	3	7	8	9	4	1	-	1	-	-

American Indian 1/, female

2003.....	4	4	-	3	4	4	6	1	1	-	-	-
2002.....	4	2	6	4	4	3	2	3	-	1	-	-
2001.....	3	4	7	3	2	4	3	1	2	-	-	-
2000.....	1	-	2	2	6	2	1	2	1	-	-	-
1999.....	2	2	4	3	3	5	2	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Asian or Pacific Islander 2/, both sexes

2003.....	6	14	27	21	24	27	14	5	3	2	-	-
2002.....	17	13	17	15	20	18	14	8	5	1	-	-
2001.....	14	14	18	16	24	24	17	10	3	-	-	-
2000.....	11	18	19	20	21	21	15	5	5	-	-	-
1999.....	9	7	14	16	17	12	13	8	6	2	-	-

Asian or Pacific Islander 2/, male

2003.....	5	7	20	13	15	18	7	4	1	2	-	-
2002.....	12	5	11	11	13	11	11	4	-	1	-	-
2001.....	11	10	16	11	10	16	10	3	1	-	-	-
2000.....	10	16	10	12	12	14	9	2	3	-	-	-
1999.....	8	4	10	12	12	8	5	5	3	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	7	7	8	9	9	7	1	2	-	-	-
2002.....	5	8	6	4	7	7	3	4	5	-	-	-
2001.....	3	4	2	5	14	8	7	7	2	-	-	-
2000.....	1	2	9	8	9	7	6	3	2	-	-	-
1999.....	1	3	4	4	5	4	8	3	3	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

All races, both sexes

2003.....	341	554	837	1,148	1,588	2,239	2,440	1,850	888	278	36	-
2002.....	321	572	738	1,151	1,756	2,366	2,341	1,864	908	255	32	1
2001.....	314	484	781	1,147	1,667	2,179	2,309	1,865	864	265	42	-
2000.....	280	486	737	1,119	1,736	2,155	2,252	1,790	875	256	31	-
1999.....	323	492	752	1,126	1,797	2,178	2,139	1,714	852	249	30	1

All races, male

2003.....	252	419	630	865	1,133	1,603	1,651	1,157	508	131	11	-
2002.....	223	425	559	845	1,244	1,659	1,540	1,163	498	118	12	1
2001.....	233	359	597	859	1,229	1,554	1,562	1,129	482	123	17	-
2000.....	212	369	547	838	1,256	1,542	1,512	1,110	478	110	12	-
1999.....	243	374	559	849	1,283	1,575	1,433	1,039	438	116	15	1

All races, female

2003.....	89	135	207	283	455	636	789	693	380	147	25	-
2002.....	98	147	179	306	512	707	801	701	410	137	20	-
2001.....	81	125	184	288	438	625	747	736	382	142	25	-
2000.....	68	117	190	281	480	613	740	680	397	146	19	-
1999.....	80	118	193	277	514	603	706	675	414	133	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

White, both sexes

2003.....	288	492	761	1,024	1,472	2,058	2,252	1,717	835	253	32	-
2002.....	280	502	668	1,031	1,594	2,177	2,167	1,722	831	229	29	1
2001.....	274	426	694	1,039	1,534	2,012	2,126	1,743	792	247	38	-
2000.....	254	439	670	1,020	1,585	1,977	2,079	1,665	830	242	27	-
1999.....	283	429	664	1,005	1,638	2,011	1,997	1,592	787	221	30	1

White, male

2003.....	221	378	584	779	1,069	1,496	1,551	1,098	484	121	9	-
2002.....	203	390	516	774	1,157	1,544	1,446	1,095	467	112	12	1
2001.....	205	323	546	798	1,148	1,462	1,467	1,069	455	121	15	-
2000.....	195	335	508	779	1,170	1,447	1,414	1,037	460	106	12	-
1999.....	210	340	503	778	1,199	1,477	1,351	985	412	107	15	1

White, female

2003.....	67	114	177	245	403	562	701	619	351	132	23	-
2002.....	77	112	152	257	437	633	721	627	364	117	17	-
2001.....	69	103	148	241	386	550	659	674	337	126	23	-
2000.....	59	104	162	241	415	530	665	628	370	136	15	-
1999.....	73	89	161	227	439	534	646	607	375	114	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

Black, both sexes

2003.....	48	50	59	109	96	156	161	115	43	22	3	-
2002.....	33	60	60	103	139	157	142	121	68	21	3	-
2001.....	35	48	77	93	105	143	152	97	62	17	3	-
2000.....	24	41	59	86	134	160	158	103	41	12	3	-
1999.....	39	59	74	107	140	142	122	110	54	25	-	-

Black, male

2003.....	28	32	34	74	50	90	83	51	18	7	1	-
2002.....	16	30	36	57	71	91	75	56	26	3	-	-
2001.....	26	27	45	50	61	74	72	44	21	2	2	-
2000.....	16	28	35	48	72	87	87	59	15	3	-	-
1999.....	32	32	46	60	70	82	68	49	18	7	-	-

Black, female

2003.....	20	18	25	35	46	66	78	64	25	15	2	-
2002.....	17	30	24	46	68	66	67	65	42	18	3	-
2001.....	9	21	32	43	44	69	80	53	41	15	1	-
2000.....	8	13	24	38	62	73	71	44	26	9	3	-
1999.....	7	27	28	47	70	60	54	61	36	18	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

American Indian 1/, both sexes

2003.....	2	5	5	3	1	5	6	2	1	-	-	-
2002.....	2	2	2	7	2	3	5	5	1	1	-	-
2001.....	2	2	2	4	4	1	6	3	-	-	-	-
2000.....	1	1	1	2	1	5	2	1	1	-	-	-
1999.....	-	1	4	5	5	1	1	2	1	-	-	-

American Indian 1/, male

2003.....	-	2	4	2	1	2	3	1	1	-	-	-
2002.....	2	1	-	6	1	2	2	2	-	1	-	-
2001.....	-	1	1	2	2	1	5	2	-	-	-	-
2000.....	-	1	1	2	1	1	2	1	1	-	-	-
1999.....	-	1	2	5	4	1	-	1	1	-	-	-

American Indian 1/, female

2003.....	2	3	1	1	-	3	3	1	-	-	-	-
2002.....	-	1	2	1	1	1	3	3	1	-	-	-
2001.....	2	1	1	2	2	-	1	1	-	-	-	-
2000.....	1	-	-	-	-	4	-	-	-	-	-	-
1999.....	-	-	2	-	1	-	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

Asian or Pacific Islander 2/, both sexes

2003.....	3	7	12	12	19	20	21	16	9	3	1	-
2002.....	6	8	8	10	21	29	27	16	8	4	-	-
2001.....	3	8	8	11	24	23	25	22	10	1	1	-
2000.....	1	5	7	11	16	13	13	21	3	2	1	-
1999.....	1	3	10	9	14	24	19	10	10	3	-	-

Asian or Pacific Islander 2/, male

2003.....	3	7	8	10	13	15	14	7	5	3	1	-
2002.....	2	4	7	8	15	22	17	10	5	2	-	-
2001.....	2	8	5	9	18	17	18	14	6	-	-	-
2000.....	1	5	3	9	13	7	9	13	2	1	-	-
1999.....	1	1	8	6	10	15	14	4	7	2	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	4	2	6	5	7	9	4	-	-	-
2002.....	4	4	1	2	6	7	10	6	3	2	-	-
2001.....	1	-	3	2	6	6	7	8	4	1	1	-
2000.....	-	-	4	2	3	6	4	8	1	1	1	-
1999.....	-	2	2	3	4	9	5	6	3	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races, both sexes

2003.....	1,105	1,337	1,361	1,349	1,487	1,449	1,061	513	173	28	3	-
2002.....	1,158	1,276	1,284	1,407	1,547	1,454	981	536	174	39	4	-
2001.....	1,121	1,174	1,290	1,354	1,469	1,506	1,060	528	157	31	5	-
2000.....	1,109	1,110	1,227	1,409	1,623	1,437	1,041	521	166	28	2	-
1999.....	1,038	1,206	1,247	1,498	1,621	1,543	949	496	144	22	1	-

All races, male

2003.....	689	741	789	741	815	745	522	214	65	8	2	-
2002.....	665	773	721	819	859	750	460	213	54	10	2	-
2001.....	673	734	736	763	826	777	511	202	48	9	2	-
2000.....	667	660	719	789	889	705	482	212	53	9	1	-
1999.....	613	670	747	858	870	785	441	192	47	7	1	-

All races, female

2003.....	416	596	572	608	672	704	539	299	108	20	1	-
2002.....	493	503	563	588	688	704	521	323	120	29	2	-
2001.....	448	440	554	591	643	729	549	326	109	22	3	-
2000.....	442	450	508	620	734	732	559	309	113	19	1	-
1999.....	425	536	500	640	751	758	508	304	97	15	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2003.....	1,008	1,225	1,271	1,247	1,390	1,368	1,007	478	164	22	3	-
2002.....	1,056	1,165	1,194	1,310	1,454	1,366	926	507	164	36	4	-
2001.....	1,030	1,073	1,197	1,250	1,378	1,433	1,007	500	148	26	4	-
2000.....	1,026	1,019	1,142	1,321	1,543	1,362	978	486	157	27	1	-
1999.....	950	1,123	1,153	1,400	1,545	1,463	900	473	131	20	1	-

White, male

2003.....	625	682	741	684	764	702	494	201	61	7	2	-
2002.....	618	700	669	767	810	705	431	204	51	9	2	-
2001.....	620	676	679	717	783	734	488	190	43	7	1	-
2000.....	609	611	673	747	846	669	459	201	51	9	-	-
1999.....	560	625	691	807	834	745	424	183	45	6	1	-

White, female

2003.....	383	543	530	563	626	666	513	277	103	15	1	-
2002.....	438	465	525	543	644	661	495	303	113	27	2	-
2001.....	410	397	518	533	595	699	519	310	105	19	3	-
2000.....	417	408	469	574	697	693	519	285	106	18	1	-
1999.....	390	498	462	593	711	718	476	290	86	14	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2003.....	77	87	66	79	73	62	41	23	7	6	-	-
2002.....	78	85	64	75	71	71	43	20	9	3	-	-
2001.....	69	76	72	83	70	54	40	24	8	5	1	-
2000.....	70	73	66	68	66	58	51	26	6	1	1	-
1999.....	70	70	72	77	58	67	40	21	9	1	-	-

Black, male

2003.....	51	47	35	48	37	35	19	4	2	1	-	-
2002.....	32	55	40	38	35	34	23	4	2	1	-	-
2001.....	41	44	45	36	34	27	15	8	4	2	1	-
2000.....	49	40	37	30	37	25	18	6	2	-	1	-
1999.....	43	35	41	42	26	30	11	8	2	-	-	-

Black, female

2003.....	26	40	31	31	36	27	22	19	5	5	-	-
2002.....	46	30	24	37	36	37	20	16	7	2	-	-
2001.....	28	32	27	47	36	27	25	16	4	3	-	-
2000.....	21	33	29	38	29	33	33	20	4	1	-	-
1999.....	27	35	31	35	32	37	29	13	7	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

American Indian 1/, both sexes

2003.....	6	4	5	5	3	2	1	3	-	-	-	-
2002.....	4	4	1	2	4	3	-	1	-	-	-	-
2001.....	7	5	3	5	3	3	2	-	-	-	-	-
2000.....	1	2	1	3	1	6	2	1	-	-	-	-
1999.....	4	3	5	4	2	2	-	-	-	1	-	-

American Indian 1/, male

2003.....	4	2	2	2	1	2	1	2	-	-	-	-
2002.....	2	4	-	1	3	1	-	-	-	-	-	-
2001.....	1	3	2	3	1	3	1	-	-	-	-	-
2000.....	1	1	-	2	-	4	1	1	-	-	-	-
1999.....	1	3	3	1	1	1	-	-	-	1	-	-

American Indian 1/, female

2003.....	2	2	3	3	2	-	-	1	-	-	-	-
2002.....	2	-	1	1	1	2	-	1	-	-	-	-
2001.....	6	2	1	2	2	-	1	-	-	-	-	-
2000.....	-	1	1	1	1	2	1	-	-	-	-	-
1999.....	3	-	2	3	1	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Asian or Pacific Islander 2/, both sexes

2003.....	14	21	19	18	21	17	12	9	2	-	-	-
2002.....	20	22	25	20	18	14	12	8	1	-	-	-
2001.....	15	20	18	16	18	16	11	4	1	-	-	-
2000.....	12	16	18	17	13	11	10	8	3	-	-	-
1999.....	14	10	17	17	16	11	9	2	4	-	-	-

Asian or Pacific Islander 2/, male

2003.....	9	10	11	7	13	6	8	7	2	-	-	-
2002.....	13	14	12	13	11	10	6	5	1	-	-	-
2001.....	11	11	10	7	8	13	7	4	1	-	-	-
2000.....	8	8	9	10	6	7	4	4	-	-	-	-
1999.....	9	7	12	8	9	9	6	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	5	11	8	11	8	11	4	2	-	-	-	-
2002.....	7	8	13	7	7	4	6	3	-	-	-	-
2001.....	4	9	8	9	10	3	4	-	-	-	-	-
2000.....	4	8	9	7	7	4	6	4	3	-	-	-
1999.....	5	3	5	9	7	2	3	1	4	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races, both sexes

2003.....	2,228	3,171	4,300	5,384	7,501	9,497	8,945	6,087	2,541	644	62	-
2002.....	2,361	3,217	4,283	5,672	7,836	9,319	8,970	6,003	2,433	574	62	-
2001.....	2,390	3,224	4,231	5,602	8,057	9,580	8,677	5,848	2,344	536	77	-
2000.....	2,350	3,156	4,128	5,932	8,109	9,743	8,705	5,716	2,329	559	70	-
1999.....	2,302	3,346	4,208	5,996	8,336	9,596	8,493	5,468	2,233	526	55	-

All races, male

2003.....	1,347	1,929	2,628	3,255	4,295	5,068	4,456	2,722	943	202	16	-
2002.....	1,388	1,947	2,523	3,391	4,512	5,047	4,506	2,624	906	186	24	-
2001.....	1,440	1,900	2,512	3,316	4,543	5,065	4,262	2,537	846	168	25	-
2000.....	1,384	1,850	2,355	3,445	4,593	5,168	4,231	2,475	783	185	15	-
1999.....	1,382	1,957	2,469	3,436	4,631	5,102	3,998	2,337	816	155	21	-

All races, female

2003.....	881	1,242	1,672	2,129	3,206	4,429	4,489	3,365	1,598	442	46	-
2002.....	973	1,270	1,760	2,281	3,324	4,272	4,464	3,379	1,527	388	38	-
2001.....	950	1,324	1,719	2,286	3,514	4,515	4,415	3,311	1,498	368	52	-
2000.....	966	1,306	1,773	2,487	3,516	4,575	4,474	3,241	1,546	374	55	-
1999.....	920	1,389	1,739	2,560	3,705	4,494	4,495	3,131	1,417	371	34	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, both sexes

2003.....	1,797	2,710	3,719	4,701	6,661	8,633	8,259	5,697	2,368	587	53	-
2002.....	1,909	2,692	3,674	4,970	7,007	8,512	8,324	5,619	2,262	534	54	-
2001.....	1,980	2,704	3,614	4,856	7,252	8,751	8,040	5,451	2,187	494	72	-
2000.....	1,957	2,693	3,562	5,251	7,307	8,944	8,129	5,292	2,145	511	59	-
1999.....	1,919	2,836	3,604	5,297	7,529	8,778	7,909	5,093	2,065	483	47	-

White, male

2003.....	1,094	1,657	2,310	2,891	3,892	4,690	4,167	2,571	881	185	15	-
2002.....	1,149	1,660	2,183	3,007	4,087	4,661	4,209	2,477	840	176	20	-
2001.....	1,209	1,607	2,163	2,910	4,130	4,672	3,969	2,381	791	154	24	-
2000.....	1,150	1,591	2,048	3,103	4,146	4,784	3,976	2,309	725	169	14	-
1999.....	1,168	1,673	2,144	3,068	4,219	4,683	3,761	2,176	750	142	17	-

White, female

2003.....	703	1,053	1,409	1,810	2,769	3,943	4,092	3,126	1,487	402	38	-
2002.....	760	1,032	1,491	1,963	2,920	3,851	4,115	3,142	1,422	358	34	-
2001.....	771	1,097	1,451	1,946	3,122	4,079	4,071	3,070	1,396	340	48	-
2000.....	807	1,102	1,514	2,148	3,161	4,160	4,153	2,983	1,420	342	45	-
1999.....	751	1,163	1,460	2,229	3,310	4,095	4,148	2,917	1,315	341	30	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2003.....	374	372	486	566	684	690	548	317	148	43	8	-
2002.....	363	444	505	587	695	651	528	326	146	34	7	-
2001.....	332	429	501	628	674	672	527	330	135	38	4	-
2000.....	328	393	482	563	678	643	466	356	159	38	8	-
1999.....	322	435	492	585	668	667	482	319	142	39	8	-

Black, male

2003.....	223	228	265	298	320	302	226	117	49	11	-	-
2002.....	190	242	281	316	369	300	232	123	52	10	4	-
2001.....	194	240	280	346	339	304	238	124	45	12	1	-
2000.....	188	223	262	281	375	313	199	132	49	13	1	-
1999.....	183	243	260	311	338	335	189	123	51	10	4	-

Black, female

2003.....	151	144	221	268	364	388	322	200	99	32	8	-
2002.....	173	202	224	271	326	351	296	203	94	24	3	-
2001.....	138	189	221	282	335	368	289	206	90	26	3	-
2000.....	140	170	220	282	303	330	267	224	110	25	7	-
1999.....	139	192	232	274	330	332	293	196	91	29	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

American Indian 1/, both sexes

2003.....	10	23	18	19	29	29	21	11	4	1	1	-
2002.....	12	16	19	17	17	23	13	6	4	2	-	-
2001.....	17	21	25	34	21	19	17	10	6	1	-	-
2000.....	11	15	19	22	24	25	23	13	7	4	-	-
1999.....	8	9	14	22	28	26	11	8	5	-	-	-

American Indian 1/, male

2003.....	6	9	9	12	13	7	9	4	1	-	1	-
2002.....	8	7	9	9	8	11	8	2	1	-	-	-
2001.....	8	11	15	17	11	7	6	4	1	-	-	-
2000.....	8	9	8	10	16	10	11	5	1	2	-	-
1999.....	5	4	8	15	14	18	3	5	3	-	-	-

American Indian 1/, female

2003.....	4	14	9	7	16	22	12	7	3	1	-	-
2002.....	4	9	10	8	9	12	5	4	3	2	-	-
2001.....	9	10	10	17	10	12	11	6	5	1	-	-
2000.....	3	6	11	12	8	15	12	8	6	2	-	-
1999.....	3	5	6	7	14	8	8	3	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Asian or Pacific Islander 2/, both sexes

2003.....	47	66	77	98	127	145	117	62	21	13	-	-
2002.....	77	65	85	98	117	133	105	52	21	4	1	-
2001.....	61	70	91	84	110	138	93	57	16	3	1	-
2000.....	54	55	65	96	100	131	87	55	18	6	3	-
1999.....	53	66	98	92	111	125	91	48	21	4	-	-

Asian or Pacific Islander 2/, male

2003.....	24	35	44	54	70	69	54	30	12	6	-	-
2002.....	41	38	50	59	48	75	57	22	13	-	-	-
2001.....	29	42	54	43	63	82	49	28	9	2	-	-
2000.....	38	27	37	51	56	61	45	29	8	1	-	-
1999.....	26	37	57	42	60	66	45	33	12	3	-	-

Asian or Pacific Islander 2/, female

2003.....	23	31	33	44	57	76	63	32	9	7	-	-
2002.....	36	27	35	39	69	58	48	30	8	4	1	-
2001.....	32	28	37	41	47	56	44	29	7	1	1	-
2000.....	16	28	28	45	44	70	42	26	10	5	3	-
1999.....	27	29	41	50	51	59	46	15	9	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

All races, both sexes

2003.....	83	98	100	83	133	153	124	75	34	3	-	-
2002.....	94	74	94	94	138	126	121	83	35	7	-	-
2001.....	91	61	82	103	108	151	127	66	24	7	-	-
2000.....	88	64	94	92	140	134	121	70	24	10	-	-
1999.....	83	87	91	120	163	149	113	70	22	5	1	-

All races, male

2003.....	47	57	62	52	76	73	59	31	7	-	-	-
2002.....	52	53	53	62	84	60	65	37	11	2	-	-
2001.....	56	40	45	65	64	75	55	26	7	1	-	-
2000.....	53	30	57	58	76	71	50	30	10	5	-	-
1999.....	50	54	61	72	88	69	50	28	16	1	-	-

All races, female

2003.....	36	41	38	31	57	80	65	44	27	3	-	-
2002.....	42	21	41	32	54	66	56	46	24	5	-	-
2001.....	35	21	37	38	44	76	72	40	17	6	-	-
2000.....	35	34	37	34	64	63	71	40	14	5	-	-
1999.....	33	33	30	48	75	80	63	42	6	4	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

White, both sexes

2003.....	75	79	91	71	125	143	115	71	33	3	-	-
2002.....	79	61	80	85	128	119	115	81	33	6	-	-
2001.....	84	52	72	99	101	138	122	64	24	7	-	-
2000.....	75	56	82	86	125	134	114	68	22	9	-	-
1999.....	74	78	85	106	154	144	110	62	21	5	1	-

White, male

2003.....	41	45	56	43	69	67	56	28	7	-	-	-
2002.....	41	43	43	57	79	57	63	37	10	2	-	-
2001.....	51	35	39	63	61	70	52	24	7	1	-	-
2000.....	45	26	48	55	67	71	47	29	9	5	-	-
1999.....	42	49	58	64	84	67	49	25	15	1	-	-

White, female

2003.....	34	34	35	28	56	76	59	43	26	3	-	-
2002.....	38	18	37	28	49	62	52	44	23	4	-	-
2001.....	33	17	33	36	40	68	70	40	17	6	-	-
2000.....	30	30	34	31	58	63	67	39	13	4	-	-
1999.....	32	29	27	42	70	77	61	37	6	4	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

Black, both sexes

2003.....	8	15	6	11	6	6	3	-	-	-	-	-
2002.....	10	10	11	6	9	5	4	1	1	1	-	-
2001.....	5	9	9	4	7	9	3	2	-	-	-	-
2000.....	10	8	9	4	11	-	5	1	2	-	-	-
1999.....	9	9	5	11	7	5	1	6	1	-	-	-

Black, male

2003.....	6	10	5	8	5	3	1	-	-	-	-	-
2002.....	7	8	8	3	5	3	1	-	1	-	-	-
2001.....	4	5	5	2	3	3	2	2	-	-	-	-
2000.....	5	4	7	2	5	-	3	-	1	-	-	-
1999.....	8	5	3	8	4	2	1	1	1	-	-	-

Black, female

2003.....	2	5	1	3	1	3	2	-	-	-	-	-
2002.....	3	2	3	3	4	2	3	1	-	1	-	-
2001.....	1	4	4	2	4	6	1	-	-	-	-	-
2000.....	5	4	2	2	6	-	2	1	1	-	-	-
1999.....	1	4	2	3	3	3	-	5	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

American Indian 1/, both sexes

2003.....	-	2	-	1	-	1	3	-	1	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	1	3	-	-	-	-	-	-	-
1999.....	-	-	-	2	1	-	-	1	-	-	-	-

American Indian 1/, male

2003.....	-	1	-	1	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	3	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

American Indian 1/, female

2003.....	-	1	-	-	-	-	3	-	1	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	2	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

Asian or Pacific Islander 2/, both sexes

2003.....	-	2	3	-	2	3	3	4	-	-	-	-
2002.....	5	3	2	3	1	2	2	1	1	-	-	-
2001.....	1	-	1	-	-	4	2	-	-	-	-	-
2000.....	3	-	2	1	1	-	2	1	-	1	-	-
1999.....	-	-	1	1	1	-	2	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	1	1	-	2	2	2	3	-	-	-	-
2002.....	4	2	2	2	-	-	1	-	-	-	-	-
2001.....	1	-	1	-	-	2	1	-	-	-	-	-
2000.....	3	-	2	-	1	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	1	2	-	-	1	1	1	-	-	-	-
2002.....	1	1	-	1	1	2	1	1	1	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	-	-	1	-	-	2	-	-	1	-	-
1999.....	-	-	1	1	1	-	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

All races, both sexes

2003.....	899	1,272	1,735	2,104	2,815	3,674	3,543	2,520	949	213	30	-
2002.....	943	1,319	1,730	2,225	3,039	3,722	3,561	2,446	881	200	22	-
2001.....	1,021	1,367	1,731	2,193	3,165	3,776	3,593	2,392	904	186	24	-
2000.....	1,015	1,434	1,736	2,386	3,212	3,984	3,551	2,330	856	193	23	-
1999.....	1,045	1,414	1,740	2,441	3,407	4,009	3,545	2,198	834	178	18	-

All races, male

2003.....	558	773	1,075	1,260	1,569	1,882	1,681	1,065	345	78	9	-
2002.....	574	823	1,015	1,334	1,652	1,898	1,696	1,013	315	51	6	-
2001.....	663	817	1,008	1,286	1,739	1,873	1,685	1,020	305	55	5	-
2000.....	646	865	984	1,342	1,763	2,018	1,616	959	274	62	4	-
1999.....	652	851	996	1,355	1,821	2,052	1,544	892	292	53	9	-

All races, female

2003.....	341	499	660	844	1,246	1,792	1,862	1,455	604	135	21	-
2002.....	369	496	715	891	1,387	1,824	1,865	1,433	566	149	16	-
2001.....	358	550	723	907	1,426	1,903	1,908	1,372	599	131	19	-
2000.....	369	569	752	1,044	1,449	1,966	1,935	1,371	582	131	19	-
1999.....	393	563	744	1,086	1,586	1,957	2,001	1,306	542	125	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

White, both sexes

2003.....	754	1,138	1,543	1,907	2,571	3,432	3,361	2,421	904	191	29	-
2002.....	777	1,150	1,549	2,015	2,784	3,497	3,376	2,347	847	192	21	-
2001.....	877	1,188	1,544	1,967	2,936	3,554	3,398	2,267	855	178	22	-
2000.....	879	1,290	1,564	2,179	2,982	3,761	3,396	2,212	817	185	21	-
1999.....	922	1,251	1,567	2,249	3,166	3,758	3,399	2,089	787	165	16	-

White, male

2003.....	473	698	966	1,163	1,457	1,777	1,607	1,030	326	69	9	-
2002.....	481	726	911	1,218	1,527	1,794	1,614	976	299	47	6	-
2001.....	570	710	899	1,166	1,615	1,766	1,599	965	289	52	5	-
2000.....	556	771	887	1,244	1,631	1,917	1,548	917	260	59	4	-
1999.....	576	751	892	1,248	1,705	1,911	1,494	839	273	47	7	-

White, female

2003.....	281	440	577	744	1,114	1,655	1,754	1,391	578	122	20	-
2002.....	296	424	638	797	1,257	1,703	1,762	1,371	548	145	15	-
2001.....	307	478	645	801	1,321	1,788	1,799	1,302	566	126	17	-
2000.....	323	519	677	935	1,351	1,844	1,848	1,295	557	126	17	-
1999.....	346	500	675	1,001	1,461	1,847	1,905	1,250	514	118	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

Black, both sexes

2003.....	127	96	150	139	172	156	122	64	29	13	1	-
2002.....	128	139	147	166	192	156	135	74	24	5	-	-
2001.....	119	139	137	177	174	148	140	83	36	7	2	-
2000.....	103	119	138	150	170	146	100	80	30	6	-	-
1999.....	99	138	126	154	177	179	109	79	35	11	2	-

Black, male

2003.....	74	55	83	70	75	63	47	18	11	4	-	-
2002.....	71	77	84	92	103	69	54	27	9	4	-	-
2001.....	80	78	76	97	94	66	59	35	11	2	-	-
2000.....	68	82	76	74	96	65	38	23	12	3	-	-
1999.....	64	81	74	90	84	99	33	33	11	5	2	-

Black, female

2003.....	53	41	67	69	97	93	75	46	18	9	1	-
2002.....	57	62	63	74	89	87	81	47	15	1	-	-
2001.....	39	61	61	80	80	82	81	48	25	5	2	-
2000.....	35	37	62	76	74	81	62	57	18	3	-	-
1999.....	35	57	52	64	93	80	76	46	24	6	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

American Indian 1/, both sexes

2003.....	1	11	6	9	11	10	5	1	1	1	-	-
2002.....	4	4	7	4	7	6	3	1	2	1	-	-
2001.....	6	6	11	18	6	11	8	4	4	-	-	-
2000.....	7	4	6	11	10	11	15	4	1	-	-	-
1999.....	1	3	6	7	11	8	4	3	2	-	-	-

American Indian 1/, male

2003.....	-	6	4	4	6	3	3	-	-	-	-	-
2002.....	2	2	5	3	3	4	-	-	-	-	-	-
2001.....	2	4	7	9	4	5	2	2	1	-	-	-
2000.....	5	2	3	3	5	6	8	1	-	-	-	-
1999.....	1	2	4	4	3	7	1	1	1	-	-	-

American Indian 1/, female

2003.....	1	5	2	5	5	7	2	1	1	1	-	-
2002.....	2	2	2	1	4	2	3	1	2	1	-	-
2001.....	4	2	4	9	2	6	6	2	3	-	-	-
2000.....	2	2	3	8	5	5	7	3	1	-	-	-
1999.....	-	1	2	3	8	1	3	2	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

Asian or Pacific Islander 2/, both sexes

2003.....	17	27	36	49	61	76	55	34	15	8	-	-
2002.....	34	26	27	40	56	63	47	24	8	2	1	-
2001.....	19	34	39	31	49	63	47	38	9	1	-	-
2000.....	26	21	28	46	50	66	40	34	8	2	2	-
1999.....	23	22	41	31	53	64	33	27	10	2	-	-

Asian or Pacific Islander 2/, male

2003.....	11	14	22	23	31	39	24	17	8	5	-	-
2002.....	20	18	15	21	19	31	28	10	7	-	-	-
2001.....	11	25	26	14	26	36	25	18	4	1	-	-
2000.....	17	10	18	21	31	30	22	18	2	-	-	-
1999.....	11	17	26	13	29	35	16	19	7	1	-	-

Asian or Pacific Islander 2/, female

2003.....	6	13	14	26	30	37	31	17	7	3	-	-
2002.....	14	8	12	19	37	32	19	14	1	2	1	-
2001.....	8	9	13	17	23	27	22	20	5	-	-	-
2000.....	9	11	10	25	19	36	18	16	6	2	2	-
1999.....	12	5	15	18	24	29	17	8	3	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Leukemia (C91-C95)

All races, both sexes

2003.....	770	1,102	1,503	1,911	2,795	3,532	3,405	2,317	1,091	350	28	-
2002.....	853	1,111	1,500	1,998	2,820	3,392	3,371	2,294	1,086	281	35	-
2001.....	845	1,073	1,462	1,963	2,923	3,541	3,155	2,256	1,008	271	50	-
2000.....	789	1,041	1,373	2,073	2,937	3,515	3,218	2,194	1,041	271	37	-
1999.....	744	1,112	1,404	2,078	2,934	3,401	3,059	2,151	984	258	31	-

All races, male

2003.....	454	683	910	1,215	1,715	2,054	1,826	1,102	416	91	6	-
2002.....	486	667	891	1,242	1,768	1,995	1,809	1,062	420	95	15	-
2001.....	467	632	920	1,216	1,746	2,023	1,676	1,041	380	89	17	-
2000.....	435	629	823	1,287	1,803	1,982	1,713	1,018	359	85	8	-
1999.....	431	636	853	1,275	1,779	1,946	1,579	964	375	75	9	-

All races, female

2003.....	316	419	593	696	1,080	1,478	1,579	1,215	675	259	22	-
2002.....	367	444	609	756	1,052	1,397	1,562	1,232	666	186	20	-
2001.....	378	441	542	747	1,177	1,518	1,479	1,215	628	182	33	-
2000.....	354	412	550	786	1,134	1,533	1,505	1,176	682	186	29	-
1999.....	313	476	551	803	1,155	1,455	1,480	1,187	609	183	22	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
White, both sexes												
2003.....	635	962	1,323	1,702	2,522	3,265	3,168	2,180	1,019	329	23	-
2002.....	715	944	1,302	1,793	2,568	3,110	3,152	2,138	1,019	260	30	-
2001.....	696	928	1,282	1,719	2,667	3,255	2,968	2,132	946	248	48	-
2000.....	669	900	1,188	1,870	2,690	3,272	3,026	2,053	967	250	32	-
1999.....	619	975	1,221	1,839	2,692	3,166	2,841	2,037	915	238	28	-
White, male												
2003.....	376	593	819	1,090	1,569	1,925	1,725	1,047	390	85	6	-
2002.....	415	580	779	1,118	1,618	1,847	1,700	1,000	396	90	12	-
2001.....	392	555	812	1,070	1,611	1,879	1,579	992	358	81	17	-
2000.....	366	550	717	1,176	1,651	1,855	1,616	949	338	76	7	-
1999.....	360	563	751	1,141	1,643	1,815	1,484	911	350	69	9	-
White, female												
2003.....	259	369	504	612	953	1,340	1,443	1,133	629	244	17	-
2002.....	300	364	523	675	950	1,263	1,452	1,138	623	170	18	-
2001.....	304	373	470	649	1,056	1,376	1,389	1,140	588	167	31	-
2000.....	303	350	471	694	1,039	1,417	1,410	1,104	629	174	25	-
1999.....	259	412	470	698	1,049	1,351	1,357	1,126	565	169	19	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Leukemia (C91-C95)

Black, both sexes

2003.....	111	113	151	171	224	219	189	115	67	16	5	-
2002.....	108	136	159	163	213	228	178	133	61	19	5	-
2001.....	112	112	137	207	208	238	155	109	55	21	1	-
2000.....	97	107	154	162	213	195	155	124	64	18	5	-
1999.....	96	106	146	201	206	196	177	96	58	18	3	-

Black, male

2003.....	67	76	75	101	120	110	79	46	23	5	-	-
2002.....	55	74	87	94	127	114	84	52	21	5	3	-
2001.....	61	63	84	127	108	111	76	43	18	7	-	-
2000.....	53	61	90	86	130	105	76	61	16	7	1	-
1999.....	56	61	81	112	112	108	70	42	20	4	-	-

Black, female

2003.....	44	37	76	70	104	109	110	69	44	11	5	-
2002.....	53	62	72	69	86	114	94	81	40	14	2	-
2001.....	51	49	53	80	100	127	79	66	37	14	1	-
2000.....	44	46	64	76	83	90	79	63	48	11	4	-
1999.....	40	45	65	89	94	88	107	54	38	14	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Leukemia (C91-C95)

American Indian 1/, both sexes

2003.....	6	4	6	4	10	6	6	6	-	-	-	-
2002.....	1	7	6	7	5	8	7	4	2	1	-	-
2001.....	6	10	8	8	7	1	4	2	1	1	-	-
2000.....	2	8	7	9	7	5	5	6	3	1	-	-
1999.....	6	4	4	7	6	6	4	3	2	-	-	-

American Indian 1/, male

2003.....	4	1	2	3	4	1	3	1	-	-	-	-
2002.....	1	3	4	4	3	4	5	1	1	-	-	-
2001.....	3	5	5	3	4	-	2	-	-	-	-	-
2000.....	2	4	3	5	6	1	3	2	1	1	-	-
1999.....	4	1	3	7	4	6	1	2	1	-	-	-

American Indian 1/, female

2003.....	2	3	4	1	6	5	3	5	-	-	-	-
2002.....	-	4	2	3	2	4	2	3	1	1	-	-
2001.....	3	5	3	5	3	1	2	2	1	1	-	-
2000.....	-	4	4	4	1	4	2	4	2	-	-	-
1999.....	2	3	1	-	2	-	3	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Leukemia (C91-C95)

Asian or Pacific Islander 2/, both sexes

2003.....	18	23	23	34	39	42	42	16	5	5	-	-
2002.....	29	24	33	35	34	46	34	19	4	1	-	-
2001.....	31	23	35	29	41	47	28	13	6	1	1	-
2000.....	21	26	24	32	27	43	32	11	7	2	-	-
1999.....	23	27	33	31	30	33	37	15	9	2	-	-

Asian or Pacific Islander 2/, male

2003.....	7	13	14	21	22	18	19	8	3	1	-	-
2002.....	15	10	21	26	20	30	20	9	2	-	-	-
2001.....	11	9	19	16	23	33	19	6	4	1	-	-
2000.....	14	14	13	20	16	21	18	6	4	1	-	-
1999.....	11	11	18	15	20	17	24	9	4	2	-	-

Asian or Pacific Islander 2/, female

2003.....	11	10	9	13	17	24	23	8	2	4	-	-
2002.....	14	14	12	9	14	16	14	10	2	1	-	-
2001.....	20	14	16	13	18	14	9	7	2	-	1	-
2000.....	7	12	11	12	11	22	14	5	3	1	-	-
1999.....	12	16	15	16	10	16	13	6	5	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races, both sexes

2003.....	473	698	961	1,280	1,749	2,127	1,870	1,165	464	78	4	-
2002.....	466	711	958	1,348	1,828	2,067	1,904	1,171	426	84	5	-
2001.....	430	721	950	1,332	1,849	2,101	1,785	1,123	400	71	3	-
2000.....	455	611	919	1,374	1,805	2,097	1,799	1,109	401	84	10	-
1999.....	424	727	967	1,347	1,825	2,023	1,765	1,040	389	85	5	-

All races, male

2003.....	287	415	580	725	929	1,052	889	520	174	33	1	-
2002.....	274	403	563	748	998	1,090	928	507	159	37	3	-
2001.....	252	410	535	741	984	1,089	837	446	152	23	3	-
2000.....	248	321	487	751	946	1,087	845	462	137	33	3	-
1999.....	246	411	557	729	940	1,026	820	450	133	26	3	-

All races, female

2003.....	186	283	381	555	820	1,075	981	645	290	45	3	-
2002.....	192	308	395	600	830	977	976	664	267	47	2	-
2001.....	178	311	415	591	865	1,012	948	677	248	48	-	-
2000.....	207	290	432	623	859	1,010	954	647	264	51	7	-
1999.....	178	316	410	618	885	997	945	590	256	59	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

White, both sexes

2003.....	330	530	761	1,016	1,434	1,782	1,613	1,016	409	64	1	-
2002.....	334	535	742	1,071	1,516	1,777	1,670	1,044	361	74	3	-
2001.....	320	536	711	1,061	1,537	1,795	1,538	978	354	60	2	-
2000.....	331	442	722	1,109	1,496	1,766	1,578	947	333	66	6	-
1999.....	299	527	725	1,093	1,510	1,697	1,549	897	338	75	2	-

White, male

2003.....	203	320	468	592	791	914	779	462	157	31	-	-
2002.....	210	310	449	610	853	959	826	459	135	36	2	-
2001.....	194	307	409	604	833	953	731	397	135	20	2	-
2000.....	181	240	392	621	793	932	758	409	116	29	3	-
1999.....	188	306	441	610	784	882	729	398	112	25	1	-

White, female

2003.....	127	210	293	424	643	868	834	554	252	33	1	-
2002.....	124	225	293	461	663	818	844	585	226	38	1	-
2001.....	126	229	302	457	704	842	807	581	219	40	-	-
2000.....	150	202	330	488	703	834	820	538	217	37	3	-
1999.....	111	221	284	483	726	815	820	499	226	50	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Black, both sexes

2003.....	128	148	179	244	282	309	234	137	52	14	2	-
2002.....	117	159	188	251	281	260	210	118	58	9	2	-
2001.....	96	167	217	239	284	276	226	135	44	10	1	-
2000.....	118	158	181	247	284	300	205	150	62	14	3	-
1999.....	117	181	215	219	278	286	195	137	48	10	3	-

Black, male

2003.....	76	87	102	119	120	126	99	53	15	2	-	-
2002.....	57	83	102	126	134	114	92	44	21	1	1	-
2001.....	49	93	115	119	134	123	100	43	16	3	1	-
2000.....	62	75	89	119	144	142	82	47	19	3	-	-
1999.....	54	95	102	101	138	125	85	47	19	1	2	-

Black, female

2003.....	52	61	77	125	162	183	135	84	37	12	2	-
2002.....	60	76	86	125	147	146	118	74	37	8	1	-
2001.....	47	74	102	120	150	153	126	92	28	7	-	-
2000.....	56	83	92	128	140	158	123	103	43	11	3	-
1999.....	63	86	113	118	140	161	110	90	29	9	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

American Indian 1/, both sexes

2003.....	3	6	6	5	8	12	7	4	2	-	1	-
2002.....	7	5	5	6	5	8	3	1	-	-	-	-
2001.....	4	5	6	8	8	6	5	4	1	-	-	-
2000.....	2	3	5	1	3	9	3	3	3	3	-	-
1999.....	1	2	4	6	10	12	3	1	1	-	-	-

American Indian 1/, male

2003.....	2	1	3	4	3	2	3	3	1	-	1	-
2002.....	5	2	-	2	2	3	3	1	-	-	-	-
2001.....	3	2	3	5	3	2	2	2	-	-	-	-
2000.....	1	3	2	1	1	3	-	2	-	1	-	-
1999.....	-	1	1	4	7	5	1	1	1	-	-	-

American Indian 1/, female

2003.....	1	5	3	1	5	10	4	1	1	-	-	-
2002.....	2	3	5	4	3	5	-	-	-	-	-	-
2001.....	1	3	3	3	5	4	3	2	1	-	-	-
2000.....	1	-	3	-	2	6	3	1	3	2	-	-
1999.....	1	1	3	2	3	7	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Asian or Pacific Islander 2/, both sexes

2003.....	12	14	15	15	25	24	16	8	1	-	-	-
2002.....	8	12	23	20	26	22	21	8	7	1	-	-
2001.....	10	13	16	24	20	24	16	6	1	1	-	-
2000.....	4	8	11	17	22	22	13	9	3	1	1	-
1999.....	7	17	23	29	27	28	18	5	2	-	-	-

Asian or Pacific Islander 2/, male

2003.....	6	7	7	10	15	10	8	2	1	-	-	-
2002.....	2	8	12	10	9	14	7	3	3	-	-	-
2001.....	6	8	8	13	14	11	4	4	1	-	-	-
2000.....	4	3	4	10	8	10	5	4	2	-	-	-
1999.....	4	9	13	14	11	14	5	4	1	-	-	-

Asian or Pacific Islander 2/, female

2003.....	6	7	8	5	10	14	8	6	-	-	-	-
2002.....	6	4	11	10	17	8	14	5	4	1	-	-
2001.....	4	5	8	11	6	13	12	2	-	1	-	-
2000.....	-	5	7	7	14	12	8	5	1	1	1	-
1999.....	3	8	10	15	16	14	13	1	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races, both sexes

2003.....	3	1	1	6	9	11	3	10	3	-	-	-
2002.....	5	2	1	7	11	12	13	9	5	2	-	-
2001.....	3	2	6	11	12	11	17	11	8	1	-	-
2000.....	3	6	6	7	15	13	16	13	7	1	-	-
1999.....	6	6	6	10	7	14	11	9	4	-	-	-

All races, male

2003.....	1	1	1	3	6	7	1	4	1	-	-	-
2002.....	2	1	1	5	10	4	8	5	1	1	-	-
2001.....	2	1	4	8	10	5	9	4	2	-	-	-
2000.....	2	5	4	7	5	10	7	6	3	-	-	-
1999.....	3	5	2	5	3	9	5	3	-	-	-	-

All races, female

2003.....	2	-	-	3	3	4	2	6	2	-	-	-
2002.....	3	1	-	2	1	8	5	4	4	1	-	-
2001.....	1	1	2	3	2	6	8	7	6	1	-	-
2000.....	1	1	2	-	10	3	9	7	4	1	-	-
1999.....	3	1	4	5	4	5	6	6	4	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

White, both sexes

2003.....	3	1	1	5	9	11	2	9	3	-	-	-
2002.....	4	2	1	6	11	9	11	9	2	2	-	-
2001.....	3	-	5	10	11	9	14	10	8	1	-	-
2000.....	3	5	6	7	14	11	15	12	6	1	-	-
1999.....	5	5	6	10	7	13	10	8	4	-	-	-

White, male

2003.....	1	1	1	3	6	7	-	4	1	-	-	-
2002.....	2	1	1	4	10	4	6	5	-	1	-	-
2001.....	2	-	4	7	10	4	8	3	2	-	-	-
2000.....	2	4	4	7	4	9	7	5	2	-	-	-
1999.....	2	4	2	5	3	8	5	3	-	-	-	-

White, female

2003.....	2	-	-	2	3	4	2	5	2	-	-	-
2002.....	2	1	-	2	1	5	5	4	2	1	-	-
2001.....	1	-	1	3	1	5	6	7	6	1	-	-
2000.....	1	1	2	-	10	2	8	7	4	1	-	-
1999.....	3	1	4	5	4	5	5	5	4	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Black, both sexes

2003.....	-	-	-	1	-	-	-	1	-	-	-	-
2002.....	-	-	-	1	-	2	1	-	2	-	-	-
2001.....	-	2	1	1	1	1	3	1	-	-	-	-
2000.....	-	1	-	-	-	2	1	1	1	-	-	-
1999.....	1	1	-	-	-	1	-	1	-	-	-	-

Black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	1	-	1	-	1	1	1	-	-	-	-
2000.....	-	1	-	-	-	1	-	1	1	-	-	-
1999.....	1	1	-	-	-	1	-	-	-	-	-	-

Black, female

2003.....	-	-	-	1	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	2	-	-	-
2001.....	-	1	1	-	1	-	2	-	-	-	-	-
2000.....	-	-	-	-	-	1	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races, both sexes

2003.....	3,377	4,446	5,374	6,299	7,978	9,798	9,082	6,421	3,159	847	129	5
2002.....	3,280	4,320	5,403	6,321	8,436	9,718	8,967	6,452	3,129	853	136	1
2001.....	3,319	4,278	5,163	6,406	8,617	9,902	8,798	6,343	3,019	845	119	-
2000.....	3,193	4,088	5,205	6,819	8,897	10,087	8,773	6,214	2,937	874	140	4
1999.....	3,230	4,249	5,614	7,265	9,569	10,451	9,205	6,485	2,955	851	112	2

All races, male

2003.....	1,869	2,525	3,048	3,544	4,256	5,042	4,279	2,648	1,055	242	28	3
2002.....	1,906	2,399	3,122	3,548	4,593	4,969	4,177	2,690	1,100	226	25	1
2001.....	1,933	2,398	2,928	3,501	4,676	4,962	4,070	2,569	1,001	239	26	-
2000.....	1,773	2,286	2,937	3,791	4,781	5,098	4,063	2,460	989	218	22	4
1999.....	1,824	2,416	3,181	3,984	5,173	5,240	4,199	2,586	986	213	21	1

All races, female

2003.....	1,508	1,921	2,326	2,755	3,722	4,756	4,803	3,773	2,104	605	101	2
2002.....	1,374	1,921	2,281	2,773	3,843	4,749	4,790	3,762	2,029	627	111	-
2001.....	1,386	1,880	2,235	2,905	3,941	4,940	4,728	3,774	2,018	606	93	-
2000.....	1,420	1,802	2,268	3,028	4,116	4,989	4,710	3,754	1,948	656	118	-
1999.....	1,406	1,833	2,433	3,281	4,396	5,211	5,006	3,899	1,969	638	91	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2003.....	2,703	3,683	4,567	5,417	7,025	8,793	8,285	5,931	2,894	762	115	5
2002.....	2,645	3,601	4,542	5,404	7,403	8,707	8,150	5,906	2,874	773	123	1
2001.....	2,661	3,522	4,313	5,469	7,537	8,900	7,974	5,841	2,777	769	103	-
2000.....	2,554	3,432	4,362	5,908	7,822	9,026	8,040	5,722	2,730	785	115	3
1999.....	2,627	3,496	4,709	6,334	8,430	9,385	8,397	5,949	2,689	769	98	2

White, male

2003.....	1,507	2,096	2,607	3,064	3,771	4,562	3,943	2,464	968	215	24	3
2002.....	1,554	2,008	2,634	3,095	4,075	4,473	3,843	2,483	1,015	206	24	1
2001.....	1,543	1,965	2,446	3,000	4,145	4,486	3,717	2,376	910	223	22	-
2000.....	1,437	1,921	2,430	3,328	4,244	4,553	3,763	2,269	919	195	20	3
1999.....	1,479	2,002	2,685	3,484	4,594	4,751	3,872	2,387	898	188	20	1

White, female

2003.....	1,196	1,587	1,960	2,353	3,254	4,231	4,342	3,467	1,926	547	91	2
2002.....	1,091	1,593	1,908	2,309	3,328	4,234	4,307	3,423	1,859	567	99	-
2001.....	1,118	1,557	1,867	2,469	3,392	4,414	4,257	3,465	1,867	546	81	-
2000.....	1,117	1,511	1,932	2,580	3,578	4,473	4,277	3,453	1,811	590	95	-
1999.....	1,148	1,494	2,024	2,850	3,836	4,634	4,525	3,562	1,791	581	78	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2003.....	572	649	683	734	774	819	654	411	220	67	9	-
2002.....	531	619	728	777	847	853	667	451	218	67	12	-
2001.....	559	626	716	789	894	810	673	418	209	66	14	-
2000.....	538	557	710	775	882	876	609	405	167	82	20	1
1999.....	513	634	788	773	948	881	668	455	223	69	14	-

Black, male

2003.....	311	378	375	400	391	386	279	146	69	16	1	-
2002.....	298	337	414	390	428	408	274	164	68	16	1	-
2001.....	332	365	398	433	450	385	273	159	73	14	4	-
2000.....	285	312	430	392	446	448	247	148	52	19	1	1
1999.....	296	365	443	423	483	403	268	156	71	22	1	-

Black, female

2003.....	261	271	308	334	383	433	375	265	151	51	8	-
2002.....	233	282	314	387	419	445	393	287	150	51	11	-
2001.....	227	261	318	356	444	425	400	259	136	52	10	-
2000.....	253	245	280	383	436	428	362	257	115	63	19	-
1999.....	217	269	345	350	465	478	400	299	152	47	13	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

American Indian 1/, both sexes

2003.....	27	25	23	31	37	35	23	15	8	8	1	-
2002.....	18	24	33	20	39	32	26	18	13	1	-	-
2001.....	19	33	27	34	33	39	24	18	10	2	-	-
2000.....	16	14	33	26	47	36	19	15	8	3	2	-
1999.....	21	35	28	38	34	37	33	15	5	1	-	-

American Indian 1/, male

2003.....	13	15	10	17	21	20	8	7	4	6	-	-
2002.....	10	10	19	13	24	17	10	4	2	-	-	-
2001.....	11	16	16	17	20	16	12	6	4	-	-	-
2000.....	10	8	17	14	21	18	4	5	3	1	-	-
1999.....	11	13	12	18	17	16	13	7	1	-	-	-

American Indian 1/, female

2003.....	14	10	13	14	16	15	15	8	4	2	1	-
2002.....	8	14	14	7	15	15	16	14	11	1	-	-
2001.....	8	17	11	17	13	23	12	12	6	2	-	-
2000.....	6	6	16	12	26	18	15	10	5	2	2	-
1999.....	10	22	16	20	17	21	20	8	4	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Asian or Pacific Islander 2/, both sexes

2003.....	75	89	101	117	142	151	120	64	37	10	4	-
2002.....	86	76	100	120	147	126	124	77	24	12	1	-
2001.....	80	97	107	114	153	153	127	66	23	8	2	-
2000.....	85	85	100	110	146	149	105	72	32	4	3	-
1999.....	69	84	89	120	157	148	107	66	38	12	-	-

Asian or Pacific Islander 2/, male

2003.....	38	36	56	63	73	74	49	31	14	5	3	-
2002.....	44	44	55	50	66	71	50	39	15	4	-	-
2001.....	47	52	68	51	61	75	68	28	14	2	-	-
2000.....	41	45	60	57	70	79	49	38	15	3	1	-
1999.....	38	36	41	59	79	70	46	36	16	3	-	-

Asian or Pacific Islander 2/, female

2003.....	37	53	45	54	69	77	71	33	23	5	1	-
2002.....	42	32	45	70	81	55	74	38	9	8	1	-
2001.....	33	45	39	63	92	78	59	38	9	6	2	-
2000.....	44	40	40	53	76	70	56	34	17	1	2	-
1999.....	31	48	48	61	78	78	61	30	22	9	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races, both sexes

2003.....	387	506	732	955	1,399	2,123	2,566	2,203	1,215	337	40	-
2002.....	387	561	632	928	1,420	2,137	2,460	2,052	1,162	366	33	-
2001.....	415	553	700	1,000	1,574	2,192	2,462	2,102	1,141	351	49	-
2000.....	397	513	711	1,007	1,539	2,142	2,503	2,103	1,143	314	40	1
1999.....	386	461	672	1,029	1,655	2,151	2,321	1,998	1,059	337	49	-

All races, male

2003.....	218	279	406	524	810	1,114	1,265	955	411	94	10	-
2002.....	196	330	350	543	772	1,145	1,187	900	398	99	8	-
2001.....	218	279	394	563	876	1,199	1,204	882	404	102	10	-
2000.....	225	270	387	587	828	1,120	1,199	881	362	86	7	1
1999.....	220	255	401	590	932	1,119	1,099	839	367	82	15	-

All races, female

2003.....	169	227	326	431	589	1,009	1,301	1,248	804	243	30	-
2002.....	191	231	282	385	648	992	1,273	1,152	764	267	25	-
2001.....	197	274	306	437	698	993	1,258	1,220	737	249	39	-
2000.....	172	243	324	420	711	1,022	1,304	1,222	781	228	33	-
1999.....	166	206	271	439	723	1,032	1,222	1,159	692	255	34	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2003.....	318	423	638	820	1,261	1,957	2,374	2,065	1,127	311	30	-
2002.....	309	470	553	812	1,279	1,961	2,261	1,922	1,076	324	28	-
2001.....	353	476	599	884	1,404	2,017	2,275	1,941	1,050	314	39	-
2000.....	333	447	611	888	1,351	1,962	2,340	1,960	1,066	291	35	-
1999.....	320	389	572	886	1,499	1,959	2,161	1,870	996	303	44	-

White, male

2003.....	182	230	353	464	736	1,039	1,176	905	379	86	8	-
2002.....	162	275	307	479	710	1,060	1,095	850	367	85	8	-
2001.....	193	243	340	506	798	1,111	1,116	813	377	94	8	-
2000.....	192	240	324	517	730	1,032	1,136	818	341	82	7	-
1999.....	186	216	353	516	856	1,027	1,035	792	352	76	14	-

White, female

2003.....	136	193	285	356	525	918	1,198	1,160	748	225	22	-
2002.....	147	195	246	333	569	901	1,166	1,072	709	239	20	-
2001.....	160	233	259	378	606	906	1,159	1,128	673	220	31	-
2000.....	141	207	287	371	621	930	1,204	1,142	725	209	28	-
1999.....	134	173	219	370	643	932	1,126	1,078	644	227	30	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes

2003.....	59	66	78	107	116	133	157	98	70	22	8	-
2002.....	64	77	67	100	112	142	160	108	68	34	4	-
2001.....	55	64	85	101	138	141	157	119	78	33	9	-
2000.....	58	57	78	96	152	150	134	118	63	20	4	1
1999.....	56	65	84	124	134	165	134	108	52	30	4	-

Black, male

2003.....	29	39	44	50	65	58	72	34	23	6	1	-
2002.....	27	50	33	51	48	65	70	41	22	11	-	-
2001.....	22	29	45	51	68	68	73	47	22	7	2	-
2000.....	31	27	49	60	71	73	46	45	18	3	-	1
1999.....	27	36	39	65	64	76	49	38	10	6	-	-

Black, female

2003.....	30	27	34	57	51	75	85	64	47	16	7	-
2002.....	37	27	34	49	64	77	90	67	46	23	4	-
2001.....	33	35	40	50	70	73	84	72	56	26	7	-
2000.....	27	30	29	36	81	77	88	73	45	17	4	-
1999.....	29	29	45	59	70	89	85	70	42	24	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

American Indian 1/, both sexes

2003.....	-	4	4	6	3	4	5	4	1	1	1	-
2002.....	2	1	3	2	7	4	4	1	1	2	-	-
2001.....	1	2	1	4	12	10	4	7	2	-	-	-
2000.....	1	2	4	4	6	5	2	6	1	1	1	-
1999.....	2	4	3	7	6	4	5	4	2	1	1	-

American Indian 1/, male

2003.....	-	3	2	1	2	3	3	2	1	-	-	-
2002.....	-	-	3	2	2	4	1	1	-	1	-	-
2001.....	1	1	1	3	4	4	2	4	1	-	-	-
2000.....	-	-	3	1	5	1	2	2	-	-	-	-
1999.....	1	2	2	2	3	3	3	2	-	-	1	-

American Indian 1/, female

2003.....	-	1	2	5	1	1	2	2	-	1	1	-
2002.....	2	1	-	-	5	-	3	-	1	1	-	-
2001.....	-	1	-	1	8	6	2	3	1	-	-	-
2000.....	1	2	1	3	1	4	-	4	1	1	1	-
1999.....	1	2	1	5	3	1	2	2	2	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Asian or Pacific Islander 2/, both sexes

2003.....	10	13	12	22	19	29	30	36	17	3	1	-
2002.....	12	13	9	14	22	30	35	21	17	6	1	-
2001.....	6	11	15	11	20	24	26	35	11	4	1	-
2000.....	5	7	18	19	30	25	27	19	13	2	-	-
1999.....	8	3	13	12	16	23	21	16	9	3	-	-

Asian or Pacific Islander 2/, male

2003.....	7	7	7	9	7	14	14	14	8	2	1	-
2002.....	7	5	7	11	12	16	21	8	9	2	-	-
2001.....	2	6	8	3	6	16	13	18	4	1	-	-
2000.....	2	3	11	9	22	14	15	16	3	1	-	-
1999.....	6	1	7	7	9	13	12	7	5	-	-	-

Asian or Pacific Islander 2/, female

2003.....	3	6	5	13	12	15	16	22	9	1	-	-
2002.....	5	8	2	3	10	14	14	13	8	4	1	-
2001.....	4	5	7	8	14	8	13	17	7	3	1	-
2000.....	3	4	7	10	8	11	12	3	10	1	-	-
1999.....	2	2	6	5	7	10	9	9	4	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Anemias (D50-D64)

All races, both sexes

2003.....	104	121	155	184	300	447	689	784	721	342	72	-
2002.....	119	121	143	184	324	474	687	787	665	322	78	-
2001.....	120	119	130	186	306	525	701	808	659	294	90	-
2000.....	117	127	133	210	305	482	689	790	638	290	78	-
1999.....	98	105	126	175	311	497	667	775	647	316	82	-

All races, male

2003.....	48	67	89	104	147	203	274	270	196	75	8	-
2002.....	52	57	74	77	167	204	250	256	182	81	13	-
2001.....	58	53	58	91	146	220	276	276	177	52	15	-
2000.....	56	66	64	95	141	198	267	237	183	58	14	-
1999.....	48	43	62	92	134	217	267	261	172	73	12	-

All races, female

2003.....	56	54	66	80	153	244	415	514	525	267	64	-
2002.....	67	64	69	107	157	270	437	531	483	241	65	-
2001.....	62	66	72	95	160	305	425	532	482	242	75	-
2000.....	61	61	69	115	164	284	422	553	455	232	64	-
1999.....	50	62	64	83	177	280	400	514	475	243	70	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Anemias (D50-D64)

White, both sexes

2003.....	44	56	97	135	239	379	606	718	638	306	65	-
2002.....	50	71	90	141	258	387	603	708	585	284	73	-
2001.....	61	69	99	146	247	446	613	728	599	266	69	-
2000.....	56	84	85	160	250	414	617	718	566	261	68	-
1999.....	41	46	76	135	254	428	604	692	583	277	68	-

White, male

2003.....	22	32	53	78	111	175	249	249	174	64	8	-
2002.....	26	36	47	61	128	173	216	231	157	68	12	-
2001.....	28	31	44	74	113	193	248	259	159	46	11	-
2000.....	27	45	42	81	118	169	238	212	156	52	13	-
1999.....	20	16	32	75	107	190	244	230	160	66	12	-

White, female

2003.....	22	24	44	57	128	204	357	469	464	242	57	-
2002.....	24	35	43	80	130	214	387	477	428	216	61	-
2001.....	33	38	55	72	134	253	365	469	440	220	58	-
2000.....	29	39	43	79	132	245	379	506	410	209	55	-
1999.....	21	30	44	60	147	238	360	462	423	211	56	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Anemias (D50-D64)

Black, both sexes

2003.....	58	64	52	41	54	64	78	59	75	30	5	-
2002.....	65	47	47	41	59	80	74	70	68	34	4	-
2001.....	58	46	29	36	55	72	79	66	50	24	18	-
2000.....	59	40	43	43	46	58	64	64	61	28	10	-
1999.....	52	55	46	33	49	64	57	67	57	30	12	-

Black, male

2003.....	26	35	32	23	33	28	25	19	19	9	-	-
2002.....	25	20	26	16	35	28	31	21	22	9	1	-
2001.....	30	21	14	15	31	26	25	15	13	6	3	-
2000.....	29	18	19	13	18	23	23	20	22	5	1	-
1999.....	26	25	29	14	23	23	19	25	9	4	-	-

Black, female

2003.....	32	29	20	18	21	36	53	40	56	21	5	-
2002.....	40	27	21	25	24	52	43	49	46	25	3	-
2001.....	28	25	15	21	24	46	54	51	37	18	15	-
2000.....	30	22	24	30	28	35	41	44	39	23	9	-
1999.....	26	30	17	19	26	41	38	42	48	26	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Anemias (D50-D64)

American Indian 1/, both sexes

2003.....	1	-	2	4	1	1	1	-	1	-	-	-
2002.....	2	2	-	-	-	1	1	1	1	1	1	-
2001.....	-	1	-	-	1	4	3	6	-	3	1	-
2000.....	-	1	-	2	2	6	2	1	3	-	-	-
1999.....	2	1	3	1	2	-	-	2	-	1	2	-

American Indian 1/, male

2003.....	-	-	1	1	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	1	-	-
2001.....	-	1	-	-	1	1	-	2	-	-	-	-
2000.....	-	1	-	1	-	5	1	-	-	-	-	-
1999.....	-	1	1	-	-	-	-	1	-	1	-	-

American Indian 1/, female

2003.....	1	-	1	3	-	1	1	-	1	-	-	-
2002.....	2	1	-	-	-	1	-	1	1	-	1	-
2001.....	-	-	-	-	-	3	3	4	-	3	1	-
2000.....	-	-	-	1	2	1	1	1	3	-	-	-
1999.....	2	-	2	1	2	-	-	1	-	-	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)												
Asian or Pacific Islander 2/, both sexes												
2003.....	1	1	4	4	6	3	4	7	7	6	2	-
2002.....	2	1	6	2	7	6	9	8	11	3	-	-
2001.....	1	3	2	4	3	3	6	8	10	1	2	-
2000.....	2	2	5	5	7	4	6	7	8	1	-	-
1999.....	3	3	1	6	6	5	6	14	7	8	-	-
Asian or Pacific Islander 2/, male												
2003.....	-	-	3	2	2	-	-	2	3	2	-	-
2002.....	1	-	1	-	4	3	2	4	3	3	-	-
2001.....	-	-	-	2	1	-	3	-	5	-	1	-
2000.....	-	2	3	-	5	1	5	5	5	1	-	-
1999.....	2	1	-	3	4	4	4	5	3	2	-	-
Asian or Pacific Islander 2/, female												
2003.....	1	1	1	2	4	3	4	5	4	4	2	-
2002.....	1	1	5	2	3	3	7	4	8	-	-	-
2001.....	1	3	2	2	2	3	3	8	5	1	1	-
2000.....	2	-	2	5	2	3	1	2	3	-	-	-
1999.....	1	2	1	3	2	1	2	9	4	6	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diabetes mellitus (E10-E14)

All races, both sexes

2003.....	3,413	4,806	5,925	7,310	9,346	11,574	11,725	8,758	4,572	1,424	210	3
2002.....	3,390	4,510	5,512	7,249	9,460	11,703	11,579	8,797	4,375	1,357	195	1
2001.....	3,280	4,188	5,382	7,195	9,536	11,546	11,259	8,366	4,259	1,350	196	1
2000.....	3,037	3,992	5,194	7,054	9,620	11,607	10,577	8,002	4,104	1,250	200	-
1999.....	2,793	3,865	5,232	7,215	9,693	11,269	10,488	7,893	3,943	1,165	177	3

All races, male

2003.....	1,987	2,715	3,329	3,929	4,758	5,547	5,132	3,260	1,318	306	39	3
2002.....	1,977	2,616	3,067	3,742	4,675	5,402	4,893	3,248	1,254	314	28	1
2001.....	1,920	2,362	2,879	3,734	4,703	5,244	4,674	2,966	1,188	269	34	1
2000.....	1,697	2,173	2,770	3,556	4,670	5,288	4,450	2,789	1,165	260	20	-
1999.....	1,602	2,120	2,736	3,630	4,649	5,190	4,349	2,816	1,071	210	28	2

All races, female

2003.....	1,426	2,091	2,596	3,381	4,588	6,027	6,593	5,498	3,254	1,118	171	-
2002.....	1,413	1,894	2,445	3,507	4,785	6,301	6,686	5,549	3,121	1,043	167	-
2001.....	1,360	1,826	2,503	3,461	4,833	6,302	6,585	5,400	3,071	1,081	162	-
2000.....	1,340	1,819	2,424	3,498	4,950	6,319	6,127	5,213	2,939	990	180	-
1999.....	1,191	1,745	2,496	3,585	5,044	6,079	6,139	5,077	2,872	955	149	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diabetes mellitus (E10-E14)

White, both sexes

2003.....	2,420	3,552	4,435	5,570	7,364	9,481	9,877	7,507	3,926	1,208	162	3
2002.....	2,419	3,329	4,111	5,439	7,503	9,611	9,822	7,532	3,787	1,144	147	-
2001.....	2,368	3,066	4,027	5,501	7,584	9,488	9,582	7,224	3,685	1,145	143	1
2000.....	2,170	2,967	3,912	5,393	7,682	9,561	8,917	6,905	3,525	1,067	148	-
1999.....	1,988	2,754	3,855	5,547	7,683	9,315	8,902	6,746	3,379	995	137	1

White, male

2003.....	1,471	2,082	2,572	3,106	3,896	4,731	4,479	2,889	1,157	276	35	3
2002.....	1,462	2,004	2,367	2,928	3,872	4,630	4,290	2,891	1,141	274	23	-
2001.....	1,428	1,801	2,230	2,986	3,931	4,452	4,068	2,665	1,046	229	27	1
2000.....	1,269	1,662	2,161	2,849	3,890	4,508	3,911	2,473	1,023	228	16	-
1999.....	1,191	1,582	2,102	2,906	3,881	4,455	3,838	2,474	924	178	22	1

White, female

2003.....	949	1,470	1,863	2,464	3,468	4,750	5,398	4,618	2,769	932	127	-
2002.....	957	1,325	1,744	2,511	3,631	4,981	5,532	4,641	2,646	870	124	-
2001.....	940	1,265	1,797	2,515	3,653	5,036	5,514	4,559	2,639	916	116	-
2000.....	901	1,305	1,751	2,544	3,792	5,053	5,006	4,432	2,502	839	132	-
1999.....	797	1,172	1,753	2,641	3,802	4,860	5,064	4,272	2,455	817	115	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diabetes mellitus (E10-E14)

Black, both sexes

2003.....	862	1,102	1,265	1,488	1,672	1,763	1,536	1,035	543	185	45	-
2002.....	854	1,010	1,214	1,522	1,653	1,748	1,501	1,054	517	195	45	1
2001.....	807	988	1,166	1,460	1,685	1,757	1,411	988	509	179	49	-
2000.....	762	894	1,121	1,441	1,677	1,781	1,434	962	507	160	49	-
1999.....	716	954	1,197	1,444	1,693	1,652	1,370	982	496	144	37	2

Black, male

2003.....	436	551	631	693	708	671	525	289	121	22	3	-
2002.....	447	514	600	663	669	644	506	261	90	33	5	1
2001.....	421	490	553	638	658	663	499	240	121	29	7	-
2000.....	374	438	524	593	671	659	454	260	109	21	2	-
1999.....	358	461	534	631	637	595	429	281	123	23	5	1

Black, female

2003.....	426	551	634	795	964	1,092	1,011	746	422	163	42	-
2002.....	407	496	614	859	984	1,104	995	793	427	162	40	-
2001.....	386	498	613	822	1,027	1,094	912	748	388	150	42	-
2000.....	388	456	597	848	1,006	1,122	980	702	398	139	47	-
1999.....	358	493	663	813	1,056	1,057	941	701	373	121	32	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diabetes mellitus (E10-E14)

American Indian 1/, both sexes

2003.....	69	68	94	102	104	91	82	48	15	10	-	-
2002.....	61	83	84	113	110	94	68	32	16	2	-	-
2001.....	50	62	80	76	95	90	66	37	16	3	-	-
2000.....	53	59	75	77	109	81	52	36	13	2	2	-
1999.....	49	73	92	105	114	94	52	51	16	5	-	-

American Indian 1/, male

2003.....	42	33	41	44	51	39	33	14	9	2	-	-
2002.....	31	41	44	48	48	34	23	11	4	1	-	-
2001.....	30	28	37	33	35	41	22	14	2	-	-	-
2000.....	23	26	42	33	37	37	13	15	5	1	1	-
1999.....	32	33	45	40	42	43	19	16	4	1	-	-

American Indian 1/, female

2003.....	27	35	53	58	53	52	49	34	6	8	-	-
2002.....	30	42	40	65	62	60	45	21	12	1	-	-
2001.....	20	34	43	43	60	49	44	23	14	3	-	-
2000.....	30	33	33	44	72	44	39	21	8	1	1	-
1999.....	17	40	47	65	72	51	33	35	12	4	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diabetes mellitus (E10-E14)

Asian or Pacific Islander 2/, both sexes

2003.....	62	84	131	150	206	239	230	168	88	21	3	-
2002.....	56	88	103	175	194	250	188	179	55	16	3	-
2001.....	55	72	109	158	172	211	200	117	49	23	4	-
2000.....	52	72	86	143	152	184	174	99	59	21	1	-
1999.....	40	84	88	119	203	208	164	114	52	21	3	-

Asian or Pacific Islander 2/, male

2003.....	38	49	85	86	103	106	95	68	31	6	1	-
2002.....	37	57	56	103	86	94	74	85	19	6	-	-
2001.....	41	43	59	77	79	88	85	47	19	11	-	-
2000.....	31	47	43	81	72	84	72	41	28	10	1	-
1999.....	21	44	55	53	89	97	63	45	20	8	1	-

Asian or Pacific Islander 2/, female

2003.....	24	35	46	64	103	133	135	100	57	15	2	-
2002.....	19	31	47	72	108	156	114	94	36	10	3	-
2001.....	14	29	50	81	93	123	115	70	30	12	4	-
2000.....	21	25	43	62	80	100	102	58	31	11	-	-
1999.....	19	40	33	66	114	111	101	69	32	13	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

All races, both sexes

2003.....	61	77	91	125	215	428	564	727	570	301	74	-
2002.....	63	77	91	150	232	452	680	834	667	312	93	-
2001.....	62	64	89	131	243	439	661	798	684	324	83	-
2000.....	53	70	103	192	268	505	780	947	781	302	90	-
1999.....	61	76	94	149	280	472	776	944	822	380	94	-

All races, male

2003.....	29	38	41	63	103	175	225	243	139	49	11	-
2002.....	28	43	50	70	116	193	241	283	169	49	13	-
2001.....	29	35	46	63	111	172	254	228	171	55	3	-
2000.....	21	45	52	87	134	207	264	289	182	52	13	-
1999.....	30	42	56	72	149	205	289	301	196	71	9	-

All races, female

2003.....	32	39	50	62	112	253	339	484	431	252	63	-
2002.....	35	34	41	80	116	259	439	551	498	263	80	-
2001.....	33	29	43	68	132	267	407	570	513	269	80	-
2000.....	32	25	51	105	134	298	516	658	599	250	77	-
1999.....	31	34	38	77	131	267	487	643	626	309	85	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

White, both sexes

2003.....	38	60	67	101	176	369	479	646	498	258	55	-
2002.....	49	54	67	114	189	387	589	726	591	274	77	-
2001.....	39	45	72	98	205	365	557	716	604	286	76	-
2000.....	36	52	78	159	211	425	687	836	712	255	71	-
1999.....	48	46	72	115	210	392	658	833	751	326	80	-

White, male

2003.....	19	26	28	53	79	144	187	218	120	42	7	-
2002.....	20	27	32	55	95	169	200	250	154	42	10	-
2001.....	14	22	33	43	90	145	214	203	145	48	2	-
2000.....	13	30	35	70	102	171	233	256	167	41	11	-
1999.....	23	21	43	52	106	165	242	265	175	55	7	-

White, female

2003.....	19	34	39	48	97	225	292	428	378	216	48	-
2002.....	29	27	35	59	94	218	389	476	437	232	67	-
2001.....	25	23	39	55	115	220	343	513	459	238	74	-
2000.....	23	22	43	89	109	254	454	580	545	214	60	-
1999.....	25	25	29	63	104	227	416	568	576	271	73	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

Black, both sexes

2003.....	21	17	22	19	35	53	77	73	66	35	15	-
2002.....	13	23	23	33	38	61	79	90	68	33	13	-
2001.....	21	17	13	28	33	62	88	72	69	30	7	-
2000.....	16	18	23	29	49	68	80	96	61	41	16	-
1999.....	12	26	21	29	61	69	107	97	64	47	14	-

Black, male

2003.....	10	12	12	9	22	30	33	22	16	5	3	-
2002.....	8	16	17	14	17	21	36	27	12	3	2	-
2001.....	14	12	10	18	20	25	31	24	21	5	1	-
2000.....	8	15	15	15	29	32	27	31	12	11	1	-
1999.....	7	18	12	17	38	34	40	30	19	13	2	-

Black, female

2003.....	11	5	10	10	13	23	44	51	50	30	12	-
2002.....	5	7	6	19	21	40	43	63	56	30	11	-
2001.....	7	5	3	10	13	37	57	48	48	25	6	-
2000.....	8	3	8	14	20	36	53	65	49	30	15	-
1999.....	5	8	9	12	23	35	67	67	45	34	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

American Indian 1/, both sexes

2003.....	2	-	-	1	3	5	2	2	2	2	1	-
2002.....	1	-	-	1	3	1	5	6	3	3	1	-
2001.....	2	1	2	3	3	3	6	5	2	1	-	-
2000.....	1	-	-	2	4	5	3	8	3	1	3	-
1999.....	1	3	1	3	3	3	4	4	3	1	-	-

American Indian 1/, male

2003.....	-	-	-	-	2	-	2	-	-	-	-	-
2002.....	-	-	-	-	3	-	1	2	-	2	-	-
2001.....	1	-	2	1	1	-	3	-	-	-	-	-
2000.....	-	-	-	1	1	2	1	-	1	-	1	-
1999.....	-	2	1	1	1	2	3	1	-	-	-	-

American Indian 1/, female

2003.....	2	-	-	1	1	5	-	2	2	2	1	-
2002.....	1	-	-	1	-	1	4	4	3	1	1	-
2001.....	1	1	-	2	2	3	3	5	2	1	-	-
2000.....	1	-	-	1	3	3	2	8	2	1	2	-
1999.....	1	1	-	2	2	1	1	3	3	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	2	4	1	1	6	6	4	6	3	-
2002.....	-	-	1	2	2	3	7	12	5	2	2	-
2001.....	-	1	2	2	2	9	10	5	9	7	-	-
2000.....	-	-	2	2	4	7	10	7	5	5	-	-
1999.....	-	1	-	2	6	8	7	10	4	6	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	1	1	-	1	3	3	3	2	1	-
2002.....	-	-	1	1	1	3	4	4	3	2	1	-
2001.....	-	1	1	1	-	2	6	1	5	2	-	-
2000.....	-	-	2	1	2	2	3	2	2	-	-	-
1999.....	-	1	-	2	4	4	4	5	2	3	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	1	3	1	-	3	3	1	4	2	-
2002.....	-	-	-	1	1	-	3	8	2	-	1	-
2001.....	-	-	1	1	2	7	4	4	4	5	-	-
2000.....	-	-	-	1	2	5	7	5	3	5	-	-
1999.....	-	-	-	-	2	4	3	5	2	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malnutrition (E40-E46)

All races, both sexes

2003.....	58	74	86	118	202	413	539	684	537	284	68	-
2002.....	55	73	83	136	218	419	648	778	616	283	87	-
2001.....	57	62	84	123	224	406	619	756	643	302	71	-
2000.....	46	61	97	178	245	478	721	888	712	280	84	-
1999.....	57	66	88	141	266	439	720	883	755	353	86	-

All races, male

2003.....	28	37	38	59	96	166	213	222	130	46	10	-
2002.....	24	39	45	64	110	176	230	267	163	45	13	-
2001.....	25	33	42	57	100	156	244	220	164	50	2	-
2000.....	18	39	47	79	128	194	240	275	165	46	13	-
1999.....	27	38	54	68	139	186	266	282	175	65	8	-

All races, female

2003.....	30	37	48	59	106	247	326	462	407	238	58	-
2002.....	31	34	38	72	108	243	418	511	453	238	74	-
2001.....	32	29	42	66	124	250	375	536	479	252	69	-
2000.....	28	22	50	99	117	284	481	613	547	234	71	-
1999.....	30	28	34	73	127	253	454	601	580	288	78	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malnutrition (E40-E46)

White, both sexes

2003.....	35	57	63	94	165	355	455	603	469	243	50	-
2002.....	42	51	61	105	180	356	561	675	544	247	72	-
2001.....	37	45	68	93	189	338	521	679	566	266	64	-
2000.....	31	45	74	146	189	405	634	781	650	235	67	-
1999.....	46	41	66	107	198	366	610	779	689	302	72	-

White, male

2003.....	18	25	26	49	73	136	176	197	112	40	7	-
2002.....	17	24	29	51	91	153	192	235	148	38	10	-
2001.....	13	22	30	40	81	130	205	195	139	44	1	-
2000.....	11	26	32	63	96	162	209	242	152	35	11	-
1999.....	21	20	41	48	98	152	221	247	154	49	6	-

White, female

2003.....	17	32	37	45	92	219	279	406	357	203	43	-
2002.....	25	27	32	54	89	203	369	440	396	209	62	-
2001.....	24	23	38	53	108	208	316	484	427	222	63	-
2000.....	20	19	42	83	93	243	425	539	498	200	56	-
1999.....	25	21	25	59	100	214	389	532	535	253	66	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malnutrition (E40-E46)

Black, both sexes

2003.....	21	17	21	19	33	52	76	73	62	33	15	-
2002.....	12	22	22	28	33	59	76	86	65	31	12	-
2001.....	18	15	12	26	30	58	82	67	66	28	7	-
2000.....	15	16	21	28	49	63	74	94	57	39	14	-
1999.....	11	21	21	29	59	62	100	90	59	44	14	-

Black, male

2003.....	10	12	11	9	21	29	32	22	15	4	3	-
2002.....	7	15	16	12	15	20	33	26	12	3	2	-
2001.....	11	10	9	16	18	24	30	24	20	4	1	-
2000.....	7	13	13	14	29	30	27	31	10	11	1	-
1999.....	6	15	12	17	36	28	38	29	19	13	2	-

Black, female

2003.....	11	5	10	10	12	23	44	51	47	29	12	-
2002.....	5	7	6	16	18	39	43	60	53	28	10	-
2001.....	7	5	3	10	12	34	52	43	46	24	6	-
2000.....	8	3	8	14	20	33	47	63	47	28	13	-
1999.....	5	6	9	12	23	34	62	61	40	31	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malnutrition (E40-E46)

American Indian 1/, both sexes

2003.....	2	-	-	1	3	5	2	2	2	2	1	-
2002.....	1	-	-	1	3	1	4	5	3	3	1	-
2001.....	2	1	2	3	3	3	6	5	2	1	-	-
2000.....	-	-	-	2	3	4	3	6	2	1	3	-
1999.....	-	3	1	3	3	3	4	4	3	1	-	-

American Indian 1/, male

2003.....	-	-	-	-	2	-	2	-	-	-	-	-
2002.....	-	-	-	-	3	-	1	2	-	2	-	-
2001.....	1	-	2	1	1	-	3	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	-	1	-	1	-
1999.....	-	2	1	1	1	2	3	1	-	-	-	-

American Indian 1/, female

2003.....	2	-	-	1	1	5	-	2	2	2	1	-
2002.....	1	-	-	1	-	1	3	3	3	1	1	-
2001.....	1	1	-	2	2	3	3	5	2	1	-	-
2000.....	-	-	-	1	2	3	2	6	1	1	2	-
1999.....	-	1	-	2	2	1	1	3	3	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malnutrition (E40-E46)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	2	4	1	1	6	6	4	6	2	-
2002.....	-	-	-	2	2	3	7	12	4	2	2	-
2001.....	-	1	2	1	2	7	10	5	9	7	-	-
2000.....	-	-	2	2	4	6	10	7	3	5	-	-
1999.....	-	1	-	2	6	8	6	10	4	6	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	1	1	-	1	3	3	3	2	-	-
2002.....	-	-	-	1	1	3	4	4	3	2	1	-
2001.....	-	1	1	-	-	2	6	1	5	2	-	-
2000.....	-	-	2	1	2	1	3	2	2	-	-	-
1999.....	-	1	-	2	4	4	4	5	2	3	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	1	3	1	-	3	3	1	4	2	-
2002.....	-	-	-	1	1	-	3	8	1	-	1	-
2001.....	-	-	1	1	2	5	4	4	4	5	-	-
2000.....	-	-	-	1	2	5	7	5	1	5	-	-
1999.....	-	-	-	-	2	4	2	5	2	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

All races, both sexes

2003.....	3	3	5	7	13	15	25	43	33	17	6	-
2002.....	8	4	8	14	14	33	32	56	51	29	6	-
2001.....	5	2	5	8	19	33	42	42	41	22	12	-
2000.....	7	9	6	14	23	27	59	59	69	22	6	-
1999.....	4	10	6	8	14	33	56	61	67	27	8	-

All races, male

2003.....	1	1	3	4	7	9	12	21	9	3	1	-
2002.....	4	4	5	6	6	17	11	16	6	4	-	-
2001.....	4	2	4	6	11	16	10	8	7	5	1	-
2000.....	3	6	5	8	6	13	24	14	17	6	-	-
1999.....	3	4	2	4	10	19	23	19	21	6	1	-

All races, female

2003.....	2	2	2	3	6	6	13	22	24	14	5	-
2002.....	4	-	3	8	8	16	21	40	45	25	6	-
2001.....	1	-	1	2	8	17	32	34	34	17	11	-
2000.....	4	3	1	6	17	14	35	45	52	16	6	-
1999.....	1	6	4	4	4	14	33	42	46	21	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

White, both sexes

2003.....	3	3	4	7	11	14	24	43	29	15	5	-
2002.....	7	3	6	9	9	31	28	51	47	27	5	-
2001.....	2	-	4	5	16	27	36	37	38	20	12	-
2000.....	5	7	4	13	22	20	53	55	62	20	4	-
1999.....	2	5	6	8	12	26	48	54	62	24	8	-

White, male

2003.....	1	1	2	4	6	8	11	21	8	2	-	-
2002.....	3	3	3	4	4	16	8	15	6	4	-	-
2001.....	1	-	3	3	9	15	9	8	6	4	1	-
2000.....	2	4	3	7	6	9	24	14	15	6	-	-
1999.....	2	1	2	4	8	13	21	18	21	6	1	-

White, female

2003.....	2	2	2	3	5	6	13	22	21	13	5	-
2002.....	4	-	3	5	5	15	20	36	41	23	5	-
2001.....	1	-	1	2	7	12	27	29	32	16	11	-
2000.....	3	3	1	6	16	11	29	41	47	14	4	-
1999.....	-	4	4	4	4	13	27	36	41	18	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

Black, both sexes

2003.....	-	-	1	-	2	1	1	-	4	2	-	-
2002.....	1	1	1	5	5	2	3	4	3	2	1	-
2001.....	3	2	1	2	3	4	6	5	3	2	-	-
2000.....	1	2	2	1	-	5	6	2	4	2	2	-
1999.....	1	5	-	-	2	7	7	7	5	3	-	-

Black, male

2003.....	-	-	1	-	1	1	1	-	1	1	-	-
2002.....	1	1	1	2	2	1	3	1	-	-	-	-
2001.....	3	2	1	2	2	1	1	-	1	1	-	-
2000.....	1	2	2	1	-	2	-	-	2	-	-	-
1999.....	1	3	-	-	2	6	2	1	-	-	-	-

Black, female

2003.....	-	-	-	-	1	-	-	-	3	1	-	-
2002.....	-	-	-	3	3	1	-	3	3	2	1	-
2001.....	-	-	-	-	1	3	5	5	2	1	-	-
2000.....	-	-	-	-	-	3	6	2	2	2	2	-
1999.....	-	2	-	-	-	1	5	6	5	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	-	-	1	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	1	-	2	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	2	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningitis (G00,G03)

All races, both sexes

2003.....	58	55	53	57	39	55	35	33	18	1	-	1
2002.....	53	48	40	46	63	47	37	36	16	3	-	1
2001.....	50	58	55	46	48	46	50	25	12	4	-	-
2000.....	55	43	47	48	60	68	55	32	16	2	-	-
1999.....	46	41	49	49	66	65	57	40	20	2	-	-

All races, male

2003.....	38	30	31	27	18	26	14	13	5	1	-	1
2002.....	30	29	19	26	36	21	20	11	8	-	-	1
2001.....	30	30	31	24	27	26	21	11	3	1	-	-
2000.....	31	26	28	26	34	41	21	13	5	1	-	-
1999.....	29	24	21	28	32	26	23	13	7	-	-	-

All races, female

2003.....	20	25	22	30	21	29	21	20	13	-	-	-
2002.....	23	19	21	20	27	26	17	25	8	3	-	-
2001.....	20	28	24	22	21	20	29	14	9	3	-	-
2000.....	24	17	19	22	26	27	34	19	11	1	-	-
1999.....	17	17	28	21	34	39	34	27	13	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningitis (G00,G03)

White, both sexes

2003.....	37	36	36	46	30	48	30	28	13	1	-	1
2002.....	32	33	25	38	50	37	31	35	12	3	-	1
2001.....	35	40	44	35	43	40	42	20	9	2	-	-
2000.....	36	29	36	32	47	56	48	24	14	2	-	-
1999.....	33	32	39	38	51	47	48	36	12	2	-	-

White, male

2003.....	23	22	21	23	14	25	12	11	4	1	-	1
2002.....	18	20	12	22	29	17	19	10	5	-	-	1
2001.....	21	21	24	17	24	21	18	7	2	-	-	-
2000.....	22	20	22	19	29	33	19	9	4	1	-	-
1999.....	19	20	17	21	25	17	21	12	4	-	-	-

White, female

2003.....	14	14	15	23	16	23	18	17	9	-	-	-
2002.....	14	13	13	16	21	20	12	25	7	3	-	-
2001.....	14	19	20	18	19	19	24	13	7	2	-	-
2000.....	14	9	14	13	18	23	29	15	10	1	-	-
1999.....	14	12	22	17	26	30	27	24	8	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningitis (G00,G03)

Black, both sexes

2003.....	17	18	17	10	8	5	4	4	5	-	-	-
2002.....	19	12	11	8	13	9	4	1	3	-	-	-
2001.....	13	16	9	9	5	5	6	4	2	2	-	-
2000.....	17	13	9	15	12	8	5	7	2	-	-	-
1999.....	13	8	10	9	12	17	8	4	8	-	-	-

Black, male

2003.....	11	8	10	4	4	-	2	1	1	-	-	-
2002.....	11	7	5	4	7	4	1	1	2	-	-	-
2001.....	8	8	6	6	3	4	2	4	1	1	-	-
2000.....	8	6	5	6	5	5	2	4	1	-	-	-
1999.....	10	4	4	6	6	8	2	1	3	-	-	-

Black, female

2003.....	6	10	7	6	4	5	2	3	4	-	-	-
2002.....	8	5	6	4	6	5	3	-	1	-	-	-
2001.....	5	8	3	3	2	1	4	-	1	1	-	-
2000.....	9	7	4	9	7	3	3	3	1	-	-	-
1999.....	3	4	6	3	6	9	6	3	5	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningitis (G00,G03)

Asian or Pacific Islander 2/, both sexes

2003.....	2	-	-	-	1	2	1	1	-	-	-	-
2002.....	1	2	3	-	-	1	1	-	1	-	-	-
2001.....	-	1	2	-	-	1	1	1	1	-	-	-
2000.....	2	1	2	1	1	2	1	1	-	-	-	-
1999.....	-	1	-	1	3	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	2	-	-	-	-	1	-	1	-	-	-	-
2002.....	-	1	1	-	-	-	-	-	1	-	-	-
2001.....	-	-	1	-	-	1	1	-	-	-	-	-
2000.....	1	-	1	1	-	2	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	1	1	1	-	-	-	-	-
2002.....	1	1	2	-	-	1	1	-	-	-	-	-
2001.....	-	1	1	-	-	-	-	1	1	-	-	-
2000.....	1	1	1	-	1	-	1	1	-	-	-	-
1999.....	-	1	-	1	2	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

All races, both sexes

2003.....	46	112	240	624	1,706	3,671	5,052	4,157	1,900	425	31	-
2002.....	39	95	223	626	1,600	3,368	4,771	3,959	1,781	440	32	-
2001.....	32	100	201	554	1,593	3,446	4,679	3,786	1,655	425	43	-
2000.....	33	82	188	626	1,482	3,322	4,330	3,567	1,627	369	27	1
1999.....	35	73	169	539	1,481	3,066	4,044	3,332	1,492	313	31	-

All races, male

2003.....	23	65	154	402	1,115	2,300	2,891	2,210	846	143	14	-
2002.....	26	57	136	433	1,054	2,065	2,746	2,109	793	147	10	-
2001.....	25	69	145	362	1,032	2,104	2,733	2,012	759	134	10	-
2000.....	19	55	115	398	940	2,052	2,478	1,879	726	115	8	1
1999.....	25	44	117	338	965	1,940	2,367	1,699	639	110	6	-

All races, female

2003.....	23	47	86	222	591	1,371	2,161	1,947	1,054	282	17	-
2002.....	13	38	87	193	546	1,303	2,025	1,850	988	293	22	-
2001.....	7	31	56	192	561	1,342	1,946	1,774	896	291	33	-
2000.....	14	27	73	228	542	1,270	1,852	1,688	901	254	19	-
1999.....	10	29	52	201	516	1,126	1,677	1,633	853	203	25	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

White, both sexes

2003.....	37	97	210	566	1,592	3,472	4,823	3,963	1,823	405	26	-
2002.....	38	82	208	567	1,494	3,193	4,587	3,795	1,708	415	30	-
2001.....	30	90	180	513	1,498	3,288	4,484	3,628	1,585	402	37	-
2000.....	29	75	176	585	1,397	3,163	4,143	3,418	1,560	350	26	1
1999.....	31	65	150	498	1,397	2,910	3,877	3,192	1,426	294	29	-

White, male

2003.....	19	56	133	368	1,043	2,179	2,767	2,097	806	133	12	-
2002.....	25	50	129	395	989	1,963	2,641	2,021	758	137	10	-
2001.....	23	62	135	339	972	2,019	2,622	1,934	725	127	9	-
2000.....	16	49	109	372	890	1,943	2,364	1,804	689	109	7	1
1999.....	24	41	106	314	922	1,842	2,281	1,625	606	102	6	-

White, female

2003.....	18	41	77	198	549	1,293	2,056	1,866	1,017	272	14	-
2002.....	13	32	79	172	505	1,230	1,946	1,774	950	278	20	-
2001.....	7	28	45	174	526	1,269	1,862	1,694	860	275	28	-
2000.....	13	26	67	213	507	1,220	1,779	1,614	871	241	19	-
1999.....	7	24	44	184	475	1,068	1,596	1,567	820	192	23	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

Black, both sexes

2003.....	7	11	22	39	81	134	151	113	53	11	4	-
2002.....	1	8	11	43	71	120	125	105	49	19	1	-
2001.....	2	5	15	25	66	109	129	116	54	18	4	-
2000.....	4	5	9	28	66	118	133	102	47	15	1	-
1999.....	4	7	15	33	64	109	112	94	52	15	2	-

Black, male

2003.....	3	7	14	22	51	81	80	62	25	5	1	-
2002.....	1	3	7	30	48	66	72	58	19	6	-	-
2001.....	2	4	9	15	44	55	72	55	29	4	-	-
2000.....	3	4	5	19	41	80	78	44	24	6	1	-
1999.....	1	3	10	22	37	67	56	47	22	5	-	-

Black, female

2003.....	4	4	8	17	30	53	71	51	28	6	3	-
2002.....	-	5	4	13	23	54	53	47	30	13	1	-
2001.....	-	1	6	10	22	54	57	61	25	14	4	-
2000.....	1	1	4	9	25	38	55	58	23	9	-	-
1999.....	3	4	5	11	27	42	56	47	30	10	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

American Indian 1/, both sexes

2003.....	-	-	4	3	10	9	17	11	5	1	-	-
2002.....	-	2	-	1	13	17	5	9	5	-	-	-
2001.....	-	-	3	5	5	10	4	7	1	1	-	-
2000.....	-	1	1	3	6	9	11	6	5	2	-	-
1999.....	-	-	-	1	3	5	11	3	2	1	-	-

American Indian 1/, male

2003.....	-	-	3	2	5	5	7	9	1	-	-	-
2002.....	-	2	-	1	9	12	4	6	3	-	-	-
2001.....	-	-	-	2	4	7	3	2	-	-	-	-
2000.....	-	1	-	1	3	5	7	4	4	-	-	-
1999.....	-	-	-	-	-	4	10	1	1	-	-	-

American Indian 1/, female

2003.....	-	-	1	1	5	4	10	2	4	1	-	-
2002.....	-	-	-	-	4	5	1	3	2	-	-	-
2001.....	-	-	3	3	1	3	1	5	1	1	-	-
2000.....	-	-	1	2	3	4	4	2	1	2	-	-
1999.....	-	-	-	1	3	1	1	2	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

Asian or Pacific Islander 2/, both sexes

2003.....	2	4	4	16	23	56	61	70	19	8	1	-
2002.....	-	3	4	15	22	38	54	50	19	6	1	-
2001.....	-	5	3	11	24	39	62	35	15	4	2	-
2000.....	-	1	2	10	13	32	43	41	15	2	-	-
1999.....	-	1	4	7	17	42	44	43	12	3	-	-

Asian or Pacific Islander 2/, male

2003.....	1	2	4	10	16	35	37	42	14	5	1	-
2002.....	-	2	-	7	8	24	29	24	13	4	-	-
2001.....	-	3	1	6	12	23	36	21	5	3	1	-
2000.....	-	1	1	6	6	24	29	27	9	-	-	-
1999.....	-	-	1	2	6	27	20	26	10	3	-	-

Asian or Pacific Islander 2/, female

2003.....	1	2	-	6	7	21	24	28	5	3	-	-
2002.....	-	1	4	8	14	14	25	26	6	2	1	-
2001.....	-	2	2	5	12	16	26	14	10	1	1	-
2000.....	-	-	1	4	7	8	14	14	6	2	-	-
1999.....	-	1	3	5	11	15	24	17	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alzheimer's disease (G30)

All races, both sexes

2003.....	57	152	402	979	2,857	7,348	13,809	17,368	13,774	5,704	975	-
2002.....	40	135	375	939	2,663	7,078	13,057	16,267	12,267	5,140	878	-
2001.....	56	165	364	858	2,564	6,568	11,974	14,843	11,168	4,474	796	-
2000.....	47	142	349	868	2,563	6,319	10,934	13,337	10,257	4,014	701	2
1999.....	40	125	323	830	2,374	5,769	10,067	11,862	9,041	3,473	604	-

All races, male

2003.....	25	76	167	456	1,228	2,855	4,751	4,916	2,933	824	91	-
2002.....	22	55	173	408	1,139	2,793	4,434	4,569	2,570	732	79	-
2001.....	27	61	167	397	1,075	2,616	4,179	4,225	2,312	608	79	-
2000.....	24	60	150	420	1,134	2,433	3,795	3,687	2,101	564	54	1
1999.....	18	68	154	389	1,048	2,332	3,572	3,382	1,861	505	48	-

All races, female

2003.....	32	76	235	523	1,629	4,493	9,058	12,452	10,841	4,880	884	-
2002.....	18	80	202	531	1,524	4,285	8,623	11,698	9,697	4,408	799	-
2001.....	29	104	197	461	1,489	3,952	7,795	10,618	8,856	3,866	717	-
2000.....	23	82	199	448	1,429	3,886	7,139	9,650	8,156	3,450	647	1
1999.....	22	57	169	441	1,326	3,437	6,495	8,480	7,180	2,968	556	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alzheimer's disease (G30)

White, both sexes

2003.....	55	142	376	861	2,591	6,745	12,880	16,313	12,967	5,351	873	-
2002.....	37	127	341	845	2,437	6,554	12,237	15,299	11,560	4,817	783	-
2001.....	54	158	341	767	2,369	6,032	11,194	13,961	10,565	4,182	707	-
2000.....	41	132	318	796	2,343	5,864	10,260	12,562	9,723	3,765	633	1
1999.....	35	113	300	757	2,169	5,363	9,470	11,216	8,611	3,267	552	-

White, male

2003.....	24	72	152	405	1,117	2,637	4,431	4,643	2,755	757	81	-
2002.....	19	51	159	363	1,046	2,600	4,146	4,321	2,418	676	64	-
2001.....	25	56	154	353	996	2,423	3,904	3,968	2,178	576	67	-
2000.....	21	55	138	389	1,043	2,271	3,583	3,466	1,974	525	45	1
1999.....	16	61	144	355	961	2,180	3,384	3,198	1,763	469	41	-

White, female

2003.....	31	70	224	456	1,474	4,108	8,449	11,670	10,212	4,594	792	-
2002.....	18	76	182	482	1,391	3,954	8,091	10,978	9,142	4,141	719	-
2001.....	29	102	187	414	1,373	3,609	7,290	9,993	8,387	3,606	640	-
2000.....	20	77	180	407	1,300	3,593	6,677	9,096	7,749	3,240	588	-
1999.....	19	52	156	402	1,208	3,183	6,086	8,018	6,848	2,798	511	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alzheimer's disease (G30)

Black, both sexes

2003.....	2	8	21	100	228	537	819	903	701	307	90	-
2002.....	3	3	29	80	203	473	714	843	615	292	81	-
2001.....	2	6	21	81	173	500	678	784	525	261	80	-
2000.....	4	7	23	67	194	400	608	682	464	219	58	1
1999.....	3	10	19	62	187	364	526	569	367	184	48	-

Black, male

2003.....	1	3	14	45	99	204	279	234	149	50	8	-
2002.....	3	2	12	38	88	179	248	202	119	52	12	-
2001.....	2	5	13	39	72	182	231	219	110	25	11	-
2000.....	2	5	11	30	83	137	185	181	99	36	6	-
1999.....	2	6	9	31	81	132	160	158	75	31	7	-

Black, female

2003.....	1	5	7	55	129	333	540	669	552	257	82	-
2002.....	-	1	17	42	115	294	466	641	496	240	69	-
2001.....	-	1	8	42	101	318	447	565	415	236	69	-
2000.....	2	2	12	37	111	263	423	501	365	183	52	1
1999.....	1	4	10	31	106	232	366	411	292	153	41	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alzheimer's disease (G30)

American Indian 1/, both sexes

2003.....	-	1	-	6	9	15	26	34	22	6	3	-
2002.....	-	-	1	7	7	13	29	28	22	5	3	-
2001.....	-	-	1	3	7	8	32	19	14	8	-	-
2000.....	-	1	2	2	12	21	18	19	15	5	-	-
1999.....	-	-	3	5	8	14	15	20	14	5	2	-

American Indian 1/, male

2003.....	-	-	-	1	3	1	8	7	2	-	-	-
2002.....	-	-	1	4	1	5	10	8	5	-	1	-
2001.....	-	-	-	2	2	3	14	6	1	-	-	-
2000.....	-	-	-	1	3	6	7	4	4	1	-	-
1999.....	-	-	1	1	4	6	5	1	5	-	-	-

American Indian 1/, female

2003.....	-	1	-	5	6	14	18	27	20	6	3	-
2002.....	-	-	-	3	6	8	19	20	17	5	2	-
2001.....	-	-	1	1	5	5	18	13	13	8	-	-
2000.....	-	1	2	1	9	15	11	15	11	4	-	-
1999.....	-	-	2	4	4	8	10	19	9	5	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alzheimer's disease (G30)

Asian or Pacific Islander 2/, both sexes

2003.....	-	1	5	12	29	51	84	118	84	40	9	-
2002.....	-	5	4	7	16	38	77	97	70	26	11	-
2001.....	-	1	1	7	15	28	70	79	64	23	9	-
2000.....	2	2	6	3	14	34	48	74	55	25	10	-
1999.....	2	2	1	6	10	28	56	57	49	17	2	-

Asian or Pacific Islander 2/, male

2003.....	-	1	1	5	9	13	33	32	27	17	2	-
2002.....	-	2	1	3	4	9	30	38	28	4	2	-
2001.....	-	-	-	3	5	8	30	32	23	7	1	-
2000.....	1	-	1	-	5	19	20	36	24	2	3	-
1999.....	-	1	-	2	2	14	23	25	18	5	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	4	7	20	38	51	86	57	23	7	-
2002.....	-	3	3	4	12	29	47	59	42	22	9	-
2001.....	-	1	1	4	10	20	40	47	41	16	8	-
2000.....	1	2	5	3	9	15	28	38	31	23	7	-
1999.....	2	1	1	4	8	14	33	32	31	12	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

All races, both sexes

2003.....	27,188	35,465	44,032	55,941	81,279	121,274	158,084	160,879	117,184	47,760	10,626	38
2002.....	27,084	35,004	43,426	57,731	86,340	126,160	161,255	163,421	117,826	47,620	10,542	51
2001.....	26,682	33,335	42,889	59,066	89,343	130,631	161,364	164,000	116,097	47,232	10,601	72
2000.....	25,798	33,454	44,028	61,769	94,418	135,296	160,906	165,313	117,092	47,427	10,618	62
1999.....	25,158	33,344	44,489	64,775	98,653	138,680	163,872	166,070	116,257	47,656	10,527	49

All races, male

2003.....	19,045	24,105	28,699	34,660	46,852	63,584	72,284	60,672	33,792	9,930	1,509	29
2002.....	18,930	23,721	28,316	35,603	49,716	65,751	73,019	61,344	33,439	9,860	1,521	36
2001.....	18,442	22,489	27,831	36,221	51,535	67,972	72,817	59,914	32,127	9,565	1,491	53
2000.....	17,991	22,817	28,606	37,721	54,789	70,196	72,601	60,379	32,104	9,636	1,539	43
1999.....	17,595	22,653	28,872	39,978	56,926	72,052	73,515	60,030	31,573	9,558	1,502	38

All races, female

2003.....	8,143	11,360	15,333	21,281	34,427	57,690	85,800	100,207	83,392	37,830	9,117	9
2002.....	8,154	11,283	15,110	22,128	36,624	60,409	88,236	102,077	84,387	37,760	9,021	15
2001.....	8,240	10,846	15,058	22,845	37,808	62,659	88,547	104,086	83,970	37,667	9,110	19
2000.....	7,807	10,637	15,422	24,048	39,629	65,100	88,305	104,934	84,988	37,791	9,079	19
1999.....	7,563	10,691	15,617	24,797	41,727	66,628	90,357	106,040	84,684	38,098	9,025	11

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

White, both sexes

2003.....	19,822	27,384	34,591	45,019	68,111	105,565	141,763	147,197	107,725	43,755	9,230	31
2002.....	20,030	27,144	34,290	46,798	73,096	109,955	144,939	149,652	108,250	43,572	9,204	35
2001.....	19,791	25,782	33,849	47,754	75,867	114,196	145,321	150,129	107,073	43,220	9,266	45
2000.....	19,428	26,035	34,836	50,692	80,400	118,684	145,286	151,317	108,139	43,327	9,357	45
1999.....	19,030	25,836	35,412	53,485	84,103	121,928	148,314	151,947	107,568	43,687	9,350	32

White, male

2003.....	14,424	19,129	23,149	28,570	40,000	56,232	65,416	56,024	31,234	9,030	1,285	22
2002.....	14,521	18,890	22,892	29,526	42,976	58,282	66,354	56,699	30,711	8,978	1,274	26
2001.....	14,225	17,886	22,557	30,079	44,549	60,332	66,225	55,249	29,612	8,672	1,269	34
2000.....	14,011	18,204	23,256	31,747	47,612	62,471	66,225	55,631	29,586	8,639	1,320	34
1999.....	13,739	18,051	23,510	33,786	49,508	64,277	67,297	55,083	29,040	8,663	1,304	25

White, female

2003.....	5,398	8,255	11,442	16,449	28,111	49,333	76,347	91,173	76,491	34,725	7,945	9
2002.....	5,509	8,254	11,398	17,272	30,120	51,673	78,585	92,953	77,539	34,594	7,930	9
2001.....	5,566	7,896	11,292	17,675	31,318	53,864	79,096	94,880	77,461	34,548	7,997	11
2000.....	5,417	7,831	11,580	18,945	32,788	56,213	79,061	95,686	78,553	34,688	8,037	11
1999.....	5,291	7,785	11,902	19,699	34,595	57,651	81,017	96,864	78,528	35,024	8,046	7

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2003.....	6,610	7,135	8,233	9,377	11,254	13,221	13,415	11,138	7,770	3,402	1,211	5
2002.....	6,346	6,979	8,029	9,428	11,265	13,644	13,588	11,343	7,920	3,425	1,159	15
2001.....	6,171	6,683	7,970	9,795	11,509	13,950	13,377	11,562	7,643	3,467	1,156	25
2000.....	5,675	6,597	8,068	9,612	12,167	14,047	13,174	11,860	7,551	3,492	1,133	17
1999.....	5,503	6,683	7,971	9,892	12,600	14,302	13,109	11,910	7,348	3,490	1,033	16

Black, male

2003.....	4,091	4,349	4,794	5,173	5,785	6,112	5,466	3,549	1,887	674	170	5
2002.....	3,915	4,254	4,720	5,185	5,718	6,104	5,356	3,572	2,044	657	190	9
2001.....	3,748	4,013	4,594	5,284	5,906	6,336	5,338	3,665	1,934	702	189	18
2000.....	3,503	4,074	4,613	5,104	6,104	6,348	5,217	3,778	1,942	758	188	9
1999.....	3,438	4,049	4,681	5,392	6,328	6,415	5,055	3,894	1,953	744	169	12

Black, female

2003.....	2,519	2,786	3,439	4,204	5,469	7,109	7,949	7,589	5,883	2,728	1,041	-
2002.....	2,431	2,725	3,309	4,243	5,547	7,540	8,232	7,771	5,876	2,768	969	6
2001.....	2,423	2,670	3,376	4,511	5,603	7,614	8,039	7,897	5,709	2,765	967	7
2000.....	2,172	2,523	3,455	4,508	6,063	7,699	7,957	8,082	5,609	2,734	945	8
1999.....	2,065	2,634	3,290	4,500	6,272	7,887	8,054	8,016	5,395	2,746	864	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

American Indian 1/, both sexes

2003.....	217	248	344	366	393	420	419	336	215	67	34	2
2002.....	191	235	282	339	384	437	408	309	204	89	25	1
2001.....	179	230	261	349	391	392	391	318	173	75	26	-
2000.....	179	224	274	367	351	436	388	322	211	90	18	-
1999.....	159	214	257	329	401	448	376	324	188	66	29	-

American Indian 1/, male

2003.....	145	167	202	203	230	211	188	137	83	18	13	2
2002.....	137	145	186	193	215	227	169	142	58	21	8	1
2001.....	117	154	171	192	228	191	164	127	57	18	3	-
2000.....	118	142	165	215	209	220	176	123	60	23	2	-
1999.....	108	136	144	204	228	241	154	110	55	17	7	-

American Indian 1/, female

2003.....	72	81	142	163	163	209	231	199	132	49	21	-
2002.....	54	90	96	146	169	210	239	167	146	68	17	-
2001.....	62	76	90	157	163	201	227	191	116	57	23	-
2000.....	61	82	109	152	142	216	212	199	151	67	16	-
1999.....	51	78	113	125	173	207	222	214	133	49	22	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

Asian or Pacific Islander 2/, both sexes

2003.....	539	698	864	1,179	1,521	2,068	2,487	2,208	1,474	536	151	-
2002.....	517	646	825	1,166	1,595	2,124	2,320	2,117	1,452	534	154	-
2001.....	541	640	809	1,168	1,576	2,093	2,275	1,991	1,208	470	153	2
2000.....	516	598	850	1,098	1,500	2,129	2,058	1,814	1,191	518	110	-
1999.....	466	611	849	1,069	1,549	2,002	2,073	1,889	1,153	413	115	1

Asian or Pacific Islander 2/, male

2003.....	385	460	554	714	837	1,029	1,214	962	588	208	41	-
2002.....	357	432	518	699	807	1,138	1,140	931	626	204	49	-
2001.....	352	436	509	666	852	1,113	1,090	873	524	173	30	1
2000.....	359	397	572	655	864	1,157	983	847	516	216	29	-
1999.....	310	417	537	596	862	1,119	1,009	943	525	134	22	1

Asian or Pacific Islander 2/, female

2003.....	154	238	310	465	684	1,039	1,273	1,246	886	328	110	-
2002.....	160	214	307	467	788	986	1,180	1,186	826	330	105	-
2001.....	189	204	300	502	724	980	1,185	1,118	684	297	123	1
2000.....	157	201	278	443	636	972	1,075	967	675	302	81	-
1999.....	156	194	312	473	687	883	1,064	946	628	279	93	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races, both sexes

2003.....	22,212	29,217	35,843	44,470	62,793	90,796	116,535	118,106	86,773	35,782	8,135	35
2002.....	22,391	28,895	35,339	45,948	66,599	94,844	118,737	119,565	86,848	35,664	8,096	47
2001.....	22,009	27,564	34,922	47,131	69,168	98,099	118,893	119,848	85,878	35,479	8,234	66
2000.....	21,167	27,676	35,723	49,178	73,227	101,764	118,296	121,003	86,424	35,605	8,210	59
1999.....	20,898	27,605	36,562	52,206	77,047	104,841	121,311	122,126	85,752	35,797	8,185	47

All races, male

2003.....	16,140	20,538	23,984	28,270	37,298	49,241	55,405	46,400	25,955	7,662	1,210	27
2002.....	16,256	20,199	23,701	29,045	39,613	51,211	56,006	46,515	25,549	7,663	1,208	34
2001.....	15,789	19,200	23,319	29,643	40,994	52,960	55,616	45,298	24,606	7,394	1,175	47
2000.....	15,331	19,456	23,829	30,890	43,699	54,421	55,272	45,756	24,475	7,430	1,235	41
1999.....	15,143	19,370	24,292	33,118	45,718	56,178	56,292	45,667	23,999	7,442	1,192	37

All races, female

2003.....	6,072	8,679	11,859	16,200	25,495	41,555	61,130	71,706	60,818	28,120	6,925	8
2002.....	6,135	8,696	11,638	16,903	26,986	43,633	62,731	73,050	61,299	28,001	6,888	13
2001.....	6,220	8,364	11,603	17,488	28,174	45,139	63,277	74,550	61,272	28,085	7,059	19
2000.....	5,836	8,220	11,894	18,288	29,528	47,343	63,024	75,247	61,949	28,175	6,975	18
1999.....	5,755	8,235	12,270	19,088	31,329	48,663	65,019	76,459	61,753	28,355	6,993	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

White, both sexes

2003.....	16,629	23,022	28,626	36,181	53,052	79,469	104,888	108,344	80,010	32,867	7,098	28
2002.....	17,008	22,793	28,296	37,663	56,829	83,021	107,101	109,795	80,022	32,672	7,076	32
2001.....	16,826	21,672	28,047	38,550	59,214	86,105	107,420	110,058	79,431	32,522	7,230	42
2000.....	16,346	21,927	28,654	40,751	62,698	89,644	107,189	111,070	80,035	32,666	7,279	44
1999.....	16,193	21,743	29,531	43,471	66,107	92,559	110,133	112,009	79,531	32,885	7,289	32

White, male

2003.....	12,523	16,624	19,670	23,530	32,089	43,802	50,349	42,981	24,092	7,002	1,026	20
2002.....	12,786	16,361	19,428	24,359	34,514	45,630	51,104	43,126	23,549	6,992	1,024	24
2001.....	12,520	15,516	19,208	24,899	35,716	47,230	50,776	41,956	22,737	6,722	1,000	31
2000.....	12,213	15,794	19,628	26,246	38,218	48,669	50,626	42,311	22,657	6,708	1,059	33
1999.....	12,092	15,652	20,043	28,215	40,032	50,391	51,748	42,048	22,135	6,760	1,037	25

White, female

2003.....	4,106	6,398	8,956	12,651	20,963	35,667	54,539	65,363	55,918	25,865	6,072	8
2002.....	4,222	6,432	8,868	13,304	22,315	37,391	55,997	66,669	56,473	25,680	6,052	8
2001.....	4,306	6,156	8,839	13,651	23,498	38,875	56,644	68,102	56,694	25,800	6,230	11
2000.....	4,133	6,133	9,026	14,505	24,480	40,975	56,563	68,759	57,378	25,958	6,220	11
1999.....	4,101	6,091	9,488	15,256	26,075	42,168	58,385	69,961	57,396	26,125	6,252	7

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Black, both sexes												
2003.....	5,024	5,504	6,317	7,176	8,429	9,628	9,633	7,982	5,563	2,495	897	5
2002.....	4,872	5,450	6,246	7,196	8,405	9,984	9,773	8,100	5,688	2,554	883	14
2001.....	4,664	5,245	6,103	7,500	8,620	10,252	9,672	8,243	5,499	2,561	878	22
2000.....	4,324	5,133	6,252	7,367	9,223	10,333	9,469	8,478	5,414	2,503	834	15
1999.....	4,250	5,256	6,219	7,711	9,529	10,523	9,515	8,571	5,280	2,569	790	15
Black, male												
2003.....	3,194	3,431	3,724	4,050	4,445	4,564	4,039	2,620	1,390	503	139	5
2002.....	3,078	3,383	3,741	4,017	4,340	4,562	3,977	2,620	1,515	509	137	9
2001.....	2,907	3,216	3,594	4,092	4,512	4,767	3,964	2,658	1,469	533	152	15
2000.....	2,746	3,237	3,645	3,973	4,699	4,752	3,827	2,776	1,408	548	153	8
1999.....	2,724	3,284	3,720	4,282	4,873	4,795	3,738	2,864	1,446	561	129	12
Black, female												
2003.....	1,830	2,073	2,593	3,126	3,984	5,064	5,594	5,362	4,173	1,992	758	-
2002.....	1,794	2,067	2,505	3,179	4,065	5,422	5,796	5,480	4,173	2,045	746	5
2001.....	1,757	2,029	2,509	3,408	4,108	5,485	5,708	5,585	4,030	2,028	726	7
2000.....	1,578	1,896	2,607	3,394	4,524	5,581	5,642	5,702	4,006	1,955	681	7
1999.....	1,526	1,972	2,499	3,429	4,656	5,728	5,777	5,707	3,834	2,008	661	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

American Indian 1/, both sexes

2003.....	173	212	280	299	293	317	306	257	151	54	28	2
2002.....	151	188	224	256	286	341	292	231	145	63	20	1
2001.....	150	197	219	260	294	289	279	224	121	53	17	-
2000.....	146	180	214	291	257	326	271	231	150	65	13	-
1999.....	135	176	209	268	316	338	278	225	135	49	20	-

American Indian 1/, male

2003.....	122	146	169	169	180	161	147	109	63	14	11	2
2002.....	115	120	157	158	166	183	131	111	42	16	8	1
2001.....	99	138	149	152	180	158	127	92	38	14	2	-
2000.....	102	118	134	175	161	180	126	91	46	18	-	-
1999.....	93	120	121	172	181	184	119	80	44	11	6	-

American Indian 1/, female

2003.....	51	66	111	130	113	156	159	148	88	40	17	-
2002.....	36	68	67	98	120	158	161	120	103	47	12	-
2001.....	51	59	70	108	114	131	152	132	83	39	15	-
2000.....	44	62	80	116	96	146	145	140	104	47	13	-
1999.....	42	56	88	96	135	154	159	145	91	38	14	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Asian or Pacific Islander 2/, both sexes

2003.....	386	479	620	814	1,019	1,382	1,708	1,523	1,049	366	112	-
2002.....	360	464	573	833	1,079	1,498	1,571	1,439	993	375	117	-
2001.....	369	450	553	821	1,040	1,453	1,522	1,323	827	343	109	2
2000.....	351	436	603	769	1,049	1,461	1,367	1,224	825	371	84	-
1999.....	320	430	603	756	1,095	1,421	1,385	1,321	806	294	86	-

Asian or Pacific Islander 2/, male

2003.....	301	337	421	521	584	714	870	690	410	143	34	-
2002.....	277	335	375	511	593	836	794	658	443	146	39	-
2001.....	263	330	368	500	586	805	749	592	362	125	21	1
2000.....	270	307	422	496	621	820	693	578	364	156	23	-
1999.....	234	314	408	449	632	808	687	675	374	110	20	-

Asian or Pacific Islander 2/, female

2003.....	85	142	199	293	435	668	838	833	639	223	78	-
2002.....	83	129	198	322	486	662	777	781	550	229	78	-
2001.....	106	120	185	321	454	648	773	731	465	218	88	1
2000.....	81	129	181	273	428	641	674	646	461	215	61	-
1999.....	86	116	195	307	463	613	698	646	432	184	66	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races, both sexes

2003.....	117	151	216	295	398	603	630	557	291	93	23	-
2002.....	140	190	243	272	431	620	607	462	263	82	12	-
2001.....	125	168	193	307	437	599	589	468	235	67	10	-
2000.....	133	183	237	327	492	601	575	416	231	54	11	-
1999.....	144	189	222	338	516	614	613	442	235	63	8	-

All races, male

2003.....	39	62	84	105	136	199	185	160	62	10	2	-
2002.....	55	73	83	84	161	177	164	100	55	8	1	-
2001.....	41	73	62	106	125	182	148	106	50	10	-	-
2000.....	55	72	94	98	139	177	144	96	40	13	1	-
1999.....	50	64	69	107	183	164	161	89	26	8	-	-

All races, female

2003.....	78	89	132	190	262	404	445	397	229	83	21	-
2002.....	85	117	160	188	270	443	443	362	208	74	11	-
2001.....	84	95	131	201	312	417	441	362	185	57	10	-
2000.....	78	111	143	229	353	424	431	320	191	41	10	-
1999.....	94	125	153	231	333	450	452	353	209	55	8	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, both sexes

2003.....	78	109	168	259	352	568	604	532	284	89	22	-
2002.....	110	149	199	230	400	570	582	438	256	79	11	-
2001.....	87	127	166	261	388	558	567	447	228	62	10	-
2000.....	107	148	202	293	441	575	554	400	216	50	11	-
1999.....	108	156	185	302	474	578	583	421	227	57	7	-

White, male

2003.....	27	43	64	91	125	193	182	154	60	10	2	-
2002.....	42	56	70	69	150	163	160	97	54	8	-	-
2001.....	32	61	54	88	114	172	141	101	48	8	-	-
2000.....	48	62	85	88	122	169	140	93	36	11	1	-
1999.....	40	53	58	100	170	151	155	87	26	7	-	-

White, female

2003.....	51	66	104	168	227	375	422	378	224	79	20	-
2002.....	68	93	129	161	250	407	422	341	202	71	11	-
2001.....	55	66	112	173	274	386	426	346	180	54	10	-
2000.....	59	86	117	205	319	406	414	307	180	39	10	-
1999.....	68	103	127	202	304	427	428	334	201	50	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2003.....	31	34	38	30	36	23	17	18	4	3	1	-
2002.....	23	30	33	30	19	28	14	12	3	2	1	-
2001.....	33	31	22	31	38	28	12	13	4	5	-	-
2000.....	20	26	24	31	27	21	15	12	11	4	-	-
1999.....	29	29	23	25	31	24	16	16	5	4	1	-

Black, male

2003.....	11	17	16	13	8	6	2	4	2	-	-	-
2002.....	10	12	10	13	8	8	3	2	-	-	1	-
2001.....	9	10	7	11	6	5	3	4	-	2	-	-
2000.....	5	7	7	8	8	7	1	3	3	2	-	-
1999.....	10	9	8	6	10	10	4	1	-	-	-	-

Black, female

2003.....	20	17	22	17	28	17	15	14	2	3	1	-
2002.....	13	18	23	17	11	20	11	10	3	2	-	-
2001.....	24	21	15	20	32	23	9	9	4	3	-	-
2000.....	15	19	17	23	19	14	14	9	8	2	-	-
1999.....	19	20	15	19	21	14	12	15	5	4	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

American Indian 1/, both sexes

2003.....	1	-	3	-	4	1	-	1	-	-	-	-
2002.....	1	2	3	3	2	4	4	1	-	-	-	-
2001.....	1	2	2	4	3	2	2	1	-	-	-	-
2000.....	-	3	3	1	3	1	2	2	-	-	-	-
1999.....	1	1	4	2	4	1	2	-	1	-	-	-

American Indian 1/, male

2003.....	-	-	2	-	1	-	-	-	-	-	-	-
2002.....	1	-	2	-	-	3	1	1	-	-	-	-
2001.....	-	-	1	1	2	2	1	1	-	-	-	-
2000.....	-	1	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	1	-	1	-	3	1	-	1	-	-	-	-
2002.....	-	2	1	3	2	1	3	-	-	-	-	-
2001.....	1	2	1	3	1	-	1	-	-	-	-	-
2000.....	-	2	3	1	2	1	1	2	-	-	-	-
1999.....	1	1	4	2	4	1	2	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Asian or Pacific Islander 2/, both sexes

2003.....	7	8	7	6	6	11	9	6	3	1	-	-
2002.....	6	9	8	9	10	18	7	11	4	1	-	-
2001.....	4	8	3	11	8	11	8	7	3	-	-	-
2000.....	6	6	8	2	21	4	4	2	4	-	-	-
1999.....	6	3	10	9	7	11	12	5	2	2	-	-

Asian or Pacific Islander 2/, male

2003.....	1	2	2	1	2	-	1	2	-	-	-	-
2002.....	2	5	1	2	3	3	-	-	1	-	-	-
2001.....	-	2	-	6	3	3	3	-	2	-	-	-
2000.....	2	2	2	2	8	1	2	-	1	-	-	-
1999.....	-	2	3	1	3	3	2	1	-	1	-	-

Asian or Pacific Islander 2/, female

2003.....	6	6	5	5	4	11	8	4	3	1	-	-
2002.....	4	4	7	7	7	15	7	11	3	1	-	-
2001.....	4	6	3	5	5	8	5	7	1	-	-	-
2000.....	4	4	6	-	13	3	2	2	3	-	-	-
1999.....	6	1	7	8	4	8	10	4	2	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

All races, both sexes

2003.....	1,784	1,852	1,799	1,892	2,153	2,945	3,620	3,976	3,285	1,590	338	-
2002.....	1,616	1,661	1,659	1,807	2,088	2,816	3,505	3,716	3,115	1,442	314	4
2001.....	1,414	1,380	1,389	1,557	2,062	2,719	3,400	3,658	2,958	1,398	331	11
2000.....	1,152	1,249	1,423	1,570	2,035	2,694	3,268	3,684	2,886	1,330	286	3
1999.....	1,041	1,260	1,277	1,553	1,964	2,672	3,158	3,498	2,710	1,188	283	1

All races, male

2003.....	1,195	1,222	1,086	1,141	1,123	1,326	1,301	1,143	669	206	36	-
2002.....	1,094	1,107	1,052	1,046	1,042	1,259	1,250	1,004	629	194	44	4
2001.....	931	879	870	906	1,055	1,188	1,226	957	565	197	27	9
2000.....	811	789	895	909	1,062	1,135	1,134	960	545	181	21	2
1999.....	673	800	760	858	958	1,185	1,067	912	499	160	23	1

All races, female

2003.....	589	630	713	751	1,030	1,619	2,319	2,833	2,616	1,384	302	-
2002.....	522	554	607	761	1,046	1,557	2,255	2,712	2,486	1,248	270	-
2001.....	483	501	519	651	1,007	1,531	2,174	2,701	2,393	1,201	304	2
2000.....	341	460	528	661	973	1,559	2,134	2,724	2,341	1,149	265	1
1999.....	368	460	517	695	1,006	1,487	2,091	2,586	2,211	1,028	260	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

White, both sexes

2003.....	1,043	1,150	1,185	1,235	1,518	2,199	2,982	3,362	2,859	1,399	269	-
2002.....	951	1,036	1,040	1,185	1,474	2,109	2,785	3,134	2,701	1,265	261	1
2001.....	810	843	870	999	1,435	2,051	2,754	3,073	2,611	1,225	262	6
2000.....	647	721	867	1,032	1,379	2,012	2,614	3,122	2,487	1,132	236	1
1999.....	590	728	801	997	1,329	1,976	2,541	2,983	2,371	1,017	231	-

White, male

2003.....	762	804	750	761	799	1,003	1,075	960	559	175	30	-
2002.....	690	730	671	698	754	952	998	844	533	161	31	1
2001.....	574	571	573	613	760	905	983	786	493	162	14	5
2000.....	480	485	564	616	749	855	893	813	456	147	18	1
1999.....	409	488	499	556	667	883	852	757	430	129	19	-

White, female

2003.....	281	346	435	474	719	1,196	1,907	2,402	2,300	1,224	239	-
2002.....	261	306	369	487	720	1,157	1,787	2,290	2,168	1,104	230	-
2001.....	236	272	297	386	675	1,146	1,771	2,287	2,118	1,063	248	1
2000.....	167	236	303	416	630	1,157	1,721	2,309	2,031	985	218	-
1999.....	181	240	302	441	662	1,093	1,689	2,226	1,941	888	212	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

Black, both sexes

2003.....	710	662	572	601	583	666	574	535	369	170	65	-
2002.....	637	592	581	576	571	642	641	515	357	160	47	2
2001.....	571	506	487	517	578	605	571	526	315	156	65	5
2000.....	475	507	524	494	612	625	584	509	360	172	46	2
1999.....	439	499	446	521	584	637	558	456	299	156	46	1

Black, male

2003.....	411	393	310	352	304	284	202	151	93	26	6	-
2002.....	384	357	359	317	269	273	219	139	76	24	10	2
2001.....	338	288	275	274	272	252	214	149	62	28	12	4
2000.....	308	291	308	269	287	248	212	134	77	24	2	1
1999.....	254	292	245	281	266	272	191	132	60	30	3	1

Black, female

2003.....	299	269	262	249	279	382	372	384	276	144	59	-
2002.....	253	235	222	259	302	369	422	376	281	136	37	-
2001.....	233	218	212	243	306	353	357	377	253	128	53	1
2000.....	167	216	216	225	325	377	372	375	283	148	44	1
1999.....	185	207	201	240	318	365	367	324	239	126	43	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

American Indian 1/, both sexes

2003.....	11	13	12	13	5	7	7	9	9	3	-	-
2002.....	3	3	7	10	3	8	5	5	6	-	-	1
2001.....	6	6	4	7	3	10	6	3	5	4	-	-
2000.....	10	5	6	6	5	5	11	6	2	2	-	-
1999.....	2	9	5	3	7	6	9	7	2	-	-	-

American Indian 1/, male

2003.....	5	9	8	6	2	4	2	-	3	-	-	-
2002.....	2	1	4	4	-	4	3	-	1	-	-	1
2001.....	3	3	3	5	2	4	2	-	1	-	-	-
2000.....	7	2	3	4	4	3	5	1	1	1	-	-
1999.....	2	5	-	2	5	3	2	1	1	-	-	-

American Indian 1/, female

2003.....	6	4	4	7	3	3	5	9	6	3	-	-
2002.....	1	2	3	6	3	4	2	5	5	-	-	-
2001.....	3	3	1	2	1	6	4	3	4	4	-	-
2000.....	3	3	3	2	1	2	6	5	1	1	-	-
1999.....	-	4	5	1	2	3	7	6	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Asian or Pacific Islander 2/, both sexes												
2003.....	20	27	30	43	47	73	57	70	48	18	4	-
2002.....	25	30	31	36	40	57	74	62	51	17	6	-
2001.....	27	25	28	34	46	53	69	56	27	13	4	-
2000.....	20	16	26	38	39	52	59	47	37	24	4	-
1999.....	10	24	25	32	44	53	50	52	38	15	6	-
Asian or Pacific Islander 2/, male												
2003.....	17	16	18	22	18	35	22	32	14	5	-	-
2002.....	18	19	18	27	19	30	30	21	19	9	3	-
2001.....	16	17	19	14	21	27	27	22	9	7	1	-
2000.....	16	11	20	20	22	29	24	12	11	9	1	-
1999.....	8	15	16	19	20	27	22	22	8	1	1	-
Asian or Pacific Islander 2/, female												
2003.....	3	11	12	21	29	38	35	38	34	13	4	-
2002.....	7	11	13	9	21	27	44	41	32	8	3	-
2001.....	11	8	9	20	25	26	42	34	18	6	3	-
2000.....	4	5	6	18	17	23	35	35	26	15	3	-
1999.....	2	9	9	13	24	26	28	30	30	14	5	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

All races, both sexes

2003.....	114	136	131	154	247	386	517	560	456	167	26	-
2002.....	90	134	139	156	246	363	503	511	386	150	32	-
2001.....	82	109	138	169	248	374	481	499	348	155	24	-
2000.....	79	103	105	149	268	376	487	494	361	153	24	-
1999.....	76	116	146	201	269	487	600	600	411	162	36	-

All races, male

2003.....	67	86	73	75	114	174	208	207	128	45	6	-
2002.....	56	82	76	67	131	158	199	171	108	38	4	-
2001.....	53	61	65	94	112	158	181	174	102	35	3	-
2000.....	44	57	51	74	132	152	198	188	109	37	7	-
1999.....	50	66	80	99	124	221	239	205	123	41	5	-

All races, female

2003.....	47	50	58	79	133	212	309	353	328	122	20	-
2002.....	34	52	63	89	115	205	304	340	278	112	28	-
2001.....	29	48	73	75	136	216	300	325	246	120	21	-
2000.....	35	46	54	75	136	224	289	306	252	116	17	-
1999.....	26	50	66	102	145	266	361	395	288	121	31	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

White, both sexes

2003.....	28	47	64	75	154	279	406	442	391	144	17	-
2002.....	21	44	63	73	145	254	401	401	319	127	24	-
2001.....	35	45	71	87	152	258	384	395	295	132	16	-
2000.....	33	46	44	72	173	263	361	399	305	129	12	-
1999.....	31	47	63	101	148	343	463	473	351	129	29	-

White, male

2003.....	18	30	38	43	75	135	172	161	115	39	2	-
2002.....	11	30	35	32	82	116	160	135	90	32	2	-
2001.....	21	24	32	50	68	114	152	146	90	29	2	-
2000.....	19	25	21	38	88	120	153	150	90	28	2	-
1999.....	19	26	34	54	69	159	199	167	108	31	5	-

White, female

2003.....	10	17	26	32	79	144	234	281	276	105	15	-
2002.....	10	14	28	41	63	138	241	266	229	95	22	-
2001.....	14	21	39	37	84	144	232	249	205	103	14	-
2000.....	14	21	23	34	85	143	208	249	215	101	10	-
1999.....	12	21	29	47	79	184	264	306	243	98	24	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

Black, both sexes

2003.....	82	84	65	71	83	97	95	100	54	21	7	-
2002.....	69	89	71	75	92	101	88	99	60	22	8	-
2001.....	47	62	64	78	88	106	84	84	51	21	7	-
2000.....	46	55	58	75	89	99	110	82	50	19	10	-
1999.....	40	66	78	91	114	122	122	114	52	32	7	-

Black, male

2003.....	46	54	33	27	36	34	33	35	7	4	2	-
2002.....	45	51	37	29	42	37	30	32	17	6	2	-
2001.....	32	36	31	42	40	39	24	22	11	5	1	-
2000.....	25	31	27	36	41	29	40	29	16	6	4	-
1999.....	28	37	44	41	51	50	34	29	13	9	-	-

Black, female

2003.....	36	30	32	44	47	63	62	65	47	17	5	-
2002.....	24	38	34	46	50	64	58	67	43	16	6	-
2001.....	15	26	33	36	48	67	60	62	40	16	6	-
2000.....	21	24	31	39	48	70	70	53	34	13	6	-
1999.....	12	29	34	50	63	72	88	85	39	23	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

American Indian 1/, both sexes

2003.....	-	1	1	-	1	1	1	-	1	-	1	-
2002.....	-	-	-	1	-	-	1	2	1	-	-	-
2001.....	-	-	2	-	-	1	1	2	-	-	-	-
2000.....	-	1	-	1	-	1	1	4	-	-	-	-
1999.....	-	1	-	2	4	4	-	1	1	-	-	-

American Indian 1/, male

2003.....	-	-	1	-	1	-	1	-	1	-	1	-
2002.....	-	-	-	1	-	-	-	1	-	-	-	-
2001.....	-	-	2	-	-	-	-	1	-	-	-	-
2000.....	-	1	-	-	-	-	1	1	-	-	-	-
1999.....	-	1	-	1	2	2	-	1	-	-	-	-

American Indian 1/, female

2003.....	-	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	1	1	-	-	-
2001.....	-	-	-	-	-	1	1	1	-	-	-	-
2000.....	-	-	-	1	-	1	-	3	-	-	-	-
1999.....	-	-	-	1	2	2	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

Asian or Pacific Islander 2/, both sexes

2003.....	4	4	1	8	9	9	15	18	10	2	1	-
2002.....	-	1	5	7	9	8	13	9	6	1	-	-
2001.....	-	2	1	4	8	9	12	18	2	2	1	-
2000.....	-	1	3	1	6	13	15	9	6	5	2	-
1999.....	5	2	5	7	3	18	15	12	7	1	-	-

Asian or Pacific Islander 2/, male

2003.....	3	2	1	5	2	5	2	11	5	2	1	-
2002.....	-	1	4	5	7	5	9	3	1	-	-	-
2001.....	-	1	-	2	4	5	5	5	1	1	-	-
2000.....	-	-	3	-	3	3	4	8	3	3	1	-
1999.....	3	2	2	3	2	10	6	8	2	1	-	-

Asian or Pacific Islander 2/, female

2003.....	1	2	-	3	7	4	13	7	5	-	-	-
2002.....	-	-	1	2	2	3	4	6	5	1	-	-
2001.....	-	1	1	2	4	4	7	13	1	1	1	-
2000.....	-	1	-	1	3	10	11	1	3	2	1	-
1999.....	2	-	3	4	1	8	9	4	5	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

All races, both sexes

2003.....	15,372	20,907	26,470	32,648	46,232	66,062	83,308	81,998	58,794	23,939	5,433	29
2002.....	15,683	20,997	26,265	34,336	49,531	69,769	85,622	84,082	59,567	24,211	5,528	37
2001.....	15,612	20,270	26,304	35,502	51,961	73,082	86,772	84,880	59,558	24,206	5,665	48
2000.....	15,252	20,724	27,013	37,257	55,628	76,569	87,123	86,890	60,269	24,477	5,683	48
1999.....	15,251	20,575	27,802	40,080	58,760	79,379	89,917	88,158	60,502	24,983	5,733	39

All races, male

2003.....	11,820	15,342	18,425	21,547	28,452	37,191	41,382	33,563	18,269	5,232	808	22
2002.....	11,969	15,350	18,306	22,573	30,563	39,154	42,085	34,073	18,199	5,353	821	27
2001.....	11,818	14,794	18,212	23,181	31,891	40,894	42,059	33,466	17,702	5,124	814	33
2000.....	11,548	15,154	18,641	24,225	34,329	42,452	42,295	34,024	17,603	5,216	847	34
1999.....	11,614	15,021	19,102	26,184	36,003	44,041	43,264	34,168	17,472	5,245	841	31

All races, female

2003.....	3,552	5,565	8,045	11,101	17,780	28,871	41,926	48,435	40,525	18,707	4,625	7
2002.....	3,714	5,647	7,959	11,763	18,968	30,615	43,537	50,009	41,368	18,858	4,707	10
2001.....	3,794	5,476	8,092	12,321	20,070	32,188	44,713	51,414	41,856	19,082	4,851	15
2000.....	3,704	5,570	8,372	13,032	21,299	34,117	44,828	52,866	42,666	19,261	4,836	14
1999.....	3,637	5,554	8,700	13,896	22,757	35,338	46,653	53,990	43,030	19,738	4,892	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

White, both sexes

2003.....	12,085	17,028	21,621	27,015	39,524	58,115	75,130	75,347	54,228	21,994	4,730	23
2002.....	12,445	17,082	21,495	28,561	42,708	61,414	77,489	77,341	54,938	22,175	4,834	28
2001.....	12,441	16,436	21,493	29,456	44,934	64,552	78,643	78,104	55,084	22,157	4,976	30
2000.....	12,265	16,873	22,082	31,304	48,138	67,860	79,215	79,909	55,931	22,506	5,046	36
1999.....	12,272	16,733	22,933	33,894	50,959	70,631	81,999	81,066	56,125	22,985	5,122	27

White, male

2003.....	9,555	12,813	15,409	18,245	24,721	33,242	37,693	31,182	17,003	4,783	688	16
2002.....	9,771	12,781	15,313	19,219	26,909	35,116	38,536	31,656	16,828	4,882	701	21
2001.....	9,682	12,284	15,253	19,729	28,066	36,733	38,597	31,119	16,373	4,664	698	21
2000.....	9,559	12,586	15,635	20,885	30,317	38,199	38,901	31,567	16,346	4,726	729	27
1999.....	9,583	12,487	16,061	22,652	31,841	39,837	39,986	31,607	16,126	4,783	737	21

White, female

2003.....	2,530	4,215	6,212	8,770	14,803	24,873	37,437	44,165	37,225	17,211	4,042	7
2002.....	2,674	4,301	6,182	9,342	15,799	26,298	38,953	45,685	38,110	17,293	4,133	7
2001.....	2,759	4,152	6,240	9,727	16,868	27,819	40,046	46,985	38,711	17,493	4,278	9
2000.....	2,706	4,287	6,447	10,419	17,821	29,661	40,314	48,342	39,585	17,780	4,317	9
1999.....	2,689	4,246	6,872	11,242	19,118	30,794	42,013	49,459	39,999	18,202	4,385	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

Black, both sexes

2003.....	2,881	3,377	4,168	4,795	5,720	6,698	6,682	5,361	3,732	1,645	608	4
2002.....	2,866	3,441	4,166	4,920	5,815	6,962	6,757	5,534	3,808	1,717	602	9
2001.....	2,789	3,349	4,203	5,210	6,025	7,193	6,783	5,626	3,775	1,772	600	16
2000.....	2,624	3,383	4,302	5,124	6,495	7,298	6,668	5,891	3,644	1,676	568	12
1999.....	2,642	3,392	4,240	5,395	6,699	7,402	6,669	5,931	3,697	1,762	546	12

Black, male

2003.....	1,938	2,158	2,544	2,750	3,129	3,289	2,897	1,792	935	335	89	4
2002.....	1,896	2,219	2,574	2,821	3,070	3,255	2,841	1,850	1,022	357	88	6
2001.....	1,847	2,133	2,536	2,927	3,247	3,406	2,797	1,820	1,046	362	99	11
2000.....	1,701	2,227	2,561	2,796	3,409	3,451	2,748	1,939	957	369	104	7
1999.....	1,767	2,197	2,609	3,038	3,505	3,433	2,652	1,999	1,022	375	86	10

Black, female

2003.....	943	1,219	1,624	2,045	2,591	3,409	3,785	3,569	2,797	1,310	519	-
2002.....	970	1,222	1,592	2,099	2,745	3,707	3,916	3,684	2,786	1,360	514	3
2001.....	942	1,216	1,667	2,283	2,778	3,787	3,986	3,806	2,729	1,410	501	5
2000.....	923	1,156	1,741	2,328	3,086	3,847	3,920	3,952	2,687	1,307	464	5
1999.....	875	1,195	1,631	2,357	3,194	3,969	4,017	3,932	2,675	1,387	460	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

American Indian 1/, both sexes

2003.....	120	147	193	225	225	247	218	185	97	35	16	2
2002.....	112	144	179	205	221	254	211	154	101	45	10	-
2001.....	119	149	175	204	224	207	199	153	83	31	11	-
2000.....	101	135	167	229	185	246	192	156	113	45	7	-
1999.....	101	133	153	206	241	248	205	166	86	33	9	-

American Indian 1/, male

2003.....	93	104	125	138	143	133	115	80	38	11	6	2
2002.....	87	97	127	132	129	135	97	78	31	11	2	-
2001.....	82	111	122	121	138	112	90	67	25	10	2	-
2000.....	73	91	110	143	114	138	93	64	38	15	-	-
1999.....	73	95	96	134	141	140	94	59	30	6	2	-

American Indian 1/, female

2003.....	27	43	68	87	82	114	103	105	59	24	10	-
2002.....	25	47	52	73	92	119	114	76	70	34	8	-
2001.....	37	38	53	83	86	95	109	86	58	21	9	-
2000.....	28	44	57	86	71	108	99	92	75	30	7	-
1999.....	28	38	57	72	100	108	111	107	56	27	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

Asian or Pacific Islander 2/, both sexes

2003.....	286	355	488	613	763	1,002	1,278	1,105	737	265	79	-
2002.....	260	330	425	650	787	1,139	1,165	1,053	720	274	82	-
2001.....	263	336	433	632	778	1,130	1,147	997	616	246	78	2
2000.....	262	333	462	600	810	1,165	1,048	934	581	250	62	-
1999.....	236	317	476	585	861	1,098	1,044	995	594	203	56	-

Asian or Pacific Islander 2/, male

2003.....	234	267	347	414	459	527	677	509	293	103	25	-
2002.....	215	253	292	401	455	648	611	489	318	103	30	-
2001.....	207	266	301	404	440	643	575	460	258	88	15	1
2000.....	215	250	335	401	489	664	553	454	262	106	14	-
1999.....	191	242	336	360	516	631	532	503	294	81	16	-

Asian or Pacific Islander 2/, female

2003.....	52	88	141	199	304	475	601	596	444	162	54	-
2002.....	45	77	133	249	332	491	554	564	402	171	52	-
2001.....	56	70	132	228	338	487	572	537	358	158	63	1
2000.....	47	83	127	199	321	501	495	480	319	144	48	-
1999.....	45	75	140	225	345	467	512	492	300	122	40	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

All races, both sexes

2003.....	6,328	8,769	10,998	13,108	17,815	24,265	28,843	27,055	17,795	6,527	1,357	5
2002.....	6,564	8,758	11,175	14,230	19,773	26,154	30,436	28,423	18,352	6,623	1,320	5
2001.....	6,720	8,704	11,126	14,690	20,690	27,766	31,488	29,242	18,525	6,606	1,354	10
2000.....	6,647	9,030	11,983	15,898	22,561	30,059	32,151	29,922	18,701	6,755	1,367	8
1999.....	6,598	9,123	12,283	17,333	24,108	31,053	33,658	30,711	18,472	6,765	1,349	7

All races, male

2003.....	4,854	6,386	7,586	8,531	10,759	13,324	14,026	10,866	5,704	1,469	217	4
2002.....	4,929	6,287	7,679	9,214	12,050	14,316	14,603	11,478	5,741	1,559	213	4
2001.....	5,012	6,271	7,587	9,446	12,517	15,151	14,974	11,411	5,778	1,576	219	6
2000.....	4,946	6,490	8,148	10,177	13,646	16,295	15,336	11,765	5,701	1,539	238	5
1999.....	5,039	6,654	8,375	11,154	14,554	16,991	15,952	12,066	5,566	1,523	202	2

All races, female

2003.....	1,474	2,383	3,412	4,577	7,056	10,941	14,817	16,189	12,091	5,058	1,140	1
2002.....	1,635	2,471	3,496	5,016	7,723	11,838	15,833	16,945	12,611	5,064	1,107	1
2001.....	1,708	2,433	3,539	5,244	8,173	12,615	16,514	17,831	12,747	5,030	1,135	4
2000.....	1,701	2,540	3,835	5,721	8,915	13,764	16,815	18,157	13,000	5,216	1,129	3
1999.....	1,559	2,469	3,908	6,179	9,554	14,062	17,706	18,645	12,906	5,242	1,147	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

White, both sexes

2003.....	5,096	7,321	9,248	11,079	15,416	21,479	25,961	24,802	16,359	5,905	1,148	4
2002.....	5,374	7,341	9,380	12,067	17,184	23,108	27,457	25,992	16,780	5,985	1,116	4
2001.....	5,484	7,257	9,369	12,411	18,058	24,674	28,512	26,793	16,980	5,981	1,171	6
2000.....	5,458	7,574	10,085	13,575	19,710	26,733	29,233	27,383	17,270	6,112	1,174	7
1999.....	5,500	7,659	10,425	14,982	21,124	27,778	30,704	28,102	17,028	6,167	1,171	5

White, male

2003.....	4,019	5,478	6,526	7,397	9,464	11,962	12,739	10,067	5,274	1,330	177	3
2002.....	4,146	5,400	6,625	8,015	10,737	12,884	13,315	10,585	5,264	1,406	175	3
2001.....	4,225	5,373	6,564	8,205	11,150	13,690	13,744	10,533	5,289	1,410	187	5
2000.....	4,206	5,558	7,028	8,918	12,171	14,738	14,094	10,840	5,246	1,363	210	4
1999.....	4,319	5,690	7,277	9,830	13,039	15,479	14,755	11,087	5,116	1,375	166	1

White, female

2003.....	1,077	1,843	2,722	3,682	5,952	9,517	13,222	14,735	11,085	4,575	971	1
2002.....	1,228	1,941	2,755	4,052	6,447	10,224	14,142	15,407	11,516	4,579	941	1
2001.....	1,259	1,884	2,805	4,206	6,908	10,984	14,768	16,260	11,691	4,571	984	1
2000.....	1,252	2,016	3,057	4,657	7,539	11,995	15,139	16,543	12,024	4,749	964	3
1999.....	1,181	1,969	3,148	5,152	8,085	12,299	15,949	17,015	11,912	4,792	1,005	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

Black, both sexes

2003.....	1,071	1,254	1,471	1,719	2,042	2,359	2,352	1,828	1,197	545	183	1
2002.....	1,058	1,255	1,569	1,823	2,207	2,527	2,461	2,002	1,291	572	180	1
2001.....	1,068	1,265	1,518	1,955	2,261	2,604	2,478	2,028	1,309	544	157	4
2000.....	1,038	1,277	1,658	1,982	2,465	2,761	2,475	2,151	1,207	542	178	1
1999.....	971	1,271	1,611	2,027	2,569	2,773	2,525	2,183	1,254	540	159	2

Black, male

2003.....	702	771	871	935	1,076	1,131	997	603	325	106	34	1
2002.....	678	769	897	994	1,109	1,157	1,016	690	356	123	25	1
2001.....	658	768	858	1,042	1,148	1,181	987	679	377	134	28	1
2000.....	626	798	950	1,041	1,254	1,254	1,016	727	348	128	27	1
1999.....	624	819	937	1,120	1,284	1,246	984	765	356	128	30	1

Black, female

2003.....	369	483	600	784	966	1,228	1,355	1,225	872	439	149	-
2002.....	380	486	672	829	1,098	1,370	1,445	1,312	935	449	155	-
2001.....	410	497	660	913	1,113	1,423	1,491	1,349	932	410	129	3
2000.....	412	479	708	941	1,211	1,507	1,459	1,424	859	414	151	-
1999.....	347	452	674	907	1,285	1,527	1,541	1,418	898	412	129	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

American Indian 1/, both sexes

2003.....	45	58	94	92	80	89	83	56	25	10	2	-
2002.....	41	55	77	79	86	97	69	51	34	9	1	-
2001.....	57	53	72	76	86	69	70	62	28	15	5	-
2000.....	42	54	59	97	66	96	70	50	38	15	2	-
1999.....	39	61	55	90	93	92	85	64	24	11	4	-

American Indian 1/, male

2003.....	38	38	63	58	47	48	42	25	11	3	-	-
2002.....	34	42	57	49	41	44	29	25	11	3	1	-
2001.....	41	40	49	44	52	38	35	33	11	6	-	-
2000.....	30	38	35	64	37	54	33	22	12	7	-	-
1999.....	27	45	32	60	53	45	39	28	7	1	1	-

American Indian 1/, female

2003.....	7	20	31	34	33	41	41	31	14	7	2	-
2002.....	7	13	20	30	45	53	40	26	23	6	-	-
2001.....	16	13	23	32	34	31	35	29	17	9	5	-
2000.....	12	16	24	33	29	42	37	28	26	8	2	-
1999.....	12	16	23	30	40	47	46	36	17	10	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

Asian or Pacific Islander 2/, both sexes

2003.....	116	136	185	218	277	338	447	369	214	67	24	-
2002.....	91	107	149	261	296	422	449	378	247	57	23	-
2001.....	111	129	167	248	285	419	428	359	208	66	21	-
2000.....	109	125	181	244	320	469	373	338	186	86	13	-
1999.....	88	132	192	234	322	410	344	362	166	47	15	-

Asian or Pacific Islander 2/, male

2003.....	95	99	126	141	172	183	248	171	94	30	6	-
2002.....	71	76	100	156	163	231	243	178	110	27	12	-
2001.....	88	90	116	155	167	242	208	166	101	26	4	-
2000.....	84	96	135	154	184	249	193	176	95	41	1	-
1999.....	69	100	129	144	178	221	174	186	87	19	5	-

Asian or Pacific Islander 2/, female

2003.....	21	37	59	77	105	155	199	198	120	37	18	-
2002.....	20	31	49	105	133	191	206	200	137	30	11	-
2001.....	23	39	51	93	118	177	220	193	107	40	17	-
2000.....	25	29	46	90	136	220	180	162	91	45	12	-
1999.....	19	32	63	90	144	189	170	176	79	28	10	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

All races, both sexes

2003.....	171	198	205	267	312	389	448	447	337	129	28	1
2002.....	226	210	250	289	342	420	478	452	339	153	27	-
2001.....	187	200	263	281	364	397	474	455	317	142	31	-
2000.....	189	219	265	294	350	440	466	439	318	136	35	1
1999.....	160	209	239	294	380	406	449	474	296	116	22	-

All races, male

2003.....	122	137	141	173	188	220	194	159	93	21	9	1
2002.....	162	153	167	193	197	216	228	172	87	35	4	-
2001.....	141	134	186	180	213	206	218	165	72	25	7	-
2000.....	141	163	178	194	204	232	214	153	68	28	5	1
1999.....	109	159	156	191	219	229	194	170	75	19	2	-

All races, female

2003.....	49	61	64	94	124	169	254	288	244	108	19	-
2002.....	64	57	83	96	145	204	250	280	252	118	23	-
2001.....	46	66	77	101	151	191	256	290	245	117	24	-
2000.....	48	56	87	100	146	208	252	286	250	108	30	-
1999.....	51	50	83	103	161	177	255	304	221	97	20	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

White, both sexes

2003.....	132	150	161	214	259	332	397	406	307	124	25	1
2002.....	171	172	195	237	287	369	427	409	312	143	23	-
2001.....	137	154	201	219	312	345	426	406	298	125	26	-
2000.....	145	160	207	236	294	373	417	400	288	130	32	-
1999.....	120	162	195	247	316	354	396	438	279	104	21	-

White, male

2003.....	97	102	111	141	161	185	166	144	86	21	6	1
2002.....	123	126	133	159	164	193	203	162	78	32	4	-
2001.....	103	107	143	135	187	179	203	148	66	23	7	-
2000.....	107	124	142	154	171	199	193	142	60	27	4	-
1999.....	85	122	124	160	183	206	169	159	69	17	2	-

White, female

2003.....	35	48	50	73	98	147	231	262	221	103	19	-
2002.....	48	46	62	78	123	176	224	247	234	111	19	-
2001.....	34	47	58	84	125	166	223	258	232	102	19	-
2000.....	38	36	65	82	123	174	224	258	228	103	28	-
1999.....	35	40	71	87	133	148	227	279	210	87	19	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Black, both sexes

2003.....	32	43	39	47	47	51	46	36	26	4	2	-
2002.....	51	32	47	47	53	45	45	38	21	8	4	-
2001.....	42	40	58	49	46	40	40	45	12	16	5	-
2000.....	40	51	52	47	46	56	43	35	26	6	3	1
1999.....	39	43	42	39	54	49	48	28	15	10	1	-

Black, male

2003.....	22	33	26	30	24	31	26	14	7	-	2	-
2002.....	35	21	29	32	32	20	21	9	8	3	-	-
2001.....	33	22	40	36	23	24	11	15	4	2	-	-
2000.....	31	35	32	32	27	28	18	9	7	1	1	1
1999.....	23	34	31	26	30	21	20	10	5	2	-	-

Black, female

2003.....	10	10	13	17	23	20	20	22	19	4	-	-
2002.....	16	11	18	15	21	25	24	29	13	5	4	-
2001.....	9	18	18	13	23	16	29	30	8	14	5	-
2000.....	9	16	20	15	19	28	25	26	19	5	2	-
1999.....	16	9	11	13	24	28	28	18	10	8	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

American Indian 1/, both sexes

2003.....	4	4	4	4	5	3	3	2	2	-	1	-
2002.....	3	3	5	3	2	4	4	1	5	2	-	-
2001.....	7	5	2	9	5	6	2	3	3	-	-	-
2000.....	4	4	5	8	7	4	4	1	2	-	-	-
1999.....	1	3	2	6	8	3	3	6	1	1	-	-

American Indian 1/, male

2003.....	2	1	4	1	2	1	-	-	-	-	1	-
2002.....	3	3	4	1	1	2	3	-	1	-	-	-
2001.....	5	5	1	7	3	3	-	1	-	-	-	-
2000.....	3	2	3	6	4	3	1	1	1	-	-	-
1999.....	1	2	1	3	5	2	3	1	-	-	-	-

American Indian 1/, female

2003.....	2	3	-	3	3	2	3	2	2	-	-	-
2002.....	-	-	1	2	1	2	1	1	4	2	-	-
2001.....	2	-	1	2	2	3	2	2	3	-	-	-
2000.....	1	2	2	2	3	1	3	-	1	-	-	-
1999.....	-	1	1	3	3	1	-	5	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Asian or Pacific Islander 2/, both sexes

2003.....	3	1	1	2	1	3	2	3	2	1	-	-
2002.....	1	3	3	2	-	2	2	4	1	-	-	-
2001.....	1	1	2	4	1	6	6	1	4	1	-	-
2000.....	-	4	1	3	3	7	2	3	2	-	-	-
1999.....	-	1	-	2	2	-	2	2	1	1	-	-

Asian or Pacific Islander 2/, male

2003.....	1	1	-	1	1	3	2	1	-	-	-	-
2002.....	1	3	1	1	-	1	1	1	-	-	-	-
2001.....	-	-	2	2	-	-	4	1	2	-	-	-
2000.....	-	2	1	2	2	2	2	1	-	-	-	-
1999.....	-	1	-	2	1	-	2	-	1	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	-	1	1	-	-	-	2	2	1	-	-
2002.....	-	-	2	1	-	1	1	3	1	-	-	-
2001.....	1	1	-	2	1	6	2	-	2	1	-	-
2000.....	-	2	-	1	1	5	-	2	2	-	-	-
1999.....	-	-	-	-	1	-	-	2	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

All races, both sexes

2003.....	8,873	11,940	15,267	19,273	28,105	41,408	54,017	54,496	40,662	17,283	4,048	23
2002.....	8,893	12,029	14,840	19,817	29,416	43,195	54,708	55,207	40,876	17,435	4,181	32
2001.....	8,705	11,366	14,915	20,531	30,907	44,919	54,810	55,183	40,716	17,458	4,280	38
2000.....	8,416	11,475	14,765	21,065	32,717	46,070	54,506	56,529	41,250	17,586	4,281	39
1999.....	8,493	11,243	15,280	22,453	34,272	47,920	55,810	56,973	41,734	18,102	4,362	32

All races, male

2003.....	6,844	8,819	10,698	12,843	17,505	23,647	27,162	22,538	12,472	3,742	582	17
2002.....	6,878	8,910	10,460	13,166	18,316	24,622	27,254	22,423	12,371	3,759	604	23
2001.....	6,665	8,389	10,439	13,555	19,161	25,537	26,867	21,890	11,852	3,523	588	27
2000.....	6,461	8,501	10,315	13,854	20,479	25,925	26,745	22,106	11,834	3,649	604	28
1999.....	6,466	8,208	10,571	14,839	21,230	26,821	27,118	21,932	11,831	3,703	637	29

All races, female

2003.....	2,029	3,121	4,569	6,430	10,600	17,761	26,855	31,958	28,190	13,541	3,466	6
2002.....	2,015	3,119	4,380	6,651	11,100	18,573	27,454	32,784	28,505	13,676	3,577	9
2001.....	2,040	2,977	4,476	6,976	11,746	19,382	27,943	33,293	28,864	13,935	3,692	11
2000.....	1,955	2,974	4,450	7,211	12,238	20,145	27,761	34,423	29,416	13,937	3,677	11
1999.....	2,027	3,035	4,709	7,614	13,042	21,099	28,692	35,041	29,903	14,399	3,725	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

White, both sexes

2003.....	6,857	9,557	12,212	15,722	23,849	36,304	48,772	50,139	37,562	15,965	3,557	18
2002.....	6,900	9,569	11,920	16,257	25,237	37,937	49,605	50,940	37,846	16,047	3,695	24
2001.....	6,820	9,025	11,923	16,826	26,564	39,533	49,705	50,905	37,806	16,051	3,779	24
2000.....	6,662	9,139	11,790	17,493	28,134	40,754	49,565	52,126	38,373	16,264	3,840	29
1999.....	6,652	8,912	12,313	18,665	29,519	42,499	50,899	52,526	38,818	16,714	3,930	22

White, male

2003.....	5,439	7,233	8,772	10,707	15,096	21,095	24,788	20,971	11,643	3,432	505	12
2002.....	5,502	7,255	8,555	11,045	16,008	22,039	25,018	20,909	11,486	3,444	522	18
2001.....	5,354	6,804	8,546	11,389	16,729	22,864	24,650	20,438	11,018	3,231	504	16
2000.....	5,246	6,904	8,465	11,813	17,975	23,262	24,614	20,585	11,040	3,336	515	23
1999.....	5,179	6,675	8,660	12,662	18,619	24,152	25,062	20,361	10,941	3,391	569	20

White, female

2003.....	1,418	2,324	3,440	5,015	8,753	15,209	23,984	29,168	25,919	12,533	3,052	6
2002.....	1,398	2,314	3,365	5,212	9,229	15,898	24,587	30,031	26,360	12,603	3,173	6
2001.....	1,466	2,221	3,377	5,437	9,835	16,669	25,055	30,467	26,788	12,820	3,275	8
2000.....	1,416	2,235	3,325	5,680	10,159	17,492	24,951	31,541	27,333	12,928	3,325	6
1999.....	1,473	2,237	3,653	6,003	10,900	18,347	25,837	32,165	27,877	13,323	3,361	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Black, both sexes												
2003.....	1,778	2,080	2,658	3,029	3,631	4,288	4,284	3,497	2,509	1,096	423	3
2002.....	1,757	2,154	2,550	3,050	3,555	4,390	4,251	3,494	2,496	1,137	418	8
2001.....	1,679	2,044	2,627	3,206	3,718	4,549	4,265	3,553	2,454	1,212	438	12
2000.....	1,546	2,055	2,592	3,095	3,984	4,481	4,150	3,705	2,411	1,128	387	10
1999.....	1,632	2,078	2,587	3,329	4,076	4,580	4,096	3,720	2,428	1,212	386	10
Black, male												
2003.....	1,214	1,354	1,647	1,785	2,029	2,127	1,874	1,175	603	229	53	3
2002.....	1,183	1,429	1,648	1,795	1,929	2,078	1,804	1,151	658	231	63	5
2001.....	1,156	1,343	1,638	1,849	2,076	2,201	1,799	1,126	665	226	71	10
2000.....	1,044	1,394	1,579	1,723	2,128	2,169	1,714	1,203	602	240	76	5
1999.....	1,120	1,344	1,641	1,892	2,191	2,166	1,648	1,224	661	245	56	9
Black, female												
2003.....	564	726	1,011	1,244	1,602	2,161	2,410	2,322	1,906	867	370	-
2002.....	574	725	902	1,255	1,626	2,312	2,447	2,343	1,838	906	355	3
2001.....	523	701	989	1,357	1,642	2,348	2,466	2,427	1,789	986	367	2
2000.....	502	661	1,013	1,372	1,856	2,312	2,436	2,502	1,809	888	311	5
1999.....	512	734	946	1,437	1,885	2,414	2,448	2,496	1,767	967	330	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

American Indian 1/, both sexes

2003.....	71	85	95	129	140	155	132	127	70	25	13	2
2002.....	68	86	97	123	133	153	138	102	62	34	9	-
2001.....	55	91	101	119	133	132	127	88	52	16	6	-
2000.....	55	77	103	124	112	146	118	105	73	30	5	-
1999.....	61	69	96	110	140	153	117	96	61	21	5	-

American Indian 1/, male

2003.....	53	65	58	79	94	84	73	55	27	8	5	2
2002.....	50	52	66	82	87	89	65	53	19	8	1	-
2001.....	36	66	72	70	83	71	55	33	14	4	2	-
2000.....	40	51	72	73	73	81	59	41	25	8	-	-
1999.....	45	48	63	71	83	93	52	30	23	5	1	-

American Indian 1/, female

2003.....	18	20	37	50	46	71	59	72	43	17	8	-
2002.....	18	34	31	41	46	64	73	49	43	26	8	-
2001.....	19	25	29	49	50	61	72	55	38	12	4	-
2000.....	15	26	31	51	39	65	59	64	48	22	5	-
1999.....	16	21	33	39	57	60	65	66	38	16	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Asian or Pacific Islander 2/, both sexes

2003.....	167	218	302	393	485	661	829	733	521	197	55	-
2002.....	168	220	273	387	491	715	714	671	472	217	59	-
2001.....	151	206	264	380	492	705	713	637	404	179	57	2
2000.....	153	204	280	353	487	689	673	593	393	164	49	-
1999.....	148	184	284	349	537	688	698	631	427	155	41	-

Asian or Pacific Islander 2/, male

2003.....	138	167	221	272	286	341	427	337	199	73	19	-
2002.....	143	174	191	244	292	416	367	310	208	76	18	-
2001.....	119	176	183	247	273	401	363	293	155	62	11	1
2000.....	131	152	199	245	303	413	358	277	167	65	13	-
1999.....	122	141	207	214	337	410	356	317	206	62	11	-

Asian or Pacific Islander 2/, female

2003.....	29	51	81	121	199	320	402	396	322	124	36	-
2002.....	25	46	82	143	199	299	347	361	264	141	41	-
2001.....	32	30	81	133	219	304	350	344	249	117	46	1
2000.....	22	52	81	108	184	276	315	316	226	99	36	-
1999.....	26	43	77	135	200	278	342	314	221	93	30	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races, both sexes

2003.....	3,533	4,208	4,902	5,194	6,300	8,253	10,086	9,600	7,316	3,135	823	14
2002.....	3,285	4,189	4,604	5,163	6,654	8,649	9,969	10,007	7,555	3,361	873	23
2001.....	3,209	3,924	4,514	5,272	6,733	9,041	10,155	9,984	7,556	3,323	873	25
2000.....	3,086	3,841	4,401	5,309	7,258	9,271	10,274	10,397	7,953	3,429	885	19
1999.....	2,945	3,747	4,636	5,691	7,704	9,666	10,655	10,745	8,204	3,591	863	24

All races, male

2003.....	2,837	3,203	3,491	3,490	3,932	4,543	4,790	3,759	2,106	700	112	12
2002.....	2,605	3,250	3,299	3,498	4,119	4,746	4,670	3,814	2,165	722	134	17
2001.....	2,549	2,978	3,267	3,483	4,185	4,897	4,774	3,630	2,167	616	108	19
2000.....	2,444	2,978	3,147	3,557	4,512	5,086	4,639	3,740	2,205	691	122	15
1999.....	2,314	2,842	3,281	3,842	4,704	5,200	4,763	3,895	2,196	692	122	22

All races, female

2003.....	696	1,005	1,411	1,704	2,368	3,710	5,296	5,841	5,210	2,435	711	2
2002.....	680	939	1,305	1,665	2,535	3,903	5,299	6,193	5,390	2,639	739	6
2001.....	660	946	1,247	1,789	2,548	4,144	5,381	6,354	5,389	2,707	765	6
2000.....	642	863	1,254	1,752	2,746	4,185	5,635	6,657	5,748	2,738	763	4
1999.....	631	905	1,355	1,849	3,000	4,466	5,892	6,850	6,008	2,899	741	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

White, both sexes

2003.....	2,653	3,255	3,672	3,976	5,000	6,822	8,750	8,563	6,593	2,837	707	10
2002.....	2,446	3,170	3,512	3,898	5,324	7,130	8,653	8,950	6,825	3,024	740	16
2001.....	2,413	2,897	3,335	3,941	5,375	7,447	8,869	8,927	6,867	2,978	745	16
2000.....	2,339	2,858	3,255	4,093	5,765	7,717	8,892	9,245	7,202	3,112	760	13
1999.....	2,183	2,765	3,446	4,342	6,130	8,014	9,303	9,568	7,430	3,248	749	15

White, male

2003.....	2,180	2,530	2,664	2,718	3,127	3,805	4,169	3,376	1,902	623	93	8
2002.....	1,999	2,504	2,563	2,678	3,362	3,952	4,048	3,427	1,953	646	111	12
2001.....	1,968	2,270	2,466	2,665	3,379	4,090	4,206	3,272	1,958	548	86	11
2000.....	1,903	2,252	2,408	2,811	3,636	4,249	4,037	3,348	1,997	611	93	12
1999.....	1,763	2,156	2,477	3,001	3,800	4,348	4,178	3,478	1,964	615	108	13

White, female

2003.....	473	725	1,008	1,258	1,873	3,017	4,581	5,187	4,691	2,214	614	2
2002.....	447	666	949	1,220	1,962	3,178	4,605	5,523	4,872	2,378	629	4
2001.....	445	627	869	1,276	1,996	3,357	4,663	5,655	4,909	2,430	659	5
2000.....	436	606	847	1,282	2,129	3,468	4,855	5,897	5,205	2,501	667	1
1999.....	420	609	969	1,341	2,330	3,666	5,125	6,090	5,466	2,633	641	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Black, both sexes

2003.....	775	840	1,086	1,062	1,128	1,258	1,114	880	620	264	98	3
2002.....	754	901	984	1,125	1,175	1,350	1,148	903	626	291	118	7
2001.....	717	911	1,054	1,186	1,208	1,405	1,113	922	602	299	113	8
2000.....	681	882	1,029	1,083	1,325	1,349	1,204	1,010	663	280	114	6
1999.....	680	889	1,053	1,212	1,396	1,466	1,173	1,019	671	304	103	9

Black, male

2003.....	573	580	726	655	697	660	515	320	160	67	13	3
2002.....	530	652	659	722	660	699	546	308	170	63	18	5
2001.....	517	620	708	722	716	711	484	295	174	51	17	7
2000.....	489	646	662	648	769	716	505	340	175	67	26	3
1999.....	485	612	697	760	799	737	499	350	188	64	9	9

Black, female

2003.....	202	260	360	407	431	598	599	560	460	197	85	-
2002.....	224	249	325	403	515	651	602	595	456	228	100	2
2001.....	200	291	346	464	492	694	629	627	428	248	96	1
2000.....	192	236	367	435	556	633	699	670	488	213	88	3
1999.....	195	277	356	452	597	729	674	669	483	240	94	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

American Indian 1/, both sexes

2003.....	31	35	26	45	49	39	31	27	18	7	7	1
2002.....	27	37	23	38	43	32	34	23	22	7	3	-
2001.....	29	31	46	45	38	37	34	19	17	5	2	-
2000.....	20	27	25	31	24	33	25	30	15	5	1	-
1999.....	24	24	28	33	37	45	34	25	19	4	-	-

American Indian 1/, male

2003.....	24	26	16	29	37	17	20	10	9	2	3	1
2002.....	23	24	14	28	27	17	15	13	7	3	-	-
2001.....	22	18	37	26	24	17	17	7	5	-	2	-
2000.....	14	21	18	18	15	18	15	9	6	1	-	-
1999.....	17	18	22	23	21	29	15	9	6	1	-	-

American Indian 1/, female

2003.....	7	9	10	16	12	22	11	17	9	5	4	-
2002.....	4	13	9	10	16	15	19	10	15	4	3	-
2001.....	7	13	9	19	14	20	17	12	12	5	-	-
2000.....	6	6	7	13	9	15	10	21	9	4	1	-
1999.....	7	6	6	10	16	16	19	16	13	3	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Asian or Pacific Islander 2/, both sexes

2003.....	74	78	118	111	123	134	191	130	85	27	11	-
2002.....	58	81	85	102	112	137	134	131	82	39	12	-
2001.....	50	85	79	100	112	152	139	116	70	41	13	1
2000.....	46	74	92	102	144	172	153	112	73	32	10	-
1999.....	58	69	109	104	141	141	145	133	84	35	11	-

Asian or Pacific Islander 2/, male

2003.....	60	67	85	88	71	61	86	53	35	8	3	-
2002.....	53	70	63	70	70	78	61	66	35	10	5	-
2001.....	42	70	56	70	66	79	67	56	30	17	3	1
2000.....	38	59	59	80	92	103	82	43	27	12	3	-
1999.....	49	56	85	58	84	86	71	58	38	12	5	-

Asian or Pacific Islander 2/, female

2003.....	14	11	33	23	52	73	105	77	50	19	8	-
2002.....	5	11	22	32	42	59	73	65	47	29	7	-
2001.....	8	15	23	30	46	73	72	60	40	24	10	-
2000.....	8	15	33	22	52	69	71	69	46	20	7	-
1999.....	9	13	24	46	57	55	74	75	46	23	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races, both sexes

2003.....	5,340	7,732	10,365	14,079	21,805	33,155	43,931	44,896	33,346	14,148	3,225	9
2002.....	5,608	7,840	10,236	14,654	22,762	34,546	44,739	45,200	33,321	14,074	3,308	9
2001.....	5,496	7,442	10,401	15,259	24,174	35,878	44,655	45,199	33,160	14,135	3,407	13
2000.....	5,330	7,634	10,364	15,756	25,459	36,799	44,232	46,132	33,297	14,157	3,396	20
1999.....	5,548	7,496	10,644	16,762	26,568	38,254	45,155	46,228	33,530	14,511	3,499	8

All races, male

2003.....	4,007	5,616	7,207	9,353	13,573	19,104	22,372	18,779	10,366	3,042	470	5
2002.....	4,273	5,660	7,161	9,668	14,197	19,876	22,584	18,609	10,206	3,037	470	6
2001.....	4,116	5,411	7,172	10,072	14,976	20,640	22,093	18,260	9,685	2,907	480	8
2000.....	4,017	5,523	7,168	10,297	15,967	20,839	22,106	18,366	9,629	2,958	482	13
1999.....	4,152	5,366	7,290	10,997	16,526	21,621	22,355	18,037	9,635	3,011	515	7

All races, female

2003.....	1,333	2,116	3,158	4,726	8,232	14,051	21,559	26,117	22,980	11,106	2,755	4
2002.....	1,335	2,180	3,075	4,986	8,565	14,670	22,155	26,591	23,115	11,037	2,838	3
2001.....	1,380	2,031	3,229	5,187	9,198	15,238	22,562	26,939	23,475	11,228	2,927	5
2000.....	1,313	2,111	3,196	5,459	9,492	15,960	22,126	27,766	23,668	11,199	2,914	7
1999.....	1,396	2,130	3,354	5,765	10,042	16,633	22,800	28,191	23,895	11,500	2,984	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, both sexes

2003.....	4,204	6,302	8,540	11,746	18,849	29,482	40,022	41,576	30,969	13,128	2,850	8
2002.....	4,454	6,399	8,408	12,359	19,913	30,807	40,952	41,990	31,021	13,023	2,955	8
2001.....	4,407	6,128	8,588	12,885	21,189	32,086	40,836	41,978	30,939	13,073	3,034	8
2000.....	4,323	6,281	8,535	13,400	22,369	33,037	40,673	42,881	31,171	13,152	3,080	16
1999.....	4,469	6,147	8,867	14,323	23,389	34,485	41,596	42,958	31,388	13,466	3,181	7

White, male

2003.....	3,259	4,703	6,108	7,989	11,969	17,290	20,619	17,595	9,741	2,809	412	4
2002.....	3,503	4,751	5,992	8,367	12,646	18,087	20,970	17,482	9,533	2,798	411	6
2001.....	3,386	4,534	6,080	8,724	13,350	18,774	20,444	17,166	9,060	2,683	418	5
2000.....	3,343	4,652	6,057	9,002	14,339	19,013	20,577	17,237	9,043	2,725	422	11
1999.....	3,416	4,519	6,183	9,661	14,819	19,804	20,884	16,883	8,977	2,776	461	7

White, female

2003.....	945	1,599	2,432	3,757	6,880	12,192	19,403	23,981	21,228	10,319	2,438	4
2002.....	951	1,648	2,416	3,992	7,267	12,720	19,982	24,508	21,488	10,225	2,544	2
2001.....	1,021	1,594	2,508	4,161	7,839	13,312	20,392	24,812	21,879	10,390	2,616	3
2000.....	980	1,629	2,478	4,398	8,030	14,024	20,096	25,644	22,128	10,427	2,658	5
1999.....	1,053	1,628	2,684	4,662	8,570	14,681	20,712	26,075	22,411	10,690	2,720	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, both sexes

2003.....	1,003	1,240	1,572	1,967	2,503	3,030	3,170	2,617	1,889	832	325	-
2002.....	1,003	1,253	1,566	1,925	2,380	3,040	3,103	2,591	1,870	846	300	1
2001.....	962	1,133	1,573	2,020	2,510	3,144	3,152	2,631	1,852	913	325	4
2000.....	865	1,173	1,563	2,012	2,659	3,132	2,946	2,695	1,748	848	273	4
1999.....	952	1,189	1,534	2,117	2,680	3,114	2,923	2,701	1,757	908	283	1

Black, male

2003.....	641	774	921	1,130	1,332	1,467	1,359	855	443	162	40	-
2002.....	653	777	989	1,073	1,269	1,379	1,258	843	488	168	45	-
2001.....	639	723	930	1,127	1,360	1,490	1,315	831	491	175	54	3
2000.....	555	748	917	1,075	1,359	1,453	1,209	863	427	173	50	2
1999.....	635	732	944	1,132	1,392	1,429	1,149	874	473	181	47	-

Black, female

2003.....	362	466	651	837	1,171	1,563	1,811	1,762	1,446	670	285	-
2002.....	350	476	577	852	1,111	1,661	1,845	1,748	1,382	678	255	1
2001.....	323	410	643	893	1,150	1,654	1,837	1,800	1,361	738	271	1
2000.....	310	425	646	937	1,300	1,679	1,737	1,832	1,321	675	223	2
1999.....	317	457	590	985	1,288	1,685	1,774	1,827	1,284	727	236	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

American Indian 1/, both sexes

2003.....	40	50	69	84	91	116	101	100	52	18	6	1
2002.....	41	49	74	85	90	121	104	79	40	27	6	-
2001.....	26	60	55	74	95	95	93	69	35	11	4	-
2000.....	35	50	78	93	88	113	93	75	58	25	4	-
1999.....	37	45	68	77	103	108	83	71	42	17	5	-

American Indian 1/, male

2003.....	29	39	42	50	57	67	53	45	18	6	2	1
2002.....	27	28	52	54	60	72	50	40	12	5	1	-
2001.....	14	48	35	44	59	54	38	26	9	4	-	-
2000.....	26	30	54	55	58	63	44	32	19	7	-	-
1999.....	28	30	41	48	62	64	37	21	17	4	1	-

American Indian 1/, female

2003.....	11	11	27	34	34	49	48	55	34	12	4	-
2002.....	14	21	22	31	30	49	54	39	28	22	5	-
2001.....	12	12	20	30	36	41	55	43	26	7	4	-
2000.....	9	20	24	38	30	50	49	43	39	18	4	-
1999.....	9	15	27	29	41	44	46	50	25	13	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Asian or Pacific Islander 2/, both sexes

2003.....	93	140	184	282	362	527	638	603	436	170	44	-
2002.....	110	139	188	285	379	578	580	540	390	178	47	-
2001.....	101	121	185	280	380	553	574	521	334	138	44	1
2000.....	107	130	188	251	343	517	520	481	320	132	39	-
1999.....	90	115	175	245	396	547	553	498	343	120	30	-

Asian or Pacific Islander 2/, male

2003.....	78	100	136	184	215	280	341	284	164	65	16	-
2002.....	90	104	128	174	222	338	306	244	173	66	13	-
2001.....	77	106	127	177	207	322	296	237	125	45	8	-
2000.....	93	93	140	165	211	310	276	234	140	53	10	-
1999.....	73	85	122	156	253	324	285	259	168	50	6	-

Asian or Pacific Islander 2/, female

2003.....	15	40	48	98	147	247	297	319	272	105	28	-
2002.....	20	35	60	111	157	240	274	296	217	112	34	-
2001.....	24	15	58	103	173	231	278	284	209	93	36	1
2000.....	14	37	48	86	132	207	244	247	180	79	29	-
1999.....	17	30	53	89	143	223	268	239	175	70	24	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

All races, both sexes

2003.....	4,825	6,171	7,227	9,481	13,763	20,800	28,460	31,015	23,947	9,993	2,315	6
2002.....	4,862	5,913	7,033	9,377	14,303	21,276	28,500	30,794	23,517	9,779	2,210	6
2001.....	4,776	5,637	6,898	9,596	14,460	21,325	27,651	30,343	22,779	9,653	2,204	7
2000.....	4,551	5,417	6,945	9,875	14,804	21,524	26,843	29,519	22,677	9,591	2,206	8
1999.....	4,386	5,465	7,115	10,034	15,538	21,689	27,023	29,428	21,894	9,401	2,125	7

All races, male

2003.....	3,019	3,826	4,316	5,402	7,473	10,351	12,329	11,327	6,827	2,169	358	5
2002.....	3,082	3,587	4,184	5,275	7,716	10,463	12,308	11,167	6,558	2,070	338	3
2001.....	2,946	3,393	4,110	5,356	7,811	10,538	12,002	10,595	6,187	2,028	331	5
2000.....	2,873	3,384	4,148	5,584	8,037	10,505	11,501	10,488	6,178	1,983	359	5
1999.....	2,756	3,419	4,281	5,870	8,450	10,567	11,561	10,293	5,879	1,988	323	5

All races, female

2003.....	1,806	2,345	2,911	4,079	6,290	10,449	16,131	19,688	17,120	7,824	1,957	1
2002.....	1,780	2,326	2,849	4,102	6,587	10,813	16,192	19,627	16,959	7,709	1,872	3
2001.....	1,830	2,244	2,788	4,240	6,649	10,787	15,649	19,748	16,592	7,625	1,873	2
2000.....	1,678	2,033	2,797	4,291	6,767	11,019	15,342	19,031	16,499	7,608	1,847	3
1999.....	1,630	2,046	2,834	4,164	7,088	11,122	15,462	19,135	16,015	7,413	1,802	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

White, both sexes

2003.....	3,395	4,688	5,588	7,597	11,504	18,308	25,766	28,661	22,248	9,241	2,060	5
2002.....	3,481	4,482	5,499	7,614	12,102	18,674	25,844	28,481	21,808	9,026	1,946	3
2001.....	3,453	4,221	5,447	7,747	12,305	18,686	25,072	28,039	21,213	8,946	1,966	6
2000.....	3,294	4,139	5,459	8,050	12,567	18,934	24,445	27,240	21,096	8,849	1,974	7
1999.....	3,192	4,079	5,549	8,177	13,197	19,031	24,547	27,066	20,457	8,697	1,900	5

White, male

2003.....	2,161	2,934	3,409	4,390	6,369	9,229	11,227	10,524	6,355	1,995	304	4
2002.....	2,272	2,764	3,339	4,341	6,619	9,283	11,250	10,394	6,044	1,909	290	2
2001.....	2,211	2,576	3,296	4,419	6,708	9,306	10,903	9,804	5,733	1,859	286	5
2000.....	2,107	2,636	3,323	4,619	6,942	9,326	10,539	9,688	5,729	1,796	309	5
1999.....	2,041	2,598	3,391	4,853	7,285	9,361	10,556	9,430	5,445	1,810	276	4

White, female

2003.....	1,234	1,754	2,179	3,207	5,135	9,079	14,539	18,137	15,893	7,246	1,756	1
2002.....	1,209	1,718	2,160	3,273	5,483	9,391	14,594	18,087	15,764	7,117	1,656	1
2001.....	1,242	1,645	2,151	3,328	5,597	9,380	14,169	18,235	15,480	7,087	1,680	1
2000.....	1,187	1,503	2,136	3,431	5,625	9,608	13,906	17,552	15,367	7,053	1,665	2
1999.....	1,151	1,481	2,158	3,324	5,912	9,670	13,991	17,636	15,012	6,887	1,624	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

Black, both sexes

2003.....	1,320	1,347	1,474	1,679	2,007	2,144	2,265	1,968	1,404	656	216	1
2002.....	1,277	1,298	1,395	1,595	1,908	2,251	2,273	1,940	1,460	653	225	3
2001.....	1,224	1,297	1,327	1,664	1,891	2,320	2,222	1,994	1,354	607	206	1
2000.....	1,159	1,162	1,344	1,643	2,000	2,290	2,092	1,984	1,349	632	210	1
1999.....	1,100	1,270	1,432	1,679	2,101	2,338	2,150	2,054	1,227	615	190	2

Black, male

2003.....	788	809	821	908	968	951	905	638	353	138	42	1
2002.....	743	744	761	837	951	989	884	597	400	122	36	1
2001.....	681	749	745	838	947	1,065	926	663	350	136	40	-
2000.....	707	681	742	864	954	1,017	826	671	355	147	43	-
1999.....	665	749	814	916	1,041	1,030	857	703	351	147	40	1

Black, female

2003.....	532	538	653	771	1,039	1,193	1,360	1,330	1,051	518	174	-
2002.....	534	554	634	758	957	1,262	1,389	1,343	1,060	531	189	2
2001.....	543	548	582	826	944	1,255	1,296	1,331	1,004	471	166	1
2000.....	452	481	602	779	1,046	1,273	1,266	1,313	994	485	167	1
1999.....	435	521	618	763	1,060	1,308	1,293	1,351	876	468	150	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

American Indian 1/, both sexes

2003.....	41	51	71	61	58	61	80	62	44	16	11	-
2002.....	35	39	35	37	60	75	71	69	37	18	10	-
2001.....	24	40	36	45	64	69	71	65	33	18	6	-
2000.....	35	36	38	54	64	73	65	63	35	18	6	-
1999.....	31	32	47	55	60	79	62	51	45	16	11	-

American Indian 1/, male

2003.....	24	33	33	25	33	24	29	29	21	3	4	-
2002.....	25	22	24	21	37	41	30	31	10	5	6	-
2001.....	14	24	21	25	38	40	34	23	12	4	-	-
2000.....	22	23	21	28	42	39	26	25	7	2	-	-
1999.....	18	19	25	35	33	39	23	19	13	5	4	-

American Indian 1/, female

2003.....	17	18	38	36	25	37	51	33	23	13	7	-
2002.....	10	17	11	16	23	34	41	38	27	13	4	-
2001.....	10	16	15	20	26	29	37	42	21	14	6	-
2000.....	13	13	17	26	22	34	39	38	28	16	6	-
1999.....	13	13	22	20	27	40	39	32	32	11	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

Asian or Pacific Islander 2/, both sexes

2003.....	69	85	94	144	194	287	349	324	251	80	28	-
2002.....	69	94	104	131	233	276	312	304	212	82	29	-
2001.....	75	79	88	140	200	250	286	245	179	82	26	-
2000.....	63	80	104	128	173	227	241	232	197	92	16	-
1999.....	63	84	87	123	180	241	264	257	165	73	24	-

Asian or Pacific Islander 2/, male

2003.....	46	50	53	79	103	147	168	136	98	33	8	-
2002.....	42	57	60	76	109	150	144	145	104	34	6	-
2001.....	40	44	48	74	118	127	139	105	92	29	5	-
2000.....	37	44	62	73	99	123	110	104	87	38	7	-
1999.....	32	53	51	66	91	137	125	141	70	26	3	-

Asian or Pacific Islander 2/, female

2003.....	23	35	41	65	91	140	181	188	153	47	20	-
2002.....	27	37	44	55	124	126	168	159	108	48	23	-
2001.....	35	35	40	66	82	123	147	140	87	53	21	-
2000.....	26	36	42	55	74	104	131	128	110	54	9	-
1999.....	31	31	36	57	89	104	139	116	95	47	21	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

All races, both sexes

2003.....	94	95	114	102	130	172	146	104	32	3	1	-
2002.....	90	73	95	107	131	161	145	76	36	8	1	-
2001.....	93	72	88	102	145	162	154	79	30	2	-	-
2000.....	97	88	77	105	119	166	136	83	30	5	1	-
1999.....	75	93	78	103	137	155	134	92	21	5	1	-

All races, male

2003.....	65	60	63	57	62	89	77	43	15	1	-	-
2002.....	62	42	49	66	67	78	75	31	15	2	1	-
2001.....	61	39	56	49	83	94	83	37	11	1	-	-
2000.....	61	52	44	59	63	95	74	40	9	2	-	-
1999.....	54	62	41	51	67	83	71	46	10	-	-	-

All races, female

2003.....	29	35	51	45	68	83	69	61	17	2	1	-
2002.....	28	31	46	41	64	83	70	45	21	6	-	-
2001.....	32	33	32	53	62	68	71	42	19	1	-	-
2000.....	36	36	33	46	56	71	62	43	21	3	1	-
1999.....	21	31	37	52	70	72	63	46	11	5	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

White, both sexes

2003.....	61	66	83	83	104	150	134	98	32	3	1	-
2002.....	60	46	79	82	110	145	138	70	33	8	1	-
2001.....	61	53	64	74	117	134	140	76	27	2	-	-
2000.....	65	59	63	89	102	148	122	74	26	5	-	-
1999.....	48	58	59	82	114	142	122	87	20	5	-	-

White, male

2003.....	41	40	51	47	50	79	72	42	15	1	-	-
2002.....	45	25	43	52	58	74	71	28	14	2	1	-
2001.....	42	28	42	35	63	82	75	35	10	1	-	-
2000.....	44	38	38	47	57	87	66	37	8	2	-	-
1999.....	33	42	31	45	57	81	66	45	10	-	-	-

White, female

2003.....	20	26	32	36	54	71	62	56	17	2	1	-
2002.....	15	21	36	30	52	71	67	42	19	6	-	-
2001.....	19	25	22	39	54	52	65	41	17	1	-	-
2000.....	21	21	25	42	45	61	56	37	18	3	-	-
1999.....	15	16	28	37	57	61	56	42	10	5	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

Black, both sexes

2003.....	32	27	26	17	21	19	10	4	-	-	-	-
2002.....	28	25	16	23	16	14	5	5	3	-	-	-
2001.....	30	19	22	21	22	27	12	2	3	-	-	-
2000.....	27	26	13	15	16	16	10	6	4	-	1	-
1999.....	23	30	18	21	21	10	7	5	1	-	-	-

Black, male

2003.....	23	18	9	9	9	8	5	-	-	-	-	-
2002.....	15	16	6	12	8	3	3	3	1	-	-	-
2001.....	18	11	13	10	16	12	7	1	1	-	-	-
2000.....	15	12	6	11	5	6	5	1	1	-	-	-
1999.....	19	18	9	6	9	1	4	1	-	-	-	-

Black, female

2003.....	9	9	17	8	12	11	5	4	-	-	-	-
2002.....	13	9	10	11	8	11	2	2	2	-	-	-
2001.....	12	8	9	11	6	15	5	1	2	-	-	-
2000.....	12	14	7	4	11	10	5	5	3	-	1	-
1999.....	4	12	9	15	12	9	3	4	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

American Indian 1/, both sexes

2003.....	-	-	2	1	1	1	-	1	-	-	-	-
2002.....	-	1	-	-	-	2	-	-	-	-	-	-
2001.....	1	-	-	2	-	-	-	-	-	-	-	-
2000.....	3	1	1	-	-	1	1	1	-	-	-	-
1999.....	1	1	-	-	-	-	2	-	-	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	2	-	-	-	-	-	-	-	-
2000.....	2	1	-	-	-	1	1	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-

American Indian 1/, female

2003.....	-	-	2	1	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	1	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	1	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

Asian or Pacific Islander 2/, both sexes

2003.....	1	2	3	1	4	2	2	1	-	-	-	-
2002.....	2	1	-	2	5	-	2	1	-	-	-	-
2001.....	1	-	2	5	6	1	2	1	-	-	-	-
2000.....	2	2	-	1	1	1	3	2	-	-	-	-
1999.....	3	4	1	-	2	3	3	-	-	-	1	-

Asian or Pacific Islander 2/, male

2003.....	1	2	3	1	3	1	-	-	-	-	-	-
2002.....	2	1	-	2	1	-	1	-	-	-	-	-
2001.....	1	-	1	2	4	-	1	1	-	-	-	-
2000.....	-	1	-	1	1	1	2	2	-	-	-	-
1999.....	1	2	1	-	1	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	1	1	2	1	-	-	-	-
2002.....	-	-	-	-	4	-	1	1	-	-	-	-
2001.....	-	-	1	3	2	1	1	-	-	-	-	-
2000.....	2	1	-	-	-	-	1	-	-	-	-	-
1999.....	2	2	-	-	1	2	3	-	-	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races, both sexes

2003.....	48	59	45	80	71	70	83	45	27	5	-	-
2002.....	51	66	52	55	88	84	73	65	25	6	-	-
2001.....	54	54	52	47	73	87	67	49	28	7	1	1
2000.....	40	47	42	52	79	82	63	43	24	8	1	-
1999.....	46	63	62	69	77	88	72	56	25	4	-	-

All races, male

2003.....	30	32	25	44	40	34	31	13	8	1	-	-
2002.....	29	34	35	29	42	34	26	25	6	2	-	-
2001.....	29	28	33	28	31	39	22	14	8	2	-	1
2000.....	18	31	21	24	43	42	31	17	7	1	1	-
1999.....	25	40	30	34	35	38	27	19	11	-	-	-

All races, female

2003.....	18	27	20	36	31	36	52	32	19	4	-	-
2002.....	22	32	17	26	46	50	47	40	19	4	-	-
2001.....	25	26	19	19	42	48	45	35	20	5	1	-
2000.....	22	16	21	28	36	40	32	26	17	7	-	-
1999.....	21	23	32	35	42	50	45	37	14	4	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

White, both sexes

2003.....	38	51	39	71	62	63	73	41	26	5	-	-
2002.....	36	49	42	48	75	67	64	62	23	5	-	-
2001.....	38	42	43	36	68	80	61	43	27	7	-	1
2000.....	33	39	33	41	67	69	61	38	21	8	1	-
1999.....	35	53	53	57	65	77	65	50	22	4	-	-

White, male

2003.....	25	26	23	41	36	31	30	13	8	1	-	-
2002.....	21	26	29	25	38	30	22	25	5	1	-	-
2001.....	21	22	26	20	31	35	20	13	8	2	-	1
2000.....	15	28	17	15	37	34	31	15	6	1	1	-
1999.....	18	34	27	27	30	35	25	17	10	-	-	-

White, female

2003.....	13	25	16	30	26	32	43	28	18	4	-	-
2002.....	15	23	13	23	37	37	42	37	18	4	-	-
2001.....	17	20	17	16	37	45	41	30	19	5	-	-
2000.....	18	11	16	26	30	35	30	23	15	7	-	-
1999.....	17	19	26	30	35	42	40	33	12	4	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Black, both sexes

2003.....	7	7	5	7	9	5	8	4	1	-	-	-
2002.....	14	14	8	7	12	14	6	3	2	-	-	-
2001.....	16	11	8	9	4	6	5	5	1	-	1	-
2000.....	7	7	8	10	8	8	2	5	2	-	-	-
1999.....	11	9	8	10	10	10	6	4	2	-	-	-

Black, male

2003.....	4	5	2	1	4	2	1	-	-	-	-	-
2002.....	7	8	5	4	3	4	3	-	1	-	-	-
2001.....	8	5	6	8	-	3	2	-	-	-	-	-
2000.....	3	2	4	8	3	5	-	2	-	-	-	-
1999.....	7	6	3	6	5	2	2	-	-	-	-	-

Black, female

2003.....	3	2	3	6	5	3	7	4	1	-	-	-
2002.....	7	6	3	3	9	10	3	3	1	-	-	-
2001.....	8	6	2	1	4	3	3	5	1	-	1	-
2000.....	4	5	4	2	5	3	2	3	2	-	-	-
1999.....	4	3	5	4	5	8	4	4	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

American Indian 1/, both sexes

2003.....	1	1	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	2	-	-	-	-	-	-
1999.....	-	-	1	-	2	-	-	-	-	-	-	-

American Indian 1/, male

2003.....	1	1	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	2	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Asian or Pacific Islander 2/, both sexes

2003.....	2	-	1	1	-	2	2	-	-	-	-	-
2002.....	-	3	2	-	1	1	3	-	-	1	-	-
2001.....	-	-	1	2	1	1	1	1	-	-	-	-
2000.....	-	1	1	1	3	3	-	-	1	-	-	-
1999.....	-	1	-	2	-	1	1	2	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	1	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	-	-	1	-	-
2001.....	-	-	1	-	-	1	-	1	-	-	-	-
2000.....	-	1	-	1	2	2	-	-	1	-	-	-
1999.....	-	-	-	1	-	1	-	2	1	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	-	1	-	-	1	2	-	-	-	-	-
2002.....	-	3	1	-	-	1	2	-	-	-	-	-
2001.....	-	-	-	2	1	-	1	-	-	-	-	-
2000.....	-	-	1	-	1	1	-	-	-	-	-	-
1999.....	-	1	-	1	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
All races, both sexes												
2003.....	623	1,007	1,462	2,302	3,836	6,697	10,439	12,844	11,082	5,071	1,222	-
2002.....	638	906	1,436	2,192	3,976	6,603	10,307	12,620	10,776	5,031	1,200	-
2001.....	601	926	1,448	2,270	4,004	6,879	10,255	12,793	10,735	4,971	1,209	1
2000.....	528	828	1,383	2,284	4,119	6,936	9,884	12,315	10,543	4,969	1,208	4
1999.....	484	853	1,254	2,240	4,284	6,926	9,845	12,169	10,197	4,850	1,111	1
All races, male												
2003.....	364	605	849	1,281	2,032	3,262	4,396	4,668	3,115	1,111	206	-
2002.....	407	526	819	1,200	2,054	3,192	4,381	4,423	3,000	1,035	178	-
2001.....	361	556	791	1,235	2,109	3,300	4,372	4,320	2,886	1,020	172	-
2000.....	305	479	785	1,243	2,171	3,286	4,212	4,256	2,819	982	191	2
1999.....	300	506	722	1,232	2,294	3,264	4,054	4,155	2,636	991	177	1
All races, female												
2003.....	259	402	613	1,021	1,804	3,435	6,043	8,176	7,967	3,960	1,016	-
2002.....	231	380	617	992	1,922	3,411	5,926	8,197	7,776	3,996	1,022	-
2001.....	240	370	657	1,035	1,895	3,579	5,883	8,473	7,849	3,951	1,037	1
2000.....	223	349	598	1,041	1,948	3,650	5,672	8,059	7,724	3,987	1,017	2
1999.....	184	347	532	1,008	1,990	3,662	5,791	8,014	7,561	3,859	934	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
White, both sexes												
2003.....	406	750	1,109	1,873	3,245	5,926	9,516	11,935	10,361	4,723	1,094	-
2002.....	428	678	1,134	1,789	3,371	5,822	9,377	11,737	10,055	4,722	1,075	-
2001.....	418	693	1,149	1,849	3,428	6,108	9,358	11,903	10,063	4,645	1,092	1
2000.....	378	627	1,088	1,870	3,514	6,178	9,038	11,433	9,847	4,628	1,098	4
1999.....	326	626	1,006	1,827	3,684	6,102	8,979	11,241	9,583	4,490	1,019	-
White, male												
2003.....	239	459	653	1,059	1,761	2,943	4,033	4,358	2,939	1,033	174	-
2002.....	280	402	652	984	1,762	2,852	4,034	4,138	2,800	977	156	-
2001.....	259	429	647	1,021	1,828	2,961	4,001	4,038	2,703	947	151	-
2000.....	224	383	624	1,049	1,892	2,959	3,872	3,961	2,618	905	166	2
1999.....	206	387	585	1,012	2,026	2,918	3,725	3,820	2,459	907	153	-
White, female												
2003.....	167	291	456	814	1,484	2,983	5,483	7,577	7,422	3,690	920	-
2002.....	148	276	482	805	1,609	2,970	5,343	7,599	7,255	3,745	919	-
2001.....	159	264	502	828	1,600	3,147	5,357	7,865	7,360	3,698	941	1
2000.....	154	244	464	821	1,622	3,219	5,166	7,472	7,229	3,723	932	2
1999.....	120	239	421	815	1,658	3,184	5,254	7,421	7,124	3,583	866	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Heart failure (I50)

Black, both sexes

2003.....	204	241	328	392	536	674	802	788	620	308	107	-
2002.....	195	207	283	363	535	690	823	772	623	274	109	-
2001.....	174	223	275	382	528	697	789	787	589	295	100	-
2000.....	143	189	266	373	544	674	742	783	608	292	102	-
1999.....	149	215	236	380	547	731	750	825	515	316	78	1

Black, male

2003.....	115	138	186	207	242	273	308	260	143	63	26	-
2002.....	116	111	158	191	262	298	310	227	156	47	17	-
2001.....	96	121	130	195	256	303	320	246	148	63	20	-
2000.....	78	90	143	175	248	285	288	255	167	60	22	-
1999.....	89	114	129	198	245	309	274	286	141	69	19	1

Black, female

2003.....	89	103	142	185	294	401	494	528	477	245	81	-
2002.....	79	96	125	172	273	392	513	545	467	227	92	-
2001.....	78	102	145	187	272	394	469	541	441	232	80	-
2000.....	65	99	123	198	296	389	454	528	441	232	80	-
1999.....	60	101	107	182	302	422	476	539	374	247	59	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Heart failure (I50)

American Indian 1/, both sexes

2003.....	6	9	15	14	16	24	32	25	20	4	8	-
2002.....	9	9	7	19	20	28	24	27	19	9	5	-
2001.....	5	5	7	13	20	22	29	29	14	9	5	-
2000.....	5	4	10	17	25	29	29	28	14	8	2	-
1999.....	5	2	5	17	17	31	27	18	24	11	6	-

American Indian 1/, male

2003.....	4	5	8	5	9	6	12	12	6	1	2	-
2002.....	7	6	5	12	10	14	8	15	4	2	2	-
2001.....	4	4	4	6	11	13	13	6	4	3	-	-
2000.....	3	1	5	8	15	16	13	12	2	1	-	-
1999.....	2	-	3	12	7	12	9	5	5	3	3	-

American Indian 1/, female

2003.....	2	4	7	9	7	18	20	13	14	3	6	-
2002.....	2	3	2	7	10	14	16	12	15	7	3	-
2001.....	1	1	3	7	9	9	16	23	10	6	5	-
2000.....	2	3	5	9	10	13	16	16	12	7	2	-
1999.....	3	2	2	5	10	19	18	13	19	8	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Heart failure (I50)

Asian or Pacific Islander 2/, both sexes

2003.....	7	7	10	23	39	73	89	96	81	36	13	-
2002.....	6	12	12	21	50	63	83	84	79	26	11	-
2001.....	4	5	17	26	28	52	79	74	69	22	12	-
2000.....	2	8	19	24	36	55	75	71	74	41	6	-
1999.....	4	10	7	16	36	62	89	85	75	33	8	-

Asian or Pacific Islander 2/, male

2003.....	6	3	2	10	20	40	43	38	27	14	4	-
2002.....	4	7	4	13	20	28	29	43	40	9	3	-
2001.....	2	2	10	13	14	23	38	30	31	7	1	-
2000.....	-	5	13	11	16	26	39	28	32	16	3	-
1999.....	3	5	5	10	16	25	46	44	31	12	2	-

Asian or Pacific Islander 2/, female

2003.....	1	4	8	13	19	33	46	58	54	22	9	-
2002.....	2	5	8	8	30	35	54	41	39	17	8	-
2001.....	2	3	7	13	14	29	41	44	38	15	11	-
2000.....	2	3	6	13	20	29	36	43	42	25	3	-
1999.....	1	5	2	6	20	37	43	41	44	21	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races, both sexes

2003.....	4,060	5,010	5,606	6,997	9,726	13,861	17,792	18,022	12,806	4,914	1,092	6
2002.....	4,083	4,868	5,450	7,023	10,108	14,428	17,975	18,033	12,680	4,734	1,009	6
2001.....	4,028	4,585	5,310	7,177	10,238	14,197	17,175	17,422	11,986	4,673	994	5
2000.....	3,886	4,454	5,443	7,434	10,487	14,340	16,760	17,078	12,080	4,609	996	4
1999.....	3,781	4,456	5,721	7,622	11,040	14,520	16,972	17,111	11,651	4,542	1,013	6

All races, male

2003.....	2,560	3,129	3,379	4,020	5,339	6,966	7,825	6,603	3,689	1,056	152	5
2002.....	2,584	2,985	3,281	3,980	5,553	7,159	7,826	6,688	3,537	1,031	159	3
2001.....	2,495	2,770	3,230	4,044	5,588	7,105	7,525	6,224	3,282	1,005	159	4
2000.....	2,489	2,822	3,298	4,258	5,760	7,082	7,184	6,175	3,343	998	167	3
1999.....	2,377	2,811	3,488	4,553	6,054	7,182	7,409	6,073	3,222	997	146	4

All races, female

2003.....	1,500	1,881	2,227	2,977	4,387	6,895	9,967	11,419	9,117	3,858	940	1
2002.....	1,499	1,883	2,169	3,043	4,555	7,269	10,149	11,345	9,143	3,703	850	3
2001.....	1,533	1,815	2,080	3,133	4,650	7,092	9,650	11,198	8,704	3,668	835	1
2000.....	1,397	1,632	2,145	3,176	4,727	7,258	9,576	10,903	8,737	3,611	829	1
1999.....	1,404	1,645	2,233	3,069	4,986	7,338	9,563	11,038	8,429	3,545	867	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, both sexes

2003.....	2,890	3,821	4,357	5,570	8,093	12,169	16,043	16,587	11,829	4,510	965	5
2002.....	2,957	3,709	4,244	5,695	8,546	12,640	16,265	16,612	11,697	4,291	870	3
2001.....	2,936	3,433	4,191	5,788	8,692	12,364	15,513	16,017	11,096	4,292	874	4
2000.....	2,818	3,414	4,275	6,050	8,884	12,539	15,224	15,695	11,202	4,208	875	3
1999.....	2,783	3,342	4,431	6,211	9,334	12,710	15,381	15,688	10,832	4,198	881	5

White, male

2003.....	1,856	2,409	2,682	3,243	4,522	6,176	7,092	6,111	3,393	960	130	4
2002.....	1,926	2,311	2,615	3,280	4,761	6,327	7,123	6,203	3,225	929	133	2
2001.....	1,889	2,097	2,581	3,343	4,786	6,228	6,807	5,718	3,012	909	135	4
2000.....	1,824	2,187	2,644	3,508	4,956	6,246	6,570	5,675	3,097	888	142	3
1999.....	1,784	2,135	2,748	3,769	5,172	6,327	6,740	5,548	2,966	903	123	4

White, female

2003.....	1,034	1,412	1,675	2,327	3,571	5,993	8,951	10,476	8,436	3,550	835	1
2002.....	1,031	1,398	1,629	2,415	3,785	6,313	9,142	10,409	8,472	3,362	737	1
2001.....	1,047	1,336	1,610	2,445	3,906	6,136	8,706	10,299	8,084	3,383	739	-
2000.....	994	1,227	1,631	2,542	3,928	6,293	8,654	10,020	8,105	3,320	733	-
1999.....	999	1,207	1,683	2,442	4,162	6,383	8,641	10,140	7,866	3,295	758	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes

2003.....	1,077	1,072	1,115	1,263	1,441	1,446	1,445	1,172	783	348	109	1
2002.....	1,040	1,052	1,088	1,202	1,345	1,533	1,439	1,160	832	379	116	3
2001.....	1,004	1,044	1,022	1,252	1,337	1,590	1,416	1,200	761	312	105	1
2000.....	982	940	1,057	1,245	1,432	1,592	1,338	1,190	735	340	107	1
1999.....	917	1,016	1,170	1,268	1,523	1,587	1,387	1,220	709	299	112	1

Black, male

2003.....	646	648	624	691	713	668	591	378	210	75	16	1
2002.....	605	609	592	630	678	684	568	367	242	75	19	1
2001.....	559	612	596	625	675	747	597	416	201	73	20	-
2000.....	611	577	589	670	698	721	533	413	187	87	21	-
1999.....	550	611	673	706	782	718	577	416	210	78	21	-

Black, female

2003.....	431	424	491	572	728	778	854	794	573	273	93	-
2002.....	435	443	496	572	667	849	871	793	590	304	97	2
2001.....	445	432	426	627	662	843	819	784	560	239	85	1
2000.....	371	363	468	575	734	871	805	777	548	253	86	1
1999.....	367	405	497	562	741	869	810	804	499	221	91	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

American Indian 1/, both sexes

2003.....	34	41	54	45	41	36	48	36	24	12	3	-
2002.....	25	29	28	18	40	43	47	42	18	9	5	-
2001.....	18	34	29	30	44	47	42	36	19	9	1	-
2000.....	27	31	27	37	38	41	35	34	21	10	4	-
1999.....	25	29	41	38	41	48	33	33	21	5	5	-

American Indian 1/, male

2003.....	19	27	25	19	24	17	17	16	15	2	2	-
2002.....	17	16	19	9	27	26	22	16	6	3	4	-
2001.....	10	19	17	17	27	27	21	17	8	1	-	-
2000.....	17	21	16	20	26	21	12	13	5	1	-	-
1999.....	15	19	22	23	26	27	13	14	8	2	1	-

American Indian 1/, female

2003.....	15	14	29	26	17	19	31	20	9	10	1	-
2002.....	8	13	9	9	13	17	25	26	12	6	1	-
2001.....	8	15	12	13	17	20	21	19	11	8	1	-
2000.....	10	10	11	17	12	20	23	21	16	9	4	-
1999.....	10	10	19	15	15	21	20	19	13	3	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Asian or Pacific Islander 2/, both sexes												
2003.....	59	76	80	119	151	210	256	227	170	44	15	-
2002.....	61	78	90	108	177	212	224	219	133	55	18	-
2001.....	70	74	68	107	165	196	204	169	110	60	14	-
2000.....	59	69	84	102	133	168	163	159	122	51	10	-
1999.....	56	69	79	105	142	175	171	170	89	40	15	-
Asian or Pacific Islander 2/, male												
2003.....	39	45	48	67	80	105	125	98	71	19	4	-
2002.....	36	49	55	61	87	122	113	102	64	24	3	-
2001.....	37	42	36	59	100	103	100	73	61	22	4	-
2000.....	37	37	49	60	80	94	69	74	54	22	4	-
1999.....	28	46	45	55	74	110	79	95	38	14	1	-
Asian or Pacific Islander 2/, female												
2003.....	20	31	32	52	71	105	131	129	99	25	11	-
2002.....	25	29	35	47	90	90	111	117	69	31	15	-
2001.....	33	32	32	48	65	93	104	96	49	38	10	-
2000.....	22	32	35	42	53	74	94	85	68	29	6	-
1999.....	28	23	34	50	68	65	92	75	51	26	14	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races, both sexes

2003.....	631	785	970	1,272	1,832	2,864	3,785	3,949	3,233	1,348	374	-
2002.....	541	681	845	1,126	1,796	2,592	3,546	3,801	2,915	1,276	293	-
2001.....	546	630	832	1,167	1,665	2,638	3,355	3,606	2,573	1,136	257	-
2000.....	526	597	843	1,094	1,690	2,507	3,116	3,273	2,375	1,017	242	-
1999.....	470	538	780	1,146	1,653	2,400	2,924	2,949	2,208	942	218	-

All races, male

2003.....	395	457	532	676	859	1,163	1,359	1,163	773	240	42	-
2002.....	322	398	452	563	856	1,103	1,302	1,182	707	215	28	-
2001.....	330	354	430	603	776	1,100	1,182	1,129	610	192	28	-
2000.....	331	352	466	504	804	1,049	1,155	991	569	155	32	-
1999.....	286	316	416	589	783	996	1,071	878	521	161	20	-

All races, female

2003.....	236	328	438	596	973	1,701	2,426	2,786	2,460	1,108	332	-
2002.....	219	283	393	563	940	1,489	2,244	2,619	2,208	1,061	265	-
2001.....	216	276	402	564	889	1,538	2,173	2,477	1,963	944	229	-
2000.....	195	245	377	590	886	1,458	1,961	2,282	1,806	862	210	-
1999.....	184	222	364	557	870	1,404	1,853	2,071	1,687	781	198	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

White, both sexes

2003.....	358	438	601	817	1,248	2,185	3,124	3,420	2,801	1,174	310	-
2002.....	302	389	536	743	1,284	1,967	2,895	3,203	2,502	1,121	249	-
2001.....	280	356	507	753	1,186	2,049	2,727	3,058	2,243	998	211	-
2000.....	269	345	506	706	1,200	1,925	2,573	2,772	2,056	856	196	-
1999.....	235	304	460	755	1,194	1,819	2,365	2,470	1,904	797	178	-

White, male

2003.....	226	270	330	454	608	910	1,120	998	664	204	33	-
2002.....	188	241	301	379	627	849	1,052	1,014	605	185	22	-
2001.....	172	206	270	395	567	864	956	956	534	161	21	-
2000.....	184	212	291	340	586	823	960	839	483	123	24	-
1999.....	146	193	255	403	579	766	864	722	443	125	16	-

White, female

2003.....	132	168	271	363	640	1,275	2,004	2,422	2,137	970	277	-
2002.....	114	148	235	364	657	1,118	1,843	2,189	1,897	936	227	-
2001.....	108	150	237	358	619	1,185	1,771	2,102	1,709	837	190	-
2000.....	85	133	215	366	614	1,102	1,613	1,933	1,573	733	172	-
1999.....	89	111	205	352	615	1,053	1,501	1,748	1,461	672	162	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Black, both sexes

2003.....	260	324	343	409	521	600	573	449	369	155	59	-
2002.....	228	274	280	344	459	561	577	514	354	132	40	-
2001.....	252	256	302	374	418	517	547	481	288	126	43	-
2000.....	238	242	317	365	441	520	462	440	283	142	41	-
1999.....	226	223	290	344	423	532	480	418	267	125	35	-

Black, male

2003.....	162	173	188	203	227	224	200	136	87	32	8	-
2002.....	128	146	130	163	201	229	215	139	85	26	6	-
2001.....	151	139	149	192	184	205	197	148	61	27	7	-
2000.....	134	137	166	154	191	197	163	127	75	27	5	-
1999.....	135	115	146	166	183	208	178	131	62	30	4	-

Black, female

2003.....	98	151	155	206	294	376	373	313	282	123	51	-
2002.....	100	128	150	181	258	332	362	375	269	106	34	-
2001.....	101	117	153	182	234	312	350	333	227	99	36	-
2000.....	104	105	151	211	250	323	299	313	208	115	36	-
1999.....	91	108	144	178	240	324	302	287	205	95	31	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

American Indian 1/, both sexes

2003.....	5	7	10	8	19	7	8	8	10	1	-	-
2002.....	3	4	7	10	8	8	9	8	3	8	1	-
2001.....	1	2	4	6	10	7	9	7	6	2	1	-
2000.....	3	4	10	4	9	9	7	13	4	2	2	-
1999.....	1	1	8	5	5	5	6	10	6	2	-	-

American Indian 1/, male

2003.....	3	6	6	3	8	6	3	1	3	-	-	-
2002.....	3	2	5	2	5	2	4	1	1	-	-	-
2001.....	1	1	2	2	6	2	2	3	2	-	-	-
2000.....	2	1	4	1	2	3	2	3	1	1	1	-
1999.....	1	-	4	3	2	2	1	3	2	2	-	-

American Indian 1/, female

2003.....	2	1	4	5	11	1	5	7	7	1	-	-
2002.....	-	2	2	8	3	6	5	7	2	8	1	-
2001.....	-	1	2	4	4	5	7	4	4	2	1	-
2000.....	1	3	6	3	7	6	5	10	3	1	1	-
1999.....	-	1	4	2	3	3	5	7	4	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Asian or Pacific Islander 2/, both sexes

2003.....	8	16	16	38	44	72	80	72	53	18	5	-
2002.....	8	14	22	29	45	56	65	76	56	15	3	-
2001.....	13	16	19	34	51	65	72	60	36	10	2	-
2000.....	16	6	10	19	40	53	74	48	32	17	3	-
1999.....	8	10	22	42	31	44	73	51	31	18	5	-

Asian or Pacific Islander 2/, male

2003.....	4	8	8	16	16	23	36	28	19	4	1	-
2002.....	3	9	16	19	23	23	31	28	16	4	-	-
2001.....	6	8	9	14	19	29	27	22	13	4	-	-
2000.....	11	2	5	9	25	26	30	22	10	4	2	-
1999.....	4	8	11	17	19	20	28	22	14	4	-	-

Asian or Pacific Islander 2/, female

2003.....	4	8	8	22	28	49	44	44	34	14	4	-
2002.....	5	5	6	10	22	33	34	48	40	11	3	-
2001.....	7	8	10	20	32	36	45	38	23	6	2	-
2000.....	5	4	5	10	15	27	44	26	22	13	1	-
1999.....	4	2	11	25	12	24	45	29	17	14	5	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cerebrovascular diseases (I60-I69)

All races, both sexes

2003.....	3,550	4,392	5,554	7,765	12,943	21,917	30,930	32,139	22,367	8,493	1,580	2
2002.....	3,440	4,293	5,604	8,026	13,966	22,911	31,978	33,197	23,110	8,521	1,584	3
2001.....	3,403	4,093	5,515	8,170	14,428	23,772	32,050	33,465	22,650	8,398	1,553	4
2000.....	3,386	4,145	5,811	8,671	14,978	24,678	32,342	34,030	23,205	8,530	1,611	2
1999.....	3,108	4,152	5,500	8,694	15,398	25,007	32,420	33,841	23,149	8,571	1,519	2

All races, male

2003.....	1,961	2,394	3,084	4,138	6,530	9,993	12,241	10,471	5,634	1,575	179	1
2002.....	1,876	2,373	3,083	4,305	6,902	10,283	12,372	10,884	5,713	1,510	209	1
2001.....	1,851	2,251	3,026	4,345	7,304	10,671	12,630	10,731	5,482	1,506	211	4
2000.....	1,834	2,294	3,202	4,513	7,529	11,192	12,754	10,864	5,588	1,564	196	1
1999.....	1,714	2,256	3,049	4,522	7,698	11,349	12,739	10,818	5,569	1,467	198	1

All races, female

2003.....	1,589	1,998	2,470	3,627	6,413	11,924	18,689	21,668	16,733	6,918	1,401	1
2002.....	1,564	1,920	2,521	3,721	7,064	12,628	19,606	22,313	17,397	7,011	1,375	2
2001.....	1,552	1,842	2,489	3,825	7,124	13,101	19,420	22,734	17,168	6,892	1,342	-
2000.....	1,552	1,851	2,609	4,158	7,449	13,486	19,588	23,166	17,617	6,966	1,415	1
1999.....	1,394	1,896	2,451	4,172	7,700	13,658	19,681	23,023	17,580	7,104	1,321	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cerebrovascular diseases (I60-I69)

White, both sexes

2003.....	2,248	3,045	3,974	5,928	10,537	18,762	27,567	29,266	20,465	7,753	1,355	2
2002.....	2,203	3,052	4,094	6,112	11,460	19,741	28,557	30,288	21,155	7,769	1,388	2
2001.....	2,139	2,907	3,967	6,188	11,859	20,546	28,719	30,444	20,767	7,672	1,342	1
2000.....	2,287	2,930	4,309	6,775	12,449	21,385	28,982	30,998	21,356	7,711	1,397	1
1999.....	2,074	2,939	4,001	6,876	12,701	21,726	29,191	30,860	21,385	7,852	1,351	-

White, male

2003.....	1,257	1,633	2,222	3,213	5,358	8,619	10,969	9,587	5,144	1,406	158	1
2002.....	1,202	1,684	2,253	3,284	5,737	8,921	11,116	9,970	5,202	1,356	164	1
2001.....	1,179	1,594	2,196	3,337	6,056	9,291	11,373	9,749	5,019	1,351	184	1
2000.....	1,244	1,607	2,391	3,540	6,288	9,746	11,464	9,892	5,095	1,367	172	1
1999.....	1,147	1,613	2,230	3,601	6,399	9,885	11,528	9,838	5,084	1,336	177	-

White, female

2003.....	991	1,412	1,752	2,715	5,179	10,143	16,598	19,679	15,321	6,347	1,197	1
2002.....	1,001	1,368	1,841	2,828	5,723	10,820	17,441	20,318	15,953	6,413	1,224	1
2001.....	960	1,313	1,771	2,851	5,803	11,255	17,346	20,695	15,748	6,321	1,158	-
2000.....	1,043	1,323	1,918	3,235	6,161	11,639	17,518	21,106	16,261	6,344	1,225	-
1999.....	927	1,326	1,771	3,275	6,302	11,841	17,663	21,022	16,301	6,516	1,174	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cerebrovascular diseases (I60-I69)

Black, both sexes

2003.....	1,140	1,143	1,338	1,501	1,937	2,546	2,667	2,288	1,538	601	196	-
2002.....	1,075	1,059	1,265	1,597	2,033	2,603	2,723	2,322	1,559	611	168	1
2001.....	1,104	1,002	1,308	1,651	2,063	2,650	2,645	2,412	1,538	609	172	3
2000.....	958	1,043	1,248	1,582	2,107	2,674	2,725	2,492	1,514	688	197	1
1999.....	896	1,028	1,267	1,538	2,275	2,733	2,618	2,453	1,461	623	144	1

Black, male

2003.....	620	646	735	751	936	1,102	986	659	345	113	15	-
2002.....	593	597	701	857	985	1,094	955	679	351	100	38	-
2001.....	590	560	705	860	999	1,129	957	733	320	116	20	3
2000.....	520	594	664	818	1,017	1,153	1,024	730	369	145	21	-
1999.....	498	547	704	787	1,081	1,177	939	743	365	110	18	-

Black, female

2003.....	520	497	603	750	1,001	1,444	1,681	1,629	1,193	488	181	-
2002.....	482	462	564	740	1,048	1,509	1,768	1,643	1,208	511	130	1
2001.....	514	442	603	791	1,064	1,521	1,688	1,679	1,218	493	152	-
2000.....	438	449	584	764	1,090	1,521	1,701	1,762	1,145	543	176	1
1999.....	398	481	563	751	1,194	1,556	1,679	1,710	1,096	513	126	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cerebrovascular diseases (I60-I69)

American Indian 1/, both sexes

2003.....	32	23	40	46	67	75	92	54	45	7	3	-
2002.....	27	34	37	49	67	69	89	59	47	15	4	-
2001.....	23	26	28	67	74	82	78	68	39	17	7	-
2000.....	25	32	47	44	64	86	92	66	47	16	-	-
1999.....	19	31	32	43	62	90	76	72	39	11	5	-

American Indian 1/, male

2003.....	14	13	20	20	34	34	32	19	15	2	-	-
2002.....	14	19	16	20	29	33	28	25	14	4	-	-
2001.....	13	11	12	29	35	25	26	24	13	4	1	-
2000.....	12	17	26	27	38	30	39	24	12	3	-	-
1999.....	11	12	14	23	36	48	27	20	9	3	1	-

American Indian 1/, female

2003.....	18	10	20	26	33	41	60	35	30	5	3	-
2002.....	13	15	21	29	38	36	61	34	33	11	4	-
2001.....	10	15	16	38	39	57	52	44	26	13	6	-
2000.....	13	15	21	17	26	56	53	42	35	13	-	-
1999.....	8	19	18	20	26	42	49	52	30	8	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cerebrovascular diseases (I60-I69)

Asian or Pacific Islander 2/, both sexes

2003.....	130	181	202	290	402	534	604	531	319	132	26	-
2002.....	135	148	208	268	406	498	609	528	349	126	24	-
2001.....	137	158	212	264	432	494	608	541	306	100	32	-
2000.....	116	140	207	270	358	533	543	474	288	115	17	-
1999.....	119	154	200	237	360	458	535	456	264	85	19	1

Asian or Pacific Islander 2/, male

2003.....	70	102	107	154	202	238	254	206	130	54	6	-
2002.....	67	73	113	144	151	235	273	210	146	50	7	-
2001.....	69	86	113	119	214	226	274	225	130	35	6	-
2000.....	58	76	121	128	186	263	227	218	112	49	3	-
1999.....	58	84	101	111	182	239	245	217	111	18	2	1

Asian or Pacific Islander 2/, female

2003.....	60	79	95	136	200	296	350	325	189	78	20	-
2002.....	68	75	95	124	255	263	336	318	203	76	17	-
2001.....	68	72	99	145	218	268	334	316	176	65	26	-
2000.....	58	64	86	142	172	270	316	256	176	66	14	-
1999.....	61	70	99	126	178	219	290	239	153	67	17	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerosis (I70)

All races, both sexes

2003.....	120	171	295	436	775	1,482	2,355	2,946	2,582	1,375	385	1
2002.....	121	207	280	494	806	1,521	2,541	3,090	2,802	1,416	415	1
2001.....	121	198	275	471	864	1,624	2,532	3,214	2,786	1,471	417	1
2000.....	105	170	281	495	928	1,685	2,592	3,188	2,870	1,518	429	-
1999.....	129	172	314	485	991	1,770	2,615	3,362	2,987	1,581	460	-

All races, male

2003.....	77	113	185	245	404	742	992	1,009	668	272	53	1
2002.....	75	127	178	282	446	748	1,055	1,088	755	281	53	1
2001.....	75	118	168	273	470	765	1,026	1,123	710	285	54	1
2000.....	71	108	186	290	510	838	1,097	1,090	738	283	55	-
1999.....	84	114	193	277	518	845	1,082	1,100	773	316	66	-

All races, female

2003.....	43	58	110	191	371	740	1,363	1,937	1,914	1,103	332	-
2002.....	46	80	102	212	360	773	1,486	2,002	2,047	1,135	362	-
2001.....	46	80	107	198	394	859	1,506	2,091	2,076	1,186	363	-
2000.....	34	62	95	205	418	847	1,495	2,098	2,132	1,235	374	-
1999.....	45	58	121	208	473	925	1,533	2,262	2,214	1,265	394	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerosis (I70)

White, both sexes

2003.....	90	139	246	365	666	1,338	2,146	2,755	2,405	1,287	342	1
2002.....	85	170	227	435	697	1,346	2,320	2,915	2,637	1,331	371	1
2001.....	97	166	216	406	760	1,475	2,339	3,034	2,624	1,374	377	1
2000.....	73	140	226	431	815	1,535	2,375	2,984	2,693	1,431	383	-
1999.....	103	140	262	413	887	1,596	2,423	3,139	2,796	1,475	413	-

White, male

2003.....	59	93	156	206	343	663	899	943	625	257	47	1
2002.....	50	106	144	251	389	663	967	1,030	710	266	46	1
2001.....	59	101	136	237	418	692	946	1,067	666	268	44	1
2000.....	50	93	154	252	461	775	1,008	1,015	687	261	48	-
1999.....	70	90	164	238	463	755	1,000	1,022	722	291	56	-

White, female

2003.....	31	46	90	159	323	675	1,247	1,812	1,780	1,030	295	-
2002.....	35	64	83	184	308	683	1,353	1,885	1,927	1,065	325	-
2001.....	38	65	80	169	342	783	1,393	1,967	1,958	1,106	333	-
2000.....	23	47	72	179	354	760	1,367	1,969	2,006	1,170	335	-
1999.....	33	50	98	175	424	841	1,423	2,117	2,074	1,184	357	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerosis (I70)

Black, both sexes

2003.....	28	30	42	62	96	122	179	153	149	78	35	-
2002.....	31	29	48	49	86	156	196	151	137	68	38	-
2001.....	20	28	56	58	90	132	158	151	139	87	32	-
2000.....	28	28	51	51	97	132	191	173	153	76	40	-
1999.....	23	27	50	63	88	154	162	190	162	92	40	-

Black, male

2003.....	17	19	25	33	55	63	80	50	33	12	5	-
2002.....	22	17	32	25	46	76	78	49	36	13	5	-
2001.....	13	14	29	30	45	65	61	45	35	13	7	-
2000.....	17	14	29	33	42	54	79	63	41	19	5	-
1999.....	11	20	27	34	46	78	65	68	39	23	10	-

Black, female

2003.....	11	11	17	29	41	59	99	103	116	66	30	-
2002.....	9	12	16	24	40	80	118	102	101	55	33	-
2001.....	7	14	27	28	45	67	97	106	104	74	25	-
2000.....	11	14	22	18	55	78	112	110	112	57	35	-
1999.....	12	7	23	29	42	76	97	122	123	69	30	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerosis (I70)

American Indian 1/, both sexes

2003.....	2	1	4	2	6	2	4	3	5	1	2	-
2002.....	4	4	3	8	9	7	9	2	4	2	-	-
2001.....	2	3	1	2	6	4	10	8	6	1	1	-
2000.....	-	2	1	8	8	3	10	8	3	4	2	-
1999.....	2	-	2	5	6	4	8	5	5	3	3	-

American Indian 1/, male

2003.....	1	-	2	2	3	1	3	2	-	-	1	-
2002.....	2	2	2	4	3	1	4	1	1	-	-	-
2001.....	2	2	1	2	3	1	5	2	3	-	-	-
2000.....	-	1	1	3	3	1	5	3	-	-	1	-
1999.....	2	-	2	3	3	3	3	1	-	1	-	-

American Indian 1/, female

2003.....	1	1	2	-	3	1	1	1	5	1	1	-
2002.....	2	2	1	4	6	6	5	1	3	2	-	-
2001.....	-	1	-	-	3	3	5	6	3	1	1	-
2000.....	-	1	-	5	5	2	5	5	3	4	1	-
1999.....	-	-	-	2	3	1	5	4	5	2	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerosis (I70)

Asian or Pacific Islander 2/, both sexes

2003.....	-	1	3	7	7	20	26	35	23	9	6	-
2002.....	1	4	2	2	14	12	16	22	24	15	6	-
2001.....	2	1	2	5	8	13	25	21	17	9	7	-
2000.....	4	-	3	5	8	15	16	23	21	7	4	-
1999.....	1	5	-	4	10	16	22	28	24	11	4	-

Asian or Pacific Islander 2/, male

2003.....	-	1	2	4	3	15	10	14	10	3	-	-
2002.....	1	2	-	2	8	8	6	8	8	2	2	-
2001.....	1	1	2	4	4	7	14	9	6	4	3	-
2000.....	4	-	2	2	4	8	5	9	10	3	1	-
1999.....	1	4	-	2	6	9	14	9	12	1	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	1	3	4	5	16	21	13	6	6	-
2002.....	-	2	2	-	6	4	10	14	16	13	4	-
2001.....	1	-	-	1	4	6	11	12	11	5	4	-
2000.....	-	-	1	3	4	7	11	14	11	4	3	-
1999.....	-	1	-	2	4	7	8	19	12	10	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

All races, both sexes

2003.....	675	900	1,370	1,998	2,936	4,215	4,479	3,739	2,229	762	152	-
2002.....	591	928	1,358	2,137	3,173	4,292	4,453	3,768	2,151	743	154	-
2001.....	603	850	1,345	2,127	3,218	4,498	4,534	3,867	2,210	748	140	1
2000.....	614	866	1,370	2,331	3,595	4,662	4,560	3,819	2,218	757	126	1
1999.....	553	877	1,333	2,244	3,564	4,662	4,602	3,792	2,161	765	145	-

All races, male

2003.....	472	603	914	1,331	1,761	2,445	2,287	1,629	762	181	25	-
2002.....	401	624	902	1,408	1,899	2,406	2,284	1,675	715	191	23	-
2001.....	397	566	888	1,357	1,991	2,476	2,363	1,633	719	188	23	1
2000.....	424	607	923	1,524	2,247	2,696	2,323	1,678	734	204	21	1
1999.....	368	597	922	1,472	2,209	2,684	2,331	1,567	711	172	26	-

All races, female

2003.....	203	297	456	667	1,175	1,770	2,192	2,110	1,467	581	127	-
2002.....	190	304	456	729	1,274	1,886	2,169	2,093	1,436	552	131	-
2001.....	206	284	457	770	1,227	2,022	2,171	2,234	1,491	560	117	-
2000.....	190	259	447	807	1,348	1,966	2,237	2,141	1,484	553	105	-
1999.....	185	280	411	772	1,355	1,978	2,271	2,225	1,450	593	119	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

White, both sexes

2003.....	497	740	1,144	1,728	2,608	3,811	4,038	3,412	2,044	674	125	-
2002.....	432	740	1,137	1,845	2,826	3,880	4,066	3,451	1,934	679	120	-
2001.....	449	681	1,112	1,857	2,848	4,021	4,116	3,535	2,008	654	106	1
2000.....	453	693	1,141	2,029	3,238	4,195	4,167	3,493	1,999	663	102	-
1999.....	425	710	1,158	1,970	3,214	4,228	4,202	3,469	1,952	678	119	-

White, male

2003.....	359	509	771	1,167	1,602	2,238	2,079	1,515	709	161	21	-
2002.....	295	498	766	1,253	1,709	2,219	2,115	1,559	645	179	18	-
2001.....	295	469	747	1,211	1,792	2,255	2,174	1,521	656	170	20	1
2000.....	320	498	792	1,369	2,059	2,458	2,167	1,574	664	180	17	-
1999.....	284	503	818	1,329	2,035	2,480	2,157	1,453	656	151	18	-

White, female

2003.....	138	231	373	561	1,006	1,573	1,959	1,897	1,335	513	104	-
2002.....	137	242	371	592	1,117	1,661	1,951	1,892	1,289	500	102	-
2001.....	154	212	365	646	1,056	1,766	1,942	2,014	1,352	484	86	-
2000.....	133	195	349	660	1,179	1,737	2,000	1,919	1,335	483	85	-
1999.....	141	207	340	641	1,179	1,748	2,045	2,016	1,296	527	101	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2003.....	158	134	193	229	271	325	363	266	151	73	24	-
2002.....	140	167	190	242	282	340	319	256	182	60	30	-
2001.....	131	152	201	212	318	399	355	275	179	84	31	-
2000.....	127	151	200	247	299	388	327	277	187	83	21	1
1999.....	108	149	145	236	285	360	334	278	178	81	24	-

Black, male

2003.....	98	80	122	136	122	159	161	84	32	14	3	-
2002.....	94	111	116	123	146	143	131	85	57	9	4	-
2001.....	87	84	117	110	166	170	159	81	49	13	3	-
2000.....	86	92	109	126	155	192	124	82	49	19	4	1
1999.....	70	83	84	123	145	157	135	88	41	20	8	-

Black, female

2003.....	60	54	71	93	149	166	202	182	119	59	21	-
2002.....	46	56	74	119	136	197	188	171	125	51	26	-
2001.....	44	68	84	102	152	229	196	194	130	71	28	-
2000.....	41	59	91	121	144	196	203	195	138	64	17	-
1999.....	38	66	61	113	140	203	199	190	137	61	16	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

American Indian 1/, both sexes

2003.....	5	5	10	11	8	19	9	14	4	4	1	-
2002.....	6	5	11	16	14	12	9	9	5	1	-	-
2001.....	3	2	9	14	7	10	15	11	1	2	-	-
2000.....	5	6	2	20	13	12	8	4	7	3	1	-
1999.....	2	6	6	8	12	11	8	12	3	1	1	-

American Indian 1/, male

2003.....	5	2	5	9	5	9	3	6	2	2	1	-
2002.....	3	2	6	9	12	8	2	4	-	1	-	-
2001.....	2	2	7	7	4	5	4	6	1	-	-	-
2000.....	2	5	-	9	5	6	4	2	1	1	-	-
1999.....	1	4	3	3	6	4	4	6	-	-	-	-

American Indian 1/, female

2003.....	-	3	5	2	3	10	6	8	2	2	-	-
2002.....	3	3	5	7	2	4	7	5	5	-	-	-
2001.....	1	-	2	7	3	5	11	5	-	2	-	-
2000.....	3	1	2	11	8	6	4	2	6	2	1	-
1999.....	1	2	3	5	6	7	4	6	3	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

Asian or Pacific Islander 2/, both sexes

2003.....	15	21	23	30	49	60	69	47	30	11	2	-
2002.....	13	16	20	34	51	60	59	52	30	3	4	-
2001.....	20	15	23	44	45	68	48	46	22	8	3	-
2000.....	29	16	27	35	45	67	58	45	25	8	2	-
1999.....	18	12	24	30	53	63	58	33	28	5	1	-

Asian or Pacific Islander 2/, male

2003.....	10	12	16	19	32	39	44	24	19	4	-	-
2002.....	9	13	14	23	32	36	36	27	13	2	1	-
2001.....	13	11	17	29	29	46	26	25	13	5	-	-
2000.....	16	12	22	20	28	40	28	20	20	4	-	-
1999.....	13	7	17	17	23	43	35	20	14	1	-	-

Asian or Pacific Islander 2/, female

2003.....	5	9	7	11	17	21	25	23	11	7	2	-
2002.....	4	3	6	11	19	24	23	25	17	1	3	-
2001.....	7	4	6	15	16	22	22	21	9	3	3	-
2000.....	13	4	5	15	17	27	30	25	5	4	2	-
1999.....	5	5	7	13	30	20	23	13	14	4	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

All races, both sexes

2003.....	452	590	887	1,317	1,954	2,726	2,745	2,059	965	245	29	-
2002.....	383	598	862	1,404	2,078	2,779	2,739	2,013	914	239	21	-
2001.....	374	532	862	1,430	2,087	2,899	2,852	2,122	1,022	242	26	1
2000.....	394	585	908	1,552	2,460	3,043	2,792	2,134	985	213	25	-
1999.....	355	570	867	1,578	2,423	3,095	2,841	2,118	935	257	27	-

All races, male

2003.....	347	448	659	912	1,262	1,667	1,525	1,033	411	74	7	-
2002.....	291	458	634	1,020	1,349	1,688	1,497	1,034	368	74	5	-
2001.....	287	398	633	981	1,401	1,722	1,591	1,000	406	78	7	1
2000.....	301	438	673	1,097	1,633	1,885	1,558	1,063	425	68	6	-
1999.....	270	430	658	1,114	1,598	1,905	1,596	1,008	398	83	6	-

All races, female

2003.....	105	142	228	405	692	1,059	1,220	1,026	554	171	22	-
2002.....	92	140	228	384	729	1,091	1,242	979	546	165	16	-
2001.....	87	134	229	449	686	1,177	1,261	1,122	616	164	19	-
2000.....	93	147	235	455	827	1,158	1,234	1,071	560	145	19	-
1999.....	85	140	209	464	825	1,190	1,245	1,110	537	174	21	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

White, both sexes

2003.....	334	482	761	1,176	1,773	2,521	2,542	1,921	915	219	25	-
2002.....	272	482	749	1,261	1,891	2,580	2,548	1,877	843	219	16	-
2001.....	270	431	736	1,276	1,905	2,639	2,654	1,987	943	221	24	1
2000.....	303	475	782	1,381	2,270	2,795	2,603	1,988	900	188	23	-
1999.....	273	473	777	1,420	2,232	2,871	2,658	1,971	871	233	22	-

White, male

2003.....	266	379	570	827	1,170	1,561	1,421	977	388	67	6	-
2002.....	210	370	556	936	1,238	1,585	1,415	985	345	72	4	-
2001.....	212	333	544	892	1,293	1,588	1,499	950	385	73	7	1
2000.....	237	364	592	1,006	1,527	1,747	1,480	1,015	389	62	6	-
1999.....	208	369	600	1,025	1,500	1,805	1,506	946	376	74	5	-

White, female

2003.....	68	103	191	349	603	960	1,121	944	527	152	19	-
2002.....	62	112	193	325	653	995	1,133	892	498	147	12	-
2001.....	58	98	192	384	612	1,051	1,155	1,037	558	148	17	-
2000.....	66	111	190	375	743	1,048	1,123	973	511	126	17	-
1999.....	65	104	177	395	732	1,066	1,152	1,025	495	159	17	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

Black, both sexes

2003.....	102	90	107	112	135	152	142	99	36	22	2	-
2002.....	99	102	95	109	137	145	143	96	49	20	5	-
2001.....	88	88	104	107	137	198	157	98	64	16	2	-
2000.....	69	98	102	129	152	190	143	114	64	20	1	-
1999.....	71	85	70	128	146	166	140	118	44	22	5	-

Black, male

2003.....	70	57	76	66	62	72	67	33	11	5	-	-
2002.....	74	76	66	59	74	68	54	28	15	2	1	-
2001.....	64	55	72	59	79	93	68	28	13	2	-	-
2000.....	52	63	62	68	83	101	54	35	21	4	-	-
1999.....	52	53	44	74	75	63	62	44	12	8	1	-

Black, female

2003.....	32	33	31	46	73	80	75	66	25	17	2	-
2002.....	25	26	29	50	63	77	89	68	34	18	4	-
2001.....	24	33	32	48	58	105	89	70	51	14	2	-
2000.....	17	35	40	61	69	89	89	79	43	16	1	-
1999.....	19	32	26	54	71	103	78	74	32	14	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

American Indian 1/, both sexes

2003.....	2	2	4	4	4	12	4	5	-	1	1	-
2002.....	2	1	5	10	9	6	7	3	2	-	-	-
2001.....	1	2	6	9	3	7	8	6	-	1	-	-
2000.....	-	1	2	13	8	9	3	-	3	1	-	-
1999.....	-	3	2	4	8	8	3	6	1	-	-	-

American Indian 1/, male

2003.....	2	1	3	3	2	6	1	2	-	1	1	-
2002.....	1	-	2	7	8	5	2	2	-	-	-	-
2001.....	1	2	4	4	3	5	3	4	-	-	-	-
2000.....	-	1	-	7	4	5	2	-	1	-	-	-
1999.....	-	2	1	1	5	4	2	3	-	-	-	-

American Indian 1/, female

2003.....	-	1	1	1	2	6	3	3	-	-	-	-
2002.....	1	1	3	3	1	1	5	1	2	-	-	-
2001.....	-	-	2	5	-	2	5	2	-	1	-	-
2000.....	-	-	2	6	4	4	1	-	2	1	-	-
1999.....	-	1	1	3	3	4	1	3	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

Asian or Pacific Islander 2/, both sexes

2003.....	14	16	15	25	42	41	57	34	14	3	1	-
2002.....	10	13	13	24	41	48	41	37	20	-	-	-
2001.....	15	11	16	38	42	55	33	31	15	4	-	-
2000.....	22	11	22	29	30	49	43	32	18	4	1	-
1999.....	11	9	18	26	37	50	40	23	19	2	-	-

Asian or Pacific Islander 2/, male

2003.....	9	11	10	16	28	28	36	21	12	1	-	-
2002.....	6	12	10	18	29	30	26	19	8	-	-	-
2001.....	10	8	13	26	26	36	21	18	8	3	-	-
2000.....	12	10	19	16	19	32	22	13	14	2	-	-
1999.....	10	6	13	14	18	33	26	15	10	1	-	-

Asian or Pacific Islander 2/, female

2003.....	5	5	5	9	14	13	21	13	2	2	1	-
2002.....	4	1	3	6	12	18	15	18	12	-	-	-
2001.....	5	3	3	12	16	19	12	13	7	1	-	-
2000.....	10	1	3	13	11	17	21	19	4	2	1	-
1999.....	1	3	5	12	19	17	14	8	9	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races, both sexes

2003.....	223	310	483	681	982	1,489	1,734	1,680	1,264	517	123	-
2002.....	208	330	496	733	1,095	1,513	1,714	1,755	1,237	504	133	-
2001.....	229	318	483	697	1,131	1,599	1,682	1,745	1,188	506	114	-
2000.....	220	281	462	779	1,135	1,619	1,768	1,685	1,233	544	101	1
1999.....	198	307	466	666	1,141	1,567	1,761	1,674	1,226	508	118	-

All races, male

2003.....	125	155	255	419	499	778	762	596	351	107	18	-
2002.....	110	166	268	388	550	718	787	641	347	117	18	-
2001.....	110	168	255	376	590	754	772	633	313	110	16	-
2000.....	123	169	250	427	614	811	765	615	309	136	15	1
1999.....	98	167	264	358	611	779	735	559	313	89	20	-

All races, female

2003.....	98	155	228	262	483	711	972	1,084	913	410	105	-
2002.....	98	164	228	345	545	795	927	1,114	890	387	115	-
2001.....	119	150	228	321	541	845	910	1,112	875	396	98	-
2000.....	97	112	212	352	521	808	1,003	1,070	924	408	86	-
1999.....	100	140	202	308	530	788	1,026	1,115	913	419	98	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, both sexes

2003.....	163	258	383	552	835	1,290	1,496	1,491	1,129	455	100	-
2002.....	160	258	388	584	935	1,300	1,518	1,574	1,091	460	104	-
2001.....	179	250	376	581	943	1,382	1,462	1,548	1,065	433	82	-
2000.....	150	218	359	648	968	1,400	1,564	1,505	1,099	475	79	-
1999.....	152	237	381	550	982	1,357	1,544	1,498	1,081	445	97	-

White, male

2003.....	93	130	201	340	432	677	658	538	321	94	15	-
2002.....	85	128	210	317	471	634	700	574	300	107	14	-
2001.....	83	136	203	319	499	667	675	571	271	97	13	-
2000.....	83	134	200	363	532	711	687	559	275	118	11	-
1999.....	76	134	218	304	535	675	651	507	280	77	13	-

White, female

2003.....	70	128	182	212	403	613	838	953	808	361	85	-
2002.....	75	130	178	267	464	666	818	1,000	791	353	90	-
2001.....	96	114	173	262	444	715	787	977	794	336	69	-
2000.....	67	84	159	285	436	689	877	946	824	357	68	-
1999.....	76	103	163	246	447	682	893	991	801	368	84	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2003.....	56	44	86	117	136	173	221	167	115	51	22	-
2002.....	41	65	95	133	145	195	176	160	133	40	25	-
2001.....	43	64	97	105	181	201	198	177	115	68	29	-
2000.....	58	53	98	118	147	198	184	163	123	63	20	1
1999.....	37	64	75	108	139	194	194	160	134	59	19	-

Black, male

2003.....	28	23	46	70	60	87	94	51	21	9	3	-
2002.....	20	35	50	64	72	75	77	57	42	7	3	-
2001.....	23	29	45	51	87	77	91	53	36	11	3	-
2000.....	34	29	47	58	72	91	70	47	28	15	4	1
1999.....	18	30	40	49	70	94	73	44	29	12	7	-

Black, female

2003.....	28	21	40	47	76	86	127	116	94	42	19	-
2002.....	21	30	45	69	73	120	99	103	91	33	22	-
2001.....	20	35	52	54	94	124	107	124	79	57	26	-
2000.....	24	24	51	60	75	107	114	116	95	48	16	-
1999.....	19	34	35	59	69	100	121	116	105	47	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

American Indian 1/, both sexes

2003.....	3	3	6	7	4	7	5	9	4	3	-	-
2002.....	4	4	6	6	5	6	2	6	3	1	-	-
2001.....	2	-	3	5	4	3	7	5	1	1	-	-
2000.....	5	5	-	7	5	3	5	4	4	2	1	-
1999.....	2	3	4	4	4	3	5	6	2	1	1	-

American Indian 1/, male

2003.....	3	1	2	6	3	3	2	4	2	1	-	-
2002.....	2	2	4	2	4	3	-	2	-	1	-	-
2001.....	1	-	3	3	1	-	1	2	1	-	-	-
2000.....	2	4	-	2	1	1	2	2	-	1	-	-
1999.....	1	2	2	2	1	-	2	3	-	-	-	-

American Indian 1/, female

2003.....	-	2	4	1	1	4	3	5	2	2	-	-
2002.....	2	2	2	4	1	3	2	4	3	-	-	-
2001.....	1	-	-	2	3	3	6	3	-	1	-	-
2000.....	3	1	-	5	4	2	3	2	4	1	1	-
1999.....	1	1	2	2	3	3	3	3	2	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Asian or Pacific Islander 2/, both sexes

2003.....	1	5	8	5	7	19	12	13	16	8	1	-
2002.....	3	3	7	10	10	12	18	15	10	3	4	-
2001.....	5	4	7	6	3	13	15	15	7	4	3	-
2000.....	7	5	5	6	15	18	15	13	7	4	1	-
1999.....	7	3	6	4	16	13	18	10	9	3	1	-

Asian or Pacific Islander 2/, male

2003.....	1	1	6	3	4	11	8	3	7	3	-	-
2002.....	3	1	4	5	3	6	10	8	5	2	1	-
2001.....	3	3	4	3	3	10	5	7	5	2	-	-
2000.....	4	2	3	4	9	8	6	7	6	2	-	-
1999.....	3	1	4	3	5	10	9	5	4	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	4	2	2	3	8	4	10	9	5	1	-
2002.....	-	2	3	5	7	6	8	7	5	1	3	-
2001.....	2	1	3	3	-	3	10	8	2	2	3	-
2000.....	3	3	2	2	6	10	9	6	1	2	1	-
1999.....	4	2	2	1	11	3	9	5	5	3	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

All races, both sexes

2003.....	282	272	297	319	425	518	643	577	382	148	26	-
2002.....	280	302	293	322	375	552	701	576	390	134	23	-
2001.....	232	272	291	366	475	572	611	568	414	140	27	1
2000.....	266	231	285	326	469	573	588	571	343	149	21	-
1999.....	176	208	237	312	380	498	619	509	322	116	25	1

All races, male

2003.....	165	166	154	160	188	214	223	163	82	27	4	-
2002.....	147	168	141	160	163	216	267	173	86	25	1	-
2001.....	135	149	155	174	231	254	234	164	97	19	6	1
2000.....	150	126	138	170	220	245	227	175	72	26	-	-
1999.....	96	106	118	163	187	225	238	155	66	19	8	-

All races, female

2003.....	117	106	143	159	237	304	420	414	300	121	22	-
2002.....	133	134	152	162	212	336	434	403	304	109	22	-
2001.....	97	123	136	192	244	318	377	404	317	121	21	-
2000.....	116	105	147	156	249	328	361	396	271	123	21	-
1999.....	80	102	119	149	193	273	381	354	256	97	17	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2003.....	221	212	230	255	350	429	564	514	339	130	21	-
2002.....	214	238	219	260	315	472	609	515	354	120	21	-
2001.....	177	213	219	294	398	480	551	519	369	127	23	-
2000.....	203	175	222	242	395	488	529	516	309	132	21	-
1999.....	131	177	190	243	325	429	549	451	290	107	24	1

White, male

2003.....	127	139	113	126	156	178	189	143	73	23	3	-
2002.....	115	130	109	127	137	186	226	154	74	21	1	-
2001.....	102	118	113	140	193	215	211	149	87	15	4	-
2000.....	117	94	107	131	192	214	205	158	64	22	-	-
1999.....	69	92	100	124	163	191	210	138	59	18	8	-

White, female

2003.....	94	73	117	129	194	251	375	371	266	107	18	-
2002.....	99	108	110	133	178	286	383	361	280	99	20	-
2001.....	75	95	106	154	205	265	340	370	282	112	19	-
2000.....	86	81	115	111	203	274	324	358	245	110	21	-
1999.....	62	85	90	119	162	238	339	313	231	89	16	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Black, both sexes

2003.....	55	57	62	59	70	81	70	53	41	16	5	-
2002.....	61	62	68	60	54	75	85	54	33	12	2	-
2001.....	48	55	68	64	67	81	55	42	40	13	3	1
2000.....	60	53	56	79	68	82	50	51	30	15	-	-
1999.....	41	27	42	62	47	64	61	51	29	8	1	-

Black, male

2003.....	34	26	38	31	31	34	28	18	8	4	1	-
2002.....	28	36	30	31	23	27	37	15	11	4	-	-
2001.....	29	30	42	31	31	35	19	12	8	4	2	1
2000.....	32	29	27	36	26	28	16	14	8	4	-	-
1999.....	24	11	17	36	20	31	24	13	5	1	-	-

Black, female

2003.....	21	31	24	28	39	47	42	35	33	12	4	-
2002.....	33	26	38	29	31	48	48	39	22	8	2	-
2001.....	19	25	26	33	36	46	36	30	32	9	1	-
2000.....	28	24	29	43	42	54	34	37	22	11	-	-
1999.....	17	16	25	26	27	33	37	38	24	7	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

American Indian 1/, both sexes

2003.....	2	2	1	1	-	3	1	1	-	1	-	-
2002.....	2	2	3	1	2	-	3	2	1	1	-	-
2001.....	4	2	1	2	4	2	-	1	1	-	-	-
2000.....	1	-	2	3	4	1	3	-	1	-	-	-
1999.....	2	-	2	1	-	1	5	-	-	-	-	-

American Indian 1/, male

2003.....	2	-	1	-	-	1	1	-	-	-	-	-
2002.....	2	2	-	1	2	-	2	1	-	-	-	-
2001.....	3	-	-	1	4	-	-	1	-	-	-	-
2000.....	-	-	1	2	1	1	2	-	-	-	-	-
1999.....	2	-	-	1	-	-	1	-	-	-	-	-

American Indian 1/, female

2003.....	-	2	-	1	-	2	-	1	-	1	-	-
2002.....	-	-	3	-	-	-	1	1	1	1	-	-
2001.....	1	2	1	1	-	2	-	-	1	-	-	-
2000.....	1	-	1	1	3	-	1	-	1	-	-	-
1999.....	-	-	2	-	-	1	4	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Asian or Pacific Islander 2/, both sexes

2003.....	4	1	4	4	5	5	8	9	2	1	-	-
2002.....	3	-	3	1	4	5	4	5	2	1	-	-
2001.....	3	2	3	6	6	9	5	6	4	-	1	-
2000.....	2	3	5	2	2	2	6	4	3	2	-	-
1999.....	2	4	3	6	8	4	4	7	3	1	-	-

Asian or Pacific Islander 2/, male

2003.....	2	1	2	3	1	1	5	2	1	-	-	-
2002.....	2	-	2	1	1	3	2	3	1	-	-	-
2001.....	1	1	-	2	3	4	4	2	2	-	-	-
2000.....	1	3	3	1	1	2	4	3	-	-	-	-
1999.....	1	3	1	2	4	3	3	4	2	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	-	2	1	4	4	3	7	1	1	-	-
2002.....	1	-	1	-	3	2	2	2	1	1	-	-
2001.....	2	1	3	4	3	5	1	4	2	-	1	-
2000.....	1	-	2	1	1	-	2	1	3	2	-	-
1999.....	1	1	2	4	4	1	1	3	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza and pneumonia (J10-J18)

All races, both sexes

2003.....	1,222	1,292	1,838	2,588	4,243	7,698	11,744	13,653	11,200	5,172	1,372	2
2002.....	1,031	1,271	1,716	2,420	4,427	7,905	12,079	13,977	11,490	5,172	1,356	3
2001.....	991	1,155	1,549	2,323	4,327	7,510	11,167	13,382	10,692	4,836	1,281	5
2000.....	991	1,198	1,681	2,533	4,656	7,912	11,909	13,752	11,318	5,116	1,361	3
1999.....	899	1,098	1,527	2,382	4,479	7,776	11,416	13,499	11,151	5,159	1,420	2

All races, male

2003.....	751	739	1,032	1,506	2,376	4,032	5,597	5,592	3,715	1,360	283	1
2002.....	619	742	996	1,384	2,466	4,114	5,781	5,716	3,875	1,352	256	2
2001.....	602	686	901	1,338	2,394	3,995	5,299	5,373	3,587	1,282	260	4
2000.....	602	710	970	1,447	2,566	4,084	5,667	5,575	3,724	1,373	273	2
1999.....	559	611	851	1,394	2,499	4,020	5,438	5,501	3,595	1,295	285	1

All races, female

2003.....	471	553	806	1,082	1,867	3,666	6,147	8,061	7,485	3,812	1,089	1
2002.....	412	529	720	1,036	1,961	3,791	6,298	8,261	7,615	3,820	1,100	1
2001.....	389	469	648	985	1,933	3,515	5,868	8,009	7,105	3,554	1,021	1
2000.....	389	488	711	1,086	2,090	3,828	6,242	8,177	7,594	3,743	1,088	1
1999.....	340	487	676	988	1,980	3,756	5,978	7,998	7,556	3,864	1,135	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza and pneumonia (J10-J18)

White, both sexes

2003.....	912	996	1,456	2,107	3,596	6,784	10,638	12,611	10,319	4,775	1,213	1
2002.....	783	981	1,376	2,018	3,812	6,969	10,929	12,839	10,607	4,770	1,224	3
2001.....	745	906	1,235	1,921	3,723	6,543	10,078	12,309	9,852	4,424	1,114	4
2000.....	736	932	1,326	2,030	3,939	7,028	10,783	12,677	10,480	4,728	1,216	1
1999.....	666	859	1,200	1,994	3,838	6,962	10,443	12,417	10,312	4,771	1,285	2

White, male

2003.....	564	562	785	1,211	2,001	3,535	5,042	5,169	3,387	1,227	241	-
2002.....	470	568	790	1,150	2,141	3,624	5,219	5,245	3,561	1,222	232	2
2001.....	434	536	703	1,090	2,054	3,452	4,773	4,931	3,263	1,160	224	4
2000.....	434	536	766	1,145	2,169	3,623	5,119	5,128	3,415	1,261	241	1
1999.....	404	471	666	1,164	2,124	3,598	4,968	5,038	3,284	1,192	261	1

White, female

2003.....	348	434	671	896	1,595	3,249	5,596	7,442	6,932	3,548	972	1
2002.....	313	413	586	868	1,671	3,345	5,710	7,594	7,046	3,548	992	1
2001.....	311	370	532	831	1,669	3,091	5,305	7,378	6,589	3,264	890	-
2000.....	302	396	560	885	1,770	3,405	5,664	7,549	7,065	3,467	975	-
1999.....	262	388	534	830	1,714	3,364	5,475	7,379	7,028	3,579	1,024	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza and pneumonia (J10-J18)

Black, both sexes

2003.....	272	248	321	397	530	720	810	758	649	280	114	1
2002.....	211	257	303	334	514	718	885	860	656	313	108	-
2001.....	215	220	260	318	488	770	827	795	637	322	139	-
2000.....	228	240	298	416	581	700	869	827	657	315	122	2
1999.....	208	213	285	333	556	665	761	884	679	313	115	-

Black, male

2003.....	165	152	206	250	306	389	386	286	223	76	30	1
2002.....	124	159	182	193	269	377	421	321	210	93	19	-
2001.....	145	133	165	197	282	438	385	292	228	83	27	-
2000.....	148	157	170	249	322	365	408	325	219	86	25	1
1999.....	140	128	161	200	323	337	358	358	227	81	20	-

Black, female

2003.....	107	96	115	147	224	331	424	472	426	204	84	-
2002.....	87	98	121	141	245	341	464	539	446	220	89	-
2001.....	70	87	95	121	206	332	442	503	409	239	112	-
2000.....	80	83	128	167	259	335	461	502	438	229	97	1
1999.....	68	85	124	133	233	328	403	526	452	232	95	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza and pneumonia (J10-J18)

American Indian 1/, both sexes

2003.....	17	21	28	30	26	31	51	51	42	11	13	-
2002.....	13	18	7	19	16	35	46	37	39	17	6	-
2001.....	14	9	16	19	23	37	48	41	31	17	2	-
2000.....	13	6	18	20	25	37	34	36	36	9	3	-
1999.....	12	12	18	15	22	37	35	46	38	18	4	-

American Indian 1/, male

2003.....	9	10	17	15	16	14	26	13	14	8	2	-
2002.....	7	8	5	11	8	19	14	16	14	4	2	-
2001.....	9	4	9	11	12	18	24	19	10	8	1	-
2000.....	11	3	13	15	14	18	15	11	16	4	-	-
1999.....	9	6	7	6	10	24	15	16	14	6	-	-

American Indian 1/, female

2003.....	8	11	11	15	10	17	25	38	28	3	11	-
2002.....	6	10	2	8	8	16	32	21	25	13	4	-
2001.....	5	5	7	8	11	19	24	22	21	9	1	-
2000.....	2	3	5	5	11	19	19	25	20	5	3	-
1999.....	3	6	11	9	12	13	20	30	24	12	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza and pneumonia (J10-J18)

Asian or Pacific Islander 2/, both sexes

2003.....	21	27	33	54	91	163	245	233	190	106	32	-
2002.....	24	15	30	49	85	183	219	241	188	72	18	-
2001.....	17	20	38	65	93	160	214	237	172	73	26	1
2000.....	14	20	39	67	111	147	223	212	145	64	20	-
1999.....	13	14	24	40	63	112	177	152	122	57	16	-

Asian or Pacific Islander 2/, male

2003.....	13	15	24	30	53	94	143	124	91	49	10	-
2002.....	18	7	19	30	48	94	127	134	90	33	3	-
2001.....	14	13	24	40	46	87	117	131	86	31	8	-
2000.....	9	14	21	38	61	78	125	111	74	22	7	-
1999.....	6	6	17	24	42	61	97	89	70	16	4	-

Asian or Pacific Islander 2/, female

2003.....	8	12	9	24	38	69	102	109	99	57	22	-
2002.....	6	8	11	19	37	89	92	107	98	39	15	-
2001.....	3	7	14	25	47	73	97	106	86	42	18	1
2000.....	5	6	18	29	50	69	98	101	71	42	13	-
1999.....	7	8	7	16	21	51	80	63	52	41	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

All races, both sexes

2003.....	32	47	38	72	111	200	257	341	303	128	31	-
2002.....	9	13	17	19	37	59	118	174	144	59	27	-
2001.....	7	8	13	10	11	18	38	37	39	14	6	-
2000.....	24	39	54	76	123	189	303	372	305	130	48	-
1999.....	13	29	42	60	104	154	297	368	303	162	48	-

All races, male

2003.....	18	26	16	38	51	89	100	132	84	26	3	-
2002.....	5	7	13	13	19	28	53	66	45	10	3	-
2001.....	4	4	5	3	5	6	14	15	8	4	2	-
2000.....	14	20	24	43	63	81	115	127	86	35	6	-
1999.....	7	13	24	27	47	71	131	127	70	35	6	-

All races, female

2003.....	14	21	22	34	60	111	157	209	219	102	28	-
2002.....	4	6	4	6	18	31	65	108	99	49	24	-
2001.....	3	4	8	7	6	12	24	22	31	10	4	-
2000.....	10	19	30	33	60	108	188	245	219	95	42	-
1999.....	6	16	18	33	57	83	166	241	233	127	42	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

White, both sexes

2003.....	27	41	35	64	106	191	248	330	292	123	29	-
2002.....	8	11	14	13	34	57	115	172	136	58	26	-
2001.....	7	7	10	8	11	18	34	34	38	13	6	-
2000.....	22	37	50	70	110	180	289	353	295	125	47	-
1999.....	10	26	39	55	93	146	283	356	298	157	46	-

White, male

2003.....	16	22	14	35	49	85	97	129	83	24	3	-
2002.....	5	5	11	9	19	27	51	65	44	10	3	-
2001.....	4	3	3	1	5	6	14	14	7	4	2	-
2000.....	12	18	23	40	59	75	112	119	83	32	6	-
1999.....	6	13	23	24	43	70	124	124	70	34	6	-

White, female

2003.....	11	19	21	29	57	106	151	201	209	99	26	-
2002.....	3	6	3	4	15	30	64	107	92	48	23	-
2001.....	3	4	7	7	6	12	20	20	31	9	4	-
2000.....	10	19	27	30	51	105	177	234	212	93	41	-
1999.....	4	13	16	31	50	76	159	232	228	123	40	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

Black, both sexes

2003.....	5	5	2	6	4	5	8	8	8	3	1	-
2002.....	-	1	3	4	2	1	3	1	5	1	1	-
2001.....	-	1	2	2	-	-	4	3	1	1	-	-
2000.....	2	2	4	5	11	5	8	17	8	4	1	-
1999.....	2	3	2	3	10	5	10	8	4	4	2	-

Black, male

2003.....	2	4	1	3	1	2	2	2	-	1	-	-
2002.....	-	1	2	3	-	-	2	1	1	-	-	-
2001.....	-	1	2	2	-	-	-	1	1	-	-	-
2000.....	2	2	1	2	3	4	1	7	1	3	-	-
1999.....	1	-	1	2	4	1	5	2	-	1	-	-

Black, female

2003.....	3	1	1	3	3	3	6	6	8	2	1	-
2002.....	-	-	1	1	2	1	1	-	4	1	1	-
2001.....	-	-	-	-	-	-	4	2	-	1	-	-
2000.....	-	-	3	3	8	1	7	10	7	1	1	-
1999.....	1	3	1	1	6	4	5	6	4	3	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

American Indian 1/, both sexes

2003.....	-	-	-	-	1	2	1	1	2	1	1	-
2002.....	-	1	-	1	1	1	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	1	1	-	-	-
1999.....	-	-	-	-	1	1	1	-	-	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	1	-	1	-	-	1	-	-
2002.....	-	1	-	1	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

American Indian 1/, female

2003.....	-	-	-	-	-	2	-	1	2	-	1	-
2002.....	-	-	-	-	1	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	1	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)												
Asian or Pacific Islander 2/, both sexes												
2003.....	-	1	1	2	-	2	-	2	1	1	-	-
2002.....	1	-	-	1	-	-	-	1	2	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	2	3	6	1	1	1	-	-
1999.....	1	-	1	2	-	2	3	4	1	1	-	-
Asian or Pacific Islander 2/, male												
2003.....	-	-	1	-	-	2	-	1	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	2	2	1	1	-	-	-
1999.....	-	-	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2003.....	-	1	-	2	-	-	-	1	-	1	-	-
2002.....	1	-	-	1	-	-	-	1	2	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	4	-	-	1	-	-
1999.....	1	-	1	1	-	2	2	3	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

All races, both sexes

2003.....	1,190	1,245	1,800	2,516	4,132	7,498	11,487	13,312	10,897	5,044	1,341	2
2002.....	1,022	1,258	1,699	2,401	4,390	7,846	11,961	13,803	11,346	5,113	1,329	3
2001.....	984	1,147	1,536	2,313	4,316	7,492	11,129	13,345	10,653	4,822	1,275	5
2000.....	967	1,159	1,627	2,457	4,533	7,723	11,606	13,380	11,013	4,986	1,313	3
1999.....	886	1,069	1,485	2,322	4,375	7,622	11,119	13,131	10,848	4,997	1,372	2

All races, male

2003.....	733	713	1,016	1,468	2,325	3,943	5,497	5,460	3,631	1,334	280	1
2002.....	614	735	983	1,371	2,447	4,086	5,728	5,650	3,830	1,342	253	2
2001.....	598	682	896	1,335	2,389	3,989	5,285	5,358	3,579	1,278	258	4
2000.....	588	690	946	1,404	2,503	4,003	5,552	5,448	3,638	1,338	267	2
1999.....	552	598	827	1,367	2,452	3,949	5,307	5,374	3,525	1,260	279	1

All races, female

2003.....	457	532	784	1,048	1,807	3,555	5,990	7,852	7,266	3,710	1,061	1
2002.....	408	523	716	1,030	1,943	3,760	6,233	8,153	7,516	3,771	1,076	1
2001.....	386	465	640	978	1,927	3,503	5,844	7,987	7,074	3,544	1,017	1
2000.....	379	469	681	1,053	2,030	3,720	6,054	7,932	7,375	3,648	1,046	1
1999.....	334	471	658	955	1,923	3,673	5,812	7,757	7,323	3,737	1,093	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

White, both sexes

2003.....	885	955	1,421	2,043	3,490	6,593	10,390	12,281	10,027	4,652	1,184	1
2002.....	775	970	1,362	2,005	3,778	6,912	10,814	12,667	10,471	4,712	1,198	3
2001.....	738	899	1,225	1,913	3,712	6,525	10,044	12,275	9,814	4,411	1,108	4
2000.....	714	895	1,276	1,960	3,829	6,848	10,494	12,324	10,185	4,603	1,169	1
1999.....	656	833	1,161	1,939	3,745	6,816	10,160	12,061	10,014	4,614	1,239	2

White, male

2003.....	548	540	771	1,176	1,952	3,450	4,945	5,040	3,304	1,203	238	-
2002.....	465	563	779	1,141	2,122	3,597	5,168	5,180	3,517	1,212	229	2
2001.....	430	533	700	1,089	2,049	3,446	4,759	4,917	3,256	1,156	222	4
2000.....	422	518	743	1,105	2,110	3,548	5,007	5,009	3,332	1,229	235	1
1999.....	398	458	643	1,140	2,081	3,528	4,844	4,914	3,214	1,158	255	1

White, female

2003.....	337	415	650	867	1,538	3,143	5,445	7,241	6,723	3,449	946	1
2002.....	310	407	583	864	1,656	3,315	5,646	7,487	6,954	3,500	969	1
2001.....	308	366	525	824	1,663	3,079	5,285	7,358	6,558	3,255	886	-
2000.....	292	377	533	855	1,719	3,300	5,487	7,315	6,853	3,374	934	-
1999.....	258	375	518	799	1,664	3,288	5,316	7,147	6,800	3,456	984	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

Black, both sexes

2003.....	267	243	319	391	526	715	802	750	641	277	113	1
2002.....	211	256	300	330	512	717	882	859	651	312	107	-
2001.....	215	219	258	316	488	770	823	792	636	321	139	-
2000.....	226	238	294	411	570	695	861	810	649	311	121	2
1999.....	206	210	283	330	546	660	751	876	675	309	113	-

Black, male

2003.....	163	148	205	247	305	387	384	284	223	75	30	1
2002.....	124	158	180	190	269	377	419	320	209	93	19	-
2001.....	145	132	163	195	282	438	385	291	227	83	27	-
2000.....	146	155	169	247	319	361	407	318	218	83	25	1
1999.....	139	128	160	198	319	336	353	356	227	80	20	-

Black, female

2003.....	104	95	114	144	221	328	418	466	418	202	83	-
2002.....	87	98	120	140	243	340	463	539	442	219	88	-
2001.....	70	87	95	121	206	332	438	501	409	238	112	-
2000.....	80	83	125	164	251	334	454	492	431	228	96	1
1999.....	67	82	123	132	227	324	398	520	448	229	93	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

American Indian 1/, both sexes

2003.....	17	21	28	30	25	29	50	50	40	10	12	-
2002.....	13	17	7	18	15	34	46	37	38	17	6	-
2001.....	14	9	16	19	23	37	48	41	31	17	2	-
2000.....	13	6	18	20	25	36	34	35	35	9	3	-
1999.....	12	12	18	15	21	36	34	46	38	18	4	-

American Indian 1/, male

2003.....	9	10	17	15	15	14	25	13	14	7	2	-
2002.....	7	7	5	10	8	18	14	16	14	4	2	-
2001.....	9	4	9	11	12	18	24	19	10	8	1	-
2000.....	11	3	13	15	14	18	15	11	15	4	-	-
1999.....	9	6	7	6	10	24	14	16	14	6	-	-

American Indian 1/, female

2003.....	8	11	11	15	10	15	25	37	26	3	10	-
2002.....	6	10	2	8	7	16	32	21	24	13	4	-
2001.....	5	5	7	8	11	19	24	22	21	9	1	-
2000.....	2	3	5	5	11	18	19	24	20	5	3	-
1999.....	3	6	11	9	11	12	20	30	24	12	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

Asian or Pacific Islander 2/, both sexes

2003.....	21	26	32	52	91	161	245	231	189	105	32	-
2002.....	23	15	30	48	85	183	219	240	186	72	18	-
2001.....	17	20	37	65	93	160	214	237	172	73	26	1
2000.....	14	20	39	66	109	144	217	211	144	63	20	-
1999.....	12	14	23	38	63	110	174	148	121	56	16	-

Asian or Pacific Islander 2/, male

2003.....	13	15	23	30	53	92	143	123	90	49	10	-
2002.....	18	7	19	30	48	94	127	134	90	33	3	-
2001.....	14	13	24	40	46	87	117	131	86	31	8	-
2000.....	9	14	21	37	60	76	123	110	73	22	7	-
1999.....	6	6	17	23	42	61	96	88	70	16	4	-

Asian or Pacific Islander 2/, female

2003.....	8	11	9	22	38	69	102	108	99	56	22	-
2002.....	5	8	11	18	37	89	92	106	96	39	15	-
2001.....	3	7	13	25	47	73	97	106	86	42	18	1
2000.....	5	6	18	29	49	68	94	101	71	41	13	-
1999.....	6	8	6	15	21	49	78	60	51	40	12	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute lower respiratory infections (J20-J22)

All races, both sexes

2003.....	12	11	10	13	17	35	45	57	60	32	15	-
2002.....	12	11	7	10	23	21	45	70	42	27	13	-
2001.....	10	4	12	10	18	24	34	60	55	25	5	-
2000.....	4	6	4	19	24	41	41	80	77	23	12	-
1999.....	8	8	13	14	21	44	63	86	96	46	21	-

All races, male

2003.....	6	6	6	11	6	17	18	14	15	4	2	-
2002.....	9	3	3	4	9	11	16	21	11	5	2	-
2001.....	6	1	5	3	7	10	10	15	10	4	1	-
2000.....	2	2	1	12	10	21	14	26	15	6	-	-
1999.....	3	4	6	6	11	22	29	19	18	6	5	-

All races, female

2003.....	6	5	4	2	11	18	27	43	45	28	13	-
2002.....	3	8	4	6	14	10	29	49	31	22	11	-
2001.....	4	3	7	7	11	14	24	45	45	21	4	-
2000.....	2	4	3	7	14	20	27	54	62	17	12	-
1999.....	5	4	7	8	10	22	34	67	78	40	16	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute lower respiratory infections (J20-J22)

White, both sexes

2003.....	11	9	9	11	15	34	42	56	59	30	14	-
2002.....	10	9	5	9	19	19	40	64	41	27	12	-
2001.....	6	3	11	9	13	21	33	58	52	23	5	-
2000.....	4	6	4	18	23	38	39	75	72	22	12	-
1999.....	5	6	11	11	16	41	58	81	93	44	19	-

White, male

2003.....	6	5	5	9	6	16	17	14	15	4	2	-
2002.....	8	3	1	3	9	11	14	19	11	5	2	-
2001.....	3	1	4	3	5	8	10	14	9	4	1	-
2000.....	2	2	1	11	9	21	12	24	12	6	-	-
1999.....	3	3	6	4	8	21	27	16	17	6	4	-

White, female

2003.....	5	4	4	2	9	18	25	42	44	26	12	-
2002.....	2	6	4	6	10	8	26	45	30	22	10	-
2001.....	3	2	7	6	8	13	23	44	43	19	4	-
2000.....	2	4	3	7	14	17	27	51	60	16	12	-
1999.....	2	3	5	7	8	20	31	65	76	38	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute lower respiratory infections (J20-J22)

Black, both sexes

2003.....	1	2	-	2	2	1	2	1	1	2	1	-
2002.....	2	1	2	1	4	2	5	4	-	-	1	-
2001.....	4	1	-	-	4	-	-	2	3	2	-	-
2000.....	-	-	-	1	1	3	2	5	4	1	-	-
1999.....	3	1	1	3	5	2	4	4	2	2	2	-

Black, male

2003.....	-	1	-	2	-	1	1	-	-	-	-	-
2002.....	1	-	2	1	-	-	2	2	-	-	-	-
2001.....	3	-	-	-	1	-	-	1	1	-	-	-
2000.....	-	-	-	1	1	-	2	2	2	-	-	-
1999.....	-	1	-	2	3	-	1	2	-	-	1	-

Black, female

2003.....	1	1	-	-	2	-	1	1	1	2	1	-
2002.....	1	1	-	-	4	2	3	2	-	-	1	-
2001.....	1	1	-	-	3	-	-	1	2	2	-	-
2000.....	-	-	-	-	-	3	-	3	2	1	-	-
1999.....	3	-	1	1	2	2	3	2	2	2	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

All races, both sexes

2003.....	9	9	8	11	11	19	26	38	36	19	8	-
2002.....	11	8	5	10	13	15	24	39	31	15	9	-
2001.....	8	3	11	9	14	18	21	38	36	13	4	-
2000.....	1	4	3	15	16	30	23	45	44	15	7	-
1999.....	7	6	12	13	16	28	41	54	55	30	11	-

All races, male

2003.....	5	5	6	9	3	7	10	6	9	3	1	-
2002.....	8	3	2	4	6	8	10	12	7	3	1	-
2001.....	5	1	5	3	5	6	4	9	5	2	1	-
2000.....	1	2	1	9	5	18	9	14	10	5	-	-
1999.....	2	3	5	6	9	15	19	12	12	5	2	-

All races, female

2003.....	4	4	2	2	8	12	16	32	27	16	7	-
2002.....	3	5	3	6	7	7	14	27	24	12	8	-
2001.....	3	2	6	6	9	12	17	29	31	11	3	-
2000.....	-	2	2	6	11	12	14	31	34	10	7	-
1999.....	5	3	7	7	7	13	22	42	43	25	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2003.....	9	7	7	9	9	19	24	37	35	18	8	-
2002.....	9	6	4	9	12	14	22	34	30	15	8	-
2001.....	4	2	10	8	10	15	21	37	33	11	4	-
2000.....	1	4	3	14	15	27	22	42	42	14	7	-
1999.....	4	4	10	10	11	25	38	51	53	29	11	-

White, male

2003.....	5	4	5	7	3	7	9	6	9	3	1	-
2002.....	7	3	1	3	6	8	9	10	7	3	1	-
2001.....	2	1	4	3	3	4	4	9	4	2	1	-
2000.....	1	2	1	8	4	18	8	14	8	5	-	-
1999.....	2	2	5	4	6	14	18	10	11	5	2	-

White, female

2003.....	4	3	2	2	6	12	15	31	26	15	7	-
2002.....	2	3	3	6	6	6	13	24	23	12	7	-
2001.....	2	1	6	5	7	11	17	28	29	9	3	-
2000.....	-	2	2	6	11	9	14	28	34	9	7	-
1999.....	2	2	5	6	5	11	20	41	42	24	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Black, both sexes

2003.....	-	2	-	2	2	-	1	1	1	1	-	-
2002.....	2	1	1	1	1	1	2	4	-	-	1	-
2001.....	4	1	-	-	3	-	-	1	3	2	-	-
2000.....	-	-	-	1	1	3	1	3	2	1	-	-
1999.....	3	1	1	3	5	2	2	3	1	1	-	-

Black, male

2003.....	-	1	-	2	-	-	1	-	-	-	-	-
2002.....	1	-	1	1	-	-	1	2	-	-	-	-
2001.....	3	-	-	-	1	-	-	-	1	-	-	-
2000.....	-	-	-	1	1	-	1	-	2	-	-	-
1999.....	-	1	-	2	3	-	-	2	-	-	-	-

Black, female

2003.....	-	1	-	-	2	-	-	1	1	1	-	-
2002.....	1	1	-	-	1	1	1	2	-	-	1	-
2001.....	1	1	-	-	2	-	-	1	2	2	-	-
2000.....	-	-	-	-	-	3	-	3	-	1	-	-
1999.....	3	-	1	1	2	2	2	1	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Unspecified acute lower respiratory infection (J22)

All races, both sexes

2003.....	3	2	2	2	6	16	19	19	24	13	7	-
2002.....	1	3	2	-	10	6	21	31	11	12	4	-
2001.....	2	1	1	1	4	6	13	22	19	12	1	-
2000.....	3	2	1	4	8	11	18	35	33	8	5	-
1999.....	1	2	1	1	5	16	22	32	41	16	10	-

All races, male

2003.....	1	1	-	2	3	10	8	8	6	1	1	-
2002.....	1	-	1	-	3	3	6	9	4	2	1	-
2001.....	1	-	-	-	2	4	6	6	5	2	-	-
2000.....	1	-	-	3	5	3	5	12	5	1	-	-
1999.....	1	1	1	-	2	7	10	7	6	1	3	-

All races, female

2003.....	2	1	2	-	3	6	11	11	18	12	6	-
2002.....	-	3	1	-	7	3	15	22	7	10	3	-
2001.....	1	1	1	1	2	2	7	16	14	10	1	-
2000.....	2	2	1	1	3	8	13	23	28	7	5	-
1999.....	-	1	-	1	3	9	12	25	35	15	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Unspecified acute lower respiratory infection (J22)

White, both sexes

2003.....	2	2	2	2	6	15	18	19	24	12	6	-
2002.....	1	3	1	-	7	5	18	30	11	12	4	-
2001.....	2	1	1	1	3	6	12	21	19	12	1	-
2000.....	3	2	1	4	8	11	17	33	30	8	5	-
1999.....	1	2	1	1	5	16	20	30	40	15	8	-

White, male

2003.....	1	1	-	2	3	9	8	8	6	1	1	-
2002.....	1	-	-	-	3	3	5	9	4	2	1	-
2001.....	1	-	-	-	2	4	6	5	5	2	-	-
2000.....	1	-	-	3	5	3	4	10	4	1	-	-
1999.....	1	1	1	-	2	7	9	6	6	1	2	-

White, female

2003.....	1	1	2	-	3	6	10	11	18	11	5	-
2002.....	-	3	1	-	4	2	13	21	7	10	3	-
2001.....	1	1	1	1	1	2	6	16	14	10	1	-
2000.....	2	2	1	1	3	8	13	23	26	7	5	-
1999.....	-	1	-	1	3	9	11	24	34	14	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Unspecified acute lower respiratory infection (J22)

Black, both sexes

2003.....	1	-	-	-	-	1	1	-	-	1	1	-
2002.....	-	-	1	-	3	1	3	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	1	2	2	-	-	-
1999.....	-	-	-	-	-	-	2	1	1	1	2	-

Black, male

2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	1	2	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	1	-

Black, female

2003.....	1	-	-	-	-	-	1	-	-	1	1	-
2002.....	-	-	-	-	3	1	2	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	1	1	1	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

All races, both sexes

2003.....	2,261	4,344	7,733	11,805	18,114	24,593	24,693	18,193	8,874	2,516	351	2
2002.....	2,249	4,220	7,060	11,405	18,383	24,495	24,746	17,971	8,508	2,430	375	7
2001.....	2,197	4,176	6,990	11,838	18,913	24,442	23,320	17,542	8,125	2,345	379	1
2000.....	2,077	3,985	6,754	11,833	19,324	24,309	23,413	17,085	7,823	2,238	350	2
1999.....	2,031	4,179	7,118	12,441	20,203	24,978	23,657	16,831	7,510	2,154	338	4

All races, male

2003.....	1,145	2,252	3,991	6,001	9,332	12,266	11,697	8,055	3,547	812	98	2
2002.....	1,178	2,141	3,642	5,838	9,501	12,457	11,975	8,140	3,448	762	86	5
2001.....	1,111	2,159	3,543	6,046	9,831	12,433	11,258	7,806	3,279	733	94	1
2000.....	1,044	2,072	3,492	6,149	10,061	12,305	11,533	7,843	3,165	776	92	2
1999.....	1,004	2,169	3,789	6,642	10,835	13,021	11,684	7,853	3,154	779	108	3

All races, female

2003.....	1,116	2,092	3,742	5,804	8,782	12,327	12,996	10,138	5,327	1,704	253	-
2002.....	1,071	2,079	3,418	5,567	8,882	12,038	12,771	9,831	5,060	1,668	289	2
2001.....	1,086	2,017	3,447	5,792	9,082	12,009	12,062	9,736	4,846	1,612	285	-
2000.....	1,033	1,913	3,262	5,684	9,263	12,004	11,880	9,242	4,658	1,462	258	-
1999.....	1,027	2,010	3,329	5,799	9,368	11,957	11,973	8,978	4,356	1,375	230	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2003.....	1,861	3,777	6,993	10,843	16,763	22,999	23,332	17,255	8,377	2,345	312	2
2002.....	1,842	3,709	6,344	10,447	17,054	22,937	23,321	17,020	8,047	2,258	337	5
2001.....	1,821	3,666	6,301	10,809	17,576	22,961	22,035	16,625	7,617	2,193	337	-
2000.....	1,702	3,469	6,063	10,793	17,972	22,832	22,193	16,186	7,360	2,070	310	1
1999.....	1,697	3,656	6,400	11,383	18,728	23,314	22,384	15,941	7,073	2,022	306	3

White, male

2003.....	946	1,942	3,569	5,471	8,531	11,312	10,880	7,556	3,301	744	86	2
2002.....	964	1,856	3,224	5,292	8,723	11,514	11,167	7,615	3,232	691	73	4
2001.....	911	1,887	3,156	5,465	9,034	11,524	10,518	7,317	3,039	666	76	-
2000.....	849	1,791	3,102	5,523	9,263	11,418	10,836	7,340	2,938	702	78	1
1999.....	827	1,885	3,362	6,017	9,956	12,010	10,934	7,376	2,922	716	97	2

White, female

2003.....	915	1,835	3,424	5,372	8,232	11,687	12,452	9,699	5,076	1,601	226	-
2002.....	878	1,853	3,120	5,155	8,331	11,423	12,154	9,405	4,815	1,567	264	1
2001.....	910	1,779	3,145	5,344	8,542	11,437	11,517	9,308	4,578	1,527	261	-
2000.....	853	1,678	2,961	5,270	8,709	11,414	11,357	8,846	4,422	1,368	232	-
1999.....	870	1,771	3,038	5,366	8,772	11,304	11,450	8,565	4,151	1,306	209	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2003.....	360	494	624	825	1,109	1,245	1,071	681	365	129	35	-
2002.....	366	461	615	811	1,105	1,257	1,141	717	352	140	31	1
2001.....	334	448	583	873	1,112	1,191	1,021	700	377	117	30	-
2000.....	340	445	628	881	1,133	1,191	961	671	360	133	31	-
1999.....	304	473	618	905	1,207	1,358	1,022	701	343	111	25	1

Black, male

2003.....	175	271	351	455	647	716	639	349	171	45	12	-
2002.....	192	256	359	466	641	742	625	379	158	55	10	1
2001.....	174	237	326	500	660	710	584	363	166	51	12	-
2000.....	169	240	355	532	677	691	532	360	173	60	8	-
1999.....	160	255	374	537	718	815	598	365	172	52	10	1

Black, female

2003.....	185	223	273	370	462	529	432	332	194	84	23	-
2002.....	174	205	256	345	464	515	516	338	194	85	21	-
2001.....	160	211	257	373	452	481	437	337	211	66	18	-
2000.....	171	205	273	349	456	500	429	311	187	73	23	-
1999.....	144	218	244	368	489	543	424	336	171	59	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

American Indian 1/, both sexes

2003.....	23	38	54	59	71	99	64	50	16	6	-	-
2002.....	15	22	50	64	71	75	64	47	20	2	1	-
2001.....	10	28	44	54	68	82	47	44	19	3	2	1
2000.....	7	36	35	58	66	66	60	49	15	3	4	-
1999.....	11	18	45	44	79	88	51	36	10	1	2	-

American Indian 1/, male

2003.....	13	20	27	26	41	57	34	25	5	4	-	-
2002.....	7	11	24	30	34	40	37	20	8	-	-	-
2001.....	5	16	18	20	36	42	21	21	8	1	1	1
2000.....	5	19	22	24	26	39	38	24	8	1	3	-
1999.....	5	7	21	25	43	39	24	18	4	-	-	-

American Indian 1/, female

2003.....	10	18	27	33	30	42	30	25	11	2	-	-
2002.....	8	11	26	34	37	35	27	27	12	2	1	-
2001.....	5	12	26	34	32	40	26	23	11	2	1	-
2000.....	2	17	13	34	40	27	22	25	7	2	1	-
1999.....	6	11	24	19	36	49	27	18	6	1	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

Asian or Pacific Islander 2/, both sexes

2003.....	17	35	62	78	171	250	226	207	116	36	4	-
2002.....	26	28	51	83	153	226	220	187	89	30	6	1
2001.....	32	34	62	102	157	208	217	173	112	32	10	-
2000.....	28	35	28	101	153	220	199	179	88	32	5	1
1999.....	19	32	55	109	189	218	200	153	84	20	5	-

Asian or Pacific Islander 2/, male

2003.....	11	19	44	49	113	181	144	125	70	19	-	-
2002.....	15	18	35	50	103	161	146	126	50	16	3	-
2001.....	21	19	43	61	101	157	135	105	66	15	5	-
2000.....	21	22	13	70	95	157	127	119	46	13	3	1
1999.....	12	22	32	63	118	157	128	94	56	11	1	-

Asian or Pacific Islander 2/, female

2003.....	6	16	18	29	58	69	82	82	46	17	4	-
2002.....	11	10	16	33	50	65	74	61	39	14	3	1
2001.....	11	15	19	41	56	51	82	68	46	17	5	-
2000.....	7	13	15	31	58	63	72	60	42	19	2	-
1999.....	7	10	23	46	71	61	72	59	28	9	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

All races, both sexes

2003.....	18	21	26	40	64	107	139	151	129	51	14	-
2002.....	20	22	32	51	84	131	155	162	114	69	27	-
2001.....	21	25	36	56	77	129	132	188	123	67	23	-
2000.....	19	20	47	67	91	156	177	236	161	80	24	-
1999.....	24	35	62	63	110	139	200	193	148	82	23	-

All races, male

2003.....	10	12	14	17	29	59	54	51	35	13	2	-
2002.....	14	10	18	32	42	69	58	62	35	20	5	-
2001.....	9	16	20	35	41	67	48	70	33	14	2	-
2000.....	8	9	23	41	38	84	90	92	46	19	1	-
1999.....	10	24	27	36	58	73	84	71	45	20	6	-

All races, female

2003.....	8	9	12	23	35	48	85	100	94	38	12	-
2002.....	6	12	14	19	42	62	97	100	79	49	22	-
2001.....	12	9	16	21	36	62	84	118	90	53	21	-
2000.....	11	11	24	26	53	72	87	144	115	61	23	-
1999.....	14	11	35	27	52	66	116	122	103	62	17	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

White, both sexes

2003.....	13	14	24	38	56	100	125	139	121	46	13	-
2002.....	17	18	28	46	81	125	142	156	108	66	26	-
2001.....	20	24	33	49	67	118	124	182	116	63	22	-
2000.....	15	17	44	58	82	144	166	229	157	77	23	-
1999.....	15	31	57	58	97	123	183	181	143	79	20	-

White, male

2003.....	8	10	13	17	23	54	48	46	34	13	2	-
2002.....	11	8	15	28	40	65	53	61	32	20	4	-
2001.....	8	16	18	31	36	60	43	67	31	11	2	-
2000.....	7	8	20	37	34	78	87	88	46	18	1	-
1999.....	6	21	24	34	51	65	73	67	43	19	5	-

White, female

2003.....	5	4	11	21	33	46	77	93	87	33	11	-
2002.....	6	10	13	18	41	60	89	95	76	46	22	-
2001.....	12	8	15	18	31	58	81	115	85	52	20	-
2000.....	8	9	24	21	48	66	79	141	111	59	22	-
1999.....	9	10	33	24	46	58	110	114	100	60	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Black, both sexes

2003.....	5	6	2	2	4	3	9	9	5	4	1	-
2002.....	3	3	2	4	2	4	9	6	5	2	1	-
2001.....	1	1	3	7	8	8	3	5	3	3	-	-
2000.....	4	1	3	7	9	10	10	5	3	3	1	-
1999.....	8	4	4	4	11	12	14	8	3	1	2	-

Black, male

2003.....	2	2	1	-	2	3	4	3	-	-	-	-
2002.....	3	1	1	3	1	3	2	1	3	-	1	-
2001.....	1	-	2	4	4	4	2	2	-	3	-	-
2000.....	1	-	3	2	4	4	3	2	-	1	-	-
1999.....	4	3	2	1	6	6	10	1	1	-	1	-

Black, female

2003.....	3	4	1	2	2	-	5	6	5	4	1	-
2002.....	-	2	1	1	1	1	7	5	2	2	-	-
2001.....	-	1	1	3	4	4	1	3	3	-	-	-
2000.....	3	1	-	5	5	6	7	3	3	2	1	-
1999.....	4	1	2	3	5	6	4	7	2	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

American Indian 1/, both sexes

2003.....	-	1	-	-	-	1	1	1	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	1	-	1	-	-	1	-

American Indian 1/, male

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	-	1	-	-	-	1	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	1	-	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	-	-	4	3	4	2	3	1	-	-
2002.....	-	-	2	1	1	2	4	-	1	1	-	-
2001.....	-	-	-	-	2	2	5	1	4	1	1	-
2000.....	-	2	-	2	-	2	1	2	1	-	-	-
1999.....	-	-	1	-	2	3	3	3	2	2	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	-	4	2	2	1	1	-	-	-
2002.....	-	-	2	1	1	1	3	-	-	-	-	-
2001.....	-	-	-	-	1	2	3	1	2	-	-	-
2000.....	-	1	-	2	-	2	-	2	-	-	-	-
1999.....	-	-	1	-	1	2	1	3	1	1	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	-	1	2	1	2	1	-	-
2002.....	-	-	-	-	-	1	1	-	1	1	-	-
2001.....	-	-	-	-	1	-	2	-	2	1	1	-
2000.....	-	1	-	-	-	-	1	-	1	-	-	-
1999.....	-	-	-	-	1	1	2	-	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

All races, both sexes

2003.....	293	630	1,125	1,618	2,445	3,124	2,796	1,740	672	143	19	2
2002.....	298	643	1,060	1,665	2,723	3,202	2,907	1,826	726	157	15	1
2001.....	307	705	1,124	1,892	2,912	3,481	2,856	1,827	727	161	21	-
2000.....	312	680	1,127	1,976	3,144	3,522	2,989	1,832	685	150	18	-
1999.....	332	723	1,296	2,246	3,410	3,768	3,187	1,785	622	146	27	-

All races, male

2003.....	174	339	593	827	1,314	1,608	1,387	792	300	48	5	2
2002.....	172	334	591	875	1,460	1,723	1,505	866	329	52	4	1
2001.....	177	395	571	988	1,566	1,809	1,446	929	326	54	6	-
2000.....	179	366	577	1,077	1,661	1,839	1,534	898	317	58	5	-
1999.....	183	394	691	1,230	1,906	2,027	1,650	886	295	68	9	-

All races, female

2003.....	119	291	532	791	1,131	1,516	1,409	948	372	95	14	-
2002.....	126	309	469	790	1,263	1,479	1,402	960	397	105	11	-
2001.....	130	310	553	904	1,346	1,672	1,410	898	401	107	15	-
2000.....	133	314	550	899	1,483	1,683	1,455	934	368	92	13	-
1999.....	149	329	605	1,016	1,504	1,741	1,537	899	327	78	18	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

White, both sexes

2003.....	258	569	1,026	1,508	2,287	2,934	2,640	1,657	622	129	17	2
2002.....	261	586	974	1,536	2,545	3,031	2,750	1,731	689	153	12	1
2001.....	256	624	1,039	1,771	2,726	3,302	2,732	1,744	681	150	20	-
2000.....	262	609	1,036	1,819	2,964	3,342	2,851	1,731	644	134	16	-
1999.....	286	654	1,197	2,104	3,216	3,547	3,050	1,692	589	135	22	-

White, male

2003.....	148	300	531	765	1,215	1,480	1,294	741	264	42	4	2
2002.....	145	299	529	780	1,354	1,606	1,410	804	313	50	2	1
2001.....	146	344	515	920	1,442	1,698	1,368	874	305	51	6	-
2000.....	146	325	522	966	1,553	1,735	1,453	827	293	47	3	-
1999.....	148	348	627	1,135	1,765	1,895	1,554	836	273	63	7	-

White, female

2003.....	110	269	495	743	1,072	1,454	1,346	916	358	87	13	-
2002.....	116	287	445	756	1,191	1,425	1,340	927	376	103	10	-
2001.....	110	280	524	851	1,284	1,604	1,364	870	376	99	14	-
2000.....	116	284	514	853	1,411	1,607	1,398	904	351	87	13	-
1999.....	138	306	570	969	1,451	1,652	1,496	856	316	72	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

Black, both sexes

2003.....	29	57	85	97	124	148	120	57	37	11	1	-
2002.....	31	49	68	112	156	137	133	69	32	4	3	-
2001.....	46	70	72	107	149	126	97	65	40	9	1	-
2000.....	47	63	84	135	146	138	109	74	31	14	-	-
1999.....	43	60	86	127	154	176	108	76	28	11	3	-

Black, male

2003.....	20	37	53	54	75	95	68	36	27	4	1	-
2002.....	23	30	47	82	92	90	80	42	14	2	2	-
2001.....	27	44	47	62	94	74	61	40	18	2	-	-
2000.....	30	36	52	97	90	74	63	47	20	9	-	-
1999.....	33	38	56	88	109	103	75	38	17	5	2	-

Black, female

2003.....	9	20	32	43	49	53	52	21	10	7	-	-
2002.....	8	19	21	30	64	47	53	27	18	2	1	-
2001.....	19	26	25	45	55	52	36	25	22	7	1	-
2000.....	17	27	32	38	56	64	46	27	11	5	-	-
1999.....	10	22	30	39	45	73	33	38	11	6	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

American Indian 1/, both sexes

2003.....	3	2	6	2	7	11	7	5	2	-	-	-
2002.....	2	5	5	6	8	9	4	6	1	-	-	-
2001.....	-	5	5	7	12	15	4	3	1	-	-	-
2000.....	1	6	5	8	11	13	5	5	1	-	1	-
1999.....	2	4	3	7	9	9	4	4	-	-	-	-

American Indian 1/, male

2003.....	3	1	4	-	4	10	3	2	-	-	-	-
2002.....	-	3	4	4	4	7	4	4	-	-	-	-
2001.....	-	5	2	2	10	6	1	3	-	-	-	-
2000.....	1	3	2	3	3	7	2	4	1	-	1	-
1999.....	1	3	1	2	6	4	2	2	-	-	-	-

American Indian 1/, female

2003.....	-	1	2	2	3	1	4	3	2	-	-	-
2002.....	2	2	1	2	4	2	-	2	1	-	-	-
2001.....	-	-	3	5	2	9	3	-	1	-	-	-
2000.....	-	3	3	5	8	6	3	1	-	-	-	-
1999.....	1	1	2	5	3	5	2	2	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

Asian or Pacific Islander 2/, both sexes

2003.....	3	2	8	11	27	31	29	21	11	3	1	-
2002.....	4	3	13	11	14	25	20	20	4	-	-	-
2001.....	5	6	8	7	25	38	23	15	5	2	-	-
2000.....	2	2	2	14	23	29	24	22	9	2	1	-
1999.....	1	5	10	8	31	36	25	13	5	-	2	-

Asian or Pacific Islander 2/, male

2003.....	3	1	5	8	20	23	22	13	9	2	-	-
2002.....	4	2	11	9	10	20	11	16	2	-	-	-
2001.....	4	2	7	4	20	31	16	12	3	1	-	-
2000.....	2	2	1	11	15	23	16	20	3	2	1	-
1999.....	1	5	7	5	26	25	19	10	5	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	1	3	3	7	8	7	8	2	1	1	-
2002.....	-	1	2	2	4	5	9	4	2	-	-	-
2001.....	1	4	1	3	5	7	7	3	2	1	-	-
2000.....	-	-	1	3	8	6	8	2	6	-	-	-
1999.....	-	-	3	3	5	11	6	3	-	-	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Asthma (J45-J46)

All races, both sexes

2003.....	315	299	263	262	270	364	388	335	236	75	25	-
2002.....	305	288	248	257	326	401	411	345	225	86	19	1
2001.....	324	265	288	288	346	384	418	371	239	77	22	-
2000.....	296	270	270	328	406	420	429	388	227	81	11	-
1999.....	295	343	240	353	420	454	471	381	224	97	18	-

All races, male

2003.....	109	117	97	80	93	116	118	78	57	14	5	-
2002.....	120	108	94	89	109	131	125	97	41	10	4	-
2001.....	116	84	98	104	116	116	109	86	57	11	3	-
2000.....	110	106	94	121	132	120	123	104	39	15	-	-
1999.....	101	119	88	115	152	144	129	81	52	17	2	-

All races, female

2003.....	206	182	166	182	177	248	270	257	179	61	20	-
2002.....	185	180	154	168	217	270	286	248	184	76	15	1
2001.....	208	181	190	184	230	268	309	285	182	66	19	-
2000.....	186	164	176	207	274	300	306	284	188	66	11	-
1999.....	194	224	152	238	268	310	342	300	172	80	16	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Asthma (J45-J46)

White, both sexes

2003.....	180	196	183	194	195	291	328	303	211	71	22	-
2002.....	174	174	177	182	242	340	361	316	204	75	14	-
2001.....	211	172	177	215	262	312	361	325	213	71	18	-
2000.....	182	162	185	231	316	348	378	338	195	69	8	-
1999.....	183	229	161	254	316	382	416	327	194	88	17	-

White, male

2003.....	61	76	60	60	67	83	101	69	52	13	5	-
2002.....	71	63	65	58	79	106	106	87	38	7	2	-
2001.....	69	57	54	76	84	91	89	76	47	9	1	-
2000.....	71	57	62	79	106	102	105	87	30	9	-	-
1999.....	63	81	55	77	106	115	116	61	40	14	2	-

White, female

2003.....	119	120	123	134	128	208	227	234	159	58	17	-
2002.....	103	111	112	124	163	234	255	229	166	68	12	-
2001.....	142	115	123	139	178	221	272	249	166	62	17	-
2000.....	111	105	123	152	210	246	273	251	165	60	8	-
1999.....	120	148	106	177	210	267	300	266	154	74	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Asthma (J45-J46)

Black, both sexes

2003.....	123	82	59	63	57	48	41	18	17	4	2	-
2002.....	119	106	64	57	75	43	34	21	10	7	2	-
2001.....	99	84	88	58	70	59	41	34	13	3	3	-
2000.....	100	89	79	80	73	60	35	32	27	9	3	-
1999.....	102	96	66	76	82	55	39	37	22	5	-	-

Black, male

2003.....	44	27	25	20	18	20	11	3	3	1	-	-
2002.....	44	40	27	25	26	14	10	7	1	2	-	-
2001.....	42	23	29	18	29	18	12	8	3	1	1	-
2000.....	30	38	29	33	21	10	11	9	8	5	-	-
1999.....	34	30	27	26	36	20	7	8	7	1	-	-

Black, female

2003.....	79	55	34	43	39	28	30	15	14	3	2	-
2002.....	75	66	37	32	49	29	24	14	9	5	2	-
2001.....	57	61	59	40	41	41	29	26	10	2	2	-
2000.....	70	51	50	47	52	50	24	23	19	4	3	-
1999.....	68	66	39	50	46	35	32	29	15	4	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
American Indian 1/, both sexes												
2003.....	4	3	5	1	2	4	3	1	1	-	-	-
2002.....	-	1	2	5	2	2	2	-	1	-	-	-
2001.....	2	5	2	1	3	1	1	2	1	1	-	-
2000.....	-	6	1	3	3	1	1	2	-	-	-	-
1999.....	2	5	2	2	4	4	-	1	1	-	-	-
American Indian 1/, male												
2003.....	-	2	3	-	-	2	2	1	-	-	-	-
2002.....	-	-	-	2	1	1	2	-	-	-	-	-
2001.....	-	2	2	1	-	-	-	-	1	-	-	-
2000.....	-	3	-	1	-	1	1	1	-	-	-	-
1999.....	-	2	-	1	2	2	-	1	1	-	-	-
American Indian 1/, female												
2003.....	4	1	2	1	2	2	1	-	1	-	-	-
2002.....	-	1	2	3	1	1	-	-	1	-	-	-
2001.....	2	3	-	-	3	1	1	2	-	1	-	-
2000.....	-	3	1	2	3	-	-	1	-	-	-	-
1999.....	2	3	2	1	2	2	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Asthma (J45-J46)

Asian or Pacific Islander 2/, both sexes

2003.....	8	18	16	4	16	21	16	13	7	-	1	-
2002.....	12	7	5	13	7	16	14	8	10	4	3	1
2001.....	12	4	21	14	11	12	15	10	12	2	1	-
2000.....	14	13	5	14	14	11	15	16	5	3	-	-
1999.....	8	13	11	21	18	13	16	16	7	4	1	-

Asian or Pacific Islander 2/, male

2003.....	4	12	9	-	8	11	4	5	2	-	-	-
2002.....	5	5	2	4	3	10	7	3	2	1	2	-
2001.....	5	2	13	9	3	7	8	2	6	1	1	-
2000.....	9	8	3	8	5	7	6	7	1	1	-	-
1999.....	4	6	6	11	8	7	6	11	4	2	-	-

Asian or Pacific Islander 2/, female

2003.....	4	6	7	4	8	10	12	8	5	-	1	-
2002.....	7	2	3	9	4	6	7	5	8	3	1	1
2001.....	7	2	8	5	8	5	7	8	6	1	-	-
2000.....	5	5	2	6	9	4	9	9	4	2	-	-
1999.....	4	7	5	10	10	6	10	5	3	2	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

All races, both sexes

2003.....	1,635	3,394	6,319	9,885	15,335	20,998	21,370	15,967	7,837	2,247	293	-
2002.....	1,626	3,267	5,720	9,432	15,250	20,761	21,273	15,638	7,443	2,118	314	5
2001.....	1,545	3,181	5,542	9,602	15,578	20,448	19,914	15,156	7,036	2,040	313	1
2000.....	1,450	3,015	5,310	9,462	15,683	20,211	19,818	14,629	6,750	1,927	297	2
1999.....	1,380	3,078	5,520	9,779	16,263	20,617	19,799	14,472	6,516	1,829	270	4

All races, male

2003.....	852	1,784	3,287	5,077	7,896	10,483	10,138	7,134	3,155	737	86	-
2002.....	872	1,689	2,939	4,842	7,890	10,534	10,287	7,115	3,043	680	73	4
2001.....	809	1,664	2,854	4,919	8,108	10,441	9,655	6,721	2,863	654	83	1
2000.....	747	1,591	2,798	4,910	8,230	10,262	9,786	6,749	2,763	684	86	2
1999.....	710	1,632	2,983	5,261	8,719	10,777	9,821	6,815	2,762	674	91	3

All races, female

2003.....	783	1,610	3,032	4,808	7,439	10,515	11,232	8,833	4,682	1,510	207	-
2002.....	754	1,578	2,781	4,590	7,360	10,227	10,986	8,523	4,400	1,438	241	1
2001.....	736	1,517	2,688	4,683	7,470	10,007	10,259	8,435	4,173	1,386	230	-
2000.....	703	1,424	2,512	4,552	7,453	9,949	10,032	7,880	3,987	1,243	211	-
1999.....	670	1,446	2,537	4,518	7,544	9,840	9,978	7,657	3,754	1,155	179	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

White, both sexes

2003.....	1,410	2,998	5,760	9,103	14,225	19,674	20,239	15,156	7,423	2,099	260	-
2002.....	1,390	2,931	5,165	8,683	14,186	19,441	20,068	14,817	7,046	1,964	285	4
2001.....	1,334	2,846	5,052	8,774	14,521	19,229	18,818	14,374	6,607	1,909	277	-
2000.....	1,243	2,681	4,798	8,685	14,610	18,998	18,798	13,888	6,364	1,790	263	1
1999.....	1,213	2,742	4,985	8,967	15,099	19,262	18,735	13,741	6,147	1,720	247	3

White, male

2003.....	729	1,556	2,965	4,629	7,226	9,695	9,437	6,700	2,951	676	75	-
2002.....	737	1,486	2,615	4,426	7,250	9,737	9,598	6,663	2,849	614	65	3
2001.....	688	1,470	2,569	4,438	7,472	9,675	9,018	6,300	2,656	595	67	-
2000.....	625	1,401	2,498	4,441	7,570	9,503	9,191	6,338	2,569	628	74	1
1999.....	610	1,435	2,656	4,771	8,034	9,935	9,191	6,412	2,566	620	83	2

White, female

2003.....	681	1,442	2,795	4,474	6,999	9,979	10,802	8,456	4,472	1,423	185	-
2002.....	653	1,445	2,550	4,257	6,936	9,704	10,470	8,154	4,197	1,350	220	1
2001.....	646	1,376	2,483	4,336	7,049	9,554	9,800	8,074	3,951	1,314	210	-
2000.....	618	1,280	2,300	4,244	7,040	9,495	9,607	7,550	3,795	1,162	189	-
1999.....	603	1,307	2,329	4,196	7,065	9,327	9,544	7,329	3,581	1,100	164	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Black, both sexes

2003.....	203	349	478	663	924	1,046	901	597	306	110	31	-
2002.....	213	303	481	638	872	1,073	965	621	305	127	25	1
2001.....	188	293	420	701	885	998	880	596	321	102	26	-
2000.....	189	292	462	659	905	983	807	560	299	107	27	-
1999.....	151	313	462	698	960	1,115	861	580	290	94	20	1

Black, male

2003.....	109	205	272	381	552	598	556	307	141	40	11	-
2002.....	122	185	284	356	522	635	533	329	140	51	7	1
2001.....	104	170	248	416	533	614	509	313	145	45	11	-
2000.....	108	166	271	400	562	603	455	302	145	45	8	-
1999.....	89	184	289	422	567	686	506	318	147	46	7	1

Black, female

2003.....	94	144	206	282	372	448	345	290	165	70	20	-
2002.....	91	118	197	282	350	438	432	292	165	76	18	-
2001.....	84	123	172	285	352	384	371	283	176	57	15	-
2000.....	81	126	191	259	343	380	352	258	154	62	19	-
1999.....	62	129	173	276	393	429	355	262	143	48	13	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

American Indian 1/, both sexes

2003.....	16	32	43	56	62	83	53	43	13	6	-	-
2002.....	13	15	43	53	61	64	58	41	18	2	1	-
2001.....	8	18	37	46	53	65	42	39	17	2	2	1
2000.....	6	24	29	47	52	52	54	42	14	3	3	-
1999.....	6	9	40	34	66	74	47	30	9	1	1	-

American Indian 1/, male

2003.....	10	17	20	26	37	45	29	21	5	4	-	-
2002.....	7	7	20	24	29	32	31	16	8	-	-	-
2001.....	5	9	14	17	26	35	20	18	7	1	1	1
2000.....	4	13	20	20	23	31	35	19	7	1	2	-
1999.....	4	2	20	21	35	33	22	15	3	-	-	-

American Indian 1/, female

2003.....	6	15	23	30	25	38	24	22	8	2	-	-
2002.....	6	8	23	29	32	32	27	25	10	2	1	-
2001.....	3	9	23	29	27	30	22	21	10	1	1	-
2000.....	2	11	9	27	29	21	19	23	7	2	1	-
1999.....	2	7	20	13	31	41	25	15	6	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Asian or Pacific Islander 2/, both sexes

2003.....	6	15	38	63	124	195	177	171	95	32	2	-
2002.....	10	18	31	58	131	183	182	159	74	25	3	-
2001.....	15	24	33	81	119	156	174	147	91	27	8	-
2000.....	12	18	21	71	116	178	159	139	73	27	4	1
1999.....	10	14	33	80	138	166	156	121	70	14	2	-

Asian or Pacific Islander 2/, male

2003.....	4	6	30	41	81	145	116	106	58	17	-	-
2002.....	6	11	20	36	89	130	125	107	46	15	1	-
2001.....	12	15	23	48	77	117	108	90	55	13	4	-
2000.....	10	11	9	49	75	125	105	90	42	10	2	1
1999.....	7	11	18	47	83	123	102	70	46	8	1	-

Asian or Pacific Islander 2/, female

2003.....	2	9	8	22	43	50	61	65	37	15	2	-
2002.....	4	7	11	22	42	53	57	52	28	10	2	-
2001.....	3	9	10	33	42	39	66	57	36	14	4	-
2000.....	2	7	12	22	41	53	54	49	31	17	2	-
1999.....	3	3	15	33	55	43	54	51	24	6	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

All races, both sexes

2003.....	10	27	43	91	142	220	267	203	84	12	2	-
2002.....	12	26	49	80	143	256	224	217	76	17	2	-
2001.....	8	27	37	91	175	252	264	188	66	16	2	-
2000.....	7	21	36	85	196	249	235	217	80	20	-	-
1999.....	12	19	38	104	170	251	252	180	54	18	1	-

All races, male

2003.....	9	26	41	90	132	213	256	192	80	12	2	-
2002.....	9	24	45	77	137	245	212	207	75	17	1	-
2001.....	7	26	36	90	171	239	255	181	63	16	2	-
2000.....	7	19	35	79	186	241	226	208	79	19	-	-
1999.....	12	19	38	99	170	243	243	172	53	17	1	-

All races, female

2003.....	1	1	2	1	10	7	11	11	4	-	-	-
2002.....	3	2	4	3	6	11	12	10	1	-	1	-
2001.....	1	1	1	1	4	13	9	7	3	-	-	-
2000.....	-	2	1	6	10	8	9	9	1	1	-	-
1999.....	-	-	-	5	-	8	9	8	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

White, both sexes

2003.....	9	25	42	85	136	208	252	198	80	12	2	-
2002.....	11	24	43	77	136	239	212	212	71	14	1	-
2001.....	6	24	31	87	165	244	249	177	64	16	2	-
2000.....	6	21	34	80	186	234	222	210	76	18	-	-
1999.....	8	19	28	95	160	239	245	168	53	15	-	-

White, male

2003.....	8	24	40	84	126	201	242	188	76	12	2	-
2002.....	8	24	40	74	130	229	201	203	70	14	-	-
2001.....	6	23	30	86	161	231	240	170	61	16	2	-
2000.....	6	19	33	74	176	227	214	201	75	17	-	-
1999.....	8	19	28	91	160	231	236	161	52	14	-	-

White, female

2003.....	1	1	2	1	10	7	10	10	4	-	-	-
2002.....	3	-	3	3	6	10	11	9	1	-	1	-
2001.....	-	1	1	1	4	13	9	7	3	-	-	-
2000.....	-	2	1	6	10	7	8	9	1	1	-	-
1999.....	-	-	-	4	-	8	9	7	1	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

Black, both sexes

2003.....	1	1	-	6	5	12	11	5	4	-	-	-
2002.....	1	2	4	3	7	12	11	5	5	3	1	-
2001.....	2	3	5	4	8	8	13	10	2	-	-	-
2000.....	1	-	2	5	10	13	10	7	3	1	-	-
1999.....	4	-	8	8	8	9	7	11	1	3	1	-

Black, male

2003.....	1	1	-	6	5	12	10	4	4	-	-	-
2002.....	1	-	4	3	7	11	10	4	5	3	1	-
2001.....	1	3	5	4	8	8	13	10	2	-	-	-
2000.....	1	-	2	5	10	12	9	7	3	1	-	-
1999.....	4	-	8	7	8	9	7	10	1	3	1	-

Black, female

2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	2	-	-	-	1	1	1	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	1	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

All races, both sexes

2003.....	204	317	456	610	1,144	2,245	3,544	3,908	2,945	1,204	250	-
2002.....	208	271	400	627	1,166	2,323	3,577	4,100	2,995	1,219	229	-
2001.....	236	259	400	589	1,241	2,259	3,500	3,958	2,960	1,160	264	-
2000.....	209	257	341	642	1,257	2,258	3,247	3,825	2,779	1,111	239	-
1999.....	155	259	327	582	1,172	2,068	2,962	3,443	2,586	1,053	212	-

All races, male

2003.....	129	196	275	388	681	1,285	1,896	1,861	1,223	407	67	-
2002.....	136	162	231	380	703	1,332	1,911	1,994	1,267	415	55	-
2001.....	151	160	224	362	744	1,253	1,892	1,893	1,180	379	59	-
2000.....	126	153	201	405	790	1,291	1,747	1,839	1,116	367	65	-
1999.....	82	151	214	367	716	1,252	1,623	1,659	1,006	372	46	-

All races, female

2003.....	75	121	181	222	463	960	1,648	2,047	1,722	797	183	-
2002.....	72	109	169	247	463	991	1,666	2,106	1,728	804	174	-
2001.....	85	99	176	227	497	1,006	1,608	2,065	1,780	781	205	-
2000.....	83	104	140	237	467	967	1,500	1,986	1,663	744	174	-
1999.....	73	108	113	215	456	816	1,339	1,784	1,580	681	166	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

White, both sexes

2003.....	174	268	347	496	982	1,968	3,167	3,590	2,697	1,101	217	-
2002.....	158	228	326	505	1,001	2,042	3,266	3,783	2,774	1,105	206	-
2001.....	184	221	329	490	1,051	2,003	3,169	3,620	2,707	1,054	222	-
2000.....	166	214	271	530	1,050	2,006	2,942	3,532	2,562	1,026	211	-
1999.....	121	204	269	479	982	1,820	2,672	3,175	2,390	947	192	-

White, male

2003.....	112	165	210	323	582	1,129	1,703	1,720	1,127	368	62	-
2002.....	102	133	194	309	602	1,173	1,758	1,855	1,165	379	54	-
2001.....	118	139	182	307	638	1,118	1,731	1,745	1,091	350	53	-
2000.....	99	131	160	329	653	1,139	1,607	1,720	1,028	341	60	-
1999.....	62	118	173	298	597	1,099	1,463	1,532	933	336	44	-

White, female

2003.....	62	103	137	173	400	839	1,464	1,870	1,570	733	155	-
2002.....	56	95	132	196	399	869	1,508	1,928	1,609	726	152	-
2001.....	66	82	147	183	413	885	1,438	1,875	1,616	704	169	-
2000.....	67	83	111	201	397	867	1,335	1,812	1,534	685	151	-
1999.....	59	86	96	181	385	721	1,209	1,643	1,457	611	148	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Black, both sexes

2003.....	27	42	100	105	130	234	300	253	205	78	24	-
2002.....	44	36	63	110	140	244	259	259	186	94	17	-
2001.....	48	34	67	86	166	222	281	274	203	93	34	-
2000.....	41	39	63	102	188	219	253	236	180	74	21	-
1999.....	31	51	54	91	175	227	246	236	176	95	17	-

Black, male

2003.....	15	27	60	59	80	130	146	102	71	27	4	-
2002.....	30	25	32	67	85	136	120	105	82	30	1	-
2001.....	31	20	40	49	92	121	135	118	69	23	5	-
2000.....	27	20	35	67	123	132	114	90	67	21	4	-
1999.....	18	30	38	60	110	137	126	105	58	30	2	-

Black, female

2003.....	12	15	40	46	50	104	154	151	134	51	20	-
2002.....	14	11	31	43	55	108	139	154	104	64	16	-
2001.....	17	14	27	37	74	101	146	156	134	70	29	-
2000.....	14	19	28	35	65	87	139	146	113	53	17	-
1999.....	13	21	16	31	65	90	120	131	118	65	15	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

American Indian 1/, both sexes

2003.....	-	3	4	4	9	7	15	11	7	2	3	-
2002.....	4	2	6	2	6	10	13	11	5	5	1	-
2001.....	3	-	1	4	6	9	12	8	7	1	2	-
2000.....	1	4	3	4	5	6	8	8	5	1	1	-
1999.....	1	3	2	5	4	3	5	12	3	2	-	-

American Indian 1/, male

2003.....	-	1	2	2	3	4	7	5	5	2	1	-
2002.....	2	2	2	-	5	6	7	8	2	3	-	-
2001.....	1	-	1	2	2	2	6	3	1	-	-	-
2000.....	-	2	2	4	3	5	3	4	2	1	-	-
1999.....	1	3	1	4	-	3	2	8	2	-	-	-

American Indian 1/, female

2003.....	-	2	2	2	6	3	8	6	2	-	2	-
2002.....	2	-	4	2	1	4	6	3	3	2	1	-
2001.....	2	-	-	2	4	7	6	5	6	1	2	-
2000.....	1	2	1	-	2	1	5	4	3	-	1	-
1999.....	-	-	1	1	4	-	3	4	1	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Asian or Pacific Islander 2/, both sexes

2003.....	3	4	5	5	23	36	62	54	36	23	6	-
2002.....	2	5	5	10	19	27	39	47	30	15	5	-
2001.....	1	4	3	9	18	25	38	56	43	12	6	-
2000.....	1	-	4	6	14	27	44	49	32	10	6	-
1999.....	2	1	2	7	11	18	39	20	17	9	3	-

Asian or Pacific Islander 2/, male

2003.....	2	3	3	4	16	22	40	34	20	10	-	-
2002.....	2	2	3	4	11	17	26	26	18	3	-	-
2001.....	1	1	1	4	12	12	20	27	19	6	1	-
2000.....	-	-	4	5	11	15	23	25	19	4	1	-
1999.....	1	-	2	5	9	13	32	14	13	6	-	-

Asian or Pacific Islander 2/, female

2003.....	1	1	2	1	7	14	22	20	16	13	6	-
2002.....	-	3	2	6	8	10	13	21	12	12	5	-
2001.....	-	3	2	5	6	13	18	29	24	6	5	-
2000.....	1	-	-	1	3	12	21	24	13	6	5	-
1999.....	1	1	-	2	2	5	7	6	4	3	3	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races, both sexes

2003.....	762	1,037	1,405	2,050	3,106	4,272	4,538	3,721	1,926	723	140	1
2002.....	766	1,055	1,385	1,971	3,266	4,212	4,446	3,445	1,908	639	138	-
2001.....	814	1,030	1,402	2,174	3,336	4,590	4,612	3,627	1,895	643	118	3
2000.....	805	1,044	1,367	2,184	3,264	4,487	4,413	3,394	1,850	629	135	3
1999.....	694	904	1,356	2,143	3,266	4,369	4,277	3,498	1,823	667	142	-

All races, male

2003.....	403	573	794	1,174	1,675	2,226	2,169	1,531	633	187	33	-
2002.....	393	573	766	1,084	1,845	2,166	2,095	1,395	625	136	20	-
2001.....	442	530	792	1,197	1,828	2,330	2,123	1,415	618	143	21	1
2000.....	431	572	752	1,179	1,782	2,353	2,062	1,409	612	156	28	2
1999.....	368	467	746	1,199	1,806	2,255	2,073	1,417	551	178	29	-

All races, female

2003.....	359	464	611	876	1,431	2,046	2,369	2,190	1,293	536	107	1
2002.....	373	482	619	887	1,421	2,046	2,351	2,050	1,283	503	118	-
2001.....	372	500	610	977	1,508	2,260	2,489	2,212	1,277	500	97	2
2000.....	374	472	615	1,005	1,482	2,134	2,351	1,985	1,238	473	107	1
1999.....	326	437	610	944	1,460	2,114	2,204	2,081	1,272	489	113	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2003.....	590	843	1,197	1,800	2,805	3,877	4,260	3,505	1,799	667	123	1
2002.....	596	860	1,156	1,729	2,918	3,878	4,100	3,219	1,785	595	126	-
2001.....	603	845	1,177	1,894	2,997	4,212	4,295	3,389	1,751	573	105	1
2000.....	635	867	1,150	1,888	2,948	4,123	4,085	3,193	1,711	576	120	1
1999.....	548	723	1,160	1,874	2,913	4,000	3,968	3,251	1,702	597	124	-

White, male

2003.....	322	469	682	1,037	1,527	2,022	2,043	1,442	583	170	33	-
2002.....	310	477	639	977	1,657	2,000	1,945	1,316	578	125	18	-
2001.....	340	450	661	1,059	1,672	2,156	1,987	1,327	571	124	19	-
2000.....	341	479	639	1,035	1,615	2,167	1,921	1,328	557	138	21	1
1999.....	296	366	648	1,069	1,637	2,083	1,936	1,318	508	156	22	-

White, female

2003.....	268	374	515	763	1,278	1,855	2,217	2,063	1,216	497	90	1
2002.....	286	383	517	752	1,261	1,878	2,155	1,903	1,207	470	108	-
2001.....	263	395	516	835	1,325	2,056	2,308	2,062	1,180	449	86	1
2000.....	294	388	511	853	1,333	1,956	2,164	1,865	1,154	438	99	-
1999.....	252	357	512	805	1,276	1,917	2,032	1,933	1,194	441	102	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2003.....	157	166	170	193	220	302	197	154	86	50	17	-
2002.....	144	165	193	195	253	236	261	159	99	35	11	-
2001.....	186	155	190	224	268	289	247	171	116	57	13	2
2000.....	146	157	170	254	250	280	252	148	111	42	13	2
1999.....	126	154	165	226	281	296	229	192	89	54	15	-

Black, male

2003.....	72	90	96	104	115	155	88	66	29	14	-	-
2002.....	73	83	107	89	133	116	113	50	33	9	2	-
2001.....	87	67	106	110	117	130	106	65	34	13	2	1
2000.....	74	81	85	125	121	143	98	54	41	15	5	1
1999.....	63	81	84	109	130	140	96	74	30	14	6	-

Black, female

2003.....	85	76	74	89	105	147	109	88	57	36	17	-
2002.....	71	82	86	106	120	120	148	109	66	26	9	-
2001.....	99	88	84	114	151	159	141	106	82	44	11	1
2000.....	72	76	85	129	129	137	154	94	70	27	8	1
1999.....	63	73	81	117	151	156	133	118	59	40	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

American Indian 1/, both sexes

2003.....	8	8	18	15	25	25	18	14	12	-	-	-
2002.....	11	10	12	11	22	16	13	20	8	-	-	-
2001.....	12	10	17	16	21	23	20	17	4	6	-	-
2000.....	11	6	18	15	24	17	20	11	8	1	-	-
1999.....	11	6	8	14	14	24	22	15	10	2	2	-

American Indian 1/, male

2003.....	4	4	9	7	10	14	7	3	6	-	-	-
2002.....	3	6	7	5	10	10	7	11	3	-	-	-
2001.....	6	2	10	7	13	11	11	5	1	2	-	-
2000.....	9	3	11	6	9	10	6	5	3	1	-	-
1999.....	5	6	3	6	9	12	7	7	3	1	1	-

American Indian 1/, female

2003.....	4	4	9	8	15	11	11	11	6	-	-	-
2002.....	8	4	5	6	12	6	6	9	5	-	-	-
2001.....	6	8	7	9	8	12	9	12	3	4	-	-
2000.....	2	3	7	9	15	7	14	6	5	-	-	-
1999.....	6	-	5	8	5	12	15	8	7	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Asian or Pacific Islander 2/, both sexes

2003.....	7	20	20	42	56	68	63	48	29	6	-	-
2002.....	15	20	24	36	73	82	72	47	16	9	1	-
2001.....	13	20	18	40	50	66	50	50	24	7	-	-
2000.....	13	14	29	27	42	67	56	42	20	10	2	-
1999.....	9	21	23	29	58	49	58	40	22	14	1	-

Asian or Pacific Islander 2/, male

2003.....	5	10	7	26	23	35	31	20	15	3	-	-
2002.....	7	7	13	13	45	40	30	18	11	2	-	-
2001.....	9	11	15	21	26	33	19	18	12	4	-	-
2000.....	7	9	17	13	37	33	37	22	11	2	2	-
1999.....	4	14	11	15	30	20	34	18	10	7	-	-

Asian or Pacific Islander 2/, female

2003.....	2	10	13	16	33	33	32	28	14	3	-	-
2002.....	8	13	11	23	28	42	42	29	5	7	1	-
2001.....	4	9	3	19	24	33	31	32	12	3	-	-
2000.....	6	5	12	14	5	34	19	20	9	8	-	-
1999.....	5	7	12	14	28	29	24	22	12	7	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

All races, both sexes

2003.....	159	183	200	228	359	545	734	685	391	138	30	-
2002.....	154	209	202	254	411	592	699	714	407	148	33	-
2001.....	147	183	235	294	424	615	823	788	458	184	40	1
2000.....	177	200	243	299	449	642	773	801	498	177	33	-
1999.....	142	186	236	314	522	665	835	771	470	164	34	2

All races, male

2003.....	110	105	132	141	205	273	333	262	126	25	5	-
2002.....	106	129	118	142	236	287	335	279	116	39	5	-
2001.....	95	123	150	179	218	315	334	277	125	47	5	1
2000.....	131	134	150	172	270	341	326	291	150	49	9	-
1999.....	102	111	154	196	308	331	381	279	138	29	2	1

All races, female

2003.....	49	78	68	87	154	272	401	423	265	113	25	-
2002.....	48	80	84	112	175	305	364	435	291	109	28	-
2001.....	52	60	85	115	206	300	489	511	333	137	35	-
2000.....	46	66	93	127	179	301	447	510	348	128	24	-
1999.....	40	75	82	118	214	334	454	492	332	135	32	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

White, both sexes

2003.....	116	148	164	197	317	475	655	630	365	127	24	-
2002.....	118	164	161	213	363	524	652	661	380	138	28	-
2001.....	112	138	193	245	374	548	755	731	431	171	39	1
2000.....	132	162	208	250	397	569	717	731	468	164	29	-
1999.....	106	152	198	259	457	600	760	713	439	150	28	1

White, male

2003.....	76	83	108	120	179	238	290	240	115	21	3	-
2002.....	77	99	95	119	205	253	311	260	108	36	5	-
2001.....	68	91	122	148	184	277	303	257	115	43	5	1
2000.....	96	110	130	146	234	300	298	263	138	42	8	-
1999.....	76	88	127	160	272	302	348	255	133	24	2	1

White, female

2003.....	40	65	56	77	138	237	365	390	250	106	21	-
2002.....	41	65	66	94	158	271	341	401	272	102	23	-
2001.....	44	47	71	97	190	271	452	474	316	128	34	-
2000.....	36	52	78	104	163	269	419	468	330	122	21	-
1999.....	30	64	71	99	185	298	412	458	306	126	26	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Peptic ulcer (K25-K28)

Black, both sexes

2003.....	39	33	28	20	34	51	57	34	16	4	4	-
2002.....	33	35	33	30	39	51	26	38	14	8	4	-
2001.....	32	35	34	33	41	45	41	38	20	9	1	-
2000.....	38	35	30	37	43	53	39	50	23	8	4	-
1999.....	32	32	34	45	48	43	56	41	24	11	4	1

Black, male

2003.....	31	22	20	15	22	25	32	13	5	1	-	-
2002.....	27	24	21	17	24	26	14	14	3	2	-	-
2001.....	25	26	22	21	27	23	19	15	5	1	-	-
2000.....	28	21	19	18	30	27	19	18	5	3	1	-
1999.....	23	22	25	29	26	19	21	18	3	3	-	-

Black, female

2003.....	8	11	8	5	12	26	25	21	11	3	4	-
2002.....	6	11	12	13	15	25	12	24	11	6	4	-
2001.....	7	9	12	12	14	22	22	23	15	8	1	-
2000.....	10	14	11	19	13	26	20	32	18	5	3	-
1999.....	9	10	9	16	22	24	35	23	21	8	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

American Indian 1/, both sexes

2003.....	2	1	-	2	-	5	-	1	-	1	-	-
2002.....	-	4	3	2	2	2	1	2	1	-	-	-
2001.....	1	4	2	4	3	4	2	-	1	-	-	-
2000.....	3	1	3	4	1	-	3	1	-	-	-	-
1999.....	1	-	-	1	1	3	-	-	-	-	1	-

American Indian 1/, male

2003.....	1	-	-	2	-	1	-	-	-	-	-	-
2002.....	-	3	-	2	1	2	1	1	-	-	-	-
2001.....	1	1	2	2	3	3	-	-	1	-	-	-
2000.....	3	1	1	3	-	-	2	-	-	-	-	-
1999.....	1	-	-	-	1	2	-	-	-	-	-	-

American Indian 1/, female

2003.....	1	1	-	-	-	4	-	1	-	1	-	-
2002.....	-	1	3	-	1	-	-	1	1	-	-	-
2001.....	-	3	-	2	-	1	2	-	-	-	-	-
2000.....	-	-	2	1	1	-	1	1	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

Asian or Pacific Islander 2/, both sexes

2003.....	2	1	8	9	8	14	22	20	10	6	2	-
2002.....	3	6	5	9	7	15	20	13	12	2	1	-
2001.....	2	6	6	12	6	18	25	19	6	4	-	-
2000.....	4	2	2	8	8	20	14	19	7	5	-	-
1999.....	3	2	4	9	16	19	19	17	7	3	1	-

Asian or Pacific Islander 2/, male

2003.....	2	-	4	4	4	9	11	9	6	3	2	-
2002.....	2	3	2	4	6	6	9	4	5	1	-	-
2001.....	1	5	4	8	4	12	12	5	4	3	-	-
2000.....	4	2	-	5	6	14	7	10	7	4	-	-
1999.....	2	1	2	7	9	8	12	6	2	2	-	-

Asian or Pacific Islander 2/, female

2003.....	-	1	4	5	4	5	11	11	4	3	-	-
2002.....	1	3	3	5	1	9	11	9	7	1	1	-
2001.....	1	1	2	4	2	6	13	14	2	1	-	-
2000.....	-	-	2	3	2	6	7	9	-	1	-	-
1999.....	1	1	2	2	7	11	7	11	5	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

All races, both sexes

2003.....	25	18	25	29	35	56	76	42	28	7	1	-
2002.....	30	28	36	32	57	61	59	56	25	10	3	-
2001.....	15	22	23	34	55	37	53	51	30	11	1	-
2000.....	25	28	32	26	45	53	68	39	24	4	1	-
1999.....	24	22	26	30	39	43	53	40	18	3	1	-

All races, male

2003.....	16	13	17	20	26	28	47	21	10	3	-	-
2002.....	20	18	27	24	32	35	32	22	6	1	2	-
2001.....	11	13	17	23	34	22	30	20	12	4	1	-
2000.....	17	20	23	15	25	30	30	15	9	-	-	-
1999.....	19	17	17	18	20	28	19	23	9	2	1	-

All races, female

2003.....	9	5	8	9	9	28	29	21	18	4	1	-
2002.....	10	10	9	8	25	26	27	34	19	9	1	-
2001.....	4	9	6	11	21	15	23	31	18	7	-	-
2000.....	8	8	9	11	20	23	38	24	15	4	1	-
1999.....	5	5	9	12	19	15	34	17	9	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

White, both sexes

2003.....	19	15	17	23	30	46	71	39	25	7	1	-
2002.....	23	25	31	26	45	56	56	52	22	10	2	-
2001.....	14	13	16	28	52	33	44	44	28	10	1	-
2000.....	19	21	29	20	34	42	58	38	22	4	1	-
1999.....	22	15	15	25	32	38	46	37	17	3	1	-

White, male

2003.....	13	11	12	16	22	23	44	20	10	3	-	-
2002.....	16	17	23	19	25	32	30	19	4	1	1	-
2001.....	10	10	12	19	33	20	26	14	11	3	1	-
2000.....	12	14	20	13	18	26	25	14	8	-	-	-
1999.....	17	12	12	15	15	24	15	21	9	2	1	-

White, female

2003.....	6	4	5	7	8	23	27	19	15	4	1	-
2002.....	7	8	8	7	20	24	26	33	18	9	1	-
2001.....	4	3	4	9	19	13	18	30	17	7	-	-
2000.....	7	7	9	7	16	16	33	24	14	4	1	-
1999.....	5	3	3	10	17	14	31	16	8	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

Black, both sexes

2003.....	3	3	7	4	5	9	2	3	2	-	-	-
2002.....	6	3	4	5	9	3	2	4	3	-	1	-
2001.....	1	9	7	6	1	3	8	3	2	1	-	-
2000.....	4	7	3	6	10	9	7	1	2	-	-	-
1999.....	2	7	10	3	7	5	6	2	1	-	-	-

Black, male

2003.....	1	2	4	3	4	4	1	1	-	-	-	-
2002.....	4	1	3	4	5	2	2	3	2	-	1	-
2001.....	1	3	5	4	-	2	3	3	1	1	-	-
2000.....	3	6	3	2	7	3	3	1	1	-	-	-
1999.....	2	5	4	2	5	4	4	2	-	-	-	-

Black, female

2003.....	2	1	3	1	1	5	1	2	2	-	-	-
2002.....	2	2	1	1	4	1	-	1	1	-	-	-
2001.....	-	6	2	2	1	1	5	-	1	-	-	-
2000.....	1	1	-	4	3	6	4	-	1	-	-	-
1999.....	-	2	6	1	2	1	2	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

Asian or Pacific Islander 2/, both sexes

2003.....	3	-	-	2	-	1	3	-	1	-	-	-
2002.....	-	-	1	1	3	2	1	-	-	-	-	-
2001.....	-	-	-	-	2	1	1	4	-	-	-	-
2000.....	2	-	-	-	1	1	3	-	-	-	-	-
1999.....	-	-	-	2	-	-	1	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	2	-	-	1	-	1	2	-	-	-	-	-
2002.....	-	-	1	1	2	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	1	3	-	-	-	-
2000.....	2	-	-	-	-	1	2	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	-	-	1	-	-	1	-	1	-	-	-
2002.....	-	-	-	-	1	1	1	-	-	-	-	-
2001.....	-	-	-	-	1	1	-	1	-	-	-	-
2000.....	-	-	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	1	-	-	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hernia (K40-K46)

All races, both sexes

2003.....	60	76	85	101	136	203	259	296	202	56	9	-
2002.....	44	55	71	90	135	208	281	320	172	85	11	-
2001.....	58	57	74	81	132	185	258	269	190	88	6	-
2000.....	58	51	62	84	126	224	242	294	188	63	10	-
1999.....	39	36	71	77	138	186	265	269	196	56	8	-

All races, male

2003.....	35	30	42	48	63	93	106	95	62	17	4	-
2002.....	19	27	29	40	63	84	105	97	45	20	-	-
2001.....	31	23	41	40	54	71	102	88	48	19	1	-
2000.....	32	20	29	39	53	91	104	96	42	12	4	-
1999.....	20	21	39	34	59	75	117	96	51	8	2	-

All races, female

2003.....	25	46	43	53	73	110	153	201	140	39	5	-
2002.....	25	28	42	50	72	124	176	223	127	65	11	-
2001.....	27	34	33	41	78	114	156	181	142	69	5	-
2000.....	26	31	33	45	73	133	138	198	146	51	6	-
1999.....	19	15	32	43	79	111	148	173	145	48	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
White, both sexes												
2003.....	48	60	71	88	120	185	230	276	184	55	7	-
2002.....	34	47	55	71	121	197	258	296	167	83	10	-
2001.....	47	45	62	69	116	164	235	251	179	77	4	-
2000.....	47	45	54	68	117	198	223	270	176	57	9	-
1999.....	30	32	62	60	123	163	245	248	183	54	6	-
White, male												
2003.....	30	27	35	42	54	86	94	87	57	17	2	-
2002.....	17	24	23	31	55	80	98	89	42	19	-	-
2001.....	23	16	34	36	49	64	91	75	45	18	1	-
2000.....	25	18	25	31	51	85	95	88	39	10	4	-
1999.....	17	21	36	29	57	68	109	86	44	7	2	-
White, female												
2003.....	18	33	36	46	66	99	136	189	127	38	5	-
2002.....	17	23	32	40	66	117	160	207	125	64	10	-
2001.....	24	29	28	33	67	100	144	176	134	59	3	-
2000.....	22	27	29	37	66	113	128	182	137	47	5	-
1999.....	13	11	26	31	66	95	136	162	139	47	4	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hernia (K40-K46)

Black, both sexes

2003.....	11	16	12	12	12	17	22	14	15	1	1	-
2002.....	8	8	13	15	12	10	20	19	4	2	1	-
2001.....	11	12	11	10	15	17	17	16	9	9	2	-
2000.....	8	5	8	14	7	22	16	24	12	3	1	-
1999.....	7	4	8	12	12	20	18	15	12	2	2	-

Black, male

2003.....	5	3	7	6	7	6	9	6	4	-	1	-
2002.....	2	3	4	8	7	4	7	6	2	1	-	-
2001.....	8	7	6	2	5	6	9	11	3	1	-	-
2000.....	5	1	4	7	2	4	8	8	3	-	-	-
1999.....	3	-	2	5	2	5	8	7	6	1	-	-

Black, female

2003.....	6	13	5	6	5	11	13	8	11	1	-	-
2002.....	6	5	9	7	5	6	13	13	2	1	1	-
2001.....	3	5	5	8	10	11	8	5	6	8	2	-
2000.....	3	4	4	7	5	18	8	16	9	3	1	-
1999.....	4	4	6	7	10	15	10	8	6	1	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hernia (K40-K46)

American Indian 1/, both sexes

2003.....	1	-	1	1	3	-	1	3	2	-	1	-
2002.....	2	-	1	3	-	1	3	1	-	-	-	-
2001.....	-	-	1	1	-	-	1	1	1	-	-	-
2000.....	2	1	-	2	1	1	1	-	-	-	-	-
1999.....	1	-	1	1	1	-	-	2	-	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	1	-	-	2	1	-	1	-
2002.....	-	-	1	1	-	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	1	-	-	-	-
2000.....	1	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	1	-	1	1	2	-	1	1	1	-	-	-
2002.....	2	-	-	2	-	1	3	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	1	-	-	-
2000.....	1	-	-	1	1	1	1	-	-	-	-	-
1999.....	1	-	-	1	1	-	-	2	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	100 years and over	Age not stated
	years	years	years	years	years	years	years	years	years	years		

Hernia (K40-K46)

Asian or Pacific Islander 2/, both sexes

2003.....	-	-	1	-	1	1	6	3	1	-	-	-
2002.....	-	-	2	1	2	-	-	4	1	-	-	-
2001.....	-	-	-	1	1	4	5	1	1	2	-	-
2000.....	1	-	-	-	1	3	2	-	-	3	-	-
1999.....	1	-	-	4	2	3	2	4	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	-	-	1	1	3	-	-	-	-	-
2002.....	-	-	1	-	1	-	-	2	1	-	-	-
2001.....	-	-	-	1	-	1	2	1	-	-	-	-
2000.....	1	-	-	-	-	2	1	-	-	2	-	-
1999.....	-	-	-	-	-	2	-	3	1	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	1	-	-	-	3	3	1	-	-	-
2002.....	-	-	1	1	1	-	-	2	-	-	-	-
2001.....	-	-	-	-	1	3	3	-	1	2	-	-
2000.....	-	-	-	-	1	1	1	-	-	1	-	-
1999.....	1	-	-	4	2	1	2	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All races, both sexes

2003.....	4,009	3,476	2,952	2,729	2,677	2,314	1,544	712	204	29	1	3
2002.....	3,812	3,258	2,839	2,670	2,711	2,412	1,588	728	230	24	3	4
2001.....	3,702	2,999	2,751	2,752	2,734	2,325	1,476	714	221	43	1	6
2000.....	3,280	2,947	2,827	2,751	2,731	2,367	1,462	742	203	30	4	3
1999.....	3,140	2,803	2,834	2,833	2,809	2,396	1,505	702	226	33	4	2

All races, male

2003.....	3,004	2,485	2,052	1,736	1,529	1,181	720	304	80	13	-	3
2002.....	2,843	2,323	1,905	1,652	1,545	1,211	742	310	77	8	-	4
2001.....	2,714	2,161	1,867	1,749	1,586	1,151	682	307	89	19	1	6
2000.....	2,460	2,066	1,939	1,729	1,588	1,240	634	325	83	12	1	3
1999.....	2,383	2,051	1,892	1,837	1,617	1,208	697	293	86	14	2	2

All races, female

2003.....	1,005	991	900	993	1,148	1,133	824	408	124	16	1	-
2002.....	969	935	934	1,018	1,166	1,201	846	418	153	16	3	-
2001.....	988	838	884	1,003	1,148	1,174	794	407	132	24	-	-
2000.....	820	881	888	1,022	1,143	1,127	828	417	120	18	3	-
1999.....	757	752	942	996	1,192	1,188	808	409	140	19	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

2003.....	3,331	2,960	2,559	2,423	2,460	2,182	1,453	670	191	26	1	3
2002.....	3,178	2,795	2,464	2,383	2,494	2,261	1,483	685	215	23	3	4
2001.....	3,031	2,513	2,411	2,442	2,497	2,165	1,396	675	208	40	1	5
2000.....	2,698	2,497	2,451	2,434	2,491	2,223	1,368	697	188	29	4	2
1999.....	2,634	2,338	2,422	2,509	2,564	2,220	1,420	654	208	30	4	2

White, male

2003.....	2,516	2,122	1,792	1,550	1,412	1,114	672	286	74	11	-	3
2002.....	2,400	2,014	1,658	1,474	1,427	1,148	695	293	71	8	-	4
2001.....	2,247	1,814	1,628	1,546	1,457	1,076	652	293	82	18	1	5
2000.....	2,045	1,775	1,700	1,553	1,453	1,162	600	305	76	12	1	2
1999.....	2,019	1,725	1,647	1,633	1,481	1,118	663	275	79	13	2	2

White, female

2003.....	815	838	767	873	1,048	1,068	781	384	117	15	1	-
2002.....	778	781	806	909	1,067	1,113	788	392	144	15	3	-
2001.....	784	699	783	896	1,040	1,089	744	382	126	22	-	-
2000.....	653	722	751	881	1,038	1,061	768	392	112	17	3	-
1999.....	615	613	775	876	1,083	1,102	757	379	129	17	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Black, both sexes

2003.....	554	417	296	242	164	83	58	31	9	2	-	-
2002.....	524	367	293	216	149	117	61	29	12	-	-	-
2001.....	539	393	264	249	168	121	54	22	10	2	-	1
2000.....	484	349	284	249	185	103	62	30	13	1	-	1
1999.....	406	370	329	251	187	130	65	38	14	2	-	-

Black, male

2003.....	411	301	202	149	88	43	30	16	4	1	-	-
2002.....	368	248	197	140	87	50	28	11	4	-	-	-
2001.....	390	280	189	170	100	58	18	7	4	-	-	1
2000.....	350	234	180	146	112	58	23	12	5	-	-	1
1999.....	297	260	200	167	110	67	29	14	4	-	-	-

Black, female

2003.....	143	116	94	93	76	40	28	15	5	1	-	-
2002.....	156	119	96	76	62	67	33	18	8	-	-	-
2001.....	149	113	75	79	68	63	36	15	6	2	-	-
2000.....	134	115	104	103	73	45	39	18	8	1	-	-
1999.....	109	110	129	84	77	63	36	24	10	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

American Indian 1/, both sexes

2003.....	95	67	67	32	18	17	6	2	2	-	-	-
2002.....	73	63	49	36	31	11	14	3	-	-	-	-
2001.....	92	56	44	31	27	11	7	3	1	-	-	-
2000.....	75	69	54	39	20	19	6	2	-	-	-	-
1999.....	70	68	55	38	20	13	2	2	-	-	-	-

American Indian 1/, male

2003.....	57	38	34	18	8	7	2	-	2	-	-	-
2002.....	51	37	28	21	17	5	7	1	-	-	-	-
2001.....	51	36	32	19	10	7	1	1	1	-	-	-
2000.....	49	39	32	15	6	11	1	1	-	-	-	-
1999.....	45	43	27	18	10	6	1	1	-	-	-	-

American Indian 1/, female

2003.....	38	29	33	14	10	10	4	2	-	-	-	-
2002.....	22	26	21	15	14	6	7	2	-	-	-	-
2001.....	41	20	12	12	17	4	6	2	-	-	-	-
2000.....	26	30	22	24	14	8	5	1	-	-	-	-
1999.....	25	25	28	20	10	7	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Asian or Pacific Islander 2/, both sexes

2003.....	29	32	30	32	35	32	27	9	2	1	-	-
2002.....	37	33	33	35	37	23	30	11	3	1	-	-
2001.....	40	37	32	30	42	28	19	14	2	1	-	-
2000.....	23	32	38	29	35	22	26	13	2	-	-	-
1999.....	30	27	28	35	38	33	18	8	4	1	-	-

Asian or Pacific Islander 2/, male

2003.....	20	24	24	19	21	17	16	2	-	1	-	-
2002.....	24	24	22	17	14	8	12	5	2	-	-	-
2001.....	26	31	18	14	19	10	11	6	2	1	-	-
2000.....	16	18	27	15	17	9	10	7	2	-	-	-
1999.....	22	23	18	19	16	17	4	3	3	1	-	-

Asian or Pacific Islander 2/, female

2003.....	9	8	6	13	14	15	11	7	2	-	-	-
2002.....	13	9	11	18	23	15	18	6	1	1	-	-
2001.....	14	6	14	16	23	18	8	8	-	-	-	-
2000.....	7	14	11	14	18	13	16	6	-	-	-	-
1999.....	8	4	10	16	22	16	14	5	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

All races, both sexes

2003.....	2,238	1,837	1,410	1,075	713	491	238	67	19	-	-	2
2002.....	2,139	1,712	1,371	1,023	826	495	235	79	16	3	-	2
2001.....	2,082	1,582	1,331	1,062	786	515	202	86	12	6	-	5
2000.....	1,888	1,575	1,394	1,085	806	533	218	76	17	2	-	2
1999.....	1,837	1,511	1,348	1,104	831	533	256	94	21	3	-	2

All races, male

2003.....	1,704	1,386	1,073	800	543	349	169	40	12	-	-	2
2002.....	1,624	1,277	1,060	771	596	334	161	54	10	1	-	2
2001.....	1,536	1,235	1,007	789	590	370	135	60	8	4	-	5
2000.....	1,458	1,172	1,044	821	612	419	142	52	12	1	-	2
1999.....	1,426	1,179	1,015	867	630	364	174	59	13	1	-	2

All races, female

2003.....	534	451	337	275	170	142	69	27	7	-	-	-
2002.....	515	435	311	252	230	161	74	25	6	2	-	-
2001.....	546	347	324	273	196	145	67	26	4	2	-	-
2000.....	430	403	350	264	194	114	76	24	5	1	-	-
1999.....	411	332	333	237	201	169	82	35	8	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

White, both sexes

2003.....	1,876	1,590	1,207	941	645	462	220	62	17	-	-	2
2002.....	1,808	1,466	1,190	903	749	450	215	71	14	3	-	2
2001.....	1,709	1,332	1,166	918	711	472	188	74	10	6	-	5
2000.....	1,576	1,329	1,204	961	728	487	194	70	16	2	-	1
1999.....	1,547	1,261	1,144	958	740	475	233	81	18	3	-	2

White, male

2003.....	1,444	1,209	934	704	494	329	153	37	11	-	-	2
2002.....	1,392	1,120	927	684	542	306	149	48	8	1	-	2
2001.....	1,287	1,045	886	683	533	343	126	53	6	4	-	5
2000.....	1,237	1,013	917	735	553	387	130	48	11	1	-	1
1999.....	1,217	993	883	753	564	325	161	50	12	1	-	2

White, female

2003.....	432	381	273	237	151	133	67	25	6	-	-	-
2002.....	416	346	263	219	207	144	66	23	6	2	-	-
2001.....	422	287	280	235	178	129	62	21	4	2	-	-
2000.....	339	316	287	226	175	100	64	22	5	1	-	-
1999.....	330	268	261	205	176	150	72	31	6	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

Black, both sexes

2003.....	274	192	153	107	52	18	15	4	1	-	-	-
2002.....	255	186	138	90	55	36	15	7	1	-	-	-
2001.....	295	200	128	120	58	33	11	8	1	-	-	-
2000.....	249	187	138	107	63	35	15	4	1	-	-	1
1999.....	227	194	167	117	77	45	19	11	3	-	-	-

Black, male

2003.....	204	139	108	76	38	11	13	3	1	-	-	-
2002.....	180	117	103	66	39	23	9	5	1	-	-	-
2001.....	206	150	92	90	45	19	7	4	1	-	-	-
2000.....	177	124	87	74	50	24	9	3	1	-	-	1
1999.....	164	147	110	93	55	30	11	7	1	-	-	-

Black, female

2003.....	70	53	45	31	14	7	2	1	-	-	-	-
2002.....	75	69	35	24	16	13	6	2	-	-	-	-
2001.....	89	50	36	30	13	14	4	4	-	-	-	-
2000.....	72	63	51	33	13	11	6	1	-	-	-	-
1999.....	63	47	57	24	22	15	8	4	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

American Indian 1/, both sexes

2003.....	75	43	38	18	9	7	-	1	-	-	-	-
2002.....	55	43	31	20	14	5	1	-	-	-	-	-
2001.....	61	34	27	18	10	3	1	1	1	-	-	-
2000.....	50	44	35	15	8	8	2	-	-	-	-	-
1999.....	53	43	32	19	8	7	1	1	-	-	-	-

American Indian 1/, male

2003.....	45	29	22	13	4	5	-	-	-	-	-	-
2002.....	38	26	19	13	10	3	-	-	-	-	-	-
2001.....	32	26	21	11	6	2	-	1	1	-	-	-
2000.....	33	25	26	11	3	5	1	-	-	-	-	-
1999.....	39	27	18	13	6	4	1	1	-	-	-	-

American Indian 1/, female

2003.....	30	14	16	5	5	2	-	1	-	-	-	-
2002.....	17	17	12	7	4	2	1	-	-	-	-	-
2001.....	29	8	6	7	4	1	1	-	-	-	-	-
2000.....	17	19	9	4	5	3	1	-	-	-	-	-
1999.....	14	16	14	6	2	3	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

Asian or Pacific Islander 2/, both sexes

2003.....	13	12	12	9	7	4	3	-	1	-	-	-
2002.....	21	17	12	10	8	4	4	1	1	-	-	-
2001.....	17	16	10	6	7	7	2	3	-	-	-	-
2000.....	13	15	17	2	7	3	7	2	-	-	-	-
1999.....	10	13	5	10	6	6	3	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	11	9	9	7	7	4	3	-	-	-	-	-
2002.....	14	14	11	8	5	2	3	1	1	-	-	-
2001.....	11	14	8	5	6	6	2	2	-	-	-	-
2000.....	11	10	14	1	6	3	2	1	-	-	-	-
1999.....	6	12	4	8	5	5	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	3	3	2	-	-	-	-	1	-	-	-
2002.....	7	3	1	2	3	2	1	-	-	-	-	-
2001.....	6	2	2	1	1	1	-	1	-	-	-	-
2000.....	2	5	3	1	1	-	5	1	-	-	-	-
1999.....	4	1	1	2	1	1	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

All races, both sexes

2003.....	1,771	1,639	1,542	1,654	1,964	1,823	1,306	645	185	29	1	1
2002.....	1,673	1,546	1,468	1,647	1,885	1,917	1,353	649	214	21	3	2
2001.....	1,620	1,417	1,420	1,690	1,948	1,810	1,274	628	209	37	1	1
2000.....	1,392	1,372	1,433	1,666	1,925	1,834	1,244	666	186	28	4	1
1999.....	1,303	1,292	1,486	1,729	1,978	1,863	1,249	608	205	30	4	-

All races, male

2003.....	1,300	1,099	979	936	986	832	551	264	68	13	-	1
2002.....	1,219	1,046	845	881	949	877	581	256	67	7	-	2
2001.....	1,178	926	860	960	996	781	547	247	81	15	1	1
2000.....	1,002	894	895	908	976	821	492	273	71	11	1	1
1999.....	957	872	877	970	987	844	523	234	73	13	2	-

All races, female

2003.....	471	540	563	718	978	991	755	381	117	16	1	-
2002.....	454	500	623	766	936	1,040	772	393	147	14	3	-
2001.....	442	491	560	730	952	1,029	727	381	128	22	-	-
2000.....	390	478	538	758	949	1,013	752	393	115	17	3	-
1999.....	346	420	609	759	991	1,019	726	374	132	17	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2003.....	1,455	1,370	1,352	1,482	1,815	1,720	1,233	608	174	26	1	1
2002.....	1,370	1,329	1,274	1,480	1,745	1,811	1,268	614	201	20	3	2
2001.....	1,322	1,181	1,245	1,524	1,786	1,693	1,208	601	198	34	1	-
2000.....	1,122	1,168	1,247	1,473	1,763	1,736	1,174	627	172	27	4	1
1999.....	1,087	1,077	1,278	1,551	1,824	1,745	1,187	573	190	27	4	-

White, male

2003.....	1,072	913	858	846	918	785	519	249	63	11	-	1
2002.....	1,008	894	731	790	885	842	546	245	63	7	-	2
2001.....	960	769	742	863	924	733	526	240	76	14	1	-
2000.....	808	762	783	818	900	775	470	257	65	11	1	1
1999.....	802	732	764	880	917	793	502	225	67	12	2	-

White, female

2003.....	383	457	494	636	897	935	714	359	111	15	1	-
2002.....	362	435	543	690	860	969	722	369	138	13	3	-
2001.....	362	412	503	661	862	960	682	361	122	20	-	-
2000.....	314	406	464	655	863	961	704	370	107	16	3	-
1999.....	285	345	514	671	907	952	685	348	123	15	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Black, both sexes

2003.....	280	225	143	135	112	65	43	27	8	2	-	-
2002.....	269	181	155	126	94	81	46	22	11	-	-	-
2001.....	244	193	136	129	110	88	43	14	9	2	-	1
2000.....	235	162	146	142	122	68	47	26	12	1	-	-
1999.....	179	176	162	134	110	85	46	27	11	2	-	-

Black, male

2003.....	207	162	94	73	50	32	17	13	3	1	-	-
2002.....	188	131	94	74	48	27	19	6	3	-	-	-
2001.....	184	130	97	80	55	39	11	3	3	-	-	1
2000.....	173	110	93	72	62	34	14	9	4	-	-	-
1999.....	133	113	90	74	55	37	18	7	3	-	-	-

Black, female

2003.....	73	63	49	62	62	33	26	14	5	1	-	-
2002.....	81	50	61	52	46	54	27	16	8	-	-	-
2001.....	60	63	39	49	55	49	32	11	6	2	-	-
2000.....	62	52	53	70	60	34	33	17	8	1	-	-
1999.....	46	63	72	60	55	48	28	20	8	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

American Indian 1/, both sexes

2003.....	20	24	29	14	9	10	6	1	2	-	-	-
2002.....	18	20	18	16	17	6	13	3	-	-	-	-
2001.....	31	22	17	13	17	8	6	2	-	-	-	-
2000.....	25	25	19	24	12	11	4	2	-	-	-	-
1999.....	17	25	23	19	12	6	1	1	-	-	-	-

American Indian 1/, male

2003.....	12	9	12	5	4	2	2	-	2	-	-	-
2002.....	13	11	9	8	7	2	7	1	-	-	-	-
2001.....	19	10	11	8	4	5	1	-	-	-	-	-
2000.....	16	14	6	4	3	6	-	1	-	-	-	-
1999.....	6	16	9	5	4	2	-	-	-	-	-	-

American Indian 1/, female

2003.....	8	15	17	9	5	8	4	1	-	-	-	-
2002.....	5	9	9	8	10	4	6	2	-	-	-	-
2001.....	12	12	6	5	13	3	5	2	-	-	-	-
2000.....	9	11	13	20	9	5	4	1	-	-	-	-
1999.....	11	9	14	14	8	4	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Asian or Pacific Islander 2/, both sexes

2003.....	16	20	18	23	28	28	24	9	1	1	-	-
2002.....	16	16	21	25	29	19	26	10	2	1	-	-
2001.....	23	21	22	24	35	21	17	11	2	1	-	-
2000.....	10	17	21	27	28	19	19	11	2	-	-	-
1999.....	20	14	23	25	32	27	15	7	4	1	-	-

Asian or Pacific Islander 2/, male

2003.....	9	15	15	12	14	13	13	2	-	1	-	-
2002.....	10	10	11	9	9	6	9	4	1	-	-	-
2001.....	15	17	10	9	13	4	9	4	2	1	-	-
2000.....	5	8	13	14	11	6	8	6	2	-	-	-
1999.....	16	11	14	11	11	12	3	2	3	1	-	-

Asian or Pacific Islander 2/, female

2003.....	7	5	3	11	14	15	11	7	1	-	-	-
2002.....	6	6	10	16	20	13	17	6	1	1	-	-
2001.....	8	4	12	15	22	17	8	7	-	-	-	-
2000.....	5	9	8	13	17	13	11	5	-	-	-	-
1999.....	4	3	9	14	21	15	12	5	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races, both sexes

2003.....	80	77	105	193	296	440	614	498	371	133	25	-
2002.....	77	93	123	194	310	420	572	525	351	150	28	-
2001.....	68	94	148	193	333	432	513	542	364	106	31	-
2000.....	66	80	109	184	266	419	507	528	358	143	24	-
1999.....	69	95	108	171	304	417	531	521	324	120	27	-

All races, male

2003.....	44	45	61	101	164	233	286	199	99	27	3	-
2002.....	48	59	63	103	155	229	271	195	112	26	2	-
2001.....	40	61	81	96	173	231	233	209	111	21	8	-
2000.....	30	48	62	100	142	224	237	204	105	27	3	-
1999.....	30	54	66	89	163	208	239	203	101	22	4	-

All races, female

2003.....	36	32	44	92	132	207	328	299	272	106	22	-
2002.....	29	34	60	91	155	191	301	330	239	124	26	-
2001.....	28	33	67	97	160	201	280	333	253	85	23	-
2000.....	36	32	47	84	124	195	270	324	253	116	21	-
1999.....	39	41	42	82	141	209	292	318	223	98	23	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2003.....	52	59	83	154	267	390	552	455	335	121	22	-
2002.....	55	77	100	166	265	361	510	483	327	142	24	-
2001.....	50	72	119	150	283	379	456	493	335	97	29	-
2000.....	49	64	87	151	215	373	454	489	335	135	23	-
1999.....	54	72	81	134	264	378	486	481	299	113	25	-

White, male

2003.....	31	35	47	82	150	208	258	184	90	25	2	-
2002.....	35	50	51	90	136	204	243	179	105	25	2	-
2001.....	31	48	64	73	152	209	211	187	101	19	7	-
2000.....	23	41	51	87	119	203	220	193	99	26	2	-
1999.....	24	38	53	72	145	191	227	184	93	22	4	-

White, female

2003.....	21	24	36	72	117	182	294	271	245	96	20	-
2002.....	20	27	49	76	129	157	267	304	222	117	22	-
2001.....	19	24	55	77	131	170	245	306	234	78	22	-
2000.....	26	23	36	64	96	170	234	296	236	109	21	-
1999.....	30	34	28	62	119	187	259	297	206	91	21	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2003.....	22	15	18	29	21	35	43	35	21	12	3	-
2002.....	19	12	17	22	34	45	49	26	14	6	4	-
2001.....	17	20	22	32	46	35	43	33	21	8	2	-
2000.....	14	13	16	28	37	37	39	33	18	6	-	-
1999.....	13	14	21	29	30	26	39	25	15	6	2	-

Black, male

2003.....	11	8	10	15	8	18	19	11	6	2	1	-
2002.....	10	6	11	9	10	17	21	8	3	-	-	-
2001.....	9	12	12	16	18	10	11	15	6	2	1	-
2000.....	6	5	9	10	15	15	12	9	3	-	-	-
1999.....	4	10	9	12	13	12	11	9	4	-	-	-

Black, female

2003.....	11	7	8	14	13	17	24	24	15	10	2	-
2002.....	9	6	6	13	24	28	28	18	11	6	4	-
2001.....	8	8	10	16	28	25	32	18	15	6	1	-
2000.....	8	8	7	18	22	22	27	24	15	6	-	-
1999.....	9	4	12	17	17	14	28	16	11	6	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

American Indian 1/, both sexes

2003.....	2	1	3	4	2	4	5	2	5	-	-	-
2002.....	2	1	3	3	3	3	2	4	2	-	-	-
2001.....	-	1	1	1	-	5	1	4	-	-	-	-
2000.....	1	1	1	-	2	5	3	3	1	1	-	-
1999.....	2	1	2	4	2	4	-	1	4	1	-	-

American Indian 1/, male

2003.....	-	1	3	2	2	2	2	-	-	-	-	-
2002.....	2	1	-	2	3	1	1	1	-	-	-	-
2001.....	-	1	1	1	-	3	-	3	-	-	-	-
2000.....	-	1	-	-	1	2	1	-	1	-	-	-
1999.....	2	-	2	2	1	1	-	-	1	-	-	-

American Indian 1/, female

2003.....	2	-	-	2	-	2	3	2	5	-	-	-
2002.....	-	-	3	1	-	2	1	3	2	-	-	-
2001.....	-	-	-	-	-	2	1	1	-	-	-	-
2000.....	1	-	1	-	1	3	2	3	-	1	-	-
1999.....	-	1	-	2	1	3	-	1	3	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Asian or Pacific Islander 2/, both sexes

2003.....	4	2	1	6	6	11	14	6	10	-	-	-
2002.....	1	3	3	3	8	11	11	12	8	2	-	-
2001.....	1	1	6	10	4	13	13	12	8	1	-	-
2000.....	2	2	5	5	12	4	11	3	4	1	1	-
1999.....	-	8	4	4	8	9	6	14	6	-	-	-

Asian or Pacific Islander 2/, male

2003.....	2	1	1	2	4	5	7	4	3	-	-	-
2002.....	1	2	1	2	6	7	6	7	4	1	-	-
2001.....	-	-	4	6	3	9	11	4	4	-	-	-
2000.....	1	1	2	3	7	4	4	2	2	1	1	-
1999.....	-	6	2	3	4	4	1	10	3	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	1	-	4	2	6	7	2	7	-	-	-
2002.....	-	1	2	1	2	4	5	5	4	1	-	-
2001.....	1	1	2	4	1	4	2	8	4	1	-	-
2000.....	1	1	3	2	5	-	7	1	2	-	-	-
1999.....	-	2	2	1	4	5	5	4	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races, both sexes

2003.....	1,166	1,638	2,168	2,996	4,349	6,298	7,795	7,277	4,584	1,641	314	2
2002.....	1,103	1,505	1,950	2,820	4,344	6,338	7,558	6,959	4,450	1,549	298	-
2001.....	1,073	1,404	1,880	3,026	4,330	6,105	6,992	6,612	4,225	1,528	303	1
2000.....	964	1,269	1,831	2,811	4,179	5,893	6,565	6,150	3,915	1,443	269	2
1999.....	872	1,176	1,688	2,694	4,147	5,707	6,220	5,825	3,703	1,388	254	-

All races, male

2003.....	711	936	1,129	1,565	2,220	3,188	3,861	3,232	1,795	492	81	2
2002.....	632	797	1,027	1,491	2,311	3,248	3,672	3,088	1,693	495	82	-
2001.....	608	751	963	1,610	2,216	3,074	3,411	2,996	1,619	435	64	-
2000.....	549	697	954	1,444	2,120	3,039	3,174	2,672	1,473	453	73	2
1999.....	481	627	903	1,366	2,138	2,986	3,045	2,584	1,353	424	67	-

All races, female

2003.....	455	702	1,039	1,431	2,129	3,110	3,934	4,045	2,789	1,149	233	-
2002.....	471	708	923	1,329	2,033	3,090	3,886	3,871	2,757	1,054	216	-
2001.....	465	653	917	1,416	2,114	3,031	3,581	3,616	2,606	1,093	239	1
2000.....	415	572	877	1,367	2,059	2,854	3,391	3,478	2,442	990	196	-
1999.....	391	549	785	1,328	2,009	2,721	3,175	3,241	2,350	964	187	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, both sexes

2003.....	677	1,054	1,463	2,113	3,265	5,141	6,627	6,406	4,028	1,430	258	1
2002.....	619	943	1,305	2,007	3,296	5,165	6,494	6,095	3,909	1,328	243	-
2001.....	637	881	1,220	2,124	3,334	4,984	5,952	5,803	3,697	1,337	251	1
2000.....	560	789	1,220	1,966	3,178	4,857	5,613	5,390	3,447	1,258	216	2
1999.....	505	743	1,125	1,908	3,179	4,679	5,255	4,989	3,249	1,192	200	-

White, male

2003.....	403	604	774	1,142	1,706	2,694	3,339	2,933	1,590	423	67	1
2002.....	358	503	683	1,096	1,818	2,731	3,241	2,759	1,496	429	62	-
2001.....	375	488	656	1,173	1,777	2,603	2,962	2,683	1,448	387	56	-
2000.....	321	444	662	1,071	1,698	2,567	2,786	2,398	1,320	387	66	2
1999.....	282	425	618	1,005	1,706	2,492	2,627	2,243	1,212	384	55	-

White, female

2003.....	274	450	689	971	1,559	2,447	3,288	3,473	2,438	1,007	191	-
2002.....	261	440	622	911	1,478	2,434	3,253	3,336	2,413	899	181	-
2001.....	262	393	564	951	1,557	2,381	2,990	3,120	2,249	950	195	1
2000.....	239	345	558	895	1,480	2,290	2,827	2,992	2,127	871	150	-
1999.....	223	318	507	903	1,473	2,187	2,628	2,746	2,037	808	145	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2003.....	453	542	641	799	966	1,005	1,027	776	496	190	48	1
2002.....	454	502	580	746	944	1,036	923	764	472	199	52	-
2001.....	397	461	593	821	890	992	918	714	459	176	50	-
2000.....	369	436	555	756	897	931	858	671	413	171	44	-
1999.....	347	382	505	704	865	917	852	732	399	174	54	-

Black, male

2003.....	290	309	325	382	474	420	452	254	172	58	13	1
2002.....	258	269	309	358	448	452	368	282	168	55	19	-
2001.....	211	241	273	398	389	407	386	262	143	39	7	-
2000.....	207	227	262	334	382	421	344	234	126	60	5	-
1999.....	190	172	257	315	385	438	375	293	114	30	12	-

Black, female

2003.....	163	233	316	417	492	585	575	522	324	132	35	-
2002.....	196	233	271	388	496	584	555	482	304	144	33	-
2001.....	186	220	320	423	501	585	532	452	316	137	43	-
2000.....	162	209	293	422	515	510	514	437	287	111	39	-
1999.....	157	210	248	389	480	479	477	439	285	144	42	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

American Indian 1/, both sexes

2003.....	14	18	24	24	22	49	27	23	14	5	-	-
2002.....	12	22	21	24	32	23	23	14	12	2	2	-
2001.....	15	26	25	24	30	32	20	15	10	2	1	-
2000.....	17	17	22	36	29	22	24	11	13	2	1	-
1999.....	7	17	19	31	24	26	16	19	6	7	-	-

American Indian 1/, male

2003.....	8	11	10	11	8	29	14	10	4	3	-	-
2002.....	7	10	10	10	9	7	13	4	3	1	1	-
2001.....	10	6	10	7	12	17	7	10	4	-	-	-
2000.....	9	9	12	18	10	7	9	5	6	2	-	-
1999.....	4	9	8	13	10	11	1	5	1	2	-	-

American Indian 1/, female

2003.....	6	7	14	13	14	20	13	13	10	2	-	-
2002.....	5	12	11	14	23	16	10	10	9	1	1	-
2001.....	5	20	15	17	18	15	13	5	6	2	1	-
2000.....	8	8	10	18	19	15	15	6	7	-	1	-
1999.....	3	8	11	18	14	15	15	14	5	5	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Asian or Pacific Islander 2/, both sexes

2003.....	22	24	40	60	96	103	114	72	46	16	8	-
2002.....	18	38	44	43	72	114	118	86	57	20	1	-
2001.....	24	36	42	57	76	97	102	80	59	13	1	-
2000.....	18	27	34	53	75	83	70	78	42	12	8	-
1999.....	13	34	39	51	79	85	97	85	49	15	-	-

Asian or Pacific Islander 2/, male

2003.....	10	12	20	30	32	45	56	35	29	8	1	-
2002.....	9	15	25	27	36	58	50	43	26	10	-	-
2001.....	12	16	24	32	38	47	56	41	24	9	1	-
2000.....	12	17	18	21	30	44	35	35	21	4	2	-
1999.....	5	21	20	33	37	45	42	43	26	8	-	-

Asian or Pacific Islander 2/, female

2003.....	12	12	20	30	64	58	58	37	17	8	7	-
2002.....	9	23	19	16	36	56	68	43	31	10	1	-
2001.....	12	20	18	25	38	50	46	39	35	4	-	-
2000.....	6	10	16	32	45	39	35	43	21	8	6	-
1999.....	8	13	19	18	42	40	55	42	23	7	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races, both sexes

2003.....	4	5	9	11	19	18	32	17	5	2	2	-
2002.....	4	6	10	13	14	19	32	23	13	5	1	-
2001.....	5	9	4	13	14	28	20	24	12	1	1	-
2000.....	11	2	8	13	16	16	36	27	13	8	1	-
1999.....	5	5	11	10	27	37	25	22	7	5	-	-

All races, male

2003.....	1	3	4	6	9	7	13	2	3	-	-	-
2002.....	1	5	6	9	5	5	12	10	2	1	1	-
2001.....	4	6	3	6	4	10	9	10	5	-	-	-
2000.....	9	2	6	7	5	11	15	13	5	4	-	-
1999.....	3	2	4	5	12	15	17	9	1	4	-	-

All races, female

2003.....	3	2	5	5	10	11	19	15	2	2	2	-
2002.....	3	1	4	4	9	14	20	13	11	4	-	-
2001.....	1	3	1	7	10	18	11	14	7	1	1	-
2000.....	2	-	2	6	11	5	21	14	8	4	1	-
1999.....	2	3	7	5	15	22	8	13	6	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

White, both sexes

2003.....	4	3	6	9	17	16	26	14	5	2	-	-
2002.....	2	5	9	9	12	17	25	17	10	5	1	-
2001.....	4	7	4	9	12	25	18	20	9	1	1	-
2000.....	4	2	8	8	14	12	34	23	12	8	1	-
1999.....	3	3	6	9	23	32	20	17	7	5	-	-

White, male

2003.....	1	2	1	4	8	5	9	2	3	-	-	-
2002.....	1	4	6	6	5	4	10	8	1	1	1	-
2001.....	4	4	3	3	3	10	9	8	4	-	-	-
2000.....	3	2	6	4	5	9	14	10	4	4	-	-
1999.....	2	1	2	4	10	13	13	5	1	4	-	-

White, female

2003.....	3	1	5	5	9	11	17	12	2	2	-	-
2002.....	1	1	3	3	7	13	15	9	9	4	-	-
2001.....	-	3	1	6	9	15	9	12	5	1	1	-
2000.....	1	-	2	4	9	3	20	13	8	4	1	-
1999.....	1	2	4	5	13	19	7	12	6	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Black, both sexes												
2003.....	-	1	2	2	2	2	4	1	-	-	1	-
2002.....	2	1	-	3	2	2	4	5	2	-	-	-
2001.....	1	1	-	4	2	3	2	2	2	-	-	-
2000.....	7	-	-	4	2	3	2	2	-	-	-	-
1999.....	2	1	4	-	3	3	3	4	-	-	-	-
Black, male												
2003.....	-	-	2	2	1	2	3	-	-	-	-	-
2002.....	-	1	-	2	-	1	1	2	-	-	-	-
2001.....	-	1	-	3	1	-	-	1	-	-	-	-
2000.....	6	-	-	2	-	1	1	1	-	-	-	-
1999.....	1	-	2	-	2	1	2	3	-	-	-	-
Black, female												
2003.....	-	1	-	-	1	-	1	1	-	-	1	-
2002.....	2	-	-	1	2	1	3	3	2	-	-	-
2001.....	1	-	-	1	1	3	2	1	2	-	-	-
2000.....	1	-	-	2	2	2	1	1	-	-	-	-
1999.....	1	1	2	-	1	2	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

American Indian 1/, both sexes

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	1	-	-	-	1	-	-	-	-	-	-

American Indian 1/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Asian or Pacific Islander 2/, both sexes

2003.....	-	1	1	-	-	-	2	1	-	-	1	-
2002.....	-	-	1	1	-	-	2	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	2	1	-	-	-
2000.....	-	-	-	1	-	1	-	1	1	-	-	-
1999.....	-	-	1	1	1	1	2	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	1	1	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	1	-	-	-
2000.....	-	-	-	1	-	1	-	1	1	-	-	-
1999.....	-	-	-	1	-	1	2	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	-	-	-	-	1	1	-	-	1	-
2002.....	-	-	1	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races, both sexes

2003.....	30	22	28	42	60	77	88	72	37	16	5	-
2002.....	25	33	19	29	56	86	92	74	38	14	1	-
2001.....	26	17	23	47	64	69	78	70	38	13	1	-
2000.....	21	31	28	43	60	78	94	78	23	15	2	-
1999.....	28	26	34	40	61	104	96	83	37	10	2	-

All races, male

2003.....	23	10	12	24	24	39	41	32	20	6	1	-
2002.....	15	20	7	14	35	50	42	32	19	2	-	-
2001.....	15	10	15	23	40	39	45	32	10	6	-	-
2000.....	14	16	15	24	30	45	38	40	9	6	-	-
1999.....	15	13	19	20	25	58	49	38	17	5	-	-

All races, female

2003.....	7	12	16	18	36	38	47	40	17	10	4	-
2002.....	10	13	12	15	21	36	50	42	19	12	1	-
2001.....	11	7	8	24	24	30	33	38	28	7	1	-
2000.....	7	15	13	19	30	33	56	38	14	9	2	-
1999.....	13	13	15	20	36	46	47	45	20	5	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2003.....	19	21	21	36	43	62	75	64	28	15	4	-
2002.....	15	24	15	28	51	68	81	60	35	12	-	-
2001.....	13	13	14	35	58	61	68	66	36	12	-	-
2000.....	11	24	19	32	53	67	83	64	17	12	1	-
1999.....	18	18	25	27	51	91	82	70	29	8	2	-

White, male

2003.....	16	10	9	21	17	30	35	28	15	6	1	-
2002.....	8	16	7	14	31	40	38	28	18	2	-	-
2001.....	7	7	9	17	37	34	41	31	10	6	-	-
2000.....	7	13	10	18	27	39	36	34	6	4	-	-
1999.....	10	9	15	15	20	52	42	32	14	4	-	-

White, female

2003.....	3	11	12	15	26	32	40	36	13	9	3	-
2002.....	7	8	8	14	20	28	43	32	17	10	-	-
2001.....	6	6	5	18	21	27	27	35	26	6	-	-
2000.....	4	11	9	14	26	28	47	30	11	8	1	-
1999.....	8	9	10	12	31	39	40	38	15	4	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Black, both sexes

2003.....	9	1	5	4	11	11	8	3	8	-	1	-
2002.....	9	5	3	1	4	13	8	12	2	2	1	-
2001.....	9	2	7	10	4	6	7	3	2	1	1	-
2000.....	9	5	7	9	5	9	10	9	4	2	-	-
1999.....	9	6	5	11	7	11	9	12	5	2	-	-

Black, male

2003.....	6	-	2	2	5	6	4	1	4	-	-	-
2002.....	6	3	-	-	4	8	3	3	1	-	-	-
2001.....	7	2	5	5	2	3	2	1	-	-	-	-
2000.....	6	2	5	6	2	4	2	4	2	1	-	-
1999.....	5	3	3	4	3	6	4	5	2	1	-	-

Black, female

2003.....	3	1	3	2	6	5	4	2	4	-	1	-
2002.....	3	2	3	1	-	5	5	9	1	2	1	-
2001.....	2	-	2	5	2	3	5	2	2	1	1	-
2000.....	3	3	2	3	3	5	8	5	2	1	-	-
1999.....	4	3	2	7	4	5	5	7	3	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Asian or Pacific Islander 2/, both sexes

2003.....	1	-	2	2	4	2	5	5	1	1	-	-
2002.....	1	3	1	-	1	4	2	2	1	-	-	-
2001.....	3	1	2	2	2	-	2	1	-	-	-	-
2000.....	-	1	2	1	2	1	1	5	2	1	-	-
1999.....	1	1	3	1	3	2	5	1	3	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	1	1	-	1	2	3	1	-	-	-
2002.....	1	-	-	-	-	1	1	1	-	-	-	-
2001.....	1	1	1	1	1	-	1	-	-	-	-	-
2000.....	-	-	-	-	1	1	-	2	1	1	-	-
1999.....	-	-	1	-	2	-	3	1	1	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	-	1	1	4	1	3	2	-	1	-	-
2002.....	-	3	1	-	1	3	1	1	1	-	-	-
2001.....	2	-	1	1	1	-	1	1	-	-	-	-
2000.....	-	1	2	1	1	-	1	3	1	-	-	-
1999.....	1	1	2	1	1	2	2	-	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Renal failure (N17-N19)

All races, both sexes

2003.....	1,131	1,611	2,130	2,939	4,267	6,198	7,672	7,184	4,541	1,621	307	2
2002.....	1,072	1,464	1,919	2,774	4,272	6,228	7,429	6,856	4,398	1,530	296	-
2001.....	1,041	1,376	1,850	2,963	4,248	6,007	6,891	6,517	4,174	1,514	300	1
2000.....	932	1,234	1,789	2,754	4,094	5,794	6,429	6,040	3,878	1,420	265	2
1999.....	836	1,144	1,640	2,641	4,056	5,560	6,092	5,718	3,656	1,372	252	-

All races, male

2003.....	686	923	1,113	1,532	2,187	3,140	3,807	3,198	1,771	486	80	2
2002.....	615	772	1,014	1,468	2,271	3,190	3,618	3,046	1,672	492	81	-
2001.....	588	735	945	1,581	2,171	3,025	3,356	2,953	1,604	429	64	-
2000.....	526	678	929	1,412	2,080	2,981	3,121	2,617	1,458	443	73	2
1999.....	462	612	880	1,340	2,100	2,912	2,976	2,535	1,334	415	67	-

All races, female

2003.....	445	688	1,017	1,407	2,080	3,058	3,865	3,986	2,770	1,135	227	-
2002.....	457	692	905	1,306	2,001	3,038	3,811	3,810	2,726	1,038	215	-
2001.....	453	641	905	1,382	2,077	2,982	3,535	3,564	2,570	1,085	236	1
2000.....	406	556	860	1,342	2,014	2,813	3,308	3,423	2,420	977	192	-
1999.....	374	532	760	1,301	1,956	2,648	3,116	3,183	2,322	957	185	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Renal failure (N17-N19)

White, both sexes

2003.....	653	1,030	1,435	2,067	3,203	5,058	6,524	6,324	3,994	1,411	254	1
2002.....	601	912	1,279	1,966	3,231	5,076	6,385	6,013	3,863	1,311	242	-
2001.....	619	860	1,199	2,077	3,260	4,898	5,865	5,717	3,651	1,324	250	1
2000.....	545	761	1,188	1,925	3,103	4,774	5,491	5,299	3,417	1,238	214	2
1999.....	482	722	1,093	1,870	3,102	4,551	5,150	4,900	3,212	1,178	198	-

White, male

2003.....	385	592	764	1,116	1,681	2,657	3,295	2,903	1,571	417	66	1
2002.....	349	483	670	1,076	1,782	2,685	3,193	2,723	1,477	426	61	-
2001.....	363	477	644	1,153	1,736	2,559	2,912	2,644	1,434	381	56	-
2000.....	311	428	643	1,048	1,661	2,517	2,736	2,352	1,309	379	66	2
1999.....	269	415	601	986	1,675	2,426	2,570	2,204	1,196	376	55	-

White, female

2003.....	268	438	671	951	1,522	2,401	3,229	3,421	2,423	994	188	-
2002.....	252	429	609	890	1,449	2,391	3,192	3,290	2,386	885	181	-
2001.....	256	383	555	924	1,524	2,339	2,953	3,073	2,217	943	194	1
2000.....	234	333	545	877	1,442	2,257	2,755	2,947	2,108	859	148	-
1999.....	213	307	492	884	1,427	2,125	2,580	2,696	2,016	802	143	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Renal failure (N17-N19)

Black, both sexes

2003.....	444	540	634	790	953	992	1,014	772	488	190	46	1
2002.....	442	496	577	742	938	1,021	910	746	468	197	51	-
2001.....	387	457	586	807	884	982	908	708	455	175	48	-
2000.....	353	431	548	743	889	918	846	659	409	169	43	-
1999.....	335	374	494	692	855	902	836	716	392	172	54	-

Black, male

2003.....	284	309	321	376	468	412	445	253	168	58	13	1
2002.....	251	265	309	356	444	443	364	277	167	55	19	-
2001.....	204	238	268	390	386	404	383	259	143	39	7	-
2000.....	195	225	257	326	380	416	341	229	124	59	5	-
1999.....	184	169	252	310	380	431	368	285	112	29	12	-

Black, female

2003.....	160	231	313	414	485	580	569	519	320	132	33	-
2002.....	191	231	268	386	494	578	546	469	301	142	32	-
2001.....	183	219	318	417	498	578	525	449	312	136	41	-
2000.....	158	206	291	417	509	502	505	430	285	110	38	-
1999.....	151	205	242	382	475	471	468	431	280	143	42	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Renal failure (N17-N19)

American Indian 1/, both sexes

2003.....	13	18	24	24	20	47	27	22	14	5	-	-
2002.....	12	21	21	24	32	22	21	14	11	2	2	-
2001.....	14	24	25	24	30	30	19	15	10	2	1	-
2000.....	16	16	21	35	29	21	24	10	13	2	-	-
1999.....	7	15	18	30	24	25	16	19	6	7	-	-

American Indian 1/, male

2003.....	7	11	10	11	6	27	14	10	4	3	-	-
2002.....	7	9	10	10	9	6	13	4	2	1	1	-
2001.....	10	5	10	7	12	15	6	10	4	-	-	-
2000.....	8	8	11	18	10	6	9	4	6	2	-	-
1999.....	4	7	8	12	10	11	1	5	1	2	-	-

American Indian 1/, female

2003.....	6	7	14	13	14	20	13	12	10	2	-	-
2002.....	5	12	11	14	23	16	8	10	9	1	1	-
2001.....	4	19	15	17	18	15	13	5	6	2	1	-
2000.....	8	8	10	17	19	15	15	6	7	-	-	-
1999.....	3	8	10	18	14	14	15	14	5	5	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Renal failure (N17-N19)

Asian or Pacific Islander 2/, both sexes

2003.....	21	23	37	58	91	101	107	66	45	15	7	-
2002.....	17	35	42	42	71	109	113	83	56	20	1	-
2001.....	21	35	40	55	74	97	99	77	58	13	1	-
2000.....	18	26	32	51	73	81	68	72	39	11	8	-
1999.....	12	33	35	49	75	82	90	83	46	15	-	-

Asian or Pacific Islander 2/, male

2003.....	10	11	18	29	32	44	53	32	28	8	1	-
2002.....	8	15	25	26	36	56	48	42	26	10	-	-
2001.....	11	15	23	31	37	47	55	40	23	9	1	-
2000.....	12	17	18	20	29	42	35	32	19	3	2	-
1999.....	5	21	19	32	35	44	37	41	25	8	-	-

Asian or Pacific Islander 2/, female

2003.....	11	12	19	29	59	57	54	34	17	7	6	-
2002.....	9	20	17	16	35	53	65	41	30	10	1	-
2001.....	10	20	17	24	37	50	44	37	35	4	-	-
2000.....	6	9	14	31	44	39	33	40	20	8	6	-
1999.....	7	12	16	17	40	38	53	42	21	7	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of kidney (N25,N27)

All races, both sexes

2003.....	1	-	1	4	3	5	3	4	1	2	-	-
2002.....	2	2	2	4	2	5	5	6	1	-	-	-
2001.....	1	2	3	3	4	1	3	1	1	-	1	-
2000.....	-	2	6	1	9	5	6	5	1	-	1	-
1999.....	3	1	3	3	3	6	7	2	3	1	-	-

All races, male

2003.....	1	-	-	3	-	2	-	-	1	-	-	-
2002.....	1	-	-	-	-	3	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	1	1	-	-	-	-
2000.....	-	1	4	1	5	2	-	2	1	-	-	-
1999.....	1	-	-	1	1	1	3	2	1	-	-	-

All races, female

2003.....	-	-	1	1	3	3	3	4	-	2	-	-
2002.....	1	2	2	4	2	2	5	6	1	-	-	-
2001.....	-	2	3	3	3	1	2	-	1	-	1	-
2000.....	-	1	2	-	4	3	6	3	-	-	1	-
1999.....	2	1	3	2	2	5	4	-	2	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of kidney (N25,N27)

White, both sexes

2003.....	1	-	1	1	2	5	2	4	1	2	-	-
2002.....	1	2	2	4	2	4	3	5	1	-	-	-
2001.....	1	1	3	3	4	-	1	-	1	-	-	-
2000.....	-	2	5	1	8	4	5	4	1	-	-	-
1999.....	2	-	1	2	3	5	3	2	1	1	-	-

White, male

2003.....	1	-	-	1	-	2	-	-	1	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	1	3	1	5	2	-	2	1	-	-	-
1999.....	1	-	-	-	1	1	2	2	1	-	-	-

White, female

2003.....	-	-	1	-	2	3	2	4	-	2	-	-
2002.....	1	2	2	4	2	2	3	5	1	-	-	-
2001.....	-	1	3	3	3	-	1	-	1	-	-	-
2000.....	-	1	2	-	3	2	5	2	-	-	-	-
1999.....	1	-	1	2	2	4	1	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of kidney (N25,N27)

Black, both sexes

2003.....	-	-	-	3	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	1	-	-	-	1	1	1	-	-	1	-
2000.....	-	-	-	-	1	1	-	1	-	-	1	-
1999.....	1	1	2	1	-	1	4	-	2	-	-	-

Black, male

2003.....	-	-	-	2	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-

Black, female

2003.....	-	-	-	1	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	1	-	-	-	1	-	-	-	-	1	-
2000.....	-	-	-	-	1	1	-	1	-	-	1	-
1999.....	1	1	2	-	-	1	3	-	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races, both sexes

2003.....	30	34	28	56	49	95	152	141	91	40	6	-
2002.....	21	26	33	48	62	118	120	146	105	36	3	-
2001.....	25	23	23	35	52	126	123	129	100	33	3	-
2000.....	21	25	34	38	66	114	134	145	114	44	17	-
1999.....	33	30	29	52	81	109	133	154	102	49	6	-

All races, male

2003.....	11	11	10	15	16	25	49	44	27	7	-	-
2002.....	10	10	11	19	28	44	28	42	25	7	-	-
2001.....	5	6	6	19	21	36	40	32	16	8	-	-
2000.....	9	5	10	13	13	34	40	32	30	5	2	-
1999.....	11	14	13	19	25	37	46	45	24	12	-	-

All races, female

2003.....	19	23	18	41	33	70	103	97	64	33	6	-
2002.....	11	16	22	29	34	74	92	104	80	29	3	-
2001.....	20	17	17	16	31	90	83	97	84	25	3	-
2000.....	12	20	24	25	53	80	94	113	84	39	15	-
1999.....	22	16	16	33	56	72	87	109	78	37	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2003.....	24	26	20	48	45	83	132	132	82	36	4	-
2002.....	16	22	24	41	60	105	112	134	98	31	3	-
2001.....	19	18	19	26	47	115	104	117	92	33	3	-
2000.....	13	22	26	31	51	96	114	134	104	38	15	-
1999.....	25	26	22	43	67	99	124	144	97	45	5	-

White, male

2003.....	9	9	6	13	15	22	40	44	25	7	-	-
2002.....	9	9	8	13	27	38	25	37	25	6	-	-
2001.....	4	6	3	12	18	30	35	28	14	8	-	-
2000.....	6	5	8	9	8	30	35	32	27	4	2	-
1999.....	8	12	10	16	18	35	42	43	23	12	-	-

White, female

2003.....	15	17	14	35	30	61	92	88	57	29	4	-
2002.....	7	13	16	28	33	67	87	97	73	25	3	-
2001.....	15	12	16	14	29	85	69	89	78	25	3	-
2000.....	7	17	18	22	43	66	79	102	77	34	13	-
1999.....	17	14	12	27	49	64	82	101	74	33	5	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Black, both sexes

2003.....	5	7	7	6	4	9	15	8	8	2	2	-
2002.....	2	4	6	7	1	11	4	11	5	5	-	-
2001.....	6	3	4	8	5	8	14	10	6	-	-	-
2000.....	8	2	6	7	12	16	12	10	8	6	1	-
1999.....	7	4	7	7	13	8	8	9	5	3	1	-

Black, male

2003.....	2	2	3	2	1	3	8	-	2	-	-	-
2002.....	-	1	2	6	-	4	1	4	-	1	-	-
2001.....	1	-	3	6	3	3	3	3	1	-	-	-
2000.....	3	-	2	4	4	4	3	-	3	1	-	-
1999.....	3	2	3	3	7	1	3	2	1	-	-	-

Black, female

2003.....	3	5	4	4	3	6	7	8	6	2	2	-
2002.....	2	3	4	1	1	7	3	7	5	4	-	-
2001.....	5	3	1	2	2	5	11	7	5	-	-	-
2000.....	5	2	4	3	8	12	9	10	5	5	1	-
1999.....	4	2	4	4	6	7	5	7	4	3	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Asian or Pacific Islander 2/, both sexes

2003.....	1	-	1	2	-	2	5	1	1	1	-	-
2002.....	2	-	2	-	1	1	3	1	1	-	-	-
2001.....	-	2	-	1	-	2	4	2	1	-	-	-
2000.....	-	1	2	-	2	2	7	1	1	-	-	-
1999.....	1	-	-	-	1	2	1	1	-	1	-	-

Asian or Pacific Islander 2/, male

2003.....	-	-	1	-	-	-	1	-	-	-	-	-
2002.....	1	-	1	-	1	1	1	1	-	-	-	-
2001.....	-	-	-	1	-	2	1	1	1	-	-	-
2000.....	-	-	-	-	1	-	2	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	-	-	2	-	2	4	1	1	1	-	-
2002.....	1	-	1	-	-	-	2	-	1	-	-	-
2001.....	-	2	-	-	-	-	3	1	-	-	-	-
2000.....	-	1	2	-	1	2	5	1	1	-	-	-
1999.....	1	-	-	-	1	1	-	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

All races, both sexes

2003.....	7	3	7	6	10	17	12	18	9	4	-	-
2002.....	7	4	2	4	4	17	12	22	11	3	-	-
2001.....	3	4	8	8	10	13	18	23	11	4	-	-
2000.....	3	4	4	11	10	17	12	11	12	3	1	-
1999.....	2	6	7	4	11	15	17	6	6	5	1	-

All races, male

2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2003.....	7	3	7	6	10	17	12	18	9	4	-	-
2002.....	7	4	2	4	4	17	12	22	11	3	-	-
2001.....	3	4	8	8	10	13	18	23	11	4	-	-
2000.....	3	4	4	11	10	17	12	11	12	3	1	-
1999.....	2	6	7	4	11	15	17	6	6	5	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

White, both sexes

2003.....	4	2	6	5	8	15	10	17	9	4	-	-
2002.....	5	3	2	4	3	16	12	19	10	3	-	-
2001.....	2	4	4	6	8	11	16	22	8	4	-	-
2000.....	3	2	4	9	8	15	11	9	12	2	1	-
1999.....	2	6	6	2	8	10	16	6	6	5	-	-

White, male

2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2003.....	4	2	6	5	8	15	10	17	9	4	-	-
2002.....	5	3	2	4	3	16	12	19	10	3	-	-
2001.....	2	4	4	6	8	11	16	22	8	4	-	-
2000.....	3	2	4	9	8	15	11	9	12	2	1	-
1999.....	2	6	6	2	8	10	16	6	6	5	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Black, both sexes

2003.....	3	1	1	-	2	1	1	1	-	-	-	-
2002.....	2	1	-	-	1	1	-	2	1	-	-	-
2001.....	1	-	4	2	2	2	2	-	3	-	-	-
2000.....	-	2	-	2	2	2	1	2	-	-	-	-
1999.....	-	-	1	2	3	5	1	-	-	-	1	-

Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2003.....	3	1	1	-	2	1	1	1	-	-	-	-
2002.....	2	1	-	-	1	1	-	2	1	-	-	-
2001.....	1	-	4	2	2	2	2	-	3	-	-	-
2000.....	-	2	-	2	2	2	1	2	-	-	-	-
1999.....	-	-	1	2	3	5	1	-	-	-	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, both sexes

2003.....	343	376	290	223	190	224	240	170	100	40	9	-
2002.....	363	351	283	224	216	232	257	184	86	33	2	1
2001.....	333	298	283	214	234	244	221	182	87	38	10	-
2000.....	325	296	273	248	216	242	246	178	97	39	6	1
1999.....	305	304	253	251	218	249	242	187	100	34	5	2

All races, male

2003.....	172	189	147	108	90	86	92	57	27	12	1	-
2002.....	186	170	151	107	106	105	96	65	22	9	1	1
2001.....	167	146	130	109	119	110	93	70	17	9	4	-
2000.....	190	130	134	120	102	99	98	74	41	13	2	-
1999.....	162	157	126	106	103	102	102	60	35	6	1	1

All races, female

2003.....	171	187	143	115	100	138	148	113	73	28	8	-
2002.....	177	181	132	117	110	127	161	119	64	24	1	-
2001.....	166	152	153	105	115	134	128	112	70	29	6	-
2000.....	135	166	139	128	114	143	148	104	56	26	4	1
1999.....	143	147	127	145	115	147	140	127	65	28	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2003.....	305	335	261	191	169	204	220	160	90	36	9	-
2002.....	314	304	248	202	191	212	232	170	83	30	2	1
2001.....	293	267	260	193	210	226	203	172	81	34	6	-
2000.....	294	255	241	219	191	225	232	168	94	34	5	1
1999.....	265	275	225	225	191	221	224	170	92	34	5	2

White, male

2003.....	155	169	129	92	77	78	79	55	24	10	1	-
2002.....	157	146	132	99	91	95	83	61	22	8	1	1
2001.....	148	130	119	103	110	106	81	64	14	9	1	-
2000.....	168	114	123	99	89	94	94	70	40	12	1	-
1999.....	140	143	114	95	91	83	96	53	33	6	1	1

White, female

2003.....	150	166	132	99	92	126	141	105	66	26	8	-
2002.....	157	158	116	103	100	117	149	109	61	22	1	-
2001.....	145	137	141	90	100	120	122	108	67	25	5	-
2000.....	126	141	118	120	102	131	138	98	54	22	4	1
1999.....	125	132	111	130	100	138	128	117	59	28	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2003.....	28	34	26	24	16	15	15	8	8	4	-	-
2002.....	40	41	28	20	20	14	18	13	2	3	-	-
2001.....	33	27	21	15	21	17	11	7	5	4	4	-
2000.....	28	32	27	25	18	14	11	7	3	5	1	-
1999.....	33	25	21	21	25	22	16	16	8	-	-	-

Black, male

2003.....	10	18	15	14	11	5	11	2	2	2	-	-
2002.....	22	22	18	8	13	7	9	4	-	1	-	-
2001.....	17	13	10	4	8	4	8	4	2	-	3	-
2000.....	19	11	7	18	8	5	4	3	1	1	1	-
1999.....	17	12	10	6	10	14	5	7	2	-	-	-

Black, female

2003.....	18	16	11	10	5	10	4	6	6	2	-	-
2002.....	18	19	10	12	7	7	9	9	2	2	-	-
2001.....	16	14	11	11	13	13	3	3	3	4	1	-
2000.....	9	21	20	7	10	9	7	4	2	4	-	-
1999.....	16	13	11	15	15	8	11	9	6	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Asian or Pacific Islander 2/, both sexes

2003.....	8	5	2	6	4	5	5	-	2	-	-	-
2002.....	9	4	5	2	5	5	3	1	1	-	-	-
2001.....	6	3	2	6	2	1	5	3	1	-	-	-
2000.....	2	7	2	2	4	2	2	3	-	-	-	-
1999.....	4	1	5	4	2	5	2	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	6	1	2	2	2	3	2	-	1	-	-	-
2002.....	7	1	1	-	2	3	2	-	-	-	-	-
2001.....	2	3	1	2	-	-	4	2	1	-	-	-
2000.....	2	4	1	2	3	-	-	1	-	-	-	-
1999.....	2	-	2	4	2	4	1	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	4	-	4	2	2	3	-	1	-	-	-
2002.....	2	3	4	2	3	2	1	1	1	-	-	-
2001.....	4	-	1	4	2	1	1	1	-	-	-	-
2000.....	-	3	1	-	1	2	2	2	-	-	-	-
1999.....	2	1	3	-	-	1	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races, both sexes

2003.....	1,296	1,077	990	1,033	1,489	1,989	3,017	3,723	3,702	2,225	915	112
2002.....	1,304	1,042	960	995	1,403	2,103	2,833	3,350	3,406	1,944	854	109
2001.....	1,622	1,192	1,100	1,207	1,567	2,209	2,794	3,239	3,220	1,882	748	115
2000.....	1,415	1,152	1,101	1,353	1,687	2,235	2,794	3,286	3,014	1,798	726	115
1999.....	983	908	952	1,140	1,558	1,945	2,468	2,818	2,799	1,592	589	75

All races, male

2003.....	829	682	612	618	813	967	1,193	1,148	919	385	118	83
2002.....	828	652	621	563	761	1,010	1,218	1,103	857	379	100	77
2001.....	1,066	758	699	685	812	1,014	1,170	1,017	760	392	105	78
2000.....	966	743	686	832	876	1,100	1,144	1,078	733	331	100	90
1999.....	650	578	603	665	848	949	981	894	699	306	85	56

All races, female

2003.....	467	395	378	415	676	1,022	1,824	2,575	2,783	1,840	797	29
2002.....	476	390	339	432	642	1,093	1,615	2,247	2,549	1,565	754	32
2001.....	556	434	401	522	755	1,195	1,624	2,222	2,460	1,490	643	37
2000.....	449	409	415	521	811	1,135	1,650	2,208	2,281	1,467	626	25
1999.....	333	330	349	475	710	996	1,487	1,924	2,100	1,286	504	19

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2003.....	985	855	790	846	1,249	1,755	2,733	3,444	3,401	2,063	811	96
2002.....	954	815	783	806	1,212	1,844	2,556	3,077	3,153	1,801	755	83
2001.....	1,286	956	892	975	1,329	1,956	2,509	2,985	2,978	1,732	673	88
2000.....	1,087	907	876	1,072	1,446	1,965	2,509	3,015	2,802	1,680	647	83
1999.....	723	694	753	919	1,335	1,680	2,193	2,582	2,596	1,466	520	61

White, male

2003.....	644	535	486	504	690	861	1,074	1,057	840	354	101	71
2002.....	602	507	509	458	659	871	1,094	1,025	792	347	77	58
2001.....	857	600	558	552	704	914	1,063	933	696	358	95	61
2000.....	744	574	551	658	750	974	1,036	988	687	308	88	66
1999.....	478	437	482	538	731	824	865	807	640	280	79	47

White, female

2003.....	341	320	304	342	559	894	1,659	2,387	2,561	1,709	710	25
2002.....	352	308	274	348	553	973	1,462	2,052	2,361	1,454	678	25
2001.....	429	356	334	423	625	1,042	1,446	2,052	2,282	1,374	578	27
2000.....	343	333	325	414	696	991	1,473	2,027	2,115	1,372	559	17
1999.....	245	257	271	381	604	856	1,328	1,775	1,956	1,186	441	14

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2003.....	290	205	186	171	229	205	240	242	257	135	89	11
2002.....	329	205	157	168	172	230	248	237	214	121	88	21
2001.....	294	203	172	201	207	222	246	214	214	131	65	22
2000.....	298	216	208	256	223	249	260	239	184	95	78	28
1999.....	236	196	182	202	201	239	251	210	183	114	65	11

Black, male

2003.....	176	132	118	105	118	91	95	80	69	23	14	8
2002.....	213	131	101	92	93	124	114	67	53	28	21	14
2001.....	175	139	125	116	94	88	88	71	54	28	10	13
2000.....	203	152	123	159	114	116	99	76	39	14	11	21
1999.....	155	132	113	116	105	114	101	74	54	23	6	7

Black, female

2003.....	114	73	68	66	111	114	145	162	188	112	75	3
2002.....	116	74	56	76	79	106	134	170	161	93	67	7
2001.....	119	64	47	85	113	134	158	143	160	103	55	9
2000.....	95	64	85	97	109	133	161	163	145	81	67	7
1999.....	81	64	69	86	96	125	150	136	129	91	59	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

American Indian 1/, both sexes

2003.....	13	8	8	4	1	5	9	8	12	5	5	2
2002.....	11	8	13	8	11	7	6	4	10	7	3	3
2001.....	13	14	12	15	7	6	10	7	9	5	3	4
2000.....	13	13	6	7	4	7	6	11	8	8	-	1
1999.....	11	7	7	8	12	11	16	9	8	6	-	1

American Indian 1/, male

2003.....	3	6	5	2	-	4	5	3	1	1	1	2
2002.....	5	4	8	5	6	4	2	1	1	1	-	3
2001.....	12	8	3	8	4	3	6	1	1	1	-	3
2000.....	9	8	5	4	3	4	2	4	2	3	-	1
1999.....	10	4	3	5	5	5	10	2	3	2	-	1

American Indian 1/, female

2003.....	10	2	3	2	1	1	4	5	11	4	4	-
2002.....	6	4	5	3	5	3	4	3	9	6	3	-
2001.....	1	6	9	7	3	3	4	6	8	4	3	1
2000.....	4	5	1	3	1	3	4	7	6	5	-	-
1999.....	1	3	4	3	7	6	6	7	5	4	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Asian or Pacific Islander 2/, both sexes

2003.....	8	9	6	12	10	24	35	29	32	22	10	3
2002.....	10	14	7	13	8	22	23	32	29	15	8	2
2001.....	29	19	24	16	24	25	29	33	19	14	7	1
2000.....	17	16	11	18	14	14	19	21	20	15	1	3
1999.....	13	11	10	11	10	15	8	17	12	6	4	2

Asian or Pacific Islander 2/, male

2003.....	6	9	3	7	5	11	19	8	9	7	2	2
2002.....	8	10	3	8	3	11	8	10	11	3	2	2
2001.....	22	11	13	9	10	9	13	12	9	5	-	1
2000.....	10	9	7	11	9	6	7	10	5	6	1	2
1999.....	7	5	5	6	7	6	5	11	2	1	-	1

Asian or Pacific Islander 2/, female

2003.....	2	-	3	5	5	13	16	21	23	15	8	1
2002.....	2	4	4	5	5	11	15	22	18	12	6	-
2001.....	7	8	11	7	14	16	16	21	10	9	7	-
2000.....	7	7	4	7	5	8	12	11	15	9	-	1
1999.....	6	6	5	5	3	9	3	6	10	5	4	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, both sexes

2003.....	8,635	8,940	9,794	11,160	15,315	22,635	30,320	32,746	24,949	10,915	2,490	17
2002.....	7,992	8,431	8,947	10,576	15,113	22,098	29,520	31,671	24,416	10,231	2,256	13
2001.....	7,536	7,549	8,499	10,676	15,127	21,941	27,770	30,312	22,714	9,872	2,134	17
2000.....	7,099	7,152	8,004	10,108	14,899	21,425	26,594	28,517	21,457	9,192	2,126	11
1999.....	6,506	6,676	7,714	10,272	14,893	20,657	24,941	27,390	20,562	8,680	1,967	10

All races, male

2003.....	5,136	5,027	5,247	5,689	7,304	9,807	11,586	10,145	5,900	1,838	295	13
2002.....	4,810	4,675	4,853	5,453	7,214	9,509	11,158	9,835	5,691	1,711	315	11
2001.....	4,427	4,112	4,540	5,480	7,252	9,406	10,411	9,265	5,258	1,740	257	16
2000.....	4,100	3,967	4,269	5,174	7,147	9,237	9,865	8,712	4,966	1,593	265	11
1999.....	3,762	3,762	4,108	5,154	7,178	8,976	9,519	8,208	4,722	1,443	250	5

All races, female

2003.....	3,499	3,913	4,547	5,471	8,011	12,828	18,734	22,601	19,049	9,077	2,195	4
2002.....	3,182	3,756	4,094	5,123	7,899	12,589	18,362	21,836	18,725	8,520	1,941	2
2001.....	3,109	3,437	3,959	5,196	7,875	12,535	17,359	21,047	17,456	8,132	1,877	1
2000.....	2,999	3,185	3,735	4,934	7,752	12,188	16,729	19,805	16,491	7,599	1,861	-
1999.....	2,744	2,914	3,606	5,118	7,715	11,681	15,422	19,182	15,840	7,237	1,717	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2003.....	6,608	7,205	8,002	9,374	13,174	19,995	27,379	30,194	22,976	9,981	2,159	13
2002.....	6,166	6,807	7,303	8,877	12,996	19,545	26,667	29,107	22,602	9,336	1,965	8
2001.....	5,783	6,014	6,915	8,959	13,096	19,411	25,187	27,782	20,985	9,088	1,845	11
2000.....	5,526	5,694	6,488	8,481	12,879	19,016	24,061	26,146	19,763	8,434	1,861	8
1999.....	5,006	5,325	6,302	8,560	12,876	18,283	22,704	25,180	19,125	7,962	1,738	6

White, male

2003.....	3,989	4,092	4,303	4,777	6,287	8,656	10,422	9,364	5,450	1,656	244	9
2002.....	3,788	3,798	3,986	4,578	6,215	8,403	10,075	9,032	5,213	1,526	258	8
2001.....	3,455	3,278	3,736	4,613	6,306	8,308	9,477	8,497	4,817	1,566	208	10
2000.....	3,245	3,198	3,443	4,349	6,184	8,195	8,912	7,958	4,509	1,441	225	8
1999.....	2,951	3,029	3,384	4,320	6,243	7,917	8,649	7,474	4,317	1,287	215	3

White, female

2003.....	2,619	3,113	3,699	4,597	6,887	11,339	16,957	20,830	17,526	8,325	1,915	4
2002.....	2,378	3,009	3,317	4,299	6,781	11,142	16,592	20,075	17,389	7,810	1,707	-
2001.....	2,328	2,736	3,179	4,346	6,790	11,103	15,710	19,285	16,168	7,522	1,637	1
2000.....	2,281	2,496	3,045	4,132	6,695	10,821	15,149	18,188	15,254	6,993	1,636	-
1999.....	2,055	2,296	2,918	4,240	6,633	10,366	14,055	17,706	14,808	6,675	1,523	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2003.....	1,806	1,518	1,559	1,542	1,823	2,246	2,468	2,144	1,681	803	293	4
2002.....	1,645	1,432	1,423	1,466	1,811	2,152	2,454	2,179	1,585	781	255	4
2001.....	1,594	1,356	1,363	1,515	1,761	2,190	2,260	2,172	1,495	688	263	6
2000.....	1,413	1,287	1,338	1,423	1,753	2,096	2,192	2,067	1,474	659	240	3
1999.....	1,335	1,208	1,257	1,513	1,772	2,070	1,926	1,918	1,281	636	199	4

Black, male

2003.....	1,015	806	823	778	882	968	962	632	355	147	40	4
2002.....	911	781	738	743	864	913	907	654	393	140	44	2
2001.....	886	734	696	775	812	959	822	625	349	142	42	6
2000.....	767	695	725	724	837	907	822	622	362	130	35	3
1999.....	727	655	636	730	808	914	729	612	336	130	24	2

Black, female

2003.....	791	712	736	764	941	1,278	1,506	1,512	1,326	656	253	-
2002.....	734	651	685	723	947	1,239	1,547	1,525	1,192	641	211	2
2001.....	708	622	667	740	949	1,231	1,438	1,547	1,146	546	221	-
2000.....	646	592	613	699	916	1,189	1,370	1,445	1,112	529	205	-
1999.....	608	553	621	783	964	1,156	1,197	1,306	945	506	175	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

American Indian 1/, both sexes

2003.....	123	101	88	94	71	88	108	82	58	31	9	-
2002.....	97	76	99	87	92	87	71	88	39	21	5	-
2001.....	84	78	96	76	79	91	77	97	43	20	9	-
2000.....	82	74	60	77	71	74	78	62	43	17	3	-
1999.....	92	76	57	63	71	76	66	58	32	21	12	-

American Indian 1/, male

2003.....	79	55	44	50	33	40	35	22	13	2	2	-
2002.....	58	38	56	48	38	39	20	33	14	6	3	-
2001.....	51	46	47	40	44	27	24	27	14	8	2	-
2000.....	44	31	30	37	34	27	33	24	17	2	2	-
1999.....	52	41	32	26	31	33	25	16	11	5	2	-

American Indian 1/, female

2003.....	44	46	44	44	38	48	73	60	45	29	7	-
2002.....	39	38	43	39	54	48	51	55	25	15	2	-
2001.....	33	32	49	36	35	64	53	70	29	12	7	-
2000.....	38	43	30	40	37	47	45	38	26	15	1	-
1999.....	40	35	25	37	40	43	41	42	21	16	10	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Asian or Pacific Islander 2/, both sexes

2003.....	98	116	145	150	247	306	365	326	234	100	29	-
2002.....	84	116	122	146	214	314	328	297	190	93	31	1
2001.....	75	101	125	126	191	249	246	261	191	76	17	-
2000.....	78	97	118	127	196	239	263	242	177	82	22	-
1999.....	73	67	98	136	174	228	245	234	124	61	18	-

Asian or Pacific Islander 2/, male

2003.....	53	74	77	84	102	143	167	127	82	33	9	-
2002.....	53	58	73	84	97	154	156	116	71	39	10	1
2001.....	35	54	61	52	90	112	88	116	78	24	5	-
2000.....	44	43	71	64	92	108	98	108	78	20	3	-
1999.....	32	37	56	78	96	112	116	106	58	21	9	-

Asian or Pacific Islander 2/, female

2003.....	45	42	68	66	145	163	198	199	152	67	20	-
2002.....	31	58	49	62	117	160	172	181	119	54	21	-
2001.....	40	47	64	74	101	137	158	145	113	52	12	-
2000.....	34	54	47	63	104	131	165	134	99	62	19	-
1999.....	41	30	42	58	78	116	129	128	66	40	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races, both sexes

2003.....	6,972	5,186	3,984	3,711	4,370	5,875	7,233	6,839	4,352	1,620	335	76
2002.....	6,434	4,557	3,788	3,672	4,414	5,938	6,966	6,507	4,255	1,595	294	85
2001.....	5,875	4,159	3,499	3,506	4,329	5,972	6,716	6,266	4,074	1,519	312	120
2000.....	5,408	4,021	3,484	3,375	4,323	5,596	6,162	5,933	3,939	1,419	304	85
1999.....	4,943	3,944	3,341	3,692	4,516	5,772	6,510	6,138	3,954	1,380	257	106

All races, male

2003.....	4,871	3,613	2,711	2,365	2,556	3,251	3,600	2,991	1,595	440	67	70
2002.....	4,552	3,255	2,526	2,369	2,662	3,268	3,419	2,795	1,488	480	54	74
2001.....	4,224	2,872	2,356	2,200	2,551	3,310	3,342	2,730	1,486	381	60	104
2000.....	3,960	2,804	2,329	2,117	2,522	3,039	3,024	2,563	1,391	397	62	83
1999.....	3,606	2,753	2,262	2,336	2,656	3,096	3,276	2,647	1,397	376	60	94

All races, female

2003.....	2,101	1,573	1,273	1,346	1,814	2,624	3,633	3,848	2,757	1,180	268	6
2002.....	1,882	1,302	1,262	1,303	1,752	2,670	3,547	3,712	2,767	1,115	240	11
2001.....	1,651	1,287	1,143	1,306	1,778	2,662	3,374	3,536	2,588	1,138	252	16
2000.....	1,448	1,217	1,155	1,258	1,801	2,557	3,138	3,370	2,548	1,022	242	2
1999.....	1,337	1,191	1,079	1,356	1,860	2,676	3,234	3,491	2,557	1,004	197	12

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2003.....	5,667	4,328	3,373	3,126	3,826	5,311	6,649	6,451	4,088	1,544	307	70
2002.....	5,285	3,789	3,196	3,073	3,894	5,361	6,455	6,120	3,991	1,499	267	78
2001.....	4,814	3,464	2,903	2,995	3,783	5,387	6,201	5,858	3,819	1,419	283	103
2000.....	4,347	3,334	2,941	2,855	3,762	5,039	5,701	5,570	3,693	1,317	272	79
1999.....	3,999	3,248	2,714	3,130	3,897	5,143	5,987	5,696	3,716	1,292	234	89

White, male

2003.....	3,973	3,009	2,299	1,997	2,262	2,940	3,295	2,819	1,484	416	60	64
2002.....	3,715	2,681	2,126	1,965	2,351	2,939	3,151	2,617	1,374	448	45	67
2001.....	3,454	2,371	1,948	1,880	2,239	2,978	3,072	2,542	1,386	348	53	91
2000.....	3,179	2,297	1,945	1,770	2,207	2,712	2,781	2,400	1,290	357	53	77
1999.....	2,917	2,248	1,852	1,975	2,292	2,740	3,018	2,443	1,292	351	54	79

White, female

2003.....	1,694	1,319	1,074	1,129	1,564	2,371	3,354	3,632	2,604	1,128	247	6
2002.....	1,570	1,108	1,070	1,108	1,543	2,422	3,304	3,503	2,617	1,051	222	11
2001.....	1,360	1,093	955	1,115	1,544	2,409	3,129	3,316	2,433	1,071	230	12
2000.....	1,168	1,037	996	1,085	1,555	2,327	2,920	3,170	2,403	960	219	2
1999.....	1,082	1,000	862	1,155	1,605	2,403	2,969	3,253	2,424	941	180	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2003.....	1,061	691	478	455	407	398	414	280	190	59	20	5
2002.....	947	626	467	445	387	436	370	291	197	76	20	4
2001.....	900	576	488	421	431	450	386	299	200	74	25	14
2000.....	879	566	431	417	451	424	354	280	191	80	28	5
1999.....	786	569	497	459	492	506	430	351	200	81	19	16

Black, male

2003.....	745	500	336	302	226	230	207	122	72	17	5	5
2002.....	699	478	330	304	235	256	196	131	82	25	5	4
2001.....	663	421	343	272	245	257	205	124	74	21	6	12
2000.....	658	434	316	285	258	253	185	121	79	32	7	5
1999.....	594	422	333	300	300	285	203	163	80	22	5	14

Black, female

2003.....	316	191	142	153	181	168	207	158	118	42	15	-
2002.....	248	148	137	141	152	180	174	160	115	51	15	-
2001.....	237	155	145	149	186	193	181	175	126	53	19	2
2000.....	221	132	115	132	193	171	169	159	112	48	21	-
1999.....	192	147	164	159	192	221	227	188	120	59	14	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

American Indian 1/, both sexes

2003.....	114	74	54	49	35	35	27	24	15	3	2	1
2002.....	98	66	46	47	32	31	26	7	17	5	-	1
2001.....	80	57	41	33	28	40	23	21	14	7	-	-
2000.....	89	55	39	31	32	29	22	13	6	3	2	1
1999.....	67	52	53	46	36	36	23	16	5	3	2	-

American Indian 1/, male

2003.....	72	46	36	29	22	16	13	10	7	-	1	1
2002.....	71	50	22	36	20	17	13	4	6	1	-	1
2001.....	60	42	26	17	16	22	12	9	4	1	-	-
2000.....	70	34	22	17	19	20	12	9	2	1	-	1
1999.....	47	42	33	29	22	18	11	7	1	1	1	-

American Indian 1/, female

2003.....	42	28	18	20	13	19	14	14	8	3	1	-
2002.....	27	16	24	11	12	14	13	3	11	4	-	-
2001.....	20	15	15	16	12	18	11	12	10	6	-	-
2000.....	19	21	17	14	13	9	10	4	4	2	2	-
1999.....	20	10	20	17	14	18	12	9	4	2	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Asian or Pacific Islander 2/, both sexes

2003.....	130	93	79	81	102	131	143	84	59	14	6	-
2002.....	104	76	79	107	101	110	115	89	50	15	7	2
2001.....	81	62	67	57	87	95	106	88	41	19	4	3
2000.....	93	66	73	72	78	104	85	70	49	19	2	-
1999.....	91	75	77	57	91	87	70	75	33	4	2	1

Asian or Pacific Islander 2/, male

2003.....	81	58	40	37	46	65	85	40	32	7	1	-
2002.....	67	46	48	64	56	56	59	43	26	6	4	2
2001.....	47	38	39	31	51	53	53	55	22	11	1	1
2000.....	53	39	46	45	38	54	46	33	20	7	2	-
1999.....	48	41	44	32	42	53	44	34	24	2	-	1

Asian or Pacific Islander 2/, female

2003.....	49	35	39	44	56	66	58	44	27	7	5	-
2002.....	37	30	31	43	45	54	56	46	24	9	3	-
2001.....	34	24	28	26	36	42	53	33	19	8	3	2
2000.....	40	27	27	27	40	50	39	37	29	12	-	-
1999.....	43	34	33	25	49	34	26	41	9	2	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

All races, both sexes

2003.....	3,098	2,507	1,934	1,578	1,632	1,782	1,595	994	350	55	2	20
2002.....	2,970	2,229	1,936	1,597	1,754	1,855	1,578	936	330	47	9	39
2001.....	2,872	2,152	1,709	1,567	1,715	1,943	1,564	903	304	63	2	43
2000.....	2,717	2,115	1,768	1,599	1,716	1,860	1,482	985	298	49	3	33
1999.....	2,524	2,131	1,681	1,727	1,852	1,902	1,566	964	311	50	5	37

All races, male

2003.....	2,198	1,778	1,309	1,019	956	1,019	866	534	187	30	-	18
2002.....	2,089	1,615	1,262	1,039	1,062	1,060	857	527	180	26	4	36
2001.....	2,032	1,467	1,164	974	1,041	1,106	877	492	163	28	1	37
2000.....	1,922	1,456	1,161	996	1,025	1,050	808	539	167	27	1	33
1999.....	1,782	1,454	1,107	1,091	1,071	1,089	870	551	179	25	3	34

All races, female

2003.....	900	729	625	559	676	763	729	460	163	25	2	2
2002.....	881	614	674	558	692	795	721	409	150	21	5	3
2001.....	840	685	545	593	674	837	687	411	141	35	1	6
2000.....	795	659	607	603	691	810	674	446	131	22	2	-
1999.....	742	677	574	636	781	813	696	413	132	25	2	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

White, both sexes

2003.....	2,564	2,109	1,644	1,338	1,439	1,586	1,462	924	322	51	1	19
2002.....	2,494	1,900	1,658	1,356	1,565	1,684	1,453	870	309	44	6	34
2001.....	2,432	1,834	1,443	1,346	1,511	1,767	1,450	843	287	58	2	35
2000.....	2,245	1,796	1,523	1,375	1,503	1,683	1,377	914	275	44	3	31
1999.....	2,102	1,802	1,412	1,497	1,637	1,716	1,438	888	281	45	4	28

White, male

2003.....	1,832	1,504	1,129	873	860	908	792	495	168	27	-	17
2002.....	1,753	1,371	1,087	875	943	949	780	484	163	25	4	31
2001.....	1,735	1,246	984	836	921	997	817	453	155	27	1	31
2000.....	1,598	1,226	987	845	899	942	743	503	154	25	1	31
1999.....	1,494	1,230	931	950	946	977	796	503	163	20	2	26

White, female

2003.....	732	605	515	465	579	678	670	429	154	24	1	2
2002.....	741	529	571	481	622	735	673	386	146	19	2	3
2001.....	697	588	459	510	590	770	633	390	132	31	1	4
2000.....	647	570	536	530	604	741	634	411	121	19	2	-
1999.....	608	572	481	547	691	739	642	385	118	25	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

Black, both sexes

2003.....	394	299	211	176	135	134	86	48	22	2	-	1
2002.....	346	251	200	163	126	117	92	44	13	2	3	3
2001.....	338	252	208	172	141	127	73	38	14	5	-	8
2000.....	361	244	180	165	160	109	74	50	16	3	-	1
1999.....	323	246	190	163	155	138	98	59	22	5	1	9

Black, male

2003.....	286	216	143	116	73	82	51	27	13	1	-	1
2002.....	253	195	138	115	85	83	58	32	10	1	-	3
2001.....	237	183	152	111	78	84	39	23	7	1	-	6
2000.....	258	188	133	120	100	73	47	27	10	2	-	1
1999.....	236	173	134	105	93	85	56	41	11	5	1	8

Black, female

2003.....	108	83	68	60	62	52	35	21	9	1	-	-
2002.....	93	56	62	48	41	34	34	12	3	1	3	-
2001.....	101	69	56	61	63	43	34	15	7	4	-	2
2000.....	103	56	47	45	60	36	27	23	6	1	-	-
1999.....	87	73	56	58	62	53	42	18	11	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

American Indian 1/, both sexes

2003.....	56	38	27	19	17	9	7	8	1	-	-	-
2002.....	63	31	32	20	15	11	7	2	2	1	-	-
2001.....	44	34	16	20	11	16	9	4	1	-	-	-
2000.....	43	26	24	19	15	14	5	3	2	1	-	1
1999.....	39	36	23	27	17	14	9	1	-	-	-	-

American Indian 1/, male

2003.....	34	23	14	10	7	3	2	3	1	-	-	-
2002.....	46	23	14	15	8	8	6	1	2	-	-	-
2001.....	32	23	9	11	7	9	6	2	-	-	-	-
2000.....	32	12	13	9	10	9	4	2	1	-	-	1
1999.....	27	28	14	14	12	9	5	-	-	-	-	-

American Indian 1/, female

2003.....	22	15	13	9	10	6	5	5	-	-	-	-
2002.....	17	8	18	5	7	3	1	1	-	1	-	-
2001.....	12	11	7	9	4	7	3	2	1	-	-	-
2000.....	11	14	11	10	5	5	1	1	1	1	-	-
1999.....	12	8	9	13	5	5	4	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

Asian or Pacific Islander 2/, both sexes

2003.....	84	61	52	45	41	53	40	14	5	2	1	-
2002.....	67	47	46	58	48	43	26	20	6	-	-	2
2001.....	58	32	42	29	52	33	32	18	2	-	-	-
2000.....	68	49	41	40	38	54	26	18	5	1	-	-
1999.....	60	47	56	40	43	34	21	16	8	-	-	-

Asian or Pacific Islander 2/, male

2003.....	46	35	23	20	16	26	21	9	5	2	-	-
2002.....	37	26	23	34	26	20	13	10	5	-	-	2
2001.....	28	15	19	16	35	16	15	14	1	-	-	-
2000.....	34	30	28	22	16	26	14	7	2	-	-	-
1999.....	25	23	28	22	20	18	13	7	5	-	-	-

Asian or Pacific Islander 2/, female

2003.....	38	26	29	25	25	27	19	5	-	-	1	-
2002.....	30	21	23	24	22	23	13	10	1	-	-	-
2001.....	30	17	23	13	17	17	17	4	1	-	-	-
2000.....	34	19	13	18	22	28	12	11	3	1	-	-
1999.....	35	24	28	18	23	16	8	9	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races, both sexes

2003.....	2,788	2,232	1,726	1,444	1,523	1,680	1,530	961	343	53	1	16
2002.....	2,697	2,004	1,760	1,469	1,644	1,763	1,508	909	321	47	8	35
2001.....	2,553	1,939	1,526	1,409	1,581	1,823	1,489	868	294	59	2	33
2000.....	2,425	1,917	1,589	1,436	1,602	1,764	1,409	953	285	47	3	26
1999.....	2,225	1,885	1,485	1,565	1,711	1,772	1,471	920	291	46	4	30

All races, male

2003.....	1,937	1,553	1,130	908	870	944	823	511	182	29	-	14
2002.....	1,861	1,423	1,117	923	965	992	801	512	176	26	4	33
2001.....	1,774	1,296	1,013	846	940	1,027	832	478	157	28	1	27
2000.....	1,689	1,293	1,013	870	932	980	759	517	161	27	1	26
1999.....	1,537	1,250	950	969	962	1,012	815	524	164	24	2	27

All races, female

2003.....	851	679	596	536	653	736	707	450	161	24	1	2
2002.....	836	581	643	546	679	771	707	397	145	21	4	2
2001.....	779	643	513	563	641	796	657	390	137	31	1	6
2000.....	736	624	576	566	670	784	650	436	124	20	2	-
1999.....	688	635	535	596	749	760	656	396	127	22	2	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2003.....	2,295	1,861	1,458	1,226	1,338	1,497	1,404	892	316	49	-	15
2002.....	2,256	1,708	1,501	1,248	1,466	1,605	1,391	846	300	44	5	30
2001.....	2,153	1,654	1,282	1,213	1,386	1,661	1,382	813	279	55	2	29
2000.....	1,989	1,622	1,362	1,234	1,402	1,598	1,308	885	262	43	3	24
1999.....	1,847	1,587	1,244	1,356	1,512	1,603	1,355	849	263	41	4	22

White, male

2003.....	1,607	1,302	968	781	779	841	755	473	164	26	-	13
2002.....	1,557	1,210	957	777	857	894	730	471	159	25	4	28
2001.....	1,509	1,106	846	730	828	925	776	441	150	27	1	25
2000.....	1,392	1,085	852	736	816	879	696	482	148	25	1	24
1999.....	1,284	1,051	798	841	851	907	748	480	150	19	2	20

White, female

2003.....	688	559	490	445	559	656	649	419	152	23	-	2
2002.....	699	498	544	471	609	711	661	375	141	19	1	2
2001.....	644	548	436	483	558	736	606	372	129	28	1	4
2000.....	597	537	510	498	586	719	612	403	114	18	2	-
1999.....	563	536	446	515	661	696	607	369	113	22	2	2

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2003.....	364	277	194	161	130	125	81	48	21	2	-	1
2002.....	318	224	187	150	118	107	86	43	13	2	3	3
2001.....	308	226	189	153	136	116	70	37	13	4	-	4
2000.....	342	225	169	150	151	102	71	47	16	2	-	1
1999.....	288	224	169	150	145	128	90	54	21	5	-	8

Black, male

2003.....	258	197	127	102	70	77	47	27	12	1	-	1
2002.....	227	170	126	102	77	73	53	31	10	1	-	3
2001.....	214	158	140	94	73	78	38	23	7	1	-	2
2000.....	242	171	124	107	94	68	46	26	10	2	-	1
1999.....	207	155	116	96	84	82	52	37	10	5	-	7

Black, female

2003.....	106	80	67	59	60	48	34	21	9	1	-	-
2002.....	91	54	61	48	41	34	33	12	3	1	3	-
2001.....	94	68	49	59	63	38	32	14	6	3	-	2
2000.....	100	54	45	43	57	34	25	21	6	-	-	-
1999.....	81	69	53	54	61	46	38	17	11	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

American Indian 1/, both sexes

2003.....	53	35	26	17	15	9	7	7	1	-	-	-
2002.....	62	30	32	18	13	11	7	2	2	1	-	-
2001.....	38	29	15	18	9	16	7	4	1	-	-	-
2000.....	37	24	23	17	14	13	5	3	2	1	-	1
1999.....	35	30	21	23	14	14	9	1	-	-	-	-

American Indian 1/, male

2003.....	33	21	14	8	5	3	2	2	1	-	-	-
2002.....	45	22	14	13	6	8	6	1	2	-	-	-
2001.....	27	18	8	9	6	9	4	2	-	-	-	-
2000.....	26	10	13	9	9	8	4	2	1	-	-	1
1999.....	25	23	13	12	9	9	5	-	-	-	-	-

American Indian 1/, female

2003.....	20	14	12	9	10	6	5	5	-	-	-	-
2002.....	17	8	18	5	7	3	1	1	-	1	-	-
2001.....	11	11	7	9	3	7	3	2	1	-	-	-
2000.....	11	14	10	8	5	5	1	1	1	1	-	-
1999.....	10	7	8	11	5	5	4	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Asian or Pacific Islander 2/, both sexes

2003.....	76	59	48	40	40	49	38	14	5	2	1	-
2002.....	61	42	40	53	47	40	24	18	6	-	-	2
2001.....	54	30	40	25	50	30	30	14	1	-	-	-
2000.....	57	46	35	35	35	51	25	18	5	1	-	-
1999.....	55	44	51	36	40	27	17	16	7	-	-	-

Asian or Pacific Islander 2/, male

2003.....	39	33	21	17	16	23	19	9	5	2	-	-
2002.....	32	21	20	31	25	17	12	9	5	-	-	2
2001.....	24	14	19	13	33	15	14	12	-	-	-	-
2000.....	29	27	24	18	13	25	13	7	2	-	-	-
1999.....	21	21	23	20	18	14	10	7	4	-	-	-

Asian or Pacific Islander 2/, female

2003.....	37	26	27	23	24	26	19	5	-	-	1	-
2002.....	29	21	20	22	22	23	12	9	1	-	-	-
2001.....	30	16	21	12	17	15	16	2	1	-	-	-
2000.....	28	19	11	17	22	26	12	11	3	1	-	-
1999.....	34	23	28	16	22	13	7	9	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races, both sexes

2003.....	97	87	60	42	39	42	38	18	5	1	1	4
2002.....	90	61	48	33	37	35	31	11	5	-	1	4
2001.....	105	67	66	64	53	59	39	20	7	2	-	10
2000.....	109	64	58	52	48	51	40	22	12	1	-	7
1999.....	125	101	81	79	66	92	67	34	16	3	1	7

All races, male

2003.....	78	68	45	35	28	25	27	11	4	1	-	4
2002.....	78	53	39	26	30	25	25	6	2	-	-	3
2001.....	84	56	48	50	34	38	18	6	4	-	-	10
2000.....	74	48	50	38	37	33	24	14	5	-	-	7
1999.....	99	77	58	52	45	51	37	20	13	1	1	7

All races, female

2003.....	19	19	15	7	11	17	11	7	1	-	1	-
2002.....	12	8	9	7	7	10	6	5	3	-	1	1
2001.....	21	11	18	14	19	21	21	14	3	2	-	-
2000.....	35	16	8	14	11	18	16	8	7	1	-	-
1999.....	26	24	23	27	21	41	30	14	3	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2003.....	77	74	52	34	33	37	33	17	4	1	1	4
2002.....	76	45	38	25	32	28	28	9	5	-	1	4
2001.....	88	56	51	48	49	52	34	17	5	2	-	6
2000.....	92	50	51	43	42	44	36	20	12	-	-	7
1999.....	104	90	71	66	57	79	60	31	14	3	-	6

White, male

2003.....	62	57	38	28	24	22	23	10	3	1	-	4
2002.....	65	39	32	20	25	18	23	5	2	-	-	3
2001.....	72	46	42	37	30	35	15	4	3	-	-	6
2000.....	62	36	46	30	32	30	22	14	5	-	-	7
1999.....	82	69	50	46	38	45	33	18	11	1	-	6

White, female

2003.....	15	17	14	6	9	15	10	7	1	-	1	-
2002.....	11	6	6	5	7	10	5	4	3	-	1	1
2001.....	16	10	9	11	19	17	19	13	2	2	-	-
2000.....	30	14	5	13	10	14	14	6	7	-	-	-
1999.....	22	21	21	20	19	34	27	13	3	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Black, both sexes

2003.....	17	12	7	6	4	4	3	-	1	-	-	-
2002.....	13	12	6	5	4	5	3	-	-	-	-	-
2001.....	13	9	13	12	3	5	1	-	1	-	-	4
2000.....	10	11	4	7	3	4	3	2	-	1	-	-
1999.....	18	11	6	8	6	8	3	3	1	-	1	1

Black, male

2003.....	15	10	7	6	3	2	2	-	1	-	-	-
2002.....	12	10	6	5	4	5	2	-	-	-	-	-
2001.....	9	8	6	10	3	3	-	-	-	-	-	4
2000.....	7	9	3	7	2	2	1	-	-	-	-	-
1999.....	15	8	4	4	5	2	1	2	1	-	1	1

Black, female

2003.....	2	2	-	-	1	2	1	-	-	-	-	-
2002.....	1	2	-	-	-	-	1	-	-	-	-	-
2001.....	4	1	7	2	-	2	1	-	1	-	-	-
2000.....	3	2	1	-	1	2	2	2	-	1	-	-
1999.....	3	3	2	4	1	6	2	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Asian or Pacific Islander 2/, both sexes

2003.....	2	-	1	2	1	1	2	-	-	-	-	-
2002.....	1	3	4	2	-	2	-	2	-	-	-	-
2001.....	2	-	2	3	1	2	2	3	1	-	-	-
2000.....	4	2	2	1	3	3	1	-	-	-	-	-
1999.....	2	-	3	2	2	5	4	-	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	1	-	-	1	-	1	2	-	-	-	-	-
2002.....	1	3	1	-	-	2	-	1	-	-	-	-
2001.....	2	-	-	2	1	-	1	2	1	-	-	-
2000.....	2	2	1	1	3	1	1	-	-	-	-	-
1999.....	2	-	3	-	1	4	3	-	1	-	-	-

Asian or Pacific Islander 2/, female

2003.....	1	-	1	1	1	-	-	-	-	-	-	-
2002.....	-	-	3	2	-	-	-	1	-	-	-	-
2001.....	-	-	2	1	-	2	1	1	-	-	-	-
2000.....	2	-	1	-	-	2	-	-	-	-	-	-
1999.....	-	-	-	2	1	1	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races, both sexes

2003.....	213	188	148	92	70	60	27	15	2	1	-	-
2002.....	183	164	128	95	73	57	39	16	4	-	-	-
2001.....	214	146	117	94	81	61	36	15	3	2	-	-
2000.....	183	134	121	111	66	45	33	10	1	1	-	-
1999.....	174	145	115	83	75	38	28	10	4	1	-	-

All races, male

2003.....	183	157	134	76	58	50	16	12	1	-	-	-
2002.....	150	139	106	90	67	43	31	9	2	-	-	-
2001.....	174	115	103	78	67	41	27	8	2	-	-	-
2000.....	159	115	98	88	56	37	25	8	1	-	-	-
1999.....	146	127	99	70	64	26	18	7	2	-	-	-

All races, female

2003.....	30	31	14	16	12	10	11	3	1	1	-	-
2002.....	33	25	22	5	6	14	8	7	2	-	-	-
2001.....	40	31	14	16	14	20	9	7	1	2	-	-
2000.....	24	19	23	23	10	8	8	2	-	1	-	-
1999.....	28	18	16	13	11	12	10	3	2	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2003.....	192	174	134	78	68	52	25	15	2	1	-	-
2002.....	162	147	119	83	67	51	34	15	4	-	-	-
2001.....	191	124	110	85	76	54	34	13	3	1	-	-
2000.....	164	124	110	98	59	41	33	9	1	1	-	-
1999.....	151	125	97	75	68	34	23	8	4	1	-	-

White, male

2003.....	163	145	123	64	57	45	14	12	1	-	-	-
2002.....	131	122	98	78	61	37	27	8	2	-	-	-
2001.....	154	94	96	69	63	37	26	8	2	-	-	-
2000.....	144	105	89	79	51	33	25	7	1	-	-	-
1999.....	128	110	83	63	57	25	15	5	2	-	-	-

White, female

2003.....	29	29	11	14	11	7	11	3	1	1	-	-
2002.....	31	25	21	5	6	14	7	7	2	-	-	-
2001.....	37	30	14	16	13	17	8	5	1	1	-	-
2000.....	20	19	21	19	8	8	8	2	-	1	-	-
1999.....	23	15	14	12	11	9	8	3	2	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Black, both sexes

2003.....	13	10	10	9	1	5	2	-	-	-	-	-
2002.....	15	15	7	8	4	5	3	1	-	-	-	-
2001.....	17	17	6	7	2	6	2	1	-	1	-	-
2000.....	9	8	7	8	6	3	-	1	-	-	-	-
1999.....	17	11	15	5	4	2	5	2	-	-	-	-

Black, male

2003.....	13	9	9	8	-	3	2	-	-	-	-	-
2002.....	14	15	6	8	4	5	3	1	-	-	-	-
2001.....	14	17	6	7	2	3	1	-	-	-	-	-
2000.....	9	8	6	6	4	3	-	1	-	-	-	-
1999.....	14	10	14	5	4	1	3	2	-	-	-	-

Black, female

2003.....	-	1	1	1	1	2	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	3	1	1	-	1	-	-
2000.....	-	-	1	2	2	-	-	-	-	-	-	-
1999.....	3	1	1	-	-	1	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Asian or Pacific Islander 2/, both sexes

2003.....	6	2	3	3	-	3	-	-	-	-	-	-
2002.....	5	2	2	3	1	1	2	-	-	-	-	-
2001.....	2	2	-	1	1	1	-	1	-	-	-	-
2000.....	7	1	4	4	-	-	-	-	-	-	-	-
1999.....	3	3	2	2	1	2	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	6	2	2	2	-	2	-	-	-	-	-	-
2002.....	4	2	2	3	1	1	1	-	-	-	-	-
2001.....	2	1	-	1	1	1	-	-	-	-	-	-
2000.....	3	1	3	3	-	-	-	-	-	-	-	-
1999.....	2	2	2	2	1	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	-	1	1	-	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	4	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	1	-	-	-	2	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

All races, both sexes

2003.....	3,874	2,679	2,050	2,133	2,738	4,093	5,638	5,845	4,002	1,565	333	56
2002.....	3,464	2,328	1,852	2,075	2,660	4,083	5,388	5,571	3,925	1,548	285	46
2001.....	3,003	2,007	1,790	1,939	2,614	4,029	5,152	5,363	3,770	1,456	310	77
2000.....	2,691	1,906	1,716	1,776	2,607	3,736	4,680	4,948	3,641	1,370	301	52
1999.....	2,419	1,813	1,660	1,965	2,664	3,870	4,944	5,174	3,643	1,330	252	69

All races, male

2003.....	2,673	1,835	1,402	1,346	1,600	2,232	2,734	2,457	1,408	410	67	52
2002.....	2,463	1,640	1,264	1,330	1,600	2,208	2,562	2,268	1,308	454	50	38
2001.....	2,192	1,405	1,192	1,226	1,510	2,204	2,465	2,238	1,323	353	59	67
2000.....	2,038	1,348	1,168	1,121	1,497	1,989	2,216	2,024	1,224	370	61	50
1999.....	1,824	1,299	1,155	1,245	1,585	2,007	2,406	2,096	1,218	351	57	60

All races, female

2003.....	1,201	844	648	787	1,138	1,861	2,904	3,388	2,594	1,155	266	4
2002.....	1,001	688	588	745	1,060	1,875	2,826	3,303	2,617	1,094	235	8
2001.....	811	602	598	713	1,104	1,825	2,687	3,125	2,447	1,103	251	10
2000.....	653	558	548	655	1,110	1,747	2,464	2,924	2,417	1,000	240	2
1999.....	595	514	505	720	1,079	1,863	2,538	3,078	2,425	979	195	9

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

White, both sexes

2003.....	3,103	2,219	1,729	1,788	2,387	3,725	5,187	5,527	3,766	1,493	306	51
2002.....	2,791	1,889	1,538	1,717	2,329	3,677	5,002	5,250	3,682	1,455	261	44
2001.....	2,382	1,630	1,460	1,649	2,272	3,620	4,751	5,015	3,532	1,361	281	68
2000.....	2,102	1,538	1,418	1,480	2,259	3,356	4,324	4,656	3,418	1,273	269	48
1999.....	1,897	1,446	1,302	1,633	2,260	3,427	4,549	4,808	3,435	1,247	230	61

White, male

2003.....	2,141	1,505	1,170	1,124	1,402	2,032	2,503	2,324	1,316	389	60	47
2002.....	1,962	1,310	1,039	1,090	1,408	1,990	2,371	2,133	1,211	423	41	36
2001.....	1,719	1,125	964	1,044	1,318	1,981	2,255	2,089	1,231	321	52	60
2000.....	1,581	1,071	958	925	1,308	1,770	2,038	1,897	1,136	332	52	46
1999.....	1,423	1,018	921	1,025	1,346	1,763	2,222	1,940	1,129	331	52	53

White, female

2003.....	962	714	559	664	985	1,693	2,684	3,203	2,450	1,104	246	4
2002.....	829	579	499	627	921	1,687	2,631	3,117	2,471	1,032	220	8
2001.....	663	505	496	605	954	1,639	2,496	2,926	2,301	1,040	229	8
2000.....	521	467	460	555	951	1,586	2,286	2,759	2,282	941	217	2
1999.....	474	428	381	608	914	1,664	2,327	2,868	2,306	916	178	8

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

Black, both sexes

2003.....	667	392	267	279	272	264	328	232	168	57	20	4
2002.....	601	375	267	282	261	319	278	247	184	74	17	1
2001.....	562	324	280	249	290	323	313	261	186	69	25	6
2000.....	518	322	251	252	291	315	280	230	175	77	28	4
1999.....	463	323	307	296	337	368	332	292	178	76	18	7

Black, male

2003.....	459	284	193	186	153	148	156	95	59	16	5	4
2002.....	446	283	192	189	150	173	138	99	72	24	5	1
2001.....	426	238	191	161	167	173	166	101	67	20	6	6
2000.....	400	246	183	165	158	180	138	94	69	30	7	4
1999.....	358	249	199	195	207	200	147	122	69	17	4	6

Black, female

2003.....	208	108	74	93	119	116	172	137	109	41	15	-
2002.....	155	92	75	93	111	146	140	148	112	50	12	-
2001.....	136	86	89	88	123	150	147	160	119	49	19	-
2000.....	118	76	68	87	133	135	142	136	106	47	21	-
1999.....	105	74	108	101	130	168	185	170	109	59	14	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

American Indian 1/, both sexes

2003.....	58	36	27	30	18	26	20	16	14	3	2	1
2002.....	35	35	14	27	17	20	19	5	15	4	-	1
2001.....	36	23	25	13	17	24	14	17	13	7	-	-
2000.....	46	29	15	12	17	15	17	10	4	2	2	-
1999.....	28	16	30	19	19	22	14	15	5	3	2	-

American Indian 1/, male

2003.....	38	23	22	19	15	13	11	7	6	-	1	1
2002.....	25	27	8	21	12	9	7	3	4	1	-	1
2001.....	28	19	17	6	9	13	6	7	4	1	-	-
2000.....	38	22	9	8	9	11	8	7	1	1	-	-
1999.....	20	14	19	15	10	9	6	7	1	1	1	-

American Indian 1/, female

2003.....	20	13	5	11	3	13	9	9	8	3	1	-
2002.....	10	8	6	6	5	11	12	2	11	3	-	-
2001.....	8	4	8	7	8	11	8	10	9	6	-	-
2000.....	8	7	6	4	8	4	9	3	3	1	2	-
1999.....	8	2	11	4	9	13	8	8	4	2	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

Asian or Pacific Islander 2/, both sexes

2003.....	46	32	27	36	61	78	103	70	54	12	5	-
2002.....	37	29	33	49	53	67	89	69	44	15	7	-
2001.....	23	30	25	28	35	62	74	70	39	19	4	3
2000.....	25	17	32	32	40	50	59	52	44	18	2	-
1999.....	31	28	21	17	48	53	49	59	25	4	2	1

Asian or Pacific Islander 2/, male

2003.....	35	23	17	17	30	39	64	31	27	5	1	-
2002.....	30	20	25	30	30	36	46	33	21	6	4	-
2001.....	19	23	20	15	16	37	38	41	21	11	1	1
2000.....	19	9	18	23	22	28	32	26	18	7	2	-
1999.....	23	18	16	10	22	35	31	27	19	2	-	1

Asian or Pacific Islander 2/, female

2003.....	11	9	10	19	31	39	39	39	27	7	4	-
2002.....	7	9	8	19	23	31	43	36	23	9	3	-
2001.....	4	7	5	13	19	25	36	29	18	8	3	2
2000.....	6	8	14	9	18	22	27	26	26	11	-	-
1999.....	8	10	5	7	26	18	18	32	6	2	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Falls (W00-W19)

All races, both sexes

2003.....	535	579	641	787	1,261	2,136	3,113	3,220	2,221	806	157	1
2002.....	530	515	574	759	1,208	2,011	2,869	3,032	2,045	784	129	1
2001.....	504	463	541	747	1,086	1,873	2,567	2,647	1,850	718	135	4
2000.....	462	448	501	595	1,065	1,641	2,200	2,352	1,682	618	120	-
1999.....	377	398	489	663	1,000	1,636	2,220	2,308	1,625	533	112	4

All races, male

2003.....	409	408	419	485	724	1,145	1,516	1,359	796	206	25	1
2002.....	394	379	392	474	728	1,079	1,365	1,262	691	231	24	-
2001.....	395	344	374	469	621	1,041	1,236	1,149	655	188	31	3
2000.....	358	326	354	378	613	895	1,052	992	592	173	27	-
1999.....	295	296	354	433	594	840	1,110	964	568	140	29	4

All races, female

2003.....	126	171	222	302	537	991	1,597	1,861	1,425	600	132	-
2002.....	136	136	182	285	480	932	1,504	1,770	1,354	553	105	1
2001.....	109	119	167	278	465	832	1,331	1,498	1,195	530	104	1
2000.....	104	122	147	217	452	746	1,148	1,360	1,090	445	93	-
1999.....	82	102	135	230	406	796	1,110	1,344	1,057	393	83	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Falls (W00-W19)

White, both sexes

2003.....	429	493	558	687	1,139	1,994	2,924	3,092	2,120	784	150	-
2002.....	448	437	507	669	1,096	1,874	2,739	2,910	1,942	754	122	1
2001.....	431	391	468	660	993	1,742	2,439	2,521	1,767	687	125	4
2000.....	377	374	428	514	966	1,539	2,088	2,262	1,613	584	111	-
1999.....	301	324	398	575	894	1,518	2,114	2,206	1,551	506	103	4

White, male

2003.....	326	349	364	416	659	1,073	1,420	1,306	757	200	25	-
2002.....	329	321	345	416	663	1,004	1,297	1,205	646	220	22	-
2001.....	334	292	322	414	571	967	1,167	1,088	622	172	27	3
2000.....	289	269	305	326	563	833	997	951	559	159	25	-
1999.....	237	239	291	372	532	771	1,057	915	535	131	28	4

White, female

2003.....	103	144	194	271	480	921	1,504	1,786	1,363	584	125	-
2002.....	119	116	162	253	433	870	1,442	1,705	1,296	534	100	1
2001.....	97	99	146	246	422	775	1,272	1,433	1,145	515	98	1
2000.....	88	105	123	188	403	706	1,091	1,311	1,054	425	86	-
1999.....	64	85	107	203	362	747	1,057	1,291	1,016	375	75	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Falls (W00-W19)

Black, both sexes

2003.....	77	68	65	72	80	79	111	74	59	14	4	-
2002.....	68	57	54	57	81	86	69	72	67	22	5	-
2001.....	66	56	55	73	72	94	81	71	51	16	7	-
2000.....	67	60	56	64	73	65	72	50	46	21	7	-
1999.....	64	59	72	81	70	75	80	59	55	21	7	-

Black, male

2003.....	57	46	42	52	37	40	48	32	19	3	-	-
2002.....	54	42	37	37	46	48	37	34	29	8	1	-
2001.....	55	41	38	47	40	52	45	30	19	8	3	-
2000.....	54	47	42	41	37	39	31	20	23	9	1	-
1999.....	49	45	48	57	43	43	36	23	20	6	1	-

Black, female

2003.....	20	22	23	20	43	39	63	42	40	11	4	-
2002.....	14	15	17	20	35	38	32	38	38	14	4	-
2001.....	11	15	17	26	32	42	36	41	32	8	4	-
2000.....	13	13	14	23	36	26	41	30	23	12	6	-
1999.....	15	14	24	24	27	32	44	36	35	15	6	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Falls (W00-W19)

American Indian 1/, both sexes

2003.....	15	8	8	8	8	11	10	9	8	2	1	1
2002.....	6	8	3	7	4	9	12	5	7	1	-	-
2001.....	2	6	6	3	5	8	4	10	7	4	-	-
2000.....	11	7	5	-	6	9	3	6	2	2	1	-
1999.....	5	3	9	1	9	5	3	8	3	3	1	-

American Indian 1/, male

2003.....	12	5	8	7	7	4	7	4	3	-	-	1
2002.....	3	7	2	4	2	3	4	3	-	-	-	-
2001.....	2	5	5	1	2	4	2	2	2	1	-	-
2000.....	10	6	1	-	2	8	2	5	1	1	-	-
1999.....	3	2	6	-	4	1	-	4	1	1	-	-

American Indian 1/, female

2003.....	3	3	-	1	1	7	3	5	5	2	1	-
2002.....	3	1	1	3	2	6	8	2	7	1	-	-
2001.....	-	1	1	2	3	4	2	8	5	3	-	-
2000.....	1	1	4	-	4	1	1	1	1	1	1	-
1999.....	2	1	3	1	5	4	3	4	2	2	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Falls (W00-W19)

Asian or Pacific Islander 2/, both sexes

2003.....	14	10	10	20	34	52	68	45	34	6	2	-
2002.....	8	13	10	26	27	42	49	45	29	7	2	-
2001.....	5	10	12	11	16	29	43	45	25	11	3	-
2000.....	7	7	12	17	20	28	37	34	21	11	1	-
1999.....	7	12	10	6	27	38	23	35	16	3	1	-

Asian or Pacific Islander 2/, male

2003.....	14	8	5	10	21	28	41	17	17	3	-	-
2002.....	8	9	8	17	17	24	27	20	16	3	1	-
2001.....	4	6	9	7	8	18	22	29	12	7	1	-
2000.....	5	4	6	11	11	15	22	16	9	4	1	-
1999.....	6	10	9	4	15	25	17	22	12	2	-	-

Asian or Pacific Islander 2/, female

2003.....	-	2	5	10	13	24	27	28	17	3	2	-
2002.....	-	4	2	9	10	18	22	25	13	4	1	-
2001.....	1	4	3	4	8	11	21	16	13	4	2	-
2000.....	2	3	6	6	9	13	15	18	12	7	-	-
1999.....	1	2	1	2	12	13	6	13	4	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental discharge of firearms (W32-W34)

All races, both sexes

2003.....	44	36	26	23	16	16	21	3	3	-	-	-
2002.....	43	36	20	17	19	18	14	5	1	1	-	-
2001.....	42	44	25	19	14	24	17	6	2	-	-	2
2000.....	40	36	14	24	17	17	7	7	4	-	-	-
1999.....	41	25	23	15	18	15	8	8	2	-	-	-

All races, male

2003.....	35	35	24	20	10	15	18	3	3	-	-	-
2002.....	39	32	18	15	17	15	11	4	1	1	-	-
2001.....	36	34	21	16	10	22	14	5	2	-	-	1
2000.....	38	30	14	20	14	16	7	7	4	-	-	-
1999.....	32	22	19	13	18	13	7	5	2	-	-	-

All races, female

2003.....	9	1	2	3	6	1	3	-	-	-	-	-
2002.....	4	4	2	2	2	3	3	1	-	-	-	-
2001.....	6	10	4	3	4	2	3	1	-	-	-	1
2000.....	2	6	-	4	3	1	-	-	-	-	-	-
1999.....	9	3	4	2	-	2	1	3	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental discharge of firearms (W32-W34)												
White, both sexes												
2003.....	41	34	25	22	12	16	17	2	2	-	-	-
2002.....	37	32	18	16	19	14	14	5	1	1	-	-
2001.....	41	38	24	18	12	23	15	6	2	-	-	2
2000.....	37	34	13	22	15	16	6	7	4	-	-	-
1999.....	36	21	20	13	17	13	7	8	2	-	-	-
White, male												
2003.....	33	34	23	19	8	15	14	2	2	-	-	-
2002.....	34	29	17	14	17	11	11	4	1	1	-	-
2001.....	36	29	20	15	8	21	12	5	2	-	-	1
2000.....	35	28	13	19	12	15	6	7	4	-	-	-
1999.....	27	19	18	11	17	11	6	5	2	-	-	-
White, female												
2003.....	8	-	2	3	4	1	3	-	-	-	-	-
2002.....	3	3	1	2	2	3	3	1	-	-	-	-
2001.....	5	9	4	3	4	2	3	1	-	-	-	1
2000.....	2	6	-	3	3	1	-	-	-	-	-	-
1999.....	9	2	2	2	-	2	1	3	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

All races, both sexes

2003.....	186	153	112	102	100	94	68	58	20	5	-	30
2002.....	169	118	94	81	88	88	79	48	25	5	1	20
2001.....	145	109	97	78	81	85	72	56	15	4	-	34
2000.....	172	120	97	80	99	71	85	37	19	8	-	24
1999.....	172	122	97	84	101	88	70	45	16	7	-	34

All races, male

2003.....	147	125	86	73	72	57	47	31	15	2	-	30
2002.....	137	93	68	64	68	56	50	31	15	4	-	18
2001.....	117	83	72	61	63	55	47	37	8	-	-	29
2000.....	143	102	77	63	70	38	50	22	7	5	-	23
1999.....	145	95	75	64	65	63	48	19	6	3	-	30

All races, female

2003.....	39	28	26	29	28	37	21	27	5	3	-	-
2002.....	32	25	26	17	20	32	29	17	10	1	1	2
2001.....	28	26	25	17	18	30	25	19	7	4	-	5
2000.....	29	18	20	17	29	33	35	15	12	3	-	1
1999.....	27	27	22	20	36	25	22	26	10	4	-	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

White, both sexes

2003.....	153	131	93	92	86	80	65	49	17	4	-	29
2002.....	133	103	78	68	71	75	68	41	22	3	-	20
2001.....	125	85	81	66	73	76	62	50	14	4	-	28
2000.....	130	105	82	70	86	58	79	34	16	6	-	23
1999.....	142	99	82	72	82	74	64	39	16	5	-	32

White, male

2003.....	119	105	70	66	63	51	45	29	12	2	-	29
2002.....	110	81	55	54	56	50	44	29	14	3	-	18
2001.....	99	63	58	52	57	50	41	33	8	-	-	25
2000.....	107	89	70	55	59	31	50	20	7	3	-	22
1999.....	122	76	63	53	53	53	44	17	6	3	-	28

White, female

2003.....	34	26	23	26	23	29	20	20	5	2	-	-
2002.....	23	22	23	14	15	25	24	12	8	-	-	2
2001.....	26	22	23	14	16	26	21	17	6	4	-	3
2000.....	23	16	12	15	27	27	29	14	9	3	-	1
1999.....	20	23	19	19	29	21	20	22	10	2	-	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

Black, both sexes

2003.....	24	19	15	5	11	6	2	4	1	-	-	1
2002.....	24	12	10	9	9	10	7	6	2	2	1	-
2001.....	17	21	11	7	6	5	8	3	1	-	-	3
2000.....	29	9	8	7	8	9	3	3	2	2	-	1
1999.....	20	19	10	9	12	11	4	3	-	2	-	1

Black, male

2003.....	22	17	13	3	7	3	2	2	1	-	-	1
2002.....	19	11	8	8	7	5	4	2	1	1	-	-
2001.....	15	17	9	7	5	4	5	3	-	-	-	3
2000.....	24	8	5	6	7	4	-	2	-	2	-	1
1999.....	18	16	7	8	9	8	3	2	-	-	-	1

Black, female

2003.....	2	2	2	2	4	3	-	2	-	-	-	-
2002.....	5	1	2	1	2	5	3	4	1	1	1	-
2001.....	2	4	2	-	1	1	3	-	1	-	-	-
2000.....	5	1	3	1	1	5	3	1	2	-	-	-
1999.....	2	3	3	1	3	3	1	1	-	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

Asian or Pacific Islander 2/, both sexes

2003.....	6	2	4	4	3	7	1	5	2	1	-	-
2002.....	9	2	5	4	8	3	3	1	1	-	-	-
2001.....	1	1	4	5	2	4	2	3	-	-	-	3
2000.....	7	3	6	2	4	4	3	-	1	-	-	-
1999.....	5	2	3	2	5	2	2	3	-	-	-	1

Asian or Pacific Islander 2/, male

2003.....	5	2	3	3	2	2	-	-	2	-	-	-
2002.....	6	1	4	2	5	1	2	-	-	-	-	-
2001.....	1	1	4	2	1	1	1	1	-	-	-	1
2000.....	7	2	1	1	3	3	-	-	-	-	-	-
1999.....	3	1	3	2	1	1	1	-	-	-	-	1

Asian or Pacific Islander 2/, female

2003.....	1	-	1	1	1	5	1	5	-	1	-	-
2002.....	3	1	1	2	3	2	1	1	1	-	-	-
2001.....	-	-	-	3	1	3	1	2	-	-	-	2
2000.....	-	1	5	1	1	1	3	-	1	-	-	-
1999.....	2	1	-	-	4	1	1	3	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races, both sexes

2003.....	256	204	188	198	221	226	234	157	67	24	5	4
2002.....	235	183	157	174	211	211	221	129	70	18	4	3
2001.....	206	215	163	190	221	225	198	159	72	18	2	3
2000.....	225	187	182	188	213	257	215	158	78	24	3	5
1999.....	170	157	162	193	223	258	229	150	74	18	3	4

All races, male

2003.....	148	138	116	113	131	124	104	72	27	8	-	3
2002.....	146	111	108	117	125	108	123	55	31	9	2	3
2001.....	137	142	94	112	117	106	98	69	30	8	1	3
2000.....	158	126	115	107	125	136	108	69	34	11	1	5
1999.....	110	98	96	111	133	128	119	67	34	9	1	4

All races, female

2003.....	108	66	72	85	90	102	130	85	40	16	5	1
2002.....	89	72	49	57	86	103	98	74	39	9	2	-
2001.....	69	73	69	78	104	119	100	90	42	10	1	-
2000.....	67	61	67	81	88	121	107	89	44	13	2	-
1999.....	60	59	66	82	90	130	110	83	40	9	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

White, both sexes

2003.....	188	156	147	152	168	175	181	119	59	19	2	4
2002.....	174	140	107	130	159	164	178	104	49	11	3	3
2001.....	149	155	122	138	163	170	151	126	53	10	1	3
2000.....	175	129	131	139	158	189	156	129	51	17	-	5
1999.....	127	123	112	136	157	187	173	108	53	13	2	3

White, male

2003.....	115	103	93	87	102	90	78	58	23	6	-	3
2002.....	105	86	75	86	93	85	105	47	20	5	1	3
2001.....	95	105	67	80	83	76	78	57	21	4	-	3
2000.....	122	89	80	76	93	100	81	61	22	8	-	5
1999.....	82	74	75	77	92	89	95	52	26	7	1	3

White, female

2003.....	73	53	54	65	66	85	103	61	36	13	2	1
2002.....	69	54	32	44	66	79	73	57	29	6	2	-
2001.....	54	50	55	58	80	94	73	69	32	6	1	-
2000.....	53	40	51	63	65	89	75	68	29	9	-	-
1999.....	45	49	37	59	65	98	78	56	27	6	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2003.....	63	45	34	42	48	51	49	36	8	4	3	-
2002.....	58	39	48	38	47	42	41	25	20	7	1	-
2001.....	51	53	37	48	54	49	43	31	18	8	1	-
2000.....	48	53	49	45	53	64	57	29	26	7	3	-
1999.....	41	34	46	52	61	67	53	41	21	5	1	1

Black, male

2003.....	32	33	20	26	27	34	23	13	4	1	-	-
2002.....	39	23	32	27	28	20	18	8	11	4	1	-
2001.....	39	32	24	29	33	26	18	11	8	4	1	-
2000.....	34	35	34	29	31	35	26	8	12	3	1	-
1999.....	27	24	20	30	39	36	23	15	8	2	-	1

Black, female

2003.....	31	12	14	16	21	17	26	23	4	3	3	-
2002.....	19	16	16	11	19	22	23	17	9	3	-	-
2001.....	12	21	13	19	21	23	25	20	10	4	-	-
2000.....	14	18	15	16	22	29	31	21	14	4	2	-
1999.....	14	10	26	22	22	31	30	26	13	3	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

American Indian 1/, both sexes

2003.....	3	1	3	2	3	-	-	-	-	-	-	-
2002.....	1	2	1	4	4	1	-	-	1	-	-	-
2001.....	5	5	2	3	2	2	-	1	-	-	-	-
2000.....	1	3	2	1	1	1	-	-	-	-	-	-
1999.....	1	-	3	5	1	3	3	-	-	-	-	-

American Indian 1/, male

2003.....	1	-	1	-	1	-	-	-	-	-	-	-
2002.....	1	2	-	4	3	1	-	-	-	-	-	-
2001.....	2	4	2	2	1	1	-	1	-	-	-	-
2000.....	1	1	1	-	1	1	-	-	-	-	-	-
1999.....	1	-	-	4	1	2	1	-	-	-	-	-

American Indian 1/, female

2003.....	2	1	2	2	2	-	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	-	-	1	-	-	-
2001.....	3	1	-	1	1	1	-	-	-	-	-	-
2000.....	-	2	1	1	-	-	-	-	-	-	-	-
1999.....	-	-	3	1	-	1	2	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Asian or Pacific Islander 2/, both sexes

2003.....	2	2	4	2	2	-	4	2	-	1	-	-
2002.....	2	2	1	2	1	4	2	-	-	-	-	-
2001.....	1	2	2	1	2	4	4	1	1	-	-	-
2000.....	1	2	-	3	1	3	2	-	1	-	-	-
1999.....	1	-	1	-	4	1	-	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	-	2	2	-	1	-	3	1	-	1	-	-
2002.....	1	-	1	-	1	2	-	-	-	-	-	-
2001.....	1	1	1	1	-	3	2	-	1	-	-	-
2000.....	1	1	-	2	-	-	1	-	-	-	-	-
1999.....	-	-	1	-	1	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	-	2	2	1	-	1	1	-	-	-	-
2002.....	1	2	-	2	-	2	2	-	-	-	-	-
2001.....	-	1	1	-	2	1	2	1	-	-	-	-
2000.....	-	1	-	1	1	3	1	-	1	-	-	-
1999.....	1	-	-	-	3	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races, both sexes

2003.....	2,083	957	413	229	174	146	138	105	47	10	4	6
2002.....	1,736	754	344	218	165	154	141	103	64	13	1	9
2001.....	1,281	530	268	161	130	153	126	91	47	12	2	10
2000.....	1,069	447	241	146	132	118	127	109	60	14	1	7
1999.....	918	449	219	158	142	138	127	100	31	12	-	12

All races, male

2003.....	1,368	584	257	137	86	71	71	34	19	4	1	5
2002.....	1,176	491	199	126	83	71	65	38	25	8	-	7
2001.....	893	341	159	88	82	83	54	28	18	2	1	10
2000.....	768	260	141	79	68	49	53	51	23	2	-	6
1999.....	650	298	150	98	74	63	53	38	10	5	-	8

All races, female

2003.....	715	373	156	92	88	75	67	71	28	6	3	1
2002.....	560	263	145	92	82	83	76	65	39	5	1	2
2001.....	388	189	109	73	48	70	72	63	29	10	1	-
2000.....	301	187	100	67	64	69	74	58	37	12	1	1
1999.....	268	151	69	60	68	75	74	62	21	7	-	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, both sexes

2003.....	1,676	787	350	167	149	133	130	94	41	9	4	4
2002.....	1,375	585	270	183	144	137	127	94	58	11	1	7
2001.....	952	416	200	135	110	131	106	80	41	10	-	9
2000.....	790	349	194	122	111	111	118	92	54	11	-	6
1999.....	685	353	157	127	113	128	117	84	29	10	-	9

White, male

2003.....	1,088	461	211	90	73	64	67	29	15	3	1	3
2002.....	911	361	139	101	69	64	54	34	25	7	-	5
2001.....	651	252	107	73	69	66	47	25	17	1	-	9
2000.....	558	187	105	60	55	46	49	43	21	1	-	5
1999.....	469	221	100	77	61	58	49	35	10	4	-	6

White, female

2003.....	588	326	139	77	76	69	63	65	26	6	3	1
2002.....	464	224	131	82	75	73	73	60	33	4	1	2
2001.....	301	164	93	62	41	65	59	55	24	9	-	-
2000.....	232	162	89	62	56	65	69	49	33	10	-	1
1999.....	216	132	57	50	52	70	68	49	19	6	-	3

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2003.....	366	152	57	52	18	13	7	10	6	1	-	2
2002.....	339	154	69	28	17	15	12	6	5	1	-	1
2001.....	310	105	60	24	17	17	17	11	6	2	2	1
2000.....	266	92	43	24	19	7	8	16	6	3	1	1
1999.....	217	85	59	26	24	10	8	14	2	2	-	3

Black, male

2003.....	255	113	45	42	12	7	3	5	4	1	-	2
2002.....	247	119	57	22	13	7	9	1	-	-	-	1
2001.....	229	81	47	15	12	14	6	3	1	1	1	1
2000.....	199	70	34	19	12	3	4	8	2	1	-	1
1999.....	170	70	47	18	12	5	4	3	-	1	-	2

Black, female

2003.....	111	39	12	10	6	6	4	5	2	-	-	-
2002.....	92	35	12	6	4	8	3	5	5	1	-	-
2001.....	81	24	13	9	5	3	11	8	5	1	1	-
2000.....	67	22	9	5	7	4	4	8	4	2	1	-
1999.....	47	15	12	8	12	5	4	11	2	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

American Indian 1/, both sexes

2003.....	25	10	4	8	1	-	-	-	-	-	-	-
2002.....	14	12	2	1	-	-	-	-	1	-	-	1
2001.....	11	5	6	1	2	-	1	-	-	-	-	-
2000.....	8	3	-	-	1	-	-	1	-	-	-	-
1999.....	6	5	1	3	1	-	-	1	-	-	-	-

American Indian 1/, male

2003.....	15	5	1	4	1	-	-	-	-	-	-	-
2002.....	11	9	1	1	-	-	-	-	-	-	-	1
2001.....	7	4	4	-	1	-	1	-	-	-	-	-
2000.....	8	2	-	-	-	-	-	-	-	-	-	-
1999.....	3	5	1	2	-	-	-	-	-	-	-	-

American Indian 1/, female

2003.....	10	5	3	4	-	-	-	-	-	-	-	-
2002.....	3	3	1	-	-	-	-	-	1	-	-	-
2001.....	4	1	2	1	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	1	-	-	1	-	-	-	-
1999.....	3	-	-	1	1	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Asian or Pacific Islander 2/, both sexes

2003.....	16	8	2	2	6	-	1	1	-	-	-	-
2002.....	8	3	3	6	4	2	2	3	-	1	-	-
2001.....	8	4	2	1	1	5	2	-	-	-	-	-
2000.....	5	3	4	-	1	-	1	-	-	-	-	-
1999.....	10	6	2	2	4	-	2	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	10	5	-	1	-	-	1	-	-	-	-	-
2002.....	7	2	2	2	1	-	2	3	-	1	-	-
2001.....	6	4	1	-	-	3	-	-	-	-	-	-
2000.....	3	1	2	-	1	-	-	-	-	-	-	-
1999.....	8	2	2	1	1	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	6	3	2	1	6	-	-	1	-	-	-	-
2002.....	1	1	1	4	3	2	-	-	-	-	-	-
2001.....	2	-	1	1	1	2	2	-	-	-	-	-
2000.....	2	2	2	-	-	-	1	-	-	-	-	-
1999.....	2	4	-	1	3	-	2	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races, both sexes

2003.....	770	750	670	794	966	1,475	2,064	2,302	1,644	720	167	15
2002.....	751	722	663	826	969	1,601	2,064	2,254	1,720	727	150	13
2001.....	825	646	696	744	1,082	1,669	2,172	2,404	1,784	704	171	24
2000.....	723	668	681	743	1,081	1,632	2,046	2,285	1,798	706	177	16
1999.....	741	662	670	852	1,180	1,735	2,290	2,563	1,895	760	137	15

All races, male

2003.....	566	545	500	518	577	820	978	958	548	190	41	13
2002.....	571	534	479	534	579	879	948	878	545	201	24	10
2001.....	614	461	472	480	617	897	1,016	950	610	155	26	21
2000.....	573	504	467	474	607	855	946	883	564	179	33	16
1999.....	592	490	461	526	701	900	1,069	1,003	598	194	27	14

All races, female

2003.....	204	205	170	276	389	655	1,086	1,344	1,096	530	126	2
2002.....	180	188	184	292	390	722	1,116	1,376	1,175	526	126	3
2001.....	211	185	224	264	465	772	1,156	1,454	1,174	549	145	3
2000.....	150	164	214	269	474	777	1,100	1,402	1,234	527	144	-
1999.....	149	172	209	326	479	835	1,221	1,560	1,297	566	110	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2003.....	616	618	556	668	833	1,327	1,870	2,171	1,527	677	150	14
2002.....	624	592	558	651	840	1,413	1,876	2,096	1,610	675	135	13
2001.....	684	545	565	632	921	1,478	1,978	2,232	1,655	650	155	22
2000.....	593	547	570	613	923	1,443	1,877	2,132	1,680	655	158	14
1999.....	606	526	533	710	997	1,507	2,074	2,363	1,784	713	125	13

White, male

2003.....	460	453	409	446	497	739	879	900	507	178	34	12
2002.....	473	432	408	419	510	776	860	814	505	187	18	10
2001.....	504	384	390	410	530	801	910	881	561	144	25	19
2000.....	470	409	385	389	526	745	855	815	523	161	27	14
1999.....	486	389	374	435	591	781	971	916	550	186	23	12

White, female

2003.....	156	165	147	222	336	588	991	1,271	1,020	499	116	2
2002.....	151	160	150	232	330	637	1,016	1,282	1,105	488	117	3
2001.....	180	161	175	222	391	677	1,068	1,351	1,094	506	130	3
2000.....	123	138	185	224	397	698	1,022	1,317	1,157	494	131	-
1999.....	120	137	159	275	406	726	1,103	1,447	1,234	527	102	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2003.....	134	106	95	107	111	115	156	107	93	38	13	1
2002.....	107	109	84	150	107	162	149	138	90	42	10	-
2001.....	117	84	116	96	139	157	162	145	110	43	15	2
2000.....	105	106	94	111	136	170	139	132	95	44	17	2
1999.....	116	122	117	126	169	203	186	175	100	46	10	2

Black, male

2003.....	91	74	72	62	68	64	77	42	30	11	5	1
2002.....	83	85	57	95	56	89	70	54	31	11	3	-
2001.....	88	63	72	62	75	76	90	54	39	7	1	2
2000.....	86	84	67	70	69	99	76	56	32	15	5	2
1999.....	89	91	76	80	103	106	80	79	41	8	3	2

Black, female

2003.....	43	32	23	45	43	51	79	65	63	27	8	-
2002.....	24	24	27	55	51	73	79	84	59	31	7	-
2001.....	29	21	44	34	64	81	72	91	71	36	14	-
2000.....	19	22	27	41	67	71	63	76	63	29	12	-
1999.....	27	31	41	46	66	97	106	96	59	38	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

American Indian 1/, both sexes

2003.....	12	16	12	11	6	14	9	7	6	1	1	-
2002.....	10	12	7	14	9	10	6	-	6	3	-	-
2001.....	16	5	10	6	8	14	9	6	6	3	-	-
2000.....	20	13	7	9	8	5	14	3	2	-	1	-
1999.....	11	6	15	9	6	13	8	6	2	-	1	-

American Indian 1/, male

2003.....	9	12	12	7	6	8	3	3	3	-	1	-
2002.....	7	9	4	11	7	5	3	-	4	1	-	-
2001.....	15	4	5	3	5	8	3	4	2	-	-	-
2000.....	14	10	6	6	5	2	6	2	-	-	-	-
1999.....	11	5	10	8	3	5	5	3	-	-	1	-

American Indian 1/, female

2003.....	3	4	-	4	-	6	6	4	3	1	-	-
2002.....	3	3	3	3	2	5	3	-	2	2	-	-
2001.....	1	1	5	3	3	6	6	2	4	3	-	-
2000.....	6	3	1	3	3	3	8	1	2	-	1	-
1999.....	-	1	5	1	3	8	3	3	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Asian or Pacific Islander 2/, both sexes

2003.....	8	10	7	8	16	19	29	17	18	4	3	-
2002.....	10	9	14	11	13	16	33	20	14	7	5	-
2001.....	8	12	5	10	14	20	23	21	13	8	1	-
2000.....	5	2	10	10	14	14	16	18	21	7	1	-
1999.....	8	8	5	7	8	12	22	19	9	1	1	-

Asian or Pacific Islander 2/, male

2003.....	6	6	7	3	6	9	19	13	8	1	1	-
2002.....	8	8	10	9	6	9	15	10	5	2	3	-
2001.....	7	10	5	5	7	12	13	11	8	4	-	-
2000.....	3	1	9	9	7	9	9	10	9	3	1	-
1999.....	6	5	1	3	4	8	13	5	7	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	4	-	5	10	10	10	4	10	3	2	-
2002.....	2	1	4	2	7	7	18	10	9	5	2	-
2001.....	1	2	-	5	7	8	10	10	5	4	1	-
2000.....	2	1	1	1	7	5	7	8	12	4	-	-
1999.....	2	3	4	4	4	4	9	14	2	1	1	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races, both sexes

2003.....	2,996	2,314	1,529	1,167	1,168	1,159	956	558	205	34	1	7
2002.....	2,835	2,186	1,432	1,197	1,266	1,231	1,028	573	211	41	1	10
2001 3/.....	2,682	1,985	1,332	1,212	1,220	1,219	973	538	200	28	3	15
2000.....	2,450	1,708	1,237	1,073	1,219	1,227	954	590	205	34	4	7
1999.....	2,264	1,678	1,218	1,187	1,284	1,280	937	580	182	39	-	16

All races, male

2003.....	2,271	1,771	1,216	958	996	985	824	484	178	28	-	7
2002.....	2,188	1,708	1,129	975	1,078	1,054	884	492	179	33	-	10
2001 3/.....	2,020	1,534	1,029	1,001	1,039	1,033	852	468	168	25	3	14
2000.....	1,842	1,306	959	870	1,018	1,062	822	507	168	26	4	7
1999.....	1,702	1,284	971	972	1,079	1,073	795	501	150	30	-	16

All races, female

2003.....	725	543	313	209	172	174	132	74	27	6	1	-
2002.....	647	478	303	222	188	177	144	81	32	8	1	-
2001 3/.....	662	451	303	211	181	186	121	70	32	3	-	1
2000.....	608	402	278	203	201	165	132	83	37	8	-	-
1999.....	562	394	247	215	205	207	142	79	32	9	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, both sexes

2003.....	2,810	2,181	1,437	1,092	1,106	1,107	922	534	204	31	1	7
2002.....	2,640	2,064	1,350	1,130	1,199	1,163	1,000	551	203	38	1	10
2001 3/.....	2,510	1,859	1,247	1,132	1,150	1,169	931	518	195	27	3	14
2000.....	2,273	1,604	1,171	1,017	1,151	1,174	916	574	198	32	2	6
1999.....	2,121	1,567	1,131	1,104	1,226	1,226	903	563	178	38	-	13

White, male

2003.....	2,141	1,673	1,143	901	944	944	797	468	177	27	-	7
2002.....	2,054	1,610	1,074	919	1,034	1,001	860	475	173	30	-	10
2001 3/.....	1,893	1,438	958	942	979	990	820	455	164	25	3	13
2000.....	1,706	1,225	908	823	961	1,020	797	495	162	24	2	6
1999.....	1,586	1,196	903	911	1,029	1,031	770	487	148	29	-	13

White, female

2003.....	669	508	294	191	162	163	125	66	27	4	1	-
2002.....	586	454	276	211	165	162	140	76	30	8	1	-
2001 3/.....	617	421	289	190	171	179	111	63	31	2	-	1
2000.....	567	379	263	194	190	154	119	79	36	8	-	-
1999.....	535	371	228	193	197	195	133	76	30	9	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Black, both sexes

2003.....	128	76	61	43	34	26	18	11	-	1	-	-
2002.....	123	69	58	33	49	39	14	16	2	2	-	-
2001 3/.....	118	82	49	43	48	30	24	8	4	1	-	-
2000.....	119	62	41	39	46	30	23	9	2	-	1	-
1999.....	84	68	50	58	38	35	17	10	4	1	-	2

Black, male

2003.....	93	57	50	32	27	24	16	8	-	-	-	-
2002.....	89	58	41	31	37	34	14	14	2	2	-	-
2001 3/.....	89	65	41	32	42	26	20	5	4	-	-	-
2000.....	90	51	31	36	40	25	15	8	2	-	1	-
1999.....	71	55	42	47	33	30	15	9	2	1	-	2

Black, female

2003.....	35	19	11	11	7	2	2	3	-	1	-	-
2002.....	34	11	17	2	12	5	-	2	-	-	-	-
2001 3/.....	29	17	8	11	6	4	4	3	-	1	-	-
2000.....	29	11	10	3	6	5	8	1	-	-	-	-
1999.....	13	13	8	11	5	5	2	1	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Asian or Pacific Islander 2/, both sexes

2003.....	47	52	24	28	26	22	16	12	1	2	-	-
2002.....	52	44	20	31	18	24	14	6	6	1	-	-
2001 3/.....	34	37	26	34	19	18	15	12	1	-	-	1
2000.....	43	29	22	14	17	22	15	7	5	2	1	1
1999.....	48	35	33	21	15	19	15	7	-	-	-	1

Asian or Pacific Islander 2/, male

2003.....	29	37	18	22	23	14	11	7	1	1	-	-
2002.....	30	33	11	22	7	14	10	3	4	1	-	-
2001 3/.....	19	25	23	26	15	15	9	8	-	-	-	1
2000.....	33	18	17	8	12	16	10	4	4	2	1	1
1999.....	38	28	24	10	12	12	8	5	-	-	-	1

Asian or Pacific Islander 2/, female

2003.....	18	15	6	6	3	8	5	5	-	1	-	-
2002.....	22	11	9	9	11	10	4	3	2	-	-	-
2001 3/.....	15	12	3	8	4	3	6	4	1	-	-	-
2000.....	10	11	5	6	5	6	5	3	1	-	-	-
1999.....	10	7	9	11	3	7	7	2	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes

2003.....	1,574	1,329	988	833	867	886	709	399	136	24	-	-
2002.....	1,484	1,295	939	846	930	918	762	392	133	25	-	2
2001.....	1,471	1,195	888	857	901	920	728	382	135	18	2	6
2000.....	1,367	1,001	832	791	909	917	694	410	129	18	1	5
1999.....	1,215	1,058	802	827	964	946	660	399	102	23	-	5

All races, male

2003.....	1,349	1,122	879	747	791	815	665	375	132	22	-	-
2002.....	1,283	1,115	815	753	867	854	722	374	127	24	-	2
2001.....	1,242	1,020	751	761	831	846	685	360	126	18	2	6
2000.....	1,146	854	714	698	830	856	649	388	119	18	1	5
1999.....	1,030	882	712	742	873	875	624	376	99	21	-	5

All races, female

2003.....	225	207	109	86	76	71	44	24	4	2	-	-
2002.....	201	180	124	93	63	64	40	18	6	1	-	-
2001.....	229	175	137	96	70	74	43	22	9	-	-	-
2000.....	221	147	118	93	79	61	45	22	10	-	-	-
1999.....	185	176	90	85	91	71	36	23	3	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2003.....	1,487	1,276	945	797	835	856	687	391	136	23	-	-
2002.....	1,395	1,238	897	817	894	878	747	380	131	23	-	2
2001.....	1,397	1,137	851	820	860	889	700	374	131	18	2	5
2000.....	1,290	941	791	759	867	894	679	403	126	17	1	4
1999.....	1,154	998	756	785	929	917	640	390	100	22	-	5

White, male

2003.....	1,275	1,079	839	717	762	787	646	368	132	22	-	-
2002.....	1,208	1,062	778	726	836	816	707	362	125	22	-	2
2001.....	1,181	969	717	731	794	818	662	353	122	18	2	5
2000.....	1,083	800	679	667	790	836	637	381	116	17	1	4
1999.....	974	827	672	707	838	849	605	368	97	20	-	5

White, female

2003.....	212	197	106	80	73	69	41	23	4	1	-	-
2002.....	187	176	119	91	58	62	40	18	6	1	-	-
2001.....	216	168	134	89	66	71	38	21	9	-	-	-
2000.....	207	141	112	92	77	58	42	22	10	-	-	-
1999.....	180	171	84	78	91	68	35	22	3	2	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, both sexes

2003.....	69	39	36	32	24	23	16	6	-	1	-	-
2002.....	60	40	33	22	34	31	11	12	2	2	-	-
2001.....	60	43	27	27	35	26	23	6	4	-	-	-
2000.....	61	39	33	28	34	19	13	6	2	-	-	-
1999.....	43	46	36	36	29	28	15	9	2	1	-	-

Black, male

2003.....	59	30	33	26	21	21	14	6	-	-	-	-
2002.....	51	37	28	20	30	29	11	12	2	2	-	-
2001.....	50	39	25	22	31	23	19	5	4	-	-	-
2000.....	48	35	27	27	33	17	10	6	2	-	-	-
1999.....	40	42	32	30	29	25	14	8	2	1	-	-

Black, female

2003.....	10	9	3	6	3	2	2	-	-	1	-	-
2002.....	9	3	5	2	4	2	-	-	-	-	-	-
2001.....	10	4	2	5	4	3	4	1	-	-	-	-
2000.....	13	4	6	1	1	2	3	-	-	-	-	-
1999.....	3	4	4	6	-	3	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races, both sexes

2003.....	1,422	985	541	334	301	273	247	159	69	10	1	7
2002.....	1,351	891	493	351	336	313	266	181	78	16	1	8
2001 3/.....	1,211	790	444	355	319	299	245	156	65	10	1	9
2000.....	1,083	707	405	282	310	310	260	180	76	16	3	2
1999.....	1,049	620	416	360	320	334	277	181	80	16	-	11

All races, male

2003.....	922	649	337	211	205	170	159	109	46	6	-	7
2002.....	905	593	314	222	211	200	162	118	52	9	-	8
2001 3/.....	778	514	278	240	208	187	167	108	42	7	1	8
2000.....	696	452	245	172	188	206	173	119	49	8	3	2
1999.....	672	402	259	230	206	198	171	125	51	9	-	11

All races, female

2003.....	500	336	204	123	96	103	88	50	23	4	1	-
2002.....	446	298	179	129	125	113	104	63	26	7	1	-
2001 3/.....	433	276	166	115	111	112	78	48	23	3	-	1
2000.....	387	255	160	110	122	104	87	61	27	8	-	-
1999.....	377	218	157	130	114	136	106	56	29	7	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

White, both sexes

2003.....	1,323	905	492	295	271	251	235	143	68	8	1	7
2002.....	1,245	826	453	313	305	285	253	171	72	15	1	8
2001 3/.....	1,113	722	396	312	290	280	231	144	64	9	1	9
2000.....	983	663	380	258	284	280	237	171	72	15	1	2
1999.....	967	569	375	319	297	309	263	173	78	16	-	8

White, male

2003.....	866	594	304	184	182	157	151	100	45	5	-	7
2002.....	846	548	296	193	198	185	153	113	48	8	-	8
2001 3/.....	712	469	241	211	185	172	158	102	42	7	1	8
2000.....	623	425	229	156	171	184	160	114	46	7	1	2
1999.....	612	369	231	204	191	182	165	119	51	9	-	8

White, female

2003.....	457	311	188	111	89	94	84	43	23	3	1	-
2002.....	399	278	157	120	107	100	100	58	24	7	1	-
2001 3/.....	401	253	155	101	105	108	73	42	22	2	-	1
2000.....	360	238	151	102	113	96	77	57	26	8	-	-
1999.....	355	200	144	115	106	127	98	54	27	7	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Black, both sexes

2003.....	59	37	25	11	10	3	2	5	-	-	-	-
2002.....	63	29	25	11	15	8	3	4	-	-	-	-
2001 3/.....	58	39	22	16	13	4	1	2	-	1	-	-
2000.....	58	23	8	11	12	11	10	3	-	-	1	-
1999.....	41	22	14	22	9	7	2	1	2	-	-	2

Black, male

2003.....	34	27	17	6	6	3	2	2	-	-	-	-
2002.....	38	21	13	11	7	5	3	2	-	-	-	-
2001 3/.....	39	26	16	10	11	3	1	-	-	-	-	-
2000.....	42	16	4	9	7	8	5	2	-	-	1	-
1999.....	31	13	10	17	4	5	1	1	-	-	-	2

Black, female

2003.....	25	10	8	5	4	-	-	3	-	-	-	-
2002.....	25	8	12	-	8	3	-	2	-	-	-	-
2001 3/.....	19	13	6	6	2	1	-	2	-	1	-	-
2000.....	16	7	4	2	5	3	5	1	-	-	-	-
1999.....	10	9	4	5	5	2	1	-	2	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Asian or Pacific Islander 2/, both sexes

2003.....	34	41	20	26	20	17	10	11	1	2	-	-
2002.....	33	32	13	26	16	19	10	6	6	1	-	-
2001 3/.....	25	24	21	25	16	15	12	10	1	-	-	-
2000.....	37	16	17	13	13	19	13	6	4	1	1	-
1999.....	36	22	26	18	12	18	12	7	-	-	-	1

Asian or Pacific Islander 2/, male

2003.....	18	27	14	20	17	9	6	7	1	1	-	-
2002.....	15	21	4	17	6	9	6	3	4	1	-	-
2001 3/.....	12	15	18	18	12	12	7	6	-	-	-	-
2000.....	27	7	12	7	9	14	8	3	3	1	1	-
1999.....	27	16	18	8	9	11	5	5	-	-	-	1

Asian or Pacific Islander 2/, female

2003.....	16	14	6	6	3	8	4	4	-	1	-	-
2002.....	18	11	9	9	10	10	4	3	2	-	-	-
2001 3/.....	13	9	3	7	4	3	5	4	1	-	-	-
2000.....	10	9	5	6	4	5	5	3	1	-	-	-
1999.....	9	6	8	10	3	7	7	2	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races, both sexes

2003.....	798	490	296	227	205	187	130	77	22	5	-	41
2002.....	760	523	318	229	192	169	127	64	24	6	1	41
2001 3/.....	1,004	628	390	310	222	183	129	71	29	4	1	32
2000.....	696	455	283	233	210	172	129	68	23	8	1	30
1999.....	670	420	302	259	214	185	119	65	24	10	-	49

All races, male

2003.....	595	352	216	156	129	110	61	32	13	1	-	29
2002.....	553	383	231	152	125	100	58	32	6	3	-	32
2001 3/.....	747	461	262	204	128	98	57	33	7	1	1	23
2000.....	511	338	199	147	132	85	69	25	10	5	-	15
1999.....	483	323	218	183	121	101	61	28	8	5	-	33

All races, female

2003.....	203	138	80	71	76	77	69	45	9	4	-	12
2002.....	207	140	87	77	67	69	69	32	18	3	1	9
2001 3/.....	257	167	128	106	94	85	72	38	22	3	-	9
2000.....	185	117	84	86	78	87	60	43	13	3	1	15
1999.....	187	97	84	76	93	84	58	37	16	5	-	16

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2003.....	493	327	203	160	144	143	101	65	18	3	-	36
2002.....	453	366	221	160	146	119	99	53	18	5	-	36
2001 3/.....	681	461	283	225	167	139	101	63	22	3	-	24
2000.....	432	309	188	150	153	138	89	57	18	7	1	25
1999.....	429	287	209	193	160	133	92	51	21	7	-	34

White, male

2003.....	354	217	147	105	89	82	46	27	11	1	-	25
2002.....	311	254	155	105	88	66	42	28	3	3	-	29
2001 3/.....	496	328	190	142	91	73	45	28	5	-	-	16
2000.....	295	224	126	86	90	67	42	20	7	4	-	13
1999.....	308	210	146	130	86	70	47	19	6	4	-	23

White, female

2003.....	139	110	56	55	55	61	55	38	7	2	-	11
2002.....	142	112	66	55	58	53	57	25	15	2	-	7
2001 3/.....	185	133	93	83	76	66	56	35	17	3	-	8
2000.....	137	85	62	64	63	71	47	37	11	3	1	12
1999.....	121	77	63	63	74	63	45	32	15	3	-	11

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2003.....	274	143	80	54	52	39	26	11	4	2	-	4
2002.....	273	132	86	55	42	48	23	11	5	1	-	3
2001 3/.....	274	141	84	71	48	40	23	7	6	1	1	8
2000.....	231	123	81	74	51	30	34	8	5	1	-	4
1999.....	198	116	79	60	44	46	21	13	3	3	-	14

Black, male

2003.....	216	123	59	44	35	27	14	5	2	-	-	3
2002.....	217	113	68	40	34	33	15	4	2	-	-	2
2001 3/.....	213	111	59	52	35	25	9	4	2	1	1	7
2000.....	193	97	63	58	37	16	23	3	3	1	-	1
1999.....	150	100	60	50	30	29	10	8	2	1	-	10

Black, female

2003.....	58	20	21	10	17	12	12	6	2	2	-	1
2002.....	56	19	18	15	8	15	8	7	3	1	-	1
2001 3/.....	61	30	25	19	13	15	14	3	4	-	-	1
2000.....	38	26	18	16	14	14	11	5	2	-	-	3
1999.....	48	16	19	10	14	17	11	5	1	2	-	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

American Indian 1/, both sexes

2003.....	9	3	1	3	2	1	-	1	-	-	-	-
2002.....	14	7	2	6	1	-	3	-	-	-	-	1
2001 3/.....	13	3	2	5	1	-	2	-	1	-	-	-
2000.....	9	2	4	2	2	2	-	2	-	-	-	1
1999.....	18	8	3	3	4	2	3	-	-	-	-	-

American Indian 1/, male

2003.....	8	1	1	2	2	-	-	-	-	-	-	-
2002.....	10	3	2	3	1	-	1	-	-	-	-	-
2001 3/.....	10	3	2	3	1	-	1	-	-	-	-	-
2000.....	7	1	3	1	2	1	-	2	-	-	-	1
1999.....	13	6	3	3	3	1	3	-	-	-	-	-

American Indian 1/, female

2003.....	1	2	-	1	-	1	-	1	-	-	-	-
2002.....	4	4	-	3	-	-	2	-	-	-	-	1
2001 3/.....	3	-	-	2	-	-	1	-	1	-	-	-
2000.....	2	1	1	1	-	1	-	-	-	-	-	-
1999.....	5	2	-	-	1	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Asian or Pacific Islander 2/, both sexes

2003.....	22	17	12	10	7	4	3	-	-	-	-	1
2002.....	20	18	9	8	3	2	2	-	1	-	1	1
2001 3/.....	36	23	21	9	6	4	3	1	-	-	-	-
2000.....	24	21	10	7	4	2	6	1	-	-	-	-
1999.....	25	9	11	3	6	4	3	1	-	-	-	1

Asian or Pacific Islander 2/, male

2003.....	17	11	9	5	3	1	1	-	-	-	-	1
2002.....	15	13	6	4	2	1	-	-	1	-	-	1
2001 3/.....	28	19	11	7	1	-	2	1	-	-	-	-
2000.....	16	16	7	2	3	1	4	-	-	-	-	-
1999.....	12	7	9	-	2	1	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	5	6	3	5	4	3	2	-	-	-	-	-
2002.....	5	5	3	4	1	1	2	-	-	-	1	-
2001 3/.....	8	4	10	2	5	4	1	-	-	-	-	-
2000.....	8	5	3	5	1	1	2	1	-	-	-	-
1999.....	13	2	2	3	4	3	2	-	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races, both sexes

2003.....	427	250	144	83	78	55	33	17	5	1	-	18
2002.....	421	263	146	92	69	72	43	14	4	1	-	13
2001.....	382	213	151	114	78	61	38	12	2	2	-	11
2000.....	377	235	132	99	85	54	27	23	4	1	-	12
1999.....	342	206	167	102	91	61	34	17	4	2	-	15

All races, male

2003.....	326	186	106	63	52	27	8	9	1	-	-	16
2002.....	316	197	104	61	40	44	20	8	-	1	-	12
2001.....	292	159	105	79	44	38	16	6	-	1	-	11
2000.....	285	172	94	58	50	23	14	6	2	1	-	7
1999.....	250	167	127	68	56	34	14	7	2	1	-	14

All races, female

2003.....	101	64	38	20	26	28	25	8	4	1	-	2
2002.....	105	66	42	31	29	28	23	6	4	-	-	1
2001.....	90	54	46	35	34	23	22	6	2	1	-	-
2000.....	92	63	38	41	35	31	13	17	2	-	-	5
1999.....	92	39	40	34	35	27	20	10	2	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2003.....	265	168	100	58	58	41	27	17	3	-	-	14
2002.....	253	190	94	62	54	55	36	13	3	1	-	11
2001.....	236	142	113	84	57	47	31	12	2	1	-	9
2000.....	229	165	95	70	64	47	17	19	4	1	-	10
1999.....	213	139	113	79	69	43	29	16	4	1	-	10

White, male

2003.....	196	116	72	43	38	20	5	9	-	-	-	13
2002.....	183	134	66	38	27	32	15	8	-	1	-	10
2001.....	173	94	76	51	29	27	10	6	-	-	-	9
2000.....	160	117	60	37	33	20	5	5	2	1	-	5
1999.....	162	106	80	50	40	20	11	6	2	1	-	9

White, female

2003.....	69	52	28	15	20	21	22	8	3	-	-	1
2002.....	70	56	28	24	27	23	21	5	3	-	-	1
2001.....	63	48	37	33	28	20	21	6	2	1	-	-
2000.....	69	48	35	33	31	27	12	14	2	-	-	5
1999.....	51	33	33	29	29	23	18	10	2	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2003.....	147	72	37	23	19	11	6	-	2	1	-	3
2002.....	150	63	46	26	14	15	6	1	1	-	-	2
2001.....	130	61	33	27	18	13	5	-	-	1	-	2
2000.....	131	56	29	25	20	7	10	4	-	-	-	1
1999.....	104	59	45	20	19	17	4	1	-	1	-	5

Black, male

2003.....	118	65	27	19	14	7	3	-	1	-	-	2
2002.....	118	56	33	21	12	11	5	-	-	-	-	2
2001.....	109	55	25	25	14	11	4	-	-	1	-	2
2000.....	113	43	26	20	17	3	9	1	-	-	-	1
1999.....	77	55	39	17	14	14	2	1	-	-	-	5

Black, female

2003.....	29	7	10	4	5	4	3	-	1	1	-	1
2002.....	32	7	13	5	2	4	1	1	1	-	-	-
2001.....	21	6	8	2	4	2	1	-	-	-	-	-
2000.....	18	13	3	5	3	4	1	3	-	-	-	-
1999.....	27	4	6	3	5	3	2	-	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Asian or Pacific Islander 2/, both sexes

2003.....	14	8	6	2	1	2	-	-	-	-	-	1
2002.....	13	9	5	3	-	2	1	-	-	-	-	-
2001.....	14	8	4	1	2	1	2	-	-	-	-	-
2000.....	15	14	6	3	1	-	-	-	-	-	-	-
1999.....	17	6	7	2	1	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	11	5	6	1	-	-	-	-	-	-	-	1
2002.....	10	6	4	2	-	1	-	-	-	-	-	-
2001.....	9	8	3	1	-	-	2	-	-	-	-	-
2000.....	11	12	6	1	-	-	-	-	-	-	-	-
1999.....	6	4	6	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	3	3	-	1	1	2	-	-	-	-	-	-
2002.....	3	3	1	1	-	1	1	-	-	-	-	-
2001.....	5	-	1	-	2	1	-	-	-	-	-	-
2000.....	4	2	-	2	1	-	-	-	-	-	-	-
1999.....	11	2	1	2	1	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races, both sexes

2003.....	371	240	152	144	127	132	97	60	17	4	-	23
2002.....	339	260	172	137	123	97	84	50	20	5	1	28
2001 3/.....	622	415	239	196	144	122	91	59	27	2	1	21
2000.....	319	220	151	134	125	118	102	45	19	7	1	18
1999.....	328	214	135	157	123	124	85	48	20	8	-	34

All races, male

2003.....	269	166	110	93	77	83	53	23	12	1	-	13
2002.....	237	186	127	91	85	56	38	24	6	2	-	20
2001 3/.....	455	302	157	125	84	60	41	27	7	-	1	12
2000.....	226	166	105	89	82	62	55	19	8	4	-	8
1999.....	233	156	91	115	65	67	47	21	6	4	-	19

All races, female

2003.....	102	74	42	51	50	49	44	37	5	3	-	10
2002.....	102	74	45	46	38	41	46	26	14	3	1	8
2001 3/.....	167	113	82	71	60	62	50	32	20	2	-	9
2000.....	93	54	46	45	43	56	47	26	11	3	1	10
1999.....	95	58	44	42	58	57	38	27	14	4	-	15

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2003.....	228	159	103	102	86	102	74	48	15	3	-	22
2002.....	200	176	127	98	92	64	63	40	15	4	-	25
2001 3/.....	445	319	170	141	110	92	70	51	20	2	-	15
2000.....	203	144	93	80	89	91	72	38	14	6	1	15
1999.....	216	148	96	114	91	90	63	35	17	6	-	24

White, male

2003.....	158	101	75	62	51	62	41	18	11	1	-	12
2002.....	128	120	89	67	61	34	27	20	3	2	-	19
2001 3/.....	323	234	114	91	62	46	35	22	5	-	-	7
2000.....	135	107	66	49	57	47	37	15	5	3	-	8
1999.....	146	104	66	80	46	50	36	13	4	3	-	14

White, female

2003.....	70	58	28	40	35	40	33	30	4	2	-	10
2002.....	72	56	38	31	31	30	36	20	12	2	-	6
2001 3/.....	122	85	56	50	48	46	35	29	15	2	-	8
2000.....	68	37	27	31	32	44	35	23	9	3	1	7
1999.....	70	44	30	34	45	40	27	22	13	3	-	10

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2003.....	127	71	43	31	33	28	20	11	2	1	-	1
2002.....	123	69	40	29	28	33	17	10	4	1	-	1
2001 3/.....	144	80	51	44	30	27	18	7	6	-	1	6
2000.....	100	67	52	49	31	23	24	4	5	1	-	3
1999.....	94	57	34	40	25	29	17	12	3	2	-	9

Black, male

2003.....	98	58	32	25	21	20	11	5	1	-	-	1
2002.....	99	57	35	19	22	22	10	4	2	-	-	-
2001 3/.....	104	56	34	27	21	14	5	4	2	-	1	5
2000.....	80	54	37	38	20	13	14	2	3	1	-	-
1999.....	73	45	21	33	16	15	8	7	2	1	-	5

Black, female

2003.....	29	13	11	6	12	8	9	6	1	1	-	-
2002.....	24	12	5	10	6	11	7	6	2	1	-	1
2001 3/.....	40	24	17	17	9	13	13	3	4	-	-	1
2000.....	20	13	15	11	11	10	10	2	2	-	-	3
1999.....	21	12	13	7	9	14	9	5	1	1	-	4

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

American Indian 1/, both sexes

2003.....	8	1	-	3	2	-	-	1	-	-	-	-
2002.....	9	6	1	5	-	-	3	-	-	-	-	1
2001 3/.....	11	1	1	3	-	-	2	-	1	-	-	-
2000.....	7	2	2	1	2	2	-	2	-	-	-	-
1999.....	10	6	1	2	2	2	2	-	-	-	-	-

American Indian 1/, male

2003.....	7	1	-	2	2	-	-	-	-	-	-	-
2002.....	5	2	1	3	-	-	1	-	-	-	-	-
2001 3/.....	9	1	1	1	-	-	1	-	-	-	-	-
2000.....	6	1	1	1	2	1	-	2	-	-	-	-
1999.....	8	4	1	2	1	1	2	-	-	-	-	-

American Indian 1/, female

2003.....	1	-	-	1	-	-	-	1	-	-	-	-
2002.....	4	4	-	2	-	-	2	-	-	-	-	1
2001 3/.....	2	-	-	2	-	-	1	-	1	-	-	-
2000.....	1	1	1	-	-	1	-	-	-	-	-	-
1999.....	2	2	-	-	1	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Asian or Pacific Islander 2/, both sexes

2003.....	8	9	6	8	6	2	3	-	-	-	-	-
2002.....	7	9	4	5	3	-	1	-	1	-	1	1
2001 3/.....	22	15	17	8	4	3	1	1	-	-	-	-
2000.....	9	7	4	4	3	2	6	1	-	-	-	-
1999.....	8	3	4	1	5	3	3	1	-	-	-	1

Asian or Pacific Islander 2/, male

2003.....	6	6	3	4	3	1	1	-	-	-	-	-
2002.....	5	7	2	2	2	-	-	-	1	-	-	1
2001 3/.....	19	11	8	6	1	-	-	1	-	-	-	-
2000.....	5	4	1	1	3	1	4	-	-	-	-	-
1999.....	6	3	3	-	2	1	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	3	3	4	3	1	2	-	-	-	-	-
2002.....	2	2	2	3	1	-	1	-	-	-	1	-
2001 3/.....	3	4	9	2	3	3	1	-	-	-	-	-
2000.....	4	3	3	3	-	1	2	1	-	-	-	-
1999.....	2	-	1	1	3	2	2	-	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races, both sexes

2003.....	490	245	123	74	57	45	51	36	12	4	-	18
2002.....	431	207	113	80	59	44	51	39	17	8	1	14
2001.....	371	152	73	58	41	53	55	22	11	9	2	18
2000.....	298	154	74	55	49	43	36	26	25	5	1	11
1999.....	265	136	86	72	42	51	45	37	18	2	1	19

All races, male

2003.....	318	153	69	37	28	26	23	15	4	2	-	14
2002.....	264	131	72	46	37	24	22	20	6	5	-	13
2001.....	229	105	41	40	24	28	29	12	6	2	-	16
2000.....	196	105	47	38	32	29	20	14	5	2	-	11
1999.....	175	94	54	47	25	31	24	20	4	1	-	13

All races, female

2003.....	172	92	54	37	29	19	28	21	8	2	-	4
2002.....	167	76	41	34	22	20	29	19	11	3	1	1
2001.....	142	47	32	18	17	25	26	10	5	7	2	2
2000.....	102	49	27	17	17	14	16	12	20	3	1	-
1999.....	90	42	32	25	17	20	21	17	14	1	1	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	403	203	96	59	44	40	45	34	12	4	-	11
2002.....	344	179	93	65	50	37	42	37	16	6	1	12
2001.....	297	129	60	46	37	48	48	20	11	8	2	15
2000.....	240	116	61	45	39	38	34	22	21	5	-	8
1999.....	203	111	75	63	34	47	42	33	16	2	1	14

White, male

2003.....	251	120	51	28	23	24	21	14	4	2	-	8
2002.....	195	110	59	34	28	19	20	19	6	4	-	11
2001.....	175	85	32	32	20	26	27	11	6	2	-	13
2000.....	154	74	38	29	24	27	18	11	4	2	-	8
1999.....	128	70	43	42	19	28	21	18	4	1	-	9

White, female

2003.....	152	83	45	31	21	16	24	20	8	2	-	3
2002.....	149	69	34	31	22	18	22	18	10	2	1	1
2001.....	122	44	28	14	17	22	21	9	5	6	2	2
2000.....	86	42	23	16	15	11	16	11	17	3	-	-
1999.....	75	41	32	21	15	19	21	15	12	1	1	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2003.....	75	31	22	9	12	3	4	2	-	-	-	7
2002.....	80	23	16	10	7	7	9	1	1	2	-	1
2001.....	67	21	11	10	2	2	3	2	-	1	-	2
2000.....	53	31	12	8	7	4	2	3	3	-	1	3
1999.....	59	20	8	8	6	4	1	3	-	-	-	5

Black, male

2003.....	58	25	15	8	4	1	1	1	-	-	-	6
2002.....	63	17	13	9	7	5	2	-	-	1	-	1
2001.....	50	19	8	8	2	-	1	1	-	-	-	2
2000.....	38	24	9	7	6	2	2	2	1	-	-	3
1999.....	46	19	8	5	5	3	1	1	-	-	-	4

Black, female

2003.....	17	6	7	1	8	2	3	1	-	-	-	1
2002.....	17	6	3	1	-	2	7	1	1	1	-	-
2001.....	17	2	3	2	-	2	2	1	-	1	-	-
2000.....	15	7	3	1	1	2	-	1	2	-	1	-
1999.....	13	1	-	3	1	1	-	2	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

American Indian 1/, both sexes

2003.....	6	5	2	1	-	1	-	-	-	-	-	-
2002.....	2	3	1	2	-	-	-	-	-	-	-	1
2001.....	4	-	1	1	-	1	1	-	-	-	-	1
2000.....	3	6	-	-	1	-	-	-	-	-	-	-
1999.....	2	2	1	1	1	-	1	-	1	-	-	-

American Indian 1/, male

2003.....	3	5	2	1	-	-	-	-	-	-	-	-
2002.....	1	2	-	1	-	-	-	-	-	-	-	1
2001.....	2	-	1	-	-	1	1	-	-	-	-	1
2000.....	2	6	-	-	-	-	-	-	-	-	-	-
1999.....	1	2	1	-	1	-	1	-	-	-	-	-

American Indian 1/, female

2003.....	3	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	1	1	1	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 2/, both sexes

2003.....	6	6	3	5	1	1	2	-	-	-	-	-
2002.....	5	2	3	3	2	-	-	1	-	-	-	-
2001.....	3	2	1	1	2	2	3	-	-	-	-	-
2000.....	2	1	1	2	2	1	-	1	1	-	-	-
1999.....	1	3	2	-	1	-	1	1	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	6	3	1	-	1	1	1	-	-	-	-	-
2002.....	5	2	-	2	2	-	-	1	-	-	-	-
2001.....	2	1	-	-	2	1	-	-	-	-	-	-
2000.....	2	1	-	2	2	-	-	1	-	-	-	-
1999.....	-	3	2	-	-	-	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	3	2	5	-	-	1	-	-	-	-	-
2002.....	-	-	3	1	-	-	-	-	-	-	-	-
2001.....	1	1	1	1	-	1	3	-	-	-	-	-
2000.....	-	-	1	-	-	1	-	-	1	-	-	-
1999.....	1	-	-	-	1	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races, both sexes

2003.....	12	14	6	4	4	4	5	1	-	-	-	-
2002.....	10	9	3	8	7	3	2	-	-	-	-	-
2001.....	13	9	3	6	4	6	5	1	-	-	-	-
2000.....	14	12	3	7	4	5	1	2	-	-	-	1
1999.....	10	6	8	9	6	6	7	4	-	-	-	-

All races, male

2003.....	11	12	4	3	4	4	4	1	-	-	-	-
2002.....	9	5	2	4	6	3	1	-	-	-	-	-
2001.....	9	8	3	6	2	6	3	1	-	-	-	-
2000.....	12	10	1	5	4	4	1	2	-	-	-	1
1999.....	6	5	6	8	6	5	7	4	-	-	-	-

All races, female

2003.....	1	2	2	1	-	-	1	-	-	-	-	-
2002.....	1	4	1	4	1	-	1	-	-	-	-	-
2001.....	4	1	-	-	2	-	2	-	-	-	-	-
2000.....	2	2	2	2	-	1	-	-	-	-	-	-
1999.....	4	1	2	1	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

White, both sexes

2003.....	11	12	5	4	4	4	5	1	-	-	-	-
2002.....	9	8	2	8	6	2	2	-	-	-	-	-
2001.....	11	6	3	6	4	6	4	1	-	-	-	-
2000.....	12	8	3	6	3	5	1	2	-	-	-	1
1999.....	10	5	7	9	4	6	6	4	-	-	-	-

White, male

2003.....	10	11	4	3	4	4	4	1	-	-	-	-
2002.....	8	5	2	4	5	2	1	-	-	-	-	-
2001.....	7	5	3	6	2	6	2	1	-	-	-	-
2000.....	10	6	1	4	3	4	1	2	-	-	-	1
1999.....	6	4	5	8	4	5	6	4	-	-	-	-

White, female

2003.....	1	1	1	1	-	-	1	-	-	-	-	-
2002.....	1	3	-	4	1	-	1	-	-	-	-	-
2001.....	4	1	-	-	2	-	2	-	-	-	-	-
2000.....	2	2	2	2	-	1	-	-	-	-	-	-
1999.....	4	1	2	1	-	1	-	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races, both sexes

2003.....	478	231	117	70	53	41	46	35	12	4	-	18
2002.....	421	198	110	72	52	41	49	39	17	8	1	14
2001.....	358	143	70	52	37	47	50	21	11	9	2	18
2000.....	284	142	71	48	45	38	35	24	25	5	1	10
1999.....	255	130	78	63	36	45	38	33	18	2	1	19

All races, male

2003.....	307	141	65	34	24	22	19	14	4	2	-	14
2002.....	255	126	70	42	31	21	21	20	6	5	-	13
2001.....	220	97	38	34	22	22	26	11	6	2	-	16
2000.....	184	95	46	33	28	25	19	12	5	2	-	10
1999.....	169	89	48	39	19	26	17	16	4	1	-	13

All races, female

2003.....	171	90	52	36	29	19	27	21	8	2	-	4
2002.....	166	72	40	30	21	20	28	19	11	3	1	1
2001.....	138	46	32	18	15	25	24	10	5	7	2	2
2000.....	100	47	25	15	17	13	16	12	20	3	1	-
1999.....	86	41	30	24	17	19	21	17	14	1	1	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2003.....	392	191	91	55	40	36	40	33	12	4	-	11
2002.....	335	171	91	57	44	35	40	37	16	6	1	12
2001.....	286	123	57	40	33	42	44	19	11	8	2	15
2000.....	228	108	58	39	36	33	33	20	21	5	-	7
1999.....	193	106	68	54	30	41	36	29	16	2	1	14

White, male

2003.....	241	109	47	25	19	20	17	13	4	2	-	8
2002.....	187	105	57	30	23	17	19	19	6	4	-	11
2001.....	168	80	29	26	18	20	25	10	6	2	-	13
2000.....	144	68	37	25	21	23	17	9	4	2	-	7
1999.....	122	66	38	34	15	23	15	14	4	1	-	9

White, female

2003.....	151	82	44	30	21	16	23	20	8	2	-	3
2002.....	148	66	34	27	21	18	21	18	10	2	1	1
2001.....	118	43	28	14	15	22	19	9	5	6	2	2
2000.....	84	40	21	14	15	10	16	11	17	3	-	-
1999.....	71	40	30	20	15	18	21	15	12	1	1	5

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2003.....	74	30	22	9	12	3	4	2	-	-	-	7
2002.....	79	22	16	10	6	6	9	1	1	2	-	1
2001.....	65	18	11	10	2	2	2	2	-	1	-	2
2000.....	52	28	12	7	6	4	2	3	3	-	1	3
1999.....	59	19	8	8	4	4	-	3	-	-	-	5

Black, male

2003.....	57	24	15	8	4	1	1	1	-	-	-	6
2002.....	62	17	13	9	6	4	2	-	-	1	-	1
2001.....	48	16	8	8	2	-	-	1	-	-	-	2
2000.....	37	21	9	6	5	2	2	2	1	-	-	3
1999.....	46	18	8	5	3	3	-	1	-	-	-	4

Black, female

2003.....	17	6	7	1	8	2	3	1	-	-	-	1
2002.....	17	5	3	1	-	2	7	1	1	1	-	-
2001.....	17	2	3	2	-	2	2	1	-	1	-	-
2000.....	15	7	3	1	1	2	-	1	2	-	1	-
1999.....	13	1	-	3	1	1	-	2	-	-	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

American Indian 1/, both sexes

2003.....	6	5	2	1	-	1	-	-	-	-	-	-
2002.....	2	3	-	2	-	-	-	-	-	-	-	1
2001.....	4	-	1	1	-	1	1	-	-	-	-	1
2000.....	3	5	-	-	1	-	-	-	-	-	-	-
1999.....	2	2	1	1	1	-	1	-	1	-	-	-

American Indian 1/, male

2003.....	3	5	2	1	-	-	-	-	-	-	-	-
2002.....	1	2	-	1	-	-	-	-	-	-	-	1
2001.....	2	-	1	-	-	1	1	-	-	-	-	1
2000.....	2	5	-	-	-	-	-	-	-	-	-	-
1999.....	1	2	1	-	1	-	1	-	-	-	-	-

American Indian 1/, female

2003.....	3	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	1	-	1	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 2/, both sexes

2003.....	6	5	2	5	1	1	2	-	-	-	-	-
2002.....	5	2	3	3	2	-	-	1	-	-	-	-
2001.....	3	2	1	1	2	2	3	-	-	-	-	-
2000.....	1	1	1	2	2	1	-	1	1	-	-	-
1999.....	1	3	1	-	1	-	1	1	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	6	3	1	-	1	1	1	-	-	-	-	-
2002.....	5	2	-	2	2	-	-	1	-	-	-	-
2001.....	2	1	-	-	2	1	-	-	-	-	-	-
2000.....	1	1	-	2	2	-	-	1	-	-	-	-
1999.....	-	3	1	-	-	-	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	-	2	1	5	-	-	1	-	-	-	-	-
2002.....	-	-	3	1	-	-	-	-	-	-	-	-
2001.....	1	1	1	1	-	1	3	-	-	-	-	-
2000.....	-	-	1	-	-	1	-	-	1	-	-	-
1999.....	1	-	-	-	1	-	-	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

All races, both sexes

2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	7	1	1	-	-	-	-
2000.....	2	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	4	2	1	-	-	-

All races, male

2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	6	1	1	-	-	-	-
2000.....	2	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	3	2	1	-	-	-

All races, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

White, both sexes

2003.....	1	4	-	-	2	2	3	-	-	-	-	-
2002.....	-	1	1	2	-	4	5	1	-	-	-	-
2001.....	1	-	-	-	3	7	1	1	-	-	-	-
2000.....	1	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	1	6	4	4	2	1	-	-	-

White, male

2003.....	1	4	-	-	2	2	3	-	-	-	-	-
2002.....	-	1	1	2	-	4	5	1	-	-	-	-
2001.....	1	-	-	-	3	6	1	1	-	-	-	-
2000.....	1	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	1	6	4	3	2	1	-	-	-

White, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races, both sexes

2003.....	128	178	202	214	320	398	396	307	152	58	13	1
2002.....	151	164	187	225	335	382	410	288	167	59	7	-
2001.....	141	178	206	247	320	439	425	315	186	56	7	1
2000.....	167	157	219	263	384	427	398	337	176	54	11	-
1999.....	123	138	177	229	357	401	410	312	153	52	12	-

All races, male

2003.....	66	101	94	103	161	174	181	129	46	8	3	1
2002.....	67	84	85	110	146	166	154	110	40	18	3	-
2001.....	68	91	105	116	149	192	191	119	55	13	2	-
2000.....	84	77	104	123	208	212	190	134	48	13	1	-
1999.....	61	77	88	102	177	193	182	128	49	17	3	-

All races, female

2003.....	62	77	108	111	159	224	215	178	106	50	10	-
2002.....	84	80	102	115	189	216	256	178	127	41	4	-
2001.....	73	87	101	131	171	247	234	196	131	43	5	1
2000.....	83	80	115	140	176	215	208	203	128	41	10	-
1999.....	62	61	89	127	180	208	228	184	104	35	9	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

White, both sexes

2003.....	89	133	164	178	267	336	360	269	135	54	11	1
2002.....	105	122	149	178	292	331	366	261	146	51	7	-
2001.....	107	137	156	198	247	387	380	267	160	50	6	1
2000.....	134	120	165	215	320	361	357	297	155	41	10	-
1999.....	91	98	142	193	299	343	356	266	137	44	8	-

White, male

2003.....	43	78	79	90	139	154	169	114	43	7	3	1
2002.....	50	65	70	84	128	147	143	97	35	14	3	-
2001.....	49	66	79	95	112	171	176	103	45	11	1	-
2000.....	70	64	80	98	175	186	177	119	44	11	1	-
1999.....	46	55	71	87	148	169	159	108	43	16	1	-

White, female

2003.....	46	55	85	88	128	182	191	155	92	47	8	-
2002.....	55	57	79	94	164	184	223	164	111	37	4	-
2001.....	58	71	77	103	135	216	204	164	115	39	5	1
2000.....	64	56	85	117	145	175	180	178	111	30	9	-
1999.....	45	43	71	106	151	174	197	158	94	28	7	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2003.....	34	39	35	34	44	56	30	31	15	4	2	-
2002.....	44	38	32	38	39	44	36	21	19	7	-	-
2001.....	31	38	47	43	63	46	41	46	23	6	1	-
2000.....	31	35	51	42	59	59	37	34	20	12	1	-
1999.....	31	34	28	32	54	53	43	37	16	7	3	-

Black, male

2003.....	21	20	15	13	15	17	10	10	3	1	-	-
2002.....	16	18	12	20	17	14	11	10	5	4	-	-
2001.....	18	22	25	21	33	17	14	14	9	2	1	-
2000.....	13	12	22	24	28	22	12	12	3	1	-	-
1999.....	15	16	14	13	27	22	18	12	6	1	1	-

Black, female

2003.....	13	19	20	21	29	39	20	21	12	3	2	-
2002.....	28	20	20	18	22	30	25	11	14	3	-	-
2001.....	13	16	22	22	30	29	27	32	14	4	-	-
2000.....	18	23	29	18	31	37	25	22	17	11	1	-
1999.....	16	18	14	19	27	31	25	25	10	6	2	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

American Indian 1/, both sexes

2003.....	1	1	3	1	2	1	2	2	1	-	-	-
2002.....	1	3	2	6	1	2	2	2	-	-	-	-
2001.....	2	1	-	1	2	3	1	-	1	-	-	-
2000.....	1	-	-	4	1	1	2	3	-	-	-	-
1999.....	1	2	1	2	1	2	3	2	-	1	1	-

American Indian 1/, male

2003.....	-	-	-	-	1	-	-	2	-	-	-	-
2002.....	1	-	1	4	-	1	-	1	-	-	-	-
2001.....	1	1	-	-	-	2	-	-	-	-	-	-
2000.....	1	-	-	-	1	1	1	1	-	-	-	-
1999.....	-	2	1	1	1	1	3	1	-	-	1	-

American Indian 1/, female

2003.....	1	1	3	1	1	1	2	-	1	-	-	-
2002.....	-	3	1	2	1	1	2	1	-	-	-	-
2001.....	1	-	-	1	2	1	1	-	1	-	-	-
2000.....	-	-	-	4	-	-	1	2	-	-	-	-
1999.....	1	-	-	1	-	1	-	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Asian or Pacific Islander 2/, both sexes

2003.....	4	5	-	1	7	5	4	5	1	-	-	-
2002.....	1	1	4	3	3	5	6	4	2	1	-	-
2001.....	1	2	3	5	8	3	3	2	2	-	-	-
2000.....	1	2	3	2	4	6	2	3	1	1	-	-
1999.....	-	4	6	2	3	3	8	7	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	2	3	-	-	6	3	2	3	-	-	-	-
2002.....	-	1	2	2	1	4	-	2	-	-	-	-
2001.....	-	2	1	-	4	2	1	2	1	-	-	-
2000.....	-	1	2	1	4	3	-	2	1	1	-	-
1999.....	-	4	2	1	1	1	2	7	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	2	2	-	1	1	2	2	2	1	-	-	-
2002.....	1	-	2	1	2	1	6	2	2	1	-	-
2001.....	1	-	2	5	4	1	2	-	1	-	-	-
2000.....	1	1	1	1	-	3	2	1	-	-	-	-
1999.....	-	-	4	1	2	2	6	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

All races, both sexes

2003.....	3,591	2,891	2,277	1,641	1,130	770	413	149	38	7	-	11
2002 5/.....	3,427	2,734	2,154	1,582	1,253	806	384	154	39	8	-	8
2001 5/.....	3,305	2,495	2,104	1,645	1,201	816	388	166	40	14	1	16
2000 5/.....	2,940	2,455	2,087	1,644	1,266	831	392	162	55	7	3	8
1999 5/.....	2,853	2,374	2,065	1,676	1,250	876	415	175	43	11	1	8

All races, male

2003.....	2,787	2,233	1,764	1,249	871	558	308	90	20	3	-	10
2002 5/.....	2,686	2,123	1,714	1,240	918	575	270	104	26	5	-	8
2001 5/.....	2,523	1,969	1,638	1,257	924	592	266	115	27	8	1	15
2000 5/.....	2,315	1,874	1,615	1,273	976	641	274	109	32	4	1	8
1999 5/.....	2,266	1,880	1,601	1,317	953	637	299	114	24	3	-	7

All races, female

2003.....	804	658	513	392	259	212	105	59	18	4	-	1
2002 5/.....	741	611	440	342	335	231	114	50	13	3	-	-
2001 5/.....	782	526	466	388	277	224	122	51	13	6	-	1
2000 5/.....	625	581	472	371	290	190	118	53	23	3	2	-
1999 5/.....	587	494	464	359	297	239	116	61	19	8	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

White, both sexes

2003.....	2,965	2,452	1,888	1,420	993	702	372	136	35	6	-	8
2002 5/.....	2,857	2,274	1,784	1,360	1,088	720	341	137	35	8	-	6
2001 5/.....	2,643	2,035	1,775	1,373	1,065	726	357	142	35	14	1	13
2000 5/.....	2,402	2,009	1,730	1,396	1,110	742	352	146	49	6	3	6
1999 5/.....	2,327	1,915	1,702	1,409	1,072	763	369	151	38	11	1	8

White, male

2003.....	2,300	1,905	1,459	1,085	768	507	272	81	18	3	-	7
2002 5/.....	2,266	1,780	1,426	1,060	790	515	240	89	23	5	-	6
2001 5/.....	2,050	1,599	1,380	1,041	815	528	245	99	23	8	1	12
2000 5/.....	1,910	1,554	1,345	1,084	853	572	249	98	27	3	1	6
1999 5/.....	1,869	1,514	1,339	1,098	814	554	270	97	23	3	-	7

White, female

2003.....	665	547	429	335	225	195	100	55	17	3	-	1
2002 5/.....	591	494	358	300	298	205	101	48	12	3	-	-
2001 5/.....	593	436	395	332	250	198	112	43	12	6	-	1
2000 5/.....	492	455	385	312	257	170	103	48	22	3	2	-
1999 5/.....	458	401	363	311	258	209	99	54	15	8	1	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Black, both sexes

2003.....	488	354	316	181	108	53	34	12	1	1	-	3
2002 5/.....	457	377	291	176	132	73	35	14	2	-	-	1
2001 5/.....	551	381	275	234	109	78	27	16	3	-	-	3
2000 5/.....	451	359	286	216	132	76	29	13	6	1	-	2
1999 5/.....	427	385	310	225	157	92	40	20	4	-	-	-

Black, male

2003.....	390	265	255	134	84	38	29	9	1	-	-	3
2002 5/.....	340	282	230	146	105	51	25	12	1	-	-	1
2001 5/.....	406	306	218	188	88	56	18	9	2	-	-	3
2000 5/.....	343	265	213	165	108	60	20	9	5	1	-	2
1999 5/.....	324	312	229	186	122	66	26	13	1	-	-	-

Black, female

2003.....	98	89	61	47	24	15	5	3	-	1	-	-
2002 5/.....	117	95	61	30	27	22	10	2	1	-	-	-
2001 5/.....	145	75	57	46	21	22	9	7	1	-	-	-
2000 5/.....	108	94	73	51	24	16	9	4	1	-	-	-
1999 5/.....	103	73	81	39	35	26	14	7	3	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

American Indian 1/, both sexes

2003.....	113	68	54	31	20	11	2	1	-	-	-	-
2002 5/.....	83	59	58	30	22	6	3	1	-	-	-	-
2001 5/.....	86	54	40	30	18	5	2	4	2	-	-	-
2000 5/.....	70	63	47	28	14	8	4	1	-	-	-	-
1999 5/.....	83	60	46	28	12	14	2	2	-	-	-	-

American Indian 1/, male

2003.....	75	50	34	23	10	9	2	-	-	-	-	-
2002 5/.....	59	40	40	21	17	4	1	1	-	-	-	-
2001 5/.....	50	43	29	22	13	2	1	4	2	-	-	-
2000 5/.....	47	39	36	21	7	5	3	1	-	-	-	-
1999 5/.....	63	41	28	21	9	11	1	2	-	-	-	-

American Indian 1/, female

2003.....	38	18	20	8	10	2	-	1	-	-	-	-
2002 5/.....	24	19	18	9	5	2	2	-	-	-	-	-
2001 5/.....	36	11	11	8	5	3	1	-	-	-	-	-
2000 5/.....	23	24	11	7	7	3	1	-	-	-	-	-
1999 5/.....	20	19	18	7	3	3	1	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Asian or Pacific Islander 2/, both sexes

2003.....	25	17	19	9	9	4	5	-	2	-	-	-
2002 5/.....	30	24	21	16	11	7	5	2	2	-	-	1
2001 5/.....	25	25	14	8	9	7	2	4	-	-	-	-
2000 5/.....	17	24	24	4	10	5	7	2	-	-	-	-
1999 5/.....	16	14	7	14	9	7	4	2	1	-	-	-

Asian or Pacific Islander 2/, male

2003.....	22	13	16	7	9	4	5	-	1	-	-	-
2002 5/.....	21	21	18	13	6	5	4	2	2	-	-	1
2001 5/.....	17	21	11	6	8	6	2	3	-	-	-	-
2000 5/.....	15	16	21	3	8	4	2	1	-	-	-	-
1999 5/.....	10	13	5	12	8	6	2	2	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	3	4	3	2	-	-	-	-	1	-	-	-
2002 5/.....	9	3	3	3	5	2	1	-	-	-	-	-
2001 5/.....	8	4	3	2	1	1	-	1	-	-	-	-
2000 5/.....	2	8	3	1	2	1	5	1	-	-	-	-
1999 5/.....	6	1	2	2	1	1	2	-	1	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

All races, both sexes

2003.....	3,228	1,546	693	441	314	288	249	187	79	13	4	12
2002 5/.....	2,731	1,235	593	365	288	264	220	159	84	15	2	10
2001 5/.....	2,136	901	460	296	257	254	185	136	69	17	4	16
2000 5/.....	1,815	794	393	244	242	235	195	150	75	15	-	14
1999 5/.....	1,582	760	401	301	255	264	203	134	49	17	1	18

All races, male

2003.....	2,022	883	383	234	150	145	108	72	35	4	1	11
2002 5/.....	1,752	724	316	181	130	120	95	62	33	7	-	7
2001 5/.....	1,353	530	230	164	129	116	75	45	23	5	1	16
2000 5/.....	1,188	433	198	127	108	109	78	63	25	4	-	13
1999 5/.....	1,012	451	237	164	122	104	80	51	16	7	-	11

All races, female

2003.....	1,206	663	310	207	164	143	141	115	44	9	3	1
2002 5/.....	979	511	277	184	158	144	125	97	51	8	2	3
2001 5/.....	783	371	230	132	128	138	110	91	46	12	3	-
2000 5/.....	627	361	195	117	134	126	117	87	50	11	-	1
1999 5/.....	570	309	164	137	133	160	123	83	33	10	1	7

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

White, both sexes

2003.....	2,621	1,286	585	371	282	276	237	175	76	12	4	7
2002 5/.....	2,177	1,017	485	319	264	242	208	145	79	13	2	8
2001 5/.....	1,680	742	373	256	231	234	173	124	64	13	2	15
2000 5/.....	1,406	655	335	209	223	224	183	135	68	11	-	10
1999 5/.....	1,232	614	326	261	228	248	194	117	48	15	1	13

White, male

2003.....	1,604	704	307	184	133	137	103	66	32	3	1	6
2002 5/.....	1,349	562	233	146	117	109	84	57	33	6	-	5
2001 5/.....	1,021	410	164	140	111	99	72	40	23	3	-	15
2000 5/.....	888	332	158	103	93	105	73	57	21	2	-	9
1999 5/.....	750	338	175	136	109	94	76	47	16	6	-	7

White, female

2003.....	1,017	582	278	187	149	139	134	109	44	9	3	1
2002 5/.....	828	455	252	173	147	133	124	88	46	7	2	3
2001 5/.....	659	332	209	116	120	135	101	84	41	10	2	-
2000 5/.....	518	323	177	106	130	119	110	78	47	9	-	1
1999 5/.....	482	276	151	125	119	154	118	70	32	9	1	6

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Black, both sexes

2003.....	549	231	92	61	24	12	10	10	3	1	-	5
2002 5/.....	525	203	102	35	17	17	9	8	4	1	-	1
2001 5/.....	435	145	79	32	22	17	8	10	5	3	2	1
2000 5/.....	392	130	53	33	17	11	11	14	6	4	-	4
1999 5/.....	326	128	70	35	22	12	8	15	1	2	-	5

Black, male

2003.....	381	166	68	47	14	8	4	5	3	1	-	5
2002 5/.....	384	153	81	28	10	9	8	2	-	-	-	1
2001 5/.....	318	109	62	21	16	14	1	5	-	1	1	1
2000 5/.....	289	97	38	23	13	4	5	6	3	2	-	4
1999 5/.....	245	104	57	25	12	7	4	4	-	1	-	4

Black, female

2003.....	168	65	24	14	10	4	6	5	-	-	-	-
2002 5/.....	141	50	21	7	7	8	1	6	4	1	-	-
2001 5/.....	117	36	17	11	6	3	7	5	5	2	1	-
2000 5/.....	103	33	15	10	4	7	6	8	3	2	-	-
1999 5/.....	81	24	13	10	10	5	4	11	1	1	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

American Indian 1/, both sexes

2003.....	35	13	9	5	1	-	-	-	-	-	-	-
2002 5/.....	17	8	2	2	1	-	-	2	-	-	-	1
2001 5/.....	12	6	5	3	2	1	1	1	-	-	-	-
2000 5/.....	10	7	-	-	1	-	-	1	-	-	-	-
1999 5/.....	10	8	-	-	1	1	-	1	-	-	-	-

American Indian 1/, male

2003.....	22	6	4	1	1	-	-	-	-	-	-	-
2002 5/.....	12	5	1	2	1	-	-	-	-	-	-	1
2001 5/.....	8	5	2	1	1	1	1	-	-	-	-	-
2000 5/.....	7	4	-	-	1	-	-	-	-	-	-	-
1999 5/.....	5	6	-	-	-	1	-	-	-	-	-	-

American Indian 1/, female

2003.....	13	7	5	4	-	-	-	-	-	-	-	-
2002 5/.....	5	3	1	-	-	-	-	2	-	-	-	-
2001 5/.....	4	1	3	2	1	-	-	1	-	-	-	-
2000 5/.....	3	3	-	-	-	-	-	1	-	-	-	-
1999 5/.....	5	2	-	-	1	-	-	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5, F12.7-F12.9,F13.0-F13.5,F13.7-F13.9, F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1, G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Asian or Pacific Islander 2/, both sexes

2003.....	23	16	7	4	7	-	2	2	-	-	-	-
2002 5/.....	12	7	4	9	6	5	3	4	1	1	-	-
2001 5/.....	9	8	3	5	2	2	3	1	-	1	-	-
2000 5/.....	7	2	5	2	1	-	1	-	1	-	-	-
1999 5/.....	14	10	5	5	4	3	1	1	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	15	7	4	2	2	-	1	1	-	-	-	-
2002 5/.....	7	4	1	5	2	2	3	3	-	1	-	-
2001 5/.....	6	6	2	2	1	2	1	-	-	1	-	-
2000 5/.....	4	-	2	1	1	-	-	-	1	-	-	-
1999 5/.....	12	3	5	3	1	2	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2003.....	8	9	3	2	5	-	1	1	-	-	-	-
2002 5/.....	5	3	3	4	4	3	-	1	1	-	-	-
2001 5/.....	3	2	1	3	1	-	2	1	-	-	-	-
2000 5/.....	3	2	3	1	-	-	1	-	-	-	-	-
1999 5/.....	2	7	-	2	3	1	1	1	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

All races, both sexes

2003.....	2,075	1,637	1,167	945	967	963	768	420	144	25	-	18
2002.....	1,963	1,608	1,111	966	1,027	1,011	822	411	138	27	-	15
2001.....	1,917	1,471	1,073	999	999	1,014	789	402	139	20	2	19
2000.....	1,805	1,289	983	923	1,018	994	729	443	137	19	1	18
1999.....	1,626	1,299	1,007	955	1,079	1,029	709	428	108	25	-	20

All races, male

2003.....	1,736	1,362	1,016	835	859	863	695	388	136	22	-	16
2002.....	1,652	1,354	942	836	932	916	755	386	128	26	-	14
2001.....	1,588	1,231	885	865	889	915	719	373	128	19	2	18
2000.....	1,488	1,071	825	783	901	900	671	404	125	19	1	13
1999.....	1,335	1,080	871	833	953	928	652	392	103	22	-	19

All races, female

2003.....	339	275	151	110	108	100	73	32	8	3	-	2
2002.....	311	254	169	130	95	95	67	25	10	1	-	1
2001.....	329	240	188	134	110	99	70	29	11	1	-	1
2000.....	317	218	158	140	117	94	58	39	12	-	-	5
1999.....	291	219	136	122	126	101	57	36	5	3	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

White, both sexes

2003.....	1,820	1,497	1,078	883	911	918	736	411	141	23	-	14
2002.....	1,699	1,471	1,014	906	974	949	800	398	135	25	-	13
2001.....	1,694	1,328	997	931	935	968	751	393	135	19	2	16
2000.....	1,573	1,152	904	858	951	963	703	432	134	18	1	15
1999.....	1,430	1,166	902	888	1,019	980	682	418	106	23	-	15

White, male

2003.....	1,528	1,246	941	784	814	827	669	380	134	22	-	13
2002.....	1,438	1,233	866	785	886	861	735	374	126	24	-	12
2001.....	1,406	1,102	821	806	835	875	687	365	124	18	2	15
2000.....	1,293	955	755	728	840	876	649	396	122	18	1	10
1999.....	1,185	959	781	778	899	886	628	383	101	21	-	14

White, female

2003.....	292	251	137	99	97	91	67	31	7	1	-	1
2002.....	261	238	148	121	88	88	65	24	9	1	-	1
2001.....	288	226	176	125	100	93	64	28	11	1	-	1
2000.....	280	197	149	130	111	87	54	36	12	-	-	5
1999.....	245	207	121	110	120	94	54	35	5	2	-	1

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Black, both sexes

2003.....	220	115	74	56	47	35	25	7	3	2	-	3
2002.....	216	109	81	48	50	51	17	13	3	2	-	2
2001.....	193	115	61	55	55	40	31	7	4	1	-	2
2000.....	198	100	63	56	58	26	24	10	2	-	-	1
1999.....	153	111	85	58	51	47	21	10	2	2	-	5

Black, male

2003.....	180	98	61	46	37	29	20	7	2	-	-	2
2002.....	174	97	62	41	44	45	16	12	2	2	-	2
2001.....	161	104	51	48	47	35	26	6	4	1	-	2
2000.....	167	83	54	49	54	20	20	7	2	-	-	1
1999.....	123	102	73	49	46	41	18	9	2	1	-	5

Black, female

2003.....	40	17	13	10	10	6	5	-	1	2	-	1
2002.....	42	12	19	7	6	6	1	1	1	-	-	-
2001.....	32	11	10	7	8	5	5	1	-	-	-	-
2000.....	31	17	9	7	4	6	4	3	-	-	-	-
1999.....	30	9	12	9	5	6	3	1	-	1	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Asian or Pacific Islander 2/, both sexes

2003.....	28	20	11	4	7	7	6	1	-	-	-	1
2002.....	32	22	12	8	2	7	5	-	-	-	-	-
2001.....	23	23	9	10	5	4	5	2	-	-	-	1
2000.....	22	28	11	4	5	4	2	1	1	1	-	1
1999.....	29	19	15	5	4	2	3	-	-	-	-	-

Asian or Pacific Islander 2/, male

2003.....	22	15	10	3	6	5	5	-	-	-	-	1
2002.....	25	19	11	7	1	6	4	-	-	-	-	-
2001.....	16	20	8	9	3	3	4	2	-	-	-	1
2000.....	18	24	11	2	3	3	2	1	1	1	-	1
1999.....	17	16	13	2	3	1	3	-	-	-	-	-

Worktable 291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2003

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Asian or Pacific Islander 2/, female

2003.....	6	5	1	1	1	2	1	1	-	-	-	-
2002.....	7	3	1	1	1	1	1	-	-	-	-	-
2001.....	7	3	1	1	2	1	1	-	-	-	-	-
2000.....	4	4	-	2	2	1	-	-	-	-	-	-
1999.....	12	3	2	3	1	1	-	-	-	-	-	-

- Quantity zero.

... Category not applicable.

1/ Includes Aleuts and Eskimos.

2/ Includes Chinese, Filipino, Hawaiian, Japanese, and other Asian or Pacific Islander.

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

4/ Included in selected categories.

5/ Figures may differ from those previously shown as they are based on a modified list of causes; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003."

Source: CDC/NCHS, National Vital Statistics System, Mortality.