CAUSE OF DEATH (BASED ON T TENTH REVISION INTERNATION CLASSIFICATION OF DISEASES 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	AL	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
·	AGES	TEAR	TEMICS	12/11(3	TEARS	TEARS	TEARS	12/1103	TEARS	TEARS	TEARS
All causes											
All origins	2,391,399	27,937	5,249	3,474	4,121	13,778	16,878	17,703	23,363	36,804	52,452
Male	1,175,460	15,646	2,975	1,964	2,528	9,780	12,634	12,542	15,734	23,681	33,437
Female	1,215,939	12,291	2,274	1,510	1,593	3,998	4,244	5,161	7,629	13,123	19,015
Hispanic	103,740	4,412	882	464	550	1,871	2,634	2,527	2,870	3,624	4,174
Male	57,991	2,411	481	266	326	1,432	2,115	1,956	2,128	2,601	2,875
Female	45,749	2,001	401	198	224	439	519	571	742	1,023	1,299
Mexican	57,425	3,159	637	326	371	1,300	1,856	1,711	1,741	2,069	2,361
Male	33,099	1,721	353	191	226	1,008	1,516	1,351	1,328	1,541	1,638
Female	24,326	1,438	284	135	145	292	340	360	413	528	723
Puerto Rican	13,909	426	71	48	51	180	234	269	412	593	727
Male	7,807	225	43	24	30	125	175	193	270	388	485
Female	6,102	201	28	24	21	55	59	76	142	205	242
Cuban	11,029	66	12	7	14	29	35	48	69	149	170
Male	5,561	34	5	5	8	18	26	38	49	110	126
Female	5,468	32	7	2	6	11	9	10	20	39	44
Central and South America	n 9,064	295	73	45	52	183	301	330	370	429	447
Male	4,708	156	36	27	31	141	236	258	272	284	300
Female	4,356	139	37	18	21	42	65	72	98	145	147
Other and unknown Hispani	c 12,313	466	89	38	62	179	208	169	278	384	469
Male	6,816	275	44	19	31	140	162	116	209	278	326
Female	5,497	191	45	19	31	39	46	53	69	106	143

PAGE

2

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS All causes Non-Hispanic 1/ 2,279,325 23,126 4,343 3,001 3,559 11,861 14,182 15,091 20,380 32,997 47,952 Male 1,112,718 12,998 2,479 1,692 2,193 8,323 10,474 10,525 13,522 20,947 30,325 Female 1,166,607 10,128 1,864 1,309 1,366 3,538 3,708 4,566 6,858 12,050 17,627 8,490 Non-Hispanic white 1,953,197 13,553 2,819 1,990 2,498 9,378 10,048 13,938 23,345 34,285 Male 944,913 7,718 1,605 1,115 1,528 5,799 6,878 7,056 9,393 15,144 22,147 Female 1,008,284 5,835 1,214 875 970 2,691 2,500 2,992 4,545 8,201 12,138 281,979 1,283 2,813 5,602 Non-Hispanic black 8,589 869 902 4,167 4,390 8,563 12,207 Male 143,883 4,765 734 508 573 2,133 3,153 3,595 5,104 7,280 3,038 Female 138,096 3,824 549 361 329 680 1,014 1,352 2,007 3,459 4,927 Origin not stated 2/ 8,334 399 24 9 12 46 62 85 113 183 326 Male 4,751 237 15 6 9 25 45 61 84 133 237 Female 3,583 162 9 3 3 21 17 24 29 50 89

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 3 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Salmonella infections (A01-A02) All origins 38 3 2 2 Male 21 2 2 1 Female 17 1 1 Hispanic Male 3 Female 2 Mexican Male Female Puerto Rican Male Female Cuban 1 Male Female 1 Central and South American Male Female Other and unknown Hispanic 1 Male 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Salmonella infections (A01-A02) Non-Hispanic 1/ 34 3 2 2 Male 18 2 1 Female 16 1 3 Non-Hispanic white 28 1 2 Male 13 1 Female 15 1 1 Non-Hispanic black 4 Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Shigellosis and amebiasis (A03,A06) All origins 10 1 1 1 Male 6 1 1 1 Female Hispanic 1 1 1 Male 3 1 1 1 Female 3 Mexican 1 1 Male 1 Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic 1 1 Male 1 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Shigellosis and amebiasis (A0	3,A06)										
Non-Hispanic 1/	6	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white	6	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Certain other intestinal infections (A04,A07-A09) All origins 1,108 12 8 2 1 1 4 3 3 2 5 Male 420 8 4 2 1 1 2 2 3 1 3 Female 688 4 4 2 1 1 2 Hispanic 34 3 1 1 1 2 Male 17 1 1 1 Female 17 1 1 19 3 Mexican 1 1 9 Male Female 10 1 Puerto Rican 3 1 Male 2 1 Female 1 Cuban 5

Source: CDC/NCHS, National Vital Statistics System, Mortality

2

3

3

2

1

4

2

2

CAUSE OF DEATH (BASED ON THE

Male

Male

Male

Female

Female

Female

Central and South American

Other and unknown Hispanic

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Certain other intestinal infections (A04,A07-A09) Non-Hispanic 1/ 1,071 9 7 2 1 3 2 3 2 5 6 Male 403 4 2 1 1 1 3 1 3 2 Female 668 3 3 1 1 2 990 5 2 3 2 Non-Hispanic white 6 1 2 Male 373 3 3 1 2 Female 617 1 3 1 Non-Hispanic black 74 3 Male 25 2 1 Female 49 Origin not stated 2/ 3 Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	ALL	UNDER 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44
POPULATION, AND SEX	AGES	YEAR	YEARS								
Tuberculosis (A16-A19)											
All origins	930	2	2	1	2	1	10	3	16	25	32
Male	584	-	2	-	2	1	3	1	10	20	23
Female	346	2	-	1	-	-	7	2	6	5	9
Hispanic	125	1	-	-	1	1	3	1	2	7	6
Male	78	-	-	-	1	1	2	1	2	5	4
Female	47	1	-	-	-	-	1	-	-	2	2
Mexican	85	1	-	-	1	1	2	1	2	4	5
Male	56	-	-	-	1	1	1	1	2	3	3
Female	29	1	-	-	-	-	1	-	-	1	2
Puerto Rican	10	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-
Cuban	10	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-
Central and South American	12	-	-	-	-	-	1	-	-	3	-
Male	9	-	-	-	-	-	1	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	1	-
Other and unknown Hispanic	8	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	_	-	-	-	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Tuberculosis (A16-A19) Non-Hispanic 1/ 788 1 2 1 1 6 2 14 18 24 Male 492 2 1 8 15 17 Female 296 1 1 6 2 6 3 7 5 Non-Hispanic white 430 1 1 6 Male 263 1 4 5 Female 167 1 1 1 Non-Hispanic black 239 4 1 8 10 14 Male 154 4 8 10 Female 85 4 1 2 4 Origin not stated 2/ 17 1 2 Male 14 1 2 Female 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Respiratory tuberculosis (A16) All origins 764 1 1 8 2 12 19 1 26 Male 486 1 1 2 8 15 18 Female 278 1 6 2 4 4 8 Hispanic 93 2 1 5 5 3 Male 55 1 1 3 2 Female 38 1 2 2 4 Mexican 58 1 2 Male 36 Female 22 1 1 2 Puerto Rican 9 1 Male 3 1 Female 6 Cuban 9 Male 3 Female 6 Central and South American 1 3 11 Male 8 1 2 Female 3 1 Other and unknown Hispanic 6 Male 5

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Respiratory tuberculosis (A16) Non-Hispanic 1/ 657 1 1 5 2 11 14 19 1 7 Male 419 1 1 12 13 Female 238 1 5 2 4 2 6 371 3 Non-Hispanic white 1 1 6 Male 234 1 3 5 Female 137 1 1 Non-Hispanic black 194 4 9 10 Male 122 3 7 6 Female 72 4 2 4 Origin not stated 2/ 14 1 2 Male 12 1 2

Source: CDC/NCHS, National Vital Statistics System, Mortality

2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 35-39 40-44 ALL 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other tuberculosis (A17-A19) 166 1 1 1 1 2 1 4 6 6 All origins 1 Male 98 1 1 1 1 1 2 5 5 Female 68 1 1 1 2 1 1 Hispanic 32 1 1 1 1 1 1 2 1 Male 23 1 1 1 1 2 1 9 Female 1 27 2 Mexican 1 1 1 1 1 1 20 2 Male 1 Female 7 1 Puerto Rican 1 Male 1 Female Cuban 1 Male Female 1 Central and South American 1 Male 1 Female Other and unknown Hispanic 2 Male 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other tuberculosis (A17-A19) Non-Hispanic 1/ 131 3 5 1 1 1 4 Male 73 1 1 3 4 2 Female 58 1 1 1 1 Non-Hispanic white 59 2 Male 29 1 Female 30 1 Non-Hispanic black 1 45 Male 32 4 Female 13 Origin not stated 2/ 3 Male 2

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 15 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Whooping cough (A37)											
All origins	7	7	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-
Female	6	6	-	-	-	-	-	-	-	-	-
Hispanic	3	3	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	3	3	-	-	-	-	-	-	-	-	-
Mexican	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	2	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Whooping cough (A37)											
Non-Hispanic 1/	4	4	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-
Female	3	3	-	-	-	-	-	-	-	-	-
Non-Hispanic white	3	3	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-
Non-Hispanic black	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 17 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Scarlet fever and erysipelas (A38,A46) 5 All origins Male Female 5 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 18 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS Scarlet fever and erysipelas (A38,A46) Non-Hispanic 1/ 5 Male Female 5 Non-Hispanic white 4 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Meningococcal infection (A39)											
All origins	227	32	30	13	12	28	11	11	12	12	15
Male	101	14	19	7	6	13	6	8	4	3	6
Female	126	18	11	6	6	15	5	3	8	9	9
Hispanic	27	7	1	1	2	-	2	2	4	3	2
Male	11	5	1	1	1	-	1	1	1	-	-
Female	16	2	-	-	1	-	1	1	3	3	2
Mexican	16	4	1	-	-	-	2	2	3	1	2
Male	6	2	1	-	-	-	1	1	1	-	-
Female	10	2	-	-	-	-	1	1	2	1	2
Puerto Rican	2	1	-	-	-	-	-	-	-	1	-
Male	1	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-
Cuban	4	-	-	-	2	-	-	-	1	-	-
Male	1	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	1	-	-
Central and South American	4	1	-	1	-	-	-	-	-	1	-
Male	2	1	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-
Other and unknown Hispanic	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Meningococcal infection (A39) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Septicemia (A40-A41)											
All origins	30,680	283	87	47	30	32	89	106	169	314	498
Male	13,395	175	48	22	14	17	50	53	94	170	290
Female	17,285	108	39	25	16	15	39	53	75	144	208
Hispanic	1,215	43	14	6	2	5	4	7	19	30	44
Male	605	27	8	3	1	3	2	3	11	20	31
Female	610	16	6	3	1	2	2	4	8	10	13
Mexican	632	34	10	5	1	4	1	3	7	16	29
Male	340	20	6	2	-	2	1	2	5	11	22
Female	292	14	4	3	1	2	-	1	2	5	7
Puerto Rican	212	2	2	1	-	-	1	3	4	7	6
Male	97	2	1	1	-	-	-	1	2	5	4
Female	115	-	1	-	-	-	1	2	2	2	2
Cuban	137	-	-	-	-	-	-	-	1	1	1
Male	67	-	-	-	-	-	-	-	1	1	-
Female	70	-	-	-	-	-	-	-	-	-	1
Central and South American	85	1	-	-	1	1	1	1	2	3	6
Male	33	-	-	-	1	1	1	-	1	3	5
Female	52	1	-	-	-	-	-	1	1	-	1
Other and unknown Hispanic	149	6	2	-	-	-	1	-	5	3	2
Male	68	5	1	-	-	-	-	-	2	-	-
Female	81	1	1	-	_	_	1	_	3	3	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Septicemia (A40-A41) Non-Hispanic 1/ 29,351 12,737 Male Female 16,614 23,023 Non-Hispanic white Male 9,863 Female 13,160 Non-Hispanic black 5,772 Male 2,594 Female 3,178 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 23 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Syphilis (A50-A53)											
All origins	33	2	-	-	-	-	-	-	1	-	1
Male	13	-	-	-	-	-	-	-	1	-	-
Female	20	2	-	-	-	-	-	-	-	-	1
Hispanic	5	2	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	4	2	-	-	-	-	-	-	-	-	-
Mexican	3	2	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Syphilis (A50-A53)											
Non-Hispanic 1/	27	-	-	-	-	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	-	-	1
Non-Hispanic white	14	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	1
Non-Hispanic black	12	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 25 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute poliomyelitis (A80)											
All origins	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	_	-	-	_	_	-	-	-	-	-	_

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 26 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20 - 24 YEARS	25-29 YEARS	30 - 34 YEARS	35-39 YEARS	40-44 YEARS
Acute poliomyelitis (A80)											
Non-Hispanic 1/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 27 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Arthropod-borne viral encephalitis (A83-A84,A85.2) All origins 3 Male 3 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 28 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS Arthropod-borne viral encephalitis (A83-A84,A85.2) Non-Hispanic 1/ 3 Male 3 Female Non-Hispanic white 3 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 29 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Measles (B05)											
All origins	2	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-
Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 30 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Non-Hispanic 1/	2	1	_	_	_	_	_	_	_	_	_
Male	_	-									
		-	-	_	_	-	-	-	-	-	_
Female	2	1	-	-	-	-	-	-	-	-	-
Non-Hispanic white	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5 - 9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Viral hepatitis (B15-B19)											
All origins	4,853	2	3	1	2	3	11	19	69	204	518
Male	3,043	2	2	-	2	3	7	9	44	139	381
Female	1,810	-	1	1	-	-	4	10	25	65	137
Hispanic	656	-	1	-	-	-	3	3	13	38	66
Male	406	-	1	-	-	-	1	-	11	30	54
Female	250	-	-	-	-	-	2	3	2	8	12
Mexican	345	-	1	-	-	-	3	2	10	20	39
Male	214	-	1	-	-	-	1	-	9	16	33
Female	131	-	-	-	-	-	2	2	1	4	6
Puerto Rican	149	-	-	-	-	-	-	1	2	7	15
Male	108	-	-	-	-	-	-	-	2	5	14
Female	41	-	-	-	-	-	-	1	-	2	1
Cuban	50	-	-	-	-	-	-	-	-	1	3
Male	23	-	-	-	-	-	-	-	-	1	3
Female	27	-	-	-	-	-	-	-	-	-	-
Central and South American	45	-	-	-	-	-	-	-	-	2	1
Male	24	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	1	1
Other and unknown Hispanic	67	-	-	-	-	-	-	-	1	8	8
Male	37	-	-	-	-	-	-	-	-	7	4
Female	30	_	-	_	_	_	-	-	1	1	4

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Viral hepatitis (B15-B19)											
Non-Hispanic 1/	4,176	2	2	1	2	3	8	16	55	166	447
Male	2,620	2	1	-	2	3	6	9	32	109	323
Female	1,556	-	1	1	-	-	2	7	23	57	124
Non-Hispanic white	3,176	1	2	1	1	3	3	8	43	131	333
Male	2,026	1	1	-	1	3	3	4	25	88	245
Female	1,150	-	1	1	-	-	-	4	18	43	88
Non-Hispanic black	718	1	-	-	1	-	4	6	11	25	94
Male	433	1	-	-	1	-	2	3	6	15	64
Female	285	-	-	-	-	-	2	3	5	10	30
Origin not stated 2/	21	-	-	-	-	-	-	-	1	-	5
Male	17	-	-	-	-	-	-	-	1	-	4
Female	4	-	-	-	-	-	-	-	-	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Human immunodeficiency virus (HIV) disease (B20-B24) All origins 14.802 1.966 2.981 3.251 Male 11.234 1.410 2.224 2.482 Female 3.568 Hispanic 1,958 Male 1,525 Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Human immunodeficiency virus (HIV) disease (B20-B24) Non-Hispanic 1/ 12,638 11 19 33 23 27 138 640 1,661 2,535 2,770 Male 9,538 8 12 18 15 12 63 412 1,172 1,867 2,117 Female 3,100 3 7 15 8 15 75 228 489 668 653 4,744 2 188 Non-Hispanic white 2 7 8 6 28 606 1,044 1,038 Male 4,043 1 4 6 4 18 139 504 880 888 Female 701 1 3 2 2 10 49 102 164 150 Non-Hispanic black 7,751 9 17 26 15 21 109 446 1,033 1,458 1,696 Male 5,377 6 11 14 9 8 44 268 651 957 1,200 Female 2,374 12 13 178 382 501 496 Origin not stated 2/ 206 8 13 24 48 Male 171 5 11 19 37 Female 35 1 3 2 5 11

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 35 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malaria (B50-B54)											
All origins	7	1	-	-	-	-	-	2	-	-	1
Male	4	1	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	1
Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	_	_	_	_	-	_	_	-	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
118181 18 (830-834)											
Non-Hispanic 1/	7	1	-	-	-	-	-	2	-	-	1
Male	4	1	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	1
Non-Hispanic white	4	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1
Non-Hispanic black	3	1	-	-	-	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

Female

Male

Female

Other and unknown Hispanic

	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS	
			seases and	their	sequelae	(A00,A05,A	\20-A36,A42	2-A44 , A48- <i>i</i>	A49,A54-A79	9,A81-A82,A	A85.0-A8	
l origins	7,279	203	81	40	39	43	47	92	150	234	311	
Male	3,655	119	51	20	24	23	26	46	85	149	184	
Female	3,624	84	30	20	15	20	21	46	65	85	127	
spanic	497	36	13	8	6	5	8	17	22	37	30	
Male	260	22	10	4	3	3	3	10	12	29	20	
Female	237	14	3	4	3	2	5	7	10	8	10	
exican	288	25	7	6	3	4	5	12	15	16	16	
Male	153	16	4	4	1	2	2	6	8	13	12	
Female	135	9	3	2	2	2	3	6	7	3	4	
uerto Rican	62	2	2	-	1	-	-	2	3	6	8	
Male	33	1	2	-	-	-	-	1	1	5	4	
Female	29	1	-	-	1	-	-	1	2	1	4	
uban	36	-	-	-	-	-	-	-	-	3	3	
Male	18	-	-	-	-	-	-	-	-	3	3	
Female	18	-	-	-	-	-	-	-	-	-	-	
entral and South American	60	4	2	1	1	1	3	1	1	11	1	
Male	25	1	2	-	1	1	1	1	-	7	-	
1	CEE FOR NON-HISPANIC IPULATION, AND SEX Ther and unspecified infection of the second	### ACE FOR NON-HISPANIC ALL AGES THE AGES T	ALL UNDER 1 PULATION, AND SEX Cher and unspecified infectious and parasitic di 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) Clorigins 7,279 203 Male 3,655 119 Female 3,624 84 Spanic 497 36 Male 260 22 Female 237 14 Dexican 288 25 Male 153 16 Female 135 9 Puerto Rican 62 24 Male 54 Female 25 Male 16 Female 27 Male 18 Female 28 49 40 40 40 40 40 40 40 40 40	ALL UNDER 1 1-4 PULATION, AND SEX AGES YEAR YEARS There and unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There and unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There and unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There and unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B25-B99) There are an unspecified infectious and parasitic diseases and 1,A85.8,A86-B04,B06-B09,B06-B09,B06-B09,B06-B09,B06-B09,B06-B09,B06-B0	ALL UNDER 1 1-4 5-9 PULATION, AND SEX AGES YEAR YEARS YEARS PULATION, AND SEX AGES YEAR YEARS YEARS YEARS PULATION, AND SEX AGES YEAR YEARS YEARS PULATION, AND SEX AGES AGES	ALL UNDER 1 1-4 5-9 10-14 PPULATION, AND SEX AGES YEARS YEARS YEARS YEARS PRINTED AGES AGES YEARS YEARS YEARS AGES AGES AGES AGES AGES AGES AGES AGE	AGES POR NON-HISPANIC AGES YEAR 1 1-4 5-9 10-14 15-19 POPULATION, AND SEX AGES YEAR YEARS YEARS YEARS YEARS AGES AND THEIR POPULATION, AND SEX AGES AGES AND THEIR POPULATION, AND SEX YEARS AGES AGES AND THEIR POPULATION, AND SEX YEARS AGES AND THEIR POPULATION, AND SEX YEARS AGES AND THEIR POPULATION, AND SEX YEARS AGES AGES AND THEIR POPULATION, AND SEX YEARS AGES AND THEIR POPULATION, AND SEX YEARS AGES AND THEIR POPULATION, AND SEX YEARS AGES AGES AND THEIR POPULATION, AND SEX YEARS AND THEIR POPULATION, AN	ALL WINDER 1 1-4 5-9 10-14 15-19 20-24 PPULATION, AND SEX AGES YEARS YEARS YEARS YEARS PPULATION, AND SEX AGES YEARS YEARS YEARS YEARS PPULATION, AND SEX YEARS YEARS YEARS PPULATION, AND SEX YEARS PRINTED TO THE PROPERTY OF THE PULL TO THE PULL TION, AND SEX YEARS PEARS AGE 1, AGES AGES AGES AGES AGES AGES AGES AGES	AGES OF NON-HISPANIC AGES YEAR YEARS YEARS YEARS YEARS YEARS YEARS PROPURING AND SEX AGES YEARS YEARS YEARS YEARS YEARS PROPURING AND SEX AGES YEARS YEARS YEARS YEARS YEARS PROPURING AND SEX AGES AGES AGES AGES AGES AGES AGES AGES	ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 PPULATION, AND SEX AGES YEAR YEARS PULL TION, AND SEX Ther and unspecified infectious and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06-B09.B25-B49.B55-B99) The and unspecified infectious and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06-B09.B25-B49.B55-B99) The and unspecified infectious and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06-B09.B25-B49.B55-B99) The and unspecified infectious and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06-B09.B25-B49.B55-B99) The and unspecified infectious and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06-B09.B25-B49.B55-B99) The and unspecified infectious and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06-B09.B25-B99) The analysis and parasitic diseases and their sequelae (A00.A05.A20-A36.A42-A44.A48-A49.A54-A751.A85.8.A86-B04.B06.B09.B26-B09.	No.	PURIATION, AND SEX MAGES MAGES

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLA: 199: RACI	TH REVISION INTERNATIONAL SSIFICATION OF DISEASES, 2), HISPANIC ORIGIN, E FOR NON-HISPANIC ULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS	
	er and unspecified infection, A85.8,A86-B04,B06-B09,B25			seases and	their	sequelae	(A00, A05,	A20-A36,A4	2-A44,A48-	A49,A54-A79	9,A81-A82,	A85.0-A8	
Non	-Hispanic 1/	6,753	166	68	32	31	38	39	75	127	197	278	
	Male	3,379	96	41	16	20	20	23	36	72	120	162	
	Female	3,374	70	27	16	11	18	16	39	55	77	116	
Noi	n-Hispanic white	5,288	101	43	24	18	27	24	39	65	121	166	
	Male	2,644	64	27	14	10	13	15	17	38	82	108	
	Female	2,644	37	16	10	8	14	9	22	27	39	58	
Noi	n-Hispanic black	1,247	57	23	7	10	7	13	31	58	73	96	
	Male	617	29	14	2	8	5	7	14	34	36	45	
	Female	630	28	9	5	2	2	6	17	24	37	51	
0ri	gin not stated 2/	29	1	-	-	2	-	-	-	1	-	3	
	Male	16	1	-	-	1	-	-	-	1	-	2	
	Female	13	-	-	-	1	-	-	-	-	-	1	

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms (C00-C97)										
All origins	549,838	68	418	509	503	745	979	1,429	2,576	5,764	10,968
Male	285,832	36	227	279	276	448	575	755	1,204	2,467	5,004
Female	264,006	32	191	230	227	297	404	674	1,372	3,297	5,964
Hispanic	20,233	16	72	87	89	94	165	189	277	471	733
Male	10,668	11	34	45	47	55	103	98	124	198	306
Female	9,565	5	38	42	42	39	62	91	153	273	427
Mexican	10,784	14	47	63	54	59	117	123	175	275	403
Male	5,751	9	23	36	29	28	74	60	85	117	166
Female	5,033	5	24	27	25	31	43	63	90	158	237
Puerto Rican	2,592	1	7	8	7	16	14	18	23	59	89
Male	1,414	1	5	3	3	12	10	8	7	20	37
Female	1,178	-	2	5	4	4	4	10	16	39	52
Cuban	2,362	-	1	2	4	4	3	4	10	27	41
Male	1,296	-	1	2	3	2	1	2	4	13	23
Female	1,066	-	-	-	1	2	2	2	6	14	18
Central and South American	2,117	-	10	10	17	9	21	37	54	74	124
Male	916	-	4	3	10	8	12	23	19	29	48
Female	1,201	-	6	7	7	1	9	14	35	45	76
Other and unknown Hispanic	2,378	1	7	4	7	6	10	7	15	36	76
Male	1,291	1	1	1	2	5	6	5	9	19	32
Female	1,087	-	6	3	5	1	4	2	6	17	44

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasms (C00-C97) Non-Hispanic 1/ 528,218 52 343 422 413 650 812 1,236 2,289 5,281 10,201 Male 274,419 25 190 234 229 393 471 655 1,077 2,266 4,680 Female 253,799 27 153 188 184 257 341 581 1,212 3,015 5,521 281 476 1,729 4,025 Non-Hispanic white 456,315 37 311 297 594 953 7,823 Male 236,384 20 160 180 165 283 346 510 831 1,789 3,636 Female 219,931 17 121 131 132 193 248 443 898 2,236 4,187 Non-Hispanic black 61,409 47 95 136 178 227 1,049 2,000 11 433 Male 32,542 3 24 40 54 84 97 113 181 382 873 Female 28,867 8 23 41 114 252 667 1,127 Origin not stated 2/ 1,387 3 1 1 2 4 10 12 34 Male 745 3 1 2 3 3 18 Female 642 1 1 1 2 7 9 16

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 1 OF 2 PAGE 41

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30 - 34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of lip,	oral cav	ity and pha	rynx (C00)-C14)							
All origins	7,486	-	2	1	5	9	18	14	29	96	195
Male	4,944	-	1	-	3	6	14	10	17	58	146
Female	2,542	-	1	1	2	3	4	4	12	38	49
Hispanic	282	-	-	-	1	-	-	-	1	9	9
Male	204	-	-	-	1	-	-	-	-	4	5
Female	78	-	-	-	-	-	-	-	1	5	4
Mexican	138	-	-	-	-	-	-	-	-	8	2
Male	92	-	-	-	-	-	-	-	-	3	1
Female	46	-	-	-	-	-	-	-	-	5	1
Puerto Rican	60	-	-	-	1	-	-	-	1	-	3
Male	46	-	-	-	1	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	1	-	2
Cuban	29	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-
Central and South American	21	-	-	-	-	-	-	-	-	1	2
Male	16	-	-	-	-	-	-	-	-	1	2
Female	5	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	34	-	-	-	-	-	-	-	-	-	2
Male	28	-	-	-	-	-	-	-	-	-	1
Female	6	-	_	_	_	_	-	-	-	_	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) Non-Hispanic 1/ 7,181 2 4 9 18 14 28 86 186 1 4,725 Male 2 6 14 10 17 53 141 1 Female 2,456 1 1 2 3 4 4 11 33 45 5,826 2 2 5 9 10 19 Non-Hispanic white 1 51 125 Male 3,746 1 3 6 6 11 30 95 Female 2,080 1 1 1 2 8 21 30 Non-Hispanic black 1,131 4 6 25 42 Male 819 3 3 17 30 Female 312 1 2 8 12 Origin not stated 2/ 23 1 Male 15 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

8

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of esophagus (C15) All origins 11,918 10 27 75 218 Male 9,094 7 22 65 190 Female 2,824 3 5 10 28 Hispanic 2 6 10 367 Male 294 2 5 10 Female 73 1 3 5 Mexican 190 1 3 5 Male 150 Female 40 Puerto Rican 70 3 Male 59 3 Female 11 Cuban 37 1 Male 31 1 Female 6 Central and South American 20 1 1 1 Male 16 1 1 1 Female 4 Other and unknown Hispanic 50 1 1

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

38

12

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of esophagus (C15) Non-Hispanic 1/ 11,520 10 25 69 207 8,775 7 Male 20 60 179 9 Female 2,745 3 5 28 9,703 9 21 Non-Hispanic white 61 152 Male 7,440 6 16 55 133 Female 2,263 3 6 19 Non-Hispanic black 1,629 3 6 47 Male 1,187 4 40 Female 442 2 7 Origin not stated 2/ 31 1 Male 25 1 Female 6

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 30-34 40-44 ALL 5-9 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of stomach (C16) All origins 12,711 Male 7.459 Female 5.252 Hispanic 1,055 Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YEAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of stomach (C16) Non-Hispanic 1/ 11,615 4 5 33 65 155 265 1 Male 6,827 1 2 3 14 37 87 155 4,788 2 Female 2 19 28 68 110 8,712 3 15 35 Non-Hispanic white 1 95 142 Male 5,171 1 7 23 57 85 Female 3,541 8 12 38 57 Non-Hispanic black 2,183 1 14 22 40 87 Male 1,252 7 12 20 52 Female 931 1 7 10 20 35 Origin not stated 2/ 41 Male 25 Female 16

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of colo	on, rectum	and anus (C18-C21)								
All origins	57,155	1	-	-	1	11	38	87	196	439	894
Male	28,240	-	-	-	1	5	22	57	103	226	452
Female	28,915	1	-	-	-	6	16	30	93	213	442
Hispanic	2,031	-	-	-	-	1	6	7	24	33	53
Male	1,062	-	-	-	-	1	4	3	11	15	24
Female	969	-	-	-	-	-	2	4	13	18	29
Mexican	982	-	-	-	-	-	4	5	13	16	27
Male	559	-	-	-	-	-	3	2	7	9	17
Female	423	-	-	-	-	-	1	3	6	7	10
Puerto Rican	300	-	-	-	-	-	2	-	3	8	7
Male	145	-	-	-	-	-	1	-	2	3	2
Female	155	-	-	-	-	-	1	-	1	5	5
Cuban	299	-	-	-	-	-	-	-	2	2	5
Male	148	-	-	-	-	-	-	-	-	1	1
Female	151	-	-	-	-	-	-	-	2	1	4
Central and South American	181	-	-	-	-	-	-	1	4	3	11
Male	70	-	-	-	-	-	-	1	1	-	3
Female	111	-	-	-	-	-	-	-	3	3	8
Other and unknown Hispanic	269	-	-	-	-	1	-	1	2	4	3
Male	140	-	-	-	-	1	-	-	1	2	1
Female	129	-	-	-	-	-	-	1	1	2	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasms of colon, rectum and anus (C18-C21) Non-Hispanic 1/ 54,976 1 1 10 31 80 172 404 837 Male 27,103 4 18 54 92 211 425 1 Female 27,873 1 6 13 26 80 193 412 47,275 303 Non-Hispanic white 1 6 20 56 126 630 Male 23,446 1 2 10 38 67 166 334 Female 23,829 4 10 18 59 137 296 Non-Hispanic black 6,709 10 20 35 90 178 Male 3,160 2 7 13 17 42 81 Female 3,549 2 18 48 97

1

1

2

2

4

3

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

148

75

73

Origin not stated 2/

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLA 199 RAC	TH REVISION INTERNATIONAL SSIFICATION OF DISEASES, 2), HISPANIC ORIGIN, E FOR NON-HISPANIC ULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
М	alignant neoplasms of liver	r and int	rahepatic b	ile ducts	(C22)							
All	origins	12,382	8	29	5	6	19	24	33	45	127	317
	Male	7,689	5	20	5	4	12	12	22	26	88	230
	Female	4,693	3	9	-	2	7	12	11	19	39	87
His	panic	1,084	1	4	1	1	2	2	3	4	13	30
	Male	688	1	3	1	1	1	2	3	2	6	21
	Female	396	-	1	-	-	1	-	-	2	7	9
Me	xican	654	1	3	1	1	2	2	2	3	5	15
	Male	407	1	3	1	1	1	2	2	2	2	11
	Female	247	-	-	-	-	1	-	-	1	3	4
Pu	erto Rican	144	-	-	-	-	-	-	-	-	2	1
	Male	115	-	-	-	-	-	-	-	-	2	-
	Female	29	-	-	-	-	-	-	-	-	-	1
Cu	ban	85	-	-	-	-	-	-	-	-	2	2
	Male	44	-	-	-	-	-	-	-	-	-	1
	Female	41	-	-	-	-	-	-	-	-	2	1
Ce	ntral and South American	110	-	-	-	-	-	-	1	1	4	6
	Male	56	-	-	-	-	-	-	1	-	2	4
	Female	54	-	-	-	-	-	-	-	1	2	2
0t	her and unknown Hispanic	91	-	1	-	-	-	-	-	-	-	6
	Male	66	-	-	-	-	-	-	-	-	-	5
	Female	25	-	1	-	-	-	-	-	-	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasms of liver and intrahepatic bile ducts (C22) Non-Hispanic 1/ 11,265 7 23 4 5 17 22 30 41 114 284 6,976 Male 4 15 4 3 11 10 19 24 82 208 Female 4,289 3 8 6 12 11 17 32 76 8,890 5 12 Non-Hispanic white 20 2 2 13 14 15 60 178 Male 5,455 13 2 9 8 8 39 134 Female 3,435 3 7 3 6 7 21 44 Non-Hispanic black 1,498 3 3 2 9 12 31 60 Male 939 2 2 7 21 44 Female 559 2 3 10 16 Origin not stated 2/ 33 2 3

1

2

2

Source: CDC/NCHS, National Vital Statistics System, Mortality

25

8

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of pancreas (C25) All origins 29,082 1 3 9 46 178 416 Male 14.176 1 7 26 107 262 Female 14,906 1 2 2 20 71 154 Hispanic 1 5 18 26 1,135 Male 563 2 12 18 8 Female 572 1 3 6 13 Mexican 656 1 20 347 9 13 Male Female 309 1 2 4 7 Puerto Rican 110 1 Male 51 Female 59 1 Cuban 125 2 2 Male 56 1 2 Female 69 1 Central and South American 119 1 1 2 Male 53 1 1 Female 66 1 1 Other and unknown Hispanic 125 1 2 Male 1 2 56

Source: CDC/NCHS, National Vital Statistics System, Mortality

69

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of pancreas (C25) Non-Hispanic 1/ 27,869 1 3 8 41 160 387 Male 13,572 1 7 24 95 243 2 Female 14,297 1 1 17 65 144 23,903 2 28 Non-Hispanic white 6 119 302 Male 11,684 6 17 73 193 Female 12,219 11 46 109 Non-Hispanic black 3,368 10 76 36 Male 1,601 19 44 Female 1,767 17 32 Origin not stated 2/ 78 3 Male 41 1 Female 37 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of larynx (C32) All origins 3,815 20 6 58 Male 2,992 5 18 46 Female 823 1 2 12 Hispanic 171 5 4 Male 149 Female 22 1 2 Mexican 81 72 2 Male Female 9 Puerto Rican 39 2 Male 33 2 Female 6 Cuban 27 Male 24 3 Female Central and South American 8 Male 6 Female 2 Other and unknown Hispanic 16 1 Male 14 Female 2 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YEAR YEARS YEARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS Malignant neoplasm of larynx (C32) Non-Hispanic 1/ 3,627 6 20 53 Male 2,832 5 18 42 795 Female 1 2 11 2,924 Non-Hispanic white 4 14 26 Male 2,263 3 14 20 Female 661 1 6 Non-Hispanic black 657 2 6 26 Male 532 2 4 21 Female 125 5 Origin not stated 2/ 17 Male 11 Female 6

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of trac	chea, bron	chus and lu	ng (C33-C	34)							
All origins	152,156	-	-	3	2	6	14	43	136	668	2,056
Male	89,453	-	-	2	2	5	8	24	73	323	1,151
Female	62,703	-	-	1	-	1	6	19	63	345	905
Hispanic	3,545	-	-	-	-	-	3	3	14	32	66
Male	2,373	-	-	-	-	-	2	3	7	17	41
Female	1,172	-	-	-	-	-	1	-	7	15	25
Mexican	1,757	-	-	-	-	-	3	2	8	14	34
Male	1,173	-	-	-	-	-	2	2	5	6	16
Female	584	-	-	-	-	-	1	-	3	8	18
Puerto Rican	506	-	-	-	-	-	-	-	-	4	8
Male	338	-	-	-	-	-	-	-	-	3	6
Female	168	-	-	-	-	-	-	-	-	1	2
Cuban	525	-	-	-	-	-	-	-	-	2	12
Male	383	-	-	-	-	-	-	-	-	1	10
Female	142	-	-	-	-	-	-	-	-	1	2
Central and South American	291	-	-	-	-	-	-	-	5	8	7
Male	176	-	-	-	-	-	-	-	1	4	6
Female	115	-	-	-	-	-	-	-	4	4	1
Other and unknown Hispanic	466	-	-	-	-	-	-	1	1	4	5
Male	303	-	-	-	-	-	-	1	1	3	3
Female	163	-	-	-	-	-	-	-	-	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	, ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of tr	rachea, bron	chus and lu	ing (C33-C	34)							
Non-Hispanic 1/	148,206	-	-	3	2	6	11	40	121	635	1,983
Male	86,822	-	-	2	2	5	6	21	66	306	1,105
Female	61,384	-	-	1	-	1	5	19	55	329	878
Non-Hispanic white	129,967	-	-	2	1	3	6	30	101	479	1,546
Male	75,338	-	-	1	1	3	3	14	57	231	847
Female	54,629	-	-	1	-	-	3	16	44	248	699
Non-Hispanic black	15,834	-	-	1	1	3	5	5	15	134	397
Male	10,039	-	-	1	1	2	3	4	7	63	238
Female	5,795	-	-	-	-	1	2	1	8	71	159
Origin not stated 2/	405	-	-	-	-	-	-	-	1	1	7
Male	258	-	-	-	-	-	-	-	-	-	5
Female	147	-	-	-	-	-	-	-	1	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant melanoma of skin	(C43)										
All origins	7,215	1	-	1	-	9	30	62	132	239	386
Male	4,529	1	-	-	-	7	16	38	72	151	246
Female	2,686	-	-	1	-	2	14	24	60	88	140
Hispanic	133	1	-	1	-	1	1	1	4	6	8
Male	79	1	-	-	-	1	-	1	3	4	5
Female	54	-	-	1	-	-	1	-	1	2	3
Mexican	71	-	-	1	-	1	-	1	3	6	6
Male	45	-	-	-	-	1	-	1	2	4	4
Female	26	-	-	1	-	-	-	-	1	2	2
Puerto Rican	10	1	-	-	-	-	-	-	-	-	1
Male	6	1	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1
Cuban	18	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-
Central and South American	16	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	18	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	1	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant melanoma of skin (C43) Non-Hispanic 1/ 7,062 8 29 61 128 233 378 37 Male 4,438 6 16 69 147 241 2 Female 2,624 13 24 59 86 137 6,936 7 27 60 230 374 Non-Hispanic white 123 Male 4,370 5 14 36 66 145 240 Female 2,566 2 13 24 57 85 134 Non-Hispanic black 1 3 3 94 1 Male 47 1 1 Female 47 2 3 Origin not stated 2/ 20 Male 12 Female 8

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of breas	t (C50)										
All origins	41,528	-	1	-	1	3	10	86	353	986	1,755
Male	384	-	-	-	-	-	-	-	3	6	9
Female	41,144	-	1	-	1	3	10	86	350	980	1,746
Hispanic	1,551	-	-	-	-	1	1	13	34	74	123
Male	13	-	-	-	-	-	-	-	-	-	-
Female	1,538	-	-	-	-	1	1	13	34	74	123
Mexican	796	-	-	-	-	1	1	10	22	45	73
Male	6	-	-	-	-	-	-	-	-	-	-
Female	790	-	-	-	-	1	1	10	22	45	73
Puerto Rican	197	-	-	-	-	-	-	2	2	8	12
Male	3	-	-	-	-	-	-	-	-	-	-
Female	194	-	-	-	-	-	-	2	2	8	12
Cuban	171	-	-	-	-	-	-	-	1	2	4
Male	-	-	-	-	-	-	-	-	-	-	-
Female	171	-	-	-	-	-	-	-	1	2	4
Central and South American	198	-	-	-	-	-	-	1	8	10	21
Male	1	-	-	-	-	-	-	-	-	-	-
Female	197	-	-	-	-	-	-	1	8	10	21
Other and unknown Hispanic	189	-	-	-	-	-	-	-	1	9	13
Male	3	-	-	-	-	-	-	-	-	-	-
Female	186	-	-	-	-	-	-	-	1	9	13

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YEAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of breast (C50) Non-Hispanic 1/ 39,881 1 2 9 72 318 911 1,626 1 370 9 Male 3 6 2 905 Female 39,511 1 9 72 315 1,617 33,790 2 Non-Hispanic white 1 4 46 211 627 1,175 Male 299 2 8 Female 33,491 1 2 4 46 209 627 1,167 Non-Hispanic black 5,386 100 259 402 24 Male 67 6 1 Female 5,319 24 99 253 401 Origin not stated 2/ 1 1 1 6 Male 1 Female 95 1 1 1 6

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of cervix uteri (C53) All origins 4,205 2 10 62 155 292 407 Female 4,205 2 10 62 155 292 407 Hispanic 378 2 7 24 34 53 Female 378 2 7 24 34 53 4 Mexican 215 1 16 20 29 29 Female 215 1 16 20 Puerto Rican 6 5 53 1 53 6 5 Female Cuban 16 1 1 1 Female 16 1 1 1 Central and South American 67 1 1 4 7 12 7 Female 67 1 1 4 12 Other and unknown Hispanic 27 6

6

Source: CDC/NCHS, National Vital Statistics System, Mortality

27

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of cervix uteri (C53) Non-Hispanic 1/ 3,817 2 55 131 258 353 2 Female 3,817 55 131 258 353 47 Non-Hispanic white 2,854 100 188 263 Female 2,854 47 188 100 263 Non-Hispanic black 825 2 27 60 80 Female 825 2 27 60 80 Origin not stated 2/ 10 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of corp	ous uteri	and uterus,	part uns	pecified	(C54-C55)						
All origins	6,468	-	-	-	-	-	2	11	12	38	104
Female	6,468	-	-	-	-	-	2	11	12	38	104
Hispanic	272	-	-	-	-	-	1	2	1	4	11
Female	272	-	-	-	-	-	1	2	1	4	11
Mexican	137	-	-	-	-	-	1	1	-	3	6
Female	137	-	-	-	-	-	1	1	-	3	6
Puerto Rican	37	-	-	-	-	-	-	1	1	1	1
Female	37	-	-	-	-	-	-	1	1	1	1
Cuban	46	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-
Central and South American	26	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	2
Other and unknown Hispanic	26	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	2

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 64
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YEARS YFARS Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55) Non-Hispanic 1/ 6,188 11 34 93 Female 6,188 9 11 34 93 9 Non-Hispanic white 5,114 6 28 68 Female 5,114 9 28 6 68 Non-Hispanic black 980 5 22 Female 980 5 22

Source: CDC/NCHS, National Vital Statistics System, Mortality

8

Origin not stated 2/
Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of ovary (C56) All origins 13,627 1 2 9 18 26 51 162 334 Female 13,627 1 2 9 18 26 51 162 334 Hispanic 478 1 1 1 5 1 11 18 Female 478 1 1 1 5 1 11 18 7 7 Mexican 269 1 1 1 5 7 7 Female 269 1 1 1 Puerto Rican 2 4 54 54 2 4 Female Cuban 56 1 1 Female 56 1 1 Central and South American 51 1 1 4 Female 51 1 1 4 Other and unknown Hispanic 48 2 2

Source: CDC/NCHS, National Vital Statistics System, Mortality

48

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YEARS YFARS Malignant neoplasm of ovary (C56) Non-Hispanic 1/ 13,103 2 8 17 21 50 150 316 Female 13,103 2 8 17 21 50 150 316 11,795 7 Non-Hispanic white 13 15 36 124 270 Female 11,795 2 7 15 124 270 13 36 Non-Hispanic black 1,065 1 3 6 16 31 Female 1,065 1 6 16 31 Origin not stated 2/ 1 46 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YEAR YEARS YEARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS Malignant neoplasm of prostate (C61) All origins 31,729 1 1 1 5 28 1 Male 31,729 1 1 1 1 5 28 Hispanic 1,050 1 1 2 Male 1,050 1 2 Mexican 510 1 Male 510 Puerto Rican 142 Male 142 Cuban 157 Male 157 Central and South American 90 1 Male 90 1 Other and unknown Hispanic 151 1 1

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

151

Male

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 68 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YEARS YEARS Malignant neoplasm of prostate (C61) Non-Hispanic 1/ 30,614 1 5 26 1 30,614 Male 1 1 5 26 24,936 Non-Hispanic white 1 4 19 24,936 19 Male 1 1 4 Non-Hispanic black 5,315 1 6 5,315 1 6 Origin not stated 2/ 65 Male

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30 - 34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of kidm	ney and re	nal pelvis	(C64-C65)								
All origins	11,116	2	19	23	10	8	9	25	29	105	237
Male	6,867	1	11	12	4	4	7	15	22	70	166
Female	4,249	1	8	11	6	4	2	10	7	35	71
Hispanic	547	-	1	4	-	1	-	1	3	9	22
Male	353	-	1	2	-	-	-	-	2	7	16
Female	194	-	-	2	-	1	-	1	1	2	6
Mexican	346	-	-	2	-	1	-	1	3	7	14
Male	223	-	-	1	-	-	-	-	2	6	11
Female	123	-	-	1	-	1	-	1	1	1	3
Puerto Rican	50	-	1	1	-	-	-	-	-	1	2
Male	34	-	1	1	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	1	-
Cuban	43	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1
Central and South American	33	-	-	1	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	1	-	-	-	-	-	-	-
Other and unknown Hispanic	75	-	-	-	-	-	-	-	-	-	5
Male	49	-	-	-	-	-	-	-	-	-	3
Female	26	-	-	-	-	-	-	-	-	-	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasms of kidney and renal pelvis (C64-C65) Non-Hispanic 1/ 10,542 2 18 19 10 7 9 23 26 94 215 6,501 7 Male 1 10 10 4 4 15 20 62 150 4,041 3 Female 1 8 9 2 8 6 32 65 9,319 2 13 15 Non-Hispanic white 11 15 6 4 65 173 Male 5,754 7 8 4 7 11 43 122 Female 3,565 1 4 6 4 22 51 Non-Hispanic black 1,028 3 4 10 10 27 34 Male 618 2 4 8 19 23 Female 410 3 2 8 11 Origin not stated 2/ 27 2

1

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

13

14

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 35-39 40-44 ALL 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of bladder (C67) All origins 11,911 6 10 32 1 1 68 Male 8,105 1 3 6 20 43 Female 3,806 1 3 4 12 25 Hispanic 301 2 1 3 Male 222 2 1 3 79 Female 2 Mexican 150 1 104 Male 1 Female 46 Puerto Rican 31 1 Male 22 1 Female 9 Cuban 46 Male 38 Female 8 Central and South American 31 1 Male 24 1 Female 7 Other and unknown Hispanic 43 1 Male 1 34 9

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YEAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Malignant neoplasm of bladder (C67) Non-Hispanic 1/ 11,582 6 8 31 65 1 1 Male 7,865 1 3 4 19 40 Female 3,717 1 3 12 25 5 Non-Hispanic white 10,548 1 3 23 54 Male 7,290 1 13 33 Female 3,258 3 10 21 Non-Hispanic black 906 8 9 Male 487 6 6 Female 419 Origin not stated 2/ 28 Male 18 Female 10

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of men	inges, bra	in and othe	r parts o	of central	nervous	system (C	C70-C72)				
All origins	12,765	10	113	184	124	115	103	177	267	443	622
Male	7,001	8	56	96	58	69	62	104	172	268	370
Female	5,764	2	57	88	66	46	41	73	95	175	252
Hispanic	568	4	27	30	18	2	16	20	11	31	34
Male	293	4	11	13	9	2	8	12	5	20	25
Female	275	-	16	17	9	-	8	8	6	11	9
Mexican	329	4	17	20	10	1	12	14	8	19	19
Male	173	4	8	10	6	1	6	7	3	12	12
Female	156	-	9	10	4	-	6	7	5	7	7
Puerto Rican	49	-	4	4	2	-	1	2	1	3	3
Male	26	-	2	1	-	-	1	2	-	3	3
Female	23	-	2	3	2	-	-	-	1	-	-
Cuban	49	-	-	1	-	-	-	2	-	-	3
Male	28	-	-	1	-	-	-	1	-	-	3
Female	21	-	-	-	-	-	-	1	-	-	-
Central and South American	83	-	4	2	6	1	2	2	1	8	5
Male	36	-	1	1	3	1	1	2	1	4	5
Female	47	-	3	1	3	-	1	-	-	4	-
Other and unknown Hispanic	58	-	2	3	-	-	1	-	1	1	4
Male	30	-	-	-	-	-	-	-	1	1	2
Female	28	-	2	3	-	-	1	-	-	-	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of men	inges, brai	n and other	parts o	f central	nervous	system (C	70-C72)				
Non-Hispanic 1/	12,173	6	86	154	106	112	87	157	256	412	588
Male	6,696	4	45	83	49	67	54	92	167	248	345
Female	5,477	2	41	71	57	45	33	65	89	164	243
Non-Hispanic white	11,238	3	70	110	85	83	69	145	227	377	525
Male	6,193	2	39	63	41	48	43	85	148	229	304
Female	5,045	1	31	47	44	35	26	60	79	148	221
Non-Hispanic black	754	3	11	36	18	23	14	10	26	26	50
Male	399	2	4	17	7	15	9	6	17	15	32
Female	355	1	7	19	11	8	5	4	9	11	18
Origin not stated 2/	24	-	-	-	-	1	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	1	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of lymp	phoid, hem	atopoietic	and relat	ed tissue	(C81-C96	5)					
All origins	56,318	30	137	142	209	303	417	423	537	837	1,076
Male	29,721	12	75	84	122	180	257	241	316	497	647
Female	26,597	18	62	58	87	123	160	182	221	340	429
Hispanic	2,538	8	28	28	47	45	83	63	72	88	102
Male	1,319	3	16	19	23	23	50	34	41	57	51
Female	1,219	5	12	9	24	22	33	29	31	31	51
Mexican	1,381	7	19	22	30	28	62	37	43	48	53
Male	737	2	11	17	13	11	37	21	26	34	28
Female	644	5	8	5	17	17	25	16	17	14	25
Puerto Rican	325	-	1	2	4	8	7	6	8	13	17
Male	172	-	1	1	2	5	5	2	4	5	7
Female	153	-	-	1	2	3	2	4	4	8	10
Cuban	299	-	1	-	1	3	1	1	7	6	7
Male	153	-	1	-	1	1	-	1	4	4	4
Female	146	-	-	-	-	2	1	-	3	2	3
Central and South American	291	-	6	4	8	4	8	19	10	16	20
Male	128	-	3	1	5	4	4	10	6	9	8
Female	163	-	3	3	3	-	4	9	4	7	12
Other and unknown Hispanic	242	1	1	-	4	2	5	-	4	5	5
Male	129	1	-	-	2	2	4	-	1	5	4
Female	113	-	1	-	2	-	1	-	3	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of	lymphoid, hema	atopoietic	and relat	ed tissue	(C81-C96	5)					
Non-Hispanic 1/	53,669	22	108	114	162	258	334	360	463	749	970
Male	28,347	9	58	65	99	157	207	207	274	440	592
Female	25,322	13	50	49	63	101	127	153	189	309	378
Non-Hispanic white	47,608	16	91	86	113	188	249	271	347	590	762
Male	25,219	9	48	51	69	114	163	153	204	351	465
Female	22,389	7	43	35	44	74	86	118	143	239	297
Non-Hispanic black	5,038	4	14	21	40	53	80	71	86	134	173
Male	2,566	-	8	11	27	32	41	43	54	74	106
Female	2,472	4	6	10	13	21	39	28	32	60	67
Origin not stated 2/	111	-	1	-	-	-	-	-	2	-	4
Male	55	-	1	-	-	-	-	-	1	-	4
Female	56	-	-	-	-	-	-	-	1	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hodgkin's disease (C81)											
All origins	1,403	-	1	-	9	31	63	77	71	90	75
Male	785	-	1	-	3	20	35	42	44	56	47
Female	618	-	-	-	6	11	28	35	27	34	28
Hispanic	113	-	-	-	3	2	6	2	5	5	7
Male	65	-	-	-	2	2	4	1	2	4	4
Female	48	-	-	-	1	-	2	1	3	1	3
Mexican	62	-	-	-	2	1	4	1	1	2	4
Male	36	-	-	-	1	1	3	1	1	2	2
Female	26	-	-	-	1	-	1	-	-	-	2
Puerto Rican	16	-	-	-	-	-	-	1	2	1	2
Male	7	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	1	2	-	1
Cuban	12	-	-	-	1	-	-	-	1	-	1
Male	10	-	-	-	1	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-
Central and South American	13	-	-	-	-	-	2	-	-	1	-
Male	9	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	1	-
Other and unknown Hispanic	10	-	-	-	-	1	-	-	1	1	-
Male	3	-	-	-	-	1	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	1	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Hodgkin's disease (C81) Non-Hispanic 1/ 1,286 6 29 57 75 66 85 68 1 719 Male 1 1 18 31 41 42 52 43 Female 567 5 11 26 34 24 33 25 58 75 Non-Hispanic white 1,149 1 5 24 43 58 59 Male 643 1 1 15 24 31 38 45 39 Female 506 4 9 19 27 20 30 20 Non-Hispanic black 5 14 17 8 10 8 125 Male 3 7 10 7 4 Female 51 2 4 3 Origin not stated 2/ 4 Male 1 Female 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 40-44 ALL 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Non-Hodgkin's lymphoma (C82-C85) All origins 22,802 Male 11,794 Female 11,008 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Non-Hodgkin's lymphoma (C82-C85) Non-Hispanic 1/ 21,792 Male 11,271 Female 10,521 19,956 Non-Hispanic white Male 10,259 Female 9,697 Non-Hispanic black 1,427 Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Leukemia (C91-C95)											
All origins	21,014	27	119	130	175	221	261	234	266	374	425
Male	11,519	11	65	75	98	130	161	128	148	204	231
Female	9,495	16	54	55	77	91	100	106	118	170	194
Hispanic	994	8	26	27	42	38	63	49	37	52	46
Male	534	3	16	18	20	17	40	26	23	28	20
Female	460	5	10	9	22	21	23	23	14	24	26
Mexican	542	7	18	21	26	26	48	30	25	26	26
Male	290	2	11	16	11	9	29	15	15	16	11
Female	252	5	7	5	15	17	19	15	10	10	15
Puerto Rican	125	-	1	2	4	6	7	4	2	9	9
Male	72	-	1	1	2	3	5	2	2	2	4
Female	53	-	-	1	2	3	2	2	-	7	5
Cuban	123	-	1	-	-	2	-	1	2	5	1
Male	65	-	1	-	-	1	-	1	1	3	-
Female	58	-	-	-	-	1	-	-	1	2	1
Central and South American	124	-	5	4	8	3	4	14	5	9	8
Male	59	-	3	1	5	3	2	8	4	4	3
Female	65	-	2	3	3	-	2	6	1	5	5
Other and unknown Hispanic	80	1	1	-	4	1	4	-	3	3	2
Male	48	1	-	-	2	1	4	-	1	3	2
Female	32	-	1	-	2	-	-	-	2	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Leukemia (C91-C95) Non-Hispanic 1/ 19,981 10,968 Male Female 9,013 Non-Hispanic white 17,885 Male 9,853 Female 8,032 Non-Hispanic black 1,694 Male Female Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 83 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Multiple myeloma and imm	unoprolife	rative neop	olasms (C8	8,C90)							
All origins	11,016	-	-	-	-	-	-	3	19	47	109
Male	5,584	-	-	-	-	-	-	2	14	33	62
Female	5,432	-	-	-	-	-	-	1	5	14	47
Hispanic	468	-	-	-	-	-	-	2	2	6	6
Male	225	-	-	-	-	-	-	2	2	6	3
Female	243	-	-	-	-	-	-	-	-	-	3
Mexican	249	-	-	-	-	-	-	2	1	4	3
Male	121	-	-	-	-	-	-	2	1	4	1
Female	128	-	-	-	-	-	-	-	-	-	2
Puerto Rican	61	-	-	-	-	-	-	-	-	1	-
Male	30	-	-	-	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	-	-	-	-
Cuban	58	-	-	-	-	-	-	-	1	-	1
Male	29	-	-	-	-	-	-	-	1	-	1
Female	29	-	-	-	-	-	-	-	-	-	-
Central and South American	51	-	-	-	-	-	-	-	-	1	2
Male	20	-	-	-	-	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	-	-	1
Other and unknown Hispanic	49	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES,

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Multiple myeloma and im	nmunoprolife	rative neop	lasms (C8	8,C90)							
Non-Hispanic 1/	10,532	-	-	-	-	-	-	1	17	41	103
Male	5,352	-	-	-	-	-	-	-	12	27	59
Female	5,180	-	-	-	-	-	-	1	5	14	44
Non-Hispanic white	8,547	-	-	-	-	-	-	-	10	29	63
Male	4,430	-	-	-	-	-	-	-	6	20	36
Female	4,117	-	-	-	-	-	-	-	4	9	27
Non-Hispanic black	1,786	-	-	-	-	-	-	1	6	10	39
Male	824	-	-	-	-	-	-	-	6	6	23
Female	962	-	-	-	-	-	-	1	-	4	16
Origin not stated 2/	16	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 85 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other and unspecified m	alignant ne	oplasms of	lymphoid,	hematopo	ietic and	related	tissue (C	96)			
All origins	83	1	1	-	-	-	-	-	1	2	2
Male	39	1	-	-	-	-	-	-	1	-	1
Female	44	-	1	-	-	-	-	-	-	2	1
Hispanic	5	-	1	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	1	-	-
Female	3	-	1	-	-	-	-	-	-	-	-
Mexican	3	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-
Puerto Rican	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	1	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 86 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other and unspecified mal	lignant nec	plasms of	lymphoid,	hematopo	ietic and	related	tissue (C	96)			
Non-Hispanic 1/	78	1	-	-	-	-	-	-	-	2	2
Male	37	1	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	2	1
Non-Hispanic white	71	1	-	-	-	-	-	-	-	2	1
Male	34	1	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	2	-
Non-Hispanic black	6	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other and unspecified m	nalignant	neoplasms	(C17,C23-C	C24,C26-C3	1,C37-C4	1,C44-C49,	,C51-C52,C	57-C60,C62	-C63,C66,C	68-C69,C73	-C80,C97
All origins	66,251	14	116	150	141	246	273	307	455	833	1,481
Male	33,449	8	64	80	80	158	171	204	286	464	836
Female	32,802	6	52	70	61	88	102	103	169	369	645
Hispanic	2,747	2	11	23	22	39	45	48	51	68	107
Male	1,399	2	3	10	13	27	35	33	30	36	54
Female	1,348	-	8	13	9	12	10	15	21	32	53
Mexican	1,543	2	7	17	13	23	28	32	32	47	64
Male	798	2	1	7	9	14	24	20	19	23	32
Female	745	-	6	10	4	9	4	12	13	24	32
Puerto Rican	283	-	1	1	-	8	4	4	2	5	14
Male	152	-	1	-	-	7	3	3	1	2	7
Female	131	-	-	1	-	1	1	1	1	3	7
Cuban	262	-	-	1	3	1	1	-	-	4	1
Male	141	-	-	1	2	1	-	-	-	3	1
Female	121	-	-	-	1	-	1	-	-	1	-
Central and South American	316	-	-	3	3	4	10	10	11	6	15
Male	128	-	-	1	2	3	7	8	5	3	7
Female	188	-	-	2	1	1	3	2	6	3	8
Other and unknown Hispanic	343	-	3	1	3	3	2	2	6	6	13
Male	180	-	1	1	-	2	1	2	5	5	7
Female	163	-	2	-	3	1	1	-	1	1	6

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20 - 24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other and unspecified ma	lignant	neoplasms	(C17,C23-	C24,C26-C3	1,C37-C41	, C44-C49	,C51-C52,C	57-C60,C62	-C63,C66,C0	68-C69,C73	-C80,C97
Non-Hispanic 1/	63,328	12	105	127	118	207	227	257	398	761	1,369
Male	31,956	6	61	70	67	131	135	169	254	427	779
Female	31,372	6	44	57	51	76	92	88	144	334	590
Non-Hispanic white	54,977	10	87	95	83	158	166	207	306	587	1,039
Male	27,780	5	52	55	49	94	96	143	195	339	604
Female	27,197	5	35	40	34	64	70	64	111	248	435
Non-Hispanic black	7,009	1	14	22	27	39	44	41	65	144	277
Male	3,514	-	8	9	13	29	27	20	40	70	149
Female	3,495	1	6	13	14	10	17	21	25	74	128
Origin not stated 2/	176	-	-	-	1	-	1	2	6	4	5
Male	94	-	-	-	-	-	1	2	2	1	3
Female	82	-	-	-	1	-	-	-	4	3	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
In situ neoplasms, benign neo	oplasms and	d neoplasms	of uncer	tain or u	ınknown be	havior (D	00-D48)				
All origins	13,227	58	63	64	37	52	46	55	116	140	202
Male	6,520	30	33	30	19	26	28	32	64	79	118
Female	6,707	28	30	34	18	26	18	23	52	61	84
Hispanic	497	8	12	11	6	7	5	8	17	6	24
Male	233	5	4	5	3	5	3	7	7	4	17
Female	264	3	8	6	3	2	2	1	10	2	7
Mexican	265	6	7	9	5	4	2	5	9	3	15
Male	132	4	3	5	2	4	1	5	5	2	12
Female	133	2	4	4	3	-	1	-	4	1	3
Puerto Rican	60	1	3	-	-	1	1	2	-	1	3
Male	32	-	-	-	-	1	-	1	-	1	1
Female	28	1	3	-	-	-	1	1	-	-	2
Cuban	58	-	2	-	-	-	-	1	-	1	1
Male	26	-	1	-	-	-	-	1	-	1	1
Female	32	-	1	-	-	-	-	-	-	-	-
Central and South American	57	1	-	-	1	1	1	-	6	1	2
Male	23	1	-	-	1	-	1	-	2	-	1
Female	34	-	-	-	-	1	-	-	4	1	1
Other and unknown Hispanic	57	-	-	2	-	1	1	-	2	-	3
Male	20	-	-	-	-	-	1	-	-	-	2
Female	37	-	-	2	-	1	-	-	2	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) Non-Hispanic 1/ 12,699 6,276 Male 6,423 Female Non-Hispanic white 11,335 Male 5,653 Female 5,682 Non-Hispanic black 1,161 Male Female Origin not stated 2/

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Anemias (D50-D64)											
All origins	4,503	12	30	24	29	43	78	64	96	90	104
Male	1,746	8	17	8	15	24	43	33	48	47	54
Female	2,757	4	13	16	14	19	35	31	48	43	50
Hispanic	164	-	6	2	1	2	4	4	4	5	5
Male	79	-	4	1	-	1	2	2	3	3	3
Female	85	-	2	1	1	1	2	2	1	2	2
Mexican	88	-	5	-	1	-	2	-	2	2	3
Male	39	-	3	-	-	-	1	-	1	1	2
Female	49	-	2	-	1	-	1	-	1	1	1
Puerto Rican	22	-	1	2	-	-	-	1	-	1	-
Male	11	-	1	1	-	-	-	-	-	1	-
Female	11	-	-	1	-	-	-	1	-	-	-
Cuban	17	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-
Central and South American	22	-	-	-	-	2	2	3	2	2	-
Male	12	-	-	-	-	1	1	2	2	1	-
Female	10	-	-	-	-	1	1	1	-	1	-
Other and unknown Hispanic	15	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-
Female	9	-	_	_	_	_	-	_	_	_	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Anemias (D50-D64)											
Non-Hispanic 1/	4,321	12	24	22	28	41	74	60	92	85	98
Male	1,659	8	13	7	15	23	41	31	45	44	51
Female	2,662	4	11	15	13	18	33	29	47	41	47
Non-Hispanic white	3,222	6	4	5	12	16	12	3	7	17	20
Male	1,162	4	2	-	6	8	3	-	2	7	11
Female	2,060	2	2	5	6	8	9	3	5	10	9
Non-Hispanic black	1,010	3	20	17	16	23	61	55	84	66	76
Male	453	2	11	7	9	13	37	29	43	35	38
Female	557	1	9	10	7	10	24	26	41	31	38
Origin not stated 2/	18	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	ALL	UNDER 1	1-4	5-9	10-14	15-19	20-24	25 - 29	30-34	35-39	40-44
POPULATION, AND SEX	AGES	YEAR	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS
Diabetes mellitus (E10-E14)											
All origins	68,399	2	8	9	23	51	104	189	393	730	1,212
Male	31,150	-	4	4	12	25	55	94	225	437	702
Female	37,249	2	4	5	11	26	49	95	168	293	510
Hispanic	5,182	-	-	-	4	2	12	14	29	56	101
Male	2,335	-	-	-	2	-	8	8	13	31	53
Female	2,847	-	-	-	2	2	4	6	16	25	48
Mexican	3,274	-	-	-	2	1	9	8	18	31	63
Male	1,444	-	-	-	-	-	6	5	11	17	35
Female	1,830	-	-	-	2	1	3	3	7	14	28
Puerto Rican	662	-	-	-	-	-	2	4	8	11	17
Male	307	-	-	-	-	-	2	2	1	7	6
Female	355	-	-	-	-	-	-	2	7	4	11
Cuban	438	-	-	-	-	-	-	-	-	2	3
Male	205	-	-	-	-	-	-	-	-	1	2
Female	233	-	-	-	-	-	-	-	-	1	1
Central and South American	277	-	-	-	-	1	1	1	2	4	9
Male	119	-	-	-	-	-	-	-	-	1	7
Female	158	-	-	-	-	1	1	1	2	3	2
Other and unknown Hispanic	531	-	-	-	2	-	-	1	1	8	9
Male	260	-	-	-	2	-	-	1	1	5	3
Female	271	-	-	_	_	-	-	-	-	3	6

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diabetes mellitus (E10-E14)											
Non-Hispanic 1/	63,017	2	8	9	19	48	92	173	361	672	1,109
Male	28,719	-	4	4	10	25	47	85	210	406	647
Female	34,298	2	4	5	9	23	45	88	151	266	462
Non-Hispanic white	49,347	2	6	8	11	36	58	122	241	462	769
Male	23,177	-	3	4	5	21	31	63	140	289	458
Female	26,170	2	3	4	6	15	27	59	101	173	311
Non-Hispanic black	11,830	-	2	1	8	12	31	48	112	192	310
Male	4,718	-	1	-	5	4	15	21	64	107	169
Female	7,112	-	1	1	3	8	16	27	48	85	141
Origin not stated 2/	200	-	-	-	-	1	-	2	3	2	2
Male	96	-	-	-	-	-	-	1	2	-	2
Female	104	-	-	-	-	1	-	1	1	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Nutritional deficiencies (E40	-E64)										
All origins	4,289	9	2	1	-	2	6	4	8	20	44
Male	1,486	4	2	1	-	2	3	1	3	8	20
Female	2,803	5	-	-	-	-	3	3	5	12	24
Hispanic	138	-	-	-	-	-	-	-	-	1	3
Male	54	-	-	-	-	-	-	-	-	1	1
Female	84	-	-	-	-	-	-	-	-	-	2
Mexican	96	-	-	-	-	-	-	-	-	1	3
Male	38	-	-	-	-	-	-	-	-	1	1
Female	58	-	-	-	-	-	-	-	-	-	2
Puerto Rican	11	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-
Cuban	6	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-
Central and South American	3	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	22	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	_	_	_	_	-	-	_	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Nutritional deficiencies (E40-E64) Non-Hispanic 1/ 4,142 9 2 2 6 4 8 19 41 1 2 7 Male 1,428 4 2 1 3 1 3 19 5 Female 2,714 5 3 3 12 22 3,486 5 2 3 Non-Hispanic white 1 3 6 14 31 Male 1,142 1 2 3 4 14 Female 2,344 3 3 10 17 Non-Hispanic black 585 3 5 10 Male 250 2 2 3 5 Female 335 2 5 Origin not stated 2/ 9 Male 4 Female 5

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malnutrition (E40-E46)											
All origins	3,985	7	2	1	-	2	5	4	7	20	41
Male	1,371	4	2	1	-	2	2	1	3	8	19
Female	2,614	3	-	-	-	-	3	3	4	12	22
Hispanic	132	-	-	-	-	-	-	-	-	1	3
Male	50	-	-	-	-	-	-	-	-	1	1
Female	82	-	-	-	-	-	-	-	-	-	2
Mexican	92	-	-	-	-	-	-	-	-	1	3
Male	34	-	-	-	-	-	-	-	-	1	1
Female	58	-	-	-	-	-	-	-	-	-	2
Puerto Rican	11	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-
Cuban	6	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-
Central and South American	3	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	20	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	_	_	_	_	_	-	_	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malnutrition (E40-E46)											
Non-Hispanic 1/	3,844	7	2	1	-	2	5	4	7	19	38
Male	1,317	4	2	1	-	2	2	1	3	7	18
Female	2,527	3	-	-	-	-	3	3	4	12	20
Non-Hispanic white	3,232	4	1	-	-	2	3	3	5	14	29
Male	1,047	2	1	-	-	2	1	-	3	4	13
Female	2,185	2	-	-	-	-	2	3	2	10	16
Non-Hispanic black	543	3	1	1	-	-	2	1	2	5	9
Male	234	2	1	1	-	-	1	1	-	3	5
Female	309	1	-	-	-	-	1	-	2	2	4
Origin not stated 2/	9	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	_	_	_	-	-	-	-	_

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 99
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other nutritional deficiencies (E50-E64) 304 2 3 All origins 1 1 Male 115 1 1 Female 189 2 1 2 Hispanic 6 Male Female 2 Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic 2 Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

2

Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 100 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other nutritional deficiencies (E50-E64) Non-Hispanic 1/ 298 2 3 1 Male 111 1 1 Female 187 2 1 2 1 Non-Hispanic white 254 1 2 Male 95 1 Female 159 1 1 1 Non-Hispanic black 42 1 Male 16 Female 26 1 Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 101 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Meningitis (G00,G03)											
All origins	850	116	45	14	13	19	22	21	27	33	43
Male	442	61	26	9	9	10	16	8	16	18	27
Female	408	55	19	5	4	9	6	13	11	15	16
Hispanic	76	17	7	1	-	3	2	2	4	2	2
Male	41	13	5	1	-	-	1	1	2	1	2
Female	35	4	2	-	-	3	1	1	2	1	-
Mexican	50	13	7	-	-	2	2	2	3	-	2
Male	29	10	5	-	-	-	1	1	2	-	2
Female	21	3	2	-	-	2	1	1	1	-	-
Puerto Rican	7	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-
Cuban	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-
Central and South American	7	2	-	1	-	-	-	-	-	-	-
Male	2	1	-	1	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	10	2	-	-	-	1	-	-	-	2	-
Male	6	2	-	-	-	-	-	-	-	1	-
Female	4	-	_	_	_	1	-	-	-	1	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Meningitis (G00,G03) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 103 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Parkinson's disease (G20-G21)											
All origins	14,593	1	-	-	-	-	2	1	1	1	4
Male	8,262	-	-	-	-	-	2	1	1	1	2
Female	6,331	1	-	-	-	-	-	-	-	-	2
Hispanic	389	-	-	-	-	-	-	-	-	-	1
Male	222	-	-	-	-	-	-	-	-	-	1
Female	167	-	-	-	-	-	-	-	-	-	-
Mexican	201	-	-	-	-	-	-	-	-	-	-
Male	113	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-
Puerto Rican	37	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-
Cuban	65	-	-	-	-	-	-	-	-	-	-
Male	37	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-
Central and South American	16	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	70	-	-	-	-	-	-	-	-	-	-
Male	39	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 104
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Parkinson's disease (G20-G21)											
Non-Hispanic 1/	14,177	1	-	-	-	-	2	1	1	1	3
Male	8,029	-	-	-	-	-	2	1	1	1	1
Female	6,148	1	-	-	-	-	-	-	-	-	2
Non-Hispanic white	13,475	1	-	-	-	-	2	1	-	1	3
Male	7,648	-	-	-	-	-	2	1	-	1	1
Female	5,827	1	-	-	-	-	-	-	-	-	2
Non-Hispanic black	502	-	-	-	-	-	-	-	-	-	-
Male	269	-	-	-	-	-	-	-	-	-	-
Female	233	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	27	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 105 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	ALL	UNDER 1	1-4	5-9	10-14	15-19	20-24	25-29	30 - 34	35-39	40-44
POPULATION, AND SEX	AGES	YEAR	YEARS	YEARS	YEARS						
Alzheimer's disease (G30)											
All origins	44,536	-	-	-	-	-	1	-	-	1	6
Male	13,391	-	-	-	-	-	1	-	-	-	3
Female	31,145	-	-	-	-	-	-	-	-	1	3
Hispanic	981	-	-	-	-	-	1	-	-	-	-
Male	314	-	-	-	-	-	1	-	-	-	-
Female	667	-	-	-	-	-	-	-	-	-	-
Mexican	445	-	-	-	-	-	-	-	-	-	-
Male	156	-	-	-	-	-	-	-	-	-	-
Female	289	-	-	-	-	-	-	-	-	-	-
Puerto Rican	108	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-
Female	80	-	-	-	-	-	-	-	-	-	-
Cuban	203	-	-	-	-	-	-	-	-	-	-
Male	60	-	-	-	-	-	-	-	-	-	-
Female	143	-	-	-	-	-	-	-	-	-	-
Central and South American	57	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	1	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	168	-	-	-	-	-	-	-	-	-	-
Male	55	-	-	-	-	-	-	-	-	-	-
Female	113	-	-	-	-	-	_	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 106 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Alzheimer's disease (G30)											
Non-Hispanic 1/	43,471	-	-	-	-	-	-	-	-	1	6
Male	13,052	-	-	-	-	-	-	-	-	-	3
Female	30,419	-	-	-	-	-	-	-	-	1	3
Non-Hispanic white	40,835	-	-	-	-	-	-	-	-	1	4
Male	12,253	-	-	-	-	-	-	-	-	-	2
Female	28,582	-	-	-	-	-	-	-	-	1	2
Non-Hispanic black	2,325	-	-	-	-	-	-	-	-	-	1
Male	689	-	-	-	-	-	-	-	-	-	-
Female	1,636	-	-	-	-	-	-	-	-	-	1
Origin not stated 2/	84	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Major cardiovascular diseases (I00-I78) All origins 950,314 1.270 2.603 5.918 11.092 Male 441.827 1.673 3.898 7,573 Female 508.487 2,020 3.519 Hispanic 33,665 Male 17,304 Female 16,361 Mexican 17,674 Male 9,367 Female 8,307 Puerto Rican 4,513 Male 2,336 Female 2,177 Cuban 4,746 Male 2.256 Female 2.490 Central and South American 2.799 Male 1.285 Female 1,514 Other and unknown Hispanic 3.933 Male 2.060

Source: CDC/NCHS, National Vital Statistics System, Mortality

1.873

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Major cardiovascular diseases (I00-I78) Non-Hispanic 1/ 913,476 528 192 122 184 479 673 1,116 2,337 5,465 10,404 Male 422,824 298 106 70 99 303 391 659 1,483 3,581 7,068 Female 490,652 230 86 52 85 176 282 457 854 1,884 3,336 Non-Hispanic white 793,945 324 121 86 117 318 412 683 1,501 3,684 7,123 Male 366,310 189 70 50 60 196 234 408 992 2,505 5,101 Female 427,635 135 51 36 57 122 178 275 509 1,179 2,022 Non-Hispanic black 103,629 172 60 58 147 241 390 741 1,615 2,995 35 Male 47,982 91 32 19 34 98 142 228 431 955 1,761 Female 55,647 81 28 16 24 49 99 162 310 660 1,234 Origin not stated 2/ 3,173 1 1 2 1 5 7 27 57 Male 1,699 1 1 1 3 5 21 46 Female 1,474 1 1 1 2 2 6 11

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Diseases of heart (I00-I09,I11,I13,I20-I51) All origins 4,703 725,192 2.074 8.897 Male 351,617 1.370 3.252 6.388 Female 373,575 1,451 2.509 Hispanic 25,866 Male 13,554 Female 12,312 Mexican 13,216 Male 7,195 Female 6,021 Puerto Rican 3,666 Male 1,906 Female 1,760 Cuban 3,909 Male 1,860 2,049 Female Central and South American 2.110 Male Female 1.126 Other and unknown Hispanic 2.965 Male 1.609

Source: CDC/NCHS, National Vital Statistics System, Mortality

1.356

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Diseases of heart (I00-I09,I11,I13,I20-I51) Non-Hispanic 1/ 696,719 427 149 98 142 409 541 894 1,876 4,358 8,409 Male 336,615 230 84 55 78 265 324 538 1,224 2,997 6,012 Female 360,104 197 65 43 64 144 217 356 652 1,361 2,397 607,704 2,998 Non-Hispanic white 259 92 65 91 269 316 542 1,213 5,938 Male 293,206 145 55 36 48 167 190 331 821 2,134 4,456 Female 314,498 114 37 29 43 102 126 211 392 864 1,482 Non-Hispanic black 77,713 141 45 128 208 318 595 1,246 2,260 49 32 Male 37,077 71 27 18 27 90 122 191 355 773 1,392 Female 40,636 70 22 18 38 127 240 473 868 Origin not stated 2/ 2,607 1 1 1 2 1 5 5 25 47 Male 1,448 1 1 1 3 4 20 39 Female 1,159 1 1 1 2 1 5 8

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 1 OF 2 PAGE 111

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute rheumatic fever and	l chronic	rheumatic h	eart dise	ases (I00	-109)						
All origins	3,676	6	1	2	3	5	8	12	30	47	77
Male	1,042	2	1	1	1	4	6	5	12	19	30
Female	2,634	4	-	1	2	1	2	7	18	28	47
Hispanic	166	2	-	-	-	-	3	1	3	9	7
Male	38	-	-	-	-	-	3	-	3	2	1
Female	128	2	-	-	-	-	-	1	-	7	6
Mexican	93	2	-	-	-	-	3	-	1	4	4
Male	23	-	-	-	-	-	3	-	1	1	1
Female	70	2	-	-	-	-	-	-	-	3	3
Puerto Rican	17	-	-	-	-	-	-	-	1	3	2
Male	4	-	-	-	-	-	-	-	1	1	-
Female	13	-	-	-	-	-	-	-	-	2	2
Cuban	13	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1
Central and South American	17	-	-	-	-	-	-	1	1	2	-
Male	3	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	1	-	2	-
Other and unknown Hispanic	26	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) Non-Hispanic 1/ 3,498 4 1 3 5 5 11 27 37 69 999 2 5 Male 1 1 1 4 3 9 16 29 2,499 Female 1 2 1 2 6 18 21 40 3,130 2 2 2 3 Non-Hispanic white 1 1 14 24 44 Male 892 1 1 11 20 Female 2,238 1 2 1 3 9 13 24 Non-Hispanic black 269 2 9 11 19 Male 86 2 2 2 8

7

7

1

1

11

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

183

12

5

Female

Male

Female

Origin not stated 2/

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hypertensive heart diseas	e (I11)										
All origins	22,702	1	-	-	-	6	27	50	160	347	617
Male	9,316	-	-	-	-	6	19	41	108	232	428
Female	13,386	1	-	-	-	-	8	9	52	115	189
Hispanic	933	-	-	-	-	1	2	5	13	31	30
Male	455	-	-	-	-	1	1	5	11	24	21
Female	478	-	-	-	-	-	1	-	2	7	9
Mexican	452	-	-	-	-	1	2	4	8	21	19
Male	221	-	-	-	-	1	1	4	7	18	15
Female	231	-	-	-	-	-	1	-	1	3	4
Puerto Rican	159	-	-	-	-	-	-	1	2	4	8
Male	76	-	-	-	-	-	-	1	1	3	5
Female	83	-	-	-	-	-	-	-	1	1	3
Cuban	124	-	-	-	-	-	-	-	1	-	2
Male	66	-	-	-	-	-	-	-	1	-	1
Female	58	-	-	-	-	-	-	-	-	-	1
Central and South American	109	-	-	-	-	-	-	-	1	5	1
Male	50	-	-	-	-	-	-	-	1	3	-
Female	59	-	-	-	-	-	-	-	-	2	1
Other and unknown Hispanic	89	-	-	-	-	-	-	-	1	1	-
Male	42	-	-	-	-	-	-	-	1	-	-
Female	47	-	-	-	-	-	-	-	-	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Hypertensive heart disease (I11) Non-Hispanic 1/ 21,639 1 5 25 45 147 316 582 8,790 97 Male 5 18 36 208 402 9 Female 12,849 50 108 180 8 19 72 Non-Hispanic white 15,654 1 1 139 289 Male 6,013 1 6 17 58 101 222 Female 9,641 1 2 14 38 67 Non-Hispanic black 5,556 3 17 26 72 169 278 Male 2,571 3 12 19 36 101 166 Female 2,985 36 112 Origin not stated 2/ 130 5 Male 71 5 Female 59

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 115 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Hypertensive heart and renal disease (I13) All origins 3,327 2 2 4 5 16 43 71 Male 1,390 1 1 2 3 13 24 43 Female 1,937 1 1 2 2 3 19 28 Hispanic 144 5 4 1 Male 2 65 Female 79 1 3 5 2 Mexican 85 1 37 2 2 Male Female 48 3 Puerto Rican 22 Male 11 Female 11 Cuban 10 Male 5 Female 5 Central and South American 9 1 Male 3 1 Female 6 Other and unknown Hispanic 18 1 Male 9 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

9

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Hypertensive heart and renal disease (I13) Non-Hispanic 1/ 3,171 2 2 5 15 37 65 Male 1,318 1 1 2 3 13 21 38 1 2 2 Female 1,853 1 2 16 27 2,091 5 Non-Hispanic white 1 1 8 11 Male 843 1 5 4 7 Female 1,248 4 4 Non-Hispanic black 989 2 4 10 28 54 Male 426 1 2 8 16 31 Female 563 1 2 2 23 Origin not stated 2/ 12 1 2 7 Male 1 1 Female 5 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Ischemic heart diseases (I20-I25) 27 5 13 9 85 245 769 2,387 All origins 529,659 52 5.364 Male 267,268 13 3 10 7 36 57 169 570 1.801 4.169 Female 262.391 14 2 3 2 16 28 76 199 586 1.195 Hispanic 19,936 2 2 6 5 24 63 128 238 Male 10,604 2 5 3 23 49 112 198 Female 9,332 2 1 2 1 14 16 40 2 3 2 20 37 Mexican 10,073 1 80 139 2 19 72 Male 5,595 1 28 113 Female 4,478 1 1 1 1 9 8 26 Puerto Rican 2,820 2 2 10 16 26 Male 1,474 1 2 6 11 24 Female 1,346 1 4 5 2 Cuban 3,212 2 8 15 Male 1,511 2 7 13 1,701 Female 1 2 2 8 Central and South American 1.612 1 12 32 2 Male 763 1 8 11 28 Female 849 1 4 Other and unknown Hispanic 2.219 1 1 2 6 12 26 2 5 Male 1.261 1 11 20 958

1

1

6

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 30-34 35-39 40-44 ALL 5-9 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Ischemic heart diseases (I20-I25) Non-Hispanic 1/ 507,700 25 5 11 9 45 80 220 706 2,245 5,099 Male 255,524 13 3 8 7 31 54 145 521 1,677 3,946 75 Female 252,176 12 2 3 2 14 26 185 568 1,153 447,960 1,724 Non-Hispanic white 16 4 5 31 45 155 510 3,883 Male 225,909 8 21 36 100 382 1,310 3,106 Female 222,051 8 10 9 55 128 414 777 Non-Hispanic black 51,368 8 12 32 53 160 1,086 466 Male 24,736 4 2 8 16 38 110 317 725 Female 26,632 2 4 16 15 149 361 Origin not stated 2/ 2,023 1 1 14 27 Male 1,140 1 12 25 Female 883 1 2 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 40-44 ALL 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute myocardial infarction (I21-I22) All origins 199.454 2.314 Male 104,169 1.763 Female 95.285 Hispanic 7,158 Male 3,836 Female 3,322 Mexican 3,884 Male 2,170 Female 1,714 Puerto Rican Male Female Cuban 1,134 Male Female Central and South American Male Female Other and unknown Hispanic

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute myocardial infarction (I21-I22) Non-Hispanic 1/ 191,729 11 2 5 6 23 33 91 311 883 2,207 Male 100,014 6 1 3 4 14 23 58 221 664 1,674 91,715 Female 5 1 2 9 10 33 90 219 533 169,545 3 17 227 710 Non-Hispanic white 2 3 18 66 1,736 Male 89,142 3 11 15 41 161 546 1,361 Female 80,403 1 6 25 66 164 375 Non-Hispanic black 19,120 3 13 22 421 6 66 159 Male 9,106 3 3 7 16 45 104 270 Female 10,014 2 3 6 151 Origin not stated 2/ 567 6 11 Male 319 5 11

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

248

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 121 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other acute ischemic heart diseases (I24) 3,232 3 1 3 2 4 25 All origins 1 56 Male 1.665 1 1 1 3 1 3 19 44 Female 1,567 2 1 1 6 12 Hispanic 53 1 1 Male 26 1 1 Female 27 Mexican 35 1 18 Male Female 17 Puerto Rican 7 Male 2 Female 5 Cuban 4 Male Female 4 Central and South American 2 Male 1 Female 1 Other and unknown Hispanic 5 1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

5

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other acute ischemic heart diseases (I24) Non-Hispanic 1/ 3,167 3 3 2 4 25 55 1 Male 1,635 1 1 3 1 3 19 43 Female 1,532 2 1 1 6 12 2,689 3 3 39 Non-Hispanic white 1 1 2 13 Male 1,361 1 3 1 3 10 30 Female 1,328 3 9 Non-Hispanic black 429 1 12 14 Male 246 9 11 Female 183 3 3 Origin not stated 2/ 12 Male 4 Female 8

40-44

YEARS

2,994

35-39

YEARS

1,427

432

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX UNDER 1 ALL 1-4 5 - 9 10-14 15-19 20-24 25-29 30-34 YEARS AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS Other forms of chronic ischemic heart disease (I20,I25) All origins 326,973 11 2 7 3 27 47 142

		,			-	-					-,	_,
	Male	161,434	6	1	6	3	20	30	100	329	1,075	2,362
	Female	165,539	5	1	1	-	7	17	42	103	352	632
Hisp	anic	12,725	-	-	1	-	4	3	14	41	82	141
	Male	6,742	-	-	1	-	3	2	13	32	74	119
	Female	5,983	-	-	-	-	1	1	1	9	8	22
Mex	ican	6,154	-	-	1	-	1	1	12	23	48	78
	Male	3,407	-	-	1	-	-	-	11	18	44	66
	Female	2,747	-	-	-	-	1	1	1	5	4	12
Pue	rto Rican	1,896	-	-	-	-	-	1	2	8	11	14
	Male	987	-	-	-	-	-	1	2	5	8	12
	Female	909	-	-	-	-	-	-	-	3	3	2
Cub	an	2,074	-	-	-	-	-	-	-	1	6	11
	Male	981	-	-	-	-	-	-	-	1	6	9
	Female	1,093	-	-	-	-	-	-	-	-	-	2
Cen	tral and South American	1,102	-	-	-	-	2	1	-	4	8	23
	Male	526	-	-	-	-	2	1	-	4	8	19
	Female	576	-	-	-	-	-	-	-	-	-	4
0th	er and unknown Hispanic	1,499	-	-	-	-	1	-	-	5	9	15
	Male	841	-	-	-	-	1	-	-	4	8	13
	Female	658	-	-	-	-	-	-	-	1	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 POPULATION, AND SEX YEARS AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other forms of chronic ischemic heart disease (I20,I25) Non-Hispanic 1/ 312,804 11 2 6 3 22 44 127 391 1,337 2,837 Male 153,875 6 5 3 17 28 86 297 994 2,229 1 Female 158,929 5 1 5 16 41 94 343 608 275,726 8 2 87 1,001 Non-Hispanic white 1 4 14 24 280 2,108 Male 135,406 2 10 18 58 218 754 1,715 Female 140,320 4 6 29 62 247 393 Non-Hispanic black 31,819 19 31 295 6 651 Male 15,384 5 9 22 65 204 444 Female 16,435 10 9 28 91 207 Origin not stated 2/ 1,444 1 8 16

7

1

14

2

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

817

627

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerotic cardi	iovascular	disease, s	o describ	ed (I25.0)						
All origins	71,926	-	1	-	1	-	13	32	128	453	1,035
Male	36,633	-	-	-	1	-	10	21	98	355	837
Female	35,293	-	1	-	-	-	3	11	30	98	198
Hispanic	3,063	-	-	-	-	-	1	4	10	23	50
Male	1,847	-	-	-	-	-	1	3	9	22	41
Female	1,216	-	-	-	-	-	-	1	1	1	9
Mexican	1,592	-	-	-	-	-	-	3	7	13	30
Male	982	-	-	-	-	-	-	2	7	13	26
Female	610	-	-	-	-	-	-	1	-	-	4
Puerto Rican	460	-	-	-	-	-	1	1	3	1	5
Male	282	-	-	-	-	-	1	1	2	1	4
Female	178	-	-	-	-	-	-	-	1	-	1
Cuban	232	-	-	-	-	-	-	-	-	-	2
Male	124	-	-	-	-	-	-	-	-	-	1
Female	108	-	-	-	-	-	-	-	-	-	1
Central and South American	289	-	-	-	-	-	-	-	-	4	9
Male	167	-	-	-	-	-	-	-	-	4	8
Female	122	-	-	-	-	-	-	-	-	-	1
Other and unknown Hispanic	490	-	-	-	-	-	-	-	-	5	4
Male	292	-	-	-	-	-	-	-	-	4	2
Female	198	-	-	-	-	-	-	-	-	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerotic card	iovascular	disease, s	o describ	ed (I25.0)						
Non-Hispanic 1/	68,373	-	1	-	1	-	12	28	118	427	977
Male	34,451	-	-	-	1	-	9	18	89	330	788
Female	33,922	-	1	-	-	-	3	10	29	97	189
Non-Hispanic white	56,375	-	-	-	1	-	5	16	80	310	707
Male	27,943	-	-	-	1	-	5	11	62	244	594
Female	28,432	-	-	-	-	-	-	5	18	66	113
Non-Hispanic black	10,643	-	1	-	-	-	7	8	33	110	248
Male	5,692	-	-	-	-	-	4	5	22	80	174
Female	4,951	-	1	-	-	-	3	3	11	30	74
Origin not stated 2/	490	-	-	-	-	-	-	-	-	3	8
Male	335	-	-	-	-	-	-	-	-	3	8
Female	155	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other forms of c	hronic isc	hemic heart	disease	(120,125.	1-I25.9)						
All origins	255,047	11	1	7	2	27	34	110	304	974	1,959
Male	124,801	6	1	6	2	20	20	79	231	720	1,525
Female	130,246	5	-	1	-	7	14	31	73	254	434
Hispanic	9,662	-	-	1	-	4	2	10	31	59	91
Male	4,895	-	-	1	-	3	1	10	23	52	78
Female	4,767	-	-	-	-	1	1	-	8	7	13
Mexican	4,562	-	-	1	-	1	1	9	16	35	48
Male	2,425	-	-	1	-	-	-	9	11	31	40
Female	2,137	-	-	-	-	1	1	-	5	4	8
Puerto Rican	1,436	-	-	-	-	-	-	1	5	10	9
Male	705	-	-	-	-	-	-	1	3	7	8
Female	731	-	-	-	-	-	-	-	2	3	1
Cuban	1,842	-	-	-	-	-	-	-	1	6	9
Male	857	-	-	-	-	-	-	-	1	6	8
Female	985	-	-	-	-	-	-	-	-	-	1
Central and South American	813	-	-	-	-	2	1	-	4	4	14
Male	359	-	-	-	-	2	1	-	4	4	11
Female	454	-	-	-	-	-	-	-	-	-	3
Other and unknown Hispanic	1,009	-	-	-	-	1	-	-	5	4	11
Male	549	-	-	-	-	1	-	-	4	4	11
Female	460	-	-	-	-	-	-	-	1	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	, ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other forms of	chronic isch	nemic heart	disease	(120,125.	1-I25.9)						
Non-Hispanic 1/	244,431	11	1	6	2	22	32	99	273	910	1,860
Male	119,424	6	1	5	2	17	19	68	208	664	1,441
Female	125,007	5	-	1	-	5	13	31	65	246	419
Non-Hispanic white	219,351	8	1	4	1	14	19	71	200	691	1,401
Male	107,463	4	1	4	1	10	13	47	156	510	1,121
Female	111,888	4	-	-	-	4	6	24	44	181	280
Non-Hispanic black	21,176	2	-	2	1	6	12	23	60	185	403
Male	9,692	1	-	1	1	5	5	17	43	124	270
Female	11,484	1	-	1	-	1	7	6	17	61	133
Origin not stated 2/	954	-	-	-	-	1	-	1	-	5	8
Male	482	-	-	-	-	-	-	1	-	4	6
Female	472	-	-	-	-	1	-	-	-	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other heart diseases (I26-I51) 1,879 All origins 165,828 1.099 2.768 Male 72.601 1.176 1.718 Female 93,227 1.050 Hispanic 4,687 Male 2,392 Female 2,295 Mexican 2,513 Male 1,319 Female 1,194 Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other heart diseases (I26-I51) Non-Hispanic 1/ 160,711 1,723 2,594 69,984 Male 1,075 1,597 90,727 Female 138,869 Non-Hispanic white 1,103 1,711 Male 59,549 1,101 Female 79,320 Non-Hispanic black 19,531 Male 9,258 Female 10,273 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 131 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute and subacute endocarditis (I33) 1,116 All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute and subacute endocarditis (I33) Non-Hispanic 1/ 1,049 2 1 4 10 13 50 54 2 7 Male 594 3 11 32 37 2 Female 455 1 1 3 18 17 1 3 9 Non-Hispanic white 814 1 8 28 31 Male 466 5 19 20 Female 348 1 3 9 11 Non-Hispanic black 208 4 20 21 Male 115 12 15 Female 93 8 6 Origin not stated 2/ 4 1 Male 3 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diseases of pericardium	and acut	e myocardit	is (I30-I	31, I40)							
All origins	890	27	25	11	10	12	20	19	31	54	60
Male	442	13	14	5	2	8	12	14	20	32	32
Female	448	14	11	6	8	4	8	5	11	22	28
Hispanic	64	2	3	2	2	3	7	2	6	7	2
Male	30	1	1	2	2	3	2	2	4	5	2
Female	34	1	2	-	-	-	5	-	2	2	-
Mexican	38	1	1	2	2	2	6	2	4	3	2
Male	20	-	1	2	2	2	2	2	2	3	2
Female	18	1	-	-	-	-	4	-	2	-	-
Puerto Rican	9	-	-	-	-	-	-	-	2	-	-
Male	4	-	-	-	-	-	-	-	2	-	-
Female	5	-	-	-	-	-	-	-	-	-	-
Cuban	4	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-
Central and South American	7	-	-	-	-	1	1	-	-	2	-
Male	2	-	-	-	-	1	-	-	-	1	-
Female	5	-	-	-	-	-	1	-	-	1	-
Other and unknown Hispanic	6	1	2	-	-	-	-	-	-	2	-
Male	3	1	-	-	-	-	-	-	-	1	-
Female	3	-	2	-	-	-	-	-	-	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diseases of pericardium	and acute	e myocardit	is (I30-I	31,I40)							
Non-Hispanic 1/	824	25	22	9	8	9	13	17	24	47	57
Male	410	12	13	3	-	5	10	12	15	27	29
Female	414	13	9	6	8	4	3	5	9	20	28
Non-Hispanic white	639	10	10	5	4	6	8	11	15	34	36
Male	320	6	4	2	-	4	6	8	9	19	21
Female	319	4	6	3	4	2	2	3	6	15	15
Non-Hispanic black	158	10	11	4	4	3	3	6	8	12	18
Male	76	4	9	1	-	1	2	4	6	7	6
Female	82	6	2	3	4	2	1	2	2	5	12
Origin not stated 2/	2	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	1	-	1
Female	_	-	-	-	=	_	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Heart failure (I50)											
All origins	54,913	11	10	5	7	14	17	27	38	88	184
Male	20,755	5	7	3	3	12	11	13	19	52	112
Female	34,158	6	3	2	4	2	6	14	19	36	72
Hispanic	1,242	3	4	-	1	1	2	2	2	7	11
Male	522	1	3	-	-	1	2	2	1	5	7
Female	720	2	1	-	1	-	-	-	1	2	4
Mexican	678	3	4	-	1	-	2	1	1	6	1
Male	293	1	3	-	-	-	2	1	-	4	-
Female	385	2	1	-	1	-	-	-	1	2	1
Puerto Rican	170	-	-	-	-	-	-	-	1	-	2
Male	73	-	-	-	-	-	-	-	1	-	-
Female	97	-	-	-	-	-	-	-	-	-	2
Cuban	94	-	-	-	-	-	-	-	-	-	-
Male	35	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-
Central and South American	84	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	216	-	-	-	-	1	-	1	-	1	7
Male	91	-	-	-	-	1	-	1	-	1	6
Female	125	_	-	_	-	_	-	-	_	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Heart failure (I50) Non-Hispanic 1/ 53,528 8 6 5 6 13 15 25 36 80 173 20,170 Male 4 4 3 3 11 9 11 18 47 105 2 Female 33,358 2 3 2 6 14 18 33 68 47,977 5 8 7 19 Non-Hispanic white 4 3 5 11 50 106 Male 17,903 2 3 7 4 10 29 71 Female 30,074 2 2 1 7 9 21 35 Non-Hispanic black 4,961 4 13 17 28 65 Male 2,005 2 3 5 7 8 16 34 Female 2,956 1 6 9 31 Origin not stated 2/ 143 1 Male 63 Female 80 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other forms of h	eart disease	(126-128,	I34-I38,I4	12-149,151	l)						
All origins	108,909	450	141	83	132	370	440	622	1,012	1,683	2,464
Male	50,765	235	78	46	78	230	257	364	613	1,058	1,533
Female	58,144	215	63	37	54	140	183	258	399	625	931
Hispanic	3,318	87	25	14	15	40	44	58	100	129	143
Male	1,798	38	15	7	10	21	31	44	70	83	100
Female	1,520	49	10	7	5	19	13	14	30	46	43
Mexican	1,760	62	19	11	8	28	26	41	66	75	89
Male	981	26	13	4	7	16	19	33	47	46	63
Female	779	36	6	7	1	12	7	8	19	29	26
Puerto Rican	456	8	3	-	2	4	7	9	14	19	28
Male	255	4	2	-	1	-	4	5	8	11	20
Female	201	4	1	-	1	4	3	4	6	8	8
Cuban	446	3	-	-	1	1	1	-	1	6	2
Male	235	2	-	-	-	-	1	-	-	4	1
Female	211	1	-	-	1	1	-	-	1	2	1
Central and South America	n 269	9	1	-	2	5	5	6	9	18	12
Male	132	3	-	-	1	3	3	6	7	14	7
Female	137	6	1	-	1	2	2	-	2	4	5
Other and unknown Hispani	c 387	5	2	3	2	2	5	2	10	11	12
Male	195	3	-	3	1	2	4	-	8	8	9
Female	192	2	2	-	1	-	1	2	2	3	3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES.

CLASSIFICATION OF DISEASE 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ES, ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other forms of	heart disease	(126-128,1	34-I38,I4	2-149,151)						
Non-Hispanic 1/	105,310	362	115	69	116	329	395	561	908	1,546	2,310
Male	48,810	197	63	39	67	208	225	319	540	969	1,426
Female	56,500	165	52	30	49	121	170	242	368	577	884
Non-Hispanic white	89,439	224	73	48	75	220	242	330	569	991	1,538
Male	40,860	127	45	26	40	133	135	192	345	641	989
Female	48,579	97	28	22	35	87	107	138	224	350	549
Non-Hispanic black	14,204	119	35	20	35	102	141	211	315	512	719
Male	7,062	60	16	12	24	72	82	119	181	300	407
Female	7,142	59	19	8	11	30	59	92	134	212	312
Origin not stated 2/	281	1	1	-	1	1	1	3	4	8	11
Male	157	-	-	-	1	1	1	1	3	6	7
Female	124	1	1	-	-	-	-	2	1	2	4

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30 - 34 YEARS	35-39 YEARS	40-44 YEARS
Essential (primary) hypert	ension and	hypertensi	ve renal	disease ([110,112)						
All origins	16,968	3	3	-	1	3	10	24	54	105	208
Male	6,488	2	1	-	1	2	6	13	35	69	126
Female	10,480	1	2	-	-	1	4	11	19	36	82
Hispanic	753	1	1	-	-	1	2	4	4	3	16
Male	340	-	-	-	-	1	1	2	3	3	11
Female	413	1	1	-	-	-	1	2	1	-	5
Mexican	416	1	1	-	-	1	1	1	4	1	11
Male	188	-	-	-	-	1	-	1	3	1	7
Female	228	1	1	-	-	-	1	-	1	-	4
Puerto Rican	107	-	-	-	-	-	1	2	-	1	3
Male	51	-	-	-	-	-	1	1	-	1	2
Female	56	-	-	-	-	-	-	1	-	-	1
Cuban	94	-	-	-	-	-	-	-	-	-	-
Male	47	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-
Central and South American	62	-	-	-	-	-	-	-	-	1	-
Male	25	-	-	-	-	-	-	-	-	1	-
Female	37	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	74	-	-	-	-	-	-	1	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	1	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Essential (primary) hypertension and hypertensive renal disease (I10,I12) Non-Hispanic 1/ 16,158 2 2 1 2 8 20 49 101 191 2 Male 6,123 1 1 5 11 31 65 114 1 10,035 9 Female 1 1 3 18 36 77 12,048 1 2 20 78 Non-Hispanic white 3 8 34 Male 4,395 1 5 17 26 51 Female 7,653 1 3 3 8 27 Non-Hispanic black 3,708 5 11 26 65 110 Male 1,550 4 5 12 37 60 Female 2,158 6 14 28 50 Origin not stated 2/ 57 1 1 1 Male 25 1 1 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

32

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Cerebrovascular diseases (I60-I69) All origins 167,366 1.658 Male 64.485 Female 102.881 Hispanic 5,907 Male 2,809 Female 3,098 Mexican 3,418 Male 1,666 Female 1,752 Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Cerebrovascular diseases (I60-I69) Non-Hispanic 1/ 161,057 1,497 Male 61,499 Female 99,558 138,716 Non-Hispanic white Male 51,968 Female 86,748 Non-Hispanic black 18,731 Male 7,823 Female 10,908 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 143 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerosis (I70)											
All origins	14,979	2	-	-	-	1	1	1	7	16	29
Male	5,443	1	-	-	-	-	-	1	4	11	21
Female	9,536	1	-	-	-	1	1	-	3	5	8
Hispanic	410	-	-	-	-	-	-	-	-	1	2
Male	185	-	-	-	-	-	-	-	-	-	1
Female	225	-	-	-	-	-	-	-	-	1	1
Mexican	254	-	-	-	-	-	-	-	-	1	2
Male	120	-	-	-	-	-	-	-	-	-	1
Female	134	-	-	-	-	-	-	-	-	1	1
Puerto Rican	30	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-
Cuban	45	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-
Central and South American	19	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	62	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	_	-	_	-	-	_	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerosis (I70)											
Non-Hispanic 1/	14,519	2	-	-	-	1	1	1	7	15	27
Male	5,236	1	-	-	-	-	-	1	4	11	20
Female	9,283	1	-	-	-	1	1	-	3	4	7
Non-Hispanic white	13,286	1	-	-	-	1	1	1	5	13	17
Male	4,727	1	-	-	-	-	-	1	3	10	13
Female	8,559	-	-	-	-	1	1	-	2	3	4
Non-Hispanic black	1,061	1	-	-	-	-	-	-	2	2	7
Male	430	-	-	-	-	-	-	-	1	1	5
Female	631	1	-	-	-	-	-	-	1	1	2
Origin not stated 2/	50	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 40-44 ALL 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other diseases of circulatory system (I71-I78) All origins 25,809 Male 13,794 Female 12,015 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other diseases of circulatory system (I71-I78) Non-Hispanic 1/ 25,023 10 2 2 7 10 30 45 84 165 280 Male 13,351 8 1 6 8 17 32 64 108 166 11,672 Female 2 1 1 2 13 13 20 57 114 22,191 8 9 32 125 Non-Hispanic white 2 5 24 62 198 Male 12,014 6 7 13 23 48 83 121 Female 10,177 2 1 2 11 9 14 42 77 Non-Hispanic black 2,416 1 12 18 35 77 Male 1,102 2 1 4 8 15 21 40 Female 1,314 37 Origin not stated 2/ 57 Male 27 Female 30

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 40-44 ALL 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Aortic aneurysm and dissection (I71) All origins 15,807 Male 9.615 Female 6.192 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Aortic aneurysm and dissection (I71) Non-Hispanic 1/ 15,393 1 2 6 21 30 57 124 182 1 9,342 Male 1 2 5 12 23 51 91 124 7 Female 6,051 1 1 9 6 33 58 13,934 1 6 22 42 95 122 Non-Hispanic white 1 1 17 Male 8,551 1 5 17 37 70 85 Female 5,383 1 1 5 5 25 37 Non-Hispanic black 1,172 4 8 13 24 56 Male 615 3 13 17 35 Female 557 2 7 21 Origin not stated 2/ 30 Male 17 Female 13

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 149 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other diseases of arterio	es, arteri	oles and ca	pillaries	(172-178	()						
All origins	10,002	11	1	2	6	6	12	18	30	44	105
Male	4,179	8	-	1	4	4	8	10	14	19	45
Female	5,823	3	1	1	2	2	4	8	16	25	60
Hispanic	345	2	-	-	1	2	3	3	3	3	7
Male	160	1	-	-	-	1	3	1	1	2	3
Female	185	1	-	-	1	1	-	2	2	1	4
Mexican	205	-	-	-	1	2	3	3	2	2	5
Male	96	-	-	-	-	1	3	1	1	2	3
Female	109	-	-	-	1	1	-	2	1	-	2
Puerto Rican	30	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	1	-
Cuban	38	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-
Central and South American	29	1	-	-	-	-	-	-	1	-	2
Male	14	-	-	-	-	-	-	-	-	-	-
Female	15	1	-	-	-	-	-	-	1	-	2
Other and unknown Hispanic	43	1	-	-	-	-	-	-	-	-	-
Male	21	1	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX YEARS AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other diseases of arteries, arterioles and capillaries (I72-I78) Non-Hispanic 1/ 9,630 9 1 2 5 4 9 15 27 41 98 4,009 9 Male 7 4 3 5 13 17 42 1 Female 5,621 1 1 1 4 6 14 24 56 8,257 3 10 20 Non-Hispanic white 1 2 4 30 76 Male 3,463 3 2 6 11 13 36 Female 4,794 1 1 1 9 17 40 Non-Hispanic black 1,244 1 11 21 Male 487 2 1 2 5

2

3

7

16

Source: CDC/NCHS, National Vital Statistics System, Mortality

757

27

10

17

Female

Male

Female

Origin not stated 2/

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 40-44 ALL 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other disorders of circulatory system (I80-I99) All origins 4,025 Male 1,714 Female 2,311 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other disorders of circulatory system (I80-I99) Non-Hispanic 1/ 3,825 21 1 9 17 30 56 92 164 1 Male 1,623 13 1 4 7 14 26 50 98 Female 2,202 8 1 5 10 16 30 42 66 17 3 9 21 Non-Hispanic white 3,149 1 36 60 108 Male 1,310 9 2 10 16 32 63 Female 1,839 8 1 1 11 20 28 45 Non-Hispanic black 6 9 19 29 52 613 Male 281 3 2 10 16 33 Female 332 4 9 13 19 Origin not stated 2/ 14 1 2 1 2 Male 8 1 Female 6 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Influenza and pneumonia (J10-	J18)										
All origins	63,730	320	130	46	47	73	106	135	204	462	601
Male	27,718	171	66	27	26	34	56	76	121	251	341
Female	36,012	149	64	19	21	39	50	59	83	211	260
Hispanic	2,246	64	17	5	5	10	15	10	14	46	28
Male	1,092	40	10	4	2	5	6	6	9	34	20
Female	1,154	24	7	1	3	5	9	4	5	12	8
Mexican	1,085	46	12	4	3	6	10	7	9	27	15
Male	561	28	7	4	1	3	5	5	7	21	9
Female	524	18	5	-	2	3	5	2	2	6	6
Puerto Rican	354	5	1	1	-	2	3	1	1	10	5
Male	157	4	1	-	-	1	1	-	-	6	4
Female	197	1	-	1	-	1	2	1	1	4	1
Cuban	310	1	-	-	-	-	-	-	-	2	-
Male	134	1	-	-	-	-	-	-	-	2	-
Female	176	-	-	-	-	-	-	-	-	-	-
Central and South American	188	4	3	-	-	1	1	-	2	2	4
Male	86	2	2	-	-	-	-	-	1	1	4
Female	102	2	1	-	-	1	1	-	1	1	-
Other and unknown Hispanic	309	8	1	-	2	1	1	2	2	5	4
Male	154	5	-	-	1	1	-	1	1	4	3
Female	155	3	1	-	1	-	1	1	1	1	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Influenza and pneumonia (J10-J18) Non-Hispanic 1/ 61,218 26,489 Male 34,729 Female Non-Hispanic white 54,280 Male 23,097 Female 31,183 Non-Hispanic black 5,794 Male 2,786 Female 3,008 Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Influenza (J10-J11)											
All origins	1,665	13	12	6	5	7	4	2	7	8	8
Male	602	6	9	4	1	1	1	-	5	5	4
Female	1,063	7	3	2	4	6	3	2	2	3	4
Hispanic	45	4	3	-	-	1	1	-	-	1	1
Male	21	-	3	-	-	-	-	-	-	1	1
Female	24	4	-	-	-	1	1	-	-	-	-
Mexican	26	3	2	-	-	-	1	-	-	-	1
Male	13	-	2	-	-	-	-	-	-	-	1
Female	13	3	-	-	-	-	1	-	-	-	-
Puerto Rican	3	1	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	1	-
Female	1	1	-	-	-	-	-	-	-	-	-
Cuban	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Central and South American	3	-	1	-	-	1	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-
Other and unknown Hispanic	12	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS Influenza (J10-J11) 1,612 Non-Hispanic 1/ 9 9 6 5 6 3 2 7 7 7 580 Male 6 6 4 1 1 1 5 4 3 1,032 2 5 2 2 Female 3 3 4 2 3 4 1,522 3 5 6 5 3 3 Non-Hispanic white 4 1 5 Male 547 3 1 2 2 Female 975 2 3 4 1 1 3 Non-Hispanic black 1 3 2 65 Male 24 3 1 1 Female 41 1 2 1 Origin not stated 2/ 8 Male 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	ALL	UNDER 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44
POPULATION, AND SEX	AGES	YEAR	YEARS								
Pneumonia (J12-J18)											
All origins	62,065	307	118	40	42	66	102	133	197	454	593
Male	27,116	165	57	23	25	33	55	76	116	246	337
Female	34,949	142	61	17	17	33	47	57	81	208	256
Hispanic	2,201	60	14	5	5	9	14	10	14	45	27
Male	1,071	40	7	4	2	5	6	6	9	33	19
Female	1,130	20	7	1	3	4	8	4	5	12	8
Mexican	1,059	43	10	4	3	6	9	7	9	27	14
Male	548	28	5	4	1	3	5	5	7	21	8
Female	511	15	5	-	2	3	4	2	2	6	6
Puerto Rican	351	4	1	1	-	2	3	1	1	9	5
Male	155	4	1	-	-	1	1	-	-	5	4
Female	196	-	-	1	-	1	2	1	1	4	1
Cuban	309	1	-	-	-	-	-	-	-	2	-
Male	134	1	-	-	-	-	-	-	-	2	-
Female	175	-	-	-	-	-	-	-	-	-	-
Central and South American	185	4	2	-	-	-	1	-	2	2	4
Male	84	2	1	-	-	-	-	-	1	1	4
Female	101	2	1	-	-	-	1	-	1	1	-
Other and unknown Hispanic	297	8	1	-	2	1	1	2	2	5	4
Male	150	5	-	-	1	1	-	1	1	4	3
Female	147	3	1	-	1	-	1	1	1	1	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Pneumonia (J12-J18) Non-Hispanic 1/ 59,606 25,909 Male Female 33,697 52,758 Non-Hispanic white Male 22,550 Female 30,208 Non-Hispanic black 5,729 Male 2,762 Female 2,967 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 159 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other acute lower respiratory infections (J20-J22) All origins 540 63 25 2 3 3 2 2 2 6 5 Male 196 36 15 2 1 2 1 3 2 Female 344 27 10 2 1 2 2 1 3 3 Hispanic 21 10 3 5 Male 10 2 5 Female 11 1 8 3 Mexican 13 3 Male 6 Female 7 5 1 Puerto Rican 2 1 Male 2 1 Female Cuban 4 Male 1 Female 3 Central and South American 1 1 Male 1 1 Female Other and unknown Hispanic 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

Male Female 01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 160 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other acute lower respiratory infections (J20-J22) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 161 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 35-39 40-44 ALL 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute bronchitis and bronchiolitis (J20-J21) All origins 388 62 24 2 2 3 2 1 2 6 4 Male 154 36 14 2 1 1 1 3 1 Female 234 26 10 2 2 2 1 3 3 Hispanic 20 10 3 1 5 Male 10 2 5 Female 10 1 13 8 3 Mexican 3 Male 6 Female 7 5 1 Puerto Rican 2 1 Male 2 1 Female Cuban 3 Male 1 Female 2 Central and South American 1 1 Male 1 1 Female Other and unknown Hispanic 1 Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Acute bronchitis and bronchiolitis (J20-J21) Non-Hispanic 1/ 367 52 21 2 1 3 2 1 2 5 4 Male 143 31 12 1 1 1 1 2 1 2 3 Female 224 21 9 1 3 299 25 12 2 2 2 5 Non-Hispanic white 1 1 1 2 Male 108 14 6 1 1 1 1 2 Female 191 11 6 1 1 2 3 2 Non-Hispanic black 24 1 55 Male 26 15 1 Female 29 9 3 1 1 Origin not stated 2/ 1 1 Male 1 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 1 OF 2 PAGE 163

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Unspecified acute lower res	spiratory	infection ([J22)								
All origins	152	1	1	-	1	-	-	1	-	-	1
Male	42	-	1	-	-	-	-	1	-	-	1
Female	110	1	-	-	1	-	-	-	-	-	-
Hispanic	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Cuban	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 1 OF 2 PAGE 164

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Unspecified acute lower re	espiratory	infection ((J22)								
Non-Hispanic 1/	151	1	1	-	1	-	-	1	-	-	1
Male	42	-	1	-	-	-	-	1	-	-	1
Female	109	1	-	-	1	-	-	-	-	-	-
Non-Hispanic white	141	1	-	-	1	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	1
Female	104	1	-	-	1	-	-	-	-	-	-
Non-Hispanic black	9	-	1	-	-	-	-	1	-	-	-
Male	4	-	1	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	_	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 40-44 ALL 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Chronic lower respiratory diseases (J40-J47) All origins 124,181 Male 62.415 Female 61,766 Hispanic 2,859 Male 1,547 Female 1,312 Mexican 1,297 Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Chronic lower respiratory diseases (J40-J47) Non-Hispanic 1/ 120,968 Male 60,669 60,299 Female Non-Hispanic white 111,620 Male 55,314 Female 56,306 Non-Hispanic black 7,840 Male 4,452 Female 3,388 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 167 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 35-39 40-44 ALL 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Bronchitis, chronic and unspecified (J40-J42) All origins 1,172 20 19 3 1 3 4 3 8 10 8 Male 498 9 11 2 1 2 2 3 3 2 7 Female 674 11 8 1 1 2 3 5 6 Hispanic 3 3 1 2 4 3 52 Male 17 1 2 1 1 2 3 Female 35 1 2 2 3 2 2 2 Mexican 30 10 1 Male Female 20 2 1 1 1 2 Puerto Rican 3 Male Female 3 Cuban 7 1 Male 3 1 Female 4 Central and South American 3 1 Male 3 1 Female Other and unknown Hispanic 9 1 2 Male 1

2

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

8

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Bronchitis, chronic and unspecified (J40-J42) Non-Hispanic 1/ 1,117 17 16 3 3 1 8 6 5 478 2 Male 8 9 2 2 3 2 1 9 7 2 5 Female 639 1 1 1 4 4 1,002 8 12 3 3 Non-Hispanic white 3 6 6 3 Male 421 6 2 2 1 Female 581 3 6 1 4 4 2 Non-Hispanic black 88 2 2 Male 43 3 Female 45 2 Origin not stated 2/ 3 Male 3 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Emphysema (J43)											
All origins	17,787	1	-	1	-	-	1	1	8	19	78
Male	9,480	-	-	-	-	-	1	-	6	9	40
Female	8,307	1	-	1	-	-	-	1	2	10	38
Hispanic	292	-	-	1	-	-	-	-	2	2	4
Male	183	-	-	-	-	-	-	-	1	-	2
Female	109	-	-	1	-	-	-	-	1	2	2
Mexican	137	-	-	1	-	-	-	-	-	1	2
Male	83	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	1	-	-	-	-	-	1	1
Puerto Rican	28	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	1
Cuban	59	-	-	-	-	-	-	-	-	-	-
Male	37	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-
Central and South American	16	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-
Other and unknown Hispanic	52	-	-	-	-	-	-	-	1	-	1
Male	32	-	-	-	-	-	-	-	1	-	1
Female	20	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Emphysema (J43)											
Non-Hispanic 1/	17,441	1	-	-	-	-	1	1	6	16	72
Male	9,260	-	-	-	-	-	1	-	5	8	37
Female	8,181	1	-	-	-	-	-	1	1	8	35
Non-Hispanic white	16,338	-	-	-	-	-	-	1	4	13	51
Male	8,540	-	-	-	-	-	-	-	3	5	28
Female	7,798	-	-	-	-	-	-	1	1	8	23
Non-Hispanic black	923	1	-	-	-	-	-	-	-	2	20
Male	594	-	-	-	-	-	-	-	-	2	9
Female	329	1	-	-	-	-	-	-	-	-	11
Origin not stated 2/	54	-	-	-	-	-	-	-	-	1	2
Male	37	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	_	_	_	-	-	-	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30 - 34 YEARS	35 - 39 YEARS	40-44 YEARS
Asthma (J45-J46)											
All origins	4,657	5	27	40	86	95	87	121	141	206	241
Male	1,620	3	16	25	60	53	55	64	64	94	90
Female	3,037	2	11	15	26	42	32	57	77	112	151
Hispanic	320	-	3	4	10	8	6	12	9	24	22
Male	119	-	-	2	7	4	4	6	5	9	13
Female	201	-	3	2	3	4	2	6	4	15	9
Mexican	107	-	1	2	6	4	2	5	2	9	5
Male	42	-	-	2	4	2	2	3	1	4	4
Female	65	-	1	-	2	2	-	2	1	5	1
Puerto Rican	110	-	2	1	2	2	3	4	4	8	8
Male	41	-	-	-	2	-	2	3	1	3	4
Female	69	-	2	1	-	2	1	1	3	5	4
Cuban	33	-	-	-	-	-	-	-	1	1	3
Male	10	-	-	-	-	-	-	-	1	-	1
Female	23	-	-	-	-	-	-	-	-	1	2
Central and South American	42	-	-	-	1	2	-	2	2	5	3
Male	17	-	-	-	-	2	-	-	2	1	3
Female	25	-	-	-	1	-	-	2	-	4	-
Other and unknown Hispanic	28	-	-	1	1	-	1	1	-	1	3
Male	9	-	-	-	1	-	-	-	-	1	1
Female	19	-	-	1	-	-	1	1	-	-	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Asthma (J45-J46)											
Non-Hispanic 1/	4,324	5	24	36	76	87	81	109	129	182	218
Male	1,495	3	16	23	53	49	51	58	58	85	77
Female	2,829	2	8	13	23	38	30	51	71	97	141
Non-Hispanic white	3,011	1	8	17	30	51	41	44	63	99	123
Male	930	1	5	10	21	28	24	22	26	44	36
Female	2,081	-	3	7	9	23	17	22	37	55	87
Non-Hispanic black	1,134	4	14	19	46	33	38	62	64	77	89
Male	474	2	10	13	32	20	26	34	32	36	36
Female	660	2	4	6	14	13	12	28	32	41	53
Origin not stated 2/	13	-	-	-	-	-	-	-	3	-	1
Male	6	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	2	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other chronic lower respira	atory dise	ases (J44,J	47)								
All origins	100,565	10	8	5	3	9	10	20	28	67	261
Male	50,817	7	3	2	3	5	5	5	10	36	136
Female	49,748	3	5	3	-	4	5	15	18	31	125
Hispanic	2,195	3	2	2	-	-	1	3	3	4	16
Male	1,228	3	1	1	-	-	1	-	1	4	6
Female	967	-	1	1	-	-	-	3	2	-	10
Mexican	1,023	2	1	1	-	-	1	-	2	1	8
Male	595	2	1	-	-	-	1	-	1	1	1
Female	428	-	-	1	-	-	-	-	1	-	7
Puerto Rican	294	-	-	1	-	-	-	2	-	-	4
Male	151	-	-	1	-	-	-	-	-	-	2
Female	143	-	-	-	-	-	-	2	-	-	2
Cuban	389	-	-	-	-	-	-	-	-	-	-
Male	231	-	-	-	-	-	-	-	-	-	-
Female	158	-	-	-	-	-	-	-	-	-	-
Central and South American	125	1	1	-	-	-	-	-	-	3	1
Male	57	1	-	-	-	-	-	-	-	3	1
Female	68	-	1	-	-	-	-	-	-	-	-
Other and unknown Hispanic	364	-	-	-	-	-	-	1	1	-	3
Male	194	-	-	-	-	-	-	-	-	-	2
Female	170	-	-	-	-	-	-	1	1	-	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other chronic lower respiratory diseases (J44,J47) Non-Hispanic 1/ 98,086 7 6 3 3 9 9 17 25 63 245 5 Male 49,436 4 2 1 3 5 4 9 32 130 Female 48,650 3 4 2 4 5 12 16 31 115 91,269 2 12 Non-Hispanic white 1 3 3 6 6 16 49 189 Male 45,423 2 2 4 9 26 97 Female 45,846 1 3 4 8 23 92 Non-Hispanic black 5,695 3 14 54 Male 3,341 4 3 6 32 Female 2,354 7 8 22 Origin not stated 2/ 284

Source: CDC/NCHS, National Vital Statistics System, Mortality

153

131

Male

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 175 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Pneumoconioses and chemical effects (J60-J66,J68) All origins 1,113 3 1 3 4 Male 1,079 3 1 2 3 Female 34 1 1 Hispanic 13 1 Male 11 1 Female 2 9 Mexican 1 Male 1 Female Puerto Rican Male Female Cuban 4 Male 3 Female 1 Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 176 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Pneumoconioses and chemical effects (J60-J66,J68) Non-Hispanic 1/ 1,100 3 3 3 1 Male 1,068 3 1 2 2 Female 32 1 1 1,024 Non-Hispanic white 1 2 Male 995 1 1 Female 29 1 Non-Hispanic black 67 3 2 1 Male 64 3 1 1 Female 1 Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Pneumonitis due to solids and	d liquids	(J69)									
All origins	15,268	16	12	4	12	9	23	33	45	59	107
Male	7,755	9	3	-	6	7	15	21	26	35	62
Female	7,513	7	9	4	6	2	8	12	19	24	45
Hispanic	413	2	2	2	3	1	1	3	1	1	2
Male	195	1	1	-	1	-	-	2	1	1	-
Female	218	1	1	2	2	1	1	1	-	-	2
Mexican	199	2	-	1	1	-	-	2	1	1	-
Male	102	1	-	-	1	-	-	1	1	1	-
Female	97	1	-	1	-	-	-	1	-	-	-
Puerto Rican	56	-	1	1	1	1	1	1	-	-	2
Male	23	-	-	-	-	-	-	1	-	-	-
Female	33	-	1	1	1	1	1	-	-	-	2
Cuban	79	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	-
Central and South American	19	-	-	-	1	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	1	-	-	-	-	-	-
Other and unknown Hispanic	60	-	1	-	-	-	-	-	-	-	-
Male	36	-	1	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Pneumonitis due to solids and liquids (J69) Non-Hispanic 1/ 14,812 Male 7,534 7,278 Female Non-Hispanic white 13,139 Male 6,642 Female 6,497 Non-Hispanic black 1,495 Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX YEARS AGES YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98) All origins 25,031 Male 12.110 Female 12,921 Hispanic 1,157 Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98) Non-Hispanic 1/ 23,802 Male 11,483 Female 12,319 20,959 Non-Hispanic white Male 10,154 Female 10,805 Non-Hispanic black 2,339 Male 1,081 Female 1,258 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 181 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	ALL	UNDER 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44
POPULATION, AND SEX	AGES	YEAR	YEARS								
Peptic ulcer (K25-K28)											
All origins	4,604	2	2	-	-	1	1	7	14	38	85
Male	2,189	-	-	-	-	1	1	5	7	27	49
Female	2,415	2	2	-	-	-	-	2	7	11	36
Hispanic	172	-	1	-	-	-	-	4	4	1	3
Male	101	-	-	-	-	-	-	3	2	1	2
Female	71	-	1	-	-	-	-	1	2	-	1
Mexican	99	-	1	-	-	-	-	4	2	-	2
Male	58	-	-	-	-	-	-	3	-	-	1
Female	41	-	1	-	-	-	-	1	2	-	1
Puerto Rican	19	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-
Cuban	14	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-
Central and South American	15	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	25	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Peptic ulcer (K25-K28)											
Non-Hispanic 1/	4,415	2	1	-	-	1	1	3	10	37	81
Male	2,079	-	-	-	-	1	1	2	5	26	46
Female	2,336	2	1	-	-	-	-	1	5	11	35
Non-Hispanic white	3,859	1	1	-	-	1	1	2	5	20	53
Male	1,783	-	-	-	-	1	1	2	1	15	26
Female	2,076	1	1	-	-	-	-	-	4	5	27
Non-Hispanic black	443	1	-	-	-	-	-	-	4	16	27
Male	240	-	-	-	-	-	-	-	3	11	19
Female	203	1	-	-	-	-	-	-	1	5	8
Origin not stated 2/	17	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 183
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 35-39 40-44 ALL 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Diseases of appendix (K35-K38) All origins 390 3 2 12 14 2 3 3 3 13 11 Male 236 1 2 6 9 2 3 3 2 10 7 Female 154 2 6 5 1 3 4 Hispanic 29 1 5 3 2 4 2 Male 20 1 2 2 9 Female 3 1 5 2 3 Mexican 21 14 3 Male Female 7 Puerto Rican 3 Male 2 Female 1 Cuban Male Female Central and South American 5 1 1 Male 4 1 1 Female 1 Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 184
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Diseases of appendix (K35-K38) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 185 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hernia (K40-K46)											
All origins	1,464	44	3	1	1	1	2	6	10	15	24
Male	597	26	2	1	-	1	2	5	3	10	13
Female	867	18	1	-	1	-	-	1	7	5	11
Hispanic	66	3	1	-	-	-	-	-	-	2	1
Male	27	2	1	-	-	-	-	-	-	2	-
Female	39	1	-	-	-	-	-	-	-	-	1
Mexican	45	3	1	-	-	-	-	-	-	2	1
Male	17	2	1	-	-	-	-	-	-	2	-
Female	28	1	-	-	-	-	-	-	-	-	1
Puerto Rican	4	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-
Cuban	6	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-
Central and South American	2	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	9	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	_	-	_	-	-	-	-	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Hernia (K40-K46) 1,390 Non-Hispanic 1/ 41 2 1 1 1 2 6 10 13 23 Male 566 24 1 1 1 2 5 3 8 13 Female 824 17 1 1 1 5 10 1,227 Non-Hispanic white 31 2 1 1 1 1 4 6 12 15 Male 505 18 1 1 1 3 7 11 Female 722 13 1 1 4 5 4 Non-Hispanic black 139 9 1 8 Male 6 1 2 53 Female 86 3 3 6 Origin not stated 2/ 8 Male 4 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Chronic liver disease and cirrhosis (K70,K73-K74) All origins 26,259 1,103 2.199 Male 17.115 1.598 Female 9.144 Hispanic 2,972 Male 2,143 Female Mexican 1,873 1,372 Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Chronic liver disease and cirrhosis (K70,K73-K74) Non-Hispanic 1/ 23,185 5 2 1 5 13 63 263 909 1,858 1 Male 14,896 4 1 6 35 159 560 1,321 Female 8,289 1 2 5 28 104 349 537 3 45 Non-Hispanic white 19,602 4 1 4 198 712 1,436 Male 12,613 29 122 445 1,033 Female 6,989 1 3 16 76 267 403 Non-Hispanic black 2,793 2 8 16 38 129 324 Male 1,812 5 22 75 218 Female 981 2 2 11 16 106 Origin not stated 2/ 102 1 4 13 Male 76 1 2 10 Female 26 2 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 25-29 40-44 ALL 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Alcoholic liver disease (K70) 11,958 8 37 222 763 1,423 All origins Male 8,922 6 24 141 491 1.029 Female 3.036 2 13 81 272 394 Hispanic 1,547 2 4 46 128 214 2 Male 1,294 37 101 175 9 39 Female 253 27 2 2 28 90 Mexican 981 139 829 22 71 Male 116 Female 152 6 19 23 Puerto Rican 225 12 28 Male 190 9 24 Female 35 3 4 Cuban 31 1 Male 29 1 Female 2 Central and South American 12 8 118 1 16 Male 102 1 10 8 14 Female 16 2 2 Other and unknown Hispanic 192 1 6 17 31 5 Male 1 12 21 144 5

1

10

Source: CDC/NCHS, National Vital Statistics System, Mortality

48

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Alcoholic liver disease (K70) Non-Hispanic 1/ 10,353 6 33 175 631 1,200 Male 7,580 4 20 103 388 847 2,773 Female 2 13 72 243 353 8,470 3 27 492 Non-Hispanic white 128 922 Male 6,288 17 78 307 660 Female 2,182 10 50 185 262 Non-Hispanic black 1,445 84 197 24 Male 1,001 2 2 12 49 129 Female 444 2 12 35 68 Origin not stated 2/ 58 1 4 9 7 Male 48 1 2 Female 10 2 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other chronic liver disease	and cirr	hosis (K73-	K74)								
All origins	14,301	5	2	1	1	9	9	33	115	340	776
Male	8,193	4	-	1	-	3	2	17	75	225	569
Female	6,108	1	2	-	1	6	7	16	40	115	207
Hispanic	1,425	-	-	-	-	4	2	3	27	62	114
Male	849	-	-	-	-	3	-	2	19	53	92
Female	576	-	-	-	-	1	2	1	8	9	22
Mexican	892	-	-	-	-	2	1	3	20	45	75
Male	543	-	-	-	-	1	-	2	15	40	57
Female	349	-	-	-	-	1	1	1	5	5	18
Puerto Rican	197	-	-	-	-	1	1	-	2	6	18
Male	135	-	-	-	-	1	-	-	1	5	17
Female	62	-	-	-	-	-	1	-	1	1	1
Cuban	89	-	-	-	-	-	-	-	1	-	3
Male	42	-	-	-	-	-	-	-	1	-	3
Female	47	-	-	-	-	-	-	-	-	-	-
Central and South American	101	-	-	-	-	1	-	-	2	6	7
Male	49	-	-	-	-	1	-	-	1	5	6
Female	52	-	-	-	-	-	-	-	1	1	1
Other and unknown Hispanic	146	-	-	-	-	-	-	-	2	5	11
Male	80	-	-	-	-	-	-	-	1	3	9

1

2

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other chronic liver disease and cirrhosis (K73-K74) Non-Hispanic 1/ 12,832 5 2 1 1 5 7 30 88 278 658 Male 7,316 4 1 2 15 56 172 474 5 Female 5,516 1 2 1 5 15 32 106 184 3 220 Non-Hispanic white 11,132 4 1 1 18 70 514 Male 6,325 1 12 44 138 373 Female 4,807 1 3 6 26 82 141 Non-Hispanic black 1,348 2 12 45 127 14 Male 811 3 10 26 89 Female 537 2 9 19 38 Origin not stated 2/ 44 4 Male 28 3

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

16

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 193 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 25-29 30-34 40-44 ALL 1-4 10-14 20-24 35 - 39POPULATION, AND SEX YEARS AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Cholelithiasis and other disorders of gallbladder (K80-K82) All origins 2,830 2 1 4 5 11 25 6 34 Male 1,248 1 1 2 3 14 16 Female 1,582 1 2 3 5 7 11 18 Hispanic 185 1 2 6 5 Male 84 1 3 3 Female 101 1 3 2 5 2 Mexican 109 3 Male 52 1 Female 57 2 1 Puerto Rican 9 1 Male 2 Female 7 1 Cuban 20 Male 10 Female 10

1

1

2

1

1

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

19

5

14

28

15

13

Central and South American

Other and unknown Hispanic

Male

Male

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Cholelithiasis and other disorders of gallbladder (K80-K82) Non-Hispanic 1/ 2,639 2 6 9 19 29 1 4 4 Male 1,163 1 1 2 3 3 11 13 1,476 2 Female 1 3 4 6 8 16 2,299 3 3 Non-Hispanic white 1 3 11 15 Male 1,017 1 1 3 7 7 Female 1,282 2 3 6 4 8 Non-Hispanic black 257 1 2 7 13 Male 102 1 2 3 6 Female 155 4 Origin not stated 2/ 6 Male 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

5

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) All origins 35,525 Male 17.016 Female 18,509 Hispanic 1,584 Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) Non-Hispanic 1/ 33,830 Male 16,143 Female 17,687 Non-Hispanic white 26,418 Male 12,793 Female 13,625 Non-Hispanic black 6,659 Male 2,984 Female 3,675 Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 1 OF 2 PAGE 197

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEAS 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX		UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute and rapidly prog	ressive nephri	tic and nep	hrotic sy	ndrome (N	N00-N01,N6	04)					
All origins	182	3	2	3	2	-	2	3	2	2	3
Male	90	3	2	1	1	-	2	2	2	-	2
Female	92	-	-	2	1	-	-	1	-	2	1
Hispanic	12	1	-	1	1	-	-	-	-	-	-
Male	4	1	-	-	1	-	-	-	-	-	-
Female	8	-	-	1	-	-	-	-	-	-	-
Mexican	8	1	-	1	1	-	-	-	-	-	-
Male	4	1	-	-	1	-	-	-	-	-	-
Female	4	-	-	1	-	-	-	-	-	-	-
Puerto Rican	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-
Cuban	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Central and South Ameri	can 1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispa	nic -	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute and rapidly progress	sive nephri	tic and nep	hrotic sy	ndrome (N	00-N01,N6	94)					
Non-Hispanic 1/	170	2	2	2	1	-	2	3	2	2	3
Male	86	2	2	1	-	-	2	2	2	-	2
Female	84	-	-	1	1	-	-	1	-	2	1
Non-Hispanic white	132	2	2	2	-	-	1	1	1	2	1
Male	63	2	2	1	-	-	1	-	1	-	1
Female	69	-	-	1	-	-	-	1	-	2	-
Non-Hispanic black	28	-	-	-	1	-	1	2	-	-	2
Male	16	-	-	-	-	-	1	2	-	-	1
Female	12	-	-	-	1	-	-	-	-	-	1
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 1 OF 2 PAGE 199

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Chronic glomerulonephritis, 05-N07,N26)	nephritis	and nephr	opathy not	: specifi	ed as acı	ite or chr	onic,and	renal scle	rosis unsp	ecified (No	92-N03,N
All origins	589	-	1	1	-	3	4	10	5	15	13
Male	304	-	-	1	-	3	3	7	3	9	9
Female	285	-	1	-	-	-	1	3	2	6	4
Hispanic	33	-	1	-	-	-	1	2	1	-	2
Male	18	-	-	-	-	-	1	2	1	-	2
Female	15	-	1	-	-	-	-	-	-	-	-
Mexican	16	-	1	-	-	-	-	2	1	-	-
Male	6	-	-	-	-	-	-	2	1	-	-
Female	10	-	1	-	-	-	-	-	-	-	-
Puerto Rican	5	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Cuban	2	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-
Central and South American	5	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	5	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	2
Female	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 200 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992 RACE	SIFICATION OF DISEASES,), HISPANIC ORIGIN, FOR NON-HISPANIC LATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
	ronic glomerulonephritis, 07,N26)	nephritis	and nephro	opathy not	specifie	ed as acut	e or chr	onic,and r	renal scler	osis unspe	ecified (NG	2-N03,N
Non-	Hispanic 1/	555	-	-	1	-	3	3	8	4	15	11
	Male	285	-	-	1	-	3	2	5	2	9	7
	Female	270	-	-	-	-	-	1	3	2	6	4
Non	-Hispanic white	436	-	-	1	-	2	1	6	2	8	5
	Male	227	-	-	1	-	2	-	4	1	4	4
	Female	209	-	-	-	-	-	1	2	1	4	1
Non	-Hispanic black	92	-	-	-	-	1	2	-	2	6	5
	Male	46	-	-	-	-	1	2	-	1	4	2
	Female	46	-	-	-	-	-	-	-	1	2	3
0rig	in not stated 2/	1	-	-	-	-	-	-	-	-	-	-
	Male	1	-	-	-	-	-	-	-	-	-	-
	Female	-	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Renal failure (N17-N19)											
All origins	34,719	163	12	9	16	21	42	83	147	250	423
Male	16,611	87	7	6	9	8	17	34	76	143	239
Female	18,108	76	5	3	7	13	25	49	71	107	184
Hispanic	1,535	19	2	3	4	1	4	8	10	23	37
Male	790	14	2	3	3	-	1	2	5	12	17
Female	745	5	-	-	1	1	3	6	5	11	20
Mexican	838	14	1	3	2	1	2	5	5	13	17
Male	447	11	1	3	2	-	-	1	3	8	7
Female	391	3	-	-	-	1	2	4	2	5	10
Puerto Rican	227	2	-	-	-	-	1	1	4	4	9
Male	109	2	-	-	-	-	1	1	2	-	3
Female	118	-	-	-	-	-	-	-	2	4	6
Cuban	157	-	-	-	-	-	-	-	-	-	1
Male	86	-	-	-	-	-	-	-	-	-	-
Female	71	-	-	-	-	-	-	-	-	-	1
Central and South American	123	1	-	-	1	-	1	2	1	2	4
Male	56	-	-	-	1	-	-	-	-	1	3
Female	67	1	-	-	-	-	1	2	1	1	1
Other and unknown Hispanic	190	2	1	-	1	-	-	-	-	4	6
Male	92	1	1	-	-	-	-	-	-	3	4
Female	98	1	-	-	1	_	-	-	-	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Renal failure (N17-N19) Non-Hispanic 1/ 33,074 15,763 Male Female 17,311 25,831 Non-Hispanic white Male 12,496 Female 13,335 Non-Hispanic black 6,527 Male 2,920 Female 3,607 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 203 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other disorders of kidney (N25,N27) All origins 35 1 1 1 Male 11 1 Female 24 1 1 Hispanic Male 2 Female 2 Mexican Male Female Puerto Rican 1 Male 1 Female Cuban 1 Male Female 1 Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 204
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other disorders of kidney (N25,N27) Non-Hispanic 1/ 31 1 1 1 9 Male 1 Female 22 1 1 19 Non-Hispanic white 1 1 1 Male 1 Female 12 1 1 Non-Hispanic black 12 Male 2 Female 10 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 205 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 35-39 40-44 ALL 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Infections of kidney (N10-N12, N13.6, N15.1) All origins 846 3 3 5 4 13 15 1 1 1 Male 263 1 1 2 5 Female 583 2 1 1 2 1 5 4 11 10 Hispanic 1 4 3 55 Male 21 1 1 Female 34 1 3 2 1 2 Mexican 32 13 Male 1 Female 19 1 1 Puerto Rican 5 1 1 1 Male 1 1 Female 4 1 1 Cuban 5 Male 4 Female 1 Central and South American 8 1 Male 3 Female 5 1 Other and unknown Hispanic 5 1 Male Female 5 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Infections of kidney (N10-N12, N13.6, N15.1) Non-Hispanic 1/ 786 3 3 4 8 12 1 1 1 4 Male 240 1 1 4 2 Female 546 1 1 1 4 4 8 8 1 3 9 Non-Hispanic white 689 1 1 1 1 4 3 Male 209 1 1 2 Female 480 1 1 1 3 4 3 7 Non-Hispanic black 2 3 3 85 Male 29 2 Female 56 2 3 1 Origin not stated 2/ 5 1 Male 2 1 Female 3

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 207 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hyperplasia of prostate (N40)											
All origins	430	-	-	-	-	-	-	-	-	-	-
Male	430	-	-	-	-	-	-	-	-	-	-
Hispanic	11	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-
Mexican	8	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-
Cuban	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-
Central and South American	2	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 208 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hyperplasia of prostate (N40)											
Non-Hispanic 1/	417	-	-	-	-	-	-	-	-	-	-
Male	417	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white	377	-	-	-	-	-	-	-	-	-	-
Male	377	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	38	-	-	-	-	-	-	-	-	-	-
Male	38	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	2	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 209 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX YEARS AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Inflammatory diseases of female pelvic organs (N70-N76) All origins 108 2 1 1 2 3 3 4 8 Female 108 2 1 1 2 3 3 4 8 Hispanic 7 1 Female 1 Mexican 1 Female 1 Puerto Rican Female Cuban Female Central and South American Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

1

Other and unknown Hispanic

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 210 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 40-44 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YEARS YEARS Inflammatory diseases of female pelvic organs (N70-N76) Non-Hispanic 1/ 101 2 1 3 3 4 7 1 2 Female 101 1 1 3 3 4 7 1 2 Non-Hispanic white 73 1 1 1 2 Female 73 1 2 1 1 1 2 Non-Hispanic black 26 2 5 Female 26 2 5 Origin not stated 2/ Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 211 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Pregnancy, childbirth and the puerperium (000-099) All origins 406 33 64 89 92 88 33 Female 406 33 64 89 92 88 33 Hispanic 70 6 13 18 8 21 4 . . . Female 70 6 13 18 8 21 4 Mexican 40 3 6 13 12 2 3 Female 40 6 13 12 2 Puerto Rican 7 1 2 2 1 1 7 1 Female 1 Cuban 1 1 Female 1 1 . . . Central and South American 15 4 3 1 6 1 . . . Female 15 4 3 1 6 1

2

2

1

1

1

1

1

1

2

2

Source: CDC/NCHS, National Vital Statistics System, Mortality

Other and unknown Hispanic

Female

7

7

. . .

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 40-44 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 ALL YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YEARS YFARS Pregnancy, childbirth and the puerperium (000-099) Non-Hispanic 1/ 333 27 49 70 84 67 29 . . . Female 333 27 49 70 84 67 29 29 Non-Hispanic white 152 8 18 45 35 13 . . . Female 8 29 45 152 18 35 13 Non-Hispanic black 158 18 28 37 33 27 13 . . . Female 158 18 28 37 33 13 Origin not stated 2/ Female 2

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 213 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Pregnancy with abortive outcome (000-007) All origins 31 3 5 5 10 6 2 Female 31 3 5 5 10 6 2 Hispanic 2 2 Female 2 2 Mexican 2 1 1 . . . Female 2 1 1 Puerto Rican 1 1 Female Cuban Female . . . Central and South American 1 1 Female 1 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

. . .

. . .

. . .

Other and unknown Hispanic

ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 01/07/24 PLATE 1 OF 2 PAGE 214

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 40-44 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 ALL POPULATION, AND SEX YEARS AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YEARS YEARS Pregnancy with abortive outcome (000-007) Non-Hispanic 1/ 26 1 5 10 2 . . . Female 26 1 5 10 4 2 . . . Non-Hispanic white 5 11 2 . . . Female 5 11 1 2 Non-Hispanic black 14 1 3 . . . Female 14 1 3 Origin not stated 2/ Female

. . .

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other complications of preg	gnancy, ch	ildbirth an	d the pue	rperium (010-099)						
All origins	375			-	-	30	59	84	82	82	31
Female	375			-	-	30	59	84	82	82	31
Hispanic	66			-	-	4	13	18	8	19	4
Female	66			-	-	4	13	18	8	19	4
Mexican	38			-	-	2	6	13	4	11	2
Female	38			-	-	2	6	13	4	11	2
Puerto Rican	6			-	-	-	2	1	2	-	1
Female	6			-	-	-	2	1	2	-	1
Cuban	1			-	-	-	-	-	-	1	-
Female	1			-	-	-	-	-	-	1	-
Central and South American	14			-	-	-	4	3	1	5	1
Female	14			-	-	-	4	3	1	5	1
Other and unknown Hispanic	7			-	-	2	1	1	1	2	-
Female	7			_	_	2	1	1	1	2	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS Other complications of pregnancy, childbirth and the puerperium (010-099) Non-Hispanic 1/ 307 26 45 65 74 63 27 . . . Female 307 26 45 65 74 63 27 . . . Non-Hispanic white 141 8 16 28 40 34 11 . . . Female 8 28 40 141 16 34 11 Non-Hispanic black 144 17 26 33 29 24 13 . . . Female 144 17 26 33 29 24 13 Origin not stated 2/

Source: CDC/NCHS, National Vital Statistics System, Mortality

. . .

. . .

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

1 R	LASSIFICATION OF DISEASES, L992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
C	Certain conditions originatin	g in the p	perinatal p	eriod (P0	0-P96)							
A	All origins	14,259	14,084	92	25	18	11	5	2	5	2	1
	Male	8,044	7,941	56	14	9	6	4	-	2	2	-
	Female	6,215	6,143	36	11	9	5	1	2	3	-	1
Н	Hispanic	2,153	2,127	18	2	-	3	1	1	-	-	-
	Male	1,192	1,181	5	2	-	2	1	-	-	-	-
	Female	961	946	13	-	-	1	-	1	-	-	-
	Mexican	1,485	1,467	14	1	-	2	1	-	-	-	-
	Male	825	819	3	1	-	1	1	-	-	-	-
	Female	660	648	11	-	-	1	-	-	-	-	-
	Puerto Rican	238	235	1	1	-	-	-	1	-	-	-
	Male	124	122	1	1	-	-	-	-	-	-	-
	Female	114	113	-	-	-	-	-	1	-	-	-
	Cuban	40	39	1	-	-	-	-	-	-	-	-
	Male	20	20	-	-	-	-	-	-	-	-	-
	Female	20	19	1	-	-	-	-	-	-	-	-
	Central and South American	145	143	1	-	-	1	-	-	-	-	-
	Male	82	81	-	-	-	1	-	-	-	-	-
	Female	63	62	1	-	-	-	-	-	-	-	-
	Other and unknown Hispanic	245	243	1	-	-	-	-	-	-	-	-
	Male	141	139	1	-	-	-	-	-	-	-	-
	Female	104	104	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Certain conditions originating in the perinatal period (P00-P96) Non-Hispanic 1/ 11,815 11,668 73 23 18 8 1 5 2 1 Male 6,680 6,589 50 12 9 4 3 2 2 Female 5,135 5,079 23 11 9 4 1 1 3 1 6,384 17 8 3 5 2 Non-Hispanic white 6,286 37 14 1 1 Male 3,627 3,570 24 8 3 2 Female 2,757 2,716 13 4 1 3 1 Non-Hispanic black 5,000 4,963 29 Male 2,818 2,794 19 Female 2,182 2,169 10 2 Origin not stated 2/ 291 289 Male 172 171 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

119

118

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Congenital malformations, de	formations	and chromo	somal abn	ormalitie	es (Q00-Q9	99)					
All origins	10,393	5,473	549	207	221	222	212	212	253	279	273
Male	5,346	2,875	310	102	113	139	123	110	132	152	144
Female	5,047	2,598	239	105	108	83	89	102	121	127	129
Hispanic	1,498	1,072	101	31	26	27	29	28	26	15	19
Male	756	534	63	16	13	17	13	15	10	8	7
Female	742	538	38	15	13	10	16	13	16	7	12
Mexican	1,087	818	69	19	19	21	19	18	13	8	13
Male	553	409	41	9	11	11	9	8	6	5	5
Female	534	409	28	10	8	10	10	10	7	3	8
Puerto Rican	117	70	8	3	1	3	6	3	1	1	3
Male	55	31	7	3	-	3	4	1	1	-	1
Female	62	39	1	-	1	-	2	2	-	1	2
Cuban	32	12	1	1	1	-	-	-	1	1	-
Male	17	7	1	-	1	-	-	-	-	1	-
Female	15	5	-	1	-	-	-	-	1	-	-
Central and South American	117	73	11	3	2	2	1	7	3	2	1
Male	61	34	6	2	-	2	-	6	2	2	-
Female	56	39	5	1	2	-	1	1	1	-	1
Other and unknown Hispanic	145	99	12	5	3	1	3	-	8	3	2
Male	70	53	8	2	1	1	-	-	1	-	1
Female	75	46	4	3	2	-	3	-	7	3	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 40-44 ALL 1-4 5-9 10-14 20-24 35 - 39POPULATION, AND SEX YEARS YEARS AGES YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) Non-Hispanic 1/ 8,825 4,349 Male 4,550 2,311 Female 4,275 2,038 6,782 Non-Hispanic white 3,127 Male 3,498 1,693 Female 3,284 1,434 Non-Hispanic black 1,680 Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Origin not stated 2/

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Symptoms, signs and abnormal	clinical	and laborat	ory findi	ngs, not	elsewhere	classifi	ed (R00-R	99)			
All origins	26,672	3,538	236	65	72	228	381	446	649	940	1,122
Male	12,877	2,072	136	35	42	153	269	287	434	599	740
Female	13,795	1,466	100	30	30	75	112	159	215	341	382
Hispanic	1,281	415	30	6	10	30	53	58	80	80	68
Male	768	237	14	5	5	21	43	42	57	59	51
Female	513	178	16	1	5	9	10	16	23	21	17
Mexican	700	289	22	4	4	17	31	41	38	35	26
Male	444	166	11	3	1	13	25	33	29	29	20
Female	256	123	11	1	3	4	6	8	9	6	6
Puerto Rican	192	52	1	1	2	5	9	4	17	20	17
Male	111	25	-	1	2	3	7	2	13	15	12
Female	81	27	1	-	-	2	2	2	4	5	5
Cuban	94	5	-	-	-	2	1	5	1	7	9
Male	50	2	-	-	-	2	1	3	1	5	7
Female	44	3	-	-	-	-	-	2	-	2	2
Central and South American	118	16	3	1	2	5	9	6	14	12	10
Male	69	9	2	1	1	3	7	3	10	7	8
Female	49	7	1	-	1	2	2	3	4	5	2
Other and unknown Hispanic	177	53	4	-	2	1	3	2	10	6	6
Male	94	35	1	-	1	-	3	1	4	3	4
Female	83	18	3	-	1	1	-	1	6	3	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 YEARS YEARS POPULATION, AND SEX AGES YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) Non-Hispanic 1/ 25,181 3,097 1,037 Male 11,962 1,817 Female 13,219 1,280 20,200 Non-Hispanic white 1,812 Male 9,249 1,083 Female 10,951 Non-Hispanic black 4,508 1,149 Male 2,460 Female 2,048 Origin not stated 2/

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other diseases (Residual -N13.5,N13.7-N13.9,N14,N15.0						193 , K00 - K2	22,K29-K31	, K50-K66 , K	71-K72,K75	-K76,K83-M	99,N13.0
All origins	171,683	1,148	647	436	485	785	986	1,235	1,827	3,203	4,786
Male	69,784	680	354	229	254	483	562	717	1,073	1,912	2,923
Female	101,899	468	293	207	231	302	424	518	754	1,291	1,863
Hispanic	7,291	200	106	50	57	95	123	166	243	306	384
Male	3,655	124	50	28	26	59	69	103	172	218	261
Female	3,636	76	56	22	31	36	54	63	71	88	123
Mexican	4,087	137	80	34	42	63	83	109	147	196	250
Male	2,105	79	38	20	18	40	47	64	106	140	167
Female	1,982	58	42	14	24	23	36	45	41	56	83
Puerto Rican	983	18	7	6	6	7	18	26	43	44	52
Male	541	14	5	2	3	2	13	17	26	31	37
Female	442	4	2	4	3	5	5	9	17	13	15
Cuban	684	4	-	1	-	3	4	1	5	8	12
Male	267	1	-	1	-	2	1	-	4	5	6
Female	417	3	-	-	-	1	3	1	1	3	6
Central and South American	545	22	7	6	3	9	11	15	22	25	27
Male	262	15	3	3	2	6	4	12	18	16	18
Female	283	7	4	3	1	3	7	3	4	9	9
Other and unknown Hispanic	992	19	12	3	6	13	7	15	26	33	43
Male	480	15	4	2	3	9	4	10	18	26	33

3

1

4

3

8

7

10

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

512

4

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX		UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other diseases (Residu -N13.5,N13.7-N13.9,N14,N15						193,K00-K2	2,K29-K31	, K50 - K66 , K	71-K72,K75-	-K76,K83-M	99,N13.0
Non-Hispanic 1/	163,832	935	537	386	428	688	859	1,068	1,577	2,878	4,369
Male	65,835	546	302	201	228	423	490	613	896	1,679	2,637
Female	97,997	389	235	185	200	265	369	455	681	1,199	1,732
Non-Hispanic white	141,002	554	375	265	304	513	587	749	1,096	1,949	3,087
Male	55,517	321	214	136	155	332	349	456	654	1,194	1,950
Female	85,485	233	161	129	149	181	238	293	442	755	1,137
Non-Hispanic black	20,137	332	129	100	101	144	241	282	422	839	1,158
Male	8,967	192	67	56	62	75	127	135	209	434	606
Female	11,170	140	62	44	39	69	114	147	213	405	552
Origin not stated 2/	560	13	4	-	-	2	4	1	7	19	33
Male	294	10	2	-	-	1	3	1	5	15	25
Female	266	3	2	-	-	1	1	-	2	4	8

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 30 - 3435 - 39POPULATION, AND SEX YEARS YEARS AGES YFAR YFARS YFARS YFARS YFARS YFARS YFARS YFARS Accidents (unintentional injuries) (V01-X59, Y85-Y86) All origins 1,898 97,860 1.459 6.688 6.968 5.881 6.009 7.366 7.865 1.632 Male 63.535 1.127 1.077 4.624 5.416 4.542 5.411 4.555 5.837 Female 34.325 2.064 1.552 1.326 1.467 1.955 2.028 Hispanic 8,650 1,146 Male 6,578 Female 2,072 Mexican 5,689 Male 4,396 Female 1,293 Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 40-44 ALL 1-4 5-9 10-14 20-24 35 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Accidents (unintentional injuries) (V01-X59,Y85-Y86) Non-Hispanic 1/ 88,669 1,550 1,286 1,435 5,865 5,793 4,913 5,121 6,481 7,048 710 Male 56,551 414 926 787 952 4,003 4,425 3,745 3,805 4,687 5,184 Female 32,118 296 624 499 483 1,862 1,368 1,168 1,316 1,794 1,864 Non-Hispanic white 73,326 435 1,042 843 1,065 4,840 4,570 3,854 4,001 5,186 5,658 Male 46,113 258 617 505 700 3,250 3,499 2,953 2,979 3,770 4,179 Female 27,213 177 425 338 365 1,590 1,071 901 1,022 1,416 1,479 Non-Hispanic black 428 392 764 841 897 1,063 1,182 12,533 251 309 930 Male 8,629 140 260 254 213 568 713 647 666 754 874 Female 3,904 168 138 96 196 217 194 231 309 308 Origin not stated 2/ 541 1 7 3 5 23 29 32 34 41 54 Male 406 4 3 5 13 24 23 30 32 38 Female 135 1 3 10 5 9 4 9 16

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX YFAR YEARS YEARS YFARS YFARS YFARS AGES YFARS YFARS YFARS YFARS Transport accidents (V01-V99,Y85) All origins 46,423 1.078 5.452 5.237 3.922 3.512 3.869 3.684 Male 31.642 3.593 3.965 2.952 2.559 2.741 2.635 Female 14.781 1,859 1.272 1.128 1.049 Hispanic 4,880 Male 3,630 Female 1,250 Mexican 3,432 Male 2,600 Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 40-44 ALL 1-4 5-9 10-14 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Transport accidents (V01-V99,Y85) Non-Hispanic 1/ 41,330 149 553 786 933 4,820 4,372 3,260 3,012 3,461 3,305 Male 27,850 79 313 464 585 3,133 3,263 2,415 2,163 2,422 2,342 Female 13,480 70 240 322 348 1,687 1,109 845 849 1,039 963 Non-Hispanic white 33,803 98 349 557 734 4,044 3,449 2,528 2,361 2,737 2,688 Male 22,682 58 192 325 457 2,595 2,586 1,876 1,685 1,917 1,910 Female 11,121 40 157 232 277 1,449 863 652 676 820 778 Non-Hispanic black 5,810 199 571 691 569 506 583 491 44 168 161 Male 4,097 18 96 124 105 400 509 436 381 409 360 Female 1,713 26 72 75 171 182 133 125 174 131 Origin not stated 2/ 213 2 3 3 17 23 22 15 16 13 Male 162 2 3 3 8 18 15 14 12 10 Female 51 9 5 7 1 4 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES,

Male

Female

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Motor vehicle accidents (V87.0-V87.8,V88.0-V88.8,V89.0		09.0,V09.2,	V12-V14,V	19.0-V19.	2,V19.4-V	/19.6,V20-	V79,V80.3	-V80.5,V81	.0-V81.1,V	82.0-V82.1	V83-V86,
All origins	42,401	184	650	802	969	5,198	4,930	3,604	3,174	3,460	3,278
Male	28,552	97	366	469	593	3,412	3,715	2,690	2,303	2,424	2,311
Female	13,849	87	284	333	376	1,786	1,215	914	871	1,036	967
Hispanic	4,525	40	148	100	131	587	797	593	442	354	330
Male	3,337	22	82	59	77	431	646	480	348	275	255
Female	1,188	18	66	41	54	156	151	113	94	79	75
Mexican	3,205	37	115	72	91	435	617	430	319	239	224
Male	2,408	21	66	43	55	324	509	345	251	189	178
Female	797	16	49	29	36	111	108	85	68	50	46
Puerto Rican	289	1	9	7	10	46	38	30	25	24	24
Male	205	-	4	5	9	29	24	26	21	20	17
Female	84	1	5	2	1	17	14	4	4	4	7
Cuban	159	-	3	1	3	14	14	10	9	14	9
Male	110	-	1	-	2	11	13	9	6	10	9
Female	49	-	2	1	1	3	1	1	3	4	-
Central and South American	471	2	7	10	8	44	76	92	53	45	34
Male	341	1	3	6	4	29	61	79	43	33	22
Female	130	1	4	4	4	15	15	13	10	12	12
Other and unknown Hispanic	401	-	14	10	19	48	52	31	36	32	39

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE

ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
	99.0,V09.2,	V12-V14,V	19.0-V19.	2,V19.4-V	19.6,V20-	V79,V80.3	-V80.5,V81	.0-V81.1,V	82.0-V82.1	V83-V86,
37,684	144	500	699	835	4,594	4,113	2,992	2,720	3,090	2,936
25,072	75	282	407	513	2,973	3,054	2,198	1,943	2,137	2,047
12,612	69	218	292	322	1,621	1,059	794	777	953	889
30,743	94	315	491	653	3,860	3,239	2,301	2,124	2,426	2,371
20,362	55	175	284	399	2,467	2,417	1,693	1,508	1,678	1,660
10,381	39	140	207	254	1,393	822	608	616	748	711
5,388	43	152	182	149	544	665	540	466	534	451
3,763	17	84	109	94	379	490	412	349	371	325
1,625	26	68	73	55	165	175	128	117	163	126
192	-	2	3	3	17	20	19	12	16	12
143	-	2	3	3	8	15	12	12	12	9
49	-	-	-	-	9	5	7	-	4	3
	AGES (V02-V04, V0), V89.2) 37,684 25,072 12,612 30,743 20,362 10,381 5,388 3,763 1,625 192 143	AGES YEAR (V02-V04, V09.0, V09.2, 7) (V089.2) 37,684 144 25,072 75 12,612 69 30,743 94 20,362 55 10,381 39 5,388 43 3,763 17 1,625 26 192 - 143 -	AGES YEAR YEARS (V02-V04, V09.0, V09.2, V12-V14, V0, V89.2) 37,684 144 500 25,072 75 282 12,612 69 218 30,743 94 315 20,362 55 175 10,381 39 140 5,388 43 152 3,763 17 84 1,625 26 68 192 - 2 143 - 2	AGES YEAR YEARS YEARS (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.0, V89.2) 37,684 144 500 699 25,072 75 282 407 12,612 69 218 292 30,743 94 315 491 20,362 55 175 284 10,381 39 140 207 5,388 43 152 182 3,763 17 84 109 1,625 26 68 73 192 - 2 3 143 - 2 3	AGES YEAR YEARS YEARS YEARS YEARS (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V0, V89.2) 37,684 144 500 699 835 25,072 75 282 407 513 12,612 69 218 292 322 30,743 94 315 491 653 20,362 55 175 284 399 10,381 39 140 207 254 5,388 43 152 182 149 3,763 17 84 109 94 1,625 26 68 73 55 192 - 2 3 3 143 - 2 3 3	AGES YEAR YEARS YEARS YEARS YEARS YEARS (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20- 0, V89.2) 37,684 144 500 699 835 4,594 25,072 75 282 407 513 2,973 12,612 69 218 292 322 1,621 30,743 94 315 491 653 3,860 20,362 55 175 284 399 2,467 10,381 39 140 207 254 1,393 5,388 43 152 182 149 544 3,763 17 84 109 94 379 1,625 26 68 73 55 165 192 - 2 3 3 17 143 - 2 3 3 3 8	AGES YEAR YEARS YEARS YEARS YEARS YEARS YEARS YEARS (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3 (V89.2) 37,684 144 500 699 835 4,594 4,113 25,072 75 282 407 513 2,973 3,054 12,612 69 218 292 322 1,621 1,059 30,743 94 315 491 653 3,860 3,239 20,362 55 175 284 399 2,467 2,417 10,381 39 140 207 254 1,393 822 5,388 43 152 182 149 544 665 3,763 17 84 109 94 379 490 1,625 26 68 73 55 165 175 192 - 2 3 3 3 17 20 143 - 2 2 3 3 3 17 20 143 - 2 3 3 3 15	AGES YEAR YEARS (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0, V89.2) 37,684 144 500 699 835 4,594 4,113 2,992 25,072 75 282 407 513 2,973 3,054 2,198 12,612 69 218 292 322 1,621 1,059 794 30,743 94 315 491 653 3,860 3,239 2,301 20,362 55 175 284 399 2,467 2,417 1,693 10,381 39 140 207 254 1,393 822 608 5,388 43 152 182 149 544 665 540 3,763 17 84 109 94 379 490 412 1,625 26 68 73 55 165 175 128 192 - 2 3 3 3 17 20 19 143 - 2 2 3 3 3 17 20 19 143	AGES YEAR YEARS YE	AGES YEAR YEARS (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1 V0, V89.2) 37,684 144 500 699 835 4,594 4,113 2,992 2,720 3,090 25,072 75 282 407 513 2,973 3,054 2,198 1,943 2,137 12,612 69 218 292 322 1,621 1,059 794 777 953 30,743 94 315 491 653 3,860 3,239 2,301 2,124 2,426 20,362 55 175 284 399 2,467 2,417 1,693 1,508 1,678 10,381 39 140 207 254 1,393 822 608 616 748 5,388 43 152 182 149 544 665 540 466 534 3,763 17 84 109 94 379 490 412 349 371 1,625 26 68 73 55 165 175 128 117 163 192 - 2 3 3 3 17 20 19 12 16 143 - 2 2 3 3 3 17 20 19 12 16

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other land transport acci 1.9,V82.2-V82.9,V87.9,V88.9,V			'09.1,V09.	3-V09.9,V	710-V11,V1	.5-V18,V19	9.3,V19.8-	V19.9,V80.	9-V80.2,V8	0.6-V80.9, [\]	V81.2-V8
All origins	2,118	4	48	62	66	172	190	152	165	205	203
Male	1,559	4	26	43	48	119	160	126	118	157	162
Female	559	-	22	19	18	53	30	26	47	48	41
Hispanic	264	-	8	6	8	20	35	29	31	28	29
Male	220	-	3	5	7	16	29	28	25	24	23
Female	44	-	5	1	1	4	6	1	6	4	6
Mexican	175	-	4	4	4	14	30	19	23	13	21
Male	148	-	2	4	3	12	25	19	18	11	16
Female	27	-	2	-	1	2	5	-	5	2	5
Puerto Rican	18	-	-	1	1	1	1	2	3	2	3
Male	16	-	-	-	1	1	1	2	3	2	3
Female	2	-	-	1	-	-	-	-	-	-	-
Cuban	17	-	-	-	-	-	-	1	2	3	-
Male	14	-	-	-	-	-	-	1	1	3	-
Female	3	-	-	-	-	-	-	-	1	-	-
Central and South American	34	-	1	1	2	2	2	5	3	6	2
Male	28	-	1	1	2	1	2	4	3	5	1
Female	6	-	-	-	-	1	-	1	-	1	1
Other and unknown Hispanic	20	-	3	-	1	3	2	2	-	4	3
Male	14	-	-	-	1	2	1	2	-	3	3
Female	6	-	3	_	_	1	1	_	_	1	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS			
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V1.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)														
Non-Hispanic 1/	1,836	4	40	56	58	152	152	121	131	177	173			
Male	1,323	4	23	38	41	103	128	96	91	133	138			
Female	513	-	17	18	17	49	24	25	40	44	35			
Non-Hispanic white	1,480	3	25	39	47	122	117	96	99	141	141			
Male	1,051	3	12	25	32	81	96	75	68	106	112			
Female	429	-	13	14	15	41	21	21	31	35	29			
Non-Hispanic black	248	1	12	14	6	18	15	18	24	29	23			
Male	192	1	9	13	6	14	13	16	18	22	19			
Female	56	-	3	1	-	4	2	2	6	7	4			
Origin not stated 2/	18	-	-	-	-	-	3	2	3	-	1			
Male	16	-	-	-	-	-	3	2	2	-	1			
Female	2	-	-	-	-	-	-	-	1	-	-			

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Water, air and space, and	d other and	d unspecifio	ed transp	ort accid	ents and	their seq	µuelae (V9	0-V99,Y85)			
All origins	1,904	1	14	33	43	82	117	166	173	204	203
Male	1,531	-	8	20	33	62	90	136	138	160	162
Female	373	1	6	13	10	20	27	30	35	44	41
Hispanic	91	-	1	2	3	8	10	18	12	10	7
Male	73	-	-	1	2	5	9	14	9	8	5
Female	18	-	1	1	1	3	1	4	3	2	2
Mexican	52	-	1	2	1	3	8	10	8	7	2
Male	44	-	-	1	1	2	8	9	6	6	1
Female	8	-	1	1	-	1	-	1	2	1	1
Puerto Rican	9	-	-	-	1	2	-	1	1	1	2
Male	7	-	-	-	1	1	-	1	1	1	1
Female	2	-	-	-	-	1	-	-	-	-	1
Cuban	6	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-
Central and South American	14	-	-	-	-	2	1	3	3	1	1
Male	10	-	-	-	-	1	1	2	2	-	1
Female	4	-	-	-	-	1	-	1	1	1	-
Other and unknown Hispanic	10	-	-	-	1	1	1	4	-	1	1
Male	6	-	-	-	-	1	-	2	-	1	1
Female	4	-	-	-	1	-	1	2	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 30-34 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 25-29 35 - 39YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85) Non-Hispanic 1/ 1,810 Male 1,455 Female 1,580 Non-Hispanic white Male 1,269 Female Non-Hispanic black Male Female Origin not stated 2/

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Nontransport accidents (W00-X59,Y86) All origins 51,437 1,186 3,497 1.236 1.731 1.959 2.497 4.181 Male 31.893 1,031 1.451 1.983 2.670 3.202 1.603 Female 19.544 Hispanic 3,770 Male 2,948 Female Mexican 2,257 Male 1,796 Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Nontransport accidents (W00-X59,Y86) Non-Hispanic 1/ 47,339 561 997 500 502 1,045 1,421 1,653 2,109 3,020 3,743 28,701 Male 335 613 323 367 870 1,162 1,330 1,642 2,265 2,842 Female 18,638 226 384 177 135 175 259 323 467 755 901 39,523 796 2,970 Non-Hispanic white 337 693 286 331 1,121 1,326 1,640 2,449 Male 23,431 200 425 180 243 655 913 1,077 1,294 1,853 2,269 Female 16,092 137 268 106 88 141 208 249 346 596 701 Non-Hispanic black 6,723 207 260 193 148 193 239 272 391 480 691 Male 4,532 122 164 130 108 168 204 211 285 345 514 Female 2,191 85 40 25 35 61 106 135 177 Origin not stated 2/ 328 1 5 2 6 6 10 19 25 41 Male 244 2 2 5 6 8 16 20 28 Female 84 1 3 1 2 3 5 13

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Falls (W00-W19)											
All origins	13,162	12	55	25	28	112	130	156	187	276	369
Male	7,109	8	37	19	21	95	114	137	157	225	303
Female	6,053	4	18	6	7	17	16	19	30	51	66
Hispanic	652	4	9	1	2	13	32	35	33	33	37
Male	464	4	6	1	2	12	31	34	31	30	32
Female	188	-	3	-	-	1	1	1	2	3	5
Mexican	362	4	7	-	1	9	24	25	18	18	25
Male	270	4	4	-	1	9	23	24	17	16	23
Female	92	-	3	-	-	-	1	1	1	2	2
Puerto Rican	62	-	-	1	-	1	-	1	1	5	5
Male	40	-	-	1	-	1	-	1	1	4	4
Female	22	-	-	-	-	-	-	-	-	1	1
Cuban	53	-	-	-	-	-	1	-	-	-	-
Male	32	-	-	-	-	-	1	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-
Central and South American	77	-	1	-	1	1	4	8	10	6	3
Male	56	-	1	-	1	-	4	8	9	6	3
Female	21	-	-	-	-	1	-	-	1	-	-
Other and unknown Hispanic	98	-	1	-	-	2	3	1	4	4	4
Male	66	-	1	-	-	2	3	1	4	4	2
Female	32	-	-	_	-	_	-	-	-	-	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Falls (W00-W19) Non-Hispanic 1/ 12,447 Male 6,602 Female 5,845 Non-Hispanic white 11,295 Male 5,860 Female 5,435 Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 40-44 ALL 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Accidental discharge of firearms (W32-W34) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX YEARS AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Accidental discharge of firearms (W32-W34) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Accidental drowning and submersion (W65-W74) All origins 3,529 Male 2,794 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Accidental drowning and submersion (W65-W74) Non-Hispanic 1/ 3,029 2,370 Male Female 2,230 Non-Hispanic white Male 1,708 Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Accidental exposure to smoke, fire and flames (X00-X09) All origins 3,348 Male 2,025 Female 1,323 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Other and unknown Hispanic

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Accidental exposure to smoke, fire and flames (X00-X09) Non-Hispanic 1/ 3,133 Male 1,879 Female 1,254 2,172 Non-Hispanic white Male 1,325 Female Non-Hispanic black Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Origin not stated 2/

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Accidental poisoning and exposure to noxious substances (X40-X49) All origins 12,186 1.403 2.082 2.467 Male 8.887 1.071 1.523 1.826 Female 3.299 Hispanic 1,502 Male 1,271 Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

Male

Female

Other and unknown Hispanic

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 YEARS YEARS POPULATION, AND SEX AGES YFAR YFARS YFARS YFARS YFARS YFARS YFARS YFARS Accidental poisoning and exposure to noxious substances (X40-X49) Non-Hispanic 1/ 10,542 8 31 10 23 219 595 790 1,165 1,753 2,154 Male 7,510 7 14 5 15 167 460 618 864 1,248 1,569 Female 3,032 1 17 5 8 52 135 172 301 505 585 6 190 Non-Hispanic white 8,311 17 3 18 504 663 918 1,416 1,680 Male 5,921 6 1 12 145 391 523 694 1,023 1,242 Female 2,390 1 11 6 45 113 140 224 393 438 Non-Hispanic black 2,008 12 2 18 75 112 216 295 434 Male 1,450 1 7 2 15 85 146 199 304 Female 558 3 14 27 70 96 130 Origin not stated 2/ 142 1 4 7 10 18 39 Male 106 1 4 6 8 13 26

1

2

5

13

Source: CDC/NCHS, National Vital Statistics System, Mortality

36

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other and unspecified	nontransport	accidents	and their	sequelae	(W20-W31	L,W35-W64,	W75-W99,X	10-X39,X50-	-X59,Y86)		
All origins	18,388	523	293	144	172	291	371	421	466	622	830
Male	10,371	314	183	93	128	238	327	361	387	513	668
Female	8,017	209	110	51	44	53	44	60	79	109	162
Hispanic	878	65	53	18	13	35	63	54	47	68	52
Male	620	35	37	12	7	28	60	51	39	60	41
Female	258	30	16	6	6	7	3	3	8	8	11
Mexican	548	46	38	11	8	26	48	41	30	49	37
Male	398	24	29	8	4	21	45	38	24	44	30
Female	150	22	9	3	4	5	3	3	6	5	7
Puerto Rican	88	11	4	1	2	5	2	3	4	7	3
Male	61	7	2	-	2	3	2	3	4	6	2
Female	27	4	2	1	-	2	-	-	-	1	1
Cuban	49	-	-	1	-	-	-	1	2	1	2
Male	31	-	-	1	-	-	-	1	2	1	2
Female	18	-	-	-	-	-	-	-	-	-	-
Central and South America	n 82	5	6	2	1	3	6	9	9	5	5
Male	59	2	3	1	-	3	6	9	7	5	4
Female	23	3	3	1	1	-	-	-	2	-	1
Other and unknown Hispani	c 111	3	5	3	2	1	7	-	2	6	5
Male	71	2	3	2	1	1	7	-	2	4	3
Female	40	1	2	1	1	-	-	-	-	2	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 25-29 30-34 35 - 39YEARS YEARS POPULATION, AND SEX AGES YFAR YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86) Non-Hispanic 1/ 17,430 Male 9,694 Female 7,736 14,930 Non-Hispanic white Male 8,122 Female 6,808 Non-Hispanic black 2,183 Male 1,365

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

Male

Female

Origin not stated 2/

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Intentional self-harm (suicide) (X60-X84,Y87.0) All origins 29.199 1.615 2.286 2.398 2.708 3.286 3.180 Male 23.458 1.347 1.979 2.005 2.189 2.590 2.429 Female 5,741 . . . Hispanic 1,695 . . . Male 1,430 . . . Female . . . Mexican . . . Male . . . Female . . . Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 15-19 20-24 25-29 30-34 35-39 40-44 ALL 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Intentional self-harm (suicide) (X60-X84,Y87.0) Non-Hispanic 1/ 27,379 1 211 1,434 2,066 2,167 2,498 3,048 3,020 . . . 21,929 Male 1 170 1,198 1,785 1,809 2,007 2,395 2,294 Female 5,450 41 236 281 358 491 653 726 2,770 Non-Hispanic white 24,534 1 173 1,169 1,661 1,776 2,151 2,699 Male 19,620 1 140 978 1,438 1,478 1,727 2,098 2,112 Female 4,914 33 191 223 298 424 601 658 Non-Hispanic black 1,924 29 176 287 258 239 244 186 Male 1,630 22 152 229 203 215 138 . . . Female 294 7 31 29 29 48 . . . Origin not stated 2/ 125 1 1 6 4 11 10 19 . . . 7 Male 99 1 1 3 2 9 5 16 . . . Female 26 3 2 2 2 3 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Intentional self-harm (suic	ide) by d	ischarge of	firearms	(X72-X74)						
All origins	16,599		-	-	103	975	1,340	1,292	1,329	1,617	1,480
Male	14,479		-	-	80	867	1,216	1,137	1,133	1,349	1,237
Female	2,120		-	-	23	108	124	155	196	268	243
Hispanic	794		-	-	13	94	125	104	78	91	57
Male	705		-	-	7	81	115	92	72	76	51
Female	89		-	-	6	13	10	12	6	15	6
Mexican	467		-	-	10	60	75	71	52	54	34
Male	425		-	-	4	53	71	65	49	47	31
Female	42		-	-	6	7	4	6	3	7	3
Puerto Rican	64		-	-	-	6	10	10	9	7	2
Male	58		-	-	-	5	9	9	8	6	2
Female	6		-	-	-	1	1	1	1	1	-
Cuban	64		-	-	-	1	2	4	2	5	5
Male	54		-	-	-	-	2	3	2	4	4
Female	10		-	-	-	1	-	1	-	1	1
Central and South American	74		-	-	-	9	16	8	7	9	7
Male	63		-	-	-	8	14	7	6	6	6
Female	11		-	-	-	1	2	1	1	3	1
Other and unknown Hispanic	125		-	-	3	18	22	11	8	16	9
Male	105		-	-	3	15	19	8	7	13	8
Female	20		-	-	-	3	3	3	1	3	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Intentional self-harm (suicide) by discharge of firearms (X72-X74) Non-Hispanic 1/ 15,755 90 879 1,213 1,184 1,247 1,520 1,417 . . . Male 13,732 73 784 1,100 1,043 1,058 1,268 1,180 Female 2,023 17 95 141 189 252 237 113 726 977 1,094 1,357 Non-Hispanic white 14,316 81 958 1,319 . . . Male 12,441 65 645 869 854 922 1,120 1,093 Female 1,875 16 81 89 123 172 237 226 Non-Hispanic black 118 205 153 115 123 74 1,112 Male 1,007 4 107 187 142 105 112 66 . . . Female 105 11 18 11 10 11 8 . . . Origin not stated 2/ 50 2 2 4 4 6 6 . . . Male 42 2 1 2 3 5 6 . . .

1

2

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

8

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Intentional self-harm (suic	cide) by o	ther and un	specified	means and	d their s	equelae ((X60-X71,X	75-X84,Y87	.0)		
All origins	12,600		-	2	139	640	946	1,106	1,379	1,669	1,700
Male	8,979		-	2	112	480	763	868	1,056	1,241	1,192
Female	3,621		-	-	27	160	183	238	323	428	508
Hispanic	901		-	-	17	81	91	116	125	137	84
Male	725		-	-	14	65	77	95	105	112	68
Female	176		-	-	3	16	14	21	20	25	16
Mexican	456		-	-	13	45	57	67	69	74	43
Male	378		-	-	12	40	50	57	58	63	32
Female	78		-	-	1	5	7	10	11	11	11
Puerto Rican	132		-	-	1	8	12	16	13	22	15
Male	105		-	-	-	6	9	14	11	18	11
Female	27		-	-	1	2	3	2	2	4	4
Cuban	78		-	-	-	2	2	5	4	7	5
Male	64		-	-	-	-	1	4	4	5	5
Female	14		-	-	-	2	1	1	-	2	-
Central and South American	104		-	-	2	9	9	12	14	20	10
Male	79		-	-	2	7	8	9	11	15	9
Female	25		-	-	-	2	1	3	3	5	1
Other and unknown Hispanic	131		-	-	1	17	11	16	25	14	11
Male	99		-	-	-	12	9	11	21	11	11

1

5

2

4

3

Source: CDC/NCHS, National Vital Statistics System, Mortality

32

Female

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATI CLASSIFICATION OF DISEAS 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX		UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Intentional self-harm	(suicide) by c	other and ur	ıspecified	means and	d their	sequelae	(X60-X71,X	75-X84,Y87	.0)		
Non-Hispanic 1/	11,624		-	1	121	555	853	983	1,251	1,528	1,603
Male	8,197		-	1	97	414	685	766	949	1,127	1,114
Female	3,427		-	-	24	141	168	217	302	401	489
Non-Hispanic white	10,218		-	1	92	443	703	799	1,057	1,342	1,451
Male	7,179		-	1	75	333	569	624	805	978	1,019
Female	3,039		-	-	17	110	134	175	252	364	432
Non-Hispanic black	812		-	-	24	58	82	105	124	121	112
Male	623		-	-	18	45	69	87	98	103	72
Female	189		-	-	6	13	13	18	26	18	40
Origin not stated 2/	75		-	1	1	4	2	7	3	4	13
Male	57		-	1	1	1	1	7	2	2	10
Female	18		-	-	-	3	1	-	1	2	3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX YFAR YEARS YEARS YFARS YFARS YFARS AGES YFARS YFARS YFARS YFARS Assault (homicide) (X85-Y09, Y87.1) All origins 2,093 16.889 2.905 2.362 1.869 1.751 1.455 Male 12,785 1.748 2.443 1.922 1.424 1.257 1.050 Female 4.104 Hispanic 2,864 Male 2,399 Female Mexican 1,916 Male 1,636 Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 20-24 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Assault (homicide) (X85-Y09, Y87.1) Non-Hispanic 1/ 13,873 270 323 155 186 1,601 2,259 1,870 1,532 1,487 1,256 Male 10,275 158 168 79 112 1,316 1,872 1,484 1,140 1,043 887 Female 3,598 112 155 76 74 285 387 386 392 444 369 409 Non-Hispanic white 5,748 143 156 78 98 553 531 604 696 631 Male 3,747 94 83 41 52 283 361 360 393 455 426 Female 2,001 49 73 37 46 126 192 171 211 241 205 Non-Hispanic black 7,536 80 1,117 1,262 860 730 575 114 151 65 1,635 Male 6,131 58 74 969 1,464 1,070 701 548 432 34 55 Female 1,405 77 25 148 171 192 159 182 143 Origin not stated 2/ 152 8 1 2 10 9 10 18 17 14 Male 111 3 1 7 6 10 14 13 10 Female 41 5 2 3 3 4 4 4

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Assault (homicide) by discharge of firearms (X93-X95) All origins 1,708 10,828 2.330 1.795 1.253 1.054 Male 8.944 1.498 2.071 1.041 1.568 Female 1.884 Hispanic 1,939 Male 1,709 Female Mexican 1,328 Male 1,183 Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Assault (homicide) by discharge of firearms (X93-X95) Non-Hispanic 1/ 8,821 1,288 1,827 1,441 1,035 Male 7,177 1,118 1,616 1,239 Female 1,644 2,995 Non-Hispanic white Male 2,119 Female Non-Hispanic black 5,508 1,447 1,053 Male 4,826 1,334 Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Assault (homicide) by other	r and unsp	ecified mea	nns and th	eir seque	lae (X85-	X92,X96-Y	(09,Y87.1)				
All origins	6,061	323	326	125	83	385	575	567	616	697	641
Male	3,841	182	168	58	37	250	372	354	383	450	443
Female	2,220	141	158	67	46	135	203	213	233	247	198
Hispanic	925	51	47	18	19	72	139	138	113	99	75
Male	690	24	22	10	10	52	114	109	90	72	64
Female	235	27	25	8	9	20	25	29	23	27	11
Mexican	588	40	31	14	10	49	93	99	77	49	36
Male	453	19	15	7	7	37	77	83	62	42	30
Female	135	21	16	7	3	12	16	16	15	7	6
Puerto Rican	103	4	6	1	6	4	11	13	13	16	9
Male	63	1	3	-	2	3	10	10	7	5	7
Female	40	3	3	1	4	1	1	3	6	11	2
Cuban	27	1	1	1	-	-	-	3	4	4	3
Male	23	1	-	1	-	-	-	2	3	3	3
Female	4	-	1	-	-	-	-	1	1	1	-
Central and South American	110	1	3	1	1	12	22	15	9	20	14
Male	84	1	-	1	1	8	18	8	9	17	13
Female	26	-	3	-	-	4	4	7	-	3	1
Other and unknown Hispanic	97	5	6	1	2	7	13	8	10	10	13
Male	67	2	4	1	-	4	9	6	9	5	11
Female	30	3	2	-	2	3	4	2	1	5	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES,

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Assault (homicide) by other	r and unspe	ecified mea	ns and th	eir seque	lae (X85-	X92, X96-Y	09,Y87.1)				
Non-Hispanic 1/	5,052	264	278	105	64	313	432	429	497	589	557
Male	3,098	155	145	48	27	198	256	245	288	372	374
Female	1,954	109	133	57	37	115	176	184	209	217	183
Non-Hispanic white	2,753	141	135	55	41	153	218	195	246	291	287
Male	1,628	92	72	26	17	86	115	107	131	178	189
Female	1,125	49	63	29	24	67	103	88	115	113	98
Non-Hispanic black	2,028	112	133	43	21	138	188	209	218	267	242
Male	1,305	57	67	20	9	95	130	123	136	174	170
Female	723	55	66	23	12	43	58	86	82	93	72
Origin not stated 2/	84	8	1	2	-	-	4	-	6	9	9
Male	53	3	1	-	-	-	2	-	5	6	5
Female	31	5	-	2	-	-	2	-	1	3	4

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 261 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 25-29 30-34 40-44 ALL 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Legal intervention (Y35,Y89.0) 398 1 47 66 70 72 49 All origins 20 Male 392 1 20 47 69 69 48 66 Female 6 1 3 1 Hispanic 58 5 8 10 13 10 4 5 8 Male 58 10 13 10 Female 44 5 7 9 6 4 Mexican 5 9 6 4 Male Female Puerto Rican 3 1 2 Male 3 1 2 Female Cuban Male Female Central and South American 2 4 1 Male 4 2 1 Female Other and unknown Hispanic 7 1 3 2 2 Male 1 3 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 YEARS POPULATION, AND SEX AGES YEAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Legal intervention (Y35,Y89.0) Non-Hispanic 1/ 338 1 15 39 55 57 61 45 Male 332 1 15 39 55 56 58 44 Female 6 1 3 1 197 5 28 29 Non-Hispanic white 15 34 36 Male 192 5 15 28 34 33 28 Female 5 3 1 Non-Hispanic black 125 10 21 24 18 21 15 Male 124 10 21 24 17 21 15 Female Origin not stated 2/ 2 1 1 Male 2 1 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 25-29 40-44 ALL 1-4 5-9 10-14 20-24 30 - 3435 - 39POPULATION, AND SEX AGES YEARS YEARS YFARS YFARS YFARS YFAR YFARS YFARS YFARS YFARS Events of undetermined intent (Y10-Y34, Y87.2, Y89.9) All origins 3,917 Male 2,692 Female 1,225 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 5-9 10-14 YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Events of undetermined intent (Y10-Y34, Y87.2, Y89.9) Non-Hispanic 1/ 3,630 Male 2,463 Female 1,167 2,848 Non-Hispanic white Male 1,887 Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 25-29 30-34 40-44 ALL 1-4 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Discharge of firearms, undetermined intent (Y22-Y24) All origins 324 3 59 32 19 28 29 12 68 Male 277 3 12 61 50 28 16 24 21 Female 47 7 9 4 3 4 8 Hispanic 31 1 1 9 9 2 1 2 1 2 Male 26 1 1 8 8 1 1 Female 5 1 1 1 1 7 2 2 Mexican 22 1 1 18 6 6 2 Male 1 Female 4 1 Puerto Rican Male Female Cuban Male

1

1

1

1

1

1

1

1

1

1

1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

3

3

6

5

1

CAUSE OF DEATH (BASED ON THE

Female

Male

Male

Female

Female

Central and South American

Other and unknown Hispanic

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Discharge of firearms, undetermined intent (Y22-Y24) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Origin not stated 2/
Male
Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES	UNDER 1 YEAR	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25 - 29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other and unspecified event	s of unde	termined in	tent and	their seq	uelae (Y1	L0-Y21,Y25	5-Y34,Y87.	2,Y89.9)			
All origins	3,593	55	40	12	25	91	212	281	411	616	622
Male	2,415	33	18	6	20	71	163	202	277	422	420
Female	1,178	22	22	6	5	20	49	79	134	194	202
Hispanic	218	6	8	1	2	11	23	30	36	34	24
Male	174	3	5	1	1	9	21	25	27	27	20
Female	44	3	3	-	1	2	2	5	9	7	4
Mexican	110	5	6	1	2	7	17	14	11	14	9
Male	87	3	4	1	1	7	15	11	8	11	8
Female	23	2	2	-	1	-	2	3	3	3	1
Puerto Rican	49	-	-	-	-	2	2	7	13	9	8
Male	39	-	-	-	-	1	2	7	9	6	7
Female	10	-	-	-	-	1	-	-	4	3	1
Cuban	4	-	-	-	-	-	-	-	2	-	1
Male	3	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	1	-	-
Central and South American	21	1	1	-	-	1	2	4	4	2	1
Male	17	-	-	-	-	1	2	3	4	2	1
Female	4	1	1	-	-	-	-	1	-	-	-
Other and unknown Hispanic	34	-	1	-	-	1	2	5	6	9	5
Male	28	-	1	-	-	-	2	4	5	8	3
Female	6	-	-	-	-	1	-	1	1	1	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 25-29 30-34 40-44 ALL 1-4 5-9 10-14 15 - 1920-24 35 - 39YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9) Non-Hispanic 1/ 3,337 Male 2,212 Female 1,125 2,631 Non-Hispanic white Male 1,700 Female Non-Hispanic black Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Origin not stated 2/

Male

Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 269 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Operations of war and their sequelae (Y36,Y89.1) All origins 23 1 Male 22 1 Female 1 Hispanic 2 Male 2 Female Mexican 1 Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic 1 Male 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 270 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 ALL YEARS POPULATION, AND SEX AGES YFAR YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Operations of war and their sequelae (Y36,Y89.1) Non-Hispanic 1/ 21 1 Male 20 1 Female 1 Non-Hispanic white 20 1 Male 19 1 Female 1 Non-Hispanic black Male Female Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 1 OF 2 PAGE 271 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 25-29 30-34 40-44 ALL 1-4 10-14 20-24 35 - 39POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Complications of medical and surgical care (Y40-Y84,Y88) All origins 2,823 Male 1,308 Female 1,515 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC UNDER 1 5-9 15-19 20-24 25-29 30-34 35-39 40-44 ALL 1-4 10-14 POPULATION, AND SEX AGES YFAR YEARS YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS Complications of medical and surgical care (Y40-Y84,Y88) Non-Hispanic 1/ 2,714 Male 1,253 Female 1,461 2,212 Non-Hispanic white Male 1,025 Female 1,187 Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60 - 64 YEARS	65 - 69 YEARS	70 - 74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
All causes										
All origins	68,120	84,854	104,884	134,095	186,695	265,905	338,349	360,241	646,141	356
Male	43,303	52,356	63,441	79,283	107,529	147,391	174,853	166,117	209,989	277
Female	24,817	32,498	41,443	54,812	79,166	118,514	163,496	194,124	436,152	79
Hispanic	4,767	5,109	5,495	6,824	8,843	10,358	11,081	9,831	17,358	66
Male	3,213	3,284	3,409	4,081	5,026	5,757	5,860	4,457	6,253	60
Female	1,554	1,825	2,086	2,743	3,817	4,601	5,221	5,374	11,105	6
Mexican	2,643	2,840	2,989	3,673	4,870	5,798	5,987	4,860	8,214	20
Male	1,793	1,826	1,823	2,167	2,726	3,182	3,266	2,267	3,156	20
Female	850	1,014	1,166	1,506	2,144	2,616	2,721	2,593	5,058	-
Puerto Rican	840	948	934	1,136	1,249	1,343	1,382	1,195	1,870	1
Male	585	645	625	718	704	739	675	514	644	-
Female	255	303	309	418	545	604	707	681	1,226	1
Cuban	254	259	414	657	955	1,172	1,505	1,699	3,507	8
Male	176	178	273	417	628	721	818	785	1,141	5
Female	78	81	141	240	327	451	687	914	2,366	3
Central and South American	523	498	557	629	793	808	788	738	1,203	2
Male	316	289	313	355	383	377	328	261	343	2
Female	207	209	244	274	410	431	460	477	860	-
Other and unknown Hispanic	507	564	601	729	976	1,237	1,419	1,339	2,564	35
Male	343	346	375	424	585	738	773	630	969	33
Female	164	218	226	305	391	499	646	709	1,595	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
All causes										
Non-Hispanic 1/	62,981	79,321	98,934	126,725	177,240	254,712	326,279	349,451	627,035	155
Male	39,813	48,754	59,720	74,820	102,119	141,091	168,453	161,229	203,128	113
Female	23,168	30,567	39,214	51,905	75,121	113,621	157,826	188,222	423,907	42
Non-Hispanic white	45,974	60,834	78,170	102,623	148,130	219,360	288,018	316,326	573,327	91
Male	29,448	37,703	47,570	61,122	86,299	122,856	149,671	146,857	184,938	66
Female	16,526	23,131	30,600	41,501	61,831	96,504	138,347	169,469	388,389	25
Non-Hispanic black	15,131	16,246	18,176	20,887	25,217	30,350	32,618	28,016	45,895	58
Male	9,266	9,760	10,627	11,866	13,720	15,539	15,706	11,846	14,628	42
Female	5,865	6,486	7,549	9,021	11,497	14,811	16,912	16,170	31,267	16
Origin not stated 2/	372	424	455	546	612	835	989	959	1,748	135
Male	277	318	312	382	384	543	540	431	608	104
Female	95	106	143	164	228	292	449	528	1,140	31

E 2 OF 2 PAGE

3

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS YEARS AND OVER STATED Salmonella infections (A01-A02) All origins 2 1 2 2 5 8 4 7 Male 2 2 2 3 2 3 2 Female 1 2 6 1 5 Hispanic 1 1 1 Male 1 1 Female 1 Mexican 1 1 Male Female Puerto Rican Male Female Cuban 1 Male Female 1 Central and South American Male Female Other and unknown Hispanic 1 Male 1 Female

PAGE

4

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Salmonella infections (A01-A02) Non-Hispanic 1/ 2 1 1 5 7 3 7 1 2 2 Male 1 1 3 2 2 2 5 Female 1 5 2 2 7 Non-Hispanic white 1 4 6 Male 2 2 2 Female 2 5 Non-Hispanic black Male Female Origin not stated 2/ Male Female

PAGE 5

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
FORULATION, AND SEX	TEARS	TEARS	AND OVER	SIMIED						
Shigellosis and amebiasis (A03	3,A06)									
All origins	-	-	-	1	-	1	1	-	4	-
Male	-	-	-	1	-	1	-	-	1	-
Female	-	-	-	-	-	-	1	-	3	-
Hispanic	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-
Mexican	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

PAGE 6

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Shigellosis and amebiasis (A03	3,A06)									
Non-Hispanic 1/	-	-	-	1	-	1	-	-	4	-
Male	-	-	-	1	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	3	-
Non-Hispanic white	-	-	-	1	-	1	-	-	4	-
Male	-	-	-	1	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	3	-
Non-Hispanic black	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

PAGE

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Certain other intestinal infections (A04,A07-A09) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

OF 2 PAGE

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Certain other intestinal infections (A04,A07-A09) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Tuberculosis (A16-A19)										
All origins	51	55	67	63	70	103	134	119	173	1
Male	42	45	48	48	49	67	71	66	85	1
Female	9	10	19	15	21	36	63	53	88	-
Hispanic	10	9	9	11	11	13	17	10	13	-
Male	8	8	3	8	8	8	9	4	6	-
Female	2	1	6	3	3	5	8	6	7	-
Mexican	7	8	6	7	8	9	12	7	4	-
Male	6	7	3	5	6	5	7	3	2	-
Female	1	1	3	2	2	4	5	4	2	-
Puerto Rican	-	-	1	-	1	1	1	1	4	-
Male	-	-	-	-	1	-	-	-	2	-
Female	-	-	1	-	-	1	1	1	2	-
Cuban	1	-	-	1	1	1	2	1	3	-
Male	-	-	-	1	1	1	-	-	-	-
Female	1	-	-	-	-	-	2	1	3	-
Central and South American	1	1	1	1	1	1	-	1	1	-
Male	1	1	-	1	-	1	-	1	1	-
Female	-	-	1	-	1	-	-	-	-	-
Other and unknown Hispanic	1	-	1	2	-	1	2	-	1	-
Male	1	-	-	1	-	1	2	-	1	-
Female	-	-	1	1	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992). HISPANIC ORIGIN.										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Tuberculosis (A16-A19)										
Non-Hispanic 1/	39	43	57	52	59	89	116	106	158	-
Male	32	34	44	40	41	58	61	61	78	-
Female	7	9	13	12	18	31	55	45	80	-
Non-Hispanic white	14	16	27	26	37	47	68	74	108	-
Male	12	15	23	18	25	32	33	44	51	-
Female	2	1	4	8	12	15	35	30	57	-
Non-Hispanic black	20	21	22	17	15	27	31	19	26	-
Male	17	16	15	16	11	17	17	8	13	-
Female	3	5	7	1	4	10	14	11	13	-
Origin not stated 2/	2	3	1	-	-	1	1	3	2	1
Male	2	3	1	-	-	1	1	1	1	1
Female	-	-	-	-	-	-	-	2	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Respiratory tuberculosis (A1	16)									
All origins	38	38	53	56	58	81	112	104	153	1
Male	32	32	39	44	41	56	60	60	76	1
Female	6	6	14	12	17	25	52	44	77	-
Hispanic	6	4	6	10	7	11	16	8	12	-
Male	5	4	2	7	5	8	8	3	5	-
Female	1	-	4	3	2	3	8	5	7	-
Mexican	4	4	4	6	5	7	11	5	4	-
Male	3	4	2	4	4	5	6	2	2	-
Female	1	-	2	2	1	2	5	3	2	-
Puerto Rican	-	-	1	-	-	1	1	1	4	-
Male	-	-	-	-	-	-	-	-	2	-
Female	-	-	1	-	-	1	1	1	2	-
Cuban	-	-	-	1	1	1	2	1	3	-
Male	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	3	-
Central and South American	1	-	1	1	1	1	-	1	1	-
Male	1	-	-	1	-	1	-	1	1	-
Female	-	-	1	-	1	-	-	-	-	-
Other and unknown Hispanic	1	-	-	2	-	1	2	-	-	-
Male	1	-	-	1	-	1	2	-	-	-
Female	-	-	-	1	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Respiratory tuberculosis (A16) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other tuberculosis (A17-A19))									
All origins	13	17	14	7	12	22	22	15	20	-
Male	10	13	9	4	8	11	11	6	9	-
Female	3	4	5	3	4	11	11	9	11	-
Hispanic	4	5	3	1	4	2	1	2	1	-
Male	3	4	1	1	3	-	1	1	1	-
Female	1	1	2	-	1	2	-	1	-	-
Mexican	3	4	2	1	3	2	1	2	-	-
Male	3	3	1	1	2	-	1	1	-	-
Female	-	1	1	-	1	2	-	1	-	-
Puerto Rican	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Central and South American	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	1	-
Female	-	-	1	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Other tuberculosis (A17-A19) Non-Hispanic 1/ 8 11 11 6 8 20 21 12 19 8 8 Male 6 8 3 5 11 10 5 3 3 3 9 Female 3 11 11 2 9 9 8 Non-Hispanic white 1 6 14 Male 1 2 3 6 5 1 Female 1 3 3 9 7 Non-Hispanic black 2 8 Male 4 5 2 2 5 3 3 Female 2 3 Origin not stated 2/ 1 1 1 Male 1 1 Female 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Whooping cough (A37)										
All orders										
All origins	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	_	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	_	_	_	_	_	_	_	_	_	-
Female	_	_	-	_	_	-	_	_	_	_
Other and unknown Hispanic	_	_	_	_	_	_	_	_	_	_
Male				·	·		·	_	-	
	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Non-Hispanic 1/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX		50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Scarlet fever and erysipelas	(A38,A46)									
All origins	-	-	-	-	1	1	1	-	2	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	2	-
Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Scarlet fever and erysipelas	(A38,A46)									
Non-Hispanic 1/	-	-	-	-	1	1	1	-	2	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	2	-
Non-Hispanic white	-	-	-	-	1	-	1	-	2	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	2	-
Non-Hispanic black	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Meningococcal infection (A39)										
All origins	11	6	3	1	9	-	7	6	8	-
Male	1	3	-	-	5	-	2	2	2	-
Female	10	3	3	1	4	-	5	4	6	-
Hispanic	2	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-
Mexican	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Central and South American	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Meningococcal infection (A39)										
Non-Hispanic 1/	8	6	3	1	9	-	6	6	8	-
Male	1	3	-	-	5	-	2	2	2	-
Female	7	3	3	1	4	-	4	4	6	-
Non-Hispanic white	7	5	3	1	9	-	6	5	8	-
Male	1	2	-	-	5	-	2	2	2	-
Female	6	3	3	1	4	-	4	3	6	-
Non-Hispanic black	1	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Origin not stated 2/	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Septicemia (A40-A41)										
All origins	731	954	1,178	1,536	2,352	3,398	4,686	5,024	9,166	-
Male	406	513	613	799	1,218	1,701	2,252	2,174	2,786	-
Female	325	441	565	737	1,134	1,697	2,434	2,850	6,380	-
Hispanic	35	57	63	68	119	146	162	134	257	-
Male	22	34	37	41	67	67	90	54	84	-
Female	13	23	26	27	52	79	72	80	173	-
Mexican	20	26	33	28	67	82	90	73	103	-
Male	11	17	21	16	44	36	47	33	44	-
Female	9	9	12	12	23	46	43	40	59	-
Puerto Rican	8	13	15	22	22	24	17	24	41	-
Male	6	4	7	16	10	11	8	9	10	-
Female	2	9	8	6	12	13	9	15	31	-
Cuban	-	-	2	4	4	18	24	22	60	-
Male	-	-	2	2	3	14	17	8	19	-
Female	-	-	-	2	1	4	7	14	41	-
Central and South American	2	3	3	7	10	10	11	6	17	-
Male	1	1	3	5	3	2	3	1	2	-
Female	1	2	-	2	7	8	8	5	15	-
Other and unknown Hispanic	5	15	10	7	16	12	20	9	36	-
Male	4	12	4	2	7	4	15	3	9	-
Female	1	3	6	5	9	8	5	6	27	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992). HISPANIC ORIGIN.										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Septicemia (A40-A41)										
Non-Hispanic 1/	695	890	1,108	1,462	2,224	3,242	4,505	4,873	8,875	-
Male	384	474	573	757	1,147	1,630	2,149	2,112	2,690	-
Female	311	416	535	705	1,077	1,612	2,356	2,761	6,185	-
Non-Hispanic white	421	594	749	1,043	1,646	2,494	3,628	4,089	7,479	-
Male	234	317	381	523	847	1,264	1,762	1,776	2,257	-
Female	187	277	368	520	799	1,230	1,866	2,313	5,222	-
Non-Hispanic black	251	267	329	387	520	692	792	721	1,281	-
Male	136	143	173	219	274	335	343	305	380	-
Female	115	124	156	168	246	357	449	416	901	-
Origin not stated 2/	1	7	7	6	9	10	19	17	34	-
Male	-	5	3	1	4	4	13	8	12	-
Female	1	2	4	5	5	6	6	9	22	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Syphilis (A50-A53)										
All origins	3	2	1	3	2	4	6	4	4	-
Male	2	1	1	2	2	3	-	-	1	-
Female	1	1	-	1	-	1	6	4	3	-
Hispanic	2	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-
Mexican	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Syphilis (A50-A53)										
Non-Hispanic 1/	1	1	1	3	2	4	6	3	4	-
Male	1	-	1	2	2	3	-	-	1	-
Female	-	1	-	1	-	1	6	3	3	-
Non-Hispanic white	1	1	-	3	2	1	2	1	1	-
Male	1	-	-	2	2	-	-	-	-	-
Female	-	1	-	1	-	1	2	1	1	-
Non-Hispanic black	-	-	1	-	-	3	3	2	3	-
Male	-	-	1	-	-	3	-	-	1	-
Female	-	-	-	-	-	-	3	2	2	-
Origin not stated 2/	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Acute poliomyelitis (A80)										
All origins	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Non-Hispanic 1/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Arthropod-borne viral encephalitis (A83-A84,A85.2) All origins 1 1 1 Male 1 1 1 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT 50-54 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Arthropod-borne viral encephalitis (A83-A84,A85.2) Non-Hispanic 1/ 1 1 1 Male 1 1 1 Female 1 Non-Hispanic white 1 1 Male 1 1 1 Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
FORULATION, AND SEX	TEARS	TEARS	AND OVER	SIMIED						
Measles (B05)										
All origins	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-
Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	_
Female	-	-	-	-	-	-	-	-	-	_
Cuban	-	-	-	-	-	-	-	-	-	_
Male	_	_	_	_	-	-	_	_	_	_
Female	-	_	-	-	-	-	-	_	_	_
Central and South American	_	_	-	_	-	-	-	_	_	_
Male	_	_	-	_	-	-	-	_	_	_
Female	_	_	-	_	-	-	-	_	_	_
Other and unknown Hispanic	_	_	_	_	_	_	_	_	_	_
Male	_	_	_	_	_	_	_	_	_	_
	_	_	-	-	-	2	-	_	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
FORULATION, AND SEX	TEARS	TEARS	AND OVER	SIMIED						
Measles (B05)										
Non-Hispanic 1/	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-
Non-Hispanic white	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-
Non-Hispanic black	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASE TENTH REVISION INTER CLASSIFICATION OF DI 1992), HISPANIC ORIC RACE FOR NON-HISPANI POPULATION, AND SEX	RNATIONAL ISEASES, GIN,	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Viral hepatitis (B15	5-B19)									
All origins	871	635	438	348	427	454	389	262	197	-
Male	651	487	298	193	217	209	185	132	82	-
Female	220	148	140	155	210	245	204	130	115	-
Hispanic	99	104	69	58	68	61	42	23	8	-
Male	76	77	43	32	32	27	14	4	4	-
Female	23	27	26	26	36	34	28	19	4	-
Mexican	51	53	37	40	30	28	17	10	4	-
Male	37	38	24	22	14	11	6	1	1	-
Female	14	15	13	18	16	17	11	9	3	-
Puerto Rican	29	37	17	6	11	11	8	4	1	-
Male	23	33	12	3	4	7	3	1	1	-
Female	6	4	5	3	7	4	5	3	-	-
Cuban	2	2	2	6	4	12	11	5	2	-
Male	2	1	2	4	1	6	2	-	1	-
Female	-	1	-	2	3	6	9	5	1	-
Central and South A	American 7	1	5	4	12	6	3	3	1	-
Male	6	-	2	2	5	3	2	2	1	-
Female	1	1	3	2	7	3	1	1	-	-
Other and unknown H	Hispanic 10	11	8	2	11	4	3	1	-	-
Male	8	5	3	1	8	-	1	-	-	-
Female	2	6	5	1	3	4	2	1	-	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Viral hepatitis (B15-B19)										
Non-Hispanic 1/	765	529	366	290	358	392	346	239	189	-
Male	570	408	253	161	184	181	170	128	78	-
Female	195	121	113	129	174	211	176	111	111	-
Non-Hispanic white	580	378	254	211	262	305	285	212	163	-
Male	423	294	180	128	144	151	146	122	67	-
Female	157	84	74	83	118	154	139	90	96	-
Non-Hispanic black	160	117	73	45	59	50	36	17	19	-
Male	125	90	44	21	21	17	13	2	8	-
Female	35	27	29	24	38	33	23	15	11	-
Origin not stated 2/	7	2	3	-	1	1	1	-	-	-
Male	5	2	2	-	1	1	1	-	-	-
Female	2	-	1	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Human immunodeficiency virus (HIV) disease (B20-B24) All origins 2,440 1.467 Male 1,936 1,198 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Human immunodeficiency virus (HIV) disease (B20-B24) Non-Hispanic 1/ 2,108 1,269 1,673 Male 1,042 Female Non-Hispanic white Male Female Non-Hispanic black 1,322 Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malaria (B50-B54)										
All origins	-	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-
Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malaria (B50-B54)										
Non-Hispanic 1/	-	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-
Non-Hispanic white	-	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All o	rigins	392	444	427	471	652	791	927	812	1,123	-
	Male	231	247	220	262	351	409	460	367	381	-
	Female	161	197	207	209	301	382	467	445	742	-
Hispa	nic	33	35	27	35	47	33	43	25	37	-
	Male	17	23	9	20	24	12	21	8	10	-
	Female	16	12	18	15	23	21	22	17	27	-
Mexi	can	24	20	21	18	26	18	20	14	18	-
	Male	10	14	6	13	15	5	12	4	6	-
	Female	14	6	15	5	11	13	8	10	12	-
Puer	to Rican	1	6	2	5	6	4	6	4	4	-
	Male	1	4	1	2	4	3	1	2	1	-
	Female	-	2	1	3	2	1	5	2	3	-
Cuba	n	1	1	1	5	3	3	7	2	7	-
	Male	1	1	1	2	1	-	2	1	3	-
	Female	-	-	-	3	2	3	5	1	4	-
Cent	ral and South American	4	3	1	4	4	5	5	4	4	-
	Male	2	2	1	2	-	2	2	-	-	-
	Female	2	1	-	2	4	3	3	4	4	-
0the	r and unknown Hispanic	3	5	2	3	8	3	5	1	4	-
	Male	3	2	-	1	4	2	4	1	-	-
	Female	-	3	2	2	4	1	1	-	4	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
YEARS	YEARS	AND OVER	STATED						

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 1/	356	407	397	434	603	755	880	786	1,084	-
Male	213	222	208	241	326	396	438	359	370	-
Female	143	185	189	193	277	359	442	427	714	-
Non-Hispanic white	227	282	300	328	488	638	751	687	959	-
Male	145	158	166	179	266	342	382	299	319	-
Female	82	124	134	149	222	296	369	388	640	-
Non-Hispanic black	116	103	83	86	96	96	105	81	106	-
Male	62	52	34	48	47	47	46	48	39	-
Female	54	51	49	38	49	49	59	33	67	-
Origin not stated 2/	3	2	3	2	2	3	4	1	2	-
Male	1	2	3	1	1	1	1	-	1	-
Female	2	-	-	1	1	2	3	1	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasms (C00-C97)										
All origins	18,412	28,269	38,637	50,430	66,814	85,524	89,874	72,896	75,014	9
Male	9,151	14,483	20,891	27,942	37,315	47,229	48,184	36,876	32,484	6
Female	9,261	13,786	17,746	22,488	29,499	38,295	41,690	36,020	42,530	3
Hispanic	1,093	1,423	1,635	2,051	2,598	2,796	2,586	1,863	1,995	-
Male	539	698	901	1,159	1,469	1,576	1,436	967	902	-
Female	554	725	734	892	1,129	1,220	1,150	896	1,093	-
Mexican	586	766	852	1,079	1,399	1,527	1,391	899	955	-
Male	298	366	466	598	804	861	806	475	450	-
Female	288	400	386	481	595	666	585	424	505	-
Puerto Rican	150	251	246	289	321	351	305	209	228	-
Male	84	148	156	179	178	195	165	108	95	-
Female	66	103	90	110	143	156	140	101	133	-
Cuban	81	81	146	235	330	355	348	335	355	-
Male	46	45	79	144	199	221	187	172	152	-
Female	35	36	67	91	131	134	161	163	203	-
Central and South American	158	170	201	225	261	228	208	155	155	-
Male	52	68	82	113	120	109	97	60	59	-
Female	106	102	119	112	141	119	111	95	96	-
Other and unknown Hispanic	118	155	190	223	287	335	334	265	302	-
Male	59	71	118	125	168	190	181	152	146	-
Female	59	84	72	98	119	145	153	113	156	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Non-Hispanic 1/	17,265	26,771	36,902	48,244	64,056	82,516	87,052	70,852	72,852	9
Male	8,581	13,742	19,936	26,708	35,752	45,519	46,616	35,827	31,512	6
Female	8,684	13,029	16,966	21,536	28,304	36,997	40,436	35,025	41,340	3
Non-Hispanic white	13,404	21,540	30,407	40,381	54,833	72,286	77,288	64,007	65,636	7
Male	6,622	10,966	16,371	22,287	30,626	39,892	41,239	32,296	28,161	4
Female	6,782	10,574	14,036	18,094	24,207	32,394	36,049	31,711	37,475	3
Non-Hispanic black	3,278	4,471	5,587	6,735	7,888	8,737	8,347	5,845	6,260	1
Male	1,687	2,411	3,117	3,816	4,423	4,805	4,603	3,000	2,828	1
Female	1,591	2,060	2,470	2,919	3,465	3,932	3,744	2,845	3,432	-
Origin not stated 2/	54	75	100	135	160	212	236	181	167	-
Male	31	43	54	75	94	134	132	82	70	-
Female	23	32	46	60	66	78	104	99	97	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) All origins 1,007 Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasms of lip, oral cavity and pharynx (C00-C14) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasm of esophagus (C15) All origins 1,139 1,392 1,739 1,784 1,202 1.955 1.114 Male 1.112 1.396 1,529 1.328 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 60-64 65-69 70-74 75-79 85 YEARS 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant neoplasm of esophagus (C15) Non-Hispanic 1/ 447 768 1,100 1,339 1,677 1,910 1,725 1,168 1,074 1 Male 390 663 932 1,064 1,349 1,492 1,281 787 550 1 275 Female 57 105 168 328 418 444 381 524 1,390 1,068 Non-Hispanic white 340 596 860 1,063 1,648 1,508 987 Male 303 524 739 859 1,148 1,303 1,127 730 497 Female 37 72 121 204 242 345 381 338 490 Non-Hispanic black 99 162 222 250 257 229 198 73 Male 79 130 177 188 176 164 141 42 42 Female 20 32 45 62 57 40 31 Origin not stated 2/ 3 3 4 6 6 5 1 2 Male 2 3 4 5 5 5 Female 1 1 1 1 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of stoma	ch (C16)									
All origins	450	651	808	1,052	1,359	1,793	2,064	1,796	2,083	-
Male	303	443	558	694	893	1,112	1,250	989	851	-
Female	147	208	250	358	466	681	814	807	1,232	-
Hispanic	50	71	83	109	134	137	137	102	105	-
Male	35	41	55	71	86	76	87	54	35	-
Female	15	30	28	38	48	61	50	48	70	-
Mexican	30	44	39	63	86	75	82	52	41	-
Male	21	27	21	40	55	47	59	31	17	-
Female	9	17	18	23	31	28	23	21	24	-
Puerto Rican	7	13	13	9	9	23	16	11	18	-
Male	4	7	10	5	6	10	8	6	8	-
Female	3	6	3	4	3	13	8	5	10	-
Cuban	-	2	6	6	9	10	9	13	11	-
Male	-	1	6	4	5	5	4	6	3	-
Female	-	1	-	2	4	5	5	7	8	-
Central and South American	8	7	15	21	17	18	17	13	20	-
Male	5	4	11	15	10	9	8	5	2	-
Female	3	3	4	6	7	9	9	8	18	-
Other and unknown Hispanic	5	5	10	10	13	11	13	13	15	-
Male	5	2	7	7	10	5	8	6	5	-
Female	-	3	3	3	3	6	5	7	10	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant neoplasm of stomach (C16) Non-Hispanic 1/ 1,221 1,646 1,924 1,691 1,973 Male 1,029 1,162 Female 1,159 1,251 1,499 1,360 Non-Hispanic white 1,579 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Malignant neoplasms of colon, rectum and anus (C18-C21) All origins 1,623 2,434 3,309 6,095 9,473 8,456 4.658 8.154 11,286 Male 1,374 1.932 3.431 4.452 4.857 3.897 2.689 3,866 Female 1.060 1.377 1.969 2.664 3,702 4.616 4,559 7.420 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Malignant neoplasms of colon, rectum and anus (C18-C21) Non-Hispanic 1/ 1,506 2,298 3,168 4,436 5,811 7,829 9,180 8,210 11,002 Male 816 1,297 1,854 2,570 3,271 4,267 4,699 3,768 3,756 Female 690 1,001 1,314 1,866 2,540 3,562 4,481 4,442 7,246 2,540 4,874 Non-Hispanic white 1,152 1,804 3,644 6,769 8,085 7,342 9,923 Male 634 1,045 1,500 2,138 2,791 3,768 4,150 3,406 3,396 Female 518 759 1,040 1,506 2,083 3,001 3,935 3,936 6,527 Non-Hispanic black 301 410 547 701 808 916 775 955 958 Male 158 216 315 377 420 430 475 311 296 Female 143 194 232 324 388 486 483 464 659 Origin not stated 2/ 4 4 9 10 13 15 19 35 32 Male 1 3 6 6 8 6 11 19 12 Female 3 1 3 4 5 9 8 16 20

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Malignant neoplasms of liver and intrahepatic bile ducts (C22) All origins 1,511 1.163 1.558 1.933 1.914 1.270 Male 1.041 1.243 1.103 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasms of liver and intrahepatic bile ducts (C22) Non-Hispanic 1/ 1,036 1,385 1,773 1,777 1,403 1,191 Male 1,137 1,029 Female 1,457 1,231 Non-Hispanic white 1,089 1,524 1,048 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55 - 59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of pancro	eas (C25)									
All origins	887	1,466	2,039	2,668	3,632	4,661	4,931	4,002	4,142	1
Male	575	907	1,204	1,517	1,976	2,371	2,281	1,659	1,283	-
Female	312	559	835	1,151	1,656	2,290	2,650	2,343	2,859	1
Hispanic	63	74	95	123	155	183	172	100	120	-
Male	43	41	58	59	85	97	68	40	40	-
Female	20	33	37	64	70	86	104	60	80	-
Mexican	36	42	61	69	85	108	95	57	65	-
Male	26	24	37	37	49	58	43	23	26	-
Female	10	18	24	32	36	50	52	34	39	-
Puerto Rican	7	7	13	11	17	14	17	11	12	-
Male	6	5	8	4	8	9	2	5	4	-
Female	1	2	5	7	9	5	15	6	8	-
Cuban	3	7	5	8	22	18	24	13	21	-
Male	3	3	2	2	14	8	12	4	5	-
Female	-	4	3	6	8	10	12	9	16	-
Central and South American	6	8	12	20	14	18	18	11	8	-
Male	2	5	9	9	7	10	4	5	-	-
Female	4	3	3	11	7	8	14	6	8	-
Other and unknown Hispanic	11	10	4	15	17	25	18	8	14	-
Male	6	4	2	7	7	12	7	3	5	-
Female	5	6	2	8	10	13	11	5	9	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasm of pancreas (C25) Non-Hispanic 1/ 820 1,387 1,942 2,538 3,467 4,461 4,747 3,891 4,015 1 Male 528 863 1,145 1,453 1,887 2,263 2,206 1,616 1,241 Female 292 524 797 1,085 1,580 2,198 2,541 2,275 2,774 1 Non-Hispanic white 663 1,108 1,564 2,111 2,926 3,858 4,160 3,465 3,590 1 Male 430 684 920 1,224 1,613 1,992 1,958 1,454 1,119 Female 233 424 644 887 1,313 1,866 2,202 2,011 2,471 Non-Hispanic black 247 327 342 459 513 486 368 141 360 Male 88 158 200 186 230 227 210 134 97 Female 89 127 156 229 286 276 226 271 Origin not stated 2/ 4 5 2 7 10 17 12 11 7 7 Male 4 3 1 5 4 11 3 2 Female 2 1 2 6 6 5 8 5

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of laryn:	x (C32)									
All origins	151	254	410	494	584	655	529	363	291	-
Male	130	216	339	396	458	490	397	284	213	-
Female	21	38	71	98	126	165	132	79	78	-
Hispanic	15	10	17	12	30	25	24	16	17	-
Male	15	9	16	11	26	22	20	11	15	-
Female	-	1	1	1	4	3	4	5	2	-
Mexican	6	3	11	5	11	12	11	9	11	-
Male	6	3	10	5	11	11	8	5	11	-
Female	-	-	1	-	-	1	3	4	-	-
Puerto Rican	6	4	3	2	9	4	5	3	1	-
Male	6	4	3	1	7	3	5	2	-	-
Female	-	-	-	1	2	1	-	1	1	-
Cuban	2	1	1	4	7	4	5	1	2	-
Male	2	1	1	4	5	4	4	1	2	-
Female	-	-	-	-	2	-	1	-	-	-
Central and South American	-	2	1	1	1	1	1	-	1	-
Male	-	1	1	1	1	-	1	-	1	-
Female	-	1	-	-	-	1	-	-	-	-
Other and unknown Hispanic	1	-	1	-	2	4	2	3	2	-
Male	1	-	1	-	2	4	2	3	1	-
Female	-	-	-	-	-	-	-	-	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant neoplasm of larynx (C32) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Malignant neoplasms of trachea, bronchus and lung (C33-C34) All origins 3,916 12,165 7,518 17.179 23.239 28.609 26,556 17.705 12,339 Male 2.346 4,505 7.465 10.548 14.186 17.101 15.461 9.936 6,315 Female 1,570 3.013 4,700 6.631 9.053 11.508 11.095 7,769 6.024 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Malignant neoplasms of trachea, bronchus and lung (C33-C34) Non-Hispanic 1/ 3,795 7,292 11,829 16,720 22,620 27,916 25,928 17,315 11,988 2 Male 2,266 4,350 7,224 10,214 13,750 16,647 15,037 9,687 6,132 2 Female 1,529 2,942 4,605 6,506 8,870 11,269 10,891 7,628 5,856 10,004 Non-Hispanic white 2,910 5,922 14,325 19,694 24,883 23,335 15,818 10,907 1 Male 1,708 3,447 6,023 8,631 11,872 14,724 13,441 8,778 5,556 1 Female 1,202 2,475 3,981 5,694 7,822 10,159 9,894 7,040 5,351 Non-Hispanic black 2,532 2,579 1,246 907 795 1,234 1,628 2,115 2,237 Male 1,088 1,412 1,637 1,630 1,371 756 488 512 Female 283 408 540 703 895 949 866 490 419 Origin not stated 2/ 10 29 33 38 63 79 75 42 27 Male 6 21 22 30 42 50 47 20 15 Female 8 11 8 21 29 28 22 12

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	AND OVER	STATED							
Malignant melanoma of skin	(C43)									
All origins	518	602	615	648	791	861	945	694	681	-
Male	337	380	425	426	526	570	596	410	328	-
Female	181	222	190	222	265	291	349	284	353	-
Hispanic	9	13	17	13	15	12	15	5	11	-
Male	7	9	11	8	10	7	5	4	3	-
Female	2	4	6	5	5	5	10	1	8	-
Mexican	8	7	4	8	5	7	8	1	5	-
Male	6	5	3	5	3	5	3	1	2	-
Female	2	2	1	3	2	2	5	-	3	-
Puerto Rican	-	2	2	2	2	-	-	-	-	-
Male	-	1	2	1	1	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	-	-
Cuban	-	1	3	2	3	3	2	1	2	-
Male	-	1	1	1	2	1	-	1	-	-
Female	-	-	2	1	1	2	2	-	2	-
Central and South American	-	2	3	1	3	1	2	1	2	-
Male	-	1	2	1	3	1	-	-	1	-
Female	-	1	1	-	-	-	2	1	1	-
Other and unknown Hispanic	1	1	5	-	2	1	3	2	2	-
Male	1	1	3	-	1	-	2	2	-	-
Female	-	-	2	-	1	1	1	-	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant melanoma of skin (C43) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of breas	t (C50)									
All origins	2,619	3,555	3,682	3,717	4,147	4,962	5,140	4,511	6,000	-
Male	11	30	36	34	47	56	56	42	54	-
Female	2,608	3,525	3,646	3,683	4,100	4,906	5,084	4,469	5,946	-
Hispanic	160	202	164	157	146	144	115	94	123	-
Male	1	2	3	-	2	3	1	1	-	-
Female	159	200	161	157	144	141	114	93	123	-
Mexican	80	110	79	83	72	77	55	40	48	-
Male	1	1	1	-	-	1	1	1	-	-
Female	79	109	78	83	72	76	54	39	48	-
Puerto Rican	16	35	27	13	22	16	16	12	16	-
Male	-	1	-	-	1	1	-	-	-	-
Female	16	34	27	13	21	15	16	12	16	-
Cuban	16	6	14	18	21	20	18	23	28	-
Male	-	-	-	-	-	-	-	-	-	-
Female	16	6	14	18	21	20	18	23	28	-
Central and South American	32	24	25	21	17	15	10	8	6	-
Male	-	-	-	-	-	1	-	-	-	-
Female	32	24	25	21	17	14	10	8	6	-
Other and unknown Hispanic	16	27	19	22	14	16	16	11	25	-
Male	-	-	2	-	1	-	-	-	-	-
Female	16	27	17	22	13	16	16	11	25	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of breast	(C50)									
Non-Hispanic 1/	2,453	3,347	3,507	3,549	3,991	4,812	5,011	4,408	5,863	-
Male	10	28	33	34	45	52	55	41	54	-
Female	2,443	3,319	3,474	3,515	3,946	4,760	4,956	4,367	5,809	-
Non-Hispanic white	1,820	2,650	2,838	2,941	3,396	4,227	4,474	4,014	5,364	-
Male	8	20	26	25	38	41	48	36	47	-
Female	1,812	2,630	2,812	2,916	3,358	4,186	4,426	3,978	5,317	-
Non-Hispanic black	545	605	577	518	522	515	490	362	461	-
Male	1	8	7	9	6	10	6	5	7	-
Female	544	597	570	509	516	505	484	357	454	-
Origin not stated 2/	6	6	11	11	10	6	14	9	14	-
Male	-	-	-	-	-	1	-	-	-	-
Female	6	6	11	11	10	5	14	9	14	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant neoplasm of cervix uteri (C53) All origins Female Hispanic Female Mexican Female Puerto Rican Female Cuban Female Central and South American Female Other and unknown Hispanic

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YFARS YEARS AND OVER STATED Malignant neoplasm of cervix uteri (C53) Non-Hispanic 1/ Female Non-Hispanic white Female Non-Hispanic black Female Origin not stated 2/ Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55) All origins 1,061 Female 1.061 Hispanic Female Mexican Female Puerto Rican Female Cuban Female Central and South American Female Other and unknown Hispanic

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT 50-54 55-59 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YEARS AND OVER STATED YFARS YFARS Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55) Non-Hispanic 1/ 172 263 398 558 790 929 1,029 925 976 558 790 1,029 Female 172 263 398 929 925 976 Non-Hispanic white 440 140 213 313 587 764 874 813 858 Female 440 587 764 874 140 213 313 813 858 Non-Hispanic black 18 39 76 113 187 159 142 103 111 Female 18 39 76 113 187 159 142 103 111 Origin not stated 2/ 2 1 Female 1 2 2 1 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant neoplasm of ovary (C56) All origins 1,081 1,345 1,986 1,732 1.626 2.143 1,562 Female 1,081 1.345 1.626 1.986 2.143 1,732 1.562 Hispanic Female Mexican Female Puerto Rican Female Cuban Female Central and South American Female Other and unknown Hispanic Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of ovar	y (C56)									
Non-Hispanic 1/	585	881	1,035	1,281	1,570	1,904	2,073	1,683	1,527	-
Female	585	881	1,035	1,281	1,570	1,904	2,073	1,683	1,527	-
Non-Hispanic white	506	762	920	1,141	1,400	1,719	1,890	1,566	1,424	-
Female	506	762	920	1,141	1,400	1,719	1,890	1,566	1,424	-
Non-Hispanic black	54	89	94	117	135	166	159	97	90	-
Female	54	89	94	117	135	166	159	97	90	-
Origin not stated 2/	2	1	5	5	5	7	11	8	1	-
Female	2	1	5	5	5	7	11	8	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Malignant neoplasm of prostate (C61) All origins 1,310 2,711 4,605 6,903 6.532 8,620 Male 1.310 2,711 4.605 6.532 6,903 8.620 Hispanic Male Mexican Male Puerto Rican Male Cuban Male Central and South American Male Other and unknown Hispanic Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Malignant neoplasm of prostate (C61) 4,431 Non-Hispanic 1/ 97 254 595 1,253 2,573 6,308 6,717 8,352 1 4,431 6,717 Male 97 254 595 1,253 2,573 6,308 8,352 1 1,930 3,483 Non-Hispanic white 65 179 430 880 5,134 5,716 7,093 1 880 1,930 3,483 5,716 7,093 Male 65 179 430 5,134 Non-Hispanic black 31 74 157 355 619 905 1,092 935 1,140 31 74 157 355 619 905 1,092 935 1,140 Origin not stated 2/ 15 9 17 14 Male 1 5 3 14 15 9 17

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Malignant neoplasms of kidney and renal pelvis (C64-C65) All origins 1,152 1,374 1,702 1,314 1.696 1.305 Male 1.064 1.016 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasms of kidney and renal pelvis (C64-C65) Non-Hispanic 1/ 1,082 1,295 1,632 1,624 1,268 1,261 Male 1,015 Female 1,464 1,470 Non-Hispanic white 1,149 1,166 1,144 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasm of bladd	er (C67)									
All origins	140	323	492	752	1,126	1,797	2,178	2,139	2,845	1
Male	107	243	374	559	849	1,283	1,575	1,433	1,608	1
Female	33	80	118	193	277	514	603	706	1,237	-
Hispanic	5	17	21	30	35	53	41	38	55	-
Male	4	12	18	25	30	39	28	24	36	-
Female	1	5	3	5	5	14	13	14	19	-
Mexican	-	9	11	15	22	30	23	16	21	-
Male	-	7	8	13	18	20	15	9	11	-
Female	-	2	3	2	4	10	8	7	10	-
Puerto Rican	-	2	4	3	2	3	1	7	8	-
Male	-	2	4	3	2	2	1	3	4	-
Female	-	-	-	-	-	1	-	4	4	-
Cuban	2	2	2	4	7	5	6	7	11	-
Male	2	-	2	3	6	4	4	7	10	-
Female	-	2	-	1	1	1	2	-	1	-
Central and South American	3	4	1	1	-	7	5	3	6	-
Male	2	3	1	1	-	7	3	1	5	-
Female	1	1	-	-	-	-	2	2	1	-
Other and unknown Hispanic	-	-	3	7	4	8	6	5	9	-
Male	-	-	3	5	4	6	5	4	6	-
Female	-	-	-	2	-	2	1	1	3	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Malignant neoplasm of bladder (C67) Non-Hispanic 1/ 135 304 470 720 1,088 1,744 2,129 2,099 2,780 1 Male 103 229 355 533 816 1,244 1,542 1,408 1,567 1 75 187 Female 32 115 272 500 587 691 1,213 407 635 1,587 1,964 1,957 Non-Hispanic white 109 266 969 2,567 1 Male 83 198 321 479 746 1,161 1,445 1,326 1,480 1 Female 26 68 156 223 426 519 631 1,087 Non-Hispanic black 23 37 59 71 105 139 140 122 188 Male 17 30 32 44 59 69 81 68 73 Female 7 27 27 46 70 59 115 Origin not stated 2/ 2 1 2 3 8 2 10 Male 2 1 1 3 5 1 5 Female 1 3 1 5

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malignant neoplasms of meni	nges, brair	n and othe	r parts of	central n	ervous sys	tem (C70-C	72)			
All origins	842	1,038	1,206	1,247	1,498	1,621	1,543	949	663	-
Male	507	613	670	747	858	870	785	441	247	-
Female	335	425	536	500	640	751	758	508	416	-
Hispanic	37	51	47	52	47	57	38	29	17	-
Male	18	24	24	36	25	23	19	12	3	-
Female	19	27	23	16	22	34	19	17	14	-
Mexican	23	33	23	30	25	32	18	12	9	-
Male	10	18	12	20	15	12	13	4	-	-
Female	13	15	11	10	10	20	5	8	9	-
Puerto Rican	5	5	1	5	1	7	2	3	-	-
Male	2	2	-	4	1	1	1	3	-	-
Female	3	3	1	1	-	6	1	-	-	-
Cuban	-	1	5	7	7	7	5	6	5	-
Male	-	-	3	5	1	5	2	4	3	-
Female	-	1	2	2	6	2	3	2	2	-
Central and South American	4	7	11	5	9	6	3	5	2	-
Male	2	2	3	2	5	3	-	-	-	-
Female	2	5	8	3	4	3	3	5	2	-
Other and unknown Hispanic	5	5	7	5	5	5	10	3	1	-
Male	4	2	6	5	3	2	3	1	-	-
Female	1	3	1	-	2	3	7	2	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72) Non-Hispanic 1/ 1,157 1,191 1,449 1,561 1,501 Male Female 1,074 1,097 1,351 1,488 1,423 Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) All origins 3,346 5,996 8,493 1.648 2.302 4.208 8.336 9.596 8,282 Male 1.382 1.957 3.436 4.631 3.998 2.469 5.102 3,329 Female 1.389 1.739 2.560 3,705 4,494 4,495 4.953 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) Non-Hispanic 1/ 1,503 2,139 3,158 3,985 5,698 8,006 9,299 8,262 8,079 Male 900 1,290 1,851 2,344 3,284 4,456 4,953 3,900 3,261 Female 603 849 1,307 1,641 2,414 3,550 4,346 4,362 4,818 2,654 8,487 Non-Hispanic white 1,220 1,760 3,388 5,004 7,204 7,685 7,493 Male 733 1,079 1,569 2,022 2,919 4,048 4,538 3,664 3,020 Female 487 681 1,085 1,366 2,085 3,156 3,949 4,021 4,473 Non-Hispanic black 318 431 486 582 665 477 502 236 665 Male 138 180 242 258 309 336 188 186 333 Female 98 138 189 228 273 329 332 289 316 Origin not stated 2/ 4 5 6 9 11 11 24 19 15 Male 2 1 4 1 7 8 12 9 5 Female 4 2 8 4 3 12 10 10

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hodgkin's disease (C81)										
All origins	82	83	87	91	120	163	149	113	98	-
Male	48	50	54	61	72	88	69	50	45	-
Female	34	33	33	30	48	75	80	63	53	-
Hispanic	6	11	7	10	14	12	9	9	5	-
Male	3	8	4	5	7	6	5	6	2	-
Female	3	3	3	5	7	6	4	3	3	-
Mexican	4	6	3	3	11	7	5	5	3	-
Male	2	4	1	2	4	4	3	3	2	-
Female	2	2	2	1	7	3	2	2	1	-
Puerto Rican	-	1	2	2	1	2	2	-	-	-
Male	-	1	2	1	1	-	-	-	-	-
Female	-	-	-	1	-	2	2	-	-	-
Cuban	1	2	-	-	1	1	1	2	1	-
Male	1	2	-	-	1	1	1	1	-	-
Female	-	-	-	-	-	-	-	1	1	-
Central and South American	-	1	1	3	1	1	-	2	1	-
Male	-	1	1	2	1	1	-	2	-	-
Female	-	-	-	1	-	-	-	-	1	-
Other and unknown Hispanic	1	1	1	2	-	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-
Female	1	1	1	2	_	1	_	_	-	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hodgkin's disease (C81)										
Non-Hispanic 1/	75	72	79	80	106	151	140	103	93	-
Male	45	42	49	56	65	82	64	44	43	-
Female	30	30	30	24	41	69	76	59	50	-
Non-Hispanic white	66	63	70	74	92	142	135	100	84	-
Male	39	34	44	53	57	78	62	43	39	-
Female	27	29	26	21	35	64	73	57	45	-
Non-Hispanic black	9	9	9	5	11	7	5	1	7	-
Male	6	8	5	3	8	4	2	1	2	-
Female	3	1	4	2	3	3	3	-	5	-
Origin not stated 2/	1	-	1	1	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Non-Hodgkin's lymphoma (C82-C85) All origins 1,414 2,441 3,407 3,545 1,045 1.740 4.009 3,228 Male 1.355 1.821 1.544 2.052 1.246 Female 1,086 1.586 1.957 2,001 1.982 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 60-64 70-74 75-79 85 YEARS AGE NOT 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Non-Hodgkin's lymphoma (C82-C85) Non-Hispanic 1/ 630 976 1,350 1,646 2,318 3,257 3,880 3,447 3,142 Male 416 609 812 942 1,296 1,748 1,986 1,504 1,216 704 Female 214 367 538 1,022 1,509 1,894 1,943 1,926 1,479 Non-Hispanic white 517 855 1,188 2,129 3,016 3,632 3,303 2,975 Male 344 535 713 841 1,190 1,632 1,847 1,455 1,138 Female 173 320 475 638 939 1,384 1,785 1,848 1,837 Non-Hispanic black 97 138 121 177 179 107 125 92 153 Male 58 62 81 72 90 84 99 50 Female 34 35 57 49 63 93 80 75 75 Origin not stated 2/ 2 2 2 3 6 6 12 10 6 7 Male 1 1 1 1 3 4 7 2 Female 1 1 1 2 3 2 5 3 4

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Leukemia (C91-C95)										
All origins	626	744	1,112	1,404	2,078	2,934	3,401	3,059	3,424	_
Male	346	431	636	853	1,275	1,779	1,946	1,579	1,423	_
Female	280	313	476	551	803	1,155	1,455	1,480	2,001	_
Hispanic	50	50	62	53	86	93	85	64	63	_
Male	30	22	31	35	45	62	45	30	23	_
Female	20	28	31	18	41	31	40	34	40	-
Mexican	32	27	32	27	38	37	39	29	28	-
Male	21	13	16	17	18	25	21	15	9	-
Female	11	14	16	10	20	12	18	14	19	-
Puerto Rican	6	8	7	8	16	16	9	5	6	-
Male	6	3	4	4	12	11	6	2	2	-
Female	-	5	3	4	4	5	3	3	4	-
Cuban	4	1	9	9	16	20	18	18	16	-
Male	2	1	4	6	8	12	10	8	7	-
Female	2	-	5	3	8	8	8	10	9	-
Central and South American	8	11	7	4	10	8	5	6	5	-
Male	1	4	2	3	3	5	2	3	3	-
Female	7	7	5	1	7	3	3	3	2	-
Other and unknown Hispanic	-	3	7	5	6	12	14	6	8	-
Male	-	1	5	5	4	9	6	2	2	-
Female	-	2	2	-	2	3	8	4	6	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Leukemia (C91-C95)										
Non-Hispanic 1/	575	691	1,048	1,347	1,989	2,837	3,308	2,990	3,356	-
Male	315	409	603	818	1,227	1,713	1,898	1,548	1,400	-
Female	260	282	445	529	762	1,124	1,410	1,442	1,956	-
Non-Hispanic white	473	567	911	1,164	1,751	2,597	3,073	2,775	3,152	-
Male	255	338	530	716	1,094	1,579	1,767	1,453	1,316	-
Female	218	229	381	448	657	1,018	1,306	1,322	1,836	-
Non-Hispanic black	79	95	106	146	200	205	196	175	173	-
Male	47	56	61	81	111	111	108	70	66	-
Female	32	39	45	65	89	94	88	105	107	-
Origin not stated 2/	1	3	2	4	3	4	8	5	5	-
Male	1	-	2	-	3	4	3	1	-	-
Female	-	3	-	4	-	-	5	4	5	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Multiple myeloma and immunoproliferative neoplasms (C88,C90) All origins 1,347 1,825 1,765 2.023 1,519 Male 1.026 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Multiple myeloma and immunoproliferative neoplasms (C88,C90) Non-Hispanic 1/ 1,275 1,755 1,959 1,711 1,475 Male Female 1,022 1,443 1,497 Non-Hispanic white 1,636 1,270 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96) 3 6 6 6 10 7 14 11 13 All origins Male 1 3 5 2 5 3 9 5 3 Female 2 3 1 5 4 5 6 10 Hispanic 1 2 Male 1 Female 1 1 1 Mexican 1 Male Female 1 Puerto Rican 1 Male Female 1 Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96) Non-Hispanic 1/ 3 6 6 6 10 6 12 11 13 3 3 Male 1 2 5 3 8 5 3 5 3 Female 4 4 6 10 5 10 12 Non-Hispanic white 2 6 6 11 10 Male 3 5 3 Female 3 3 9 Non-Hispanic black Male 1 Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Origin not stated 2/
Male
Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
All other and unspecified m	alignant no	eoplasms (C17,C23-C2	4,C26-C31,	C37-C41,C4	4-C49,C51-	C52,C57-C6	0,C62-C63	, C66 , C68-C6	9,C73-C80,C97
All origins	2,247	3,230	4,249	5,614	7,265	9,569	10,451	9,205	10,403	2
Male	1,274	1,824	2,416	3,181	3,984	5,173	5,240	4,199	3,806	1
Female	973	1,406	1,833	2,433	3,281	4,396	5,211	5,006	6,597	1
Hispanic	144	170	216	255	325	346	336	251	288	-
Male	75	87	115	141	161	190	159	119	109	-
Female	69	83	101	114	164	156	177	132	179	-
Mexican	71	100	116	145	186	197	192	134	137	-
Male	37	48	70	77	97	104	96	65	53	-
Female	34	52	46	68	89	93	96	69	84	-
Puerto Rican	26	26	22	30	20	40	29	25	26	-
Male	15	16	13	20	8	22	14	12	8	-
Female	11	10	9	10	12	18	15	13	18	-
Cuban	7	6	22	28	30	38	40	34	46	-
Male	6	4	11	16	17	28	18	16	17	-
Female	1	2	11	12	13	10	22	18	29	-
Central and South American	22	23	33	26	41	28	33	21	27	-
Male	8	9	7	13	18	8	10	9	10	-
Female	14	14	26	13	23	20	23	12	17	-
Other and unknown Hispanic	18	15	23	26	48	43	42	37	52	-
Male	9	10	14	15	21	28	21	17	21	-
Female	9	5	9	11	27	15	21	20	31	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 50-54 55-59 60-64 65-69 75 - 7980-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS YFARS STATED All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97 2,095 3,054 4,021 5,340 6,926 9,195 10,091 8,929 10,094 Non-Hispanic 1/ Male 1,194 1,734 2,295 3,030 3,816 4,963 5,065 4,067 3,692 1 Female 901 1,320 1,726 2,310 3,110 4,232 5,026 4,862 6,402 1 3,273 Non-Hispanic white 1,624 2,453 4,449 6,006 8,070 9,032 8,126 9,204 Male 915 1,390 1,885 2,538 3,321 4,393 4,580 3,742 3,383 1 Female 709 1,063 1,388 1,911 2,685 3,677 4,452 4,384 5,821 Non-Hispanic black 778 766 938 757 416 512 629 876 663 Male 244 295 361 440 418 477 401 266 247 Female 172 217 268 338 348 461 475 397 510 Origin not stated 2/ 8 6 12 19 14 28 24 25 21 Male 5 3 6 10 7 20 16 13 5 Female 3 3 6 9 7 8 8 12 16

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

C URIGIN,										
ISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT
D SEX	YEARS	AND OVER	STATED							

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins	276	386	461	672	1,029	1,655	2,151	2,321	3,443	-
Male	142	220	255	401	590	932	1,119	1,099	1,303	-
Female	134	166	206	271	439	723	1,032	1,222	2,140	-
Hispanic	15	24	31	29	39	70	57	54	74	-
Male	9	17	14	16	13	34	24	23	23	-
Female	6	7	17	13	26	36	33	31	51	-
Mexican	10	14	13	18	19	37	34	24	31	-
Male	7	11	6	10	6	18	14	9	8	-
Female	3	3	7	8	13	19	20	15	23	-
Puerto Rican	1	5	4	4	6	8	7	5	8	-
Male	-	4	3	3	3	4	4	3	4	-
Female	1	1	1	1	3	4	3	2	4	-
Cuban	1	1	3	1	2	11	8	11	15	-
Male	1	-	-	-	2	6	2	4	7	-
Female	-	1	3	1	-	5	6	7	8	-
Central and South American	3	1	5	5	7	6	3	6	8	-
Male	1	1	3	3	-	4	2	2	1	-
Female	2	-	2	2	7	2	1	4	7	-
Other and unknown Hispanic	-	3	6	1	5	8	5	8	12	-
Male	-	1	2	-	2	2	2	5	3	-
Female	-	2	4	1	3	6	3	3	9	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) Non-Hispanic 1/ 261 359 428 642 988 1,584 2,088 2,261 3,362 Male 133 201 239 384 576 898 1,093 1,075 1,278 Female 128 158 189 258 412 686 995 1,186 2,084 542 1,430 2,102 Non-Hispanic white 199 295 357 846 1,899 3,133 Male 107 169 200 336 503 823 1,002 1,011 1,209 Female 92 126 206 343 607 897 1,091 1,924 Non-Hispanic black 54 84 124 194 50 132 162 133 Male 19 25 36 39 65 63 75 49 54 Female 31 29 28 45 59 69 87 84 140 Origin not stated 2/ 3 2 1 2 1 6 6 7 Male 2 2 1 1 2 1 2

1

1

4

5

5

Source: CDC/NCHS, National Vital Statistics System, Mortality

1

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Anemias (D50-D64)										
All origins	134	98	105	126	175	311	497	667	1,820	-
Male	68	48	43	62	92	134	217	267	518	-
Female	66	50	62	64	83	177	280	400	1,302	-
Hispanic	6	3	5	5	7	20	16	19	50	-
Male	4	-	-	3	4	13	6	11	19	-
Female	2	3	5	2	3	7	10	8	31	-
Mexican	1	-	4	4	4	12	14	11	23	-
Male	-	-	-	3	3	8	5	6	6	-
Female	1	-	4	1	1	4	9	5	17	-
Puerto Rican	1	2	1	-	1	1	2	2	7	-
Male	1	-	-	-	1	1	1	1	3	-
Female	-	2	1	-	-	-	1	1	4	-
Cuban	2	-	-	-	-	4	-	3	7	-
Male	2	-	-	-	-	3	-	2	3	-
Female	-	-	-	-	-	1	-	1	4	-
Central and South American	2	-	-	1	2	1	-	2	3	-
Male	1	-	-	-	-	1	-	2	1	-
Female	1	-	-	1	2	-	-	-	2	-
Other and unknown Hispanic	-	1	-	-	-	2	-	1	10	-
Male	-	-	-	-	-	-	-	-	6	-
Female	-	1	-	-	-	2	-	1	4	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992). HISPANIC ORIGIN.										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Anemias (D50-D64)										
Non-Hispanic 1/	126	95	100	120	165	290	478	647	1,764	-
Male	63	48	43	58	87	120	209	256	497	-
Female	63	47	57	62	78	170	269	391	1,267	-
Non-Hispanic white	48	38	41	71	126	234	410	585	1,567	-
Male	26	20	16	29	70	93	183	233	449	-
Female	22	18	25	42	56	141	227	352	1,118	-
Non-Hispanic black	73	52	55	45	33	48	63	56	164	-
Male	35	26	25	28	14	23	22	19	37	-
Female	38	26	30	17	19	25	41	37	127	-
Origin not stated 2/	2	-	-	1	3	1	3	1	6	-
Male	1	-	-	1	1	1	2	-	2	-
Female	1	-	-	-	2	-	1	1	4	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						
Diabetes mellitus (E10-E14)										
All origins	1,942	2,793	3,865	5,232	7,215	9,693	11,269	10,488	13,178	3
Male	1,189	1,602	2,120	2,736	3,630	4,649	5,190	4,349	4,125	2
Female	753	1,191	1,745	2,496	3,585	5,044	6,079	6,139	9,053	1
Hispanic	169	244	368	559	721	830	782	587	704	-
Male	105	147	202	278	331	378	336	226	217	-
Female	64	97	166	281	390	452	446	361	487	-
Mexican	124	163	247	349	482	548	489	360	380	-
Male	76	98	127	161	214	240	204	135	115	-
Female	48	65	120	188	268	308	285	225	265	-
Puerto Rican	22	37	53	93	88	91	86	77	73	-
Male	16	21	33	55	45	37	36	28	18	-
Female	6	16	20	38	43	54	50	49	55	-
Cuban	7	7	16	31	47	59	78	65	123	-
Male	4	7	11	17	22	33	42	27	39	-
Female	3	-	5	14	25	26	36	38	84	-
Central and South American	7	10	18	27	36	52	45	27	37	-
Male	4	3	11	15	20	23	15	7	13	-
Female	3	7	7	12	16	29	30	20	24	-
Other and unknown Hispanic	9	27	34	59	68	80	84	58	91	-
Male	5	18	20	30	30	45	39	29	32	-
Female	4	9	14	29	38	35	45	29	59	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Diabetes mellitus (E10-E14)										
Non-Hispanic 1/	1,769	2,536	3,484	4,654	6,477	8,841	10,455	9,870	12,436	2
Male	1,082	1,443	1,911	2,447	3,288	4,260	4,842	4,113	3,894	1
Female	687	1,093	1,573	2,207	3,189	4,581	5,613	5,757	8,542	1
Non-Hispanic white	1,197	1,743	2,382	3,292	4,820	6,844	8,523	8,298	10,533	-
Male	760	1,041	1,379	1,821	2,568	3,498	4,116	3,606	3,374	-
Female	437	702	1,003	1,471	2,252	3,346	4,407	4,692	7,159	-
Non-Hispanic black	513	706	948	1,188	1,438	1,682	1,635	1,360	1,642	2
Male	290	351	457	529	630	632	588	428	426	1
Female	223	355	491	659	808	1,050	1,047	932	1,216	1
Origin not stated 2/	4	13	13	19	17	22	32	31	38	1
Male	2	12	7	11	11	11	12	10	14	1
Female	2	1	6	8	6	11	20	21	24	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Nutritional deficiencies (E40	-E64)									
All origins	45	61	76	94	149	280	472	776	2,240	-
Male	22	30	42	56	72	149	205	289	577	-
Female	23	31	34	38	77	131	267	487	1,663	-
Hispanic	1	5	5	2	4	13	16	15	73	-
Male	1	4	1	1	1	7	10	3	24	-
Female	-	1	4	1	3	6	6	12	49	-
Mexican	1	3	5	1	-	8	13	12	49	-
Male	1	3	1	-	-	4	8	3	16	-
Female	-	-	4	1	-	4	5	9	33	-
Puerto Rican	-	2	-	-	2	2	1	-	4	-
Male	-	1	-	-	1	1	-	-	1	-
Female	-	1	-	-	1	1	1	-	3	-
Cuban	-	-	-	-	-	-	-	-	6	-
Male	-	-	-	-	-	-	-	-	3	-
Female	-	-	-	-	-	-	-	-	3	-
Central and South American	-	-	-	-	-	-	-	1	2	-
Male	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	1	1	-
Other and unknown Hispanic	-	-	-	1	2	3	2	2	12	-
Male	-	-	-	1	-	2	2	-	3	-
Female	-	-	-	-	2	1	-	2	9	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Nutritional deficiencies (E40-E64) Non-Hispanic 1/ 2,162 Male Female 1,610 Non-Hispanic white 1,916 Male Female 1,438 Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55 - 59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malnutrition (E40-E46)										
All origins	42	57	66	88	141	266	439	720	2,077	-
Male	21	27	38	54	68	139	186	266	530	-
Female	21	30	28	34	73	127	253	454	1,547	-
Hispanic	1	5	5	2	4	13	15	15	68	-
Male	1	4	1	1	1	7	9	3	21	-
Female	-	1	4	1	3	6	6	12	47	-
Mexican	1	3	5	1	-	8	12	12	46	-
Male	1	3	1	-	-	4	7	3	13	-
Female	-	-	4	1	-	4	5	9	33	-
Puerto Rican	-	2	-	-	2	2	1	-	4	-
Male	-	1	-	-	1	1	-	-	1	-
Female	-	1	-	-	1	1	1	-	3	-
Cuban	-	-	-	-	-	-	-	-	6	-
Male	-	-	-	-	-	-	-	-	3	-
Female	-	-	-	-	-	-	-	-	3	-
Central and South American	-	-	-	-	-	-	-	1	2	-
Male	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	1	1	-
Other and unknown Hispanic	-	-	-	1	2	3	2	2	10	-
Male	-	-	-	1	-	2	2	-	3	-
Female	-	-	-	-	2	1	-	2	7	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Malnutrition (E40-E46)										
Non-Hispanic 1/	41	52	60	86	137	252	423	704	2,004	-
Male	20	23	36	53	67	132	176	262	508	-
Female	21	29	24	33	70	120	247	442	1,496	-
Non-Hispanic white	26	41	35	64	103	184	350	595	1,773	-
Male	14	17	18	40	47	91	142	217	435	-
Female	12	24	17	24	56	93	208	378	1,338	-
Non-Hispanic black	13	11	21	21	29	59	62	99	204	-
Male	5	6	15	12	17	36	28	38	63	-
Female	8	5	6	9	12	23	34	61	141	-
Origin not stated 2/	-	-	1	-	-	1	1	1	5	-
Male	-	-	1	-	-	-	1	1	1	-
Female	-	-	-	-	-	1	-	-	4	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Other nutritional deficiencies (E50-E64) All origins 3 4 10 6 8 14 33 163 56 Male 1 3 4 2 4 10 19 23 47 Female 2 1 6 4 4 14 33 116 Hispanic 1 5 3 Male 1 2 Female 1 3 Mexican 3 Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic 2 Male 2

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Other nutritional deficiencies (E50-E64) Non-Hispanic 1/ 3 4 10 6 8 14 32 56 158 3 Male 1 4 2 4 10 18 23 44 Female 1 6 4 4 4 14 33 114 2 5 8 25 Non-Hispanic white 1 6 12 48 143 Male 1 2 4 8 12 21 43 Female 4 13 27 100 Non-Hispanic black 1 2 15 Male 1 3 2 6 2 1 Female 14 Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Meningitis (G00,G03)										
All origins	62	46	41	49	49	66	65	57	62	-
Male	39	29	24	21	28	32	26	23	20	-
Female	23	17	17	28	21	34	39	34	42	-
Hispanic	8	4	3	4	6	3	4	-	4	-
Male	4	2	3	-	2	3	-	-	1	-
Female	4	2	-	4	4	-	4	-	3	-
Mexican	4	3	2	1	2	2	2	-	3	-
Male	2	1	2	-	-	2	-	-	1	-
Female	2	2	-	1	2	-	2	-	2	-
Puerto Rican	1	1	-	1	2	1	-	-	-	-
Male	-	1	-	-	2	1	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-
Cuban	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-
Central and South American	1	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	2	-	-	-
Other and unknown Hispanic	2	-	1	1	1	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Meningitis (G00,G03)										
Non-Hispanic 1/	54	42	37	45	43	63	61	57	58	-
Male	35	27	20	21	26	29	26	23	19	-
Female	19	15	17	24	17	34	35	34	39	-
Non-Hispanic white	31	29	28	35	32	48	44	48	46	-
Male	18	17	16	17	19	22	17	21	15	-
Female	13	12	12	18	13	26	27	27	31	-
Non-Hispanic black	22	13	8	10	9	12	16	8	12	-
Male	17	10	4	4	6	6	8	2	4	-
Female	5	3	4	6	3	6	8	6	8	-
Origin not stated 2/	-	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Parkinson's disease (G20-G21)										
All origins	8	35	73	169	539	1,481	3,066	4,044	5,168	-
Male	5	25	44	117	338	965	1,940	2,367	2,454	-
Female	3	10	29	52	201	516	1,126	1,677	2,714	-
Hispanic	-	2	2	6	24	60	97	87	110	-
Male	-	2	1	4	19	33	60	53	49	-
Female	-	-	1	2	5	27	37	34	61	-
Mexican	-	1	1	3	15	34	45	50	52	-
Male	-	1	-	2	12	21	25	27	25	-
Female	-	-	1	1	3	13	20	23	27	-
Puerto Rican	-	-	1	-	4	4	13	5	9	-
Male	-	-	1	-	4	-	9	5	5	-
Female	-	-	-	-	-	4	4	-	4	-
Cuban	-	-	-	1	3	7	20	13	21	-
Male	-	-	-	1	2	5	12	9	8	-
Female	-	-	-	-	1	2	8	4	13	-
Central and South American	-	1	-	-	2	5	1	5	2	-
Male	-	1	-	-	1	2	1	1	2	-
Female	-	-	-	-	1	3	-	4	-	-
Other and unknown Hispanic	-	-	-	2	-	10	18	14	26	-
Male	-	-	-	1	-	5	13	11	9	-
Female	_	-	_	1	-	5	5	3	17	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Parkinson's disease (G20-G21)										
Non-Hispanic 1/	8	33	71	163	511	1,419	2,967	3,950	5,046	-
Male	5	23	43	113	318	930	1,878	2,311	2,402	-
Female	3	10	28	50	193	489	1,089	1,639	2,644	-
Non-Hispanic white	8	29	63	144	471	1,336	2,811	3,783	4,822	-
Male	5	22	40	102	294	888	1,780	2,225	2,287	-
Female	3	7	23	42	177	448	1,031	1,558	2,535	-
Non-Hispanic black	-	4	7	15	32	63	109	112	160	-
Male	-	1	3	10	22	36	67	56	74	-
Female	-	3	4	5	10	27	42	56	86	-
Origin not stated 2/	-	-	-	-	4	2	2	7	12	-
Male	-	-	-	-	1	2	2	3	3	-
Female	-	-	-	-	3	-	-	4	9	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						
Alzheimer's disease (G30)										
All origins	20	40	125	323	830	2,374	5,769	10,067	24,980	-
Male	10	18	68	154	389	1,048	2,332	3,572	5,796	-
Female	10	22	57	169	441	1,326	3,437	6,495	19,184	-
Hispanic	-	3	2	12	25	75	166	196	501	-
Male	-	-	-	5	17	34	67	61	129	-
Female	-	3	2	7	8	41	99	135	372	-
Mexican	-	2	-	3	8	39	80	87	226	-
Male	-	-	-	1	5	18	39	26	67	-
Female	-	2	-	2	3	21	41	61	159	-
Puerto Rican	-	-	1	4	5	11	19	23	45	-
Male	-	-	-	2	3	2	4	8	9	-
Female	-	-	1	2	2	9	15	15	36	-
Cuban	-	-	1	2	6	8	32	39	115	-
Male	-	-	-	-	4	4	13	12	27	-
Female	-	-	1	2	2	4	19	27	88	-
Central and South American	-	1	-	1	3	5	7	10	29	-
Male	-	-	-	-	2	1	2	5	4	-
Female	-	1	-	1	1	4	5	5	25	-
Other and unknown Hispanic	-	-	-	2	3	12	28	37	86	-
Male	-	-	-	2	3	9	9	10	22	-
Female	-	-	-	-	-	3	19	27	64	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
TOTOLATION, AND SEA	TEARS	TEARS	TEARS	TEARS	TEARS	12/11(3	TEARS	TEARS	AND OVER	31/(120
Alzheimer's disease (G30)										
Non-Hispanic 1/	20	37	123	311	804	2,295	5,594	9,853	24,427	-
Male	10	18	68	149	372	1,012	2,262	3,506	5,652	-
Female	10	19	55	162	432	1,283	3,332	6,347	18,775	-
Non-Hispanic white	18	33	111	288	731	2,090	5,195	9,258	23,106	-
Male	10	16	61	139	338	925	2,113	3,319	5,330	-
Female	8	17	50	149	393	1,165	3,082	5,939	17,776	-
Non-Hispanic black	2	2	10	19	62	187	358	525	1,159	-
Male	-	2	6	9	31	81	130	160	270	-
Female	2	-	4	10	31	106	228	365	889	-
Origin not stated 2/	-	-	-	-	1	4	9	18	52	-
Male	-	-	-	-	-	2	3	5	15	-
Female	-	-	-	-	1	2	6	13	37	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Major cardiovascular diseases	(100-178)									
All origins	17,354	25,158	33,344	44,489	64,775	98,653	138,680	163,872	340,510	49
Male	12,150	17,595	22,653	28,872	39,978	56,926	72,052	73,515	102,663	38
Female	5,204	7,563	10,691	15,617	24,797	41,727	66,628	90,357	237,847	11
Hispanic	994	1,392	1,696	2,277	3,147	4,003	4,708	4,583	9,031	6
Male	696	1,004	1,124	1,436	1,890	2,322	2,523	1,996	3,058	6
Female	298	388	572	841	1,257	1,681	2,185	2,587	5,973	-
Mexican	572	808	915	1,207	1,744	2,213	2,536	2,227	4,270	2
Male	389	575	587	756	1,001	1,267	1,416	1,014	1,553	2
Female	183	233	328	451	743	946	1,120	1,213	2,717	-
Puerto Rican	149	228	299	433	453	541	635	585	958	-
Male	104	172	199	274	263	304	310	244	319	-
Female	45	56	100	159	190	237	325	341	639	-
Cuban	72	84	131	210	356	459	656	819	1,902	1
Male	55	68	103	141	260	291	361	376	561	1
Female	17	16	28	69	96	168	295	443	1,341	-
Central and South American	111	148	183	208	278	324	328	364	651	-
Male	81	98	117	130	147	152	128	115	172	-
Female	30	50	66	78	131	172	200	249	479	-
Other and unknown Hispanic	90	124	168	219	316	466	553	588	1,250	3
Male	67	91	118	135	219	308	308	247	453	3
Female	23	33	50	84	97	158	245	341	797	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Major cardiovascular diseases	(100-178)									
Non-Hispanic 1/	16,278	23,657	31,488	41,985	61,383	94,260	133,532	158,822	330,537	34
Male	11,392	16,515	21,413	27,262	37,928	54,345	69,278	71,310	99,298	25
Female	4,886	7,142	10,075	14,723	23,455	39,915	64,254	87,512	231,239	9
Non-Hispanic white	11,537	17,597	24,071	33,023	50,223	79,888	116,966	143,429	302,824	18
Male	8,494	12,706	16,870	21,981	31,816	47,037	61,605	65,165	90,818	13
Female	3,043	4,891	7,201	11,042	18,407	32,851	55,361	78,264	212,006	5
Non-Hispanic black	4,273	5,448	6,612	7,875	9,789	12,453	14,150	12,979	23,581	15
Male	2,585	3,399	4,001	4,608	5,329	6,238	6,332	4,997	6,691	11
Female	1,688	2,049	2,611	3,267	4,460	6,215	7,818	7,982	16,890	4
Origin not stated 2/	82	109	160	227	245	390	440	467	942	9
Male	62	76	116	174	160	259	251	209	307	7
Female	20	33	44	53	85	131	189	258	635	2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 50-54 55-59 60-64 65-69 75 - 7980-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS YFARS STATED Diseases of heart (I00-I09,I11,I13,I20-I51) All origins 47 14.096 20.898 27.605 36.562 52.206 77.047 104.841 121.311 251,860 Male 10.346 24.292 33.118 45.718 37 15.143 19.370 56.178 56.292 78,300 Female 3,750 5.755 8.235 12.270 19.088 31.329 48.663 65.019 173,560 10 Hispanic 716 6 1,090 1,331 1,824 2,464 3,093 3,618 3,428 6,968 Male 531 817 924 1,522 1,833 1,957 1,526 2,324 6 1,167 Female 185 273 407 657 942 1,260 1,661 1,902 4,644 2 Mexican 406 620 709 957 1,322 1,664 1,893 1,623 3,176 459 476 Male 301 609 785 983 1,077 759 1,140 Female 105 161 233 348 537 681 816 864 2,036 Puerto Rican 112 195 244 350 382 449 519 460 775 Male 79 149 170 221 222 251 251 196 250 Female 33 46 74 129 160 198 268 264 525 Cuban 57 72 107 174 303 375 544 656 1,576 1 Male 43 59 84 123 223 234 292 309 460 1 Female 14 13 23 51 80 141 252 347 1.116 Central and South American 79 100 139 163 213 246 244 274 513 Male 59 71 97 97 113 117 94 89 144 Female 20 29 42 66 100 129 150 185 369 Other and unknown Hispanic 103 132 180 244 359 415 928 62 418 3 179 Male 49 79 97 117 248 173 330 3 243 63 598

Source: CDC/NCHS, National Vital Statistics System, Mortality

13

24

35

65

111

175

242

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Diseases of heart (I00-I09,I11,I13,I20-I51) Non-Hispanic 1/ 13,314 19,715 26,140 34,544 49,540 73,621 100,854 117,505 244,151 32 Male 9,762 14,257 18,345 22,976 31,461 43,657 54,006 54,599 75,721 24 Female 3,552 5,458 7,795 11,568 18,079 29,964 46,848 62,906 168,430 8 Non-Hispanic white 9,762 15,061 20,346 27,604 40,913 62,837 88,727 106,460 224,193 18 Male 7,501 11,246 14,675 18,791 26,627 38,072 48,308 50,114 69,476 13 Female 2,261 3,815 5,671 8,813 14,286 24,765 40,419 56,346 154,717 5 Non-Hispanic black 5,202 14 3,239 4,209 6,144 7,622 9,397 10,398 9,410 17,056 Male 2,024 2,692 3,245 4,226 4,790 4,725 3,691 4,945 11 3,662 Female 1,215 1,517 1,957 2,482 3,396 4,607 5,673 5,719 12,111 3 Origin not stated 2/ 66 93 134 194 202 333 369 378 741 9 Male 53 69 101 149 135 228 215 167 255 7 Female 13 24 33 45 67 105 154 211 486 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Other and unknown Hispanic

Male

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hypertensive heart disease	e (I11)									
All origins	889	1,041	1,260	1,277	1,553	1,964	2,672	3,158	7,679	1
Male	586	673	800	760	858	958	1,185	1,067	1,594	1
Female	303	368	460	517	695	1,006	1,487	2,091	6,085	-
Hispanic	43	56	77	69	95	79	103	112	217	-
Male	34	35	51	45	52	43	44	38	50	-
Female	9	21	26	24	43	36	59	74	167	-
Mexican	20	31	32	26	32	37	52	49	118	-
Male	15	20	19	17	19	18	18	20	29	-
Female	5	11	13	9	13	19	34	29	89	-
Puerto Rican	7	12	15	15	23	13	13	14	32	-
Male	5	7	11	8	12	8	6	4	5	-
Female	2	5	4	7	11	5	7	10	27	-
Cuban	5	3	7	11	16	11	23	21	24	-
Male	5	1	6	8	10	7	13	6	8	-
Female	-	2	1	3	6	4	10	15	16	-
Central and South American	8	5	16	10	17	10	7	9	20	-
Male	7	3	10	7	7	6	2	2	2	-
Female	1	2	6	3	10	4	5	7	18	-
Other and unknown Hispanic	3	5	7	7	7	8	8	19	23	-
Male	2	4	5	5	4	4	5	6	6	-
Female	1	1	2	2	3	4	3	13	17	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80 - 84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	AND OVER	STATED
Hypertensive heart disease	e (I11)									
Non-Hispanic 1/	837	978	1,170	1,193	1,446	1,869	2,559	3,032	7,433	1
Male	545	635	742	703	798	904	1,135	1,022	1,539	1
Female	292	343	428	490	648	965	1,424	2,010	5,894	-
Non-Hispanic white	423	531	648	723	904	1,243	1,868	2,426	6,359	-
Male	303	373	436	446	504	618	835	812	1,281	-
Female	120	158	212	277	400	625	1,033	1,614	5,078	-
Non-Hispanic black	403	435	491	441	508	576	632	550	954	1
Male	235	252	287	241	274	262	270	188	224	1
Female	168	183	204	200	234	314	362	362	730	-
Origin not stated 2/	9	7	13	15	12	16	10	14	29	-
Male	7	3	7	12	8	11	6	7	5	-
Female	2	4	6	3	4	5	4	7	24	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hypertensive heart and rem	nal disease	(I13)								
All origins	80	76	116	146	201	269	487	600	1,209	-
Male	50	50	66	80	99	124	221	239	374	-
Female	30	26	50	66	102	145	266	361	835	-
Hispanic	8	3	9	10	17	15	20	19	33	-
Male	5	1	5	7	6	8	8	6	13	-
Female	3	2	4	3	11	7	12	13	20	-
Mexican	4	1	6	5	12	8	10	12	19	-
Male	3	-	3	3	4	4	5	4	7	-
Female	1	1	3	2	8	4	5	8	12	-
Puerto Rican	-	2	-	2	2	4	5	2	5	-
Male	-	1	-	2	1	3	-	1	3	-
Female	-	1	-	-	1	1	5	1	2	-
Cuban	3	-	1	1	-	1	-	2	2	-
Male	2	-	1	1	-	-	-	-	1	-
Female	1	-	-	-	-	1	-	2	1	-
Central and South American	-	-	2	1	-	1	1	-	3	-
Male	-	-	1	-	-	-	-	-	1	-
Female	-	-	1	1	-	1	1	-	2	-
Other and unknown Hispanic	1	-	-	1	3	1	4	3	4	-
Male	-	-	-	1	1	1	3	1	1	-
Female	1	-	-	-	2	-	1	2	3	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Hypertensive heart and renal disease (I13) Non-Hispanic 1/ 1,174 Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Ischemic heart diseases (120-125)									
All origins	9,524	15,251	20,575	27,802	40,080	58,760	79,379	89,917	179,376	39
Male	7,447	11,614	15,021	19,102	26,184	36,003	44,041	43,264	57,726	31
Female	2,077	3,637	5,554	8,700	13,896	22,757	35,338	46,653	121,650	8
Hispanic	502	826	1,018	1,445	1,971	2,521	2,937	2,718	5,525	5
Male	391	651	740	937	1,252	1,518	1,629	1,240	1,849	5
Female	111	175	278	508	719	1,003	1,308	1,478	3,676	-
Mexican	292	459	541	759	1,079	1,368	1,545	1,275	2,470	1
Male	230	358	382	494	662	817	902	608	904	1
Female	62	101	159	265	417	551	643	667	1,566	-
Puerto Rican	78	144	183	285	298	352	430	381	613	-
Male	58	115	129	175	178	200	216	164	195	-
Female	20	29	54	110	120	152	214	217	418	-
Cuban	30	59	87	138	226	320	444	527	1,355	1
Male	22	51	73	98	168	197	242	255	382	1
Female	8	8	14	40	58	123	202	272	973	-
Central and South American	55	77	102	129	164	203	201	212	414	-
Male	42	59	75	80	91	101	77	67	121	-
Female	13	18	27	49	73	102	124	145	293	-
Other and unknown Hispanic	47	87	105	134	204	278	317	323	673	3
Male	39	68	81	90	153	203	192	146	247	3
Female	8	19	24	44	51	75	125	177	426	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Ischemic heart diseases (I20-I25) Non-Hispanic 1/ 8,980 14,351 19,458 26,208 37,948 55,971 76,140 86,895 173,278 26 Male 7,021 10,907 14,202 18,047 24,820 34,297 42,234 41,893 55,678 20 Female 1,959 3,444 5,256 8,161 13,128 21,674 33,906 45,002 117,600 6 Non-Hispanic white 7,017 11,408 15,660 21,406 31,841 48,288 67,511 79,083 159,351 15 Male 5,680 8,905 11,699 15,055 21,341 30,212 38,098 38,665 51,267 11 Female 1,337 2,503 3,961 6,351 10,500 18,076 29,413 40,418 108,084 Non-Hispanic black 1,748 4,183 2,614 3,359 5,331 6,598 7,306 6,584 11,808 11 Male 1,160 1,744 2,173 2,564 2,996 3,441 3,380 2,611 3,435 9 Female 588 870 1,186 1,619 2,335 3,157 3,926 3,973 8,373 2 Origin not stated 2/ 42 74 99 149 161 268 302 304 573 8 Male 35 56 79 118 112 188 178 131 199 6 7 Female 18 20 31 49 80 124 173 374 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Acute myocardial infarction (I21-I22) 4,225 All origins 6.598 9.123 12.283 17.333 24.108 31.053 33.658 57,297 Male 3,261 6.654 11.154 14.554 16.991 15.952 19,357 5.039 8.375 Female 1.559 2.469 3,908 6,179 9.554 14.062 17,706 37,940 Hispanic 1,131 1,750 Male Female 1,158 Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Acute myocardial infarction (I21-I22) Non-Hispanic 1/ 3,996 8,730 11,679 16,535 23,079 29,839 32,627 55,400 4 6.268 Male 3,092 4,795 6,374 7,989 10,658 13,940 16,288 15,497 18,712 1 Female 904 1,473 2,356 3,690 5,877 9,139 13,551 17,130 36,688 3 7,279 14,202 Non-Hispanic white 3,229 5,175 9,837 20,137 26,593 29,696 50,606 2 Male 2,591 4,080 5,420 6,899 9,342 12,449 14,792 14,312 17,114 Female 638 1,095 1,859 2,938 4,860 7,688 11,801 15,384 33,492 Non-Hispanic black 1,262 1,600 2,011 2,531 2 685 968 2,751 2,505 4,109 Male 621 812 931 1,113 1,262 1,236 975 1,270 434 Female 251 347 450 669 898 1,269 1,515 1,530 2,839 Origin not stated 2/ 13 18 41 37 48 80 83 80 147 3 Male 10 12 32 27 26 56 48 38 53 1 Female 3 6 9 10 22 24 35 42 94 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Other acute ischemic heart diseases (I24) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Other acute ischemic heart diseases (I24) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS YFARS STATED Other forms of chronic ischemic heart disease (I20,I25) 5,207 11,243 All origins 8.493 15.280 22.453 34.272 47.920 55.810 121,171 Male 4.117 8.208 14.839 27.118 6.466 10.571 21.230 26.821 38.103 Female 1,090 2.027 3.035 4.709 7.614 13.042 21.099 28.692 83.068 Hispanic 1,216 1,765 1,569 1,796 3,764 Male 1,255 Female 2,509 Mexican 1,651 Male Female 1,061 Puerto Rican Male Female Cuban Male Female Central and South American

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

Male

Female

Female

Other and unknown Hispanic

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Other forms of chronic ischemic heart disease (I20,I25) Non-Hispanic 1/ 4,893 7,929 10,524 14,300 21,125 32,516 45,906 53,822 116,987 22 Male 3,861 6,006 7,672 9,907 13,974 20,139 25,724 26,204 36,703 19 Female 1,032 1,923 2,852 4,393 7,151 12,377 20,182 27,618 80,284 3 Non-Hispanic white 3,733 6,119 8,223 11,383 17,398 27,839 40,575 48,994 107,920 13 Male 3,048 4,743 6,159 8,036 11,842 17,581 23,107 24,186 33,909 11 Female 685 1,376 2,064 3,347 5,556 10,258 17,468 24,808 74,011 Non-Hispanic black 1,029 2,055 3,281 4,013 4,506 4,031 9 1,607 2,542 7,645 Male 700 1,100 1,328 1,603 1,857 2,149 2,123 1,616 2,148 8 Female 329 507 727 939 1,424 1,864 2,383 2,415 5,497 Origin not stated 2/ 29 56 57 111 112 187 218 223 420 5 Male 25 44 46 90 86 132 130 92 145 5 Female 4 12 11 21 26 55 88 131 275

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Atherosclerotic cardiovascular disease, so described (I25.0) 1,792 3,747 7,704 All origins 2.945 4.636 5.691 9.666 10.655 23,403 Male 1,438 2.314 2.842 3.842 4.704 5.200 4.763 3.281 6,905 Female 1.355 1.849 3.000 4,466 5,892 16.498 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS 50-54 55-59 60-64 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Atherosclerotic cardiovascular disease, so described (I25.0) Non-Hispanic 1/ 1,671 2,716 3,472 4,282 5,281 7,243 9,198 10,267 22,661 18 Male 1,339 2,118 2,635 3,028 3,538 4,384 4,924 4,580 6,654 16 Female 332 598 837 1,254 1,743 2,859 4,274 5,687 16,007 2 1,191 2,508 8,937 20,283 10 Non-Hispanic white 1,964 3,118 3,954 5,711 7,580 Male 984 1,574 1,960 2,248 2,715 3,512 4,093 4,004 5,928 8 Female 207 390 548 1,239 2,199 3,487 4,933 14,355 Non-Hispanic black 672 876 1,028 1,194 1,359 1,435 2,073 8 433 1,158 Male 479 604 674 746 772 717 493 601 8 313 Female 120 193 272 354 448 587 718 665 1,472 Origin not stated 2/ 11 31 28 61 46 78 84 64 73 3 Male 10 25 23 53 36 61 56 31 26 3 Female 1 6 5 8 10 17 28 33 47

CAUSE OF DEATH (BASED ON THE

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55 - 59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
All other forms of ch	ıronic ische	emic heart	disease (120,125.1-	125.9)					
All origins	3,415	5,548	7,496	10,644	16,762	26,568	38,254	45,155	97,768	8
Male	2,679	4,152	5,366	7,290	10,997	16,526	21,621	22,355	31,198	7
Female	736	1,396	2,130	3,354	5,765	10,042	16,633	22,800	66,570	1
Hispanic	175	310	415	576	852	1,186	1,412	1,441	3,095	2
Male	142	245	306	374	511	700	747	670	1,030	2
Female	33	65	109	202	341	486	665	771	2,065	-
Mexican	102	164	212	268	455	623	683	631	1,312	1
Male	81	128	150	175	267	362	388	313	468	1
Female	21	36	62	93	188	261	295	318	844	-
Puerto Rican	26	56	80	141	136	178	207	218	369	-
Male	20	45	63	82	73	96	89	95	123	-
Female	6	11	17	59	63	82	118	123	246	-
Cuban	17	33	48	64	115	170	266	297	816	-
Male	15	27	37	48	82	101	145	145	242	-
Female	2	6	11	16	33	69	121	152	574	-
Central and South American	16	29	35	54	73	90	106	132	253	-
Male	15	22	25	36	38	48	39	41	73	-
Female	1	7	10	18	35	42	67	91	180	-
Other and unknown Hispanic	14	28	40	49	73	125	150	163	345	1
Male	11	23	31	33	51	93	86	76	124	1
Female	3	5	9	16	22	32	64	87	221	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED All other forms of chronic ischemic heart disease (I20,I25.1-I25.9) Non-Hispanic 1/ 3,222 5,213 7,052 10,018 15,844 25,273 36,708 43,555 94,326 4 Male 2,522 3,888 5,037 6,879 10,436 15,755 20,800 21,624 30,049 3 Female 700 1,325 2,015 3,139 5,408 9,518 15,908 21,931 64,277 1 Non-Hispanic white 2,542 4,155 5,715 8,265 13,444 22,128 32,995 40,057 87,637 3 Male 2,064 3,169 4,199 5,788 9,127 14,069 19,014 20,182 27,981 3 Female 478 986 1,516 2,477 4,317 8,059 13,981 19,875 59,656 Non-Hispanic black 596 935 1,179 2,087 3,071 2,873 1,514 2,654 5,572 Male 387 621 724 929 1,111 1,377 1,406 1,123 1,547 Female 209 314 455 585 976 1,277 1,665 1,750 4,025 1 Origin not stated 2/ 18 25 29 50 66 109 134 159 347 2 Male 15 19 23 37 50 71 74 61 119 2

13

16

38

60

98

228

Source: CDC/NCHS, National Vital Statistics System, Mortality

3

6

6

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Other heart diseases (I26-I51) 5,465 All origins 3.502 4,386 7.115 10.034 15.538 21.689 27.023 62,848 Male 2.223 2.756 3.419 4.281 5.870 8.450 10.567 11.561 18.483 Female 1,279 1.630 2.046 2.834 4.164 7,088 11.122 15,462 44.365 Hispanic 1,180 Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Other heart diseases (I26-I51) Non-Hispanic 1/ 3,337 4,183 5,229 6,801 9,652 15,033 21,099 26,397 61,532 5 Male 2,118 2,619 3,279 4,091 5,650 8,162 10,263 11,292 18,021 3 Female 1,219 1,564 1,950 2,710 4,002 6,871 10,836 15,105 43,511 2 3,857 23,937 Non-Hispanic white 2,238 3,000 5,251 7,809 12,720 18,469 56,837 3 Male 1,482 1,913 2,469 3,211 4,641 7,015 9,074 10,291 16,511 Female 756 1,087 1,388 2,040 3,168 5,705 9,395 13,646 40,326 Non-Hispanic black 1,016 1,257 1,421 1,668 2,079 4,064 2 1,091 2,316 2,138 Male 589 739 806 909 1,027 1,016 854 1,234 Female 427 433 518 615 759 1,052 1,300 1,284 2,830 Origin not stated 2/ 12 11 21 28 26 46 54 60 136 1 Male 8 9 14 18 14 28 31 29 50 1 Female 4 2 7 10 12 18 23 31 86

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Acute and subacute endocarditis (I33) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Acute and subacute endocarditis (I33) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Diseases of pericardium and acute myocarditis (I30-I31,I40) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Diseases of pericardium and acute myocarditis (I30-I31,I40) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Heart failure (I50)										
All origins	298	484	853	1,254	2,240	4,284	6,926	9,845	28,327	1
Male	186	300	506	722	1,232	2,294	3,264	4,054	7,959	1
Female	112	184	347	532	1,008	1,990	3,662	5,791	20,368	-
Hispanic	10	19	32	55	71	131	167	186	538	-
Male	6	10	19	28	40	72	87	72	166	-
Female	4	9	13	27	31	59	80	114	372	-
Mexican	7	13	21	33	38	71	91	114	271	-
Male	3	6	12	17	20	40	51	49	84	-
Female	4	7	9	16	18	31	40	65	187	-
Puerto Rican	1	5	7	9	16	29	22	15	63	-
Male	1	4	5	5	8	14	8	6	21	-
Female	-	1	2	4	8	15	14	9	42	-
Cuban	2	-	-	1	3	5	12	16	55	-
Male	2	-	-	1	2	3	5	5	17	-
Female	-	-	-	-	1	2	7	11	38	-
Central and South American	-	-	3	3	7	4	13	15	38	-
Male	-	-	1	-	4	2	6	4	12	-
Female	-	-	2	3	3	2	7	11	26	-
Other and unknown Hispanic	-	1	1	9	7	22	29	26	111	-
Male	-	-	1	5	6	13	17	8	32	-
Female	_	1	-	4	1	9	12	18	79	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50 - 54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Heart failure (I50)										
Non-Hispanic 1/	286	464	818	1,189	2,164	4,144	6,740	9,637	27,718	1
Male	179	290	485	688	1,189	2,217	3,168	3,972	7,766	1
Female	107	174	333	501	975	1,927	3,572	5,665	19,952	-
Non-Hispanic white	173	307	594	943	1,751	3,551	5,924	8,778	25,738	-
Male	110	196	369	553	969	1,951	2,828	3,644	7,149	-
Female	63	111	225	390	782	1,600	3,096	5,134	18,589	-
Non-Hispanic black	109	148	212	234	380	542	724	744	1,724	1
Male	67	89	111	127	198	244	304	273	513	1
Female	42	59	101	107	182	298	420	471	1,211	-
Origin not stated 2/	2	1	3	10	5	9	19	22	71	-
Male	1	-	2	6	3	5	9	10	27	-
Female	1	1	1	4	2	4	10	12	44	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED All other forms of heart disease (I26-I28, I34-I38, I42-I49, I51) 3,077 4,456 All origins 3.781 5.721 7.622 11.040 14,520 16.972 34,317 Male 1.957 2.377 2.811 3.488 4.553 6.054 7.182 7.409 10.438 Female 1.120 1.404 1.645 2.233 3,069 4,986 7.338 9.563 23.879 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED All other forms of heart disease (I26-I28, I34-I38, I42-I49, I51) Non-Hispanic 1/ 2,936 3,603 4,263 5,477 7,325 10,687 14,129 16,561 33,614 4 Male 1,863 2,255 2,698 3,333 4,379 5,850 6,981 7,225 10,171 2 Female 1,073 1,348 1,565 2,144 2,946 4,837 7,148 9,336 23,443 2 Non-Hispanic white 1,988 2,615 3,160 4,200 5,928 9,002 12,339 14,979 30,915 3 Male 1,319 1,671 2,030 2,600 3,603 4,984 6,137 6,559 9,282 Female 669 1,130 1,600 2,325 4,018 6,202 8,420 21,633 Non-Hispanic black 873 909 1,006 1,257 1,506 1,572 1,381 1,162 2,328 Male 502 543 604 668 699 769 709 575 720 Female 371 366 402 494 558 737 863 806 1,608 1 Origin not stated 2/ 9 10 17 18 21 36 35 38 65 1 Male 6 9 11 12 11 23 22 19 23 1 Female 1 6 6 10 13 13 19 42

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Essential (primary) hypertension and hypertensive renal disease (I10,I12) All origins 1,146 1,653 2,400 2,924 6,317 Male 1.071 1,580 Female 1,404 1,853 4.737 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Essential (primary) hypertension and hypertensive renal disease (I10,I12) Non-Hispanic 1/ 299 435 506 726 1,071 1,562 2,287 2,786 6,110 Male 180 267 303 386 545 740 940 1,018 1,513 Female 119 168 203 340 526 822 1,347 1,768 4,597 275 411 1,105 2,230 Non-Hispanic white 143 200 683 1,710 5,145 Male 93 127 181 229 361 536 714 812 1,240 Female 50 73 94 182 322 569 996 1,418 3,905 Non-Hispanic black 144 226 221 285 342 422 528 478 842 Male 83 135 114 142 165 183 204 177 226 Female 91 107 143 177 239 301 616 Origin not stated 2/ 1 3 1 4 4 3 16 22 Male 2 1 2 1 10 6 Female 1 1 2 3 3 6 16

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Cerebrovascular diseases (I60-I69) All origins 2,455 3,108 4.152 5.500 8.694 15.398 25.007 32.420 67,080 Male 1.282 1.714 2.256 4.522 7.698 12.739 3.049 11.349 18,052 Female 1.173 1.394 1.896 2.451 4.172 7,700 13.658 19.681 49.028 Hispanic 1,536 Male Female 1,009 Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Cerebrovascular diseases (I60-I69) Non-Hispanic 1/ 2,213 2,858 3,848 5,143 8,145 14,669 24,125 31,518 65,404 2 Male 1,139 1,558 2,088 2,834 4,234 7,308 10,892 12,383 17,489 1 Female 1,074 1,300 1,760 2,309 3,911 7,361 13,233 19,135 47,915 1 Non-Hispanic white 1,302 1,841 2,649 3,662 6,342 11,990 20,866 28,312 59,818 Male 664 998 1,452 2,027 3,320 6,017 9,436 11,182 15,886 Female 638 843 1,197 1,635 3,022 5,973 11,430 17,130 43,932 Non-Hispanic black 882 1,017 1,529 4,644 780 1,252 2,263 2,715 2,599 Male 409 491 541 694 782 1,075 1,169 931 1,225 Female 371 391 476 558 747 1,188 1,546 1,668 3,419 Origin not stated 2/ 13 15 18 26 25 42 59 140 7 Male 7 11 18 15 22 34 21 36 Female 6 8 7 8 10 20 25 32 104

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Atherosclerosis (I70)										
All origins	56	129	172	314	485	991	1,770	2,615	8,390	-
Male	37	84	114	193	277	518	845	1,082	2,255	-
Female	19	45	58	121	208	473	925	1,533	6,135	-
Hispanic	5	9	11	14	15	39	60	70	184	-
Male	2	5	7	9	9	22	30	30	70	-
Female	3	4	4	5	6	17	30	40	114	-
Mexican	3	8	6	9	10	27	40	46	102	-
Male	2	4	4	6	6	13	20	22	42	-
Female	1	4	2	3	4	14	20	24	60	-
Puerto Rican	-	1	3	2	2	3	5	6	8	-
Male	-	1	2	2	-	2	2	1	3	-
Female	-	-	1	-	2	1	3	5	5	-
Cuban	2	-	1	-	2	2	6	6	26	-
Male	-	-	-	-	2	2	5	2	9	-
Female	2	-	1	-	-	-	1	4	17	-
Central and South American	-	-	-	1	-	3	1	4	10	-
Male	-	-	-	-	-	2	-	1	4	-
Female	-	-	-	1	-	1	1	3	6	-
Other and unknown Hispanic	-	-	1	2	1	4	8	8	38	-
Male	-	-	1	1	1	3	3	4	12	-
Female	-	-	-	1	-	1	5	4	26	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						
Atherosclerosis (I70)										
Non-Hispanic 1/	49	120	160	297	465	951	1,706	2,536	8,181	-
Male	33	79	106	181	266	495	814	1,047	2,178	-
Female	16	41	54	116	199	456	892	1,489	6,003	-
Non-Hispanic white	41	94	129	245	394	847	1,534	2,346	7,617	-
Male	28	65	83	152	227	440	724	966	2,014	-
Female	13	29	46	93	167	407	810	1,380	5,603	-
Non-Hispanic black	6	23	26	50	62	88	152	160	482	-
Male	4	11	19	27	34	46	78	64	140	-
Female	2	12	7	23	28	42	74	96	342	-
Origin not stated 2/	2	-	1	3	5	1	4	9	25	-
Male	2	-	1	3	2	1	1	5	7	-
Female	-	-	-	-	3	-	3	4	18	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Other diseases of circulatory system (I71-I78) All origins 2,244 1.333 3.564 4.662 4.602 6,863 Male 1.472 2.209 2.331 2.684 2.476 Female 1.355 1,978 2,271 4.387 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Other diseases of circulatory system (I71-I78) Non-Hispanic 1/ 403 529 834 1,275 2,162 3,457 4,560 4,477 6,691 Male 278 354 571 885 1,422 2,145 2,626 2,263 2,397 390 Female 125 175 263 740 1,312 1,934 2,214 4,294 289 1,891 4,081 Non-Hispanic white 401 672 1,101 3,109 4,129 6,051 Male 208 270 479 782 1,281 1,972 2,423 2,091 2,202 Female 81 131 193 319 610 1,137 1,706 1,990 3,849 Non-Hispanic black 104 108 146 144 283 332 557 234 357 Male 65 70 82 83 122 144 156 134 155 Female 39 38 64 61 112 139 201 198 402 Origin not stated 2/ 1 4 3 9 10 5 11 14 Male 1 3 6 7 1 6 3 Female 1 3 3 3 4 5 11

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Aortic aneurysm and dissection (I71) All origins 1,578 2,423 2,841 3.095 3,337 Male 1.114 1,598 1.905 1.596 1.495 Female 1.190 1,245 1.842 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Aortic aneurysm and dissection (I71) Non-Hispanic 1/ 273 338 543 835 1,528 2,365 3,036 2,775 3,276 Male 207 257 412 633 1,079 1,556 1,873 1,559 1,457 Female 66 81 131 202 449 809 1,163 1,216 1,819 1,373 2,175 2,814 2,593 Non-Hispanic white 184 256 450 745 3,037 Male 145 195 353 575 992 1,459 1,774 1,469 1,364 Female 39 61 97 170 381 716 1,040 1,124 1,673 Non-Hispanic black 71 70 126 145 139 189 81 164 Male 52 52 44 73 74 62 62 65 Female 24 19 30 26 71 102 77 124 Origin not stated 2/ 3 1 8 5 2 8 3 Male 1 1 5 5 1 3 1 Female 2 3 1 5 2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other diseases of arteries, arterioles and capillaries (I72-I78) All origins 1,141 1,761 1.567 3,526 Male Female 1,026 2.545 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other diseases of arteries, arterioles and capillaries (I72-I78) Non-Hispanic 1/ 1,092 1,524 1,702 3,415 Male 2,475 Female Non-Hispanic white 1,315 1,488 3,014 Male Female 2,176 Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55 - 59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other disorders of circulator	y system (]	180-199)								
All origins	202	176	208	237	312	380	498	619	972	1
Male	102	96	106	118	163	187	225	238	248	-
Female	100	80	102	119	149	193	273	381	724	1
Hispanic	20	14	9	15	15	27	8	18	35	-
Male	9	6	4	7	5	12	6	7	12	-
Female	11	8	5	8	10	15	2	11	23	-
Mexican	10	5	4	9	10	10	7	9	15	-
Male	2	1	3	4	4	3	6	3	6	-
Female	8	4	1	5	6	7	1	6	9	-
Puerto Rican	3	2	3	2	2	6	-	2	6	-
Male	3	1	-	1	-	3	-	1	4	-
Female	-	1	3	1	2	3	-	1	2	-
Cuban	1	3	-	1	-	6	1	2	4	-
Male	1	1	-	1	-	3	-	1	-	-
Female	-	2	-	-	-	3	1	1	4	-
Central and South American	2	2	1	2	2	2	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-
Female	1	1	1	2	1	1	-	-	-	-
Other and unknown Hispanic	4	2	1	1	1	3	-	5	10	-
Male	2	2	1	1	-	2	-	2	2	-
Female	2	-	-	-	1	1	-	3	8	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Other disorders of circulatory system (I80-I99) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Influenza and pneumonia (J10	J18)									
All origins	798	899	1,098	1,527	2,382	4,479	7,776	11,416	31,229	2
Male	500	559	611	851	1,394	2,499	4,020	5,438	10,676	1
Female	298	340	487	676	988	1,980	3,756	5,978	20,553	1
Hispanic	48	56	58	97	143	185	252	317	876	-
Male	32	39	32	48	82	105	129	155	334	-
Female	16	17	26	49	61	80	123	162	542	-
Mexican	27	33	24	52	81	95	115	148	371	-
Male	18	21	13	22	45	57	65	75	155	-
Female	9	12	11	30	36	38	50	73	216	-
Puerto Rican	10	9	18	18	25	36	51	48	110	-
Male	6	8	10	10	13	24	23	16	30	-
Female	4	1	8	8	12	12	28	32	80	-
Cuban	4	5	6	7	12	17	42	53	161	-
Male	3	4	3	4	7	7	18	23	62	-
Female	1	1	3	3	5	10	24	30	99	-
Central and South American	5	-	1	9	10	16	17	29	84	-
Male	3	-	-	4	6	7	10	17	29	-
Female	2	-	1	5	4	9	7	12	55	-
Other and unknown Hispanic	2	9	9	11	15	21	27	39	150	-
Male	2	6	6	8	11	10	13	24	58	-
Female	-	3	3	3	4	11	14	15	92	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Influenza and pneumonia (J10-J18) Non-Hispanic 1/ 741 836 1,029 1,417 2,223 4,276 7,494 11,059 30,247 1 Male 461 513 573 794 1,303 2,384 3,875 5,263 10,298 Female 280 323 456 623 920 1,892 3,619 5,796 19,949 1 797 27,838 Non-Hispanic white 503 606 1,097 1,843 3,638 6,686 10,099 1 Male 309 361 436 613 1,075 2,012 3,454 4,801 9,410 Female 194 245 361 484 768 1,626 3,232 5,298 18,428 Non-Hispanic black 220 205 208 278 327 1,963 553 659 751 Male 140 137 125 157 198 320 352 676 336 Female 80 68 83 121 129 233 323 399 1,287 Origin not stated 2/ 9 7 11 13 16 18 30 40 106 1 7 Male 7 6 9 9 10 16 20 44 1 Female 5 4 7 8 14 20 62

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Influenza (J10-J11)										
All origins	13	13	29	42	60	104	154	297	881	-
Male	8	7	13	24	27	47	71	131	238	-
Female	5	6	16	18	33	57	83	166	643	-
Hispanic	-	2	2	3	4	4	5	4	10	-
Male	-	2	2	2	4	3	1	1	1	-
Female	-	-	-	1	-	1	4	3	9	-
Mexican	-	2	2	1	3	2	2	2	5	-
Male	-	2	2	-	3	2	-	-	1	-
Female	-	-	-	1	-	-	2	2	4	-
Puerto Rican	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-
Central and South American	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	1	2	3	2	4	-
Male	-	-	-	-	1	1	1	1	-	-
Female	_	_	-	-	-	1	2	1	4	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Influenza (J10-J11)										
Non-Hispanic 1/	13	11	27	39	55	99	149	290	868	-
Male	8	5	11	22	23	43	70	130	237	-
Female	5	6	16	17	32	56	79	160	631	-
Non-Hispanic white	12	8	24	36	51	88	141	276	844	-
Male	7	4	11	21	20	39	69	123	233	-
Female	5	4	13	15	31	49	72	153	611	-
Non-Hispanic black	1	2	3	2	2	10	5	10	18	-
Male	1	1	-	1	2	4	1	5	3	-
Female	-	1	3	1	-	6	4	5	15	-
Origin not stated 2/	-	-	-	-	1	1	-	3	3	-
Male	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	3	3	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Pneumonia (J12-J18)										
All origins	785	886	1,069	1,485	2,322	4,375	7,622	11,119	30,348	2
Male	492	552	598	827	1,367	2,452	3,949	5,307	10,438	1
Female	293	334	471	658	955	1,923	3,673	5,812	19,910	1
Hispanic	48	54	56	94	139	181	247	313	866	-
Male	32	37	30	46	78	102	128	154	333	-
Female	16	17	26	48	61	79	119	159	533	-
Mexican	27	31	22	51	78	93	113	146	366	-
Male	18	19	11	22	42	55	65	75	154	-
Female	9	12	11	29	36	38	48	71	212	-
Puerto Rican	10	9	18	17	25	36	51	48	110	-
Male	6	8	10	9	13	24	23	16	30	-
Female	4	1	8	8	12	12	28	32	80	-
Cuban	4	5	6	7	12	17	42	53	160	-
Male	3	4	3	4	7	7	18	23	62	-
Female	1	1	3	3	5	10	24	30	98	-
Central and South American	5	-	1	8	10	16	17	29	84	-
Male	3	-	-	3	6	7	10	17	29	-
Female	2	-	1	5	4	9	7	12	55	-
Other and unknown Hispanic	2	9	9	11	14	19	24	37	146	-
Male	2	6	6	8	10	9	12	23	58	-
Female	-	3	3	3	4	10	12	14	88	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	AND OVER	STATED							
Pneumonia (J12-J18)										
Non-Hispanic 1/	728	825	1,002	1,378	2,168	4,177	7,345	10,769	29,379	1
Male	453	508	562	772	1,280	2,341	3,805	5,133	10,061	-
Female	275	317	440	606	888	1,836	3,540	5,636	19,318	1
Non-Hispanic white	491	598	773	1,061	1,792	3,550	6,545	9,823	26,994	1
Male	302	357	425	592	1,055	1,973	3,385	4,678	9,177	-
Female	189	241	348	469	737	1,577	3,160	5,145	17,817	1
Non-Hispanic black	219	203	205	276	325	543	654	741	1,945	-
Male	139	136	125	156	196	316	335	347	673	-
Female	80	67	80	120	129	227	319	394	1,272	-
Origin not stated 2/	9	7	11	13	15	17	30	37	103	1
Male	7	7	6	9	9	9	16	20	44	1
Female	2	-	5	4	6	8	14	17	59	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other acute lower respiratory	infections	(J20-J22))							
All origins	7	8	8	13	14	21	44	63	249	-
Male	5	3	4	6	6	11	22	29	48	-
Female	2	5	4	7	8	10	22	34	201	-
Hispanic	-	1	-	-	-	1	-	-	5	-
Male	-	-	-	-	-	1	-	-	1	-
Female	-	1	-	-	-	-	-	-	4	-
Mexican	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-
Puerto Rican	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	1	-	-	-	1	-	-	2	-
Male	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	2	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Other acute lower respiratory infections (J20-J22) 7 Non-Hispanic 1/ 7 8 13 14 20 44 63 244 3 Male 5 4 6 6 10 22 29 47 7 8 22 197 Female 4 4 10 34 6 232 Non-Hispanic white 6 4 11 11 15 41 58 Male 5 3 6 4 7 21 27 42 Female 1 8 20 31 190 Non-Hispanic black 3 3 5 10 Male 2 3 3 Female 3 2 7 Origin not stated 2/ Male Female

ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 01/07/24 PLATE 2 OF 2 PAGE 161

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Acute bronchitis and bronchiolitis (J20-J21) All origins 7 7 6 12 13 16 41 150 28 Male 5 2 3 5 6 9 15 19 31 7 Female 2 5 3 7 7 13 22 119 Hispanic 1 1 4 Male 1 1 3 Female 1 Mexican 1 Male Female Puerto Rican 1 Male 1 Female Cuban 1 1 1 Male 1 1 Female 1 Central and South American Male Female Other and unknown Hispanic 1 Male 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 YEARS YEARS POPULATION, AND SEX YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Acute bronchitis and bronchiolitis (J20-J21) 7 Non-Hispanic 1/ 6 6 12 13 15 28 41 146 2 5 Male 5 3 6 8 15 19 30 3 7 7 7 Female 4 13 22 116 3 4 10 10 10 25 140 Non-Hispanic white 6 38 Male 5 5 4 5 14 18 27 Female 1 5 6 5 11 20 113 Non-Hispanic black 3 3 5 5 Male 2 3 2 Female 3 2 3 Origin not stated 2/ Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 163 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YEARS YFARS YFARS AND OVER STATED Unspecified acute lower respiratory infection (J22) All origins 1 2 1 5 16 99 1 22 Male 1 1 1 2 7 10 17 9 Female 1 1 3 12 82 Hispanic 1 Male Female 1 Mexican Male Female Puerto Rican Male Female Cuban 1 Male Female 1 Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 YEARS POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Unspecified acute lower respiratory infection (J22) Non-Hispanic 1/ 1 2 1 5 16 22 98 1 7 Male 1 1 1 2 10 17 3 9 Female 1 1 12 81 1 2 5 92 Non-Hispanic white 1 1 16 20 Male 1 1 2 9 15 Female 1 3 9 11 77 Non-Hispanic black 5 Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Chronic lower respiratory diseases (J40-J47) All origins 1,079 12,441 2.031 4.179 7.118 20.203 24.978 23.657 26,833 Male 1,004 2.169 3.789 6.642 10.835 13.021 11.684 11,894 Female 1.027 2,010 3.329 5,799 9.368 11.957 11,973 14.939 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Chronic lower respiratory diseases (J40-J47) Non-Hispanic 1/ 1,023 1,955 4,075 6,943 12,145 19,757 24,425 23,153 25,995 3 Male 523 963 2,113 3,684 6,482 10,558 12,709 11,415 11,472 2 Female 500 992 1,962 3,259 5,663 9,199 11,716 11,738 14,523 1 3,555 22,770 Non-Hispanic white 747 1,625 6,231 11,099 18,301 21,892 24,521 Male 386 788 1,830 3,261 5,865 9,694 11,705 10,672 10,695 1 Female 361 837 1,725 2,970 5,234 8,607 11,065 11,220 13,826 Non-Hispanic black 300 472 899 1,191 1,351 1,013 256 613 1,166 Male 126 158 254 370 533 705 810 592 595 Female 130 142 218 243 366 486 541 421 571 Origin not stated 2/ 4 13 9 31 34 59 70 52 74 1 Male 3 10 6 20 24 40 34 24 34 1 Female 1 3 3 11 10 19 36 28 40

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45 - 49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Bronchitis, chronic and uns	pecified (J	I40-J42)								
All origins	14	24	35	62	63	110	139	200	446	-
Male	9	10	24	27	36	58	73	84	142	-
Female	5	14	11	35	27	52	66	116	304	-
Hispanic	-	-	-	2	2	2	4	9	17	-
Male	-	-	-	-	2	1	1	2	5	-
Female	-	-	-	2	-	1	3	7	12	-
Mexican	-	-	-	1	-	2	1	4	11	-
Male	-	-	-	-	-	1	-	2	3	-
Female	-	-	-	1	-	1	1	2	8	-
Puerto Rican	-	-	-	-	-	-	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-
Cuban	-	-	-	-	1	-	-	3	2	-
Male	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	3	1	-
Central and South American	-	-	-	-	-	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	1	1	-	2	-	2	-
Male	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	2	-	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Bronchitis, chronic and unspecified (J40-J42) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Emphysema (J43)										2
All origins	136	332	723	1,296	2,246	3,410	3,768	3,187	2,580	-
Male	85	183	394	691	1,230	1,906	2,027	1,650	1,258	-
Female	51	149	329	605	1,016	1,504	1,741	1,537	1,322	-
Hispanic	5	7	6	15	28	46	56	57	63	-
Male	4	4	3	13	18	33	34	35	36	-
Female	1	3	3	2	10	13	22	22	27	-
Mexican	2	3	2	6	9	19	26	32	34	-
Male	2	2	-	4	5	13	16	20	20	-
Female	-	1	2	2	4	6	10	12	14	-
Puerto Rican	1	-	1	5	4	4	5	3	3	-
Male	1	-	1	5	3	3	3	3	1	-
Female	-	-	-	-	1	1	2	-	2	-
Cuban	1	2	2	2	6	9	9	12	16	-
Male	1	1	2	2	5	6	4	6	10	-
Female	-	1	-	-	1	3	5	6	6	-
Central and South American	-	-	-	-	2	2	3	4	4	-
Male	-	-	-	-	1	2	3	3	2	-
Female	-	-	-	-	1	-	-	1	2	-
Other and unknown Hispanic	1	2	1	2	7	12	13	6	6	-
Male	-	1	-	2	4	9	8	3	3	-
Female	1	1	1	-	3	3	5	3	3	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Emphysema (J43)										
Non-Hispanic 1/	130	320	716	1,276	2,213	3,358	3,701	3,124	2,506	-
Male	80	175	390	673	1,208	1,868	1,986	1,612	1,217	-
Female	50	145	326	603	1,005	1,490	1,715	1,512	1,289	-
Non-Hispanic white	98	274	648	1,177	2,073	3,165	3,482	2,987	2,365	-
Male	59	140	344	609	1,115	1,728	1,855	1,516	1,138	-
Female	39	134	304	568	958	1,437	1,627	1,471	1,227	-
Non-Hispanic black	32	43	60	86	126	153	175	108	117	-
Male	21	33	38	56	87	108	103	75	62	-
Female	11	10	22	30	39	45	72	33	55	-
Origin not stated 2/	1	5	1	5	5	6	11	6	11	-
Male	1	4	1	5	4	5	7	3	5	-
Female	-	1	-	-	1	1	4	3	6	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Asthma (J45-J46)										
All origins	312	295	343	240	353	420	454	471	720	-
Male	96	101	119	88	115	152	144	129	152	-
Female	216	194	224	152	238	268	310	342	568	-
Hispanic	25	23	23	16	32	28	27	23	25	-
Male	9	11	9	3	10	9	7	5	6	-
Female	16	12	14	13	22	19	20	18	19	-
Mexican	10	6	10	4	8	8	10	5	10	-
Male	2	3	4	2	2	1	3	1	2	-
Female	8	3	6	2	6	7	7	4	8	-
Puerto Rican	11	8	9	5	13	6	6	9	9	-
Male	6	4	3	-	4	2	2	2	3	-
Female	5	4	6	5	9	4	4	7	6	-
Cuban	2	3	1	2	4	5	3	4	4	-
Male	1	2	-	1	2	1	-	1	-	-
Female	1	1	1	1	2	4	3	3	4	-
Central and South American	2	4	1	1	4	5	5	3	2	-
Male	-	1	1	-	1	3	1	1	1	-
Female	2	3	-	1	3	2	4	2	1	-
Other and unknown Hispanic	-	2	2	4	3	4	3	2	-	-
Male	-	1	1	-	1	2	1	-	-	-
Female	-	1	1	4	2	2	2	2	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Asthma (J45-J46)										
Non-Hispanic 1/	286	271	319	224	320	391	427	447	692	-
Male	86	90	110	85	104	142	137	123	145	-
Female	200	181	209	139	216	249	290	324	547	-
Non-Hispanic white	167	160	205	146	223	288	355	392	598	-
Male	45	52	72	52	67	97	108	110	110	-
Female	122	108	133	94	156	191	247	282	488	-
Non-Hispanic black	110	101	96	66	76	81	55	39	64	-
Male	38	34	30	27	26	35	20	7	16	-
Female	72	67	66	39	50	46	35	32	48	-
Origin not stated 2/	1	1	1	-	1	1	-	1	3	-
Male	1	-	-	-	1	1	-	1	1	-
Female	-	1	1	-	-	-	-	-	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other chronic lower respiratory diseases (J44,J47) All origins 3,078 9,779 1.380 5.520 16.263 20.617 19.799 23,087 Male 1.632 2.983 5.261 8.719 10.777 9.821 10,342 Female 1.446 2.537 4,518 7,544 9,840 9,978 12.745 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other chronic lower respiratory diseases (J44,J47) Non-Hispanic 1/ 593 3,006 5,383 9,552 15,900 20,162 19,391 22,369 3 1.340 Male 348 688 1,590 2,899 5,137 8,491 10,514 9,598 9,974 2 Female 245 652 1,416 2,484 4,415 7,409 9,648 9,793 12,395 1 2,671 Non-Hispanic white 473 1,176 4,853 8,748 14,753 18,814 18,339 21,153 Male 277 590 1,393 2,576 4,652 7,819 9,678 8,975 9,319 1 Female 196 586 1,278 2,277 4,096 6,934 9,136 9,364 11,834 Non-Hispanic black 148 457 693 1,109 971 110 313 946 852 Male 87 184 285 419 556 681 500 64 514 Female 46 61 129 172 274 390 428 352 457 Origin not stated 2/ 2 7 6 26 27 52 59 45 59 1 Male 1 6 4 15 18 34 27 20 27 1 Female 1 1 2 11 9 18 32 25 32

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Pneumoconioses and chemical e	ffects (J60	9-J66,J68)								
All origins	3	12	19	38	104	170	251	252	253	-
Male	3	12	19	38	99	170	243	243	243	-
Female	-	-	-	-	5	-	8	9	10	-
Hispanic	-	1	-	1	2	3	2	-	3	-
Male	-	1	-	1	2	3	2	-	1	-
Female	-	-	-	-	-	-	-	-	2	-
Mexican	-	1	-	1	1	1	2	-	2	-
Male	-	1	-	1	1	1	2	-	1	-
Female	-	-	-	-	-	-	-	-	1	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	1	2	-	-	1	-
Male	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 60-64 65-69 70-74 75-79 85 YEARS 50-54 55-59 80-84 YEARS POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Pneumoconioses and chemical effects (J60-J66,J68) Non-Hispanic 1/ 3 11 19 37 102 167 249 252 250 37 97 Male 3 11 19 167 241 243 242 8 Female 5 8 9 7 19 27 93 157 237 233 Non-Hispanic white 3 245 Male 3 19 27 89 157 229 236 226 Female 8 9 7 Non-Hispanic black 4 8 8 8 9 16 Male 4 8 7 8 9 15 Female 1 Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54 V54B5	55-59	60-64	65-69	70-74 V5AB5	75-79 YFARS	80-84 VEAR	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	AND OVER	STATED
Pneumonitis due to solids ar	nd liquids (J	169)								
All origins	129	155	259	327	582	1,172	2,068	2,962	7,294	-
Male	83	82	151	214	367	716	1,252	1,623	3,083	-
Female	46	73	108	113	215	456	816	1,339	4,211	-
Hispanic	4	4	8	18	21	36	57	61	186	-
Male	3	1	4	10	12	19	34	31	74	-
Female	1	3	4	8	9	17	23	30	112	-
Mexican	3	3	4	11	10	16	34	29	81	-
Male	3	-	1	8	7	7	24	13	34	-
Female	-	3	3	3	3	9	10	16	47	-
Puerto Rican	1	-	4	3	3	7	3	8	19	-
Male	-	-	3	2	-	5	1	5	6	-
Female	1	-	1	1	3	2	2	3	13	-
Cuban	-	1	-	1	2	4	8	14	49	-
Male	-	1	-	-	1	2	2	6	15	-
Female	-	-	-	1	1	2	6	8	34	-
Central and South American	-	-	-	1	3	2	3	2	7	-
Male	-	-	-	-	1	1	2	2	1	-
Female	-	-	-	1	2	1	1	-	6	-
Other and unknown Hispanic	-	-	-	2	3	7	9	8	30	-
Male	-	-	-	-	3	4	5	5	18	-
Female	-	-	-	2	-	3	4	3	12	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Pneumonitis due to solids and liquids (J69) Non-Hispanic 1/ 124 150 250 309 560 1,131 2,004 2,891 7,092 204 Male 79 80 146 354 692 1,214 1,589 3,000 105 Female 45 70 104 206 439 790 1,302 4,092 251 2,603 Non-Hispanic white 93 116 195 458 942 1,758 6,508 Male 62 60 113 163 286 574 1,063 1,429 2,767 Female 31 56 82 88 172 368 695 1,174 3,741 Non-Hispanic black 28 31 54 174 225 244 521 Male 18 30 38 59 109 135 126 193 Female 13 13 21 16 65 90 328 Origin not stated 2/ 1 1 1 1 5 7 10 16 Male 1 1 1 1 5 4 3 9 7 Female 3 7

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98) All origins 4,277 1.356 2.143 3.266 4.369 6,130 Male 1.199 1.806 2.255 2.073 2,175 Female 1,460 2.114 2,204 3.955 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98) Non-Hispanic 1/ 484 843 1,249 2,008 3,119 4,183 4,132 5,926 636 Male 246 343 431 686 1,128 1,729 2,157 1,996 2,096 Female 238 293 412 563 880 1,390 2,026 2,136 3,830 1,743 3,825 Non-Hispanic white 335 491 662 1,056 2,768 3,817 5,471 Male 171 272 330 589 999 1,562 1,988 1,860 1,926 Female 164 219 332 467 744 1,206 1,829 1,965 3,545 Non-Hispanic black 125 154 163 223 280 295 228 349 133 Male 70 62 81 83 108 129 139 96 123 Female 63 63 73 80 115 151 156 132 226 Origin not stated 2/ 2 3 8 3 4 9 12 12 13 7 7 Male 1 2 5 2 2 10 4 Female 1 1 3 1 2 2 5 2 9

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Peptic ulcer (K25-K28)										
All origins	113	142	186	236	314	522	665	835	1,439	2
Male	67	102	111	154	196	308	331	381	448	1
Female	46	40	75	82	118	214	334	454	991	1
Hispanic	7	8	14	11	16	21	20	24	38	-
Male	3	7	9	9	10	12	14	13	16	-
Female	4	1	5	2	6	9	6	11	22	-
Mexican	5	3	7	6	8	14	12	15	20	-
Male	2	2	4	4	6	8	8	10	10	-
Female	3	1	3	2	2	6	4	5	10	-
Puerto Rican	1	1	1	1	4	2	1	1	5	-
Male	1	1	1	1	2	1	1	1	3	-
Female	-	-	-	-	2	1	-	-	2	-
Cuban	-	-	-	1	-	2	2	4	5	-
Male	-	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	-	-	1	4	5	-
Central and South American	1	-	3	2	1	1	2	1	2	-
Male	-	-	3	2	-	1	2	-	-	-
Female	1	-	-	-	1	-	-	1	2	-
Other and unknown Hispanic	-	4	3	1	3	2	3	3	6	-
Male	-	4	1	1	2	-	2	2	3	-
Female	-	-	2	-	1	2	1	1	3	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Peptic ulcer (K25-K28)										
Non-Hispanic 1/	106	133	170	223	297	499	642	809	1,398	2
Male	64	94	101	143	186	295	315	367	432	1
Female	42	39	69	80	111	204	327	442	966	1
Non-Hispanic white	73	98	136	186	242	435	578	735	1,291	1
Male	42	69	78	117	150	259	287	335	399	1
Female	31	29	58	69	92	176	291	400	892	-
Non-Hispanic black	28	32	32	33	45	47	42	56	79	1
Male	20	23	22	24	29	26	18	21	24	-
Female	8	9	10	9	16	21	24	35	55	1
Origin not stated 2/	-	1	2	2	1	2	3	2	3	-
Male	-	1	1	2	-	1	2	1	-	-
Female	-	-	1	-	1	1	1	1	3	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
FOFULATION, AND SEX	TEARS	TEARS	AND OVER	STATED						
Diseases of appendix (K35-K38)	ı									
All origins	25	24	22	26	30	39	43	53	62	-
Male	18	19	17	17	18	20	28	19	35	-
Female	7	5	5	9	12	19	15	34	27	-
Hispanic	2	2	1	2	2	1	2	1	1	-
Male	1	2	1	1	2	-	1	-	1	-
Female	1	-	-	1	-	1	1	1	-	-
Mexican	-	2	1	1	1	-	2	1	-	-
Male	-	2	1	-	1	-	1	-	-	-
Female	-	-	-	1	-	-	1	1	-	-
Puerto Rican	1	-	-	1	-	-	-	-	1	-
Male	-	-	-	1	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	1	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Diseases of appendix (K35-K38)										
Non-Hispanic 1/	22	22	21	24	28	37	41	52	61	-
Male	16	17	16	16	16	19	27	19	34	-
Female	6	5	5	8	12	18	14	33	27	-
Non-Hispanic white	13	20	14	13	23	31	36	45	57	-
Male	9	15	11	11	13	15	23	15	32	-
Female	4	5	3	2	10	16	13	30	25	-
Non-Hispanic black	8	2	7	10	3	6	5	6	3	-
Male	7	2	5	4	2	4	4	4	2	-
Female	1	-	2	6	1	2	1	2	1	-
Origin not stated 2/	1	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	AND OVER	STATED							
Hernia (K40-K46)										
All origins	16	39	36	71	77	138	186	265	529	-
Male	12	20	21	39	34	59	75	117	157	-
Female	4	19	15	32	43	79	111	148	372	-
Hispanic	1	3	-	5	7	9	6	11	17	-
Male	1	2	-	4	1	5	2	4	3	-
Female	-	1	-	1	6	4	4	7	14	-
Mexican	-	2	-	4	6	4	5	7	10	-
Male	-	1	-	3	1	1	2	2	2	-
Female	-	1	-	1	5	3	3	5	8	-
Puerto Rican	-	-	-	-	-	1	-	2	1	-
Male	-	-	-	-	-	-	-	2	-	-
Female	-	-	-	-	-	1	-	-	1	-
Cuban	-	-	-	1	-	1	1	1	2	-
Male	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-
Central and South American	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-
Other and unknown Hispanic	1	1	-	-	1	2	-	1	3	-
Male	1	1	-	-	-	2	-	-	1	-
Female	-	-	-	-	1	-	-	1	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hernia (K40-K46)										
Non-Hispanic 1/	14	36	36	66	70	128	177	254	509	-
Male	10	18	21	35	33	53	72	113	153	-
Female	4	18	15	31	37	75	105	141	356	-
Non-Hispanic white	9	27	32	57	53	114	156	234	471	-
Male	7	15	21	32	28	51	66	105	135	-
Female	2	12	11	25	25	63	90	129	336	-
Non-Hispanic black	5	7	4	8	12	12	19	18	31	-
Male	3	3	-	2	5	2	5	8	14	-
Female	2	4	4	6	7	10	14	10	17	-
Origin not stated 2/	1	-	-	-	-	1	3	-	3	-
Male	1	-	-	-	-	1	1	-	1	-
Female	-	-	-	-	-	-	2	-	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Chronic liver disease and cirrhosis (K70,K73-K74) All origins 3,228 2,803 1,505 3,140 2.834 2.833 2.809 2.396 Male 2,446 2.383 2.051 1.892 1.837 1.617 1.208 Female 1.192 1.188 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Chronic liver disease and cirrhosis (K70,K73-K74) Non-Hispanic 1/ 2,791 2,756 2,467 2,511 2,514 2,530 2,201 1,405 888 2 Male 2,080 2,075 1,784 1,660 1,631 1,450 1,111 654 363 2 Female 711 681 683 851 883 1,080 1,090 751 525 2,170 2,013 2,104 2,191 2,291 2,027 1,320 Non-Hispanic white 2,260 821 2 Male 1,647 1,718 1,467 1,418 1,428 1,318 1,021 620 338 Female 523 542 546 686 763 973 1,006 700 483 Non-Hispanic black 507 398 363 328 250 182 52 128 65 Male 357 292 254 199 166 107 29 17 Female 150 106 109 129 75 61 36 35 Origin not stated 2/ 14 20 11 10 9 6 8 2 4 Male 12 15 10 8 8 5 4 1 Female 2 5 1 2 1 1 4 2 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Alcoholic liver disease (K70	9)									
All origins	1,965	1,837	1,511	1,348	1,104	831	533	256	118	2
Male	1,501	1,426	1,179	1,015	867	630	364	174	73	2
Female	464	411	332	333	237	201	169	82	45	-
Hispanic	270	229	196	160	124	89	51	18	16	-
Male	232	194	174	133	103	77	38	14	10	-
Female	38	35	22	27	21	12	13	4	6	-
Mexican	167	133	126	88	83	62	38	12	11	-
Male	145	118	113	72	70	55	27	10	6	-
Female	22	15	13	16	13	7	11	2	5	-
Puerto Rican	41	39	38	32	16	11	5	2	1	-
Male	35	32	32	25	16	10	5	1	1	-
Female	6	7	6	7	-	1	-	1	-	-
Cuban	6	5	2	9	3	1	1	1	2	-
Male	5	4	2	9	3	1	1	1	2	-
Female	1	1	-	-	-	-	-	-	-	-
Central and South American	26	11	16	8	8	7	3	2	-	-
Male	23	11	15	8	5	4	2	1	-	-
Female	3	-	1	-	3	3	1	1	-	-
Other and unknown Hispanic	30	41	14	23	14	8	4	1	2	-
Male	24	29	12	19	9	7	3	1	1	-
Female	6	12	2	4	5	1	1	-	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Alcoholic liver disease (K70) Non-Hispanic 1/ 1,686 1,597 1,309 1,182 Male 1,261 1,224 1,000 Female 1,335 1,065 Non-Hispanic white 1,314 Male 1,021 1,020 Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Other chronic liver disease and cirrhosis (K73-K74) All origins 1,263 1,292 1,978 1,249 1.303 1.486 1.729 1.863 Male Female 1,019 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Other chronic liver disease and cirrhosis (K73-K74) 1,792 Non-Hispanic 1/ 1,105 1,159 1,158 1,329 1,539 1,722 1,167 Male Female 1,124 1,640 1,105 Non-Hispanic white 1,362 1,606 Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Cholelithiasis and other disorders of gallbladder (K80-K82) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Cholelithiasis and other disorders of gallbladder (K80-K82) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) All origins 1,176 2,694 4,147 5,707 6,220 1.688 11,170 Male 1.366 2.986 3.045 4.428 2.138 Female 1.328 2,009 2,721 3.175 6.742 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) Non-Hispanic 1/ 557 811 1,073 1,548 2,514 3,937 5,459 6,015 10,797 Male 334 443 565 819 1,276 2,036 2,849 2,959 4,254 Female 223 368 508 729 1,238 1,901 2,610 3,056 6,543 991 1,738 5,064 Non-Hispanic white 272 445 645 2,980 4,440 9,268 Male 162 244 365 538 922 1,610 2,359 2,548 3,721 Female 110 201 280 453 816 1,370 2,081 2,516 5,547 Non-Hispanic black 346 378 501 698 856 911 1,351 269 839 Male 163 190 171 255 310 381 435 368 448 Female 106 207 246 388 475 476 471 903 Origin not stated 2/ 1 4 7 6 8 13 9 18 42 Male 1 4 4 5 6 8 4 7 18 Female 3 1 2 5 5 11 24

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04) All origins 6 5 5 11 10 27 37 25 34 Male 3 3 2 4 5 12 15 17 14 Female 3 2 3 7 5 15 22 8 20 Hispanic 1 2 3 1 1 1 Male 1 1 Female 1 3 1 1 2 Mexican 1 1 1 Male Female 2 1 Puerto Rican 1 1 Male Female 1 1 Cuban 1 Male Female 1 Central and South American 1 Male Female 1 Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 YEARS POPULATION, AND SEX YFARS YFARS YFARS YEARS YFARS AND OVER YFARS YFARS STATED Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04) Non-Hispanic 1/ 5 5 26 36 25 33 3 Male 3 3 5 12 15 17 13 2 2 Female 6 14 21 8 20 3 3 3 22 28 Non-Hispanic white 6 31 20 Male 1 4 10 13 13 9 Female 1 3 12 18 7 19 Non-Hispanic black 3 3 4 Male 1 2 2 3 Female 2 1 Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All origins	16	28	26	34	40	61	104	96	132	-
Male	10	15	13	19	20	25	58	49	60	-
Female	6	13	13	15	20	36	46	47	72	-
Hispanic	2	-	3	3	5	3	2	3	5	-
Male	1	-	2	2	4	-	1	-	2	-
Female	1	-	1	1	1	3	1	3	3	-
Mexican	-	-	1	1	3	3	1	1	2	-
Male	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	1	3	1	1	2	-
Puerto Rican	-	-	1	-	1	-	1	1	-	-
Male	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-
Cuban	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-
Central and South American	1	-	-	1	-	-	-	1	2	-
Male	-	-	-	1	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	1	1	-
Other and unknown Hispanic	1	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 50-54 55-59 60-64 65-69 75 - 7980-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N 05-N07,N26) Non-Hispanic 1/ 14 28 23 30 35 58 102 93 127 9 Male 15 11 16 16 25 57 49 58 Female 5 13 12 14 19 33 45 44 69 21 104 Non-Hispanic white 12 18 15 24 48 89 80 Male 8 10 12 13 20 51 42 48 Female 8 9 11 28 38 38 56 Non-Hispanic black 9 10 7 11 19 Male 5 3 3 6 8 Female 4 3 7 4 11 Origin not stated 2/ Male 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70 - 74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Renal failure (N17-N19)										
All origins	586	836	1,144	1,640	2,641	4,056	5,560	6,092	10,998	-
Male	352	462	612	880	1,340	2,100	2,912	2,976	4,351	-
Female	234	374	532	760	1,301	1,956	2,648	3,116	6,647	-
Hispanic	47	57	93	129	163	193	234	183	325	-
Male	29	34	56	76	80	94	131	78	153	-
Female	18	23	37	53	83	99	103	105	172	-
Mexican	25	41	57	82	77	106	125	98	164	-
Male	16	26	35	50	34	55	75	41	79	-
Female	9	15	22	32	43	51	50	57	85	-
Puerto Rican	11	10	16	21	38	28	25	22	35	-
Male	7	4	9	11	20	15	10	9	15	-
Female	4	6	7	10	18	13	15	13	20	-
Cuban	-	1	7	6	15	17	31	21	58	-
Male	-	1	4	5	9	8	22	10	27	-
Female	-	-	3	1	6	9	9	11	31	-
Central and South American	3	4	7	7	14	15	23	13	25	-
Male	2	2	5	4	9	7	7	5	10	-
Female	1	2	2	3	5	8	16	8	15	-
Other and unknown Hispanic	8	1	6	13	19	27	30	29	43	-
Male	4	1	3	6	8	9	17	13	22	-
Female	4	-	3	7	11	18	13	16	21	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Renal failure (N17-N19)										
Non-Hispanic 1/	538	775	1,044	1,506	2,470	3,850	5,317	5,891	10,631	-
Male	322	424	552	800	1,254	1,998	2,777	2,891	4,180	-
Female	216	351	492	706	1,216	1,852	2,540	3,000	6,451	-
Non-Hispanic white	257	422	627	965	1,707	2,907	4,317	4,962	9,132	-
Male	152	231	357	525	905	1,579	2,295	2,492	3,661	-
Female	105	191	270	440	802	1,328	2,022	2,470	5,471	-
Non-Hispanic black	266	334	370	490	687	846	896	824	1,326	-
Male	161	184	168	250	306	376	428	361	437	-
Female	105	150	202	240	381	470	468	463	889	-
Origin not stated 2/	1	4	7	5	8	13	9	18	42	-
Male	1	4	4	4	6	8	4	7	18	-
Female	-	-	3	1	2	5	5	11	24	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 203 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS YEARS AND OVER STATED Other disorders of kidney (N25,N27) All origins 3 1 3 3 3 6 7 6 Male 1 1 1 1 3 3 Female 2 1 3 2 2 5 3 Hispanic 1 2 Male 1 Female 1 Mexican 1 Male Female Puerto Rican 1 Male 1 Female Cuban 1 Male Female 1 Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other disorders of kidney (N25,N27)									
Non-Hispanic 1/	-	3	1	3	2	3	4	6	6	-
Male	-	1	-	-	1	1	-	2	3	-
Female	-	2	1	3	1	2	4	4	3	-
Non-Hispanic white	-	2	-	1	1	3	3	2	4	-
Male	-	1	-	-	-	1	-	1	3	-
Female	-	1	-	1	1	2	3	1	1	-
Non-Hispanic black	-	1	1	2	1	-	1	4	2	-
Male	-	-	-	-	1	-	-	1	-	-
Female	-	1	1	2	-	-	1	3	2	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49 VEADS	50-54 VFARS	55-59	60-64	65-69	70-74	75-79 X5486	80 - 84 VEARC	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	AND OVER	STATED
Infections of kidney (N10-N12,	N13.6,N15.	1)								
All origins	22	33	30	29	52	81	109	133	311	-
Male	8	11	14	13	19	25	37	46	81	-
Female	14	22	16	16	33	56	72	87	230	-
Hispanic	5	1	6	3	3	5	6	9	9	-
Male	2	-	3	2	2	2	2	4	2	-
Female	3	1	3	1	1	3	4	5	7	-
Mexican	3	-	2	2	2	3	6	4	7	-
Male	1	-	2	1	2	1	2	1	2	-
Female	2	-	-	1	-	2	4	3	5	-
Puerto Rican	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-
Cuban	-	-	1	-	-	1	-	3	-	-
Male	-	-	-	-	-	1	-	3	-	-
Female	-	-	1	-	-	-	-	-	-	-
Central and South American	1	1	2	1	-	1	-	-	1	-
Male	1	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	-	1	-	-	1	-
Other and unknown Hispanic	-	-	1	-	-	-	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	2	1	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Infections of kidney (N10-N12, N13.6, N15.1) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hyperplasia of prostate (N40)										
All origins	-	1	4	9	15	32	59	101	209	-
Male	-	1	4	9	15	32	59	101	209	-
Hispanic	-	-	1	-	2	1	-	4	3	-
Male	-	-	1	-	2	1	-	4	3	-
Mexican	-	-	-	-	2	-	-	4	2	-
Male	-	-	-	-	2	-	-	4	2	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-
Central and South American	-	-	1	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	1	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Hyperplasia of prostate (N40)										
Non-Hispanic 1/	-	-	3	9	13	30	59	97	206	-
Male	-	-	3	9	13	30	59	97	206	-
Non-Hispanic white	-	-	2	6	10	26	56	87	190	-
Male	-	-	2	6	10	26	56	87	190	-
Non-Hispanic black	-	-	1	3	3	3	3	10	15	-
Male	-	-	1	3	3	3	3	10	15	-
Origin not stated 2/	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	_	-	-

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 209 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT 50-54 55-59 POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Inflammatory diseases of female pelvic organs (N70-N76) All origins 2 6 7 4 11 15 17 18 Female 2 7 11 15 17 18 Hispanic 1 1 1 1 2 Female 1 1 1 2 Mexican 1 Female 1 Puerto Rican Female Cuban Female Central and South American Female Other and unknown Hispanic 1

1

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT 50-54 YEARS YEARS POPULATION, AND SEX YFARS YFARS YEARS YFARS YEARS YFARS AND OVER STATED Inflammatory diseases of female pelvic organs (N70-N76) Non-Hispanic 1/ 3 1 7 11 14 17 16 1 7 Female 3 4 11 14 17 16 Non-Hispanic white 2 9 1 6 8 16 15 Female 1 2 8 9 15 16 Non-Hispanic black 3 1 1 Female 3 1 1 Origin not stated 2/ Female

ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 01/07/24 PLATE 2 OF 2 PAGE 211

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Pregnancy, childbirth and the puerperium (000-099) All origins 5 2 . . . Female 5 2 Hispanic . Female Mexican . Female . Puerto Rican . Female . Cuban . Female Central and South American Female Other and unknown Hispanic Female

. . .

. . .

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 212 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 70-74 45-49 55-59 60-64 65-69 75-79 80-84 85 YEARS AGE NOT 50-54 POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS YEARS AND OVER STATED Pregnancy, childbirth and the puerperium (000-099) Non-Hispanic 1/ 5 2 2 Female 5 Non-Hispanic white . Female 4 . Non-Hispanic black 2 Female 2 Origin not stated 2/ Female .

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 213 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Pregnancy with abortive outcome (000-007) All origins Female Hispanic . Female Mexican . Female . Puerto Rican . Female . Cuban . Female Central and South American Female Other and unknown Hispanic

. . .

. . .

Source: CDC/NCHS, National Vital Statistics System, Mortality

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 214 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 70-74 45-49 50-54 55-59 60-64 65-69 75-79 80-84 85 YEARS AGE NOT POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS YEARS AND OVER STATED Pregnancy with abortive outcome (000-007) Non-Hispanic 1/ Female . Non-Hispanic white . Female . Non-Hispanic black . Female Origin not stated 2/ Female .

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 2 OF 2 PAGE 215

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE

Female

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other complications of pregr	nancy, chil	dbirth and	d the puer	perium (01	0-099)					
All origins	5	2							-	-
Female	5	2							-	-
Hispanic	-								-	-
Female	-								-	-
Mexican	-								-	-
Female	-								-	-
Puerto Rican	-								-	-
Female	-								-	-
Cuban	-								-	-
Female	-								-	-
Central and South American	-								-	-
Female	-								-	-
Other and unknown Hispanic	-								-	-

. . .

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999 PLATE 2 OF 2 PAGE 216

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES,

Origin not stated 2/

Female

1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other complications of preg	nancy, chil	dbirth and	I the puer	perium (01	0-099)					
Non-Hispanic 1/	5	2							-	-
Female	5	2	• • •						-	-
Non-Hispanic white	4								-	-
Female	4								-	-
Non-Hispanic black	-	2							-	-
Female	-	2							-	-

. . .

. . .

. . .

. . .

. . .

. . .

. . .

. . .

. . .

. . .

Source: CDC/NCHS, National Vital Statistics System, Mortality

. . .

. . .

. . .

. . .

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 217 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Certain conditions originating in the perinatal period (P00-P96) All origins 2 1 2 1 1 1 5 1 Male 2 1 1 1 1 Female 1 1 1 1 Hispanic 1 Male 1 Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic 1 Male 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 80-84 POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Certain conditions originating in the perinatal period (P00-P96) Non-Hispanic 1/ 2 2 1 1 3 1 1 1 Male 2 1 1 1 1 3 Female 1 1 1 2 1 Non-Hispanic white 1 1 1 1 1 Male 1 1 1 1 Female 1 1 Non-Hispanic black Male 2 Female Origin not stated 2/ 1 Male Female 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS YFARS STATED Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) All origins 1,168 1,140 1,558 1,945 2,468 7,798 Male 1,984 Female 1,487 5.814 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female

Source: CDC/NCHS, National Vital Statistics System, Mortality

Other and unknown Hispanic

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99) Non-Hispanic 1/ 1,087 1,090 1,511 1,896 2,414 7,658 Male 1,944 Female 1,454 5,714 1,288 2,141 Non-Hispanic white 1,634 7,033 Male 1,771 Female 1,296 5,262 Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins	5,877	6,506	6,676	7,714	10,272	14,893	20,657	24,941	58,599	10
Male	3,510	3,762	3,762	4,108	5,154	7,178	8,976	9,519	14,623	5
Female	2,367	2,744	2,914	3,606	5,118	7,715	11,681	15,422	43,976	5
Hispanic	437	400	395	447	513	620	627	638	1,482	2
Male	296	245	230	251	250	314	282	248	428	1
Female	141	155	165	196	263	306	345	390	1,054	1
Mexican	249	236	219	244	271	363	330	313	721	-
Male	168	142	127	141	134	179	149	121	225	-
Female	81	94	92	103	137	184	181	192	496	-
Puerto Rican	85	70	70	69	79	81	74	72	155	1
Male	63	49	46	38	40	46	32	28	49	-
Female	22	21	24	31	39	35	42	44	106	1
Cuban	12	12	15	43	48	45	82	111	277	1
Male	6	5	9	20	25	23	36	49	73	1
Female	6	7	6	23	23	22	46	62	204	-
Central and South American	37	30	36	43	45	45	45	41	76	-
Male	24	18	20	28	13	21	14	13	14	-
Female	13	12	16	15	32	24	31	28	62	-
Other and unknown Hispanic	54	52	55	48	70	86	96	101	253	-
Male	35	31	28	24	38	45	51	37	67	-
Female	19	21	27	24	32	41	45	64	186	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 50-54 55-59 60-64 65-69 75 - 7980-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0 -N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98) 5,397 6,070 6,247 9,721 14,235 19,977 24,255 56,968 5 Non-Hispanic 1/ 7.232 Male 3,186 3,490 3,509 3,832 4,882 6,841 8,672 9,252 14,153 3 Female 2,211 2,580 2,738 3,400 4,839 7,394 11,305 15,003 42,815 4,912 Non-Hispanic white 3,858 4,579 5,837 8,027 12,240 17,624 22,039 52,406 1 Male 2,344 2,685 2,786 3,120 4,054 5,923 7,621 8,392 12,830 1 Female 1,514 1,894 2,126 2,717 3,973 6,317 10,003 13,647 39,576 Non-Hispanic black 1,195 1,498 1,399 1,329 1,242 1,753 2,053 1,910 4,006 Male 765 646 624 725 793 909 721 1,097 2 Female 634 607 549 618 773 960 1,144 1,189 2,909 2 Origin not stated 2/ 43 36 34 35 38 38 53 48 149 3 Male 28 27 23 25 22 23 22 19 42 1

10

16

15

31

29

107

2

Source: CDC/NCHS, National Vital Statistics System, Mortality

15

9

11

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS YFARS STATED Accidents (unintentional injuries) (V01-X59, Y85-Y86) All origins 3,944 5,772 6.696 4,943 3.341 3.692 4.516 6.510 11,729 Male 4,996 2.753 2.336 2.656 3.276 4.480 3.606 2.262 3.096 Female 1,700 1.337 1.191 1.079 1.356 1.860 2.676 3,234 7.249 Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Accidents (unintentional injuries) (V01-X59, Y85-Y86) Non-Hispanic 1/ 6,099 3,652 3,101 3,430 4,308 5,567 6,329 11,432 26 4.523 Male 4,510 3,281 2,537 2,090 2,161 2,520 2,984 3,174 4,343 23 Female 1,589 1,242 1,115 1,011 1,269 1,788 2,583 3,155 7,089 3 4,789 2,968 2,875 Non-Hispanic white 3,603 2,487 3,693 4,946 5,811 10,647 13 Male 3,531 2,611 2,040 1,690 1,805 2,157 2,632 2,919 4,006 12 Female 1,258 992 928 797 1,070 1,536 2,314 2,892 6,641 Non-Hispanic black 1,138 765 562 485 454 489 499 646 12 426 Male 864 577 418 324 297 299 282 201 268 10 Female 274 188 144 161 157 190 217 225 378 2 Origin not stated 2/ 51 40 42 24 17 13 22 30 37 36 Male 43 34 33 21 13 10 12 20 17 31 Female 8 6 9 3 4 3 10 10 20 5

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Transport accidents (V01-V99, Y85) All origins 2,131 1,902 3.121 2,524 1.681 1.727 1.852 1.566 1,330 Male 2,234 1.782 1.454 1.107 1.091 1,071 1.089 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT	
POPULATION, AND SEX	YEARS	AND OVER	STATED								
Transport accidents (V01-V99,Y85)											
Non-Hispanic 1/	2,859	2,316	1,967	1,568	1,589	1,749	1,820	1,509	1,291	11	
Male	2,037	1,629	1,346	1,033	1,002	1,007	1,040	829	738	10	
Female	822	687	621	535	587	742	780	680	553	1	
Non-Hispanic white	2,307	1,904	1,643	1,303	1,361	1,536	1,636	1,384	1,180	4	
Male	1,644	1,348	1,127	859	863	882	929	757	668	4	
Female	663	556	516	444	498	654	707	627	512	-	
Non-Hispanic black	443	315	244	186	161	153	136	96	86	7	
Male	327	230	171	132	103	93	84	55	58	6	
Female	116	85	73	54	58	60	52	41	28	1	
Origin not stated 2/	16	9	16	8	6	7	6	13	3	15	
Male	13	7	14	7	4	5	3	10	1	13	
Female	3	2	2	1	2	2	3	3	2	2	

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All origins	2,747	2,225	1,885	1,485	1,565	1,711	1,772	1,471	1,261	30
Male	1,936	1,537	1,250	950	969	962	1,012	815	714	27
Female	811	688	635	535	596	749	760	656	547	3
Hispanic	224	181	141	97	119	85	72	41	33	10
Male	165	129	87	59	75	49	42	29	17	10
Female	59	52	54	38	44	36	30	12	16	-
Mexican	156	115	82	58	75	55	37	25	17	6
Male	120	87	53	39	46	34	16	18	8	6
Female	36	28	29	19	29	21	21	7	9	-
Puerto Rican	13	17	15	7	12	3	6	1	1	-
Male	8	10	10	7	8	1	5	1	-	-
Female	5	7	5	-	4	2	1	-	1	-
Cuban	7	9	13	8	14	9	11	7	4	-
Male	6	5	8	3	9	5	7	5	1	-
Female	1	4	5	5	5	4	4	2	3	-
Central and South American	29	15	14	14	7	6	10	-	4	1
Male	18	12	7	6	4	2	7	-	3	1
Female	11	3	7	8	3	4	3	-	1	-
Other and unknown Hispanic	19	25	17	10	11	12	8	8	7	3
Male	13	15	9	4	8	7	7	5	5	3
Female	6	10	8	6	3	5	1	3	2	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES. 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 50-54 55-59 60-64 65-69 75 - 7980-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2) 2,509 2,037 1,728 1,382 1,440 1,619 1,694 1,418 1,225 9 Non-Hispanic 1/ 777 Male 1,760 1,403 1,149 885 890 908 967 696 8 497 711 Female 749 634 579 550 727 641 529 1 1,434 1,233 1,422 Non-Hispanic white 2,011 1,667 1,145 1,527 1,305 1,122 3 Male 1,410 1,155 954 735 764 797 863 712 633 3 Female 601 512 480 410 469 625 664 593 489 Non-Hispanic black 403 282 222 165 148 143 88 79 6 126 Male 296 203 114 94 84 81 52 5 Female 107 79 69 51 59 45 37 27 1 Origin not stated 2/ 14 7 16 6 6 7 6 12 3 11 Male 11 5 14 6 4 5 3 9 1 9

2

2

3

3

2

2

Source: CDC/NCHS, National Vital Statistics System, Mortality

3

2

2

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

1332), HISI/MIC ORIGIN,										
RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75 - 79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	AND OVER	STATED						

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V8 1.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins	179	125	101	81	79	66	92	67	54	7
Male	135	99	77	58	52	45	51	37	35	7
Female	44	26	24	23	27	21	41	30	19	-
Hispanic	16	15	2	6	13	8	4	3	2	1
Male	13	14	2	6	10	7	4	2	1	1
Female	3	1	-	-	3	1	-	1	1	-
Mexican	11	12	1	4	5	4	4	-	1	1
Male	9	11	1	4	3	4	4	-	1	1
Female	2	1	-	-	2	-	-	-	-	-
Puerto Rican	1	-	1	-	2	-	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-
Cuban	-	1	-	2	4	2	-	2	-	-
Male	-	1	-	2	4	1	-	1	-	-
Female	-	-	-	-	-	1	-	1	-	-
Central and South American	4	2	-	-	-	2	-	1	1	-
Male	3	2	-	-	-	2	-	1	-	-
Female	1	-	-	-	-	-	-	-	1	-
Other and unknown Hispanic	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE

	TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED		
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-1.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)													
	Non-Hispanic 1/	162	109	99	73	66	58	88	63	52	2		
	Male	121	84	75	51	42	38	47	34	34	2		
	Female	41	25	24	22	24	20	41	29	18	-		
	Non-Hispanic white	130	89	88	63	53	49	75	56	46	1		
	Male	95	68	67	43	36	31	41	30	29	1		
	Female	35	21	21	20	17	18	34	26	17	-		
	Non-Hispanic black	23	17	11	6	8	6	8	3	5	1		
	Male	18	14	8	4	4	5	2	1	4	1		
	Female	5	3	3	2	4	1	6	2	1	-		
	Origin not stated 2/	1	1	-	2	-	-	-	1	-	4		
	Male	1	1	-	1	-	-	-	1	-	4		
	Female	-	-	-	1	-	-	-	-	-	-		

CAUSE OF DEATH (BASED ON THE

Male Female

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS	AGE NOT
POPULATION, AND SEX	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	YEARS	AND OVER	STATED
Water, air and space, and	other and	unspecified	transpor	t accident	s and the	ir sequela	e (V90-V99	, Y85)		
All origins	195	174	145	115	83	75	38	28	15	-
Male	163	146	127	99	70	64	26	18	9	-
Female	32	28	18	16	13	11	12	10	6	-
Hispanic	6	3	5	2	-	3	-	-	1	-
Male	6	3	5	2	-	3	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	3	2	3	2	-	-	-	-	-	-
Male	3	2	3	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	1	-	1	-	-	2	-	-	1	-
Male	1	-	1	-	-	2	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	1	1	-	-	-	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 85 YEARS AGE NOT 50-54 55-59 60-64 65-69 75 - 7980-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS STATED Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/

Source: CDC/NCHS, National Vital Statistics System, Mortality

Male

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
TOTOLATION, AND SEA	ILAKS	AND OVER	STATED							
Nontransport accidents (W00-	-X59,Y86)									
All origins	3,575	2,419	1,813	1,660	1,965	2,664	3,870	4,944	10,399	69
Male	2,762	1,824	1,299	1,155	1,245	1,585	2,007	2,406	3,722	60
Female	813	595	514	505	720	1,079	1,863	2,538	6,677	9
Hispanic	300	181	102	111	113	99	107	107	224	33
Male	259	145	89	84	77	67	54	51	101	29
Female	41	36	13	27	36	32	53	56	123	4
Mexican	170	86	63	60	66	54	55	48	105	9
Male	149	69	54	43	43	38	29	27	45	9
Female	21	17	9	17	23	16	26	21	60	-
Puerto Rican	51	35	16	20	14	6	6	12	24	-
Male	40	26	14	17	11	4	2	4	9	-
Female	11	9	2	3	3	2	4	8	15	-
Cuban	10	8	5	6	5	12	8	15	41	6
Male	9	6	5	6	5	7	5	4	19	3
Female	1	2	-	-	-	5	3	11	22	3
Central and South American	18	25	7	6	9	10	11	8	8	-
Male	17	22	6	4	5	6	2	4	1	-
Female	1	3	1	2	4	4	9	4	7	-
Other and unknown Hispanic	51	27	11	19	19	17	27	24	46	18
Male	44	22	10	14	13	12	16	12	27	17
Female	7	5	1	5	6	5	11	12	19	1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 65-69 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Nontransport accidents (W00-X59,Y86) Non-Hispanic 1/ 3,240 2,207 1,685 1,533 1,841 2,559 3,747 4,820 10,141 15 Male 2,473 1,652 1,191 1,057 1,159 1,513 1,944 2,345 3,605 13 Female 767 555 494 476 682 1,046 1,803 2,475 6,536 2 1,325 4,427 9 Non-Hispanic white 2,482 1,699 1,184 1,514 2,157 3,310 9,467 Male 1,887 1,263 913 831 942 1,275 1,703 2,162 3,338 8 Female 595 436 412 353 572 882 1,607 2,265 6,129 Non-Hispanic black 695 450 318 299 293 336 560 5 363 330 Male 537 347 247 192 194 206 198 146 210 4 Female 158 103 71 107 99 130 165 184 350 1 Origin not stated 2/ 35 31 26 16 11 6 16 17 34 21 Male 30 27 19 14 9 5 9 10 16 18 7 Female 5 4 7 2 2 1 7 18 3

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Falls (W00-W19)										
All origins	447	377	398	489	663	1,000	1,636	2,220	4,578	4
Male	366	295	296	354	433	594	840	1,110	1,701	4
Female	81	82	102	135	230	406	796	1,110	2,877	-
Hispanic	34	31	30	33	47	37	55	61	123	2
Male	28	28	28	27	28	24	29	31	56	2
Female	6	3	2	6	19	13	26	30	67	-
Mexican	20	16	18	20	30	19	28	29	51	-
Male	17	15	16	16	19	13	16	16	21	-
Female	3	1	2	4	11	6	12	13	30	-
Puerto Rican	4	4	4	7	2	2	3	7	15	-
Male	3	3	4	6	2	1	1	2	6	-
Female	1	1	-	1	-	1	2	5	9	-
Cuban	2	3	1	1	1	8	6	8	22	-
Male	2	3	1	1	1	5	4	3	11	-
Female	-	-	-	-	-	3	2	5	11	-
Central and South American	4	7	3	1	7	1	6	6	8	-
Male	4	6	3	1	3	1	1	4	1	-
Female	-	1	-	-	4	-	5	2	7	-
Other and unknown Hispanic	4	1	4	4	7	7	12	11	27	2
Male	2	1	4	3	3	4	7	6	17	2
Female	2	-	-	1	4	3	5	5	10	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,										
RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Falls (W00-W19)										
Non-Hispanic 1/	408	345	363	450	611	961	1,571	2,150	4,440	1
Male	333	266	264	323	400	569	805	1,073	1,639	1
Female	75	79	99	127	211	392	766	1,077	2,801	-
Non-Hispanic white	323	270	291	362	524	855	1,456	2,044	4,231	1
Male	265	209	209	262	339	507	739	1,020	1,549	1
Female	58	61	82	100	185	348	717	1,024	2,682	-
Non-Hispanic black	73	63	57	70	80	70	73	80	140	-
Male	59	48	43	47	57	43	41	36	49	-
Female	14	15	14	23	23	27	32	44	91	-
Origin not stated 2/	5	1	5	6	5	2	10	9	15	1
Male	5	1	4	4	5	1	6	6	6	1
Female	_	_	1	2	_	1	4	3	9	_

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Accidental discharge of f	irearms (W3	2-W34)								
All origins	51	41	25	23	15	18	15	8	10	-
Male	46	32	22	19	13	18	13	7	7	-
Female	5	9	3	4	2	-	2	1	3	-
Hispanic	4	-	1	-	2	-	-	-	-	-
Male	3	-	1	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Mexican	3	-	1	-	1	-	-	-	-	-
Male	2	-	1	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Puerto Rican	1	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 YEARS POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Accidental discharge of firearms (W32-W34) Non-Hispanic 1/ 47 41 24 23 13 18 15 8 10 7 Male 43 32 21 19 11 18 13 9 3 2 3 Female 4 4 2 20 20 17 13 10 Non-Hispanic white 41 36 11 Male 38 27 18 18 9 17 11 7 Female 3 9 2 3 Non-Hispanic black 5 4 3 2 1 Male 5 3 2 1 Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Accidental drowning and submersion (W65-W74) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Accidental drowning and submersion (W65-W74) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Accidental exposure to smoke, fire and flames (X00-X09) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 60-64 65-69 70-74 75-79 85 YEARS 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Accidental exposure to smoke, fire and flames (X00-X09) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Accidental poisoning and exposure to noxious substances (X40-X49) All origins 1,926 Male 1,456 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 65-69 70-74 75-79 85 YEARS 50-54 55-59 60-64 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Accidental poisoning and exposure to noxious substances (X40-X49) Non-Hispanic 1/ 1,715 1,275 Male Female 1,290 Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other and unspecified non	ntransport	accidents	and their	sequelae	(W20-W31,W	35-W64,W75-	W99,X10-X3	9,X50-X59	,Y86)	
All origins	755	741	662	670	852	1,180	1,735	2,290	5,355	15
Male	584	592	490	461	526	701	900	1,069	1,822	14
Female	171	149	172	209	326	479	835	1,221	3,533	1
Hispanic	49	33	27	42	41	49	35	37	94	3
Male	44	26	23	25	27	33	14	16	39	3
Female	5	7	4	17	14	16	21	21	55	-
Mexican	29	16	17	20	20	28	18	13	51	2
Male	27	13	15	10	10	18	7	8	21	2
Female	2	3	2	10	10	10	11	5	30	-
Puerto Rican	6	4	3	6	8	4	2	5	8	-
Male	6	3	2	5	6	3	1	2	2	-
Female	-	1	1	1	2	1	1	3	6	-
Cuban	2	2	3	4	3	3	2	6	17	-
Male	2	1	3	4	3	2	1	1	7	-
Female	-	1	-	-	-	1	1	5	10	-
Central and South American	5	8	1	3	2	7	3	2	-	-
Male	4	7	1	1	2	4	-	-	-	-
Female	1	1	-	2	-	3	3	2	-	-
Other and unknown Hispanic	7	3	3	9	8	7	10	11	18	1
Male	5	2	2	5	6	6	5	5	9	1

4

2

1

5

6

9

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

1 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 70-74 50-54 55-59 60-64 65-69 75 - 7980-84 85 YEARS POPULATION, AND SEX YFARS YFARS YFARS AND OVER YFARS YFARS YFARS YFARS YFARS STATED Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86) Non-Hispanic 1/ 1,128 1,695 2,248 5,245 Male 1,049 1,775 Female 1,199 3,470 1,470 2,034 Non-Hispanic white 4,876 Male 1,628 Female 1,082 3,248 Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Intentional self-harm (suicide) (X60-X84,Y87.0) All origins 2,817 2,264 1.678 1.218 1.187 1.284 1.280 Male 2.152 1.702 1.284 1,079 1.073 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Intentional self-harm (suicide) (X60-X84,Y87.0) Non-Hispanic 1/ 2,683 2,159 1,608 1,165 1,133 1,255 1,240 906 776 9 Male 2,038 1,625 1,225 927 923 1,055 1,039 768 661 9 Female 645 534 383 238 210 200 201 138 115 2,019 1,501 1,079 1,050 1,199 754 Non-Hispanic white 2,468 1,186 872 6 Male 1,876 1,512 1,141 860 862 1,007 997 743 644 6 Female 592 507 360 219 188 192 189 129 110 Non-Hispanic black 144 82 67 49 58 35 17 15 2 36 Male 114 69 54 41 47 31 30 12 2 Female 30 13 13 8 5 5 2 3 Origin not stated 2/ 13 16 5 6 8 2 5 3 3 5 Male 10 12 4 5 8 2 4 3 2 5 Female 3 4 1 1 1 1

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Intentional self-harm (suicide) by discharge of firearms (X72-X74) All origins 1,462 1,058 1.215 Male 1,221 1.030 Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 65-69 70-74 75-79 85 YEARS AGE NOT 50-54 55-59 60-64 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Intentional self-harm (suicide) by discharge of firearms (X72-X74) Non-Hispanic 1/ 1,395 1,178 1,028 Male 1,164 1,000 Female 1,295 Non-Hispanic white 1,118 Male 1,084 Female Non-Hispanic black Male Female Origin not stated 2/ Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Intentional self-harm (suic	ide) by otl	her and un	specified r	means and	their sequ	elae (X60-	X71,X75-X84	4,Y87.0)		
All origins	1,355	1,049	620	416	360	320	334	277	277	11
Male	931	672	402	259	230	206	198	171	185	11
Female	424	377	218	157	130	114	136	106	92	-
Hispanic	57	57	36	28	22	14	11	13	11	1
Male	49	39	29	20	19	9	6	10	7	1
Female	8	18	7	8	3	5	5	3	4	-
Mexican	26	26	10	7	7	5	3	2	2	-
Male	22	18	7	5	6	3	2	2	1	-
Female	4	8	3	2	1	2	1	-	1	-
Puerto Rican	10	14	5	6	2	1	2	3	2	-
Male	10	11	4	3	2	1	1	2	2	-
Female	-	3	1	3	-	-	1	1	-	-
Cuban	3	3	10	8	8	5	4	5	7	-
Male	3	2	9	8	8	5	3	3	4	-
Female	-	1	1	-	-	-	1	2	3	-
Central and South American	4	9	5	3	3	1	1	1	-	1
Male	3	5	5	1	2	-	-	1	-	1
Female	1	4	-	2	1	1	1	-	-	-
Other and unknown Hispanic	14	5	6	4	2	2	1	2	-	-
Male	11	3	4	3	1	-	-	2	-	-
Female	3	2	2	1	1	2	1	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC

CLASSIFICATION OF DISEAS 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	SES, 45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75 - 79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Intentional self-harm	(suicide) by o	ther and u	unspecified	means and	their seq	uelae (X60	-X71,X75-X	(84,Y87.0)		
Non-Hispanic 1/	1,288	981	580	387	333	306	322	261	265	6
Male	874	625	370	239	206	197	192	158	177	6
Female	414	356	210	148	127	109	130	103	88	-
Non-Hispanic white	1,173	901	531	346	292	285	297	247	255	3
Male	792	567	339	211	180	184	176	152	171	3
Female	381	334	192	135	112	101	121	95	84	-
Non-Hispanic black	68	40	21	14	22	7	7	2	3	2
Male	51	30	12	10	17	2	5	1	1	2
Female	17	10	9	4	5	5	2	1	2	-
Origin not stated 2/	10	11	4	1	5	-	1	3	1	4
Male	8	8	3	-	5	-	-	3	1	4
Female	2	3	1	1	-	-	1	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Assault (homicide) (X85-Y09,Y	87.1)									
All origins	998	670	420	302	259	214	185	119	99	49
Male	746	483	323	218	183	121	101	61	41	33
Female	252	187	97	84	76	93	84	58	58	16
Hispanic	133	64	47	22	20	13	6	5	3	5
Male	100	50	37	19	17	9	5	3	2	5
Female	33	14	10	3	3	4	1	2	1	-
Mexican	84	40	24	12	8	7	4	2	-	2
Male	64	32	21	10	7	5	3	1	-	2
Female	20	8	3	2	1	2	1	1	-	-
Puerto Rican	14	5	3	1	6	2	1	-	1	-
Male	11	4	3	1	5	1	1	-	-	-
Female	3	1	-	-	1	1	-	-	1	-
Cuban	5	1	7	5	3	2	-	1	-	-
Male	4	1	4	4	3	2	-	-	-	-
Female	1	-	3	1	-	-	-	1	-	-
Central and South American	20	5	9	1	2	1	1	-	-	-
Male	16	4	7	1	2	-	1	-	-	-
Female	4	1	2	-	-	1	-	-	-	-
Other and unknown Hispanic	10	13	4	3	1	1	-	2	2	3
Male	5	9	2	3	-	1	-	2	2	3
Female	5	4	2	_	1	_	_	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 50-54 55-59 60-64 65-69 70-74 75-79 85 YEARS 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Assault (homicide) (X85-Y09, Y87.1) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Assault (homicide) by discharge of firearms (X93-X95) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 55-59 60-64 65-69 70-74 75-79 85 YEARS 50-54 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Assault (homicide) by discharge of firearms (X93-X95) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 70-74 75-79 50-54 55-59 60-64 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS YFARS STATED Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE

Male

Male

Male

Female

Female

Female

Central and South American

Other and unknown Hispanic

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL

Female

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED	
TOTOLATION, AND SEA	TEARS	TEMICS	TEMICS	TEMICS	TEARS	TEARS	TEARS	TEARS	AND OVER	31/(120	
Assault (homicide) by other	and unspec	ified means	and their	sequelae	(X85-X92,	, X96-Y09 , Y8	7.1)				
Non-Hispanic 1/	389	287	186	121	141	112	121	82	73	12	
Male	275	200	135	77	99	58	65	45	30	6	
Female	114	87	51	44	42	54	56	37	43	6	
Non-Hispanic white	218	180	121	82	99	80	87	60	56	8	
Male	154	117	84	52	65	39	48	34	19	3	
Female	64	63	37	30	34	41	39	26	37	5	
Non-Hispanic black	148	89	56	34	39	25	29	17	16	4	
Male	106	69	44	21	32	16	15	8	10	3	
Female	42	20	12	13	7	9	14	9	6	1	
Origin not stated 2/	5	9	4	4	3	-	-	-	1	19	
Male	3	9	2	4	3	-	-	-	-	10	

1

9

2

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 261 RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Legal intervention (Y35,Y89.0)										
All origins	19	31	9	11	2	-	1	-	-	-
Male	19	30	9	11	2	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-
Hispanic	1	4	-	2	1	-	-	-	-	-
Male	1	4	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Mexican	1	3	-	1	1	-	-	-	-	-
Male	1	3	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Puerto Rican	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Cuban	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Other and unknown Hispanic	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-
Female	-	_	-	_	_	_	_	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Non-Hispanic 1/	18	27	9	9	1	-	1	-	-	-
Male	18	26	9	9	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-
Non-Hispanic white	14	21	7	6	1	-	1	-	-	-
Male	14	20	7	6	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-
Non-Hispanic black	4	6	2	3	-	-	-	-	-	-
Male	4	6	2	3	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Origin not stated 2/	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Events of undetermined intent (Y10-Y34, Y87.2, Y89.9) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC AGE NOT 45-49 60-64 65-69 70-74 75-79 85 YEARS 50-54 55-59 80-84 POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS STATED Events of undetermined intent (Y10-Y34, Y87.2, Y89.9) Non-Hispanic 1/ Male Female Non-Hispanic white Male Female Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Discharge of firearms, undetermined intent (Y22-Y24) All origins 18 10 6 8 9 6 6 7 4 Male 15 6 5 6 8 6 5 7 4 Female 3 4 1 2 1 1 Hispanic 2 2 2 Male 1 1 Female Mexican 1 Male Female Puerto Rican Male Female Cuban Male Female Central and South American 1 Male 1 Female Other and unknown Hispanic 1 1 Male 1 1

Source: CDC/NCHS, National Vital Statistics System, Mortality

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT 50-54 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Discharge of firearms, undetermined intent (Y22-Y24) Non-Hispanic 1/ 16 10 5 8 6 6 4 Male 14 6 4 6 6 6 5 7 4 Female 2 4 1 10 4 Non-Hispanic white 11 4 6 6 Male 9 6 6 4 Female 4 Non-Hispanic black 2 Male 2 Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE

TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Other and unspecified event	s of undete	ermined in	tent and t	heir seque	lae (Y10-Y	21,Y25-Y34	, Y87.2, Y89	.9)		
All origins	510	255	130	78	63	36	45	38	54	19
Male	342	169	89	48	39	19	26	17	21	13
Female	168	86	41	30	24	17	19	21	33	6
Hispanic	16	10	7	5	-	1	2	1	-	1
Male	14	8	6	4	-	-	2	-	-	1
Female	2	2	1	1	-	1	-	1	-	-
Mexican	9	3	4	4	-	1	2	1	-	-
Male	8	2	3	3	-	-	2	-	-	-
Female	1	1	1	1	-	1	-	1	-	-
Puerto Rican	3	4	-	1	-	-	-	-	-	-
Male	2	4	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-
Cuban	-	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-
Central and South American	2	2	1	-	-	-	-	-	-	-
Male	2	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-
Other and unknown Hispanic	2	1	1	-	-	-	-	-	-	1
Male	2	1	1	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-

70-74

YFARS

75-79

YFARS

80-84

YEARS

85 YEARS

AND OVER

AGE NOT

STATED

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX 45-49 55-59 60-64 50-54 65-69 YEARS YFARS YFARS YFARS YFARS Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 1/	491	240	121	73	61	33	43	37	52	10
Male	326	157	81	44	38	17	24	17	20	8
Female	165	83	40	29	23	16	19	20	32	2
Non-Hispanic white	379	178	98	63	53	28	39	35	46	6
Male	238	110	59	34	33	14	21	15	18	5
Female	141	68	39	29	20	14	18	20	28	1
Non-Hispanic black	103	59	19	8	7	3	4	-	3	4
Male	82	46	18	8	5	2	3	-	1	3
Female	21	13	1	-	2	1	1	-	2	1
Origin not stated 2/	3	5	2	-	2	2	-	-	2	8
Male	2	4	2	-	1	2	-	-	1	4
Female	1	1	-	-	1	-	-	-	1	4

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 269
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 85 YEARS 50-54 55-59 65-69 80-84 AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS YFARS YFARS AND OVER STATED Operations of war and their sequelae (Y36,Y89.1) All origins 2 1 2 6 4 4 3 Male 2 1 2 6 3 3 Female 1 Hispanic 1 Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic 1 Male 1 Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS AGE NOT 50-54 YEARS POPULATION, AND SEX YEARS YFARS YFARS YFARS YFARS YEARS YFARS AND OVER STATED Operations of war and their sequelae (Y36,Y89.1) Non-Hispanic 1/ 2 1 2 6 4 3 2 2 2 2 Male 1 6 4 2 Female 2 2 Non-Hispanic white 1 1 6 4 3 Male 2 1 6 Female 1 Non-Hispanic black Male Female Origin not stated 2/ Male Female

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC 45-49 60-64 70-74 75-79 50-54 55-59 65-69 80-84 85 YEARS AGE NOT POPULATION, AND SEX YFARS YFARS YFARS YFARS YFARS YFARS AND OVER YFARS YFARS STATED Complications of medical and surgical care (Y40-Y84,Y88) All origins Male Female Hispanic Male Female Mexican Male Female Puerto Rican Male Female Cuban Male Female Central and South American Male Female Other and unknown Hispanic Male Female

01/07/24 ORIG210A. DEATHS FROM 113 SELECTED CAUSES, BY 5-YEAR AGE GROUPS, SPECIFIED HISPANIC ORIGIN, PLATE 2 OF 2 PAGE 272
RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 1999

(RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER	AGE NOT STATED
Complications of medical and	surgical c	are (Y40-)	(84,Y88)							
Non-Hispanic 1/	86	116	131	167	220	350	394	404	515	-
Male	35	56	73	83	98	176	187	179	193	-
Female	51	60	58	84	122	174	207	225	322	-
Non-Hispanic white	64	84	91	133	185	293	337	351	441	-
Male	24	41	51	66	83	147	164	157	164	-
Female	40	43	40	67	102	146	173	194	277	-
Non-Hispanic black	20	31	34	27	31	53	52	42	63	-
Male	11	15	16	14	13	27	21	17	20	-
Female	9	16	18	13	18	26	31	25	43	-
Origin not stated 2/	-	1	1	-	1	-	-	1	2	-
Male	-	1	-	-	-	-	-	1	-	-
Female	-	-	1	-	1	-	-	-	2	-

⁻ QUANTITY ZERO.

^{...} CATEGORY NOT APPLICABLE.

^{1/} INCLUDES RACES OTHER THAN WHITE AND BLACK.

^{2/} INCLUDES DEATHS FOR WHICH HISPANIC ORIGIN WAS NOT REPORTED ON THE DEATH CERTIFICATE.